

ON MINIMAL RATIONAL ELLIPTIC SURFACES

ANTONIO LAFACE AND DAMIANO TESTA

ABSTRACT. We construct 13 projective \mathbb{Q} -factorial Fano toric varieties and show that for any minimal rational elliptic surface X there is one such toric variety Z_X and a divisor class $\delta_X \in \text{Cl}(Z_X)$ such that the number of (-1) -curves of X equals the dimension of the Riemann-Roch space of δ_X . As an application we give the number of (-1) -curves of any such elliptic fibration of Halphen index 2.

INTRODUCTION

Let X be a smooth projective rational surface which admits a morphism $\pi: X \rightarrow \mathbb{P}^1$ whose general fiber is a smooth curve of genus one. We say that π is *minimal* if it does not contract any (-1) -curve. Denote by K_X the canonical divisor class on X ; by [3, 12.1] the map π is induced by a complete linear system of the form $|-mK_X|$, with $m > 0$. As a consequence, the anticanonical divisor on X is semiample with $(-K_X)^2 = 0$. It is possible to show [4] that π is the unique elliptic fibration on X , that is π admits a unique multiple fiber and that this multiplicity is m . The number m is the *Halphen index* of X . It is well known that the Mori cone of X is finitely generated if and only if the Cox ring is finitely generated and these two conditions are equivalent to requiring X to have a finite number of (-2) -curves and (-1) -curves [1, 10]. While the former are exactly the prime components of the reducible fibers of π , thus easily enumerated, less is known about the number of (-1) -curves of X .

The aim of this note is to provide a combinatorial approach to determine the number of such curves. More precisely, we associate to any such surface X a \mathbb{Q} -factorial projective toric variety Z_X in the following way. Let F be the free abelian group with basis $\{f_C\}$ indexed by the prime components C of the reducible fibers of π . By the genus formula it follows that such components are exactly the (-2) -curves of X . The linear map

$$P_X: F \rightarrow \frac{K_X^\perp}{\langle K_X \rangle} = E_8 \cong \mathbb{Z}^8 \quad \text{determined by} \quad f_C \mapsto [C]$$

defines the 1-skeleton of the fan of a projective toric variety Z_X . It is possible to show that there is a unique complete toric variety Z_X with this skeleton (see Remark 1.4). Recall that, given $E = \text{Hom}(F, \mathbb{Z})$ the dual of F , the cokernel Q_X of the dual map P_X^* is the class map $Q_X: E \rightarrow \text{Cl}(Z_X)$. This allows us to define a

Date: September 20, 2021.

2000 Mathematics Subject Classification. Primary 14J26.

Key words and phrases. Rational surfaces, elliptic fibrations, toric varieties.

The first author was supported by Proyecto FONDECYT Regular, N. 1150732.

The second author was supported by EPSRC grant EP/K019279/1.

homomorphism

$$\alpha: \text{Pic}(X) \rightarrow \text{Cl}(Z_X)$$

by mapping any divisor class $[D]$ to the element $(\Gamma \mapsto D \cdot \Gamma)$ of E and then applying Q_X . In what follows we will denote by $=_{\text{free}}$ equality up to torsion. Our main result is the following theorem.

Theorem 1. *Let $\pi: X \rightarrow \mathbb{P}^1$ be a minimal elliptic fibration on a rational surface with Halphen index m and let Z_X be the associated toric variety.*

- (1) *All the (-1) -curves of X are mapped to the same class $\delta_X \in \text{Cl}(Z_X)$ by α . The set of (-1) -curves of X is in bijection with the set of integral points of the Riemann-Roch polyhedron*

$$\Delta(\delta_X) := Q_X^{-1}(\delta_X) \cap (E_{\mathbb{Q}})_{\geq 0}.$$

Moreover the coordinates of an integral point of the polyhedron $\Delta(\delta_X)$ are the intersection multiplicities of the corresponding (-1) -curve with the prime components of the fibers of π .

- (2) *For any reducible fiber F_i of π there exists an element $w_i \in \text{Cl}(Z_X)$ such that for any prime component C of F_i*

$$Q_X(C) =_{\text{free}} n_C w_i$$

where n_C is the positive integer which appear as a coefficient of the root C in the unique integer relation between the roots of the affine Dynkin diagram corresponding to F_i .

- (3) *Let F_1, \dots, F_r be the reducible fibers of π , let $w_1, \dots, w_r \in \text{Cl}(Z_X)$ be as in (2) and let F be the multiple fiber of π . Then the following holds*

$$\delta_X =_{\text{free}} \begin{cases} m(w_1 + \dots + w_r) & \text{if } F \text{ is not in } \{F_1, \dots, F_r\}, \\ m(w_1 + \dots + w_r) - (m-1)w_i & \text{if } F = F_i. \end{cases}$$

The paper is organised as follows. In Section 1 we prove Theorem 1. Section 2 is devoted to rational elliptic surfaces of Halphen index two and the enumeration of their (-1) -curves as an application of Theorem 1. Finally in the Appendix we list the degree matrices of the thirteen \mathbb{Q} -factorial Fano toric varieties related to root lattices and which appear in the theory exposed in this note.

1. (-1) -CURVES AND LATTICE POINTS

Let $\pi: X \rightarrow \mathbb{P}^1$ be a minimal elliptic fibration on a smooth projective rational surface. Denote by E_8 the lattice $K_X^\perp / \langle K_X \rangle$ and by $p: K_X^\perp \rightarrow E_8$ the canonical quotient map; as the notation suggests, the lattice E_8 is isometric to the negative definite E_8 -lattice. We let F be the free abelian group with basis indexed by the (-2) -curves on X and we let $\text{cl}_X: F \rightarrow K_X^\perp$ be the homomorphism assigning to each basis vector the corresponding (-2) -class; we denote by P_X the composition $p \circ \text{cl}_X$. The images of the rays of the standard basis of F under the homomorphism P_X define a one-dimensional fan in E_8 . We will show in Remark 1.4 that there exists a unique projective \mathbb{Q} -factorial toric variety Z_X whose one-dimensional cones are the cones of the fan defined by P_X . We summarise the situation in the following

commutative diagram with exact rows

$$(1.1) \quad \begin{array}{ccccccc} F & \xrightarrow{\text{cl}_X} & K_X^\perp & \xrightarrow{\tau} & A & \longrightarrow & 0 \\ \parallel & & \downarrow p & & \downarrow & & \\ F & \xrightarrow{P_X} & E_8 & \xrightarrow{\bar{\tau}} & A/\langle h \rangle & \longrightarrow & 0 \end{array}$$

where A is the cokernel of cl_X and $h := \tau(K_X)$.

Remark 1.1. The image of the homomorphism P_X is a root sublattice Λ of E_8 . According to [9, Thm. 3.2 and Thm. 3.3] there is a unique isometric embedding of Λ into E_8 modulo the action of the Weyl group of E_8 unless Λ is one of the following lattices: A_7 , $2A_3$, $A_5 + A_1$, $A_3 + 2A_1$, $4A_1$. In particular, if the rank of Λ is eight, then the embedding is unique.

Let $E = \text{Hom}(F, \mathbb{Z})$ be the dual of F and let $Q_X: E \rightarrow \text{Cl}(Z_X)$ be the cokernel of the dual map $P_X^*: E_8^* \rightarrow E$. Dualising the exact sequences in (1.1) we obtain the following commutative diagram with exact rows

$$(1.2) \quad \begin{array}{ccccccc} (K_X^\perp)^* & \xrightarrow{\text{cl}_X^*} & E & \longrightarrow & \frac{\text{Cl}(Z_X)}{\langle \delta_X \rangle} & \longrightarrow & 0 \\ \uparrow p^* & & \parallel & & \uparrow \tau & & \\ (E_8)^* & \xrightarrow{P_X^*} & E & \xrightarrow{Q_X} & \text{Cl}(Z_X) & \longrightarrow & 0 \end{array}$$

where $\delta_X \in \text{Cl}(Z_X)$ generates the kernel of τ and is defined by

$$\delta_X = (Q_X \circ \text{cl}_X^*)(m),$$

where $m \in (K_X^\perp)^*$ generates the cokernel of p^* , that is m is any projection of the homomorphism $i: \mathbb{Z} \rightarrow K_X^\perp$ defined by $1 \mapsto -K_X$. Observe that δ_X does not depend on the choice of m .

Given a class $w \in \text{Cl}(Z_X)$ we recall that the *Riemann-Roch polytope* of w is the convex polytope

$$\Delta(w) := Q_Z^{-1}(w) \cap (E_{\mathbb{Q}})_{\geq 0},$$

where $(E_{\mathbb{Q}})_{\geq 0}$ is the positive orthant of the rational vector space $E_{\mathbb{Q}} = E \otimes_{\mathbb{Z}} \mathbb{Q}$. Assume now that D is a (-1) -curve of X and denote by e_D the linear function $F \rightarrow \mathbb{Z}$ defined by $f \mapsto \text{cl}_X(f) \cdot D$, where the product is induced by the intersection pairing of X . We therefore obtain a function

$$\varepsilon: \{(-1)\text{-curves of } X\} \longrightarrow E \quad D \longmapsto e_D.$$

The following lemma is needed in the proof of Lemma 1.3 (see also [6, Proposition 3.3] for another proof).

Lemma 1.2. *Let $\pi: X \rightarrow \mathbb{P}^1$ be a minimal elliptic fibration on a rational surface and let D be a divisor on X such that $D^2 = D \cdot K_X = -1$ and $D \cdot C \geq 0$ for any (-2) -curve of X . Then D is linearly equivalent to a (-1) -curve.*

Proof. Since the elliptic fibration π is minimal, the anticanonical divisor $-K_X$ is linearly equivalent to a multiple of a fiber and thus it is nef. By the Riemann-Roch theorem either D or $K_X - D$ is linearly equivalent to an effective divisor. The equality $-K_X \cdot (K_X - D) = -1$ and the fact that the divisor $-K_X$ is nef allow us

to conclude that the first possibility occurs. Thus, without loss of generality, we can assume D to be effective.

By the genus formula and the fact that $-K_X$ is nef, the only curves of X which are orthogonal to $-K_X$ are (-2) -curves, while the only ones which have intersection 1 with $-K_X$ are the (-1) -curves. Thus the equality $-K_X \cdot D = 1$ implies $D = \Gamma + D_1$, where Γ is a (-1) -curve and D_1 is a non-negative sum of (-2) -curves. From $(\Gamma + D_1)^2 = D^2 = -1$ we deduce the following

$$0 \leq (D_1 + \Gamma) \cdot D_1 = -\Gamma \cdot D_1 \leq 0,$$

where the first inequality is due to the hypothesis on $D = (D_1 + \Gamma)$ and the definition of D_1 , while the second inequality is due to the fact that Γ is not a component of D_1 . Thus $(D_1)^2 = \Gamma \cdot D_1 = 0$ and, since $-K_X \cdot D_1 = 0$, by the Hodge index theorem we conclude that D_1 is linearly equivalent to $-\alpha K_X$ for some rational number α . Using the identities $-1 = D^2 = (\Gamma - \alpha K_X)^2 = -1 + 2\alpha$ we deduce that $\alpha = 0$ and thus $D = \Gamma$ is a (-1) -curve. \square

Lemma 1.3. *The function ε induces a bijection between the set of (-1) -curves on X and the set of integral points of the polytope $\Delta(\delta_X)$.*

Proof. We begin by showing that the image of ε is contained in $\Delta(\delta_X)$. Indeed, if D is a (-1) -curve of X the homomorphism $e_D: F \rightarrow \mathbb{Z}$ factors through K_X^\perp by definition and it defines a projection $m_D: K_X^\perp \rightarrow \mathbb{Z}$ of $i: \mathbb{Z} \rightarrow K_X^\perp$ since $-K_X \cdot D = 1$. Thus $Q_X(e_D) = \delta_X$ by our definition of δ_X . Moreover, for each element f of the standard basis of F , we have

$$e_D(f) = \text{cl}_X(f) \cdot C \geq 0,$$

where the last inequality is due to the fact that $\text{cl}_X(f)$ and D correspond to distinct irreducible curves on X . We deduce that the image of ε is contained in $\Delta(\delta_X)$.

Now, we show that ε is injective. Let D, D' be (-1) -curves on X and suppose that $e_D = e_{D'}$ holds. It follows that the difference $D - D'$ is proportional to K_X . We deduce that $0 = (D - D')^2 = -2 - 2D \cdot D'$, so that D and D' coincide.

Finally, let e be an integral point of $\Delta(\delta_X)$. Therefore, we write $e = e_D + P_X^*(\gamma)$, where D is a (-1) -curve on X and γ is an element of E_8^* . We deduce that e factors through cl_X inducing a retraction $m_e: K_X^\perp \rightarrow \mathbb{Z}$ of i . Using the perfect pairing

$$K_X^\perp \times \frac{\text{Pic}(X)}{\langle -K_X \rangle} \longrightarrow \mathbb{Z},$$

we see that m_e is multiplication by an element $[D] + \mathbb{Z}(-K_X)$, where $D \cdot (-K_X) = 1$ and for each (-2) -curve Γ on X the inequality $D \cdot \Gamma \geq 0$ holds. Let r be an integer such that $(D + rK_X)^2 = D^2 - 2r = -1$; such an integer r exists since D^2 is odd by the adjunction formula. By Lemma 1.2, the class of $D + rK_X$ is the class of a (-1) -curve and we are done. \square

We now make explicit the homomorphism $Q: E \rightarrow \text{Cl}(Z)$. We begin with a sublattice Λ of E_8 spanned by roots of E_8 . Let $\Lambda = \Lambda_1 \oplus \dots \oplus \Lambda_r$ be a decomposition of Λ as a direct sum of irreducible root lattices. We fix a subset $\{\lambda_1, \dots, \lambda_s\}$ of the lattice Λ consisting of simple roots chosen as follows. First of all we choose a basis of Λ_1 indecomposable roots together with the root corresponding to the affine vertex in the extended Dynkin diagram of Λ_1 , equivalently, the last root is the opposite of the longest root of Λ_1 . Then we proceed similarly with Λ_2 and so

on. Let F be the free abelian group of rank s with basis f_1, \dots, f_s . Let $P: F \rightarrow E_8$ be the homomorphism defined by $f_1 \mapsto \lambda_1, \dots, f_s \mapsto \lambda_s$ and let $E = \text{Hom}(F, \mathbb{Z})$ be the dual lattice of F . The homomorphism P defines a toric variety Z and an exact sequence

$$E_8^* \xrightarrow{P^*} E \xrightarrow{Q} \text{Cl}(Z) \longrightarrow 0.$$

as in (1.2). Let $\pi: X \rightarrow \mathbb{P}^1$ be a minimal rational elliptic surface with singular fibers of type $\tilde{\Lambda}_1, \dots, \tilde{\Lambda}_r$. We denote the (-2) -curves of X by C_1, \dots, C_s where the numbering is compatible with the numbering of the roots $\lambda_1, \dots, \lambda_s$ given above. It follows from the definitions that the homomorphisms P and P_X coincide, and thus that also Q and Q_X coincide. With the above notation the matrix Q has the following form

$$(1.3) \quad Q = \begin{bmatrix} Q_1 & 0 & \cdots & 0 \\ 0 & Q_2 & \cdots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & Q_r \\ \hline & & & \text{torsion} \end{bmatrix}$$

where each Q_i is the row vector of length $\text{rk } \Lambda_i + 1$ defined by the unique relation between the simple roots of the chosen basis of Λ_i together with the root corresponding to its affine vertex. In these coordinates the matrix Q defines a map $E \rightarrow \mathbb{Z}^r \oplus (E_8/\Lambda)$.

Proof of Theorem 1. The first part of (1) follows from Lemma 1.3. For the second part observe that a (-1) -curve D gives an element $e_D \in E$ by taking intersection with D . Thus the statement follows from the fact that the canonical basis of F consists of the prime components of the fibers of π .

To prove items (2) and (3) we begin by recalling that the class $\delta_X \in \text{Cl}(Z_X)$ is defined by any projection $m: K_X^\perp \rightarrow \mathbb{Z}$ of the inclusion map $i: \mathbb{Z} \rightarrow K_X^\perp$ given by $1 \mapsto -K_X$. In particular such a projection is obtained by intersecting with a (-1) -curve D of X , that is $m(C) = C \cdot D$ for any $C \in K_X^\perp$. Due to the above presentation of the matrix Q the integers in the row corresponding to the vector Q_i are exactly the multiplicities of the unique primitive vector of self-intersection zero in the root lattice $\tilde{\Lambda}_i \subset K_X^\perp$ whose class is thus the class of $-mK_X$ whenever the corresponding (-2) -curves form a fiber of π or it is $-K_X$ whenever these curves form the support of the non-reduced fiber. Thus items (2) and (3) follow. \square

Remark 1.4. Due to (1.3) the free part of the grading of any variable of the Cox ring of the toric variety Z is one of the canonical vectors w_1, \dots, w_r of \mathbb{Z}^r and any such vector occurs as a degree of at least two variables. As a consequence, by [2, Proposition 2.4.2.6], there is a unique projective variety Z with the given Cox ring and such that

$$\text{SAmple}(Z) = \text{Mov}(Z) = \text{Eff}(Z) = \text{cone}(w_1, \dots, w_r).$$

It follows that any class in the relative interior of the above cone defines the same \mathbb{Q} -factorial projective toric variety Z . This is the case for the anticanonical class $-K_Z$ so that the variety Z is Fano.

Remark 1.5. Recall [8, Lemma 8.16] that if Z is a toric variety with Cox ring $\mathcal{R}(Z)$ and grading matrix $Q := [w_1, \dots, w_n]$, then the multigraded Hilbert series of $\mathcal{R}(Z)$ is

$$H(\mathcal{R}(Z), t) = \frac{1}{(1 - t^{w_1}) \cdots (1 - t^{w_n})},$$

where t^w is the character of the quasi-torus $\text{Spec}(\mathbb{C}[\text{Cl}(Z)])$ corresponding to $w \in \text{Cl}(Z)$. Hence Table 2.3 and Theorem 1 allow us to determine the number of (-1) -curves on any such surface.

Example 1.6. Let X be a minimal rational elliptic surface of type D_8 of Halphen index 2 and whose multiple fiber is irreducible. According to the table appearing in the Appendix, the Hilbert series of the Cox ring of the corresponding toric variety Z is

$$\frac{1}{(1-t)^2(1-tu)^2(1-t^2)^3(1-t^2u)^2}$$

where the two variables are in $\mathbb{C}[t, u]/\langle u^2 - 1 \rangle$. Expanding the above series, a direct calculation shows that the coefficient of the term t^2 is 9, while that of t^2u is 6. Both cases are realized as shown in the next paragraphs.

Case 1. Choose nine points p_1, \dots, p_9 on a smooth plane cubic curve C with flex at p_1 such that the classes of $p_2 - p_1, \dots, p_9 - p_1$ are the following elements of $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$: 00, 00, 01, 01, 01, 10, 10, 11, 11. For example we can take $C := V(x^3 - y^2z - xz^2)$, $p_1 = (0, 1, 0)$, $p_3 = (-1, 0, 1)$, $p_6 = (0, 0, 1)$, $p_8 = (1, 0, 1)$ and the remaining points to be infinitely near to the given ones according to the classes $p_i - p_1$ written above. The corresponding pencil of sextics curves is generated by $(x^3 - y^2z - xz^2)^2$ and $x^2y^2(x-z)(x+z)$. The classes of the (-1) -curves of X are the rows of the following matrix.

$$\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & -1 & -1 & -1 & -1 & -1 & 0 & 0 \\ 2 & 0 & 0 & -1 & -1 & -1 & 0 & 0 & -1 & -1 \\ 4 & -2 & -2 & -1 & -1 & -1 & -2 & 0 & -1 & -1 \\ 4 & -2 & -2 & -1 & -1 & -1 & -1 & -1 & -2 & 0 \end{bmatrix}$$

The first column is the degree of the plane model, while the remaining columns are the negative of the multiplicities at the nine points. For example the first row is an exceptional divisor while the last row is a plane quartic with three double points and through other five simple points.

Case 2. Choose nine points p_1, \dots, p_9 on a smooth plane cubic curve C with flex at p_1 such that the classes of $p_2 - p_1, \dots, p_9 - p_1$ are the following elements of $\mathbb{Z}/4\mathbb{Z}$: 0, 0, 2, 2, 2, 1, 1, 3, 3. For example we can take $C := V(y^2z + xyz + yz^2 - x^3 - x^2z)$, $p_1 = (0, 1, 0)$, $p_3 = (-1, 0, 1)$, $p_6 = (0, 0, 1)$, $p_8 = (0, -1, 1)$ and the remaining points to be infinitely near to the given ones according to the classes $p_i - p_1$ written above. The corresponding pencil of sextics curves is generated by $(y^2z + xyz + yz^2 - x^3 - x^2z)^2$ and $x^2(x+z)^2y(x+y+z)$. The classes of the (-1) -curves of X are the rows of the following matrix.

$$\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & -1 & -1 & -1 & -1 & 0 & -1 & 0 \end{bmatrix}$$

2. RATIONAL ELLIPTIC SURFACES

Let X be a smooth projective rational surface with $-K_X$ nef and $(K_X)^2 = 0$. Given a curve C in the linear system $|-K_X|$ we can construct the homomorphism of groups

$$\alpha: K_X^\perp \rightarrow \text{Pic}^0(C) \quad w \mapsto i^*w,$$

where $i: C \rightarrow X$ is the inclusion. We denote by G the image of α and by $i: \mathbb{Z} \rightarrow K_X^\perp$ the homomorphism defined by $1 \mapsto -K_X$ and we let H be the subgroup of G generated by the images via α of the classes of all the irreducible components of C . Let $\kappa: \mathbb{Z} \rightarrow H$ be the homomorphism defined by $1 \mapsto h = \alpha(-K_X)$. Using the isomorphism between the lattices E_8 and $K_X^\perp/\mathbb{Z}(-K_X)$ we define $p: K_X^\perp \rightarrow E_8$ to be the corresponding projection. We summarize these definitions in the commutative diagram

$$(2.1) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{i} & K_X^\perp & \xrightarrow{p} & E_8 & \longrightarrow & 0 \\ & & \downarrow \kappa & & \downarrow \alpha & & \downarrow \bar{\alpha} & & \\ 0 & \longrightarrow & \langle h \rangle & \longrightarrow & G & \longrightarrow & G/\langle h \rangle & \longrightarrow & 0 \\ & & \downarrow \hookrightarrow & & \parallel & & \downarrow \beta & & \\ 0 & \longrightarrow & H & \longrightarrow & G & \longrightarrow & G/H & \longrightarrow & 0 \end{array}$$

with exact rows.

We recall the proofs of the following well-known lemmas for the sake of completeness. See also [4] and [5].

Lemma 2.1. *The order of the element h is finite if and only if the surface X is a minimal rational elliptic surface. Moreover, if h has finite order m , then the linear system $|-mK_X|$ is the elliptic pencil.*

Proof. Given $C \in |-K_X|$ there is an exact sequence of sheaves

$$(2.2) \quad 0 \longrightarrow \mathcal{O}_X(K_X) \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_C \longrightarrow 0.$$

Let $n > 0$ be an integer. By taking tensor product with $\mathcal{O}_X(-nK_X)$ and passing to the long exact sequence in cohomology, we see that the inequality $h^0(X, -nK_X) > 1$ holds if and only if $\alpha(-nK_X)$ is trivial in $\text{Pic}^0(C)$, that is if and only if the class $h = \alpha(-K_X)$ is torsion. This is equivalent to requiring that h has finite order. \square

Lemma 2.2. *Let R be a root of E_8 . Then one of the following possibilities occurs:*

- (1) if $\bar{\alpha}(R) = 0$ then $p^{-1}(R)$ contains an effective class;
- (2) otherwise the only effective classes in $p^{-1}(R)$ are components of C .

Proof. Assume that we are in case (1). Let D' be a class in $p^{-1}(R)$, so that $D'^2 = -2$ and $D' \cdot K_X = 0$, and let m be an integer such that $\alpha(D') = mh$. We denote by D the class $D' - mK_X$, so that $\alpha(D)$ is trivial. By abuse of notation, denote by D also a divisor of X whose class is D and form the exact sequence of sheaves

$$0 \longrightarrow \mathcal{O}_X(D + K_X) \longrightarrow \mathcal{O}_X(D) \longrightarrow \mathcal{O}_C(D) \longrightarrow 0.$$

Since $\alpha(D)$ is trivial, the first two cohomology groups of $\mathcal{O}_C(D)$ have dimension one. The above exact sequence and the vanishing of the Euler characteristic of $\mathcal{O}_X(D)$ immediately imply that at least one group $H^0(X, D)$, $H^2(X, D + K_X)^\vee \cong H^0(X, -D)$ is non-trivial. Thus either D or $-D$ is linearly equivalent to an effective divisor \bar{D} . If D is linearly equivalent to \bar{D} , then we are done. Otherwise there exists a positive integer n such that $D - nK_X \sim -\bar{D} - nK_X$ is linearly equivalent to an effective divisor and again the statement follows.

To prove (2) it suffices to observe that any effective class of $p^{-1}(R)$ is the class of a non-negative sum of (-2) -curves and that if Γ is a (-2) -curve of X not contained in C , then Γ is disjoint from C and hence $\alpha([\Gamma]) = 0$. \square

Given a smooth rational surface X with $-K_X$ nef and $K_X^2 = 0$, denote by Λ_X the sublattice of K_X^\perp spanned by the classes of (-2) -curves of X :

$$\Lambda_X = \langle R : R \text{ is the class of a } (-2)\text{-curve} \rangle.$$

Remark 2.3. By the adjunction formula Λ_X is a negative semidefinite sublattice of K_X^\perp and the quotient $\Lambda_X / \langle K_X \rangle$ is a negative definite lattice. By Lemma 2.2 each fiber of the quotient homomorphism $\Lambda_X \rightarrow \Lambda_X / \langle K_X \rangle$ contains at most a finite number of classes of (-2) -curves. Since the image of the set of classes of (-2) -curves of X is a subset of the finite set of roots of $\Lambda_X / \langle K_X \rangle \subseteq E_8$ we conclude that X contains finitely many (-2) -curves.

We assume now that C is a smooth plane cubic, p_1, \dots, p_9 are nine points on C , not necessarily distinct, and p_1 is a flex. Denote by X the blow up of \mathbb{P}^2 at the nine points p_1, \dots, p_9 , by E_1, \dots, E_9 the corresponding sequence of (-1) -classes in $\text{Pic}(X)$ and by L the class of the pull-back of a line. Recall from (2.1) that the homomorphism α is given by

$$(2.3) \quad \alpha: K_X^\perp \rightarrow G \quad dL - (m_1 E_1 + \dots + m_9 E_9) \mapsto \sum_{i=2}^9 m_i p_i$$

since $L|_C \sim 3p_1$ and p_1 is the origin of the elliptic curve (C, p_1) . The *characteristic sequence* of the pair given by the points p_1, \dots, p_9 and the smooth plane cubic curve C is the sequence $[p_1, \dots, p_9]$ in $\text{Pic}^9(C)$.

Proposition 2.4. *Let Λ be a sublattice of E_8 of type ADE and of finite index. Then there exists a minimal rational elliptic surface X such that $\Lambda_X / \langle K_X \rangle$ is isomorphic to Λ . An example of such a surface for any such Λ is given in Table (2.1).*

Lattice	Group	Characteristic sequence
E_8	$\langle 0 \rangle$	$[0, 0, 0, 0, 0, 0, 0, 0]$
D_8	$\mathbb{Z}/2$	$[0, 0, 0, 0, 0, 1, 1, 1]$
$E_7 + A_1$	$\mathbb{Z}/2$	$[0, 0, 0, 0, 0, 0, 1, 1]$
A_8	$\mathbb{Z}/3$	$[0, 0, 0, 0, 1, 1, 1, 2]$
$E_6 + A_2$	$\mathbb{Z}/3$	$[0, 0, 0, 0, 0, 0, 1, 1]$

$A_7 + A_1$	$\mathbb{Z}/4$	$[0, 0, 0, 0, 1, 1, 2, 2, 2]$
$D_5 + A_3$	$\mathbb{Z}/4$	$[0, 0, 0, 0, 0, 1, 1, 1, 1]$
$2A_4$	$\mathbb{Z}/5$	$[0, 0, 0, 0, 1, 1, 1, 1, 1]$
$A_5 + A_2 + A_1$	$\mathbb{Z}/6$	$[0, 0, 0, 1, 1, 1, 1, 1, 1]$
$D_6 + 2A_1$	$\mathbb{Z}/2 \oplus \mathbb{Z}/2$	$[0, 0, 0, 0, 0, 10, 10, 01, 01]$
$2D_4$	$\mathbb{Z}/2 \oplus \mathbb{Z}/2$	$[0, 0, 0, 10, 10, 01, 01, 11, 11]$
$2A_3 + 2A_1$	$\mathbb{Z}/2 \oplus \mathbb{Z}/4$	$[0, 0, 0, 10, 10, 01, 01, 01, 01]$
$4A_2$	$\mathbb{Z}/3 \oplus \mathbb{Z}/3$	$[0, 0, 0, 10, 10, 10, 01, 01, 01]$

Table 2.1: Characteristic sequences for maximal rank lattices

Proof. By hypothesis the group $G := E_8/\Lambda$ is finite. Moreover, by the classification of sublattices of E_8 generated by roots, the lattice Λ is isomorphic to one of the lattices of first column of the above table. Assume now that there exists a characteristic sequence $[p_1, \dots, p_9]$ of points on a smooth plane cubic curve C such that Λ is contained in the kernel of the homomorphism α of (2.3) and moreover α is surjective. By Lemma 2.1 and the fact that G is a finite group we deduce that the surface X is elliptic and minimal. Moreover the prime components of the unique elliptic fibration $\pi: X \rightarrow \mathbb{P}^1$ form a Coxeter-Dynkin diagram of type $\tilde{\Lambda}$ by Lemma 2.2, which gives the statement.

Hence it suffices to find a characteristic sequence realizing each of the thirteen lattices; this is shown in the third column of the table. \square

As an application of our results we determine the number of (-1) -curves on any minimal elliptic surface of Halphen index 2. In what follows, given a root lattice Λ we denote by $\tilde{\Lambda}$ the corresponding affine root lattice and by v_Λ a primitive vector of null square of $\tilde{\Lambda}$.

Lemma 2.5. *Let $v: \Lambda \rightarrow E_8$ be an embedding of a root lattice of rank eight into E_8 . Then the embeddings $\tilde{v}: \tilde{\Lambda} \rightarrow \tilde{E}_8$, which are compatible with j , are classified by the pairs (m, ξ) , where m is a positive integer and $\xi \in \text{Ext}_{\mathbb{Z}}^1(E_8/\Lambda, \mathbb{Z}/m\mathbb{Z})$.*

Proof. Observe that there are two isometries $\tilde{\Lambda} \cong \Lambda \oplus \langle v_\Lambda \rangle$ and $\tilde{E}_8 \cong E_8 \oplus \langle v_{E_8} \rangle$. Any isometric embedding $\tilde{v}: \tilde{\Lambda} \rightarrow \tilde{E}_8$ must map v_Λ to an integer multiple mv_{E_8} of v_{E_8} being these the only vectors of null square of both lattices. Moreover, up to sign change, we can assume m to be a positive integer. Then the statement is consequence of the following commutative diagram with exact rows and columns.

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{\cdot m} & \mathbb{Z} & \longrightarrow & \mathbb{Z}/m\mathbb{Z} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \Lambda \oplus \mathbb{Z} & \xrightarrow{\tilde{v}} & E_8 \oplus \mathbb{Z} & \xrightarrow{\tilde{\beta}} & G \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \alpha \\
 0 & \longrightarrow & \Lambda & \xrightarrow{v} & E_8 & \xrightarrow{\beta} & E_8/\Lambda \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Indeed, given such an embedding $\tilde{\iota}$, its cokernel $\tilde{\beta}$ gives an extension of E_8/Λ by $\mathbb{Z}/m\mathbb{Z}$ represented by the third column of the above diagram. Observe that the cokernel $\tilde{\beta}$ is unique since $\tilde{\iota}$ is a full-rank homomorphism of free abelian groups. Moreover equivalent embeddings, that is embeddings which differ up to isometries of the domain $\Lambda \oplus \mathbb{Z}$ and of the codomain $E_8 \oplus \mathbb{Z}$, give isomorphic extensions of E_8/Λ by $\mathbb{Z}/m\mathbb{Z}$.

On the other hand, given an extension of E_8/Λ by $\mathbb{Z}/m\mathbb{Z}$, since $E_8 \oplus \mathbb{Z}$ is a free abelian group, one can lift the map β to a map $\tilde{\beta}$ whose restriction to \mathbb{Z} surjects onto $\mathbb{Z}/m\mathbb{Z}$. The kernel $\tilde{\iota}$ of $\tilde{\beta}$ gives an embedding of $\Lambda \oplus \mathbb{Z}$ into $E_8 \oplus \mathbb{Z}$. \square

Proposition 2.6. *Let $\pi: X \rightarrow \mathbb{P}^1$ be a minimal elliptic fibration of Halphen index 2 on a rational surface. Then the number of (-1) -curves of X is given in Table (2.6).*

Type	Number of (-1) -curves
E_8	3
D_8	6, 9
$E_7 + A_1$	8, 10
A_8	15
$E_6 + A_2$	18
$A_7 + A_1$	24, 30
$D_5 + A_3$	28, 32
$2A_4$	45
$A_5 + A_2 + A_1$	60, 66
$D_6 + 2A_1$	26, 28
$2D_4$	28
$2A_3 + 2A_1$	108, 112
$4A_2$	144

TABLE 2.2. Number of (-1) -curves on Halphen elliptic surfaces of index 2.

Proof. Given a maximal rank sublattice Λ of E_8 , a positive integer m and an element $\xi \in \text{Ext}_{\mathbb{Z}}^1(E_8/\Lambda, \mathbb{Z}/m\mathbb{Z})$, in order to construct a Halphen fibration of index m whose configuration of reducible fibers is of type $\tilde{\Lambda}$ we choose a characteristic sequence $[p_1, \dots, p_9]$ of points on a smooth cubic curve C which satisfies the following conditions. If ξ is the following abelian extension

$$0 \longrightarrow \mathbb{Z}/m\mathbb{Z} \longrightarrow G \xrightarrow{\alpha} E_8/\Lambda \longrightarrow 0$$

we realize G as a subgroup of the elliptic curve (C, p_1) generated by nine points $p_1, \dots, p_9 \in C$ such that the element $h = p_1 + \dots + p_9$ of order m generates the kernel of α and $[p_1, \dots, p_9] \pmod{\langle h \rangle} \equiv [\bar{p}_1, \dots, \bar{p}_9]$, where the second sequence is the characteristic sequence in Table 2.1 associated to Λ .

Given such a characteristic sequence we reconstruct the set of (-2) -curves of the blow-up X of \mathbb{P}^2 at the nine points in the following way. By Lemma 2.2 each root R of Λ lifts to the class of an effective curve of X . The set \mathcal{S} of such lifted classes spans a cone in $\text{Pic}_{\mathbb{Q}}(X)$. Fix an ample class A of $\text{Pic}(X)$ and form the subset \mathcal{S}_1 of \mathcal{S} consisting of elements having minimal intersection with A . Then construct the set \mathcal{S}_n inductively as follows. Let \mathcal{S}_n be the subset of $\mathcal{S} \setminus \bigcup_{i \leq n-1} \mathcal{S}_i$ consisting of elements having minimal intersection with A and non-negative intersection with

any element of $\bigcup_{i \leq n-1} \mathcal{S}_i$. It follows that the set of (-2) -curves of X is

$$\mathcal{C} = \bigcup_i \mathcal{S}_i.$$

According to Lemma 2.5 the classes of these curves span a root sublattice of $K_X^\perp \cong \tilde{E}_8$ isometric to $\tilde{\Lambda}$. By means of these curves we define the lifting $\text{cl}_X: F \rightarrow K_X^\perp$ of the map $P: F \rightarrow K_X^\perp / \langle K_X \rangle = E_8$. We thus get the class δ_X of the corresponding toric variety, whose Riemann-Roch dimension is the number of (-1) -curves of X .

Now assume $m = 2$. Computing the multigraded Hilbert series of the toric varieties given in the appendix and excluding the cases where the trivial extension of E_8/Λ by $\mathbb{Z}/2\mathbb{Z}$ needs more than two generators, one directly checks that all the remaining cases are realized. \square

APPENDIX

We provide here the matrix Q defining the linear map $E \rightarrow \text{Cl}(Z)$ for each type of lattice analysed in this note.

Type	Class group	Grading matrix
E_8	\mathbb{Z}	$[1 \ 2 \ 3 \ 4 \ 5 \ 6 \ 4 \ 3 \ 2]$
D_8	$\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$	$\begin{bmatrix} 1 & 1 & 1 & 1 & 2 & 2 & 2 & 2 & 2 \\ 0 & \bar{0} & \bar{1} & \bar{1} & \bar{0} & \bar{0} & \bar{0} & \bar{1} & \bar{1} \end{bmatrix}$
A_8	$\mathbb{Z} \oplus \mathbb{Z}/3\mathbb{Z}$	$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & \bar{1} & \bar{2} & \bar{0} & \bar{1} & \bar{2} & \bar{0} & \bar{1} & \bar{2} \end{bmatrix}$
$E_7 + A_1$	$\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z}$	$\begin{bmatrix} 1 & 2 & 3 & 4 & 3 & 2 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ \bar{0} & \bar{0} & \bar{0} & \bar{0} & \bar{1} & \bar{1} & \bar{1} & \bar{0} & \bar{0} & \bar{1} \end{bmatrix}$
$E_6 + A_2$	$\mathbb{Z}^2 \oplus \mathbb{Z}/3\mathbb{Z}$	$\begin{bmatrix} 1 & 2 & 3 & 2 & 1 & 2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ \bar{0} & \bar{0} & \bar{0} & \bar{1} & \bar{1} & \bar{2} & \bar{1} & \bar{0} & \bar{1} & \bar{2} \end{bmatrix}$
$A_7 + A_1$	$\mathbb{Z}^2 \oplus \mathbb{Z}/4\mathbb{Z}$	$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ \bar{0} & \bar{1} & \bar{2} & \bar{3} & \bar{0} & \bar{1} & \bar{2} & \bar{3} & \bar{0} & \bar{2} \end{bmatrix}$
$D_5 + A_3$	$\mathbb{Z}^2 \oplus \mathbb{Z}/4\mathbb{Z}$	$\begin{bmatrix} 1 & 1 & 1 & 1 & 2 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ \bar{0} & \bar{1} & \bar{2} & \bar{3} & \bar{0} & \bar{2} & \bar{0} & \bar{1} & \bar{2} & \bar{3} \end{bmatrix}$
$2A_4$	$\mathbb{Z}^2 \oplus \mathbb{Z}/5\mathbb{Z}$	$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 \\ \bar{0} & \bar{1} & \bar{2} & \bar{3} & \bar{4} & \bar{0} & \bar{1} & \bar{2} & \bar{3} & \bar{4} \end{bmatrix}$
$2D_4$	$\mathbb{Z}^2 \oplus (\mathbb{Z}/2\mathbb{Z})^2$	$\begin{bmatrix} 1 & 1 & 1 & 1 & 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 2 \\ \bar{0} & \bar{1} & \bar{0} & \bar{1} & \bar{0} & \bar{0} & \bar{1} & \bar{0} & \bar{1} & \bar{0} \\ \bar{0} & \bar{0} & \bar{1} & \bar{1} & \bar{0} & \bar{0} & \bar{0} & \bar{1} & \bar{1} & \bar{0} \end{bmatrix}$

$A_5 + A_2 + A_1$	$\mathbb{Z}^3 \oplus \mathbb{Z}/6\mathbb{Z}$	$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ \bar{0} & \bar{1} & \bar{2} & \bar{3} & \bar{4} & \bar{5} & \bar{0} & \bar{2} & \bar{4} & \bar{0} & \bar{3} \end{bmatrix}$
$D_6 + 2A_1$	$\mathbb{Z}^3 \oplus (\mathbb{Z}/2\mathbb{Z})^2$	$\begin{bmatrix} 1 & 1 & 1 & 1 & 2 & 2 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ \bar{0} & \bar{1} & \bar{0} & \bar{1} & \bar{0} & \bar{0} & \bar{1} & \bar{0} & \bar{1} & \bar{0} & \bar{0} \\ \bar{0} & \bar{0} & \bar{1} & \bar{1} & \bar{0} & \bar{0} & \bar{0} & \bar{0} & \bar{1} & \bar{0} & \bar{1} \end{bmatrix}$
$2A_3 + 2A_1$	$\mathbb{Z}^4 \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$	$\begin{bmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ \bar{0} & \bar{1} & \bar{2} & \bar{3} & \bar{0} & \bar{1} & \bar{2} & \bar{3} & \bar{0} & \bar{2} & \bar{0} & \bar{0} \\ \bar{0} & \bar{1} & \bar{0} & \bar{1} & \bar{0} & \bar{0} & \bar{0} & \bar{0} & \bar{0} & \bar{1} & \bar{0} & \bar{1} \end{bmatrix}$
$4A_2$	$\mathbb{Z}^4 \oplus (\mathbb{Z}/3\mathbb{Z})^2$	$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ \bar{0} & \bar{1} & \bar{2} & \bar{0} & \bar{0} & \bar{0} & \bar{0} & \bar{1} & \bar{2} & \bar{0} & \bar{1} & \bar{2} \\ \bar{0} & \bar{0} & \bar{0} & \bar{0} & \bar{1} & \bar{2} & \bar{0} & \bar{1} & \bar{2} & \bar{0} & \bar{2} & \bar{1} \end{bmatrix}$

Table 2.3: Grading matrices

REFERENCES

- [1] Michela Artebani and Antonio Laface, *Cox rings of surfaces and the anticanonical Itaka dimension*, *Advances in Mathematics* **226** (2011), no. 6, 5252 - 5267. $\uparrow 1$
- [2] Ivan Arzhantsev, Ulrich Derenthal, Jürgen Hausen, and Antonio Laface, *Cox rings*, *Cambridge Studies in Advanced Mathematics*, vol. 144, Cambridge University Press, Cambridge, 2014. $\uparrow 5$
- [3] Wolf P. Barth, Klaus Hulek, Chris A. M. Peters, and Antonius Van de Ven, *Compact complex surfaces*, 2nd ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, vol. 4, Springer-Verlag, Berlin, 2004. MR2030225 (2004m:14070) $\uparrow 1$
- [4] François R. Cossec and Igor V. Dolgachev, *Enriques surfaces. I*, *Progress in Mathematics*, vol. 76, Birkhäuser Boston Inc., Boston, MA, 1989. MR986969 (90h:14052) $\uparrow 1, 7$
- [5] Brian Harbourne, *Anticanonical rational surfaces*, *Trans. Amer. Math. Soc.* **349** (1997), no. 3, 1191–1208, DOI 10.1090/S0002-9947-97-01722-4. MR1373636 (97f:14007) $\uparrow 7$
- [6] Mustapha Lahyane and Brian Harbourne, *Irreducibility of -1 -classes on anticanonical rational surfaces and finite generation of the effective monoid*, *Pacific J. Math.* **218** (2005), no. 1, 101–114, DOI 10.2140/pjm.2005.218.101. MR2224591 (2007b:14082) $\uparrow 3$
- [7] Rick Miranda and Ulf Persson, *On extremal rational elliptic surfaces*, *Math. Z.* **193** (1986), no. 4, 537–558, DOI 10.1007/BF01160474. MR867347 (88a:14044) \uparrow
- [8] Ezra Miller and Bernd Sturmfels, *Combinatorial commutative algebra*, *Graduate Texts in Mathematics*, vol. 227, Springer-Verlag, New York, 2005. MR2110098 (2006d:13001) $\uparrow 6$
- [9] Keiji Oguiso and Tetsuji Shioda, *The Mordell-Weil lattice of a rational elliptic surface*, *Comment. Math. Univ. St. Paul.* **40** (1991), no. 1, 83–99. MR1104782 (92g:14036) $\uparrow 3$
- [10] Burt Totaro, *The cone conjecture for Calabi-Yau pairs in dimension 2*, *Duke Math. J.* **154** (2010), no. 2, 241–263. $\uparrow 1$

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD DE CONCEPCIÓN, CASILLA 160-C, CONCEPCIÓN,
CHILE

E-mail address: antonio.laface@gmail.com

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL, UNITED KINGDOM

E-mail address: adomani@gmail.com