

# Predicting the extinction of Ebola spreading in Liberia due to mitigation strategies

L. D. Valdez,<sup>1</sup> Hênio Henrique Aragão Rêgo,<sup>2</sup> H. E. Stanley,<sup>3</sup> and L. A. Braunstein<sup>1,3</sup>

<sup>1</sup>*Departamento de Física, Facultad de Ciencias Exactas y Naturales,*

*Universidad Nacional de Mar del Plata,*

*and Instituto de Investigaciones Físicas de Mar del Plata (IFIMAR-CONICET),*

*Deán Funes 3350, 7600 Mar del Plata, Argentina*

<sup>2</sup>*Departamento de Física, Instituto Federal de Educação,*

*Ciência e Tecnologia do Maranhão, São Luís, MA, 65030-005, Brazil*

<sup>3</sup>*Center for Polymer Studies, Boston University,*

*Boston, Massachusetts 02215, USA.*

## Abstract

The Ebola virus is spreading throughout West Africa, causing thousands of deaths. When developing strategies to slow or halt the epidemic, mathematical models that reproduce the spread of this virus are essential for understanding which variables are relevant. We study the propagation of the Ebola virus through Liberia by using a model in which people travel from one county to another in order to quantify the effectiveness of different strategies. For the initial months in which the Ebola virus spreads, we find that the arrival times of the disease into the counties predicted by our model are in good agreement with World Health Organization data. We also find that reducing mobility delays the arrival of Ebola virus in each county by only a few weeks, and as a consequence, is insufficient to contain the epidemic. Finally we study the effect of a strategy focused on increasing safe burials and hospitalization under two scenarios: (i) one implemented in mid-July 2014 and (ii) one in mid-August, which was the actual time that strong interventions began in Liberia. We find that if scenario (i) had been pursued the lifetime of the epidemic would have been reduced by three months and the total number of infected individuals reduced by 80%, when compared to scenario (ii). Our projection under scenario (ii) is that the spreading will cease by mid-summer 2015.

## I. INTRODUCTION

*For a fleeting moment last spring, the epidemic sweeping West Africa might have been stopped. But the opportunity to control the virus, which has now caused more than 7,800 deaths, was lost [1].*

One of the deadliest and most persistent of epidemics [2], the current Ebola outbreak in Western Africa as of 31 December 2014 has caused 20,171 cases and approximately 7,889 deaths in three different countries—Guinea, Sierra Leona and Liberia—according to the World Health Organization [3]. The total numbers increase when cases and deaths from other affected countries in which the outbreak has been officially declared over [4] are added. Although the epidemic is now primarily restricted to those three countries, because there are no concrete signs that the spread of the infection is under control [1, 5, 6], the fear that it could spread to other countries remains.

Cultural, economic, and political factors in that region of Western Africa [2, 7–11] have hampered the effectiveness of the intervention strategies used by the health authorities. Because of a lack of reliable information about local patterns of the spreading of the Ebola virus disease (EVD) [12–14], the strategies currently being used, including the mobilization of resources, the creation of new Ebola treatment centers (ETC), the development of safe burial procedures, and the international coordination of the efforts [15] have been only partially successful.

Population mobility—the movement of individuals seeking safer areas, better health infrastructures, or food supplies—strongly affects disease propagation and plays a major role in epidemic spreading and in the effectiveness of any intervention scheme [16]. Thus understanding these patterns of movement is essential when planning health interventions to contain outbreaks within a certain region.

In recent years a number of mobility studies have been published [17–19], including Wesolowski et al. [16], who used mobile network data to analyze mobility patterns within the context of the ongoing Ebola outbreak. In response to this work there is an expanding effort to model and simulate the spatio-temporal evolution of Ebola spreading [20] in order to predict its local dynamics and understand its complexity.

In this manuscript we present a stochastic compartmental model and a set of differential equations, which are the quasi-deterministic representation of the stochastic model, for

the spreading of EVD in Liberia that incorporates population mobility between regions within the country. Unlike other models used to describe Ebola outbreaks [5, 21–24], our model enables us to explain how the spreading inside Liberia is due to mobility between different counties. As a result, our findings show that reducing mobility among counties only delays the spread of the disease, and has no practical effect in containing the Ebola epidemic. We also show how an earlier and more energetic response to the spreading would have been extremely effective in containing and decreasing the impact of the disease on the healthy population.

## II. MODEL

In our model we classify individuals as susceptible (S), exposed (E), i.e., infected but not infectious, infected (I), hospitalized (H), recovered (R), i.e., either cured or dead with a safe burial that does not transmit the disease, or dead (F) with an unsafe burial that transmits the disease. Additionally, we classify infected and hospitalized individuals according to their fate: those who are infected and will be hospitalized and will die ( $I_{DH}$ ), those who are infected, won't be hospitalized and will die ( $I_{DNH}$ ), those who are infected, will be hospitalized and will recover ( $I_{RH}$ ), those who are infected, won't be hospitalized and will recover ( $I_{RNH}$ ), those who are hospitalized and will die ( $H_D$ ), and those who are hospitalized and will recover ( $H_R$ ).

Figure 1 shows a schematic presentation of the model indicating the compartmental states (red boxes) and the transition rates among the states (connecting arrows). The  $I$ ,  $I = I_{DH} + I_{DNH} + I_{RH} + I_{RNH}$  represents the total number of infected individuals, and  $H = H_R + H_D$  the total number of those hospitalized. In Table I we show the different parameters used to calculate the transition rates among the different compartmental states in our model, while Table S1 (see Supplementary Information, Sec. A) lists the  $N_T = 12$  transitions between states and their rates  $\lambda_i$  with  $i = 1, \dots, N_T$ .




FIG. 1: A schematic of the transitions between different states of our model for the EVD spreading in West Africa 2014 and their respective transition rates. In the model, the population is divided into ten compartmental states (See Table S1): Susceptible ( $S$ ) individuals who in contact with infected individuals can become exposed ( $E$ ). These  $E$  individuals after the incubation period become infected and follow four different scenarios: (i) Infected individuals that will be cured —recovered— without hospitalization ( $I_{RNH}$ ); (ii) Infected individuals who will be cured ( $I_{RH}$ ) after spending a period on a hospital ( $H_R$ ); (iii) Infected individuals without being hospitalized ( $I_{DNH}$ ) who will die and may infect other individuals in their funerals ( $F$ ); and (iv) Infected individuals ( $I_{DH}$ ) that even after spending a period in a hospital ( $H_D$ ) will die and may also spread the infection in the funerals ( $F$ ). Recovered individual ( $R$ ), can be cured or dead.

To explain how geographical mobility spreads the disease, we assume that there is a flow of individuals between all  $N_{co} = 15$  counties of Liberia. We also assume that only susceptible or exposed individuals can travel between counties. Thus the deterministic evolution equations for the number of individuals in each state in county  $c$  in our model

Transition Parameters	Value	References
Mean duration of the incubation period ( $1/\alpha$ )	7 days	[25–27]
Mean time from the onset to the hospitalization ( $1/\gamma_H$ )	5 days	[28]
Mean duration from onset to death ( $1/\gamma_D$ )	9.6 days	[28]
Mean time from onset to the end for the cured ( $1/\gamma_I$ )	10 days	[27, 29]
Mean time from death to traditional burial ( $1/\gamma_F$ )	2 days	[21]
Proportion of cases hospitalized ( $\theta$ )	50%	[5]
Fatality Ratio ( $\delta$ )	50 %	[5]
Mean time from hospitalization to end for cured ( $1/\gamma_I$ )	5 days	[21]
Mean time from hospitalization to dead ( $1/\gamma_{HD}$ )	4.6 days	[21]

TABLE I: **Transition parameters used to calculate the transition rates in our epidemic model.** Table describing the different parameters used to calculate the transition rates among the ten different compartmental states in our model.

are

$$\frac{d S^c}{d t} = -\frac{1}{N_c}(\beta_I S^c I^c + \beta_H S^c H^c + \beta_F S^c F^c) + \sigma_c(\bar{S}) ; \quad (1)$$

$$\frac{d E^c}{d t} = \frac{1}{N_c}(\beta_I S^c I^c + \beta_H S^c H^c + \beta_F S^c F^c) - \alpha E^c + \sigma_c(\bar{E}) ; \quad (2)$$

$$\frac{d I_{DH}^c}{d t} = \alpha \delta \theta E^c - \gamma_H I_{DH}^c ; \quad (3)$$

$$\frac{d I_{DNH}^c}{d t} = \alpha \delta (1 - \theta) E^c - \gamma_D I_{DNH}^c ; \quad (4)$$

$$\frac{d I_{RH}^c}{d t} = \alpha (1 - \delta) \theta E^c - \gamma_H I_{RH}^c ; \quad (5)$$

$$\frac{d I_{RNH}^c}{d t} = \alpha (1 - \delta) (1 - \theta) E^c - \gamma_I I_{RNH}^c ; \quad (6)$$

$$\frac{d H_D^c}{d t} = \gamma_H I_{DH}^c - \gamma_{HD} H_D^c ; \quad (7)$$

$$\frac{d H_R^c}{d t} = \gamma_H I_{RH}^c - \gamma_{HI} H_R^c ; \quad (8)$$

$$\frac{d F^c}{d t} = \gamma_D I_{DNH}^c + \gamma_{HD} H_D^c - \gamma_F F^c ; \quad (9)$$

$$\frac{d R^c}{d t} = \gamma_I I_{RNH}^c + \gamma_{HI} H_R^c + \gamma_F F^c , \quad (10)$$

where  $\sigma_c$  is the total rate of mobility in each county  $c$  and is given by

$$\sigma_c(\bar{x}) = \sum_{c \neq j} \frac{x^j}{N_j} r_{j c} - \frac{x^c}{N_c} \sum_j r_{c j}, \quad (11)$$

where  $x^j$  ( $x^c$ ) is the number of individuals (susceptible or exposed), in county  $j$  ( $c$ ),  $N_j$  ( $N_c$ ) the total population of county  $j$  ( $c$ ), and  $r_{j c}$  and  $r_{c j}$  the mobility rates from county  $j \rightarrow c$  and from county  $c \rightarrow j$ , respectively. Note that due to mobility the population in each county changes, but since this evolution is much slower than the dynamics of the disease spreading, we consider  $N_c$  to be constant [30]. In addition, in this model we disregard the mobility inside each county, i.e., we assume that the population is fully mixed. Because recovered individuals are unable to transmit the disease or be reinfected, they do not affect the results of our model and we disregard their movements between counties.

In the context of complex network research, this model of mobility between counties breaks the traditional full-mixing approach because each county can be thought of as a node of a metapopulation network [31] in which the weight of each link is proportional to the mobility flow.

Note that if in Eqs. (1)–(10) we drop the index  $c$  and disregard the flow mobility we are no longer taking the counties into account, and we have a scenario that represents the spread throughout the entire country.

### III. METHODS

We generate a stochastic compartmental model based on the Gillespie algorithm. At each iteration of the simulation we draw a random number  $\tau$  (which represents the waiting time until the next transition) from an exponential distribution  $\text{Exp}(\Delta)$ , with parameter

$$\Delta = \sum_{i=1}^{N_T} \sum_{j=1}^{N_{co}} \lambda_i^j + \sum_{i=1}^{N_{co}} \sum_{j=1}^{N_{co}} (E^j + S^j) r_{j i} / N_j. \quad (12)$$

Here the first term  $\lambda_i^j$  is the rate of transition between states  $i$  in county  $j$  given in Table S1, and the second term corresponds to the mobility rates given in Eq. (11) with  $x = E$  and  $x = S$ .

To estimate the transmission coefficients  $\beta_I$ ,  $\beta_H$ , and  $\beta_F$  we calibrate the system of differential equations using least-squares optimization with the data of the total cases

from Liberia in the March–August period [3], and we apply a temporal shift, which we will explain below. We compute the least-square values using a set of parameters generated using Latin hypercube sampling (LHS) in the parameter space  $[0, 1]^3$ , which we divide into  $10^6$  cubes of the same size. For each cube we choose a random point as a candidate for  $(\beta_I, \beta_H, \beta_F)$  in order to compute the standard deviation between the data and the system of differential equations obtained from this point.

At the beginning of the epidemic there are very few cases (infected individuals), thus the evolution of the disease is in a stochastic regime in which the dispersion of the number of new cases is comparable to its mean value (see Ref. [32]). When the number of infected individuals increases to a certain level, however, the epidemic evolves toward a quasi-deterministic regime and the evolution of the states of the stochastic simulation is the same as the states obtained using the solution of the evolution equations (Eqs. 1-10). Nevertheless, due to fluctuations in the initial stochastic regime, a random temporal displacement of the quasi-deterministic growth of the number of accumulated cases is generated.

Thus to remove this stochastic temporal shift and to compare the three aspects—the simulations, the numerical solution, and the data—we set the initial time at  $t = 0$  when the total number of cases is above a cutoff  $s_c$  [32–34]. For the calibration of the transmission coefficients, we use  $s_c = 200$  and in the least square method, we give the data above this cutoff 50% of the weight because we are assuming that above  $s_c$  the evolution of the disease spreading is quasi-deterministic. Finally, after we compute the sum of square residuals for each point in the parameter space, we apply the Akaike information criterion [35] to obtain a model-averaged estimate of the transmission coefficients.

To estimate the migration between counties inside Liberia, we utilize the model of West African regional transportation patterns developed by Wesolowski *et al.* [16, 19], which used, among other sources, recent mobile-phone data for Senegal and population-mobility data collected from surveys. Although this movement data is “historical” and does not reflect how local population behavior may have changed in response to the current crisis, we assume the patterns of mobility obtained in the Wesolowski model [16, 36] still represent, on average, a good approximation of how the population in Liberia was moving before the declaration of the outbreak. In other words, we are assuming that individuals seek locations that are familiar to them. The mobility data for the Wesolowski model

were provided by Flowminder [16, 36, 37], and the case data used to calibrate the model were those supplied in reports generated by the World Health Organization (WHO) [3].

## IV. RESULTS

### A. Calibration

According to WHO data [3], the first index case (patient zero) was diagnosed in Lofa on 17 March 2014. Thus our initial conditions in Lofa are (i) one infected individual in that county and (ii) the rest of the population susceptible. The estimated rates of transmission in  $\text{day}^{-1}$  obtained using the method presented in Sec. III with a cutoff of  $s_c = 200$  are  $\beta_I = 0.14 [0, 0.26]$  in the community,  $\beta_H = 0.29 [0, 0.92]$  in the hospitals, and  $\beta_F = 0.40 [0, 0.99]$  at the funerals, where the intervals correspond to the values used to obtain the average rates of transmission obtained from the Akaike criterion. From these rates, we construct the next-generation matrix [38, 39] (see Supplementary Information, Sec. B) in order to compute the reproductive number  $R_0$ , defined as the average number of people in a susceptible population one infected individual infects during his or her infectious period. This parameter is fundamental when predicting whether a disease can reach a macroscopic fraction of individuals [40]. For a critical value of  $R_0 = 1$  there is a phase transition below which no epidemic takes place, and the disease is only a small outbreak, while for  $R_0 > 1$  the probability that an epidemic spreading develops is greater than zero [40]. For the values of rates of transmission given above, we find that the reproductive number of the current EVD outbreak is  $R_0 = 2.11$ , well above the critical threshold  $R_0 = 1$ . We run our stochastic simulations for these estimated values in order to compare the total number of cases with the data given by WHO before the interventions began in the middle of August [15]. In Figure 2(a) we plot the number of cumulative cases as a function of time for 1000 realizations of our stochastic model and compare the results with WHO data [3] without any shift correction. The individual realizations have the same shape as the data but due to the stochasticity at the beginning of the outbreak the exponential increase in the number of cases occurs at different moments.






FIG. 2: **Cumulative number of cases in Liberia for the parameters given in Table I.** Cumulative number of cases obtained with our stochastic model with the transition presented in Table S1 and Eq. (11) in Liberia with 1000 realizations (gray lines) and the data (symbols) without temporal shift (a) and (b) with a temporal shift using  $s_c = 200$ . The transmission coefficients  $\beta_I = 0.14$ ,  $\beta_H = 0.29$  and  $\beta_F = 0.40$  were obtained as explained in Section III. From the WHO’s data the index case is located at Lofa on March 17 2014.

In Figure 2(b) we plot the cumulative number of cases as a function of time with the initial conditions explained above when a temporal shift is applied to the stochastic simulations. The agreement between the simulations and the data indicates that our model can successfully represent the dynamics of the spreading of the current Ebola outbreak in Liberia.

### B. The geographical spread of Ebola cases across Liberia due to mobility

The mobility among the 15 counties allows us to compute the arrival time  $t_a$  in each county, assuming that the index case was in Lofa on 17 March 2014. Figure 3 shows the violin plots of the arrival times  $t_a$  of the disease as it spreads from Lofa County into the other 14 Liberian counties and compares our results with those supplied in the WHO reports (circles).




FIG. 3: Figure a: Violin plots representing the distribution of arrival time  $t_a$  to each county considering the mobility flow of individuals among counties [16] without any restriction on the mobility. The results are obtained from our stochastic model with the estimated transmission coefficients over 1000 realizations. Error bars indicate the 95% confidence interval. From WHO's reports the index case (patient zero) was located at Lofa at 17 of March 2014. The circles represent the values of  $t_a$  reported by WHO. The very early case of Margibi is below the 5% probability, and it is explained in Ref. [41]. Figure b: Violin plots representing the distribution of arrival time  $t_a$  to each county reducing the mobility among counties by 80%.

Comparing the results of our predictions of the arrival times of the first case as it spreads to the other counties with the WHO data (see Fig. 3a), all counties except Margibi and Grand Gedeh fall into a 95% confidence interval (Grand Kru is almost 95%). This could be caused by (i) an underestimation of the number of cases in the WHO data [3] due to a lack of information [1], or (ii) because the data recorded is actually the time of reporting and not the time of onset. Although the data is thus an indirect measure of the arrival time  $t_a$  in each county, there is good agreement between simulations and data concerning the order of arrival of the disease from Lofa to the other 14 counties.

When the disease began to spread, population mobility decreased. This was in part due to imposed regulations attempting to contain the disease but also due to the population's fear of contagion. We reflect this in our model by decreasing the mobility. Figure 3(b) shows the arrival times produced by our model when, as a strategy for slowing the spread, the mobility is reduced by 80%. Note that this reduction delays the arrival of EVD in each county by only a few weeks. This suggests that reducing the mobility of the

individuals between counties will not stop the spread but will slow it sufficiently that other strategies can be developed and applied. Reducing mobility is also insufficient when considering international transmission of the disease [23] and more aggressive interventions are needed. We believe that an increase in the percentage of infected individuals receiving hospitalization in ETCs is needed, and also an increase in the percentage of safe burials following procedures that do not transmit the disease.

### C. Interventions and time to extinction

We test this statement and propose the following strategy to contain the disease and reduce its transmission: we reduce the mobility by 80%, we increase the number of burials following procedures that do not transmit the disease, and we increase the rate of hospitalization in ETCs. Because health workers in ETCs are appropriately trained, we expect that the probability that they will be infected is greatly reduced and the transmission coefficient  $\beta_H$  decreases. A sufficiently rapid response to the EVD by the ETCs requires that  $\beta_H$  be decreased exponentially to a final value of  $10^{-3}$ , and hospitalization  $\theta$  must be increased exponentially to reach  $\theta = 1$ . On the other hand, when considering efforts to change local burial customs, we propose that  $\beta_F$  decreases linearly to approach zero. This assumption is based on the fact that this strategy cannot be as aggressive as the other interventions because it takes a longer period of time to gather and train burial teams.

We apply these changes to simulate a two-month period and the final result is that  $R_0$  decreases from 2.11 to  $R_0 = 0.69$  (see Supplementary Information, Sec. B), below the epidemic threshold. We consider two scenarios, (i) implementing the strategy beginning August 15 (the middle of the month indicated by WHO [42] for the outbreak of EVD in West Africa) in which all symptomatic individuals are admitted to ETCs and safe burial procedures begin to apply, or (ii) implementing the same strategy, but beginning July 15 in order to study how delaying the implementation of the strategies affected containment. Our goal is to demonstrate that if the international response had been more rapid, the spreading disease would have been contained by early March 2015 instead of the end of May 2015.




FIG. 4: Evolution of the number of cases (black) and deaths (red) when a reduction of 80% in the mobility rates is applied. The value of  $\beta_H$  decreases exponentially to reach the value  $10^{-3}$  and  $\beta_F$  decreases linearly to reach a 0% of their original values. Also the hospitalization fraction increases exponentially to reach  $\theta = 1$ . All reductions in the transmission coefficient were applied during two months, for (a) beginning at July 15th and (b) August 15th. Solid lines were obtained from the evolution equations (1-10) and the symbols are the data. Figures c) and d) are the distribution of time to extinction in of the EVD epidemic obtained from the stochastic simulations, when the strategy is applied from middle July and from middle August 2014, respectively. We show these distributions from December 1st, 2014 to December 1st, 2015.

In Figs. 4(a) and 4(b) we can see that applying strategies that focus on reducing the number of cases produced in hospitals and funerals causes the cumulative number of cases to drop to a lowered plateau than the one predicted when no strategies are applied [43]. Note that the figures also indicate that when the proposed strategies in our model are used a crossover from exponential to quasi-linear behavior in the cumulative number of cases and deaths occurs within a period of approximately one month. Figure 4(a) shows that if our strategy had been applied in the middle of July the cumulative number of

cases and deaths would have been approximately 80% lower than the reported number that resulted when the strategies were instituted in the middle of August. Figure 4(b) shows that if we apply the strategy of our model to that actual mid-August starting time, it accurately predicts the actual trends of cases and deaths reported in the WHO data.

Our model is designed to quantify how the two different strategy implementation times affect the extinction time of the EVD epidemic. Figures 4(c) and 4(d) show the extinction time distributions, i.e, when  $E = I = H = F = 0$ , when the strategy is implemented in July 2014 and August 2014, respectively. We find that the median of this distribution when the strategy is implemented in July is 6 March 2015 and when it is implemented in August the median is 25 May 2015. Implementation in mid-August allows the generation of 8,000 new cases of the disease, but implementation in mid-July reduces the time to disease extinction by three months and allows the generation of only 1,700 new cases. The mid-August implementation faces a larger number of cases, the disease progression has a greater inertia against the strategy, and the cumulative number of cases requires a longer time to go from an exponential regime to a sub exponential regime. Thus if the health authorities and the international community had acted sooner the number of infected people would have been much lower.

## V. CONCLUSIONS

In this manuscript we study the spreading of the Ebola virus using stochastic and deterministic compartmental models that incorporate the mobility of individuals between the counties in Liberia. We find that our model describes well the arrival of the disease into each of the counties, that reducing population mobility has little effect on geographical containment of the disease, and that reducing population mobility must be accompanied by other intervention strategies. We thus examine the effect of an intervention strategy that focuses on both an increase in hospitalization and an increase in safer burial practices (which were the actual measures taken to contain the disease). Our study indicates that the intervention implemented in August 2014 reduced the total number of infected individuals significantly when compared to a scenario in which there is no strategy implementation, and it predicts that the epidemic will be extinct by mid-summer 2015. We also use our model to consider the difference in outcome had the strategy been implemented

one month earlier. We find that the cumulative number of cases and deaths would have been significantly lower and that the epidemic would have ended three months earlier. This indicates that a rapid and early intervention that reduces the disease transmission in hospitals and at funerals is the most important response to any possible re-emerging Ebola epidemic.

Although our model simplifies the dynamics of epidemic spreading, it provides an adequate picture of the geographical expansion of the disease and the evolution in the number of cases and deaths. In future research we will incorporate more aspects of population mobility and intervention strategies carried out by health authorities, which will enable us to describe in greater detail the evolution of an epidemic and the efficacy of different strategies.

Finally, the methods used in this manuscript to study Liberia can also be applied to Guinea and Sierra Leone, as soon as high quality data about the development of the epidemic in those countries become available.

## Supplementary Information

### A. Table of the transitions

Transition	Transition rate ( $\lambda_i$ )
$(S, E) \rightarrow (S - 1, E + 1)$	$\frac{1}{N}(\beta_F S I + \beta_H S H + \beta_F S F)$
$(E, I_{DH}) \rightarrow (E - 1, I_{DH} + 1)$	$\alpha \theta \delta E$
$(E, I_{DNH}) \rightarrow (E - 1, I_{DNH} + 1)$	$\alpha (1 - \theta) \delta E$
$(E, I_{RH}) \rightarrow (E - 1, I_{RH} + 1)$	$\alpha \theta (1 - \delta) E$
$(E, I_{RNH}) \rightarrow (E - 1, I_{RNH} + 1)$	$\alpha (1 - \theta) (1 - \delta) E$
$(I_{DH}, H_D) \rightarrow (I_{DH} - 1, H_D + 1)$	$\gamma_H I_{DH}$
$(I_{DNH}, F) \rightarrow (I_{DNH} - 1, F + 1)$	$\gamma_D I_{DNH}$
$(I_{RH}, H_R) \rightarrow (I_{RH} - 1, H_R + 1)$	$\gamma_H I_{RH}$
$(I_{RNH}, R) \rightarrow (I_{RNH} - 1, R + 1)$	$\gamma_I I_{RNH}$
$(H_D, F) \rightarrow (H_D - 1, F + 1)$	$\gamma_{HD} H_D$
$(H_R, R) \rightarrow (H_R - 1, R + 1)$	$\gamma_{HI} H_R$
$(F, R) \rightarrow (F - 1, R + 1)$	$\gamma_F F$

TABLE S1: **Table of the transition with their respective transition rates for our model.** Table representing the transition rates between different compartmental states in our model. The capital letters represents number of: susceptible individuals ( $S$ ), number of exposed individuals ( $E$ ), individuals infected who will be hospitalized and die ( $I_{DH}$ ), individuals infected who won't be non hospitalized and will die ( $I_{DNH}$ ), individuals infected who will be hospitalized and recovered ( $I_{RH}$ ), individuals infected who won't be hospitalized and will recover ( $I_{RNH}$ ), individuals hospitalized who will die ( $H_D$ ), individuals hospitalized who will recover ( $H_R$ ). Here  $R$  is the number of individuals cured or dead and  $F$  is the number of individuals in the funerals who will have unsafe burials and can infect. Here  $\beta_I$ ,  $\beta_H$  and  $\beta_F$  are the transmission coefficients in the community in the hospital and in the funerals respectively,  $\delta$  is the fatality ratio and  $\theta$  the fraction of the hospitalized ones. The inverse of the mean time period of the incubation is  $1/\alpha$ . The mean time period from symptoms to hospitalization is  $1/\gamma_H$ , from symptoms for non hospitalized individuals to dead is  $1/\gamma_D$ , from symptoms for hospitalized individuals to dead is  $1/\gamma_{HD}$ , from symptoms for hospitalized individuals to recovery is  $1/\gamma_{HI}$  and from dead to recover is  $1/\gamma_F$ . The flow of mobility for individuals in county  $i \rightarrow j$  is explained in Eq. (11)

## B. Estimation of $R_0$

In order to compute the reproduction number  $R_0$ , we construct the next-generation matrix, following van den Driessche et al. [38] and Diekmann et al. [39].

First, from the Jacobian matrix of the system of Eqs. (1-10) we construct the “transmission matrix”  $F$ , and the “transition matrix”  $V$ , obtaining

$$\mathbf{F} = \begin{pmatrix} \mathbf{F}_1 & \mathbf{0} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{F}_2 & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{F}_3 & \dots & \mathbf{0} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \dots & \mathbf{F}_{15} \end{pmatrix}$$

where

$$\mathbf{F}_i = \begin{pmatrix} 0 & \beta_I & \beta_I & \beta_I & \beta_I & \beta_H & \beta_H & \beta_F \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

and

$$\mathbf{V} = \begin{pmatrix} \mathbf{V}_{1,1} & \mathbf{V}_{1,2} & \mathbf{V}_{1,3} & \dots & \mathbf{V}_{1,15} \\ \mathbf{V}_{2,1} & \mathbf{V}_{2,2} & \mathbf{V}_{2,3} & \dots & \mathbf{V}_{2,15} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{V}_{15,1} & \mathbf{V}_{15,2} & \mathbf{V}_{15,3} & \dots & \mathbf{V}_{15,15} \end{pmatrix}$$

where



$$\mathbf{V}_{i,i} = \begin{pmatrix} \alpha + \sum_u r_{iu}/N_i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\alpha\delta\theta & \gamma_H & 0 & 0 & 0 & 0 & 0 & 0 \\ -\alpha\delta(1-\theta) & 0 & \gamma_D & 0 & 0 & 0 & 0 & 0 \\ -\alpha(1-\delta)\theta & 0 & 0 & \gamma_H & 0 & 0 & 0 & 0 \\ -\alpha(1-\delta)(1-\theta) & 0 & 0 & 0 & \gamma_I & 0 & 0 & 0 \\ 0 & -\gamma_H & 0 & 0 & 0 & \gamma_{HD} & 0 & 0 \\ 0 & 0 & 0 & -\gamma_H & 0 & 0 & \gamma_{HI} & 0 \\ 0 & 0 & -\gamma_D & 0 & 0 & -\gamma_{HD} & 0 & \gamma_F \end{pmatrix}$$

$$\mathbf{V}_{i,j} = \begin{pmatrix} -r_{ji}/N_j & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

with  $i \neq j$

Note that the mobility rates are only in the transition matrix. Using these matrices we construct the next generation matrix, defined as  $\mathbf{FV}^{-1}$ . Finally, the reproduction number is given by the spectral radio of the next generation matrix,  $R_0 = \rho(\mathbf{FV}^{-1})$ , i.e., its highest eigenvalue. Note that when the mobility rates go to zero,  $R_0$  decreases since in this limit, an infected individual in a given county, can transmits the disease only to those susceptible individuals who belong to the same county and cannot interact with people from other counties.

- 
- [1] Sack, K., Fink, S., Belluck, P., Nossiter, A. & Berhulak, D. How Ebola Roared Back. *The New York Times* (2014). URL <http://www.nytimes.com/2014/12/30/health/how-ebola-roared-back.html>, (Accessed: 4th February 2015).

- [2] Frieden, T. R., Damon, I., Bell, B. P., Kenyon, T. & Nichol, S. Ebola 2014 - New Challenges, New Global Response and Responsibility. *The New England Journal of Medicine* **371**, 1177–1180 (2014).
- [3] World Health Organization. Ebola response roadmap - Situation report 31-12-2014 (2014). URL "<http://www.who.int/csr/disease/ebola/situation-reports/en/>".
- [4] Center for Disease Control and Prevention. 2014 Ebola Outbreak in West Africa (2014). URL <http://www.cdc.gov/vhf/ebola/outbreaks/2014-west-africa/index.html>, (Accessed: 1st January 2015).
- [5] Rivers, C. M., Lofgren, E. T., Marathe, M., Eubank, S. & Lewis, B. L. Modeling the Impact of Interventions on an Epidemic of Ebola in Sierra Leone and Liberia. *PLOS Currents Outbreaks* (2014). URL <http://dx.doi.org/10.1371/currents.outbreaks.fd38dd85078565450b0be3fcd78f5ccf>.
- [6] Reuters / S. Nebehay. Ebola still spreading in western Sierra Leone, Guinea's forest: U.N (2014). URL <http://www.reuters.com/article/2014/12/09/us-health-ebola-who-idUSKBN0JM24X20141209>, (Accessed: 4th February 2015).
- [7] Aljazeera America / A. Mukpo. The biggest concern of the Ebola outbreak is political, not medical (2014). URL "<http://america.aljazeera.com/opinions/2014/8/ebola-virus-liberiasierraleonepolitics>". (Accessed: 4th February 2015).
- [8] The Guardian / A. Konneh. Ebola isn't just a health crisis - it's a social and economic one too (2014). URL <http://www.theguardian.com/commentisfree/2014/oct/10/ebola-liberia-catastrophe-genera> (Accessed: 4th February 2015).
- [9] Global Issues / T. Brewer. Water and Sanitation Report Card: Slow Progress, Inadequate Funding (2014). URL <http://www.globalissues.org/news/2014/11/24/20342>, (Accessed: 4th February 2015).
- [10] The Washington Post / L.H. Sun, B. Dennis and J. Achenbach. Ebola's lessons, painfully learned at great cost in dollars and human lives (2014). URL <http://www.washingtonpost.com/national/health-science/ebolas-lessons-painfully-learned> (Accessed: 4th February 2015).

- [11] World Bulletin / News Desk. 2014: Liberia’s Ebola nightmare (2014). URL <http://www.worldbulletin.net/news/151832/2014-liberias-ebola-nightmare>, (Accessed: 4th February 2015).
- [12] Scientific American / S. Yasmin. Ebola Infections Fewer Than Predicted by Disease Models (2014). URL <http://www.scientificamerican.com/article/ebola-infections-fewer-than-predicted-by-d>. (Accessed: 4th February 2015).
- [13] News 24. Ebola: Bad data an issue (2014). URL <http://www.news24.com/Africa/News/Ebola-Bad-data-an-issue-20141208>, (Accessed: 4th February 2015).
- [14] The New Zealand Herald. In Ebola outbreak, bad data adds another problem (2014). URL [http://www.nzherald.co.nz/world/news/article.cfm?c\\_id=2&objectid=11371063](http://www.nzherald.co.nz/world/news/article.cfm?c_id=2&objectid=11371063), (Accessed: 4th February 2015).
- [15] World Health Organization. Ebola Virus Disease Outbreak Response Plan in West Africa (2014). URL <http://www.who.int/csr/disease/ebola/outbreak-response-plan/en/>.
- [16] Wesolowski, A. *et al.* Commentary: Containing the Ebola outbreak—the potential and challenge of mobile network data. *PLOS Currents Outbreaks* (2014). URL <http://dx.doi.org/10.1371/currents.outbreaks.0177e7fcf52217b8b634376e2f3efc5e>.
- [17] González, M. C., Hidalgo, C. A. & Barabási, A.-L. Understanding individual human mobility patterns. *Nature* **453**, 779–782 (2008).
- [18] Tatem, A. J. *et al.* The use of mobile phone data for the estimation of the travel patterns and imported Plasmodium falciparum rates among Zanzibar residents. *Malaria Journal* **8**, 287 (2009).
- [19] Wesolowski, A. *et al.* Quantifying travel behavior for infectious disease research: a comparison of data from surveys and mobile phones. *Scientific Reports* **4** (2014). URL <http://dx.doi.org/10.1038/srep05678>.
- [20] Merler, S. *et al.* Spatiotemporal spread of the 2014 outbreak of Ebola virus disease in Liberia and the effectiveness of non-pharmaceutical interventions: a computational modelling analysis. *The Lancet Infectious Diseases* **15**, 204–211 (2015).
- [21] Legrand, J., Grais, R., Boelle, P., Valleron, A. & Flahault, A. Understanding the dynamics of Ebola epidemics. *Epidemiology and Infection* **135**, 610–621 (2007).

- [22] Gomes, M. F. *et al.* Assessing the international spreading risk associated with the 2014 West African Ebola outbreak. *PLOS Currents Outbreaks* (2014).
- [23] Poletto, C. *et al.* Assessing the impact of travel restrictions on international spread of the 2014 West African Ebola epidemic. *Euro Surveillance* **19**, pii:20936 (2014).
- [24] Volz, E. & Pond, S. Phylodynamic analysis of Ebola virus in the 2014 Sierra Leone epidemic. *PLoS Currents Outbreaks* **10** (2014). URL <http://dx.doi.org/10.1371/currents.outbreaks.6f7025f1271821d4c815385b08f5f80e>.
- [25] Bwaka, M. A. *et al.* Ebola hemorrhagic fever in Kikwit, Democratic Republic of the Congo: clinical observations in 103 patients. *Journal of Infectious Diseases* **179**, S1–S7 (1999).
- [26] Ndambi, R. *et al.* Epidemiologic and clinical aspects of the Ebola virus epidemic in Mosango, Democratic Republic of the Congo, 1995. *Journal of Infectious Diseases* **179**, S8–S10 (1999).
- [27] Dowell, S. F. *et al.* Transmission of Ebola hemorrhagic fever: a study of risk factors in family members, Kikwit, Democratic Republic of the Congo, 1995. *Journal of Infectious Diseases* **179**, S87–S91 (1999).
- [28] Khan, A. S. *et al.* The reemergence of Ebola hemorrhagic fever, Democratic Republic of the Congo, 1995. *Journal of Infectious Diseases* **179**, S76–S86 (1999).
- [29] Rowe, A. K. *et al.* Clinical, virologic, and immunologic follow-up of convalescent Ebola hemorrhagic fever patients and their household contacts, Kikwit, Democratic Republic of the Congo. *Journal of Infectious Diseases* **179**, S28–S35 (1999).
- [30] In our model without restriction on the mobility, the population in each county changes less than a 5% in one year.
- [31] Colizza, V. & Vespignani, A. Invasion threshold in heterogeneous metapopulation networks. *Physical Review Letters* **99**, 148701 (2007).
- [32] Barthélemy, M., Barrat, A., Pastor-Satorras, R. & Vespignani, A. Dynamical patterns of epidemic outbreaks in complex heterogeneous networks. *Journal of Theoretical Biology* **235**, 275 – 288 (2005).
- [33] Volz, E. SIR dynamics in random networks with heterogeneous connectivity. *Journal of Mathematical Biology* **56**, 293–310 (2008).
- [34] Valdez, L. D., Macri, P. A. & Braunstein, L. A. Temporal Percolation of the Susceptible Network in an Epidemic Spreading. *PLoS ONE* **7**, e44188 (2012). URL

<http://dx.doi.org/10.1371/journal.pone.0044188>.

- [35] Burnham, K. P. & Anderson, D. R. *Model selection and multimodel inference: a practical information-theoretic approach* (Springer, 2002).
- [36] Flowminder Foundation (2014). URL <http://www.flowminder.org/>.
- [37] The WorldPop project. Mobile phone flow maps (2014). URL <http://www.worldpop.org.uk/ebola/>, (Accessed: 4th February 2015).
- [38] Van den Driessche, P. & Watmough, J. Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Mathematical Biosciences* **180**, 29–48 (2002).
- [39] Diekmann, O., Heesterbeek, J. & Roberts, M. The construction of next-generation matrices for compartmental epidemic models. *Journal of the Royal Society Interface* **7**, 873–885 (2010).
- [40] Anderson, R. M. & May, R. M. *Infectious Diseases of Humans: Dynamics and Control* (Oxford University Press, Oxford, 1992).
- [41] Health Minister of Liberia told the Observer that a woman whose Ebola test was positive, traveled from Lofa to Margibi County by the end of March. URL "<http://www.liberianobserver.com/security-health/2-5-test-positive-ebola-liberia>" (Accessed: 4th February 2015).
- [42] WHO (2014) WHO Statement on the Meeting of the International Health Regulations Emergency Committee regarding the 2014 Ebola outbreak in West Africa.
- [43] Meltzer, M. I. *et al.* Estimating the future number of cases in the ebola epidemic—Liberia and Sierra Leone, 2014–2015. *MMWR Surveill Summ* **63**, 1–14 (2014).

### *Acknowledgments*

HES thanks the NSF (grants CMMI 1125290 and CHE-1213217) and the Keck Foundation for financial support. LDV and LAB wish to thank to UNMdP and FONCyT (Pict 0429/2013) for financial support.

### *Contributions*

All authors designed the research, analyzed data, discussed results, and contributed to writing the manuscript. LDV implemented and performed numerical experiments and simulations.