Errors-in-variables models with dependent measurements ∗†

Mark Rudelson^{*} and Shuheng Zhou[†] [⋆]Department of Mathematics, †Department of Statistics, University of Michigan, Ann Arbor, MI 48109

Department of Statistics, Tech Report 538, July 31, 2015

August 13, 2018

Abstract

Suppose that we observe $y \in \mathbb{R}^f$ and $X \in \mathbb{R}^{f \times m}$ in the following errors-in-variables model:

$$
y = X_0 \beta^* + \epsilon
$$

$$
X = X_0 + W
$$

where X_0 is a $f \times m$ design matrix with independent subgaussian row vectors, $\epsilon \in \mathbb{R}^f$ is a noise vector and W is a mean zero $f \times m$ random noise matrix with independent subgaussian column vectors, independent of X_0 and ϵ . This model is significantly different from those analyzed in the literature in the sense that we allow the measurement error for each covariate to be a dependent vector across its f observations. Such error structures appear in the science literature when modeling the trial-to-trial fluctuations in response strength shared across a set of neurons.

Under sparsity and restrictive eigenvalue type of conditions, we show that one is able to recover a sparse vector $\beta^* \in \mathbb{R}^m$ from the model given a single observation matrix X and the response vector y. We establish consistency in estimating β^* and obtain the rates of convergence in the ℓ_q norm, where $q = 1, 2$ for the Lasso-type estimator, and for $q \in [1, 2]$ for a Dantzig-type conic programming estimator. We show error bounds which approach that of the regular Lasso and the Dantzig selector in case the errors in W are tending to 0.

1 Introduction

The matrix variate normal model has a long history in psychology and social sciences, and is becoming increasingly popular in biology and genomics, neuroscience, econometric theory, image and signal processing, wireless communication, and machine learning in recent years, see for example [Dawid](#page-58-0) [\(1981](#page-58-0)); [Gupta and Varga](#page-58-1) [\(1992\)](#page-58-1); [Dutilleul](#page-58-2) [\(1999](#page-58-2)); [Werner et al.](#page-60-0) [\(2008](#page-60-0)); [Bonilla et al.](#page-57-0) [\(2008](#page-57-0)); [Yu et al.](#page-60-1) [\(2009](#page-60-1)); [Efron](#page-58-3)

[∗]Mark Rudelson is partially supported by NSF grant DMS 1161372 and USAF Grant FA9550-14-1-0009. Shuheng Zhou was supported in part by NSF under Grant DMS-1316731 and Elizabeth Caroline Crosby Funding from the Advance Program at the University of Michigan. This manuscript was submitted for peer review in August 1, 2015; minor typos are being corrected in this version.

[†]**Keywords.** Errors-in-variable models, measurement error data, subgaussian concentration, matrix variate distributions.

[\(2009](#page-58-3)); [Allen and Tibshirani](#page-57-1) [\(2010\)](#page-57-1); [Kalaitzis et al.](#page-58-4) [\(2013\)](#page-58-4), and the references therein. We call the random matrix X which contains f rows and m columns a single data matrix, or one instance from the matrix variate normal distribution. We say that an $f \times m$ random matrix X follows a matrix normal distribution with a separable covariance matrix $\Sigma_X = A \otimes B$, which we write $X_{f \times m} \sim \mathcal{N}_{f,m}(M, A_{m \times m} \otimes B_{f \times f})$. This is equivalent to say vec $\{X\}$ follows a multivariate normal distribution with mean vec $\{M\}$ and covariance $\Sigma_X = A \otimes B$. Here, vec $\{X\}$ is formed by stacking the columns of X into a vector in \mathbb{R}^{m} . Intuitively, A describes the covariance between columns of X while B describes the covariance between rows of X. See [Dawid](#page-58-0) [\(1981](#page-58-0)); [Gupta and Varga](#page-58-1) [\(1992](#page-58-1)) for more characterization and examples.

In this paper, we introduce the related Kronecker Sum models to encode the covariance structure of a matrix variate distribution. The proposed models and methods incorporate ideas from recent advances in graphical models, high-dimensional regression model with observation errors, and matrix decomposition. Let $A_{m \times m}$, $B_{f \times f}$ be symmetric positive definite covariance matrices. Denote the Kronecker sum of $A = (a_{ij})$ and $B = (b_{ij})$ by

$$
\Sigma = A \oplus B := A \otimes I_f + I_m \otimes B
$$
\n
$$
= \begin{bmatrix}\n a_{11}I_f + B & a_{12}I_f & \dots & a_{1m}I_f \\
 a_{21}I_f & a_{22}I_f + B & \dots & a_{2m}I_f \\
 \dots & \dots & \dots & \dots \\
 a_{m1}I_f & a_{m2}I_f & \dots & a_{mm}I_f + B\n\end{bmatrix}_{(mf)\times(mf)}
$$

where I_f is an $f \times f$ identity matrix. This covariance model arises naturally from the context of errors-invariables regression model defined as follows. Suppose that we observe $y \in \mathbb{R}^f$ and $X \in \mathbb{R}^{f \times m}$ in the following model:

$$
y = X_0 \beta^* + \epsilon \tag{1a}
$$

$$
X = X_0 + W \tag{1b}
$$

where X_0 is a $f \times m$ design matrix with independent row vectors, $\epsilon \in \mathbb{R}^f$ is a noise vector and W is a mean zero $f \times m$ random noise matrix, independent of X_0 and ϵ , with independent column vectors $\omega^1, \ldots, \omega^m$. In particular, we are interested in the additive model of $X = X_0 + W$ such that

$$
\text{vec}\left\{X\right\} \sim \mathcal{N}(0,\Sigma) \quad \text{where} \quad \Sigma = A \oplus B := A \otimes I_f + I_m \otimes B \tag{2}
$$

where we use one covariance component $A \otimes I_f$ to describe the covariance of matrix $X_0 \in \mathbb{R}^{f \times m}$, which is considered as the *signal* matrix, and the other component $I_m \otimes B$ to describe that of the *noise matrix* $W \in \mathbf{R}^{f \times m}$, where $\mathbb{E} \omega^j \otimes \omega^j = B$ for all j, where ω^j denotes the j^{th} column vector of W. Our focus is on deriving the statistical properties of two estimators for estimating β^* in [\(1a\)](#page-1-0) and [\(1b\)](#page-1-0) despite the presence of the additive error W in the observation matrix X . We will show that our theory and analysis works with a model much more general than that in [\(2\)](#page-1-1), which we will define in Section [1.1.](#page-2-0)

Before we go on to define our estimators, we now use an example to motiviate [\(2\)](#page-1-1) and its subgaussian generalization in Definition [1.2.](#page-2-1) Suppose that there are f patients in a particular study, for which we use X_0 to model the "systolic blood pressure" and W to model the seasonal effects. In this case, X models the fact that among the f patients we measure, each patient has its own row vector of observed set of blood pressures across time, and each column vector in W models the seasonal variation on top of the true signal at a particular day/time. Thus we consider X as measurement of X_0 with W being the observation error. That is, we model the seasonal effects on blood pressures across a set of patients in a particular study with a vector of dependent entries. Thus W is a matrix which consists of repeated independent sampling of spatially dependent vectors, if we regard the individuals as having spatial coordinates, for example, through their geographic locations. We will come back to discuss this example in Section [1.3.](#page-4-0)

1.1 The model and the method

We first need to define an independent isotropic vector with *subgaussian* marginals as in Definition [1.1.](#page-2-2) **Definition 1.1.** Let Y be a random vector in \mathbb{R}^p

- *1. Y is called isotropic if for every* $y \in \mathbf{R}^p$, $\mathbb{E}\left(\vert\left\langle Y,y\right\rangle\vert^2\right)=\Vert y\Vert_2^2$ 2 *.*
- *2. Y is* ψ_2 *with a constant* α *if for every* $y \in \mathbb{R}^p$,

$$
\| \langle Y, y \rangle \|_{\psi_2} := \inf \{ t : \mathbb{E} \left(\exp(\langle Y, y \rangle^2 / t^2) \right) \leq 2 \} \leq \alpha \| y \|_2.
$$
 (3)

The ψ_2 *condition on a scalar random variable* V *is equivalent to the subgaussian tail decay of* V, which *means* $\mathbb{P}(|V| > t) \leq 2 \exp(-t^2/c^2)$, *for all* $t > 0$.

Throughout this paper, we use ψ_2 vector, a vector with subgaussian marginals and subgaussian vector interchangeably.

Definition 1.2. Let Z be an $f \times m$ random matrix with independent entries Z_{ij} satisfying $\mathbb{E}Z_{ij} = 0$, $1 = \mathbb{E}Z_{ij}^2 \leq \|Z_{ij}\|_{\psi_2} \leq K$. Let Z_1, Z_2 be independent copies of Z. Let $X = X_0 + W$ such that

- 1. $X_0 = Z_1 A^{1/2}$ is the design matrix with independent subgaussian row vectors, and
- 2. $W = B^{1/2} Z_2$ *is a random noise matrix with independent subgaussian column vectors.*

Assumption (A1) allows the covariance model in [\(2\)](#page-1-1) and its subgaussian variant in Definition [1.2](#page-2-1) to be identifiable.

(A1) We assume $tr(A) = m$ is a known parameter, where $tr(A)$ denotes the trace of matrix A.

In the kronecker sum model, we could assume we know $\text{tr}(B)$, in order not to assume knowing $\text{tr}(A)$. Assuming one or the other is known is unavoidable as the covariance model is not identifiable otherwise. Moreover, by knowing $tr(A)$, we can construct an estimator for $tr(B)$:

$$
\widehat{\text{tr}}(B) = \frac{1}{m} \left(\|X\|_F^2 - f \text{tr}(A) \right)_+ \quad \text{and define} \quad \widehat{\tau}_B := \frac{1}{f} \widehat{\text{tr}}(B) \ge 0 \tag{4}
$$

where $(a)_+ = a \vee 0$. We first introduce the Lasso-type estimator, adapted from those as considered in [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)).

Suppose that $\widehat{tr}(B)$ is an estimator for $tr(B)/f$; for example, as constructed in [\(4\)](#page-2-3). Let

$$
\widehat{\Gamma} = \frac{1}{f} X^T X - \frac{1}{f} \widehat{\text{tr}}(B) I_m \quad \text{and} \quad \widehat{\gamma} = \frac{1}{f} X^T y. \tag{5}
$$

For a chosen penalization parameter $\lambda \geq 0$, and parameters b_0 and d, we consider the following regularized estimation with the ℓ_1 -norm penalty,

$$
\widehat{\beta} = \underset{\beta: ||\beta||_1 \le b_0 \sqrt{d}}{\arg \min} \frac{1}{2} \beta^T \widehat{\Gamma} \beta - \langle \widehat{\gamma}, \beta \rangle + \lambda ||\beta||_1,\tag{6}
$$

which is a variation of the Lasso [Tibshirani](#page-59-1) [\(1996](#page-59-1)) or the Basis Pursuit [Chen et al.](#page-58-5) [\(1998](#page-58-5)) estimator. Although in our analysis, we set $b_0 \ge ||\beta^*||_2$ and $d = |\text{supp}(\beta^*)|$ for simplicity. In practice, both b_0 and d are understood to be parameters chosen to provide an upper bound on the ℓ_2 norm and the sparsity of the true $\beta^*.$

Recently, [Belloni et al.](#page-57-2) [\(2014\)](#page-57-2) discussed the following conic programming compensated matrix uncertainly (MU) selector, which is a variant of the Dantzig selector Candès and Tao [\(2007](#page-57-3)); [Rosenbaum and Tsybakov](#page-59-2) [\(2010](#page-59-2), [2013\)](#page-59-3). Adapted to our setting, it is defined as follows. Let $\lambda, \mu, \tau > 0$,

$$
\widehat{\beta} = \arg \min \{ ||\beta||_1 + \lambda t : (\beta, t) \in \Upsilon \} \text{ where}
$$
\n
$$
\Upsilon = \{ (\beta, t) : \beta \in \mathbb{R}^m, \left\| \widehat{\gamma} - \widehat{\Gamma}\beta \right\|_{\infty} \le \mu t + \tau, \|\beta\|_2 \le t \}
$$
\n(7)

where $\hat{\gamma}$ and $\hat{\Gamma}$ are as defined in [\(5\)](#page-2-4) with $\mu \sim \sqrt{\frac{\log m}{f}}$ $\frac{gm}{f}$, $\tau \sim \sqrt{\frac{\log m}{f}}$ $\frac{gm}{f}$. We refer to this estimator as the Conic programming estimator from now on.

1.2 Our contributions

We provide a unified analysis of the rates of convergence for both the Lasso-type estimator [\(6\)](#page-2-5) as well as the Conic Programming estimator [\(7\)](#page-3-0), which is a Dantzig selector-type, although under slightly different conditions. We will show the rates of convergence in the ℓ_q norm for $q = 1, 2$ for estimating a sparse vector $\beta^* \in \mathbb{R}^m$ in the model [\(1a\)](#page-1-0) and [\(1b\)](#page-1-0) using the Lasso-type estimator [\(6\)](#page-2-5) in Theorems [2](#page-8-0) and [4,](#page-11-0) and the Conic Programming estimator [\(7\)](#page-3-0) in Theorems [3](#page-9-0) and [5](#page-11-1) for $1 \le q \le 2$. For the Conic Programming estimator, we also show bounds on the predictive errors. The bounds we derive in both Theorems [2](#page-8-0) and [3](#page-9-0) focus on cases where the errors in W are not too small in their magnitudes in the sense that $\tau_B := \text{tr}(B)/f$ is bounded from below. For the extreme case when τ_B approaches 0, one hopes to recover bounds close to those for the regular Lasso or the Dantzig selector as the effect of the noise in matrix W on the procedure becomes negligible. We show in Theorems [4](#page-11-0) and [5](#page-11-1) that this is indeed the case. These results are new to the best of our knowledge.

In Theorems [2](#page-8-0) to [5,](#page-11-1) we consider the regression model in [\(1a\)](#page-1-0) and [\(1b\)](#page-1-0) with subgaussian random design, where $X_0 = Z_1 A^{1/2}$ is a subgaussian random matrix with independent row vectors, and $W = B^{1/2} Z_2$ is a $f \times m$ random noise matrix with independent column vectors where Z_1, Z_2 are independent subgaussian random matrices with independent entries (cf. Definition [1.2\)](#page-2-1). This model is significantly different from those analyzed in the literature. For example, unlike the present work, the authors in [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)) apply Theorem [8](#page-14-0) which states a general result on statistical convergence properties of the estimator [\(6\)](#page-2-5) to cases where W is composed of independent subgaussian row vectors, when the row vectors of X_0 are either independent or follow a Gaussian vector auto-regressive model. See also [Rosenbaum and Tsybakov](#page-59-2) [\(2010](#page-59-2), [2013\)](#page-59-3); [Chen and Caramanis](#page-58-6) [\(2013\)](#page-58-6); [Belloni et al.](#page-57-2) [\(2014](#page-57-2)) for the corresponding results on the compensated MU selectors, variant on the Orthogonal Matching Pursuit algorithm and the Conic Programming estimator [\(7\)](#page-3-0).

The second key difference between our framework and the existing work is that we assume that only one observation matrix X with the single measurement error matrix W is available. Assuming $(A1)$ allows us to estimate $\mathbb{E}W^TW$ as required in the estimation procedure [\(5\)](#page-2-4) directly, given the knowledge that W is composed of independent column vectors. In contrast, existing work needs to assume that the covariance matrix $\Sigma_W := \frac{1}{f} E W^T W$ of the independent row vectors of W or its functionals are either known a priori, or can

be estimated from an dataset independent of X, or from replicated X measuring the same X_0 ; see for example [Rosenbaum and Tsybakov](#page-59-2) [\(2010](#page-59-2), [2013](#page-59-3)); [Belloni et al.](#page-57-2) [\(2014\)](#page-57-2); [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)); [Carroll et al.](#page-57-4) [\(2006](#page-57-4)). Such repeated measurements are not always available or are costly to obtain in practice [Carroll et al.](#page-57-4) [\(2006](#page-57-4)).

A noticeable exception is the work of [Chen and Caramanis](#page-58-6) [\(2013](#page-58-6)), which deals with the scenario when the noise covariance is not assumed to be known. We now elaborate on their result, which is a variant of the orthogonal matching pursuit (OMP) algorithm [Tropp](#page-60-2) [\(2004](#page-60-2)); [Tropp and Gilbert](#page-59-4) [\(2007](#page-59-4)). Their support recovery result, that is, recovering the support set of β^* , applies only to the case when both signal matrix and the measurement error matrix have isotropic subgaussian row vectors; that is, they assume independence among both rows and columns in X (X_0 and W); moreover, their algorithm requires the knowledge of the sparsity parameter d, which is the number of non-zero entries in β^* , as well as a $\beta_{\rm min}$ condition: $\min_{j \in \text{supp } \beta^*} \left| \beta_j^* \right|$ $\left(= \Omega \left(\sqrt{\frac{\log m}{f}} (\|\beta^*\|_2 + 1) \right)$. They recover essentially the same ℓ_2 -error bounds as in [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)) and the current work when the covariance Σ_W is known.

In summary, oblivion in Σ_W and a general dependency condition in the data matrix X are not simultaneously allowed in existing work. In contrast, while we assume that X_0 is composed of independent subgaussian row vectors, we allow rows of W to be dependent, which brings dependency to the row vectors of the observation matrix X . In the current paper, we focus on the proof-of-the-concept on using the kronecker sum covariance and additive model to model two way dependency in data matrix X , and derive bounds in statistical convergence for [\(6\)](#page-2-5) and [\(7\)](#page-3-0). In some sense, we are considering a parsimonious model for fitting observation data with two-way dependencies; that is, we use the signal matrix to encode columnwise dependency among covariates in X , and error matrix W to explain its row-wise dependency. When replicates of X or W are available, we are able to study more sophisticated models and inference problems to be described in Section [1.3.](#page-4-0)

1.3 Discussion

The key modeling question is: would each row vector in W for a particular patient across all time points be a correlated normal or subgaussian vector as well? It is our conjecture that combining the newly developed techniques, namely, the concentration of measure inequalities we have derived in the current framework with techniques from existing work, we can handle the case when W follows a matrix normal distribution with a separable covariance matrix $\Sigma_W = C \otimes B$, where C is an $m \times m$ positive semi-definite covariance matrix. Moreover, for this type of "seasonal effects" as the measurement errors, the time varying covariance model would make more sense to model W, which we elaborate in the second example.

As a second example, in neuroscience applications, population coding refers to the information contained in the combined activity of multiple neurons [Kass et al.](#page-58-7) [\(2005](#page-58-7)). The relationship between population encod-ing and correlations is complicated and is an area of active investigation, see for example [Ruff and Cohen](#page-59-5) [\(2014](#page-59-5)); [Cohen and Kohn](#page-58-8) [\(2011](#page-58-8)) It becomes more often that repeated measurements (trials) simultaneously recorded across a set of neurons and over an ensemble of stimuli are available. In this context, one can imagine using a random matrix $X_0 \sim N_{f,m}(\mu, A \otimes B)$ which follows a matrix-variate normal distribution, or its subgaussian correspondent, to model the ensemble of mean response variables, e.g., the membrane potential, corresponding to the cross-trial average over a set of experiments. Here we use A to model the task correlations and B to model the baseline correlation structure among all pairs of neurons at the *signal* level. It has been observed that the onset of stimulus and task events not only change the cross-trial mean response in μ , but also alter the structure and correlation of the *noise* for a set of neurons, which correspond to the trial-to-trial fluctuations of the neuron responses. We use W to model such task-specific trial-to-trial fluctuations of a set of neurons recorded over the time-course of a variety of tasks. Models as in [\(1a\)](#page-1-0) and [\(1b\)](#page-1-0) are useful in predicting the response of set of neurons based on the current and past mean responses of all neurons. Moreover, we could incorporate non-i.i.d. non-Gaussian $W = [w_1, \ldots, w_m]$ where $w_t = B^{1/2}(t)z(t)$, where $z(1), \ldots, z(m)$ are independent isotropic subgaussian random vectors and $B(t) > 0$ for all t, to model the time-varying correlated noise as observed in the trial-to-trial fluctuations. It is possible to combine the techniques developed in the present paper with those in [Zhou et al.](#page-60-3) [\(2010](#page-60-3)); [Zhou](#page-60-4) [\(2014](#page-60-4)) to develop estimators for A, B and the time varying $B(t)$ which is itself an interesting topic, however, beyond the scope of the current work.

We leave the investigation of this more general modeling framework and relevant statistical questions to future work. We refer to [Carroll et al.](#page-57-4) [\(2006](#page-57-4)) for an excellent survey of the classical as well as modern developments in measurement error models. In future work, we will also extend the estimation methods to the settings where the covariates are measured with multiplicative errors which are shown to be re-ducible to the additive error problem as studied in the present work; see [Rosenbaum and Tsybakov](#page-59-3) [\(2013](#page-59-3)); [Loh and Wainwright](#page-59-0) [\(2012\)](#page-59-0). Moreover, we are interested in applying the analysis and concentration of measure results developed in the current paper and in our ongoing work to the more general contexts and settings where measurement error models are introduced and investigated; see for example [Dempster et al.](#page-58-9) [\(1977](#page-58-9)); [Carroll et al.](#page-57-5) [\(1985](#page-57-5)); [Stefanski](#page-59-6) [\(1985\)](#page-59-6); [Hwang](#page-58-10) [\(1986\)](#page-58-10); [Fuller](#page-58-11) [\(1987](#page-58-11)); [Stefanski](#page-59-7) [\(1990](#page-59-7)); [Carroll and Wand](#page-57-6) [\(1991](#page-57-6)); [Carroll et al.](#page-57-7) [\(1993](#page-57-7)); [Cook and Stefanski](#page-58-12) [\(1994](#page-58-12)); [Stefanski and Cook](#page-59-8) [\(1995](#page-59-8)); [Iturria et al.](#page-58-13) [\(1999](#page-58-13)); [Liang et al.](#page-58-14) [\(1999\)](#page-58-14); [Strimmer](#page-59-9) [\(2003](#page-59-9)); [Xu and You](#page-60-5) [\(2007\)](#page-60-5); [Hall and Ma](#page-58-15) [\(2007](#page-58-15)); [Liang and Li](#page-58-16) [\(2009](#page-58-16)); [Ma and Li](#page-59-10) [\(2010](#page-57-1)); [Allen and Tibshirani](#page-57-1) (2010); Städler et al. [\(2014\)](#page-59-11); [Søresen et al.](#page-59-12) [\(2014b](#page-59-12), a) and the references therein.

2 Assumptions and preliminary results

We will now define some parameters related to the restricted and sparse eigenvalue conditions that are needed to state our main results. We also state a preliminary result in Lemma [1](#page-6-0) regarding the relationships between the two conditions in Definitions [2.1](#page-5-0) and [2.2.](#page-5-1)

Definition 2.1. (Restricted eigenvalue condition $RE(s_0, k_0, A)$). Let $1 \le s_0 \le p$, and let k_0 be a positive *number.* We say that a $p \times q$ *matrix* A satisfies $RE(s_0, k_0, A)$ *condition with parameter* $K(s_0, k_0, A)$ *if for* $any \nu \neq 0,$ kauka al-

$$
\frac{1}{K(s_0, k_0, A)} := \min_{\substack{J \subseteq \{1, \ldots, p\}, \\ |J| \le s_0}} \min_{\|v_{Jc}\|_1 \le k_0 \|v_J\|_1} \quad \frac{\|Av\|_2}{\|v_J\|_2} > 0. \tag{8}
$$

It is clear that when s_0 and k_0 become smaller, this condition is easier to satisfy. We also consider the following variation of the baseline RE condition.

Definition 2.2. (Lower-RE condition) [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)) *The matrix* Γ *satisfies a Lower-*RE *condition with curvature* $\alpha > 0$ *and tolerance* $\tau > 0$ *if*

$$
\theta^T \Gamma \theta \ge \alpha \left\| \theta \right\|_2^2 - \tau \left\| \theta \right\|_1^2 \ \forall \theta \in \mathbf{R}^m.
$$

As α becomes smaller, or as τ becomes larger, the Lower-RE condition is easier to be satisfied.

Lemma 1. *Suppose that the Lower-RE condition holds for* $\Gamma := A^T A$ *with* $\alpha, \tau > 0$ *such that* $\tau(1 +$ $(k_0)^2 s_0 \le \alpha/2$. Then the $\mathsf{RE}(s_0, k_0, A)$ condition holds for A with

$$
\frac{1}{K(s_0, k_0, A)} \ge \sqrt{\frac{\alpha}{2}} > 0.
$$

Assume that $\mathsf{RE}((k_0+1)^2,k_0,A)$ holds. Then the Lower- RE condition holds for $\Gamma=A^TA$ with

$$
\alpha = \frac{1}{(k_0 + 1)K^2(s_0, k_0, A)} > 0
$$

where $s_0 = (k_0 + 1)^2$ *, and* $\tau > 0$ *which satisfies*

$$
\lambda_{\min}(\Gamma) \ge \alpha - \tau s_0/4.
$$
\nThe condition above holds for any $\tau \ge \frac{4}{(k_0+1)^3 K^2(s_0, k_0, A)} - \frac{4\lambda_{\min}(\Gamma)}{(k_0+1)^2}.$

The first part of Lemma [1](#page-6-0) means that, if k_0 is fixed, then smaller values of τ guarantee RE(s_0, k_0, A) holds with larger s_0 , that is, a stronger RE condition. The second part of the Lemma implies that a weak RE condition implies that the Lower-RE (LRE) holds with a large τ . On the other hand, if one assumes $RE((k_0 + 1)^2, k_0, A)$ holds with a large value of k_0 (in other words, a strong RE condition), this would imply LRE with a small τ . In short, the two conditions are similar but require tweaking the parameters. Weaker RE condition implies LRE condition holds with a larger τ , and Lower-RE condition with a smaller τ , that is, stronger LRE implies stronger RE. We prove Lemma [1](#page-6-0) in Section [8.](#page-20-0)

Definition 2.3. (Upper-RE condition) [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)) *The matrix* Γ *satisfies an upper-*RE *condition with curvature* $\bar{\alpha} > 0$ *and tolerance* $\tau > 0$ *if*

$$
\theta^T \Gamma \theta \leq \bar{\alpha} \left\| \theta \right\|_2^2 + \tau \left\| \theta \right\|_1^2 \ \forall \theta \in \mathbf{R}^m.
$$

Definition 2.4. Define the largest and smallest d-sparse eigenvalue of a $p \times q$ matrix A to be

$$
\rho_{\max}(d, A) := \max_{t \neq 0; d-sparse} \|At\|_2^2 / \|t\|_2^2, \text{ where } d < p,\tag{10}
$$

and
$$
\rho_{\min}(d, A) := \min_{t \neq 0; d-sparse} ||At||_2^2 / ||t||_2^2.
$$
 (11)

The rest of the paper is organized as follows. In Section [3,](#page-7-0) we present two main results Theorems [2](#page-8-0) and [3.](#page-9-0) We state results which improve upon Theorems [2](#page-8-0) and Theorem [3](#page-9-0) in Section [4,](#page-10-0) when the measurement errors in W are *small* in their magnitudes in the sense of $tr(B)$ being small. In Section [5,](#page-13-0) we outline the proof of the main theorems. In particular, In Section [5,](#page-13-0) we outline the proof for Theorems [2,](#page-8-0) [3,](#page-9-0) [4,](#page-11-0)and [5](#page-11-1) in Section [5,](#page-13-0) [5.1,](#page-15-0) [5.3](#page-17-0) and [5.4](#page-17-1) respectively. In Section [6,](#page-18-0) we show a deterministic result as well as its application to the random matrix $\hat{\Gamma} - A$ for $\hat{\Gamma}$ as in [\(5\)](#page-2-4) with regards to the upper and Lower RE conditions. In section [7,](#page-19-0) we show the concentration properties of the gram matrices XX^T and X^TX after we correct them with the corresponding *population* error terms defined by $tr(A)I_f$ and $tr(B)I_m$ respectively. These results might be of independent interests. The technical details of the proof are collected at the end of the paper. We prove Theorem [2](#page-8-0) in Section [9.](#page-22-0) We prove Theorem [3](#page-9-0) in Section [10.](#page-23-0) We prove Theorem [4](#page-11-0) and [5](#page-11-1) in Section [11](#page-24-0) and Section [12](#page-25-0) respectively. The paper concludes with a discussion of the results in Section [13.](#page-26-0) Additional proofs and theoretical results are collected in the Appendix.

Notation. Let e_1, \ldots, e_p be the canonical basis of \mathbb{R}^p . For a set $J \subset \{1, \ldots, p\}$, denote $E_J = \text{span}\{e_j :$ $j \in J$. For a matrix A, we use $||A||_2$ to denote its operator norm. For a set $V \subset \mathbb{R}^p$, we let conv V denote the convex hull of V. For a finite set Y, the cardinality is denoted by |Y|. Let B_1^p $_1^p$, B_2^p $_2^p$ and S^{p-1} be the unit ℓ_1 ball, the unit Euclidean ball and the unit sphere respectively. For a matrix $A = (a_{ij})_{1 \le i,j \le m}$, let $||A||_{\text{max}} = \max_{i,j} |a_{ij}|$ denote the entry-wise max norm. Let $||A||_1 = \max_j \sum_{i=1}^m |a_{ij}|$ denote the matrix ℓ_1 norm. The Frobenius norm is given by $||A||_F^2 = \sum_i \sum_j a_{ij}^2$. Let |A| denote the determinant and tr(A) be the trace of A. Let $\lambda_{\max}(A)$ and $\lambda_{\min}(A)$ be the largest and smallest eigenvalues, and $\kappa(A)$ be the condition number for matrix A. The operator or ℓ_2 norm $||A||_2^2$ $\frac{2}{2}$ is given by $\lambda_{\text{max}}(AA^T)$.

For a matrix A, denote by $r(A)$ the effective rank $\text{tr}(A)/||A||_2$. Let $||A||_F^2$ $_{F}^{2}/\Vert A\Vert_{2}^{2}$ $\frac{2}{2}$ denote the stable rank for matrix A. We write $diag(A)$ for a diagonal matrix with the same diagonal as A. For a symmetric matrix A, let $\Upsilon(A) = (v_{ij})$ where $v_{ij} = \mathbb{I}(a_{ij} \neq 0)$, where $\mathbb{I}(\cdot)$ is the indicator function. Let I be the identity matrix. We let C be a constant which may change from line to line. For two numbers $a, b, a \wedge b := \min(a, b)$ and $a \vee b := \max(a, b)$. We write $a \leq b$ if $ca \leq b \leq Ca$ for some positive absolute constants c, C which are independent of n, f, m or sparsity parameters. Let $(a)_+ := a \vee 0$. We write $a = O(b)$ if $a \leq Cb$ for some positive absolute constants C which are independent of n, f, m or sparsity parameters. These absolute constants C, C_1, c, c_1, \ldots may change line by line.

3 Main results

In this section, we will state our main results in Theorems [2](#page-8-0) and [3](#page-9-0) where we consider the regression model in [\(1a\)](#page-1-0) and [\(1b\)](#page-1-0) with random matrices $X_0, W \in \mathbb{R}^{f \times m}$ as defined in Definition [1.2.](#page-2-1)

For the Lasso-type estimator, we are interested in the case where the smallest eigenvalue of the column-wise covariance matrix A does not approach 0 too quickly and the effective rank of the row-wise covariance matrix B is bounded from below (cf. (14)). For the Conic Programming estimator, we impose a restricted eigenvalue condition as formulated in [Bickel et al.](#page-57-8) [\(2009](#page-57-8)); [Rudelson and Zhou](#page-59-14) [\(2013](#page-59-14)) on A and assume that the sparsity of β^* is bounded by $o(\sqrt{f/\log m})$. These conditions will be relaxed in Section [4](#page-10-0) where we allow τ_B to approach 0.

Before stating our main result for the Lasso-type estimator in Theorem [2,](#page-8-0) we need to introduce some more notation and assumptions. Let $a_{\text{max}} = \max_i a_{ii}$ and $b_{\text{max}} = \max_i b_{ii}$ be the maximum diagonal entries of A and B respectively. In general, under (A1), one can think of $\lambda_{\min}(A) \leq 1$ and for $s \geq 1$,

$$
1 \le a_{\max} \le \rho_{\max}(s, A) \le \lambda_{\max}(A),
$$

where $\lambda_{\text{max}}(A)$ denotes the maximum eigenvalue of A.

- (A2) The minimal eigenvalue $\lambda_{\min}(A)$ of the covariance matrix A is bounded: $1 \ge \lambda_{\min}(A) > 0$.
- (A3) Moreover, we assume that the condition number $\kappa(A)$ is upper bounded by $O\left(\sqrt{\frac{f}{\log A}}\right)$ $\log m$) and $\tau_B =$ $O(\lambda_{\max}(A)).$

Throughout the rest of the paper, $s_0 \geq 1$ is understood to be the largest integer chosen such that the following inequality still holds:

$$
\sqrt{s_0}\varpi(s_0) \le \frac{\lambda_{\min}(A)}{32C} \sqrt{\frac{f}{\log m}} \text{ where } \varpi(s_0) := \rho_{\max}(s_0, A) + \tau_B \tag{12}
$$

where we denote by $\tau_B = \text{tr}(B)/f$ and C is to be defined. Denote by

$$
M_A = \frac{64C\varpi(s_0)}{\lambda_{\min}(A)} \ge 64C. \tag{13}
$$

Throughout this paper, for the Lasso-type estimator, we will use the expression

$$
\tau := \frac{\alpha}{s_0}, \text{ where } \alpha = \lambda_{\min}(A)/2;
$$

(A2) thus ensures that the Lower-RE condition as in Definition [2.2](#page-5-1) is not vacuous. (A3) ensures that [\(12\)](#page-7-1) holds for some $s_0 \geq 1$.

Theorem 2. (**Estimation for the Lasso-type estimator**) *Set* $1 \leq f \leq m$ *. Suppose* m *is sufficiently large. Suppose (A1), (A2) and (A3) hold. Consider the regression model in* [\(1a\)](#page-1-0) *and* [\(1b\)](#page-1-0) *with independent random matrices* X_0 , W *as in Definition* [1.2,](#page-2-1) and an error vector $\epsilon \in \mathbb{R}^f$ independent of X_0 , W, with *independent entries* ϵ_j *satisfying* $\mathbb{E}\epsilon_j = 0$ *and* $\|\epsilon_j\|_{\psi_2} \leq M_{\epsilon}$. Let C_0 , $c' > 0$ be some absolute constants. Let $D_2 := 2(\|A\|_2 + \|B\|_2)$. Suppose that $\|B\|_F^2$ $\frac{2}{F}/\|B\|_2^2 \geq \log m$. Suppose that $c'K^4 \leq 1$ and

$$
r(B) := \frac{\text{tr}(B)}{\|B\|_2} \ge 16c'K^4 \frac{f}{\log m} \log \frac{\mathcal{V}m \log m}{f}
$$
 (14)

where V is a constant which depends on $\lambda_{\min}(A)$ *,* $\rho_{\max}(s_0, A)$ *and* $\text{tr}(B)/f$ *. Let* b_0 , ϕ *be numbers which satisfy*

$$
\frac{M_{\epsilon}^2}{K^2 b_0^2} \le \phi \le 1. \tag{15}
$$

Assume that the sparsity of β^* satisfies for some $0 < \phi \leq 1$

$$
d := |\text{supp}(\beta^*)| \le \frac{c'\phi K^4}{128M_A^2} \frac{f}{\log m} < f/2. \tag{16}
$$

Let $\widehat{\beta}$ *be an optimal solution to the Lasso-type estimator as in* [\(6\)](#page-2-5) *with*

$$
\lambda \ge 4\psi \sqrt{\frac{\log m}{f}} \quad \text{where} \quad \psi := C_0 D_2 K \left(K \left\| \beta^* \right\|_2 + M_{\epsilon} \right) \tag{17}
$$

Then for any d-sparse vectors $\beta^* \in \mathbb{R}^m$, such that $\phi b_0^2 \leq ||\beta^*||_2^2 \leq b_0^2$, we have with probability at least $1 - 16/m^3$,

$$
\left\|\widehat{\beta}-\beta^*\right\|_2 \le \frac{20}{\alpha}\lambda\sqrt{d} \quad \text{and} \quad \left\|\widehat{\beta}-\beta^*\right\|_1 \le \frac{80}{\alpha}\lambda d.
$$

We give an outline of the proof of Theorem [2](#page-8-0) in Section [5.1.](#page-15-0) We prove Theorem 2 in Section [9.](#page-22-0)

Discussions. Denote the Signal-to-noise ratio by

$$
\textsf{S/N}:= K^2\, \|\beta^*\|_2^2/M_{\epsilon}^2 \;\; \text{where} \;\; \textsf{S}:= K^2\, \|\beta^*\|_2^2 \;\; \text{and} \;\; \textsf{N}:= M_{\epsilon}^2.
$$

The two conditions on b_0 , ϕ imply that $N \leq \phi S$. Notice that this could be restrictive if ϕ is small.

We will show in Section [5.1](#page-15-0) that condition [\(15\)](#page-8-2) is not needed in order for the error bounds in terms of the ℓ_p , $p = 1, 2$ $p = 1, 2$ norm of $\beta - \beta^*$, as shown in the Theorem 2 statement to hold. It was indeed introduced so as to simplify the expression for the condition on d as shown in [\(16\)](#page-8-3). There we provide a slightly more general condition on d in [\(41\)](#page-16-0), where [\(15\)](#page-8-2) is not required. In summary, we prove that Theorem [2](#page-8-0) holds with $N = M_{\epsilon}^2$ and $S = \phi K^2 b_0^2$ in arbitrary orders, so long as condition [\(14\)](#page-8-1) holds and

$$
d = O\left(\frac{1}{M_A^2} \frac{f}{\log m}\right).
$$

For both cases, we require that $\lambda \asymp (\|A\|_2 + \|B\|_2)K\sqrt{S+N}\sqrt{\frac{\log m}{f}}$ $\frac{gm}{f}$ as expressed in [\(17\)](#page-8-4). That is, when either the noise level M_{ϵ} or the signal strength $K \|\beta^*\|$ increases, we need to increase λ correspondingly; moreover, when N dominates the signal $K^2 ||\beta^*||_2^2$ $_2^2$, we have for $d \approx \frac{1}{M_A^2}$ $\frac{f}{\log m}$,

$$
\left\|\widehat{\beta}-\beta^*\right\|_2/\left\|\beta^*\right\|_2\leq \frac{20}{\alpha}D_2K^2\sqrt{\frac{\mathsf{N}}{\mathsf{S}}}\frac{1}{M_A}\asymp D_2K^2\sqrt{\frac{\mathsf{N}}{\mathsf{S}}}\frac{1}{\varpi(s_0)}
$$

which eventually becomes a vacuous bound when $N \gg S$. We will present an improved bound in Theorem [4.](#page-11-0) We further elaborate on the relationships among the noise, the measurement error and the signal strength in Section [4.2.](#page-12-0)

Theorem 3. *Suppose* (A1) holds. Set $0 < \delta < 1$. *Suppose that* $f < m \ll \exp(f)$ and $1 \le d_0 < f$. Let $\lambda > 0$ be the same parameter as in [\(7\)](#page-3-0). Assume that $\mathsf{RE}(2d_0,3(1+\lambda),A^{1/2})$ holds. Suppose that $||B||_F^2$ $\frac{2}{F}$ / $||B||_2^2 \ge \log m$. Suppose that the sparsity of β^* is bounded by

$$
d_0 := |\text{supp}(\beta^*)| \le c_0 \sqrt{f/\log m}
$$
\n(18)

for some constant $c_0 > 0$ *; Suppose* $k_0 := 1 + \lambda$

$$
f \geq \frac{2000dK^4}{\delta^2} \log \left(\frac{60em}{d\delta} \right) \text{ where } \tag{19}
$$

$$
d = 2d_0 + 2d_0 a_{\text{max}} \frac{16K^2 (2d_0, 3k_0, A^{1/2})(3k_0)^2 (3k_0 + 1)}{\delta^2}.
$$
 (20)

Consider the regression model in [\(1a\)](#page-1-0) *and* [\(1b\)](#page-1-0) *with* X_0 , W *as in Definition* [1.2](#page-2-1) *and an error vector* $\epsilon \in \mathbb{R}^f$, *independent of* X_0, W , with independent entries ϵ_j *satisfying* $\mathbb{E} \epsilon_j = 0$ and $\|\epsilon_j\|_{\psi_2} \leq M_\epsilon$. Let $\widehat{\beta}$ be an *optimal solution to the Conic Programming estimator as in* [\(7\)](#page-3-0) *with input* $(\hat{\gamma}, \hat{\Gamma})$ *as defined in* [\(5\)](#page-2-4)*, where* $\text{tr}(B)$ *is as defined in* [\(4\)](#page-2-3)*. Choose for* $D_2 = 2(||A||_2 + ||B||_2)$ *and* $D_0 = \sqrt{\tau_B} + \sqrt{a_{\text{max}}}$ *,*

$$
\mu \approx D_2 K^2 \sqrt{\frac{\log m}{f}}
$$
 and $\tau \approx D_0 K M_{\epsilon} \sqrt{\frac{\log m}{f}}$.

Then with probability at least $1 - \frac{c'}{m^2} - 2 \exp(-\delta^2 f/2000K^4)$ *, for* $2 \ge q \ge 1$ *,*

$$
\left\|\widehat{\beta} - \beta^*\right\|_q \leq CD_2 K^2 d_0^{1/q} \sqrt{\frac{\log m}{f}} \left(\|\beta^*\|_2 + \frac{M_\epsilon}{K}\right). \tag{21}
$$

Under the same assumptions, the predictive risk admits the following bounds with the same probability as above,

$$
\frac{1}{f} \left\| X(\widehat{\beta} - \beta^*) \right\|_2^2 \le C'D_2^2 K^4 d_0 \frac{\log m}{f} \left(\|\beta\|_2^* + \frac{M_{\epsilon}}{K} \right)^2
$$

where $c', C_0, C, C' > 0$ are some absolute constants.

We give an outline of the proof of Theorem [3](#page-9-0) in Section [5](#page-13-0) while leaving the detailed proof in Section [10.](#page-23-0)

Discussions. Similar results have been derived in [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)); [Belloni et al.](#page-57-2) [\(2014](#page-57-2)), however, under different assumptions on the distribution of the noise matrix W . When W is a random matrix with i.i.d. subgaussian noise, our results will essentially recover the results in [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)) and [Belloni et al.](#page-57-2) [\(2014](#page-57-2)). The choice of λ for the Lasso estimator and parameters μ, τ for the DS-type estimator satisfy

$$
\lambda \asymp \mu \left\| \beta^* \right\|_2 + \tau
$$

This relationship is made clear through Theorem [8](#page-14-0) regarding the Lasso-type estimator, which follows from Theorem 1 [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)), Lemmas [6,](#page-13-1) [11,](#page-16-1) [14,](#page-17-2) and [16,](#page-18-1) which are the key results in proving Theorems [2,](#page-8-0) [3,](#page-9-0) [4,](#page-11-0) and [5.](#page-11-1) Finally, we note that following Theorem 2 as in [Belloni et al.](#page-57-2) [\(2014](#page-57-2)), one can show that without the relatively restrictive sparsity condition (18) , a bound similar to that in (21) holds, however with $\|\beta^*\|_2$ being replaced by $\|\beta^*\|_1$, so long as the sample size satisfies the requirement as in [\(27\)](#page-12-1).

4 Improved bounds when the measurement errors are small

Throughout our analysis of Theorems [2](#page-8-0) and [3,](#page-9-0) we focused on the case when the errors in W are sufficiently large in the sense that $\tau_B = \text{tr}(B)/f > 0$ is bounded from below; for example, this is explicitly indicated by the lower bound on the effective rank $r(B) = \text{tr}(B)/||B||_2$, when $||B||_2$ is bounded away from 0. More precisely, by the condition on the effective rank as in [\(14\)](#page-8-1), we have

$$
\tau_B = \frac{\text{tr}(B)}{f} \quad \geq \quad 16c' K^4 \frac{\|B\|_2}{\log m} \log \frac{\mathcal{V}m \log m}{f} \quad \text{where} \quad \mathcal{V} = 3e M_A^3/2.
$$

The bounds we derive in this section focus on cases where the measurement errors in W are *small* in their magnitudes in the sense of τ_B being small. For the extreme case when τ_B approaches 0, one hopes to recover a bound close to the regular Lasso or the Dantzig selector as the effect of the noise on the procedure should become negligible. We show in Theorems [4](#page-11-0) and [5](#page-11-1) that this is indeed the case. First, we define some contants which we use throughout the rest of the paper. Denote by

$$
D_0 = \sqrt{\tau_B} + a_{\text{max}}^{1/2}, \quad D'_0 = ||B||_2^{1/2} + a_{\text{max}}^{1/2}, \text{ and } \tau_B^+ := (\tau_B^{+/2})^2
$$
 (22)

where
$$
\tau_B^{+/2} := \sqrt{\tau_B} + \frac{D_{\text{oracle}}}{\sqrt{m}}
$$
 and $D_{\text{oracle}} = 2(||A||_2^{1/2} + ||B||_2^{1/2})$. (23)

We first state a more refined result for the Lasso-type estimator.

Theorem 4. *Suppose all conditions in Theorem [2](#page-8-0) hold, except that we drop* [\(15\)](#page-8-2) *and replace* [\(17\)](#page-8-4) *with*

$$
\lambda \ge 2\psi \sqrt{\frac{\log m}{f}} \quad \text{where} \quad \psi := 2C_0 D_0' K \left(\tau_B^{+/2} K \left\| \beta^* \right\|_2 + M_\epsilon \right). \tag{24}
$$

Suppose that for $0 < \phi \leq 1$ *and* $C_A := \frac{1}{128M_A^2}$

$$
d := |\text{supp}(\beta^*)| \le C_A \frac{f}{\log m} \left\{ c'C_{\phi} \wedge 2 \right\} \quad \text{where}
$$
\n
$$
C_{\phi} := \frac{\|B\|_2 + a_{\max}}{D^2} D_{\phi} \quad \text{for} \quad D_{\phi} = \frac{M_{\epsilon}^2 K^2}{b_0^2} + \tau_B^+ K^4 \phi,
$$
\n(25)

 $D = \rho_{\text{max}}(s_0, A) + \tau_B$, and $c', \phi, b_0, M_\epsilon$ and K as defined in Theorem [2.](#page-8-0)

Then for any d-sparse vectors $\beta^* \in \mathbb{R}^m$, such that $\phi b_0^2 \leq ||\beta^*||_2^2 \leq b_0^2$, we have with probability at least $1 - 16/m^3$,

$$
\left\|\widehat{\beta}-\beta^*\right\|_2 \le \frac{20}{\alpha}\lambda\sqrt{d} \quad \text{and} \quad \left\|\widehat{\beta}-\beta^*\right\|_1 \le \frac{80}{\alpha}\lambda d.
$$

We give an outline for the proof of Theorem [4](#page-11-0) in Section [5.3,](#page-17-0) and show the actual proof in Section [11.](#page-24-0) **Remark 4.1.** *Let us redefine the Signal-to-noise ratio by*

$$
\begin{array}{rcl}\n\mathbf{S/M} & := & \frac{K^2 \left\| \beta^* \right\|_2^2}{\tau_B^+ K^2 \left\| \beta^* \right\|_2^2 + M_\epsilon^2} \quad \text{where} \\
\mathbf{S} & := & K^2 \left\| \beta^* \right\|_2^2 \quad \text{and} \quad \mathbf{M} := M_\epsilon^2 + \tau_B^+ K^2 \left\| \beta^* \right\|_2^2\n\end{array}
$$

We now only require that $\lambda \asymp (a_{\text{max}}^{1/2} + ||B||_2^{1/2})$ $\frac{1}{2}$) $K\sqrt{M}\sqrt{\frac{\log m}{f}}$ $\frac{g\cdot m}{f}.$ That is, when either the noise level M_ϵ or the measurement error strength in terms of $\tau_B^{+/2} K\|\beta^*\|_2$ increases, we need to increase the penalty parameter λ correspondingly; moreover, when $d \asymp \frac{1}{M_A^2}$ f $\log m$

$$
\frac{\left\|\widehat{\beta}-\beta^*\right\|_2}{\left\|\beta^*\right\|_2}\leq \frac{20}{\alpha}D_0' K^2\sqrt{\frac{M}{\mathbb{S}}} \frac{1}{M_A}\asymp D_0' K^2\sqrt{\frac{M}{\mathbb{S}}} \frac{1}{\varpi(s_0)},
$$

which eventually becomes a vacuous bound when M ≫ S*.*

4.1 A Corollary for Theorem [3](#page-9-0)

We next state in Theorem [5](#page-11-1) an improved bound for the Conic programming estimator [\(7\)](#page-3-0), which improves upon Theorem [3](#page-9-0) when τ_B is small.

Theorem 5. *Suppose all conditions in Theorem [3](#page-9-0) hold, except that we replace the condition on* d *as in* [\(18\)](#page-9-1) *with the following. Suppose that the sample size* f *and the size of the support of* β ∗ *satisfy the following*

requirements: for $C_6 \geq D_{\text{oracle}}$ *and* $r_{m,m} = 2C_0 \sqrt{\frac{\log m}{fm}}$,

$$
d_0 = O\left(\tau_B^{-} \sqrt{\frac{f}{\log m}}\right) \quad \text{where} \quad \tau_B^{-} \le \frac{1}{\tau_B^{1/2} + 2C_6 K r_{m,m}^{1/2}} \tag{26}
$$

$$
and \t f \geq \frac{2000dK^4}{\delta^2} \log \left(\frac{60em}{d\delta} \right) \text{ where } \t (27)
$$

$$
d = 2d_0 + 2d_0 a_{\text{max}} \frac{16K^2 (2d_0, 3k_0, A^{1/2})(3k_0)^2 (3k_0 + 1)}{\delta^2}.
$$
 (28)

Let $\widehat{\beta}$ *be an optimal solution to the Conic Programming estimator as in* [\(7\)](#page-3-0) *with input* $(\widehat{\gamma}, \widehat{\Gamma})$ *as defined in* [\(5\)](#page-2-4), where $\widehat{tr}(B)$ *is as defined in* [\(4\)](#page-2-3)*. Suppose*

$$
\tau \approx D_0 M_{\epsilon} r_{m,f} \quad \text{where} \quad r_{m,f} = C_0 K \sqrt{\frac{\log m}{f}} \quad \text{and} \tag{29}
$$

$$
\mu \approx D'_0 \tilde{\tau}_B^{1/2} K r_{m,f} \quad \text{where} \quad \tilde{\tau}_B^{1/2} := \hat{\tau}_B^{1/2} + C_6 K r_{mm}^{1/2}.
$$
 (30)

Then with probability at least $1 - \frac{c''}{m^2} - 2 \exp(-\delta^2 f/2000K^4)$, for $2 \ge q \ge 1$, and $\tau_B^{\dagger/2} = (\tau_B^{\dagger/2} +$ $\frac{3}{2}C_6Kr_{m,m}^{1/2})$

$$
\left\|\widehat{\beta} - \beta^*\right\|_q \le C'D_0' K^2 d_0^{1/q} \sqrt{\frac{\log m}{f}} \left(\tau_B^{\dagger/2} \left\|\beta^*\right\|_2 + \frac{M_\epsilon}{K}\right);\tag{31}
$$

Under the same assumptions, the predictive risk admits the following bounds

$$
\frac{1}{f} \| X(\hat{\beta} - \beta^*) \|_2^2 \le C''(\|B\|_2 + a_{\max}) K^2 d_0 \frac{\log m}{f} \left(\tau_B^{\ddagger} K^2 \|\beta^*\|_2^2 + M_{\epsilon}^2 \right)
$$

with the same probability as above, where c'' , C' , $C'' > 0$ *are some absolute constants, and* $\tau_B^{\ddagger} \approx 2\tau_B +$ $3C_6^2K^2r_{m,m}.$

4.2 Discussions

In particular, when $\tau_B \to 0$, Theorem [5](#page-11-1) allows us to recover a rate close that of the Dantzig selector with an exact recovery if $\tau_B = 0$ is known a priori; see Section [13.](#page-26-0) Moreover the constraint [\(18\)](#page-9-1) on the sparsity parameter d_0 appearing in Theorem [3](#page-9-0) can now be relaxed as in [\(26\)](#page-12-1). Roughly speaking, one can think of d_0 being bounded as follows for the Conic programming estimator [\(7\)](#page-3-0):

$$
d_0 = O\left(\tau_B^- \sqrt{\frac{f}{\log m}} \wedge \frac{f}{\log(m/d_0)}\right) \quad \text{where} \quad \tau_B^- \asymp \frac{1}{\tau_B^{1/2}} \tag{32}
$$

That is, when τ_B decreases, we allow larger values of d_0 ; however, when $\tau_B \to 0$, the sparsity level of $d = O(f/\log(m/d))$ starts to dominate, which enables the Conic Programming estimator to achieve results similar to the Dantzig Selector when the design matrix X_0 is a subgaussian random matrix satis-fying the Restricted Eigenvalue conditions; See for example Candès and Tao [\(2007\)](#page-57-3); [Bickel et al.](#page-57-8) [\(2009](#page-57-8)); [Rudelson and Zhou](#page-59-14) [\(2013](#page-59-14)).

The condition on d (and D_{ϕ}) for the Lasso estimator as defined in [\(25\)](#page-11-2) suggests that as $\tau_B \to 0$, and thus $\tau_B^+ \to 0$ the requirement on the sparsity parameter d becomes slightly more stringent when $K^2 M_e^2/b_0^2 \approx 1$ and much more restrictive when $K^2 M_{\epsilon}^2 / b_0^2 = o(1)$; however, suppose we require

$$
M_{\epsilon}^{2} = \Omega(\tau_{B}^{+}K^{2} \, || \beta^{*} ||_{2}^{2}),
$$

that is, the stochastic error ϵ in the response variable y as in [\(1a\)](#page-1-0) does not converge to 0 as quickly as the measurement error W in [\(1b\)](#page-1-0) does, then the sparsity constraint becomes essentially unchanged as $\tau_B^+ \to 0$. In this case, essentially, we require that for some $c'' := c'C_{\phi}$

$$
d \leq C_A' \frac{f}{\log m} \left\{ \frac{c'' K^2 M_{\epsilon}^2}{b_0^2} \wedge 1 \right\} \text{ where } D_{\phi} \approx \frac{K^2 M_{\epsilon}^2}{b_0^2} \text{ and } C_A' := \frac{1}{64 M_A^2},
$$

given that $\tau_B^+ K^4 \phi \leq \frac{\tau_B^+ K^4 \|\beta^*\|_2^2}{b_0^2} \ll \frac{K^2 M_{\epsilon}^2}{b_0^2}.$

These tradeoffs are somehow different from the behavior of the Conic programming estimator (cf [\(32\)](#page-12-2)).

5 Proof of theorems

We first consider the following large deviation bound on $\left\|\hat{\gamma} - \hat{\Gamma}\beta^*\right\|$ as stated in Lemma [6.](#page-13-1) This entity appears in the constraint set in the conic programming estimator [\(7\)](#page-3-0), and is directly related to the choice of λ for the lasso-type estimator in view of Theorem [8.](#page-14-0) Events B_0 and B_{10} are defined in Section [B.2](#page-29-0) in the Appendix.

Lemma 6. *Suppose* (A1) holds. Let $X = X_0 + W$, where X_0, W are as defined in Theorem [2.](#page-8-0) Suppose that

$$
||B||_F^2 / ||B||_2^2 \ge \log m \quad \text{where} \quad m \ge 16.
$$

Let Γ *and* $\widehat{\gamma}$ *be as in* [\(5\)](#page-2-4). On event \mathcal{B}_0 , we have for $D_2 = 2(||A||_2 + ||B||_2)$ and some absolute constant C_0

$$
\left\|\widehat{\gamma} - \widehat{\Gamma}\beta^*\right\|_{\infty} \leq \psi \sqrt{\frac{\log m}{f}} \text{ where } \psi = C_0 D_2 K \left(K\left\|\beta^*\right\|_2 + M_{\epsilon}\right)
$$

is as defined in Theorem [2.](#page-8-0) Then $\mathbb{P}(\mathcal{B}_0) \geq 1 - 16/m^3$.

Lemma 7. Let $m \geq 2$. Let X be defined as in Definition [1.2](#page-2-1) and $\hat{\tau}_B$ be as defined in [\(4\)](#page-2-3). Denote by $\tau_B = \text{tr}(B)/f$ and $\tau_A = \text{tr}(A)/m$ *. Suppose that* $f \vee (r(A)r(B)) > \log m$ *. Denote by* \mathcal{B}_6 *the event such that*

$$
|\widehat{\tau}_B - \tau_B| \leq 2C_0 K^2 \sqrt{\frac{\log m}{mf}} \left(\frac{\|A\|_F}{\sqrt{m}} + \frac{\|B\|_F}{\sqrt{f}} \right) =: D_1 K^2 r_{m,m}
$$

where $D_1 := \frac{\|A\|_F}{\sqrt{m}} + \frac{\|B\|_F}{\sqrt{f}}$ and $r_{m,m} = 2C_0 \sqrt{\frac{\log m}{mf}}$.

Then $\mathbb{P}(\mathcal{B}_6) \geq 1 - \frac{3}{m^3}$ *. If we replace* $\sqrt{\log m}$ *with* $\log m$ *in the definition of event* \mathcal{B}_6 *, then we can drop the condition on* f *or* $r(A)r(B) = \frac{\text{tr}(A)}{\|A\|_2}$ $\mathrm{tr}(B)$ $\frac{dE(B)}{\|B\|_2}$ to achieve the same bound on event \mathcal{B}_6 .

We prove Lemma [7](#page-13-2) in Section [B.3](#page-29-1) in the Appendix. We prove Lemma [6](#page-13-1) in Section [C.1.](#page-31-0) We mention in passing that Lemma [6](#page-13-1) is essential in proving Theorem [3](#page-9-0) as well.

We state variations on this inequality in Lemma 14 and the remark which immediately follows. **Theorem 8.** *Consider the regression model in* [\(1a\)](#page-1-0) *and* [\(1b\)](#page-1-0)*. Let* $d \leq f/2$ *. Let* $\hat{\gamma}$ *,* $\hat{\Gamma}$ *be as constructed in* [\(5\)](#page-2-4)*. Suppose that the matrix* Γ *satisfies the Lower-RE condition with curvature* $\alpha > 0$ *and tolerance* $\tau > 0$ *,*

$$
\sqrt{d}\tau \le \min\left\{\frac{\alpha}{32\sqrt{d}}, \frac{\lambda}{4b_0}\right\} \tag{33}
$$

where d, b_0 and λ are as defined in [\(6\)](#page-2-5). Then for any d -sparse vectors $\beta^* \in \mathbf{R}^m$, such that $\|\beta^*\|_2 \le b_0$ and

$$
\left\|\widehat{\gamma} - \widehat{\Gamma}\beta^*\right\|_{\infty} \le \frac{1}{2}\lambda, \text{ the following bounds hold:}
$$
\n(34)

$$
\left\|\widehat{\beta} - \beta^*\right\|_2 \le \frac{20}{\alpha} \lambda \sqrt{d}, \quad \text{and} \quad \left\|\widehat{\beta} - \beta^*\right\|_1 \le \frac{80}{\alpha} \lambda d \tag{35}
$$

where $\widehat{\beta}$ *is an optimal solution to the Lasso-type estimator as in* [\(6\)](#page-2-5)*.*

We defer the proof of Theorem [8](#page-14-0) to Section [D,](#page-35-0) for clarity of presentation. In section [5.1,](#page-15-0) we provide two Lemmas [9](#page-15-1) and [10](#page-16-2) in checking the RE conditions as well condition [\(33\)](#page-14-1). One can then combine with Theorem [8,](#page-14-0) Lemmas [6,](#page-13-1) [9](#page-15-1) and [10](#page-16-2) to prove Theorem [2.](#page-8-0) In more details, Lemma [9](#page-15-1) checks the Lower and the Upper RE conditions on the modified gram matrix:

$$
\widehat{\Gamma}_A := X^T X - \widehat{\text{tr}}(B) I_m \tag{36}
$$

while Lemma [10](#page-16-2) checks condition [\(33\)](#page-14-1) as stated in Theorem [8](#page-14-0) for curvature α and tolerance τ as derived in Lemma [9.](#page-15-1) Finally Lemma [6](#page-13-1) ensures that [\(34\)](#page-14-2) holds with high probability for λ chosen as in [\(17\)](#page-8-4). We defer stating these lemmas in Section [5.1.](#page-15-0) The full proof of Theorem [2](#page-8-0) appears in Section [9.](#page-22-0)

For Theorem [3,](#page-9-0) our first goal is to show that the following holds with high probability

$$
\left\|\widehat{\gamma} - \widehat{\Gamma}\beta^*\right\|_{\infty} = \left\|\frac{1}{f}X^T(y - X\beta^*) + \frac{1}{f}\widehat{\text{tr}}(B)\beta^*\right\|_{\infty} \le \mu \left\|\beta^*\right\|_2 + \tau,
$$

where μ, τ are as chosen in [\(43\)](#page-16-3). This forms the basis for proving the ℓ_q convergence, where $q \in [1, 2]$, for the Conic Programming estimator [\(7\)](#page-3-0). This follows immediately from Lemma [6.](#page-13-1) More explicitly, we will state it in Lemma [11.](#page-16-1) Before we proceed, we first need to introduce some notation and definitions. Let $X_0 = Z_1 A^{1/2}$ be defined as in Definition [1.2.](#page-2-1) Let $k_0 = 1 + \lambda$. First we need to define the ℓ_q -sensitivity parameter for $\Psi := \frac{1}{f} X_0^T X_0$ following [Belloni et al.](#page-57-2) [\(2014](#page-57-2)):

$$
\kappa_q(d_0, k_0) = \min_{J: |J| \le d_0} \min_{\Delta \in \text{Cone}_J(k_0)} \frac{\|\Psi \Delta\|_{\infty}}{\|\Delta\|_q} \quad \text{where}
$$
\n(37)

$$
ConeJ(k0) = \{x \in \mathbf{R}^m \mid s.t. \quad ||x_{Jc}||_1 \le k_0 ||x_{J}||_1\}.
$$
 (38)

See also [Gautier and Tsybakov](#page-58-17) [\(2011\)](#page-58-17). Let (β, t) be the optimal solution to [\(7\)](#page-3-0) and denote by $v = \beta - \beta^*$. We will state the following auxiliary lemmas, the first of which is deterministic in nature. The two lemmas reflect the two geometrical constraints on the optimal solution to [\(7\)](#page-3-0). The optimal solution β satisfies:

1. *v* obeys the following cone constraint: $||v_{S}||_1 \le k_0 ||v_S||_1$ and $\hat{t} \le \frac{1}{\lambda}$ $\frac{1}{\lambda} ||v||_1 + ||\beta^*||_2.$ 2. $\|\Psi v\|_{\infty}$ is upper bounded by a quantity at the order of $O\left(\mu(\|\beta^*\|_2 + \|v\|_1) + \tau\right)$

Now combining Lemma 6 of [Belloni et al.](#page-57-2) [\(2014](#page-57-2)) and an earlier result of the two authors (cf. Theo-rem [25](#page-23-1) [Rudelson and Zhou](#page-59-14) [\(2013\)](#page-59-14)), we can show that the $\mathsf{RE}(2d_0,3(1+\lambda),A^{1/2})$ condition and the sample requirement as in [\(27\)](#page-12-1) are enough to ensure that the ℓ_q -sensitivity parameter satisfies the following lower bound for all $1 \le q \le 2$: for some contant c,

$$
\kappa_q(d_0, k_0) \geq cd_0^{-1/q} \text{ which ensures that for } v = \widehat{\beta} - \beta^*,
$$

$$
\|\Psi v\|_{\infty} \geq \kappa_q(d_0, k_0) \|v\|_q \geq cd_0^{-1/q} \|v\|_q \text{ where } \Psi = \frac{1}{f} X_0^T X_0.
$$
 (39)

Combining [\(39\)](#page-15-2) with Lemmas [11,](#page-16-1) [12](#page-16-4) and [13](#page-16-5) gives us both the lower and upper bounds on $\|\Psi v\|_{\infty}$, with the lower bound being $\kappa_q(d_0, k_0) ||v||_q$ and the upper bound as specified in Lemma [13.](#page-16-5) Following some algebraic manipulation, this yields the bound on the $||v||_q$ for all $1 \le q \le 2$. We state Lemmas [11](#page-16-1) to [13](#page-16-5) in Section [5.2](#page-16-6) while leaving the proof for Theorem [3](#page-9-0) in Section [10.](#page-23-0)

5.1 Additional technical results for Theorem [2](#page-8-0)

The main focus of the current section is to apply Theorem [8](#page-14-0) to show Theorem [2,](#page-8-0) which applies to the general subgaussian model as considered in the present work. We first state Lemma [9,](#page-15-1) which follows immediately from Corollary [19.](#page-18-2) First, we replace (A3) with (A3') which reveals some additional information regarding the constant hidden inside the $O(·)$ notation.

(A3') Suppose (A3) holds; moreover, for $D_2 = 2(||A||_2 + ||B||_2)$, $mf \ge 1024C_0^2D_2^2K^4 \log m/\lambda_{\min}^2(A)$ or equivalently,

$$
\frac{\lambda_{\min}(A)}{\|A\|_2 + \|B\|_2} > C_K \sqrt{\frac{\log m}{mf}}
$$
 for some large enough contain C_K .

Lemma 9. (Lower and Upper-RE conditions) *Suppose (A1), (A2) and (A3') hold. Denote by* $V :=$ $3eM_A^3/2$, where M_A is as defined in [\(13\)](#page-8-5). Let s_0 be as defined in [\(12\)](#page-7-1). Suppose that for some $c' > 0$,

$$
\frac{\text{tr}(B)}{\|B\|_2} \ge c' K^4 \frac{s_0}{\varepsilon^2} \log \left(\frac{3em}{s_0 \varepsilon} \right) \quad \text{where } \varepsilon = \frac{1}{2M_A}.
$$
\n(40)

Let A_0 *be the event that the modified gram matrix* $\widehat{\Gamma}_A$ *as defined in* [\(36\)](#page-14-3) *satisfies the Lower as well as Upper* RE *conditions with*

curvature
$$
\alpha = \frac{1}{2} \lambda_{\min}(A)
$$
, smoothness $\bar{\alpha} = 3 \lambda_{\max}(A)/2$,
and tolerance $\frac{512C^2 \varpi(s_0)^2 \log m}{\lambda_{\min}(A)} \leq \tau := \frac{\alpha}{s_0} \leq \frac{1024C^2 \varpi^2(s_0 + 1) \log m}{\lambda_{\min}(A)}$

for α , $\bar{\alpha}$ *and* τ *as defined in Definitions* [2.2](#page-5-1) *and* [2.3,](#page-6-1) *and* C , s_0 , $\varpi(s_0)$ *in* [\(12\)](#page-7-1)*. Then* $\mathbb{P}(\mathcal{A}_0) \geq 1$ − $4 \exp \left(-\frac{c_3 f}{M_A^2 \log m} \log \left(\frac{\mathcal{V} m \log m}{f}\right)\right) - 2 \exp \left(-\frac{4c_2 f}{M_A^2 K^4}\right)$ $(-6/m^3)$.

Lemma 10. *Suppose all conditions in Lemma [9](#page-15-1) hold. Suppose that* $s_0 \geq 3$ *and*

$$
d := |\text{supp}(\beta^*)| \le C_A \frac{f}{\log m} \left\{ c' D_\phi \wedge 2 \right\} \quad \text{where } C_A := \frac{1}{128M_A^2},
$$

$$
D_\phi = \left(\frac{K^2 M_\epsilon^2}{b_0^2} + K^4 \phi \right) \ge K^4 \phi \ge \phi
$$
 (41)

where $c', \phi, b_0, M_\epsilon$ and K are as defined in Theorem [2,](#page-8-0) where we assume that $\|\beta^*\|_2^2 \ge \phi b_0^2$ for some $0 <$ $\phi \leq 1$ *. Then the following condition holds*

$$
d \le \frac{s_0}{32} \bigwedge \left(\frac{s_0}{\alpha}\right)^2 \frac{\log m}{f} \left(\frac{\psi}{b_0}\right)^2 \tag{42}
$$

where ψ *is as defined in* [\(17\)](#page-8-4) *and* $\alpha = \lambda_{\min}(A)/2$.

We prove Lemmas [9](#page-15-1) and [10](#page-16-2) in Sections [D.1](#page-37-0) and [D.2](#page-38-0) respectively. **Remark 5.1.** *Clearly for* d, b_0, ϕ *as bounded in Theorem [2,](#page-8-0) we have by assumption* [\(15\)](#page-8-2) *the following upper and lower bound on* D_{ϕ} *:*

$$
2K^{4}\phi \ge D_{\phi} := \left(\frac{M_{\epsilon}^{2}K^{2}}{b_{0}^{2}} + K^{4}\phi\right) \ge K^{4}\phi.
$$

In this regime, the conditions on d *as in* [\(41\)](#page-16-0) *can be conveniently expressed as in* [\(16\)](#page-8-3)*.*

5.2 Technical lemmas for Theorem [3](#page-9-0)

We state the technical lemmas needed for proving Theorem [3.](#page-9-0) The proof for Lemma [12](#page-16-4) follows directly from that in [Belloni et al.](#page-57-2) [\(2014](#page-57-2)) in view of Lemma [11.](#page-16-1)

Lemma 11. *Suppose all conditions in Lemma* [6](#page-13-1) *hold. Then on event* \mathcal{B}_0 *as defined therein, the pair* (β, t) = $(\beta^*, \|\beta^*\|_2)$ belongs to the feasible set of the minimization problem [\(7\)](#page-3-0) with $r_{m,f} := C_0 K \sqrt{\frac{\log m}{f}}$ $\frac{gm}{f}$

$$
\mu \approx 2D_2Kr_{m,f} \quad \text{and} \quad \tau \approx D_0M_{\epsilon}r_{m,f} \tag{43}
$$

where $D_0 = (\sqrt{\tau_B} + \sqrt{a_{\text{max}}})$ *and* $D_2 = 2(||A||_2 + ||B||_2)$ *as in Theorem [3.](#page-9-0)* **Lemma 12.** *Let* $\mu, \tau > 0$ *be set. Suppose that the pair* $(\beta, t) = (\beta^*, \|\beta^*\|_2)$ *belongs to the feasible set of the minimization problem* [\(7\)](#page-3-0)*, for which* (β, t) *is an optimal solution. Denote by* $v = \beta - \beta^*$ *. Then*

$$
||v_{Sc}||_1 \le (1+\lambda) ||v_S||_1
$$
 and $\hat{t} \le \frac{1}{\lambda} ||v||_1 + ||\beta^*||_2$.

Lemma 13. *On event* $\mathcal{B}_0 \cap \mathcal{B}_{10}$ *,*

$$
\|\Psi v\|_{\infty} \leq \mu_1 \|\beta^*\|_2 + \mu_2 \|v\|_1 + \tau
$$

where $\mu_1 = 2\mu$, $\mu_2 = \mu(\frac{1}{\lambda} + 1)$ *and* $\tau' = 2\tau$ *for* μ, τ *as defined in* [\(43\)](#page-16-3)*.*

We prove Lemmas [11,](#page-16-1) [12](#page-16-4) and [13](#page-16-5) in Section [E.1.](#page-40-0)

5.3 Improved bounds for the Lasso-type estimator

We give an outline illustrating where the improvement for the lasso error bounds as stated in Theorem [4](#page-11-0) come from. We emphasize the impact of this improvement over sparsity parameter d_0 . The proof for Theorem [4](#page-11-0) follows exactly the same line of arguments as in Theorem [2](#page-8-0) except that we now use the improved bound on the error term $\left\|\hat{\gamma} - \hat{\Gamma}\beta^*\right\|_{\infty}$ given in Lemma [14](#page-17-2) instead of that in Lemma [6](#page-13-1) which is used in proving Theorems [2](#page-8-0) and [3.](#page-9-0) See Section [11](#page-24-0) for details, as well as the proof for Theorem [4](#page-11-0) and the following two lemmas.

Lemma 14. *Suppose all conditions in Lemma [6](#page-13-1) hold. Let* $D_0, D'_0, D_{\text{oracle}}$ *, and* $\tau_B^{+/2}$ $\tau_B^{+/2} := \tau_B^{1/2} + \frac{D_\text{oracle}}{\sqrt{m}}$ be *as defined in* [\(22\)](#page-10-1) *and* [\(23\)](#page-10-1)*. On event* B_0 *,*

$$
\left\|\hat{\gamma} - \widehat{\Gamma}\beta^*\right\|_{\infty} \leq \psi \sqrt{\frac{\log m}{f}}
$$
\n(44)

where $\psi:= C_0 K \left(D_0' \tau_B^{+/2} K \left\| \beta^* \right\|_2 + D_0 M_\epsilon \right)$. Then $\mathbb{P}\left(\mathcal{B}_0 \right) \geq 1-16/m^3$.

Moreover, we replace Lemma [10](#page-16-2) with Lemma [15,](#page-17-3) the proof of which follows from Lemma 10 with d now being bounded as in [\(25\)](#page-11-2) and ψ being redefined as immediately above in [\(44\)](#page-17-4).

Lemma 15. *Suppose all conditions in Lemma [9](#page-15-1) hold. Suppose that* [\(25\)](#page-11-2) *holds. Then* [\(42\)](#page-16-7) *holds with* ψ *as defined in Theorem [4](#page-11-0) and* $\alpha = \lambda_{\min}(A)/2$.

5.4 Improved bounds for the DS-type estimator

An "oracle" rate for the Conic programming estimator [\(7\)](#page-3-0) is defined as follows. Recall the following notation: $r_{m,f} = C_0 K \sqrt{\frac{\log m}{f}}$ $\frac{gm}{f}$. The trick is that we assume that we know the noise level in W by knowing $\tau_B := \text{tr}(B)/f$, then we can set

$$
\mu \approx D_0'(\tau_B^{1/2} + D_{\text{oracle}}/\sqrt{m}) K r_{m,f} \quad \text{while retaining} \quad \tau \approx D_0 M r_{m,f}
$$

in view of the improved error bounds over $\left\|\hat{\gamma} - \hat{\Gamma}\beta^*\right\|_{\infty}$ as given in Lemma [14.](#page-17-2) Without knowing this parameter, we could rely on the estimate from $\hat{\tau}_B$ as in [\(4\)](#page-2-3), which is what we do next. For a chosen parameter C_6 , we use $\hat{\tau}_{B}^{1/2} + C_6 K r_{m,m}^{1/2}$ to replace $\tau_{B}^{+/2}$ $\sigma_B^{+/2} := \tau_B^{1/2} + D_{\text{oracle}}/\sqrt{m}$ and set

$$
\mu \ge C_0 D_0'(\hat{\tau}_B^{1/2} + C_6 K r_{m,m}^{1/2}) K^2 \sqrt{\frac{\log m}{f}} \quad \text{where} \quad C_6 \ge D_{\text{oracle}},
$$
\n
$$
r_{m,m} = C_0 \sqrt{\frac{\log m}{mf}} > C_0 \frac{\sqrt{\log m}}{m} \text{ and } D_{\text{oracle}} := 2(||A||_2^{1/2} + ||B||_2^{1/2}).
$$

Notice that we know neither D'_0 nor D_{oracle} , where recall $D'_0 = \sqrt{||B||_2} + a_{\text{max}}^{1/2}$. However, assuming that we normalize the column norms of design matrix X to be roughly at the same scale, we have

 $D'_0 \approx 1$ while $D_{\text{oracle}}/\sqrt{m} = o(1)$ in case $||A||_2$, $||B||_2 \le M$

for some large enough constant M. This is crucial in deriving and putting the faster rates of convergence in estimating $\hat{\beta}$ and in predictive error $||Xv||_2^2$ when $\tau_B = o(1)$ in perspective, in view of Lemmas [16](#page-18-1) and [18.](#page-18-3) Lemma [16](#page-18-1) follows directly from Lemma [14.](#page-17-2)

Lemma 16. *Suppose all conditions in Lemma [14](#page-17-2) hold. Let* $D_0 = (\sqrt{\tau_B} + \sqrt{a_{\text{max}}}) \approx 1$ *under (A1). Then on event* \mathcal{B}_0 , the pair $(\beta, t) = (\beta^*, \|\beta^*\|_2)$ belongs to the feasible set Υ of the minimization problem [\(7\)](#page-3-0) with

$$
\mu \ge D_0' \tau_B^{+/2} K r_{m,f} \quad \text{and} \quad \tau \ge D_0 M_{\epsilon} r_{m,f}. \tag{45}
$$

where $\tau^{+/2}_B$ $\sigma_B^{+/2} := \tau_B^{1/2} + \frac{D_{\text{oracle}}}{\sqrt{m}}$ is as defined in [\(23\)](#page-10-1). **Lemma 17.** *On event* \mathcal{B}_6 *and* (A1), *the choice of* $\widetilde{\tau}_B^{1/2}$ $\lim_{B} \frac{A}{2}$ *as in* [\(30\)](#page-12-3) *satisfies for* $m \ge 16$ *and* $C_0 \ge 1$ *,*

$$
\tau_B^{+/2} \leq \tilde{\tau}_B^{1/2} \leq \tau_B^{1/2} + \frac{3}{2} C_6 K r_{mm}^{1/2} =: \tau_B^{\dagger/2} \tag{46}
$$

$$
\widetilde{\tau}_B \leq 2\tau_B + 3C_6^2 K^2 r_{mm} \asymp \tau_B^{\ddagger} \text{ and moreover } \widetilde{\tau}_B^{1/2} \tau_B^- \leq 1 \tag{47}
$$

We next state an updated result in Lemma [18.](#page-18-3)

Lemma 18. *On event* $\mathcal{B}_0 \cap \mathcal{B}_{10}$ *, the solution* β *to* [\(7\)](#page-3-0) *with* μ, τ *as in* [\(30\)](#page-12-3) *and* [\(29\)](#page-12-3)*, satisfies for* $v := \beta - \beta^*$

$$
\left\| \frac{1}{f} X_0^T X_0 v \right\|_{\infty} \le \mu_1 \left\| \beta^* \right\|_2 + \mu_2 \left\| v \right\|_1 + \tau'
$$

where $\mu_1 = 2\mu$, $\mu_2 = 2\mu(1 + \frac{1}{2\lambda})$ *and* $\tau' = 2\tau$.

6 Lower and Upper RE conditions

The goal of this section is to show that for Δ defined in [\(51\)](#page-18-4), the presumption in Lemmas [32](#page-46-0) and [34](#page-48-0) as restated in [\(48\)](#page-18-5) holds with high probability (cf Theorem [20\)](#page-18-6). We first state a deterministic result showing that the Lower and Upper RE conditions hold for Γ_A under condition [\(48\)](#page-18-5) in Corollary [19.](#page-18-2) This allows us to prove Lemma [9](#page-15-1) in Sections [D.1.](#page-37-0) See Sections [G](#page-45-0) and [H,](#page-49-0) where we show that Corollary [19](#page-18-2) follows immediately from the geometric analysis result as stated in Lemma [34.](#page-48-0)

Corollary 19. Let $1/8 > \delta > 0$. Let $1 \leq \zeta < m/2$. Let $A_{m \times m}$ be a symmetric positive semidefinite *covariance matrice. Let* $\widehat{\Gamma}_A$ *be an* $m \times m$ *symmetric matrix and* $\Delta = \widehat{\Gamma}_A - A$ *. Let* $E = \bigcup_{|J| \leq \zeta} E_J$ *, where* $E_J = \text{span}\{e_j : j \in J\}$. Suppose that $\forall u, v \in E \cap S^{m-1}$

$$
\left|u^T \Delta v\right| \le \delta \le \frac{1}{8} \lambda_{\min}(A). \tag{48}
$$

Then the Lower and Upper RE *conditions holds: for all* $v \in \mathbb{R}^m$,

$$
v^T \widehat{\Gamma}_A v \ge \frac{1}{2} \lambda_{\min}(A) \|v\|_2^2 - \frac{\lambda_{\min}(A)}{2\zeta} \|v\|_1^2 \tag{49}
$$

$$
v^T \widehat{\Gamma}_A v \leq \frac{3}{2} \lambda_{\max}(A) \|v\|_2^2 + \frac{\lambda_{\min}(A)}{2\zeta} \|v\|_1^2. \tag{50}
$$

Theorem 20. *Let* $A_{m \times m}$, $B_{f \times f}$ *be symmetric positive definite covariance matrices. Let* $E = \bigcup_{|J| \leq \zeta} E_J$ *for* $1 \leq \zeta < m/2$. Let Z, X be $f \times m$ random matrices defined as in Theorem [2.](#page-8-0) Let $\hat{\tau}_B$ be defined as in [\(4\)](#page-2-3). *Let*

$$
\Delta := \widehat{\Gamma}_A - A := \frac{1}{f} X^T X - \widehat{\tau}_B I_m - A. \tag{51}
$$

Suppose that for some absolute constant $c' > 0$ and $0 < \varepsilon \leq \frac{1}{C}$

$$
\frac{\text{tr}(B)}{\|B\|_2} \ge \left(c' K^4 \frac{\zeta}{\varepsilon^2} \log \left(\frac{3em}{\zeta \varepsilon} \right) \right) \bigvee \log m \tag{52}
$$

where $C = C_0 / \sqrt{c'}$ for C_0 as chosen to satisfy [\(86\)](#page-39-0).

Then with probability at least $1 - 4 \exp \left(-c_2 \varepsilon^2 \frac{\text{tr}(B)}{K^4 \|B\|_2}\right)$ $-2 \exp \left(-c_2 \varepsilon^2 \frac{f}{K^4}\right)$ $-6/m³$ *, where* $c_2 ≥ 2$ *, we have for all* $u, v \in E \cap S^{m-1}$ *and* $\varpi(\zeta) = \tau_B + \rho_{\max}(\zeta, A)$ *, and* $D_1 \leq \frac{\|A\|_F}{\sqrt{m}} + \frac{\|B\|_F}{\sqrt{f}}$ *,*

$$
|u^T \Delta v| \le 8C \varpi(\zeta)\varepsilon + 4C_0 D_1 K^2 \sqrt{\frac{\log m}{mf}}.
$$

We prove Theorem [20](#page-18-6) in Section [I.](#page-49-1) As a corollary of Theorem [20,](#page-18-6) we will state Corollary [23](#page-20-1) in Section [7.](#page-19-0)

7 Concentration bounds for error-corrected gram matrices

In this section, we show an upper bound on the operator norm convergence as well as an isometry property for estimating B using the corrected gram matrix $\widetilde{B} := \frac{1}{m}(XX^T - \text{tr}(A)I_f)$. Theorem [21](#page-19-1) and Corollary [22](#page-19-2) state that for the matrix $B \succ 0$ with the smaller dimension, \tilde{B} tends to stay positive definite after this error correction step with an overwhelming probability, where we rely on f being dominated by the effective rank of the positive definite matrix A. When we subtract a diagonal matrix $\tau_B I_m$ from the gram matrix $\frac{1}{f} X^T X$ to form an estimator, we clearly introduce a large number of negative eigenvalues when $f \ll m$. This in general is a bad idea. However, the sparse eigenvalues for A can stay pretty close to those of A as we will show in Corollary [23.](#page-20-1)

Theorem 21. Let $\varepsilon > 0$. Let X be defined as in Definition [1.2.](#page-2-1) Suppose that for some $c' > 0$ and $0 < \varepsilon < 1/2$,

$$
\frac{\text{tr}(A)}{\|A\|_2} \ge c' f K^4 \frac{\log(3/\varepsilon)}{\varepsilon^2}.
$$
\n(53)

Then with probability at least $1 - 2 \exp\left(-c\epsilon^2 \frac{m}{K^4}\right) - 4 \exp\left(-c_5 \epsilon^2 \frac{\text{tr}(A)}{K^4 ||A||_2}\right)$ *,*

$$
\left\| \frac{1}{m} X X^T - \frac{\text{tr}(A) I_f}{m} - B \right\|_2 \le C_2 \varepsilon (\tau_A + \|B\|_2)
$$

where C_2 , c_5 are absolute constants depending on c' , C, where $C > 4 \max(\frac{1}{cc'}, \frac{1}{\sqrt{4c'}})$ $\frac{1}{4cc'}$) is a large enough *constant.*

Corollary 22. *Suppose all conditions in Theorem [21](#page-19-1) hold. Suppose*

$$
\frac{\text{tr}(A)}{\|A\|_2} \ge c' f K^4 \frac{C_3^2}{\delta^2} \log\left(\frac{3C_3}{\delta}\right). \tag{54}
$$

where $C_3=C_2\left(\frac{\tau_A}{\lambda_{\min}(B)}\vee 1\right)$ for C_2 as in Theorem [21.](#page-19-1) Then with the probability as stated in Theorem [21,](#page-19-1)

$$
(1+2\delta)B \succ \frac{XX^T}{m} - \frac{\text{tr}(A)I_f}{m} \succ (1-2\delta)B \succ 0
$$

where for the last inequality to hold, we assume that $\lambda_{\min}(B) > 0$ *.*

Next we show a large deviation bound on the sparse eigenvalues of the error corrected \widetilde{A} : $\widetilde{A} := \frac{1}{f} X^T X$ $\tau_B I_m$.

Corollary 23. *Let* X *be defined as in Definition [1.2.](#page-2-1) Let* $\widetilde{A} := \frac{1}{f} X^T X - \tau_B I_m$ *. Suppose*

$$
\frac{\text{tr}(B)}{\|B\|_2} \ge c' k K^4 \frac{\log(3em/k\varepsilon)}{\varepsilon^2}.
$$
\n(55)

Then with probability at least $1 - 2 \exp(-c_4 \varepsilon^2 \frac{f}{K^4}) - 4 \exp(-c_4 \varepsilon^2 \frac{\text{tr}(B)}{K^4 ||B||_2}),$

$$
\rho_{\max}(k,\widetilde{A}) \le \rho_{\max}(k,A)(1+10\varepsilon) + C_4\varepsilon\tau_B
$$

where C_4 is an absolute constant. Moreover, suppose for $C_5 = C_4 \left(\frac{\tau_B}{\rho_{\min}(k,A)} \vee 1 \right)$

$$
\frac{\text{tr}(B)}{\|B\|_2} \ge c' k K^4 \frac{C_5^2}{\delta^2} \log(\frac{12C_5em}{k\delta}).\tag{56}
$$

Then with the probability as stated immediately above, we have

$$
\rho_{\min}(k, A) \ge \rho_{\min}(k, A)(1 - 2\delta).
$$

We prove Theorem [21](#page-19-1) in Section [J.](#page-51-0) We also prove the concentration of measure bounds on error-corrected gram matrices in Corollaries [22](#page-19-2) and [23](#page-20-1) in Sections [J.1](#page-53-0) and [J.2](#page-54-0) respectively.

8 Proof of Lemma [1](#page-6-0)

We define $Cone(d_0, k_0)$, where $0 < d_0 < m$ and k_0 is a positive number, as the set of vectors in \mathbb{R}^m which satisfy the following cone constraint:

Cone
$$
(d_0, k_0) = \{x \in \mathbb{R}^m \mid \exists I \in \{1, ..., p\}, |I| = d_0 \text{ s.t. } ||x_{I^c}||_1 \le k_0 ||x_I||_1\}.
$$

For each vector $x \in \mathbb{R}^p$, let T_0 denote the locations of the s_0 largest coefficients of x in absolute values. The following elementary estimate [Rudelson and Zhou](#page-59-14) [\(2013](#page-59-14)) will be used in conjunction with the RE condition. **Lemma 24.** *For each vector* $x \in \text{Cone}(s_0, k_0)$, let T_0 *denotes the locations of the* s_0 *largest coefficients of* x *in absolute values. Then*

$$
||x_{T_0}||_2 \ge \frac{||x||_2}{\sqrt{1+k_0}}.\tag{57}
$$

Proof of Lemma [1.](#page-6-0) Part I: Suppose that the Lower-RE condition holds for $\Gamma := A^T A$. Let $x \in$ $Cone(s_0, k_0)$. Then

$$
||x||_1 \le (1 + k_0) ||x_{T_0}||_1 \le (1 + k_0) \sqrt{s_0} ||x_{T_0}||_2.
$$

Thus for $x \in \text{Cone}(s_0, k_0) \cap S^{p-1}$ and $\tau (1 + k_0)^2 s_0 \leq \alpha/2$, we have

$$
||Ax||_2 = (x^T A^T A x)^{1/2} \ge (\alpha ||x||_2^2 - \tau ||x||_1^2)^{1/2}
$$

\n
$$
\ge (\alpha ||x||_2^2 - \tau (1 + k_0)^2 s_0 ||x_{T_0}||_2^2)^{1/2}
$$

\n
$$
\ge (\alpha - \tau (1 + k_0)^2 s_0)^{1/2} \ge \sqrt{\frac{\alpha}{2}}.
$$

Thus the $RE(s_0, k_0, A)$ condition holds with

$$
\frac{1}{K(s_0, k_0, A)} \ := \ \min_{x \in \text{Cone}(s_0, k_0)} \frac{\|Ax\|_2}{\|x_{T_0}\|_2} \ge \sqrt{\frac{\alpha}{2}}
$$

where we use the fact that for any $J \in \{1, ..., p\}$ such that $|J| \leq s_0$, $||x_J||_2 \leq ||x_{T_0}||_2$. We now show the other direction.

Part II. Assume that $RE(4R^2, 2R - 1, A)$ holds for some integer $R > 1$. Assume that for some $R > 1$

$$
||x||_1 \le R ||x||_2.
$$

Let $(x_i^*)_{i=1}^p$ be non-increasing arrangement of $(|x_i|)_{i=1}^p$. Then

$$
||x||_1 \leq R \left(\sum_{j=1}^s (x_j^*)^2 + \sum_{j=s+1}^\infty \left(\frac{||x||_1}{j} \right)^2 \right)^{1/2}
$$

$$
\leq R \left(||x_J^*||_2^2 + ||x||_1^2 \frac{1}{s} \right)^{1/2} \leq R \left(||x_J^*||_2 + ||x||_1 \frac{1}{\sqrt{s}} \right)
$$

where $J := \{1, \ldots, s\}$. Choose $s = 4R^2$. Then

$$
||x||_1 \leq R ||x^*_J||_2 + \frac{1}{2} ||x||_1
$$

Thus we have

$$
||x||_1 \le 2R ||x_J^*||_2 \le 2R ||x_J^*||_1 \text{ and hence}
$$
\n(58)
\n
$$
||x_{J^c}^*||_1 \le (2R-1) ||x_J^*||_1.
$$
\n(59)

.

Then $x \in \text{Cone}(4R^2, 2R - 1)$. Then for all $x \in S^{p-1}$ such that $||x||_1 \le R ||x||_2$, we have for $k_0 = 2R - 1$ and $s_0 := 4R^2$,

$$
x^{T}\Gamma x \ge \frac{\|x_{T_0}\|_2^2}{K^2(s_0, k_0, A)} \ge \frac{\|x\|_2^2}{\sqrt{s_0}K^2(s_0, k_0, A)} =: \alpha \|x\|_2^2
$$

where we use the fact that $(1 + k_0) \|x_{T_0}\|_2^2 \ge \|x\|_2^2$ $^{2}_{2}$ by Lemma [24](#page-20-2) with x_{T_0} as defined therein. Otherwise, suppose that $||x||_1 \ge R ||x||_2$. Then for a given $\tau > 0$,

$$
\alpha \|x\|_2^2 - \tau \|x\|_1^2 \le \left(\frac{1}{\sqrt{s_0} K^2(s_0, k_0, A)} - \tau R^2\right) \|x\|_2^2. \tag{60}
$$

Thus we have by the choice of τ as in [\(29\)](#page-12-3) and [\(60\)](#page-21-0)

$$
x^{T} \Gamma x \ge \lambda_{\min}(\Gamma) \|x\|_{2}^{2} \ge \left(\frac{1}{\sqrt{s_{0}} K^{2}(s_{0}, k_{0}, A)} - \tau R^{2}) \|x\|_{2}^{2} \right)
$$

$$
\ge \alpha \|x\|_{2}^{2} - \tau \|x\|_{1}^{2}.
$$

The Lemma thus holds. \square

9 Proof of Theorem [2](#page-8-0)

First we note that it is sufficient to have [\(14\)](#page-8-1) in order for [\(40\)](#page-15-3) to hold. (14) guarantees that for $V =$ $3eM_A^3/2$

$$
r(B) := \frac{\text{tr}(B)}{\|B\|_2} \ge 16c'K^4 \frac{f}{\log m} \log \frac{\mathcal{V}m \log m}{f}
$$

\n
$$
\ge 16c'K^4 \frac{f}{\log m} \log \left(\frac{3emM_A^3 \log m}{2f}\right)
$$

\n
$$
= c'K^4 \frac{1}{\varepsilon^2} \frac{4}{M_A^2} \frac{f}{\log m} \log \left(\frac{6emM_A}{\frac{4}{M_A^2}(f/\log m)}\right)
$$

\n
$$
\ge c'K^4 \frac{1}{\varepsilon^2} s_0 \log \left(\frac{6emM_A}{s_0}\right) = c'K^4 \frac{s_0}{\varepsilon^2} \log \left(\frac{3em}{s_0 \varepsilon}\right)
$$
(61)

where $\varepsilon = \frac{1}{2M}$ $\frac{1}{2M_A} \leq \frac{1}{128}$ $\frac{1}{128C}$, and the last inequality holds given that $k \log(c m/k)$ on the RHS of [\(61\)](#page-22-1) is a monotonically increasing function of k , and

$$
s_0 \le \frac{4f}{M_A^2 \log m} \text{ and } M_A = \frac{64C(\rho_{\text{max}}(s_0, A) + \tau_B)}{\lambda_{\text{min}}(A)} \ge 64C.
$$

Next we check that the choice of d as in [\(16\)](#page-8-3) ensures that [\(41\)](#page-16-0) holds. Indeed, for $c'K^4 \leq 1$, we have

$$
d \leq C_A(c'K^4 \wedge 1) \frac{\phi f}{\log m} \leq C_A\left(c'D_{\phi} \wedge 1\right) \frac{f}{\log m}.
$$

By Lemma [9,](#page-15-1) we have on event \mathcal{A}_0 , the modified gram matrix $\widehat{\Gamma}_A := \frac{1}{f}(X^T X - \widehat{\text{tr}}(B)I_m)$ satisfies the Lower RE conditions with

curvature
$$
\alpha = \frac{1}{2} \lambda_{\min}(A)
$$
 and tolerance $\tau = \frac{\lambda_{\min}(A)}{2s_0} = \frac{\alpha}{s_0}.$ (62)

Theorem [2](#page-8-0) follows from Theorem [8,](#page-14-0) so long as we can show that condition [\(33\)](#page-14-1) holds for $\lambda \ge 4\psi \sqrt{\frac{\log m}{f}}$ f where the parameter ψ is as defined [\(17\)](#page-8-4), and α and $\tau = \frac{\alpha}{s_0}$ are as defined immediately above. Combining [\(62\)](#page-22-2) and [\(33\)](#page-14-1), we need to show [\(42\)](#page-16-7) holds. This is precisely the content of Lemma [10.](#page-16-2) This is the end of the proof for Theorem [2](#page-8-0) \Box

10 Proof of Theorem [3](#page-9-0)

For the set $Cone_J(k_0)$ as in (57),

$$
\kappa_{\mathsf{RE}}(d_0, k_0) \ := \ \min_{J: |J| \leq d_0} \min_{\Delta \in \operatorname{Cone}_{J}(k_0)} \frac{|\Delta^T \Psi \Delta|}{\|\Delta_J\|_2^2} = \left(\frac{1}{K(d_0, k_0, (1/\sqrt{f})Z_1 A^{1/2})}\right)^2.
$$

Recall the following Theorem [25](#page-23-1) from [Rudelson and Zhou](#page-59-14) [\(2013\)](#page-59-14).

Theorem 25. [Rudelson and Zhou](#page-59-14) [\(2013](#page-59-14)) *Set* $0 < \delta < 1$ *,* $k_0 > 0$ *, and* $0 < d_0 < p$ *. Let* $A^{1/2}$ *be an* $m \times m$ *matrix satisfying* $RE(d_0, 3k_0, A^{1/2})$ *condition as in Definition [2.1.](#page-5-0) Let d be as defined in* [\(63\)](#page-23-2)

$$
d = d_0 + d_0 \max_j \left\| A^{1/2} e_j \right\|_2^2 \frac{16K^2 (d_0, 3k_0, A^{1/2}) (3k_0)^2 (3k_0 + 1)}{\delta^2}.
$$
 (63)

Let Ψ *be an* $n \times m$ *matrix whose rows are independent isotropic* ψ_2 *random vectors in* \mathbb{R}^m *with constant* α *. Suppose the sample size satisfies*

$$
n \ge \frac{2000d\alpha^4}{\delta^2} \log \left(\frac{60em}{d\delta} \right). \tag{64}
$$

Then with probability at least $1 - 2 \exp(-\delta^2 n/2000\alpha^4)$, $\mathsf{RE}(d_0, k_0, (1/\sqrt{n})\Psi A^{1/2})$ *condition holds for matrix* $(1/\sqrt{n}) \Psi A$ *with*

$$
0 < K(d_0, k_0, \left(1/\sqrt{n}\right) \Psi A^{1/2}) \le \frac{K(d_0, k_0, A^{1/2})}{1 - \delta}.\tag{65}
$$

Proof of Theorem [3.](#page-9-0) Suppose $RE(2d_0, 3k_0, A^{1/2})$ holds. Then for d as defined in [\(28\)](#page-12-1) and $f =$ $\Omega(dK^4\log(m/d))$, we have with probability at least $1-2\exp(\delta^2 f/2000K^4)$, the RE $(2d_0, k_0, \frac{1}{\sqrt{3}})$ $\bar{z}^{Z_1 A^{1/2}}$ condition holds with

$$
\kappa_{\sf RE}(2d_0, k_0) = \left(\frac{1}{K(2d_0, k_0, (1/\sqrt{f})Z_1A^{1/2})}\right)^2 \ge \left(\frac{1}{2K(2d_0, k_0, A^{1/2})}\right)^2
$$

by Theorem [25.](#page-23-1)

The rest of the proof follows from [Belloni et al.](#page-57-2) [\(2014](#page-57-2)) Theorem 1 and thus we only provide a sketch. In more details, in view of the lemmas shown in Section [5,](#page-13-0) we need

$$
\kappa_q(d_0, k_0) \geq cd_0^{-1/q}
$$

to hold for some constant c for $\Psi := \frac{1}{f} X_0^T X_0$. It is shown in Appendix C in [Belloni et al.](#page-57-2) [\(2014\)](#page-57-2) that under the RE $(2d_0, k_0, \frac{1}{\sqrt{2}})$ $\frac{f}{f}Z_1A^{1/2}$) condition, for any $d_0 \leq m/2$ and $1 \leq q \leq 2$, we have

$$
\kappa_1(d_0, k_0) \geq cd_0^{-1} \kappa_{\mathsf{RE}}(d_0, k_0), \n\kappa_q(d_0, k_0) \geq c(q) d_0^{-1/q} \kappa_{\mathsf{RE}}(2d_0, k_0)
$$
\n(66)

where $c(q) > 0$ depends on k_0 and q. The theorem is thus proved following exactly the same line of arguments as in the proof of Theorem 1 in [Belloni et al.](#page-57-2) [\(2014](#page-57-2)) in view of the ℓ_q sensitivity condition

derived immediately above, in view of Lemmas [11,](#page-16-1) [12](#page-16-4) and [13.](#page-16-5) Indeed, we have for $v := \beta - \beta^*$, we have by definition of ℓ_q sensitivity as in [\(37\)](#page-14-4)

$$
c(q)d_0^{-1/q}\kappa_{\mathsf{RE}}(2d_0, k_0) \|v\|_q \le \kappa_q(d_0, k_0) \|v\|_q \le \left\|\frac{1}{f}X_0^T X_0 v\right\|_\infty
$$

\n
$$
\le \mu_1 \|\beta^*\|_2 + \mu_2 \|v\|_1 + \tau
$$

\n
$$
\le \mu_1 \|\beta^*\|_2 + \mu_2(2+\lambda) \|v_S\|_1 + \tau
$$

\n
$$
\le \mu_1 \|\beta^*\|_2 + \mu_2(2+\lambda)d_0^{1-1/q} \|v_S\|_q + \tau
$$

\n
$$
\le \mu_1 \|\beta^*\|_2 + \mu_2(2+\lambda)d_0^{1-1/q} \|v\|_q + \tau.
$$
 (67)

Thus we have for $d_0 = c_0 \sqrt{f / \log m}$ where c_0 is sufficiently small,

$$
d_0^{-1/q}(c(q)\kappa_{\mathsf{RE}}(2d_0, k_0) - \mu_2(2+\lambda)d_0) \|v\|_q \le \mu_1 \|\beta^*\|_2 + \tau
$$

\nhence
\n
$$
\|v\|_q \le C(4D_2r_{m,f}K \|\beta^*\|_2 + 2D_0M_{\epsilon}r_{m,f})d_0^{1/q}
$$

\n
$$
\le 4CD_2r_{m,f}(K \|\beta^*\|_2 + M_{\epsilon})d_0^{1/q}
$$

for some constant $C = 1/(c(q)\kappa_{RE}(2d_0, k_0) - \mu_2(2+\lambda)d_0) \ge 1/(2c(q)\kappa_{RE}(2d_0, k_0))$ given that

$$
\mu_2(2+\lambda)d_0 = 2D_2Kr_{m,f}(\frac{1}{\lambda}+1)(2+\lambda)c_0\sqrt{f/\log m} = 2c_0C_0D_2K^2(2+\lambda)(\frac{1}{\lambda}+1)
$$

is sufficiently small and thus (21) holds. The prediction error bound follows exactly the same line of arguments as in [Belloni et al.](#page-57-2) [\(2014](#page-57-2)) which we omit here. See proof of Theorem [5](#page-11-1) in Section [F](#page-43-0) for details. \Box

11 Proof of Theorem [4](#page-11-0)

The proof is identical to the proof of Theorem [2](#page-8-0) up till (62) , except that we replace the condition on d as in the theorem statement by (25) : that is,

$$
d := |\text{supp}(\beta^*)| \le C_A \frac{f}{\log m} \left\{ c'C_{\phi} \wedge 2 \right\} \quad \text{where } C_A := \frac{1}{128M_A^2},
$$

$$
C_{\phi} := \frac{\|B\|_2 + a_{\max}}{D^2} \left(\frac{K^2 M_{\epsilon}^2}{b_0^2} + \tau_B^+ K^4 \phi \right) \ge \frac{\|B\|_2 + a_{\max}}{D^2} \tau_B^+
$$

where $c', \phi, b_0, M_\epsilon$ and K are as defined in Theorem [2,](#page-8-0) where we assume that $b_0^2 \ge ||\beta^*||_2^2 \ge \phi b_0^2$ for some $0 <$ $\phi \leq 1$. Theorem [4](#page-11-0) follows from Theorem [8,](#page-14-0) so long as we can show that condition [\(33\)](#page-14-1) holds for $\lambda \ge 2\psi\sqrt{\frac{\log m}{f}}$ where the parameter ψ is as defined [\(44\)](#page-17-4), and α and $\tau = \frac{\alpha}{s_0}$ $\frac{\alpha}{s_0}$ are as defined in [\(62\)](#page-22-2). Combining (62) and (33) , we need to show (42) holds. This is precisely the content of Lemma [15.](#page-17-3) This is the end of the proof for Theorem [4.](#page-11-0) \Box

12 Proof of Theorem [5](#page-11-1)

Throughout this proof, we assume that $\mathcal{B}_0 \cap \mathcal{B}_{10}$ holds. The rest of the proof follows that of Theorem [3,](#page-9-0) except for the last part. Let μ_1, μ_2, τ be as defined in Lemma [13.](#page-16-5) We have for $\mu_2 := 2\mu(1 + \frac{1}{2\lambda})$ where $\mu = D'_0 K r_{m,f} \tilde{\tau}_B^{1/2}$ $\int_{B}^{1/2}$, and $d_0 = c_0 \tau_B^- \sqrt{f/\log m}$,

$$
\mu_2(2+\lambda)d_0 = 2C_0D'_0K^2\tilde{\tau}_B^{1/2}(\frac{1}{2\lambda}+1)(2+\lambda)c_0\tau_B^-
$$
\n
$$
\leq 2c_0C_0D'_0K^2(2+\lambda)(\frac{1}{2\lambda}+1) \leq \frac{1}{2}c(q)\kappa_{\text{RE}}(2d_0,k_0)
$$
\n(68)

which holds when c_0 is sufficiently small, where by [\(47\)](#page-18-7) $\tau_B^- \tilde{\tau}_B^{1/2} \le 1$. Hence

$$
\mu_2 d_0 \le \frac{c(q)\kappa_{\sf RE}(2d_0, k_0)}{2(2+\lambda)}
$$

Thus for c_0 sufficiently small, $\mu_1 = 2\mu$, by [\(66\)](#page-23-3), [\(68\)](#page-25-1), [\(67\)](#page-24-1) and [\(46\)](#page-18-7),

$$
d_0^{-1/q} \frac{1}{2} (c(q)\kappa_{\mathsf{RE}}(2d_0, k_0)) \|v\|_q
$$

= $d_0^{-1/q} (c(q)\kappa_{\mathsf{RE}}(2d_0, k_0) - \mu_2(2+\lambda)d_0) \|v\|_q$

$$
\leq (\kappa_q(d_0, k_0) - \mu_2(2+\lambda)d_0^{1-1/q}) \|v\|_q \leq \mu_1 \|\beta^*\|_2 + \tau
$$

$$
\leq 2D'_0 r_{m,f} K^2 ((\tau_B^{1/2} + (3/2)C_6Kr_{m,m}^{1/2}) \|\beta^*\|_2 + M_{\epsilon}/K)
$$
 (69)

and thus [\(31\)](#page-12-4) holds, following the proof in Theorem [3.](#page-9-0) The prediction error bound follows exactly the same line of arguments as in [Belloni et al.](#page-57-2) [\(2014\)](#page-57-2), which we now include for the sake completeness. Following (31) , we have by (69) ,

$$
||v||_1 \leq C_{11}d_0(\mu_1 ||\beta^*||_2 + \tau) \text{ where } C_{11} = 2/(c(q)\kappa_{\mathsf{RE}}(2d_0, k_0))
$$

and hence $\mu_2 ||v||_1 \leq C_{11}\mu_2 d_0(\mu_1 ||\beta^*||_2 + \tau)$
 $\leq C_{11} \frac{1}{2(2+\lambda)} (c(q)\kappa_{\mathsf{RE}}(2d_0, k_0)) (\mu_1 ||\beta^*||_2 + \tau)$
 $= \frac{1}{2+\lambda}(\mu_1 ||\beta^*||_2 + \tau)$

Thus we have by (69) , the bounds immediately above, and (47)

$$
\frac{1}{f} \| X(\hat{\beta} - \beta^*) \|_2^2 \leq \|v\|_1 \| \frac{1}{f} X_0^T X_0 v \|_{\infty} \n\leq C_{11} d_0(\mu_1 \| \beta^* \|_2 + \tau) (\mu_1 \| \beta^* \|_2 + \mu_2 \|v\|_1 + 2\tau) \n\leq C_{11} d_0(\mu_1 \| \beta^* \|_2 + \tau) (1 + \frac{1}{2 + \lambda}) (\mu_1 \| \beta^* \|_2 + 2\tau) \n= C'(D'_0)^2 K^4 d_0 \frac{\log m}{f} \left(\tilde{\tau}_B^{1/2} \| \beta^* \|_2 + \frac{M_{\epsilon}}{K} \right)^2 \n\leq C'' (\|B\|_2 + a_{\text{max}}) K^2 d_0 \frac{\log m}{f} \left((2\tau_B + 3C_6^2 K^2 r_{m,m}) K^2 \| \beta^* \|_2^2 + M_{\epsilon}^2 \right)
$$

where $(D'_0)^2 \leq 2||B||_2 + 2a_{\text{max}}$. The theorem is thus proved. \square

13 Conclusion

In view of the main Theorems [2](#page-8-0) and [3,](#page-9-0) at this point, we do not really think one estimator is preferable to the other. While the rates we obtain for both estimators are at the same order for $q = 1, 2$, the conditions under which these rates are obtained are somewhat different. Lasso estimator allows large values of sparsity, while Conic-programming estimator conceptually is more adaptive by not fixing an upper bound on $\|\beta^*\|_2$ a priori, the cost of which seems to be a more stringent requirement on the sparsity level. The lasso-type procedure can recover a sparse model using $O(\log m)$ number of measurements per nonzero component despite the measurement error in X and the stochastic noise ϵ while the Dantzig selector-type allows only $d \leq \sqrt{f / \log m}$ to achieve the error rate at the same order as the Lasso-type estimator.

However, we show in Theorem [5](#page-11-1) in Section [5.4](#page-17-1) that this restriction on the sparsity can be relaxed for the Conic programming estimator [\(7\)](#page-3-0), when we make a different choice for the parameter μ based on a more refined analysis. Eventually, as $\tau_B \to 0$, this relaxation on d as in [\(32\)](#page-12-2) enables the Conic Programming estimator to achieve bounds which are essentially identical to the Dantzig Selector when the design matrix X_0 is a subgaussian random matrix satisfying the Restricted Eigenvalue conditions; See for exam-ple Candès and Tao [\(2007\)](#page-57-3); [Bickel et al.](#page-57-8) [\(2009](#page-57-8)); [Rudelson and Zhou](#page-59-14) [\(2013\)](#page-59-14). For the Lasso estimator, when we require that the stochastic error ϵ in the response variable y as in [\(1a\)](#page-1-0) does not converge to 0 as quickly as the measurement error W in [\(1b\)](#page-1-0) does, then the sparsity constraint becomes essentially unchanged as $\tau_B \to 0$. These tradeoffs are somehow different from the behavior of the Conic programming estimator versus the Lasso estimator; however, we believe the differences are minor.

We now state a slightly sharper bound than those in Lemma [14](#page-17-2) which provides a significant improvement on the error bounds in case $\tau_B = o(1)$ while $||A||_2 \geq 1$ for the Lasso-type estimator in [\(6\)](#page-2-5) as well as the Conic programming estimator [\(7\)](#page-3-0). Recall $D'_0 := \sqrt{||B||_2} + a_{\text{max}}^{1/2}$. By [\(74\)](#page-32-0),

$$
\left\|\hat{\gamma} - \hat{\Gamma}\beta^*\right\|_{\infty} \leq D'_0 K \tau_B^{1/2} \left\|\beta^*\right\|_2 r_{m,f} + \frac{2D_1 K}{\sqrt{m}} \left\|\beta^*\right\|_{\infty} r_{m,f} + D_0 M_{\epsilon} r_{m,f}
$$

When $\tau_B \to 0$, we have for $D_0 = \sqrt{\tau_B} + a_{\text{max}}^{1/2} \to a_{\text{max}}^{1/2}$

$$
\left\|\widehat{\gamma}-\widehat{\Gamma}\beta^*\right\|_{\infty} \;\; = \;\; O\left(D_1K\frac{1}{\sqrt{m}}\left\|\beta^*\right\|_{\infty}+D_0KM_{\epsilon}\right)K\sqrt{\frac{\log m}{f}}
$$

where $D_1 = \frac{\|A\|_F}{\sqrt{m}} + \frac{\|B\|_F}{\sqrt{f}} \to \|A\|_2^{1/2}$ $\frac{1}{2}$ ^{1/2} under (A1), given that $||B||_F / \sqrt{f} \le \tau_B^{1/2}$ $\mathbb{E}_{B}^{1/2}$ $||B||_2^{1/2} \rightarrow 0$, and the first term inside the bracket comes from the estimation error in $\hat{tr}(B)/f$, which can be made go away if we were to assume that $tr(B)$ is also known. In this case, the error term involving $\|\beta^*\|_2$ in [\(17\)](#page-8-4) vanishes, and we only need to set

$$
\lambda \ge 2\psi \sqrt{\frac{\log m}{f}} \quad \text{where} \quad \psi \asymp D_0 K M_{\epsilon} + ||A||_2^{1/2} K^2 m^{-1/2} ||\beta^*||_{\infty} \,. \tag{70}
$$

Moreover, suppose that $tr(B)$ is given, then one can drop the second term in ψ as in [\(70\)](#page-26-1) and hence recover the lasso bound when the design matrix X is assumed to be free of measurement errors.

Finally, we note that the bounds corresponding to the Upper RE condition as stated in Corollary [19,](#page-18-2) Theorem [20](#page-18-6) and Lemma [9](#page-15-1) are not needed for Theorem [2.](#page-8-0) They are useful to ensure algorithmic convergence and to bound the optimization error for the gradient descent-type of algorithms as considered in [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)), when one is interested in approximately solving the non-convex optimization function [\(6\)](#page-2-5). Our numerical results validate such algorithmic and statistical convergence properties.

Acknowledgements

The authors are grateful for the helpful discussions with Prof. Rob Kass.

A Outline

In Sections [B](#page-27-0) and [B.2,](#page-29-0) we present variations of the Hanson-Wright inequality as recently derived in [Rudelson and Vershynin](#page-59-15) [\(2013](#page-59-15)) (cf. Lemma [27\)](#page-27-1), concentration of measure bounds and stochastic error bounds in Lemma [29.](#page-29-2)

In Sections [C](#page-31-1) and [E,](#page-40-1) we prove the technical lemmas for Theorems [2](#page-8-0) and [3](#page-9-0) respectively. In Section [F,](#page-43-0) we prove the Lemmas needed for Proof of Theorem [5.](#page-11-1) In order to prove Corollary [19,](#page-18-2) we need to first state some geometric analysis results Section [G.](#page-45-0) We prove Corollary [19](#page-18-2) in Section [H](#page-49-0) and Theorem [20](#page-18-6) in Section [I.](#page-49-1) Results presented in Section [7](#page-19-0) are proved in Section [J.](#page-51-0) In particular, we prove Theorem [21](#page-19-1) in Section [J.](#page-51-0) We also prove the concentration of measure bounds on error-corrected gram matrices in Corollaries [22](#page-19-2) and [23](#page-20-1) in Sections [J.1](#page-53-0) and [J.2](#page-54-0) respectively. The results appearing in Section [J](#page-51-0) are proved in Section [K.](#page-55-0)

B Some auxiliary results

We first need to state the following form of the Hanson-Wright inequality as recently derived in Rudelson and Vershynin [Rudelson and Vershynin](#page-59-15) [\(2013\)](#page-59-15), and an auxiliary result in Lemma [27](#page-27-1) which may be of independent interests.

Theorem 26. *Let* $X = (X_1, \ldots, X_m) \in \mathbb{R}^m$ *be a random vector with independent components* X_i *which* satisfy $\mathbb{E}(X_i) = 0$ and $||X_i||_{\psi_2} \leq K$. Let A be an $m \times m$ matrix. Then, for every $t > 0$,

$$
\mathbb{P}\left(\left|X^T A X - \mathbb{E}\left(X^T A X\right)\right| > t\right) \leq 2 \exp\left[-c \min\left(\frac{t^2}{K^4 \left\|A\right\|_F^2}, \frac{t}{K^2 \left\|A\right\|_2}\right)\right].
$$

We note that following the proof of Theorem [26,](#page-27-2) it is clear that the following holds: Let $X = (X_1, \ldots, X_m)$ \mathbb{R}^m be a random vector as defined in Theorem [26.](#page-27-2) Let Y, Y' be independent copies of X. Let A be an $m \times m$ matrix. Then, for every $t > 0$,

$$
\mathbb{P}\left(\left|Y^{T}AY'\right|>t\right) \leq 2\exp\left[-c\min\left(\frac{t^{2}}{K^{4}\left\|A\right\|_{F}^{2}},\frac{t}{K^{2}\left\|A\right\|_{2}}\right)\right].
$$
\n(71)

We next need to state Lemma [27,](#page-27-1) which we prove in Section [B.1.](#page-28-0)

Lemma 27. Let $u, w \in S^{f-1}$. Let $A \succ 0$ be a $m \times m$ symmetric positive definite matrix. Let Z be an $f \times m$ *random matrix with independent entries* Z_{ij} *satisfying* $EZ_{ij} = 0$ *and* $||Z_{ij}||_{\psi_2} \le K$ *. Let* Z_1, Z_2 *be* *independent copies of Z. Then for every* $t > 0$ *,*

$$
\mathbb{P}\left(\left|u^T Z_1 A^{1/2} Z_2^T w\right| > t\right) \leq 2 \exp\left(-c \min\left(\frac{t^2}{K^4 \text{tr}(A)}, \frac{t}{K^2 \|A\|_2^{1/2}}\right)\right),
$$

$$
\mathbb{P}\left(\left|u^T Z A Z^T w - \mathbb{E}u^T Z A Z^T w\right| > t\right) \leq 2 \exp\left(-c \min\left(\frac{t^2}{K^4 \|A\|_F^2}, \frac{t}{K^2 \|A\|_2}\right)\right)
$$

where c *is the same constant as defined in Theorem [26.](#page-27-2)*

B.1 Proof of Lemma [27](#page-27-1)

Lemma [28](#page-28-1) is a well-known fact.

Lemma 28. *Let* $A_{uw} := (u \otimes w) \otimes A$ *where* $u, w \in \mathbb{S}^{p-1}$ *where* $p \geq 2$ *. Then* $||A_{uw}||_2 \leq ||A||_2$ *and* $||A_{uw}||_F \leq$ $||A||_F$.

Proof of Lemma [27.](#page-27-1) Let $z_1, \ldots, z_f, z'_1, \ldots, z'_f \in \mathbb{R}^m$ be the row vectors Z_1, Z_2 respectively. Notice that we can write the quadratic form as follows:

$$
u^T Z_1 A^{1/2} Z_2^T w = \sum_{i,j=1,m} u_i w_j z_i A^{1/2} z'_j
$$

\n
$$
= \operatorname{vec} \{ Z_1^T \}^T ((u \otimes w) \otimes A^{1/2}) \operatorname{vec} \{ Z_2^T \}
$$

\n
$$
=:\operatorname{vec} \{ Z_1^T \}^T A_{uw}^{1/2} \operatorname{vec} \{ Z_2^T \},
$$

\n
$$
u^T Z A Z^T w = \operatorname{vec} \{ Z^T \}^T ((u \otimes w) \otimes A) \operatorname{vec} \{ Z^T \}
$$

\n
$$
=:\operatorname{vec} \{ Z^T \}^T A_{uw} \operatorname{vec} \{ Z^T \}
$$

where clearly by independence of Z_1, Z_2 ,

$$
\begin{array}{rcl}\n\text{Evec}\left\{\left. Z_{1}^{\mathrm{T}}\right\}^{\mathrm{T}}\left((\mathrm{u}\otimes\mathrm{w})\otimes\mathrm{A}^{1/2}\right)\mathrm{vec}\left\{\left. Z_{2}^{\mathrm{T}}\right\} & = & 0, \text{ and} \\
\text{Evec}\left\{\left. Z\right\}^{\mathrm{T}}\left((\mathrm{u}\otimes\mathrm{u})\otimes\mathrm{A}\right)\mathrm{vec}\left\{\left. Z\right\} \right. & = & \text{tr}\left((\mathrm{u}\otimes\mathrm{u})\otimes\mathrm{A}\right)=\mathrm{tr}(A).\n\end{array}
$$

Thus we invoke [\(71\)](#page-27-3) and Lemma [28](#page-28-1) to show the concentration bounds on event $\{|u^T Z_1 A^{1/2} Z_2^T w| > t\}$:

$$
\mathbb{P}\left(\left|u^T Z_1 A^{1/2} Z_2^T w\right| > t\right) \leq 2 \exp\left(-\min\left(\frac{t^2}{K^4 \left\|A_{uw}^{1/2}\right\|_F^2}, \frac{t}{K^2 \left\|A_{uw}^{1/2}\right\|_2}\right)\right)
$$

$$
\leq 2 \exp\left(-\min\left(\frac{t^2}{K^4 \text{tr}(A)}, \frac{t}{K^2 \left\|A^{1/2}\right\|_2}\right)\right).
$$

Similarly, we have by Theorem [26](#page-27-2) and Lemma [28,](#page-28-1)

$$
\mathbb{P}\left(\left|u^T Z A Z^T w - \mathbb{E} u^T Z A Z^T w\right| > t\right)
$$
\n
$$
\leq 2 \exp\left(-c \min\left(\frac{t^2}{K^4 \left\|A_{uw}\right\|_F^2}, \frac{t}{K^2 \left\|A_{uw}\right\|_2}\right)\right)
$$
\n
$$
\leq 2 \exp\left(-c \min\left(\frac{t^2}{K^4 \left\|A\right\|_F^2}, \frac{t}{K^2 \left\|A\right\|_2}\right)\right).
$$

The Lemma thus holds. \square

B.2 Stochastic error terms

The following large deviation bounds in Lemmas [29](#page-29-2) and [7](#page-13-2) are the key results in proving Lemmas [6](#page-13-1) and [13.](#page-16-5) Let C_0 satisfy [\(86\)](#page-39-0) for c as defined in Theorem [26.](#page-27-2) Throughout this section, we denote by:

$$
r_{m,f}=C_0K\sqrt{\frac{\log m}{f}}\quad\text{and}\quad r_{m,m}=2C_0\sqrt{\frac{\log m}{mf}}.
$$

We also define some events \mathcal{B}_4 , \mathcal{B}_5 , \mathcal{B}_6 , \mathcal{B}_{10} ; Denote by $\mathcal{B}_0 := B_4 \cap \mathcal{B}_5 \cap \mathcal{B}_6$, which we use throughout this paper.

Lemma 29. Assume that the stable rank of B, $||B||_F^2$ $\frac{2}{\sqrt{F}}$ / $||B||_2^2 \ge \log m$. Let Z , X_0 and W as defined in *Theorem* [2.](#page-8-0) Let Z_0, Z_1 and Z_2 be independent copies of Z. Let $\epsilon^T \sim Y M_{\epsilon}/K$ where $Y := e_1^T Z_0^T$. Let $\tau_B = \frac{\text{tr}(B)}{f}$ f *. Denote by* B⁴ *the event such that*

$$
\frac{1}{f} \left\| A^{\frac{1}{2}} Z_1^T \epsilon \right\|_{\infty} \leq r_{m,f} M_{\epsilon} a_{\max}^{1/2}
$$

and
$$
\frac{1}{f} \left\| Z_2^T B^{\frac{1}{2}} \epsilon \right\|_{\infty} \leq r_{m,f} M_{\epsilon} \sqrt{\tau_B}.
$$

Then $\mathbb{P}(\mathcal{B}_4) \geq 1 - 4/m^3$ *. Moreover, denote by* \mathcal{B}_5 *the event such that*

$$
\frac{1}{f} \left\| (Z^T B Z - \text{tr}(B) I_m) \beta^* \right\|_{\infty} \leq r_{m,f} K \left\| \beta^* \right\|_2 \frac{\left\| B \right\|_F}{\sqrt{f}}
$$

and
$$
\frac{1}{f} \left\| X_0^T W \beta^* \right\|_{\infty} \leq r_{m,f} K \left\| \beta^* \right\|_2 \sqrt{\tau_B} a_{\text{max}}^{1/2}.
$$

Then $\mathbb{P}(\mathcal{B}_5) \geq 1 - 4/m^3$ *.*

Finally, denote by B_{10} *the event such that*

$$
\frac{1}{f} \left\| (Z^T B Z - \text{tr}(B) I_m) \right\|_{\text{max}} \le r_{m,f} K \frac{\|B\|_F}{\sqrt{f}}
$$

and
$$
\frac{1}{f} \left\| X_0^T W \right\|_{\text{max}} \le r_{m,f} K \left\| \beta^* \right\|_2 \sqrt{\tau_B} a_{\text{max}}^{1/2}.
$$

Then $\mathbb{P}(\mathcal{B}_{10}) \geq 1 - 4/m^2$ *.*

We prove Lemmas [29](#page-29-2) in Section **[B.3.](#page-29-1)**

B.3 Stochastic error bounds

s Following Lemma [27,](#page-27-1) we have for all $t > 0$, $B \succ 0$ being an $f \times f$ symmetric positive definite matrix, and $v, w \in \mathbb{R}^m$

$$
\mathbb{P}\left(\left|v^T Z_1^T B^{1/2} Z_2 w\right| > t\right) \leq 2 \exp\left[-c \min\left(\frac{t^2}{K^4 \text{tr}(B)}, \frac{t}{K^2 \left\|B\right\|_2^{1/2}}\right)\right]
$$
(72)

$$
\mathbb{P}\left(\left|v^T Z^T B Z w - \mathbb{E} v^T Z^T B Z w\right| > t\right) \leq 2 \exp\left(-c \min\left(\frac{t^2}{K^4 \left\|B\right\|_F^2}, \frac{t}{K^2 \left\|B\right\|_2}\right)\right).
$$

B.4 Proof for Lemma [29](#page-29-2)

Let $e_1, \ldots, e_m \in \mathbb{R}^m$ be the canonical basis spanning \mathbb{R}^m . Let $x_1, \ldots, x_m, x'_1, \ldots, x'_m \in \mathbb{R}^f$ be the column vectors Z_1, Z_2 respectively. Let $Y \sim e_1^T Z_0^T$. Let $w_i = \frac{A^{1/2} e_i}{\|A^{1/2} e_i\|}$ $\frac{A^{4/2}e_i}{\|A^{1/2}e_i\|_2}$ for all *i*. Clearly the condition on the stable rank of B guarantees that

$$
f \ge r(B) = \frac{\text{tr}(B)}{\|B\|_2} = \frac{\text{tr}(B) \|B\|_2}{\|B\|_2^2} \ge \|B\|_F^2 / \|B\|_2^2 \ge \log m.
$$

By [\(71\)](#page-27-3), we obtain for $t' = C_0 M_e K \sqrt{\text{tr}(B) \log m}$ and $t = C_0 K^2 \sqrt{\log m} \text{tr}(B)^{1/2}$:

$$
\mathbb{P}\left(\exists j, \left| \epsilon^T B^{1/2} Z_2 e_j \right| > t'\right) =
$$
\n
$$
\mathbb{P}\left(\exists j, \frac{M_\epsilon}{K} \left| e_1^T Z_0^T B^{1/2} Z_2 e_j \right| > C_0 M_\epsilon K \sqrt{\log m} \text{tr}(B)^{\frac{1}{2}}\right)
$$
\n
$$
\leq \exp(\log m) \mathbb{P}\left(\left| Y^T B^{1/2} x'_j \right| > C_0 K^2 \sqrt{\log m} \text{tr}(B)^{\frac{1}{2}}\right) \leq 2/m^3
$$

where the last inequality holds by the union bound, given that $\frac{\text{tr}(B)}{\|B\|}$ $\frac{\text{tr}(B)}{\|B\|_2} \ge \log m$, and for all j

$$
\begin{array}{lcl} \displaystyle &\mathbb{P}\left(\left|Y^{T}B^{1/2}x_{j}'\right|>t\right) &\leq & 2\exp\left(-c\min\left(\frac{t^{2}}{K^{4}\mathrm{tr}(B)},\frac{t}{K^{2}\left\|B\right\|_{2}^{1/2}}\right)\right),\\ \\ &\leq & 2\exp\left(-c\min\left(C_{0}^{2}\log m,\frac{C_{0}\log^{1/2}m\sqrt{\mathrm{tr}(B)}}{\left\|B\right\|_{2}^{1/2}}\right)\right)\\ \\ &\leq & 2\exp\left(-c\min(C_{0}^{2},C_{0})\log m\right)\leq 2\exp\left(-4\log m\right). \end{array}
$$

Let $v, w \in S^{m-1}$. Thus we have by Lemma [27,](#page-27-1) for $t_0 = C_0 M_e K \sqrt{f \log m}$ and $\tau = C_0 K^2 \sqrt{f \log m}$, $w_j = \frac{A^{1/2}e_j}{\|\mathbf{A}^{1/2}e_j\|}$ $\frac{A^{\prime\prime}e_j}{\|A^{1/2}e_j\|_2}$ and $f \ge \log m$,

$$
\mathbb{P}(\exists j, |\epsilon^T Z_1 w_j| > t_0) \le \mathbb{P}\left(\exists j, \frac{M_{\epsilon}}{K} |Y^T Z_1 w_j| > C_0 M_{\epsilon} K \sqrt{f \log m}\right)
$$

\n
$$
\le m \mathbb{P}\left(|Y^T Z_1 w_j| > C_0 K^2 \sqrt{f \log m}\right)
$$

\n
$$
= \exp(\log m) \mathbb{P}\left(|e_1^T Z_0^T Z_1 w_j| > \tau\right) \le 2 \exp\left(-c \min\left(\frac{\tau^2}{K^4 f}, \frac{\tau}{K^2}\right)\right),
$$

\n
$$
\le 2 \exp\left(-c \min\left(\frac{(C_0 K^2 \sqrt{f \log m})^2}{K^4 f}, \frac{C_0 K^2 \sqrt{f \log m}}{K^2}\right) + \log m\right)
$$

\n
$$
\le 2m \exp\left(-c \min\left(C_0^2 \log m, C_0 \log^{1/2} m \sqrt{f}\right)\right)
$$

\n
$$
\le 2m \exp\left(-c \min(C_0^2, C_0) \log m\right) \le 2 \exp(-3 \log m).
$$

Therefore we have with probability at least $1 - 4/m^3$,

$$
\begin{aligned}\n\left\|Z_2^T B^{\frac{1}{2}} \epsilon\right\|_{\infty} &:= \max_{j=1,\dots,m} \left\langle \epsilon^T B^{1/2} Z_2, e_j \right\rangle \le t' = C_0 M_{\epsilon} K \sqrt{\text{tr}(B) \log m} \\
\left\|A^{\frac{1}{2}} Z_1^T \epsilon\right\|_{\infty} &:= \max_{j=1,\dots,m} \left\langle A^{1/2} e_j, Z_1^T \epsilon \right\rangle \le \max_{j=1,\dots,m} \left\|A^{1/2} e_j\right\|_{2} \max_{j=1,\dots,m} \left\langle w_j, Z_1^T \epsilon \right\rangle \\
&\le a_{\text{max}}^{1/2} t_0 = a_{\text{max}}^{1/2} C_0 M_{\epsilon} K \sqrt{f \log m}.\n\end{aligned}
$$

The "moreover" part follows exactly the same arguments as above. Denote by $\bar{\beta}^* := \beta^* / ||\beta^*||_2 \in E \cap S^{m-1}$ and $w_i := A^{1/2} \tilde{e_i} / \|A^{1/2} e_i\|_2$. By [\(72\)](#page-29-3)

$$
\mathbb{P}\left(\exists i, \langle w_i, Z_1^T B^{1/2} Z_2 \overline{\beta}^* \rangle \ge C_0 K^2 \sqrt{\log m} \text{tr}(B)^{1/2}\right)
$$

\n
$$
\le \sum_{i=1}^m \mathbb{P}\left(\langle w_i, Z_1^T B^{1/2} Z_2 \overline{\beta}^* \rangle \ge C_0 K^2 \sqrt{\log m} \text{tr}(B)\right)
$$

\n
$$
\le 2 \exp\left(-c \min\left(C_0^2 \log m, C_0 \log m\right) + \log m\right) \le 2/m^3.
$$

Now for $t = C_0 K^2 \sqrt{\log m} ||B||_F$, and $||B||_F / ||B||_2 \ge \sqrt{\log m}$,

$$
\mathbb{P}\left(\exists e_i : \langle e_i, (Z^T BZ - \text{tr}(B)I_m)\overline{\beta}^* \rangle \ge C_0 K^2 \sqrt{\log m} \|B\|_F\right)
$$

$$
\le 2m \exp\left[-c \min\left(\frac{t^2}{K^4 \|B\|_F^2}, \frac{t}{K^2 \|B\|_2}\right)\right] \le 2/m^3.
$$

By the two inequalities immediately above, we have with probability at least $1 - 4/m³$,

$$
||X_0^T W \beta^*||_{\infty} = ||A^{1/2} Z_1^T B^{1/2} Z_2 \beta^*||_{\infty}
$$

\n
$$
\leq ||\beta^*||_2 \max_{e_i} ||A^{1/2} e_i||_2 \left(\sup_{w_i} \langle w_i, Z_1^T B^{1/2} Z_2 \bar{\beta}^* \rangle \right)
$$

\n
$$
\leq C_0 K^2 ||\beta^*||_2 \sqrt{\log m} a_{\max}^{1/2} \sqrt{\text{tr}(B)}
$$

and

$$
\begin{aligned} \left\| (Z^T B Z - \text{tr}(B) I_m) \beta^* \right\|_{\infty} &= \left\| (Z^T B Z - \text{tr}(B) I_m) \bar{\beta}^* \right\|_{\infty} \left\| \beta^* \right\|_2 \\ &= \|\beta^* \right\|_2 \left(\sup_{e_i} \left\langle e_i, (Z^T B Z - \text{tr}(B) I_m) \bar{\beta}^* \right\rangle \right) \\ &\leq C_0 K^2 \left\| \beta^* \right\|_2 \sqrt{\log m} \left\| B \right\|_F. \end{aligned}
$$

The last two bounds follow exactly the same arguments as above, except that we replace β^* with e_j , $j =$ 1, ..., *m* and apply the union bounds to m^2 events instead of *m*, and thus $\mathbb{P}(\mathcal{B}_{10}) \ge 1 - 4/m^2$, \Box

C Proofs for the Lasso-type estimator

C.1 Proof of Lemma [6](#page-13-1)

Clearly the condition on the stable rank of B guarantees that

$$
f \ge r(B) = \frac{\text{tr}(B)}{\|B\|_2} = \frac{\text{tr}(B) \|B\|_2}{\|B\|_2^2} \ge \|B\|_F^2 / \|B\|_2^2 \ge \log m.
$$

Thus the conditions in Lemmas [29](#page-29-2) and [7](#page-13-2) hold. First notice that

$$
\widehat{\gamma} = \frac{1}{f} \left(X_0^T X_0 \beta^* + W^T X_0 \beta^* + X_0^T \epsilon + W^T \epsilon \right)
$$

$$
(\frac{1}{f} X^T X - \frac{\widehat{\text{tr}}(B)}{f} I_m) \beta^* = \frac{1}{f} (X_0^T X_0 + W^T X_0 + X_0^T W + W^T W - \frac{\widehat{\text{tr}}(B)}{f} I_m) \beta^*
$$

Thus

$$
\begin{split}\n\left\|\widehat{\gamma} - \widehat{\Gamma}\beta^*\right\|_{\infty} &\leq \quad \left\|\widehat{\gamma} - \frac{1}{f}\left(X^TX - \widehat{\text{tr}}(B)I_m\right)\beta^*\right\|_{\infty} \\
&= \quad \frac{1}{f}\left\|X_0^T\epsilon + W^T\epsilon - \left(W^TW + X_0^TW - \widehat{\text{tr}}(B)I_m\right)\beta^*\right\|_{\infty} \\
&\leq \quad \frac{1}{f}\left\|X_0^T\epsilon + W^T\epsilon\right\|_{\infty} + \frac{1}{f}\left\|\left(W^TW - \widehat{\text{tr}}(B)I_m\right)\beta^*\right\|_{\infty} + \left\|\frac{1}{f}X_0^TW\beta^*\right\|_{\infty} \\
&\leq \quad \frac{1}{f}\left\|X_0^T\epsilon + W^T\epsilon\right\|_{\infty} + \frac{1}{f}(\left\|\left(Z^T BZ - \text{tr}(B)I_m\right)\beta^*\right\|_{\infty}) + \frac{1}{f}\left\|X_0^TW\beta^*\right\|_{\infty} \\
&\quad + \frac{1}{f}\left|\widehat{\text{tr}}(B) - \text{tr}(B)\right|\left\|\beta^*\right\|_{\infty} =: U_1 + U_2 + U_3 + U_4\n\end{split}
$$

By Lemma [29](#page-29-2) we have on B_4 for $D_0 := \sqrt{\tau_B} + a_{\text{max}}^{1/2}$,

$$
U_1 = \frac{1}{f} \| X_0^T \epsilon + W^T \epsilon \|_{\infty} = \frac{1}{f} \| A^{\frac{1}{2}} Z_1^T \epsilon + Z_2^T B^{\frac{1}{2}} \epsilon \|_{\infty} \le r_{m,f} M_{\epsilon} D_0
$$

and on event \mathcal{B}_5 for $D'_0 := \sqrt{\|B\|_2} + a_{\text{max}}^{1/2}$,

$$
U_2 + U_3 = \frac{1}{f} ||(Z^T B Z - \text{tr}(B)I_m)\beta^*||_{\infty} + \frac{1}{f} ||X_0^T W \beta^*||_{\infty}
$$

$$
\leq r_{m,f} K ||\beta^*||_2 \left(\frac{||B||_F}{\sqrt{f}} + \sqrt{\tau_B} a_{\text{max}}^{1/2} \right) \leq K r_{m,f} ||\beta^*||_2 \tau_B^{1/2} D_0'
$$

where recall $||B||_F \leq \sqrt{\text{tr}(B)} ||B||_2^{1/2}$ ^{1/2}. Denote by $B_0 := B_4 \cap B_5 \cap B_6$. We have on B_0 and under (A1), by Lemmas [29](#page-29-2) and $\overline{7}$ $\overline{7}$ $\overline{7}$ and \overline{D}_1 defined therein,

$$
\left\|\hat{\gamma} - \hat{\Gamma}\beta^*\right\|_{\infty} \leq U_1 + U_2 + U_3 + U_4
$$

\n
$$
\leq r_{m,f} M_{\epsilon} D_0 + D'_0 \tau_B^{1/2} K r_{m,f} \left\|\beta^*\right\|_2 + \frac{1}{f} \left|\hat{\text{tr}}(B) - \text{tr}(B)\right| \|\beta^*\|_{\infty}
$$

\n
$$
\leq D_0 M_{\epsilon} r_{m,f} + D'_0 K \tau_B^{1/2} \|\beta^*\|_2 r_{m,f} + D_1 K^2 \|\beta^*\|_{\infty} r_{m,m}
$$
(73)

$$
\leq D_0 M_{\epsilon} r_{m,f} + D'_0 K \tau_B^{1/2} \|\beta^*\|_2 r_{m,f} + 2D_1 K \frac{1}{\sqrt{m}} \|\beta^*\|_{\infty} r_{m,f}
$$
(74)

$$
\leq r_{m,f} \left(\left(\frac{3}{4} D_2 + D_2 \frac{1}{\sqrt{m}} \right) K \|\beta^*\|_2 + D_0 M_{\epsilon} \right)
$$

where $2D_1 \leq 2 ||A||_2 + 2 ||B||_2 = D_2$, for $(D'_0)^2 \leq 2 ||B||_2 + 2a_{\text{max}}$

$$
D_0 \leq D'_0 \leq \sqrt{2(||B||_2 + a_{\max})} \leq 2(a_{\max} + ||B||_2) = D_2,
$$

and
$$
D'_0 \tau_B^{1/2} \leq (||B||_2^{1/2} + a_{\max}^{1/2})\tau_B^{1/2} \leq \tau_B + \frac{1}{2}(||B||_2 + a_{\max}) \leq \frac{3}{4}D_2
$$

given that under (A1): $\tau_A = 1$, $||A||_2 \ge a_{\text{max}} \ge a_{\text{max}}^{\frac{1}{2}} \ge 1$. Hence the lemma holds for $m \ge 16$ and $\psi = C_0 D_2 K (K ||\beta^*||_2 + M_\epsilon)$. Finally, we have by the union bound, $\mathbb{P}(\mathcal{B}_0) \geq 1 - 16/m^3$. \Box

C.2 Proof of Lemma [7](#page-13-2)

First we write

$$
XX^{T} - \text{tr}(A)I_{f} = (Z_{1}A^{1/2} + B^{1/2}Z_{2})(Z_{1}A^{1/2} + B^{1/2}Z_{2})^{T} - \text{tr}(A)I_{f}
$$

\n
$$
= (Z_{1}A^{1/2} + B^{1/2}Z_{2})(Z_{2}^{T}B^{1/2} + A^{1/2}Z_{1}^{T}) - \text{tr}(A)I_{f}
$$

\n
$$
= Z_{1}A^{1/2}Z_{2}^{T}B^{1/2} + B^{1/2}Z_{2}Z_{2}^{T}B^{1/2}
$$

\n
$$
+ B^{1/2}Z_{2}A^{1/2}Z_{1}^{T} + Z_{1}AZ_{1}^{T} - \text{tr}(A)I_{f}.
$$

Thus we have for $\check{\operatorname{tr}}(B) := \frac{1}{m} (||X||_F^2 - f \operatorname{tr}(A))$

$$
\frac{1}{f}(\tilde{tr}(B) - tr(B)) := \frac{1}{mf} (||X||_F^2 - ftr(A) - mtr(B))
$$
\n
$$
= \frac{1}{mf} (tr(XX^T) - ftr(A) - mtr(B))
$$
\n
$$
= \frac{2}{mf} tr(Z_1 A^{1/2} Z_2^T B^{1/2}) + \left(\frac{tr(B^{1/2} Z_2 Z_2^T B^{1/2})}{mf} - \frac{tr(B)}{f} \right)
$$
\n
$$
+ \frac{tr(Z_1 A Z_1^T)}{mf} - \frac{tr(A)}{m}
$$

By constructing a new matrix $A_f = I_f \otimes A$ which is block diagonal with f identical submatrices A along its diagonal, we prove the following large deviation bound: for $t_1 = C_0 K^2 ||A||_F \sqrt{f \log m}$ and $f > \log m$,

$$
\mathbb{P}\left(\left|\text{tr}(Z_1 A Z_1^T) - f \text{tr}(A)\right| \ge t_1\right) = \mathbb{P}\left(\left|\text{vec}\left\{Z_1\right\}^T (\text{I} \otimes A)\text{vec}\left\{Z_1\right\} - \text{ftr}(A)\right| \ge t_1\right)
$$
\n
$$
\le \exp\left(-c \min\left(\frac{t_1^2}{K^4 \left\|A_f\right\|_F^2}, \frac{t_1}{K^2 \left\|A_f\right\|_2}\right)\right)
$$
\n
$$
\le 2 \exp\left(-c \min\left(\frac{(C_0 K^2 \sqrt{f \log m} \|A\|_F)^2}{K^4 f \|A\|_F^2}, \frac{C_0 K^2 \sqrt{f \log m} \|A\|_F}{K^2 \|A\|_2}\right)\right)
$$
\n
$$
\le 2 \exp\left(-4 \log m\right)
$$

where the first inequality holds by Theorem [26](#page-27-2) and the second inequality holds given that $||A_f||_F^2 = f||A||_F$ and $||A_f||_2^2 = ||A||_2$. Similarly, by constructing a new matrix $B_m = I_m \otimes B$ which is block diagonal with m identical submatrices B along its diagonal, we prove the following large deviation bound: for $t_2 = C_0 K^2 ||B||_F \sqrt{m \log m}$ and $m \geq 2$,

$$
\mathbb{P}\left(\left|\text{tr}(Z_2^T B Z_2) - m \text{tr}(B)\right| \ge t_2\right) = \mathbb{P}\left(\left|\text{vec}\{Z_2\}^T (I_m \otimes B)\text{vec}\{Z_2\} - m \text{tr}(B)\right| \ge t_2\right)
$$
\n
$$
\le \exp\left(-c \min\left(\frac{t_2^2}{K^4 m \|B\|_F^2}, \frac{t_2}{K^2 \|B\|_2}\right)\right)
$$
\n
$$
\le 2 \exp\left(-c \min\left(\frac{(C_0 K^2 \sqrt{m \log m \|B\|_F})^2}{K^4 m \|B\|_F^2}, \frac{C_0 K^2 \sqrt{m \log m \|B\|_F}}{K^2 \|B\|_2}\right)\right)
$$
\n
$$
\le 2 \exp(-4 \log m).
$$

Finally, we have by [\(71\)](#page-27-3) for $t_0 = C_0 K^2 \sqrt{\text{tr}(A)\text{tr}(B) \log m}$,

$$
\mathbb{P}\left(\left|\text{vec}\left\{Z_{1}\right\}^{\mathrm{T}}\mathbf{B}^{1/2} \otimes \mathbf{A}^{1/2}\text{vec}\left\{Z_{2}\right\}\right| > t_{0}\right)
$$
\n
$$
\leq 2 \exp\left(-c \min\left(\frac{t_{0}^{2}}{K^{4} \left\|B^{1/2} \otimes A^{1/2}\right\|_{F}^{2}}, \frac{t_{0}}{K^{2} \left\|B^{1/2} \otimes A^{1/2}\right\|_{2}}\right)\right)
$$
\n
$$
= 2 \exp\left(-c \min\left(\frac{(C_{0}\sqrt{\text{tr}(A)\text{tr}(B)\log m})^{2}}{\text{tr}(A)\text{tr}(B)}, \frac{C_{0}\sqrt{\text{tr}(A)\text{tr}(B)\log m}}{\left\|B\right\|_{2}^{1/2} \left\|A\right\|_{2}^{1/2}}\right)\right)
$$
\n
$$
\leq 2 \exp(-4 \log m)
$$

where we used and the fact that $r(A)r(B) \ge \log m$, $||B^{1/2} \otimes A^{1/2}||_2 = ||B||_2^{1/2}$ $\frac{1}{2}$ $||A||_2^{1/2}$ $\frac{1}{2}$ and

$$
\left\|B^{1/2}\otimes A^{1/2}\right\|_{F}^{2} = \text{tr}((B^{1/2}\otimes A^{1/2})(B^{1/2}\otimes A^{1/2})) = \text{tr}(B\otimes A) = \text{tr}(A)\text{tr}(B).
$$

Thus we have with probability $1 - 6/m^4$,

$$
\frac{1}{f} |\dot{\mathbf{tr}}(B) - \mathbf{tr}(B)| = \frac{1}{mf} |\mathbf{tr}(XX^T) - f\mathbf{tr}(A) - m\mathbf{tr}(B)|
$$
\n
$$
\leq \frac{2}{mf} |\text{vec}\{Z_1\}^T (B^{1/2} \otimes A^{1/2}) \text{vec}\{Z_2\}|
$$
\n
$$
+ \left| \frac{\mathbf{tr}(Z_2^T B Z_2)}{mf} - \frac{\mathbf{tr}(B)}{f} \right| + \left| \frac{\mathbf{tr}(Z_1 A Z_1^T)}{mf} - \frac{\mathbf{tr}(A)}{m} \right|
$$
\n
$$
\leq \frac{1}{mf} (2t_0 + t_1 + t_2) = \frac{\sqrt{\log m}}{\sqrt{mf}} C_0 K^2 \left(\frac{\|A\|_F}{\sqrt{m}} + 2\sqrt{\tau_A \tau_B} + \frac{\|B\|_F}{\sqrt{f}} \right)
$$
\n
$$
\leq 2C_0 \frac{\sqrt{\log m}}{\sqrt{mf}} K^2 D_1 =: D_1 K^2 r_{m,m}
$$

where recall $r_{m,m} = 2C_0 \frac{\sqrt{3}}{2}$ $\frac{\sqrt{\log m}}{\sqrt{mf}}, D_1 = \frac{\|A\|_F}{\sqrt{m}} + \frac{\|B\|_F}{\sqrt{f}},$ and

$$
2\sqrt{\tau_A \tau_B} \le \tau_A + \tau_B \le \frac{\|A\|_F}{\sqrt{m}} + \frac{\|B\|_F}{\sqrt{f}}.
$$

To see this, recall

$$
m\tau_A = \sum_{i=1}^m \lambda_i(A) \le \sqrt{m} \left(\sum_{i=1}^m \lambda_i^2(A) \right)^{1/2} = \sqrt{m} \|A\|_F
$$

\n
$$
f\tau_B = \sum_{i=1}^f \lambda_i(B) \le \sqrt{f} \left(\sum_{i=1}^f \lambda_i^2(B) \right)^{1/2} = \sqrt{f} \|B\|_F
$$
\n(75)

where $\lambda_i(A), i = 1, \ldots, m$ and $\lambda_i(B), i = 1, \ldots, f$ denote the eigenvalues of positive semidefinite covariance matrices A and B respectively.

Denote by B_6 the following event

$$
\left\{ \frac{1}{f} \left| \check{\operatorname{tr}}(B) - \operatorname{tr}(B) \right| \le D_1 K^2 r_{m,m} \right\}
$$

Clearly $\hat{\text{tr}}(B) := (\check{\text{tr}}(B))_+$ by definition [\(4\)](#page-2-3). As a consequence, on \mathcal{B}_6 , $\hat{\text{tr}}(B) = \check{\text{tr}}(B) > 0$ when $\tau_B > 0$ $D_1 K^2 r_{m,m}$; hence

$$
\frac{1}{f} \left| \widehat{\text{tr}}(B) - \text{tr}(B) \right| = \frac{1}{f} \left| \widecheck{\text{tr}}(B) - \text{tr}(B) \right| \le D_1 K^2 r_{m,m}.
$$

Otherwise, it is possible that $\check{tr}(B) < 0$. However, suppose we set

$$
\widehat{\tau}_B := \frac{1}{f}\widehat{\text{tr}}(B) := \frac{1}{f}(\widecheck{\text{tr}}(B) \vee 0),
$$

then we can also guarantee that

$$
|\widehat{\tau}_B - \tau_B| = |\tau_B| \le D_1 K^2 r_{m,m} \quad \text{in case} \quad \tau_B \le D_1 K^2 r_{m,m}.
$$

The lemma is thus proved. \square

D Proof of Theorem [8](#page-14-0)

Denote by $\beta = \beta^*$. Let $S := \text{supp } \beta$, $d = |S|$ and

$$
v=\widehat{\beta}-\beta.
$$

where $\hat{\beta}$ is as defined in [\(6\)](#page-2-5). We first show Lemma [30,](#page-35-1) followed by the proof of Theorem [8.](#page-14-0) **Lemma 30.** *[Bickel et al.](#page-57-8) [\(2009](#page-57-8)); [Loh and Wainwright](#page-59-0) [\(2012](#page-59-0)) Suppose that* [\(34\)](#page-14-2) *holds. Suppose that there exists a parameter* ψ *such that*

$$
\sqrt{d}\tau \le \frac{\psi}{b_0} \sqrt{\frac{\log m}{f}}, \quad \text{and} \quad \lambda \ge 4\psi \sqrt{\frac{\log m}{f}}
$$

where b_0 , λ *are as defined in* [\(6\)](#page-2-5). Then $||v_{Sc}||_1 \leq 3 ||v_S||_1$.

Proof. By the optimality of $\widehat{\beta}$, we have

$$
\lambda_n \|\beta\|_1 - \lambda_n \left\|\widehat{\beta}\right\|_1 \geq \frac{1}{2}\widehat{\beta}\widehat{\Gamma}\widehat{\beta} - \frac{1}{2}\widehat{\beta}\widehat{\Gamma}\beta - \langle \widehat{\gamma}, v \rangle
$$

=
$$
\frac{1}{2}v\widehat{\Gamma}v + \langle v, \widehat{\Gamma}\beta \rangle - \langle v, \widehat{\gamma} \rangle
$$

=
$$
\frac{1}{2}v\widehat{\Gamma}v - \langle v, \widehat{\gamma} - \widehat{\Gamma}\beta \rangle
$$

Hence, we have for $\lambda \ge 4\psi \sqrt{\frac{\log m}{f}}$ $\frac{\mathrm{g}\, m}{f},$

$$
\frac{1}{2}v\widehat{\Gamma}v \leq \langle v,\widehat{\gamma}-\widehat{\Gamma}\beta\rangle + \lambda_n \left(\|\beta\|_1 - \left\|\widehat{\beta}\right\|_1\right) \leq \lambda_n \left(\|\beta\|_1 - \left\|\widehat{\beta}\right\|_1\right) + \left\|\widehat{\gamma}-\widehat{\Gamma}\beta\right\|_{\infty} \|v\|_1
$$
\n(76)

Hence

$$
v\widehat{\Gamma}v \leq \lambda_n \left(2\left\|\beta\right\|_1 - 2\left\|\widehat{\beta}\right\|_1 \right) + 2\psi \sqrt{\frac{\log m}{f}} \left\|v\right\|_1
$$
\n
$$
\leq \lambda_n \left(2\left\|\beta\right\|_1 - 2\left\|\widehat{\beta}\right\|_1 + \frac{1}{2}\left\|v\right\|_1 \right) \tag{77}
$$

$$
\leq \lambda_n \frac{1}{2} \left(5 \left\| v_S \right\|_1 - 3 \left\| v_{S^c} \right\|_1 \right). \tag{78}
$$

where by the triangle inequality, and $\beta_{S^c} = 0$, we have

$$
2\|\beta\|_{1} - 2\left\|\widehat{\beta}\right\|_{1} + \frac{1}{2}\|v\|_{1} = 2\|\beta_{S}\|_{1} - 2\left\|\widehat{\beta}_{S}\right\|_{1} - 2\|v_{S^{c}}\|_{1} + \frac{1}{2}\|v_{S}\|_{1} + \frac{1}{2}\|v_{S^{c}}\|_{1}
$$

\n
$$
\leq 2\|v_{S}\|_{1} - 2\|v_{S^{c}}\|_{1} + \frac{1}{2}\|v_{S}\|_{1} + \frac{1}{2}\|v_{S^{c}}\|_{1}
$$

\n
$$
\leq \frac{1}{2}(5\|v_{S}\|_{1} - 3\|v_{S^{c}}\|_{1}). \tag{79}
$$

We now give a lower bound on the LHS of [\(76\)](#page-35-2), applying the lower-RE condition as in Definition [2.2,](#page-5-1)

$$
v^T \hat{\Gamma} v \geq \alpha \|v\|_2^2 - \tau \|v\|_1^2 \geq -\tau \|v\|_1^2
$$

\nthus $-v^T \hat{\Gamma} v \leq \|v\|_1^2 \tau \leq \|v\|_1 2b_0 \sqrt{d\tau}$
\n $\leq \|v\|_1 2b_0 \frac{\psi}{b_0} \sqrt{\frac{\log m}{f}} = \|v\|_1 2\psi \sqrt{\frac{\log m}{f}}$
\n $\leq \frac{1}{2} \lambda (\|v_S\|_1 + \|v_{S^c}\|_1)$ (80)

where we use the assumption that

$$
\sqrt{d}\tau \leq \frac{\psi}{b_0}\sqrt{\frac{\log m}{f}}, \quad \text{ and } \ \left\|v\right\|_1 \leq \left\|\widehat{\beta}\right\|_1 + \left\|\beta\right\|_1 \leq 2b_0\sqrt{d}
$$

which holds by the triangle inequality and the fact that both $\hat{\beta}$ and β have ℓ_1 norm being bounded by $b_0\sqrt{d}$. Hence by (78) and (80)

$$
0 \leq -v\widehat{\Gamma}v + \frac{5}{2}\lambda \|v_S\|_1 - \frac{3}{2}\lambda \|v_{S^c}\|_1
$$
\n
$$
\leq \frac{1}{2}\lambda \|v_S\|_1 + \frac{1}{2}\lambda \|v_{S^c}\|_1 + \frac{5}{2}\lambda \|v_S\|_1 - \frac{3}{2}\lambda \|v_{S^c}\|_1
$$
\n
$$
\leq 3\lambda \|v_S\|_1 - \lambda \|v_{S^c}\|_1
$$
\n(82)

Thus we have

$$
||v_{S^c}||_1 \le 3 ||v_S||_1
$$

Thus Lemma [30](#page-35-1) holds. \square

Proof of Theorem [8.](#page-14-0) Following the conclusion of Lemma [30,](#page-35-1) we have

$$
||v||_1 \le 4 ||v_S||_1 \le 4\sqrt{d} ||v||_2.
$$
\n(83)

Moreover, we have by the lower-RE condition as in Definition [2.2](#page-5-1)

$$
v^T \hat{\Gamma} v \ge \alpha \|v\|_2^2 - \tau \|v\|_1^2 \ge (\alpha - 16d\tau) \|v\|_2^2 \ge \frac{1}{2} \alpha \|v\|_2^2 \tag{84}
$$

where the last inequality follows from the assumption that $16d\tau \leq \alpha/2$.

Combining the bounds in (84) , (83) and (77) , we have

$$
\frac{1}{2}\alpha \left\|v\right\|_{2}^{2} \leq v^{T}\widehat{\Gamma}v \leq \lambda_{n} \left(2\left\|\beta\right\|_{1} - 2\left\|\widehat{\beta}\right\|_{1}\right) + 2\psi\sqrt{\frac{\log m}{f}} \left\|v\right\|_{1}
$$

$$
\leq \frac{5}{2}\lambda \left\|v_{S}\right\|_{1} \leq 10\lambda\sqrt{d} \left\|v\right\|_{2}
$$

And thus we have $||v||_2 \le 20\lambda\sqrt{d}$. The theorem is thus proved. \Box

D.1 Proof of Lemma [9](#page-15-1)

In view of Remark [D.1,](#page-38-1) Condition [\(40\)](#page-15-3) implies that [\(52\)](#page-19-3) in Theorem [20](#page-18-6) holds for $\zeta = s_0$ and $\varepsilon = \frac{1}{2M}$ $\frac{1}{2M_A}$. Now, by Theorem [20,](#page-18-6) we have $\forall u, v \in E \cap S^{m-1}$, under (A1) and (A3), condition [\(48\)](#page-18-5) holds under event \mathcal{A}_0 , and so long as $mf \geq 1024C_0^2D_2^2K^4\log m/\lambda_{\min}(A)^2$,

$$
\begin{array}{rcl} \left| u^T \Delta v \right| & \leq & 8C \varpi(s_0) \varepsilon + 2C_0 D_2 K^2 \sqrt{\frac{\log m}{m f}} =: \delta \text{ with } \delta \leq \frac{1}{8} \lambda_{\min}(A) \leq \frac{1}{8} \\ & \text{which holds for all } \varepsilon \leq \frac{1}{2} \frac{\lambda_{\min}(A)}{64C \varpi(s_0)} := \frac{1}{2M_A} \leq \frac{1}{128C} \end{array}
$$

with $\mathbb{P}\left(\mathcal{A}_0\right) \geq 1-4\exp\left(-c_2\varepsilon^2 \frac{\text{tr}(B)}{K^4\|B\|_2}\right)$ $-2 \exp \left(-c_2 \varepsilon^2 \frac{f}{K^4}\right)$ $-6/m^3$. Hence, by Corollary [19,](#page-18-2) $\forall \theta \in \mathbb{R}^m$, $\theta^T \widehat{\Gamma}_A \theta \ge \alpha \, \|\theta\|_2^2 - \tau \, \|\theta\|_1^2$ ²₁ and $\theta^T \widehat{\Gamma}_A \theta \leq \bar{\alpha} \|\theta\|_2^2 + \tau \|\theta\|_1^2$ 1

where $\alpha = \frac{1}{2}\lambda_{\min}(A)$ and $\bar{\alpha} = \frac{3}{2}\lambda_{\max}(A)$ and

$$
\frac{512C^2\varpi(s_0)^2\log m}{\lambda_{\min}(A)} \leq \tau = \frac{\alpha}{s_0} \leq \frac{2\alpha}{s_0 + 1}
$$

$$
\leq \frac{1024C^2\varpi^2(s_0 + 1)\log m}{\lambda_{\min}(A)}.
$$

where we plugged in s_0 as defined in [\(12\)](#page-7-1). The lemma is thus proved in view of Remark [D.1.](#page-38-1) \Box

Remark D.1. *Clearly the condition on* $\text{tr}(B)/\|B\|_2$ *as stated in Lemma [9](#page-15-1) ensures that we have for* $\varepsilon = \frac{1}{2M_A}$ and $s_0 \asymp \frac{4f}{M_A^2 \log m}$

$$
\varepsilon^2 \frac{\text{tr}(B)}{K^4 \|B\|_2} \geq \frac{\varepsilon^2}{K^4} c' K^4 \frac{s_0}{\varepsilon^2} \log\left(\frac{3em}{s_0 \varepsilon}\right)
$$

$$
\geq \frac{1}{4M_A^2 K^4} 4c' K^4 M_A^2 s_0 \log\left(\frac{6emM_A}{s_0}\right)
$$

$$
\geq c' s_0 \log\left(\frac{6emM_A}{s_0}\right)
$$

and hence

$$
\exp\left(-c_2\varepsilon^2 \frac{\operatorname{tr}(B)}{K^4 \|B\|_2}\right) \le \exp\left(-c'c_2s_0\log\left(\frac{6emM_A}{s_0}\right)\right)
$$

$$
\approx \exp\left(-c_3\frac{4f}{M_A^2\log m}\log\left(\frac{3eM_A^3m\log m}{2f}\right)\right)
$$

D.2 Proof of Lemma [10](#page-16-2)

Let

$$
M_+ = \frac{64C\varpi(s_0+1)}{\lambda_{\min}(A)} \text{ where } \varpi(s_0+1) = \rho_{\max}(s_0+1, A) + \tau_B =: D
$$

By definition of s_0 , we have

$$
\sqrt{s_0 + 1} \varpi(s_0 + 1) \ge \frac{\lambda_{\min}(A)}{32C} \sqrt{\frac{f}{\log m}}
$$
 and hence

$$
s_0 + 1 \ge \frac{\lambda_{\min}^2(A)}{1024C^2 \varpi^2(s_0 + 1)} \frac{f}{\log m} = \left(\frac{\alpha}{16CD}\right)^2 \frac{f}{\log m} \ge \frac{1}{M_A^2} \frac{f}{\log m}
$$

The first inequality in [\(33\)](#page-14-1) holds given that $M_+ \leq 2 M_A$ and hence

$$
d \le \frac{1}{64 M_A^2} \frac{f}{\log m} \le \frac{1}{16 M_+^2} \frac{f}{\log m} \le \frac{s_0 + 1}{64} \le \frac{s_0}{32}
$$

Moreover, for $D = \rho_{\text{max}}(s_0 + 1, A) + \tau_B \leq D_2$ and $C = C_0 / \sqrt{c'}$, we have

$$
d \leq C_A c' D_{\phi} \frac{f}{\log m} \leq \frac{1}{128M_A^2} \left(\frac{C_0 D_2}{CD}\right)^2 D_{\phi} \frac{f}{\log m}
$$

$$
\leq \frac{1}{2} \left(\frac{1}{16CD}\right)^2 4C_0^2 D_2^2 D_{\phi} \frac{f}{M_A^2 \log m}
$$

$$
\leq \frac{1}{2} \frac{(s_0 + 1)^2 \log m}{\alpha^2} \left(\frac{\psi}{b_0}\right)^2 \leq \frac{(s_0)^2 \log m}{\alpha^2} \left(\frac{\psi}{b_0}\right)^2
$$

where assuming that $s_0 \geq 3$, we have

$$
\frac{2s_0^2}{\alpha^2} \ge \left(\frac{s_0+1}{\alpha}\right)^2 \ge \frac{\alpha^2}{(16CD)^4} \left(\frac{f}{\log m}\right)^2
$$
\n
$$
\left(\frac{\psi}{b_0}\right)^2 = 4C_0^2 D_2^2 \frac{K^2}{b_0^2} (M_\epsilon + K \|\beta^*\|_2)^2
$$
\n
$$
\ge 4C_0^2 D_2^2 D_\phi = 4C_0^2 D_2^2 \left(\frac{K^2 M_\epsilon^2}{b_0^2} + K^4 \phi\right).
$$
\n(85)

We have shown that [\(42\)](#page-16-7) indeed holds, and the lemma is thus proved. \square

Remark D.2. *Throughout this paper, we assume that* C_0 *is a large enough constant such that for c as defined in Theorem [26,](#page-27-2)*

$$
c\min\{C_0^2, C_0\} \ge 4. \tag{86}
$$

By definition of s_0 *, we have for* $\varpi^2(s_0) \geq 1$ *,*

$$
s_0 \varpi^2(s_0) \le \frac{c' \lambda_{\min}^2(A)}{1024C_0^2} \frac{f}{\log m} \text{ and hence}
$$

$$
s_0 \le \frac{c' \lambda_{\min}^2(A)}{1024C_0^2} \frac{f}{\log m} \le \frac{\lambda_{\min}^2(A)}{1024C_0^2} \frac{f}{\log m} =: \check{s}_0.
$$

Remark D.3. *The proof shows that one can take* $C = C_0 / \sqrt{c'}$ *, and take*

$$
\mathcal{V} = 3eM_A^3/2 = \frac{3e64^3C^3\varpi^3(s_0)}{2\lambda_{\min}^3(A)} \le \frac{3e64^3C_0^3\varpi^3(s_0)}{2(c')^{3/2}\lambda_{\min}^3(A)}.
$$

Hence a sufficient condition on $r(B)$ *is:*

$$
r(B) \ge 16c'K^4 \frac{f}{\log m} \left(3\log \frac{64C_0 \varpi(\check{s}_0)}{\sqrt{c'}\lambda_{\min}(A)} + \log \frac{3em\log m}{2f} \right). \tag{87}
$$

It remains to prove Lemmas [14](#page-17-2) and [15.](#page-17-3)

Proof of Lemma [14.](#page-17-2) Suppose that event \mathcal{B}_0 holds. By [\(74\)](#page-32-0) and that fact that $2D_1 := 2(\frac{\|A\|_F}{\sqrt{m}} +$ $\frac{\|B\|_F}{\sqrt{f}}$) $\leq 2(\|A\|_2^{1/2} + \|B\|_2^{1/2})$ $2^{1/2}) (\sqrt{\tau_A} + \sqrt{\tau_B}) \le D_{\text{oracle}} D_0',$ where recall $D_0' = ||B||_2^{1/2} + a_{\text{max}}^{1/2},$

$$
\left\| \hat{\gamma} - \hat{\Gamma}\beta^* \right\|_{\infty} \leq D'_0 K \tau_B^{1/2} \left\| \beta^* \right\|_2 r_{m,f} + 2D_1 K \frac{1}{\sqrt{m}} \left\| \beta^* \right\|_{\infty} r_{m,f} + D_0 M_{\epsilon} r_{m,f}
$$

$$
\leq D'_0 K \left\| \beta^* \right\|_2 r_{m,f} \left(\tau_B^{1/2} + \frac{D_{\text{oracle}}}{\sqrt{m}} \right) + D_0 M_{\epsilon} r_{m,f}
$$

$$
\leq D'_0 \left(\tau_B^{1/2} + \frac{D_{\text{oracle}}}{\sqrt{m}} \right) K \left\| \beta^* \right\|_2 r_{m,f} + D_0 M_{\epsilon} r_{m,f}
$$

The lemma is thus proved. \square

Proof of Lemma [15.](#page-17-3) Recall that we require

$$
d \le C_A \left\{ c'C_{\phi} \wedge 2 \right\} \frac{f}{\log m} \text{ where } C_{\phi} = \frac{\|B\|_2 + a_{\max}}{D^2} D_{\phi}
$$

where $C_A = \frac{1}{128M_A^2}$ and $b_0^2 \ge ||\beta^*||_2^2 \ge \phi b_0^2$.

The proof for $d \leq s_0/32$ follows exactly that of Lemma [10.](#page-16-2) In order to show the second inequality, we follow the same line of arguments where we need to replace one inequality. By definition of D'_0 , we have $||B||_2 + a_{\text{max}} \le (D'_0)^2 \le 2(||B||_2 + a_{\text{max}})$. Now suppose that for $C_\phi = \frac{||B||_2 + a_{\text{max}}}{D^2} D_\phi$

$$
d \quad := \quad C_A c' C_\phi \frac{f}{\log m} \le C_A \frac{f}{\log m} \left(\frac{C_0 D_0'}{CD}\right)^2 D_\phi
$$

where $1 \le D = \rho_{\text{max}}(s_0 + 1, A) + \tau_B \le D_2$ and $C = C_0 / \sqrt{c'}$.

$$
d \leq C_{A}c'C_{\phi} \frac{f}{\log m} \leq \frac{1}{128M_{A}^{2}} \left(\frac{C_{0}D_{0}'}{CD}\right)^{2} D_{\phi} \frac{f}{\log m}
$$

$$
\leq \frac{1}{2} \left(\frac{1}{16CD}\right)^{2} 4C_{0}^{2} (D_{0}')^{2} D_{\phi} \frac{f}{M_{A}^{2} \log m}
$$

$$
\leq \frac{1}{2} \frac{(s_{0} + 1)^{2} \log m}{\alpha^{2}} \left(\frac{\psi}{b_{0}}\right)^{2} \leq \frac{(s_{0})^{2} \log m}{\alpha^{2}} \left(\frac{\psi}{b_{0}}\right)^{2}
$$

where assuming that $s_0 \geq 3$, we have the following inequality by definition of s_0 and $\alpha = \lambda_{\min}(A)/2$

$$
\frac{2s_0^2}{\alpha^2} \ge \left(\frac{s_0+1}{\alpha}\right)^2 \ge \frac{\alpha^2}{(16CD)^4} \left(\frac{f}{\log m}\right)^2
$$

which is identical in the proof of Lemma [10,](#page-16-2) while we replace (85) with

$$
4C_0^2(D'_0)^2D_\phi = 4C_0^2(D'_0)^2\left(\frac{K^2M_\epsilon^2}{b_0^2} + \tau_B^+K^4\phi\right)
$$

$$
\leq 4C_0^2(D'_0)^2\frac{K^2}{b_0^2}\left(M_\epsilon + \tau_B^{+/2}K\left\|\beta^*\right\|_2\right)^2 \leq \left(\frac{\psi}{b_0}\right)^2
$$

where $D_{\phi} := \frac{K^2 M_{\epsilon}^2}{b_0^2} + \tau_B^+ K^4 \phi$ and $\psi = 2C_0 \left(D_0' K^2 (\tau_B^{1/2} + \frac{D_{\text{oracle}}}{\sqrt{m}}) ||\beta^*||_2 + D_0 M_{\epsilon} K \right)$ as in [\(44\)](#page-17-4). \square

E Proofs for the Conic Programming estimator

E.1 Proof of Lemmas [11](#page-16-1) and [13](#page-16-5)

We next provide proofs for Lemmas [11](#page-16-1) and [13](#page-16-5) in this section.

Proof of Lemma [11.](#page-16-1) Suppose event B_0 holds. Then by the proof of Lemma [6,](#page-13-1)

$$
\left\| \frac{1}{f} X^T (y - X\beta^*) + \frac{1}{f} \widehat{\text{tr}}(B) \beta^* \right\|_{\infty} = \left\| \widehat{\gamma} - \widehat{\Gamma} \beta^* \right\|_{\infty}
$$

$$
\leq 2C_0 D_2 K^2 \left\| \beta^* \right\|_2 \sqrt{\frac{\log m}{f}} + C_0 D_0 K M_{\epsilon} \sqrt{\frac{\log m}{f}}
$$

$$
=: \mu \left\| \beta^* \right\|_2 + \tau
$$

The lemma follows immediately for the chosen μ, τ as in [\(43\)](#page-16-3) given that $(\beta^*, ||\beta^*||_2) \in \Upsilon$. \Box

Proof of Lemma [12.](#page-16-4) By optimality of $(\widehat{\beta}, \widehat{t})$, we have

$$
\left\|\widehat{\beta}\right\|_{1} + \lambda \left\|\widehat{\beta}\right\|_{2} \le \left\|\widehat{\beta}\right\|_{1} + \lambda \widehat{t} \le \left\|\beta^*\right\|_{1} + \lambda \left\|\beta^*\right\|_{2}
$$

Thus we have for $S := \text{supp}(\beta^*)$,

$$
\left\|\widehat{\beta}\right\|_{1} = \left\|\widehat{\beta}_{S^{c}}\right\|_{1} + \left\|\widehat{\beta}_{S}\right\|_{1} \leq \left\|\beta^{*}\right\|_{1} + \lambda(\left\|\beta^{*}\right\|_{2} - \left\|\widehat{\beta}\right\|_{2})
$$

Now by the triangle inequality,

$$
\begin{aligned}\n\left\|\hat{\beta}_{S^c}\right\|_1 &= \|v_{S^c}\|_1 \leq \| \beta_S^*\|_1 - \left\|\hat{\beta}_S\right\|_1 + \lambda(\|\beta^*\|_2 - \left\|\hat{\beta}\right\|_2) \\
&\leq \|v_S\|_1 + \lambda(\|\beta^*\|_2 - \left\|\hat{\beta}\right\|_2) \\
&\leq \|v_S\|_1 + \lambda(\|\beta^*\|_2 - \left\|\hat{\beta}_S\right\|_2) \\
&= \|v_S\|_1 + \lambda \|v_S\|_2 \leq (1 + \lambda) \|v_S\|_1.\n\end{aligned}
$$

The lemma thus holds given

$$
\widehat{t} \ \leq \ \frac{1}{\lambda} (\|\beta^*\|_1 - \left\|\widehat{\beta}\right\|_1) + \|\beta^*\|_2 \leq \frac{1}{\lambda} \left\|v\right\|_1 + \|\beta^*\|_2
$$

 \Box

Proof of Lemma [13.](#page-16-5) Recall the following shorthand notation:

$$
D_0 = (\sqrt{\tau_B} + \sqrt{a_{\text{max}}})
$$
 and $D_2 = 2(||A||_2 + ||B||_2)$

First we rewrite an upper bound for $v = \beta - \beta^*$, $D = \text{tr}(B)$ and $D = \text{tr}(B)$

$$
\begin{aligned}\n\left\|X_0^T X_0 v\right\|_{\infty} &= \left\|(X - W)^T X_0 (\widehat{\beta} - \beta^*)\right\|_{\infty} \le \left\|X^T X_0 (\widehat{\beta} - \beta^*)\right\|_{\infty} + \left\|W^T X_0 v\right\|_{\infty} \\
&\le \left\|X^T (X \widehat{\beta} - y) - \widehat{D}\widehat{\beta}\right\|_{\infty} + \left\|X^T \epsilon\right\|_{\infty} + \left\|(X^T W - D)\widehat{\beta}\right\|_{\infty} \\
&+ \left\|(\widehat{D} - D)\widehat{\beta}\right\|_{\infty} + \left\|W^T X_0 v\right\|_{\infty}\n\end{aligned}
$$

where

$$
\left\|X^T X_0(\hat{\beta} - \beta^*)\right\|_{\infty} \leq \left\|X^T (X_0 \hat{\beta} - y + \epsilon)\right\|_{\infty}
$$

\n
$$
= \left\|X^T ((X - W)\hat{\beta} - y)\right\|_{\infty} + \left\|X^T \epsilon\right\|_{\infty}
$$

\n
$$
\leq \left\|X^T (X \hat{\beta} - y) - \hat{D}\hat{\beta}\right\|_{\infty} + \left\|X^T \epsilon\right\|_{\infty}
$$

\n
$$
+ \left\|(X^T W - D)\hat{\beta}\right\|_{\infty} + \left\|(\hat{D} - D)\hat{\beta}\right\|_{\infty}.
$$

On event \mathcal{B}_0 , we have by Lemma [12](#page-16-4) and the fact that $\widehat{\beta} \in \Upsilon$

$$
I := \left\|\hat{\gamma} - \hat{\Gamma}\hat{\beta}\right\|_{\infty} = \left\|\frac{1}{f}X^{T}(y - X\hat{\beta}) + \frac{1}{f}\hat{D}\hat{\beta}\right\|_{\infty} \leq \mu\hat{t} + \tau
$$

$$
\leq \mu(\frac{1}{\lambda}\|v\|_{1} + \|\beta^{*}\|_{2}) + \tau
$$

$$
= 2D_{2}Kr_{m,f}(\frac{1}{\lambda}\|v\|_{1} + \|\beta^{*}\|_{2}) + D_{0}r_{m,f}M_{\epsilon}
$$

and on event \mathcal{B}_4 ,

$$
II := \frac{1}{f} ||X^T \epsilon||_{\infty} \le \frac{1}{f} (||X_0^T \epsilon||_{\infty} + ||W^T \epsilon||_{\infty})
$$

$$
\le r_{m,f} M_{\epsilon}(a_{\max}^{1/2} + \sqrt{\tau_B}) = D_0 r_{m,f} M_{\epsilon}
$$

Thus on event \mathcal{B}_0 , we have

$$
I + II \le 2D_2 K r_{m,f}(\frac{1}{\lambda} ||v||_1 + ||\beta^*||_2) + 2D_0 r_{m,f} M_{\epsilon} = \mu((\frac{1}{\lambda} ||v||_1 + ||\beta^*||_2) + 2\tau.
$$

Now on event \mathcal{B}_6 , we have for $2D_1 \le D_2$

$$
\begin{array}{lcl} IV := \left\| (\widehat{D} - D)\widehat{\beta} \right\|_{\infty} & \leq & \left| \widehat{D} - D \right| \left\| \widehat{\beta} \right\|_{\infty} \leq 2 D_1 K \frac{1}{\sqrt{m}} r_{m,f}(\left\| \beta^* \right\|_{\infty} + \left\| v \right\|_{\infty}) \\ & \leq & D_2 K \frac{1}{\sqrt{m}} r_{m,f}(\left\| \beta^* \right\|_2 + \left\| v \right\|_1) \end{array}
$$

On event $\mathcal{B}_5 \cap \mathcal{B}_{10}$, we have

$$
III := \frac{1}{f} \left\| (X^T W - D)\hat{\beta} \right\|_{\infty} \leq \frac{1}{f} \left\| (X^T W - D)\beta^* \right\|_{\infty} + \frac{1}{f} \left\| (X^T W - D)v \right\|_{\infty} \n\leq \frac{1}{f} \left\| X_0^T W \beta^* \right\|_{\infty} + \left\| (W^T W - D)\beta^* \right\|_{\infty} \n+ \frac{1}{f} \left(\left\| (Z^T BZ - \text{tr}(B)I_m) \right\|_{\max} + \frac{1}{f} \left\| X_0^T W \right\|_{\max} \right) \left\| v \right\|_1 \n\leq r_{m,f} K \left(\frac{\|B\|_F}{\sqrt{f}} + \sqrt{\tau_B} a_{\max}^{1/2} \right) (\|v\|_1 + \|\beta^*\|_2) \n\text{and } V = \frac{1}{f} \left\| W^T X_0 v \right\|_{\infty} \leq \frac{1}{f} \left\| W^T X_0 \right\|_{\max} \left\| v \right\|_1 \leq r_{m,f} K \sqrt{\tau_B} a_{\max}^{1/2} \left\| v \right\|_1.
$$

Thus we have on $\mathcal{B}_0 \cap \mathcal{B}_{10}$, for $D_0 \leq D_2$ and $\tau_A = 1$

$$
III + IV + V \leq r_{m,f} K \left(\|B\|_2 + \tau_B + a_{\max} + \frac{2}{\sqrt{m}} (\|A\|_2 + \|B\|_2) \right) (\|v\|_1 + \|\beta^*\|_2)
$$

\n
$$
\leq r_{m,f} K (4 \|B\|_2 + 3 \|A\|_2) (\|v\|_1 + \|\beta^*\|_2)
$$

\n
$$
\leq 2D_2 K r_{m,f} (\|v\|_1 + \|\beta^*\|_2)
$$

\n
$$
\leq \mu (\|v\|_1 + \|\beta^*\|_2)
$$

Thus we have

$$
\left\| \frac{1}{f} X_0^T X_0 v \right\|_{\infty} \leq I + II + III + IV + V
$$

\n
$$
\leq \mu (\frac{1}{\lambda} \|v\|_1 + \|\beta^*\|_2) + 2D_0 M_{\epsilon} r_{m,f} + \mu (\|v\|_1 + \|\beta^*\|_2)
$$

\n
$$
\leq 2\mu \| \beta^* \|_2 + \mu (\frac{1}{\lambda} + 1) \|v\|_1 + 2\tau.
$$

The lemma thus holds. \square

F Proof for Theorem [5](#page-11-1)

We prove Lemmas [16](#page-18-1) to [18](#page-18-3) in this section.

Proof of Lemma [16.](#page-18-1) Suppose event B_0 holds. Then by the proof of Lemma [14,](#page-17-2) we have for $D'_0 =$ $||B||_2^{1/2} + a_{\text{max}}^{1/2}$ and $\tau_B^{+/2} = \sqrt{\tau_B} + \frac{D_{\text{oracle}}}{\sqrt{m}}$, where $D_{\text{oracle}} = 2(||B||_2^{1/2} + ||A||_2^{1/2})$ $\binom{1}{2}$,

$$
\left\|\widehat{\gamma} - \widehat{\Gamma}\beta^*\right\|_{\infty} \leq D'_0 \tau_B^{+/2} K r_{m,f} \left\|\beta^*\right\|_2 + D_0 M_{\epsilon} r_{m,f}.
$$

The lemma follows immediately for μ , τ as chosen in [\(45\)](#page-18-8). \Box

Proof of Lemma [17.](#page-18-9) We first show [\(46\)](#page-18-7) and [\(47\)](#page-18-7). Recall $r_{m,m} := 2C_0 \sqrt{\frac{\log m}{mf}} \ge 2C_0 \frac{\log^{1/2} m}{m}$. By Lemma [7,](#page-13-2) we have on event \mathcal{B}_6 ,

$$
|\hat{\tau}_B - \tau_B| \leq D_1 K^2 r_{m,m}.
$$

Moreover, we have under (A1) $1 = \tau_A \leq D_1 := \frac{\|A\|_F}{m^{1/2}} + \frac{\|B\|_F}{f^{1/2}}$ $\frac{D \parallel F}{f^{1/2}}$ in view of [\(75\)](#page-34-0). And

$$
D_1 \le ||A||_2 + ||B||_2 \le (\frac{D_{\text{oracle}}}{2})^2
$$

and hence

$$
\sqrt{D_1} \le \frac{D_{\text{oracle}}}{2} = ||B||_2^{1/2} + ||A||_2^{1/2}.
$$

By definition and construction, we have τ_B , $\hat{\tau}_B \ge 0$,

$$
\left| \widehat{\tau}_{B}^{1/2} - \tau_{B}^{1/2} \right| \leq \widehat{\tau}_{B}^{1/2} + \tau_{B}^{1/2};
$$

hence
$$
\left| \widehat{\tau}_{B}^{1/2} - \tau_{B}^{1/2} \right|^{2} \leq \left| (\widehat{\tau}_{B}^{1/2} + \tau_{B}^{1/2}) (\widehat{\tau}_{B}^{1/2} - \tau_{B}^{1/2}) \right| = |\widehat{\tau}_{B} - \tau_{B}|
$$

Thus, on event \mathcal{B}_6 , we have

and

$$
\left|\widehat{\tau}_{B}^{1/2} - \tau_{B}^{1/2}\right| \leq \sqrt{|\widehat{\tau}_{B} - \tau_{B}|} \leq \sqrt{D_1} Kr_{m,m}^{1/2} \leq \frac{D_{\text{oracle}}}{2} Kr_{m,m}^{1/2}
$$

Thus we have for $C_6 \ge D_{\text{oracle}} \ge 2\sqrt{D_1}$ and $D_{\text{oracle}} = 2(||A||_2^{1/2} + ||B||_2^{1/2})$ $\binom{1}{2}$,

$$
\hat{\tau}_{B}^{1/2} - \frac{D_{\text{oracle}}}{2} K r_{mm}^{1/2} \le \tau_{B}^{1/2} \le \hat{\tau}_{B}^{1/2} + \frac{D_{\text{oracle}}}{2} K r_{mm}^{1/2} \tag{88}
$$

Thus we have for $\tau_B^{+/2}$ $B_B⁺²$ as defined in [\(23\)](#page-10-1), [\(88\)](#page-43-1) and the fact that

$$
r_{m,m}^{1/2} := \sqrt{2C_0} \frac{(\log m)^{1/4}}{\sqrt{m}} \ge 2/\sqrt{m} \text{ for } m \ge 16 \text{ and } C_0 \ge 1,
$$

the following inequalities hold: for $K \geq 1$,

$$
\tau_B^{+/2} := \tau_B^{1/2} + D_{\text{oracle}} m^{-1/2}
$$
\n
$$
\leq \hat{\tau}_B^{1/2} + \frac{D_{\text{oracle}}}{2} K r_{mm}^{1/2} + \frac{D_{\text{oracle}}}{2} r_{m,m}^{1/2}
$$
\n
$$
\leq \hat{\tau}_B^{1/2} + D_{\text{oracle}} K r_{mm}^{1/2} \leq \tilde{\tau}_B^{1/2}
$$
\n(89)

where the last inequality holds by the choice of $\tilde{\tau}_{B}^{1/2} \ge \tilde{\tau}_{B}^{1/2} + D_{\text{oracle}} Kr_{mm}^{1/2}$ as in [\(30\)](#page-12-3). Moreover, we have on event B_6 , by (88)

$$
\widetilde{\tau}_{B}^{1/2} := \widehat{\tau}_{B}^{1/2} + C_6 K r_{mm}^{1/2} \le \tau_{B}^{1/2} + \frac{D_{\text{oracle}}}{2} K r_{mm}^{1/2} + C_6 K r_{mm}^{1/2}
$$
\n
$$
\le \tau_{B}^{1/2} + \frac{3}{2} C_6 K r_{mm}^{1/2}
$$
\n
$$
\widetilde{\tau}_{B} := (\widehat{\tau}_{B}^{1/2} + C_6 K r_{mm}^{1/2})^2 \le 2 \widehat{\tau}_{B} + 2 C_6^2 K^2 r_{mm}
$$
\n
$$
\le 2 \tau_{B} + 2 D_1 K^2 r_{m,m} + 2 C_6^2 K^2 r_{mm}
$$
\n
$$
\le 2 \tau_{B} + \frac{D_{\text{oracle}}^2}{2} K^2 r_{m,m} + 2 C_6^2 K^2 r_{mm} \le 2 \tau_{B} + 3 C_6^2 K^2 r_{mm}
$$

and thus [\(46\)](#page-18-7) and [\(47\)](#page-18-7) hold given that $2D_1 \le D_{\text{oracle}}^2/2 \le C_6^2/2$. Finally, we have

$$
\widetilde{\tau}_{B}^{1/2} \tau_{B}^{-} \leq (\tau_{B}^{1/2} + \frac{3}{2} C_{6} K r_{mm}^{1/2}) \tau_{B}^{-} \leq \frac{\tau_{B}^{1/2} + \frac{3}{2} C_{6} K r_{mm}^{1/2}}{\tau_{B}^{1/2} + 2 C_{6} K r_{m,m}^{1/2}} \leq 1
$$

for τ_B^- as defined in [\(26\)](#page-12-1). \Box

Remark F.1. *The set* Υ *in our setting is equivalent to the following: for* μ , τ *as defined in* [\(30\)](#page-12-3) *and* $\beta \in \mathbb{R}^m$,

$$
\Upsilon = \left\{ (\beta, t) \; : \; \left\| \frac{1}{f} X^T (y - X\beta) + \frac{1}{f} \widehat{\text{tr}}(B)\beta \right\|_{\infty} \le \mu t + \tau, \|\beta\|_2 \le t \right\}. \tag{90}
$$

Proof of Lemma [18.](#page-18-3) For the rest of the proof, we will follow the notation in the proof for Lemma [13.](#page-16-5) Notice that the bounds as stated in Lemma [12](#page-16-4) remain true with τ, μ chosen as in [\(45\)](#page-18-8), so long as $(\beta^*, \| \beta^* \|_2) \in$ Υ. This indeed holds by Lemma [16:](#page-18-1) for τ [\(29\)](#page-12-3) and μ [\(30\)](#page-12-3) as chosen in Theorem [5,](#page-11-1) we have by [\(89\)](#page-44-0),

$$
\mu \approx D_0' \tilde{\tau}_B^{1/2} K r_{m,f} \ge D_0' K r_{m,f} \tau_B^{+/2}
$$

where $\tau_B^{+/2} = (\sqrt{\tau_B} + \frac{D_{\text{oracle}}}{\sqrt{m}})$, which ensures that $(\beta^*, ||\beta^*||_2) \in \Upsilon$ by Lemma [16.](#page-18-1)

On event \mathcal{B}_0 , we have by Lemma [12](#page-16-4) and the fact that $\hat{\beta} \in \Upsilon$ as in [\(90\)](#page-44-1)

$$
I + II := \left\| \widehat{\gamma} - \widehat{\Gamma}\widehat{\beta} \right\|_{\infty} + \frac{1}{f} \left\| X^T \epsilon \right\|_{\infty}
$$

\n
$$
\leq \left\| \frac{1}{f} X^T (y - X\widehat{\beta}) + \frac{1}{f} \widehat{D}\widehat{\beta} \right\|_{\infty} + \tau \leq \mu \widehat{t} + 2\tau
$$

\n
$$
\leq \mu (\frac{1}{\lambda} \|v\|_1 + \|\beta^*\|_2) + 2\tau
$$

for μ , τ as chosen in [\(30\)](#page-12-3) and [\(29\)](#page-12-3) respectively. Now on event \mathcal{B}_6 , we have

$$
IV := \left\| (\widehat{D} - D)\widehat{\beta} \right\|_{\infty} \leq \left\| \widehat{D} - D \right\| \left\| \widehat{\beta} \right\|_{\infty} \leq 2D_1 K \frac{1}{\sqrt{m}} r_{m,f}(\|\beta^*\|_{\infty} + \|v\|_{\infty})
$$

$$
\leq D'_0 \frac{D_{\text{oracle}}}{\sqrt{m}} K r_{m,f}(\|\beta^*\|_2 + \|v\|_1)
$$

where $2D_1 \leq D_{\text{oracle}} D'_0$ for $1 \leq D'_0 := ||B||_2^{1/2} + a_{\text{max}}^{1/2}$ and $D_{\text{oracle}} = 2 (||B||_2^{1/2} + ||A||_2^{1/2})$ 2 , where $a_{\text{max}} \geq \tau_A = 1$ under (A1). Hence

$$
III + IV + V \leq r_{m,f} K \sqrt{\tau_B} \left(\|B\|_2^{1/2} + a_{\text{max}}^{1/2} \right) \left(\|v\|_1 + \|\beta^*\|_2 \right)
$$

+2D₁K $\frac{1}{\sqrt{m}} r_{m,f} (\|\beta^*\|_2 + \|v\|_1) + r_{m,f} K \sqrt{\tau_B} a_{\text{max}}^{1/2} \|v\|_1$

$$
\leq D'_0 K r_{m,f} (\|v\|_1 + \|\beta^*\|_2) (\sqrt{\tau_B} + \frac{D_{\text{oracle}}}{\sqrt{m}}) + r_{m,f} K \sqrt{\tau_B} a_{\text{max}}^{1/2} \|v\|_1
$$

$$
\leq D'_0 K r_{m,f} \tau_B^{+/2} (\|v\|_1 + \|\beta^*\|_2) + D'_0 K r_{m,f} \sqrt{\tau_B} \|v\|_1
$$

$$
\leq C_0 D'_0 K^2 \sqrt{\frac{\log m}{f}} (\tau_B^{1/2} + \frac{D_{\text{oracle}}}{\sqrt{m}}) (2 \|v\|_1 + \|\beta^*\|_2)
$$

$$
\leq \mu (2 \|v\|_1 + \|\beta^*\|_2)
$$

for μ as defined in [\(30\)](#page-12-3) in view of [\(89\)](#page-44-0). Thus we have

$$
I + II + III + IV + V \leq \mu(\frac{1}{\lambda} ||v||_1 + ||\beta^*||_2) + 2\tau + \mu(2 ||v||_1 + ||\beta^*||_2)
$$

= $2\mu((1 + \frac{1}{2\lambda}) ||v||_1 + ||\beta^*||_2) + 2\tau$

and the improved bounds as stated in the Lemma thus holds. \square

G Some geometric analysis results

Let us define the following set of vectors in \mathbb{R}^m :

$$
Cone(s_0) := \{v : ||v||_1 \le \sqrt{s_0} ||v||_2\}
$$

For each vector $x \in \mathbb{R}^m$, let T_0 denote the locations of the s_0 largest coefficients of x in absolute values. Any vector $x \in S^{m-1}$ satisfies:

$$
\left\|x_{T_0^c}\right\|_{\infty} \leq \left\|x_{T_0}\right\|_{1} / s_0 \leq \frac{\left\|x_{T_0}\right\|_{2}}{\sqrt{s_0}} \tag{91}
$$

We need to state the following result from [Mendelson et al.](#page-59-16) [\(2008\)](#page-59-16). Let S^{m-1} be the unit sphere in \mathbb{R}^m , for $1 \leq s \leq m$,

$$
U_s := \{ x \in \mathbf{R}^m : |\operatorname{supp}(x)| \le s \} \tag{92}
$$

The sets U_s is an union of the s-sparse vectors. The following three lemmas are well-known and mostly standard; See [Mendelson et al.](#page-59-16) [\(2008\)](#page-59-16) and [Loh and Wainwright](#page-59-0) [\(2012\)](#page-59-0).

Lemma 31. *For every* $1 \leq s_0 \leq m$ *and every* $I \subset \{1, \ldots, m\}$ *with* $|I| \leq s_0$ *,*

$$
\sqrt{|I|}B_1^m\cap S^{m-1}\subset 2\operatorname{conv}\left(U_{s_0}\cap S^{m-1}\right)=:2\operatorname{conv}\left(\bigcup_{|J|\leq s_0}E_J\cap S^{m-1}\right)
$$

and moreover, for $\rho \in (0, 1]$ *.*

$$
\sqrt{|I|}B_1^m \cap \rho B_2^m \subset (1+\rho)\operatorname{conv}\left(U_{s_0} \cap B_2^m\right) =: (1+\rho)\operatorname{conv}\left(\bigcup_{|J| \leq s_0} E_J \cap S^{m-1}\right)
$$

Proof. Fix $x \in \mathbb{R}^m$. Let x_{T_0} denote the subvector of x confined to the locations of its s_0 largest coefficients in absolute values; moreover, we use it to represent its 0-extended version $x' \in \mathbb{R}^p$ such that $x'_{T^c} = 0$ and $x'_{T_0} = x_{T_0}$. Throughout this proof, T_0 is understood to be the locations of the s_0 largest coefficients in absolute values in x .

Moreover, let $(x_i^*)_{i=1}^m$ be non-increasing rearrangement of $(|x_i|)_{i=1}^m$. Denote by

$$
L = \sqrt{s_0} B_1^m \cap \rho B_2^m
$$

$$
R = 2 \operatorname{conv} \left(\bigcup_{|J| \le s} E_J \cap B_2^m \right) = 2 \operatorname{conv} \left(E \cap B_2^m \right)
$$

Any vector $x \in \mathbb{R}^m$ satisfies:

$$
\left\|x_{T_0^c}\right\|_{\infty} \leq \left\|x_{T_0}\right\|_{1} / s_0 \leq \frac{\left\|x_{T_0}\right\|_{2}}{\sqrt{s_0}} \tag{93}
$$

It follows that for any $\rho > 0$, $s_0 \ge 1$ and for all $z \in L$, we have the i^{th} largest coordinate in absolute value in z is at most $\sqrt{s_0}/i$,

$$
\sup_{z \in L} \langle x, z \rangle \leq \max_{\|z\|_2 \leq \rho} \langle x_{T_0}, z \rangle + \max_{\|z\|_1 \leq \sqrt{s_0}} \langle x_{T_0^c}, z \rangle
$$

$$
\leq \rho \|x_{T_0}\|_2 + \|x_{T_0^c}\|_{\infty} \sqrt{s_0}
$$

$$
\leq \|x_{T_0}\|_2 (\rho + 1)
$$

where clearly $\max_{\|z\|_2\leq \rho} \langle x_{T_0}, z \rangle = \rho \sum_{i=1}^{s_0} (x_i^*)^{1/2}$. And denote by $S^J := S^{m-1} \cap E_J$,

$$
\sup_{z \in R} \langle x, z \rangle = (1 + \rho) \max_{J:|J| \le s_0} \max_{z \in S^J} \langle x, z \rangle
$$

$$
= (1 + \rho) \|x_{T_0}\|_2
$$

given that for a convex function $\langle x, z \rangle$, the maximum happens at an extreme point, and in this case, it happens for z such that z is supported on T_0 , such that $z_{T_0} = \frac{x_{T_0}}{\ln x}$ $\frac{x_{T_0}}{\|x_{T_0}\|_2}$, and $z_{T_0^c} = 0$. \Box

Lemma 32. Let $1/5 > \delta > 0$. Let $E = \bigcup_{|J| \le s_0} E_J$ for $0 < s_0 < m/2$ and $k_0 > 0$. Let Δ be a $m \times m$ *matrix such that*

$$
|u^T \Delta v| \le \delta \quad \forall u, v \in E \cap S^{m-1}
$$
\n(94)

Then for all $v \in (\sqrt{s_0}B_1^m \cap B_2^m)$, we have

$$
\left| v^T \Delta v \right| \leq 4\delta. \tag{95}
$$

Proof. First notice that

$$
\max_{\upsilon \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} \left|\upsilon^T \Delta \upsilon\right| \le \max_{w, u \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} \left|w^T \Delta u\right| \tag{96}
$$

 $\overline{}$ $\overline{}$

Now that we have decoupled u and w on the RHS of [\(96\)](#page-47-0), we first fix u. Then for any fixed $u \in S^{m-1}$ and matrix $\Delta \in \mathbb{R}^{m \times m}$, $f(w) = |w^T \Delta u|$ is a convex function of w, and hence for $w \in (\sqrt{s_0} B_1^m \cap B_2^m) \subset$ 2 conv $\left(\bigcup_{|J|\leq s_0} E_J \cap S^{m-1}\right)$,

$$
\max_{w \in (\sqrt{s_0}B_1^m \cap B_2^m)} |w^T \Delta u| \le 2 \max_{w \in \text{conv}(E \cap S^{m-1})} |w^T \Delta u|
$$

= $2 \max_{w \in E \cap S^{m-1}} |w^T \Delta u|$

where the maximum occurs at an extreme point of the set conv $(E \cap S^{m-1})$, because of the convexity of the function $f(w)$,

Clearly the RHS of [\(96\)](#page-47-0) is bounded by

$$
\max_{u,w \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} |w^T \Delta u| = \max_{u \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} \max_{w \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} \max_{w \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} |w^T \Delta u|
$$

$$
\leq 2 \max_{u \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} \max_{w \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} |w^T \Delta u|
$$

$$
= 2 \max_{u \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} g(u)
$$

where the function g of $u \in (\sqrt{s_0}B_1^m \cap B_2^m)$ is defined as

$$
g(u) = \max_{w \in (E \cap S^{m-1})} |w^T \Delta u|
$$

which is convex since it is the maximum of a function $f_w(u) := |w^T \Delta u|$ which is convex in u for each $w \in (E \cap S^{m-1})$. Thus we have for $u \in (\sqrt{s_0}B_1^m \cap B_2^m) \subset 2$ conv $\left(\bigcup_{|J| \le s_0} E_J \cap S^{m-1}\right) =$: 2 conv $(E \cap S^{m-1})$

$$
\max_{u \in \left(\sqrt{s_0}B_1^m \cap B_2^m\right)} g(u) \le 2 \max_{u \in \text{conv}(E \cap S^{m-1})} g(u)
$$
\n
$$
= 2 \max_{u \in E \cap S^{m-1}} g(u) \tag{97}
$$

$$
= 2 \max_{u \in E \cap S^{m-1}} \max_{w \in E \cap S^{m-1}} |w^T \Delta u| \le 4\delta
$$
 (98)

where [\(97\)](#page-47-1) holds given that the maximum occurs at an extreme point of the set conv $(E \cap B_2^m)$, because of the convexity of the function $g(u)$. \Box

Corollary 33. *Suppose all conditions in Lemma* [32](#page-46-0) *hold. Then* $\forall v \in \text{Cone}(s_0)$ *,*

$$
\left|v^T \Delta v\right| \le 4\delta \|v\|_2^2. \tag{99}
$$

Proof. It is sufficient to show that $\forall v \in \text{Cone}(s_0) \cap S^{m-1}$,

$$
\left|v^T \Delta v\right| \ \leq \ 4\delta.
$$

Denote by Cone := $Cone(s_0)$. Clearly this set of vectors satisfy:

$$
\operatorname{Cone} \cap S^{m-1} \subset \left(\sqrt{s_0} B_1^m \cap B_2^m\right)
$$

Thus [\(99\)](#page-47-2) follows from [\(95\)](#page-46-1). \Box

Remark G.1. *Suppose we relax the definition of* $Cone(s_0)$ *to be:*

$$
Cone(s_0) := \{v : ||v||_1 \le 2\sqrt{s_0} ||v||_2\}
$$

Clearly, $Cone(s_0, 1) \subset Cone(s_0)$ *. given that* $\forall u \in Cone(s_0, 1)$ *, we have*

$$
||u||_1 \le 2 ||u_{T_0}||_1 \le 2\sqrt{s_0} ||u_{T_0}||_2 \le 2\sqrt{s_0} ||u||_2
$$

Lemma 34. *Suppose all conditions in Lemma [32](#page-46-0) hold. Then for all* $v \in \mathbb{R}^m$,

$$
\left| \nu^T \Delta \nu \right| \le 4\delta (\| \nu \|_2^2 + \frac{1}{s_0} \| \nu \|_1^2)
$$
 (100)

Proof. The lemma follows given that $\forall v \in \mathbb{R}^m$, one of the following must hold:

$$
\text{if } v \in \text{Cone}(s_0) \quad \left| v^T \Delta v \right| \le 4\delta \left\| v \right\|_2^2 \tag{101}
$$

otherwise
$$
|v^T \Delta v| \le \frac{4\delta}{s_0} ||v||_1^2,
$$
 (102)

leading to the same conclusion in [\(100\)](#page-48-1). We have shown [\(101\)](#page-48-2) in Lemma [32.](#page-46-0) Let $Cone(s_0)^c$ be the complement set of $\text{Cone}(s_0)^c$ in \mathbb{R}^m . That is, we focus now on the set of vectors such that

Cone
$$
(s_0)^c := \{v : ||v||_1 \ge \sqrt{s_0} ||v||_2\}
$$

and show that for $u = \sqrt{s_0} \frac{v}{||v||}$ $\frac{v}{\|v\|_1}$

$$
\frac{|v^T \Delta v|}{\|v\|_1^2} := \frac{1}{s_0} |u^T \Delta u| \le \frac{1}{s_0} \delta
$$

where the last inequality holds by Lemma [32](#page-46-0) given that

$$
u\in (\sqrt{s_0}B_1^m\cap B_2^m)\subset 2\operatorname{conv}\left(\bigcup_{|J|\leq s_0}E_J\cap B_2^m\right)
$$

and thus

$$
\frac{\left|v^T \Delta v\right|}{\left\|v\right\|_1^2} \leq \frac{1}{s_0} \sup_{u \in \sqrt{s_0} B_1^m \cap B_2^m} \left|u^T \Delta u\right| \leq \frac{1}{s_0} 4\delta
$$

 \Box

H Proof of Corollary [19](#page-18-2)

First we show that for all $v \in \mathbb{R}^m$, [\(103\)](#page-49-2) holds. It is sufficient to check that the condition [\(94\)](#page-46-2) in Lemma [32](#page-46-0) holds. Then, [\(103\)](#page-49-2) follows from Lemma [34:](#page-48-0) for $v \in \mathbb{R}^m$,

$$
\left|v^T \Delta v\right| \le 4\delta(\|v\|_2^2 + \frac{1}{\zeta} \|v\|_1^2) \le \frac{1}{2}\lambda_{\min}(A)(\|v\|_2^2 + \frac{1}{\zeta} \|v\|_1^2). \tag{103}
$$

The Lower and Upper RE conditions thus immediately follow. The Corollary is thus proved. \Box

I Proof of Theorem [20](#page-18-6)

We first state the following preliminary results in Lemmas [35](#page-49-3) and [36;](#page-49-4) their proofs appear in Section [K.](#page-55-0) Throughout this section, the choice of $C = C_0/\sqrt{c'}$ satisfies the conditions on C in Lemmas [35](#page-49-3) and [36,](#page-49-4) where recall $\min\{C_0, C_0^2\} \ge 4/c$ for c as defined in Theorem [26.](#page-27-2) For a set $J \subset \{1, \ldots, m\}$, denote $F_J = A^{1/2} E_J$ where recall $E_J = \text{span}\{e_j : j \in J\}.$

Lemma 35. *Suppose all conditions in Theorem [20](#page-18-6) hold. Let*

$$
E = \bigcup_{|J|=k} E_J \cap S^{m-1}.
$$

Suppose that for some $c' > 0$ *and* $\varepsilon \leq \frac{1}{C}$ $\frac{1}{C}$ *, where* $C = C_0 / \sqrt{c'}$ *,*

$$
r(B) := \frac{\text{tr}(B)}{\|B\|_2} \ge c' k K^4 \frac{\log(3em/k\varepsilon)}{\varepsilon^2}.
$$
 (104)

Then for all vectors $u, v \in E \cap S^{m-1}$, *on event* \mathcal{B}_1 , *where* $\mathbb{P}\left(\mathcal{B}_1\right) \geq 1-2\exp\left(-c_2\varepsilon^2\frac{\text{tr}(B)}{K^4\|B\|_2}\right)$ *for* $c_2 \geq 2$ *,*

$$
\left| u^T Z^T B Z v - \mathbb{E} u^T Z^T B Z v \right| \leq 4C \varepsilon \text{tr}(B).
$$

Lemma 36. *Suppose that* $\epsilon \leq 1/C$ *, where* C *is as defined in Lemma [35.](#page-49-3) Suppose that* [\(104\)](#page-49-5) *holds. Let*

$$
E = \bigcup_{|J|=k} E_J \quad and \quad F = \bigcup_{|J|=k} F_J. \tag{105}
$$

Then on event \mathcal{B}_2 , where $\mathbb{P}(\mathcal{B}_2) \geq 1 - 2 \exp \left(-c_2 \varepsilon^2 \frac{\text{tr}(B)}{K^4 \|B\|_2}\right)$ *for* $c_2 \geq 2$ *, we have for all vectors* $u \in$ $E \cap S^{m-1}$ and $w \in F \cap S^{m-1}$,

$$
\left| w^T Z_1^T B^{1/2} Z_2 u \right| \le \frac{C \varepsilon \text{tr}(B)}{(1 - \varepsilon)^2 \|B\|_2^{1/2}} \le 4C \varepsilon \text{tr}(B) / \|B\|_2^{1/2}
$$

where Z_1 , Z_2 *are independent copies of* Z *, as defined in Theorem [20.](#page-18-6)*

In fact, the same conclusion holds for all $y, w \in F \cap S^{m-1}$; and in particular, for $B = I$, we have the following.

Corollary 37. *Suppose all conditions in Lemma* [35](#page-49-3) *hold. Suppose that* $F = A^{1/2}E$ *for* E *as defined in Lemma [35.](#page-49-3) Let*

$$
f \ge c' k K^4 \frac{\log(3em/k\varepsilon)}{\varepsilon^2}.
$$
 (106)

Then on event \mathcal{B}_3 , where $\mathbb{P}(\mathcal{B}_3) \geq 1 - 2 \exp(-c_2 \varepsilon^2 f \frac{1}{K^4})$, we have for all vectors $w, y \in F \cap S^{m-1}$ and $\varepsilon \leq 1/C$ for *C* is as defined in Lemma [35,](#page-49-3)

$$
\left| y^T \left(\frac{1}{f} Z^T Z - I \right) w \right| \leq 4C\varepsilon. \tag{107}
$$

We prove Lemmas [35](#page-49-3) and [36](#page-49-4) and Corollary [37](#page-50-0) in Section [K.](#page-55-0) We are now ready to prove Theorem [20.](#page-18-6)

Proof of Theorem [20.](#page-18-6) Recall the following for $X_0 = Z_1 A^{1/2}$,

$$
\Delta := \widehat{\Gamma}_A - A := \frac{1}{f} X^T X - \frac{1}{f} \widehat{\text{tr}}(B) I_m - A
$$

=
$$
(\frac{1}{f} X_0^T X_0 - A) + \frac{1}{f} (W^T X_0 + X_0^T W) + \frac{1}{f} (W^T W - \widehat{\text{tr}}(B) I_m).
$$

Notice that

$$
\begin{split}\n\left|u^{T}(\widehat{\Gamma}_{A}-A)v\right| &= \left|u^{T}(X^{T}X-\widehat{\text{tr}}(B)I_{m}-A)v\right| \\
&\leq \left|u^{T}(\frac{1}{f}X_{0}^{T}X_{0}-A)v\right| + \left|u^{T}\frac{1}{f}(W^{T}X_{0}+X_{0}^{T}W)v\right| + \left|u^{T}(\frac{1}{f}W^{T}W-\frac{\widehat{\text{tr}}(B)}{f}I_{m})v\right| \\
&\leq \left|u^{T}A^{1/2}\frac{1}{f}Z_{1}^{T}Z_{1}A^{1/2}v - u^{T}Av\right| + \left|u^{T}\frac{1}{f}(W^{T}X_{0}+X_{0}^{T}W)v\right| \\
&\quad + \left|u^{T}(\frac{1}{f}Z_{2}^{T}BZ_{2}-\tau_{B}I_{m})v\right| + \frac{1}{f}|\widehat{\text{tr}}(B)-\text{tr}(B)|\left|u^{T}v\right| =: I + II + III + IV.\n\end{split}
$$

For $u \in E \cap S^{m-1}$, define $h(u) := \frac{A^{1/2}u}{\|A^{1/2}u\|}$ $\frac{A^{1/2}u}{\|A^{1/2}u\|_2}$. The conditions in [\(104\)](#page-49-5) and [\(106\)](#page-50-1) hold for k. We first bound the middle term as follows. Fix $u, v \in E \cap S^{m-1}$ Then on event B_2 , for $\Upsilon = Z_1^T B^{1/2} Z_2$,

$$
\begin{split} \left| u^T (W^T X_0 + X_0^T W) v \right| &= \left| u^T Z_2^T B^{1/2} Z_1 A^{1/2} v + u^T A^{1/2} Z_1^T B^{1/2} Z_2 v \right| \\ &\leq \left| u^T \Upsilon^T h(v) \right| \left\| A^{1/2} v \right\|_2 + \left| h(u)^T \Upsilon v \right| \left\| A^{1/2} u \right\|_2 \\ &\leq \left| 2 \max_{w \in F \cap S^{m-1}, v \in E \cap S^{m-1}} \left| w^T \Upsilon v \right| \rho_{\text{max}}^{1/2}(k, A) \\ &\leq \left| 8C \varepsilon \text{tr}(B) \left(\frac{\rho_{\text{max}}(k, A)}{\|B\|_2} \right)^{1/2} \right|. \end{split}
$$

We now use Lemma [35](#page-49-3) to bound both I and III. We have for C as defined in Lemma [35,](#page-49-3) on event $\mathcal{B}_1 \cap \mathcal{B}_3$,

$$
\left|u^T(Z_2^T B Z_2 - \text{tr}(B) I_m)v\right| \leq 4C\varepsilon \text{tr}(B).
$$

Moreover, by Corollary [37,](#page-50-0) we have on event B_3 , for all $u, v \in E \cap S^{m-1}$,

$$
\begin{array}{rcl} \left| u^T (\frac{1}{f} X_0^T X_0 - A) v \right| & = & \left| u^T A^{1/2} Z^T Z A^{1/2} v - u^T A v \right| \\ & = & \left| h(u)^T (\frac{1}{f} Z^T Z - I) h(v) \right| \left\| A^{1/2} u \right\|_2 \left\| A^{1/2} v \right\|_2 \\ & \leq & \frac{1}{f} \max_{w, y \in F \cap S^{m-1}} \left| w^T (Z^T Z - I) y \right| \rho_{\text{max}}(k, A) \\ & \leq & 4C \varepsilon \rho_{\text{max}}(k, A). \end{array}
$$

Thus we have on event $\mathcal{B}_1 \cap \mathcal{B}_2 \cap \mathcal{B}_3$ and for $\tau_B := \text{tr}(B)/f$

$$
I + II + III \le 4C\varepsilon \left(\rho_{\max}(k, A) + 2\tau_B \left(\frac{\rho_{\max}(k, A)}{\|B\|_2} \right)^{1/2} + \tau_B \right)
$$

$$
\le 8C\varepsilon (\tau_B + \rho_{\max}(k, A)).
$$

On event \mathcal{B}_6 , we have for D_1 as defined in Lemma [7,](#page-13-2)

$$
IV \leq |\hat{\tau}_B - \tau_B| \leq 2C_0 D_1 K^2 \sqrt{\frac{\log m}{fm}}.
$$

The theorem thus holds by the union bound. \square

J Proof for Theorem [21](#page-19-1)

We first state the following bounds in [\(108\)](#page-51-1) before we prove Theorem [21.](#page-19-1) On event A_2 , where $\mathbb{P}(A_2) \geq$ $1 - 2 \exp \left(-c_3 \varepsilon^2 \frac{\text{tr}(A)}{K^4 ||A||_2}\right)$ $\breve{}$

$$
\forall u, w \in S^{f-1} \quad \left| u^T Z_1 A^{1/2} Z_2^T w \right| \le \frac{4C \varepsilon \text{tr}(A)}{\|A\|_2^{1/2}}.
$$
\n(108)

To see this, first note that by Lemma [27,](#page-27-1) we have for $t = C \varepsilon \text{tr}(A) / ||A||_2^{1/2}$ $\frac{1}{2}$ and $\varepsilon \leq 1/2$,

$$
\mathbb{P}\left(\left|u^T Z_1 A^{1/2} Z_2^T w\right| > t\right) \leq 2 \exp\left(-c \min\left(\frac{C^2 \varepsilon^2 \text{tr}(A)}{K^4 \|A\|_2}, \frac{C \varepsilon \text{tr}(A)}{K^2 \|A\|_2}\right)\right)
$$

$$
\leq 2 \exp\left(-c \min\left(C^2, 2C\right) \frac{\varepsilon^2 \text{tr}(A)}{K^4 \|A\|_2}\right)
$$

where recall

$$
C' = cc' \min (2C, C^2) > 4.
$$

Before we proceed, we state the following well-known result on *volumetric estimate*; see e.g. [Milman and Schechtman](#page-59-17) [\(1986](#page-59-17)).

Lemma 38. *Given* $m \geq 1$ *and* $\varepsilon > 0$. *There exists an* ε *-net* $\Pi \subset B_2^m$ *of* B_2^m *with respect to the Euclidean metric such that* $B_2^m \subset (1 - \varepsilon)^{-1}$ conv Π *and* $|\Pi| \le (1 + 2/\varepsilon)^m$. Similarly, there exists an ε -net of the *sphere* S^{m-1} , $\Pi' \subset S^{m-1}$ *such that* $|\Pi'| \leq (1 + 2/\varepsilon)^m$ *.*

Choose an ε -net $\Pi \subset S^{f-1}$ such that $|\Pi| \leq (1+2/\varepsilon)^f = \exp(f \log(3/\varepsilon))$. The existence of such Π is guaranteed by Lemma [38.](#page-51-2) By the union bound and Lemma [27,](#page-27-1) we have for some $C \geq 2$ and $c' \geq 1$ large enough such that

$$
\mathbb{P}\left(\exists u, w \in \Pi s.t. \left|u^T Z_1 A^{1/2} Z_2^T w\right| \ge C\varepsilon \frac{\text{tr}(A)}{\|A\|_2^{1/2}}\right) \le 2 \exp\left(-c_3 \frac{\varepsilon^2 \text{tr}(A)}{K^4 \|A\|_2}\right).
$$

Hence, [\(108\)](#page-51-1) follows from a standard approximation argument.

Lemma 39. Let $\varepsilon > 0$. Let Z as defined in Definition [1.2.](#page-2-1) Assume that

$$
\frac{\text{tr}(A)}{\|A\|} \ge c' f \frac{\log(3/\varepsilon)}{\varepsilon^2}
$$

.

.

Then

$$
\mathbb{P}\left(\exists x \in S^{f-1} \left\| \left\| A^{1/2} Z^T x \right\|_2 - (\text{tr}(A))^{1/2} \right| > \varepsilon (\text{tr}(A))^{1/2} \right) \le \exp\left(-c\varepsilon^2 \frac{\text{tr}(A)}{K^4 \|A\|}\right)
$$

Proof. Let $x \in S^{f-1}$. Then $Y = Z^T x \in \mathbb{R}^m$ is a random vector with independent coordinates satisfying $\mathbb{E}Y_j = 0$ and $||Y_j||_{\psi_2} \leq CK$ for all $j \in 1...m$. The last estimate follows from Hoeffding inequality. By Theorem 2.1 [Rudelson and Vershynin](#page-59-15) [\(2013](#page-59-15)),

$$
\mathbb{P}\left(\left|\left\|A^{1/2}Y\right\|_{2} - (\text{tr}(A))^{1/2}\right| > \varepsilon(\text{tr}(A))^{1/2}\right) \leq \exp\left(-c\varepsilon^{2} \frac{\text{tr}(A)}{K^{4} \|A\|}\right).
$$

Choose an ε -net $\Pi \subset S^{f-1}$ such that $|\Pi| \leq (3/\varepsilon)^f$. By the union bound and the assumption of the Lemma,

$$
\mathbb{P}\left(\exists x \in \Pi \left\| \left\| A^{1/2} Z^T x \right\|_2 - (\text{tr}(A))^{1/2} \right| > \varepsilon (\text{tr}(A))^{1/2} \right) \leq \left\| \Pi \right\| \cdot \exp\left(-c\varepsilon^2 \frac{\text{tr}(A)}{K^4 \left\| A \right\|} \right)
$$

$$
\leq \exp\left(-c' \varepsilon^2 \frac{\text{tr}(A)}{K^4 \left\| A \right\|} \right).
$$

A standard approximation argument shows that if $\left|\left|\left|A^{1/2}Z^Tx\right|\right|_2 - (\text{tr}(A))^{1/2}\right| \leq \varepsilon (\text{tr}(A))^{1/2}$ for all $x \in \Pi$, then $\left|\left|\left|A^{1/2}Z^Tx\right|\right|_2 - (\text{tr}(A))^{1/2}\right| \leq 3\varepsilon(\text{tr}(A))^{1/2}$ for all $x \in S^{f-1}$. This finishes the proof of the Lemma. \Box

Proof of Theorem [21.](#page-19-1) First we write

$$
XXT - tr(A)If = (Z1A1/2 + B1/2Z2)(Z1A1/2 + B1/2Z2)T - tr(A)If
$$

= (Z₁A^{1/2} + B^{1/2}Z₂)(Z₂^TB^{1/2} + A^{1/2}Z₁^T) - tr(A)I_f
= Z₁A^{1/2}Z₂^TB^{1/2} + B^{1/2}Z₂Z₂^TB^{1/2} + B^{1/2}Z₂A^{1/2}Z₁^T + Z₁AZ₁^T - tr(A)I_f.

Hence,

$$
\left| \frac{u^T (XX^T)u}{m} - \frac{u^T \text{tr}(A)Iu}{m} - u^T Bu \right| \le \left| \frac{1}{m} u^T Z_1 A Z_1^T u - \frac{\text{tr}(A)}{m} u^T u \right| + \left| \frac{1}{m} u^T B^{1/2} Z_2 Z_2^T B^{1/2} u - u^T Bu \right| + \frac{2}{m} \left| u^T Z_1 A^{1/2} Z_2^T B^{1/2} u \right|.
$$

where by [\(108\)](#page-51-1), we have on event A_2 , for $\tau_A := \frac{\text{tr}(A)}{m}$ and $w := \frac{B^{1/2}u}{\|B^{1/2}u\|_2}$,

$$
\frac{2}{m} \left| u^T Z_1 A^{1/2} Z_2^T B^{1/2} u \right| = \frac{2}{m} \left| u^T Z_1 A^{1/2} Z_2^T w \right| \left\| B^{1/2} u \right\|_2
$$

$$
\leq \frac{8C \varepsilon \text{tr}(A) \left\| B^{1/2} u \right\|_2}{\left\| A \right\|_2^{1/2} m} =: 8C \varepsilon \tau_A \left\| B^{1/2} u \right\|_2 / \|A\|_2^{1/2}.
$$

Moreover, by the union bound and Lemma [39,](#page-52-0) we have on event A_1 , where $\mathbb{P}(\mathcal{A}_1) \geq 1 - \exp(c \varepsilon^2 \frac{m}{K^4})$ $\exp(c\varepsilon^2 \frac{\text{tr}(A)}{K^4 ||A||_2}),$

$$
(1-\varepsilon)\left\|B^{1/2}u\right\|_2 \leq \left\|\frac{1}{\sqrt{m}}\left\|Z_2B^{1/2}u\right\|_2 \leq (1+\varepsilon)\left\|B^{1/2}u\right\|_2
$$

$$
(1-\varepsilon)\frac{\operatorname{tr}(A)^{1/2}}{\sqrt{m}} \leq \left\|\frac{1}{\sqrt{m}}\right\|A^{1/2}Z_1^Tu\right\|_2 \leq (1+\varepsilon)\frac{\operatorname{tr}(A)^{1/2}}{\sqrt{m}}.
$$

Hence on event A_1 , we have

$$
\frac{1}{m} \left| \left\| A^{1/2} Z_1^T u \right\|_2^2 - \text{tr}(A) \right| \le \max((1+\varepsilon)^2 - 1, 1 - (1-\varepsilon)^2) \frac{\text{tr}(A)}{m},
$$

$$
\left| \frac{1}{m} \left\| Z_2^T B^{1/2} u \right\|_2^2 - u^T B u \right| \le \max((1+\varepsilon)^2 - 1, 1 - (1-\varepsilon)^2) \left\| B^{1/2} u \right\|_2^2.
$$

Thus we have for all $u \in S^{f-1}$, on event $\mathcal{A}_1 \cap \mathcal{A}_2$, for $C_2 := 4C + 3$

$$
\left| \frac{1}{m} u^{T} (XX^{T}) u - u^{T} \frac{\text{tr}(A) I_{f}}{m} u - u^{T} B u \right| \le
$$
\n
$$
\leq \left| \left\| Z_{2}^{T} B^{1/2} u \right\|_{2}^{2} / m - u^{T} B u \right| + \frac{1}{m} \left| \left\| A^{1/2} Z_{1}^{T} u \right\|_{2}^{2} - \text{tr}(A) \right| + 8C \varepsilon \tau_{A} \left\| B^{1/2} u \right\|_{2} / \|A\|_{2}^{1/2}
$$
\n
$$
\leq 3\varepsilon \left\| B^{1/2} u \right\|_{2}^{2} + 3\varepsilon \tau_{A} + 8C \varepsilon \tau_{A} \left\| B^{1/2} u \right\|_{2} / \|A\|_{2}^{1/2} \leq C_{2}\varepsilon \left\| B^{1/2} u \right\|_{2}^{2} + C_{2}\varepsilon \tau_{A}
$$

where $2\tau_A^{1/2}$ $\frac{1}{2}$ $\left\| B^{1/2} u \right\|_2 \leq \tau_A + \left\| B^{1/2} u \right\|_2^2$ $\frac{2}{2}$. The theorem thus holds. \square

J.1 Proof of Corollary [22](#page-19-2)

Lower bound: For all $u \in S^{f-1}$ and

$$
\frac{1}{m}u^{T}(XX^{T})u - u^{T}\frac{\text{tr}(A)I_{f}}{m}u
$$
\n
$$
\geq u^{T}Bu(1 - 3\varepsilon) - 3\varepsilon\tau_{A} - 8C \left\| B^{1/2}u \right\|_{2} \varepsilon\tau_{A}/\left\| A \right\|_{2}^{1/2}
$$
\n
$$
\geq u^{T}Bu(1 - 3\varepsilon - 4C\varepsilon) - 3\varepsilon\tau_{A} - 4C\varepsilon\tau_{A}
$$
\n
$$
\geq u^{T}Bu(1 - C_{2}\varepsilon) - C_{2}\varepsilon\tau_{A} \geq u^{T}Bu(1 - 2\delta)
$$

where we bound the term using the fact that $1 \leq \tau_A \leq \lambda_{\max}(B)$ and

$$
C_2 \tau_A \varepsilon \leq \delta \lambda_{\min}(B) \text{ and } \varepsilon \leq \delta \lambda_{\min}(B) / (C_2 \tau_A)
$$

$$
C_2 \varepsilon \leq \delta \text{ and } C_3 \varepsilon \leq \delta \min\left(\frac{\lambda_{\min}(B)}{\tau_A}, 1\right).
$$

By a similar argument, we can prove the upper bound on the isometry property as stated in the corollary. \Box

J.2 Proof of Corollary [23](#page-20-1)

Recall the following

$$
\begin{aligned}\n\widetilde{A} &:= X^T X - \text{tr}(B)I_m = \left(Z_1 A^{1/2} + B^{1/2} Z_2\right)^T \left(Z_1 A^{1/2} + B^{1/2} Z_2\right) - \text{tr}(B)I_m \\
&= \left(Z_2^T B^{1/2} + A^{1/2} Z_1^T\right) \left(Z_1 A^{1/2} + B^{1/2} Z_2\right) - \text{tr}(B)I_m \\
&= \left(Z_2^T B^{1/2} Z_1 A^{1/2} + A^{1/2} Z_1^T B^{1/2} Z_2\right) + A^{1/2} Z_1^T Z_1 A^{1/2} + \left(Z_2^T B Z_2 - \text{tr}(B)I_m\right)\n\end{aligned}
$$

.

Hence, for all vectors $u \in \mathbb{S}^{m-1} \cap E$

$$
\frac{u^T (X^T X)u}{f} - \frac{u^T \text{tr}(B)Iu}{f} - u^T A u \le \frac{1}{f} \left| u^T Z_2 B Z_2^T u - \text{tr}(B) u^T u \right|
$$

$$
+ \left| \frac{1}{f} u^T A^{1/2} Z_1^T Z_1 A^{1/2} u - u^T A u \right| + \frac{2}{f} \left| u^T A^{1/2} Z_1^T B^{1/2} Z_2 u \right|.
$$

By Lemma [35,](#page-49-3) we have on event B_1 ,

$$
\forall u \in E \cap S^{m-1} \quad \left| u^T Z^T B Z u - \text{tr}(B) \right| \leq 4C \varepsilon \text{tr}(B);
$$

By Lemma [36,](#page-49-4) we have on event B_2 ,

$$
\forall u \in E \cap S^{m-1} \left| u^T A^{1/2} Z_1^T B^{1/2} Z_2 u \right| \leq 4C \varepsilon \text{tr}(B) \left\| A^{1/2} u \right\|_2 / \|B\|_2^{1/2}.
$$

For all $u \in S^{m-1} \cap E$,

$$
8C\varepsilon\tau_B \left\| A^{1/2}u \right\|_2 / \|B\|_2^{1/2} \le 2(2C\varepsilon^{1/2} \frac{\tau_B}{\|B\|_2^{1/2}})(2\varepsilon^{1/2} \left\| A^{1/2}u \right\|_2)
$$

$$
\le 4C^2\varepsilon \frac{\tau_B^2}{\|B\|_2} + 4\varepsilon \left\| A^{1/2}u \right\|_2^2 \le 4C^2\varepsilon\tau_B + 4\varepsilon \left\| A^{1/2}u \right\|_2^2.
$$

And finally, we have also shown that for all $u \in E$ on event \mathcal{B}_9 ,

$$
(1 - \varepsilon) \|A^{1/2}u\|_2 \le \frac{1}{\sqrt{f}} \|Z_1 A^{1/2}u\|_2 \le (1 + \varepsilon) \|A^{1/2}u\|_2.
$$

Thus we have for all $u \in S^{m-1} \cap E$, on event $\mathcal{B}_1 \cap \mathcal{B}_2 \cap \mathcal{B}_9$,

$$
\left| \frac{u^T (X^T X) u}{f} - \frac{u^T \text{tr}(B) I u}{f} - u^T A u \right| \leq \frac{1}{f} \left| u^T Z_2^T B Z_2 u - \text{tr}(B) u^T u \right| \n+ \left| \frac{1}{f} \left\| Z_1 A^{1/2} u \right\|_2^2 - \left\| A^{1/2} u \right\|_2^2 \right| + \frac{2}{f} \left| u^T A^{1/2} Z_1^T B^{1/2} Z_2 u \right| \n\leq 4C \varepsilon \tau_B + 6\varepsilon \left\| A^{1/2} u \right\|_2^2 + 8C \varepsilon \tau_B \left\| A^{1/2} u \right\|_2 / \|B\|_2^{1/2} \n\leq 4C \varepsilon \tau_B + 6\varepsilon \left\| A^{1/2} u \right\|_2^2 + 4C^2 \varepsilon \tau_B + 4\varepsilon \left\| A^{1/2} u \right\|_2^2 \n\leq 10\varepsilon \left\| A^{1/2} u \right\|_2^2 + 4(C^2 + C) \varepsilon \tau_B.
$$
\n(109)

Upper bound: Thus we have by [\(109\)](#page-54-1) for the maximum sparse eigenvalue of \widetilde{A} at order k:

$$
\rho_{\max}(k,\widetilde{A}) := \max_{u \in E \cap S^{m-1}} \left| u^T \widetilde{A} u \right| \le \max_{u \in E \cap S^{m-1}} \left| u^T \widetilde{A} u - u^T A u \right| + \rho_{\max}(k,A)
$$

$$
\le \rho_{\max}(k,A)(1+10\varepsilon) + C_4 \varepsilon \tau_B
$$

where $C_4 = 4(C + C^2)$. The upper bound on $\rho_{\text{max}}(k, \tilde{A} - A)$ in the theorem statement thus holds.

Lower bound: Suppose $C_4 = 4(C + C^2) \vee 10$

$$
\varepsilon \le \frac{\delta}{C_4} \min\left(\frac{\rho_{\min}(k, A)}{\tau_B}, 1\right) = \frac{\delta}{C_5} \text{ and } C_4 \varepsilon \le \delta \min\left(\frac{\rho_{\min}(k, A)}{\tau_B}, 1\right).
$$

We have by [\(109\)](#page-54-1) for all $u \in S^{m-1} \cap E$, on event $\mathcal{B}_1 \cap \mathcal{B}_2 \cap \mathcal{B}_9$,

$$
\frac{1}{f}u^{T}(X^{T}X)u - u^{T}\frac{\operatorname{tr}(B)I_{m}}{f}u
$$
\n
$$
\geq u^{T}Au - \left(6\varepsilon u^{T}Au + 4C\varepsilon\tau_{B} + 8C\varepsilon\tau_{B} \left\| A^{1/2}u \right\|_{2} / \|B\|_{2}^{1/2} \right)
$$
\n
$$
\geq u^{T}Au - 6\varepsilon u^{T}Au - 4C\varepsilon\tau_{B} - 8C\varepsilon\tau_{B}^{1/2} \left\| A^{1/2}u \right\|_{2}
$$
\n
$$
\geq u^{T}Au - 10\varepsilon u^{T}Au - 4(C + C^{2})\varepsilon\tau_{B} \geq u^{T}Au(1 - 10\varepsilon - \delta)
$$
\n
$$
\geq u^{T}Au(1 - 2\delta)
$$

where $4(C + C^2)\varepsilon \tau_B \leq \delta \rho_{\min}(k, A)$ and $10\varepsilon \leq \delta$. \Box

K Proofs of Lemmas [35](#page-49-3) and [36](#page-49-4) and Corollary [37](#page-50-0)

Throughout the following proofs, we denote by $r(B) = \frac{\text{tr}(B)}{\|B\|_2}$. Let $\varepsilon \leq \frac{1}{C}$ where C is large enough so that $cc^{\prime}C^2 \geq 4$, and hence the choice of $C = C_0/\sqrt{c'}$ satisfies our need.

Proof of Lemma [35.](#page-49-3) First we prove concentration bounds for all pairs of $u, v \in \Pi'$, where $\Pi' \subset \mathbb{S}^{m-1}$ is an ε -net of E. Let $t = CK^2 \varepsilon \text{tr}(B)$. We have by Lemma [27,](#page-27-1) and the union bound,

$$
\mathbb{P}(\exists u, v \in \Pi', \, \left| u^T Z^T B Z v - \mathbb{E} u^T Z^T B Z v \right| > t) \n\leq 2 \left| \Pi' \right|^2 \exp \left[-c \min \left(\frac{t^2}{K^4 \left\| B \right\|_F^2}, \frac{t}{K^2 \left\| B \right\|_2} \right) \right] \n\leq 2 \left| \Pi' \right|^2 \exp \left[-c \min \left(C^2, \frac{C K^2}{\varepsilon} \right) \frac{\varepsilon^2 r(B)}{K^4} \right] \leq 2 \exp \left(-c_2 \varepsilon^2 r(B) / K^4 \right)
$$

where we use the fact that $||B||_F^2 \le ||B||_2 \text{ tr}(B)$, and

$$
\left|\Pi'\right| \le \binom{m}{k} (3/\varepsilon)^k \le \exp\left(k \log(3em/k\varepsilon)\right)
$$

while

$$
c \min\left(C^2, \frac{CK^2}{\varepsilon}\right) \varepsilon^2 \frac{r(B)}{K^4} = cC^2 \varepsilon^2 \frac{\text{tr}(B)}{\|B\|_2 K^4} \ge cC_0^2 k \log\left(\frac{3em}{k\varepsilon}\right) \ge 4k \log\left(\frac{3em}{k\varepsilon}\right)
$$

Denote by \mathcal{B}_2 the event such that for $\Lambda := \frac{1}{tr(B)} (Z^T B Z - I),$

$$
\sup_{u,v \in \Pi'} |v^T \Lambda u| \leq C\varepsilon =: r'_{f,k}
$$

holds. A standard approximation argument shows that under B_2 and for $\varepsilon \leq 1/2$,

$$
\sup_{x,y \in \mathbb{S}^{m-1} \cap E} |y^T \Lambda x| \le \frac{r'_{k,f}}{(1-\varepsilon)^2} \le 4C\varepsilon. \tag{110}
$$

The lemma is thus proved. \square

Proof of Lemma [36.](#page-49-4) By Lemma [27,](#page-27-1) we have for $t = C \epsilon \text{tr}(B) / ||B||_2^{1/2}$ $^{1/2}_{2}$ for $C = C_0/\sqrt{c'}$

$$
\mathbb{P}\left(\left|w^T Z_1^T B^{1/2} Z_2 u\right| > t\right) \leq \exp\left(-c \min\left(\frac{C^2 \frac{\text{tr}(B)^2}{\|B\|_2} \varepsilon^2}{K^4 \text{tr}(B)}, \frac{C \varepsilon \text{tr}(B)}{K^2 \|B\|_2}\right)\right)
$$

$$
\leq 2 \exp\left(-c \min\left(\frac{C^2 \varepsilon^2 r_B}{K^4}, \frac{C \varepsilon r_B}{K^2}\right)\right)
$$

$$
\leq 2 \exp\left(-c \min\left(C^2, \frac{C K^2}{\varepsilon}\right) \varepsilon^2 r_B / K^4\right)
$$

Choose an ε -net $\Pi' \subset S^{m-1}$ such that

$$
\Pi' = \bigcup_{|J|=k} \Pi'_J \quad \text{where} \quad \Pi'_J \subset E_J \cap S^{m-1} \tag{111}
$$

is an ε -net for $E_J \cap S^{m-1}$ and

$$
\left|\Pi'\right| \le \binom{m}{k} (3/\varepsilon)^k \le \exp\left(k \log(3em/k\varepsilon)\right).
$$

Similarly, choose ε -net Π of $F \cap S^{m-1}$ of size at most $\exp(k \log(3em/k\varepsilon))$. By the union bound and Lemma [27,](#page-27-1) and for $K^2 \geq 1$,

$$
\mathbb{P}\left(\exists w \in \Pi, u \in \Pi' \ s.t. \ \left|w^T Z_1^T B^{1/2} Z_2 u\right| \ge C \varepsilon \text{tr}(B) / \|B\|_2^{1/2}\right)
$$
\n
$$
\le \|\Pi'\| \|\Pi\| 2 \exp\left(-c \min\left(C K^2/\varepsilon, C^2\right) \varepsilon^2 r_B / K^4\right)
$$
\n
$$
\le \exp\left(2k \log(3em/k\varepsilon)\right) 2 \exp\left(-cC^2 \varepsilon^2 r_B / K^4\right)
$$
\n
$$
\le 2 \exp\left(-c_2 \varepsilon^2 r_B / K^4\right)
$$

where *C* is large enough such that $cc'C^2 := C' > 4$ and for $\varepsilon \leq \frac{1}{C}$ $\frac{1}{C}$

$$
c \min\left(CK^2/\varepsilon, C^2\right) \varepsilon^2 \frac{\operatorname{tr}(B)}{\|B\|_2 K^4} \ge C' k \log(3em/k\varepsilon) \ge 4k \log(3em/k\varepsilon).
$$

Denote by $\Upsilon := Z_1^T B^{1/2} Z_2$. A standard approximation argument shows that if

$$
\sup_{w \in \Pi, u \in \Pi'} |w^T \Upsilon u| \leq C \varepsilon \frac{\text{tr}(B)}{\|B\|_2^{1/2}} =: r_{k,f}
$$

an event which we denote by \mathcal{B}_2 , then for all $u \in E$ and $w \in F$,

$$
\left| w^T Z_1^T B^{1/2} Z_2 u \right| \le \frac{r_{k,f}}{(1-\varepsilon)^2}.
$$
\n(112)

The lemma thus holds for $c_2 \ge C'/2 \ge 2$. \Box

Proof of Corollary [37.](#page-50-0) Clearly [\(107\)](#page-50-2) implies that [\(104\)](#page-49-5) holds for $B = I$. Clearly [\(106\)](#page-50-1) holds following the analysis of Lemma [35](#page-49-3) by setting $B = I$, while replacing event B_1 with B_3 , which denotes an event such that

$$
\sup_{u,v \in \Pi} \frac{1}{f} \left| v^T (Z^T Z - I) u \right| \leq C \varepsilon
$$

The rest of the proof follows by replacing E with F everywhere. The corollary thus holds. \Box

References

- ALLEN, G. and TIBSHIRANI, R. (2010). Transposable regularized covariance models with an application to missing data imputation. *Annals of Applied Statistics* **4** 764–790.
- BELLONI, A., ROSENBAUM, M. and TSYBAKOV, A. (2014). Linear and conic programming estimators in high-dimensional errors-in-variables models. ArXiv:1408.0241.
- BICKEL, P., RITOV, Y. and TSYBAKOV, A. (2009). Simultaneous analysis of Lasso and Dantzig selector. *The Annals of Statistics* **37** 1705–1732.
- BONILLA, E., CHAI, K. and WILLIAMS, C. (2008). Multi-task gaussian process prediction. In *In Advances in Neural Information Processing Systems 20 (NIPS 2010)*.
- CANDES, E. and TAO, T. (2007). The Dantzig selector: statistical estimation when p is much larger than n. *Annals of Statistics* **35** 2313–2351.
- CARROLL, R., RUPPERT, D., STEFANSKI, L. and CRAINICEANU, C. M. (2006). *Measurement Error in Nonlinear Models (Second Edition)*. Chapman & Hall.
- CARROLL, R. and WAND, M. (1991). Semiparametric estimation in logistic measurement error models. *J. R. Statist. Soc. B* **53** 573–585.
- CARROLL, R. J., GAIL, M. H. and LUBIN, J. H. (1993). Case-control studies with errors in predictors. *Journal of American Statistical Association* **88** 177 – 191.
- CARROLL, R. J., GALLO, P. P. and GLESER, L. J. (1985). Comparison of least squares and errors-invariables regression with special reference to randomized analysis of covariance. *Journal of American Statistical Association* **80** 929 – 932.
- CHEN, S., DONOHO, D. and SAUNDERS, M. (1998). Atomic decomposition by basis pursuit. *SIAM Journal on Scientific and Statistical Computing* **20** 33–61.
- CHEN, Y. and CARAMANIS, C. (2013). Noisy and missing data regression: Distribution-oblivious support recovery. In *Proceedings of The 30th International Conference on Machine Learning ICML-13*.
- COHEN, M. and KOHN, A. (2011). Measuring and interpreting neuronal correlations. *Nature Neuroscience* **14** 809–811.
- COOK, J. R. and STEFANSKI, L. A. (1994). Simulation-extrapolation estimation in parametric measurement error models. *Journal of the American Statistical Association* **89** 1314–1328.
- DAWID, A. P. (1981). Some matrix-variate distribution theory: Notational considerations and a bayesian application. *Biometrika* **68** 265–274.
- DEMPSTER, A., LAIRD, N. and RUBIN, D. (1977). Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society, Series B* **39** 1–38.
- DUTILLEUL, P. (1999). The mle algorithm for the matrix normal distribution. *Journal of Statistical Computation and Simulation* **64** 105–123.
- EFRON, B. (2009). Are a set of microarrays independ of each other? *Ann. App. Statist.* **3** 922–942.
- FULLER, W. A. (1987). *Measurement error models*. John Wiley and Sons.
- GAUTIER, E. and TSYBAKOV, A. (2011). High-dimensional instrumental variables regression and confidence sets. ArXiv:1105.2454.
- GUPTA, A. and VARGA, T. (1992). Characterization of matrix variate normal distributions. *Journal of Multivariate Analysis* **41** 80–88.
- HALL, P. and MA, Y. (2007). Semiparametric estimators of functional measurement error models with unknown error. *Journal of the Royal Statistical Society B* **69** 429–446.
- HWANG, J. T. (1986). Multiplicative errors-in-variables models with applications to recent data released by the u.s. department of energy. *Journal of American Statistical Association* **81** 680–688.
- ITURRIA, S. J., CARROLL, R. J. and FIRTH, D. (1999). Polynomial regression and estimating functions in the presence of multiplicative measurement error. *Journal of the Royal Statistical Society, Series B, Methodological* **61** 547–561.
- KALAITZIS, A., LAFFERTY, J., LAWRENCE, N. and ZHOU, S. (2013). The bigraphical lasso. In *Proceedings of The 30th International Conference on Machine Learning ICML-13*.
- KASS, R., VENTURA, V. and BROWN, E. (2005). Statistical issues in the analysis of neuronal data. *J Neurophysiol* **94** 8–25.
- LIANG, H., HÄRDLE, W. and CARROLL, R. J. (1999). Estimation in a semiparametric partially linear errors-in-variables model. *Ann. Statist.* **27** 1519–1535.
- LIANG, H. and LI, R. (2009). Variable selection for partially linear models with measurement errors. *Journal of the American Statistical Association* **104** 234–248.
- LOH, P. and WAINWRIGHT, M. (2012). High-dimensional regression with noisy and missing data: Provable guarantees with nonconvexity. *The Annals of Statistics* **40** 1637–1664.
- MA, Y. and LI, R. (2010). Variable selection in measurement error models. *Bernoulli* **16** 274–300.
- MENDELSON, S., PAJOR, A. and TOMCZAK-JAEGERMANN, N. (2008). Uniform uncertainty principle for bernoulli and subgaussian ensembles. *Constructive Approximation* **28** 277–289.
- MILMAN, V. D. and SCHECHTMAN, G. (1986). *Asymptotic Theory of Finite Dimensional Normed Spaces. Lecture Notes in Mathematics 1200*. Springer.
- ROSENBAUM, M. and TSYBAKOV, A. (2010). Sparse recovery under matrix uncertainty. *The Annals of Statistics* **38** 2620–2651.
- ROSENBAUM, M. and TSYBAKOV, A. (2013). Improved matrix uncertainty selector. *IMS Collections* **9** 276–290.
- RUDELSON, M. and VERSHYNIN, R. (2013). Hanson-Wright inequality and sub-gaussian concentration. *Electronic Communications in Probability* **18** 1–9.
- RUDELSON, M. and ZHOU, S. (2013). Reconstruction from anisotropic random measurements. *IEEE Transactions on Information Theory* **59** 3434–3447.
- RUFF, D. and COHEN, M. (2014). Attention can either increase or decrease spike count correlations in visual cortex. *Nature Neuroscience* **17** 1591–7.
- SØRESEN, Ø., FRIGENSSI, A. and THORESEN, M. (2014a). Covariate selection in high-dimensional generalized linear models with measurement error. ArXiv:1407.1070.
- SØRESEN, Ø., FRIGENSSI, A. and THORESEN, M. (2014b). Measurement error in Lasso: Impact and likelihood bias correction. *Statistical Sinica Preprint* .
- STÄDLER, N., STEKHOVEN, D. J. and BÜHLMANN, P. (2014). Pattern alternating maximization algorithm for missing data in high-dimensional problems. *Journal of Machine Learning Research* **15** 1903–1928.
- STEFANSKI, L. A. (1985). The effects of measurement error on parameter estimation. *Biometrika* **72** 583–592.
- STEFANSKI, L. A. (1990). Rates of convergence of some estimators in a class of deconvolution problems. *Statistics and Probability Letters* **9** 229–235.
- STEFANSKI, L. A. and COOK, J. R. (1995). Simulation-extrapolation: The measurement error jackknife. *Journal of the American Statistical Association* **90** 1247–1256.
- STRIMMER, K. (2003). Modeling gene expression measurement error: a quasi-likelihood approach. *BMC Bioinformatics* **4**.
- TIBSHIRANI, R. (1996). Regression shrinkage and selection via the Lasso. *J. Roy. Statist. Soc. Ser. B* **58** 267–288.
- TROPP, J. and GILBERT, A. (2007). Signal recovery from random measurements via orthogonal matching pur- suit. *IEEE Trans. Inform. Theory* **53** 4655–4666.
- TROPP, J. A. (2004). Greed is good: Algorithmic results for sparse approximation. *IEEE Trans. Inform. Theory* **50** 2231–2241.
- WERNER, K., JANSSON, M. and STOICA, P. (2008). On estimation of covariance matrices with kronecker product structure. *IEEE Transactions on Signal Processing* **56** 478 – 491.
- XU, Q. and YOU, J. (2007). Covariate selection for linear errors-in-variables regression models. *Communications in Statistics – Theory and Methods* **36**.
- YU, K., LAFFERTY, J., ZHU, S. and GONG, Y. (2009). Large-scale collaborative prediction using a nonparametric random effects model. *Proceedings of the 26th International Conference on Machine Learning* .
- ZHOU, S. (2014). Gemini: Graph estimation with matrix variate normal instances. *Annals of Statistics* **42** 532–562.
- ZHOU, S., LAFFERTY, J. and WASSERMAN, L. (2010). Time varying undirected graphs. *Machine Learning* **80** 295–319.