# Approximation error estimates and inverse inequalities for B-splines of maximum smoothness

Stefan Takacs, Thomas Takacs

November 16, 2015

Abstract In this paper, we develop approximation error estimates as well as corresponding inverse inequalities for B-splines of maximum smoothness, where both the function to be approximated and the approximation error are measured in standard Sobolev norms and semi-norms. The presented approximation error estimates do not depend on the polynomial degree of the splines but only on the grid size.

We will see that the approximation lives in a subspace of the classical Bspline space. We show that for this subspace, there is an inverse inequality which is also independent of the polynomial degree. As the approximation error estimate and the inverse inequality show complementary behavior, the results shown in this paper can be used to construct fast iterative methods for solving problems arising from isogeometric discretizations of partial differential equations.

### 1 Introduction

The objective of this paper is to prove approximation error estimates as well as corresponding inverse estimates for B-splines of maximum smoothness. The presented approximation error estimates do not depend on the degree of the splines but only on the grid size. All bounds are given in terms of classical Sobolev norms and semi-norms.

E-mail: thomas.takacs@unipv.it

Stefan Takacs

Department of Computational Mathematics, Johannes Kepler University Linz, Austria E-mail: stefan.takacs@numa.uni-linz.ac.at

The work of this author was funded by the Austrian Science Fund (FWF): J3362-N25.

Thomas Takacs

Department of Mathematics, University of Pavia, Italy

The work of this author was partially funded by the European Research Council through the FP7 Ideas Consolidator Grant HIgeoM.

In approximation theory, B-splines have been studied for a long time and many properties are already well known. We do not go into the details of the existing results but present the results of importance for our study throughout this paper.

The emergence of Isogeometric Analysis, introduced in [\[11\]](#page-33-0), sparked new interest in the theoretical properties of B-splines. Since isogeometric Galerkin methods are aimed at solving variational formulations of differential equations, approximation properties measured in Sobolev norms need to be studied.

The results presented in this paper improve the results given in  $[13, 7, 1]$  $[13, 7, 1]$  $[13, 7, 1]$ by explicitly studying the dependence on the polynomial degree p. Such an analysis was done in [\[2\]](#page-33-4). However, the results there do not cover (for  $p \geq 2$ ) the most important case of B-splines of maximum smoothness  $k = p - 1$ . It turns out that the methods established in [\[2\]](#page-33-4) for proving those bounds are not suitable in that case. Therefore, we develop a framework based on Fourier analysis to prove rigorous bounds for  $k = p - 1$ , which has the limitation that it is only applicable for uniform grids.

Unlike the aforementioned papers we only consider approximation with Bsplines in the parameter domain within the framework of Isogeometric Analysis. A generalization of the results to NURBS as well as the introduction of a geometry mapping, as presented in [\[1\]](#page-33-3), is straightforward and does not lead to any additional insight.

Note that a detailed study of direct and inverse estimates may lead to a deeper understanding of isogeometric multigrid methods and give insight to suitable preconditioning methods. We refer to [\[10,](#page-33-5) [8\]](#page-33-6), where similar techniques were used.

#### 1.1 The main results

We now go through the main results of this paper. For simplicity, we consider the case of one dimension first, where  $\Omega = (a, b)$  with  $a < b$  is the open parameter domain. For this domain we can introduce a uniform grid by subdividing  $\Omega$  into elements (subintervals) of length h. The setup of a uniform grid is only possible if

$$
n_h := h^{-1}(b - a) \in \mathbb{N},
$$

where  $\mathbb{N} := \{1, 2, 3, \ldots\}$ . In other words, the grid size h has to be chosen such that  $n_h$ , the number of subintervals, is an integer. We will assume this implicitly throughout the paper. On these grids we can introduce spaces of spline functions.

**Definition 1** The space of spline functions on the domain  $\Omega$  of degree  $p \in \mathcal{D}$  $\mathbb{N}_0 := \{0, 1, 2, \ldots\}$  and continuity  $k \in \{-1, 0, 1, 2, \ldots\}$  over the uniform grid of size  $h$  is given by

$$
S_{p,k,h}(\Omega) := \left\{ u \in H^k(\Omega) : \ u|_{(a+hj,a+h(j+1)]} \in \mathbb{P}^p \text{ for all } j=0,\ldots,n_h-1 \right\},\
$$

where  $\mathbb{P}^p$  is the space of polynomials of degree p.

Here and in what follows,  $L^2(\Omega)$  and  $H^r(\Omega)$  denote the standard Lebesque and Sobolev spaces with norms  $\|\cdot\|_{L^2(\Omega)}, \|\cdot\|_{H^r(\Omega)}$  and semi-norms  $|\cdot|_{H^r(\Omega)}$ . Moreover, let  $(\cdot, \cdot)_{L^2(\Omega)}$  be the standard scalar product for  $L^2(\Omega)$  and

$$
(u,v)_{H^r(\Omega)} := \left(\frac{\partial^r}{\partial x^r}u, \frac{\partial^r}{\partial x^r}v\right)_{L^2(\Omega)}
$$

be the scalar product for  $H^r(\Omega)$ , where  $\frac{\partial^r}{\partial x^r}$  denotes the r-th derivative. We then have  $|u|_{H^r(\Omega)}^2 := (u, u)_{H^r(\Omega)}$  as well as

$$
||u||_{H^r(\Omega)}^2 := ||u||_{L^2(\Omega)}^2 + \sum_{s=1}^r |u|_{H^s(\Omega)}^2
$$

for all  $r \in \mathbb{N}_0 := \{0, 1, 2, \ldots\}.$ 

Using standard trace theorems, we obtain that for  $k > 0$  the space  $S_{p,k,h}(\Omega)$ is the space of all  $k-1$  times continuously differentiable functions  $(C^{k-1}(\Omega))$ functions), which are polynomials of degree  $p$  on each element of the uniform grid on  $\Omega$ . For  $k = 0$ , there is no continuity condition, i.e., the space  $S_{p,0,h}(\Omega)$ is the space of piecewise polynomials of degree  $p$ .

For  $k > p$ , the spline spaces reduce to spaces of global polynomials. So, the largest possible choice for k without having this effect is  $k = p$ . Therefore we call B-splines with  $k = p$  B-splines of maximum smoothness. As we are mostly interested in this case, here and in what follows, we will use  $S_{p,h}(\Omega)$ :=  $S_{p,p,h}(\Omega)$ .

The main result of this paper is the following.

<span id="page-2-0"></span>**Theorem 1** For all  $u \in H^1(\Omega)$ , all grid sizes h and each degree  $p \in \mathbb{N}$  with  $h p < |\Omega| = b - a$ , there is a spline approximation  $u_{p,h} \in S_{p,h}(\Omega)$  such that

<span id="page-2-1"></span>
$$
||u - u_{p,h}||_{L^{2}(\Omega)} \le \sqrt{2} \; h|u|_{H^{1}(\Omega)} \tag{1}
$$

is satisfied.

Note that, in contrast to the existing results presented in the next subsection, this theorem achieves two goals, it covers the case of maximum smoothness and gives a uniform estimate for all polynomial degrees p.

*Remark [1](#page-2-0)* Obviously  $S_{p,k,h}(\Omega) \supseteq S_{p,h}(\Omega)$  for all  $0 \leq k < p$ . So, Theorem 1 is also valid in that case. However, for this case there might be better estimates for these larger B-spline spaces. Moreover, Theorem [1](#page-2-0) is also satisfied in the case of having repeated knots, as this is just a local reduction of the continuity (which enlarges the corresponding space of spline functions).

In Section [5,](#page-21-0) we will introduce a subspace  $\widetilde{S}_{p,h}(\Omega) \subseteq S_{p,h}(\Omega)$  (cf. Definition [8\)](#page-22-0) and show that the spline approximation is even in that subspace (cf. Corollary [1\)](#page-22-1). Moreover, we show also a corresponding inverse inequality for  $S_{p,h}(\Omega)$  (cf. Theorem [4](#page-22-2) in Section [6\)](#page-22-3), i.e., we will show that

$$
|u_{p,h}|_{H^1(\Omega)} \le 2\sqrt{3}h^{-1} \|u_{p,h}\|_{L^2(\Omega)}
$$

is satisfied for all grid sizes h, each  $p \in \mathbb{N}$  and all  $u_{p,h} \in S_{p,h}(\Omega)$ .

We will moreover show that both the approximation error estimate and the inverse inequality are sharp up to constants (Corollaries [2](#page-23-0) and [3\)](#page-24-0).

<span id="page-3-1"></span>Remark 2 This inverse inequality does not extend to the whole space  $S_{p,h}(\Omega)$ . Here it is easy to find a counterexample: Let  $\Omega = (0, 1)$ . The function  $u_{p,h}$ , given by

$$
u_{p,h}(x) = \begin{cases} (1 - x/h)^p & \text{for } x \in [0, h) \\ 0 & \text{for } x \in [h, 1], \end{cases}
$$

is a member of the space  $S_{p,h}(0,1)$ . Straight-forward computations yield

$$
\frac{|u_{p,h}|_{H^1(0,1)}}{\|u_{p,h}\|_{L^2(0,1)}} = \sqrt{\frac{2p+1}{2p-1}} p h^{-1},
$$

which cannot be bounded from above by a constant times  $h^{-1}$  uniformly in p. Using a standard scaling argument, this counterexample can be extended to any  $\Omega = (a, b)$ .

The approximation error estimate and the inverse inequality are extended to higher Sobolev indices in Section [7.](#page-24-1) Corresponding results for two and more dimensions are given in Section [8.](#page-26-0) There, also the extension to Isogeometric Analysis is discussed.

#### 1.2 Known approximation error estimates

Before proving the main theorems, we start with recalling two important preexisting approximation error estimates. The first result is well-known in literature, cf. [\[13\]](#page-33-1), Theorem 6.25 or [\[7\]](#page-33-2), Theorem 7.3. In the framework of Isogeometric Analysis, such results have been used, e.g., in [\[1\]](#page-33-3), Lemma 3.3.

**Theorem 2** For each  $r \in \mathbb{N}_0$ , each  $k \in \mathbb{N}$ , each  $q \in \mathbb{N}$  and each  $p \in \mathbb{N}$ , with  $0 \leq r \leq q \leq p+1$  and  $r \leq k \leq p$ , there is a constant  $C(p, k, r, q)$  such that the following approximation error estimate holds. For all  $u \in H<sup>q</sup>(\Omega)$  and all grid sizes h, there is a spline approximation  $u_{p,k,h} \in S_{p,k,h}(\Omega)$  such that

<span id="page-3-0"></span>
$$
|u - u_{p,k,h}|_{H^r(\Omega)} \le C(p,k,r,q)h^{q-r}|u|_{H^q(\Omega)}
$$

is satisfied.

This theorem is valid for tensor-product spaces in any dimension and gives a local bound for locally quasi-uniform knot vectors. However, the dependence of the constant on the polynomial degree has not been derived.

A major step towards estimates with explicit p-dependence was presented in [\[2\]](#page-33-4), Theorem 2, where an estimate with an explicit dependence on  $p, k, r$ and q was given. However, there the continuity  $k$  is limited by the upper bound  $\frac{1}{2}(p+1)$ . In our notation, the theorem reads as follows.

**Theorem 3** There is a constant  $C > 0$  such that for each  $r \in \mathbb{N}_0$ , each  $k \in \mathbb{N}$ , each  $q \in \mathbb{N}$  and each  $p \in \mathbb{N}$  with  $0 \le r \le k \le q \le p+1$  and  $k \le \frac{1}{2}(p+1)$ and all grid sizes h, the following approximation error estimate holds. For all  $u \in H^{q}(\Omega)$ , there is a spline approximation  $u_{p,k,h} \in S_{p,k,h}(\Omega)$  such that

$$
|u - u_{p,k,h}|_{H^r(\Omega)} \le Ch^{q-r}(p-k+1)^{-(q-r)}|u|_{H^q(\Omega)}
$$

is satisfied.

Again, the original result was stated for locally quasi-uniform knots. For any  $p \geq 2$  the relevant case  $k = p$ , which we consider, is not covered by this theorem.

Similar results to Theorem [1](#page-2-0) are known in approximation theory, cf. [\[12\]](#page-33-7). There, however, different norms have been discussed. Hence we do not go into the details. In [\[9\]](#page-33-8), it was suggested and confirmed by numerical experiments that Theorem [1](#page-2-0) is satisfied. A proof was however not given.

#### 1.3 Organization of this paper

This paper is organized as follows. In Section [2,](#page-4-0) we present the main steps of the proof of Theorem [1](#page-2-0) and give some preliminaries. In the following two sections, the details of the proof are worked out. In Section [5,](#page-21-0) we introduce the reduced spline space  $S_{p,h}(\Omega)$ , discuss its properties and extend Theorem [1](#page-2-0) to that space. In the following section, Section [6,](#page-22-3) we give an inverse inequality for  $S_{p,h}(\Omega)$ . In the remainder of the paper, we generalize those results: In Section [7](#page-24-1) we consider higher Sobolev indices and in Section [8,](#page-26-0) the results are generalized to two or more dimensions.

## <span id="page-4-0"></span>2 Concept of the proof of Theorem 1 and Preliminaries

The proof of Theorem [1](#page-2-0) is based on an estimate for periodic splines, which is formulated as Lemma [9.](#page-20-0) The proof of Lemma [9](#page-20-0) is based on a telescoping argument based on a hierarchy of grids. For the proof, we require

- an estimate for the difference of the spline approximations of a given function on two consecutive grids, cf. [\(8\)](#page-8-0), and
- an estimate for the difference between the spline approximation on some finest grid and the given function, cf. Lemma [1.](#page-6-0)

As the size of the finest grid approaches 0, the constant in Lemma [1](#page-6-0) or its dependence on the spline degree  $p$  does not matter, whereas the constant in  $(8)$ directly affects the constant in the final result.

The estimate [\(8\)](#page-8-0) is shown in Section [3,](#page-8-1) cf. Lemma [8.](#page-20-1) There, the proof is done by means of Fourier analysis, which causes the restriction of the analysis to equidistant grids. The Fourier analysis follows a classical line: first, a matrixvector formulation is introduced, cf. Lemma [3,](#page-11-0) then the symbols of the involved matrices are derived, cf. Subsections [3.3](#page-13-0) and [3.4.](#page-14-0) A closed form for the symbol of the mass matrix is not available, so some statements on that matrix are derived (Lemmas [4](#page-12-0) and [6\)](#page-16-0), which are used in the proof of Lemma [8.](#page-20-1)

Having the result for two consecutive grids in the periodic case, we use the aforementioned telescoping argument to give an approximation error estimate for apprximating a general periodic  $H^1$ -function. The extension to the nonperiodic case is done by means of a periodic extension.

## 2.1 Periodic splines

To establish the theory within this paper, we need to introduce spaces of periodic splines, which we define as follows.

**Definition 2** Given a spline space  $S_{p,h}(\Omega)$  over  $\Omega = (a, b)$ , the *periodic spline* space  $\widehat{S}_{p,h}(\Omega)$  contains all functions  $u_{p,h} \in S_{p,h}(\Omega)$  that satisfy the linear periodicity condition

<span id="page-5-1"></span>
$$
\frac{\partial^l}{\partial x^l} u_{p,h}(a) = \frac{\partial^l}{\partial x^l} u_{p,h}(b) \text{ for all } l \in \mathbb{N}_0 \text{ with } l < p. \tag{2}
$$

The next step is to introduce a B-spline-like basis for this space. First, we introduce the cardinal B-splines. On R, the cardinal B-splines are defined as follows, cf. [\[13\]](#page-33-1), (4.22).

**Definition 3** The cardinal B-splines of degree  $p = 0$ ,  $\psi_0^{(i)} : \mathbb{R} \to \mathbb{R}$  coincide with the characteristic function, i.e.,

$$
\psi_0^{(i)}(x) := \begin{cases} 1 & \text{for } x \in (i, i+1], \\ 0 & \text{else,} \end{cases}
$$

where  $i \in \mathbb{Z}$ .

The cardinal B-splines  $\psi_p^{(i)} : \mathbb{R} \to \mathbb{R}$  of degree  $p \in \mathbb{N}$  are given by the recurrence formula

<span id="page-5-0"></span>
$$
\psi_p^{(i)}(x) := \frac{x - i}{p} \psi_{p-1}^{(i)}(x) + \frac{(p + i + 1) - x}{p} \psi_{p-1}^{(i+1)}(x),\tag{3}
$$

where  $i \in \mathbb{Z}$ .

From the cardinal B-splines  $\psi_p^{(i)}$ , we derive the B-splines  $\varphi_{p,h}^{(i)}$  on  $\Omega$  over a uniform grid of size  $h$  by a suitable scaling and shifting.

**Definition 4** For  $i \in \mathbb{Z}$  the uniform B-spline  $\varphi_{p,h}^{(i)}$ :  $\Omega = (a, b) \to \mathbb{R}$  of degree  $p \in \mathbb{N}_0$  and grid size h is given by

$$
\varphi_{p,h}^{(i)}(x) := \psi_p^{(i)}\left(\frac{x-a}{h}\right). \tag{4}
$$

We obtain by construction that  $\text{supp}(\varphi_{p,h}^{(i)}) \subset [ih + a, (i + p + 1)h + a].$ Hence,  $-p$  and  $n_h - 1$  with  $n_h = h^{-1}(b-a)$  are the first and last indices of the B-splines supported in  $\Omega$ , respectively, i.e.  $\text{supp}(\varphi_{p,h}^{(i)}) \cap \Omega \neq \emptyset$  is equivalent to  $-p \leq i \leq n_h - 1$ . Moreover,  $\{\varphi_{p,h}^{(i)}\}_{i=-p}^{n_h-1}$  forms a basis for  $S_{p,h}$ , see, e.g., [\[13\]](#page-33-1). Note that both  $n_h$  and the basis functions depend implicitly on the choice of  $\Omega$ , i.e., the values a and b. Throughout the paper, it is clear from the context which  $\Omega$  is chosen.

For the construction of the basis for the periodic spline space  $\widehat{S}_{p,h}(\Omega)$ , we assume that

$$
hp < |\Omega| = b - a,\tag{5}
$$

i.e., that the grid is fine enough not to have basis functions that are non-zero at both end points of the grid, cf. [\[13\]](#page-33-1).

**Definition 5** For  $\widehat{S}_{p,h}(\Omega)$ , the *B-spline-like basis*  $\{\widehat{\varphi}_{p,h}^{(i)}\}_{i=0}^{n_h-1}$  is given by

$$
\begin{aligned}\n\widehat{\varphi}_{p,h}^{(i)} &:= \varphi_{p,h}^{(i)} & \text{if } i < n_h - p, \text{ and} \\
\widehat{\varphi}_{p,h}^{(i)} &:= \varphi_{p,h}^{(i)} + \varphi_{p,h}^{(i-n_h)} & \text{if } i \ge n_h - p.\n\end{aligned}
$$

Up to indexing, this definition coincides with (8.6) and (8.7) in [\[13\]](#page-33-1). Theorem 8.2 in [\[13\]](#page-33-1) states that [\(6\)](#page-6-1) is actually a basis.

As  $\varphi_{p,h}^{(i)}$  vanishes on  $\Omega$  for all  $i \notin \{-p,\ldots,n_h-1\}$ , we have

<span id="page-6-1"></span>
$$
\widehat{\varphi}_{p,h}^{(i)} = \sum_{j \in \mathbb{Z}} \varphi_{p,h}^{(i+jn_h)},\tag{6}
$$

where  $\mathbb Z$  is the set of integers, for all  $i = 0, \ldots, n_h - 1$ . Using this definition, we directly obtain that also  $\hat{\varphi}_{p,h}^{(i)} = \hat{\varphi}_{p,h}^{(i+jn_h)}$  for any  $j \in \mathbb{Z}$ , which we will use for association throughout this ppor for ease of notation throughout this paper.

We call this basis B-spline-like, as each function is a non-negative linear combination of B-splines and it forms a partition of unity on  $\Omega$ .

#### 2.2 A non-robust approximation error estimate in the periodic case

We can extend Theorem [2](#page-3-0) for  $k = p - 1$  $k = p - 1$  to the following Lemma 1 stating that the approximation error estimate is still satisfied if we approximate periodic functions with periodic splines. First, we introduce the spaces of periodic functions as follows.

**Definition 6** For  $\Omega = (a, b)$ , the space  $\hat{H}^q(\Omega)$  is the space of all  $u \in H^q(\Omega)$ that satisfy the periodicity condition

<span id="page-6-2"></span><span id="page-6-0"></span>
$$
\frac{\partial^l}{\partial x^l}u(a) = \frac{\partial^l}{\partial x^l}u(b) \text{ for all } l \in \mathbb{N}_0 \text{ with } l < q. \tag{7}
$$

Note that standard trace theorems guarantee that the periodicity condition [\(7\)](#page-6-2) is well-defined. For this space, the following lemma holds.

**Lemma 1** For each  $r \in \mathbb{N}_0$ , each  $q \in \mathbb{N}$  and each  $p \in \mathbb{N}$  with  $0 \leq r \leq$  $q \leq p+1$ , there is a constant  $C(p,r,q)$  such that the following approximation error estimate holds. For all  $u \in \hat{H}^q(\Omega)$  and all grid sizes h, there is a spline approximation  $u_{p,h} \in \widehat{S}_{p,h}(\Omega)$  such that

$$
|u - u_{p,h}|_{H^r(\Omega)} \le C(p,r,q)h^{q-r}|u|_{H^q(\Omega)}
$$

is satisfied.

*Proof* In the following, we assume without loss of generality that  $\Omega = (0, 1)$ . The extension to any other  $\Omega = (a, b)$ , follows using a standard scaling argument.

Let w be the periodic extension of the function u to R, i.e.,  $w(x) := u(x |x|$ ). Note that the restriction of w to any finite interval is again a function in the Sobolev space  $H<sup>q</sup>$ . The following of the proof is based on the proof in § 6.4 in [\[13\]](#page-33-1). We make use of the fact that the proof uses local projections. Let  $Q_{p,h}$ :  $\dot{H}^q(\mathbb{R}) \to S_{p,h}(\mathbb{R})$  be the projection operator, as introduced in (6.40) in [\[13\]](#page-33-1). The value of the approximation  $Q_{p,h}w$  of a function w at a certain subinterval  $I_i := (i \; h, (i+1) \; h) \subseteq \Omega$  only depends on the values of the function to be approximated in a certain neighborhood  $I_i := ((i-p) \; h, (i+p+1) \; h)$ . So, from the periodicity of w, the periodicity of  $Q_{p,h}w$  follows immediately. Hence its restriction to  $(0, 1)$  is a periodic spline, i.e.  $Q_{p,h}w|_{(0,1)} \in S_{p,h}(0, 1)$ . We define  $u_{p,h}$  to be the restriction of  $Q_{p,h}w$  to  $(0,1)$ . Due to [\[13\]](#page-33-1), Theorem 6.24, the local estimate

$$
|w - Q_{p,h}w|_{H^r(I_i)} \leq \widetilde{C}(p,r,q)h^{q-r}|w|_{H^q(\widetilde{I}_i)}.
$$

is satisfied for the projector  $Q_{p,h}$  and a constant  $\widetilde{C}(p,r,q)$ , which is independent of h. By summing over all elements, we obtain

$$
|u - u_{p,h}|_{H^r(0,1)}^2 = |w - Q_{p,h}w|_{H^r(0,1)}^2 = \sum_{i=0}^{n_h-1} |w - Q_{p,h}w|_{H^r(I_i)}^2
$$
  

$$
\leq \widetilde{C}^2(p,r,q)h^{2(q-r)} \sum_{i=0}^{n_h-1} |w|_{H^q(\widetilde{I}_i)}^2 = \widetilde{C}^2(p,r,q)h^{2(q-r)} \sum_{i=0}^{n_h-1} \sum_{j=-p}^p |w|_{H^q(I_{i+j})}^2.
$$

Using the periodicity of w, we can express the last term using  $|u|_{H^q(I_l)}$  for  $l \in$  $\{0, \ldots, n_h-1\}$  only. By counting the occurrences of the summands  $|u|_{H^q(I_l)},$ we obtain

$$
\sum_{i=0}^{n_h-1} \sum_{j=-p}^p |w|_{H^q(I_{i+j})}^2 = (2p+1) \sum_{i=0}^{n_h-1} |u|_{H^q(I_i)}^2 = (2p+1)|u|_{H^q(0,1)}^2,
$$

which finishes the proof for  $C(p, r, q) = (2p + 1)^{1/2} \widetilde{C}(p, r, q)$ .

# <span id="page-8-1"></span>3 A robust approximation error estimate for two consecutive grids in the periodic case

In this section we analyze the case of approximating a periodic spline function on a fine grid by a periodic spline function on a coarser grid. In the next section, we extend these results to the approximation of general functions and to the non-periodic case. The extension to the non-periodic case is done by extending functions in  $H^1(0,1)$  to  $(-1,1)$  by reflecting them on the yaxis. So, without loss of generality, we will restrict ourselves to  $\Omega = (-1, 1)$ throughout this section. Moreover, for the construction of [\(28\)](#page-15-0), we will need that  $hp < 1$ , which is stronger than the requirement  $hp < b-a$ , cf. Theorem [1.](#page-2-0) So, throughout this section, we will use the following assumptions.

**Assumption 1** The domain is given by  $\Omega = (-1, 1)$  and the grid size is small enough such that  $hp < 1$  holds.

In the next section, we will make use of a telescoping argument. For this purpose, we have to analyze a fixed interpolation operator. So, within this section, we will show that

<span id="page-8-0"></span>
$$
\|(I - \widehat{H}_{p,h})u_{p,h}\|_{L^2(-1,1)} \le \frac{1}{\sqrt{2}}\;h|u_{p,h}|_{H^1(-1,1)}\tag{8}
$$

holds for all  $u_{p,h} \in \widehat{S}_{p,\frac{h}{2}}(-1,1)$ , where I is the identity and  $\widehat{H}_{p,h}$  is the  $H^1$ orthogonal projection operator, given by the following definition.

**Definition 7** The projection  $\widehat{H}_{p,h} : \widehat{H}^1(-1,1) \to \widehat{S}_{p,h}(-1,1)$  maps every  $u \in \hat{H}^1(-1, 1)$  to the function  $u_{p,h} \in \hat{S}_{p,h}(-1, 1)$  satisfying

<span id="page-8-3"></span><span id="page-8-2"></span>
$$
(u_{p,h}, v_{p,h})_{H^1_o(-1,1)} = (u, v_{p,h})_{H^1_o(-1,1)}
$$
\n(9)

for all  $v_{p,h} \in \hat{S}_{p,h}(-1,1)$ , where

$$
(u,v)_{H_0^1(-1,1)} := (u,v)_{H^1(-1,1)} + \left(\int_{-1}^1 u(x) dx\right) \left(\int_{-1}^1 v(x) dx\right).
$$

Within the next subsections, we will prove [\(8\)](#page-8-0). This will be done by a rigorous version of Fourier analysis. Fourier analysis is a well-known tool for analyzing convergence properties of numerical methods, cf. the work by A. Brandt, like [\[5\]](#page-33-9), and many others. It provides a framework to determine sharp bounds for the convergence rates of multigrid methods and other iterative solvers for problems arising from partial differential equations. This is different to classical analysis, which typically yields qualitative statements only. For a detailed introduction into Fourier analysis, see, e.g., [\[15\]](#page-33-10). Recently, it has also been applied in the area of Isogeometric Analysis, cf. [\[10\]](#page-33-5).

Typically, Fourier analysis is done under simplifying assumptions, like assuming uniform grids and neglecting the boundary. In this case, one refers to local Fourier analysis (or local mode analysis). This analysis can be understood as a heuristic method to study methods of interest. In a recent work, cf. [\[10\]](#page-33-5), it was understood also as a rigorous statement for a limit case.

We, however, are interested in a completely rigorous analysis. As we restrict ourselves to periodic spline spaces, the Fourier modes are the exact eigenvectors of the matrices of interest, which will allow us to diagonalize these matrices using a similarity transformation. Based on such a diagonalization, we will be able to prove [\(8\)](#page-8-0).

As a first step, we introduce a matrix-vector formulation of [\(8\)](#page-8-0).

### 3.1 A matrix-vector formulation of the estimate

Having fixed the B-spline like basis  $\{\widehat{\varphi}_{p,h}^{(i)}\}_{i=0}^{n_h-1}$ , we can write any function  $u_{p,h} \in \widehat{S}_{p,h}(-1,1)$  as a linear combination of these basis functions:

$$
u_{p,h} = \sum_{i=0}^{n_h-1} u_{p,h}^{(i)} \hat{\varphi}_{p,h}^{(i)}.
$$

The coefficients  $u_{p,h}^{(i)}$  can be collected in a coefficient vector: We define  $\underline{u}_{p,h}$ :=  $(u_{p,h}^{(i)})_{i=0}^{n_h-1}$ . So, the vector  $\underline{u}_{p,h}$  is the representation of the function  $u_{p,h}$  with respect to the B-spline like basis. Here and in what follows, we will always assume underlined quantities to be the basis representation of the corresponding function with respect to the basis  $\{\widehat{\varphi}_{p,h}^{(i)}\}_{i=0}^{n_h}$ .

By plugging such a decomposition into the standard  $L^2$ -scalar product  $(\cdot,\cdot)_{L^2(-1,1)}$ , we obtain

$$
(u_{p,h}, v_{p,h})_{L^2(-1,1)} = \sum_{i=0}^{n_h-1} \sum_{j=0}^{n_h-1} u_{p,h}^{(i)} v_{p,h}^{(j)} (\hat{\varphi}_{p,h}^{(i)}, \hat{\varphi}_{p,h}^{(j)})_{L^2(-1,1)}.
$$

As the grid is equidistant and the splines are periodic, we obtain that for all *i* and *j* the relation  $(\hat{\varphi}_{p,h}^{(i)}, \hat{\varphi}_{p,h}^{(j)})_{L^2(-1,1)} = m_{p,h}^{(i-j)}$  holds with coefficients  $m_{p,h}^{(i)} := (\widehat{\varphi}_{p,h}^{(i)}, \widehat{\varphi}_{p,h}^{(0)})_{L^2(-1,1)}$ . Those coefficients form a circulant matrix  $M_{p,h} :=$  $(m_{p,h}^{(i-j)})_{i=0,\ldots,n_h-1}^{j=0,\ldots,n_h-1}$ , which is called the mass matrix. We immediately obtain

$$
(u_{p,h}, v_{p,h})_{L^2(-1,1)} = (\underline{u}_{p,h}, \underline{v}_{p,h})_{M_{p,h}} := \underline{v}_{p,h}^T M_{p,h} \underline{u}_{p,h}
$$

and

$$
||u_{p,h}||_{L^2(-1,1)}^2 = ||u_{p,h}||_{M_{p,h}}^2 := \underline{u}_{p,h}^T M_{p,h} \underline{u}_{p,h}.
$$

Having a look onto the support of the functions  $\hat{\varphi}_{p,h}^{(0)}$ , we obtain that the bandwidth of the mass matrix is  $2p+1$ , i.e.  $m_{p,h}^{(i-j)} = 0$  for all  $i, j$  with  $|i-j| > p$ .

Analogously to the definition of the mass matrix, we can introduce the stiffness matrix, representing the  $H_0^1$ -scalar product. The stiffness matrix is given by  $K_{p,h} := (k_{p,h}^{(i-j)})_{i=0,\ldots,n_h-1}^{j=0,\ldots,n_h-1}$ , where the coefficients are given by

$$
k_{p,h}^{(i)} := \left(\widehat{\varphi}_{p,h}^{(i)}, \widehat{\varphi}_{p,h}^{(0)}\right)_{H_0^1(-1,1)}.
$$

Since the basis functions  $\hat{\varphi}_{p,h}^{(i)}$  form a partition of unity on  $\Omega = (-1,1)$ ,  $\int_{-1}^{1} \widehat{\varphi}_{p,h}^{(i)}(x) dx = h$  and further

<span id="page-10-0"></span>
$$
k_{p,h}^{(i)} = \left(\widehat{\varphi}_{p,h}^{(i)}, \widehat{\varphi}_{p,h}^{(0)}\right)_{H^1(-1,1)} + h^2.
$$
 (10)

Note that for uniform knot vectors the identity

$$
\frac{\partial}{\partial x} \varphi_{p,h}^{(j)}(x) = \frac{1}{h} \left( \varphi_{p-1,h}^{(j-1)}(x) - \varphi_{p-1,h}^{(j)}(x) \right)
$$

holds, see e.g.  $(5.36)$  in [\[13\]](#page-33-1). This statement directly carries over to the periodic splines using relation [\(6\)](#page-6-1), i.e.,

$$
\frac{\partial}{\partial x}\widehat{\varphi}_{p,h}^{(j)}(x) = \frac{1}{h}\left(\widehat{\varphi}_{p-1,h}^{(j-1)}(x) - \widehat{\varphi}_{p-1,h}^{(j)}(x)\right)
$$

also holds. By plugging this into [\(10\)](#page-10-0), the entries of the stiffness matrix can be derived directly using the entries of the mass matrix for splines of order  $p-1$ . Straight-forward calculations show that

<span id="page-10-1"></span>
$$
K_{p,h} = D_h M_{p-1,h} D_h^T + E_h,
$$
\n(11)

where the gradient matrix  $D_h := (d_h^{(i-j)})$  $\binom{(i-j)}{h}$ <sub>i=0,...,n<sub>h</sub>-1</sub> is given by the coefficients

$$
d_h^{(i)} := \frac{1}{h} \begin{cases} 1 & \text{for } i \in n_h \mathbb{Z} \\ -1 & \text{for } i \in n_h \mathbb{Z} - 1 \\ 0 & \text{else} \end{cases}
$$

the rank-one matrix  $E_h$  is given by  $E_h := h^2 \mathbf{1}_h \mathbf{1}_h^T$ , where  $\mathbf{1}_h := (1, \ldots, 1)^T \in$  $\mathbb{R}^{n_h}$  is a vector consisting only of ones, representing the constant function. Note that  $D_h$ ,  $E_h$  and, consequently,  $K_h$  are also circulant matrices.

To derive a matrix-vector formulation of [\(8\)](#page-8-0), we have to introduce a matrix that represents the canonical embedding from  $S_{p,h}(-1,1)$  into  $S_{p,\frac{h}{2}}(-1,1)$ . The following lemma is rather well-known in literature, cf. [\[6\]](#page-33-11) equation (4.3.4), and can be easily shown by induction in p.

**Lemma 2** For all  $p \in \mathbb{N}$ , all grid sizes h and all  $x \in \mathbb{R}$ ,

<span id="page-10-2"></span>
$$
\varphi_{p,h}^{(j)}(x) = 2^{-p} \sum_{l=0}^{p+1} \binom{p+1}{l} \varphi_{p,\frac{h}{2}}^{(2j+l)}(x)
$$

is satisfied for all  $j = -p, \ldots, n_h - p - 1$ .

This directly carries over to the periodic splines, i.e., we obtain

$$
\widehat{\varphi}_{p,h}^{(j)}(x) = 2^{-p} \sum_{l=0}^{p+1} {p+1 \choose l} \widehat{\varphi}_{p,\frac{h}{2}}^{(2j+l)}(x) = \sum_{i \in \mathbb{Z}} \underbrace{2^{-p} {p+1 \choose i-2j}}_{p,\frac{h}{2}} \widehat{\varphi}_{p,\frac{h}{2}}^{(i)}(x). \tag{12}
$$

Here, we use equation [\(6\)](#page-6-1) and that the binomial coefficient  $\begin{pmatrix} a \\ b \end{pmatrix}$ b vanishes for  $b \notin \{0, \ldots, a\}$ . Again, we define the matrix  $P_{p, \frac{h}{2}}$  $:= (p^{(i,j)}_{h})$  $p, \frac{h}{2}$  $\rangle_{i=0,...,n_h-1}^{j=0,...,n_h-1}$ . Here and in what follows, we make use of  $n_{\frac{h}{2}} = 2n_h$ .

<span id="page-11-0"></span>**Lemma 3** The inequality  $(8)$  is equivalent to

<span id="page-11-2"></span><span id="page-11-1"></span>
$$
||M_{p,\frac{h}{2}}^{1/2}(I - P_{p,\frac{h}{2}} K_{p,h}^{-1} P_{p,\frac{h}{2}}^T K_{p,\frac{h}{2}}) K_{p,\frac{h}{2}}^{-1/2}|| \le \frac{1}{\sqrt{2}} h,
$$
\n(13)

which is a consequence of the combination of

$$
||M_{p, \frac{h}{2}}^{1/2} M_{p-1, \frac{h}{2}}^{-1/2}|| \le 1 \qquad and \qquad (14)
$$

$$
||M_{p-1,\frac{h}{2}}^{1/2}(I - P_{p,\frac{h}{2}} K_{p,h}^{-1} P_{p,\frac{h}{2}}^T K_{p,\frac{h}{2}}) K_{p,\frac{h}{2}}^{-1/2}|| \le \frac{1}{\sqrt{2}}h. \tag{15}
$$

Here and in what follows,  $\|\cdot\|$  is the Euclidean norm and the square root  $A^{1/2}$ of a symmetric and positive definite matrix  $A$  is that symmetric and positive definite matrix that satisfies  $A^{1/2}A^{1/2} = A$ .

*Proof of Lemma [3](#page-11-0)* Using the introduced matrices  $K_{p,h}$  and  $P_{p,\frac{h}{2}}$ , we can 2 rewrite [\(9\)](#page-8-2) for the choice  $u := u_{p, \frac{h}{2}} \in S_{p, \frac{h}{2}}$  in matrix-vector form as

<span id="page-11-4"></span>
$$
(P_{p,\frac{h}{2}}\underline{u}_{p,h},P_{p,\frac{h}{2}}\underline{v}_{p,h})_{K_{p,\frac{h}{2}}}=(\underline{u}_{p,\frac{h}{2}},P_{p,\frac{h}{2}}\underline{v}_{p,h})_{K_{p,\frac{h}{2}}},
$$

which is equivalent to

$$
P_{p,\frac{h}{2}}^T K_{p,\frac{h}{2}} P_{p,\frac{h}{2}} \underline{u}_{p,h} = P_{p,\frac{h}{2}}^T K_{p,\frac{h}{2}} \underline{u}_{p,\frac{h}{2}}.
$$

This yields, using the Galerkin principle  $(P_{p, \frac{h}{2}}^T K_{p, \frac{h}{2}} P_{p, \frac{h}{2}} = K_{p,h})$ , that the coarse-grid approximation  $\underline{u}_{p,h}$  is given by

$$
\underline{u}_{p,h} = K_{p,h}^{-1} P_{p,\frac{h}{2}}^T K_{p,\frac{h}{2}} \underline{u}_{p,\frac{h}{2}}
$$

<span id="page-11-3"></span>.

By plugging this into [\(8\)](#page-8-0), we see that we have to show

P

$$
\|(I-P_{p,\frac{h}{2}}K_{p,h}^{-1}P_{p,\frac{h}{2}}^TK_{p,\frac{h}{2}}K_{p,\frac{h}{2}})u_{p,\frac{h}{2}}\|_{M_{p,\frac{h}{2}}}\leq \frac{1}{\sqrt{2}}h\|\underline{u}_{p,\frac{h}{2}}\|_{K_{p,\frac{h}{2}}}
$$

for all  $\underline{u}_{p,\frac{h}{2}} \in \mathbb{R}^{2n_h}$ . By rewriting this using a standard matrix norm, we obtain [\(13\)](#page-11-1). Using the semi-multiplicativity of matrix norms, we obtain that [\(13\)](#page-11-1) is a consequence of  $(14)$  and  $(15)$ .

Note that the stiffness matrix for some degree  $p$  depends implicitly on the mass matrix for the degree  $p - 1$ . So, analyzing [\(15\)](#page-11-3) is more convenient than analyzing [\(13\)](#page-11-1) as the inequality [\(15\)](#page-11-3) depends just on the one mass matrix  $M_{p-1,\frac{h}{2}}$ , whereas [\(13\)](#page-11-1) depends on two mass matrices:  $M_{p-1,\frac{h}{2}}$  and  $M_{p,\frac{h}{2}}$ . We will show [\(14\)](#page-11-2) in the next subsection and [\(15\)](#page-11-3) in the remainder of this section. 3.2 A lemma relating the mass matrices for different polynomial degrees

The estimate [\(14\)](#page-11-2) is a direct consequence of the following lemma.

**Lemma 4** For all  $p \in \mathbb{N}$ , grid sizes h and vectors  $\underline{u}_h \in \mathbb{R}^{n_h}$ , the inequality

<span id="page-12-0"></span>
$$
\|\underline{u}_h\|_{M_{p,h}} \le \|\underline{u}_h\|_{M_{p-1,h}}
$$

is satisfied.

Proof First we observe that the convolution formula for cardinal B-splines, cf. equation (13) in [\[10\]](#page-33-5), can be carried over to the functions  $\hat{\varphi}_{p,h}^{(i)}$ , i.e., that

$$
\hat{\varphi}_{p,h}^{(i)}(x) = h^{-1} \int_0^h \hat{\varphi}_{p-1,h}^{(i)}(x-t) \mathrm{d}t \tag{16}
$$

holds. Let  $\underline{u}_h = (u_h^{(i)}\$  $\binom{i}{h}$ <sub> $i=0$ </sub> $\ldots$ <sup>n<sub>h</sub>-1</sub>. Then, using [\(16\)](#page-11-4), we have that</sup>

$$
\|u_h\|_{M_{p,h}}^2 = \int_{-1}^1 \left(\sum_{i=0}^{n_h-1} u_h^{(i)} \hat{\varphi}_{p,h}^{(i)}(x)\right)^2 dx
$$
  
\n
$$
= \int_{-1}^1 \left(\sum_{i=0}^{n_h-1} u_h^{(i)} h^{-1} \int_0^h \hat{\varphi}_{p-1,h}^{(i)}(x-t) dt\right)^2 dx
$$
  
\n
$$
= h^{-2} \int_{-1}^1 \left(\int_0^h \left(\sum_{i=0}^{n_h-1} u_h^{(i)} \hat{\varphi}_{p-1,h}^{(i)}(x-t)\right) dt\right)^2 dx
$$
  
\n
$$
= h^{-2} \int_{-1}^1 \left(\int_0^h 1 s(x-t) dt\right)^2 dx
$$

holds, where  $s(x) := \sum_{i=0}^{n_h-1} u_h^{(i)} \hat{\varphi}_{p-1,h}^{(i)}(x-t)$ .<br>Now we apply the Gaughy Sepwary inequ

Now, we apply the Cauchy-Schwarz inequality to the inner integral and obtain

$$
\|\underline{u}_h\|_{M_{p,h}}^2 \le h^{-2} \int_{-1}^1 \left(\int_0^h 1^2 dt\right) \left(\int_0^h s^2(x-t) dt\right) dx
$$
  
=  $h^{-1} \int_{-1}^1 \int_0^h s^2(x-t) dt dx = h^{-1} \int_0^h \int_{-1}^1 s^2(x-t) dx dt.$ 

Observe that due to periodicity,  $\int_{-1}^{1} s^2(x-t) dx = \int_{-1}^{1} s^2(\xi) d\xi$  for all  $t \in [0, h]$ , which implies

$$
\|\underline{u}_h\|_{M_{p,h}}^2 \le h^{-1} \int_0^h \int_{-1}^1 s^2(\xi) d\xi dt = h^{-1} \left( \int_0^h 1 dt \right) \left( \int_{-1}^1 s^2(\xi) d\xi \right)
$$

$$
= \int_{-1}^1 \left( \sum_{i=0}^{n_h - 1} u_h^{(i)} \widehat{\varphi}_{p-1,h}^{(i)}(\xi) \right)^2 d\xi = \|\underline{u}_h\|_{M_{p-1,h}}^2,
$$

which finishes the proof.



# <span id="page-13-0"></span>3.3 Symbols of mass matrix and stiffness matrix

As the matrices  $M_{p,h}$  and  $K_{p,h}$  are circulant matrices, we can analyze them using Fourier analysis. So, we consider the Fourier vectors

$$
\underline{f}_{h,j} := (\mathbf{e}^{2ijh\pi \mathbf{i}})_{i=0}^{n_h - 1} \quad \text{for } j = 0, \dots, n_h - 1,
$$

where **i** is the imaginary unit.

We observe (using that the bandwith of the mass matrix is  $2p + 1$ ) that

$$
(M_{p,h}\underline{f}_{h,j})_i = \sum_{l=-p}^p m_{p,h}^{(l)} e^{2(i+l)jh\pi i} = \sum_{l=-p}^p m_{p,h}^{(l)} e^{2ljh\pi i} e^{2ijh\pi i}
$$

$$
= \underbrace{\sum_{l=-p}^p m_{p,h}^{(l)} e^{2ljh\pi i} (\underline{f}_{h,j})_i}_{\widehat{m}_{p,h}^{(j)} :=}
$$

for all  $i = 0, \ldots, n_h - 1$  and  $j = 0, \ldots, n_h - 1$  and consequently

<span id="page-13-1"></span>
$$
M_{p,h} \underline{f}_{h,j} = \widehat{m}_{p,h}^{(j)} \underline{f}_{h,j}
$$

is satisfied for all  $j = 0, \ldots, n_h-1$ , i.e., that  $\underline{f}_{h,j}$  is an eigenvector of  $M_{p,h}$  with corresponding eigenvalue  $\widehat{m}_{ph}^{(j)}$ . As we have identified  $n_h$  different eigenvalues, the corresponding eigenvalues define a basis of  $\mathbb{R}^{n_h}$ . Therefore, the matrix the corresponding eigenvectors define a basis of  $\mathbb{R}^{n_h}$ . Therefore, the matrix  $\mathbb{F}_h$ , obtained by collecting the vectors  $\underline{f}_{h,j}$ , i.e.,

$$
\mathbb{F}_h := \left(\underline{f}_{h,0} \,\underline{f}_{h,1} \,\cdots \,\underline{f}_{h,n_h-1}\right) = (\mathbf{e}^{2ijh\pi\mathbf{i}})_{i=0,\ldots,n_h-1}^{j=0,\ldots,n_h-1},
$$

is a non-singular matrix. As  $\mathbb{F}_h$  is the matrix built from the eigenvectors, it diagonalizes the matrix  $M_{p,h}$ , i.e.,

$$
\mathbb{F}_h^{-1} M_{p,h} \mathbb{F}_h = \widehat{M}_{p,h},\tag{17}
$$

where  $\widehat{M}_{p,h} := \text{diag}(\widehat{m}_{p,h}^{(0)}, \ldots, \widehat{m}_{p,h}^{(n_h-1)})$ . Analogously, we obtain

<span id="page-13-2"></span>
$$
\mathbb{F}_h^{-1} D_h \mathbb{F}_h = \widehat{D}_h,\tag{18}
$$

where  $\widehat{D}_h := \text{diag}(\widehat{d}_h^{(0)}, \dots, \widehat{d}_h^{(n_h-1)})$  with

<span id="page-13-4"></span>
$$
\hat{d}_h^{(j)} := h^{-1} (1 - e^{2jh\pi i}).
$$
\n(19)

Using the same construction we obtain that further

<span id="page-13-3"></span>
$$
\mathbb{F}_h^{-1} D_h^T \mathbb{F}_h = \widehat{D}_h^*.
$$
\n(20)

With  $\hat{D}_h^*$  we denote the adjoint (the conjugate transpose) of the matrix  $\hat{D}_h$ . Note that  $E_h = h^2 \mathbf{1}_h \mathbf{1}_h^T$  is a circulant matrix with rank 1. The only non-zero eigenvalue is h, with corresponding eigenvector  $\underline{\mathbf{1}}_h = \underline{f}_{h,0}$ . So, we obtain

<span id="page-14-1"></span>
$$
\mathbb{F}_h^{-1} E_h \mathbb{F}_h = \widehat{E}_h \tag{21}
$$

where  $\widehat{E}_h := \text{diag}(\widehat{e}_h^{(0)})$  $\hat{e}_h^{(0)}, \ldots, \hat{e}_h^{(n_h-1)}$  with

<span id="page-14-7"></span>
$$
\hat{e}_h^{(j)} := \begin{cases} h & \text{for } j = 0 \\ 0 & \text{otherwise.} \end{cases}
$$
 (22)

So, we can determine,  $\widehat{K}_h$ , the symbol of the stiffness matrix. Using [\(11\)](#page-10-1), [\(17\)](#page-13-1), [\(18\)](#page-13-2), [\(20\)](#page-13-3) and [\(21\)](#page-14-1), we obtain that

<span id="page-14-4"></span>
$$
\mathbb{F}_h^{-1} K_{p,h} \mathbb{F}_h = \widehat{K}_h,\tag{23}
$$

where  $\widehat{K}_h := \text{diag}(\widehat{k}_{p,h}^{(0)}, \ldots, \widehat{k}_{p,h}^{(n_h-1)})$  with

<span id="page-14-6"></span>
$$
\widehat{k}_{p,h}^{(j)} := \widehat{d}_h^{(j)} \widehat{m}_{p-1,h}^{(j)} (\widehat{d}_h^{(j)})^* + \widehat{e}_h^{(j)}.
$$
\n(24)

# <span id="page-14-0"></span>3.4 Symbol of the intergrid transfer

The following lemma characterizes the symbol of the intergrid transfer.

Lemma 5 We have

<span id="page-14-5"></span><span id="page-14-2"></span>
$$
\mathbb{F}_{\frac{h}{2}}^{-1}P_{p,\frac{h}{2}}\mathbb{F}_h = \hat{P}_{p,\frac{h}{2}},\tag{25}
$$

where  $\widehat{P}_{p,\frac{h}{2}} := (\widehat{p}_{p,\frac{h}{2}}^{(i,j)}$  $_{p,\frac{h}{2}}^{(i,j)})_{i=0,\dots,n_h-1}^{j=0,\dots,n_h-1}$  with

<span id="page-14-3"></span>
$$
\widehat{p}_{p,\frac{h}{2}}^{(i,j)} := 2^{-p-1} \begin{cases} \left(1 + e^{-2i\frac{h}{2}\pi i}\right)^{p+1} & \text{for } i - j \in \{0, n_h\} \\ 0 & \text{otherwise} \end{cases}
$$
 (26)

for all  $i = 0, \ldots, 2n_h - 1$  and all  $j = 0, \ldots, n_h - 1$ .

*Proof* The equation [\(25\)](#page-14-2) is equivalent to  $P_{\frac{h}{2}} \mathbb{F}_h = \mathbb{F}_{\frac{h}{2}} \widehat{P}_{\frac{h}{2}}$ . We obtain using [\(12\)](#page-10-2) and the definition of  $\mathbb{F}_h$  for any unit vector  $\underline{\mathbf{I}}_h^{(j)}$  with  $j = 0, \ldots, n_h - 1$  that

$$
P_{\frac{h}{2}}\mathbb{F}_{h}\underline{\mathbf{I}}_{h}^{(j)} = P_{\frac{h}{2}}\underline{f}_{h,j} = 2^{-p} \left( \sum_{r \in \mathbb{Z}} \binom{p+1}{i-2r} e^{2jrh\pi i} \right)_{i=0}^{2n_h - 1}.
$$

Because  $\frac{1}{2}(1 + e^{t\pi i})$  takes the value 0 for t being odd and 1 for t being even, we can substitute  $r$  by 2t and obtain

$$
P_{\frac{h}{2}}\mathbb{F}_{h}\mathbf{I}_{h}^{(j)} = 2^{-p-1} \left( \sum_{t \in \mathbb{Z}} \binom{p+1}{i-t} e^{2jt\frac{h}{2}\pi i} (1+e^{t\pi i}) \right)_{i=0}^{2n_{h}-1}
$$
  
\n
$$
= 2^{-p-1} \left( \sum_{k \in \mathbb{Z}} \binom{p+1}{k} e^{2j(i-k)\frac{h}{2}\pi i} (1+e^{(i-k)\pi i}) \right)_{i=0}^{2n_{h}-1}
$$
  
\n
$$
= 2^{-p-1} \sum_{k \in \mathbb{Z}} \binom{p+1}{k} \left( e^{-2jk\frac{h}{2}\pi i} \underline{f}_{\frac{h}{2},j} + e^{-2(j+n_{h})k\frac{h}{2}\pi i} \underline{f}_{\frac{h}{2},j+n_{h}} \right)
$$
  
\n
$$
= 2^{-p-1} \left( 1 + e^{-2j\frac{h}{2}\pi i} \right)^{p+1} \underline{f}_{\frac{h}{2},j} + 2^{-p-1} \left( 1 + e^{-2(j+n_{h})\frac{h}{2}\pi i} \right)^{p+1} \underline{f}_{\frac{h}{2},j+n_{h}}.
$$

This shows that the j-th column of  $P_{\frac{h}{2}}\mathbb{F}_h$  is just the combination of two columns of  $\mathbb{F}_{\frac{h}{2}}$ . Therefore, the matrix  $\widehat{P}_{\frac{h}{2}}$  has just two non-zero entries, in the j-th row: those which we have claimed in [\(26\)](#page-14-3).

For determining the symbol of  $P_{p,\frac{h}{2}}^T$ , we observe as follows. As the Fourier modes  $f_{h,j}$  are pairwise orthogonal, and  $f_{h,j}^* f_{h,j} = n_h$ , we immediately obtain  $\mathbb{F}_{h}^{*}\mathbb{F}_{h} = n_{h}I$  and, consequently,  $\mathbb{F}_{h}^{-1} = h\mathbb{F}_{h}^{*}$ . So, we obtain using [\(25\)](#page-14-2) that

<span id="page-15-1"></span>
$$
\mathbb{F}_h^{-1} P_{p,\frac{h}{2}}^T \mathbb{F}_{\frac{h}{2}} = (\mathbb{F}_{\frac{h}{2}}^* P_{p,\frac{h}{2}} \mathbb{F}_h^{-*})^* = (2 \mathbb{F}_{\frac{h}{2}}^{-1} P_{p,\frac{h}{2}} \mathbb{F}_h)^* = 2 \widehat{P}_{p,\frac{h}{2}}^*.
$$
 (27)

3.5 Some statements on the symbol of the mass matrix

A closed form for the symbol of the mass matrix is not known. Within this subsection we will show a few statements characterizing the symbol, which we will need later on. Due to [\[6,](#page-33-11)16], we have

<span id="page-15-0"></span>
$$
m_{p,h}^{(j)} = h \frac{E(2p+1, p+j)}{(2p+1)!},
$$
\n(28)

where  $j \in \{-p, \ldots, p\}$ . Here,  $E(n, k)$  are the Eulerian numbers, which satisfy the recurrence relation

$$
E(n,k) = (n-k)E(n-1,k-1) + (k+1)E(n-1,k)
$$

and the initial condition

$$
E(0,j) = \begin{cases} 1 \text{ for } j = 0 \\ 0 \text{ for } j \neq 0 \end{cases}.
$$

A similar result was also stated in [\[10\]](#page-33-5). There, the entries of the mass matrix, i.e., the  $L^2$ -products of two B-splines of order p have been shown to be equal to the function value of one B-spline of order  $p + 1$ . Using the recurrence relation [\(3\)](#page-5-0), one obtains that the result in [\[10\]](#page-33-5) is equivalent to [\(28\)](#page-15-0).

As  $m_{p,h}^{(j)} = m_{p,h}^{(-j)}$  and  $e^{\theta i} + e^{-\theta i} = 2 \cos \theta$ , we obtain

<span id="page-16-1"></span><span id="page-16-0"></span>
$$
\widehat{m}_{p,h}^{(j)} = h \sum_{l=-p}^{p} \frac{E(2p+1, p+l)}{(2p+1)!} \cos(2ljh\pi).
$$

The symbol is better characterized by the following lemma.

Lemma 6 The following two statements hold:

$$
- \widehat{m}_{p,h}^{(j)} > 0 \text{ for all } j = 0, \dots, n_h - 1 \text{ and}
$$
  
\n
$$
- \widehat{m}_{p,h}^{(j)} \leq \widehat{m}_{p,h}^{(k)} \text{ for all } j,k = 0, \dots, n_h - 1 \text{ with } \cos(2jh\pi) \leq \cos(2kh\pi).
$$
  
\nProof For  $c \in [0, 2]$  we define

*Proof* For  $c \in [0, 2]$ , we define

$$
g_p(c) := \sum_{l=-p}^{p} \frac{E(2p+1, p+l)}{(2p+1)!} \cos(l \arccos(c-1))
$$

and observe  $g_p(c) = h^{-1} \hat{m}_{ph}^{(\eta(c))}$ , where  $\eta(c) := \frac{1}{2h\pi} \arccos(c-1)$ . The statement of the lemma is now equivalent to the combination of the following two statements:

$$
- h^{-1} \widehat{m}_{p,h}^{(\eta(0))} = g_p(0) > 0
$$
 and  
- 
$$
h^{-1} \widehat{m}_{p,h}^{(\eta(c))} = g_p(c)
$$
 is monotonically increasing for  $c > 0$ .

Since we can express  $\cos(l \arccos(c-1))$  as the l-th Chebyshev polynomial,  $g_p$ is a polynomial function in c. Using the recurrence relation for the Eulerian numbers, we can derive the following recurrence formula for  $g_p$ :

$$
g_p(c) = \frac{1+cp}{1+2p}g_{p-1}(c) + \frac{(2-c)(1+c(2p-1))}{p(1+2p)}g'_{p-1}(c) + \frac{(c-2)^2c}{p(1+2p)}g''_{p-1}(c).
$$
  
We can make an exact

We can make an ansatz

$$
g_p(c) = \sum_{j=0}^p a_{p,j} c^j,
$$

where we use  $0^0 = 1$ , and derive the recurrence formula

$$
a_{p,j} = \underbrace{\frac{(1-j+p)^2}{p+2p^2}}_{A_{p,j} := a_{p-1,j-1} + \underbrace{\frac{4j(p-j)+j+p}{p+2p^2}}_{B_{p,j} := a_{p-1,j} + \underbrace{\frac{2+6j+4j^2}{p+2p^2}}_{C_{p,j} := a_{p-1,j+1}}
$$

for the coefficients  $a_{p,j}$ . For  $p=1$ , we obtain

$$
a_{1,j} = \begin{cases} \frac{1}{3} & \text{for } j \in \{0, 1\} \\ 0 & \text{otherwise.} \end{cases}
$$

As  $A_{p,j} > 0$ ,  $B_{p,j} > 0$  and  $C_{p,j} > 0$  for  $0 \leq j \leq p$ , one can show using induction in p that for all  $p \geq 1$ :

$$
\begin{cases} a_{p,j} > 0 \text{ for } j \in \{0, 1, \dots, p\} \\ a_{p,j} = 0 \text{ otherwise.} \end{cases}
$$

This immediately implies that  $g_p(0) > 0$  and that  $g_p(c)$  is monotonically increasing for  $c > 0$ , which concludes the proof. 3.6 An estimate for the projection operator

Now, we are able to prove the following lemma.

Lemma 7 The inequality [\(15\)](#page-11-3) holds.

Proof The inequality [\(15\)](#page-11-3) is equivalent to

<span id="page-17-0"></span>
$$
\underbrace{h^{-1}\|M_{p-1,\frac{h}{2}}^{1/2}(I-P_{p,\frac{h}{2}}K_{p,h}^{-1}P_{p,\frac{h}{2}}^{T}K_{p,\frac{h}{2}})K_{p,\frac{h}{2}}^{-1/2}\|}_{q:=}
$$

Using Galerkin orthogonality, we obtain  $K_{p,h} = P_{p,\frac{h}{2}}^T K_{p,\frac{h}{2}} P_{p,\frac{h}{2}}$ . Note that  $\mathcal{H} := I - P_{p, \frac{h}{2}} K_{p, h}^{-1} P_{p, \frac{h}{2}}^T K_{p, \frac{h}{2}}$  is a projection operator, so  $\mathcal{H} \mathcal{H} = \mathcal{H}$ . Moreover, observe that  $\mathcal{H} K_{p, \frac{h}{2}}^{-1} = K_{p, \frac{h}{2}}^{-1} \mathcal{H}^T$ . Using these identities and  $||W||^2 = \rho(WW^T)$ , where  $\rho$  denotes the spectral radius, we obtain

$$
q^{2} = h^{-2} \rho (M_{p-1,\frac{h}{2}}^{-1/2} \mathcal{H} K_{p,\frac{h}{2}}^{-1} \mathcal{H}^{T} M_{p-1,\frac{h}{2}}^{-1/2}) = h^{-2} \rho (K_{p,\frac{h}{2}}^{-1} M_{p-1,\frac{h}{2}}^{-1} \mathcal{H})
$$
  
=  $h^{-2} \rho (K_{p,\frac{h}{2}}^{-1} M_{p-1,\frac{h}{2}} (I - P_{p,\frac{h}{2}} (P_{p,\frac{h}{2}}^{T} K_{p,\frac{h}{2}} P_{p,\frac{h}{2}})^{-1} P_{p,\frac{h}{2}}^{T} K_{p,\frac{h}{2}})).$ 

Using [\(17\)](#page-13-1), [\(18\)](#page-13-2), [\(20\)](#page-13-3), [\(23\)](#page-14-4), [\(25\)](#page-14-2) and [\(27\)](#page-15-1), we obtain further

$$
q^{2} = h^{-2} \rho \underbrace{(\hat{K}_{p,\frac{h}{2}}^{-1} \widehat{M}_{p-1,\frac{h}{2}} (I - 2\hat{P}_{p,\frac{h}{2}} (2\hat{P}_{p,\frac{h}{2}}^{*} \hat{K}_{p,\frac{h}{2}} \hat{P}_{p,\frac{h}{2}})^{-1} \hat{P}_{p,\frac{h}{2}}^{*} \hat{K}_{p,\frac{h}{2}})}_{T} \cdot \hat{F}_{p,\frac{h}{2}}.
$$

Lemma [6](#page-16-0) states that all digonal entries of  $M_{p-1,\frac{h}{2}}$  are non-zero. It is straightforward to see that also the diagonal entries of  $\hat{K}_{p, \frac{h}{2}}$  and  $\hat{K}_{p,h} = \hat{P}^*_{p, \frac{h}{2}} \hat{K}_{p, \frac{h}{2}} \hat{P}_{p, \frac{h}{2}}$ are non-zero. So,  $T_{p,\frac{h}{2}}$  is well-defined.

Recall that Lemma [5](#page-14-5) states that the matrix  $\widehat{P}_{p, \frac{h}{2}} = (\widehat{p}_{p, \frac{h}{2}}^{(i,j)})$  $\binom{i,j}{p,\frac{h}{2}}$   $\binom{j=0,\ldots,n_h-1}{i=0,\ldots,2n_h-1}$  has a block-structure, given by

$$
\widehat{p}_{p,\frac{h}{2}}^{(i,j)} = 0 \text{ for all } i - j \notin \{0, n_h\}.
$$

Therefore and because the matrices  $M_{p-1,\frac{h}{2}}$  and  $K_{p,\frac{h}{2}}$  are diagonal, the matrix  $\widehat{T}_{p,\frac{h}{2}}=(\widehat{t}_{p,\frac{h}{2}}^{(i,j)}$  $\binom{(i,j)}{p, \frac{h}{2}}$ ,  $\binom{j=0,\ldots,2n_h-1}{i=0,\ldots,2n_h-1}$  has a block-structure, given by

$$
\hat{t}_{p,\frac{h}{2}}^{(i,j)} = 0
$$
 for all  $i - j \notin \{-n_h, 0, n_h\}.$ 

By reordering the coefficients of the matrix  $T_{p, \frac{h}{2}}$ , we obtain a block-diagonal matrix with blocks

$$
\mathcal{T}_{p,\frac{h}{2}}^{(l)} = \begin{pmatrix} \hat{t}_{p,\frac{h}{2}}^{(l,l)} & \hat{t}_{p,\frac{h}{2}}^{(l,l+n_h)} \\ \hat{t}_{p,\frac{h}{2}}^{(l+n_h,l)} & \hat{t}_{p,\frac{h}{2}}^{(l+n_h,l+n_h)} \end{pmatrix}.
$$

As this block-diagonal matrix is spectrally equivalent to  $T_{p, \frac{h}{2}}$  and the spectral radius of a block-diagonal matrix is just the maximum over the spectral radii of the blocks, we obtain

$$
q^{2} = \rho(\widehat{T}_{p,\frac{h}{2}}) = \max_{l=0,\dots,n_h-1} \rho(\mathcal{T}_{p,\frac{h}{2}}^{(l)}).
$$

So, in the following, we derive the spectral radius of  $\mathcal{T}_{h}^{(l)}$  $\overline{p}$ ,  $\frac{h}{2}$  for any particular l. Straight-forward computation yields that for  $l \in \{0, \ldots, n_h-1\}, i \in \{l, l+n_h\}$ and  $j \in \{l, l + n_h\}$ , we have

$$
\hat{t}_{p,\frac{h}{2}}^{(i,j)} = \frac{\widehat{m}_{p-1,\frac{h}{2}}^{(i)}}{\widehat{k}_{p,\frac{h}{2}}^{(i)}} \left( \delta_{i,j} - \frac{\widehat{p}_{p,\frac{h}{2}}^{(i,l)} (\widehat{p}_{p,\frac{h}{2}}^{(j,l)})^*}{\sum_{r=0}^1 (\widehat{p}_{p,\frac{h}{2}}^{(l+rn_h,l)})^* \widehat{k}_{p,\frac{h}{2}}^{(l+rn_h)} \widehat{p}_{p,\frac{h}{2}}^{(l+rn_h,l)}} \widehat{k}_{p,\frac{h}{2}}^{(j)} \right), \quad (29)
$$

where  $\delta_{i,j}$  is the Kronecker-delta, i.e.,  $\delta_{i,j} = 1$  for  $i = j$  and  $\delta_{i,j} = 0$  for  $i \neq j$ .

Now, consider *case A*:  $l \in \{1, ..., n_h-1\}$ . Here, we plug the values of  $\widehat{k}_{p, \frac{l}{2}}^{(j)}$  $\binom{(J)}{p, \frac{h}{2}},$  $\hat{d}_{\frac{h}{2}}^{(j)}$ ,  $\hat{e}_{\frac{h}{2}}^{(j)}$  (which takes the value 0 for  $j \in \{l, l + n_h\}$ ),  $\hat{p}_{p, \frac{h}{2}}^{(i,j)}$ , as given by  $(24)$  $\zeta^{(i,j)}_{p,\frac{h}{2}},$  as given by  $(24),$  $(24),$ [\(19\)](#page-13-4), [\(22\)](#page-14-7) and [\(26\)](#page-14-3), into [\(29\)](#page-16-1) and substitute  $\hat{m}_{p-1,\frac{h}{2}}^{(l+n_h)}$  $\binom{(l+n_h)}{p-1,\frac{h}{2}}$  by  $\xi \hat{m}_{p-1}^{(l)}$  $\sum_{p=1,\frac{h}{2}}^{(i)}$ . Doing so, the term  $\widehat{m}_{p-}^{(l)}$  $p-1, \frac{h}{2}$  cancels out and we obtain by straight-forward computation

<span id="page-18-0"></span>
$$
\mathcal{T}_{p,\frac{h}{2}}^{(l)} = \frac{1}{\delta} \begin{pmatrix} -z(1-z)^{p-3}\xi \\ z(1+z)^{p-3} \end{pmatrix} \begin{pmatrix} (-1)^p(1-z)^{p+1} \\ (1+z)^{p+1} \end{pmatrix}^T,
$$

where  $\delta := (1+z)^{2p} + (-1)^p (1-z)^{2p} \xi$  and  $z := e^{2l\frac{h}{2}\pi i}$ . Note that the computations are not a problem, as none of the symbols (except  $\hat{e}_{\frac{\hbar}{2}}^{(j)}$ ) takes the value 0 for case A. Moreover, for case A we have that  $z \notin \{-1, 1\}$ .

Observe that  $\mathcal{T}_{n,h}^{(l)}$  $P_{p,\frac{h}{2}}^{(\ell)}$  has rank 1. Therefore, its spectral radius equals its trace, so we obtain by straight-forward computations that

$$
\rho(\mathcal{T}_{p,\frac{h}{2}}^{(l)}) = \frac{z(1+z)^{2p-2} - (-1)^p z(1-z)^{2p-2}\xi}{(1+z)^{2p} + (-1)^p (1-z)^{2p}\xi}
$$
\n
$$
= \frac{z^{-p+1}(1+2z+z^2)^{p-1} - (-1)^p z^{-p+1}(1-2z+z^2)^{p-1}\xi}{z^{-p}(1+2z+z^2)^p + (-1)^p z^{-p}(1-2z+z^2)^p\xi}
$$
\n
$$
= \frac{(z^{-1} + 2 + z)^{p-1} - (-1)^p (z^{-1} - 2 + z)^{p-1}\xi}{(z^{-1} + 2 + z)^p + (-1)^p (z^{-1} - 2 + z)^p\xi}
$$
\n
$$
= \frac{(2+2c)^{p-1} - (-1)^p (-2+2c)^{p-1}\xi}{(2+2c)^p + (-1)^p (-2+2c)^p\xi} = \frac{(1+c)^{p-1} + (1-c)^{p-1}\xi}{\frac{2((1+c)^p + (1-c)^p\xi)}{\frac{p_p(c,\xi)}{z}}}
$$

holds, where  $c := \cos(2l\frac{h}{2}\pi)$  and, as defined above,  $\xi = \widehat{m}_{p-1,\frac{h}{2}}^{(l+n_h)}$  $\binom{(l+n_h)}{p-1,\frac{h}{2}}/\widehat{m}_{p-1}^{(l)}$  $\frac{(t)}{p-1,\frac{h}{2}}$ . Note that  $c \in (-1, 1)$  holds as we have restricted ourselves to  $l \in \{1, \ldots, n_h - 1\}$ .

Observe that Lemma [6](#page-16-0) implies that  $\xi > 0$ . Now, consider two cases:

- If  $c = \cos(2l\frac{h}{2}\pi) > 0$ , then  $\cos(2(l + n_h)\frac{h}{2}\pi) = \cos(2l\frac{h}{2}\pi + \pi) \leq 0$ . For this case Lemma [6](#page-16-0) states that  $\widehat{m}_{p-1,\frac{h}{2}}^{(l+n_h)}$  $\binom{(l+n_h)}{p-1,\frac{h}{2}} \leq \widehat{m}_{p-1}^{(l)}$  $\sum_{p=1,\frac{h}{2}}^{(l)}$ , so  $\xi \leq 1$  holds.
- Analogously,  $\xi \geq 1$  holds if  $c \leq 0$ .

To finalize the proof of case A, we need to show

$$
\Psi_p\left(\cos\left(2l\frac{h}{2}\pi\right), \frac{\widehat{m}_{p-1,\frac{h}{2}}^{(l+n_h)}}{\widehat{m}_{p-1,\frac{h}{2}}^{(l)}}\right) \le \frac{1}{2}
$$

for all  $l = 1, \ldots, n_h - 1$ . It suffices to show

$$
\Psi_p(c,\xi) \le \frac{1}{2} \tag{30}
$$

for all  $(c, \xi) \in [0, 1) \times (0, 1] \cup (-1, 0] \times [1, \infty)$  and all  $p \in \mathbb{N}$ , i.e., to show the inequality for the whole range of c and  $\xi$ , ignoring their dependence on l. As a next step, we observe that  $\Psi_p(c,\xi) = \Psi_p(-c,\xi^{-1})$ , which indicates that it suffices to show [\(30\)](#page-18-0) for all  $(c, \xi) \in [0, 1) \times (0, 1]$  and all  $p \in \mathbb{N}$ . We observe that

<span id="page-19-0"></span>
$$
\Psi_p(c,\xi) = \frac{1 + \left(\frac{1-c}{1+c}\right)^{p-1} \xi}{2\left((1+c) + (1-c)\left(\frac{1-c}{1+c}\right)^{p-1} \xi\right)}
$$

and  $\omega := \left(\frac{1-c}{1+c}\right)^{p-1} \in [0,1]$  for  $c \in [0,1]$ . So, it suffices to show that

$$
\frac{1+\omega\xi}{2((1+c)+(1-c)\omega\xi)} \le \frac{1}{2}
$$
\n(31)

for all  $(c, \xi, \omega) \in [0, 1] \times (0, 1] \times [0, 1]$  and all  $p \in \mathbb{N}$ , again ignoring the dependence of  $\omega$  on p and c.

As the denominator is always positive, [\(31\)](#page-19-0) is equivalent to

$$
1 + \omega\xi \le 1 + \omega\xi + c(1 - \omega\xi),
$$

which is obviously true for all  $(c, \xi, \omega) \in [0, 1) \times (0, 1] \times [0, 1]$ .

Now, we consider *case*  $B: l = 0$ . Here, we have to use that  $\hat{e}_{p, \frac{l}{2}}^{(0)}$  $\frac{1}{p,\frac{h}{2}} \neq 0$  and obtain – by straight-forward computation – that

$$
\mathcal{T}_{p,\frac{h}{2}}^{(0)} = \begin{pmatrix} 0 & 0\\ 0 & \frac{1}{4} \end{pmatrix}
$$

and consequently  $\rho(\mathcal{T}_{n,h}^{(0)})$  $p_1(\theta) = \frac{1}{4}$ . Also this is bounded from above by  $\frac{1}{2}$ , which finishes the proof.  $\Box$ 

# 3.7 The approximation error estimate

Now, we are able to show the approximation error estimate [\(8\)](#page-8-0).

**Lemma 8** The inequality  $(8)$ , i.e.,

<span id="page-20-1"></span>
$$
\|(I-\widehat{\Pi}_{p,h})u_{p,\frac{h}{2}}\|_{L^2(-1,1)} \leq \frac{1}{\sqrt{2}}\,h|u_{p,\frac{h}{2}}|_{H^1(-1,1)},
$$

holds for all  $u_{p,\frac{h}{2}} \in S_{p,\frac{h}{2}}(-1,1)$ .

Proof Lemma [3](#page-11-0) states that  $(8)$  is a consequence of  $(14)$  and  $(15)$ . As Lemma [4](#page-12-0) shows [\(14\)](#page-11-2) and Lemma [7](#page-17-0) shows [\(15\)](#page-11-3), this finishes the proof.  $\square$ 

# 4 The proof of Theorem 1

In the previous section, we have given a proof for the approximation error of discretized functions between two consecutive grids. Using a telescoping argument, we can extend this result to an approximation error estimate for general functions. As in the last section, we first consider the periodic case.

**Lemma 9** For all  $u \in \widehat{H}^1(-1, 1)$ , all grid sizes h and each  $p \in \mathbb{N}$ , with  $hp < 1$ ,

<span id="page-20-0"></span>
$$
\|(I - \widehat{H}_{p,h})u\|_{L^2(-1,1)} \le \sqrt{2} h|u|_{H^1(-1,1)}
$$

is satisfied, where  $\widehat{\Pi}_{p,h}$  is given as in Definition [7.](#page-8-3)

Proof Using a telescoping argument, i.e. iteratively applying the triangular inequality, and the relation  $\hat{H}_{p,2h}\hat{H}_{p,h} = \hat{H}_{p,2h}$  for the projectors, we obtain for any  $q \in \mathbb{N}$ 

$$
||(I - \widehat{\Pi}_{p,h})u||_{L^2(-1,1)} \le ||(I - \widehat{\Pi}_{p,2^{-q}h})u||_{L^2(-1,1)} +\sum_{l=0}^{q-1}||(I - \widehat{\Pi}_{p,2^{-l}h})\widehat{\Pi}_{p,2^{-l-1}h}u||_{L^2(-1,1)}.
$$

We use Lemma [1](#page-6-0) and a standard Aubin-Nitsche duality argument to estimate  $||(I - \Pi_{p,2-q_h})u||_{L^2(-1,1)}$  from above. Using [\[4\]](#page-33-13), Lemma 7.6, and Lemma [1](#page-6-0) for  $r = 1$  and  $q = 2$ , we immediately obtain

<span id="page-20-2"></span>
$$
||(I - \widehat{H}_{p,2^{-q}h})u||_{L^{2}(-1,1)} \leq \widetilde{C}(p)2^{-q}h||u||_{H^{1}(-1,1)},
$$
\n(32)

where  $\widetilde{C}(p)$  is independent of the grid size. Using [\(32\)](#page-20-2) and Lemma [8,](#page-20-1) we obtain

$$
||(I - \widehat{\Pi}_{p,h})u||_{L^2(-1,1)} \le \widetilde{C}(p) 2^{-q}h||u||_{H^1(-1,1)} + \sum_{l=0}^{q-1} \frac{1}{\sqrt{2}} 2^{-l}h|\widehat{\Pi}_{p,2^{-l-1}h}u|_{H^1(-1,1)}.
$$

Because  $\widehat{H}_{p,h}$  is  $H^1$ -orthogonal, we obtain  $|\widehat{H}_{p,2^{-l-1}h}u|_{H^1(-1,1)} \leq |u|_{H^1(-1,1)}$ and further

$$
\|(I-\widehat{\Pi}_{p,h})u\|_{L^2(-1,1)} \le \widetilde{C}(p) 2^{-q}h\|u\|_{H^1(-1,1)} + \sum_{l=0}^{q-1} \frac{1}{\sqrt{2}} 2^{-l}h|u|_{H^1(-1,1)}.
$$

The summation formula for the infinite geometric series gives

$$
\|(I-\widehat{\Pi}_{p,h})u\|_{L^2(-1,1)} \leq \widetilde{C}(p) 2^{-q}h\|u\|_{H^1(-1,1)} + 2\frac{1}{\sqrt{2}}h|u|_{H^1(-1,1)}.
$$

As this is true for all  $q \in \mathbb{N}$ , we can take the limit  $q \to \infty$  and obtain the desired result.  $\Box$ 

Having this result, we note that Theorem [1](#page-2-0) is just the extension of Lemma [9](#page-20-0) to the non-periodic case. So, we can easily prove Theorem [1.](#page-2-0)

Proof of Theorem [1](#page-2-0) In the following, we assume without loss of generality that  $\Omega = (0, 1)$ . The extension to any other  $\Omega = (a, b)$ , follows using a standard scaling argument.

Observe that any  $u \in H^1(0,1)$  can be extended to a  $w \in \hat{H}^1(-1,1)$  by defining  $w(x) := u(|x|)$ . The assumption  $hp < 1$  $hp < 1$  in Theorem 1 guarantees that Lemma [9](#page-20-0) can be applied. We set  $w_{p,h} := \widehat{H}_{p,h}w \in \widehat{S}_{p,h}(-1,1)$  as in Lemma [9,](#page-20-0) such that √

$$
||w - w_{p,h}||_{L^2(-1,1)} \le \sqrt{2} h |w|_{H^1(-1,1)}.
$$

The function  $w_{p,h}$  is symmetric, i.e.,  $w_{p,h}(x) = w_{p,h}(-x)$  holds. This can be seen by the following argument: As w satisfies  $w(x) = w(-x)$ , we have for  $\widetilde{w}_{p,h}(x) := w_{p,h}(-x)$ 

$$
||w - w_{p,h}||_{L^2(-1,1)} = ||w - \widetilde{w}_{p,h}||_{L^2(-1,1)}
$$

and as  $w_{p,h}$  was a unique minimizer, consequently  $w_{p,h}(x) = \tilde{w}_{p,h}(x)$  $w_{p,h}(-x)$  holds. By restricting  $w_{p,h}$  to  $(0,1)$ , we obtain a function  $u_{p,h} \in$  $S_{p,h}(0,1)$ . This function satisfies the desired approximation error estimate since  $|w|_{H^1(-1,1)} = \sqrt{2}|u|_{H^1(0,1)}$  and  $||w-w_{p,h}||_{L^2(-1,1)} = \sqrt{2}||u-w_{p,h}||_{L^2(0,1)}$ hold due to the symmetry of  $w$ .

### <span id="page-21-0"></span>5 Approximation error estimate for a reduced spline space

In the proof of Theorem [1](#page-2-0) we have defined  $u_{p,h}$  to be the restriction of a symmetric and periodic spline  $w_{p,h} \in \widehat{S}_{p,h}(-1,1)$  to  $(0,1)$ . So, we know more about  $u_{p,h}$  than just  $u_{p,h} \in S_{p,h}(0,1)$ . Throughout this Section we again assume  $hp < |\Omega|$ .

As we have shown in the proof of Theorem [1](#page-2-0) the spline  $w_{p,h}$  is symmetric, i.e.,  $w_{p,h}(x) = w_{p,h}(-x)$ , so we have

$$
\frac{\partial^l}{\partial x^l} w_{p,h}(x) = (-1)^l \frac{\partial^l}{\partial x^l} w_{p,h}(-x) \text{ for all } l \in \mathbb{N}_0.
$$

	dim $S_{p,h}(0,1)$	dim $\widehat{S}_{p,h}(0,1)$	dim $S_{p,h}(0,1)$
p even	$n+p$	$\, n$	n
p odd	$n+p$	$\boldsymbol{n}$	$n+1$

<span id="page-22-4"></span>**Table 1** Degrees of freedom, where *n* is the number of elements in  $(0, 1)$ .

By plugging  $x = 0$  into this relation, we obtain that all odd derivatives vanish for  $x = 0$ . By plugging  $x = 1$  into the relation, we obtain together with [\(2\)](#page-5-1) that also for  $x = 1$  all odd derivatives vanish.

So, we have shown that the approximation error estimate [\(1\)](#page-2-1) is still satisfied if we restrict the approximating spline  $u_{p,h}$  to be in the space  $\widetilde{S}_{p,h}(0,1)$ , defined as follows.

<span id="page-22-0"></span>**Definition 8** Given a spline space  $S_{p,h}(\Omega)$  over  $\Omega = (a, b)$ , the space of splines with vanishing odd derivatives  $\widetilde{S}_{p,h}(\Omega)$  is the space of all  $u_{p,h} \in S_{p,h}(\Omega)$ that satisfy the following condition:

$$
\frac{\partial^{2l+1}}{\partial x^{2l+1}}u_{p,h}(a) = \frac{\partial^{2l+1}}{\partial x^{2l+1}}u_{p,h}(b) = 0 \text{ for all } l \in \mathbb{N}_0 \text{ with } 2l+1 < p.
$$

Using a standard scaling argument, we can again extend the result for  $\Omega = (0, 1)$  to any  $\Omega = (a, b)$  and obtain the following Corollary.

<span id="page-22-1"></span>**Corollary 1** For all  $u \in H^1(\Omega)$ , all grid sizes h and all  $p \in \mathbb{N}$ , with  $hp < |\Omega|$ , there is a spline approximation  $u_{p,h} \in \widetilde{S}_{p,h}(\Omega)$  such that

$$
||u - u_{p,h}||_{L^2(\Omega)} \le \sqrt{2} \; h |u|_{H^1(\Omega)}
$$

is satisfied.

In the Appendix, we will introduce a basis for the space  $\widetilde{S}_{p,h}(\Omega)$ . Based on the bases of those spaces, we obtain that their dimensions are as given in Table [1.](#page-22-4)

### <span id="page-22-3"></span>6 An inverse inequality for the reduced spline space

For the space  $\widetilde{S}_{p,h}(\Omega)$ , a robust inverse inequality holds. Note that an extension to  $S_{p,h}(\Omega)$  is not possible (cf. Remark [2\)](#page-3-1).

<span id="page-22-2"></span>**Theorem 4** For all grid sizes h and each  $p \in \mathbb{N}$ ,

<span id="page-22-5"></span>
$$
|u_{p,h}|_{H^1(\Omega)} \le 2\sqrt{3}h^{-1} \|u_{p,h}\|_{L^2(\Omega)} \tag{33}
$$

is satisfied for all  $u_{p,h} \in \widetilde{S}_{p,h}(\Omega)$ .

*Proof* In the following, we assume without loss of generality that  $\Omega = (0, 1)$ . The extension to any other  $\Omega = (a, b)$ , follows directly using a standard scaling argument. We can extend every  $u_{p,h} \in \widetilde{S}_{p,h}(0,1)$  to  $(-1,1)$  by defining  $w_{p,h}(x) := u_{p,h}(|x|)$  and obtain  $w_{p,h} \in \widehat{S}_{p,h}(-1,1)$ . [\(33\)](#page-22-5) is equivalent to

<span id="page-23-1"></span>
$$
|w_{p,h}|_{H^1(-1,1)} \le 2\sqrt{3}h^{-1}||w_{p,h}||_{L^2(-1,1)}.
$$
\n(34)

This is shown using induction in p for all  $u \in \widetilde{S}_{p,h}(-1,1)$ . For  $p = 1$ , [\(34\)](#page-23-1) is known, cf. [\[14\]](#page-33-14), Theorem 3.91.

Now, we show that the constant does not increase for larger  $p$ . So assume  $p > 1$  to be fixed. Due to the periodicity and due to the Cauchy-Schwarz inequality,

$$
|w_{p,h}|_{H^1(-1,1)}^2 = \int_{-1}^1 (w'_{p,h})^2 dx = -\int_{-1}^1 w''_{p,h} w_{p,h} dx
$$
  
\$\leq \|w''\_{p,h}\|\_{L^2(-1,1)} \|w\_{p,h}\|\_{L^2(-1,1)} = |w'\_{p,h}|\_{H^1(-1,1)} \|w\_{p,h}\|\_{L^2(-1,1)}

is satisfied. Using the induction assumption (and  $w'_{p,h} \in \widehat{S}_{p-1,h}(-1,1)$ , cf. [\[13\]](#page-33-1), Theorem 5.9), we know that

$$
|w'_{p,h}|_{H^1(-1,1)} \le 2\sqrt{3}h^{-1} \|w'_{p,h}\|_{L^2(-1,1)} = 2\sqrt{3}h^{-1} |w_{p,h}|_{H^1(-1,1)}.
$$

Combining these results, we obtain

$$
|w_{p,h}|^2_{H^1(-1,1)} \le 2\sqrt{3}h^{-1}|w_{p,h}|_{H^1(-1,1)}||w_{p,h}||_{L^2(-1,1)}
$$

and further

$$
|w_{p,h}|_{H^1(-1,1)} \le 2\sqrt{3}h^{-1} \|w_{p,h}\|_{L^2(-1,1)}.
$$

This shows [\(34\)](#page-23-1), which concludes the proof.  $\square$ 

Remark 3 Neither Theorem 3.91 in [\[14\]](#page-33-14), nor any of the arguments in the proof of Theorem [4](#page-22-2) requires the grid to be equidistant. So, also having a general grid, estimate √

<span id="page-23-2"></span>
$$
|u_{p,h}|_{H^1(\Omega)} \le 2\sqrt{3} \; h_{\min}^{-1} \|u_{p,h}\|_{L^2(\Omega)}
$$

is satisfied for all splines  $u_{p,h}$  on  $\Omega = (a, b)$  with vanishing odd derivatives at the boundary. Here, as in any standard inverse inequality,  $h_{\min}$  is the size of the smallest element.

As we have proven both an approximation error estimate and a corresponding inverse inequality, both of them are sharp (up to constants independent of  $p$  and  $h$ ). First, we show that there is a lower bound for the approximation error. As [\(35\)](#page-23-2) is obviously true for constant functions, we show that there also exist other functions satisfying this inequality.

<span id="page-23-0"></span>**Corollary 2** For all grid sizes h and each  $p \in \mathbb{N}$ , there is a non-constant function  $u \in H^1(\Omega)$  such that

$$
\inf_{u_{p,h}\in \widetilde{S}_{p,h}(\Omega)} \|u - u_{p,h}\|_{L^2(\Omega)} \ge \frac{1}{4\sqrt{3}} \; h|u|_{H^1(\Omega)}.\tag{35}
$$

*Proof* Let  $u \in S_{p,\frac{h}{2}}(\Omega) \setminus \{0\}$  be such that  $(u,\tilde{u}_{p,h})_{L^2(\Omega)} = 0$  for all  $\tilde{u}_{p,h} \in$  $S_{p,h}(\Omega)$ . As the constant functions are in  $S_{p,h}(\Omega)$ , this orthogonality implies that  $u$  is non-constant. Using this orthogonality, we know that the infimum in [\(35\)](#page-23-2) is taken for  $u_{p,h} = 0$ . So, we obtain using Theorem [4](#page-22-2) inf $\lim_{u_{p,h} \in \widetilde{S}_{p,h}(\Omega)} ||u-\rangle$  $u_{p,h}||_{L^2(\Omega)} = ||u||_{L^2(\Omega)} \geq \frac{1}{2\nu}$  $\frac{1}{2\sqrt{3}}$  $\frac{h}{2}|u|_{H^1(\Omega)}$ , which finishes the proof.  $\square$ 

Similarly, we can give an lower bound for the inverse inequality. Again, we show the existence of a non-trivial function.

**Corollary 3** For all grid sizes h with  $2hp < |\Omega|$  and each  $p \in \mathbb{N}$ , there is a non-constant function  $u_{p,h} \in \widetilde{S}_{p,h}(\Omega)$  such that

<span id="page-24-0"></span>
$$
|u_{p,h}|_{H^1(\Omega)} \ge \frac{1}{2\sqrt{2}} \; h^{-1} \|u_{p,h}\|_{L^2(\Omega)}.
$$

*Proof* Let  $u_{p,h} \in S_{p,h}(\Omega) \setminus \{0\}$  be such that  $(u_{p,h}, \tilde{u}_{p,2h})_{L^2(\Omega)} = 0$  for all  $\tilde{u}_{p,2h} \in$  $S_{p,2h}(\Omega)$ . As the constant functions are in  $S_{p,2h}(\Omega)$ , this orthogonality implies that  $u_{p,h}$  is non-constant. Using this orthogonality and Theorem [1,](#page-2-0) we ob- $\text{tain } \|u_{p,h}\|_{L^2(\Omega)} = \text{inf}_{u_{p,2h}\in \widetilde{S}_{p,n-1}(\Omega)} \|u_{p,h} - u_{p,2h}\|_{L^2(\Omega)} \leq \sqrt{2(2h)}|u_{p,h}|_{H^1(\Omega)},$ which finishes the proof.  $\Box$ 

#### <span id="page-24-1"></span>7 An extension to higher Sobolev indices

We can easily lift the statement of Theorem [1](#page-2-0) (and also Corollary [1\)](#page-22-1) up to higher Sobolev indices.

<span id="page-24-2"></span>**Theorem 5** For all grid sizes h, each  $q \in \mathbb{N}$  and each  $p \in \mathbb{N}$  with  $0 < q \leq p+1$ and with  $h(p-q+1) < |\Omega|$ , there is for each  $u \in H^q(\Omega)$ , a spline approximation  $u_{p,h} \in \widetilde{S}_{p,h}^{(q)}(\Omega)$  such that

$$
|u - u_{p,h}|_{H^{q-1}(\Omega)} \le \sqrt{2} \, h |u|_{H^q(\Omega)},
$$

where  $\widetilde{S}_{p,h}^{(q)}(\Omega)$  is the space of all  $u_{p,h} \in S_{p,h}(\Omega)$  that satisfy the following symmetry condition:

$$
\frac{\partial^{2l+q}}{\partial x^{2l+q}}u_{p,h}(a) = \frac{\partial^{2l+q}}{\partial x^{2l+q}}u_{p,h}(b) = 0 \text{ for all } l \in \mathbb{N}_0 \text{ with } 2l+q < p.
$$

*Proof* Let again  $\Omega = (0, 1)$  without loss of generality. The proof is done by induction. From Corollary [1,](#page-22-1) we know the estimate for  $q = 1$  (as  $\widetilde{S}_{p,h}^{(1)}(0,1) =$  $\widetilde{S}_{p,h}(0,1)$  and all  $p > q - 1 = 0$ . For  $q = 1$  and  $p = q - 1 = 0$ , the estimate is a well-known result, cf. [\[13\]](#page-33-1), Theorem 6.1,  $(6.7)$ , where  $(in \text{ our notation})$  $|u - u_{0,h}|_{L^2(0,1)} \leq h|u|_{H^1(0,1)}$  has been shown.

So, now we assume to know the estimate for some  $q-1$  and show it for q. As  $u \in H^q(0,1)$ , we know that  $u' \in H^{q-1}(0,1)$ , so we can apply the induction hypothesis and obtain that there is some  $u_{p-1,n} \in \widetilde{S}_{p-1,n}^{(q-1)}(0,1)$  with

$$
|u'-u_{p-1,n}|_{H^{q-2}(0,1)} \leq \sqrt{2} \; h|u'|_{H^{q-1}(0,1)}.
$$

Define

<span id="page-25-0"></span>
$$
u_{p,h}(x) := c + \int_0^x u_{p-1,n}(\xi) d\xi.
$$
 (36)

Note that  $u_{p,h} \in S_{p,h}(0,1)$  as integrating increases both the polynomial degree and the differentiability by 1, cf. [\[13\]](#page-33-1), Theorem 5.16. After integrating, the boundary conditions on the *l*-th derivative become conditions on the  $l + 1$ -st derivative, therefore we further have  $u_{p,h} \in \widetilde{S}_{p,h}^{(q)}(0,1)$ .

Therefore, we have

$$
|u'-u'_{p,h}|_{H^{q-2}(0,1)} \leq \sqrt{2} \; h|u'|_{H^{q-1}(0,1)},
$$

which is the same as

$$
|u - u_{p,h}|_{H^{q-1}(0,1)} \le \sqrt{2} \; h |u|_{H^q(0,1)}.
$$

The bound on the grid size with respect to the degree, i.e.  $h(p - q + 1) < |Q|$ is sufficient, as the degree of  $\partial^{q-1}/\partial x^{q-1}u$  is equal to  $p-q+1$ . This finishes the proof.  $\Box$ 

Remark 4 The integration constant (integration constants for  $q > 2$ ) in [\(36\)](#page-25-0) can be used to guarantee that

$$
\int_{\Omega} \frac{\partial^l}{\partial x^l} (u(x) - u_{p,h}(x)) \mathrm{d}x = 0
$$

for all  $l \in \{0, 1, \ldots, q - 1\}.$ 

For the spaces  $\widetilde{S}_{p,h}^{(q)}(\Omega)$  there is again an inverse inequality.

**Theorem 6** For all grid sizes h, each  $q \in \mathbb{N}$  and each  $p \in \mathbb{N}$  with  $0 < q \leq p+1$ ,

<span id="page-25-1"></span>
$$
|u_{p,h}|_{H^q(\Omega)} \le 2\sqrt{3}h^{-1}|u_{p,h}|_{H^{q-1}(\Omega)} \tag{37}
$$

is satisfied for all  $u_{p,h} \in \widetilde{S}_{p,h}^{(q)}(\Omega)$ , where  $\widetilde{S}_{p,h}^{(q)}(\Omega)$  is as defined in Theorem [5.](#page-24-2)

Proof First note that [\(37\)](#page-25-1) is equivalent to

<span id="page-25-2"></span>
$$
\left| \frac{\partial^{q-1}}{\partial x^{q-1}} u_{p,h} \right|_{H^1(\Omega)} \leq 2\sqrt{3}h^{-1} \left\| \frac{\partial^{q-1}}{\partial x^{q-1}} u_{p,h} \right\|_{L^2(\Omega)}.
$$
\n(38)

As  $\frac{\partial^{q-1}}{\partial x^{q-1}} u_{p,h} \in \widetilde{S}_{p-q+1,n}^{(1)}(\Omega) = \widetilde{S}_{p-q+1,n}(\Omega)$ , cf. [\[13\]](#page-33-1), Theorem 5.9, the estimate [\(38\)](#page-25-2) follows directly from Theorem [4.](#page-22-2)

Again, as we have both an approximation error estimate and an inverse inequality, we know that both of them are sharp (cf. Corollaries [2](#page-23-0) and [3\)](#page-24-0).

<span id="page-25-3"></span>The following theorem is directly obtained from telescoping.

**Theorem 7** For all grid sizes h, each  $q \in \mathbb{N}_0$ , each  $p \in \mathbb{N}$ , each  $r \in \mathbb{N}$  with  $0 \leq r \leq q \leq p+1$  and  $h(p-r) < |\Omega|$ , there is for each  $u \in H^{q}(\Omega)$  a spline approximation  $u_{p,h} \in S_{p,h}(\Omega)$  such that

$$
|u - u_{p,h}|_{H^r(\Omega)} \le (\sqrt{2} \; h)^{q-r} |u|_{H^q(\Omega)}
$$

is satisfied.

*Proof* Theorem [5](#page-24-2) states the desired result for  $r = q - 1$ . For  $r < q - 1$ , the statement is shown by induction in  $r$ . So, we assume to know the desired result for some r, i.e., there is a spline approximation  $w_{p,h} \in S_{p,h}(\Omega)$  such that

<span id="page-26-1"></span>
$$
|u - w_{p,h}|_{H^r(\Omega)} \le (\sqrt{2} \; h)^{q-r} |u|_{H^q(\Omega)}.
$$
\n(39)

Now, we show that there is some  $u_{p,h} \in S_{p,h}(\Omega)$  such that

<span id="page-26-2"></span>
$$
|u - u_{p,h}|_{H^{r-1}(\Omega)} \le (\sqrt{2} \; h)^{q - (r-1)} |u|_{H^q(\Omega)}.
$$
\n(40)

As  $u-w_{p,h} \in H^r(\Omega)$ , Theorem [5](#page-24-2) states that there is a function  $u_{p,h} \in S_{p,h}(0,1)$ such that √

$$
|u - u_{p,h}|_{H^{r-1}(\Omega)} \le \sqrt{2} \; h |u - w_{p,h}|_{H^r(\Omega)},
$$

which shows together with the induction assumption [\(39\)](#page-26-1) the induction hy-pothesis [\(40\)](#page-26-2). Again, the bound on the grid size  $h(p-r) < |\Omega|$  follows directly from the bounds in Theorem [5.](#page-24-2)  $\Box$ 

Here, it is not known to the authors how to choose a proper subspace of  $S_{p,h}(\Omega)$  such that a complementary inverse inequality can be shown.

# <span id="page-26-0"></span>8 Extension to two and more dimensions and application in Isogeometric Analysis

Without loss of generality and to simplify the notation, we restrict ourselves to  $\Omega := (0, 1)^d$  $\Omega := (0, 1)^d$  $\Omega := (0, 1)^d$  throughout this section. We can extend Theorem 1 (and also Corollary [1\)](#page-22-1) to the following theorem for a tensor-product structured grid on  $\Omega$ . Here, we can introduce  $\widetilde{W}_{p,h}(\Omega) = \otimes_{l=1}^d \widetilde{S}_{p,h}(0,1)$ . Let  $n = n_h$ , for even p, and  $n = n_h + 1$  for odd p. Assuming that  $(\widetilde{\varphi}_{p,h}^{(0)}, \ldots, \widetilde{\varphi}_{p,h}^{(n-1)})$  is a basis of  $\widetilde{S}_{p,h}(0,1)$ , the space  $\widetilde{W}_{p,h}(\Omega)$  is given by

$$
\widetilde{W}_{p,h}(\Omega)=\left\{w\,:\,w(x_1,\ldots,x_d)=\sum_{i_1,\ldots,i_d=0}^{n-1}w_{i_1,\ldots,i_d}\widetilde{\varphi}_{p,h}^{(i_1)}(x_1)\cdots\widetilde{\varphi}_{p,h}^{(i_d)}(x_d)\right\}.
$$

<span id="page-26-3"></span>**Theorem 8** For all  $u \in H^1(\Omega)$ , all grid sizes h and each  $p \in \mathbb{N}_0$ , with  $hp < 1$ , there is a spline approximation  $w_{p,h} \in W_{p,n}(\Omega)$  such that

$$
||u - w_{p,h}||_{L^2(\Omega)} \le \sqrt{2d} \, h|u|_{H^1(\Omega)}
$$

is satisfied.

The proof is similar to the proof in [\[3\]](#page-33-15), Section 4, for the two dimensional case. To keep the paper self-contained we give a proof of this theorem.

*Proof of Theorem [8](#page-26-3)* For sake of simplicity, we restrict ourselves to  $d = 2$ . The extension to more dimensions is completely analogous. Here

$$
\widetilde{W}_{p,h}(\Omega) = \widetilde{S}_{p,h}(0,1) \otimes \widetilde{S}_{p,h}(0,1) = \left\{ w : w(x,y) = \sum_{i,j=0}^{n-1} w_{i,j} \widetilde{\varphi}_{p,h}^{(i)}(x) \widetilde{\varphi}_{p,h}^{(j)}(y) \right\}.
$$

We assume  $u \in C^{\infty}(\Omega)$  and show the desired result using a standard density argument. Using Corollary [1,](#page-22-1) we can introduce for each  $x \in (0,1)$  a function  $v(x, \cdot) \in \widetilde{S}_{p,h}(0,1)$  with

$$
||u(x, \cdot) - v(x, \cdot)||_{L^2(0,1)} \le \sqrt{2} \; h |u(x, \cdot)|_{H^1(0,1)}.
$$

By squaring and taking the integral over  $x$ , we obtain

<span id="page-27-2"></span>
$$
||u - v||_{L^{2}(\Omega)} \leq \sqrt{2} h \left\| \frac{\partial}{\partial y} u \right\|_{L^{2}(\Omega)}.
$$
\n(41)

By choosing  $v(x, \cdot)$  to be the  $L^2$ -orthogonal projection, we also have

$$
||v(x, \cdot)||_{L^2(0,1)} \leq ||u(x, \cdot)||_{L^2(0,1)}
$$

for all  $x \in (0,1)$  and consequently

<span id="page-27-1"></span>
$$
\left\|\frac{\partial}{\partial x}v(x,\cdot)\right\|_{L^2(0,1)} \le \left\|\frac{\partial}{\partial x}u(x,\cdot)\right\|_{L^2(0,1)}.\tag{42}
$$

As  $v(x, \cdot) \in \widetilde{S}_{p,h}(0,1)$ , there are coefficients  $v_j(x)$  such that

$$
v(x,y) = \sum_{j=0}^{n-1} v_j(x) \widetilde{\varphi}_{p,h}^{(j)}(y).
$$

Using Corollary [1,](#page-22-1) we can introduce for each  $j \in \{0, ..., N\}$  a function  $w_j \in$  $\widetilde{S}_{p,h}(0,1)$  with √

<span id="page-27-0"></span>
$$
||v_j - w_j||_{L^2(0,1)} \le \sqrt{2} \; h|v_j|_{H^1(0,1)}.\tag{43}
$$

Next, we introduce a function  $w$  by defining

$$
w(x,y) := \sum_{j=0}^{n-1} w_j(x) \widetilde{\varphi}_{p,h}^{(j)}(y),
$$

which is obviously a member of the space  $\widetilde{W}_{p,n}(\Omega)$ . By squaring [\(43\)](#page-27-0), multiplying it with  $\tilde{\varphi}_{p,h}^{(j)}(y)^2$ , summing over j and taking the integral, we obtain

<span id="page-27-3"></span>
$$
\int_0^1 \sum_{j=0}^{n-1} \|v_j - w_j\|_{L^2(0,1)}^2 \widetilde{\varphi}_{p,h}^{(j)}(y)^2 dy \le 2 \, h^2 \int_0^1 \sum_{j=0}^{n-1} |v_j|_{H^1(0,1)}^2 \widetilde{\varphi}_{p,h}^{(j)}(y)^2 dy.
$$

Using the definition of the norms, we obtain

$$
\int_0^1 \int_0^1 \sum_{j=0}^{n-1} (v_j(x) - w_j(x))^2 \widetilde{\varphi}_{p,h}^{(j)}(y)^2 dx dy \le 2 h^2 \int_0^1 \int_0^1 \sum_{j=0}^{n-1} v_j'(x)^2 \widetilde{\varphi}_{p,h}^{(j)}(y)^2 dx dy
$$

and further

$$
||v - w||_{L^2(\Omega)} \le \sqrt{2} h \left\| \frac{\partial}{\partial x} v \right\|_{L^2(\Omega)}.
$$

Using [\(42\)](#page-27-1), we obtain

<span id="page-28-0"></span>
$$
||v - w||_{L^{2}(\Omega)} \le \sqrt{2} h \left\| \frac{\partial}{\partial y} u \right\|_{L^{2}(\Omega)}.
$$
\n(44)

Using  $(41)$  and  $(44)$ , we obtain

$$
||u - w||_{L^2(\Omega)} \le ||u - v||_{L^2(\Omega)} + ||v - w||_{L^2(\Omega)}
$$
  
\n
$$
\le \sqrt{2} h \left\| \frac{\partial}{\partial y} u \right\|_{L^2(\Omega)} + \sqrt{2} h \left\| \frac{\partial}{\partial x} u \right\|_{L^2(\Omega)}
$$
  
\n
$$
\le 2 h |u|_{H^1(\Omega)}, \tag{45}
$$

which finishes the proof.  $\Box$ 

The extension of Theorem [4](#page-22-2) to two or more dimensions is rather easy.

**Theorem 9** For all grid sizes h and each  $p \in \mathbb{N}$ , the inequality

<span id="page-28-1"></span>
$$
|u_{p,h}|_{H^1(\Omega)} \le 2\sqrt{3d} \; h^{-1} \; \|u_{p,h}\|_{L^2(\Omega)}
$$

is satisfied for all  $u_{p,h} \in \widetilde{W}_{p,h}(\Omega)$ .

*Proof* For sake of simplicity, we restrict ourselves to  $d = 2$ . The generalization to more dimensions is completely analogous.

We have obviously

$$
|u_{p,h}|_{H^1(\Omega)}^2 = \left\| \frac{\partial}{\partial x} u_{p,h} \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial}{\partial y} u_{p,h} \right\|_{L^2(\Omega)}^2
$$
  
= 
$$
\int_0^1 |u_{p,h}(\cdot, y)|_{H^1(0,1)}^2 dy + \int_0^1 |u_{p,h}(x, \cdot)|_{H^1(0,1)}^2 dx
$$

This can be bounded from above using Theorem [4](#page-22-2) via

$$
|u_{p,h}|_{H^1(\Omega)}^2 \le 12h^{-2} \left( \int_0^1 \|u_{p,h}(\cdot,y)\|_{L^2(0,1)}^2 dy + \int_0^1 \|u_{p,h}(x,\cdot)\|_{L^2(0,1)}^2 dx \right)
$$
  
=24h<sup>-2</sup>  $||u_{p,h}||_{L^2(\Omega)}^2$ ,

which finishes the proof.



The extension to isogeometric spaces can be done following the approach presented in [\[1\]](#page-33-3), Section 3.3. In Isogeometric Analysis, we have a geometry parameterization  $\mathbf{F} : (0,1)^d \to \hat{\Omega}$ . An isogeometric function on  $\hat{\Omega}$  is then given as the composition of a B-spline on  $(0, 1)^d$  with the inverse of **F**. The following result can be shown using a standard chain rule argument.

There exists a constant  $C = C(\mathbf{F}, q)$  such that

$$
C^{-1} \|f\|_{H^{q}(\hat{\Omega})} \le \|f \circ \mathbf{F}\|_{H^{q}((0,1)^{d})} \le C \|f\|_{H^{q}(\hat{\Omega})}
$$
(46)

for all  $f \in H^q(\hat{\Omega})$ .

See [\[1\]](#page-33-3), Lemma 3.5, or [\[3\]](#page-33-15), Corollary 5.1, for related results. In both papers the statements are slightly more general, [\[1\]](#page-33-3) gives a more detailed dependence on the parameterization F whereas [\[3\]](#page-33-15) establishes bounds for anisotropic grids. Obviously, an extension to anisotropic grids can be achieved directly using the estimate [\(45\)](#page-28-0). Note that the degree and the grid size are not necessarily equal in each parameter direction.

Using this equivalence of norms, we can transfer all results from the parameter domain  $(0,1)^d$  to the physical domain  $\hat{\Omega}$ . However, we need to point out that this equivalence is not valid for seminorms. Hence, in Theorem [1](#page-2-0) (and follow-up Theorems [5,](#page-24-2) [7](#page-25-3) and [8\)](#page-26-3) the seminorms on the right hand side of the equations need to be replaced by the full norms. Moreover, the bounds depend on the geometry parameterization via the constant  $C$  in [\(46\)](#page-28-1).

A similar strategy can be followed when extending the results to NURBS. We do not go into the details here but refer to [\[1,](#page-33-3)3] for a more detailed study. In the case of NURBS the seminorms again have to be replaced by the full norms due to the quotient rule of differentiation. In that case the constants of the bounds additionally depend on the given denominator of the NURBS parameterization.

### Acknowledgements

The authors want to thank Walter Zulehner for his suggestions, which helped to improve the presentation of the results in this paper.

### Appendix

At this point, we want to give a basis for  $\widetilde{S}_{p,h}(\Omega)$  to make the reader more familiar with that space and to demonstrate that it is possible to work with it. The basis, which we introduce, is directly related to the (scaled) cardinal B-splines  $\{\varphi_{p,h}^{(i)}\}_{i=-p}^{n_h-1}$ .

<span id="page-29-1"></span><span id="page-29-0"></span>**Lemma 10** The set 
$$
\{\widetilde{\varphi}_{p,h}^{(i)}\}_{i=-\lceil \frac{p}{2} \rceil,\dots,n_h-\lfloor \frac{p}{2} \rfloor-1}
$$
 with  

$$
\widetilde{\varphi}_{p,h}^{(i)} := \sum_{l \in \{-i-p-1,i,2n_h-i-p-1\}} \varphi_{p,h}^{(l)}
$$
(47)

is a basis of  $\widetilde{S}_{p,h}(\Omega)$ .

Before we prove Lemma [10](#page-29-0) we give a more practical representation of the basis functions by removing all contributions vanishing in  $\Omega$ . We obtain for odd p that

$$
\widetilde{\varphi}_{p,h}^{(i)} = \varphi_{p,h}^{(i)}
$$
\n
$$
\widetilde{\varphi}_{p,h}^{(i)} = \varphi_{p,h}^{(i)} + \varphi_{p,h}^{(-i-p-1)}
$$
\n
$$
i = -(p+1)/2
$$
\n
$$
\widetilde{\varphi}_{p,h}^{(i)} = \varphi_{p,h}^{(i)}
$$
\n
$$
i = -(p-1)/2, \dots, -1
$$
\n
$$
\widetilde{\varphi}_{p,h}^{(i)} = \varphi_{p,h}^{(i)}
$$
\n
$$
i = 0, \dots, n_h - p
$$
\n
$$
\widetilde{\varphi}_{p,h}^{(i)} = \varphi_{p,h}^{(i)} + \varphi_{p,h}^{(2n_h - i - p - 1)}
$$
\n
$$
i = n_h - p + 1, \dots, n_h - (p+1)/2
$$
\n
$$
\widetilde{\varphi}_{p,h}^{(i)} = \varphi_{p,h}^{(i)}
$$
\n
$$
i = n_h - (p-1)/2
$$

and for even  $p$  that

$$
\widetilde{\varphi}_{p,h}^{(i)} = \varphi_{p,h}^{(i)} + \varphi_{p,h}^{(-i-p-1)} \qquad i = -p/2, \dots, -1
$$
  
\n
$$
\widetilde{\varphi}_{p,h}^{(i)} = \varphi_{p,h}^{(i)} \qquad i = 0, \dots, n_h - p - 1
$$
  
\n
$$
\widetilde{\varphi}_{p,h}^{(i)} = \varphi_{p,h}^{(i)} + \varphi_{p,h}^{(2n_h - i - p - 1)} \qquad i = n_h - p, \dots, n_h - p/2 - 1.
$$

Note that here we need that  $0 \leq n_h - p - 1$ , which is equivalent to  $hp < 1$ .

*Proof of Lemma [10](#page-29-0)* For the sake of simplicity, we consider the case  $\Omega = (0, 1)$ only. We show first that every function in [\(47\)](#page-29-1) is in  $\widetilde{S}_{p,h}(0,1)$ . Note that we have constructed  $\widetilde{S}_{p,h}(0,1)$  such that the restriction of any symmetric function in  $S_{p,h}(-1,1)$  to  $(0,1)$  is a member of  $S_{p,h}(0,1)$ . Let  $n = 1/h$ . So, consider the functions  $\{\widehat{\varphi}_{p,h}^{(j)}\}_{j=-n}^{n-1}$ , forming a basis for  $\widehat{S}_{p,h}(-1,1)$ . Here we consider a different indexing with  $j = i - n$ . Defining

$$
s_j(x) := \hat{\varphi}_{p,h}^{(j)}(x) + \hat{\varphi}_{p,h}^{(j)}(-x) = \hat{\varphi}_{p,h}^{(j)}(x) + \hat{\varphi}_{p,h}^{(-j-p-1)}(x),
$$

for  $j = -n, \ldots, n-1$ , we obtain symmetric functions in  $\widehat{S}_{p,h}(-1, 1)$ . Using the relation

$$
\widehat{\varphi}_{p,h}^{(j)}|_{(0,1)} = \sum_{k \in \mathbb{Z}} \varphi_{p,h}^{(j+2nk)},
$$

we obtain that the restriction of  $s_j$  to  $(0, 1)$  fulfills

$$
s_j|_{(0,1)} = \sum_{k \in \mathbb{Z}} \varphi_{p,h}^{(j+2nk)} + \sum_{k \in \mathbb{Z}} \varphi_{p,h}^{(-j-p-1+2nk)} = \varphi_{p,h}^{(j)} + \sum_{k \in \mathbb{Z}} \varphi_{p,h}^{(-j-p-1+2nk)},
$$

which is

$$
s_j|_{(0,1)} = \varphi_{p,h}^{(j)} + \varphi_{p,h}^{(-j-p-1)}
$$
 for  $j \in \{-n, \ldots, -1\}$ ,  
\n
$$
s_j|_{(0,1)} = \varphi_{p,h}^{(j)}
$$
 for  $j \in \{0, \ldots, n-p-1\}$ , or  
\n
$$
s_j|_{(0,1)} = \varphi_{p,h}^{(j)} + \varphi_{p,h}^{(-j-p-1+2n)}
$$
 for  $j \in \{n-p, \ldots, n-1\}$ .

In all three cases  $s_j$  equals  $\widetilde{\varphi}_{p,h}^{(j)}$  or  $2\widetilde{\varphi}_{p,h}^{(j)}$ . This shows that  $\widetilde{\varphi}_{p,h}^{(i)} \in \widetilde{S}_{p,h}(0,1)$ .<br>It is easy to see that the functions in [\(47\)](#page-29-1) are linear independent for  $i =$  $-\left[\frac{p}{2}\right], \ldots, n-\left[\frac{p}{2}\right] - 1$ . So, it remains to show that every function  $u_{p,h}$  $S_{p,h}(0,1)$  can be expressed as a linear combination of the functions in [\(47\)](#page-29-1). As we have already noticed, by construction the function  $u_{p,h}$  can be extended to  $(-1, 1)$ , by defining  $w_{p,h}(x) := u_{p,h}(|x|)$ . Note that  $w_{p,h} \in \widehat{S}_{p,h}(-1, 1)$ . So, we can express it as a linear combination of basis functions of the basis given in  $(6)$  via

$$
w_{p,h} = \sum_{j=-n}^{n-1} w_j \widehat{\varphi}_{p,h}^{(j)}.
$$

By construction,  $w_{p,h}(x) = w_{p,h}(-x)$ , so we obtain

$$
w_{p,h}(x) = \frac{1}{2} (w_{p,h}(x) + w_{p,h}(-x)) = \frac{1}{2} \sum_{j=-n}^{n-1} w_j (\hat{\varphi}_{p,h}^{(j)}(x) + \hat{\varphi}_{p,h}^{(j)}(-x))
$$
  
\n
$$
= \frac{1}{2} \sum_{j=-n}^{n-1} w_j (\hat{\varphi}_{p,h}^{(j)}(x) + \hat{\varphi}_{p,h}^{(-j-p-1)}(x))
$$
  
\n
$$
= \frac{1}{2} \sum_{j=-n}^{n-1} \sum_{k \in \mathbb{Z}} w_j (\varphi_{p,h}^{(j+2nk)}(x) + \varphi_{p,h}^{(-j-p-1+2nk)}(x))
$$
  
\n
$$
= \frac{1}{2} \sum_{j=-n}^{n-1} w_j (\varphi_{p,h}^{(-j-p-1)}(x) + \varphi_{p,h}^{(j)}(x) + \varphi_{p,h}^{(2n-j-p-1)}(x)).
$$

Again, it can be checked easily, that for all  $j, n \in \mathbb{Z}$  the term

$$
\varphi_{p,h}^{(-j-p-1)}(x)+\varphi_{p,h}^{(j)}(x)+\varphi_{p,h}^{(2n-j-p-1)}(x)
$$

is in the span of  $\{\widetilde{\varphi}_{p,h}^{(i)}\}_{i=-\lceil \frac{p}{2} \rceil,\dots,n-\lfloor \frac{p}{2} \rfloor-1}$ , which concludes the proof.  $\Box$ 

We observe that the basis forms a partition of unity. Moreover, all basis functions are obviously non-negative linear combinations of B-splines. Hence we call it a B-spline-like basis.

Fig. [1](#page-32-0) and [2](#page-32-1) depict the B-spline basis functions that span  $\tilde{S}_{p,h}(0,1)$ . Here, the basis functions that have an influence at the boundary are plotted with solid lines. The basis functions that have zero derivatives up to order  $p - 1$ at the boundary coincide with standard B-spline functions. They are plotted with dashed lines.

If we compare the pictures of the B-spline basis functions in  $\widetilde{S}_{p,h}(0,1)$ (Fig. [1](#page-32-0) and [2\)](#page-32-1) with the standard B-spline basis functions for  $S_{p,h}(0,1)$  (Fig. [3](#page-32-2)) and [4\)](#page-32-3) obtained from a classical open knot vector, we see that the latter ones have more basis functions that are associated with the boundary. This can also be seen by counting the number of degrees of freedom, cf. Table [1.](#page-22-4)



Fig. 1 B-spline-like basis functions for  $\widetilde S_{1,h}(0,1)$  and  $\widetilde S_{2,h}(0,1)$ 

<span id="page-32-0"></span>

Fig. 2 B-spline-like basis functions for  $\widetilde S_{3,h}(0,1)$  and  $\widetilde S_{4,h}(0,1)$ 

<span id="page-32-1"></span>

Fig. 3 B-spline basis functions for  $S_{1,h}(0,1)$  and  $S_{2,h}(0,1)$ 

<span id="page-32-2"></span>

<span id="page-32-3"></span>Fig. 4 B-spline basis functions for  $S_{3,h}(0,1)$  and  $S_{4,h}(0,1)$ 

# References

- <span id="page-33-3"></span>1. Yuri Bazilevs, Lourenço Beirão da Veiga, J Austin Cottrell, Thomas JR Hughes, and Giancarlo Sangalli, Isogeometric analysis: approximation, stability and error estimates for h-refined meshes, Mathematical Models and Methods in Applied Sciences 16 (2006), no. 07, 1031–1090.
- <span id="page-33-4"></span>2. Lourenço Beirão da Veiga, Annalisa Buffa, Judith Rivas, and Giancarlo Sangalli, Some estimates for  $h-p-k$ -refinement in isogeometric analysis, Numerische Mathematik 118 (2011), no. 2, 271–305.
- <span id="page-33-15"></span>3. Lourenço Beirão da Veiga, Durkbin Cho, and Giancarlo Sangalli, Anisotropic nurbs approximation in isogeometric analysis, Computer Methods in Applied Mechanics and Engineering 209 (2012), 1–11.
- <span id="page-33-13"></span>4. Dietrich Braess, Finite elements: Theory, fast solvers, and applications in solid mechanics (second editin), Cambridge University Press, 2001.
- <span id="page-33-9"></span>5. Achi Brandt, Multi-level adaptive solutions to boundary-value problems, Math. Comp. 31 (1977), 333 – 390.
- <span id="page-33-11"></span>6. Charles K Chui, An introduction to wavelets, wavelet analysis and its applications, Academic Press, Boston (1992).
- <span id="page-33-2"></span>7. Ronald A DeVore and George G Lorentz, Constructive approximation, Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen, Springer, 1993.
- <span id="page-33-6"></span>8. Marco Donatelli, Carlo Garoni, Carla Manni, Stefano Serra-Capizzano, and Hendrik Speleers, Robust and optimal multi-iterative techniques for iga galerkin linear systems, Computer Methods in Applied Mechanics and Engineering 284 (2015), no. 0, 230 – 264, Isogeometric Analysis Special Issue.
- <span id="page-33-8"></span>9. John A Evans, Yuri Bazilevs, Ivo Babuška, and Thomas JR Hughes, n-widths, supinfs, and optimality ratios for the k-version of the isogeometric finite element method, Computer Methods in Applied Mechanics and Engineering 198 (2009), no. 21, 1726– 1741.
- <span id="page-33-5"></span>10. Carlo Garoni, Carla Manni, Francesca Pelosi, Stefano Serra-Capizzano, and Hendrik Speleers, On the spectrum of stiffness matrices arising from isogeometric analysis, Numerische Mathematik 127 (2014), no. 4, 751–799 (English).
- <span id="page-33-0"></span>11. Thomas JR Hughes, J Austin Cottrell, and Yuri Bazilevs, Isogeometric analysis: CAD, finite elements, NURBS, exact geometry and mesh refinement, Computer Methods in Applied Mechanics and Engineering 194 (2005), no. 39-41, 4135 – 4195.
- <span id="page-33-7"></span>12. Nicolai Korneichuk, Exact constants in approximation theory, vol. 38, Cambridge University Press, 1991.
- <span id="page-33-1"></span>13. Larry L Schumaker, Spline functions: basic theory, Cambridge University Press, 1981.
- <span id="page-33-14"></span>14. Christoph Schwab, p- and hp- finite element methods: Theory and applications in solid and fluid mechanics, Numerical Mathematics and Scientific Computation, Clarendon Press, 1998.
- <span id="page-33-10"></span>15. Ulrich Trottenberg, Cornelius Oosterlee, and Anton Schüller, *Multigrid*, Academic Press, London, 2001.
- <span id="page-33-12"></span>16. Ren-Hong Wang, Yan Xu, and Zhi-Qiang Xu, Eulerian numbers: a spline perspective, Journal of Mathematical Analysis and Applications 370 (2010), no. 2, 486–490.