Further results on differentially 4-uniform permutations over $\mathbb{F}_{2^{2m}}$

Zhengbang Zha^{1,2}, Lei Hu^{2,3}, Siwei Sun², Jinyong Shan²

¹School of Mathematical Sciences, Luoyang Normal University, Luoyang 471022, China

²State Key Laboratory of Information Security, Institute of Information Engineering,

Chinese Academy of Sciences, Beijing 100093, China

³Beijing Center for Mathematics and Information Interdisciplinary Sciences,

Beijing 100048, China

E-mail: zhazhengbang@163.com; {hu,swsun,jyshan12}@is.ac.cn

Abstract In this paper, we present several new constructions of differentially 4-uniform permutations over $\mathbb{F}_{2^{2m}}$ by modifying the values of the inverse function on some subsets of $\mathbb{F}_{2^{2m}}$. The resulted differentially 4-uniform permutations have high nonlinearities and algebraic degrees, which provide more choices for the design of crytographic substitution boxes.

Keywords Permutation; Differentially 4-uniform function; Nonlinearity; Algebraic degree

Mathematics Subject Classification (2000) 94A60; 11T71; 14G50

1 Introduction

Many block ciphers use substitution boxes (S-boxes) to bring the confusion into the cipher, and hence the design of S-boxes plays an important role in the design of cryptographic systems. To achieve a correct inverse decryption and for ease of the software implementation, S-boxes are usually designed as permutations over a characteristic 2 finite field of even extension degree, namely $\mathbb{F}_{2^{2m}}$. To resist against linear cryptanalysis, differential cryptanalysis and other cryptanalysis like algebraic attacks, one would like the S-boxes having high nonlinearity, low differential uniformity and high algebraic degree simultaneously. The inverse function x^{2^n-2} over \mathbb{F}_{2^n} is such a function, which is used to construct the S-box of the Advanced Encryption Standard (AES) with n = 8.

The differential uniformity [15] of $f(x) \in \mathbb{F}_{2^n}[x]$ is defined by

$$\Delta_f = \max\{N(a,b)|a, b \in \mathbb{F}_{2^n}, a \neq 0\},\$$

where N(a, b) denotes the number of solutions $x \in \mathbb{F}_{2^n}$ of the equation f(x+a) + f(x) = b. It is well known that $\Delta_f = 2$ is the minimum possible value of Δ_f . Differentially 2-uniform functions are called almost perfect nonlinear (APN) functions which provide the optimal resistance to differential attacks. A recent progress on APN functions can be found in [6, 9] and the references therein. Up to now, there is only one known APN permutation, which is defined on \mathbb{F}_{2^6} [2], and the existence of more ones on $\mathbb{F}_{2^{2m}}$ remains open. Therefore, finding differentially 4-uniform permutations with good cryptographic properties is an interesting and active research topic for the goal of providing more choices for S-boxes.

Recently, there has been significant progress in finding functions with low differential uniformity [5, 8, 10, 13, 16, 17, 20, 24]. In [3, 4] Bracken *et al.* firstly studied highly nonlinear monomials and binomials which are differentially 4-uniform permutations over $\mathbb{F}_{2^{2m}}$. After that, many new differentially 4-uniform permutations over $\mathbb{F}_{2^{2m}}$ were constructed by composing the inverse function and permutations over $\mathbb{F}_{2^{2m}}$. Qu *et al.* [18, 19] applied a powerful switching method (which can be seen in [9] and was initially proposed by Dillon in his talk in the ninth international conference on finite fields and their applications) to construct differentially 4-uniform permutations and found that the number of CCZ-inequivalent differentially 4-uniform permutations over $\mathbb{F}_{2^{2m}}$ grows exponentially when m increases. Later, some new differentially 4-uniform permutations which are CCZinequivalent to the inverse function were obtained by composing the inverse function and cycles on $\mathbb{F}_{2^{2m}}$ [14, 22]. Zha *et al.* [23] presented two new families of differentially 4uniform permutations by modifying the values of the inverse function on some subfields of $\mathbb{F}_{2^{2m}}$. In [21] Tang *et al.* gave a construction providing a large number of differentially 4-uniform bijections with maximum algebraic degrees and high nonlinearities. In this paper, we will revisit differentially 4-uniform permutations in [23] which are constructed by modifying the inverse function on some subsets of $\mathbb{F}_{2^{2m}}$, and show some new differentially 4-uniform permutations which have high nonlinearities and algebraic degrees.

The rest of this paper is organized as follows. In Section 2, some preliminaries needed later as well as a brief overview of known differentially 4-uniform permutations are presented. We present some new constructions of differentially 4-uniform permutations in Section 3, and show their cryptographic properties in Section 4. Finally, we conclude the paper in Section 5.

2 Preliminaries

We define the trace map from \mathbb{F}_{2^n} onto its subfield \mathbb{F}_{2^k} (with k|n) as

$$\operatorname{Tr}_{k}^{n}(x) = x + x^{2^{k}} + x^{2^{2k}} + \dots + x^{2^{n-k}}$$

and denote the absolute trace map from \mathbb{F}_{2^n} onto the binary subfield \mathbb{F}_2 by $\operatorname{Tr}(x) = \sum_{i=0}^{n-1} x^{2^i}$.

The algebraic degree of $f(x) = \sum_{i=0}^{2^n-1} a_i x^i \in \mathbb{F}_{2^n}[x]$ is denoted by deg f, which equals to the maximal 2-weight of the exponent i with $a_i \neq 0$, where the 2-weight of an integer is the number of ones in its binary expression. It is known that deg f is upper bounded by n-1 if f is a permutation on \mathbb{F}_{2^n} . If deg $f \leq 1$, then f is called an affine function.

For a function $f : \mathbb{F}_{2^n} \to \mathbb{F}_{2^n}$ and any $(a, b) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n}^*$, the Walsh transform of f is defined as

$$f^{\mathcal{W}}(a,b) := \sum_{x \in \mathbb{F}_{2^n}} (-1)^{\operatorname{Tr}(ax+bf(x))}$$

and the Walsh spectrum of f is $\mathcal{W}_f := \{f^{\mathcal{W}}(a,b) | a \in \mathbb{F}_{2^n}, b \in \mathbb{F}_{2^n}^*\}$. The nonlinearity $\mathcal{NL}(f)$ of f is defined as

$$\mathcal{NL}(f) \stackrel{\triangle}{=} 2^{n-1} - \frac{1}{2} \max_{w \in W_f} |w|.$$

For odd n, the nonlinearity $\mathcal{NL}(f)$ is upper bounded by $2^{n-1} - 2^{\frac{n-1}{2}}$; and for even n it is conjectured that $\mathcal{NL}(f)$ is upper bounded by $2^{n-1} - 2^{\frac{n}{2}}$ [4]. We call a function maximal nonlinear if its nonlinearity attains these bounds.

Two functions $f, g : \mathbb{F}_{2^n} \to \mathbb{F}_{2^n}$ are called extended affine equivalent (EA-equivalent) if $g = A_1 \circ f \circ A_2 + A$ for some affine permutations A_1 and A_2 and an affine function A. Nonconstant EA-equivalent functions have the same algebraic degree.

Two functions f and g from \mathbb{F}_{2^n} to itself are called Carlet-Charpin-Zinoviev equivalent (CCZ-equivalent) if the graphs of f and g are affine equivalent. It is shown in [7] that EA-equivalence implies CCZ-equivalence, but not vice versa. Every permutation is CCZ-equivalent to its inverse. Differential spectrum and Walsh spectrum are CCZ-invariants while algebraic degree is not a CCZ-invariant [1, 7].

Many new classes of differentially 4-uniform permutations are introduced by using the definition of CCZ-equivalence and CCZ-invariants. One may refer to [4, 14, 18, 21] for recent progress on this topic.

Below we always denote the inverse function by x^{-1} with the convention that $0^{-1} = 0$. Let n, k and q be positive integers with n > 1, k|n and $q = 2^k$. Let ω be a primitive third root of unity in the algebraic closure of \mathbb{F}_q . Clearly, $\omega \in \mathbb{F}_q$ if k is even and $\omega \notin \mathbb{F}_q$ otherwise.

3 Some constructions of differentially 4-uniform permutations

In this section, we revisit a class of differentially 4-uniform permutations of the form $x^{-1} + t(x^q + x)^{2^n-1} + t$ in [23]. We can not only unify some previous constructions, but also present new differentially 4-uniform permutations of this form.

Let S be a subset of \mathbb{F}_{2^n} satisfying: (a) either both of 0 and 1 or neither of them belongs to S and; (b) $\frac{x}{1+x} \in S$ holds for any $x \in S \setminus \{0, 1\}$. Let $\delta_S(x)$ be a characteristic function of S, i.e., $\delta_S(x) = 1$ if $x \in S$ and $\delta_S(x) = 0$ otherwise. According to the Lagrange interpolation, we have $\delta_S(x) = 1 + \prod_{a \in S} (x + \theta)^{2^n - 1}$.

Inspired by the ideas of [19] and [21], we discuss the cryptographic properties of the function

$$f(x) = x^{-1} + \delta_S(x) \tag{1}$$

over \mathbb{F}_{2^n} in the sequel.

Proposition 1. The function f defined by (1) is a permutation over \mathbb{F}_{2^n} and its compositional inverse is $g(x) = (x + \delta_{S'}(x))^{-1}$, where S' is a subset of \mathbb{F}_{2^n} satisfying that $x^{-1} + 1 \in S'$ holds if and only if $x \in S$.

Proof. Assume $x, y \in \mathbb{F}_{2^n}$ with f(x) = f(y). Then we have $x^{-1} + \delta_S(x) = y^{-1} + \delta_S(y)$. If $\delta_S(x) = \delta_S(y)$, we get $x^{-1} = y^{-1}$, which leads to x = y. If $\delta_S(x) \neq \delta_S(y)$, we get $x^{-1} + 1 = y^{-1}$, which, excluding the cases of (x, y) = (0, 1) or (x, y) = (1, 0), implies $y = \frac{x}{1+x}$ and $\delta_S(x) \neq \delta_S(\frac{x}{1+x})$. It is a contradiction with the hypothesis on S. Then we deduce that f is a permutation over \mathbb{F}_{2^n} . We can easily check that g(f(x)) = f(g(x)) = x, which completes the proof.

Proposition 2. Let f be defined as in (1). Then f is a differentially 4-uniform permutation over \mathbb{F}_{2^n} if and only if for any $a \in \mathbb{F}_{2^n}$ with $a \neq 0, 1$, one of the following two statements holds:

1) For $\delta_S(a) = \delta_S(0)$, the cases of $\delta_S(\omega a) = \delta_S(\omega^2 a)$ and of $x^2 + ax + \frac{a^2}{1+a} = 0$ with $\delta_S(x+a) \neq \delta_S(x)$ cannot occur simultaneously; and

2) For $\delta_S(a) \neq \delta_S(0)$, the cases of $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$ and of $x^2 + ax + \frac{a^2}{1+a} = 0$ with $\delta_S(x+a) = \delta_S(x)$ cannot occur simultaneously.

Proof. Let $a, b \in \mathbb{F}_{2^n}$ with $a \neq 0$. We consider the solutions of the equation f(x+a) + f(x) = b, i.e.,

$$(x+a)^{-1} + \delta_S(x+a) + x^{-1} + \delta_S(x) = b$$
(2)

over \mathbb{F}_{2^n} . By Proposition 1 we get that f is a permutation over \mathbb{F}_{2^n} , which implies that Eq. (2) has no solution if b = 0. Below we assume $b \neq 0$ and x is a solution of (2). When x = 0 or a, we get

$$b = a^{-1} + \delta_S(a) + \delta_S(0) (:= b_0)$$

from (2). For other possible solutions namely ones with $x \neq 0$ and $x \neq a$, we divide into the following two disjoint cases to discuss.

Case I: If $\delta_S(x+a) = \delta_S(x)$, then we have $(x+a)^{-1} + x^{-1} = b$ from (2), which leads to $x^2 + ax + \frac{a}{b} = 0$.

Case II: If $\delta_S(x+a) \neq \delta_S(x)$, then we have $(x+a)^{-1} + x^{-1} = b+1$ from (2). If b = 1, we have no solution of (2). Otherwise, we obtain $x^2 + ax + \frac{a}{b+1} = 0$ from (2).

For any pair (a, b) with $b \neq b_0$, there are at most four solutions of (2) in Cases I and II. In the sequel, we consider the case of $b = b_0 \neq 0$.

If $\delta_S(a) = \delta_S(0)$, then we have $b_0 = a^{-1}$. In this case, we get two solutions ωa and $\omega^2 a$ in Case I, which need satisfying $\delta_S(\omega a) = \delta_S(\omega^2 a)$. Similarly in Case II, Eq. (2) turns to $x^2 + ax + \frac{a^2}{1+a} = 0$.

If $\delta_S(a) \neq \delta_S(0)$, then we have $b_0 = a^{-1} + 1$. In this case, we get two solutions ωa and $\omega^2 a$ in Case II, which need satisfying $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$. Similarly in Case I, Eq. (2) becomes $x^2 + ax + \frac{a^2}{1+a} = 0$.

For constructing differentially 4-uniform permutations, we want $\Delta_f \leq 4$ when $a \neq 0, 1$ and $b = b_0$. More precisely, we need to find an appropriate set S such that there are at most two solutions in Cases I and II when $a \neq 0, 1$ and $b = b_0$, which equivalent to the two statements in Proposition 2.

From the definition of S, we always get $\delta_S(0) = \delta_S(1)$ and $\delta_S(\omega) = \delta_S(\omega^2)$. Then we can easily check that there are exactly four solutions $0, 1, \omega, \omega^2$ of (2) when $a = b_0 = 1$, which implies that $\Delta_f = 4$. The desired conclusion then follows.

We note that the function $f(x) = x^{-1} + \delta_s(x)$ is a special case of the switching method studied in [19]. In Theorem 5.3 of [19], the composite inverse of G is of the form $G^{-1} = 1/(x + g(x)) = 1/(x + 1)$ if g(x) = 1; and it is 1/x if g(x) = 0. If S = supp(g) is taken, then it immediately lead to the function f. Our main contributions are characterizing new subsets S and obtaining new constructions of differentially 4-uniform permutations.

According to Proposition 2, we will give some constructions in the sequel by choosing different sets S.

3.1 The relevance with known constructions

In this subsection, we will give results related to some known constructions. First we list a lemma needed below.

Lemma 1. [11] For any $a, b \in \mathbb{F}_{2^n}$ with $a \neq 0$, the polynomial $f(x) = x^2 + ax + b$ is irreducible over \mathbb{F}_{2^n} if and only if $\operatorname{Tr}(b/a^2) = 1$.

From Proposition 2 and Lemma 1, we can obtain the following construction. The proof is trivial and we omit it here.

Theorem 1. Let $S = \mathbb{F}_{2^k}$. Then f is a differentially 4-uniform permutation over \mathbb{F}_{2^n} if k is even or k = 1, 3 and $\frac{n}{2}$ is odd.

The result of Theorem 1 includes the constructions of differentially 4-uniform permutations in Theorems 1 and 3 of [23] in the case of t = 1.

We need the following lemma which can be derived from Lemma 2 in [21]. We present here its proof for completeness.

Lemma 2. For any $a \in \mathbb{F}_{2^n}/\{0,1\}$, define the polynomial $\mu(x) := x^2 + ax + \frac{a^2}{1+a} \in \mathbb{F}_{2^n}[x]$. If $\mu(x) = 0$ has two solutions λ and ν in \mathbb{F}_{2^n} , then we have $\operatorname{Tr}(\frac{1}{\lambda+1}) = \operatorname{Tr}(\frac{1}{\nu+1}) = 0$.

Proof. If $\mu(x) = 0$ has two solutions λ, ν , from Lemma 1 we have $\text{Tr}(\frac{1}{1+a}) = 0$ and $\lambda, \nu \neq 0, 1$ since $a \neq 0, 1$. Since

$$\frac{1}{\lambda+1} \cdot \frac{1}{\nu+1} = \frac{1}{\lambda\nu+\lambda+\nu+1} = \frac{1}{a^2(1+a)^{-1}+a+1} = 1+a$$

and $\frac{1}{\lambda+1} + \frac{1}{\nu+1} = \frac{\lambda+\nu}{\lambda\nu+\lambda+\nu+1} = a + a^2$, we get that $\frac{1}{\lambda+1}$ and $\frac{1}{\nu+1}$ are the roots of equation $x^2 + (a + a^2)x + 1 + a = 0$. Note that $\operatorname{Tr}(\frac{1}{\lambda+1}) = \operatorname{Tr}(\frac{1}{\nu+1}) = 0$ if and only if there exist two values $u, v \in \mathbb{F}_{2^n}$ such that $\frac{1}{\lambda+1} = u + u^2$ and $\frac{1}{\nu+1} = v + v^2$. Assume u and v are the roots of equation $x^2 + sx + p = 0$, then $u + u^2$ and $v + v^2$ must be the roots of equation $x^2 + (s + s^2)x + p(1 + s + p) = 0$.

We choose s = a. By Lemma 1 and $\operatorname{Tr}(\frac{1}{1+a}) = 0$, there exists an element $p \in \mathbb{F}_{2^n}$ such that $p^2 + (1+a)p + (1+a) = 0$ or namely, p(1+s+p) = 1+s. Then we have $\frac{p^2}{s^2} + \frac{p}{s} + \frac{p}{s^2} + \frac{1}{s} + \frac{1}{s^2} = 0$ and $\operatorname{Tr}(\frac{p}{s^2}) = 0$, which implies that equation $x^2 + sx + p = 0$ actually has two roots in \mathbb{F}_{2^n} .

Theorem 2. Let S be a subset of \mathbb{F}_{2^n} satisfying $\frac{x}{1+x} \in S$ and $\operatorname{Tr}(x) = 1$ for any $x \in S$. Then f is a differentially 4-uniform permutation over \mathbb{F}_{2^n} .

Proof. Since $\operatorname{Tr}(x) = 1$ for any $x \in S$, we have $0 \notin S$ and $\operatorname{Tr}(\frac{1}{x+1}) = 1$ for any $x \in S$. If $\delta_S(a) = \delta_S(0)$, we can deduce that the solutions of $\mu(x) = 0$ satisfying $\delta_S(x+a) = \delta_S(x) = 0$ from Lemma 2. If $\delta_S(a) \neq \delta_S(0)$, then we get $a \in S$ and $\operatorname{Tr}(\frac{1}{a+1}) = 1$, which implies that there are no solutions of $\mu(x) = 0$ from Lemma 1. We complete the proof by Proposition 2.

We note that the compositional inverse of the permutation f defined in Theorem 2 is exactly the known differentially 4-uniform bijection presented in Construction 1 of [21].

3.2 New constructions from unions of two subfields of \mathbb{F}_{2^n}

In what follows, we introduce two new constructions by combining two suitable subfields of \mathbb{F}_{2^n} . To do this, we need the following lemma.

Lemma 3. Let k_1 and k_2 be divisors of n and $S = \mathbb{F}_{2^{k_1}} \cup \mathbb{F}_{2^{k_2}}$. Then $\delta_S(\frac{y^2}{1+y}) \neq \delta_S(x^2+xy)$ for any $x, y \in S$ with $x + y \notin S$.

Proof. Assume $x, y \in S$ with $x+y \notin S$, without loss of generality, we assume $x \in \mathbb{F}_{2^{k_1}} \setminus \mathbb{F}_{2^{k_2}}$ and $y \in \mathbb{F}_{2^{k_2}} \setminus \mathbb{F}_{2^{k_1}}$. Obviously, we have $\frac{y^2}{1+y} \in \mathbb{F}_{2^{k_2}}$ and $\delta_S(\frac{y^2}{1+y}) = 1$. If $\delta_S(x^2 + xy) = 1$, then we obtain $x^2 + xy \in \mathbb{F}_{2^{k_2}}$, which implies

$$x^2 + xy = x^{2^{1+k_2}} + x^{2^{k_2}}y.$$

Raising the above equation by 2^{k_1} -th powers, we have

$$x^{2} + xy^{2^{k_{1}}} = x^{2^{1+k_{2}}} + x^{2^{k_{2}}}y^{2^{k_{1}}}$$

Then we get

$$(x+x^{2^{k_2}})(y+y^{2^{k_1}}) = 0$$

by adding the above two equations. It leads to $x \in \mathbb{F}_{2^{k_2}}$ or $y \in \mathbb{F}_{2^{k_1}}$, which is a contradiction.

Utilizing Proposition 2 and Lemma 3, we have the following theorems.

Theorem 3. Let k_1 and k_2 be even divisors of n and $S = \mathbb{F}_{2^{k_1}} \cup \mathbb{F}_{2^{k_2}}$. Then f is a differentially 4-uniform permutation over \mathbb{F}_{2^n} .

Proof. If $\delta_S(a) = \delta_S(0)$, then we get $a \in S$. By Lemma 3, we have that $\delta_S(\frac{y^2}{1+y}) \neq \delta_S(x^2 + xy)$ for any $x, y \in S$ with $x + y \notin S$, which implies that the equation $\mu(x) = 0$ has no solution.

If $\delta_S(a) \neq \delta_S(0)$, then we have $a \notin S$. We assume $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$ is true and $\omega a \in S$ and $\omega^2 a \notin S$ without loss of generality. Since $S = \mathbb{F}_{2^{k_1}} \cup \mathbb{F}_{2^{k_2}}$ and k_1 and k_2 are even integers, we obtain $\omega^2 a \in S$ from $\omega a \in S$ and $\omega \in S$, which is a contradiction. The proof is completed.

Theorem 4. Let k_1 be an even divisor of n with $gcd(3, k_1) = 1$. Assume 6|n, n/6 is odd and $S = \mathbb{F}_{2^3} \cup \mathbb{F}_{2^{k_1}}$. Then f is a differentially 4-uniform permutation over \mathbb{F}_{2^n} .

Proof. If $\delta_S(a) = \delta_S(0)$, similarly to the proof of Theorem 3, we can show that the equation $\mu(x) = 0$ has no solution.

If $\delta_S(a) \neq \delta_S(0)$, $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$ holds only if $\omega a \in \mathbb{F}_{2^3}$ and $\omega^2 a \notin S$ or $\omega^2 a \in \mathbb{F}_{2^3}$ and $\omega a \notin S$. Without loss of generality, we consider the case of $\omega a \in \mathbb{F}_{2^3}$ and $\omega^2 a \notin S$. Then we have $a^8 = \omega^2 a$, $a^{64} = a$ and

$$\begin{aligned} \operatorname{Tr}(\frac{1}{1+a}) &= \operatorname{Tr}_1^3(\operatorname{Tr}_3^6(\operatorname{Tr}_6^n(\frac{1}{1+a}))) = \operatorname{Tr}_1^3(\operatorname{Tr}_3^6(\frac{1}{1+a})) \\ &= \operatorname{Tr}_1^3(\frac{1}{1+\omega a + (\omega a)^{-1}}) = 1 \end{aligned}$$

since $\operatorname{Tr}_1^3(\frac{1}{1+x+x^{-1}}) = 1$ for any $x \in \mathbb{F}_{2^3}$, which implies that $\mu(x) = 0$ has no solution while $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$. This completes the proof.

3.3 A construction from unions of subsets of \mathbb{F}_{2^n}

In this subsection, we introduce a new construction by combining some subsets of \mathbb{F}_{2^n} .

Theorem 5. Let k be even and $\frac{n}{k}$ be odd and let l be a divisor of k. Let S_1 be a subset of \mathbb{F}_{2^n} satisfying $\frac{x}{1+x} \in S_1$ and $\operatorname{Tr}(x) = 1$ for any $x \in S_1$. Assume $S = S_1 \cup (\mathbb{F}_q \setminus \mathbb{F}_{2^l})$ or $S = \mathbb{F}_q \setminus \mathbb{F}_{2^l}$, then f is a differentially 4-uniform permutation over \mathbb{F}_{2^n} if

- 1) l is even;
- 2) l = 1 and $k \equiv 2 \pmod{4}$; or
- 3) l = 3 and k = 6.

Proof. If $\delta_S(a) = \delta_S(0)$, we get $a \notin S$. From Lemma 2, if $\mu(x) = 0$ has two solutions λ, ν , then we have $\nu = \lambda + a$ and $\lambda, \nu \notin S_1$. Suppose $\mu(x) = 0$ with $\delta_S(\lambda) \neq \delta_S(\nu)$ holds. Without loss of generality, we assume $\delta_S(\nu) = 0$ and $\delta_S(\lambda) = 1$, which implies that $\lambda + a \notin S$ and $\lambda \in S$. We get $\lambda \in \mathbb{F}_q \setminus \mathbb{F}_{2^l}$ since $\lambda \notin S_1$. Then we obtain

$$\lambda^2 + a\lambda + \frac{a^2}{1+a} = 0 \tag{3}$$

and

$$\lambda^2 + a^q \lambda + \frac{a^{2q}}{1+a^q} = 0. \tag{4}$$

Since $a \notin S$, we have that $a \notin \mathbb{F}_q$ or $a \in \mathbb{F}_{2^l}$. If $a \notin \mathbb{F}_q$, we can deduce that $\lambda = 1 + \frac{1}{(1+a)^{q+1}}$, which implies $a^{q^2} = a$. Since n/k is odd and $a^{q^{n/k}} = a$, we obtain $a^q = a$, which is a contradiction. If $a \in \mathbb{F}_{2^l}$, then we get $\lambda + a \in \mathbb{F}_q \setminus \mathbb{F}_{2^l}$, which contradicts our first assumption $\lambda + a \notin S$. Hence, $\mu(x) = 0$ with $\delta_S(x+a) \neq \delta_S(x)$ cannot holds.

If $\delta_S(a) \neq \delta_S(0)$, then we have $a \in S$. When $a \in S_1$, then we have $\operatorname{Tr}(\frac{1}{1+a}) = 1$, which implies that there are no solutions of $\mu(x) = 0$. When $a \in \mathbb{F}_q \setminus \mathbb{F}_{2^l}$, we assume that $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$ is true.

If l is even, then we have $\omega a, \omega^2 a \in \mathbb{F}_q \setminus \mathbb{F}_{2^l}$ for $\omega^4 = \omega$. It implies that $\delta_S(\omega a) = \delta_S(\omega^2 a) = 1$, which is a contradiction.

If l = 1 and $k \equiv 2 \pmod{4}$, $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$ can be true only if $a = \omega$ or $a = \omega^2$. For $a = \omega, \omega^2$, we obtain that $\operatorname{Tr}(\frac{1}{1+a}) = \operatorname{Tr}_1^2(\frac{1}{1+a}) = 1$, which implies that $\mu(x) = 0$ has no solutions in this case.

If k = 6 and l = 3, $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$ holds only if $a^8 = \omega a$ or $a^8 = \omega^2 a$. We consider the case of $a^8 = \omega^2 a$ for example. We have $\operatorname{Tr}(\frac{1}{1+a}) = \operatorname{Tr}_1^3(\operatorname{Tr}_6^6(\operatorname{Tr}_6^n(\frac{1}{1+a}))) = 1$, which also deduce that there are no solutions of $\mu(x) = 0$.

Thus, the cases of $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$ and of $\mu(x) = 0$ cannot occur simultaneously if $\delta_S(a) \neq \delta_S(0)$. The proof is finished by Proposition 2.

Remark 1. In Theorem 5, if $\operatorname{Tr}(x) = 1$ for any $x \in \mathbb{F}_q \setminus \mathbb{F}_{2^l}$, then f is equal to one differential 4-uniform permutation defined in Theorem 2, which is CCZ-equivalent to the one of Theorem 1 in [21]. Theorem 5 exhibits some specific examples of Theorem 5.3 in [19].

3.4 Two constructions from inverse sets of affine subspaces of \mathbb{F}_{2^n}

Let $t_1 \in \mathbb{F}_{2^n}^*$ with $\operatorname{Tr}_k^n(t_1) = 0$. We consider the subset $S = \{x \in \mathbb{F}_{2^n} : x^{-q} = x^{-1} + t_1\}$, which is the set of the inverses of the elements in the non-empty affine subspace $\{x \in \mathbb{F}_{2^n} : x^q = x + t_1\}$. Obviously we have $0 \notin S$ and $\delta_S(0) = 0$.

Lemma 4. For $a \in S$, the cases of $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$ and of $\mu(x) = 0$ cannot occur simultaneously if

- 1) $q \equiv 1 \pmod{3};$
- 2) $q \equiv 2 \pmod{3}$ and $t_1 \notin \mathbb{F}_q$; or
- 3) $\frac{n}{2k}$ is odd, $t_1 \in \mathbb{F}_q$ and q = 2 or 8.

Proof. Assume $\delta_S(\omega a) \neq \delta_S(\omega^2 a)$, without loss of generality, we consider the case of $\omega a \in S$ and $\omega^2 a \notin S$. Since $S = \{x \in \mathbb{F}_{2^n} : x^{-q} = x^{-1} + t_1\}$, we get $a^{-q} = a^{-1} + \omega t_1$ when $q \equiv 1 \pmod{3}$ and $a^{-q} = \omega a^{-1} + \omega^2 t_1$ when $q \equiv 2 \pmod{3}$.

Since $a \in S$, we get $a^{-q} = a^{-1} + t_1$. When $q \equiv 1 \pmod{3}$, we have $t_1 = 0$, which is a contradiction. When $q \equiv 2 \pmod{3}$, we have $\omega^2 a^{-1} + t_1 = 0$ and $a^{-1} = \omega t_1$, which implies $t_1^q = t_1$. Therefore, we achieve the goal when $q \equiv 2 \pmod{3}$ and $t_1^q \neq t_1$. When $\frac{n}{2k}$ is odd and q = 2 or 8, we need to show that there are no solutions of $\mu(x) = 0$. From Lemma 1, a direct proof is to show that $\operatorname{Tr}_1^n(\frac{1}{1+a}) = 1$. We get

$$\begin{aligned} \operatorname{Tr}_{1}^{n}(\frac{1}{1+a}) &= \operatorname{Tr}_{1}^{n}(\frac{1}{1+(\omega t_{1})^{-1}}) = \operatorname{Tr}_{1}^{k}(\operatorname{Tr}_{k}^{2k}(\operatorname{Tr}_{2k}^{n}(\frac{1}{1+\omega t_{1}}))) \\ &= \operatorname{Tr}_{1}^{k}(\operatorname{Tr}_{k}^{2k}(\frac{1}{1+\omega t_{1}})) = \operatorname{Tr}_{1}^{k}(\frac{1}{1+t_{1}+t_{1}^{-1}}) = 1, \end{aligned}$$

which completes the proof.

Lemma 5. If $a \notin S$, there are no solutions of $\mu(x) = 0$ with $\delta_S(x+a) \neq \delta_S(x)$ if the equation

$$\begin{array}{l} (t_1^4 + t_1^2)a^{4q} + (t_1^2 + t_1)a^{4q-1} + (t_1^2 + t_1)a^{4q-2} + a^{4q-3} + a^{4q-4} + (t_1^2 + t_1)a^{3q} \\ + t_1^2a^{3q-1} + (1+t_1)a^{3q-2} + a^{3q-3} + (t_1^2 + t_1)a^{2q} + (1+t_1)a^{2q-1} + a^q + a^{q-1} \\ = 1 \end{array}$$
(5)

has no solution in $\mathbb{F}_{2^n} \setminus \mathbb{F}_q$.

Proof. We assume that $\mu(x) = 0$ has two solutions x, x + a satisfying $\delta_S(x) \neq \delta_S(x + a)$. Without loss of generality, we consider the case of $x \in S$ and $x + a \notin S$. As $x \neq 0, x \neq a$ and $x^2 + ax + \frac{a^2}{1+a} = 0$, we have $a \neq 1$ and

$$x^{-2} + \frac{1+a}{a}x^{-1} + \frac{1+a}{a^2} = 0.$$
 (6)

From $x \in S$ we get $x^{-q} = x^{-1} + t_1$. Substituting it to the q-th power of (6), we obtain

$$x^{-2} + \frac{1+a^{q}}{a^{q}}x^{-1} + t_{1}^{2} + \frac{1+a^{q}}{a^{q}}t_{1} + \frac{1+a^{q}}{a^{2q}} = 0.$$
 (7)

If $a \in \mathbb{F}_q$, we get $t_1^2 + \frac{1+a}{a}t_1 = 0$ from (6) and (7), which leads to $t_1 = \frac{1+a}{a}$. Since $(x+a)^2 + a(x+a) + \frac{a^2}{1+a} = 0$, we derive $(x+a)^{-2} + \frac{1+a}{a}(x+a)^{-1} + \frac{1+a}{a^2} = 0$ and

 $(x+a)^{-2q} + \frac{1+a}{a}(x+a)^{-q} + \frac{1+a}{a^2} = 0$ for $a \in \mathbb{F}_q$. Combining the above two equations, we may draw the conclusion that $(x+a)^{-1} \in \mathbb{F}_q$ or $x+a \in S$. Since $x+a \notin S$, we get $(x+a)^{-1} \in \mathbb{F}_q$ and $x \in \mathbb{F}_q$, which contradicts the first assumption $x \in S$. Therefore, we obtain $a \notin \mathbb{F}_q$ and

$$x^{-1} = \frac{(t_1^2 + t_1)a^{2q} + a^{2q-1} + a^{2q-2} + (1+t_1)a^q + 1}{a^{2q-1} + a^q}$$

from (6) and (7). Substituting it into (6), we can derive Eq. (5) by some trivial computation. The proof is finished. \Box

We also need the following lemma.

Lemma 6. [11] An irreducible polynomial over \mathbb{F}_q of degree n remains irreducible over \mathbb{F}_{q^l} if and only if gcd(l,n) = 1.

Now we can get many differentially 4-uniform permutations by computing the solutions of (5). We list two simple examples in the following theorems.

Theorem 6. If q = 2, $\frac{n}{2}$ is odd and $t_1 = 1$, then $S = \{\omega, \omega^2\}$ and f is a differentially 4-uniform permutation over \mathbb{F}_{2^n} .

Proof. The case of $a \in S$ is proved by Lemma 4. Now we consider the case of $a \notin S$. Since q = 2, $\frac{n}{2}$ is odd and $t_1 = 1$, Eq. (5) becomes $a^4 + a^3 + a^2 + a + 1 = 0$, which has no solution on \mathbb{F}_{2^n} . The proof is completed by Lemma 5 and Proposition 2.

Theorem 7. Let gcd(n,5) = 1 and $\frac{n}{4}$ be odd. If q = 4 and $t_1 = 1$, then $S = \{x \in \mathbb{F}_{2^n} : x^{-4} = x^{-1} + 1\}$ and f is a differentially 4-uniform permutation over \mathbb{F}_{2^n} .

Proof. Similarly to the proof of Theorem 6, we need to show that Eq. (5) has no solution in $\mathbb{F}_{2^n} \setminus \mathbb{F}_q$. Since q = 4 and $t_1 = 1$, Eq. (5) becomes $g(a) := a^{13} + a^{12} + a^{11} + a^9 + a^4 + a^3 + 1 = 0$. We remark that g(a) = 0 has no solution in \mathbb{F}_{2^4} and

$$g(a) = a^{13} + a^{12} + a^{11} + a^9 + a^4 + a^3 + 1$$

= $(a^5 + a^3 + 1)(a^4 + \omega a^3 + \omega a^2 + \omega^2 a + \omega^2)(a^4 + \omega^2 a^3 + \omega^2 a^2 + \omega a + \omega)$

has only irreducible factors with degrees 2 and 5 over \mathbb{F}_{2^4} . Since gcd(n,5) = 1 and $\frac{n}{4}$ is odd, then we get that g(a) = 0 has no solution on \mathbb{F}_{2^n} by Lemma 6. We complete the proof.

4 Cryptographic properties of functions constructed

In this section, we focus on the cryptographic properties of the function f defined by (1). It is shown in [19] that all differentially 4-uniform permutations on $\mathbb{F}_{2^{2m}}$ have algebraic degree n-1. Since f lies in a more general framework in Theorem 5.3 of [19], we have that f has the maximum possible algebraic degree n-1.

In the following, we give some lower bounds on the nonlinearities of f, present some numerical results about the differential spectra and nonlinearities of f, and discuss the CCZ-inequivalence between f and some known differential 4-uniform permutations.

4.1 Nonlinearity

Lemma 7. [12] Define the Kloosterman sum over \mathbb{F}_{2^n} as

$$K_n(\lambda) = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{\operatorname{Tr}(\lambda x + x^{-1})}, \quad \lambda \in \mathbb{F}_{2^n}.$$

The set $\{K_n(\lambda) : \lambda \in \mathbb{F}_{2^n}\}$ is exactly the set of all integers $t \equiv 0 \pmod{4}$ in the range $\left[-2^{\frac{n}{2}+1}+1, 2^{\frac{n}{2}+1}+1\right]$.

Let $\lfloor 2^{\frac{1}{2}+1} \rfloor = 2$ and $\lfloor 2^{\frac{k}{2}+1} \rfloor = t$ for k > 1, $t \equiv 0 \pmod{4}$ and $2^{\frac{k}{2}+1} - 4 < t \le 2^{\frac{k}{2}+1}$. We have the following result by Lemma 7.

Lemma 8. Let $a, b \in \mathbb{F}_{2^n}$ with $\operatorname{Tr}(b) = 1$. Let k be a positive integer with $k \mid n$. Then $|\sum_{x \in \mathbb{F}_{2^k}} (-1)^{\operatorname{Tr}(ax+bx^{-1})}| \leq \lfloor 2^{\frac{k}{2}+1} \rfloor$.

Proof. Since $\operatorname{Tr}(b) = 1$, for any $x \in \mathbb{F}_{2^k}$ we get $\operatorname{Tr}_k^n(b) \neq 0$ and

$$\operatorname{Tr}(ax + bx^{-1}) = \operatorname{Tr}_1^k(\operatorname{Tr}_k^n(ax + bx^{-1})) = \operatorname{Tr}_1^k(x\operatorname{Tr}_k^n(a) + x^{-1}\operatorname{Tr}_k^n(b)).$$

By Lemma 7, we have that

$$|\sum_{x \in \mathbb{F}_{2^k}} (-1)^{\operatorname{Tr}(ax+bx^{-1})}| = |\sum_{x \in \mathbb{F}_{2^k}} (-1)^{\operatorname{Tr}_1^k(x\operatorname{Tr}_k^n(a)+x^{-1}\operatorname{Tr}_k^n(b))}| \le \lfloor 2^{\frac{k}{2}+1} \rfloor.$$

Lemma 9. [21] For any even $n \ge 6$, we have $\mathcal{NL}(f) \ge 2^{n-1} - 2^{n/2} - |S|$.

From Lemma 9, we get a direct lower bound on the nonlinearities of f defined in Theorems 3-7. For example, we obtain $\mathcal{NL}(f) \geq 2^{n-1} - 2^{n/2} - 2$ for the function defined in Theorem 6 and $\mathcal{NL}(f) \geq 2^{n-1} - 2^{n/2} - 4$ for the function defined in Theorem 7. In the following, we give a more tightly bound on the nonlinearities of f defined in Theorems 3 and 4.

Proposition 3. If $S = \mathbb{F}_{2^{k_1}} \cup \mathbb{F}_{2^{k_2}}$, the nonlinearity of the function f defined by (1) satisfies

$$\mathcal{NL}(f) \ge 2^{n-1} - \lfloor 2^{\frac{n}{2}} \rfloor - \lfloor 2^{\frac{k_1}{2}+1} \rfloor - \lfloor 2^{\frac{k_2}{2}+1} \rfloor - \lfloor 2^{\frac{\gcd(k_1,k_2)}{2}+1} \rfloor.$$

Especially, we have

$$\mathcal{NL}(f) \ge 2^{n-1} - \lfloor 2^{\frac{n}{2}} \rfloor - \lfloor 2^{\frac{k_2}{2}+1} \rfloor$$

if $k_1 \mid k_2$ and

$$\mathcal{NL}(f) \ge 2^{n-1} - \lfloor 2^{\frac{n}{2}} \rfloor - \lfloor 2^{\frac{k_2}{2}+1} \rfloor - 6$$

if $k_1 = 3$ and $gcd(k_2, 3) = 1$.

Proof. As we know, the nonlinearity of f is defined by $\mathcal{NL}(f) = 2^{n-1} - \frac{1}{2} \max f^{\mathcal{W}}(a, b)$, where $f^{\mathcal{W}}(a, b) = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{\operatorname{Tr}(ax+bf(x))}$ for $a, b \in \mathbb{F}_{2^n}$ and $b \neq 0$. We have that

$$f^{\mathcal{W}}(a,b) = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{\operatorname{Tr}(ax+b(x^{-1}+\delta_S(x)))} \\ = \sum_{x \in \mathbb{F}_{2^n} \setminus S} (-1)^{\operatorname{Tr}(ax+bx^{-1})} + \sum_{x \in S} (-1)^{\operatorname{Tr}(ax+bx^{-1}+b)}$$

If $\operatorname{Tr}(b) = 0$, we get $|f^{\mathcal{W}}(a, b)| = |\sum_{x \in \mathbb{F}_{2^n}} (-1)^{\operatorname{Tr}(ax+bx^{-1})}| \le \lfloor 2^{\frac{n}{2}+1} \rfloor$ by Lemma 7. If $\operatorname{Tr}(b) = 1$, we obtain

$$|f^{\mathcal{W}}(a,b)| = |\sum_{x \in \mathbb{F}_{2^n}} (-1)^{\operatorname{Tr}(ax+bx^{-1})} - 2\sum_{x \in S} (-1)^{\operatorname{Tr}(ax+bx^{-1})}| \\ \leq |\sum_{x \in \mathbb{F}_{2^n}} (-1)^{\operatorname{Tr}(ax+bx^{-1})}| + 2|\sum_{x \in S} (-1)^{\operatorname{Tr}(ax+bx^{-1})}|.$$

Notice that

$$\sum_{x \in S} (-1)^{\operatorname{Tr}(ax+bx^{-1})} = \sum_{\substack{x \in \mathbb{F}_{2^{k_1}}}} (-1)^{\operatorname{Tr}(ax+bx^{-1})} + \sum_{\substack{x \in \mathbb{F}_{2^{k_2}}}} (-1)^{\operatorname{Tr}(ax+bx^{-1})} - \sum_{\substack{x \in \mathbb{F}_{2^{\gcd(k_1,k_2)}}} (-1)^{\operatorname{Tr}(ax+bx^{-1})},$$

then we obtain

$$|f^{\mathcal{W}}(a,b)| \le \lfloor 2^{\frac{n}{2}+1} \rfloor + 2(\lfloor 2^{\frac{k_1}{2}+1} \rfloor + \lfloor 2^{\frac{k_2}{2}+1} \rfloor + \lfloor 2^{\frac{\gcd(k_1,k_2)}{2}+1} \rfloor)$$

from Lemma 8, which implies that

$$\mathcal{NL}(f) \ge 2^{n-1} - \lfloor 2^{\frac{n}{2}} \rfloor - \lfloor 2^{\frac{k_1}{2}+1} \rfloor - \lfloor 2^{\frac{k_2}{2}+1} \rfloor - \lfloor 2^{\frac{\gcd(k_1,k_2)}{2}+1} \rfloor.$$

If $k_1 \mid k_2$, we can get $S = \mathbb{F}_{2^{k_2}}$ and $\mathcal{NL}(f) \geq 2^{n-1} - \lfloor 2^{\frac{n}{2}} \rfloor - \lfloor 2^{\frac{k_2}{2}+1} \rfloor$ by Lemma 8 directly. Furthermore, if $k_1 = 3$ and k_2 is even, we have

$$\mathcal{NL}(f) \ge 2^{n-1} - \lfloor 2^{\frac{n}{2}} \rfloor - \lfloor 2^{\frac{k_2}{2}+1} \rfloor - 6$$

since $|\sum_{x \in \mathbb{F}_{2^3}} (-1)^{\operatorname{Tr}(ax+bx^{-1})}| \le 4$ and $|\sum_{x \in \mathbb{F}_2} (-1)^{\operatorname{Tr}(ax+bx^{-1})}| \le 2$.

4.2 Numerical results and CCZ-inequivalence

For even $n \leq 12$, we computed the nonlinearities and differential spectra of the functions in Section 3 by using the MAGMA software system. Some of the computational results are listed in Tables 1-3, where the notation $\mathcal{D}(f)$ represents the differential spectrum of a function f, $\mathcal{B}(f)$ represents a bound on the nonlinearity of f, and the multiset M = $\{a_1^{m_1}, a_2^{m_2}, \dots, a_t^{m_t}\}$ means the elements a_i appears m_i times in M for $1 \leq i \leq t$.

By Tables 1-3, we conclude that Theorems 3-7 present several differentially 4-uniform permutations CCZ-inequivalent to the ones defined in Theorem 1. Especially, compared to the numerical results in Tables III and IV of [19], Tables 2 and 3 list some new results on nonlinearities and differential spectra when n = 10 and 12.

Table 1: Nonlinearities and differential spectra of functions in Section 3 over \mathbb{F}_{2^6}

S	f	$\mathcal{NL}(f)$	$\mathcal{D}(f)$	$\mathcal{B}(f)$
\mathbb{F}_2	Theorem 1	24	$\{0^{2079}, 2^{1890}, 4^{63}\}$	22
\mathbb{F}_{2^2}	Theorem 1	22	$\{0^{2127}, 2^{1794}, 4^{111}\}\$	20
\mathbb{F}_{2^3}	Theorem 1	22	$\{0^{2199}, 2^{1650}, 4^{183}\}$	20
$\mathbb{F}_{2^2} \cup \mathbb{F}_{2^3}$	Theorem 4	20	$\{0^{2247}, 2^{1554}, 4^{231}\}$	14
$\mathbb{F}_{2^2} \setminus \mathbb{F}_2$	Theorems 5, 6	22	$\{0^{2127}, 2^{1794}, 4^{111}\}$	22

Table 2: Nonlinearities and differential spectra of functions in Section 3 over $\mathbb{F}_{2^{10}}$

S	f	$\mathcal{NL}(f)$	$\mathcal{D}(f)$	$\mathcal{B}(f)$
\mathbb{F}_2	Theorem 1	480	$\{0^{524799}, 2^{521730}, 4^{1023}\}$	478
\mathbb{F}_{2^2}	Theorem 1	478	$\{0^{34335}, 2^{29250}, 4^{1695}\}$	476
$\mathbb{F}_{2^2} \setminus \mathbb{F}_2$	Theorems 5, 6	478	$\{0^{525879}, 2^{519570}, 4^{2103}\}$	478

Table 3: Nonlinearities and differential spectra of functions in Section 3 over $\mathbb{F}_{2^{12}}$

S	f	$\mathcal{NL}(f)$	$\mathcal{D}(f)$	$\mathcal{B}(f)$
\mathbb{F}_{2^2}	Theorem 1	1982	$\{0^{8394735}, 2^{8370210}, 4^{8175}\}$	1980
\mathbb{F}_{2^4}	Theorem 1	1978	$\{0^{8419263}, 2^{8321154}, 4^{32703}\}$	1968
\mathbb{F}_{2^6}	Theorem 1	1970	$\{0^{8511615}, 2^{8136450}, 4^{125055}\}$	1920
$\mathbb{F}_{2^4} \cup \mathbb{F}_{2^6}$	Theorem 3	1966	$\{0^{8534127}, 2^{8091426}, 4^{147567}\}$	1908
$\mathbb{F}_{2^4} \setminus \mathbb{F}_{2^2}$	Theorem 5	1978	$\{0^{8415183}, 2^{8329314}, 4^{28623}\}$	1972
$x^{-4} + x^{-1} = 1$	Theorem 7	1980	$\{0^{8399055}, 2^{8361570}, 4^{12495}\}$	1980

For n = 12, by a Magma computation, we checked that there are 1036 elements of $\mathbb{F}_{2^{12}}$ satisfying $\operatorname{Tr}(x) = \operatorname{Tr}(\frac{x}{1+x}) = 1$. That is to say, there are $2^{518} - 1$ different sets S_1 and 2^{518} different sets S in Theorem 5. For a random choice set S_1 , the functions constructed in Theorem 5 ($S = S_1 \cup (\mathbb{F}_{16} \setminus \mathbb{F}_4)$) and Theorem 2 ($S = S_1$) often have different differential spectra or nonlinearities, which implies that they are CCZ-inequivalent. We randomly choose 10000 different sets S_1 and compute the nonlinearities of the differentially 4-uniform permutations defined in Theorems 5 and 2 separately. The computational results are listed in Table 4. The notations $\operatorname{Ave}(\mathcal{NL}(f))$, $\operatorname{Max}(\mathcal{NL}(f))$ and $\operatorname{Min}(\mathcal{NL}(f))$ denote the average nonlinearity, maximal nonlinearity and minimal nonlinearity of f respectively.

Table 4: Variance of nonlinearities of 10000 samples f on $\mathbb{F}_{2^{12}}$

S	f	$\operatorname{Ave}(\mathcal{NL}(f))$	$\operatorname{Max}(\mathcal{NL}(f))$	$\operatorname{Min}(\mathcal{NL}(f))$
S_1	Theorem 2	1911.106	1982	1864
$S_1 \cup (\mathbb{F}_{16} \setminus \mathbb{F}_4)$	Theorem 5	1910.264	1978	1866

From Table 4, we found that the average nonlinearity of the functions f in Theorem

5 is less than that of the functions f in Theorem 2. This is due to

$$|S_1 \cup (\mathbb{F}_{16} \setminus \mathbb{F}_4)| \ge |S_1| + 4$$

since there are exactly four solutions of $\operatorname{Tr}(x) = 0$ with $x \in (\mathbb{F}_{16} \setminus \mathbb{F}_4)$ over $\mathbb{F}_{2^{12}}$. It seems that the nonlinearity of f gets smaller when |S| gets larger, this, however, is not always true, below we give some examples.

Example 1. For n = 12, let α be a primitive element of $\mathbb{F}_{2^{12}}$ defined by $\alpha^{12} + \alpha^7 + \alpha^4 + \alpha^3 + 1 = 0$ and $S_1 = \{\alpha^i | i \in \Pi\}$, where

 $\Pi = \{649, 3411, 2016, 2422, 437, 903, 3963, 2464, 1914, 1180, 3755, 2410, 119, 647, 3624, 841, 2833, 2709, 2352, 4092, 3812, 2696, 3166, 2950, 1784, 3475, 233, 1831, 1157, 1422, 3897, 2429, 918, 1363, 2910, 3955, 1372, 1785, 3013, 1589, 2021, 1721\}.$ *We can easily check that* $|S_1| = 42$. *The nonlinearity of the function f in Theorem 5 i*

We can easily check that $|S_1| = 42$. The nonlinearity of the function f in Theorem 5 is 1958, while the function in Theorem 2 is of nonlinearity 1956.

Example 2. For n = 12, let α be a primitive element of $\mathbb{F}_{2^{12}}$ defined in Example 1 and $S_1 = \{\alpha^i | i \in \{3351, 1475, 777, 661, 921, 977, 4076, 2037, 3359, 2414, 3616, 3033, 3401, 3697, 3459, 654, 3160, 123, 3226, 2837, 526, 2832, 1182, 4094, 3964, 3887, 1705, 2489, 1766, 4066, 589, 184, 1842, 2752\}\}$. We can easily check that $|S_1| = 34$ and the nonlinearities of f defined in Theorems 5 and 2 are both 1962.

Example 3. For n = 12, let α be a primitive element of $\mathbb{F}_{2^{12}}$ defined in Example 1 and

 $S_1 = \{ \alpha^i \mid i \in \{273, 546, 1092, 2184\} \}.$

We can check that S_1 is a subset of $\mathbb{F}_{16} \setminus \mathbb{F}_4$ and $S = \mathbb{F}_{16} \setminus \mathbb{F}_4$. The nonlinearity of the function f in Theorem 5 is 1978, while the function in Theorem 2 is of nonlinearity 1980.

5 Conclusion

In this paper, we refine a general technique for constructing a class of differentially 4uniform permutations by modifying the values of the inverse function on a subset S of $\mathbb{F}_{2^{2m}}$. By using this technique, we get many differentially 4-uniform permutations with high nonlinearities and algebraic degrees. Our numerical results support that some of them are new and have the nonlinearity approaching the maximum while the size of S is not too small.

Acknowledgement: The authors would like to thank the anonymous reviewers for their valuable comments and suggestions which improved both the quality and presentation of this paper. The work of this paper was supported by the National Basic Research Programme under Grant 2013CB834203, the National Natural Science Foundation of China (Grants 11201214 and 61472417), and the Strategic Priority Research Program of Chinese Academy of Sciences under Grant XDA06010702.

References

[1] Budaghyan L, Carlet C, Pott A. New classes of almost bent and almost perfect nonlinear polynomials. IEEE Trans Inform Theory, 2006, 52: 1141-1152

- [2] Browning K.A., Dillon J.F., McQuistan M.T., et al. An APN permutation in dimension six. In: Contemporary Mathematics, Vol. 518 (Post Proceedings of the Ninth International Conference on Finite Fields and Their Applications-Fq'9), J Am Math Soc, 2010, 33-42
- [3] Bracken C, Leander G. A highly nonlinear differentially 4 uniform power mapping that permutes fields of even degree. Finite Fields Appl, 2010, 16: 231-242
- [4] Bracken C, Tan C, Tan Y. Binomial differentially 4 uniform permutations with high nonlinearity. Finite Fields Appl, 2012, 18: 537-546
- [5] Carlet C. On known and new differentially uniform functions. In: Lecture Notes in Computer Sciences, vol. 6812. Heidelberg: Springer, 2011, 1-15
- [6] Carlet C. More constructions of APN and differentially 4-uniform functions by concatenation. Science China Mathematics, 2013, 56: 1373-1384
- [7] Carlet C, Charpin P, Zinoviev V. Codes, bent functions and permutations suitable for DES-like cryptosystems. Des Codes Cryptogr, 1998, 15: 125-156
- [8] Carlet C, Tang D, Tang X, et al. New construction of differentially 4-uniform bijections. In: Lecture Notes in Computer Sciences, vol. 1106. Heidelberg: Springer, 2013, 1-17
- [9] Edel Y, Pott A. A new almost perfect nonlinear function which is not quadratic. Advances in Mathematical Communications, 2009, 3: 59-81
- [10] Jia W, Zeng X, Li C, et al. Permutation polynomials with low differential uniformity over finite fields of odd characteristic. Science China Mathematics, 2013, 56: 1429-1440
- [11] Lidl R, Niederreiter H. Finite Fields. Encyclopedia of Mathematics and its Applications, vol. 20. Cambridge University Press, 1997
- [12] Lachaud G, Wolfmann J. The weights of the orthogonals of the extended quadratic binary Goppa codes. IEEE Trans Inform Theory, 1990, 36: 686-692
- [13] Li Y, Wang M. Constructing differentially 4-uniform permutations over $\mathbb{F}_{2^{2m}}$ from quadratic APN permutations over $\mathbb{F}_{2^{2m+1}}$. Des Codes Cryptography, 2014, 72: 249-264
- [14] Li Y, Wang M, Yu Y. Constructing differentially 4-uniform permutations over $GF(2^{2k})$ from the inverse function revisited. http://eprint.iacr.org/2013/731
- [15] Nyberg K. Differentially uniform mappings for cryptography. In: Lecture Notes in Computer Science, vol.765. New York: Springer, 1994, 134-144
- [16] Pott A, Zhou Y. Switching constructions of planar functions on finite fields. In: Lecture Notes in Computer Science, vol. 6087. Heidelberg: Springer, 2010, 135-150
- [17] Qu L, Li C, Dai Q, et al. On the differential uniformities of functions over finite fields. Science China Mathematics, 2013, 56: 1477-1484

- [18] Qu L, Tan Y, Li C, et al. More constructions of differentially 4-uniform permutations on $\mathbb{F}_{2^{2k}}$. Des Codes Cryptography, http://dx.doi.org/10.1007/s10623-014-0006-x, in press
- [19] Qu L, Tan Y, Tan C, et al. Constructing differentially 4-uniform permutations over $\mathbb{F}_{2^{2k}}$ via the switching method. IEEE Trans Inform Theory, 2013, 59: 4675-4686
- [20] Qu L, Xiong H, Li C. A negative answer to Bracken-Tan-Tan's problem on differentially 4-uniform permutations over \mathbb{F}_{2^n} . Finite Fields Appl, 2013, 24: 55-65
- [21] Tang D, Carlet C, Tang X. Differentially 4-uniform bijections by permuting the inverse functions. Des Codes Cryptography, http://dx.doi.org/10.1007/s10623-014-9992-y, in press
- [22] Yu Y, Wang M, Li Y. Constructing differential 4-uniform permutations from know ones. Chinese Journal of Electronics, 2013, 22: 495-499
- [23] Zha Z, Hu L, Sun S. Constructing new differential 4-uniform permutations from the inverse function. Finite Fields Appl, 2014, 25: 64-78
- [24] Zha Z, Wang X. Power functions with low uniformity on odd characteristic finite fields. Science China Mathematics, 2010, 53: 1931-1940