RITA GIULIANO AND MICHEL WEBER

ABSTRACT. We study the local limit theorem for weighted sums of Bernoulli variables. We show on examples that this is an important question in the general theory of the local limit theorem, and which turns up to be not well explored. The examples we consider arise from standard random models used in arithmetical number theory. We next use the characteristic function method to prove new local limit theorems for weighted sums of Bernoulli variables. Further, we give an application of the almost sure local limit theorem to a representation problem in additive number theory due to Burr, using an appropriate random model. We also give a simple example showing that the local limit theorem, in its standard form, fails to be sharp enough for estimating the probability $\mathbb{P}\{S_n \in E\}$ for infinite sets of integers E, already in the simple case where S_n is a sum of n independent standard Bernoulli random variables and E an arithmetic progression.

1. Introduction.

This work is devoted to the study of the local limit theorem and of its recent developments, in the context of some standard random models used in arithmetical number theory. It is also somehow completing the recent paper [8]. We will be mainly interested in studying the local limit theorem for weighted sums of Bernoulli variables. As it will be clarified soon, this turns up to be a fundamental question in the local limit theorem theory. We first recall some basic results and the used methods. The local limit theorem was established already three centuries ago in the binomial case by De Moivre and Laplace around 1730. Based on Stirling approximation formula of n!, it is a very precise result for moderate deviations.

Lemma 1.1. Let 0 , <math>q = 1 - p. Let X be such that $\mathbb{P}\{X = 1\} = p = 1 - \mathbb{P}\{X = 0\}$. Let X_1, X_2, \ldots be independent copies of X and let $S_n = X_1 + \ldots + X_n$. Let $0 < \gamma < 1$ and let $\beta \leq \gamma \sqrt{pq} n^{1/3}$. Then for all k such that letting $x = \frac{k - np}{\sqrt{npq}}$, $|x| \leq \beta n^{1/6}$, we have

$$\mathbb{P}\{S_n = k\} = \frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi n p q}} e^E,$$

with $|E| \leq \frac{|x|^3}{\sqrt{npq}} + \frac{|x|^4}{npq} + \frac{|x|^3}{2(npq)^{\frac{3}{2}}} + \frac{1}{4n\min(p,q)(1-\gamma)}.$

This slightly more precise formulation than the one given in Chow and Teicher [4], p. 46, is easily extrapolated from their proof. More generally, let $\widetilde{X} = \{X_n, n \ge 1\}$ be a sequence of independent, square integrable random variables taking values in a common lattice $\mathcal{L}(v_0, D) = \{v_0 + Dk, k \in \mathbb{Z}\}$, where v_0 and D > 0 are real numbers. Let also $M_n = \sum_{j=1}^n \mathbb{E} X_j$, $\Sigma_n = \sum_{j=1}^n \operatorname{Var}(X_j)$. We say that \widetilde{X} satisfies a local limit theorem if

(1.1)
$$\Delta_n := \sup_{N=v_0 n + Dk} \left| \sqrt{\Sigma_n} \mathbb{P}\{S_n = N\} - \frac{D}{\sqrt{2\pi}} e^{-\frac{(N-M_n)^2}{2\Sigma_n}} \right| = o(1).$$

This fine limit theorem has connections with Number Theory, see for instance Postnikov [20]. If \widetilde{X} is an i.i.d. sequence, then (1.1) holds if and only if the "span" D is maximal $(D = \sup \{d > d\})$

[AS.LLT]1 August 22, 2021

²⁰¹⁰ Mathematics Subject Classification: Primary: 60F15, 60G50; Secondary: 60F05.

Key words and phrases: independent random variables, lattice distributed, Bernoulli part, local limit theorem, almost sure local limit theorem, effective remainder, random walk in random scenery.

 $0; \exists a \in \mathbb{Z} : \mathbb{P}\{X \in a + d\mathbb{Z}\} = 1\}$). This is Gnedenko's well-known result, which is also optimal (Matskyavichyus [17]). Under stronger integrability conditions, the remainder term can be improved (see [14] Theorem 4.5.3), [19] Theorem 6 p.197). The general form of the local limit theorem ([14], Th. 4.2.1) for i.i.d. random variables, states

Theorem 1.2. In order that for some choice of constants a_n and b_n

$$\lim_{n \to \infty} \sup_{N \in \mathcal{L}(v_0 n, D)} \left| \frac{b_n}{\lambda} \mathbb{P}\{S_n = N\} - g\left(\frac{N - a_n}{b_n}\right) \right| = 0,$$

where g is the density of some stable distribution G with exponent $0 < \alpha \leq 2$, it is necessary and sufficient that

(i)
$$\frac{S_n - a_n}{b_n} \stackrel{\mathcal{D}}{\Rightarrow} G \quad as \ n \to \infty$$
 (ii) D is maximal

There are essentially two approaches used: the method of characteristic functions and the Bernoulli part extraction method. In the later case, this method is called the extraction method of the Bernoulli part of a random variable and was developed by McDonald [16], for proving local limit theorems in presence of the central limit theorem. Kolmogorov [15] (see also Kolmogorov's interesting comment p. 29) initiated twenty years before a similar approach in the study of Lévy's concentration function. We also mention Arratia, Barbour and Tavaré [1, 2] probabilistic approach in the study of the asymptotic behaviour of logarithmic combinatorial structures, and the recent work of Röllin and Ross [21] based on Landau-Kolmogorov inequalities.

An important problem inside the general study of the local limit theorem concerns the case when the considered sums are weighted sums of Bernoulli variables, the "simple" case when the weights are increasing covering already non-trivial examples of random models used in number theory. The purpose of the next Section is to underline this in providing a few examples of such models, which we believe, are challenging problems for probabilists.

Additionally, for weighted sums of independent or i.i.d. random variables, the Bernoulli part extraction method reduces the problem to the case of weighted sums of Bernoulli variables, thereby making this case crucial too for the application of this method.

The goal of this work is to investigate the local limit theorem for weighted sums of Bernoulli variables. In Section 3, we use the characteristic function method to prove new local limit theorems. Next in Section 4, we give an application of the almost sure local limit theorem to Burr's representation problem in additive number theory, using an appropriate random model. Finally, we also give an example showing that the standard form (1.1) of the local limit theorem, fails to be sharp enough for estimating the probability $\mathbb{P}\{S_n \in E\}$ for infinite sets of integers E; and this already in the simple case where S_n is a sum of n independent standard Bernoulli random variables and E an arithmetic progression.

2. Some Random Models in Number Theory.

2.1. A Probabilistic Model for the Dickman Function. This function originates from the study by Dickman of the asymptotic distribution of the largest prime factor $P^+(n)$ of a natural integer n. He has shown that the limit

(2.1)
$$\lim_{n \to \infty} \frac{1}{n} \# \{k; 1 \le k \le n : P^+(k) \le n^{1/u} \} = \rho(u)$$

exists, and $\rho(u)$, called the Dickman Function, is defined as the continuous solution of the differential-difference equation

$$u\rho'(u) + \rho(u-1) = 0, \qquad (u > 1)$$

with the initial condition $\rho(u) = 1$ for $0 \le u \le 1$. We have $\int_0^{\infty} \rho(v) dv = e^{\gamma}$, where γ is Euler's constant. This is a function of first importance in analytic number theory, which has been thoroughly investigated by Hensley, Hildebrand, Tenenbaum notably, see [22] for more details.

There is a probabilistic way of describing the Dickman Function. We refer to Hwang and Tsai [13]. Let $X = \{X_j, j \ge 1\}$ be a sequence of independent random variables such that

(2.2)
$$\begin{cases} \mathbb{P}\{X_j = j\} = j^{-1} \\ \mathbb{P}\{X_j = 0\} = 1 - j^{-1}. \end{cases} \quad (j \ge 1)$$

Proposition 2.1. Let $D_n = \sum_{j=1}^n X_j$. Then

$$\lim_{n \to \infty} \mathbb{P}\left\{n^{-1}D_n < x\right\} = e^{-\gamma} \int_0^x \rho(v) \mathrm{d}v \qquad (x > 0).$$

Arratia, Barbour and Tavaré [2], Corollary 2.8 proved a (restricted) local limit theorem for D_n

(2.3)
$$\lim_{n \to \infty} n \mathbb{P}\{D_n = k_n\} = e^{-\gamma} \rho(x), \quad \text{when } \lim_{n \to \infty} k_n/n = x > 0.$$

The almost sure local limit theorem was recently established in Giuliano, Szewczak and Weber in [9]. The proof is essentially based on a long and delicate study of the related correlations functions. A proof of the local limit theorem in the form (2.3) using only characteristic functions is also given, correcting the one indicated [13]. No local limit theorem for X (in the sense of (1.1)) is known.

Remarks 2.2. (i) Hensley [12] has shown that the limiting law is infinitely divisible. In the same paper, he also constructed another very interesting probabilistic model, adapted to the "psixiology" i.e. to functions Ψ, Φ linked to P^+, P^- .

(ii) Obviously D_n also reads as $D_n = \sum_{j=1}^n j\beta_j$ where $\{\beta_j, j = 1, \dots, n\}$ are independent Bernoulli random variables such that

(2.4)
$$\begin{cases} \mathbb{P}\{\beta_j = 1\} = j^{-1} \\ \mathbb{P}\{\beta_j = 0\} = 1 - j^{-1}. \end{cases} \quad (j = 1, \dots, n)$$

(iii) Let Z_1, \ldots, Z_n be independent Poisson distributed random variables with intensity $\mathbb{E} Z_j = 1/j$, and let $T_n = \sum_{j=1}^n j Z_j$. Then we have the exact formula $\mathbb{P}\{T_n = n\} = e^{-\sum_{j=1}^n 1/j}$, based on Cauchy formula for cycles of permutations ([1], formula (1.2)).

(iv) Vervaart has shown that independent Bernoulli random variables can be embedded into a Poisson process (see [23], Chapter 4).

2.2. A Diophantine Equation. Let $\mathcal{N} = \{\nu_0, \ldots, \nu_P\}$ be a finite set of integers. Consider the diophantine equation

(2.5)
$$x_1 + \ldots + x_n = y_1 + \ldots + y_n,$$

in which the unknown $x_i, y_j, 1 \leq i, j \leq n$, are subject to belong to \mathbb{N} . Let $N_n(\mathbb{N})$ denote the number of 2*n*-uples $(x_1, \ldots, x_n, y_1, \ldots, y_n) \in \mathbb{N}^{2n}$ which satisfy (2.5).

Examine the basic case $\mathbb{N} = \{0, \dots, P-1\}$ and note $N_n(P) = N_n(\mathbb{N})$. Recall the approach used in [20] §2.4. Let X be a random variable defined by

$$\mathbb{P}\{X = k\} = \begin{cases} \frac{P - |k|}{P^2} & \text{if } 0 \le |k| < P, \\ 0 & \text{if } |k| \ge P. \end{cases}$$

We easily verify that $\mathbb{E} X = 0$, $\sigma^2 = \mathbb{E} X^2 = \frac{P^2 - 1}{6}$ and $\mathbb{E} |X|^3 \leq CP^3$. Moreover, $\mathbb{E} e^{2i\pi tX} = (1/P)F_{P-1}(2\pi t)$ where F_m is the Fejér kernel,

$$F_m(u) = \frac{1}{m+1} \left(\frac{\sin\frac{m+1}{2}u}{\sin\frac{u}{2}}\right)^2.$$

Note that if u(k) is the number of solutions of the equation x-y=k, $0 \le x \le P-1$, $0 \le y \le P-1$, then u(k) = P - |k| if |k| < P, and u(k) = 0 if $|k| \ge P$. So that in turn $\mathbb{P}\{X = k\} = \frac{u(k)}{P^2}$, $k \in \mathbb{Z}$. Let X_1, \ldots, X_n be independent copies of X and note $S_n = X_1 + \ldots + X_n$. As $(x_1 - y_1) + \ldots + (x_n - y_n) = 0$ if and only if $x_1 - y_1 = k_1, \ldots, x_n - y_n = k_n$, for some integers k_j verifying $k_1 + \ldots + k_n = 0$, we have

$$\mathbb{P}\{S_n = 0\} = \sum_{\substack{k_1 + \dots + k_n = 0 \\ |k_n| \le P}} \mathbb{P}\{X_1 = k_1\} \dots \mathbb{P}\{X_n = k_n\} = \frac{N_n(P)}{P^{2n}}.$$

We have, as a direct consequence of the approximate local limit theorem with effective remainder given in [8], Corollary 1.8,

(2.6)
$$\frac{N_n(P)\sqrt{n}}{P^{2n-1}} = \sqrt{3/\pi} + \mathcal{O}\left(\frac{1}{P^2} + \frac{P}{\sqrt{n}}\right),$$

uniformly over n, P such that for $n \ge CP^2$.

Remark 2.3. (i) As $\mathbb{P}\{S_n = 0\} = \int_0^1 |\frac{\sin P \pi t}{P \sin \pi t}|^{2n} dt$, it is easy to bound from below $N_n(P)$ by CP^{2n-1}/\sqrt{n} and to get the upper bound $C_{\varepsilon}P^{2n-1+\varepsilon}/\sqrt{n}$, for any $\varepsilon > 0$, uniformly in P and n. See for instance [26], inequality (2.3).

In fact, one "can" take $\varepsilon = 0$.

Theorem 2.4 ([7], Th. 2.1). There exist absolute constants C', C'' such that for any positive integers P and n,

(2.7)
$$C' \frac{P^{2n-1}}{\sqrt{n}} \le N_n(P) \le C'' \frac{P^{2n-1}}{\sqrt{n}}.$$

The proof depends on finer bounds of the previous Fejér integrals, requiring more elaborated calculations.

Remark 2.5. We don't exactly know how the normalized ratios $\frac{N_n(P)}{P^{2n}}$ behave when n and P vary simultaneously; a question which is tightly related to the variation properties of powers of the Fejér kernels $\{F_{P_i}^n(u), j \ge 1\}$ for growing sequences $\{P_j, j \ge 1\}$.

2.3. Freiman-Pitman's Probabilistic Model of the Partition Function. This is probably the most informative example. Let $q_m(n)$, $m \leq n$, denote the number of partitions of n into distinct parts, each of which is at least m, namely the number of ways to express n as

(2.8)
$$n = i_1 + \ldots + i_r, \qquad m \le i_1 < \ldots < i_r \le n.$$

Let X_m, \ldots, X_n be independent random variables defined by

(2.9)
$$\mathbb{P}\{X_j = 0\} = \frac{1}{1 + e^{-\sigma j}}, \qquad \mathbb{P}\{X_j = j\} = \frac{e^{-\sigma j}}{1 + e^{-\sigma j}}.$$

The random variable $Y = X_m + \ldots + X_n$ can serve to modelize the partition function $q_m(n)$. There is a one-to-one correspondence between the number of partitions of n of the required type and the number of vectors (x_m, \ldots, x_n) with $x_j = 0$ or 1 such that $mx_m + \ldots + nx_n = n$. Notice that

$$e^{\sigma n} \int_{0}^{1} \prod_{\substack{j=m \\ m \leq j \leq n}}^{n} \left(1 + e^{-\sigma j} e^{2i\pi\alpha j}\right) e^{-2i\pi\alpha n} d\alpha$$

= $e^{\sigma n} \sum_{\substack{x_j \in \{0,1\} \\ m \leq j \leq n}} \int_{0}^{1} e^{-\sigma (mx_m + \dots + nx_n)} e^{-2i\pi (mx_m + \dots + nx_n - n)\alpha} d\alpha$
= $e^{\sigma n} \sum_{\substack{x_j \in \{0,1\} \\ m \leq j < n}}^{n} e^{-\sigma n} \chi\{mx_m + \dots + nx_n = n\} = q_m(n).$

Hence the formula (in which σ only appears in the right-hand side)

(2.10)
$$q_m(n) = e^{\sigma n} \int_0^1 \prod_{j=m}^n \left(1 + e^{-\sigma j} e^{2i\pi\alpha j}\right) e^{-2i\pi\alpha n} d\alpha$$

This also implies (letting $\varphi(t) = \mathbb{E} e^{2i\pi tY}$ be the characteristic function of Y)

$$\begin{aligned} q_m(n) &= e^{\sigma n} \Big(\prod_{j=m}^n (1+e^{-\sigma j}) \Big) \int_0^1 \prod_{j=m}^n \Big(\frac{1+e^{-\sigma j}e^{2i\pi\alpha j}}{1+e^{-\sigma j}} \Big) e^{-2i\pi\alpha n} \mathrm{d}\alpha \\ &= e^{\sigma n} \Big(\prod_{j=m}^n (1+e^{-\sigma j}) \Big) \int_0^1 \varphi(\alpha) e^{-2i\pi\alpha n} \mathrm{d}\alpha \ = \ e^{\sigma n} \Big(\prod_{j=m}^n (1+e^{-\sigma j}) \Big) \mathbb{P}\{Y=n\}. \end{aligned}$$

In [5] p. 387 and 389, the authors noticed that an appropriate local limit theorem would allow to write $\mathbb{P}\{Y=n\} \sim e^{-(\mathbb{E}Y-n)^2/(2\operatorname{Var}(Y))}/\sqrt{2\pi\operatorname{Var}(Y)}$. Choosing σ as being the unique solution of the equation $\mathbb{E}Y = \sum_{j=m}^{n} \frac{j}{1+e^{\sigma j}} = n$ would then give $\mathbb{P}\{Y=n\} \sim 1/B\sqrt{2\pi}$, and by reporting

$$q_m(n) \sim e^{\sigma n} \Big(\prod_{j=m}^n (1+e^{-\sigma j}) \Big) \frac{1}{B\sqrt{2\pi}}.$$

In place, Freiman and Pitman directly estimated the integral in (2.10) in a long delicate work [5]. **Remark 2.6.** By Euler's pentagonal theorem, $q_0(n)$ appears as a coefficient in the expansion of $\prod_{k < n} (1 + e^{ik\theta})$.

2.4. The basic problem illustrated by the previous examples states as follows.

Problem 2.7. Let $\{k_j, j \ge 1\}$ be an increasing sequence of positive integers and $\{p_j, j \ge 1\}$ be a sequence of reals in]0, 1[. Describe the CLT and LLT for the sequence $S_n = k_1\beta_1 + \ldots + k_n\beta_n$, $n \ge 1$, where β_j are independent Bernoulli random variables defined by

(2.11)
$$\begin{cases} \mathbb{P}\{\beta_j = 1\} = p_j \\ \mathbb{P}\{\beta_j = 0\} = 1 - p_j. \end{cases} \quad (j \ge 1)$$

In the Freiman-Pitman model, the system of independent random variables varies with the choice of the integer. And so there is, properly speaking, no central limit theorem involved and thereby no local limit theorem either, except when placing the problem in the setting of triangular arrays. Corresponding forms of the central limit theorem exist. As to suitable versions of the local limit theorem for triangular arrays *with* remainder term, we don't know whether such a result exists in the litterature. Thus it makes sense to also consider a "local" version of the previous problem.

Problem 2.8 (Finite version). To obtain effective sharp estimates of

$$\mathbb{P}\{S_n = N\}.$$

We refer to [8] where this question is investigated.

Returning to the Freiman-Pitman model, we observe that the relevant question rather concerns the search of sharp estimates of $\mathbb{P}\{S_n = 0\}$ (namely of $\int_{-1/2}^{1/2} \mathbb{E} e^{2i\pi t S_n} dt$), the random variables being centered, than working out a local limit theorem, which is quite another problem. Nevertheless, this model, as well as others previously reviewed, sheds light on limitations to the domain of validity of the local limit theorem, in a quite informative way.

Some further useful remarks are necessary. We note throughout $\{\varsigma, \varsigma_j, j \ge 1\}$ a sequence independent standard Bernoulli random variables (namely associated with $p_j \equiv 1/2$) and

$$T_n = \varsigma_1 + \ldots + \varsigma_n \qquad n \ge 1.$$

Remark 2.9 (Reduction to standard Bernoulli random variables). Let β be a Bernoulli random variable with $\mathbb{P}\{\beta = 1\} = \alpha = 1 - \mathbb{P}\{\beta = 0\}$. Assume $0 < \alpha < 1/2$. Let ε, ς be such that $\beta, \varepsilon, \varsigma$ are independent and $\mathbb{P}\{\varepsilon = 1\} = 2\alpha = 1 - \mathbb{P}\{\varepsilon = 0\}$. Trivially $\varepsilon\varsigma \stackrel{\mathcal{L}}{=} \beta$. We can thus write when $0 < p_j < 1/2, 1 \le j \le n$,

$$S_n = k_1 \varepsilon_1 \varsigma_1 + \ldots + k_n \varepsilon_n \varsigma_n.$$

Problem 2.8 reduces to first estimate (conditionnally to ε_j) a sum of the same kind

$$T'_n = k'_1\varsigma_1 + \ldots + k'_n\varsigma_n$$

with k'_i increasing, but where the Bernoulli random variables are standard.

Remark 2.10. If 1/2 < a < 1, let τ_0 be verifying $0 < \tau_0 < 2\min(\alpha, 1 - \alpha)$. Define a pair of random variables (V, ε) as follows.

$$\begin{cases} \mathbb{P}\{(V,\varepsilon) = (1,1)\} = 0 \\ \mathbb{P}\{(V,\varepsilon) = (1,0)\} = \alpha - \frac{\tau_0}{2}. \end{cases} \begin{cases} \mathbb{P}\{(V,\varepsilon) = (0,1)\} = \tau_0 \\ \mathbb{P}\{(V,\varepsilon) = (0,0)\} = 1 - \alpha - \frac{\tau_0}{2}. \end{cases}$$

Let ς be independent from (V, ε) . Then $V + \varepsilon \varsigma \stackrel{\mathcal{L}}{=} \beta$.

Remark 2.11. Fix the integer *n*. Let $T'_m = k_m \varsigma_m + \ldots + k_n \varsigma_n$, $1 \le m \le n$, and consider the parallelogram $H_m = \{h = k_{i_1} + \ldots + k_{i_r} \text{ with } i_1 < \ldots < i_r \text{ and } 1 \le r \le m\}$. Then we have the following formula

(2.12)
$$\mathbb{P}\{T'_1 = b\} = \frac{1}{2^m} \sum_{h \in H_m \cup \{0\}} \mathbb{P}\{T'_{n-m} = b - h\}.$$

3. Weighted Local Limit Theorems.

We use the characteristic function method to study the local limit theorem for the sums

$$B_{\nu} = \beta_1 + \ldots + \beta_{\nu}, \qquad \nu = 1, 2, \ldots$$

where β_j are independent random variables defined by

(3.1)
$$\mathbb{P}\{\beta_j = 0\} = \vartheta_j, \qquad \mathbb{P}\{\beta_j = k_j\} = 1 - \vartheta_j,$$

with $0 < \vartheta_j < 1$ for each j, and k_j are increasing positive weights. Let

$$\operatorname{Var}(B_{\nu}) = \sum_{j=1}^{\nu} (1 - \vartheta_j) \vartheta_j k_j^2, \qquad \vartheta = \inf_{j=1}^{\nu} \vartheta_j (1 - \vartheta_j).$$

Theorem 3.1. Let $\rho \leq k < k + \nu \leq n$ where $0 < \rho < 1$, *n* is some positive integer, and let $k_j = k + j - 1$, $j = 1, ..., \nu$. Let $1/24 < \varepsilon < 1/6$. For every $m \in \mathbb{Z}$,

$$\left| \mathbb{P}\{B_{\nu}=n\} - \frac{e^{-\frac{(\sum_{j=k}^{k+\nu-1}(1-\vartheta_j)j-n)^2}{2\operatorname{Var}(B_{\nu})}}}{\sqrt{2\pi\operatorname{Var}(B_{\nu})}} \right| \leq \frac{C}{\sqrt{\operatorname{Var}(B_{\nu})}} \left(\frac{\nu^{1/6-4\varepsilon}}{\vartheta^{1/3}\rho} + e^{-2\pi^2\vartheta\rho^2\delta^2\nu^{1/3}}\right).$$

Put

(3.2)
$$P_{\nu} = \sum_{j=1}^{\nu} (1 - \vartheta_j), \qquad M_{\nu} = \sum_{j=1}^{\nu} (1 - \vartheta_j) k_j, \qquad B_{\nu} = \sum_{j=1}^{\nu} (1 - \vartheta_j) \vartheta_j k_j^2.$$

Theorem 3.2.

$$\sup_{n \in \mathbb{Z}} \left| \mathbb{P}\{B_{\nu} = n\} - \frac{1}{\sqrt{2\pi B_{\nu}}} e^{-\frac{(n-M_{\nu})^2}{2B_{\nu}}} \right| \leq \frac{C}{P_{\nu}}$$

Further if $n_0 := \sum_{j=1}^{\nu} (1 - \vartheta_j) k_j$ is integer, then

$$\left|\mathbb{P}\{\beta_1 + \ldots + \beta_\nu = n_0\} - \frac{1}{\sqrt{2\pi B_\nu}}\right| \leq \frac{C}{P_\nu}.$$

Before passing to the proofs, we begin with making a brief analysis. Let $\varphi_j(t) = \mathbb{E} e^{2i\pi t\beta_j}$, $\varphi_{B_\nu}(t) = \mathbb{E} e^{2i\pi tB_\nu}$. By the Fourier inversion formula,

$$\mathbb{P}\{B_{\nu} = m\} = \int_{|t| \le \tau} e^{-2i\pi m t} \varphi_{B_{\nu}}(t) dt + \int_{\tau \le |t| \le \frac{1}{2}} e^{-2i\pi m t} \varphi_{B_{\nu}}(t) dt := I_{\tau}(\nu, m) + I^{\tau}(\nu, m),$$

where $\tau > 0$ will be chosen to be small, depending of m. The first integral term produces the main term and is easily tractable. The estimation of the second integral term is in fact the hard part of the problem, where all the difficulty is concentrated. It is necessary to show that

$$\left|\int_{\tau \le |t| \le \frac{1}{2}} e^{-2i\pi m t} \varphi_{B_{\nu}}(t) \mathrm{d}t\right| \ll \frac{1}{\sqrt{\mathrm{Var}(B_{\nu})}}.$$

There seems to be no other way than controlling $\int_{\tau \leq |t| \leq \frac{1}{2}} |\varphi_{B_{\nu}}(t)| dt$. From Lemma 3.3-(i) will follow that $|\varphi_{B_{\nu}}(t)| \leq \exp\left\{-2\sum_{j=1}^{\nu} \vartheta_j(1-\vartheta_j)\sin^2 \pi t k_j\right\}$. The whole matter consequently directly depends on the behaviour of the sine sum

$$\sum_{j=1}^{\nu} \vartheta_j (1 - \vartheta_j) \sin^2 \pi t k_j$$

away from 0, an obviously difficult question. Thus, answers can be expected only for specific cases.

3.1. Estimates of $I_{\tau}(\nu, m)$ **.** Recall Lemma 3 in [5]. Although stated with the choice of probability values given by (2.9), this lemma is general. For completion, we have included a slightly shorter proof.

Lemma 3.3. Let *m* be a positive real and *p* be a real such that $0 . Let <math>\beta$ be a random variable defined by $\mathbb{P}\{\beta = 0\} = p$, $\mathbb{P}\{\beta = m\} = 1 - p = q$. Let $\varphi(t) = \mathbb{E} e^{2i\pi t\beta}$. Then we have the following estimates,

- (i) For all real t, $|\varphi(t)| \le \exp\left\{-2pq\sin^2 \pi tm\right\}$
- (ii) If $q |\sin \pi t m| \le 1/3$,

$$\varphi(t) = \exp\left\{2i\pi qmt - 2\pi^2 pqm^2 t^2 + B(t)\right\},\,$$

and $|B(t)| \leq Cqm^3t^3$, the constant C being absolute.

Proof. One verifies that $|\varphi(t)|^2 = 1 - 4pq \sin^2 \pi mt$. As moreover $1 - \vartheta \leq e^{-\vartheta}$ if $\vartheta \geq 0$, (i) follows. Write now $\varphi(t) = 1 + q(e^{2i\pi mt} - 1) = 1 + u$ and notice that $|u| = 2q |\sin \pi mt|$. We use the fact that if $|\theta| \leq 2/3$, then

$$1 + \theta = \exp\{\theta - \theta^2 + B\}, \qquad |B| \le C|\theta|^3.$$

And C is an absolute constant. From the bound $|e^z - (1 + \frac{z}{1!} + \ldots + \frac{z^n}{n!})| \leq \frac{|z|^{n+1}}{(n+1)!}e^{|z|}$, valid for $z \in \mathbb{C}$ and $n \in \mathbb{N}_*$ ([18], 3.8.25), we get by applying it with $z = 2i\pi mt$,

$$\begin{aligned} |u - q (2i\pi mt - 2(\pi mt)^2)| &\leq Cqm^2 |t|^2 \\ |u^2 + (2q\pi mt)^2| &\leq Cq^2 m^3 |t|^3 \\ |u|^3 &\leq Cq^3 m^3 |t|^3. \end{aligned}$$

As we assumed $q |\sin \pi t m| \le 1/3$, we consequently find that

$$\varphi(t) = 1 + u = \exp\{u - u^2 + B\} = \exp\{2i\pi qmt - 2\pi^2 pqm^2t^2 + B(t)\},\$$
with $|B(t)| \le Cqm^3t^3$.

The next Lemma provides an estimate for the main integral term. Let $0 < \delta \leq \frac{1}{3\pi}$ and put

(3.3)
$$\tau = \frac{\delta}{(\sum_{j=1}^{\nu} (1 - \vartheta_j) k_j^3)^{1/3}}$$

Lemma 3.4. For every $n \in \mathbb{Z}$,

$$\left|\int_{-\tau}^{\tau} e^{-2i\pi nt} \varphi_{B_{\nu}}(t) \mathrm{d}t - \frac{e^{-\frac{(\sum_{j=1}^{\nu}(1-\vartheta_j)k_j-n)^2}{2\mathrm{Var}(B_{\nu})}}}{\sqrt{2\pi\mathrm{Var}(B_{\nu})}}\right| \leq C\left(\tau\delta^3 + \frac{e^{-2\pi^2\tau^2\mathrm{Var}(B_{\nu})}}{\sqrt{\mathrm{Var}(B_{\nu})}}\right).$$

Further if $n_0 := \sum_{j=1}^{\nu} (1 - \vartheta_j) k_j$ is integer, then

$$\int_{-\tau}^{\tau} e^{-2i\pi n_0 t} \varphi_{B_{\nu}}(t) \mathrm{d}t = \frac{1}{\sqrt{2\pi \mathrm{Var}(B_{\nu})}} \left(1+B\right),$$

with

$$|B| \le C \Big(\tau \delta^3 + \frac{e^{-2\pi^2 \tau^2 \operatorname{Var}(B_{\nu})}}{1 + \sqrt{\operatorname{Var}(B_{\nu})}}\Big).$$

Proof of Lemma 3.4. As $\delta \leq \frac{1}{3\pi}$, we observe that for $j = 1, \ldots, \nu$,

$$\sup_{|t| \le \tau} (1 - \vartheta_j) |\sin \pi t k_j| \le (1 - \vartheta_j) \pi \tau k_j = \frac{\delta (1 - \vartheta_j) \pi k_j}{(\sum_{j=1}^{\nu} (1 - \vartheta_j) k_j^3)^{1/3}} \le \delta (1 - \vartheta_j)^{1/3} \pi \le \frac{1}{3}$$

Lemma 3.3 thus implies,

$$\varphi_{B_{\nu}}(t) = \prod_{j=1}^{\nu} \varphi_j(t) = \exp\left\{2i\pi t \sum_{j=1}^{\nu} (1-\vartheta_j)k_j - 2\pi^2 t^2 \operatorname{Var}(B_{\nu}) + B_1(t)\right\}$$

and $|B_1(t)| \le C|t|^3 \sum_{j=1}^{\nu} (1 - \vartheta_j) k_j^3$. By (3.3),

$$\sup_{|t| \le \tau} |B_1(t)| \le C\tau^3 \sum_{j=1}^{\nu} (1 - \vartheta_j) k_j^3 = C\delta^3.$$

Noting then $\varsigma = \sum_{j=1}^{\nu} (1 - \vartheta_j) k_j - n$ and writing that

$$\varphi_{B_{\nu}}(t) = \exp\left\{2i\pi t(\varsigma+n) - 2\pi^2 t^2 \operatorname{Var}(B_{\nu}) + B_1(t)\right\},\$$

we thus deduce the following bound

$$\begin{aligned} & \left| \int_{-\tau}^{\tau} \left\{ e^{-2i\pi nt} \varphi_{B_{\nu}}(t) - e^{2i\pi t\varsigma - 2\pi^{2}t^{2}\operatorname{Var}(B_{\nu})} \right\} \mathrm{d}t \right| \\ & \leq \int_{-\tau}^{\tau} \left| e^{-2i\pi nt} \varphi_{B_{\nu}}(t) - e^{2i\pi t\varsigma - 2\pi^{2}t^{2}\operatorname{Var}(B_{\nu})} \right| \mathrm{d}t \\ & = \int_{-\tau}^{\tau} \left| e^{2i\pi t\varsigma - 2\pi^{2}t^{2}\operatorname{Var}(B_{\nu})} (e^{B(t)} - 1) \right| \mathrm{d}t \\ & \leq \int_{-\tau}^{\tau} e^{-2\pi^{2}t^{2}\operatorname{Var}(B_{\nu})} \left| e^{B(t)} - 1 \right| \mathrm{d}t \leq \int_{-\tau}^{\tau} |B(t)| \mathrm{d}t \leq C\tau \delta^{3}. \end{aligned}$$

Now we also have that

$$\int_{-\tau}^{\tau} e^{2i\pi t\varsigma - 2\pi^2 t^2 \operatorname{Var}(B_{\nu})} dt = \int_{\mathbb{R}} e^{2i\pi t\varsigma - 2\pi^2 t^2 \operatorname{Var}(B_{\nu})} dt + H$$
$$\left(t = \frac{u}{2\pi\sqrt{\operatorname{Var}(B_{\nu})}}\right) = \int_{\mathbb{R}} e^{\frac{i\varsigma u}{\sqrt{\operatorname{Var}(B_{\nu})}} - \frac{u^2}{2}} \frac{du}{2\pi\sqrt{\operatorname{Var}(B_{\nu})}} + H$$
$$= \frac{e^{-\frac{\varsigma^2}{2\operatorname{Var}(B_{\nu})}}}{\sqrt{2\pi\operatorname{Var}(B_{\nu})}} + H,$$

and recalling Boyd's estimate [18, p. 179] of Mill's ratio $R(x) = e^{x^2/2} \int_x^\infty e^{-t^2/2} dt$,

$$\frac{\pi}{\sqrt{x^2 + 2\pi} + (\pi - 1)x} \le R(x) \le \frac{\pi}{\sqrt{(\pi - 2)^2 x^2 + 2\pi} + 2x}$$

for all $x \ge 0$, we further have

$$|H| \leq \int_{|u| \geq 2\pi\tau\sqrt{\operatorname{Var}(B_{\nu})}} e^{-\frac{u^2}{2}} \frac{\mathrm{d}u}{2\pi\sqrt{\operatorname{Var}(B_{\nu})}} = \frac{e^{-2\pi^2\tau^2\operatorname{Var}(B_{\nu})}}{2\pi\sqrt{\operatorname{Var}(B_{\nu})}} R\left(2\pi\tau\sqrt{\operatorname{Var}(B_{\nu})}\right)$$
$$\leq C \frac{e^{-2\pi^2\tau^2\operatorname{Var}(B_{\nu})}}{\sqrt{\operatorname{Var}(B_{\nu})}(1+\sqrt{\operatorname{Var}(B_{\nu})})}.$$

Consequently

$$\left| \int_{-\tau}^{\tau} e^{-2i\pi nt} \varphi_{B_{\nu}}(t) \mathrm{d}t - \frac{e^{-\frac{\zeta^2}{2\mathrm{Var}(B_{\nu})}}}{\sqrt{2\pi}\mathrm{Var}(B_{\nu})} \right| \leq C \left(\tau \delta^3 + \frac{e^{-2\pi^2 \tau^2 \mathrm{Var}(B_{\nu})}}{\sqrt{\mathrm{Var}(B_{\nu})}(1 + \sqrt{\mathrm{Var}(B_{\nu})})}\right).$$

Now if there is an integer n such that $\sum_{j=1}^{\nu} (1-\vartheta_j)k_j = n$, then $\varsigma = 0$ and $\varphi_{B_{\nu}}(t) = \exp\left\{2i\pi tn - 2\pi^2 t^2 \operatorname{Var}(B_{\nu}) + B_1(t)\right\}$; whence

$$\int_{-\tau}^{\tau} e^{-2i\pi nt} \varphi_{B_{\nu}}(t) dt = \int_{-\tau}^{\tau} e^{-2\pi^{2}t^{2} \operatorname{Var}(B_{\nu})} (1 + e^{B_{1}(t)} - 1) dt$$

(3.4)
$$= (1+B) \int_{-\tau}^{\tau} e^{-2\pi^2 t^2 \operatorname{Var}(B_{\nu})} \mathrm{d}t,$$

with $|B| \leq C\tau \delta^3$. As moreover

$$\left|\int_{-\tau}^{\tau} e^{-2\pi^2 t^2 \operatorname{Var}(B_{\nu})} \mathrm{d}t - \frac{1}{\sqrt{2\pi \operatorname{Var}(B_{\nu})}}\right| \le C \frac{e^{-2\pi^2 \tau^2 \operatorname{Var}(B_{\nu})}}{\sqrt{\operatorname{Var}(B_{\nu})}(1 + \sqrt{\operatorname{Var}(B_{\nu})})},$$

we have

$$\int_{-\tau}^{\tau} e^{-2i\pi nt} \varphi_{B_{\nu}}(t) \mathrm{d}t = \frac{1}{\sqrt{2\pi \mathrm{Var}(B_{\nu})}} (1+B) (1+B_1),$$

where $|B_1| \leq Ce^{-2\pi^2 \tau^2 \operatorname{Var}(B_{\nu})} / (1 + \sqrt{\operatorname{Var}(B_{\nu})})$. We conclude to

$$\int_{-\tau}^{\tau} e^{-2i\pi nt} \varphi_{B_{\nu}}(t) \mathrm{d}t = \frac{1}{\sqrt{2\pi \mathrm{Var}(B_{\nu})}} \left(1 + B_2\right),$$

with

$$|B_2| \le C \Big(\tau \delta^3 + \frac{e^{-2\pi^2 \tau^2 \operatorname{Var}(B_{\nu})}}{1 + \sqrt{\operatorname{Var}(B_{\nu})}}\Big).$$

3.2. Estimates of $I^{\tau}(\nu, m)$. We assume here that $k_j = k + j - 1, j = 1, \dots, \nu$. Then, Lemma 3.5.

$$\int_{|t|>\tau} |\varphi_{B_{\nu}}(t)| \mathrm{d}t \le e^{-\frac{\vartheta\nu^3\tau^2}{2}}.$$

Proof. This is an immediate consequence of the following lemma

Lemma 3.6 ([5], Lemma 8). For $|t| \le 1/2$ and any positive integers m and k such that $k \ge 2$ we have m+k-1

$$\sum_{j=m}^{n+k-1} \sin^2 \pi jt \ge \frac{k}{4} \min(1, (tk)^2).$$

By Lemma 3.3-(i),

$$|\varphi_{B_{\nu}}(t)| \leq \exp\left\{-2\sum_{j=1}^{\nu}\vartheta_j(1-\vartheta_j)\sin^2\pi tk_j\right\} \leq \exp\left\{-2\vartheta\sum_{j=1}^{\nu}\sin^2\pi tk_j\right\}.$$

Thus

$$\int_{|t|>\tau} |\varphi_{B_{\nu}}(t)| \mathrm{d}t \leq \int_{|t|>\tau} e^{-\frac{\vartheta}{2}\sum_{j=1}^{\nu} \sin^2 \pi j t} \mathrm{d}t \leq \int_{|t|>\tau} e^{-\frac{\vartheta\nu}{2} \min(1,\nu|t|)^2} \mathrm{d}t \leq e^{-\frac{\vartheta\nu^3\tau^2}{2}}.$$

3.3. Proof of Theorem 3.1. By Lemma 3.4 applied with $k_j = k + j - 1$, $j = 1, ..., \nu$, for every $n \in \mathbb{Z}$,

$$\begin{aligned} \left| \int_{0}^{\tau} e^{-2i\pi nt} \varphi_{B_{\nu}}(t) \mathrm{d}t - \frac{e^{-\frac{(\sum_{j=k}^{k+\nu-1} (1-\vartheta_{j})j-n)^{2}}{2\mathrm{Var}(B_{\nu})}}}{\sqrt{2\pi}\mathrm{Var}(B_{\nu})} \right| &\leq C \left(\tau \delta^{3} + \frac{e^{-2\pi^{2}\tau^{2}\mathrm{Var}(B_{\nu})}}{\sqrt{\mathrm{Var}(B_{\nu})}}\right) \\ &\leq C \frac{\nu^{1/6-4\varepsilon}\rho^{-1} + e^{-2\pi^{2}\rho^{2}\delta^{2}\nu^{1/3}}}{\sqrt{\mathrm{Var}(B_{\nu})}}.\end{aligned}$$

By combining with Lemma 3.5 and using Fourier inversion formula,

$$\left| \mathbb{P}\{B_{\nu} = n\} - \frac{e^{-\frac{(\sum_{j=k}^{k+\nu-1}(1-\vartheta_{j})j-n)^{2}}{2\operatorname{Var}(B_{\nu})}}}{\sqrt{2\pi\operatorname{Var}(B_{\nu})}} \right| \leq C\left(\tau\delta^{3} + \frac{e^{-2\pi^{2}\tau^{2}\operatorname{Var}(B_{\nu})}}{\sqrt{\operatorname{Var}(B_{\nu})}} + e^{-\frac{\vartheta\nu^{3}\tau^{2}}{8}}\right).$$

We have the following estimates

$$\begin{cases} \text{(i)} & \frac{\delta}{n\nu^{1/3}} \leq \tau = \frac{\delta}{(\sum_{j=1}^{\nu}(1-\vartheta_j)k_j^3)^{1/3}} \leq \frac{\delta}{\rho n(\vartheta\nu)^{1/3}},\\ \text{(ii)} & \vartheta(\rho n)^2 \nu \leq \operatorname{Var}(B_{\nu}) = \sum_{j=1}^{\nu}(1-\vartheta_j)\vartheta_j k_j^2 \leq n^2\nu,\\ \text{(iii)} & \tau^2 \operatorname{Var}(B_{\nu}) \geq \frac{\vartheta\delta^2}{n^2\nu^{2/3}}(\rho n)^2 \nu = \vartheta\rho^2 \delta^2 \nu^{1/3}. \end{cases}$$

Choose $\delta = \nu^{-\varepsilon}$ with $1/24 < \varepsilon < 1/6$. Then

$$\tau\delta^3 \le \frac{\delta^4}{\rho n(\vartheta\nu)^{1/3}} = \frac{1}{\rho n\vartheta^{1/3}\nu^{1/3+4\varepsilon}} = \frac{\nu^{1/6-4\varepsilon}}{\vartheta^{1/3}\rho n\nu^{1/2}} \le \frac{\nu^{1/6-4\varepsilon}}{\vartheta^{1/3}\rho\sqrt{\operatorname{Var}(B_\nu)}}$$

We pass to the control of the error terms. For the major integral term we have,

$$\tau \delta^3 + \frac{e^{-2\pi^2 \tau^2 \operatorname{Var}(B_{\nu})}}{\sqrt{\operatorname{Var}(B_{\nu})}} \le \frac{1}{\sqrt{\operatorname{Var}(B_{\nu})}} \Big(\frac{\nu^{1/6-4\varepsilon}}{\vartheta^{1/3}\rho} + e^{-2\pi^2 \vartheta \rho^2 \delta^2 \nu^{1/3}}\Big).$$

Consequently,

$$\left| \mathbb{P}\{B_{\nu}=n\} - \frac{e^{-\frac{\left(\sum_{j=k}^{k+\nu-1}(1-\vartheta_{j})j-n\right)^{2}}{2\operatorname{Var}(B_{\nu})}}}{\sqrt{2\pi\operatorname{Var}(B_{\nu})}} \right| \leq \frac{C}{\sqrt{\operatorname{Var}(B_{\nu})}} \left(\frac{\nu^{1/6-4\varepsilon}}{\vartheta^{1/3}\rho} + e^{-2\pi^{2}\vartheta\rho^{2}\delta^{2}\nu^{1/3}}\right).$$

3.4. Other Estimates of $I^{\tau}(\nu, m)$ **.** The following lemma is relevant. Introduce for $q \geq 1$ integer,

(3.5)
$$\varphi_{\theta_1,...,\theta_{\nu},k_1,...,k_{\nu}}(q) = \varphi(q) := \frac{\left\|\sum_{j=1}^{\nu} \theta_j \cos 2\pi t k_j\right\|_{2q}}{\sum_{j=1}^{\nu} \theta_j}.$$

Lemma 3.7. For any $0 < c \le 1$,

$$\int_0^1 |\varphi_{B_\nu}(t)| \mathrm{d}t \leq \left(\frac{\varphi(q)}{c}\right)^{2q} + e^{-(1-c)\sum_{j=1}^\nu \vartheta_j(1-\vartheta_j)}.$$

Proof of Lemma 3.7. Let $\theta_j = \vartheta_j(1-\vartheta_j), j = 1..., \nu$ and note $E = \{t; |t| \leq \frac{1}{2} : \left|\sum_{j=1}^{\nu} \theta_j \cos 2\pi t k_j\right| > c \sum_{j=1}^{\nu} \theta_j \}$. At first, by using Tchebycheff's inequality,

$$\lambda\{E\} \leq \left(c\sum_{j=1}^{\nu} \theta_{j}\right)^{-2q} \int_{-\frac{1}{2}}^{\frac{1}{2}} \left|\sum_{j=1}^{\nu} \theta_{j} \cos 2\pi t k_{j}\right|^{2q} \mathrm{d}t = c^{-2q} \varphi(q)^{2q},$$

Since $|\varphi_{B_{\nu}}(t)| \leq 1$, we have

$$\int_{0}^{1} |\varphi_{B_{\nu}}(t)| \mathrm{d}t \le \lambda\{E\} + \int_{E^{c}} |\varphi_{B_{\nu}}(t)| \mathrm{d}t \le c^{-2q} \varphi(q)^{2q} + \int_{E^{c}} |\varphi_{B_{\nu}}(t)| \mathrm{d}t,$$

By Lemma 3.3, using that $2\sin^2 a = 1 - \cos 2a$, we have for all real t,

$$\begin{aligned} |\varphi_{B_{\nu}}(t)| &= \prod_{j=1}^{\nu} |\varphi_{j}(t)| \le \exp\left\{-\sum_{j=1}^{\nu} 2\theta_{j} \sin^{2} \pi t k_{j}\right\} \\ &= \exp\left\{-\sum_{j=1}^{\nu} \theta_{j}\right\} \exp\left\{\sum_{j=1}^{\nu} \theta_{j} \cos 2\pi t k_{j}\right\}. \end{aligned}$$

So that

$$\int_{E^c} |\varphi_{B_{\nu}}(t)| \mathrm{d}t \leq \int_{E^c} e^{-\sum_{j=1}^{\nu} 2\theta_j \sin^2 \pi t k_j} \mathrm{d}t$$
$$\leq e^{-\sum_{j=1}^{\nu} \theta_j} \int_{E^c} e^{|\sum_{j=1}^{\nu} \theta_j \cos 2\pi t k_j|} \mathrm{d}t$$
$$\leq e^{-(1-c)\sum_{j=1}^{\nu} \theta_j}.$$

By combining

$$\int_0^1 |\varphi_{B_\nu}(t)| \mathrm{d}t \leq \left(\frac{\varphi(q)}{c}\right)^{2q} + e^{-(1-c)\sum_{j=1}^\nu \theta_j}.$$

Remark 3.8. Assume $\vartheta_j = \vartheta$ for all j. Then $\varphi(q) = \nu^{-1} \left\| \sum_{j=1}^{\nu} \cos 2\pi t k_j \right\|_{2q}$ and further

$$\left\|\sum_{j=1}^{\nu}\cos 2\pi t k_{j}\right\|_{2q}^{2q} = N_{2q}(\mathcal{N}),$$

where $N_{2q}(\mathbb{N})$ is the number of solutions of (2.5) with corresponding set of values $\mathbb{N} = \{k_1, \ldots, k_\nu\}$. So that

$$\int_0^1 |\varphi_{B_\nu}(t)| \mathrm{d}t \leq \frac{N_{2q}(\mathcal{N})}{(c\nu)^{2q}} + e^{-(1-c)\vartheta(1-\vartheta)\nu}.$$

– In the case when $\mathcal{N} = \{1, \dots, \nu\}$, this together with Theorem 2.4 gives

$$\int_0^1 |\varphi_{B_{\nu}}(t)| \mathrm{d}t \le \frac{\nu^{2q-1}}{\sqrt{q}(c\nu)^{2q}} + e^{-(1-c)\vartheta(1-\vartheta)\nu} = \frac{1}{\nu c^{2q}\sqrt{q}} + e^{-(1-c)\vartheta(1-\vartheta)\nu}.$$

Taking $c = 1 - (2q)^{-1}$ gives

$$\int_0^1 |\varphi_{B_\nu}(t)| \mathrm{d}t \leq \frac{C}{\nu\sqrt{q}} + e^{-\vartheta(1-\vartheta)\nu/2q}.$$

Take q large, $q \sim \vartheta(1-\vartheta)\nu/3\log\nu$. It follows that

$$\int_0^1 |\varphi_{B_\nu}(t)| \mathrm{d}t \leq \frac{C \log \nu}{\nu^{3/2}}$$

whereas, in the other hand $\operatorname{Var}(B_{\nu}) = \vartheta(1-\vartheta) \sum_{j=1}^{\nu} j^2 \sim C \vartheta(1-\vartheta) \nu^3 \dots$

Remark 3.9. We also have

$$\sum_{j=1}^{\nu} \vartheta_j (1-\vartheta_j) \sin^2 \pi t k_j \geq \nu \Big(\prod_{j=1}^{\nu} \vartheta_j (1-\vartheta_j) \Big)^{1/\nu} \Big| \prod_{j=1}^{k_\nu} \sin \pi t j \Big|^{2/\nu}.$$

3.5. Proof of Theorem 3.2. Let $\delta = 1/2$. By Lemma 3.4, Lemma 3.7 and Fourier inversion formula

$$\left| \mathbb{P}\{B_{\nu} = n\} - \frac{e^{-\frac{(n-\mathbb{E}|B_{\nu}|)^{2}}{2\operatorname{Var}(B_{\nu})}}}{\sqrt{2\pi\operatorname{Var}(B_{\nu})}} \right| \leq C \left\{ \tau \delta^{3} + \frac{e^{-2\pi^{2}\tau^{2}\operatorname{Var}(B_{\nu})}}{\sqrt{\operatorname{Var}(B_{\nu})}} + \frac{1}{\sum_{j=1}^{\nu} \vartheta_{j}(1-\vartheta_{j})} \right\}$$

$$\leq C \left\{ \frac{1}{(\sum_{j=1}^{\nu} (1-\vartheta_{j})k_{j}^{3})^{1/3}} + \frac{e^{-2\pi^{2}\delta L^{2}}}{(\sum_{j=1}^{\nu} (1-\vartheta_{j})k_{j}^{2})^{1/2}} + \frac{1}{\sum_{j=1}^{\nu} \vartheta_{j}(1-\vartheta_{j})} \right\}.$$

$$\leq \frac{C}{\sum_{j=1}^{\nu} \vartheta_{j}(1-\vartheta_{j})}.$$

Now similarly,

$$\begin{aligned} \left| \mathbb{P}\{B_{\nu} = n_{0}\} - \frac{1}{\sqrt{2\pi \operatorname{Var}(B_{\nu})}} \right| &\leq \frac{B_{2}}{\sqrt{2\pi \operatorname{Var}(B_{\nu})}} + \frac{1}{\sum_{j=1}^{\nu} \vartheta_{j}(1 - \vartheta_{j})} \\ &\leq \frac{C}{(\sum_{j=1}^{\nu} (1 - \vartheta_{j})k_{j}^{2})^{1/2}} \left(\frac{1}{(\sum_{j=1}^{\nu} (1 - \vartheta_{j})k_{j}^{3})^{1/3}} + e^{-2\pi^{2}\delta L^{2}} \right) \\ &+ \frac{1}{\sum_{j=1}^{\nu} \vartheta_{j}(1 - \vartheta_{j})} \leq \frac{C}{\sum_{j=1}^{\nu} \vartheta_{j}(1 - \vartheta_{j})}. \end{aligned}$$

This achieves the proof.

4. An ASLLT related to Burr's problem.

Let $\lambda_0 < \lambda_1 < \dots$ be a sequence of positive integers, call it A, and let

$$P(A) = \Big\{ \sum_{i} \varepsilon_i \lambda_i, \ \varepsilon_i = 0 \text{ or } 1, \ a_i \in A \text{ and } \sum_{i} \varepsilon_i < \infty \Big\}.$$

Burr asked in [3] which sets S of integers are equal to P(A) for some A? He mentioned that if the complement of S grows sufficiently rapidly, then there exists such a sequence A. Hegyvari showed in [12] that if $B = \{b_i, i \ge 1\}$ is such that $7 \le b_1 < b_2 < \cdots$ and

$$(4.1) b_{n+1} \ge 5b_n for every n,$$

then there exists a sequence A such that $P(A) = \mathbb{N} \setminus B$, thereby improving substantially an earlier unpublished result of Burr. He also showed that his result cannot be improved essentially. More precisely, if B is such that

(4.2)
$$b_{n+1} \le 2b_n$$
 for every *n* large enough,

and B is a Sidon set, namely $b_i + b_j = b_k + b_\ell$ implies $i = k, j = \ell$ or $i = \ell, j = k$, then there is no sequence A for which $P(A) = \mathbb{N} \setminus B$. We refer to [3, 6, 12] for similar questions. Here we examine a variant of the initial problem. Consider the set E composed with all finite sums

(4.3)
$$\lambda_{j_1} + \ldots + \lambda_{j_n}, \qquad 0 \le j_1 \le \ldots \le j_n, \quad n \ge 1.$$

Let $0 < \eta < 1$ and let $E_{\eta} \subseteq E$ be the set composed with all finite sums $\lambda_{j_1} + \ldots + \lambda_{j_n}$ such that at most $|\eta n|$ summands may coincide.

Now let $\{x_n, n \ge 1\}$ be a sequence of integers increasing nearly linearly, so that it is a relatively "full" sequence. More precisely, we assume there are reals $a > 1, \delta > 0$ such that

(4.4)
$$x_n - na \sim \delta \sqrt{n}, \qquad n \to \infty.$$

We are interested in estimating from below the proportion of terms from this sequence which may be represented by a sum $\lambda_{j_1} + \ldots + \lambda_{j_n}$, namely which belong to E.

Theorem 4.1. Let $D = \text{g.c.d.} \{\lambda_i - \lambda_j, i > j \ge 0\}$. Let also

(4.5)
$$\rho = \sup \left\{ r : \sum_{j=0}^{\infty} \lambda_j r^j < \infty \right\}.$$

Assume that $0 < \rho \leq 1$. Then for some $0 < \eta < 1$ depending on both ρ and a,

$$\liminf_{t \to \infty} \frac{1}{\log t} \sum_{n \le t} \frac{1}{\sqrt{n}} \mathbf{1}_{\{x_n \in E_\eta\}} \ge \frac{D}{\sqrt{2\pi\sigma}} e^{-\frac{\delta^2}{2\sigma^2}}.$$

Here we have noted $\sigma^2 = (1-r) \sum_{j=0}^{\infty} \lambda_j^2 r^j - ((1-r) \sum_{j=0}^{\infty} \lambda_j r^j)^2$, and $0 < r < \rho$ is solution of the equation

$$(1-r)\sum_{j=0}^{\infty}\lambda_j r^j = a.$$

Further, there exists with probability one a random subsequence $\lambda'_j = \lambda'_j(\omega)$, j = 1, ..., tending to infinity with n, such that for all n large enough, among $\lambda'_1, \ldots, \lambda'_n$ at most $\lfloor n\eta \rfloor$ may coincide, and

$$\lim_{t \to \infty} \frac{1}{\log t} \sum_{n \le t} \frac{1}{\sqrt{n}} \mathbf{1}_{\{x_n = \lambda'_1 + \dots + \lambda'_n\}} = \frac{D}{\sqrt{2\pi\sigma}} e^{-\frac{\delta^2}{2\sigma^2}}.$$

4.1. Preliminaries. We first recall some auxiliary results on which the proof is based. Let X be a square integrable random variable with lattice distribution function F and put

(4.6)
$$\mu = \mathbb{E} X, \qquad \sigma^2 = \mathbb{E} X^2 - (\mathbb{E} X)^2.$$

Let *D* be the maximal span of *X*. Let also $\{X_k, k \ge 1\}$ be independent copies of *X*, and consider their partial sums $S_n = X_1 + \ldots + X_n$, $n \ge 1$. We assume throughout that $\sigma > 0$. Almost sure versions with rate of Gnedenko's theorem (see after (1.1)) were recently proved in [10]. Let $g(x) = \frac{D}{\sqrt{2\pi\sigma}} e^{-x^2/(2\sigma^2)}$, *x* real. By Gnedenko's local limit theorem,

(4.7)
$$\lim_{n \to \infty} \sqrt{n} \mathbb{P}\{S_n = \kappa_n\} = g(\kappa),$$

for any sequence $\{\kappa_n, n \ge 1\}$ of reals such that

(4.8)
$$\lim_{n \to \infty} \frac{\kappa_n - n\mu}{\sqrt{n}} = \kappa$$

We say that X satisfies an almost sure local limit theorem if

(4.9)
$$\lim_{N \to \infty} \frac{1}{\log N} \sum_{n=1}^{N} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} \stackrel{\text{a.s.}}{=} g(\kappa),$$

holds whenever (4.8) is satisfied. It is easily seen that (4.9) amounts to establish

(4.10)
$$\lim_{N \to \infty} \frac{1}{\log N} \sum_{n=1}^{N} \frac{B_n}{n} \stackrel{\text{a.s.}}{=} 0,$$

where we put $B_n = \sqrt{n} (\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P} \{S_n = \kappa_n\}).$

Theorem 4.2 ([10], Theorem 1). Assume that $\mathbb{E} X^{2+\varepsilon} < \infty$ for some positive ε . Then,

$$\lim_{N \to \infty} \frac{1}{\log N} \sum_{n \le N} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} \stackrel{a.s.}{=} g(\kappa),$$

for any sequence of integers $\{\kappa_n, n \ge 1\}$ such that (4.8) holds. Moreover, if (4.8) is sharpened as follows,

$$\frac{\kappa_n - n\mu}{\sqrt{n}} = \kappa + \mathcal{O}_\eta \big((\log n)^{-1/2 + \eta} \big),$$

then

$$\frac{1}{\log N} \sum_{n \le N} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} \stackrel{\text{a.s.}}{=} g(\kappa) + \mathcal{O}_\eta \left((\log N)^{-1/2 + \eta} \right) \Big).$$

4.2. Proof. We consider the following random model. Let $0 < r < \rho$ and let X be a random variable defined by

$$\mathbb{P}\{X=\lambda_j\}=(1-r)r^j, \qquad j=0,1,\ldots$$

The function $\mu(r) = (1-r) \sum_{j=0}^{\infty} \lambda_j r^j$ is continuous on $[0, \rho[$. Further $\mu(0) = 0$ and $\lim_{r\uparrow\rho} \mu(r) = \infty$. We can thus select a real $r \in]0, \rho[$ so that $\mathbb{E} X = \mu(r) = a$. Next $\mathbb{E} X^2 = (1-r) \sum_{j=0}^{\infty} \lambda_j^2 r^j < \infty$. And because $r < \rho$, $\mathbb{E} X^{2+\alpha} < \infty$ for some positive α . It is further clear that $\sigma^2 = \mathbb{E} X^2 - (\mathbb{E} X)^2$ cannot vanish unless X is a constant almost surely, since $\sigma^2 = \mathbb{E} (X - \mathbb{E} X)^2$. This case being excluded by construction, we have $\sigma > 0$. Let also $\{X_k, k \ge 1\}$ be independent copies of X, and consider their partial sums $S_n = X_1 + \ldots + X_n, n \ge 1$

Now observe that

$$\mathbb{P}\left\{X_{i_1} = X_{i_2} = \dots = X_{i_k}\right\} = \sum_{j=0}^{\infty} \mathbb{P}\left\{X = \lambda_j\right\}^k = (1-r)^k \sum_{j=0}^{\infty} r^{kj}$$
$$= \frac{(1-r)^k}{(1-r^k)} \le 2(1-r)^k,$$

if k is large, which we do assume. Thus

$$\mathbb{P}\{\exists 1 \le i_1 < \ldots < i_k \le n : X_{i_1} = X_{i_2} = \ldots = X_{i_k}\} \le 2C_n^k (1-r)^k.$$

We take $k = |n\eta|$. Since $n! \sim \sqrt{2\pi n} n^n e^{-n}$, we have for n large

$$C_{n}^{k} \leq 2\left(\frac{n}{n-k}\right)^{n-k} \left(\frac{n}{k}\right)^{k} \left(\frac{n}{2\pi(n-k)k}\right)^{1/2} \leq \left(\frac{1}{1-\eta}\right)^{n(1-\eta)} \left(\frac{1}{\eta}\right)^{n\eta} \\ = \left[\left(\frac{1}{1-\eta}\right)^{\frac{1}{\eta}-1} \left(\frac{1}{\eta}\right)\right]^{n\eta}.$$

Let

$$C_n = \left\{ \exists 1 \le i_1 < \ldots < i_{\lfloor n\eta \rfloor} : X_{i_1} = X_{i_2} = \ldots = X_{i_{\lfloor n\eta \rfloor}} \right\}, \qquad n = 1, 2, \ldots$$

These sets are non-increasing. And so

$$\mathbb{P}\Big\{\bigcup_{n=m}^{\infty} C_n\Big\} \le \mathbb{P}\{C_m\} \le \Big[\Big(\frac{1-r}{\eta(1-\eta)^{\frac{1}{\eta}-1}}\Big)\Big]^{m\eta}.$$

Since $(1-\eta)^{\frac{1}{\eta}-1} = \exp\{-\frac{1-\eta}{\eta}\log\frac{1}{1-\eta}\} \to 1$ as $\eta \uparrow 1$, and 0 < r < 1, it follows that one can select η so that

$$\frac{1-r}{\eta(1-\eta)^{\frac{1}{\eta}-1}} < 1.$$

This choice implies that

$$\mathbb{P}\big\{\limsup_{n \to \infty} C_n\big\} = 0.$$

Thus, with probability one, for all n large enough, there is no k-uple, $1 \leq i_1 < \ldots < i_k$, with $k \geq \lfloor n\eta \rfloor$, such that $X_{i_1} = X_{i_2} = \ldots = X_{i_k}$. In particular, with probability one, for all n large enough, at most $\lfloor n\eta \rfloor$ from the random variables X_i , $i \leq n$ may coincide.

Besides, using Gnedenko's theorem we have, uniformly in N,

$$\sqrt{n} \mathbb{P}\{S_n = N\} = \frac{D}{\sigma\sqrt{2\pi}} e^{-(N-na)^2/2n\sigma^2} + o(1).$$

By assumption $\lim_{n\to\infty} (x_n - na)/\sqrt{n} = \delta$, so that (4.8) is satisfied. Therefore $\sqrt{n} \mathbb{P}\{S_n = x_n\} \sim \frac{D}{\sigma\sqrt{2\pi}}e^{-\frac{\delta^2}{2\sigma^2}}$ as $n \to \infty$. Since the assumptions of Theorem 4.2 are also fulfilled, it follows that

(4.11)
$$\lim_{t \to \infty} \frac{1}{\log t} \sum_{n \le t} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = x_n\}} \stackrel{a.s.}{=} \frac{D}{\sqrt{2\pi\sigma}} e^{-\frac{\delta^2}{2\sigma^2}}$$

By picking ω in a measurable set of full measure, we find by what proceeds and (4.11), that there exists a subsequence $\lambda'_0 \leq \lambda'_1 \leq \ldots, \lambda'_n = \lambda'_n(\omega)$, such that

(4.12)
$$\lim_{t \to \infty} \frac{1}{\log t} \sum_{n \le t} \frac{1}{\sqrt{n}} \mathbf{1}_{\{x_n = \lambda'_1 + \dots + \lambda'_n\}} = \frac{D}{\sqrt{2\pi\sigma}} e^{-\frac{\delta^2}{2\sigma^2}}$$

Further, for all *n* large enough, at most $\lfloor n\eta \rfloor$ from the summands $\lambda'_1, \ldots, \lambda'_n$ may coincide. Thus $\lambda'_n \to \infty$ with *n*. But $x_n = \lambda'_1 + \ldots + \lambda'_n$ and the fact that among $\lambda'_1, \ldots, \lambda'_n$, at most $\lfloor n\eta \rfloor$ can coincide, implies that $x_n \in E_\eta$. We consequently deduce

$$\liminf_{t \to \infty} \frac{1}{\log t} \sum_{n \le t} \frac{1}{\sqrt{n}} \mathbf{1}_{\{x_n \in E_\eta\}} \ge \frac{D}{\sqrt{2\pi\sigma}} e^{-\frac{x^2}{2\sigma^2}},$$

as claimed. The second part of the Theorem is a direct consequence of (4.12).

5. A Concluding Remark.

A probably well-known fact is that the local limit theorem is not a sufficiently sharp tool for estimating the probability $\mathbb{P}\{S_n \in E\}$, where E is an infinite set of integers and $S_n = X_1 + \ldots + X_n$ a sum of independent copies of a random variable X. As we could not find in the litterature an explicit example, we mention here a very simple one given in [24] and showing that this already arises for bounded random variables and for elementary sets E, namely arithmetic progressions. Let d be some positive integer and take $E = d\mathbb{N}$. Let also $B_n = \beta_1 + \ldots + \beta_n$, where β_i are independent standard Bernoulli random variables. By using the sharpest form of the local limit theorem for standard Bernoulli random variables, derived from [19, Theorem 13, Chapter 7],

$$\sup_{z} \left| \mathbb{P} \{ B_n = z \} - \sqrt{\frac{2}{\pi n}} e^{-\frac{(2z-n)^2}{2n}} \right| = o(1/n^{3/2}),$$

one easily gets

(5.1)
$$\sup_{2 \le d \le n} \left| \mathbb{P}\{B_n \in d\mathbb{N}\} - \sqrt{\frac{2}{\pi n}} \sum_{z \equiv 0 \ (d)} e^{-\frac{(2z-n)^2}{2n}} \right| = o(\frac{\sqrt{\log n}}{n}).$$

By operating quite differently, we obtained in [26] the following uniform estimate. Let $\Theta(d, m)$ be the Theta elliptic function defined by

$$\Theta(d,m) = \sum_{\ell \in \mathbb{Z}} e^{im\pi \frac{\ell}{d} - \frac{m\pi^2 \ell^2}{2d^2}},$$

We have

(5.2)
$$\sup_{2 \le d \le n} \left| \mathbb{P} \left\{ B_n \in d\mathbb{N} \right\} - \frac{\Theta(d,n)}{d} \right| = \mathcal{O} \left(\frac{\log^{5/2} n}{n^{3/2}} \right).$$

By using Poisson summation formula, this implies that

$$\sup_{2 \le d \le n} \left| \mathbb{P} \{ B_n \in d\mathbb{N} \} - \sqrt{\frac{2}{\pi n}} \sum_{z \equiv 0 \ (d)} e^{-\frac{(2z-n)^2}{2n}} \right| = \mathcal{O} \left(\frac{\log^{5/2} n}{n^{3/2}} \right)$$

which is much better than (5.1). We refer to [24] for more details.

References

- R. Arratia, A. D. Barbour and S. Tavaré, Logarithmic Combinatorial Structures, Ann. Probab. 28 no 4, (2002), 1620–1644.
- [2] R. Arratia, A. D. Barbour and S. Tavaré, The Poisson-Dirichlet distribution and the scale-invariant Poisson process, Comb. Probab. Comput. 8, (1999), 407–416.
- [3] S. A. Burr, Combinatorial theory and its applications III, Ed. P. Erdös, A. Rényi, V.T. Sös, Coll. Math. Soc. J. Bolyai 4. North-Holland Publ. Comp. (Amsterdam-London), (1970), p.1155.
- [4] Y. S. Chow and H. Teicher, Probability Theory: Independence, Interchangability, Martingales, Third Edition, Springer Texts in Statistics, (2003), Springer-Verlag New York-Berlin-Heidelberg.
- [5] G. A. Freiman and J. Pitman, Partitions into distinct large parts, J. Austral. Math. Soc. (Series A) 57, (1994), 386–416.
- [6] J. Folkman, On the representation of integers as sums of distinct terms from a fixed sequence Canad. J. Math. 18, (1966), 643–655.
- [7] R. Giuliano and M. Weber, (2012) Sharp bounds for the number of solutions of a particular diophantine equation: a probabilistic approach via the Féjer kernel, *preprint*.
- [8] R. Giuliano and M. Weber, Approximate Local Limit Theorems with Effective Rate and Application to Random Walks in Random Scenery, arXiv:1412.3980v1, (2014).
- [9] R. Giuliano, Z. S. Szewczak and M. Weber, Almost Sure Local Limit Theorem for the Dickman distribution, arXiv:1309.1578v1.
- [10] R. Giuliano and M. Weber, (2011) Almost sure local limit theorems with rate, Stochastic Analysis and Applications 29, 779–798.
- [11] N. Hegyvári, On representation problems in the additive number theory, Acta Math. Hungar. 72 (1-2), (1996), 35-41.
- [12] D. Hensley, The convolution powers of the Dickman function, J. London Math. Soc. 33, (1986), 3595–406.
- [13] H.-K. Hwang and T.-H. Tsai, Quickselect and the Dickman function, Combinatorics, Probability and Computing 11, (2002), 353–371.
- [14] I. A. Ibragimov and Y. V. Linnik, (1971) Independent and stationary sequences of random variables, Wolters-Noordhoff Publishing Groningen, The Netherlands.
- [15] M. A. Kolmogorov, Sur les propriétés de fonctions de concentrations de M. P. Lévy, Annales de l'I. H. P. 16 no. 1, (1958), 27–34.
- [16] D. MacDonald, On local limit theorems for integer valued random variables, Theor. of Prob. Appl. 33, (1979), 352–355.
- [17] V. K. Matskyavichyus, On a lower bound for the convergence rate in a local limit theorem, Theor. Prob. & Appl. 30 (4),(1986), 810–814.

RITA GIULIANO AND MICHEL WEBER

- [18] D. S. Mitrinović, (1970) Analytic inequalities, Springer Verlag 165.
- [19] V. V. Petrov, (1975) Sums of Independent Random Variables, Ergebnisse der Math. und ihre Grenzgebiete **82**, Springer.
- [20] A. G. Postnikov, (1988) Introduction to analytic number theory, AMS Translation of mathematical monographs 68. First publ. in Russian in 1971.
- [21] A. Röllin and N. Ross, (2012) Local limit theorems via Landau-Kolmogorov inequalities, arXiv:1011.3100v2, (2012), to appear in *Bernoulli*.
- [22] G. Tenenbaum, (2008) Introduction à la théorie analytique et probabiliste des nombres, Coll. Échelles Ed. Belin Paris.
- [23] W. Vervaat, (1972) Success epochs in Bernoulli trials with applications in number theory, Math. Center Tracts No 42, Mathematisch Centrum, Amsterdam.
- [24] M. Weber, Correlation properties of divisors of Bernoulli sums, unpublished notes, (2010).
- [25] M. Weber, On localization in Kronecker's diophantine theorem, Uniform Distribution Theory, 4 No. 1, (2009), 97–116.
- [26] M. Weber, Small divisors of Bernoulli sums, Indag. Math. 18 (2), (2007), 281–293.

IRMA, UMR 7501, UNIVERSITÉ LOUIS-PASTEUR, 7 RUE RENÉ DESCARTES, 67084 STRASBOURG CEDEX, FRANCE. E-MAIL: michel.weber@math.unistra.fr

DIPARTIMENTO DI MATEMATICA, VIA F. BUONARROTI 2, 56127 PISA, ITALY. E-MAIL: giuliano@dm.unipi.it