Unique Factorization and Controllability of Tail-Biting Trellis Realizations via Controller Granule Decompositions

G. David Forney, Jr. Laboratory for Information and Decision Systems Massachusetts Institute of Technology Cambridge, MA 02139 USA Email: forney@mit.edu

*Abstract***—The Conti-Boston factorization theorem (CBFT) for linear tail-biting trellis realizations is extended to group realizations with a new and simpler proof, based on a controller granule decomposition of the behavior and known controllability results for group realizations. Further controllability results are given; e.g., a trellis realization is controllable if and only if its top (controllability) granule is trivial.**

I. INTRODUCTION

Tail-biting trellis realizations are the simplest class of realizations of codes on cyclic graphs. Decoding is generally simpler than for conventional trellis realizations [\[1\]](#page-4-0).

Koetter and Vardy [\[8\]](#page-4-1), [\[9\]](#page-4-2) developed the foundations of the theory of linear tail-biting trellis realizations. Their key result was a factorization theorem (KVFT), which shows that every reduced realization has a factorization into elementary trellises.

Recently, Conti and Boston [\[2\]](#page-4-3) have proved a stronger unique factorization theorem (CBFT): the behavior ("label code") of a reduced linear tail-biting trellis realization factors uniquely into quotient spaces of "span subcodes." This work was the main stimulus for the work reported here.

Our main result is a generalization of the CBFT to group realizations, with a new proof that we feel is even simpler and more insightful. [\[2,](#page-4-3) Remark III.3] notes that such a generalization is not straightforward.

In Section [II,](#page-0-0) we introduce a granule decomposition along the lines of the controller granule decomposition of minimal conventional trellis realizations of Forney and Trott [\[5\]](#page-4-4), [\[6\]](#page-4-5), and the span subcode decomposition of [\[2\]](#page-4-3).

In Section [III,](#page-2-0) using results of [\[3\]](#page-4-6) on the controllability of group realizations, we show that this granule decomposition yields a unique factorization of a group trellis behavior B. We develop other controllability properties not considered in [\[2\]](#page-4-3); *e.g.,* the trellis diagram of an uncontrollable group trellis realization is disconnected [\[4\]](#page-4-7). We show that the controller canonical realization based on this factorization is one-to-one, minimal, and group-theoretic, but possibly nonhomomorphic.

Our development uses only elementary group theory, principally the *fundamental theorem of homomorphisms* (FTH) and the *correspondence theorem* (CT). For a brief introduction to the necessary group theory and our notation, see [\[3\]](#page-4-6).

A. Preliminaries

A (tail-biting) trellis realization R of length n is defined by a set of *n symbol alphabets* $\{A_j, j \in \mathbb{Z}_n\}$, a set of *n state alphabets* $\{\mathcal{S}_j, j \in \mathbb{Z}_n\}$, and a set of *n constraint codes* $\{\mathcal{C}_j \subseteq \mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}, j \in \mathbb{Z}_n\}$, where index arithmetic is in \mathbb{Z}_n ; *e.g.,* $\mathcal{C}_{n-1} \subseteq \mathcal{S}_{n-1} \times \mathcal{A}_{n-1} \times \mathcal{S}_0$.

The *configuration universe* $\mathcal{U} = \prod_{j \in \mathbb{Z}_n} C_j$ is thus a subset of $S \times A \times S$, where $A = \prod_{j \in \mathbb{Z}_n} A_j$ and $S = \prod_{j \in \mathbb{Z}_n} S_j$.

In a linear trellis realization, each symbol or state alphabet is a finite-dimensional vector space over some field \mathbb{F} , and each C_j is a subspace of $S_j \times A_j \times S_{j+1}$, so U is a subspace of $S \times A \times S$. (In [\[9\]](#page-4-2) and [\[2\]](#page-4-3), it is assumed that $A_i = \mathbb{F}$ always.) In a group trellis realization, each symbol or state alphabet is a finite abelian group, and each C_i is a subgroup of $S_j \times A_j \times S_{j+1}$, so U is a subgroup of $S \times A \times S$.

The *extended behavior* $\bar{\mathfrak{B}}$ of R is the set of configurations $(s, a, s') \in \mathcal{U}$ such that $s = s'$; *i.e.*, such that the constraints of U and the equality constraints $s = s'$ are both satisfied [\[3\]](#page-4-6). Its *behavior* \mathfrak{B} is the projection of $\bar{\mathfrak{B}}$ onto $A \times S$, which is an isomorphism. The *code* $\mathcal C$ realized by $\mathcal R$ is the projection of $\mathfrak B$ or $\mathfrak B$ onto $\mathcal A$.

The (normal) *graph* of R [\[3\]](#page-4-6) is the single-cycle graph with *n* vertices corresponding to the constraint codes C_j , *n* edges corresponding to the state variables S_i , where edge S_j is incident on vertices C_{j-1} and C_j , and n half-edges corresponding to the symbol variables A_j , where half-edge A_i is incident only on vertex C_i .

II. GRANULE DECOMPOSITION

A. Partial ordering of fragments

A *proper fragment* of a trellis realization R corresponds to a *circular interval* $[j, k), j \in \mathbb{Z}_n, k \in \mathbb{Z}_n$, and will be denoted by $\mathcal{F}^{[j,k)}$. $\mathcal{F}^{[j,k)}$ includes the constraint codes $\{\mathcal{C}_{j'}, j' \in [j,k)\}\$ and the *internal state variables* $\{S_{j'}, j' \in (j, k)\}\$, and has *boundary* $\{S_i, S_k\}$. Accordingly, we define its *vertex set* as $V(\mathcal{F}^{[j,k)}) = [j,k)$, and its *edge set* as $E(\mathcal{F}^{[j,k)}) = (j,k)$. The (normal) graph of every proper fragment is cycle-free.

We define the *level* of $\mathcal{F}^{[j,k)}$ as the number $\ell = |E(\mathcal{F}^{[j,k)})|$ of its internal state variables; *i.e.*, $\ell = k - j - 1 \mod n$.

Thus $|V(\mathcal{F}^{[j,k)})| = \ell + 1$. We may denote a level- ℓ fragment $\mathcal{F}^{[j,j+\ell+1)}$ by $\mathcal{F}^{[j,j+\ell]}$. A level- $(n-1)$ fragment $\mathcal{F}^{[j,j)}$ is obtained from R by cutting the edge S_j into two half-edges; it contains all *n* constraint codes and $n - 1$ internal state variables. A level-0 fragment $\mathcal{F}^{[j,j+1)} = \mathcal{F}^{[j,j]}$ contains one constraint code C_i and no internal state variables.

We also regard the entire realization R as a fragment, whose level is n. R contains $\ell = |E(\mathcal{R})| = n$ internal state variables, and $\ell = |V(\mathcal{R})| = n$ (not $\ell + 1$) constraint codes.

As observed in [\[2\]](#page-4-3), the set $\mathfrak{F}(\mathcal{R})$ of fragments of a tail-biting trellis realization R is partially ordered by set inclusion. The maximum fragment R includes all proper fragments $\mathcal{F}^{[j,k)}$. The partial ordering of proper fragments corresponds to the partial ordering of the circular intervals $[j, k)$ by set inclusion; *i.e.,* $\mathcal{F}^{[j',k')} \leq \mathcal{F}^{[j,k]}$ iff $[j',k') \subseteq [j,k]$. The minimal fragments are the level-0 fragments $\mathcal{F}^{[j,j+1)}$.

The partial ordering of $\mathfrak{F}(\mathcal{R})$ may be illustrated by a *Hasse diagram*, as follows. A fragment $\mathcal{F}' \in \mathfrak{F}(\mathcal{R})$ is said to be *covered* by another fragment $\mathcal{F} \in \mathfrak{F}(\mathcal{R})$ if $\mathcal{F}' < \mathcal{F}$ and there is no fragment $\mathcal{F}'' \in \mathfrak{F}(\mathcal{R})$ such that $\mathcal{F}' < \mathcal{F}'' < \mathcal{F}$ [\[10\]](#page-4-8). In our setting, \mathcal{F}' is covered by \mathcal{F} if $\mathcal{F}' < \mathcal{F}$ and the level of \mathcal{F}' is one less than the level of \mathcal{F} . The set $\mathfrak{F}(\mathcal{R})$ is thus said to be *graded* by level (number of internal state variables).

The Hasse diagram of $\mathfrak{F}(\mathcal{R})$ is illustrated in Figure [1](#page-1-0) for a tail-biting trellis realization $\mathcal R$ of length $n = 4$.

As numerous authors have observed (*e.g.,* [\[9\]](#page-4-2), [\[2\]](#page-4-3)), a conventional trellis realization may be viewed as a special case of a tail-biting trellis realization in which S_0 is trivial. Correspondingly, the Hasse diagram of a conventional trellis realization is a subdiagram of the Hasse diagram for a tailbiting trellis realization R of the same length, comprising the fragments $\{\mathcal{F} \in \mathfrak{F}(\mathcal{R}) \mid \mathcal{F} \leq \mathcal{F}^{[0,0)}\}$. By cyclic rotation of the index set \mathbb{Z}_n , any level- $(n-1)$ fragment $\mathcal{F}^{[j,j)}$ may be regarded as a conventional trellis realization.

B. Subbehaviors

For every proper fragment $\mathcal{F} = \mathcal{F}^{[j,k)} \in \mathfrak{F}(\mathcal{R})$, we define the *subbehavior* $\mathfrak{B}_{\mathcal{F}} = \mathfrak{B}^{[j,k)}$ as the set of $(a, s) \in \mathfrak{B}$ that are all-zero on or outside the boundary of F . For example, $\mathfrak{B}^{[0,0)}$ is the behavior of a conventional trellis realization of length *n*. We also define $\mathfrak{B}_R = \mathfrak{B}$.

Evidently if $\mathcal{F}' \leq \mathcal{F}$, then $\mathfrak{B}_{\mathcal{F}'} \subseteq \mathfrak{B}_{\mathcal{F}}$. Thus the set $\{\mathfrak{B}_{\mathcal{F}}, \mathcal{F} \in \mathfrak{F}(\mathcal{R})\}$ has the same partial ordering as $\mathfrak{F}(\mathcal{R})$.

For a level-0 fragment $\mathcal{F}^{[j,j]}$, we have

$$
\mathfrak{B}^{[j,j]} = \{(\mathbf{a},\mathbf{0}) \mid a_j \in (\mathcal{C}_j)_{: \mathcal{A}_j}, a_{j'} = 0 \text{ if } j' \neq j\},\
$$

where $(C_j)_{:A_j} = \{a_j \in A_j \mid (0, a_j, 0) \in C_j\}$ is the *crosssection* of C_j on A_j . As in [\[3\]](#page-4-6), $(C_j)_{A_j}$ will be denoted by $\underline{\mathcal{A}}_j$, and called the *nondynamical symbol alphabet* of \mathcal{C}_j .

C. Granules

For non-level-0 fragments, we define $\mathfrak{B}_{\leq \mathcal{F}}$ as the behavior generated by all $\mathfrak{B}_{\mathcal{F}'}$ such that $\mathcal{F}' < \mathcal{F}$, as in [\[2\]](#page-4-3). In other words, $\mathfrak{B}_{\leq \mathcal{F}} = \sum_{\mathcal{F}' \leq \mathcal{F}} \mathfrak{B}_{\mathcal{F}'}$. Evidently $\mathfrak{B}_{\leq \mathcal{F}} \subseteq \mathfrak{B}_{\mathcal{F}}$.

We define the *controller granule* Γ _{*F*} as the quotient $\mathfrak{B}_{\mathcal{F}}/\mathfrak{B}_{\leq \mathcal{F}}$. In the linear case, $\mathfrak{B}_{\mathcal{F}}$ and $\mathfrak{B}_{\leq \mathcal{F}}$ are vector spaces, and their quotient $\Gamma_{\mathcal{F}}$ is a vector space of dimension $\dim \Gamma_{\mathcal{F}} =$ $\dim \mathfrak{B}_{\mathcal{F}} - \dim \mathfrak{B}_{\leq \mathcal{F}}$. In the group case, $|\Gamma_{\mathcal{F}}| = |\mathfrak{B}_{\mathcal{F}}|/|\mathfrak{B}_{\leq \mathcal{F}}|$.

For a level-0 fragment $\mathcal{F}^{[j,j+1)}$, we define the *nondynamical granule* $\Gamma_{\mathcal{F}}$ as $\mathfrak{B}^{[j,j+1)} \cong \underline{\mathcal{A}}_j$. The set $\{\Gamma_{\mathcal{F}}, \mathcal{F} \in \mathfrak{F}(\mathcal{R})\}$ thus consists of nondynamical granules at level $\ell = 0$, and controller granules at levels $\ell > 0$.

At level *n*, where $\mathcal{F} = \mathcal{R}$, we will call $\Gamma_{\mathcal{R}} = \mathcal{B}/\mathcal{B}_{\leq \mathcal{R}}$ the *top granule* of R, or the *controllability granule* of R, since as we will see $\Gamma_{\mathcal{R}}$ governs the controllability properties of \mathcal{R} .

Note that $\mathfrak{B}_{\leq \mathcal{R}} = \sum_j \mathfrak{B}^{[j,j)}$, the behavior generated by all level- $(n-1)$ subbehaviors $\mathfrak{B}^{[j,j)}$. We will call $\mathfrak{B}_{\leq \mathcal{R}}$ the *controllable subbehavior* \mathfrak{B}^c of \mathfrak{B} .

At levels $1 \leq \ell \leq n-1$, a proper fragment $\mathcal{F}^{[j,k)}$ covers precisely two fragments, namely $\mathcal{F}^{[j,k-1)}$ and $\mathcal{F}^{[j+1,k)}$. Thus $\mathfrak{B}_{\leq \mathcal{F}(j,k)} = \mathfrak{B}^{[j,k-1)} + \mathfrak{B}^{[j+1,k)}$, and the corresponding controller granule is

$$
\Gamma^{[j,k)} = \frac{\mathfrak{B}^{[j,k)}}{\mathfrak{B}^{[j,k-1)} + \mathfrak{B}^{[j+1,k)}}.
$$

Forney and Trott [\[5\]](#page-4-4), [\[6\]](#page-4-5) define a controller granule for a conventional group trellis realization similarly as $\Gamma^{[j,k)}$ = $\mathcal{C}^{[j,k)}/(\mathcal{C}^{[j,k-1)}+\mathcal{C}^{[j+1,k)}),$ where the subcode $\mathcal{C}^{[j,k)}\subseteq\mathcal{C}$ is the set of $a \in C$ that are all-zero outside the boundary of $\mathcal{F}^{[j,k)}$. The two definitions turn out to be equivalent for minimal conventional trellis realizations.

D. ℓ*-controllable behaviors*

For $0 \leq \ell \leq n-1$, we define the ℓ -controllable behavior \mathfrak{B}_{ℓ} as the behavior generated by all level- ℓ subbehaviors $\mathfrak{B}^{[j,j+\ell]}$. In other words, $\mathfrak{B}_{\ell} = \sum_{j} \mathfrak{B}^{[j,j+\ell]}$. Note that $\mathfrak{B}_{n-1} = \mathfrak{B}^{c}$, the controllable subbehavior of \mathfrak{B} . We also define $\mathfrak{B}_n = \mathfrak{B}$. $\sum_j \mathfrak{B}^{[j,j+1)} = \underline{\mathcal{A}} \times \{0\}$, where $\underline{\mathcal{A}} = \{ \mathbf{a} \in \mathcal{A} \mid (\mathbf{a}, \mathbf{0}) \in \mathfrak{B} \} =$ Evidently $\mathfrak{B}_{\ell-1} \subseteq \mathfrak{B}_{\ell}$ for $1 \leq \ell \leq n$. Moreover, $\mathfrak{B}_0 =$ $\prod_j A_j$. We call \mathfrak{B}_0 the *nondynamical behavior* of R.

We thus have a chain of subgroups $\mathfrak{B}_0 = \underline{\mathcal{A}} \times \{0\} \subseteq$ $\mathfrak{B}_1 \subset \cdots \subset \mathfrak{B}_n = \mathfrak{B}$, which is a normal series since all groups are abelian. We denote the factor groups of this chain by $Q_{\ell} = \mathfrak{B}_{\ell}/\mathfrak{B}_{\ell-1}, 1 \leq \ell \leq n$, plus $Q_0 = \mathfrak{B}_0$.

By elementary group theory, we have $|\mathfrak{B}| = \prod_{\ell} |Q_{\ell}|$; or, in the linear case, dim $\mathfrak{B} = \sum_{\ell} \dim Q_{\ell}$. If we define sets $[Q_\ell]$ of coset representatives for the cosets of $\mathfrak{B}_{\ell-1}$ in \mathfrak{B}_{ℓ} , then every $(a, s) \in \mathfrak{B}$ may be uniquely expressed as a sum of coset representatives; or, in the linear case, if we define a basis \mathcal{B}_{ℓ} for each quotient Q_{ℓ} , then every $(a, s) \in \mathfrak{B}$ may be uniquely expressed as a linear combination of basis elements.

Since Q_{ℓ} is generated by the elements of \mathfrak{B}_{ℓ} that are not in $\mathfrak{B}_{\ell-1}$, and every element of \mathfrak{B}_{ℓ} is an element of some level- ℓ subbehavior $\mathfrak{B}^{[j,j+\ell]}$, the nonzero coset representatives in $[Q_{\ell}]$ may all be taken as elements of some $\mathfrak{B}^{[j,j+\ell]} \setminus \mathfrak{B}_{\ell-1}$. We note that if $(a, s) \in \mathfrak{B}^{[j,j+\ell]} \setminus \mathfrak{B}_{\ell-1}$, then the support of s must be precisely the length- ℓ circular interval $[i+1, j+\ell]$, else $(\mathbf{a}, \mathbf{s}) \in \mathfrak{B}_{\ell-1}$.

The level- ℓ subbehaviors $\mathfrak{B}^{[j,j+\ell]}$ thus comprise a sufficient set of representatives for Qℓ. We say that *unique factorization* holds if every element of every level- ℓ behavior \mathfrak{B}_{ℓ} is a unique sum of elements of level- ℓ subbehaviors $\mathfrak{B}^{[j,j+\ell]}$, modulo $\mathfrak{B}_{\ell-1}$; *i.e.*, if \mathfrak{B}_{ℓ} modulo $\mathfrak{B}_{\ell-1}$ is the (internal) *direct sum*

$$
\mathfrak{B}_\ell=\bigoplus_{j\in\mathbb{Z}_n}\mathfrak{B}^{[j,j+\ell]} \mod \mathfrak{B}_{\ell-1}.
$$

III. CONTROLLABILITY AND UNIQUE FACTORIZATION

In previous work [\[4\]](#page-4-7), [\[3\]](#page-4-6), we have defined controllability as the property of "having independent constraints," since we have proved that a realization is observable if and only if its dual realization has this property.

We now show that for a linear or group tail-biting trellis realization R , controllability in this sense is equivalent to the property that the top granule $\Gamma_{\mathcal{R}}$ is trivial. Simultaneously, we obtain an easy proof that unique factorization holds for R, under the proviso (as in [\[8\]](#page-4-1), [\[9\]](#page-4-2), [\[2\]](#page-4-3)) that R is *reduced*; that is, R is *state-trim*— *i.e.*, $\mathfrak{B}_{|\mathcal{S}_j} = \mathcal{S}_j$ for all j— and R is *branch-trim— i.e.*, $\mathfrak{B}_{|S_i \times A_i \times S_{i+1}} = C_j$ for all j.

(Notation: in this section, we will use notation appropriate to the group case— *i.e.,* we use sizes rather than dimensions; the reader may translate to the linear case if desired.)

A. Controllability

In [\[4\]](#page-4-7), [\[3\]](#page-4-6), a realization R is called *controllable* if the the constraints of U and the equality constraints $s = s'$ are independent. More concretely, R is controllable if the image \mathcal{S}^c of the syndrome-former homomorphism $\mathcal{U} \to \mathcal{S}$ defined by $(s, a, s') \mapsto s - s'$ is equal to S. Since the kernel of this homomorphism is the extended behavior \mathfrak{B} , we have $\mathcal{U}/\bar{\mathfrak{B}} \cong \mathcal{S}^c \subseteq \mathcal{S}$, by the FTH. This yields the following **controllability test**: $|\mathcal{U}|/|\bar{\mathfrak{B}}| \leq |\mathcal{S}|$, with equality if and only if R is controllable [\[3\]](#page-4-6). In other words, since $\mathfrak{B} \cong \overline{\mathfrak{B}}$, a realization is uncontrollable if and only if its constraints are dependent in the following sense: $¹$ $¹$ $¹$ </sup>

$$
|\mathfrak{B}| > \frac{|\mathcal{U}|}{|\mathcal{S}|} = \frac{\prod_j |\mathcal{C}_j|}{\prod_j |\mathcal{S}_j|}.
$$

¹This result may be understood as follows. Ignoring state equality constraints, there are $|\mathcal{U}| = \prod_j |\mathcal{C}_j|$ possible configurations. If the state equality constraints $\{s_j = s'_j, j \in \mathbb{Z}_n\}$ are all independent of the set of code constraints $\{C_j, j \in \mathbb{Z}_n\}$, then each state equality constraint $s_j = s'_j$ reduces the number of possible configurations by a factor of $|\mathcal{S}_j|$, so $|\mathfrak{B}| = |\mathcal{U}|/|\mathcal{S}|$, where $|\mathcal{S}| = \prod_j |\mathcal{S}_j|$. If the constraints are dependent— *i.e.*, if \mathcal{R} is not controllable— then the reduction is strictly less, and $|\mathfrak{B}| > |\mathcal{U}|/|\mathcal{S}|$.

B. Disconnected trellis realizations

We now show that if the top granule $\Gamma_{\mathcal{R}} = \mathcal{B}/\mathcal{B}^c$ is nontrivial, then \mathfrak{B} consists of $|\Gamma_{\mathcal{R}}|$ disconnected subbehaviors, namely the cosets of the controllable subbehavior \mathfrak{B}^c = $\sum_j \mathfrak{B}^{[j,j)}$ in \mathfrak{B} . Similar results were proved in [\[4\]](#page-4-7) and [\[7,](#page-4-9) Appendix A]; the proof here is simpler, and does not rely on duality.

Lemma. For a linear or group trellis realization \mathcal{R} with behavior \mathfrak{B} and controllable subbehavior \mathfrak{B}^c , for any $j \in \mathbb{Z}_n$:

(a)
$$
\mathfrak{B}_{|\mathcal{S}_j}/(\mathfrak{B}^c)|_{\mathcal{S}_j} \cong \Gamma_{\mathcal{R}};
$$

(b) $\mathfrak{B}_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}/(\mathfrak{B}^c)|_{\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}} \cong \Gamma_{\mathcal{R}}.$

Proof. (a) The projections of \mathfrak{B} and \mathfrak{B}^c onto S_i have a common kernel $\mathfrak{B}^{[j,j)} = \{(\mathbf{a}, \mathbf{s}) \in \mathfrak{B} \mid s_j = 0\}.$ Thus $\mathfrak{B}_{|\mathcal{S}_j}/(\mathfrak{B}^c)|_{\mathcal{S}_j} \cong \mathfrak{B}/\mathfrak{B}^c = \overline{\Gamma}_{\mathcal{R}}$, by the CT.

(b) The projections of \mathfrak{B} and \mathfrak{B}^c onto $S_j \times A_j \times S_{j+1}$ have a common kernel $\mathfrak{B}^{[j+1,j)} = \{(\mathbf{a}, \mathbf{s}) \in \mathfrak{B} \mid (s_j, a_j, s_{j+1}) =$ $(0, 0, 0)$ }, so (b) follows also from the CT. \Box

If R is reduced, as we assume, then $\mathfrak{B}_{|S_j} = S_j$ and $\mathfrak{B}_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}} = \mathcal{C}_j$. Moreover, we may regard \mathfrak{B}^c as the behavior of the *controllable subrealization* \mathcal{R}^c of \mathcal{R} , defined as the reduced tail-biting trellis realization with state spaces $(\mathcal{S}_j)^c = (\mathfrak{B}^c)_{|\mathcal{S}_j}$, symbol spaces \mathcal{A}_j , and constraint codes $(\mathcal{C}_j)^c = (\mathfrak{B}^c)|_{\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}$. This lemma then states that $S_j/(\mathcal{S}_j)^c \cong \Gamma_{\mathcal{R}}$ and $\mathcal{C}_j/(\mathcal{C}_j)^c \cong \Gamma_{\mathcal{R}}$.

More concretely, (a) implies that, if $\Gamma_{\mathcal{R}}$ is nontrivial, then for each j, each coset $\mathfrak{B}^c + (\mathbf{a}, \mathbf{s})$ of \mathfrak{B}^c in \mathfrak{B} passes through a distinct corresponding coset $(S_j)^c + (\mathbf{s})_j$ of $(S_j)^c$ in S_j . Similarly, \mathcal{C}_j partitions into $|\Gamma_{\mathcal{R}}|$ disjoint cosets of $(\mathcal{C}_j)^c$, each representing state transitions within one coset of \mathfrak{B}^c in \mathfrak{B} . The trellis diagram of R thus consists of $|\Gamma_{\mathcal{R}}|$ disconnected subdiagrams, one representing each coset of \mathfrak{B}^c in \mathfrak{B} . Thus for any j, j' , there is no trajectory (a, s) connecting any state s_j in a given coset of $(S_j)^c$ in S_j to a state $s_{j'}$ in a coset of $(\mathcal{S}_{j'})^c$ in $\mathcal{S}_{j'}$, unless the two cosets correspond to the same coset of \mathfrak{B}^c in \mathfrak{B} .

C. First-state chain

We now show that the controller granules of R are isomorphic to factor groups of certain normal series.

Lemma (**first-state chain**). For $j \in \mathbb{Z}_n$, $1 \leq \ell \leq n-1$,

$$
\Gamma^{[j,j+\ell]} \cong \frac{(\mathfrak{B}^{[j,j+\ell]})_{|S_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}}{(\mathfrak{B}^{[j,j+\ell)})_{|S_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}} \cong \frac{(\mathfrak{B}^{[j,j+\ell]})_{|S_{j+1}}}{(\mathfrak{B}^{[j,j+\ell)})_{|S_{j+1}}}.
$$

Proof. We have $\Gamma^{[j,j+\ell]} = \mathfrak{B}^{[j,j+\ell]}/(\mathfrak{B}^{[j,j+\ell]}+\mathfrak{B}^{(j,j+\ell]})$. The projections of $\mathfrak{B}^{[j,j+\ell]}$ and $\mathfrak{B}^{[j,j+\ell)} + \mathfrak{B}^{(j,j+\ell]}$ onto $\mathcal{S}_j \times \mathcal{A}_j \times$ S_{j+1} are $(\mathfrak{B}^{[j,j+\ell]})_{|S_j \times A_j \times S_{j+1}}$ and $(\mathfrak{B}^{[j,j+\ell)})_{|S_j \times A_j \times S_{j+1}}$, respectively, and their common kernel is $\mathfrak{B}^{(j,j+\ell)} = \{(\mathbf{a}, \mathbf{s}) \in$ $\mathfrak{B}^{[j,j+\ell]} \mid (s_j, a_j, s_{j+1}) = (0,0,0) \}.$ Similarly, the projections of $(\mathfrak{B}^{[j,j+\ell]})_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}$ and $(\mathfrak{B}^{[j,j+\ell)})_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}$ onto S_{j+1} are $(\mathfrak{B}^{[j,j+\ell]})_{|S_{j+1}}$ and $(\mathfrak{B}^{[j,j+\ell)})_{|S_{j+1}}$, respectively, and their common kernel is $(\mathfrak{B}^{[j,j]})_{|S_j \times A_j \times S_{j+1}} = \{0\} \times \underline{A}_j \times \{0\}.$ Thus both isomorphisms follow from the CT.

It follows from the first isomorphism that for each C_i there is a normal series $(\mathfrak{B}^{[j,j]})_{|S_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}} = \{0\} \times \underline{\mathcal{A}}_j \times \{0\} \subseteq$ $(\mathfrak{B}^{[j,j+1]})_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}} \subseteq \cdots \subseteq (\mathfrak{B}^{[j,j)})_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}},$ whose factor groups are isomorphic to the granules $\Gamma^{[j,j+\ell]}, 0 \leq \ell \leq$ $n - 1$. This chain implies that

$$
|(\mathfrak{B}^{[j,j)})_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}| = \prod_{\ell=0}^{n-1} |\Gamma^{[j,j+\ell]}|.
$$

This result will be useful in the next section.

It follows from the second isomorphism that for each state space S_{j+1} there is a normal series $(\mathfrak{B}^{[j,j]})_{|S_{j+1}} = \{0\} \subseteq$ $(\mathfrak{B}^{[j,j+1]})_{| \mathcal{S}_{j+1}} \subseteq \cdots \subseteq (\mathfrak{B}^{[j,j)})_{| \mathcal{S}_{j+1}},$ whose factor groups are isomorphic to the granules $\Gamma^{[j,j+\ell]}, 1 \leq \ell \leq n-1$. We call this normal series the *first-state chain* at S_{i+1} , since S_{i+1} is the first possibly nonzero state in the trajectories in $\mathfrak{B}^{[j,j+\ell]}, 1 \leq$ $\ell \leq n-1$. This chain implies that

$$
|(\mathfrak{B}^{[j,j)})_{|S_{j+1}}| = \prod_{\ell=1}^{n-1} |\Gamma^{[j,j+\ell]}|.
$$

D. Controllability and unique factorization

We will now show that R is controllable if and only if $\mathfrak{B} = \mathfrak{B}^c$; *i.e.*, if and only if the top granule $\Gamma_{\mathcal{R}}$ is trivial. Moreover, the controller granule decomposition gives a unique factorization of both \mathfrak{B}^c and \mathfrak{B} .

We first state a technical lemma that shows that in the controllable subrealization \mathcal{R}^c , the number of transitions $(s_j, a_j, s_{j+1}) \in (C_j)^c$ is the number of states $s_j \in (S_j)^c$ times the number of transitions $(0, a_j, s_{j+1}) \in (\mathfrak{B}^{[j,j)})_{|S_j \times A_j \times S_{j+1}}$. **Lemma**. For all j, $|(\mathcal{C}_j)^c| = |(\mathcal{S}_j)^c| \cdot |(\mathfrak{B}^{[j,j)})_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}|$. *Proof.* The projection of \mathfrak{B}^c on \mathcal{S}_j is $(\mathcal{S}_j)^c$, and its kernel

is $\mathfrak{B}^{[j,j)}$, so $(S_j)^c \cong \mathfrak{B}^c / \mathfrak{B}^{[j,j)}$ by the FTH. The projections of \mathfrak{B}^c and $\mathfrak{B}^{[j,j)}$ on $\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}$ are $(\mathcal{C}_j)^c$ and $(\mathfrak{B}^{[j,j)})_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}$, respectively, and $\mathfrak{B}^{[j,j+1)}$ is their common kernel, so $\mathfrak{B}^{c}/\mathfrak{B}^{[j,j)} \cong (\mathcal{C}_j)^c/(\mathfrak{B}^{[j,j)})_{|S_j \times A_j \times S_{j+1}}$ by the CT.

Next, we define P^c as the product of the sizes of all controller granules up to level $n - 1$, *i.e.*, P^c \prod controller granules up to level $n-1$, *i.e.*, $P^c =$
 $n-1$ $\prod_{i=1}^n |r_i|$ and P $\prod_{i=1}^n |P^c_i|$ as the product of $\prod_{\ell=0}^{n-1} \prod_{j \in \mathbb{Z}_n} |\Gamma^{[j,j+\ell]}|$, and $P = |\Gamma_{\mathcal{R}}| P^c$ as the product of the sizes of all controller granules. We observe that since P is the number of possible sums of granule representatives, we have $|\mathfrak{B}| \leq P$, with equality if and only if unique factorization holds for B. Similarly, we have $|\mathfrak{B}^c| \leq P^c$, with equality if and only if unique factorization holds for \mathfrak{B}^c .

Theorem (**controllability and unique factorization**). Let B and \mathfrak{B}^c be the behaviors of a reduced linear or group tailbiting trellis realization R and its controllable subrealization \mathcal{R}^c , respectively. Then:

- (a) \mathcal{R}^c is controllable.
- (b) Unique factorization holds for \mathfrak{B}^c ; *i.e.*, $|\mathfrak{B}^c| = P^c$.

(c) R is controllable if and only if $\mathfrak{B} = \mathfrak{B}^c$; *i.e.*, iff the top granule $\Gamma_{\mathcal{R}}$ is trivial.

(d) Unique factorization holds for \mathfrak{B} ; *i.e.*, $|\mathfrak{B}| = P$.

Proof. (a-b) From the previous lemma, $\prod_j |(\mathcal{C}_j)^c|$ = $(\prod_j |(\mathcal{S}_j)^c|)(\prod_j |(\mathfrak{B}^{[j,j)})_{|\mathcal{S}_j \times \mathcal{A}_j \times \mathcal{S}_{j+1}}|)$. By Section [III-C,](#page-2-2) we have $|(\mathfrak{B}^{[j,j)})_{|S_j \times A_j \times S_{j+1}}| = \prod_{\ell=0}^{n-1} |\Gamma^{[j,j+\ell]}|,$ so $(\prod_j |(\mathcal{C}_j)^c|)/(\prod_j |(\mathcal{S}_j)^c|) = \prod_j \prod_{\ell=0}^{n-1} |\Gamma^{[j,j+\ell]}| = P^c$, the product of the sizes of all proper controller granules $\Gamma^{[j,j+\ell]}$. Therefore, by our controllability test, we have $|\mathfrak{B}^c| \geq P^c$, with equality if and only if \mathcal{R}^c is controllable. On the other hand, in view of the controller granule decomposition of \mathfrak{B}^c , we have $|\mathfrak{B}^c| \leq P^c$, with equality if and only if unique factorization holds for \mathfrak{B}^c . Thus $|\mathfrak{B}^c| = P^c$, \mathcal{R}^c is controllable, and unique factorization holds for \mathfrak{B}^c .

(c) By Section [III-B,](#page-2-3) \mathfrak{B} is the disjoint union of $|\Gamma_{\mathcal{R}}|$ disconnected cosets of \mathfrak{B}^c . Thus we have $|\mathfrak{B}| = |\Gamma_{\mathcal{R}}||\mathfrak{B}^c|$, $|C_j| = |\Gamma_{\mathcal{R}}||(C_j)^c|$, and $|S_j| = |\Gamma_{\mathcal{R}}||(S_j)^c|$. Therefore $\left(\prod_j|\mathcal{C}_j|\right)/\left(\prod_j|\mathcal{S}_j|\right) = P^c = |\mathfrak{B}^c| = |\mathfrak{B}|/|\Gamma_{\mathcal{R}}|$. By our controllability test, R is controllable if and only if $|\Gamma_{\mathcal{R}}| = 1$.

(d) By Section [III-B,](#page-2-3) every element of $\mathfrak B$ is uniquely expressible as the sum of an element of \mathfrak{B}^c and a coset representative in $[\Gamma_{\mathcal{R}}]$, so since unique factorization holds for \mathfrak{B}^c , it holds also for \mathfrak{B} . \Box

E. State space and constraint code sizes

Unique factorization of $\mathfrak B$ implies unique factorization of $\mathfrak{B}_{\mathcal{F}}$ for any fragment $\mathcal{F} \leq \mathcal{R}$. It follows that the size of each state space S_i and each constraint code C_i may be determined in terms of granule sizes as follows:

Corollary (**state space and constraint code sizes**). If \mathcal{R} is a reduced linear or group tail-biting trellis realization with state spaces S_i and constraint codes C_i , then:

(a)
$$
S_j \cong \mathfrak{B}/\mathfrak{B}^{[j,j)}
$$
, and

$$
|S_j| = \prod
$$

$$
|\mathcal{S}_j| = \prod_{\mathcal{F} \leq \mathcal{R}: |\mathcal{S}_j \in E(\mathcal{F})} |\Gamma_{\mathcal{F}}|;
$$

(b) $C_j \cong \mathfrak{B}/\mathfrak{B}^{[j+1,j)}$, and

$$
|\mathcal{C}_j| = \prod_{\mathcal{F} \leq \mathcal{R}: \ \mathcal{C}_j \in V(\mathcal{F})} |\Gamma_{\mathcal{F}}|.
$$

Proof. (a) If R is state-trim at S_j , then $S_j = \mathfrak{B}_{|S_j}$. Moreover, the kernel of the projection of \mathfrak{B} onto \mathcal{S}_j is $\mathfrak{B}^{[j,j)}$. Thus $\mathcal{S}_j \cong \mathfrak{B}/\mathfrak{B}^{[j,j)}$ by the FTH, so $|\mathcal{S}_j| =$ $|\mathfrak{B}|/|\mathfrak{B}^{[j,j)}| = |P/\prod_{\mathcal{F}\leq \mathcal{F}^{[j,j)}}|\Gamma_{\mathcal{F}}| = |\prod_{\mathcal{F}\nleq \mathcal{F}^{[j,j)}}|\Gamma_{\mathcal{F}}| =$ $\prod_{\mathcal{F}\leq \mathcal{R}|\mathcal{S}_j\in E(\mathcal{F})}| \Gamma_{\mathcal{F}}|,$ since $\mathcal{F}\leq \mathcal{F}^{[j,j)}$ iff $\mathcal{S}_j \notin E(\mathcal{F})$.

(b) If R is branch-trim at C_j , then $C_j = \mathfrak{B}_{|S_j \times A_j \times S_{j+1}}$. Moreover, the kernel of the projection of \mathfrak{B} onto \mathcal{C}_j is $\mathfrak{B}^{[j+1,j)}$. Thus $\mathcal{C}_j \cong \mathfrak{B}/\mathfrak{B}^{[j+1,j)}$ by the FTH, so $|C_j| = |\mathfrak{B}|/|\mathfrak{B}^{[j+1,j)}| = P/\prod$ \prod $_{\mathcal{F}\leq \mathcal{F}^{\left(j+1,j\right) }}\left\vert \Gamma_{\mathcal{F}}\right\vert \quad =% \mathcal{F}^{\left(j+1\right) }\left\vert \mathcal{F}\right\vert \quad \text{and}\quad \mathcal{F}^{\left(j\right) }=\mathcal{F}^{\left(j\right) }%$ $\mathcal{F}_{\cancel{\mathcal{I}}\mathcal{F}}[\mathcal{F}_{\mathcal{F}}] = \prod_{\mathcal{F}\leq \mathcal{R}|\mathcal{C}_{\cancel{j}}\in V(\mathcal{F})} |\Gamma_{\mathcal{F}}|, \text{ since } \mathcal{F} \leq$ $\mathcal{F}^{[j+1,j)}$ iff $\mathcal{C}_j \notin V(\mathcal{F})$.

In other words, assuming trimness, S_i factors into components isomorphic to those granules $\Gamma_{\mathcal{F}}$ such that $\mathcal{S}_i \in E(\mathcal{F})$ (*i.e.*, S_j is "active" during $\mathcal F$). Also, $\mathcal C_j$ factors into components isomorphic to those granules $\Gamma_{\mathcal{F}}$ such that $\mathcal{C}_i \in V(\mathcal{F})$ (*i.e.*, C_i is "active" during \mathcal{F}).

F. Controller canonical realization

The unique factorization result of Section [III-D](#page-3-0) implies that every reduced linear or group trellis realization is equivalent to a *controller canonical realization*, which we define as follows.

For each $\mathcal{F} \leq \mathcal{R}$, we have a one-to-one map $\Gamma_{\mathcal{F}} \to [\Gamma_{\mathcal{F}}]$ from the granule Γ_F to the set of coset representatives $[\Gamma_F] =$ $[\mathfrak{B}_{\mathcal{F}}/\mathfrak{B}_{\leq \mathcal{F}}]$. We may thus map each element of the Cartesian product $\prod_{\mathcal{F}\leq \mathcal{R}}\Gamma_\mathcal{F}$ to the sum $(\mathbf{a}, \mathbf{s}) = \sum_{\mathcal{F}\leq \mathcal{R}}(\mathbf{a}_\mathcal{F}, \mathbf{s}_\mathcal{F})$ of the corresponding coset representatives $(a_{\mathcal{F}}, s_{\mathcal{F}}) \in [\Gamma_{\mathcal{F}}]$, which is an element of $\mathfrak B$ since each coset representative is an element of B. By unique factorization, the map so defined from $\prod_{\mathcal{F}\leq \mathcal{R}} \Gamma_{\mathcal{F}}$ to \mathfrak{B} is one-to-one.

More concretely, the map $\prod_{\mathcal{F}\leq \mathcal{R}} \Gamma_{\mathcal{F}} \to \mathfrak{B}$ may be implemented as follows. We generate the trajectories in $[\Gamma_{\mathcal{F}}]$ by an *atomic trellis realization* whose state spaces S_i are equal to $\Gamma_{\mathcal{F}}$ when $\mathcal{S}_j \in E(\mathcal{F})$, and trivial otherwise. An element of $\Gamma_{\mathcal{F}}$ determines the state value $(s_{\mathcal{F}})_j$ when $\mathcal{S}_j \in E(\mathcal{F})$, and the symbol value $(a_{\mathcal{F}})_j$ when $\mathcal{C}_j \in V(\mathcal{F})$. The state value s_j is thus the sum $\sum_{\mathcal{F}\leq \mathcal{R}|\mathcal{S}_j\in E(\mathcal{F})} (\mathbf{s}_{\mathcal{F}})_j$, and the symbol value \mathbf{a}_j is the sum $\sum_{\mathcal{F} \leq \mathcal{R} | \mathcal{C}_j \in V(\mathcal{F})} (\mathbf{a}_{\mathcal{F}})_j$. The size of the aggregate state space S_j is thus $|S_j| = \prod_{\mathcal{F} \leq \mathcal{R} | S_j \in E(\mathcal{F})} |\Gamma_{\mathcal{F}}|$, as in our state space size result. Thus the controller canonical realization is a minimal realization of B. (We can also show that the number of possible transitions (s_j, a_j, s_{j+1}) is $\prod_{\mathcal{F}\leq \mathcal{R}|\mathcal{C}_j\in V(\mathcal{F})}|\Gamma_{\mathcal{F}}|$, as in our constraint code size result.)

If $\mathfrak B$ is linear, then the controller canonical realization of B is easily seen to be linear. However, for a group realization R, although the map $\prod_{\mathcal{F}\leq R} \Gamma_{\mathcal{F}} \rightarrow \mathfrak{B}$ yields a one-to-one, group-theoretic, and minimal realization of \mathfrak{B} , it may well not be isomorphic, even when R is conventional [\[5\]](#page-4-4). This issue was raised in [\[2,](#page-4-3) Remark III.3] via the following example, in which the controller canonical realization is nonhomomorphic.

Example (Conventional group trellis realization over \mathbb{Z}_4). Let R be a conventional group trellis realization of length 3 with behavior $\mathfrak{B} = \langle (112, 0120) \rangle \subseteq (\mathbb{Z}_4)^3 \times (\mathbb{Z}_4)^4$; *i.e.*, $\mathfrak{B} = \{(000, 0000), (112, 0120), (220, 0200), (332, 0320)\}\approx$ \mathbb{Z}_4 . Its ℓ -controllable subbehaviors are $\mathfrak{B}_0 = \{(000, 0000)\};$ $\mathfrak{B}_{1} \; = \; \mathfrak{B}^{[0,1]} \; = \; \{(000,0000), (220,0200)\} \; \cong \; 2\mathbb{Z}_{4} \; \cong \; \mathbb{Z}_{2};$ and $\mathfrak{B}_2 = \mathfrak{B} \cong \mathbb{Z}_4$. Its nontrivial controller granules are $\Gamma^{[0,1]} = \mathfrak{B}^{[0,1]} \cong \mathbb{Z}_2$, which is realized by a 2-state atomic trellis realization that is active during [0, 1], and $\Gamma^{[0,2]}$ = $\mathfrak{B}/\mathfrak{B}^{[0,1]} \cong \mathbb{Z}_4/2\mathbb{Z}_4 \cong \mathbb{Z}_2$, which is realized by a 2-state atomic trellis realization that is active during [0, 2].

Figure [2](#page-4-10) depicts the controller canonical realization of β via trellis diagrams for the atomic trellis realizations of $\Gamma^{[0,1]}$ = $\mathfrak{B}^{[0,1]}$ and $[\Gamma^{[0,2]}] = [\mathfrak{B}/\mathfrak{B}^{[0,1]}]$, plus a trellis diagram for \mathfrak{B} .

IV. CONCLUSION

We have generalized the CBFT to group trellis realizations, with a proof based on a controller granule decomposition of B and our controllability test for general group realizations.

It would be natural to dualize these results, using a dual observer granule decomposition. However, as discussed in [\[6\]](#page-4-5), such a dualization is not straightforward, even for minimal conventional trellis realizations. Developing a nice dual observer granule decomposition for linear and group tail-biting trellis realizations is a good goal for future research.

It would be nice also to extend these results to non-trellis realizations. However, it is known (see [\[4,](#page-4-7) Appendix A]) that unique factorization generally does not hold for nontrellis linear or group realizations, even simple cycle-free realizations. New ideas will therefore be needed.

Finally, we would like ultimately to redevelop all of the principal results of classical discrete-time linear systems theory using a purely group-theoretic approach. However, the classical theory generally assumes an infinite time axis. One possible approach would be to regard a time-invariant or periodically time-varying linear or group system on an infinite time axis as the "limit" of a sequence of covers of a linear or group tail-biting trellis realization on a sequence of finite time axes of increasing length. Such an approach would hopefully be purely algebraic, and thus might avoid the subtle topological issues discussed in [\[6\]](#page-4-5).

ACKNOWLEDGMENTS

I am grateful to Mitchell Trott and Heide Gluesing-Luerssen for our earlier collaborations, in which many of these ideas first arose, and to David Conti, for early access to and discussion of the results of [\[2\]](#page-4-3).

REFERENCES

- [1] A. R. Calderbank, G. D. Forney, Jr. and A. Vardy, "Minimal tail-biting trellises: The Golay code and more," *IEEE Trans. Inf. Theory*, vol. 45, pp. 1435–1455, July 1999.
- [2] D. Conti and N. Boston, "On the algebraic structure of linear tail-biting trellises," *IEEE Trans. Inf. Theory*, to appear, 2015. ArXiv: 1402.6404.
- [3] G. D. Forney, Jr., "Codes on graphs: Fundamentals," *IEEE Trans. Inf. Theory*, vol. 60, pp. 5809–5826, Oct. 2014.
- [4] G. D. Forney, Jr. and H. Gluesing-Luerssen, "Codes on graphs: Observability, controllability and local reducibility," *IEEE Trans. Inf. Theory*, vol. 59, pp. 223–238, Jan. 2013.
- [5] G. D. Forney, Jr. and M. D. Trott, "The dynamics of group codes: State spaces, trellis diagrams and canonical encoders," *IEEE Trans. Inf. Theory*, vol. 39, pp. 1491–1513, Sept. 1993.
- [6] G. D. Forney, Jr. and M. D. Trott, "The dynamics of group codes: Dual abelian group codes and systems," *IEEE Trans. Inf. Theory*, vol. 50, pp. 2935–2965, Dec. 2004.
- [7] H. Gluesing-Luerssen and G. D. Forney, Jr., "Local irreducibility of tailbiting trellises," *IEEE Trans. Inf. Theory*, vol. 59, pp. 6597–6610, Oct. 2013.
- [8] R. Koetter and A. Vardy, "On the theory of linear trellises," in *Information, Coding and Mathematics* (M. Blaum, P. G. Farrell, H. C. A. van Tilborg, eds.), pp. 323–354. Boston, MA: Kluwer, 2002.
- [9] R. Koetter and A. Vardy, "The structure of tail-biting trellises: Minimality and basic principles," *IEEE Trans. Inf. Theory*, vol. 49, pp. 2081– 2105, Sept. 2003.
- [10] R. P. Stanley, *Enumerative Combinatorics*, vol. 1, 2d ed. Cambridge, UK: Cambridge U. Press, 2012.