Generalized solutions of the multidimensional stochastic Burgers equation.

P. Catuogno,^{*} J.F. Colombeau [†] C. Olivera[‡]

Key words: Stochastic Burgers equation, Generalized functions, Multiplication of distributions, Generalized solutions.

MSC2000 subject classification: 60H15, 35R60, 46F99.

Abstract

We introduce a new concepts of weak solution for the conservative stochastic Burgers equation in any dimension. The definition is based on weak solution concepts introduced by various authors in order to make sense of equations which do not solutions in the sense of distributions. In one dimension the solution reduces to the classical distributional solution of the 1–D stochastic Burgers equation.

1 Introduction

The aim of this paper is to study the existence of solution to the multidimensional stochastic Burgers equation in \mathbb{R}^d of the following form:

[∗]Departamento de Matem´atica, Universidade Estadual de Campinas, Brazil. E-mail: pedrojc@ime.unicamp.br.

[†]Institut Fourier, Université de Grenoble., France. E-mail: jf.colombeau@wanadoo.fr.

[‡]Departamento de Matemática, Universidade Estadual de Campinas, Brazil. E-mail: colivera@ime.unicamp.br.

$$
\begin{cases}\n\partial_t U(t,x) = \Delta_x U(t,x) + \nabla_x ||U(t,x)||^2 + \nabla_x \dot{W}(t,x), \\
U(0,x) = \nabla_x f(x).\n\end{cases} (1)
$$

where $W(t, x)$ is a space–time white noise.

During the past few decades, the stochastic Burgers equation has found applications in diverse fields ranging from statistical physics, cosmology to fluid dynamics. The problem of Burgers turbulence, that is the study of the solutions of the Burgers equation with random initial conditions or random forcing is a central issue in the study of nonlinear systems out of equilibrium. See J. Bec, K. Khanin [\[4\]](#page-11-0) and E. Weinan [\[28\]](#page-13-0).

The multidimensional conservative stochastic Burgers equation has been studied by several authors in the case $W(t, x)$ is a noise white in time and correlated in space. See A. Dermoune [\[13\]](#page-12-0), Z. Brezniak, B. Goldys, M. Neklyudov [\[8\]](#page-11-1) and R. Iturriaga, K. Khanin [\[19\]](#page-12-1). In the case that the nonlinear term is interpreted in the sense of Wick calculus the stochastic Burgers equation has been well studied, see S. Assing [\[3\]](#page-11-2) and also H. Holden, T. Lindstrøm, B. Øksendal, J. Ubøe, T. Zhang [\[17\]](#page-12-2) and [\[18\]](#page-12-3). They showed existence and uniqueness results for the solution regarded as a stochastic process with values in a Kondratiev space of stochastic distributions.

Up to our knowledge there is no proof of existence of solution in any sense for equation [\(1\)](#page-1-0) when $d \geq 2$. In the case that $d = 1$, the problem of existence of solutions for stochastic Burgers equation is well understood, see L. Bertini, N. Cancrini, G. Jona–Lasinio [\[6\]](#page-11-3) , G. Da Prato, A. Debussche, R. Teman [\[12\]](#page-12-4), P. Catuogno, C. Olivera [\[9\]](#page-11-4) and M. Gubinelli, N.Perkowski [\[15\]](#page-12-5).

A main difficulty with the multidimensional stochastic Burgers equation is that the solutions take values in a distributional space. Therefore, it is nec-

essary to give meaning to the non-linear term $\nabla_x ||U(t, x)||^2$. In the onedimensional case the Cole–Hopf transformation is used to sort out this problem, but this does not make sense in the multidimensional case since the solution of the stochastic heat equation is not a standard stochastic process [\[27\]](#page-13-1).

We remark that it is believed that the Cole–Hopf solution $U(t, x) := \nabla_x \log Z(t, x)$, where $Z(t, x)$ denotes the solution of the stochastic heat equation with multiplicative noise, is the correct physical solution of [\(1\)](#page-1-0), see [\[6\]](#page-11-3) and [\[7\]](#page-11-5). We mention also that the Cole–Hopf solution is related to the KPZ equation, see $|16|$.

We introduce a new concept of solution in order to give sense to this expected solution of the multidimensional stochastic Burgers equation. This new concept of solution is strongly inspired by previous works of many authors using generalized solutions for equations of mathematical physics in cases in that the expected or known (for instance from numerical investigations) solutions do not make sense within distribution theory: [\[10,](#page-11-6) [11,](#page-12-7) [14,](#page-12-8) [21,](#page-13-2) [24\]](#page-13-3). In particular such generalized solutions have been considered for stochastic equations in [\[1\]](#page-11-7), [\[23\]](#page-13-4), [\[22\]](#page-13-5).

We denote by $\mathcal{D}((0,T) \times \mathbb{R}^d; \mathbb{R}^d)$ the space of the infinitely differentiable functions with compact support in $(0, T) \times \mathbb{R}^d$ and values in \mathbb{R}^d . Let us denote by \mathcal{L}_B the Burgers operator,

$$
\mathcal{L}_B \varphi = \partial_t \varphi - \Delta_x \varphi - \nabla_x ||\varphi||^2
$$

for $\varphi \in \mathcal{D}((0,T) \times \mathbb{R}^d; \mathbb{R}^d)$.

Now, we introduce the concept of weak solution for the stochastic Burgers equation.

Definition 1.1 We say that a sequence (U_n) of smooth semimartingales is an \mathbb{R}^d -weak generalized solution of the equation [\(1\)](#page-1-0) if

1. For each $\varphi \in \mathcal{D}((0,T) \times \mathbb{R}^d; \mathbb{R}^d)$,

$$
\lim_{n\to\infty}(\mathcal{L}_BU_n,\varphi)=(\nabla\dot{W},\varphi)_{\mathcal{D}'},
$$

where the convergence is in $L^2(\Omega)$.

2. $U_n(0, x) = \nabla f(x)$.

The above means that, after averaging on any smooth test function, the sequence (U_n) tends to satisfy the equation when $n \to \infty$. In general we ignore the nature of the possible limit of the sequence (U_n) : this limit can be a distribution which is not necessarily a solution of the equation in the sense of distribution theory, such as in the case of shock waves for nonconservative systems [\[2,](#page-11-8) [11\]](#page-12-7), in the case of delta waves in general and a fortiori delta–prime wave [\[24\]](#page-13-3), or it can be an object which is not a distribution, such as objects put in evidence in general relativity in [\[26\]](#page-13-6). In one dimension the limit of the sequence (U_n) that we construct exists as a distribution which is a solution in the sense of distributions.

We shall show that there exists a weak generalized solution for the multi– dimensional stochastic Burgers equation, which is a type of Cole–Hopf solution.

This paper is organized as follows. In the next section we review some facts on space–time white noise. In section 3 we show the existence of such a weak solution for the stochastic Burgers equation [\(1\)](#page-1-0). Then we recover the stronger result of existence of a distributional solution for the stochastic Burgers equation in dimension 1.

2 Preliminaries

We say that a random field $\{S(t,x) : t \in [0,T], x \in \mathbb{R}^d\}$ is a spatially dependent semimartingale if for each $x \in \mathbb{R}^d$, $\{S(t, x) : t \in [0, T]\}$ is a \mathbb{R}^d valued semimartingale in relation to the same filtration $\{\mathcal{F}_t : t \in [0,T]\}.$ If $S(t, x)$ is a C^{∞} -function of x and continuous in t almost surely (i.e., for almost all ω), it is called a smooth semimartingale. See [\[20\]](#page-12-9) for a rigorous study of spatially depend semimartingales and applications to stochastic differential equations.

A distribution valued Gaussian process with mean zero $\{W(t, x) : t \in [0, T], x \in$ \mathbb{R}^d is a space–time white noise if

$$
\mathbb{E}(\dot{W}(t,x)\dot{W}(s,y)) = \delta(t-s)\delta(x-y).
$$

More precisely :

- For any $\xi \in \mathcal{D}((0,T) \times \mathbb{R}^d)$ the random variable $\dot{W}(\xi)$ is Gaussian variable with mean zero.
- For any $\xi, \rho \in \mathcal{D}((0,T) \times \mathbb{R}^d)$ the random variables $\dot{W}(\xi), \dot{W}(\rho)$ have covariance

$$
\mathbb{E}(\dot{W}(\xi)\dot{W}(\rho)) = \int_{[0,T]\times\mathbb{R}^d} \xi(t,x) \cdot \rho(t,x) dx dt.
$$

It is not difficult to construct a space time–white noise. In fact, let $\{f_j:$ $j \in \mathbb{N}$ be a complete orthonormal basis of $L^2([0,T] \times \mathbb{R}^d)$ and $\{Z_j : j \in \mathbb{N}\}\$ be a family of independent Gaussian random variables with mean zero and variance one. Then

$$
\dot{W}(t,x) := \sum_{j=1}^{\infty} f_j(t,x) Z_j \tag{2}
$$

is a space–time white noise, where the action is defined as

$$
(\dot{W}, \xi)_{\mathcal{D}'} = \sum_{j=1}^{\infty} (\xi, f_j) Z_j.
$$

It is well know that the last action can be extended to $\xi \in L^2([0,T] \times \mathbb{R}^d)$ using Itô isometry.

The cylindrical Wiener process $\{W_t: L^2(\mathbb{R}^d) \to L^2(\Omega) : t \in [0,T]\}$ associated to \dot{W} is given by

$$
W_t(\varphi) := (\dot{W}, \varphi 1_{[0,t]}).
$$

It is clear that $W_t(\varphi)$ is a Brownian motion with variance $t\|\varphi\|^2$ for each $\varphi \in L^2(\mathbb{R}^d).$

We say that a sequence of smooth semimartingales $(Wⁿ)$ is a weak approxi*mation* to the cylindrical Wiener W if for all $\varphi \in L^2(\mathbb{R}^d)$,

$$
\lim_{n \to \infty} \int_{\mathbb{R}^d} \varphi(x) W_t^n(x) dx = W_t(\varphi), \tag{3}
$$

where the convergence is in $C([0,T], L^2(\Omega))$.

A weak approximation (W^n) to the cylindrical Wiener process W is good if for each $n \in \mathbb{N}$, $W_t^n(x)$ is a Brownian motion with quadratic variation

$$
\langle W^n(x) \rangle_t = C_n \cdot t,
$$

for all $x \in \mathbb{R}^d$ and where C_n is a constant depending on n.

A main ingredient in our approach for solving the Burgers equation is the use of regularization techniques for the white noise with respect to space, we refer the reader to [\[7\]](#page-11-5) and [\[21\]](#page-13-2) to background material.

Let $\rho : \mathbb{R}^d \to [0, \infty)$ be an infinitely differentiable symmetric function with compact support such that $\int_{\mathbb{R}^d} \rho(x) dx = 1$. We will consider the mollifiers $\rho_n(x) = n^d \rho(nx)$, with $n \in \mathbb{N}$. The regularization by ρ of the space-time white noise \dot{W} , denoted by \dot{W}_{ρ_n} , are defined to be

$$
\dot{W}_{\rho_n}(t,x) := \rho_n * \dot{W}(t,x).
$$

We observe that \dot{W}_{ρ_n} is white in time and colored in space, in fact we have that $\dot{W}_{\rho_n}(t,x)$ is a distribution valued Gaussian process with mean zero and covariance,

$$
\mathbb{E}(\dot{W}_{\rho_n}(t,x)\dot{W}_{\rho_n}(s,y)) = \delta(t-s)h_n(x-y)
$$

where $h_n : \mathbb{R}^d \to \mathbb{R}$ is given by $h_n(z) = \int_{\mathbb{R}^d} \rho_n(u) \rho_n(u+z) du$. In terms of the expansion [\(2\)](#page-4-0) we have that

$$
\dot{W}_{\rho_n}(t,x) := \sum_{j=1}^{\infty} \rho_n * f_j(t,x) Z_j.
$$

The mollified cylindrical Wiener process $W_t^{\rho_n}$ $t^{\rho_n}(x)$ associated with the spacetime white noise $W(t, x)$ is defined by

$$
W_t^{\rho_n}(x) := W_t(\rho_n(x - \cdot)).\tag{4}
$$

The distributional time derivative of $W_t^{\rho_n}$ $\dot{W}_{\rho_n}(t,x)$ is $\dot{W}_{\rho_n}(t,x)$. We have that $W_t^{\rho_n}$ $t^{\rho_n}(x)$ is a Brownian motion with quadratic variation,

$$
\langle W^{\rho_n}(x) \rangle_t = \|\rho\|^2 n^d \cdot t. \tag{5}
$$

In particular we have proved the existence of good weak approximations to cylindrical Wiener processes.

Proposition 2.1 (W^{ρ_n}) is a good weak approximation to the cylindrical Wiener process W.

3 Existence of weak solution in any dimension

Let $(Wⁿ)$ be a good weak approximation to the cylindrical Wiener W. We will denote by $H_n(t, x)$ the process $\log Z_n(t, x)$, where Z_n is the solution of the regularized stochastic heat equation in the Itô sense

$$
\begin{cases}\n dZ_n &= \Delta_x Z_n \, dt + Z_n \, dW^n, \\
 Z_n(0, x) &= e^{f(x)}.\n\end{cases} \tag{6}
$$

We observe that $W^n(t, x)$ is C^{∞} -martingale then we follow the existence and uniqueness of solution of the classical theory, see for instance [\[20\]](#page-12-9). Applying the Feynman–Kac formula we obtain the following representation of the solution to equation [\(6\)](#page-6-0) with $f \in C_b^{\infty}$ $b^{\infty}(\mathbb{R}^d)$,

$$
Z_n(t,x) = \tilde{\mathbb{E}}(e^{f(x+B_t)}e^{\int_0^t W^n(t-s,x+B_s)ds})
$$
\n(7)

where B_t is a Brownian motion independent of W^n such that $B_0 = 0$ defined in an auxiliary probability space, see [\[13\]](#page-12-0).

The sequence of smooth semimartingales (U_n) given by $U_n := \nabla_x H_n$ is called the Cole-Hopf sequence associated to $(Wⁿ)$.

Theorem 3.1 Let $f \in C_b^{\infty}$ $\mathcal{E}_b^{\infty}(\mathbb{R}^d)$. The Cole-Hopf sequence associated to $(Wⁿ)$ is a weak generalized solution of the stochastic Burgers equation [\(1\)](#page-1-0).

Proof: Let (U_n) be the Cole–Hopf sequence given by $U_n(t, x) := \nabla_x H_n(t, x) =$ $\nabla_x \log Z_n(t, x)$. Since Z_n satisfies the equation [\(6\)](#page-6-0) it is a C^{∞} -semimartingale. Then applying the classical Itô formula we have that

$$
dH_n = \frac{dZ_n}{Z_n} - \frac{d\langle Z_n \rangle}{2Z_n^2}
$$

=
$$
\frac{\Delta_x Z_n dt + Z_n dW^n}{Z_n} - \frac{Z_n^2 C_n dt}{2Z_n^2}
$$

=
$$
\frac{\Delta_x Z_n dt}{Z_n} + dW^n - \frac{C_n dt}{2}.
$$

It is easy to check that

$$
\frac{\Delta_x Z_n}{Z_n} = \Delta_x H_n + \|\nabla_x H_n\|^2.
$$

Combining the above equations we obtain

$$
dH_n = (\Delta_x H_n + \|\nabla_x H_n\|^2)dt + dW^n - \frac{C_n dt}{2},
$$
\n(8)

that is

$$
H_n(t,x) = f(x) + \int_0^t (\Delta_x H_n(s,x) + ||\nabla_x H_n(s,x)||^2) ds + W_t^n(x) - \frac{C_n t}{2}.
$$

Let $\varphi = (\varphi_1, \dots, \varphi_d) \in \mathcal{D}((0, T) \times \mathbb{R}^d; \mathbb{R}^d)$. Multiplying [\(8\)](#page-7-0) by $\partial_t \partial_{x_i} \varphi_i(t, x)$ and integrating on $[0, T] \times \mathbb{R}^d$, we obtain that

$$
\int_{[0,T]\times\mathbb{R}^d} H_n \, \partial_t \partial_{x_i} \varphi_i dx dt = - \int_{[0,T]\times\mathbb{R}^d} (\Delta_x H_n + \|\nabla_x H_n(s,x)\|^2) \partial_{x_i} \varphi_i dx dt \n- \int_{[0,T]\times\mathbb{R}^d} \partial_t \partial_{x_i} \varphi_i(t,x) dW_t^n(x) dx.
$$

Thus

$$
(\mathcal{L}_B U_n, \varphi) = \int_{[0,T] \times \mathbb{R}^d} \nabla_x \partial_t \varphi(t, x) dW_t^n(x) dx \tag{9}
$$

Then we conclude

$$
\lim_{n \to \infty} (\mathcal{L}_B U_n, \varphi) = (\nabla \dot{W}, \varphi)_{\mathcal{D}'},\tag{10}
$$

where the convergence is in $L^2(\Omega)$. \square

Remark 3.1 We observe that by definition of the solution we have

$$
\lim_{n\to\infty}(\mathcal{L}_B U_n,\varphi)=(\nabla \dot{W},\varphi)_{\mathcal{D}'}\,\,\text{in}\,\,L^2(\Omega).
$$

Via classical arguments, taking a dense and countable set of D , we can conclude that there exists a subsequence of U_n such that

$$
\lim_{n\to\infty}(\mathcal{L}_BU_n,\varphi)=(\nabla\dot{W},\varphi)_{\mathcal{D}'}
$$

where the convergence is almost surely in Ω .

3.1 Classical distributional solution in the one dimensional case.

We observe that in the one-dimensional case the sequence (Z_n) of solutions of the regularized stochastic heat equations [\(6\)](#page-6-0) converges uniformly on compacts of $(0, T) \times \mathbb{R}$ to Z, where Z is the solution of the stochastic heat

equation in the Itô sense

$$
\begin{cases}\ndZ = \Delta_x Z \, dt + Z \, dW, \\
Z(0, x) = e^{f(x)}.\n\end{cases} \tag{11}
$$

See L. Bertini and N. Cancrini [\[5\]](#page-11-9), Theorem 2.2.

We will denote by $U(t, x)$ the gradient in the sense of distributions of log $Z(t, x)$, that is, $U(t, x) = \nabla_x \log Z(t, x)$.

Corollary 3.2 U is a distributional solution of the one-dimensional stochastic Burgers equation [\(1\)](#page-1-0). That is, U verifies

1. There exists a sequence of C^{∞} -semimartingales (U_n) such that

$$
U = \lim_{n \to \infty} U_n
$$

and there exists $\nabla_x ||U||^2$ such that

$$
\nabla_x ||U||^2 := \lim_{n \to \infty} \nabla_x ||U_n||^2 \text{ in } \mathcal{D}'((0,T) \times \mathbb{R}).
$$

2. For all $\varphi \in \mathcal{D}((0,T) \times \mathbb{R}; \mathbb{R}),$

$$
(\mathcal{L}_B U, \varphi) = (\nabla \dot{W}, \varphi).
$$

3. $\nabla_x f$ is the section of U at $t = 0$ in the sense of S. Lojasiewicz, see [\[21\]](#page-13-2).

Proof: It is clear that,

$$
\lim_{n \to \infty} (U_n, \partial_t \varphi) = (U, \partial_t \varphi) \tag{12}
$$

and

$$
\lim_{n \to \infty} (U_n, \Delta_x \varphi) = (U, \Delta_x \varphi).
$$
\n(13)

We have that $\int_{[0,T]\times\mathbb{R}} \varphi(t,x)dW_t(x)dx$ defines a continuous linear functional from $\mathcal{D}((0,T)\times\mathbb{R};\mathbb{R})$ to $\mathbb R$ (see K. Schaumlöffel [\[25\]](#page-13-7)) and

$$
\lim_{n \to \infty} \int_{[0,T] \times \mathbb{R}} \nabla_x \partial_t \varphi(t, x) dW_t^n(x) dx = (\nabla \dot{W}, \varphi).
$$
 (14)

From the equation (9) and the convergences (12) , (13) and (14) we have that for all $\varphi \in \mathcal{D}((0,T) \times \mathbb{R}; \mathbb{R}),$

$$
\int_{[0,T]\times\mathbb{R}} \nabla_x ||U_n(t,x)||^2 \varphi(t,x) dt dx
$$

converges and defines a linear functional. Thus the nonlinearity

$$
(\nabla_x ||U(t,x)||^2, \varphi) := \lim_{n \to \infty} \int_{[0,T] \times \mathbb{R}} \nabla_x ||U_n(t,x)||^2 \varphi(t,x) dt dx
$$

is well defined.

Let $\varphi \in \mathcal{D}(\mathbb{R})$ and $\{\rho_{\varepsilon} : \varepsilon > 0\}$ be a strict delta–net. From the continuity of $Z(t, x),$

$$
\lim_{\varepsilon \to 0} \int_{[0,T] \times \mathbb{R}} \nabla_x \ln \ Z(t, x) \, \rho_{\varepsilon}(t) \, \varphi(x) \, dt \, dx = - \int_{\mathbb{R}} f(x) \, \nabla_x \varphi(x) \, dx
$$
\n
$$
= \int_{\mathbb{R}} \nabla_x f(x) \, \varphi(x) \, dx.
$$

Thus $\nabla_x f$ is the section of U at $t = 0$. \Box

Remark 3.2 More details on the results of this subsection can be found in [\[9\]](#page-11-4).

Acknowledgements

The authors P. Catuogno and C, Olivera are partially supported by CNPq through the grant 460713/2014–0. C. Olivera also by FAPESP through the grants 2015/04723–2 and 2015/07278–0. J.F. Colombeau is supported by FAPESP through the grant 2012/18940–7 .

References

- [1] S. Albeverio ; Z. Haba ; F. Russo. A two–space dimensional semilinear heat equation perturbed by (Gaussian) white noise. Probab. Theory Related Fields. 121 (2001), 3, 319–366.
- [2] J. Aragona ; J.F. Colombeau ; S.O. Juriaans. Nonlinear Generalized functions and jump conditions for a standard liquid–gas model. J. Math. Anal. Appl. 418 (2014), 2, 964–977.
- [3] S. Assing. A pregenerator for Burgers equation forced by conservative noise. Commun. Math. Phys. 225 (2002), 3, 611–632.
- [4] J. Bec ; K. Khanin. Burgers turbulence. Phys. Rep. 447 (2007), 1–2, 1–66.
- [5] L. Bertini ; N. Cancrini. The stochastic heat equation: Feynman–Kac formula and intermittence. Journal of Statistical Physics. 78 (1995), 5–6, 1377–1401.
- [6] L. Bertini ; N. Cancrini ; G. Jona–Lasinio. The stochastic Burgers equation. Comm. Math. Phys. 165 (1994), 2, 211–232.
- [7] L. Bertini ; G. Giacomin. Stochastic Burgers and KPZ equations from particle systems. Comm. Math. Phys. 183 (1997), 3, 571–607.
- [8] Z. Brzezniak ; B. Goldys ; M. Neklyudov. Multidimensional stochastic Burgers equation. SIAM J. Math. Anal. 46 (2014), 1, 871–889.
- [9] P. Catuogno ; C. Olivera. Strong solution of the stochastic Burgers equation. Appl. Anal. 93 (2014), 3, 646–652.
- [10] J. Colombeau. Elementary introduction to new generalized functions. Math. Studies 113, North Holland, 1985
- [11] J. Colombeau. Multiplication of distributions. Lecture Notes in Math. 1532, Springer Verlag, 1992.
- [12] G. Da Prato ; A. Debussche ; R. Temam. Stochastic Burgers equation. NoDEA Nonlinear Differential Equations Appl. 1 (1994), 4, 389–402.
- [13] A. Dermoune. Around the stochastic Burgers equation. Stochastic Anal. Appl. 15 (1997), 3, 295–311.
- [14] M. Grosser ; M. Kunzinger ; M. Oberguggenberger ; R. Steinbauer. Geometric theory of generalized functions. Kluwer, Dordrecht, 2001.
- [15] M. Gubinelli ; N. Perkowski Lectures on singular stochastic PDEs , Ensaios Matemáticos, 29. Sociedade Brasileira de Matemática, Rio de Janeiro, 2015.
- [16] M. Hairer. Solving the KPZ equation. Ann. of Math. (2) 178 (2013), 2, 559–664.
- [17] H. Holden ; T. Lindstrøm ; B. Øksendal ; J. Ubøe ; T. Zhang. The Burgers equation with a noisy force and the stochastic heat equation. Comm. Partial Differential Equations. 19 (1994), 1–2, 119–141.
- [18] H. Holden ; T. Lindstrøm ; B. Øksendal ; J. Ubøe ; T. Zhang. The stochastic Wick type Burgers equation. Stochastic Partial Differential Equations. 141–161, London Math. Soc. LNS, 216, Cambridge, 1995.
- [19] R. Iturriaga ; K. Khanin. Burgers turbulence and random Lagrangian systems. Comm. Math. Phys. 232 (2003), 3, 377–428.
- [20] H. Kunita. Stochastic flows and stochastic differential equations. Cambridge University Press, 1990.
- [21] M. Oberguggenberger. Multiplication of distributions and applications to partial differential equations. Pitman Research Notes in Math. Series 259. Ed. Longman Science and Technology, 1993.
- [22] M. Oberguggenberger; F. Russo. Nonlinear SPDEs: Colombeau solutions and pathwise limits. Stochastic analysis and related topics, VI (Geilo, 1996), Progr. Probab., 42, Birkhauser Boston, Boston, MA, 1998.
- [23] M. Oberguggenberger; F. Russo. White noise driven stochastic partial differential equations: triviality and non–triviality. Nonlinear theory of generalized functions (Vienna, 1997), 315–333, Chapman– Hall/CRC Res. Notes Math., 401, Chapman–Hall/CRC, 1999.
- [24] E. Panov ; V. Shelkovich. δ -shock waves as a new type of solutions to systems of conservation laws. J. Differential Equations 228 (2006), 49–86.
- [25] K. Schaumlöffel. White noise in space and time as the time–derivative of a cylindrical Wiener process . Stochastic Partial Differential Equations and Applications II, 225–229, LNM 1390, Springer, 1989.
- [26] R. Steinbauer. The ultra relativistic Reissner–Nordstrom field in the Colombeau algebra. J. Math. Phys. 38 (1977), 3, 1614–1622.
- [27] J. Walsh. An introduction to stochastic partial differential equations. Ecole d été de probabilités de Saint–Flour, XIV1984, 265–439, LNM 1180, Springer, 1986.
- [28] E. Weinan. Stochastic hydrodynamics. Current developments in mathematics, 2000, 109–147, Int. Press, Somerville, 2001.