Well-Posedness and Comparison Principle for Option Pricing with Switching Liquidity

T.B. Gyulov and L.G. Valkov[∗]

University of Ruse "Angel Kanchev", 7017 Ruse, Bulgaria

June 28, 2021

Abstract

We consider an integro-differential equation derived from a system of coupled parabolic PDE and an ODE which describes an European option pricing with liquidity shocks. We study the well-posedness and prove comparison principle for the corresponding initial value problem.

1 Introduction

This work is devoted to the study of an initial value problem of the following form

$$
\begin{cases}\n\frac{\partial u}{\partial \tau} - \frac{1}{2}\sigma^2 S^2 \frac{\partial^2 u}{\partial S^2} &= -\nu_{01} e^{u(S,\tau)} \left(\nu_{10} \int_0^\tau e^{-u(S,s)} ds + e^{-\gamma h(S)} \right) + \kappa, \\
u(S,0) &= \gamma h(S).\n\end{cases}
$$
\n(1)

Here $\tau \in [0, T]$, $S \in (0, +\infty)$, $h(S)$ is a given function and σ , ν_{01} , ν_{10} , κ and γ are constants.

The integro-differential equation in [\(1\)](#page-0-0) is derived from a system of coupled parabolic PDE and ODE which is suggested by M. Ludkovski and Q. Shen [\[6\]](#page-14-0) in European option pricing in a financial market switching between two states -a liquid state (0) and an illiquid (1) one. We briefly describe their model. First, it is assumed that the dynamics of the liquidity is represented by a continuous-time Markov chain (M_t) with intensity rates of the transitions $0 \to 1$ and $1 \to 0$ and determined by the constants ν_{01} and ν_{10} , respectively. During the liquid phase $(M_t = 0)$ the market dynamics follows the classical Black-Scholes model. More precisely, the price S_t of a stock is modelled by geometric Brownian motion

$$
dS_t = \mu S_t dt + \sigma S_t dW_t,
$$

with drift μ and volatility σ and a standard one-dimensional Brownian motion (W_t) which is independent of the Markov chain (M_t) (under the "real world" probability \mathbb{P}). Then the wealth process (X_t) satisfies

$$
dX_t = \mu \pi_t X_t dt + \sigma \pi_t X_t dW_t,
$$

[∗] e-mails: tgulov@uni-ruse.bg (T.Gyulov) and lvalkov@uni-ruse.bg (L.Valkov)

where π_t denotes the proportion of stock holdings in the total wealth X_t . For simplicity, it is assumed that the interest rate of the riskless asset is zero.

Respectively, in the illiquid phase $(M_t = 1)$, the market is static and trading in stock is not permitted, i.e., $dS_t = dX_t = 0$.

The presence of liquidity shocks is a source of non-traded risk and makes the market incomplete. Ludkovski and Shen investigate expected utility maximization with exponential utility function:

$$
u(x) = -e^{-\gamma x},
$$

where $\gamma > 0$ is the investor's risk aversion parameter. The value functions $\hat{U}^{i}(t, X, S), i = 0, 1$ for the optimal investment problem are defined as follows:

$$
\hat{U}^i(t, X, S) := \sup_{\pi_t} \mathbb{E}^{\mathbb{P}}_{t, X, S, i} \left[-e^{-\gamma (X_T + h(S_T))} \right], \quad i = 0, 1,
$$

where $\mathbb{E}_{t,X,S,i}^{\mathbb{P}}$ is the expectation under the measure \mathbb{P} with starting values $S_t =$ S, $X_t = X$ and $M_t = i$. The supremum above is taken over all admissible trading strategies (π_t) and the function $h(S)$ denotes the terminal payoff of a contingent claim. Standard stochastic control methods and the properties of the exponential utility function imply that the value functions can be presented by

$$
\hat{U}^{i}(t, X, S) = -e^{-\gamma X} e^{-\gamma R^{i}(t, S)}, \quad i = 0, 1,
$$

where $R^{i}(t, S)$ are the unique viscosity solutions of the system ([\[6\]](#page-14-0))

$$
\begin{cases}\nR_t^0 + \frac{1}{2}\sigma^2 S^2 R_{SS}^0 - \frac{\nu_{01}}{\gamma} e^{-\gamma (R^1 - R^0)} + \frac{d_0 + \nu_{01}}{\gamma} = 0, \\
R_t^1 - \frac{\nu_{10}}{\gamma} e^{-\gamma (R^0 - R^1)} + \frac{\nu_{10}}{\gamma} = 0,\n\end{cases} \tag{2}
$$

with the terminal condition $R^{i}(T, S) = h(S), i = 0, 1$. Here $d_0 := \mu^2/2\sigma^2$.

Let p and q denote the buyer's indifference prices corresponding to liquid and illiquid initial state respectively. They are defined as follows: $\hat{U}^0(t, X - \hat{U}^0(t, X - \hat{U}^0$ $p, S) = \hat{V}^0(t, X)$ and $\hat{U}^1(t, X - q, S) = \hat{V}^1(t, X)$ where $\hat{V}^i, i = 0, 1$ are the value functions of the Merton optimal investment problem (i.e. the case when $h(S) \equiv 0$). It can be shown that p and q satisfy a system of differential equations which is quite similar to (2) (see (15)). In fact,

$$
p = R^0 + \gamma^{-1} \ln F_0(t)
$$
 and $q = R^1 + \gamma^{-1} \ln F_1(t)$

where

$$
F_0(t) = c_1 e^{\lambda_1 t} + c_2 e^{\lambda_2 t}
$$

\n
$$
F_1(t) = \frac{1}{\nu_{01}} \left(c_1 \left(d_0 + \nu_{01} - \lambda_1 \right) e^{\lambda_1 t} + c_2 \left(d_0 + \nu_{01} - \lambda_2 \right) e^{\lambda_2 t} \right)
$$

$$
\lambda_{1,2} = \frac{d_0 + \nu_{01} + \nu_{10} \pm \sqrt{(d_0 + \nu_{01} + \nu_{10})^2 - 4d_0\nu_{10}}}{2},
$$

$$
c_1 = \frac{\lambda_2 - d_0}{\lambda_2 - \lambda_1}e^{-\lambda_1 T}, \quad \text{and} \quad c_2 = \frac{\lambda_1 - d_0}{\lambda_1 - \lambda_2}e^{-\lambda_2 T}
$$

.

Indifference pricing was first used in the pioneering paper of Hodges and Neuberger [\[3\]](#page-14-1). We refer also to [\[2\]](#page-14-2) for further applications (see [\[4\]](#page-14-3) and [\[8\]](#page-14-4) as well).

The existence of classical solutions was proved in [\[6\]](#page-14-0) when the payoff function $h(S)$ is bounded. This case is restrictive since it does not include such typical example as the call option $h = \max\{S - K, 0\}$ with strike price K. We investigate the solvability of the problem and prove the existence and uniqueness of a weak solution in suitable Sobolev weighted spaces which allows unbounded terminal payoff functions.

The integro-differential equation [\(1\)](#page-0-0) is derived from [\(2\)](#page-1-0) as follows. Denote $r^0 := \gamma R^0$, $r^1 = \gamma R^1$. The system of differential equations for r^0 and r^1 has the following from:

$$
\begin{cases}\n r_{\tau}^{0} - \frac{1}{2} \sigma^{2} S^{2} r_{SS}^{0} = -\nu_{01} e^{-\left(r^{1} - r^{0}\right)} + d_{0} + \nu_{01} \\
 r_{\tau}^{1} = -\nu_{10} e^{-\left(r^{0} - r^{1}\right)} + \nu_{10}\n\end{cases} \tag{3}
$$

where $\tau = T - t$. The ODE in [\(3\)](#page-2-0) can be solved explicitly with respect to $r¹$. Then we obtain the initial value problem [\(1\)](#page-0-0) under the substitution $u :=$ $r^0 - \nu_{10}\tau$ and $\kappa := d_0 + \nu_{01} - \nu_{10}$.

The paper is organized as follows. In Section 2 we prove a comparison principle (Theorem [2.1\)](#page-2-1) for classical solutions to the problem [\(1\)](#page-0-0). Then, in Section 3 we prove a comparison principle (Theorem [3.4\)](#page-8-0) for weak sub/super solutions. In addition, we study the existence and uniqueness of weak solutions in a suitable weighted Sobolev space (see Theorem [3.7\)](#page-11-0).

2 Comparison principle for classical solutions

In this section we consider solutions of [\(1\)](#page-0-0) satisfying

$$
|u|, |h| \le A \exp\left(\alpha \ln^2 S\right) = A S^{\alpha \ln S},\tag{4}
$$

for some positive constants A and α . Note that conditions [\(4\)](#page-2-2) include for example linear growth, polinomial and powers of S with arbitrary exponent.

We prove the following comparison principle:

Theorem 2.1. Let $u_1, u_0 \in C((0, +\infty) \times [0, T)) \cap C^{2,1}((0, +\infty) \times (0, T))$ be two clasical solutions of [\(1\)](#page-0-0) corresponding to the initial data $h = h_1$ and $h = h_0$, respectively and such that the conditions [\(4\)](#page-2-2) hold. Then

$$
\gamma \inf (h_1 - h_0) \le u_1 - u_0 \le \gamma \sup (h_1 - h_0).
$$
 (5)

We will only prove the lower bound in [\(5\)](#page-2-1) since the upper one follows immediately from it. In addition, we can assume that

$$
\underline{h} := \gamma \inf \left(h_1 - h_0 \right) > -\infty,
$$

otherwise the left inequality in [\(5\)](#page-2-1) is trivial. We will use the following auxiliary lemma

Lemma 2.2. Let u_1 and u_0 be as in Theorem [2.1](#page-2-1) and $\tau_1 \geq 0$ be such that $u_1(S,\tau) - u_0(S,\tau) \geq \underline{h}$ for any $\tau \in [0,\tau_1]$. Then, there exists a constant $\bar{\tau} > 0$ such that $u_1(S, \tau) - u_0(S, \tau) \geq \underline{h}$ for any $\tau \in [0, \tau_1 + \overline{\tau})$. In addition, $\overline{\tau}$ depends only on α defined in [\(4\)](#page-2-2) and σ .

Proof. Let u_1 and u_0 be two solutions of [\(1\)](#page-0-0) corresponding to the initial conditions $u_1(S, 0) = \gamma h_1(S)$ and $u_0(S, 0) = \gamma h_0(S)$. Denote $\tilde{u} = u_1 - u_0$, $\tilde{h} = \gamma (h_1 - h_0), u_{\xi} = \xi u_1 + (1 - \xi) u_0, h_{\xi} = \xi h_1 + (1 - \xi) h_0, \text{ for } \xi \in [0, 1]$ and define

$$
\mathcal{F}[\tau; u, g] := -\nu_{01} e^{u(\tau)} \left(\nu_{10} \int_0^{\tau} e^{-u(s)} ds + e^{-g} \right) + \kappa.
$$

Then

$$
\mathcal{F}[\tau; u_1, \gamma h_1] - \mathcal{F}[\tau; u_0, \gamma h_0] = \int_0^1 \frac{d}{d\xi} \left(\mathcal{F}[\tau; u_\xi, \gamma h_\xi] \right) d\xi \tag{6}
$$

$$
= -\nu_{01}\tilde{u}\int_0^1 e^{u_\xi(\tau)} \left(\nu_{10}\int_0^\tau e^{-u_\xi(s)}ds + e^{-\gamma h_\xi}\right)d\xi \tag{7}
$$

$$
+\nu_{01}\int_{0}^{1}e^{u_{\xi}(\tau)}\left(\nu_{10}\int_{0}^{\tau}e^{-u_{\xi}(s)}\tilde{u}(s)ds+e^{-\gamma h_{\xi}}\tilde{h}\right)d\xi
$$

$$
=-\nu_{01}\nu_{10}\int_{0}^{1}\int_{0}^{\tau}e^{u_{\xi}(\tau)-u_{\xi}(s)}\left(\tilde{u}(\tau)-\tilde{u}(s)\right)dsd\xi
$$

$$
-\nu_{01}\left(\tilde{u}(\tau)-\tilde{h}\right)\int_{0}^{1}e^{u_{\xi}(\tau)-\gamma h_{\xi}}d\xi
$$
(8)

and

$$
\tilde{u}_{\tau} - \frac{1}{2}\sigma^2 S^2 \tilde{u}_{SS} = -\nu_{01}\nu_{10} \int_0^{\tau} \left(\tilde{u}\left(\tau\right) - \tilde{u}\left(s\right)\right) ds \int_0^1 e^{u_{\xi}(\tau) - u_{\xi}(s)} d\xi \qquad (9)
$$

$$
-\nu_{01} \left(\tilde{u}\left(\tau\right) - \tilde{h}\right) \int_0^1 e^{u_{\xi}(\tau) - \gamma h_{\xi}} d\xi
$$

Next, define

$$
\omega(S,\tau) := \frac{1}{\sqrt{T_1 - \tau}} \exp\left(\frac{\left(\ln S - \frac{1}{2}\sigma^2 (T_1 - \tau)\right)^2}{2\sigma^2 (T_1 - \tau)}\right),\tag{10}
$$

where $T_1 > 0$ and $(S, \tau) \in (0, +\infty) \times [0, T_1)$. Note that $\mathcal{L}_{BS} \omega = \omega_{\tau}$ $\frac{1}{2}\sigma^2 S^2 \omega_{SS} = 0$ and ω is increasing with respect to τ in the interval $\tau \in [T_1 - 4/\sigma^2, T_1)$. Choose $T_1 > \tau_1$ in [\(10\)](#page-3-0) such that the inequality

$$
\alpha < \frac{1}{2\sigma^2 \left(T_1 - \tau \right)},
$$

holds for all $\tau \in [\tau_1, T_1)$ and $T_1 - 4/\sigma^2 < \tau_1$. It is enough to define $T_1 := \tau_1 + \overline{\tau}$,

where $0 < \bar{\tau} < \min \left\{ (2\sigma^2 \alpha)^{-1}, 4/\sigma^2 \right\}$. Next, let $\varphi_{\epsilon} = \tilde{u} + \epsilon \omega$. Then

$$
(\varphi_{\epsilon})_{\tau} - \frac{1}{2}\sigma^{2} S^{2} (\varphi_{\epsilon})_{SS} = -\nu_{01}\nu_{10} \int_{0}^{\tau} (\tilde{u}(\tau) - \tilde{u}(s)) ds \int_{0}^{1} e^{u_{\xi}(\tau) - u_{\xi}(s)} d\xi
$$

$$
-\nu_{01} (\tilde{u}(\tau) - \tilde{h}) \int_{0}^{1} e^{u_{\xi}(\tau) - \gamma h_{\xi}} d\xi
$$
(11)

$$
\geq -\nu_{01}\nu_{10}\left(\tilde{u}\left(\tau\right)-\underline{h}\right)\int_{0}^{\tau_{1}}ds\int_{0}^{1}e^{u_{\xi}(\tau)-u_{\xi}(s)}d\xi
$$
\n
$$
-\nu_{01}\nu_{10}\int_{\tau_{1}}^{\tau}\left(\tilde{u}\left(\tau\right)-\tilde{u}\left(s\right)\right)ds\int_{0}^{1}e^{u_{\xi}(\tau)-u_{\xi}(s)}d\xi
$$
\n
$$
-\nu_{01}\left(\tilde{u}\left(\tau\right)-\tilde{h}\right)\int_{0}^{1}e^{u_{\xi}(\tau)-\gamma h_{\xi}}d\xi
$$
\n(12)

We will prove that $\varphi_{\epsilon} \geq \underline{h}$ for any $\tau \in [\tau_1, T_1)$. Indeed, assume by contradiction that inf $\varphi_{\epsilon} < \underline{h}$. Note that $\varphi_{\epsilon}|_{\tau=\tau_1} > \underline{h}$ and there exist \overline{S} and \underline{S} such that $\varphi_{\epsilon} > \underline{h}$ if either $S \leq \underline{S}$ or $S \geq \overline{S}$. In fact, $\varphi_{\epsilon} \to +\infty$ uniformly when either $|\ln S| \to +\infty$ or $\tau \to T_1$. The last observations imply that φ_ϵ attains minimum in an interior point $(S_*, \tau_*) \in (\underline{S}, \overline{S}) \times (\tau_1, T_1)$ and $\varphi_{\epsilon}(S_*, \tau_*) < \underline{h}$. Then, $(\varphi_{\epsilon})_{\tau}(S_*, \tau_*) = 0$, $(\varphi_{\epsilon})_{SS}$ $(S_*, \tau_*) \geq 0$ and

$$
\tilde{u}(S_*, \tau_*) - \tilde{h} \le \tilde{u}(S_*, \tau_*) - \underline{h} = \varphi_{\epsilon}(S_*, \tau_*) - \underline{h} - \epsilon \omega(S_*, \tau_*) < 0 \tag{13}
$$
\n
$$
\tilde{u}(S_*, \tau_*) - \tilde{u}(S_*, s) = \varphi_{\epsilon}(S_*, \tau_*) - \varphi_{\epsilon}(S_*, s)
$$

$$
-\epsilon\left(\omega(S_*,\tau_*)-\omega(S_*,s)\right)<0,\quad\forall s\in[\tau_1,\tau_*],\quad(14)
$$

since ω is increasing in τ . Thus the right hand side of [\(12\)](#page-4-1) is positive, a contradiction. Hence $\varphi_{\epsilon} = \tilde{u} + \epsilon \omega \geq \underline{h}$ for any $\tau \in [\tau_1, T_1)$. Let $\epsilon \to 0$. Then $\tilde{u} = u_1 - u_0 > h$ for any $\tau \in [\tau_1, T_1)$. $\tilde{u} = u_1 - u_0 \geq \underline{h}$ for any $\tau \in [\tau_1, T_1)$.

Proof. (of Theorem [2.1\)](#page-2-1) The comparison principle follows by induction and the auxiliary Lemma [2.2:](#page-3-1) we first take $\tau_1 = 0$ and prove it in the interval $[0, 1/2\bar{\tau}]$, then let $\tau_1 = 1/2\bar{\tau}$ and consider the interval $[1/2\bar{\tau}, \bar{\tau}]$ and etc. \Box

Now, as a corollary we formulate comparison principle for the buyer's indifference prices $p(S, t)$, $q(S, t)$ which satisfy the terminal value problem

$$
\begin{cases}\n p_t + \frac{1}{2}\sigma^2 S^2 p_{SS} - \frac{\nu_{01}}{\gamma} \frac{F_1}{F_0} e^{-\gamma(q-p)} + \frac{d_0 + \nu_{01}}{\gamma} - \frac{1}{\gamma} \frac{F'_0}{F_0} = 0 \\
 q_t - \frac{\nu_{10}}{\gamma} \frac{F_0}{F_1} e^{-\gamma(p-q)} + \frac{\nu_{10}}{\gamma} - \frac{1}{\gamma} \frac{F'_1}{F_1} = 0 \\
 p(S, T) = q(S, T) = h(S).\n\end{cases}
$$
\n(15)

By classical solutions of [\(15\)](#page-4-0) we mean functions such that $p \in C((0, +\infty) \times$ $(0, T] \cap C^{2,1}((0, +\infty) \times (0, T)), q \in C((0, +\infty) \times (0, T]), q_t \in C((0, +\infty) \times (0, T)).$ Note that

$$
\gamma p = \nu_{10}(T - t) + \ln F_0(t) + u(S, T - t),\tag{16}
$$

$$
\gamma q = \nu_{10}(T - t) + \ln F_1(t) - \ln \left(\nu_{10} \int_0^{T - t} e^{-u(S, s)} ds + e^{-\gamma h(S)} \right), \quad (17)
$$

since $p(t) = \gamma^{-1} (r^0 + \ln F_0(t))$ and $q(t) = \gamma^{-1} (r^1 + \ln F_1(t))$. Then, a comparison principle in (p, q) solutions will be equivalent to a comparison principle for the (r^0, r^1) variables.

We consider growth conditions analogous to [\(4\)](#page-2-2)

$$
|p|, |h| \le A \exp\left(\alpha \ln^2 S\right) = A S^{\alpha \ln S},\tag{18}
$$

for some positive constants A and α .

Corollary 2.3. Let (p_1, q_1) and (p_0, q_0) be two classical solutions of the system [\(15\)](#page-4-0) corresponding to terminal data $h \equiv h_1(S)$ and $h \equiv h_0(S)$, respectively. If there exist some positive constants A and α such that $p_i(S, t)$ and $h_i(S)$, $i = 0, 1$ satisfy the conditions [\(18\)](#page-5-0), then

$$
\inf\left(h_1 - h_0\right) \le p_1(S, t) - p_0(S, t) \le \sup\left(h_1 - h_0\right),\tag{19}
$$

$$
\inf\left(h_1 - h_0\right) \le q_1(S, t) - q_0(S, t) \le \sup\left(h_1 - h_0\right). \tag{20}
$$

In particular, let $h(S)$ be bounded from below (or from above) by a constant, i.e. $h(S) \geq h_*$ (resp. $h(S) \leq h^*$) and $p(S,t)$, $q(S,t)$, be a classical solutions of the terminal value problem [\(15\)](#page-4-0) satisfying [\(18\)](#page-5-0). Then

$$
p(S,t) \ge h_* \text{ and } q(S,t) \ge h_* \text{ (respectively } p(S,t) \le h^* \text{ and } q(S,t) \le h^*),
$$

for any $S \in (0, +\infty)$ and any $t \in (0, T]$.

Proof. The inequalities [\(19\)](#page-5-1) follow immediately from Theorem [2.1](#page-2-1) and representation (16) . In order to prove (20) we will use (17) , i.e.

$$
q_i(\cdot, t) = \gamma^{-1} \left[\nu_{10}(T - t) + \ln F_1(t) - \ln \left(\nu_{10} \int_0^{T - t} e^{-u_i(\cdot, s)} ds + e^{-\gamma h_i(\cdot)} \right) \right],
$$

for $i = 0, 1$. Similarly to the proof of Lemma [2.2](#page-3-1) we derive

$$
q_{1}(\cdot,t) - q_{0}(\cdot,t) = -\gamma^{-1} \int_{0}^{1} \frac{d}{d\xi} \left[\ln \left(\nu_{10} \int_{0}^{T-t} e^{-u_{\xi}(\cdot,s)} ds + e^{-\gamma h_{\xi}(\cdot)} \right) \right] d\xi
$$

$$
= \gamma^{-1} \int_{0}^{1} \frac{\nu_{10} \int_{0}^{T-t} e^{-u_{\xi}(\cdot,s)} (u_{1}(\cdot,s) - u_{0}(\cdot,s)) ds}{\nu_{10} \int_{0}^{T-t} e^{-u_{\xi}(\cdot,s)} ds + e^{-\gamma h_{\xi}(\cdot)}} d\xi
$$

$$
+ (h_{1}(\cdot) - h_{0}(\cdot)) \int_{0}^{1} \frac{e^{-\gamma h_{\xi}(\cdot)}}{\nu_{10} \int_{0}^{T-t} e^{-u_{\xi}(\cdot,s)} ds + e^{-\gamma h_{\xi}(\cdot)}} d\xi
$$

Now, [\(5\)](#page-2-1) implies the estimates [\(20\)](#page-5-2).

The second part follows immediately due to the fact that $p_*(S, t) \equiv h_*$ and $q_*(S, t) \equiv h_*$ are the solutions of the problem [\(15\)](#page-4-0) with constant terminal condition $h \equiv h_*$. Indeed, if we formally substitute $p_*(S, t) \equiv h_*$ and $q_*(S, t) \equiv$ h_* in [\(15\)](#page-4-0), then we arrive at the conclusion that it is sufficient to check the following identities

$$
-\frac{\nu_{01}}{\gamma}\frac{F_1}{F_0} + \frac{d_0 + \nu_{01}}{\gamma} - \frac{1}{\gamma}\frac{F'_0}{F_0} = 0, \tag{21}
$$

$$
-\frac{\nu_{10}}{\gamma}\frac{F_0}{F_1} + \frac{\nu_{10}}{\gamma} - \frac{1}{\gamma}\frac{F_1'}{F_1} = 0,\tag{22}
$$

or equivalently

$$
F_0' = -\nu_{01}F_1 + (d_0 + \nu_{01})F_0, \tag{23}
$$

$$
F_1' = -\nu_{10} F_0 + \nu_{10} F_1, \tag{24}
$$

 \Box

which follow directly from the definition of F_0 and F_1 .

3 Existence of weak solutions

In this section we study the existence and uniqueness of weak solutions in suitable function spaces. First we introduce the weighted L^2 space

$$
L^2_w := \left\{ u : ||u||_0^2 := \int_0^{+\infty} u^2(S)w(S)dS < \infty \right\},\,
$$

given a weight function $w > 0$. Then we define a weighted Sobolev space as follows

$$
H_w^1 := \{ u : u \in L_w^2 \text{ s.t. } Su'(S) \in L_w^2 \},
$$

with norm $\|\cdot\|_1$ such that $\|u\|_1^2 = \|u\|_0^2 + \|Su'\|_0^2$ 0 .

Let $\xi : [0, +\infty) \to [0, 1]$ be increasing, infinitely continuously differentiable function and such that $\xi \equiv 0$ on $[0, 1/2]$ and $\xi \equiv 1$ on $[1, +\infty)$. We will use ξ to construct a sequence ${u_{\epsilon}}$ of compactly supported functions converging in H_w^1 to a given element $u \in H_w^1$. More precisely, the following auxiliary result holds.

Lemma 3.1. Let $\xi_{\epsilon}(x) := \xi(x/\epsilon) [1 - \xi(x\epsilon/2)], 0 < \epsilon < 1$ and $u_{\epsilon} := \xi_{\epsilon} u$. Then $u_{\epsilon} \to u$ in H_w^1 , as $\epsilon \to 0$.

Proof. Note that $(u - u_{\epsilon})' = (1 - \xi_{\epsilon}) u' - \xi'_{\epsilon} u$,

$$
S\xi'_\epsilon(S)=(S/\epsilon)\,\xi'(S/\epsilon)\,[1-\xi(S\epsilon/2)]-(S\epsilon/2)\,\xi'(S\epsilon/2)\xi(S/\epsilon)
$$

is uniformly bounded with respect to ϵ and $1 - \xi_{\epsilon} \to 0$ as well as $S\xi'_{\epsilon}(S) \to 0$ as $\epsilon \to 0$. Then the Lebesgue's dominated convergence theorem implies that $||u - u_{\epsilon}|| \to 0$ as $\epsilon \to 0$. $||u - u_{\epsilon}|| \to 0$ as $\epsilon \to 0$.

Next, let $u(S)$ be twice continuously differentiable on $(0, +\infty)$ and denote the operator $\mathcal{L}u := -\frac{1}{2}\sigma^2 S^2 u''$. Then after integration by parts we formally obtain:

$$
\begin{aligned} (\mathcal{L}u, v)_{L^2_w} &= -\frac{1}{2}\sigma^2 \int_0^{+\infty} wS^2 u'' v \, dS \\ &= \frac{1}{2}\sigma^2 \int_0^{+\infty} \left[wS^2 u' v' + \left(S \frac{w'}{w} + 2 \right) wS u' v \right] dS, \end{aligned}
$$

provided that the integrals above are well-defined, w is continuously differentiable and $wS^2u'v \to \text{as } S \to 0 \text{ and } S \to \infty$. For example, the above holds when v is continuously differentiable and with compact support.

Following the above observations we introduce the bilinear form:

$$
a(u,v) := \frac{1}{2}\sigma^2 \int_0^{+\infty} wS u' \left[Sv' + \left(S \frac{w'}{w} + 2 \right) v \right] dS. \tag{25}
$$

If the weight function w is twice continuously differentiable, and there exists a constant $C > 0$, such that

$$
\left| S \frac{w'(S)}{w(S)} \right|, \left| S^2 \frac{w''(S)}{w(S)} \right| \le C, \forall S \in (0, +\infty). \tag{26}
$$

then the bilinear form $a(u, v)$ is continuous and semi-coercive on H_w^1 , i.e.,

$$
|a(u, v)| \le c \|u\|_{1} \|v\|_{1}, \quad \forall u, v \in H_w^1 \tag{27}
$$

$$
a(u, u) \ge \alpha ||u||_1^2 - \beta ||u||_0^2, \quad \forall u \in H_w^1
$$
 (28)

for some suitable constants $c > 0$, $\alpha > 0$ and $\beta > 0$ which are independent of u and v.

We can choose such weight function that the call option payoff function $h = \max\{S - K, 0\}$ belongs to the space H_w^1 , for example, take $w := (1 + S)^{\gamma}$, where $\gamma < -3$.

In addition, we assume that

$$
\theta := \int_0^{+\infty} w(S)dS < +\infty.
$$
 (29)

This assumption guarantees that any bounded and measurable function belongs to L^2_w .

Lemma 3.2. There exists a constant $c_0 > 0$ such that

$$
|u(S)|^2 \le c_0 \|u\|_1^2 \frac{1}{S} \exp(C|\ln S|), \qquad \forall u \in H_w^1,
$$
 (30)

where C satisfies (26) .

Proof. Note that there exists a constant c_0 such that

$$
|u(1)|^2 \le c_0 \|u\|_1^2, \qquad \forall u \in H_w^1,\tag{31}
$$

due to the Sobolev embbeding theorem.

Let S be fixed and denote $v(\zeta) := u(\zeta S)$. We have

$$
||v||_1^2 = \int_0^{+\infty} w(\zeta) \left(\zeta^2 S^2 (u'(\zeta S))^2 + u^2(\zeta S)\right) d\zeta
$$
 (32)

$$
= \int_0^{+\infty} \frac{w(\zeta)}{Sw(\zeta S)} w(\zeta S) \left(\zeta^2 S^2 \left(u'(\zeta S)\right)^2 + u^2(\zeta S)\right) d\left(S\zeta\right) \tag{33}
$$

$$
\leq \frac{1}{S} \exp(C |\ln S|) \|u\|_{1}^{2}, \tag{34}
$$

since

$$
\frac{w(\zeta)}{Sw(\zeta S)} = \frac{1}{S} \exp\left(\int_{\zeta S}^{\zeta} \frac{w'(\xi)}{w(\xi)} d\xi\right) \le \frac{1}{S} \exp\left(C |\ln S|\right).
$$

Then [\(30\)](#page-7-1) follows from [\(31\)](#page-7-2) since $v(1) = u(S)$.

 \Box

The space H_w^1 is densely and continuously embbeded in L^2_w . We consider the Gelfand triples

$$
H_w^1\subset L_w^2\subset H_w^*,
$$

and

$$
L^2(0,T;H^1_w) \subset L^2(0,T;L^2_w) \subset L^2(0,T;H^*_w),
$$

where H_w^* is the dual of H_w^1 . Next, we define the set

$$
W(0,T) := \left\{ u \in L^2(0,T;H_w^1), \dot{u} \in L^2(0,T;H_w^*) \right\},\tag{35}
$$

where \dot{u} is the distributional derivative of u. It is well known (see Lions and Magenes[\[5\]](#page-14-5)) that

$$
W(0,T) \subset C([0,T], L^2_w).
$$

For simplicity we will further write $u(\tau)$ instead of $u(S, \tau)$ when this does not lead to misunderstanding. Recall that

$$
\mathcal{F}[\tau; u, \gamma h] := -\nu_{01} e^{u(\tau)} \left(\nu_{10} \int_0^{\tau} e^{-u(s)} ds + e^{-\gamma h} \right) + \kappa.
$$

Definition 3.3. A function $u \in W(0,T)$ is called weak supersolution (subso-lution) of the initial value problem [\(1\)](#page-0-0) if $u(0) \geq \gamma h$ (resp. $u(0) \leq \gamma h$) and for a.a. $\tau \in (0, T)$ the inequality

$$
\langle \dot{u}, v \rangle + a(u, v) \ge (\le) \int_0^{+\infty} w \mathcal{F}[\tau; u, \gamma h] v dS,
$$
 (36)

holds for any nonegative $v \in H_w^1$. Respectively, the function $u \in W(0,T)$ is called weak solution of the initial value problem [\(1\)](#page-0-0) if $u(0) = \gamma h$ and for a.a. $\tau \in (0, T)$ the equality

$$
\langle \dot{u}, v \rangle + a(u, v) = \int_0^{+\infty} w \mathcal{F}[\tau; u, \gamma h] v dS, \qquad \forall v \in H_w^1,
$$
 (37)

holds.

Next, we prove the following comparison principle for weak super/subsolutions satisfying growth conditions of type [\(4\)](#page-2-2).

Theorem 3.4. Let \overline{u} be a weak supersolution of the initial value problem [\(1\)](#page-0-0) with initial data $h(S) \equiv h$ and μ be a weak subsolution corresponding to the initial data $h(S) \equiv \underline{h}$ where \underline{h} and h are given and $\underline{h} \leq h$. Assume in addition, that there exist positive constants A and α such that

$$
\left|\underline{h}\right|, \left|\overline{h}\right|, \left|\underline{u}\right|, \left|\overline{u}\right| \le A \exp\left(\alpha \ln^2 S\right) = A S^{\alpha \ln S},\tag{38}
$$

for a.a. $(S, t) \in (0, +\infty) \times [0, T]$. Then $\underline{u} \leq \overline{u}$ for a.a. $(S, t) \in (0, +\infty) \times [0, T]$.

Denote $u := \overline{u} - \underline{u}$. We will prove that $u_- := \max\{-u, 0\} = 0$ almost everywhere. Similarly to [\(9\)](#page-3-2), we obtain that the following inequality holds for a.a. $\tau \in (0, T)$ and for any nonegative $v \in H_w^1$ with compact support in $(0, +\infty)$:

$$
\langle \dot{u}, v \rangle + a(u, v) \ge -\nu_{01}\nu_{10} \int_0^\infty \left(\int_0^\tau \delta(\tau, s) \left(u(S, \tau) - u(S, s) \right) ds \right) v(S) w dS
$$
\n
$$
- \nu_{01} \int_0^\infty \left(u(S, \tau) - \tilde{h}(S) \right) v(S) \delta(\tau) w dS,
$$
\n(39)

where

$$
\delta(\tau,s) := \int_0^1 e^{u_\xi(\tau) - u_\xi(s)} d\xi, \quad \delta(\tau) := \int_0^1 e^{u_\xi(\tau) - \gamma h_\xi} d\xi,
$$

 $u_{\xi} := \xi \overline{u} + (1 - \xi) \underline{u}, u(\cdot, 0) \ge \tilde{h} := \gamma \left(\overline{h} - \underline{h} \right) \ge 0$ and $h_{\xi} := \xi \overline{h} + (1 - \xi) \underline{h}$. It is sufficient to prove the following auxiliary result:

Lemma 3.5. Assume that $\tau_1 \geq 0$ is such that for any $t \in [0, \tau_1]$ the inequality $\overline{u}(t) - \underline{u}(t) \geq 0$ holds a.e. on $(0, +\infty)$. Then the same inequality holds for any $t \in [0, \tau_1 + \overline{\tau}]$, where $\overline{\tau} > 0$ is a constant which depends only on α and σ .

Proof. Let ω be defined by [\(10\)](#page-3-0) and $u_{\epsilon} := u + \epsilon \omega$ where $u = \overline{u} - \underline{u}$. Then, assume that $\bar{\tau}$ is chosen as in the proof of Lemma [2.2.](#page-3-1) We will prove that $u_{\epsilon-} := \max\{-u_{\epsilon}, 0\} \equiv 0$ for a.a. $(S, t) \in (0, +\infty) \times [\tau_1, \tau_1 + \overline{\tau}]$. Note that there exist a closed interval $I_{\epsilon} \subset (0, +\infty)$ such that $u_{\epsilon-} = 0$ on the set $((0, +\infty) \setminus I_{\epsilon}) \times$ $[\tau_1, \tau_1 + \overline{\tau}]$ due to the conditions [\(38\)](#page-8-1). Now, let $\varphi(S)$ be a smooth function with compact support in $(0, +\infty)$ such that $\varphi(S) = 1$ on the interval I_{ϵ} . Then $u_{\epsilon}\varphi \in L^2(\tau_1, \tau_1 + \bar{\tau}; H_w^1)$ and $(u_{\epsilon}\varphi) = u_{\epsilon-}$. Next, for any nonnegative $v \in H_w^1$ with compact support supp $v \subset I_\epsilon$ we have $\varphi v = v$, $a(u\varphi, v) = a(u, v)$ and then

$$
\left\langle \frac{d}{d\tau} \left(u_{\epsilon} \varphi \right), v \right\rangle + a \left(u_{\epsilon} \varphi, v \right) = \left\langle \dot{u}, \varphi v \right\rangle + \epsilon \left\langle \varphi \dot{\omega}, v \right\rangle + a(u\varphi, v) + \epsilon a(\omega \varphi, v) \quad (40)
$$

$$
= \left\langle \dot{u}, v \right\rangle + a(u, v) - \frac{1}{2} \epsilon \sigma^2 \left(2\omega' \varphi' + \omega \varphi'', v \right)_{L^2_w}
$$

$$
= \langle \dot{u}, v \rangle + a(u, v) - \frac{1}{2} \epsilon \sigma^2 \underbrace{(2\omega' \varphi' + \omega \varphi'', v)_{L^2_{w}}}_{=0}
$$
\n
$$
\geq -\nu_{01} \nu_{10} \int_0^\infty \left(\int_0^\tau \delta(\tau, s) \left(u(S, \tau) - u(S, s) \right) ds \right) v(S) w dS \tag{41}
$$

$$
- \nu_{01} \int_0^{\infty} \left(u(S,\tau) - \tilde{h}(S) \right) v(S) \delta(\tau) w dS
$$

\n
$$
\geq -\nu_{01} \nu_{10} \int_0^{\infty} \left(\int_0^{\tau_1} \delta(\tau,s) ds \right) u(S,\tau) v(S) w dS
$$

\n
$$
- \nu_{01} \nu_{10} \int_0^{\infty} \left(\int_{\tau_1}^{\tau} \delta(\tau,s) (u(S,\tau) - u(S,s)) ds \right) v(S) w dS
$$

\n
$$
- \nu_{01} \int_0^{\infty} u(S,\tau) v(S) \delta(\tau) w dS,
$$
\n(42)

i.e.,

$$
\left\langle \frac{d}{d\tau} \left(u_{\epsilon} \varphi \right), v \right\rangle + a \left(u_{\epsilon} \varphi, v \right) \geq -\nu_{01} \nu_{10} \int_{0}^{\infty} \left(\int_{0}^{\tau_{1}} \delta(\tau, s) ds \right) u_{\epsilon}(S, \tau) v(S) w dS
$$
\n
$$
- \nu_{01} \nu_{10} \int_{0}^{\infty} \left(\int_{\tau_{1}}^{\tau} \delta(\tau, s) \left(u_{\epsilon}(S, \tau) - u_{\epsilon}(S, s) \right) ds \right) v(S) w dS
$$
\n
$$
- \nu_{01} \int_{0}^{\infty} u_{\epsilon}(S, \tau) v(S) \delta(\tau) w dS,
$$
\n
$$
(43)
$$

where we have used the fact that $u_{\epsilon} > u$ and $u_{\epsilon}(S, \tau) - u_{\epsilon}(S, s) > u(S, \tau) - u_{\epsilon}(S, \tau)$ $u(S, s)$ for any $s \in [\tau_1, \tau]$ since $\omega(S, \cdot)$ is increasing on that interval. Now, take $v = u_{\epsilon-}$ and note that $u_{\epsilon} = u_{\epsilon+} - u_{\epsilon-}$, $a(u_{\epsilon}\varphi, u_{\epsilon-}) = -a(u_{\epsilon-}, u_{\epsilon-})$ and

$$
u_{\epsilon}(S,s) u_{\epsilon-}(S,\tau) \geq -u_{\epsilon-}(S,s) u_{\epsilon-}(S,\tau) \geq -\frac{1}{2} \left(u_{\epsilon-}^{2}(S,s) + u_{\epsilon-}^{2}(S,\tau) \right).
$$

After integration with respect to τ form τ_1 to $t \in [\tau_1, \tau_1 + \overline{\tau}]$ the inequality [\(43\)](#page-9-0) implies

$$
\frac{1}{2} \left\| u_{\epsilon-}(t) \right\|_{0}^{2} + a \left(u_{\epsilon-}, u_{\epsilon-} \right) \leq - \int_{\tau_1}^{t} \left(\int_{0}^{\infty} \Sigma(S, \tau) u_{\epsilon-}^{2}(S, \tau) w dS \right) d\tau, \tag{44}
$$

where

$$
\Sigma(S,\tau) := \nu_{01}\nu_{10}\left(\int_0^{\tau_1} \delta(\tau,s)ds + \frac{1}{2}\int_{\tau_1}^{\tau} \delta(\tau,s)ds - \frac{1}{2}\int_{\tau}^t \delta(s,\tau)ds\right) + \nu_{01}\delta(\tau).
$$

 $|\Sigma(S,\tau)|$ is bounded from above by a constant, say $C > 0$, when $S \in I_{\epsilon}$ and due to the semi-coercivity of the bilinear form $a(\cdot, \cdot)$ (see [\(28\)](#page-7-3)) we obtain:

$$
\frac{1}{2} \|u_{\epsilon-}(t)\|_{0}^{2} \le (C+\beta) \int_{\tau_{1}}^{t} \|u_{\epsilon-}(\tau)\|_{0}^{2} d\tau.
$$
 (45)

Hence the Gronwall inequality implies $||u_{\epsilon-}(t)||_0 = 0$ for any $t \in [\tau_1, \tau_1 + \overline{\tau}]$ since $||u_{\epsilon-}(\tau_1)||_0 = 0$. Then $u + \epsilon \omega \geq 0$ a.e. Thus $u \geq 0$ a.e. since $\epsilon > 0$ is arbitrary.

We further prove another useful estimate.

Lemma 3.6. There exists a constant $C > 0$ such that

$$
\max_{t \in [0,T]} \|u(t)\|_{0} + \|u\|_{L^{2}(0,T,H_{w}^{1})} \leq C \left(\|u(0)\|_{0} + \|\hat{u}\|_{W(0,T)} + \gamma \|h\|_{0} + 1 \right) \tag{46}
$$

for any weak subsolution u and any function $\hat{u} \in W(0,T)$ satisfying $u \geq \hat{u}$.

Proof. Let $v \in H_w^1$ be some nonnegative function. We have

$$
\langle \dot{u}, v \rangle + a(u, v) \le \int_0^{+\infty} w \mathcal{F}[\tau; u, \gamma h] v dS,\n\le -\nu_{01} \nu_{10} \left(\int_0^{\tau} [u(\tau) - u(s)] ds \right), v \right)_{L_w^2} \n- \nu_{01} (u(\tau) - \gamma h, v)_{L_w^2} \n+ (\kappa - \nu_{01} \nu_{10} \tau - \nu_{01}) (1, v)_{L_w^2}.
$$
\n(47)

Take $v = u - \hat{u}$ and integrate [\(47\)](#page-10-0) with respect to τ from 0 to t.

$$
\frac{1}{2} ||u(t)||_{0}^{2} + a(u, u) \leq \frac{1}{2} ||u(0)||_{0}^{2} + (u, \hat{u})_{L_{w}^{2}} \Big|_{0}^{t} + a(u, \hat{u}) - \int_{0}^{t} \left\langle \dot{\hat{u}}, u \right\rangle d\tau
$$
\n
$$
- \nu_{01} \int_{0}^{t} (\nu_{10}\tau + 1) ||u(\tau)||_{0}^{2} d\tau + \nu_{01}\nu_{10} \frac{1}{2} ||\int_{0}^{t} u(\tau) d\tau||_{0}^{2}
$$
\n
$$
+ C_{1} (||\hat{u}||_{L^{2}(0, t, L_{w}^{2})} + \gamma ||h||_{0} + 1) ||u||_{L^{2}(0, t, L_{w}^{2})}
$$
\n
$$
+ C_{2} (\gamma ||h||_{0} + 1) ||\hat{u}||_{L^{2}(0, t, L_{w}^{2})}.
$$
\n(48)

Then a standard argument implies the estimate [\(46\)](#page-10-1).

 \Box

Now, we prove the existence of weak solutions, provided that $h \in H_w^1$. The proof is based on the lower and upper solution method (cf. [\[7\]](#page-14-6)). However, the exponential nonlinearity in [\(1\)](#page-0-0) causes some very technical difficulties which have to be overcome.

Theorem 3.7. Assume that $h \in H_w^1$. Then there exist a weak solution u to the initial value problem [\(1\)](#page-0-0). Moreover, there exists a constant $C > 0$ independent of u such that

$$
\|\dot{u}\|_{L^{2}(0,T,L^{2}_{w})} + \|u\|_{L^{\infty}(0,T,H^{1}_{w})} \leq C \left(\|u(0)\|_{1} + 1\right) \tag{49}
$$

Proof. We will present the proof in several steps.

Step 1. Let $h \in L^2_w$ be bounded. Then there exists a weak solution u to the initial value problem [\(1\)](#page-0-0). In addition, if $u(0) = \gamma h \in H_w^1$, then the inequality [\(49\)](#page-11-1) holds with a constant C independent of $u(0)$.

Note that we can conctruct appropriate couple of a supersolution \overline{u} and a subsolution <u>u</u>. Indeed, let the constant c_0 be such that $|\gamma h| \leq c_0$ and take $u := -c_0 - Mt$ for some positive constant M. If M is great enough then u is a subsolution. Analogously, $\overline{u} := c_0 + Mt$ is a supersolution provided that $M \geq \kappa$. Next, according to [\(8\)](#page-3-3) we can choose a constant $N > 0$ such that

$$
Nu(\tau) + \mathcal{F}[\tau; u, \gamma h] = Nu(\tau) - \nu_{01} e^{u(\tau)} \left(\nu_{10} \int_0^{\tau} e^{-u(s)} ds + e^{-\gamma h} \right) + \kappa
$$

is increasing in u , i.e.

$$
Nu_1(\tau)+\mathcal{F}[\tau;u_1,\gamma h]\geq Nu_0(\tau)+\mathcal{F}[\tau;u_0,\gamma h],
$$

for all u_0 and u_1 such that $u \leq u_0 \leq u_1 \leq \overline{u}$. Now, we can construct a decreasing sequence of supersolutions $u_0 := \overline{u}, u_1, u_2, \dots$ such that u_{n+1} is the solution of the initial value problem

$$
\begin{cases} \n\dot{u}_{n+1} - \frac{1}{2}\sigma^2 S^2 u''_{n+1,SS} + N u_{n+1} = N u_n + \mathcal{F}[\tau; u_n, \gamma h],\\ \n u_{n+1}(S, 0) = \gamma h(S) \n\end{cases}
$$

and $u \leq u_n \leq \overline{u}$. A standard argument implies that u_n converges to a weak solution of the problem [\(1\)](#page-0-0). We omit the details.

Next, assume in addition that $h \in H_w^1$. Then $\dot{u} \in L^2(0,T; L_w^2)$ and $u \in$ $L^{\infty}(0,T; H^1_w)$ (see, e.g., Bonnans [\[1\]](#page-14-7)) and the following parabolic estimate holds:

$$
\|\dot{u}\|_{L^{2}(0,T,L^{2}_{w})} + \|u\|_{L^{\infty}(0,T,H^{1}_{w})} \leq c_{0} \left(\|u(0)\|_{1} + \|\mathcal{F}[\cdot;u,\gamma h]\|_{L^{2}(0,T,L^{2}_{w})} \right)
$$

We will prove the stronger estimate [\(49\)](#page-11-1). First, we have

$$
-\frac{1}{2}\sigma^2 S^2 u_{SS}'' = \mathcal{F}[\tau; u, \gamma h] - \dot{u} \in L^2(0, T, L_w^2),\tag{50}
$$

$$
-\frac{1}{2}\sigma^2 \int_0^t \left(S^2 u''_{SS}, \dot{u}\right)_{L^2_w} d\tau = \frac{1}{2}\sigma^2 \left(\frac{1}{2} \|u(t)\|_1^2 - \frac{1}{2} \|u(0)\|_1^2\right) + \frac{1}{2}\sigma^2 \int_0^t \left(S\left(S\frac{w'}{w} + 2\right)u'_S - u, \dot{u}\right)_{L^2_w} d\tau
$$
\n(51)

$$
\int_{0}^{t} \left(\mathcal{F}[\tau; u, \gamma h], \dot{u}\right)_{L_{w}^{2}} d\tau = \int_{0}^{+\infty} \left(\int_{0}^{t} \frac{d}{d\tau} \left(\mathcal{F}[\tau; u, \gamma h]\right) d\tau\right) w dS \qquad (52)
$$

$$
+ \int_{0}^{+\infty} \left(\int_{0}^{t} \left(\kappa \dot{u} + \nu_{01} \nu_{10}\right) d\tau\right) w dS
$$

$$
\leq |\kappa| \theta^{1/2} \int_{0}^{t} \|\dot{u}(\tau)\|_{0} d\tau + \nu_{01} \left(1 + \nu_{10} t\right) \theta \qquad (53)
$$

since

$$
\frac{d}{d\tau}\left(\mathcal{F}[\tau;u,\gamma h]\right) = \frac{d}{d\tau}\left[-\nu_{01}e^{u(\tau)}\left(\nu_{10}\int_0^{\tau}e^{-u(s)}ds + e^{-\gamma h}\right) + \kappa\right]
$$

$$
= -\nu_{01}e^{u(\tau)}\left(\nu_{10}\int_0^{\tau}e^{-u(s)}ds + e^{-\gamma h}\right)\dot{u} - \nu_{01}\nu_{10} \qquad (54)
$$

$$
= \mathcal{F}[\tau;u,\gamma h]\dot{u} - \kappa\dot{u} - \nu_{01}\nu_{10}. \qquad (55)
$$

and

$$
\int_0^t \frac{d}{d\tau} \left(\mathcal{F}[\tau; u, \gamma h] \right) d\tau = \mathcal{F}[t; u, \gamma h] - \mathcal{F}[0; u, \gamma h] \le \nu_{01} \tag{56}
$$

We multiply both sides of the equation $\dot{u} - 1/2\sigma^2 S^2 u''_{SS} = \mathcal{F}[\tau; u, \gamma h]$ with \dot{u} in L^2_w and integrate from 0 to T. Then [\(51\)](#page-11-2) and [\(53\)](#page-12-0) imply

$$
\int_{0}^{t} \|\dot{u}\|_{0}^{2} d\tau + \frac{1}{4} \sigma^{2} \|u(t)\|_{1}^{2} \leq -\frac{1}{2} \sigma^{2} \int_{0}^{t} \left(S \left(S \frac{w'}{w} + 2 \right) u'_{S} - u, \dot{u} \right)_{L_{w}^{2}} d\tau \quad (57)
$$

$$
+ |\kappa| \theta^{1/2} \int_{0}^{t} \|\dot{u}(\tau)\|_{0} d\tau + \frac{1}{4} \sigma^{2} \|u(0)\|_{1}^{2}
$$

$$
+ \nu_{01} (1 + \nu_{10} t) \theta
$$

$$
\leq \tilde{C} \left[\int_{0}^{t} (\|u(\tau)\|_{1} + 1) \|\dot{u}(\tau)\|_{0} d\tau + \|u(0)\|_{1}^{2} + 1 \right]
$$

for some constant $\tilde{C} > 0$. Now, a techical, but standard argument implies that [\(49\)](#page-11-1) holds.

Step 2. Let $h \in H_w^1$ be bounded from below, i.e., $u(0) = \gamma h \geq c$. Then there exists a weak solution u to the initial value problem [\(1\)](#page-0-0). In addition, the inequality [\(49\)](#page-11-1) holds.

Let $\xi_{\epsilon}(x)$ be defined as in Lemma [3.1,](#page-6-0) i.e., $\xi_{\epsilon}(x) := \xi(x/\epsilon) [1 - \xi(x\epsilon/2)].$ Step 1 implies that there exists a solution u_{ϵ} corresponding to the initial condition $u_{\epsilon}(0) = \xi_{\epsilon}(\gamma h - c) + c = \xi_{\epsilon} \gamma h + (1 - \xi_{\epsilon})c$ which is bounded. Moreover, $\xi_{\epsilon} \gamma h + (1 - \xi_{\epsilon})c$ ξ_{ϵ})c $\leq \gamma h$ increases as $\epsilon \downarrow 0$ and converges in H_w^1 to γh . Then the comparison principle from Theorem [3.4](#page-8-0) implies that the sequence u_{ϵ} is increasing as $\epsilon \downarrow$ 0. Next, the estimate [\(49\)](#page-11-1) and Lemma [3.2](#page-7-4) imply that $u_{\epsilon}(S, \tau)$ converges to a finite limit $u(S,\tau)$ for any $(S,\tau) \in (0,+\infty) \times [0,T]$. What is more, \dot{u}_{ϵ} is weakly convergent to $\dot{u}(S,\tau)$ in $L^2(0,T; L^2_w)$, u_{ϵ} is weakly- $*$ convergent to u in $L^{\infty}(0,T, H_w^1)$ and u satisfies the estimate [\(49\)](#page-11-1). Then it is sufficient to prove that $\mathcal{F}[\tau; u_{\epsilon}, \xi_{\epsilon} \gamma \bar{h} + (1 - \xi_{\epsilon}) c]$ is weakly convergent to $\mathcal{F}[\tau; u, \gamma h]$ in $L^2(0, T; H_w^*)$. First, note that

$$
\mathcal{F}[\tau; u_{\epsilon}, \xi_{\epsilon} \gamma h + (1 - \xi_{\epsilon}) c] = \dot{u}_{\epsilon} - \frac{1}{2} \sigma^2 S^2 u_{\epsilon, SS}''
$$

is bounded in $L^2(0,T; H_w^*)$ and then there exists an element $\tilde{\mathcal{F}} \in L^2(0,T; H_w^*)$ such that

$$
\mathcal{F}\left[\tau;u_{\epsilon},\xi_{\epsilon}\gamma h+(1-\xi_{\epsilon})c\right]\underset{L^{2}(0,T;H_{w}^{*})}{\rightarrow}\tilde{\mathcal{F}}.
$$

On the other hand, $\mathcal{F}[\tau; u_{\epsilon}, \xi_{\epsilon} \gamma h + (1 - \xi_{\epsilon})c]$ is bounded from above by the constant function κ . Let $v \in L^2(0,T; H^1_w)$ be some arbitrary nonnegative function. Then Fatou's lemma implies

$$
\left\langle \kappa - \tilde{\mathcal{F}}, v \right\rangle = \lim_{\epsilon \to 0} \left(\kappa - \mathcal{F}[\cdot; u_{\epsilon}, \xi_{\epsilon} \gamma h + (1 - \xi_{\epsilon}) c], v \right)_{L^{2}(0, T; L_{w}^{2})}
$$

$$
\geq \left\langle \kappa - \mathcal{F}[\cdot; u, \gamma h], v \right\rangle \geq 0,
$$
 (58)

i.e.

$$
\mathcal{F}[\cdot; u, \gamma h] \in L^2(0, T; H_w^*)
$$
 and $\mathcal{F}[\cdot; u, \gamma h] \geq \tilde{\mathcal{F}}$.

Finally, we prove that in fact

$$
\mathcal{F}[\cdot; u, \gamma h] \equiv \tilde{\mathcal{F}}, \text{ i.e., } \langle \mathcal{F}[\cdot; u, \gamma h], v \rangle = \langle \tilde{\mathcal{F}}, v \rangle \quad \forall v \in L^2(0, T; H_w^1). \tag{59}
$$

First, observe that, $v_{\epsilon} := \xi_{\epsilon} v \to v$ as $\epsilon \to 0$ in $L^2(0,T; H^1_w)$. Hence, it is sufficient to prove [\(59\)](#page-13-0) for functions v vanishing outside a set of the form $I \times [0, T]$ where $I \subset (0, +\infty)$ is some closed interval. According to estimate [\(49\)](#page-11-1) and Lemma [3.2](#page-7-4) (applied to the interval I) the functions u_{ϵ} and u are uniformly bounded on $I \times [0, T]$. Then

$$
\langle \mathcal{F}[\cdot; u, \gamma h], v \rangle = \lim_{\epsilon \to 0} \left(\mathcal{F}[\cdot; u_{\epsilon}, \xi_{\epsilon} \gamma h + (1 - \xi_{\epsilon}) c], v \right)_{L^2(0,T;L^2_w)} = \langle \tilde{\mathcal{F}}, v \rangle.
$$

Step 3. Let $h \in H_w^1$. Then there exists a weak solution u to the initial value problem (1) . In addition, the inequality (49) holds.

Consider a sequence of problems with initial condition

$$
u_N(S, 0) = \max \{ \gamma h(S), -N \}, \quad N = 1, 2,
$$

Then the corresponding solutions u_N form a decreasing sequence due to the com-parison principle and Lemma [3.2.](#page-7-4) Moreover, the pointwise limit $\lim_{N\to\infty}u_N(S,\tau)$ is finite for any (S, τ) since the inequality [\(49\)](#page-11-1) holds for each function u_N . Then the proof follows similar arguments as in Step 2. \Box

Finally, note that the uniqueness of the weak solution is a consequence of the comparison principle. More precisely, we have the following corollary.

Corollary 3.8. Assume that $h \in H_w^1$. Then there exists a unique weak solution $u \in W(0,T) \cap L^{\infty}(0,T,H_w^1)$ to the initial value problem [\(1\)](#page-0-0). Moreover, the estimate [\(49\)](#page-11-1) holds with a constant $C > 0$ independent of u.

Acknowledgement

The research is supported by the European Union under Grant Agreement number 304617 (FP7 Marie Curie Action Project Multi-INT STRIKE - Novel Methods in Computational Finance). The second author is also supported by Bulgarian National Fund of Science under Project I02/20-2014.

References

- [1] J.F. Bonnans, Numerical analysis of partial differential equations arising in finance, Master 2 "Probabilités et Finance", Paris VI and Ecole Polytechnique: revision of Aug. 29, 2013, available at [http://www.cmap.polytechnique.fr/](http://www.cmap.polytechnique.fr/~bonnans/notes/edpfin/edpfin.html)∼bonnans/notes/edpfin/edpfin.html
- [2] R. Carmona (Ed.), Indifference Pricing: Theory and Applications, Princeton University Press, Princeton NJ, 2008.
- [3] S.D. Hodges, A. Neuberger, Optimal Replication of Contingent Claims under Transaction Costs, Rev. Futures Markets 8, (1989), 222–239.
- [4] T.S.T. Leung, A Markov-modulated stochastic control problem with optimal multiple stopping with application to finance, Decision and Control (CDC), 49th IEEE Conference, IEEE, (2010), 559-566.
- [5] $J - L$. Lions, E. Magenes, *Problèmes aux Limites non Homogènes et Appli*cations, Vols. I and II., Dunod, Paris, 1968.
- [6] M. Ludkovski, Q. Shen, European option pricing with liquidity shocks, Int. J. Theor. Appl. Finan., 16, No. 7, Article ID 1350043, 30 p. (2013). ISSN 0219-0249
- [7] C. V. Pao, Nonlinear parabolic and elliptic equations. Plenum Press, New York, 1992.
- [8] T. Zhou, Indifference valuation of mortgage-backed securities in the presence of prepayment risk, Mathematical Finance 20 (2010), no. 3, 479-507.