

# EFFECTIVE WHITNEY THEOREM FOR COMPLEX POLYNOMIAL MAPPINGS OF THE PLANE

M. FARNIK & Z. JELONEK & M.A.S. RUAS

ABSTRACT. We describe the topology of a general polynomial mapping  $F = (f, g) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ , where  $\deg f = d_1$ ,  $\deg g = d_2$ . We show that the set  $C(F)$  of critical points of  $F$  is a smooth connected curve, which is topologically equivalent to a sphere with  $\frac{(d_1+d_2-3)(d_1+d_2-4)}{2}$  handles and  $d_1 + d_2 - 2$  points removed. Moreover, the discriminant of  $\Delta(F) = F(C(F))$  is a curve birationally equivalent to  $C(F)$  which has only cusps and nodes as singularities and it has

$$c(F) = d_1^2 + d_2^2 + 3d_1d_2 - 6d_1 - 6d_2 + 7$$

simple cusps and

$$d(F) = \frac{1}{2} [(d_1d_2 - 4)((d_1 + d_2 - 2)^2 - 2) - (d - 5)(d_1 + d_2 - 2) - 6]$$

nodes (here  $d = \gcd(d_1, d_2)$ ). If  $d_1 = d_2$  then  $\Delta(F)$  has  $d_1 + d_2 - 2$  smooth points at infinity, at which it is tangent to the line at infinity with multiplicity  $d_1$ . If  $d_1 > d_2$  then  $\Delta(F)$  has exactly one singular point  $P$  at infinity with  $d_1 + d_2 - 2$  branches and with delta invariant

$$\delta_P = \frac{1}{2}d_1(d_1 - d_2)(d_1 + d_2 - 2)^2 + \frac{1}{2}(-2d_1 + d_2 + d)(d_1 + d_2 - 2).$$

Finally let  $F = (f, g) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$  be an arbitrary polynomial mapping with  $\deg f \leq d_1$  and  $\deg g \leq d_2$ . Assume that  $F$  has generalized cusps at points  $a_1, \dots, a_r$ . Then  $\sum_{i=1}^r \mu_{a_i} \leq d_1^2 + d_2^2 + 3d_1d_2 - 6d_1 - 6d_2 + 7$ , where  $\mu_{a_i}$  denotes the index of a generalized cusp at the point  $a_i$ .

## 1. INTRODUCTION

In [11] Whitney showed that a general smooth mapping  $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  has only two-folds and simple cusps as singularities. The problem of counting the number of cusps of a general perturbation of a plane-to-plane real singularity was considered by Fukuda and Ishikawa in [1]. They proved that the number modulo 2 of cusps of a general perturbation  $F$  of a finitely determined map-germ  $F_0 : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$  is a topological invariant of  $F_0$ . More recently, in [9] Krzyżanowska and Szafraniec gave an algorithm to compute the number of cusps for sufficiently general fixed real polynomial mapping of the real plane.

Algebraic formulas to count the number of cusps and nodes of a general perturbation of a finitely determined holomorphic map-germ  $F_0 : (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}^2, 0)$ , were given by Gaffney and Mond in [2, 3]. In this case any two general perturbations  $F$  of  $F_0$  defined on a sufficiently small neighborhood of 0 are topologically equivalent, so the number of cusps and nodes of  $F$  is an invariant of the map-germ  $F_0$ .

Here we consider the global complex case. It is proved in [8] that two general polynomial mappings  $F : \mathbb{C}^n \rightarrow \mathbb{C}^m$  (where  $n \leq m$ ) of fixed degree have the same topology. In

---

1991 *Mathematics Subject Classification.* 14 D 06, 14 Q 20.

*Key words and phrases.* polynomials, folds, cusp singularities.

The first two authors are partially supported by the grant of Narodowe Centrum Nauki, grant number 2015/17/B/ST1/02637, the third author is partially supported by CNPq grant 305651/2011-0 and FAPESP grant 2014/00304-2.

particular they have the same number of topological invariants. We consider here the simplest case  $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$  and describe the topology of a general polynomial mapping  $F = (f, g) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ , where  $\deg f = d_1$ ,  $\deg g = d_2$ . In Section 3 we show that the set  $C(F)$  of critical points of  $F$  is a smooth connected curve which is topologically equivalent to a sphere with  $\frac{(d_1+d_2-3)(d_1+d_2-4)}{2}$  handles and  $d_1 + d_2 - 2$  points removed. In Section 4 we show that the discriminant of  $\Delta(F) = F(C(F))$  is a curve birationally equivalent to  $C(F)$  which has only cusps and nodes as singularities and it has

$$c(F) = d_1^2 + d_2^2 + 3d_1d_2 - 6d_1 - 6d_2 + 7$$

simple cusps and

$$d(F) = \frac{1}{2} [(d_1d_2 - 4)((d_1 + d_2 - 2)^2 - 2) - (d - 5)(d_1 + d_2 - 2) - 6]$$

nodes (here  $d = \gcd(d_1, d_2)$ ). If  $d_1 = d_2$  then  $\Delta(F)$  has  $d_1 + d_2 - 2$  smooth points at infinity, at which it is tangent to the line at infinity with multiplicity  $d_1$ . If  $d_1 > d_2$  then  $\Delta(F)$  has exactly one singular point  $P$  at infinity with  $d_1 + d_2 - 2$  branches and with delta invariant

$$\delta_P = \frac{1}{2}d_1(d_1 - d_2)(d_1 + d_2 - 2)^2 + \frac{1}{2}(-2d_1 + d_2 + d)(d_1 + d_2 - 2).$$

We conclude the paper with Section 5 where we introduce the notions of a generalized cusp and the index of a generalized cusp  $\mu$  (see Definitions 5.1 and 5.3). We show that if  $F = (f, g) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$  is an arbitrary polynomial mapping with  $\deg f \leq d_1$ ,  $\deg g \leq d_2$  and generalized cusps at points  $a_1, \dots, a_r$  then  $\sum_{i=1}^r \mu_{a_i} \leq d_1^2 + d_2^2 + 3d_1d_2 - 6d_1 - 6d_2 + 7$ .

Let us note that in the special case when  $\gcd(d_1, d_2) = 1$  the numbers  $c(F)$  and  $d(F)$  can be also computed by using local methods of Gaffney and Mond. Indeed, in that case we can choose a homogenous mapping  $F \in \Omega(d_1, d_2)$ , which is finitely determined and show that there is a deformation  $F_t$  of a generic mapping to  $F$  such that all cusps and nodes of  $F_t$  tend to 0 when  $t \rightarrow 0$ . In this case our formulas for  $c(F)$  and  $d(F)$  coincide with formulas of Gaffney-Mond. However in the general case this approach does not work since the appropriate homogenous mapping is not finitely determined.

## 2. GENERAL POLYNOMIAL MAPPINGS

Let  $\Omega_n(d_1, \dots, d_n)$  denote the space of polynomial mappings  $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$  of multi-degree bounded by  $d_1, \dots, d_n$ . Of course  $\Omega_n(d_1, \dots, d_n)$  has the structure of the affine space. By  $J^q(n)$  we denote the space of  $q$ -jets of polynomial mappings  $F = (f_1, \dots, f_n) : \mathbb{C}^n \rightarrow \mathbb{C}^n$ . We define it exactly as in [6]. However, if we fix coordinates in the domain and the target then we can identify  $J^q(n)$  with the space  $\mathbb{C}^n \times \mathbb{C}^n \times (\mathbb{C}^{N_q})^n$ , where  $\mathbb{C}^{N_q}$  parameterizes coefficients of polynomials of  $n$ -variables and of degree bounded by  $q$  with zero constant term (which correspond to suitable Taylor polynomials). In further applications, in most cases, we treat the space  $J^q(n)$  in this simple way. In particular for a given polynomial mapping  $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$  we can define the mapping  $j^q(F)$  as

$$j^q(F) : \mathbb{C}^n \ni x \mapsto \left( x, F(x), \left( \frac{\partial^{|\alpha|} f_i}{\partial x^\alpha}(x) \right)_{1 \leq i \leq n, 1 \leq |\alpha| \leq q} \right) \in J^q(n).$$

We start with the following observation:

**Proposition 2.1.** *Assume that  $d_i \geq q$  for  $i = 1, \dots, n$ . Let  $S_1, \dots, S_k$  be smooth algebraic submanifolds of  $J^q(n)$ . Then there is a Zariski open dense subset  $\Omega_n(d_1, \dots, d_n)(S_1, \dots, S_k) \subset$*

$\Omega_n(d_1, \dots, d_n)$  such that for every  $F \in \Omega_n(d_1, \dots, d_n)(S_1, \dots, S_k)$  we have

$$j^q(F) \pitchfork S_i, \text{ for } i = 1, \dots, k.$$

*Proof.* Consider the mapping

$$\Psi : \Omega_n(d_1, \dots, d_n) \times \mathbb{C}^n \ni (F, x) \mapsto j^q(F)(x) \in J^q(n).$$

It is easy to see that  $\Psi$  is a submersion. Fix  $1 \leq i \leq k$ . By the transversality theorem with a parameter the set of polynomials  $F \in \Omega_n(d_1, \dots, d_n)$  such that  $j^q(F)$  is transversal to  $S_i$  is dense in  $\Omega_n(d_1, \dots, d_n)$ . On the other side this set is constructible in  $\Omega_n(d_1, \dots, d_n)$ . We conclude that there is a Zariski open dense subset  $U_i \subset \Omega_n(d_1, \dots, d_n)$  such that for every  $F \in U_i$  we have  $j^q(F) \pitchfork S_i$ . Now it is enough to take  $\Omega_n(d_1, \dots, d_n)(S_1, \dots, S_k) = \bigcap_{i=1}^k U_i$ .  $\square$

**Definition 2.2.** Let  $S_k^1 \subset J^1(n)$  denote the subvariety of 1-jets of corank  $k$ . Let  $F \in \Omega_n(d_1, \dots, d_n)$ . We say that  $F$  is one-generic if  $F$  is proper and  $j^1(F) \pitchfork S_1^1$ .

By Proposition 2.1 the subset of one-generic mappings contains a Zariski open dense subset of  $\Omega_n(d_1, \dots, d_n)$ . The following is a well known result (see for instance [6]). Our proof is based on Corollary 1.11 in [7].

**Theorem 2.3.** *Let  $F \in \Omega_n(d_1, \dots, d_n)$  be one-generic. Let  $C(F)$  denote the set of critical points of  $F$ . Then there is an open and dense subset  $U \subset C(F)$  such that for every  $a \in U$  the germ  $F_a : (\mathbb{C}^n, a) \rightarrow (\mathbb{C}^n, F(a))$  is holomorphically equivalent to a two-fold.*

*Proof.* Let  $\Delta = F(C(F))$  be the discriminant of  $F$ . Take  $U = C(F) \setminus F^{-1}(\text{Sing}(\Delta))$ . The set  $U$  is a Zariski open dense subset of  $C(F)$ . Take a point  $a \in U$  and consider the germ  $F_a : (\mathbb{C}^n, a) \rightarrow (\mathbb{C}^n, F(a))$ . By the choice of the point  $a$  the germ of the discriminant of  $F_a$  is smooth. Hence by [7], Corollary 1.11, the germ  $F_a$  is biholomorphically equivalent to a  $k$ -fold:  $(\mathbb{C}^n, 0) \ni (x_1, \dots, x_n) \mapsto (x_1^k, x_2, \dots, x_n) \in (\mathbb{C}^n, 0)$ . In particular  $\text{corank}[F_a] = 1$ .

Now note that  $J^1(n) \cong \mathbb{C}^n \times \mathbb{C}^n \times M(n, n)$ , where  $M(n, n) = \{[a_{ij}], 1 \leq i, j \leq n\}$  is the set of  $n \times n$  matrices. In these coordinates the set  $S_1^1$  is given as  $\{(x, y, m) : \det[m_{ij}] = \phi(x, y, m) = 0\}$  on the open subset  $\{(x, y, m) : \text{corank}[m_{ij}] \leq 1\}$ . Since the mapping  $j^1(F)$  is transversal to  $S_1^1$  the mapping  $\phi \circ j^1(F) = kx_1^{k-1}$  has to be a submersion at 0. This is possible only for  $k = 2$ .  $\square$

For convenience of the reader we will give a short introduction to (the simplest case of) the Thom-Boardman singularities (see e.g. [6]):

**Definition 2.4.** We define the set  $S_1^k \subset J^k(n)$  as follows:

- (1) for  $k = 1$  it is the set of elements  $\sigma = [F_a]$  such that  $\text{corank } d_a F_a = 1$ ,
- (2)  $\sigma \in S_1^k$  if for a representant  $[F_a]$  of  $\sigma$  we have  $j^{k-1}(F_a)(a) \in S_1^{k-1}$ , and  $j^{k-1}(F_a)$  is transversal to  $S_1^{k-1}$  and

$$\text{corank } d_a F_a|_{S_1^{k-1}(F_a)} = 1.$$

We prove that all varieties  $S_1^k$  are smooth (at least in all cases which we need). Consequently for a mapping  $F$  as above the varieties  $S_1^k(F) = j^k(F)^{-1}(S_1^k)$  are smooth, hence the definition makes sense. In fact we show that on a dense open subset of  $J^q(n)$  the sets  $S_1^k$  are described by algebraic equations and they are locally closed smooth varieties.

## 3. PLANE MAPPINGS

Here we will study the set  $\Omega_2(d_1, d_2)$ . Let us denote coordinates in  $J^1(2)$  by

$$(x, y, f, g, f_x, f_y, g_x, g_y).$$

For a mapping  $F = (f, g) \in \Omega_2(d_1, d_2)$ , we have

$$j^1(F) = (x, y, f(x, y), g(x, y), \frac{\partial f}{\partial x}(x, y), \frac{\partial f}{\partial y}(x, y), \frac{\partial g}{\partial x}(x, y), \frac{\partial g}{\partial y}(x, y)),$$

which justifies our notation. The set  $S_1^1$  is given by the equation  $\phi(x, y, f, g, f_x, f_y, g_x, g_y) = f_x g_y - f_y g_x = 0$ . Since  $S_1^1$  describes elements of rank one it is easy to see that it is a smooth (non-closed) subvariety of  $J^1(2)$ .

Now we would like to describe the set  $S_1^2$ . We restrict our attention only to sufficiently general mappings. In the space  $J^2(2)$  we introduce coordinates

$$(x, y, f, g, f_x, f_y, g_x, g_y, f_{xx}, f_{yy}, f_{xy}, g_{xx}, g_{yy}, g_{xy}).$$

A generic mapping  $F$  satisfies  $\text{rank } d_a F \geq 1$  for every  $a$  (because  $\text{codim } S_2^1 = 4$ ). We can assume that  $F = (f, g)$  and  $\nabla_a f \neq 0$ . The critical set of  $F$  is exactly the set  $S_1^1(F)$  and it has a reduced equation  $\frac{\partial f}{\partial x}(x, y) \frac{\partial g}{\partial y}(x, y) - \frac{\partial f}{\partial y}(x, y) \frac{\partial g}{\partial x}(x, y) = 0$ , which by simplicity we write as  $f_x g_y - f_y g_x = 0$ . In particular the tangent line to  $S_1^1(f)$  is given as

$$(f_{xx} g_y + f_x g_{xy} - f_{xy} g_x - f_y g_{xx})v + (f_{xy} g_y + f_x g_{yy} - f_{yy} g_x - f_y g_{xy})w = 0.$$

Consequently the condition for  $[F_a] \in S_1^2$  is:

$$f_x g_y - f_y g_x = 0$$

and

$$(f_{xx} g_y + f_x g_{xy} - f_{xy} g_x - f_y g_{xx})f_y - (f_{xy} g_y + f_x g_{yy} - f_{yy} g_x - f_y g_{xy})f_x = 0.$$

Let us note that the last equation contains terms  $g_{xx} f_y^2$  and  $g_{yy} f_x^2$  hence for  $\nabla f \neq 0$  these two equations form a complete intersection. In general, if we omit the assumption  $\nabla f \neq 0$  the set  $S_1^2$  is given in  $J^2(2)$  by three equations:

$$L_1 := f_x g_y - f_y g_x = 0,$$

$$L_2 := (f_{xx} g_y + f_x g_{xy} - f_{xy} g_x - f_y g_{xx})f_y - (f_{xy} g_y + f_x g_{yy} - f_{yy} g_x - f_y g_{xy})f_x = 0,$$

and

$$L_3 := (f_{xx} g_y + f_x g_{xy} - f_{xy} g_x - f_y g_{xx})g_y - (f_{xy} g_y + f_x g_{yy} - f_{yy} g_x - f_y g_{xy})g_x = 0.$$

As above by symmetry the set  $S_1^2$  is smooth and locally is given as a complete intersection of either  $L_1, L_2$  or  $L_1, L_3$ .

We will denote by  $J, J_{1,1}, J_{1,2}$  curves given by  $L_1 \circ j^1(F) = 0$ ,  $L_2 \circ j^2(F) = 0$  and  $L_3 \circ j^2(F) = 0$ . We will also identify these curves with their equations.

**Definition 3.1.** Let  $F \in \Omega_2(d_1, d_2)$ . We say that  $F$  is generic if  $F$  is proper,  $j^1(F) \pitchfork S_1^1, j^2(F) \pitchfork S_1^2$ , and additionally  $j^1(F) \pitchfork S_2^1$ .

Again by Proposition 2.1 the subset of generic mappings contains a Zariski open dense subset of  $\Omega_2(d_1, d_2)$ . Thus a general mapping is generic.

**Definition 3.2.** Let  $F : (\mathbb{C}^2, a) \rightarrow (\mathbb{C}^2, F(a))$  be a holomorphic mapping. We say that  $F$  has a simple cusp at  $a$  if  $F$  is biholomorphically equivalent to the mapping  $(\mathbb{C}^2, 0) \ni (x, y) \mapsto (x, y^3 + xy) \in (\mathbb{C}^2, 0)$ .

For the convenience of the reader we give a precise description of singularities of a generic (in the sense of Definition 3.1) plane mapping:

**Theorem 3.3.** *Let  $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$  be a generic polynomial mapping. Then  $F$  has only two-folds and simple cusps as singularities.*

*Proof.* Since  $F$  is generic,  $\text{rank } F \geq 1$ . Since  $j^1(F) \pitchfork S_1^1$ , the set  $S_1^1(F)$  is smooth. Because  $\text{rank } F \geq 1$  we have  $S_1^1(F) = C(F)$  i.e. the set of critical values of  $F$ . For a point  $a \in C(F)$  we have two possibilities:

- (1)  $\text{rank } d_a F|_{C(F)} = 1$ ,
- (2)  $\text{rank } d_a F|_{C(F)} = 0$ .

In the case (1) we see that the discriminant  $F_a(C(F_a))$  of a germ  $F_a$  is smooth and we can proceed as in the proof of Theorem 2.3. Hence  $F_a$  is two-fold. Now assume (2). We can assume that  $F_a = (x, g(x, y))$ . We have  $L_1 \circ j^1(F_a) = g_y(a) = 0$  and  $L_2 \circ j^2(F) = g_{yy}(a) = 0$ . Since  $F$  is a generic mapping, the curves  $g_y = 0$  and  $g_{yy} = 0$  intersect transversally at  $a$ . In particular we have

$$\det \begin{bmatrix} g_{yx}(a) & g_{yy}(a) \\ g_{yyx}(a) & g_{yyy}(a) \end{bmatrix} \neq 0.$$

Since  $g_{yy}(a) = 0$  it is easy to see that  $g_{yyy}(a) \neq 0$  and  $g_{xy}(a) \neq 0$ . We can assume that  $a = (0, 0)$  and  $g(0, 0) = 0$ . By the Weierstrass Preparation Theorem we have

$$g(x, y) = v(x, y)(y^3 + a_1(x)y^2 + a_2(x)y + a_3(x)),$$

where  $v, a_i$  are holomorphic and  $v(0, 0) \neq 0$ ,  $a_i(0) = 0$  for  $i = 1, 2, 3$ . By the Rouché Theorem we see that  $\mu(F_a) = \mu((x, y^3 + a_1(x)y^2 + a_2(x)y + a_3(x))) = 3$  (here  $\mu(F)$  denotes the topological degree of the mapping  $F$ ). Hence  $F_a$  is a local analytic covering of degree 3. By [5], Theorem 12, p. 104, we have

$$(*) \quad y^3 + b_1(x, g)y^2 + b_2(x, g)y + b_3(x, g) = 0.$$

Here  $b_i$  are holomorphic and  $b_i(0, 0) = 0$ . Now we follow [6], p. 148. The equality (\*) can be rewritten as

$$(**) \quad (y + b_1(x, g)/3)^3 + c(x, g)(y + b_1(x, g)/3) + d(x, g) = 0,$$

where  $c, d$  are holomorphic and  $c(0, 0) = d(0, 0) = 0$ . Since  $g(0, y) = ay^3 + \dots$ , we have from (\*\*) that  $\frac{\partial d}{\partial y}(0, 0) \neq 0$ . The inequality  $\frac{\partial c^2}{\partial x \partial y}(0, 0) \neq 0$  implies  $\frac{\partial c}{\partial x}(0, 0) \neq 0$ . Moreover (\*\*) implies  $\frac{\partial d}{\partial x}(0, 0) = 0$ . Take the coordinates

$$x_1 = c(x, g), \quad y_1 = y + b_1(x, g)/3 \quad \text{and} \quad x_2 = c(x, y), \quad y_2 = -d(x, y).$$

We have  $F_a^*(x_2, y_2) = (c(x, g), -d(x, g)) = (c(x, g), (y + b_1(x, g)/3)^3 + c(x, g)(y + b_1(x, g)/3)) = (x_1, y_1^3 + x_1 y_1)$ .  $\square$

Now we compute the number of cusps of a general polynomial mapping  $F \in \Omega_2(d_1, d_2)$ . To do this we need the series of lemmas:

**Lemma 3.4.** *Let  $L_\infty$  denote the line at infinity of  $\mathbb{C}^2$ . There is a non-empty open subset  $V \subset \Omega_2(d_1, d_2)$  such that for all  $(f, g) \in V$ :*

- (1)  $\left\{ \frac{\partial f}{\partial x} = 0 \right\} \pitchfork \left\{ \frac{\partial f}{\partial y} = 0 \right\}$ ,
- (2)  $\left\{ \frac{\partial f}{\partial x} = 0 \right\} \cap \left\{ \frac{\partial f}{\partial y} = 0 \right\} \cap L_\infty = \emptyset$ .

*Proof.* The case  $d_1 = 1$  is trivial so assume  $d_1 > 1$ . Let us note that the set  $S \subset J^1(2)$  given by  $\{f_x = f_y = 0\}$  is smooth. Hence (1) follows from Proposition 2.1. To prove (2) it is enough to assume that  $f \in H_d$ , where  $H_d$  denotes the set of homogenous polynomials of two variables of degree  $d$ . Let  $\Psi : H_d \times (\mathbb{C} \times \mathbb{C}) \setminus \{0, 0\} \ni (f, x, y) \mapsto (\frac{\partial f}{\partial x}(x, y), \frac{\partial f}{\partial y}(x, y)) \in \mathbb{C}^2$ . It is easy to see that  $\Psi$  is a submersion. Indeed, if  $f = \sum a_i x^{d-i} y^i$  then  $f_x := \frac{\partial f}{\partial x}(x, y) = da_0 x^{d-1} + \dots + a_{d-1} y^{d-1}$ ,  $f_y := \frac{\partial f}{\partial y}(x, y) = a_1 x^{d-1} + \dots + da_d y^{d-1}$ . Since  $(x, y) \neq (0, 0)$  we can assume by symmetry that  $y \neq 0$ . Now  $\frac{\partial f_x}{\partial a_{d-1}} = y^{d-1}$ ,  $\frac{\partial f_x}{\partial a_d} = 0$ ,  $\frac{\partial f_y}{\partial a_d} = dy^{d-1}$ . Thus  $\frac{\partial(f_x, f_y)}{\partial(a_{d-1}, a_d)} = dy^{2(d-1)} \neq 0$ .

Hence for a general polynomial  $f \in H_d$  the mapping  $\Psi_f : (\mathbb{C} \times \mathbb{C}) \setminus \{0, 0\} \ni (x, y) \mapsto (\frac{\partial f}{\partial x}(x, y), \frac{\partial f}{\partial y}(x, y)) \in \mathbb{C}^2$  is transversal to the point  $(0, 0)$ . In particular  $\Psi_f^{-1}(0, 0)$  is either zero-dimensional or the empty set. Since  $f$  is a homogenous polynomial the first possibility is excluded. This means that  $\overline{\left\{ \frac{\partial f}{\partial x} = 0 \right\}} \cap \overline{\left\{ \frac{\partial f}{\partial y} = 0 \right\}} \cap L_\infty = \emptyset$ .  $\square$

**Lemma 3.5.** *Let  $L_\infty$  denote the line at infinity of  $\mathbb{C}^2$ . There is a non-empty open subset  $V \subset \Omega_2(d_1, d_2)$  such that for all  $F = (f, g) \in V$ :*

- (1)  $\overline{J(F)} \cap \overline{J_{1,1}(F)} \cap L_\infty = \emptyset$ ,
- (2)  $\overline{J(F)} \pitchfork L_\infty$ .

*Proof.* Since the case  $d_1 = d_2 = 1$  is trivial and the assertion does not depend on replacing  $(f, g)$  with  $(g, f)$  we may assume that  $d_2 > 1$ . We consider the (general) case when  $\deg f = d_1$  and  $\deg g = d_2$ . Hence  $\overline{J(F)} \cap L_\infty$  and  $\overline{J_{1,1}(F)} \cap L_\infty$  depend only on the homogeneous parts of  $f$  and  $g$  of degree  $d_1$  and  $d_2$  respectively. Let  $H_d$  denote the set of homogeneous polynomials of degree  $d$  in two variables. It is sufficient to show that there is an open subset  $V \subset H_{d_1, d_2} := H_{d_1} \times H_{d_2}$  such that  $\overline{J(F)} \cap \overline{J_{1,1}(F)} \cap L_\infty = \emptyset$  for all  $F = (f, g) \in V$ .

Consider the set  $X = \{(p, F) \in \mathbb{P}^1 \times H_{d_1, d_2} : J(F)(p) = J_{1,1}(F)(p) = 0\}$ . Note that  $X$  is a closed subset of  $\mathbb{P}^1 \times H_{d_1, d_2}$ , and if  $\overline{J(F)} \cap \overline{J_{1,1}(F)} \cap L_\infty \neq \emptyset$  then  $F$  belongs to the image of the projection of  $X$  on  $H_{d_1, d_2}$ . So to prove (1) it is sufficient to show that  $X$  has dimension strictly smaller than the dimension of  $H_{d_1, d_2}$ .

Let  $q = (1 : 0) \in \mathbb{P}^1$ ,  $Y := \{q\} \times H_{d_1, d_2}$  and  $X_0 = X \cap Y$ . Note that all fibers of the projection  $X \rightarrow \mathbb{P}^1$  are isomorphic to  $X_0$ . Thus  $\dim(X) = \dim(X_0) + \dim(\mathbb{P}^1)$  and to prove (1) it is sufficient to show that  $X_0$  has codimension at least 2 in  $Y$ .

Let  $p = (q, F) \in Y$  and let  $a_i$  and  $b_i$  be the parameters in  $H_{d_1, d_2}$  giving respectively the coefficients of  $f$  at  $x_1^{d_1-i} x_2^i$  and of  $g$  at  $x_1^{d_2-i} x_2^i$ . For  $0 \leq i + j \leq d_1$ , we have  $\frac{\partial^{i+j} f}{\partial x_1^i \partial x_2^j}(q) = \frac{(d_1-j)! j!}{(d_1-i-j)!} a_j(F)$  and similarly for  $g$  and  $b_j$ .

To conclude the proof of (1) we will show that the codimension of  $\{a_0 = 0\} \cap X_0$  in  $Y$  is at least 2 and  $\nabla J$  and  $\nabla J_{1,1}$  are linearly independent outside  $\{a_0 = 0\} \cap X_0$  and thus the variety  $X_0$  has codimension 2 in  $Y$ .

Let us calculate  $J(p)$ . We have  $J(p) = (f_x g_y - f_y g_x)(q, F) = (d_1 a_0 b_1 - d_2 a_1 b_0)(F)$ . Thus  $\{a_0 = 0\} \cap X_0 \subset \{a_0 = a_1 b_0 = 0\} \cap Y$  has codimension at least 2 and we may assume  $a_0(F) \neq 0$  in further calculations.

We have  $\frac{\partial J}{\partial b_1}(p) = \frac{\partial d_1 a_0 b_1 - d_2 a_1 b_0}{\partial b_1}(F) = d_1 a_0(F)$  and  $\frac{\partial J(p)}{\partial b_2} = 0$ . Now let us calculate  $\frac{\partial J_{1,1}}{\partial b_2}(p)$ . The coefficient  $b_2$  can only be obtained from  $\frac{\partial^2 g}{\partial x_2^2}$ , which is present in  $J_{1,1}$

in the summand  $-2\frac{\partial^2 g}{\partial x_2^2}(d_1\frac{\partial f}{\partial x_1})^2$ . Thus  $\frac{\partial J_{1,1}}{\partial b_2}(p) = \frac{\partial(-2d_1^2 b_2 a_0^2)}{\partial b_2}(F) = -2(d_1 a_0(F))^2$ . So  $\det \frac{\partial(J, J_{1,1})}{\partial(b_1, b_2)}(p) = -2(d_1 a_0(F))^3 \neq 0$ .

To prove (2) note that  $\overline{\{\frac{\partial J}{\partial x}(F) = 0\}} \cap \overline{\{\frac{\partial J}{\partial y}(F) = 0\}} \subset \overline{J_{1,1}(F)}$ , hence (1) implies (2).  $\square$

**Lemma 3.6.** *There is a non-empty open subset  $V_1 \subset \Omega_2(d_1, d_2)$  such that for all  $(f, g) \in V_1$  and every  $a \in \mathbb{C}^2$ : if  $\frac{\partial f}{\partial x}(a) = 0$  and  $\frac{\partial f}{\partial y}(a) = 0$ , then  $\frac{\partial g}{\partial x}(a) \neq 0$  and  $\frac{\partial g}{\partial y}(a) \neq 0$ .*

*Proof.* Let us consider two subsets in  $J^1(2)$ :  $S_1 := \{(x, y, f, g, f_x, f_y, g_x, g_y) : f_x = 0, f_y = 0, g_x = 0\}$  and  $S_2 := \{(x, y, f, g, f_x, f_y, g_x, g_y) : f_x = 0, f_y = 0, g_y = 0\}$ . By Proposition 2.1 there is a non-empty open subset  $V_1 \subset \Omega_2(d_1, d_2)$  such that for every  $F \in V_1$  the mapping  $j^1(F)$  is transversal to  $S_1$  and  $S_2$ . Since these subsets have codimension three, we see that the image of  $j^1(F)$  is disjoint with  $S_1$  and  $S_2$ .  $\square$

**Lemma 3.7.** *There is a non-empty open subset  $V_2 \subset \Omega_2(d_1, d_2)$  such that for all  $(f, g) \in V_2$  we have  $\{\frac{\partial f}{\partial x} = 0\} \cap \{\frac{\partial f}{\partial y} = 0\} \cap J_{1,2}(f, g) = \emptyset$ .*

*Proof.* Let us consider the (non-closed) subvariety  $S \subset J^2(2)$  given by equations:  $f_x = 0, f_y = 0, (f_{xx}g_y + f_{xy}g_x - f_{xy}g_x - f_y g_{xx})g_y - (f_{xy}g_y + f_x g_{yy} - f_{yy}g_x - f_y g_{xy})g_x = 0, g_x \neq 0, g_y \neq 0$ . It is easy to check that  $S$  is a smooth complete intersection and it has codimension three. The set of generic mappings  $F$  which are transversal to  $S$  contains a Zariski open dense subset  $V_2 \subset \Omega_2(d_1, d_2)$ . By construction for all  $(f, g) \in V_2$  we have  $\{\frac{\partial f}{\partial x} = 0\} \cap \{\frac{\partial f}{\partial y} = 0\} \cap J_{1,2}(f, g) = \emptyset$ .  $\square$

**Lemma 3.8.** *There is a non-empty open subset  $V_3 \subset \Omega_2(d_1, d_2)$  such that for all  $(f, g) \in V_3$  the curve  $J(f, g)$  is transversal to the curve  $J_{1,1}(f, g)$ .*

*Proof.* There is a Zariski open subset  $V_3$  which contains only generic mappings which satisfy hypotheses of all lemmas above. We can also assume that the curves  $\{\frac{\partial f}{\partial x} = 0\}$  and  $\{\frac{\partial f}{\partial y} = 0\}$  intersect transversally. We have to show that the curves  $J(f, g)$  and  $J_{1,1}(f, g)$  intersect transversally at every point  $a \in J(f, g) \cap J_{1,1}(f, g)$ . If  $\nabla f \neq 0$  then it follows from transversality of the mapping  $F$  to the set  $S_1^2$ . Hence we can assume  $\{\frac{\partial f}{\partial x}(a) = 0\}$  and  $\{\frac{\partial f}{\partial y}(a) = 0\}$ . By Lemma 3.6 we have  $\frac{\partial g}{\partial x}(a) \neq 0$  and  $\frac{\partial g}{\partial y}(a) \neq 0$ . Let us denote:  $\frac{\partial f}{\partial x}(x, y) = f_x, \frac{\partial f}{\partial y}(x, y) = f_y$ , etc. It is enough to prove that in the ring  $\mathcal{O}_a^2$  we have the equality  $I = (f_x g_y - f_y g_x, (f_{xx}g_y + f_{xy}g_x - f_{xy}g_x - f_y g_{xx})f_y - (f_{xy}g_y + f_x g_{yy} - f_{yy}g_x - f_y g_{xy})f_x) = \mathfrak{m}_a$ , where  $\mathfrak{m}_a$  denotes the maximal ideal of  $\mathcal{O}_a^2$ . Put  $L = f_x g_y - f_y g_x$ . Hence  $I = (L, L_x f_y - L_y f_x)$ . Since  $g_x(a) \neq 0, g_y(a) \neq 0$ , we have

$$\begin{aligned} I &= (L, g_x[L_x f_y - L_y f_x], g_y[L_x f_y - L_y f_x]) = (L, L_x g_x f_y - L_y g_x f_x, L_x g_y f_y - L_y g_y f_x) = \\ &= (L, L_x g_y f_x - L_y g_x f_x, L_x g_y f_y - L_y g_x f_y) = (L, f_x[L_x g_y - L_y g_x], f_y[L_x g_y - L_y g_x]). \end{aligned}$$

By Lemma 3.7 we have  $[L_x g_y - L_y g_x](a) \neq 0$ , hence  $I = (f_x, f_y) = \mathfrak{m}_a$ .  $\square$

Now we are in a position to prove:

**Theorem 3.9.** *There is a Zariski open, dense subset  $U \subset \Omega_2(d_1, d_2)$  such that for every mapping  $F \in U$  the mapping  $F$  has only two-folds and cusps as singularities and the number of cusps is equal to*

$$d_1^2 + d_2^2 + 3d_1 d_2 - 6d_1 - 6d_2 + 7.$$

Moreover, if  $d_1 > 1$  or  $d_2 > 1$  then the set  $C(F)$  of critical points of  $F$  is a smooth connected curve, which is topologically equivalent to a sphere with  $g = \frac{(d_1+d_2-3)(d_1+d_2-4)}{2}$  handles and  $d_1 + d_2 - 2$  points removed.

*Proof.* If  $d_1 = d_2 = 1$  then the theorem is obvious. Hence we can assume that  $d_1 > 1$ . Assume first that also  $d_2 > 1$ . Note that every point  $a$  of the intersection of curves  $J(f, g)$  and  $J_{1,1}(f, g)$  with  $\nabla_a f \neq 0$  is a cusp. Moreover for a general mapping  $F$  points with  $\nabla_a f = 0$  are not cusps (Lemma 3.7). By Bezout Theorem we have that in  $J(f, g) \cap J_{1,1}(f, g)$  there are exactly  $(d_1 - 1)^2$  points with  $\nabla f = 0$  and that the number of cusps of a general mapping is equal to

$$(d_1 + d_2 - 2)(2d_1 + d_2 - 4) - (d_1 - 1)^2 = d_1^2 + d_2^2 + 3d_1d_2 - 6d_1 - 6d_2 + 7.$$

If  $d_2 = 1$  then we can modify the definition of  $J^2(2)$  (and other spaces) and replace it by its subspace given by equations  $g_{xx} = g_{xy} = g_{yy} = 0$ . Now again all submersions considered by us remain to be submersions and we can proceed as above. We leave the details to the reader.

Finally by Lemma 3.5 we have that  $C(F) = S_1^1(F)$  is a smooth affine curve which is transversal to the line at infinity. This means that  $\overline{C(F)}$  is also smooth at infinity, hence it is a smooth projective curve of degree  $d = d_1 + d_2 - 2$ . Hence by the Riemann-Roch Theorem the curve  $\overline{C(F)}$  has genus  $g = \frac{(d-1)(d-2)}{2}$ . This means in particular that  $\overline{C(F)}$  is homeomorphic to a sphere with  $g = \frac{(d-1)(d-2)}{2}$  handles. Moreover, by the Bezout Theorem it has precisely  $d$  points at infinity.  $\square$

#### 4. THE DISCRIMINANT

Here we analyze the discriminant of a general mapping from  $\Omega(d_1, d_2)$ . Let us recall that the discriminant of the mapping  $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$  is the curve  $\Delta(F) := F(C(F))$ , where  $C(F)$  is the critical curve of  $F$ . We have:

**Lemma 4.1.** *There is a non-empty open subset  $U \subset \Omega_2(d_1, d_2)$  such that for every mapping  $F \in U$ :*

- (1)  $F|_{C(F)}$  is injective outside a finite set,
- (2) if  $p \in \Delta(F)$  then  $|F^{-1}(p) \cap C(F)| \leq 2$ ,
- (3) if  $|F^{-1}(p) \cap C(F)| = 2$  then the curve  $\Delta(F)$  has a normal crossing at  $p$ .

*Proof.* We may assume that  $d_1 \geq d_2$ . Let  $\Omega_2^*(d_1, d_2)$  be the set of  $(f, g) \in \Omega_2(d_1, d_2)$  such that  $g - g(0, 0)$  is not 0 and does not divide  $f - f(0, 0)$ . Note that  $\Omega_2^*(d_1, d_2)$  is a non-empty open subset of  $\Omega_2(d_1, d_2)$  and if  $F \in \Omega_2^*(d_1, d_2)$  and  $\alpha \in \Omega_2(1, 1)$  is an affine automorphism of  $\mathbb{C}^2$  then  $F \circ \alpha \in \Omega_2^*(d_1, d_2)$ .

To prove (1) consider the set  $X = \{(p, q, F) \in \mathbb{C}^2 \times \mathbb{C}^2 \times \Omega_2^*(d_1, d_2) : p \neq q, F(p) = F(q), J(F)(p) = J(F)(q) = 0\}$ . We will show that  $X$  has dimension not greater than  $\dim \Omega_2^*(d_1, d_2)$ . So the projection of  $X$  on  $\Omega_2^*(d_1, d_2)$  has finite fibers on some open subset  $U \subset \Omega_2^*(d_1, d_2)$ . Moreover if the fiber over  $F$  is finite then  $F|_{C(F)}$  is injective outside a finite set given by the fiber.

Let  $p = (0, 0)$ ,  $q = (0, 1)$ ,  $Y := \{p\} \times \{q\} \times \Omega_2^*(d_1, d_2)$  and  $X_0 = X \cap Y$ . Note that all fibers of the projection  $X \rightarrow \mathbb{C}^2 \times \mathbb{C}^2$  are isomorphic to  $X_0$ . Thus  $\dim(X) = \dim(X_0) + 4$  and to prove (1) it is sufficient to show that  $X_0$  has codimension at least 4 in  $Y$ .

Let  $(p, q, F) \in Y$  and let  $a_{ij}$  and  $b_{ij}$  be the parameters in  $\Omega_2(d_1, d_2)$  giving respectively the coefficients of  $f$  and  $g$  at  $x^i y^j$ . For  $0 \leq i + j \leq d_1$ , we have  $\frac{\partial^{i+j} f}{\partial x^i \partial y^j}(p) = i!j!a_{ij}(F)$  and  $\frac{\partial^{i+j} f}{\partial x^i \partial y^j}(q) = i! \sum_{k=j}^{d_1-i} \frac{k!}{(k-j)!} a_{ij}(F)$  and similarly for  $g$  and  $b_{ij}$ .

The condition  $F(p) = F(q)$  yields the equations  $w_1 = \sum_{j=1}^{d_1} a_{0j}(F) = 0$  and  $w_2 = \sum_{j=1}^{d_2} b_{0j}(F) = 0$ , the conditions  $J(F)(p) = J(F)(q) = 0$  give  $w_3 = (a_{10}b_{01} - a_{01}b_{10})(F) = 0$  and  $w_4 = (\sum_{j=0}^{d_1-1} a_{1j} \sum_{j=1}^{d_2} j b_{0j} - \sum_{j=1}^{d_1} j a_{0j} \sum_{j=0}^{d_2-1} b_{1j})(F) = 0$ . If  $d_2 \geq 2$  then note that the matrix  $\frac{\partial(w_1, w_2, w_3, w_4)}{\partial(a_{01}, b_{01}, a_{10}, a_{11})}$  is triangular and its determinant is equal to  $b_{01}(F) \sum_{j=1}^{d_2} b_{0j}(F)$ . Calculating similar derivations with  $a_{10}$  replaced by  $b_{10}$  or  $a_{11}$  replaced by  $b_{11}$  we obtain that  $\nabla w_1, \dots, \nabla w_4$  are independent outside  $S = \{a_{01}(F) = b_{01}(F) = 0\} \cup \{\sum_{j=1}^{d_2} a_{0j}(F) = \sum_{j=1}^{d_2} b_{0j}(F) = 0\}$ . Thus  $X_0 \setminus S$  has codimension 4 in  $Y$  and it is easy to see that  $X_0 \cap S$  has also codimension at least 4.

If  $d_2 = 1$  then we have  $w_2 = b_{01}(F) = 0$ . Since  $F = (f, g) \in \Omega_2^*(d_1, 1)$  we have  $b_{10}(F) \neq 0$  and  $x$  does not divide  $f - f(0, 0)$ , i.e.  $a_{0j}(F) \neq 0$  for some  $j \geq 1$ . Moreover we may take  $w_3 = a_{01}(F) = 0$  and  $w_4 = \sum_{j=1}^{d_1} j a_{0j}(F) = 0$ . If  $d_1 = 2$  then we obtain a contradiction thus showing that  $X$  is in fact empty. If  $d_1 \geq 3$  then calculating  $\det \frac{\partial(w_1, w_2, w_3, w_4)}{\partial(b_{01}, a_{01}, a_{02}, a_{03})} = 1$  we obtain that  $X_0$  has codimension 4.

To prove (2) consider the set  $X = \{(p, q, r, F) \in \mathbb{C}^2 \times \mathbb{C}^2 \times \mathbb{C}^2 \times \Omega_2^*(d_1, d_2) : p \neq q \neq r \neq p, F(p) = F(q) = F(r), J(F)(p) = J(F)(q) = J(F)(r) = 0\}$ . Similarly as in (1) we compute that  $X$  has codimension at least 7. It follows that the projection of  $X$  on  $\Omega_2^*(d_1, d_2)$  has empty fibers on some open subset  $U \subset \Omega_2^*(d_1, d_2)$ . Note that unlike in (1) there are two types of fibers of the projection onto  $\mathbb{C}^6$ :  $X_0 := \{((0, 0), (1, 0), (0, 1))\} \times \Omega_2^*(d_1, d_2) \cap X$  and  $X_t := \{((0, 0), (0, 1), (0, t))\} \times \Omega_2^*(d_1, d_2) \cap X$ . In both cases the computation is purely technical and similar to the computation in (1), so we leave the details to the reader.

To prove (3) note that if  $q \in C(F)$  then  $d_q F(T_q C(F))$  is spanned by the vector  $(J_{1,1}(F)(q), J_{1,2}(F)(q))$ . Thus if  $F^{-1}(p) \cap C(F) = \{q_1, q_2\}$  then  $\Delta(F)$  has a normal crossing at  $p$  if and only if  $(J_{1,1}(F)(q_1), J_{1,2}(F)(q_1))$  and  $(J_{1,1}(F)(q_2), J_{1,2}(F)(q_2))$  are independent, i.e.  $J_{1,1}(F)(q_1)J_{1,2}(F)(q_2) - J_{1,2}(F)(q_1)J_{1,1}(F)(q_2) \neq 0$ . Similarly as in (1) let us consider the set  $X = \{(p, q, F) \in \mathbb{C}^2 \times \mathbb{C}^2 \times \Omega_2^*(d_1, d_2) : p \neq q, F(p) = F(q), J(F)(p) = J(F)(q) = J_{1,1}(F)(p)J_{1,2}(F)(q) - J_{1,2}(F)(p)J_{1,1}(F)(q) = 0\}$ . One can compute that  $X$  has codimension at least 5, thus the projection of  $X$  on  $\Omega_2^*(d_1, d_2)$  has empty fibers on some open subset  $U \subset \Omega_2^*(d_1, d_2)$ .  $\square$

Hence for a general  $F$  the only singularities of  $\Delta(F)$  are cusps and nodes. We showed in Theorem 3.9 that there are exactly  $c(F) = d_1^2 + d_2^2 + 3d_1d_2 - 6d_1 - 6d_2 + 7$  cusps. Now we will compute the number  $d(F)$  of nodes of  $\Delta(F)$ . We will use the following theorem of Serre (see [10], p. 85):

**Theorem 4.2.** *If  $\Gamma$  is an irreducible curve of degree  $d$  and genus  $g$  in the complex projective plane then*

$$\frac{1}{2}(d-1)(d-2) = g + \sum_{z \in \text{Sing}(\Gamma)} \delta_z,$$

where  $\delta_z$  denotes the delta invariant of a point  $z$ .

First we compute the degree of the discriminant:

**Lemma 4.3.** *Let  $F = (f, g) \in \Omega(d_1, d_2)$  be a general mapping. If  $d_1 \geq d_2$  then  $\deg \Delta(F) = d_1(d_1 + d_2 - 2)$ .*

*Proof.* Let  $L \subset \mathbb{C}^2$  be a generic line  $\{ax + by + c = 0\}$ . Then  $L$  intersects  $\Delta(F)$  in smooth points and  $\deg \Delta(F) = \#L \cap \Delta(F)$ . If  $j : C(F) \rightarrow \Delta(F)$  is a mapping induced by  $F$  then  $\#L \cap \Delta(F) = \#j^{-1}(L \cap \Delta(F))$ . The curve  $j^{-1}(L) = \{af + bg + c = 0\}$  has no common points at infinity with  $C(F)$ . Hence by Bezout Theorem we have  $\#j^{-1}(L \cap \Delta(F)) = (\deg j^{-1}(L))(\deg C(F)) = d_1(d_1 + d_2 - 2)$ . Consequently  $\deg \Delta(F) = d_1(d_1 + d_2 - 2)$ .  $\square$

We have the following method of computing the delta invariant in the case of one analytic branch (see [10], p. 92-93):

**Theorem 4.4.** *Let  $V_0 \subset \mathbb{C}^2$  be an irreducible germ of an analytic curve with the Puiseux parametrization of the form*

$$z_1 = t^{a_0}, \quad z_2 = \sum_{i>0} \lambda_i t^{a_i}, \quad \text{where } \lambda_i \neq 0, \quad a_1 < a_2 < a_3 < \dots$$

Let  $D_j = \gcd(a_0, a_1, \dots, a_{j-1})$ . Then

$$\delta_0 = \frac{1}{2} \sum_{j \geq 1} (a_j - 1)(D_j - D_{j+1}).$$

If  $V = \bigcup_{i=1}^r V_i$  has  $r$  branches then

$$\delta(V) = \sum_{i=1}^r \delta(V_i) + \sum_{i < j} V_i \cdot V_j,$$

where  $V \cdot W$  denotes the intersection product.

The main result of this section will be based on the following:

**Theorem 4.5.** *Let  $F \in \Omega(d_1, d_2)$  be a general mapping. Let  $d_1 \geq d_2$  and  $d = \gcd(d_1, d_2)$ . Denote by  $\overline{\Delta}$  the projective closure of the discriminant  $\Delta$ . Then*

$$\sum_{z \in (\overline{\Delta} \setminus \Delta)} \delta_z = \frac{1}{2} d_1 (d_1 - d_2) (d_1 + d_2 - 2)^2 + \frac{1}{2} (-2d_1 + d_2 + d) (d_1 + d_2 - 2).$$

*Proof.* Let  $\tilde{f}(x, y, z) = z^{d_1} f(\frac{x}{z}, \frac{y}{z})$  and  $\tilde{g}(x, y, z) = z^{d_2} g(\frac{x}{z}, \frac{y}{z})$  be the homogenizations of  $f$  and  $g$  and let  $\overline{f}(x, z) = \tilde{f}(x, 1, z)$  and  $\overline{g}(x, z) = \tilde{g}(x, 1, z)$ . For a general mapping the curves  $C(F)$  and  $\{f = 0\}$  have no common points at infinity (see Lemma 4.6). Moreover by a linear change of coordinates in the domain we can ensure that  $(1 : 0 : 0) \notin \overline{C(F)}$ . Thus  $F$  extends to a neighborhood of  $\overline{C(F)} \cap L_\infty$  on which it is given by the formula

$$\overline{F}(x, z) = \left( z^{d_1 - d_2} \frac{\overline{g}(x, z)}{\overline{f}(x, z)}, \frac{z^{d_1}}{\overline{f}(x, z)} \right).$$

Let  $\{P_1, \dots, P_{d_1 + d_2 - 2}\} = \overline{C(F)} \cap L_\infty$ , fix a point  $P = P_i$ . The curve  $\overline{C(F)}$  is transversal to the line at infinity so it has a local parametrization at  $P$  of the form  $\gamma(t) := (\sum_i a_i t^i, t)$ . We have the following:

**Lemma 4.6.** *If  $F$  is a general mapping then  $\overline{f}(P) \neq 0$ ,  $\overline{g}(P) \neq 0$  and*

$$\overline{f}(\gamma(t)) = \overline{f}(P)(1 + ct + \dots), \quad \overline{g}(\gamma(t)) = \overline{g}(P)(1 + dt + \dots),$$

where  $cd \neq 0$  and  $d_2 c \neq d_1 d$ .

*Proof.* Let  $L_\infty$  be the line at  $\{z = 0\}$  in  $\mathbb{P}^2$  and let  $\tilde{J}$  be the homogenization of  $J$ . Obviously  $\tilde{J}(F) = \frac{\partial \tilde{f}}{\partial x} \frac{\partial \tilde{g}}{\partial y} - \frac{\partial \tilde{f}}{\partial y} \frac{\partial \tilde{g}}{\partial x}$ . To prove that  $\tilde{f}(P) \neq 0$  consider the set  $X = \{(p, F) \in L_\infty \times \Omega_2(d_1, d_2) : \tilde{f}(p) = \tilde{J}(p) = 0\}$ . Clearly  $X$  has codimension 2 in  $L_\infty \times \Omega_2(d_1, d_2)$  so a general fiber of the projection on  $\Omega_2(d_1, d_2)$  is empty. Similarly we obtain  $\tilde{g}(P) \neq 0$ .

Now let  $\bar{J}(x, z) = \tilde{J}(x, 1, z)$ . Since  $\bar{J}(\gamma(t)) = 0$  and  $\frac{\partial \gamma(t)}{\partial t}|_{t=0} = (a_1, 1)$  we have

$$\begin{aligned} \bar{f}(P)c \frac{\partial \bar{J}(P)}{\partial x} &= \left( \frac{\partial \bar{f}(\gamma(t))}{\partial t} \frac{\partial \bar{J}(\gamma(t))}{\partial x} \right)_{|t=0} = \\ &= \left( \frac{\partial \bar{f}(\gamma(t))}{\partial x} a_1 \frac{\partial \bar{J}(\gamma(t))}{\partial x} + \frac{\partial \bar{f}(\gamma(t))}{\partial z} \frac{\partial \bar{J}(\gamma(t))}{\partial x} \right)_{|t=0} = \\ &= \left( -\frac{\partial \bar{f}(\gamma(t))}{\partial x} \frac{\partial \bar{J}(\gamma(t))}{\partial z} + \frac{\partial \bar{f}(\gamma(t))}{\partial z} \frac{\partial \bar{J}(\gamma(t))}{\partial x} \right)_{|t=0} = \frac{\partial \bar{f}(P)}{\partial z} \frac{\partial \bar{J}(P)}{\partial x} - \frac{\partial \bar{f}(P)}{\partial x} \frac{\partial \bar{J}(P)}{\partial z} \end{aligned}$$

Consider the set

$$X = \left\{ (p, F) \in L_\infty \times \Omega_2(d_1, d_2) : \tilde{J}(F)(p) = \left( \frac{\partial \tilde{f}}{\partial z} \frac{\partial \tilde{J}}{\partial x} - \frac{\partial \tilde{f}}{\partial x} \frac{\partial \tilde{J}}{\partial z} \right) (p) = 0 \right\}.$$

Note that if  $c = 0$  then the fiber over  $P$  of the projection from  $X$  to  $L_\infty$  is non-empty. Hence it suffices to prove that  $X$  has codimension at least 2. Similarly as in the proof of Lemma 4.1 we take  $p = (0 : 1 : 0)$  and  $Y := \{p\} \times \Omega_2(d_1, d_2)$  and show that  $X \cap Y$  has codimension 2 in  $Y$ . Let  $a_{ij}$  be the parameters in  $\Omega_2(d_1, d_2)$  giving the coefficients of  $\tilde{f}$  at  $x^i y^{d_1-i-j} z^j$  (i.e. of  $f$  at  $x^i y^{d_1-i-j}$ ) and let  $a_{ij}$  describe the coefficients of  $\tilde{g}$ .

The first equation of  $X \cap Y$  is  $d_2 a_{01} b_{00} - d_1 b_{01} a_{00}$ , denote the second one by  $w$ . We have  $\frac{\partial w}{\partial a_{02}} = 2d_2 b_{00} a_{10}$ ,  $\frac{\partial w}{\partial b_{02}} = -2d_1 a_{00} a_{10}$ ,  $\frac{\partial w}{\partial a_{11}} = d_2 b_{00} a_{01}$  and  $\frac{\partial w}{\partial b_{11}} = d_1 a_{00} a_{01}$ , so the equations are independent outside the set  $\{a_{10} = a_{01} = 0\} \cup \{a_{00} = b_{00} = 0\}$ . Thus  $X \cap Y$  has codimension 2 in  $Y$ .

Finally note that if  $d_2 c = d_1 d$  then

$$d_2 \bar{g}(P) \left( \frac{\partial \bar{f}}{\partial z} \frac{\partial \bar{J}}{\partial x} - \frac{\partial \bar{f}}{\partial x} \frac{\partial \bar{J}}{\partial z} \right) (P) = d_1 \bar{f}(P) \left( \frac{\partial \bar{g}}{\partial z} \frac{\partial \bar{J}}{\partial x} - \frac{\partial \bar{g}}{\partial x} \frac{\partial \bar{J}}{\partial z} \right) (P).$$

Hence we consider the set

$$X = \left\{ (p, F) \in L_\infty \times \Omega_2(d_1, d_2) : \tilde{J}(F)(p) = d_2 \tilde{g}(p) \left( \frac{\partial \tilde{f}}{\partial z} \frac{\partial \tilde{J}}{\partial x} - \frac{\partial \tilde{f}}{\partial x} \frac{\partial \tilde{J}}{\partial z} \right) (p) - d_1 \tilde{f}(p) \left( \frac{\partial \tilde{g}}{\partial z} \frac{\partial \tilde{J}}{\partial x} - \frac{\partial \tilde{g}}{\partial x} \frac{\partial \tilde{J}}{\partial z} \right) (p) = 0 \right\}.$$

Similarly as above one can show that it has codimension 2, which concludes the proof.  $\square$

Let  $C_p$  be the branch of  $\overline{C(F)}$  at  $P$ . We find the Puiseux expansion of the branch  $\overline{F}(C_p)$  of  $\overline{\Delta(F)}$  at  $\overline{F}(P)$ . We have

$$\begin{aligned} \overline{F}(\gamma(t)) &= \left( t^{d_1-d_2} \frac{\bar{g}(\gamma(t))}{\bar{f}(\gamma(t))}, \frac{t^{d_1}}{\bar{f}(\gamma(t))} \right) = \\ &= \left( t^{d_1-d_2} (1 + (d-c)t + \dots) \frac{\bar{g}(P)}{\bar{f}(P)}, \frac{t^{d_1} (1 - ct + \dots)}{\bar{f}(P)} \right). \end{aligned}$$

If  $d_1 = d_2$  then by Lemma 4.6 we have  $d - c \neq 0$  and  $\overline{F}(C_p)$  is smooth at  $\overline{F}(P)$ . So assume  $d_1 > d_2$ . Since the function  $h(t) = \left( \frac{\bar{f}(P) \bar{g}(\gamma_P(t))}{\bar{g}(P) \bar{f}(\gamma_P(t))} \right)^{\frac{1}{d_1-d_2}} = 1 + \frac{d-c}{d_1-d_2} t + \dots$

is invertible in  $t = 0$  we can introduce a new variable  $T = th(t)$ . We have  $\overline{F}(\gamma(T)) = \left( T^{d_1-d_2} \frac{\overline{g}(P)}{\overline{f}(P)}, T^{d_1} h(t)^{-d_1} (1-ct+\dots) \frac{1}{\overline{f}(P)} \right)$ . Moreover  $h(t)^{-d_1} (1-ct+\dots) = (1-d_1 \frac{d-c}{d_1-d_2} T + \dots)(1-cT+\dots) = 1 + \frac{d_2c-d_1d}{d_1-d_2} T + \dots$ . By Lemma 4.6 we have  $d_2c - d_1d \neq 0$  and we can apply Theorem 4.4 to compute  $\delta(C_P)_{\overline{F}(P)}$ . Since  $a_0 = d_1 - d_2$ ,  $a_1 = d_1$  and  $a_2 = d_1 + 1$ , we have  $2\delta(C_P)_{\overline{F}(P)} = (d_1 - 1)(d_1 - d_2 - d) + (d_1 + 1 - 1)(d - 1) = (d_1 - 1)(d_1 - d_2 - 1) + (d - 1)$ .

To proceed further we also need:

**Lemma 4.7.** *If  $F$  is a general mapping then*

$$\overline{f}(P_i)^{d_2} \overline{g}(P_j)^{d_1} \neq \overline{f}(P_j)^{d_2} \overline{g}(P_i)^{d_1}$$

for  $i, j \in \{1, 2, \dots, d_1 + d_2 - 2\}$  and  $i \neq j$ .

*Proof.* Consider the set  $X = \{(p, q, F) \in L_\infty \times L_\infty \times \Omega_2(d_1, d_2) : p \neq q, \tilde{J}(F)(p) = \tilde{J}(F)(q) = \tilde{f}(p)^{d_2} \tilde{g}(q)^{d_1} - \tilde{f}(q)^{d_2} \tilde{g}(p)^{d_1} = 0\}$ . Since one can prove similarly as in Lemma 4.6 that  $X$  has codimension 3, there is a dense open subset  $S \subset \Omega(d_1, d_2)$  such that the projection from  $X$  has empty fibers over  $F \in S$ .  $\square$

Now we are in a position to compute  $\sum_{z \in (\overline{\Delta} \setminus \Delta)} \delta_z$ . If  $d_1 = d_2$  then  $\overline{\Delta}$  has exactly  $d_1 + d_2 - 2$  smooth points at infinity and consequently  $\sum_{z \in (\overline{\Delta} \setminus \Delta)} \delta_z = 0$ . So assume  $d_1 > d_2$ , then  $\overline{\Delta}$  has only one point at infinity  $Q = (1 : 0 : 0)$ . In  $Q$  the curve  $\overline{\Delta}$  has exactly  $r = d_1 + d_2 - 2$  branches  $V_i = \overline{F}(C_{P_i})$ . We computed above that  $2\delta(V_i)_Q = (d_1 - 1)(d_1 - d_2 - 1) + (d - 1)$ . Now we will compute  $V_i \cdot V_j$ . Let  $t_{a,b}(x, y) = (x + a, y + b)$ . By the dynamical definition of intersection there exists a neighborhood  $U$  of 0, such that for small general  $a, b$  we have

$$V_i \cdot V_j = \#(U \cap V_i \cap t_{a,b}(V_j)).$$

This means that  $V_i \cdot V_j$  is equal to the number of solutions of the following system:

$$\begin{aligned} \frac{\overline{g}(P_i)}{\overline{f}(P_i)} T^{d_1-d_2} &= \frac{\overline{g}(P_j)}{\overline{f}(P_j)} S^{d_1-d_2} + a, \\ \frac{1}{\overline{f}(P_i)} T^{d_1} (1 + \alpha_i T + \dots) &= \frac{1}{\overline{f}(P_j)} S^{d_1} (1 + \alpha_j S + \dots) + b, \end{aligned}$$

where  $a, b$  and  $S, T$  are sufficiently small. Take

$$Q : (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}^2, 0),$$

$$Q(T, S) = \left( \frac{\overline{g}(P_i)}{\overline{f}(P_i)} T^{d_1-d_2} - \frac{\overline{g}(P_j)}{\overline{f}(P_j)} S^{d_1-d_2}, \frac{1}{\overline{f}(P_i)} T^{d_1} (1 + \alpha_i T + \dots) - \frac{1}{\overline{f}(P_j)} S^{d_1} (1 + \alpha_j S + \dots) \right).$$

Thus we have  $V_i \cdot V_j = \text{mult}_0 Q$ . Note that by Lemma 4.7 the minimal homogenous polynomials of the two components of  $Q$  have no nontrivial common zeroes, hence  $V_i \cdot V_j = d_1(d_1 - d_2)$ . Consequently

$$\begin{aligned} \sum_i \delta(V_i) + \sum_{i>j} V_i \cdot V_j &= \frac{1}{2} [(d_1 - 1)(d_1 - d_2 - 1) + (d - 1)](d_1 + d_2 - 2) + \\ &\quad \frac{1}{2} d_1(d_1 - d_2)(d_1 + d_2 - 2)(d_1 + d_2 - 3) = \\ &\quad \frac{1}{2} d_1(d_1 - d_2)(d_1 + d_2 - 2)^2 + \frac{1}{2} (-2d_1 + d_2 + d)(d_1 + d_2 - 2). \end{aligned}$$

$\square$

We can now prove the following:

**Theorem 4.8.** *There is a Zariski open, dense subset  $U \subset \Omega_2(d_1, d_2)$  such that for every mapping  $F \in U$  the discriminant  $\Delta(F) = F(C(F))$  has only cusps and nodes as singularities. Let  $d = \gcd(d_1, d_2)$ . Then the number of cusps is equal to*

$$c(F) = d_1^2 + d_2^2 + 3d_1d_2 - 6d_1 - 6d_2 + 7$$

and the number of nodes is equal to

$$d(F) = \frac{1}{2} [(d_1d_2 - 4)((d_1 + d_2 - 2)^2 - 2) - (d - 5)(d_1 + d_2 - 2) - 6].$$

*Proof.* Let  $d_1 \geq d_2$  and  $D = d_1 + d_2 - 2$ . By Lemma 4.3 we have  $\deg \Delta(F) = d_1D$ . From Lemma 4.1 we know that  $\Delta(F)$  has only cusps and nodes as singularities and is birational with  $C(F)$ . Hence  $\Delta(F)$  has genus  $g = \frac{1}{2}(D - 1)(D - 2)$ . Thus by Theorem 4.2 we have

$$\frac{1}{2}(d_1D - 1)(d_1D - 2) = \frac{1}{2}(D - 1)(D - 2) + c(F) + d(F) + \sum_{z \in (\overline{\Delta} \setminus \Delta)} \delta_z.$$

Substituting

$$\sum_{z \in (\overline{\Delta} \setminus \Delta)} \delta_z = \frac{1}{2}d_1(d_1 - d_2)D^2 + \frac{1}{2}(-2d_1 + d_2 + d)D$$

from Theorem 4.5 we obtain

$$2(c(F) + d(F)) = d_1d_2D^2 - D^2 + 3D - d_1D - d_2D - dD = (d_1d_2 - 2)D^2 - (d - 1)D.$$

Thus by Theorem 3.9 we get:

$$\begin{aligned} d(F) &= \frac{1}{2} [(d_1d_2 - 2)D^2 - (d - 1)D - 2(D^2 - 2D + d_1d_2 - 1)] = \\ &= \frac{1}{2} [(d_1d_2 - 4)(D^2 - 2) - (d - 5)D - 6]. \end{aligned}$$

□

## 5. GENERALIZED CUSPS

In this section our aim is to estimate the number of cusps of non-generic mappings. We start from:

**Definition 5.1.** Let  $F : (\mathbb{C}^2, a) \rightarrow (\mathbb{C}^2, F(a))$  be a holomorphic mapping. We say that  $F$  has a generalized cusp at  $a$  if  $F_a$  is proper, the curve  $J(F) = 0$  is reduced near  $a$  and the discriminant of  $F_a$  is not smooth at  $F(a)$ .

**Remark 5.2.** If  $F_a$  is proper,  $J(F) = 0$  is reduced near  $a$  and  $J(F)$  is singular at  $a$  then it follows from Corollary 1.11 from [7] that also the discriminant of  $F_a$  is singular at  $F(a)$  and hence  $F$  has a generalized cusp at  $a$ .

Now we introduce the index of generalized cusp:

**Definition 5.3.** Let  $F = (f, g) : (\mathbb{C}^2, a) \rightarrow (\mathbb{C}^2, F(a))$  be a holomorphic mapping. Assume that  $F$  has a generalized cusp at a point  $a \in \mathbb{C}^2$ . Since the curve  $J(F) = 0$  is reduced near  $a$ , we have that the set  $\{\nabla f = 0\} \cap \{\nabla g = 0\}$  has only isolated points near  $a$ . For a general linear mapping  $T \in GL(2)$ , if  $F' = (f', g') = T \circ F$  then  $\nabla f'$  does not vanish identically on any branch of  $\{J(F) = 0\}$  near  $a$ . We say that the cusp of  $F$  at  $a$  has an index  $\mu_a := \dim_{\mathbb{C}} \mathcal{O}_a / (J(F'), J_{1,1}(F')) - \dim_{\mathbb{C}} \mathcal{O}_a / (f'_x, f'_y)$ .

**Remark 5.4.** We show below that the index  $\mu_a$  is well-defined and finite. Moreover, it is easy to see that a simple cusp has index one.

**Remark 5.5.** Using the exact sequence 1.7 from [2] we see that

$$\mu_a = \dim_{\mathbb{C}} \mathcal{O}_a / (J(F), J_{1,1}(F), J_{1,2}(F)).$$

Hence our index coincides with the classical local number of cusps defined e.g. in [2].

We have (compare with [2], [3], [4]):

**Theorem 5.6.** *Let  $F = (f, g) \in \Omega_2(d_1, d_2)$  and assume that  $F$  has a generalized cusp at  $a \in \mathbb{C}^2$ . If  $U_a$  is a sufficiently small ball around  $a$  then  $\mu_a$  is equal to the number of simple cusps in  $U_a$  of a general mapping  $F' \in \Omega_2(d'_1, d'_2)$ , where  $d'_1 \geq d_1, d'_2 \geq d_2$ , which is sufficiently close to  $F$  in the natural topology of  $\Omega_2(d'_1, d'_2)$ .*

*Proof.* We can assume that  $\nabla f$  does not vanish identically on any branch of  $\{J(F) = 0\}$  near  $a$ . In particular we have  $\dim \mathcal{O}_a / (f_x, f_y) = \dim \mathcal{O}_a / (J(F), f_x, f_y) < \infty$ .

Let  $F_i = (f_i, g_i) \in \Omega_2(d'_1, d'_2)$  be a sequence of general mappings, which is convergent to  $F$ . Consider the mappings  $\Phi = (J(F), J_{1,1}(F))$ ,  $\Phi_i = (J(F_i), J_{1,1}(F_i))$ ,  $\Psi = (\nabla f)$  and  $\Psi_i = (\nabla f_i)$ . Thus  $\Phi_i \rightarrow \Phi$  and  $\Psi_i \rightarrow \Psi$ .

Since  $a$  is a cusp of  $F$  we have  $\Phi(a) = 0$ . Moreover  $d_a(\Phi) < \infty$ , where  $d_a(\Phi)$  denotes the local topological degree of  $\Phi$  at  $a$ . Indeed, if  $J_{1,1}(F) = 0$  on some branch  $B$  of the curve  $J(F) = 0$  then the rank of  $F|_B$  would be zero and by Sard theorem  $F$  has to contract  $B$ , which is a contradiction ( $F_a$  is proper). By the Rouché Theorem we have that for large  $i$  the mapping  $\Phi_i$  has exactly  $d_a(\Phi)$  zeroes in  $U_a$  and  $\Psi_i$  has exactly  $d_a(\Psi)$  zeroes in  $U_a$  (counted with multiplicities, if  $\Psi(a) \neq 0$  we put  $d_a(\Psi) = 0$ ). However, the mappings  $F_i$  are general, in particular all zeroes of  $\Phi_i$  and  $\Psi_i$  are simple. Moreover the zeroes of  $\Phi_i$  which are not cusps of  $F_i$  are zeroes of  $\Psi_i$ . Hence  $\mu_a = d_a(\Phi) - d_a(\Psi)$  is indeed the number of simple cusps of  $F_i$  in  $U_a$ .  $\square$

**Corollary 5.7.** *Let  $F \in \Omega_2(d_1, d_2)$ . Assume that  $F$  has generalized cusps at points  $a_1, \dots, a_r$ . Then  $\sum_{i=1}^r \mu_{a_i} \leq d_1^2 + d_2^2 + 3d_1d_2 - 6d_1 - 6d_2 + 7$ .*

## REFERENCES

- [1] T. Fukuda, G. Ishikawa, *On the number of cusps of stable perturbations of a plane-to-plane singularity*, Tokyo J. Math. 10 (1987), no. 2, 375-384.
- [2] T. Gaffney, D.M.Q. Mond, *Cusps and double folds of germs of analytic maps  $\mathbb{C}^2 \rightarrow \mathbb{C}^2$* , J. London Math. Soc. (2) 43 (1991), no. 1, 185-192.
- [3] T. Gaffney, D.M.Q. Mond, *Weighted homogeneous maps from the plane to the plane*, Math. Proc. Cambridge Philos. Soc. 109 (1991), no. 3, 451-470.
- [4] T. Gaffney, *Polar multiplicities and equisingularity of map germs*, Topology 32 (1993), no. 1, 185-223.
- [5] R. Gunning, H. Rossi, *Analytic functions of several variables*, Prentice Hall (1965).
- [6] M. Golubitsky, V. Guillemin, *Stable mappings and their singularities*, GTM, Springer-Verlag (1973).
- [7] Z. Jelonek, *On finite regular and holomorphic mappings*, arXiv:1404.7466v3 [math.AG], 2014.
- [8] Z. Jelonek, *On semi-equivalence of generically-finite polynomial mappings*, Math. Z., (2016) DOI 10.1007/s00209-015-1591-8.
- [9] I. Krzyżanowska, Z. Szafraniec, *On polynomial mappings from the plane to the plane*, J. Math. Soc. Japan Vol. 66, no 3 (2014), 805-818.
- [10] J. Milnor, *Singular points of complex hypersurfaces*, Annals of Mathematics Studies, Princeton University Press, (1968).
- [11] H. Whitney, *On singularities of mappings of Euclidean spaces. I. Mappings of the plane into the plane*, Ann. of Math. (2) 62 (1955), 374-410.

(M. Farnik) INSTYTUT MATEMATYCZNY, POLSKA AKADEMIA NAUK, ŚNIADECKICH 8, 00-656 WARSZAWA,  
POLAND

*E-mail address:* `micchal.farnik@gmail.com`

(Z. Jelonek) INSTYTUT MATEMATYCZNY, POLSKA AKADEMIA NAUK, ŚNIADECKICH 8, 00-656 WARSZAWA,  
POLAND

*E-mail address:* `najelone@cyf-kr.edu.pl`

(M.A.S. Ruas) DEPARTAMENTO DE MATEMÁTICA, ICMC-USP, CAIXA POSTAL 668, 13560-970 SÃO  
CARLOS, S.P., BRASIL

*E-mail address:* `maasruas@icmc.usp.br`