Constructive sparse trigonometric approximation for functions with small mixed smoothness^{*}

V. Temlyakov †

August 8, 2018

Abstract

The paper gives a constructive method, based on greedy algorithms, that provides for the classes of functions with small mixed smoothness the best possible in the sense of order approximation error for the m-term approximation with respect to the trigonometric system.

1 Introduction

The paper is a follow up to the author's recent paper [\[16\]](#page-29-0). The main goal of this paper is to extend the results from $[16]$ on *m*-term trigonometric approximation in L_p of classes \mathbf{W}_q^r of functions with bounded in L_q mixed derivative of order r to the case of small smoothness r . The most important contribution of this paper, alike the paper [\[16\]](#page-29-0), is that it gives a constructive method, based on greedy algorithms, that provides for the classes \mathbf{W}_q^r the best possible in the sense of order approximation error $\sigma_m(\mathbf{W}_q^r)_p$. Theory of sparse approximation with respect to the trigonometric system has a long and interesting history. We give a brief description of this history with emphases

[∗]Math Subject Classifications. primary: 41A65; secondary: 42A10, 46B20.

[†]University of South Carolina, USA, and Steklov Institute of Mathematics, Russia. Research was supported by NSF grant DMS-1160841

put on methods of approximation. We introduce some notation. Denote by $\mathcal{T} := \{e^{ikx}\}_{k\in\mathbb{Z}}$ the univariate trigonometric system and by $\mathcal{T}^d := \mathcal{T} \times \cdots \times$ $\mathcal{T} = \{e^{i(\mathbf{k}, \mathbf{x})}\}_{\mathbf{k} \in \mathbb{Z}^d}$ the multivariate trigonometric system. Define best m-term approximations for a function

$$
\sigma_m(f)_p := \inf_{\{c_j\}, \{\mathbf{k}_j\}} \|f - \sum_{j=1}^m c_j e^{i(\mathbf{k}_j, \mathbf{x})} \|_p
$$

and for a class $\mathbf W$ of functions

$$
\sigma_m(\mathbf{W})_p := \sup_{f \in \mathbf{W}} \sigma_m(f)_p.
$$

We study m -term approximation problems for classes of functions with mixed smoothness. We begin with the definition of a smoothness class in the case of univariate periodic functions. Let for $r > 0$

$$
F_r(x) := 1 + 2 \sum_{k=1}^{\infty} k^{-r} \cos(kx - r\pi/2)
$$
 (1.1)

and

$$
W_q^r := \{ f : f = \varphi * F_r, \quad ||\varphi||_q \le 1 \}. \tag{1.2}
$$

In the multivariate case for $\mathbf{x} = (x_1, \dots, x_d)$ denote

$$
F_r(\mathbf{x}) := \prod_{j=1}^d F_r(x_j)
$$

and

$$
\mathbf{W}_q^r := \{ f : f = \varphi * F_r, \quad ||\varphi||_q \le 1 \}.
$$

For $f \in \mathbf{W}_q^r$ we denote $f^{(r)} := \varphi$ where φ is such that $f = \varphi * F_r$, and define

$$
||f||_{\mathbf{W}_q^r} := ||\varphi||_q.
$$

The first results that showed advantage of m -term approximation with respect to the univariate trigonometric system $\mathcal T$ over the classical approximation by the trigonometric polynomials of order m were obtained by Ismagilov [\[6\]](#page-28-0) in 1974. His results were improved by Maiorov [\[8\]](#page-29-1) in 1986 to the relation

$$
\sigma_m(F_2)_{\infty} \asymp m^{-3/2}.
$$

Note, that best approximation of F_2 in the uniform norm by trigonometric polynomials of degree m is of order m^{-1} . Both Ismagilov [\[6\]](#page-28-0) and Maiorov [\[8\]](#page-29-1) used constructive methods, based on number theoretical results. They considered the univariate m-term approximation with respect to \mathcal{T} . An interesting phenomenon specific for the multivariate m-term approximation was discovered in [\[11\]](#page-29-2) and [\[12\]](#page-29-3) in 1986. It was established that $\sigma_m(\mathbf{W}_q^r)_p$ decays faster than the Kolmogorov width $d_m(\mathbf{W}_q^r)_p$ for $1 < q < p \leq 2$. The proof of upper bounds for the $\sigma_m(\mathbf{W}_q^r)_p$, $1 < q \le p \le 2$, $r > 2(1/q - 1/p)$, in [\[11\]](#page-29-2) and [\[12\]](#page-29-3) is constructive. It is based on Theorem [1.6](#page-6-0) (see below). Theorem [1.6](#page-6-0) is often used in approximation of classes with mixed smoothness. We use it in this paper many times.

A very interesting and difficult case for the m-term approximation is the approximation in L_p , $p > 2$. Makovoz [\[9\]](#page-29-4) used in 1984 the probabilistic Rosenthal inequality for *m*-term approximation in L_p , $2 < p < \infty$. Later, in 1987, Belinskii [\[1\]](#page-28-1) used the Rosenthal inequality technique to prove the following lemma.

Lemma 1.1. Let $2 < p < \infty$. For any trigonometric polynomial

$$
t(\theta_n, x) := \sum_{j=1}^n c_j e^{ik_j x}, \quad \theta_n := \{k_j\}_{j=1}^n,
$$

and any $m \leq n$ there exists $t(\theta_m, x)$ with $\theta_m \subset \theta_n$ such that

$$
||t(\theta_n, x) - t(\theta_m, x)||_p \le C(p)(n/m)^{1/2} ||t(\theta_n, x)||_2.
$$

Lemma [1.1](#page-2-0) and its multivariate versions were used in a number of papers on *m*-term trigonometric approximation in L_p , $2 < p < \infty$ (see, for instance, [\[10\]](#page-29-5) and references therein). The use of Lemma [1.1](#page-2-0) allowed researchers to obtain the right orders of $\sigma_m(\mathbf{W})_p$ for different function classes **W** in L_p , $2 < p < \infty$. However, this way does not provide a constructive method of approximation. Other nonconstructive method for m-term trigonometric approximation, which is more powerful than the above discussed probabilistic method was suggested in [\[3\]](#page-28-2) in 1995. The method in [\[3\]](#page-28-2) is based on a nonconstructive result from finite dimensional geometry due to Gluskin [\[5\]](#page-28-3).

Breakthrough results in constructive m-term approximation were obtained by application of general theory of greedy approximation in Banach spaces. It is pointed out in [\[4\]](#page-28-4) (2002) that the Weak Chebyshev Greedy Algorithm provides a constructive proof of the inequality

$$
\sigma_m(f)_p \le C(p)m^{-1/2}||f||_A, \quad p \in [2, \infty).
$$

Here

$$
||f||_A := \sum_{\mathbf{k}} |\hat{f}(\mathbf{k})|, \quad \hat{f}(\mathbf{k}) := (2\pi)^{-d} \int_{\mathbb{T}^d} f(\mathbf{x}) e^{-i(\mathbf{k}, \mathbf{x})} d\mathbf{x}.
$$

In [\[13\]](#page-29-6) (2005) a constructive proof, based on the Weak Chebyshev Greedy Algorithm, was given for the following inequality

$$
\sigma_m(f)_{\infty} \leq C m^{-1/2} (\log(1 + N/m))^{1/2} ||f||_A,
$$

under assumption that f is a trigonometric polynomial of order N .

The following Theorem [1.1](#page-3-0) is from [\[16\]](#page-29-0). We use it in this paper. Let

$$
\Pi(\mathbf{N}, d) := \{ (a_1, \ldots, a_d) \in \mathbb{R}^d : |a_j| \le N_j, \ j = 1, \ldots, d \},
$$

where N_j are nonnegative integers and $\mathbf{N} := (N_1, \ldots, N_d)$. We denote

$$
\mathcal{T}(\mathbf{N},d) := \{t : t = \sum_{\mathbf{k} \in \Pi(\mathbf{N},d)} c_{\mathbf{k}} e^{i(\mathbf{k},\mathbf{x})}\}.
$$

Then

$$
\dim \mathcal{T}(\mathbf{N}, d) = \prod_{j=1}^{d} (2N_j + 1) =: \vartheta(\mathbf{N}).
$$

For a nonnegative integer m denote $\bar{m} := \max(m, 1)$.

Theorem 1.1. There exist constructive greedy-type approximation methods $G_m^p(\cdot)$, which provide m-term polynomials with respect to \mathcal{T}^d with the following properties: for $2 \leq p < \infty$

$$
||f - G_m^p(f)||_p \le C_1(d)(\bar{m})^{-1/2} p^{1/2} ||f||_A, \quad ||G_m^p(f)||_A \le C_2(d) ||f||_A, \quad (1.3)
$$

and for $p = \infty$, $f \in \mathcal{T}(\mathbf{N}, d)$

$$
||f - G_m^{\infty}(f)||_{\infty} \le C_3(d)(\bar{m})^{-1/2}(\ln \vartheta(\mathbf{N}))^{1/2} ||f||_A, \quad ||G_m^{\infty}(f)||_A \le C_4(d) ||f||_A.
$$
\n(1.4)

We now formulate the main results of the paper. The main results of the paper are in Section 3, where we consider m -term approximation in L_p with $p \in (2,\infty)$. Here is a typical result from Section 3.

Theorem 1.2. Let
$$
1 < q \leq 2 < p < \infty
$$
 and $\beta p' < r < 1/q$. Then we have\n
$$
\sigma_m(\mathbf{W}_q^r)_p \asymp m^{-(r-\beta)p/2} (\log m)^{(d-1)(r(p-1)-\beta p)}.
$$

The upper bounds are achieved by a constructive greedy-type algorithm.

Theorem [1.2](#page-3-1) complements the known result from [\[16\]](#page-29-0) for large smoothness: Let $1 < q \leq 2 < p < \infty$ and $r > 1/q$. Then we have

$$
\sigma_m(\mathbf{W}_q^r)_p \asymp m^{-r+\eta} (\log m)^{(d-1)(r-2\eta)}.
$$

The upper bounds are achieved by a constructive greedy-type algorithm.

In Section 3 we also consider the case $r = 1/q$ and more general smoothness classes $\mathbf{W}_q^{a,b}$, which we define momentarily. We introduce some more notations. Let $\mathbf{s} = (s_1, \ldots, s_d)$ be a vector whose coordinates are nonnegative integers

$$
\rho(\mathbf{s}) := \left\{ \mathbf{k} \in \mathbb{Z}^d : [2^{s_j - 1}] \le |k_j| < 2^{s_j}, \qquad j = 1, \dots, d \right\},
$$
\n
$$
Q_n := \bigcup_{\|\mathbf{s}\|_1 \le n} \rho(\mathbf{s}) \quad - \text{ a step hyperbolic cross,}
$$
\n
$$
\Gamma(N) := \left\{ \mathbf{k} \in \mathbb{Z}^d : \prod_{j=1}^d \max(|k_j|, 1) \le N \right\} \quad - \text{ a hyperbolic cross.}
$$

For $f \in L_1(\mathbb{T}^d)$

$$
\delta_{\mathbf{s}}(f) := \delta_{\mathbf{s}}(f, \mathbf{x}) := \sum_{\mathbf{k} \in \rho(\mathbf{s})} \hat{f}(\mathbf{k}) e^{i(\mathbf{k}, \mathbf{x})}.
$$

Let G be a finite set of points in \mathbb{Z}^d , we denote

$$
\mathcal{T}(G) := \left\{ t : t(\mathbf{x}) = \sum_{\mathbf{k} \in G} c_{\mathbf{k}} e^{i(\mathbf{k}, \mathbf{x})} \right\}.
$$

For the sake of simplicity we shall write $\mathcal{T}(\Gamma(N)) = \mathcal{T}(N)$.

Along with classes \mathbf{W}_q^r defined above it is natural to consider some more general classes. We proceed to the definition of these classes.

Define for $f \in L_1$

$$
f_l := \sum_{\|\mathbf{s}\|_1 = l} \delta_{\mathbf{s}}(f), \quad l \in \mathbb{N}_0, \quad \mathbb{N}_0 := \mathbb{N} \cup \{0\}.
$$

Consider the class

$$
\mathbf{W}_q^{a,b} := \{ f : ||f_l||_q \le 2^{-al}(\bar{l})^{(d-1)b} \}.
$$

Define

$$
||f||_{\mathbf{W}_q^{a,b}} := \sup_l ||f_l||_q 2^{al} (\bar{l})^{-(d-1)b}.
$$

It is well known that the class \mathbf{W}_q^r is embedded in the class $\mathbf{W}_q^{r,0}$. Classes $\mathbf{W}_q^{a,b}$ provide control of smoothness at two scales: a controls the power type smoothness and b controls the logarithmic scale smoothness. Similar classes with the power and logarithmic scales of smoothness are studied in the recent book of Triebel [\[17\]](#page-29-7).

In Section 2 we discuss the case $1 < q \leq p \leq 2$. We use the technique developed in [\[11\]](#page-29-2) and [\[12\]](#page-29-3). The main results of Section 2 are the following two theorems. We use the notation $\beta := \beta(q, p) := 1/q - 1/p$ and $\eta :=$ $\eta(q) := 1/q - 1/2.$

Theorem 1.3. Let $1 < q \leq p \leq 2$. We have

$$
\sigma_m(\mathbf{W}_q^{a,b})_p \asymp \begin{cases} m^{-a+\beta} (\log m)^{(d-1)(a+b-2\beta)}, & a > 2\beta, \\ m^{-a+\beta} (\log m)^{(d-1)b}, & \beta < a < 2\beta, \\ m^{-\beta} (\log m)^{(d-1)b} (\log \log m)^{1/q}, & a = 2\beta. \end{cases}
$$

Theorem 1.4. Let $1 < q \leq p \leq 2$, $r > \beta$. We have

$$
\sigma_m(\mathbf{W}_q^r)_p \asymp m^{-r+\beta} (\log m)^{(d-1)(r-2\beta)_+}.
$$

In the case $r > 2\beta$ Theorem [1.4](#page-5-0) is proved in [\[11\]](#page-29-2) and [\[12\]](#page-29-3) and, as it is pointed out in [\[10\]](#page-29-5), in the case $\beta \leq r \leq 2\beta$ the order of $\sigma_m(\mathbf{W}_q^r)_p$ is obtained in [\[2\]](#page-28-5). We present a detailed proof of Theorem [1.4](#page-5-0) in Section 2 for completeness (for instance, the author could not find the paper [\[2\]](#page-28-5)).

We formulate some known results from harmonic analysis and from the hyperbolic cross approximation theory, which will be used in our analysis.

Theorem 1.5. Let $1 < p < \infty$. There exist positive constants $C_1(p,d)$ and $C_2(p,d)$, which may depend on p and d, such that for each $f \in L_p$

$$
C_1(p,d)\|f\|_p \le \left\| \left(\sum_{\mathbf{s}} |\delta_{\mathbf{s}}(f,\mathbf{x})|^2 \right)^{1/2} \right\|_p \le C_2(p,d) \|f\|_p.
$$

Corollary 1.1. Let G be a finite set of indices s and let the operator S_G map a function $f \in L_p$ to a function

$$
S_G(f) := \sum_{\mathbf{s} \in G} \delta_{\mathbf{s}}(f).
$$

Then for $1 < p < \infty$

$$
||S_G(f)||_p \le C(p,d) ||f||_p.
$$

Corollary 1.2. Let $1 < p < \infty$. Denote $p_* := \max(p, 2)$ and $p^* := \min(p, 2)$. Then for $f \in L_p$ we have

$$
C_3(p,d) \left(\sum_{\mathbf{s}} \|\delta_{\mathbf{s}}(f)\|_{p}^{p_*} \right)^{1/p_*} \leq \|f\|_p \leq C_4(p,d) \left(\sum_{\mathbf{s}} \|\delta_{\mathbf{s}}(f)\|_{p}^{p^*} \right)^{1/p^*}.
$$

We now proceed to the problem of estimating $||f||_p$ in terms of the array $\{ \|\delta_{s}(f)\|_{q} \}.$ Here and below p and q are scalars such that $1 \leq q, p \leq \infty$. Let an array $\varepsilon = {\varepsilon_s}$ be given, where $\varepsilon_s \geq 0$, $s = (s_1, \ldots, s_d)$, and s_j are nonnegative integers, $j = 1, \ldots, d$. We denote by $G(\varepsilon, q)$ and $F(\varepsilon, q)$ the following sets of functions $(1 \leq q \leq \infty)$:

$$
G(\varepsilon, q) := \{ f \in L_q : ||\delta_{\mathbf{s}}(f)||_q \le \varepsilon_{\mathbf{s}} \quad \text{for all } \mathbf{s} \},
$$

$$
F(\varepsilon, q) := \{ f \in L_q : ||\delta_{\mathbf{s}}(f)||_q \ge \varepsilon_{\mathbf{s}} \quad \text{for all } \mathbf{s} \}.
$$

The following theorem is from [\[12\]](#page-29-3), p.29. For the special case $q = 2$ see [\[11\]](#page-29-2) and [\[12\]](#page-29-3), p.86.

Theorem 1.6. The following relations hold:

$$
\sup_{f \in G(\varepsilon, q)} ||f||_p \asymp \left(\sum_{\mathbf{s}} \varepsilon_{\mathbf{s}}^p 2^{\|\mathbf{s}\|_1 (p/q - 1)}\right)^{1/p}, \qquad 1 \le q < p < \infty; \tag{1.5}
$$

$$
\inf_{f \in F(\varepsilon, q)} \|f\|_{p} \asymp \left(\sum_{\mathbf{s}} \varepsilon_{\mathbf{s}}^{p} 2^{\|\mathbf{s}\|_{1}(p/q-1)}\right)^{1/p}, \qquad 1 < p < q \le \infty,\tag{1.6}
$$

with constants independent of ε .

We will need a corollary of Theorem [1.6](#page-6-0) (see [\[12\]](#page-29-3), Ch.1, Theorem 2.2), which we formulate as a theorem.

Theorem 1.7. Let $1 < q \leq 2$. For any $t \in \mathcal{T}(N)$ we have

$$
||t||_A := \sum_{\mathbf{k}} |\hat{t}(\mathbf{k})| \le C(q, d) N^{1/q} (\log N)^{(d-1)(1-1/q)} ||t||_q.
$$

The following Nikol'skii type inequalities are from [\[12\]](#page-29-3), Chapter 1, Section 2.

Theorem 1.8. Let $1 \leq q < p < \infty$. For any $t \in \mathcal{T}(N)$ we have

$$
||t||_p \le C(q, p, d)N^{\beta}||t||_q, \quad \beta = 1/q - 1/p.
$$

2 The case $1 < q \le p \le 2$

Proof of Theorem [1.3.](#page-5-1) In the case $1 < q = p \leq 2$ the upper bounds follow from approximation by partial sums $S_{Q_n}(\cdot)$. The corresponding lower bounds follow from the proof of the lower bounds of Theorem 2.1 from [\[12\]](#page-29-3), Chapter 4. We now assume that $\beta > 0$. The case $a > 2\beta$ in Theorem [1.3,](#page-5-1) which corresponds to the first line, was proved for classes \mathbf{W}_q^r in [\[11\]](#page-29-2) (see also [\[12\]](#page-29-3), Ch.4). In that proof assumption $f \in \mathbf{W}_q^r$ was used to claim that $||f_l||_q \ll 2^{-rl}$, which means $||f||_{\mathbf{W}_q^{a,0}} < \infty$. Thus, that proof gives the required upper bound for the class $\mathbf{W}_q^{a,0}$. That same proof gives the corresponding upper bound for the class $\mathbf{W}_q^{\hat{a},b}$ for all b. The proofs from [\[11\]](#page-29-2) and [\[12\]](#page-29-3) are constructive.

Consider now the case $\beta < a < 2\beta$. The proof of upper bounds in this case uses the ideas from [\[11\]](#page-29-2) and [\[12\]](#page-29-3). Take an $n \in \mathbb{N}$ and include in approximation the

$$
S_{Q_n}(f) := \sum_{\mathbf{s}: \|\mathbf{s}\|_1 \le n} \delta_{\mathbf{s}}(f).
$$

Choose N such that

$$
2^N \asymp 2^n n^{d-1}
$$

and for $l \in (n, N]$ include in the approximation m_l blocks $\delta_{s}(f)$, $||s||_1 = l$, with largest $\|\delta_{s}(f)\|_{p}$. Denote this set of indices s by G_{l} . Then by Theorem [1.6](#page-6-0) and the assumption $f \in \mathbf{W}_q^{a,b}$ we obtain

$$
\left(\sum_{\mathbf{s}:\|\mathbf{s}\|_1=l} \|\delta_{\mathbf{s}}(f)\|_p^q 2^{-l\beta q}\right)^{1/q} \ll \|f_l\|_q \le 2^{-al} l^{(d-1)b}.\tag{2.1}
$$

We now need the following well known simple lemma (see, for instance, [\[12\]](#page-29-3), p.92).

Lemma 2.1. Let $a_1 \ge a_2 \ge \cdots \ge a_M \ge 0$ and $1 \le q \le p \le \infty$. Then for all $m < M$ one has

$$
\left(\sum_{k=m}^{M} a_k^p\right)^{1/p} \le m^{-\beta} \left(\sum_{k=1}^{M} a_k^q\right)^{1/q}.
$$

Applying Lemma [2.1](#page-7-0) to $\{\|\delta_{\mathbf{s}}(f)\|_{p}\}\$ we obtain

$$
\left(\sum_{\mathbf{s}:\|\mathbf{s}\|_1=l,\mathbf{s}\notin G_l} \|\delta_{\mathbf{s}}(f)\|_p^p\right)^{1/p} \ll (m_l+1)^{-\beta} 2^{-(a-\beta)l} l^{(d-1)b}.\tag{2.2}
$$

Next, using the Corollary [1.2](#page-6-1) of the Littlewood-Paley theorem we derive from [\(2.2\)](#page-7-1)

$$
\|\sum_{\mathbf{s}: \|\mathbf{s}\|_1 = l, \mathbf{s} \notin G_l} \delta_{\mathbf{s}}(f)\|_p \ll \left(\sum_{\mathbf{s}: \|\mathbf{s}\|_1 = l, \mathbf{s} \notin G_l} \|\delta_{\mathbf{s}}(f)\|_p^p\right)^{1/p} \ll (m_l + 1)^{-\beta} 2^{-(a-\beta)l} l^{(d-1)b}.
$$
\n(2.3)

Denote

$$
f'_l := \sum_{\mathbf{s}: \|\mathbf{s}\|_1 = l, \mathbf{s} \notin G_l} \delta_{\mathbf{s}}(f).
$$

Let $\kappa > 0$ be such that $a - \beta < \kappa \beta < \beta$. Specify

$$
m_l := [2^{\kappa(N-l)}].
$$

Then [\(2.3\)](#page-8-0) implies

$$
\|\sum_{n
$$

The approximant

$$
A_m(f) := S_{Q_n}(f) + \sum_{n < l \le N} \sum_{\mathbf{s}: ||\mathbf{s}||_1 = l, \mathbf{s} \notin G_l} \delta_{\mathbf{s}}(f)
$$

has at most m terms

$$
m \ll |Q_n| + \sum_{n < l \le N} 2^l m_l \ll 2^N. \tag{2.5}
$$

By Theorem [1.8](#page-6-2) we have

$$
\|\sum_{l>N} f_l\|_p \ll \sum_{l>N} \|f_l\|_p \ll \sum_{l>N} \|f_l\|_q 2^{\beta l} \ll 2^{-(a-\beta)N} N^{(d-1)b}.
$$
 (2.6)

Combining (2.4) with (2.6) and taking into account (2.5) we obtain

$$
||f - A_m(f)||_p \ll m^{-(a-\beta)} (\log m)^{b(d-1)}.
$$

This completes the proof of upper bounds in the case $\beta < a < 2\beta$.

We now proceed to the case $a = 2\beta$. We begin with the upper bounds. The proof is as in the above case $\beta < a < 2\beta$. As above, we choose N such

that $2^N \approx 2^n n^{d-1}$. Then $N - n \approx \log n$. For $l \in (n, N]$ set $m_l = 2^{N-l}$. Then as above

$$
m \ll |Q_n| + \sum_{n < l \le N} 2^l m_l \ll 2^N (N - n). \tag{2.7}
$$

In the same way as [\(2.6\)](#page-8-2) was established we get

$$
\|\sum_{l>N} f_l\|_p \ll \sum_{l>N} \|f_l\|_p \ll \sum_{l>N} \|f_l\|_q 2^{\beta l} \ll 2^{-\beta N} N^{(d-1)b}.
$$
 (2.8)

By (2.3) we have

$$
||f_l'||_p \ll (m_l + 1)^{-\beta} 2^{-\beta l} l^{(d-1)b} \tag{2.9}
$$

and

$$
\|\sum_{n

$$
\ll \left(\sum_{n
$$
$$

Relations [\(2.7\)](#page-9-0), [\(2.8\)](#page-9-1) and [\(2.10\)](#page-9-2) imply the required upper bound.

We now prove the lower bounds in the case $a = 2\beta$. Let N be as above. For $l \in (n, N]$ choose an arbitrary set B_l of **s** such that $||\mathbf{s}||_1 = l$ and $|B_l|$ $m_l := 2^{N-l}$. Consider f such that $f_l = 0$ for $l \notin (n, N]$ and for $l \in (n, N]$

$$
f_l := 2^{-(2\beta+1-1/q)l} l^{(d-1)b} m_l^{-1/q} \sum_{\mathbf{s}\in B_l} \sum_{\mathbf{k}\in \rho(\mathbf{s})} e^{i(\mathbf{k}, \mathbf{x})}.
$$

Then

$$
\|f_l\|_q \ll 2^{-2\beta l} l^{(d-1)b}
$$

and therefore $||f||_{\mathbf{W}^{2\beta,b}_q} \ll 1$. We prove the lower bound for the $\sigma_m(f)_p$ with $m < 2^N(N-n)/8$. Let $K_m := {\mathbf{k}}^j\}_{j=1}^m$ be given. Denote

$$
L := \{ l \in (n, N] : |K_m \cap \cup_{\mathbf{s} \in B_l} \rho(\mathbf{s})| \le 2^N/4 \}.
$$

Then

$$
(N - n - |L|)2^N/4 \le m \le 2^N(N - n)/8,
$$

which implies

$$
|L| \ge (N - n)/2.
$$

Take $l \in L$. Denote

$$
K_m^l := K_m \cap \cup_{\mathbf{s} \in B_l} \rho(\mathbf{s})
$$

and

$$
B'_l := \{ \mathbf{s} \in B_l : |K_m^l \cap \rho(\mathbf{s})| \leq |\rho(\mathbf{s})|/2 \}.
$$

As above we derive that

$$
|B'_l|\geq |B_l|/2.
$$

Let q be any polynomial of the form

$$
g = \sum_{\mathbf{k} \in K_m} c_{\mathbf{k}} e^{i(\mathbf{k}, \mathbf{x})}.
$$

By Theorem [1.6](#page-6-0) we get

$$
||f - g||_p \gg \left(\sum_{n < l \le N} \sum_{\mathbf{s} \in B_l} \left(\|\delta_\mathbf{s}(f - g)\|_2 2^{l(1/2 - 1/p)}\right)^p\right)^{1/p}
$$
\n
$$
\gg \left(\sum_{l \in L} \sum_{\mathbf{s} \in B_l'} \left(\|\delta_\mathbf{s}(f - g)\|_2 2^{l(1/2 - 1/p)}\right)^p\right)^{1/p}
$$
\n
$$
\gg \left(\sum_{l \in L} \sum_{\mathbf{s} \in B_l'} \left(2^{-(2\beta + 1 - 1/q)l} l^{(d-1)b} 2^{-(N-l)/q} 2^{l/2} 2^{l(1/2 - 1/p)}\right)^p\right)^{1/p}
$$
\n
$$
\gg 2^{-\beta N} N^{(d-1)b} |L|^{1/p} \gg 2^{-\beta N} N^{(d-1)b} (N - n)^{1/p}.
$$

Taking into account that $2^N \approx 2^n n^{d-1}$ and $m \leq 2^N (N - n)/8$ we complete the proof of lower bounds.

Proof of Theorem [1.4.](#page-5-0) In the case $r \neq 2\beta$ the upper bounds in Theorem [1.4](#page-5-0) follow from Theorem [1.3](#page-5-1) by the embedding of \mathbf{W}_q^r into $\mathbf{W}_q^{r,0}$. It turns out that in the case $r = 2\beta$ the above way does not give a sharp upper bound. We now proof the corresponding upper bound in the case $r = 2\beta$. We begin with an analog of Lemma [2.1.](#page-7-0)

Lemma 2.2. Let $\{w_j\}_{j=1}^M$ be a set of positive weights. Let $a_1 \ge a_2 \ge \cdots \ge$ $a_M \geq 0$ and $1 \leq q \leq p \leq \infty$. Then for all $m < M$ one has

$$
\left(\sum_{k=m}^{M} a_k^p w_k\right)^{1/p} \le \left(\sum_{k=1}^{m} w_k\right)^{-\beta} \left(\sum_{k=1}^{M} a_k^q w_k\right)^{1/q}.
$$

Proof. Monotonicity of ${a_k}_{k=1}^M$ implies

$$
a_m^q \sum_{k=1}^m w_k \le \sum_{k=1}^M a_k^q w_k,
$$

and

$$
a_m \leq \left(\sum_{k=1}^m w_k\right)^{-1/q} \left(\sum_{k=1}^M a_k^q w_k\right)^{1/q}.
$$

Therefore,

$$
\left(\sum_{k=m}^{M} a_k^p w_k\right)^{1/p} \le a_m^{(p-q)/p} \left(\sum_{k=m}^{M} a_k^q w_k\right)^{1/p} \le \left(\sum_{k=1}^{m} w_k\right)^{-\beta} \left(\sum_{k=1}^{M} a_k^q w_k\right)^{1/q}.
$$

We now prove the upper bound in the case $r = 2\beta$. Let $n \in \mathbb{N}$ and, as above, N be such that $2^N \approx 2^n n^{d-1}$. For $f \in \mathbf{W}_q^r$ we include in the approximation $S_{Q_n}(f)$ and approximate

$$
g := g(f) := \sum_{\mathbf{s} \in \Delta(n,N)} \delta_{\mathbf{s}}(f), \quad \Delta(n,N) := \{\mathbf{s} : n < ||\mathbf{s}||_1 \leq N\}.
$$

Using Theorem [1.6](#page-6-0) we obtain

$$
\left(\sum_{\mathbf{s}\in\Delta(n,N)}\left(2^{\beta\|\mathbf{s}\|_1}\|\delta_\mathbf{s}(f)\|_p\right)^q\right)^{1/q}\ll\|f\|_{\mathbf{W}_q^r}.\tag{2.11}
$$

We want to apply Lemma [2.2.](#page-10-0) Consider $\{v_s\}_{s\in\Delta(n,N)}$, $v_s := ||\delta_s(f)||_p 2^{-||s||_1/p}$ with weights $w_s := 2^{\|\mathbf{s}\|_1}$. Then [\(2.11\)](#page-11-0) gives

$$
\left(\sum_{\mathbf{s}\in\Delta(n,N)}v_{\mathbf{s}}^q w_{\mathbf{s}}\right)^{1/q}\ll\|f\|_{\mathbf{W}_q^r}.
$$

Choose k largest v_s and denote the corresponding set of indices **s** by $G(k)$. By Lemma [2.2](#page-10-0) we obtain from the above estimate

$$
\left(\sum_{\mathbf{s}\in\Delta(n,N)\backslash G(k)} v_{\mathbf{s}}^p w_{\mathbf{s}}\right)^{1/p} \ll \left(\sum_{\mathbf{s}\in G(k)} w_{\mathbf{s}}\right)^{-\beta} \|f\|_{\mathbf{W}_q^r}.\tag{2.12}
$$

By the Corollary [1.2](#page-6-1) to the Littlewood-Paley Theorem we find

$$
\|\sum_{\mathbf{s}\in\Delta(n,N)\backslash G(k)}\delta_{\mathbf{s}}(f)\|_{p}\ll\left(\sum_{\mathbf{s}\in\Delta(n,N)\backslash G(k)}\|\delta_{\mathbf{s}}(f)\|_{p}^{p}\right)^{1/p}.\tag{2.13}
$$

Combining (2.13) with (2.12) we obtain

$$
\left(\sum_{\mathbf{s}\in\Delta(n,N)\backslash G(k)}\|\delta_{\mathbf{s}}(f)\|_{p}^{p}\right)^{1/p} = \left(\sum_{\mathbf{s}\in\Delta(n,N)\backslash G(k)}v_{\mathbf{s}}^{p}w_{\mathbf{s}}\right)^{1/p}
$$
\n
$$
\ll \left(\sum_{\mathbf{s}\in G(k)}w_{\mathbf{s}}\right)^{-\beta}\|f\|_{\mathbf{W}_{q}^{r}}.\tag{2.14}
$$

Choose k such that

$$
2^N \le \sum_{\mathbf{s}\in G(k)} w_\mathbf{s} < 2^{N+1}.
$$

In this way we have constructed an *m*-term approximation of f with $m \ll 2^N$ and error

$$
||f - S_{Q_n}(f) - \sum_{\mathbf{s} \in G(k)} \delta_{\mathbf{s}}(f)||_p \ll 2^{-\beta N} \ll m^{-\beta}.
$$

The upper bounds in Theorem [1.4](#page-5-0) are proved.

The lower bounds in the case $r > 2\beta$ are proved in [\[12\]](#page-29-3). The lower bounds in the case $\beta < r \leq 2\beta$ follow from the univariate case.

3 The case $1 < q \leq 2 < p < \infty$

The main goal of this section is to prove Theorem [1.2](#page-3-1) from the Introduction. We reformulate it here for convenience.

Theorem 3.1. Let $1 < q \leq 2 < p < \infty$ and $\beta p' < r < 1/q$. Then we have

$$
\sigma_m(\mathbf{W}_q^r)_p \asymp m^{-(r-\beta)p/2} (\log m)^{(d-1)(r(p-1)-\beta p)}
$$

.

The upper bounds are achieved by a constructive greedy-type algorithm.

Proof. We will prove the upper bounds for a wider class $\mathbf{W}^{r,0}_{q}$. Let $f \in \mathbf{W}^{r,0}_{q}$. Let $n \in \mathbb{N}$. We build an *m*-term approximation with $m = 2|Q_n| \approx 2^n n^{d-1}$. We include in the approximation $S_{Q_n}(f)$. We split the remainder function into two functions

$$
f - S_{Q_n}(f) = g_A + g_0.
$$

We use Theorem [1.1](#page-3-0) to approximate g_A and approximate g_0 by 0. We now describe a construction of g_A . First, we choose $N \in (n, Cn]$, $C = C(p, d)$, which will be specified later on, and include in the g_A the

$$
g_A^1 := \sum_{n < l \le N} f_l.
$$

Then by Theorem [1.7](#page-6-3) we have

$$
||g_A^1||_A = \sum_{n < l \le N} ||f_l||_A \ll \sum_{n < l \le N} ||f_l||_q 2^{l/q} l^{(d-1)(1-1/q)}
$$
\n
$$
\ll \sum_{n < l \le N} 2^{l(1/q - r)} l^{(d-1)(1-1/q)} \ll 2^{N(1/q - r)} N^{(d-1)(1-1/q)}.\tag{3.1}
$$

Next, for $l > N$ define

$$
u_l := [n^{d-1} 2^{\kappa(N-l)}]
$$

with κ satisfying

$$
\frac{1/q-r}{1-1/q} < \kappa < \frac{r-\beta}{\beta}.
$$

Such κ exists because our assumption $r > \beta p'$ is equivalent to the inequality

$$
\frac{1/q-r}{1-1/q} < \frac{r-\beta}{\beta}.
$$

Denote $G(l)$ the set of indices **s**, $\|\mathbf{s}\|_1 = l$, of cardinality $|G(l)| = u_l$, with largest $\|\delta_{s}(f_{l})\|_{2}$. Second, we include in g_{A} the

$$
g_A^2 := \sum_{l>N} \sum_{\mathbf{s} \in G(l)} \delta_{\mathbf{s}}(f_l).
$$

It is clear that there is only finite number of nonzero terms in the above sum. We have

$$
||g_A^2||_A = \sum_{l>N} ||\sum_{\mathbf{s}\in G(l)} \delta_{\mathbf{s}}(f_l)||_A \ll \sum_{l>N} ||f_l||_q 2^{l/q} |G(l)|^{1-1/q}
$$

$$
\ll \sum_{l>N} 2^{l(1/q-r)} n^{(d-1)(1-1/q)} 2^{\kappa(N-l)(1-1/q)} \\
\ll n^{(d-1)(1-1/q)} 2^{\kappa N(1-1/q)} \sum_{l>N} 2^{l(1/q-r-\kappa(1-1/q))}.
$$

By our choice of κ we have $1/q - r - \kappa(1 - 1/q) < 0$ and, therefore, we continue

$$
\ll N^{(d-1)(1-1/q)} 2^{N(1/q-r)}.\tag{3.2}
$$

By Theorem [1.1](#page-3-0) we obtain

$$
||g_A - G_{m/2}^p(g_A)||_p \ll m^{-1/2} N^{(d-1)(1-1/q)} 2^{N(1/q-r)}.
$$
 (3.3)

We now bound the $||g_0||_p$. Denote

$$
f_l^o := f_l - \sum_{\mathbf{s} \in G(l)} \delta_{\mathbf{s}}(f_l).
$$

By Theorem [1.6](#page-6-0) we get

$$
\left(\sum_{\|\mathbf{s}\|_1=l} \left(\|\delta_\mathbf{s}(f_l)\|_2 2^{\|\mathbf{s}\|_1 (1/2-1/q)}\right)^q\right)^{1/q} \ll 2^{-rl}
$$

and

$$
\left(\sum_{\|\mathbf{s}\|_1=l} \|\delta_{\mathbf{s}}(f_l)\|_2^q\right)^{1/q} \ll 2^{l(-r+1/q-1/2)}.
$$

By Lemma [2.1](#page-7-0) we obtain

$$
\left(\sum_{\mathbf{s}\notin G(l)} \|\delta_{\mathbf{s}}(f_l)\|_2^p\right)^{1/p} \ll (u_l+1)^{-\beta} 2^{l(-r+1/q-1/2)}.
$$

By Theorem [1.6](#page-6-0) we get from here

$$
||f_l^o||_p \ll \left(\sum_{\mathbf{s} \notin G(l)} \left(||\delta_{\mathbf{s}}(f_l)||_2 2^{\|\mathbf{s}\|_1 (1/2 - 1/p)}\right)^p\right)^{1/p} \ll 2^{-(r-\beta)l} (u_l + 1)^{-\beta}.
$$

Thus

$$
||g_0||_p \le \sum_{l>N} ||f_l^o||_p \ll \sum_{l>N} 2^{-(r-\beta)l} n^{-\beta(d-1)} 2^{-\beta\kappa(N-l)}
$$

$$
\ll n^{-\beta(d-1)} 2^{-\beta\kappa N} \sum_{l>N} 2^{-(r-\beta-\beta\kappa)l}.
$$
 (3.4)

By our choice of κ we have $r - \beta - \beta \kappa > 0$. Therefore, [\(3.4\)](#page-15-0) gives

$$
||g_0||_p \ll 2^{-(r-\beta)N} n^{-\beta(d-1)}.
$$
\n(3.5)

.

We now choose N from the condition

$$
2^{(1/q-r)N}n^{(d-1)(1-1/q)}m^{-1/2} \asymp 2^{-(r-\beta)N}n^{-\beta(d-1)}
$$

This is equivalent to

$$
2^N \asymp 2^{np/2} n^{(d-1)(1-p/2)}
$$

or in terms of m

$$
2^N \asymp m^{p/2} (\log m)^{(d-1)(1-p)}.\tag{3.6}
$$

As a result it gives us the following upper bound for the error of approximation

$$
||f - S_{Q_n}(f) - G_{m/2}^p(g_A)||_p \ll 2^{-(r-\beta)N} n^{-\beta(d-1)}
$$

$$
\ll m^{-(r-\beta)p/2} (\log m)^{(d-1)(r(p-1)-\beta p)}.
$$

This completes the proof of upper bounds.

We proceed to the lower bounds. For a given m chose N as in [\(3.6\)](#page-15-1). Consider the function

$$
g(\mathbf{x}) := \sum_{\|\mathbf{s}\|_1 = N} \sum_{\mathbf{k} \in \rho(\mathbf{s})} e^{i(\mathbf{k}, \mathbf{x})} = \sum_{\mathbf{k} \in \Delta Q_N} e^{i(\mathbf{k}, \mathbf{x})}, \quad \Delta Q_N := Q_N \setminus Q_{N-1}.
$$

It is known that

$$
||g||_q \asymp 2^{N(1-1/q)} N^{(d-1)/q}, \quad 1 < q < \infty.
$$
 (3.7)

We now estimate the $\sigma_m(g)_p$ from below. Take any set K_m of m frequencies k. Consider an additional function

$$
h(\mathbf{x}) := \sum_{\mathbf{k} \in \Delta Q_N \setminus K_m} e^{i(\mathbf{k}, \mathbf{x})}.
$$

For any polynomial t with frequencies from K_m we have

$$
\langle g-t, h \rangle \le ||g-t||_p ||h||_{p'} \tag{3.8}
$$

and

$$
\langle g-t, h \rangle = \langle g, h \rangle = \sum_{\mathbf{k} \in \Delta Q_N \backslash K_m} 1 = |\Delta Q_N \setminus K_m|.
$$
 (3.9)

From our choice (3.6) of N it is clear that asymptotically

$$
|\Delta Q_N \setminus K_m| \ge |\Delta Q_N| - m \gg 2^N N^{d-1}.
$$

Next, we have

$$
||h||_{p'} \le ||g||_{p'} + ||g - h||_{p'} \le ||g||_{p'} + ||g - h||_2
$$

$$
\ll 2^{N/p} N^{(d-1)/p'} + m^{1/2} \ll m^{1/2}.
$$

Thus, (3.8) and (3.9) yield

$$
\sigma_m(g)_p \gg 2^N N^{d-1} m^{-1/2}.
$$

We have from [\(3.7\)](#page-15-2)

$$
\|g^{(r)}\|_q \ll 2^{N(r+1-1/q)}N^{(d-1)/q}.
$$

Therefore,

$$
\sigma_m(\mathbf{W}_q^r)_p \gg 2^{N(1/q-r)} N^{(d-1)(1-1/q)} m^{-1/2}
$$

$$
\asymp m^{-(r-\beta)p/2} (\log m)^{(d-1)(r(p-1)-\beta p)}.
$$

This proves the lower bounds.

The above proof of Theorem [3.1](#page-12-1) gives the right order of $\sigma_m(\mathbf{W}_q^{a,b})_p$ for $\beta p' < a < 1/q$ and all $b.$ We formulate this as a theorem.

Theorem 3.2. Let $1 < q \leq 2 < p < \infty$ and $\beta p' < a < 1/q$. Then we have

$$
\sigma_m(\mathbf{W}_q^{a,b})_p \asymp m^{-(a-\beta)p/2} (\log m)^{(d-1)(b+a(p-1)-\beta p)}.
$$

The upper bounds are achieved by a constructive greedy-type algorithm.

 \Box

For $f \in \mathbf{W}^{r,0}_{q}$ one has for $r > \beta$

$$
\|\sum_{l>N} f_l\|_p \le \sum_{l>N} \|f_l\|_p \ll \sum_{l>N} \|f_l\|_q 2^{\beta l} \ll 2^{-(r-\beta)N}.
$$
 (3.10)

In the proof of Theorem [3.1](#page-12-1) we constructed g_A^2 and g_0 . It resulted in a better error estimate of the *m*-term approximation of the tail $\sum_{l>N} f_l$ than the simple bound [\(3.10\)](#page-17-0). We got the error $\ll 2^{-(r-\beta)N}n^{-\beta(d-1)}$. We obtain the same improvement of the error if, in addition to the assumption $f \in \mathbf{W}^{r,0}_{q}$, we assume that $f \in \mathbf{W}_{p}^{r-\beta,-\beta}$. For $f \in \mathbf{W}_{p}^{r-\beta,-\beta}$ we have

$$
\|\sum_{l>N} f_l\|_p \ll 2^{-(r-\beta)N} N^{-\beta(d-1)}.
$$

We formulate a theorem, which follows from the proof of Theorem [3.1.](#page-12-1)

Theorem 3.3. Let $1 < q \leq 2 < p < \infty$ and $\beta < a < 1/q$. Then we have

$$
\sigma_m(\mathbf{W}_q^{a,b} \cap \mathbf{W}_p^{a-\beta,b-\beta})_p \asymp m^{-(a-\beta)p/2} (\log m)^{(d-1)(b+a(p-1)-\beta p)}.
$$

The upper bounds are achieved by a constructive greedy-type algorithm.

We note that the class \mathbf{H}_q^r (see the definition in Section 5), $1 < q \leq 2$, is embedded into the class $\mathbf{W}^{r,\delta}_{q} \cap \mathbf{W}^{r-\beta,b-\beta}_{p}$ with $b = 1/q$. This follows from Corollary [1.2](#page-6-1) and Theorem [1.6.](#page-6-0) The following theorem holds.

Theorem 3.4. Let $1 < q \leq 2 < p < \infty$ and $\beta < a < 1/q$. Then we have

$$
\sigma_m(\mathbf{H}_q^r)_p \asymp m^{-(r-\beta)p/2} (\log m)^{(d-1)(1/q+r(p-1)-\beta p)}.
$$

The upper bounds are achieved by a constructive greedy-type algorithm.

The order of $\sigma_m(\mathbf{H}_q^r)_p$ is known (see Romanyuk [\[10\]](#page-29-5)). However, the corresponding upper bounds in [\[10\]](#page-29-5) are proved by a nonconstructive method of approximation.

We now proceed to the case $a = 1/q$.

Theorem 3.5. Let $1 < q \leq 2 < p < \infty$ and $a = 1/q$. Then we have

$$
\sigma_m(\mathbf{W}_q^{1/q,b})_p \asymp m^{-1/2} (\log m)^{(d-1)(b+1-1/q)+1}.
$$

The upper bounds are achieved by a constructive greedy-type algorithm.

Proof. The proof goes along the lines of the proof of Theorem [3.1.](#page-12-1) We use the same notation as above. We begin with the upper bounds. In the case $a = 1/q$ the bound [\(3.1\)](#page-13-0) reads

$$
||g_A^1||_A \ll (N-n)N^{(d-1)(b+1-1/q)}.
$$
\n(3.11)

We repeat the argument from the proof of Theorem [3.1](#page-12-1) for g_A^2 and g_0 with $\kappa \in (0, (a - \beta)/\beta)$. It gives

$$
||g_A^2||_A \ll N^{(d-1)(b+1-1/q)},\tag{3.12}
$$

$$
||g_0||_p \ll 2^{-(a-\beta)N} n^{(b-\beta)(d-1)}.
$$
\n(3.13)

Choosing N from (3.6) we obtain

$$
||f - S_{Q_n}(f) - G_{m/2}^p(g_A)||_p \ll (N - n)N^{(d-1)(b+1-1/q)}m^{-1/2}
$$

$$
\ll m^{-1/2}(\log m)^{(d-1)(b+1-1/q)+1}.
$$

This completes the proof of upper bounds.

We proceed to the lower bounds. For a given m chose N as in [\(3.6\)](#page-15-1). Consider the function

$$
g(\mathbf{x}) := \sum_{n < l \leq N} 2^{-l/q} l^{b(d-1)} \left(2^{l(1-1/q)} l^{(d-1)/q} \right)^{-1} \sum_{\mathbf{k} \in \Delta Q_l} e^{i(\mathbf{k}, \mathbf{x})}, \quad \Delta Q_l := Q_l \setminus Q_{l-1}.
$$

Then $||g||_{\mathbf{W}_q^{1/q,b}} \ll 1$.

We now estimate the $\sigma_m(g)_p$ from below. Take any set K_m of m frequencies k. Consider an additional function

$$
h(\mathbf{x}) := \sum_{\mathbf{k} \in \Delta(n,N) \setminus K_m} e^{i(\mathbf{k},\mathbf{x})}
$$

For any polynomial t with frequencies from K_m we have

$$
\langle g-t, h \rangle \le ||g-t||_p ||h||_{p'} \tag{3.14}
$$

.

and

$$
\langle g-t, h \rangle = \langle g, h \rangle = \sum_{\mathbf{k} \in \Delta(n, N) \backslash K_m} \hat{g}(\mathbf{k}). \tag{3.15}
$$

From our choice (3.6) of N it is clear that asymptotically

$$
\sum_{\mathbf{k}\in\Delta(n,N)\backslash K_m}\hat{g}(\mathbf{k})\gg (N-n)N^{(d-1)(b+1-1/q)}.
$$

Next, we have

$$
||h||_{p'} \le ||g||_{p'} + ||g - h||_{p'} \le ||g||_{p'} + ||g - h||_2
$$

$$
\ll 2^{N/p} N^{(d-1)/p'} + m^{1/2} \ll m^{1/2}.
$$

Thus, [\(3.14\)](#page-18-0) and [\(3.15\)](#page-18-1) yield

$$
\sigma_m(g)_p \gg (N-n)N^{(d-1)(b+1-1/q)}m^{-1/2}.
$$

Therefore,

$$
\sigma_m(\mathbf{W}_q^{1/q,b})_p \gg (N-n)N^{(d-1)(b+1-1/q)}m^{-1/2}
$$

$$
\asymp m^{-1/2}(\log m)^{(d-1)(b+1-1/q)+1}.
$$

This proves the lower bounds.

The above proof of Theorem [3.5](#page-17-1) can be adjusted to prove the following results for the $\mathbf{W}_q^{1/q}$ classes.

Theorem 3.6. Let $1 < q \leq 2 < p < \infty$. Then we have

$$
\sigma_m(\mathbf{W}_q^{1/q})_p \asymp m^{-1/2} (\log m)^{d(1-1/q)}.
$$

The upper bounds are achieved by a constructive greedy-type algorithm.

Proof. We need the following analog of Theorem [1.7.](#page-6-3)

Lemma 3.1. For $t \in \mathcal{T}(\Delta(n, N))$ we have for $1 < q \leq 2$

$$
||t||_A \ll N^{(d-1)(1-1/q)}(N-n)^{1-1/q}||t||_{\mathbf{W}_q^{1/q}}.
$$

Proof. Let $r = 1/q$ and

$$
t = \varphi * F_r, \quad \|\varphi\|_q = \|t\|_{\mathbf{W}_q^r}.
$$

 \Box

Then

$$
||t||_A \le ||\varphi||_q \left\| \sum_{\mathbf{k} \in \Delta(n,N)} \epsilon_{\mathbf{k}} \hat{F}_r(\mathbf{k}) e^{i(\mathbf{k}, \mathbf{x})} \right\|_{q'}, \quad |\epsilon_{\mathbf{k}}| = 1.
$$
 (3.16)

Using Theorem [1.6](#page-6-0) with parameters q' and 2 we obtain

$$
\left\| \sum_{\mathbf{k} \in \Delta(n,N)} \epsilon_{\mathbf{k}} \hat{F}_r(\mathbf{k}) e^{i(\mathbf{k}, \mathbf{x})} \right\|_{q'}
$$

$$
\ll \left(\sum_{n < l \le N} \left(2^{-lr} 2^{l/2} 2^{l(1/2 - 1/q')} \right)^{q'} l^{d-1} \right)^{1/q'} \ll (N - n)^{1/q'} N^{(d-1)/q'}.
$$
 (3.17)

Combining [\(3.16\)](#page-20-0) and [\(3.17\)](#page-20-1) we complete the proof of Lemma [3.1.](#page-19-0)

 \Box

We return to the proof of Theorem [3.6.](#page-19-1) The proof goes along the lines of the proof of Theorem [3.5.](#page-17-1) We use the same notation as above. We begin with the upper bounds. In our case Lemma [3.1](#page-19-0) implies the following analog of the bound [\(3.11\)](#page-18-2)

$$
||g_A^1||_A \ll (N-n)^{1-1/q} N^{(d-1)(1-1/q)}.
$$
\n(3.18)

Choosing N from (3.6) we obtain

$$
||f - S_{Q_n}(f) - G_{m/2}^p(g_A)||_p \ll (N - n)^{1 - 1/q} N^{(d-1)(1 - 1/q)} m^{-1/2}
$$

$$
\ll m^{-1/2} (\log m)^{(d-1)(1 - 1/q) + 1 - 1/q} = m^{-1/2} (\log m)^{d(1 - 1/q)}.
$$

This completes the proof of upper bounds.

The lower bounds follow from the same example (with $b = 0$) that was used in the proof of Theorem [3.5.](#page-17-1) In this case instead of $||g||_{\mathbf{W}_q^{1/q,0}} \ll 1$ we have

$$
\|g\|_{\mathbf{W}^{1/q}_q} \ll (N-n)^{1/q}
$$

which brings the bound

$$
\sigma_m(\mathbf{W}_q^{1/q})_p \gg (N - n)^{1 - 1/q} N^{(d-1)(1 - 1/q)} m^{-1/2}
$$

$$
\asymp m^{-1/2} (\log m)^{d(1 - 1/q)}.
$$

This proves the lower bounds.

4 The case $q = 1$

We begin with the case $2 \leq p < \infty$.

Theorem 4.1. For any $\epsilon > 0$ we have for $2 \leq p < \infty$

$$
\sigma_m(\mathbf{W}_1^{a,b})_p \ll \begin{cases} m^{-a+1/2}(\log m)^{(d-1)(a-1+b)+\epsilon}, & a > 1, \\ m^{-(a-\beta)p/2}(\log m)^{(d-1)b+\epsilon}, & \beta < a < 1, \\ m^{-1/2}(\log m)^{(d-1)b+1+\epsilon}, & a = 1, \end{cases} \quad \beta = 1 - 1/p,
$$

with constants in \ll allowed to depend on ϵ , d, and p.

The upper bounds are achieved by a constructive greedy-type algorithm.

Proof. For large smoothness $a > 1$ the following lemma from [\[16\]](#page-29-0) plays the key role in the proof.

Lemma 4.1. Define for $f \in L_1$

$$
f_l:=\sum_{\|\mathbf{s}\|_1=l}\delta_\mathbf{s}(f),\quad l\in\mathbb{N}_0,\quad \mathbb{N}_0:=\mathbb{N}\cup\{0\}.
$$

Consider the class

$$
\mathbf{W}_A^{a,b} := \{ f : \|f_l\|_A \le 2^{-al}(\bar{l})^{(d-1)b} \}.
$$

Then for $2 \le p < \infty$ and $0 < \mu < a$ there is a constructive method $A_m(\cdot, p, \mu)$ based on greedy algorithms, which provides the bound for $f \in W_A^{a,b}$

$$
||f - A_m(f, p, \mu)||_p \ll m^{-a - 1/2} (\log m)^{(d-1)(a+b)}, \quad 2 \le p < \infty,
$$
 (4.1)

We also need the following version of Theorem [1.7](#page-6-3) in the case $q = 1$ (see [\[12\]](#page-29-3), Chapter 1, Section 2).

Lemma 4.2. For any $\epsilon > 0$ there is $C(\epsilon, d)$ such that for each $t \in \mathcal{T}(N)$ we have

$$
||t||_A \le C(\epsilon, d)N(\log N)^{\epsilon}||t||_1.
$$

Let $f \in \mathbf{W}^{a,b}_1, a > 1$. By Lemma [4.2](#page-21-0) we get

$$
||f_l||_A\ll 2^{-l(a-1)}l^{b(d-1)+\epsilon}
$$

with a constant in \ll allowed to depend on ϵ and d. Setting $b(\epsilon) := b + \epsilon/(d - \epsilon)$ 1), we obtain

$$
\|f\|_{{\bf W}^{a-1,b(\epsilon)}_{A}}\ll 1.
$$

Lemma [4.1](#page-21-1) gives a constructive proof of

$$
\sigma_m(f)_p \ll m^{-a+1/2} (\log m)^{(d-1)(a-1+b)+\epsilon}.
$$

This proves the first inequality in Theorem [4.1.](#page-21-2)

Consider now the case $\beta < a < 1$. The argument in this case is close to the proof of Theorem [3.1.](#page-12-1) We use the same notations. We now define

$$
g_A := \sum_{n < l \le N} f_l, \qquad g_0 := \sum_{l > N} f_l.
$$

By Lemma [4.2](#page-21-0) we get

$$
||g_A||_A \ll \sum_{n < l \le N} 2^{l(1-a)} l^{b(d-1)+\epsilon} \ll 2^{N(1-a)} N^{b(d-1)+\epsilon}.\tag{4.2}
$$

By Theorem [1.1](#page-3-0) we obtain

$$
\sigma_m(g_A)_p \ll m^{-1/2} 2^{N(1-a)} N^{b(d-1)+\epsilon}.
$$
\n(4.3)

By Theorem [1.8](#page-6-2)

$$
||g_0||_p \le \sum_{l>N} ||f_l||_p \le \sum_{l>N} ||f_l||_1 2^{\beta l} \ll 2^{-(a-\beta)N} N^{b(d-1)}.
$$
 (4.4)

Close ${\cal N}$ such that

$$
m^{-1/2}2^{N(1-a)} \asymp 2^{-(a-\beta)N},
$$

that is

$$
2^N \asymp m^{p/2}.\tag{4.5}
$$

This gives the error bound

$$
\sigma_m(f)_p \ll m^{-(a-\beta)p/2} (\log m)^{b(d-1)+\epsilon}.
$$

This proves the required bound in the second case.

In the case $a = 1$ we get as in (4.2)

$$
||g_A||_A \ll N^{b(d-1)+1+\epsilon}.
$$

Choosing N from (4.5) we obtain

$$
\sigma_m(f)_p \ll m^{-1/2} (\log m)^{b(d-1)+1+\epsilon}.
$$

This completes the proof of Theorem [4.1.](#page-21-2)

We note that in the case $a \leq 1$ the corresponding lower bounds with $\epsilon = 0$ follow from the univariate case (see [\[1\]](#page-28-1)). We now prove the lower bounds for $a > 1$. It is sufficient to prove them for $p = 2$. Let m be given and n be such that $2^n n^{d-1} \simeq m$ and $m \leq c(d) |\Delta Q_n|$ with small enough $c(d) > 0$. Let

$$
\mathcal{K}_N(x) := \sum_{|k| \le N} \left(1 - |k|/N\right) e^{ikx} = \left(\sin(Nx/2)\right)^2 / \left(N(\sin(x/2))^2\right)
$$

be a univariate Fejér kernel. The Fejér kernel \mathcal{K}_N is an even nonnegative trigonometric polynomial in $\mathcal{T}(N-1)$. In the multivariate case define

$$
\mathcal{K}_{\mathbf{N}}(\mathbf{x}) := \prod_{j=1}^d \mathcal{K}_{N_j}(x_j), \qquad \mathbf{N} = (N_1, \ldots, N_d).
$$

Then the K_N are nonnegative trigonometric polynomials from $\mathcal{T}(\mathbf{N} - \mathbf{1}, d)$ which have the following property:

$$
\|\mathcal{K}_{\mathbf{N}}\|_{1} = 1. \tag{4.6}
$$

Consider the function

$$
g(\mathbf{x}) := \sum_{\|\mathbf{s}\|_1 = n} \mathcal{K}_{2^{\mathbf{s}-2}}(\mathbf{x}) e^{i(2^{\mathbf{s}} - 2^{\mathbf{s}-2}, \mathbf{x})}.
$$

Then by [\(4.6\)](#page-23-0)

$$
||g||_1 \ll n^{d-1}.\tag{4.7}
$$

 \Box

Take any set K_m of m frequencies. It is clear that for small enough $c(d)$ we have

$$
\sigma_m(g)_2^2 \ge \sum_{\mathbf{k} \in \Delta Q_n \backslash K_m} |\hat{g}(\mathbf{k})|^2 \gg |\Delta Q_n|.
$$
 (4.8)

Relations [\(4.7\)](#page-23-1) and [\(4.8\)](#page-23-2) imply

$$
\sigma_m(\mathbf{W}_1^{a,b})_2 \gg 2^{n(1/2-a)} n^{-(d-1)/2 + b(d-1)} \asymp m^{-a+1/2} (\log m)^{(d-1)(b+a-1)}.
$$

Consider now the case $q = 1, 1 < p \le 2$. We need a version of the relation [\(1.6\)](#page-6-4) from Theorem [1.6](#page-6-0) adjusted to our case.

Lemma 4.3. Let $1 < p < \infty$. For any $\epsilon > 0$ there exists a constant $C(\epsilon, d, p)$ such that for each $t \in \mathcal{T}(Q_n)$ we have

$$
\sum_{\|\mathbf{s}\|_1 \le n} \|\delta_\mathbf{s}(t)\|_p \le C(\epsilon, d, p) n^{\epsilon} 2^{\beta n} \|t\|_1, \quad \beta = 1 - 1/p.
$$

Proof. Choose $v \in (1, p)$ such that $v' := v/(v - 1) > d/\epsilon$. By the Hölder inequality and Theorem [1.6](#page-6-0) we get

$$
\sum_{\|\mathbf{s}\|_1 \le n} \|\delta_{\mathbf{s}}(t)\|_p \le \left(\sum_{\|\mathbf{s}\|_1 \le n} 1\right)^{1/v'} \left(\sum_{\|\mathbf{s}\|_1 \le n} \|\delta_{\mathbf{s}}(t)\|_p^v\right)^{1/v}
$$

$$
\le n^{\epsilon} 2^{n(1/v-1/p)} \left(\sum_{\|\mathbf{s}\|_1 \le n} \|\delta_{\mathbf{s}}(t)\|_p^v 2^{\|\mathbf{s}\|_1 (1/p-1/v)v}\right)^{1/v} \ll n^{\epsilon} 2^{n(1/v-1/p)} \|t\|_v.
$$

By Theorem [1.8](#page-6-2) continue

$$
\ll n^{\epsilon} 2^{n(1/\nu - 1/p)} 2^{n(1-1/\nu)} \|t\|_1 = n^{\epsilon} 2^{n\beta} \|t\|_1.
$$

Theorem 4.2. Let $1 < p \le 2$. For any $\epsilon > 0$ we have

$$
\sigma_m(\mathbf{W}_1^{a,b})_p \ll \begin{cases} m^{-a+\beta}(\log m)^{(d-1)(a+b-2\beta)+\epsilon}, & a > 2\beta, \\ m^{-a+\beta}(\log m)^{(d-1)b+\epsilon}, & \beta < a \le 2\beta. \end{cases}
$$

Proof. We begin with the case of large smoothness. Let $f \in \mathbf{W}_1^{a,b}, a > 2\beta$. Then by Lemma [4.3](#page-24-0)

$$
\sum_{\|\mathbf{s}\|_1 = l} \|\delta_{\mathbf{s}}(f_l)\|_p \ll 2^{-(a-\beta)l} l^{b(d-1)+\epsilon}.\tag{4.9}
$$

 \Box

As in the proof of Theorem [1.3](#page-5-1) from Section 2 denote by G_l the set of indices **s** with m_l largest $\|\delta_{\mathbf{s}}(f_l)\|_p$. Then by Corollary [1.2](#page-6-1) and Lemma [2.1](#page-7-0) we obtain for

$$
f'_l := \sum_{\mathbf{s} \notin G_l} \delta_{\mathbf{s}}(f_l),
$$

$$
||f'_l||_p \ll \left(\sum_{\mathbf{s} \notin G_l} ||\delta_{\mathbf{s}}(f_l)||_p^p\right)^{1/p}
$$

$$
\ll (m_l + 1)^{-\beta} \sum_{\|\mathbf{s}\|_1 = l} \|\delta_{\mathbf{s}}(f_l)\|_p \le (m_l + 1)^{-\beta} 2^{-(a-\beta)l} l^{b(d-1) + \epsilon}.
$$
 (4.10)

Set for $l > n$

$$
m_l:=[2^{-\kappa(l-n)}n^{d-1}],
$$

where $\kappa > 1$ is such that $a > (1 + \kappa)\beta$. We define the *m*-term approximant

$$
A_m(f) := S_{Q_n}(f) + \sum_{l>n} \sum_{\mathbf{s} \in G_l} \delta_{\mathbf{s}}(f_l).
$$

Then

$$
m \le |Q_n| + \sum_{l>n} 2^l m_l \ll 2^n n^{d-1}.
$$
\n(4.11)

For the error of approximation we obtain from [\(4.10\)](#page-25-0)

$$
||f - A_m(f)||_p \le \sum_{l>n} ||f'_l||_p \ll 2^{-(a-\beta)n} n^{(b-\beta)(d-1)+\epsilon}.
$$
 (4.12)

Relations [\(4.11\)](#page-25-1) and [\(4.12\)](#page-25-2) imply the required upper bound.

In the case $\beta < a \leq 2\beta$ the proof repeats the corresponding argument from the proof of Theorem [1.3](#page-5-1) in Section 2. Instead of (2.1) we use (4.9) . Also, in the case $a = 2\beta$ the factor $\log \log m$ is included in $(\log m)^{\epsilon}$.

The lower bounds with $\epsilon = 0$ in the case of small smoothness $a \leq 2\beta$ follow from the univariate case. We now consider the case of large smoothness $a > 2\beta$. In the case $p = 2$ it is proved in the proof of Theorem [4.1.](#page-21-2) We use the same example to prove the lower bounds for $p < 2$. Instead of [\(4.8\)](#page-23-2) by Theorem [1.6](#page-6-0) we obtain

$$
\sigma_m(g)_p^p \gg \sum_{\|\mathbf{s}\|_1 = n} \sum_{\mathbf{k} \in \rho(\mathbf{s}) \backslash K_m} \left((|\hat{g}|\mathbf{k})|^2 \right)^{1/2} 2^{n(1/2 - 1/p)} \pmb{\}}^p \gg n^{d-1} 2^{n(p-1)}.\tag{4.13}
$$

Relations [\(4.7\)](#page-23-1) and [\(4.13\)](#page-25-3) imply

$$
\sigma_m(\mathbf{W}_1^{a,b})_p \gg 2^{n(1-1/p-a)} n^{(d-1)(b+1/p-1)} \asymp m^{-a+\beta} (\log m)^{(d-1)(a+b-1)}.
$$

 \Box

Consider a class $\bar{\mathbf{W}}_q^{a,b}$, which consists of functions f with a representation

$$
f = \sum_{n=1}^{\infty} t_n
$$
, $t_n \in \mathcal{T}(Q_n)$, $||t_n||_q \leq 2^{-an} n^{b(d-1)}$.

It is easy to see that in the case $1 < q < \infty$ classes $\bar{W}_q^{a,b}$ and $W_q^{a,b}$ are equivalent. Embedding of $\mathbf{W}_q^{a,b}$ into $\bar{\mathbf{W}}_q^{a,b}$ is obvious and the opposite embedding follows from the inequality for $f \in \mathbf{W}_q^{a,b}$

$$
||f_l||_q = ||(S_{Q_l} - S_{Q_{l-1}})(f)||_q \ll \sum_{n \geq l} ||t_n||_q \ll 2^{-al}(\bar{l})^{b(d-1)}.
$$

In the case $q = 1$ classes $\bar{\mathbf{W}}_1^{a,b}$ are wider than $\mathbf{W}_1^{a,b}$. However, the results of this section hold for these classes as well.

Remark [4.1](#page-21-2). Theorems 4.1 and [4.2](#page-24-2) hold for the class $\bar{\mathbf{W}}_1^{a,b}$ instead of $\mathbf{W}_1^{a,b}$.

5 Discussion

The effect of *small smoothness* in the behavior of asymptotic characteristics of smoothness classes was discovered by Kashin [\[7\]](#page-29-8) in 1981. He proved that the rate of decay of the Kolmogorov widths $d_n(W_1^r, L_p)$ of the univariate classes W_1^r depends on r differently in the range $1 - 1/p < r < 1$ (small smoothness) and in the range $r > 1$. Belinskii [\[1\]](#page-28-1) studied the univariate m-term trigonometric approximation and observed the small smoothness effect in that setting. Romanyuk [\[10\]](#page-29-5) conducted a detailed study of m-term trigonometric approximation of classes of multivariate functions with small mixed smoothness. The Besov classes $\mathbf{B}^r_{q,\theta}$ are studied in [\[10\]](#page-29-5). Define

$$
||f||_{\mathbf{H}_q^r} := \sup_{\mathbf{s}} ||\delta_{\mathbf{s}}(f)||_q 2^{r||\mathbf{s}||_1},
$$

and for $1 \leq \theta < \infty$ define

$$
||f||_{\mathbf{B}_{q,\theta}^r} := \left(\sum_{\mathbf{s}} \left(||\delta_{\mathbf{s}}(f)||_q 2^{r||\mathbf{s}||_1}\right)^{\theta}\right)^{1/\theta}.
$$

We write $\mathbf{B}_{q,\infty}^r := \mathbf{H}_q^r$. With a little abuse of notation, denote the corresponding unit ball

$$
\mathbf{B}_{q,\theta}^r := \{f : \|f\|_{\mathbf{B}_{q,\theta}^r} \le 1\}.
$$

In case of approximation in L_p , $2 < p < \infty$, Lemma [1.1](#page-2-0) was used in [\[10\]](#page-29-5). This makes the corresponding results in [\[10\]](#page-29-5) nonconstructive. We note that the bound for the m-term approximation error in Lemma [1.1](#page-2-0) follows from Theorem [1.1](#page-2-0) and extra property $\theta_m \subset \theta_n$ in Lemma 1.1 follows from the proof of Theorem [1.1](#page-3-0) in [\[16\]](#page-29-0). Thus, Theorem [1.1](#page-3-0) makes Lemma [1.1](#page-2-0) constructive and, therefore, the nonconstructive results from [\[10\]](#page-29-5), which are based on Lemma [1.1,](#page-2-0) are made constructive in this way. Also, the use of Theorem [1.1](#page-3-0) is technically easier than the use of Lemma [1.1.](#page-2-0) For instance, in the proof of upper bounds in Theorem [3.1](#page-12-1) we estimate $\|g_A\|_A$ in a rather simple way because of additivity property of the norm $\|\cdot\|_A$ and then apply Theorem [1.1](#page-2-0) to g_A . Typically, in [\[10\]](#page-29-5) Lemma 1.1 is applied to individual dyadic blocks $\delta_{s}(f)$ with m_{s} depending on the norm of the $\delta_{s}(f)$. It would be interesting to see how much the technique, based on Theorem [1.1,](#page-3-0) could simplify the study of $\sigma_m(\mathbf{B}_{q,\theta}^r)_p$.

Let us make some comparison of our results on the \mathbf{W}_q^r classes with known results on $\mathbf{B}_{q,\theta}^r$ classes. It follows from Corollary [1.2](#page-6-1) that for $1 < q \leq 2$ we have

$$
||f||_{\mathbf{B}_{q,2}^r} \ll ||f||_{\mathbf{W}_q^r} \ll ||f||_{\mathbf{B}_{q,q}^r}.
$$

For example, in the case $\beta p' < r < 1/q$ Theorem [3.1](#page-12-1) gives

$$
\sigma_m(\mathbf{W}_q^r)_p \asymp m^{-(r-\beta)p/2} (\log m)^{(d-1)(r(p-1)-\beta p)}.\tag{5.1}
$$

The corresponding results from [\[10\]](#page-29-5) give

$$
\sigma_m(\mathbf{B}_{q,\theta}^r)_p \asymp m^{-(r-\beta)p/2} (\log m)^{(d-1)((r-1/q)(p-1)+1-1/\theta)}.\tag{5.2}
$$

In the case $\theta = q$ the right hand sides of [\(5.1\)](#page-27-0) and [\(5.2\)](#page-27-1) coincide. This means that our results for a wider class \mathbf{W}_q^r imply the corresponding results for a smaller class $\mathbf{B}_{q,q}^r$. Relations [\(5.1\)](#page-27-0) and [\(5.2\)](#page-27-1) show that $\sigma_m(\mathbf{W}_q^r)_p$ and $\sigma_m(\mathbf{B}_{q,2}^r)_p$ have different orders.

As we already pointed out in the Introduction the main novelty of the paper is in providing constructive algorithms for optimal m-term trigonometric approximation on classes with small mixed smoothness. This is achieved by using Theorem [1.1.](#page-3-0) The use of Theorem [1.1](#page-3-0) is simpler than the use of Lemma [1.1](#page-2-0) traditionally used in this area of research. In addition to traditional use of Theorem [1.6,](#page-6-0) which goes back to papers $[11]$ and $[12]$, we use other deep results from the hyperbolic cross approximation theory – Theorem [1.7,](#page-6-3) Theorem [1.8](#page-6-2) and Lemma [4.2.](#page-21-0) We also prove a new result – Lemma [4.3.](#page-24-0) These results allowed us to treat the case $q = 1$ (see Section 4).

A number of interesting unresolved problems on m-term trigonometric approximation is discussed in [\[16\]](#page-29-0), Section 6. This paper makes a progress in some of them. For instance, Theorems [1.2](#page-3-1) and [3.6](#page-19-1) cover the case $\beta p'$ $r \leq 1/q$ for constructive *m*-term approximation of \mathbf{W}_q^r classes. The case $\beta < r \leq \beta p'$ is still open. There is no progress on small smoothness classes in the case $2 \le q < p < \infty$. In the case $q = 1$ results presented in Section 4 are optimal up to a factor $(\log m)^{\epsilon}$ with arbitrarily small $\epsilon > 0$. It would be interesting to find right orders of $\sigma_m(\mathbf{W}_{1}^{a,b})_p$ and right orders of constructive m-term approximation of these classes.

The reader can find a detailed discussion of greedy algorithms in Banach spaces in [\[14\]](#page-29-9) and their applications for the m-term trigonometric approximation in [\[4\]](#page-28-4), [\[13\]](#page-29-6), [\[15\]](#page-29-10), and [\[16\]](#page-29-0).

References

- [1] E. S. Belinskii, Approximation by a "floating" system of exponentials on classes of smooth periodic functions, Matem. Sb. 132 (1987), 20- 27; English translation in Math. USSR Sb. 60 (1988).
- [2] E.S. Belinskii, Approximation by a "floating" system of exponentials on classes of periodic functions with bounded mixed derivative, Research on the theory of functions of many real variables, Yaroslavl' State University, 1988, 16–33 (in Russian).
- [3] R.A. DeVore and V.N. Temlyakov, Nonlinear approximation by trigonometric sums, J. Fourier Analysis and Applications, 2 (1995), 29–48.
- [4] S.J. Dilworth, D. Kutzarova, V.N. Temlyakov, Convergence of some Greedy Algorithms in Banach spaces, The J. Fourier Analysis and Applications 8 (2002), 489–505.
- [5] E.D. Gluskin, Extremal properties of orthogonal parallelpipeds and their application to the geometry of Banach spaces, Math USSR Sbornik 64 (1989), 85–96.
- [6] R.S. Ismagilov, Widths of sets in normed linear spaces and the approximation of functions by trigonometric polynomials, Uspekhi Mat. Nauk, 29 (1974), 161–178; English transl. in Russian Math. Surveys, 29 (1974).
- [7] B.S. Kashin, On widths of Sobolev classes of small smoothness, Vestnik MGU, Mat. i Mekh., 5 (1981), 50–54 (in Russian).
- [8] V.E. Maiorov, Trigonometric diameters of the Sobolev classes W_p^r in the space L_q , *Math. Notes* **40** (1986), 590–597.
- [9] Y. Makovoz, On trigonometric n-widths and their generalizations, J. Approx. Theory 41 (1984), 361-366.
- $[10]$ A.S. Romanyuk, Best M-term trigonometric approximations of Besov classes of periodic functions of several variables, Izvestia RAN, Ser. Mat. 67 (2003), 61–100; English transl. in Izvestiya: Mathematics $(2003), 67(2):265.$
- [11] V.N. Temlyakov, Approximation of Periodic Functions of Several Variables by Bilinear Forms, Izvestiya AN SSSR, 50 (1986), 137–155; English transl. in Math. USSR Izvestija, 28 (1987), 133–150.
- [12] V.N. Temlyakov, Approximation of functions with bounded mixed derivative, Trudy MIAN, 178 (1986), 1–112. English transl. in Proc. Steklov Inst. Math., 1 (1989).
- [13] V.N. Temlyakov, Greedy-Type Approximation in Banach Spaces and Applications, Constr. Approx. 21 (2005), 257–292.
- [14] V.N. Temlyakov, Greedy approximation, Cambridge University Press, 2011.
- [15] V.N. Temlyakov, Sparse approximation and recovery by greedy algorithms in Banach spaces, Forum of Mathematics, Sigma, 2 (2014), e12, 26 pages; IMI Preprint, 2013:09, 1–27; [arXiv:1303.6811v](http://arxiv.org/abs/1303.6811)1, 27 Mar 2013.
- [16] V.N. Temlyakov, Constructive sparse trigonometric approximation and other problems for functions with mixed smoothness, arXiv: 1412.8647v1 [math.NA] 24 Dec 2014, 1–37.
- [17] H. Triebel, Bases in Function Spaces, Sampling, Discrepancy, Numerical Integration, European Mathematical Society, Germany, 2010.