

The Role of Topology in Quantum Tomography

Michael Kech,^{1,*} Péter Vrana,^{2,†} and Michael M. Wolf^{1,‡}

¹*Department of Mathematics, Technische Universität München, 85748 Garching, Germany*

²*Department of Geometry, Budapest University of Technology and Economics,
Egry József u. 1., 1111 Budapest, Hungary*

(Dated: June 17, 2015)

We investigate quantum tomography in scenarios where prior information restricts the state space to a smooth manifold of lower dimensionality. By considering stability we provide a general framework that relates the topology of the manifold to the minimal number of binary measurement settings that is necessary to discriminate any two states on the manifold.

We apply these findings to cases where the subset of states under consideration is given by states with bounded rank, fixed spectrum, given unitary symmetry or taken from a unitary orbit. For all these cases we provide both upper and lower bounds on the minimal number of binary measurement settings necessary to discriminate any two states of these subsets.

Keywords: quantum tomography, prior information, topology, smooth embeddings

Contents

I. Introduction	2
II. Preliminaries	4
III. Topological Analysis of Measurements	5
A. Stable Measurements	5
B. Generalized Measurements and Smooth Embeddings	10
IV. Upper and Lower Bounds for Concrete Submanifolds	12
A. States of Fixed Spectrum and States of Bounded Rank	13
B. States with Unitary Symmetry	15
C. Unitarily Invariant States of Fixed Spectrum	16
D. Bob-Unitary Orbit	16
A. Technical appendix	21
1. Proof of Theorem III.3	21
2. Proof of Lemma III.5	24
3. Proof of Proposition III.10	26
4. Proof of Lemma III.13	26
5. Proof of Proposition IV.6	28
6. Proof of Proposition IV.8	30
References	30

*Electronic address: kech@ma.tum.de

†Electronic address: vranap@math.bme.hu

‡Electronic address: wolf@ma.tum.de

I. INTRODUCTION

The reconstruction of a quantum state from the outcome of an experiment, called quantum state tomography, is a task of fundamental importance in quantum information science. Already for small systems this task may be non-trivial, requiring many measurements and extensive postprocessing to reconstruct a state. With growing system size this complexity becomes exceedingly relevant [1].

There are at least three kinds of resources that can be considered in this context: (i) the number of measurement settings or, mathematically equivalent, the number of measurement outcomes if a single generalized measurement is considered, (ii) the number of samples to be measured, i.e., the sampling complexity, which takes the statistics into account and (iii) the classical post-processing that is required to interpret the data. In this work we will focus on (i).

We are interested in cases where prior information is available that effectively restricts the state space to a submanifold of lower dimensionality. This information may concern the rank of the density operator, its spectrum, symmetry, energy, associated particle number or other properties and combinations thereof.

The question behind our analysis is: what is the minimal number of binary measurement settings that is required to uniquely identify the state under the assumption that it is taken from the given submanifold? Motivated by the results in [2] our aim is a better understanding of the relation between the minimal number of required measurement settings and the topology of the considered submanifold. Such a relation is most clear in low dimensional examples: suppose the submanifold forms a Klein bottle. Then, although it is two-dimensional, it requires at least four binary measurement settings to identify every point since, loosely speaking, in less than four dimensions the Klein bottle has no realization without self-intersections.

This topological reasoning was introduced in [2] and there applied successfully for instance to the case of pure state quantum tomography. The latter has been a topic of active research in quantum information theory [2–12], closely related to the problem of phase retrieval [13, 14].

On a Hilbert space of dimension d , the set of pure states is of dimension $2d - 2$, whereas the set of all states is of dimension $d^2 - 1$. Consequently, in order to uniquely identify an arbitrary state, one has to at least perform $d^2 - 1$ different binary measurements, whereas in the case of pure states one can hope that $\mathcal{O}(d)$ measurements suffice to uniquely identify a state. In [2] it was shown that to leading order $4d$ binary measurements are necessary and sufficient to identify pure states and the compressed sensing approach of [11, 12, 15] provides an algorithm based on $\mathcal{O}(dr \log(d))$ binary measurements with which a $d \times d$ matrix of rank r can be reliably identified.

The approach we take in this paper extends the results of [2] and gives a general framework for the validity of the topological reasoning in quantum tomography. Thereby, we show that the approach is applicable in the presence of statistical fluctuations, imprecise prior information or inaccuracies in the implementation of the measurement set-up. Moreover, we provide a detailed analysis of a variety of old and new examples of submanifolds.

Outline. We consider measurements as smooth maps from a smooth submanifold of

states into Euclidean space. The methods we deploy to find bounds on the number of measurement outcomes necessary to identify a state of a given submanifold uniquely rely on the technical assumption that this smooth map is a smooth embedding.

In section III we give an operational meaning of the smooth embedding assumption and we determine the relation of quantum tomography to the embedding problem in differential topology.

First, in subsection III A, we justify the smooth embedding assumption by relating it to properties one would generally require of measurements. More precisely, we give two natural notions of stability and we show that these are in fact equivalent to the measurement being a smooth embedding. In the sense of these stability properties our approach is robust with respect to noise.

Secondly, in subsection III B, we generalize the measurement scheme by allowing for measurements on several copies of a state. We then show that any smooth embedding can be approximated by these generalized measurements. This proves that asking for the minimal number of measurement that is needed to identify all states of a given submanifold of states is equivalent to asking for the minimal dimension in which this manifold can be embedded.

Having justified our methods of finding bounds in section III, we devote section IV to applying this method in concrete scenarios. We obtain upper and lower bounds on the number of measurement outcomes necessary to identify states of certain interesting submanifolds. The lower bounds result from topological obstructions, whereas the upper bounds rely on the explicit construction of measurement schemes. The methods used in this section are very different from the the ones used in the first two sections and from this point of view section IV can be read independently.

First, we investigate states of fixed spectrum and we relate these to states of bounded rank in subsection IV A. More precisely, we present lower bounds on the number of measurements necessary to identify states of fixed spectrum and these lower bounds turn out to be very close to the upper bounds for states of bounded rank obtained in [2]. In this way we obtain good upper and lower bounds for both the states of fixed spectrum and the states with bounded rank.

In subsection IV B, we obtain lower and upper bounds for states with a unitary symmetry and we use this to obtain both lower and upper bounds for states of fixed spectrum with a unitary symmetry in subsection IV C.

Finally, in subsection IV D, we obtain upper and lower bounds for states in a bipartite system that lie in the Bob-unitary orbit of a certain pure state, i.e. we consider all states that can be reached from a given pure state by acting with a unitary matrix that just effects Bob's subsystem. Physically, this scenario may correspond to an interferometry experiment. Note that if the initial state is maximally entangled, this orbit is the set of maximally entangled states which may be interesting in its own right. Identifying a maximally entangled state is equivalent to determining the unitary matrix that acted on Bob's subsystem. So this method can also be used for process-tomography of unitary time evolutions, complementing the results in [16, 17].

Proofs of technical results can be found in the appendix.

II. PRELIMINARIES

Let \mathcal{H} be a finite dimensional Hilbert space. We denote by $\mathcal{B}(\mathcal{H})$ the complex vector space of linear operators on \mathcal{H} . $H(\mathcal{H})$ denotes the real vector space of hermitian operators on \mathcal{H} and $H(\mathcal{H})_0$ denotes the real vector space of traceless hermitian matrices, i.e. $H(\mathcal{H})_0 := \{h \in H(\mathcal{H}) : \text{tr}(h) = 0\}$. Throughout we consider these spaces as inner product spaces equipping them with the Hilbert-Schmidt inner product. Furthermore, $\mathcal{S}(\mathcal{H})$ will denote the set of quantum states on \mathcal{H} , i.e. $\mathcal{S}(\mathcal{H}) := \{\rho \in H(\mathcal{H}) : \rho \geq 0, \text{tr}(\rho) = 1\}$.

A positive operator valued measure (POVM) corresponds to a set of positive semidefinite operators $P := \{P_1, \dots, P_m\}$ in $H(\mathcal{H})$ such that

$$\sum_{i=1}^m P_i = \mathbb{1}_{\mathcal{H}}.$$

An element of P is called an effect operator. We define the dimension of P by $\dim P = m - 1$. In quantum mechanics, POVMs are used to describe general measurements [18, 19].

There is an operator system ¹ σ_P associated to each POVM P given by the complex linear span of the operators of P . For an operator system σ denote by $\sigma^{\mathbb{R}}$ the real vector space of hermitian operators in σ , i.e. $\sigma^{\mathbb{R}} := \{h \in \sigma : h^\dagger = h\}$ ². In the following we assume the effect operators of a POVM P to be linearly independent over \mathbb{C} . Note that by this convention $\dim P = \dim \sigma_P - 1$. To each operator system σ one can associate the orthogonal projection from $H(\mathcal{H})$ to $\sigma^{\mathbb{R}} \subseteq H(\mathcal{H})$. Throughout we denote this associated projection by π_σ .

Definition II.1. A POVM $P := \{P_1, \dots, P_m\}$ induces a linear map

$$\begin{aligned} h_P : H(\mathcal{H}) &\rightarrow \mathbb{R}^m \\ \rho &\mapsto (\text{tr}(P_1\rho), \dots, \text{tr}(P_m\rho)). \end{aligned}$$

P is called \mathcal{R} -complete for a subset $\mathcal{R} \subseteq \mathcal{S}(\mathcal{H})$ if $h_P|_{\mathcal{R}}$ is injective and it is called \mathcal{P} -embedding for a smooth submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ if $h_P|_{\mathcal{P}}$ is a smooth embedding ³.

Recall that the question behind our analysis concerns the minimal m for which there is a \mathcal{P} -complete POVM for a given smooth submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ that characterized the available prior information. From the dimension $D := \dim \mathcal{P}$ alone one obtains that $m \geq D$ is necessary and $m = 2D + 1$ is generally sufficient [2]. For better bounds, one has to invoke more of the (topological) structure of the manifold.

In the following all manifolds and submanifolds are assumed to be smooth. Throughout we regard both $\mathcal{S}(\mathcal{H})$ and submanifolds $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ with $\mathcal{H} \simeq \mathbb{C}^n$ as submanifolds of $H(\mathcal{H}) \simeq \mathbb{R}^{n^2}$ equipped with the subspace topology and the standard smooth structure.

¹ An operator system $\sigma \subseteq \mathcal{B}(\mathcal{H})$ is a linear subspace such that $\mathbb{1}_{\mathcal{H}} \in \sigma$ and $\sigma^\dagger = \sigma$.

² Note that $\sigma^{\mathbb{R}}$ determines σ uniquely.

³ A smooth mapping $\psi : M \rightarrow N$ is called a smooth embedding if $d\psi_x$ is injective for all $x \in M$ and ψ is a homeomorphism onto its image.

We often use this picture to identify the tangent space at a point $\rho \in \mathcal{P}$, $T_\rho \mathcal{P}$, with a linear subspace in $H(\mathcal{H})$, i.e. we think of tangent vectors $v \in T_\rho \mathcal{P}$ as hermitian operators. We assume submanifolds $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ to be closed and without boundary. In particular, this means that \mathcal{P} is an embedded submanifold by the compactness of $\mathcal{S}(\mathcal{H})$, i.e. the inclusion is a homeomorphism onto its image.

III. TOPOLOGICAL ANALYSIS OF MEASUREMENTS

A. Stable Measurements

Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a submanifold. In order for our methods for finding bounds on the dimension of \mathcal{P} -complete POVMs to apply, we need the technical requirement that these POVMs are \mathcal{P} -embeddings. In this section we justify this assumption. We develop two notions of stability for a \mathcal{P} -complete POVM and we show that these notions are equivalent to the POVM being a \mathcal{P} -embedding. These notions of stability are properties one would naturally require for \mathcal{P} -complete POVMs. Thus, under the premise of stability, \mathcal{P} -complete POVMs are \mathcal{P} -embeddings.

For a given POVM P the notions of \mathcal{P} -embedding and \mathcal{P} -completeness just depend on its associated operator system σ_P as the following proposition shows.

Proposition III.1. *Let P be a POVM, h_P be the associated linear map, $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a submanifold and let $\pi_P : H(\mathcal{H}) \rightarrow \sigma_P^{\mathbb{R}}$ be the orthogonal projection on $\sigma_P^{\mathbb{R}}$.*

1. *P is \mathcal{P} -complete if and only if $\pi_P|_{\mathcal{P}}$ is injective.*
2. *P is a \mathcal{P} -embedding if and only if h_P is \mathcal{P} -complete and $d(\pi_P|_{\mathcal{P}})_\rho = \pi_P|_{T_\rho \mathcal{P}}$ is injective for each $\rho \in \mathcal{P}$.*

Proof. Since we equipped $H(\mathcal{H})$ with the Hilbert-Schmidt inner product, by the definition of h_P , we get $\sigma_P^{\mathbb{R}} = \text{span}_{\mathbb{R}} P \subseteq \ker(h_P)^\perp$ and since $\text{rank } h_P = \dim \sigma_P^{\mathbb{R}}$, we get $\sigma_P^{\mathbb{R}} = \ker(h_P)^\perp$ by dimensional reasons. So $h_P = h_P \circ \pi_P$.

For the first statement, let P be \mathcal{P} -complete and h_P be the associated linear map. Since $h_P|_{\mathcal{P}}$ is injective and $h_P = h_P \circ \pi_P$, we get that $\pi_P|_{\mathcal{P}}$ is injective.

Conversely, let $\pi_P|_{\mathcal{P}}$ be injective. Then, since $h_P = h_P \circ \pi_P$, $h_P|_{\mathcal{P}}$ is injective because $\pi_P|_{\mathcal{P}}$ is injective and h_P is injective restricted to the image of π_P .

Noting that by linearity we have $d(h_P|_{\mathcal{P}})_\rho = dh_P|_{T_\rho \mathcal{P}} = h_P|_{T_\rho \mathcal{P}}$, the above reasoning also applies for the second statement. \square

For a submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$, \mathcal{P} -completeness and being a \mathcal{P} -embedding are the only properties of a POVM we are interested in. Thus, by proposition III.1, there is a natural equivalence relation on the set of POVMs, namely

$$P \sim P' \quad \Leftrightarrow \quad \sigma_P = \sigma_{P'}.$$

Since every n -dimensional operator system is generated by an $(n-1)$ -dimensional POVM [2], the operator systems are precisely the equivalence classes.

Since the proofs we give are easier to formulate using operator system we often state our results in terms of operator systems and then transfer them to POVMs.

Let $\Sigma(n)$ be the set of n -dimensional operator systems. For a subset $\mathcal{R} \subseteq \mathcal{S}(\mathcal{H})$ we call $\sigma \in \Sigma(n)$ \mathcal{R} -complete if $\pi_\sigma|_{\mathcal{R}}$ is injective and for a submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ we call $\sigma \in \Sigma(n)$ a \mathcal{P} -embedding if $\pi_\sigma|_{\mathcal{P}}$ is a smooth embedding.

A metric on $\Sigma(n)$, which is natural for our purpose, can be defined in terms of any norm on the corresponding linear map π_σ . For an arbitrary linear map $L : H(\mathcal{H}) \rightarrow H(\mathcal{H})$ we consider

$$\|L\|_{op} = \sup_{B \in H(\mathcal{H}), \|B\| \leq 1} \|L(B)\|$$

where $\|\cdot\|$ denotes the Hilbert-Schmidt norm. The sought metric is then given by $d(\sigma, \sigma') = \|\pi_\sigma - \pi_{\sigma'}\|_{op}$. The following definition refers to the metric topology induced on $\Sigma(n)$.

Definition III.2. (*Stability*). *Let $\mathcal{R} \subseteq \mathcal{S}(\mathcal{H})$ be a subset. An \mathcal{R} -complete operator system $\sigma \in \Sigma(n)$ is called stably \mathcal{R} -complete if there exists a neighbourhood $N \subseteq \Sigma(n)$ of σ such that every $\sigma' \in N$ is an \mathcal{R} -complete operator system. A POVM P is called stably \mathcal{R} -complete if its associated operator system σ_P is \mathcal{P} -complete.*

Remark In the following we will see that closeness of POVMs is equivalent to closeness of the associated operators systems. Thus, this definition says that a stably \mathcal{P} -complete POVM P is robust against inaccuracy in its implementation in the sense that every close enough POVM is also \mathcal{P} -complete.

The intuition behind this definition is best envisioned by thinking of operator systems as planes in $H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$, see figure 1.




FIG. 1: This figure shows a submanifold $\mathcal{P} \subseteq H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$ and two close operator systems $\sigma, \sigma' \subseteq H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$ that are \mathcal{P} -complete.

Now we are in a position to state one of the main results of this section.

Theorem III.3. (*Stable measurements are embeddings*). *Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a closed submanifold and let $\sigma \in \Sigma(n)$ be \mathcal{P} -complete. Then σ is stably \mathcal{P} -complete if and only if it is a \mathcal{P} -embedding.*

Since the proof of this theorem is rather lengthy we relegated it to appendix A 1.

Theorem III.3 is a statement about operator systems. In order to provide it with an operational meaning, we prove the corresponding stability result for POVMs in the following.

Corollary III.4. *Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a closed submanifold and let $P := \{P_1, \dots, P_m\}$ be a \mathcal{P} -complete POVM of dimension $m-1$ with associated linear map h_P . P is a \mathcal{P} -embedding if and only if there is an $\epsilon > 0$ such that every POVM Q with $\sup_{v \in H(\mathcal{H}), \|v\| \leq 1} \|(h_P - h_Q)v\| < \epsilon$ is \mathcal{P} -complete.*

Proof. Let $\mathcal{H} = \mathbb{C}^n$ and let π_P be the orthogonal projection associated to $P := \{P_1, \dots, P_m\}$.

Let $\epsilon > 0$ such that every POVM Q with $\sup_{v \in H(\mathcal{H}), \|v\| \leq 1} \|(h_P - h_Q)v\| < \epsilon$ is \mathcal{P} -complete. We show, that there is $\delta > 0$ such that for all $\sigma' \in \Sigma(m)$ with

$$\|\pi_P - \pi_{\sigma'}\|_{op} < \delta$$

there is a POVM $P' := \{P'_1, \dots, P'_m\}$ with $\sigma_{P'} = \sigma'$ and $\sup_{v \in H(\mathcal{H}), \|v\| \leq 1} \|(h_P - h_{P'})v\| < \epsilon$, hence $\pi_{\sigma'}$ is \mathcal{P} -complete by proposition III.1.

For every $\eta > 0$, we can slightly deform P to a POVM $\tilde{P} := \{\tilde{P}_1, \dots, \tilde{P}_m\}$ with full rank effect operators such that $\|P_i - \tilde{P}_i\| < \eta$ and $\sigma_P = \sigma_{\tilde{P}}$: Let $\tilde{P}_i := \frac{\sqrt{n}}{\sqrt{n+\eta}}(P_i + \frac{\eta}{\sqrt{n}}\mathbb{1}_{\mathcal{H}})$ for $i = 1, \dots, m$. Then, for $i = 1, \dots, m$, $\|P_i - \tilde{P}_i\| = \eta \frac{\sqrt{n}}{\sqrt{n+\eta}} < \eta$. Note that we also ensured that the smallest eigenvalue of \tilde{P}_i is bigger than $\eta/2$ for $i = 1, \dots, m$ and η small enough.

For some $\sigma' \in \Sigma(m)$ with

$$\|\pi_P - \pi_{\sigma'}\|_{op} < \delta$$

let $P'_i := \pi_{\sigma'}(\tilde{P}_i)$, $i = 1, \dots, m$. Then, for $i = 1, \dots, m$,

$$\|\tilde{P}_i - P'_i\| = \|\pi_P(\tilde{P}_i) - \pi_{\sigma'}(\tilde{P}_i)\| < \sqrt{n}\delta$$

and thus the P'_i are positive for $\sqrt{n}\delta < \eta/2$. Furthermore, $\sum_{i=1}^m P'_i = \pi_{\sigma'}(\sum_{i=1}^m \tilde{P}_i) = \pi_{\sigma'}(\mathbb{1}_{\mathcal{H}}) = \mathbb{1}_{\mathcal{H}}$. For small enough δ , the P'_i are linearly independent because the \tilde{P}_i are linearly independent by assumption. Thus $\sigma' = \sigma_{P'}$ by dimensional reasoning. Finally,

$$\begin{aligned} & \|h_P - h_{P'}\|_{op}^2 \\ & \leq \sum_{i=1}^m \|P_i - P'_i\|^2 = \sum_{i=1}^m \|P_i - \tilde{P}_i + \tilde{P}_i - P'_i\|^2 \\ & \leq \sum_{i=1}^m (\|P_i - \tilde{P}_i\| + \|\tilde{P}_i - P'_i\|)^2 \\ & < m(\eta^2 + n\delta^2). \end{aligned}$$

By choosing η and δ so small that $m(\eta^2 + n\delta^2) < \epsilon^2$, $h_{P'}$ is injective by assumption and thus σ' is injective by III.1. Thus, σ_P is stably \mathcal{P} -complete and III.3 concludes the proof of this direction.

Conversely, suppose σ is a \mathcal{P} -embedding. Corollary A.2 states, that there is an $\epsilon > 0$ such that every POVM Q with $\sup_{B \in H(\mathcal{H}), \|B\| \leq 1} \|h_P(B) - h_Q(B)\| < \epsilon$ is a \mathcal{P} -embedding and thus in particular \mathcal{P} -complete. \square

However, the notion of stability for measurements developed so far may not be satisfactory yet since it just considers inaccuracy in the implementation of the measurement set-up. Noisiness of the outcome, resulting from e.g. dissipation or finite statistics, or noisiness of the input, originating from e.g. inaccurate prior information, are inevitable but not considered in the definition.

In the remainder of this section we show that also from this point of view, stably \mathcal{P} -complete is a operationally meaningful property.

The idea of the following lemma, which is the essential ingredient for the second theorem of this section, is to construct a neighbourhood for every point of submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ that can be approximated by the tangent space at that point. Let $\pi^T : \mathcal{P} \rightarrow \mathcal{B}(H(\mathcal{H}))$ be the mapping that associates to each point $\rho \in \mathcal{P}$ its orthogonal projection π_ρ^T to $T_\rho \mathcal{P} \subseteq H(\mathcal{H})$ and let π^N be the analogue mapping for the normal space, i.e. $\pi_\rho^N + \pi_\rho^T = \text{id}_{H(\mathcal{H})}$ for all $\rho \in \mathcal{P}$. Furthermore, $B_x(\epsilon)$ denotes the open ball with center $x \in H(\mathcal{H})$ and radius $\epsilon > 0$, i.e. $B_x(\epsilon) := \{y \in H(\mathcal{H}) : \|x - y\| < \epsilon\}$ and d denotes the metric induced by $\|\cdot\|$.

Lemma III.5. *Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a submanifold. For every $\eta > 0$ there is an $\epsilon > 0$ such that for all $\rho \in \mathcal{P}$,*

$$\rho' \in B_\epsilon(\rho) \cap \mathcal{P} \Rightarrow \|\pi_\rho^N(\rho' - \rho)\| < \eta \|\rho - \rho'\|.$$

Since the proof of this lemma is rather technical, it is relegated to appendix A 2.

The following theorem is the second main result of this section. It is formulated in terms of operator systems but again the result transfers to POVMs. Since the interpretation of the theorem may not be obvious let us first give some intuition and motivation: Adding small perturbations to states of a submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ can be thought of as blowing up \mathcal{P} to a small tubular neighbourhood $\mathcal{P}_\epsilon = \{\rho \in \mathcal{S}(\mathcal{H}) : d(\rho, \mathcal{P}) < \epsilon\}$. The dimension of \mathcal{P}_ϵ is then equal to the dimension of $H(\mathcal{H})$. Thus, one cannot expect \mathcal{P} -complete POVMs to stay injective when allowing for small errors. However, one can hope for being able to separate points in \mathcal{P}_ϵ that are sufficiently far away, in the sense that $\pi_P(\rho) \neq \pi_P(\rho')$ for $\|\rho - \rho'\| > C\epsilon$ with $C > 0$ a constant. For a given small enough ϵ such a C obviously exists however it is not immediate that C can be chosen independent of ϵ . The following theorem asserts that for ϵ smaller than a certain fixed value, C is independent of ϵ . The existence of a C independent of ϵ means that the measurement can be made arbitrarily precise by reducing the errors.

Theorem III.6. *Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a submanifold and let P be a POVM with associated orthogonal projection π_P . P is stably \mathcal{P} -complete if and only if there exists $\epsilon_0 > 0$ and $C > 2$ such that for all ϵ with $0 < \epsilon < \epsilon_0$,*

$$\rho, \rho' \in \mathcal{P} \text{ with } \|\rho - \rho'\| > \epsilon C \Rightarrow \pi_P(B_\epsilon(\rho)) \cap \pi_P(B_\epsilon(\rho')) = \emptyset \quad .$$

Proof. Let P be stably \mathcal{P} -complete and thus a \mathcal{P} -embedding by theorem III.3 and let $\rho \in \mathcal{P}$. Let $l := \max_{\rho \in \mathcal{P}} \|\pi_P \circ \pi_\rho^T - \pi_\rho^T\|_{op}$ and let η as well as $\tilde{\epsilon}$ be as in lemma III.5. Shrink ϵ_0 such that $C\epsilon_0 < \tilde{\epsilon}$.

Note that $l < 1$ because π_P is an immersion. Without the immersion property we could not assume $l < 1$ and in fact this is the essential idea of this proof ⁴.

Noting that $B = \{(\rho, \rho') \in \mathcal{P}^2 : \|\rho - \rho'\| \geq \epsilon_0 C\}$ is compact, $\kappa := \min_{(\rho, \rho') \in B} \|\pi_P(\rho - \rho')\|$ is attained and thus $\kappa > 0$ by the injectivity of π_P . If necessary, shrink ϵ_0 such that $\epsilon_0 < \kappa/2$. Then, for $\rho' \in \mathcal{P} - (B_{C\epsilon_0}(\rho) \cap \mathcal{P})$ the claim holds because $\|\pi_P(\rho - \rho')\| \geq \kappa > 2\epsilon_0$ and $\pi_P(B_\epsilon(\rho)) = B_\epsilon(\pi_P(\rho))$.

Finally, let $\rho' \in B_{C\epsilon_0}(\rho) \cap \mathcal{P}$ and $\|\rho - \rho'\| > C\epsilon$, $0 < \epsilon < \epsilon_0$. Then,

$$\begin{aligned} \|\pi_P(\rho - \rho')\| &\geq \|\pi_P(\pi_\rho^T(\rho - \rho'))\| - \|\pi_P(\pi_\rho^N(\rho - \rho'))\| \\ &\geq \|\pi_P(\pi_\rho^T(\rho - \rho')) - \pi_\rho^T(\rho - \rho') + \pi_\rho^T(\rho - \rho')\| - \eta\|\rho - \rho'\| \\ &\geq \|\pi_\rho^T(\rho - \rho')\| - \|\pi_P(\pi_\rho^T(\rho - \rho')) - \pi_\rho^T(\rho - \rho')\| - \eta\|\rho - \rho'\| \\ &> \|\rho - \rho'\| \left(\sqrt{1 - \eta^2} - l - \eta \right) > \epsilon C \left(\sqrt{1 - \eta^2} - l - \eta \right), \end{aligned}$$

where we used the fact that, for $0 \leq l < 1$, we can choose η small enough such that $\left(\sqrt{1 - \eta^2} - l - \eta \right) > 0$. Furthermore, we can choose $C > 0$ such that $C \left(\sqrt{1 - \eta^2} - l - \eta \right) > 2$. Since $\pi_P(B_\epsilon(\rho)) = B_\epsilon(\pi_P(\rho))$, this proves the statement.

For the converse, let $\rho, \rho' \in \mathcal{P}$ and $\rho \neq \rho'$. Choosing $\epsilon = \min\{\epsilon_0, \|\rho - \rho'\|/(2C)\}$, we find $\pi_P(B_\epsilon(\rho)) \cap \pi_P(B_\epsilon(\rho')) = \emptyset$ and thus $\pi_P(\rho) \neq \pi_P(\rho')$.

Finally, assume $\pi_P|_{\mathcal{P}}$ is not an immersion at some $\rho \in \mathcal{P}$. Let $\gamma : (-1, 1) \rightarrow \mathcal{P} \subseteq H(\mathcal{H})$ be a smooth curve with $\gamma(0) = \rho$ and $\frac{d}{dt}\gamma(0) = v \in \ker \pi_P$. Let $C > 0$ as in the theorem, then

$$2/C \leq \lim_{t \rightarrow 0} \frac{\|\pi_P(\gamma(t) - \rho)\|}{\|\gamma(t) - \rho\|} \leq \lim_{t \rightarrow 0} \frac{\|\pi_P(\gamma(t) - \rho)\|}{t/2} = 2\|\pi_P(v)\| = 0,$$

a contradiction. Here we assumed $\|\gamma(t) - \rho\| > t/2$ which is clearly true for t small enough by lemma III.5. \square

Remark Note that it is essentially the constant l that determines C . For small l , i.e. in the case where the tangent spaces are not steep with respect to σ_P , we can ensure that C is close to 2 (if we make ϵ_0 small enough). On the other hand if l is close to one C has to be big and in this sense l is a measure for the stability of the POVM P .

ϵ_0 is mainly determined by the constant κ , which is more of ‘‘global’’ nature. Loosely speaking it is a measure for how bad \mathcal{P} wiggles around in $H(\mathcal{H})$.

It is worth noting, that if π_P fails to be an immersion, $C \rightarrow \infty$ for $\epsilon \rightarrow 0$ and from this point of view, stably \mathcal{P} -complete measurements are the ones that can be made arbitrarily precise.

⁴ The bigger l is, the steeper the tangent spaces can be with respect to the operator system σ_P . By the previous lemma we saw that small neighbourhoods around a point $\rho \in \mathcal{P}$ can be approximated by the tangent space at that point. We can ensure that these approximations are so good that the fluctuations around the steepest tangent space have no component orthogonal to σ_P and in this sense we can locally think of \mathcal{P} as a plane.

This theorem transfers to the corresponding theorem for POVMs as the following corollary shows.

Corollary III.7. *Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a submanifold and let P be a POVM with associated linear map h_P . P is a smooth embedding if and only if there exists $\epsilon_0 > 0$ and $C > 2$ such that for all ϵ with $0 < \epsilon < \epsilon_0$,*

$$\rho, \rho' \in \mathcal{P} \text{ with } \|\rho - \rho'\| > \epsilon C \Rightarrow h_P(B_\epsilon(\rho)) \cap h_P(B_\epsilon(\rho')) = \emptyset \quad .$$

Proof. Let π_P be the orthogonal projection associated to P . Let $\lambda := \min_{v \in \text{supp } \pi_P, \|v\|=1} \|h_P(v)\|$ and observe that $\lambda > 0$ since h_P is injective on the support of π_P . Thus $\|h_P(\rho - \rho')\| = \|h_P \circ \pi_P(\rho - \rho')\| > \lambda \|\pi_P(\rho - \rho')\|$. Then, the proposition holds for h_P by replacing C with C/λ . \square

Finally, this result also incorporates robustness against noisiness of the outcome as the following corollary shows.

Corollary III.8. *Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a submanifold and let P be a stably \mathcal{P} -complete POVM with associated linear map h_P . There exists $\epsilon_0 > 0$ and $C > 2$ such that for all ϵ with $0 < \epsilon < \epsilon_0$,*

$$\mathcal{P} \cap h_P^{-1}(B_{2\epsilon}(h_P(\rho))) \subseteq B_{C\epsilon}(\rho) \text{ for all } \rho \in \mathcal{P}.$$

Proof. Let C, ϵ_0 be as in corollary III.7. Let $\rho \in \mathcal{P}$ and $\rho' \in h_P^{-1}(B_{2\epsilon}(h_P(\rho)))$ with $\|\rho - \rho'\| > C\epsilon$. Then, $2\epsilon < \|h_P(\rho) - h_P(\rho')\| < 2\epsilon$, a contradiction. \square

B. Generalized Measurements and Smooth Embeddings

Linear measurements are clearly not sufficient to realize all smooth embeddings. More precisely, if there is a smooth embedding $\phi : \mathcal{P} \subseteq \mathcal{S}(\mathcal{H}) \rightarrow \mathbb{R}^m$, then there need not be an m -dimensional POVM that is a \mathcal{P} -embedding. For example the set $N := \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1, x \geq -0.5\}$ can clearly be embedded in \mathbb{R}^1 , but an injective orthogonal projection has to have rank two. However, the embedding cannot get arbitrarily bad because from Whitney's embedding theorem we know that there is a \mathcal{P} -embedding in Euclidean space of twice the dimension of \mathcal{P} .

In this section we generalize our approach to measurements of the type

$$\text{tr}(\rho^{\otimes n} P_i)$$

and we show that these measurements can approximate any smooth embedding. This means that if there exists a smooth embedding $\psi : \mathcal{P} \subseteq \mathcal{S}(\mathcal{H}) \rightarrow \mathbb{R}^m$, then there is POVM of dimension m that is a \mathcal{P} -embedding. Thus, the problem described in the beginning of this section can be circumvented by this generalized measurement scheme.

Let us fix some notation.

Definition III.9. A measurement $P := \{P_1, \dots, P_m\}$ on k copies is a POVM on $H(\mathcal{H})^{\otimes k}$. P induces a linear map

$$h_P : H(\mathcal{H})^{\otimes k} \rightarrow \mathbb{R}^m$$

$$\rho \mapsto (\text{tr}(P_1 \rho^{\otimes k}), \dots, \text{tr}(P_m \rho^{\otimes k})).$$

Let $i : H(\mathcal{H}) \rightarrow H(\mathcal{H})^{\otimes k}$, $\rho \mapsto \rho^{\otimes k}$. P is called \mathcal{R} -complete for a subset $\mathcal{R} \subseteq \mathcal{S}(\mathcal{H})$ if $h_P \circ i|_{\mathcal{R}} = h_P|_{i(\mathcal{R})}$ is injective and it is called a \mathcal{P} -embedding for a submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ if $h_P|_{i(\mathcal{P})}$ is a smooth embedding.

The following proposition makes the connection to the theory developed in the last section.

Proposition III.10. The mapping $i : H(\mathcal{H}) \rightarrow H(\mathcal{H})^{\otimes k}$, $\rho \mapsto \rho^{\otimes k}$ is smooth. Furthermore, for a smooth closed submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$, $i|_{\mathcal{P}}$ is a smooth embedding.

The proof of this proposition is relegated to appendix A 3.

Remark Let $\Sigma(n, k)$ be the set of n -dimensional operator systems on $\mathcal{B}(\mathcal{H})^{\otimes k}$. Each n -dimensional measurement on k copies P generates an operator system $\sigma_P = \text{span}\{P_i\}_{P_i \in P} \in \Sigma(n, k)$. If $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ is a closed submanifold, $i(\mathcal{P}) \subseteq \mathcal{S}(\mathcal{H}^{\otimes k}) = \mathcal{S}(\mathcal{H})^{\otimes k}$ is a closed submanifold by the previous proposition. So the ideas and results of the last section can be naturally applied to measurements on k copies. In particular for a submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ the notions of \mathcal{P} -embedding and \mathcal{P} -complete naturally apply to the equivalence classes $\sigma \in \Sigma(n, k)$ of measurements on k copies ($P \sim P' \Leftrightarrow \sigma_P = \sigma_{P'}$).

Theorem III.11. Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a closed submanifold and let $\sigma \in \Sigma(n, k)$ be \mathcal{P} -complete. Then σ is stably \mathcal{P} -complete if and only if it is a \mathcal{P} -embedding.

Proof. By the previous remark $i(\mathcal{P}) \subseteq \mathcal{S}(\mathcal{H})^{\otimes k}$ is a closed submanifold. Then the claim follows by applying III.3 to $i(\mathcal{P})$ and σ . \square

Choosing an orthonormal basis $\{\sigma_i\}_{i \in \{1, \dots, d^2\}}$ of $H(\mathcal{H})$ with $\sigma_1 = \mathbb{1}_{\mathcal{H}}$ gives an identification $H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$. Under this identification we can think of elements in $H(\mathcal{H})^{\otimes k}$ as elements in $P^k(\mathbb{R}^{d^2})$, the vector space of polynomial functions of degree k on $H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$.⁵

More precisely, let $\text{Sym}(H(\mathcal{H}), k) \subseteq H(\mathcal{H})^{\otimes k}$ be the vector space of symmetric elements of degree k in $H(\mathcal{H})^{\otimes k}$. Then, use the identification $H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$ to define a linear map

$$\phi : \text{Sym}(H(\mathcal{H}), k) \rightarrow P^k(\mathbb{R}^{d^2}) \tag{1}$$

by the relation

$$\phi(\eta)(x) = \text{tr} \left(\eta \left(\sum_{i=1}^{n^2} x_i \sigma_i \right)^{\otimes n} \right)$$

where $\eta \in \text{Sym}(H(\mathcal{H}), k)$ and $x \in \mathbb{R}^{d^2}$.

⁵ Note that by viewing $H(\mathcal{H})$ as a smooth manifold this corresponds to choosing a particular coordinate system (x_1, \dots, x_{d^2}) .

Lemma III.12. *The mapping ϕ is an isomorphism.*

Proof. Let $d = \dim H(\mathcal{H})$. Note that,

$$\dim \text{Sym}(H(\mathcal{H}), k) = \binom{n+k-1}{k} = \dim P^k(\mathbb{R}^{d^2}).$$

Then, by linearity of ϕ , it is enough to check that ϕ is surjective. Under the identification $H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$, a basis of $P^k(\mathbb{R}^{d^2})$ is given by polynomials of the form $x_{i_1} \dots x_{i_k}$, $i_j \in \{1, \dots, d\}$, $i_1 \leq \dots \leq i_k$. For each such polynomial $p = x_{i_1} \dots x_{i_k}$ there is a $\eta \in \text{Sym}(H(\mathcal{H}), k)$ such that $\phi(\eta)(x) = p$, namely $\eta = \sigma_{i_1} \dots \sigma_{i_k}$, where \cdot denotes the symmetric product. \square

Remark Note that every $\rho \in \mathcal{S}(\mathcal{H})$ decomposes as $\rho = \mathbb{1}_{\mathcal{H}} + \sum_{i=2}^{d^2} \sigma_i$ and thus $x_1 = 1$ on $\mathcal{S}(\mathcal{H})$. From this point of view $P^k(\mathbb{R}^{d^2})$ corresponds to $P^{\leq k}(\mathbb{R}^{d^2-1})$, the set of polynomials of degree $d \leq k$ in x_2, \dots, x_{d^2} .

The following lemma is the crucial ingredient of the main theorem of this section. Let $\mathbb{1}_{\mathcal{H}} + H(\mathcal{H})_0 := \{\mathbb{1}_{\mathcal{H}} + h : h \in H(\mathcal{H})_0\}$.

Lemma III.13. *Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H}) \subseteq \mathbb{1}_{\mathcal{H}} + H(\mathcal{H})_0 \simeq \mathbb{R}^{n \times n-1}$ be a closed submanifold and $\psi : \mathcal{P} \rightarrow \mathbb{R}^m$ be a smooth embedding. Then, there is a $k \in \mathbb{N}$ and a map $\tilde{\psi}' = (p_1, \dots, p_m)$, $p_i \in P^{\leq k}(\mathbb{R}^{n \times n-1})$, such that $\psi' = \tilde{\psi}'|_{\mathcal{P}}$ is a smooth embedding.*

The proof of this lemma can be found in appendix A 4.

Theorem III.14. *Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a closed submanifold. There is a smooth embedding of \mathcal{P} in \mathbb{R}^m if and only if, for some $k \in \mathbb{N}$, there exists a stably \mathcal{P} -complete m -dimensional measurement on k copies.*

Proof. III.11 gives one direction. For the other direction, let $\psi : \mathcal{P} \rightarrow \mathbb{R}^m$ be a smooth embedding. Then, by III.13, there is a smooth embedding $\psi' = (p_1, \dots, p_m)|_{\mathcal{P}}$, $p_i \in P^{\leq k}(\mathbb{R}^{n \times n-1})$. $\sigma = \text{span}_{\mathbb{R}}\{\mathbb{1}_{\mathcal{H}}, \phi^{-1}(p_1), \dots, \phi^{-1}(p_m)\}$ (ϕ from III.12) is clearly an operator system whose dimension is less or equal to $m+1$. σ is a \mathcal{P} -embedding because $\psi'|_{\mathcal{P}}$ is an embedding and thus stably \mathcal{P} -complete by III.11. \square

Thus, under the premise of stability, asking for the minimal dimension of a \mathcal{P} -complete POVM is equivalent to the related problem in differential topology of finding the smallest m such that \mathcal{P} can be smoothly embedded in \mathbb{R}^m .

IV. UPPER AND LOWER BOUNDS FOR CONCRETE SUBMANIFOLDS

In this section we obtain lower as well as upper bounds on the dimension of complete and stable POVMs on some interesting submanifolds of states. The procedure is to first relate the submanifolds to well-known homogeneous spaces and then to obtain or use existing non-immersion results for these. Upper bounds are obtained by directly constructing POVMs. First, we give bounds for the set of states with fixed spectrum. Thereby we also obtain bounds for the closely related set of states with bounded rank.

Then, we give a brief analysis of states with an underlying unitary symmetry which is needed in the next section, where we generalize the previous results to states of fixed spectrum with an underlying symmetry.

Finally, we obtain bounds for the set of pure states of bipartite systems, obtained from the action of the unitary group of the second system on some fixed pure state.

In the following let $\mathcal{H} = \mathbb{C}^n$.

A. States of Fixed Spectrum and States of Bounded Rank

First, we consider the set of states in $\mathcal{S}(\mathbb{C}^n)$ with fixed spectrum $s = (s_1, \dots, s_n)$ ⁶ and we denote by $D_s := \text{diag}(s_1, \dots, s_n)$ the diagonal matrix with entries from s .

The set of all states with spectrum s , $\mathcal{S}(\mathbb{C}^n)_s$, is the orbit of D_s with respect to the action G of $U(n)$ on $\mathcal{S}(\mathbb{C}^n)$ by conjugation, i.e.

$$\mathcal{S}_s^n := \{UD_sU^\dagger : U \in U(n)\}.$$

The isotropy group of ρ under this action is $U(n_1) \times \dots \times U(n_k)$, where n_i is the multiplicity of the i -th biggest eigenvalue. Note that $\sum_{j=1}^k n_j = n$. By theorem 3.62 of [20], factoring the orbit map over this isotropy group induces a diffeomorphism

$$U(n)/U(n_1) \times \dots \times U(n_k) \simeq \mathcal{S}(\mathbb{C}^n)_s.$$

Thus, \mathcal{S}_s^n can be identified with a complex flag manifold.

In [21], Walgenbach obtains lower bounds for the immersion dimension of complex flag manifolds. To present his result, we first introduce some notation.

Definition IV.1. Let $n \in \mathbb{N}$, $k \in \{0, 1, \dots, n\}$.

$\alpha(n) :=$ number of ones in the binary expansion of n ,

$\alpha_1(n) := \sum_{i=0}^{n-1} \alpha(i)$,

$\beta(n, k) := \alpha_1(n) - \alpha_1(k) - \alpha_1(n - k)$,

Let $\{n_1, \dots, n_k\}$ be a partition of n . Let K be some subset of $\{1, \dots, k\}$ and set $m = \sum_{i \in K} n_i$.

Proposition IV.2. [21] *The complex flag manifold $U(n)/U(n_1) \times \dots \times U(n_k)$ cannot be immersed in Euclidean Space of dimension $4m(n - m) - 2\beta(n, m) - 1$ and it cannot be embedded in Euclidean space of dimension $4k(n - m) - 2\beta(n, m)$.*

Next, we want to obtain upper bounds on the dimension of stably \mathcal{S}_s^n -complete POVMs. Let σ be the function that associates to each $h \in H(\mathbb{C}^n)$ its spectrum. For $A \subseteq H(\mathbb{C}^n)$, let $\text{Spec}(A) := \{D_s : s = \sigma(M), M \in A\}$ and let $G(A) := \{UMU^\dagger : U \in U(n), M \in A\}$.

⁶ By spectrum we mean the set of eigenvalues order increasingly together with their multiplicities.

Lemma IV.3. $\Delta\mathcal{S}_s^n = G(\text{Spec}(\Delta\mathcal{S}_s^n))$ and $T\mathcal{S}_s^n = G(\text{Spec}(T_{D_s}\mathcal{S}_s^n))$ as sets. Furthermore, let r be the biggest multiplicity of an eigenvalue in s , then $\text{rank}(M) < 2(n - r) + 1$ for $M \in \Delta\mathcal{S}_s^n \cup T\mathcal{S}_s^n$.

Proof. The first claim is essentially true by definition. For the second claim, let us compute the tangent space at M_s . Let $h \in H(\mathbb{C}^n)$ and consider the curve $\gamma : t \mapsto e^{iht}D_s e^{-iht}$. The derivative at $t = 0$ of this curve is then an element of $T_{D_s}\mathcal{S}_s^n$ and we find

$$\frac{d}{dt}\gamma|_{t=0} = \frac{d}{dt}|_{t=0} e^{iht}D_s e^{-iht}|_{t=0} = i[h, D_s].$$

In the canonical basis, these elements are of the form

$$\begin{pmatrix} 0 & iA & iB & \dots \\ iA^\dagger & 0 & iC & \\ iB^\dagger & iC^\dagger & 0 & \\ \vdots & & & \ddots \end{pmatrix}$$

Thus, by dimensional reasons, all elements of $T_{D_s}\mathcal{S}_s^n$ are of this form. Furthermore for $U \in U(n)$, observe that

$$U[h, D_s]U^\dagger = [UhU^\dagger, UD_sU^\dagger]$$

and thus

$$\begin{aligned} c_U : T_{D_s}\mathcal{S}_s^n &\rightarrow T_{UD_sU^\dagger}\mathcal{S}_s^n \\ v &\mapsto UvU^\dagger \end{aligned}$$

is an isomorphism. This proves the second claim. To prove the last claim, observe that for $U, V \in U(n)$

$$UD_sU^\dagger - VD_sV^\dagger = U(D_s - \lambda\mathbf{1})U^\dagger - V(D_s - \lambda\mathbf{1})V^\dagger$$

and similarly

$$[h, D_s] = [h, D_s] - [h, \lambda\mathbf{1}] = [h, D_s - \lambda\mathbf{1}].$$

Choosing λ to be the eigenvalue in s with the biggest multiplicity r , the expressions above are differences of rank $n - r$ matrices and thus maximally of rank $2(n - r)$. \square

Remark It is immediate that a POVM P , that is injective on the set of hermitian operators with rank smaller than r , \mathcal{P}_r , is an \mathcal{S}_s^n -embedding. This is because $\Delta\mathcal{P}_r = \mathcal{P}_{2r}$ and thus $\Delta\mathcal{S}_s^n \cup T\mathcal{S}_s^n = \mathcal{P}_{2r} = \Delta\mathcal{P}_r$ by IV.3.

As a consequence, the POVM constructed in [2] for states of bounded rank is also a \mathcal{S}_s^n -embedding and we obtain the following upper bounds.

Proposition IV.4. *Let m be the biggest multiplicity of an eigenvalue in the spectrum s and let $r := n - m$. Then, there is a POVM P of dimension $4r(n - r) - 1$ that is a \mathcal{S}_s^n -embedding.*

Remark Note that for $r = n/2$ the dimension of the POVM is $4n/2(n - n/2) - 1 = n^2 - 1$. Thus, it is the trivial POVM that can identify all states and hence we also get a \mathcal{S}_s^n -embedding for $r > n/2$.

The construction of the POVM is based on [22]. The idea is to use a totally non-singular matrix, like e.g. the Vandermonde-matrix, to construct a linear subspace of $M(\mathbb{C}, n)$ that just contains matrices of rank bigger than $2r$.

$l \setminus k$	2	3	4
5	22/34;39		
6	26/40;47		
7	30/50;55	48/76;83	
8	34/60;63	54/90;95	
9	38/66;71	60/98;107	84/134;143
10	42/72;79	66/110;119	92/148;159

TABLE I: Dimension/ Lower bounds on immersion dimension [21]; Upper bound on embedding dimension IV.4 for $U(l+k)/U(l) \times U(1)^k$.

As presented in the table I, these results are rather close to the lower bounds of [21]. Thus, the POVM of [2] gives good upper bounds on \mathcal{S}_s^n and in addition we have indirectly obtained good lower bounds on the dimension of a POVM that is complete with respect to the states of bounded rank.

B. States with Unitary Symmetry

Next, we shortly discuss subsets of states invariant under some unitary subgroup.

More precisely, we analyze the structure of the fix point sets of the action by conjugation G_H of some subgroup $H \subseteq U(n)$, i.e.

$$\mathcal{S}(\mathbb{C}^n)_H := \{\rho \in \mathcal{S}(\mathbb{C}^n) : U\rho U^\dagger = \rho, \forall U \in H\}.$$

Consider the sets $\mathcal{B}(\mathbb{C}^n)_H := \{B \in \mathcal{B}(\mathbb{C}^n) : UBU^\dagger = B, \forall U \in H\}$. $\mathcal{B}(\mathbb{C}^n)_H$ is a C^* algebra, since it is certainly a vector space and closed under the *-involution by the unitarity of H and thus the structure theorem [23] yields that $\mathcal{B}(\mathbb{C}^n)_H$ is unitarily equivalent to $\bigoplus_{i=1}^k M(n_i, \mathbb{C}) \otimes \mathbb{1}_{m_i}$.

Observe that the linear isomorphism

$$\begin{aligned} \iota : \mathcal{S} \left(\bigoplus_{i=1}^k M(n_i, \mathbb{C}) \right) &\rightarrow \bigoplus_{i=1}^k M(n_i, \mathbb{C}) \otimes \mathbb{1}_{m_i} \\ (\rho_1, \dots, \rho_k) &\mapsto \left(\frac{1}{m_1} \rho_1 \otimes \mathbb{1}_{m_1}, \dots, \frac{1}{m_k} \rho_k \otimes \mathbb{1}_{m_k} \right). \end{aligned} \tag{2}$$

descends to a diffeomorphism on states. Form this we immediately get the following proposition.

Proposition IV.5. *There is a POVM P with $\dim P = \dim \mathcal{S}(\mathbb{C}^n)_H$ that is stably $\mathcal{S}(\mathbb{C}^n)_H$ -complete.*

C. Unitarily Invariant States of Fixed Spectrum

Now, given some unitary subgroup $H \subseteq U(n)$, we want to identify G_H -invariant (c.f. IV B) states of fixed spectrum s^7 , i.e.

$$\mathcal{S}(\mathbb{C}^n)_{H,s} := \{\rho \in \mathcal{S}(\mathbb{C}^n) : U\rho U^\dagger = \rho, \forall U \in H, \text{spec}(\rho) = s\}.$$

Via the map ι defined in (2), there is natural action of $U_{n_1} \times \dots \times U_{n_k}$ on $B(\mathbb{C}^n)_H$ coming from its action on $\bigoplus_{i=1}^k M(n_i, \mathbb{C})$. $\mathcal{S}(\mathbb{C}^n)_H$ is then the orbit of this action on the set

$$\mathcal{D} := \{(D_{M_1}, \dots, D_{M_k}) : [(M_1)^{c_1} \cup \dots \cup (M_k)^{c_k}] = s\}.$$

Here M_i is a multiset of order n_i and $(M_i)^{c_i}$ is the union of c_i copies of M_i . By the same argument as in the previous section the orbit of some $\rho \in \mathcal{D}$ under G_H is diffeomorphic to a product of complex flag manifolds $\prod_{i=1}^k U(n_i) / \prod_{j=1}^{k_j} U(n_j^i)$. Since \mathcal{D} is clearly finite, $\mathcal{S}(\mathbb{C}^n)_{H,s}$ is a disjoint union of products of complex flag manifolds. Thus, it is enough to look at one of these components at a time to get non-immersion results.

For some component, let m_i be the number associated to the i -th factor in the product, that is constructed just like the number m for IV.2.

Proposition IV.6. *The product of complex flag manifolds $\prod_{i=1}^k U(n_i) / \prod_{j=1}^{k_j} U(n_j^i)$ cannot be immersed in Euclidean Space of dimension $\sum_{i=1}^k (4m_i(n - m_i) - 2\beta(n, m_i)) - 1$ and it cannot be embedded in Euclidean space of dimension $\sum_{i=1}^k 4m_i(n - m_i) - 2\beta(n, m_i)$.*

The proof of this result can be found in appendix A 5. Of course, IV.4 also transfers to this situation and gives upper bounds on the dimension of stably $\mathcal{S}(\mathbb{C}^n)_{H,s}$ -complete POVMs.

D. Bob-Unitary Orbit

Let $\alpha \in \mathcal{H}_A \otimes \mathcal{H}_B$, $\langle \alpha | \alpha \rangle = 1$. In this section we investigate pure states of the form

$$\mathcal{S}_B(\alpha) := \{|\beta\rangle\langle\beta| \in \mathcal{S}(\mathcal{H}_A \otimes \mathcal{H}_B) : \beta = (\mathbb{1} \otimes U)\alpha, U \in U(\mathcal{H}_B)\}.$$

Let $\{e_1, \dots, e_{\dim \mathcal{H}_A}\}$, $\{f_1, \dots, f_{\dim \mathcal{H}_B}\}$ be orthonormal bases of \mathcal{H}_A respectively \mathcal{H}_B such that

$$\alpha = \sum_{i=1}^r \alpha_i e_i \otimes f_i$$

⁷ Here s has to be compatible with the decomposition illustrated in section IV B.

is a Schmidt decomposition, where r is the Schmidt rank of α . Then, $\mathcal{S}_B(\alpha)$ is diffeomorphic to the projective Stiefel manifold $PW_{n,r}$. In order to see this, note that $(\mathbb{1} \otimes U(\mathcal{H}_B))\alpha$ is diffeomorphic to the complex Stiefel manifolds $W_{n,r} := \{m \in \mathbb{C}^{d \times r} : m^* \cdot m = \mathbb{1}\}$ via

$$i : (\mathbb{1} \otimes U(\mathcal{B}))\alpha \rightarrow W_{n,k},$$

$$M_{i,j} \left(\sum_{i=1}^r \alpha_i e_i \otimes U f_i \right) := \langle U f_i | f_j \rangle$$

Factoring both sides over the free action of the cyclic group $S^1 \subseteq \mathbb{C}$, $m \mapsto z \cdot m$, then yields the desired map [20].

In order to state the main result of this section we introduce two functions,

Definition IV.7. Let $n, k \in \mathbb{N}$ and $n \geq k$.

1. $N(n, k) := \min\{n - k < i \leq n : \binom{n}{i} \pmod{2} \equiv 1\}$,
2. $\sigma(n, k) := 2 \cdot \max\{0 \leq i < N(n, k) : \binom{nk + i - 1}{i} \pmod{2} \equiv 1\}$.

Proposition IV.8. Let $\alpha \in \mathcal{H}_A \otimes \mathcal{H}_B$, $\langle \alpha | \alpha \rangle = 1$, with Schmidt rank k and $n = \dim \mathcal{H}_B$. Then $\mathcal{S}_B(\alpha)$ cannot be immersed in Euclidean space of dimension $(2n - k)k - 1 + \sigma(n, k)$ and cannot be embedded in Euclidean space of dimension $(2n - k)k - 1 + \sigma(n, k) + 1$.

The proof of this result is very similar to [24] and can be found in appendix A 6. This non-immersion result is obtained deploying a standard approach based on the dual Stiefel-Whitney class of the tangent bundle [25].

For $k = 1$ the complex projective Stiefel manifold is just the complex Projective space, so in this case we can compare the result obtained here to the upper bounds of Milgram [26], which are known to be close to optimal. Table ?? shows these bounds for some dimensions. For $n = 2^k$, $k \in \mathbb{N}$, the dual Stiefel-Whitney classes give no obstructions, whereas they essentially equal Milgram's result in [26] for $n = 2^k + 1$, $k \in \mathbb{N}$, and hence are close to optimal in this case.

2	6	6	14	14	14	14	30	30	30	30	30	30	30	30	62
2	6	8	14	16	21	22	30	32	37	38	45	46	52	52	62

TABLE II: Lower bounds on immersions of $W_{n,1} \simeq PC^n$ for $n = 2, \dots, 17$. In the first row the result is obtained from the dual Stiefel-Whitney classes in the second row the results of [26] are presented.

In [27], it is shown that $PW_{n,n}$ and $PW_{n,n-1}$ is parallelizable for $n \neq 2$ and thus can be immersed in Euclidean space of codimension one by a result of Hirsch [28]. For $PW_{4,k}$ and $PW_{8,k}$, there are no obstructions because the dual Stiefel-Whitney classes vanish for $nk = q2^r$, $q, r \in \mathbb{Z}$ and $N(n, k) < 2^r$ [24].

The dual Stiefel-Whitney classes do not generally give good obstructions, but can be

n\r	2	3	4	5	6	7	8	9	10	11	12	13
2	2	3										
3	6	7	8									
4	6	11	14	15								
5	14	19	22	23	24							
6	14	27	30	31	34	35						
7	14	27	38	39	46	47	48					
8	14	27	38	47	54	59	62	63				
9	30	43	54	63	70	75	78	79	80			
10	30	51	54	63	86	91	94	95	98	99		
11	30	55	72	79	86	107	110	111	118	119	120	
12	30	59	78	79	102	107	126	127	134	139	142	143

TABLE III: Lower bounds on immersion dimension of $PW_{n,r}$ obtained from dual Stiefel-Whitney classes IV.8.

supplemented by other methods. In Table III these bounds are presented for some explicit scenarios. Another approach to the non-immersion problem is due to [29]. In a similar vein, another method is given to obtain non-immersion results, with the exterior powers γ_i of $KO(X)$ playing the role of the Stiefel-Whitney classes. Both of these methods are worked out and compared in [24].

Next, we give upper bounds on the dimension of an $\mathcal{S}_B(\alpha)$ -embedding, presenting two different approaches.

The first approach is based on the upper bounds obtained for states of fixed spectrum. The problem is split into determining the minor obtained by tracing over \mathcal{H}_A and afterwards determining the relative phases.

Before stating the upper bounds, let us first prove the following lemma which will be useful later on.

Lemma IV.9. *Let $\alpha := \sum_{i=1}^r \lambda_i e_i \otimes f_i$, $O \in H(\mathcal{H}_A)$, $S \in H(\mathcal{H}_B)$, $U \in U(\mathcal{H}_B)$ and $P_\alpha := \sum_{i=1}^r \lambda_i |e_i\rangle\langle f_i|$. Then $\text{tr}(O \otimes USU^\dagger|\alpha\rangle\langle\alpha|) = \text{tr}((P_\alpha U)^\dagger O^T (P_\alpha U) S)$.*

Proof. The prove of this is a straightforward computation.

$$\begin{aligned}
\text{tr}(O \otimes USU^\dagger|\alpha\rangle\langle\alpha|) &= \langle\alpha|O \otimes USU^\dagger|\alpha\rangle = \sum_{i,j=1}^r \lambda_i \lambda_j \langle e_i| \otimes \langle f_i|O \otimes USU^\dagger|e_j\rangle \otimes |f_j\rangle \\
&= \sum_{i,j=1}^r \lambda_i \lambda_j \langle e_i|O|e_j\rangle \langle f_i|USU^\dagger|f_j\rangle = \sum_{i,j=1}^r \lambda_j \langle e_j|O^T|e_i\rangle \lambda_i \langle f_i|USU^\dagger|f_j\rangle \\
&= \sum_{i,j=1}^r \langle f_i|f_j\rangle \lambda_j \langle e_j|O^T P_\alpha USU^\dagger|f_i\rangle = \sum_{i=1}^r \langle f_i|U^\dagger P_\alpha^\dagger O^T P_\alpha US|f_i\rangle \\
&= \text{tr}((P_\alpha U)^\dagger O^T (P_\alpha U) S)
\end{aligned}$$

□

The following proposition is motivated by a method to embed Lie groups in Euclidean space, introduced in [30].

Proposition IV.10. *Let $\alpha \in \mathcal{H}_A \otimes \mathcal{H}_B$ with Schmidt rank k and $n = \dim \mathcal{H}_B$. Then, there is a $\mathcal{S}_B(\alpha)$ -embedding of dimension $4r(n-r) - 1 + 4n - 5$ for $r < n/2$ and $n^2 - 1 + 4n - 5$ for $r \geq d/2$.*

Proof. First, note that by lemma IV.9 we can assume w.l.o.g that $\lambda_i \neq \lambda_j$ for $1 \leq i, j \leq r$, $i \neq j$, because this can always be achieved by choosing O appropriately.

The idea is to take advantage of the natural projection $\pi : PW_{n,r} \rightarrow \frac{U(n)}{U(1)^r \times U(n-r)}$, which just amounts to choosing $O = \mathbb{1}$ in lemma IV.9. More precisely, for $O = \mathbb{1}$ we get

$$P_\alpha O P_\alpha^\dagger = \begin{pmatrix} \lambda_1 & & & & \\ & \ddots & & & \\ & & \lambda_r & & \\ & & & 0 & \\ & & & & \ddots \end{pmatrix}$$

and thus we are in the situation discussed in the last section with the isotropy group given by $U(1)^r \times U(n-r)$. This means, the projected state can be embedded in dimension $4r(d-r) - 1$ using the POVM of IV.4. Let us call this map ϕ_1 .

Let $v \in \mathbb{C}^r$ be the vector with a one in every entry and consider the map

$$\begin{aligned} \psi : U(n)/U(n-r) &\rightarrow \mathbb{C}^n \\ U &\mapsto U(v \oplus 0). \end{aligned}$$

This is clearly well defined and also observe that ψ descends to a map $\tilde{\psi} : PW_{n,r} \simeq U(n)/(U(n-r) \times U(1)) \rightarrow P\mathbb{C}^n$. Let $U, V \in U(n)$ with $U \sim V$ in $U(n)/U(1)^r \times U(n-r)$, i.e. $U = V(D \oplus \mathbb{1})(\mathbb{1} \oplus W)$ with $D \in U(1)^r$ and $W \in U(n-r)$. Then, $Uv = \lambda Vv$ just has a solution for $U \sim V$ in $U(n)/(U(1) \times U(n-r))$. To see this, note that

$$\begin{aligned} U(v \oplus 0) &= \lambda V(v \oplus 0) \\ V(D \oplus \mathbb{1})(\mathbb{1} \oplus W)(v \oplus 0) &= \lambda V(v \oplus 0) \\ (D \oplus \mathbb{1})(v \oplus 0) &= \lambda(v \oplus 0) \\ Dv &= \lambda v \end{aligned}$$

and thus $D = \lambda \mathbb{1}$ is the only solution.

Hence, supplementing the embedding above by $\tilde{\psi}$ guarantees injectivity and the only problem left, is embedding $P\mathbb{C}^n$.

In terms of lemma IV.9 $\tilde{\psi}$ corresponds to choosing $P_\alpha O^T P_\alpha^\dagger = v^\dagger v \oplus 0$, the projective version of $v \oplus 0$. Then choose S according to the POVM of [2] to obtain an embedding in Euclidean space. Let us call this mapping ϕ_2

The map $\phi := (\phi_1, \phi_2)$ is clearly smooth as well as injective and thus a topological embedding by compactness of $\mathcal{S}_B(\alpha)$. From lemma IV.9 it is easy to see that $\frac{d}{dt}|_{t=0}(\mathbb{1} \otimes U e^{iHt})|\alpha\rangle\langle\alpha|(\mathbb{1} \otimes e^{-iHt}U^\dagger)$ for $H = h \oplus 0$ and $h \neq \mathbb{1}$ diagonal gives a $n-1$ dimensional

subspace V_U of the tangent space at $\alpha_U := (\mathbb{1} \otimes U)|\alpha\rangle\langle\alpha|(\mathbb{1} \otimes U^\dagger)$. V_U is clearly in the kernel of $d\phi_1|_{\alpha_U}$. Thus, by dimensional reasoning, it is enough to see that $d\phi_2|_{\alpha_U}$ is injective on V_U . Since the POVM of [2] can identify all rank one matrices, it is enough to see that $h \oplus 0 \mapsto [v \dagger v \oplus 0, h \oplus 0]$ is injective for $h \neq \mathbb{1}$ diagonal. This can be easily verified by a direct computation. \square

Remark This result is best for r close to n , so in particular for $PW_{n,n}$, the set of maximally entangled states. For $PW_{n,n}$ we obtain an embedding in Euclidean space of codimension $4n - 5$.

Furthermore, note that in the context of quantum process tomography, the result for maximally entangled states can be used to identify a unitary time evolution. Preparing a certain maximally entangled state, the POVM given above can identify unitary processes up to a phase, i.e. with $\mathcal{O}(n^2)$ measurements.

The second approach relies on the direct construction of an $\mathcal{S}_B(\alpha)$ -embedding.

Proposition IV.11. *Let $\alpha \in \mathcal{H}_A \otimes \mathcal{H}_B$ with Schmidt rank r and $n = \dim \mathcal{H}_B$. Then, there is a $\mathcal{S}_B(\alpha)$ -embedding of dimension $2nr + 2n - 3$.*

Proof. First, note that w.l.o.g we can assume $\lambda_1 = \dots = \lambda_r = 1$, as can be easily seen from lemma IV.9.

Then, for $O = |e_i\rangle\langle e_j|, S = |f_k\rangle\langle f_l|$ we obtain

$$\text{tr}((P_\alpha U)^\dagger O^T (P_\alpha U) S) = \langle f_i | U | f_k \rangle \langle f_j | U | f_l \rangle^*,$$

so from this point of view any linear combination of such products of elements of $P_\alpha U$ determines an operator, that need not be hermitian, and a set of such equations determines an operator system (here we think of $P_\alpha U$ as a matrix in the $\{e_i\}_{i \in \{1, \dots, r\}}, \{f_l\}_{l \in \{1, \dots, n\}}$ basis). It is worth noting that an equation not corresponding to a non-hermitian operator actually corresponds to two operators in the operator system, namely its hermitian and anti-hermitian part.

Let $M_{n(i-1)+j}(U) := \langle e_i | P_\alpha U | f_j \rangle$ for $i \in \{1, \dots, r\}, j \in \{1, \dots, n\}$ and $M_k := 0$ for $k > nr$. For $k \in \{1, \dots, nr + n - 1\}$, define operators \tilde{G}_k via the equations

$$G_k(U) := \sum_{i=1, i \leq k+1-i}^n M_i(U) M_{k+1-i}^*(U).$$

Then, the operator system σ_G spanned by the \tilde{G}_k is an $\mathcal{S}_B(\alpha)$ -embedding. It is clear that the dimension of σ_G is $2nr + 2n - 3$, noting that non-hermitian operators count twice.

Let $U, V \in U(\mathcal{H}_B)$. In order to prove injectivity, we have to show that if $G_k(U) = G_k(V)$, then there is a $\phi \in \mathbb{R}$ such that $P_\alpha U = e^{i\phi} P_\alpha V$. First, observe that for $M_1(U) = \dots = M_k(U) = 0$ we have $k \leq n$ because $P_\alpha U$ has full rank. Let m be the smallest number such that M_m does not vanish. Then the claim is clearly true for all $j < m$. Now, let

$l > m$ and assume that the claim holds for all $j \leq l$, then

$$\begin{aligned}
G_{m+l}(U) &= \sum_{i=1, i \leq m+l+1-i}^n M_i(U) M_{m+l+1-i}^*(U) \\
&= M_m(U) M_{l+1}(U)^* + \sum_{i=m+1, i \leq m+l+1-i}^n M_i(U) M_{m+l+1-i}^*(U) \\
&= M_m(V) e^{i\phi} M_{l+1}(U)^* + \sum_{i=m+1, i \leq m+l+1-i}^n M_i(V) M_{m+l+1-i}^*(V) \\
&= G_{m+l}(V),
\end{aligned}$$

thus $e^{-i\phi} M_{l+1}(U) = M_{l+1}(V)$.

To conclude the proof, we need to show that the measurement constructed above is an immersion. For $h \in H(\mathcal{H}_B)$, $U \in U(\mathcal{H}_B)$ and define a curve $\gamma(t) := (\mathbf{1} \otimes e^{iUhU^\dagger t})|\alpha\rangle\langle\alpha|(\mathbf{1} \otimes U^\dagger e^{-iUhU^\dagger t})$. The derivative of this curve yields tangent vectors at $(\mathbf{1} \otimes U)|\alpha\rangle\langle\alpha|(\mathbf{1} \otimes U^\dagger)$ and by lemma IV.9 an effect operator $O \otimes S$ maps these to

$$\begin{aligned}
&\frac{d}{dt}\Big|_{t=0} \text{tr}((P_\alpha e^{iUhU^\dagger t} U)^\dagger O^T (P_\alpha e^{-iUhU^\dagger t} U) S) \\
&= i \text{tr}(P_\alpha^\dagger O^T P_\alpha U[h, S] U^\dagger).
\end{aligned}$$

For $k \in \{1, \dots, nr + n - 1\}$, this yields the equations

$$F_k(U, h) = \sum_{i=1, i \leq m+l+1-i}^n M_i(U) M_{k+1-i}^*(Uh) - M_i(Uh) M_{k+1-i}^*(U).$$

Observe that $F_k(U, h) = F_k(U, h + c\mathbf{1})$ holds for each $c \in \mathbb{C}$. Furthermore, it is easy to see that for every $k \in \{1, \dots, nr\}$ and every $h \in H(\mathcal{H}_B)$ there is a $\lambda \in \mathbb{C}$ such that $M_k(U(\lambda\mathbf{1} + h)) = 0$ and thus we can assume w.l.o.g that $M_k(Uh) = 0$.

Let m be the smallest number such that $M_m(U)$ does not vanish and assume w.l.o.g $M_m(Uh) = 0$. Let $l \in \{1, \dots, nr + m - 1\}$. It is easy to see that the vanishing of these equations for all $i \leq l$ implies that $M_j(Uh) = 0$ for $j \leq l + 1 - k$ and thus, we obtain injectivity on a real vector space of dimension $2nr - r^2 - 1$. \square

In table IV both of these methods are compared. It is clear that the embedding in IV.11 works best for $k/m \ll 1$, because this approach does not take the orthogonality of the f_i into account. The embedding in IV.10 works best for $k/m \sim 1$, because just in this case the projected state can be determined efficiently.

Appendix A: Technical appendix

1. Proof of Theorem III.3

Before we give the proof, let us first fix some notion.

$n \setminus r$	5	9	17	65
5	24/24;40;57			
9	64/70;112;105	80/80;112;177		
17	144/166;302;201	224/238;352;337	288/288;352;609	
65	624/742;1454;777	1088/1198;2270;1297	1920/2014;3518;2337	4224/4224;4480;8577
129	1264/1510;2990;1545	2240/2478;4830;2577	4096/4318;8126;4641	12544/12670;17152;17025

TABLE IV: Dimension/Lower bounds on immersion dimension IV.8; First upper bound on embedding dimension IV.10; Second upper bound on embedding dimension IV.11 for $PW_{n,r}$.

Let $\mathcal{SO}(H(\mathcal{H}))$ be the orthogonal group on the inner product space $H(\mathcal{H})$. The generalized Pauli basis together with the identity $\mathbb{1}_{\mathcal{H}}$ gives an identification of $H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$ and the Hilbert-Schmidt inner product induces the standard inner product on \mathbb{R}^{d^2} . Thus $\mathcal{SO}(H(\mathcal{H}))$ can be identified with $\mathcal{SO}(\mathbb{R}^{d^2})$, the standard orthogonal group on \mathbb{R}^{d^2} . Denote by $\mathcal{SO}(H(\mathcal{H}))_v := \{O \in \mathcal{SO}(H(\mathcal{H})) : Ov = v\}$ the stabilizer subgroup of $v \in H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$. Note that for $O \in \mathcal{SO}(H(\mathcal{H}))_{\mathbb{1}_{\mathcal{H}}}$ and $\sigma \in \Sigma(n)$, we have $O\sigma \in \Sigma(n)$. Thus there is an action of $\mathcal{SO}(H(\mathcal{H}))_{\mathbb{1}_{\mathcal{H}}}$ on $\Sigma(n)$,

$$\begin{aligned} \Sigma(n) \times \mathcal{SO}(H(\mathcal{H}))_{\mathbb{1}_{\mathcal{H}}} &\rightarrow \Sigma(n) \times \Sigma(n) \\ (\sigma, O) &\mapsto (\sigma, O\sigma). \end{aligned}$$

The geometric intuition of thinking of operator systems as planes in $H(\mathcal{H}) \simeq \mathbb{R}^{d^2}$ is essential for the following proof. Then, for $O \in \mathcal{SO}(H(\mathcal{H}))_{\mathbb{1}_{\mathcal{H}}}$, $O\sigma$ is just a rotated plane and the intuition is that for small rotations these operator systems are close.

Proof. Let $\sigma \in \Sigma(n)$ be stably \mathcal{P} -complete and π_σ be the associated orthogonal projection. Furthermore let $SH(\mathcal{H}) := \{B \in H(\mathcal{H}) : \|B\| = 1\}$ be the unit sphere in $H(\mathcal{H})$. Assume by contradiction that σ is not a \mathcal{P} -embedding, i.e. π_σ is not an immersion. Then, since $d(\pi_\sigma)_\rho = \pi_\sigma|_{T_\rho \mathcal{P}}$, there exists a point $\rho \in \mathcal{P}$ and a smooth curve $\gamma : (-1, 1) \rightarrow \mathcal{P}$ with $\gamma(0) = \rho$ and $v = \dot{\gamma}(0) \in \sigma^\perp$, $v \in SH(\mathcal{H})$. The idea is that $\gamma(t) \approx \rho + vt$ for small t and to obtain a contradiction we construct for a point $\rho' = \gamma(t') \approx \rho + vt'$ an operator system σ' with $\pi_{\sigma'}(\rho' - \rho) = 0$. This procedure is presented in figure 2.

More precisely, we prove that for each $\delta > 0$ there is a $t \in (0, 1)$ such that $\gamma(t) \in V_\rho^\delta := \{\rho + \lambda \cdot Ov : \lambda > 0, O \in \mathcal{SO}(H(\mathcal{H})), \|\mathbb{1} - O\|_{op} < \delta\}$.

First, we prove that V_ρ^δ is open. Note that the left action

$$\begin{aligned} \mathcal{SO}(H(\mathcal{H})) \times SH(\mathcal{H}) &\rightarrow SH(\mathcal{H}) \times SH(\mathcal{H}) \\ (O, v) &\mapsto (Ov, v) \end{aligned}$$

is smooth and transitive. Thus, the orbit map $\beta_v : \mathcal{SO}(H(\mathcal{H})) \rightarrow SH(\mathcal{H})$, $O \mapsto Ov$ is smooth and factors over the natural projection $\pi : \mathcal{SO}(H(\mathcal{H})) \rightarrow \mathcal{SO}(H(\mathcal{H}))/\mathcal{SO}(H(\mathcal{H}))_v$




FIG. 2: This figure shows the curve γ with $\gamma(0) = \rho$, $\gamma(t') = \rho'$ and $\frac{d}{dt}\gamma(0) = v \in \sigma^\perp$ together with the operator system σ' that is constructed such that $\rho' - \rho \in (\sigma')^\perp$.

(theorem 3.62 of [20]), i.e. $\beta_v = \tilde{\beta}_v \circ \pi$ with $\tilde{\beta}_v$ a diffeomorphism. In particular β_v is open because π is open⁸.

Since β_v is open there is an $\eta > 0$ such that $\emptyset \neq B_{\rho+v}(\eta) \cap B_\rho(1) \subseteq V_\rho^\delta$. By possibly shrinking η we can even assume that $\overline{B_{\rho+v}(\eta)} \subseteq V_\rho^\delta$ because of the conic structure of V_ρ^δ . It follows that $B_{\rho+sv}(s\eta) \subseteq V_\rho^\delta$ for $s > 0$.

Then,

$$\frac{\|\gamma(t) - (\rho + tv)\|}{t} = \frac{\|\gamma(t) - \gamma(0) - tv\|}{t} \rightarrow 0 \text{ as } t \rightarrow 0,$$

whereas we find for the distance $d(\rho + tv, \partial B_{\rho+tv}(t\eta))/t := \inf_{\rho' \in \partial B_{\rho+tv}(t\eta)} \|\rho - \rho'\|/t = \eta$. So by continuity of the norm there is a $t > 0$ such that $\gamma((0, t)) \subseteq V_\rho^\delta$.

But then, for every $\delta > 0$, there is an $O \in \mathcal{SO}(H(\mathcal{H}))$, a $\lambda > 0$ and a $t > 0$ with

$$\begin{aligned} d(\sigma, O\sigma) &= \|\pi_\sigma - \pi_{O\sigma}\|_{op} \\ &= \|\pi_\sigma - O\pi_\sigma O^{-1}\|_{op} \\ &\leq \|\pi_\sigma - \pi_\sigma O^{-1}\|_{op} + \|\pi_\sigma - O\pi_\sigma\|_{op} \\ &\leq 2\|\mathbb{1} - O\|_{op} < 2\delta \end{aligned}$$

such that $\gamma(t) - \gamma(0) = \lambda O v \neq 0$. But then, $\pi_{O\sigma}(\gamma(t) - \gamma(0)) = O\pi_\sigma(O^{-1}(\gamma(t) - \gamma(0))) = \lambda O\pi_\sigma(v) = 0$ by assumption on v . Also note that $\langle \mathbb{1}_{\mathcal{H}}, \gamma(t) - \gamma(0) \rangle := \text{tr}[\mathbb{1}_{\mathcal{H}}(\gamma(t) - \gamma(0))] = 0$ and $\langle \mathbb{1}, T_\rho \mathcal{P} \rangle = 0$ and thus we can choose $O \in \mathcal{SO}(H(\mathcal{H}))_{\mathbb{1}_{\mathcal{H}}}$. So $O\sigma$ is an operator system but it is not \mathcal{P} -complete, contradicting the stability of σ .

Conversely, suppose σ is a \mathcal{P} -embedding. A.2 states, that there is an $\epsilon > 0$ such that every $\sigma' \in \Sigma(n)$ with $\sup_{B \in H(\mathcal{H}), \|B\| \leq 1} \|\pi_\sigma(B) - \pi_{\sigma'}(B)\| < \epsilon$ is a \mathcal{P} -embedding and thus in particular \mathcal{P} -complete.. \square

⁸ For an open set $A \subseteq \mathcal{SO}(H(\mathcal{H}))$ we find $\pi^{-1}(\pi(A)) = \mathcal{SO}(H(\mathcal{H}))_v \cdot A = \bigcup_{O \in \mathcal{SO}(H(\mathcal{H}))_v} O(A)$. So $\pi^{-1}(\pi(A))$ is open for any open set $A \subseteq \mathcal{SO}(H(\mathcal{H}))$ and thus π is open.

2. Proof of Lemma III.5

The following proof uses geometric concepts and is based on the identification of the tangent spaces with planes in $H(\mathcal{H})$.

Proof. π^T is smooth, as can be easily seen in local coordinates. The mapping

$$\begin{aligned} \psi : \mathcal{P} \times \mathcal{P} &\rightarrow \mathbb{R} \\ (\rho', \rho) &\mapsto \|\pi_{\rho'}^T - \pi_{\rho}^T\|_{op} \end{aligned}$$

is clearly continuous as a composition of continuous mappings and thus, for every $\eta > 0$, there is an open neighbourhood N_{ρ_0} of $\rho_0 \in \mathcal{P}$ such that $\psi(\rho, \rho_0) < \eta/4$ for all $\rho \in N_{\rho_0}$. Let $\nu_0 > 0$ and let $B_{5\nu_0}(\rho_0)$ be the open ball of radius $5\nu_0$ around ρ_0 , such that $B_{5\nu_0}(\rho_0) \cap \mathcal{P}$ is contained in N_{ρ_0} .

Let $\rho \in B_{\nu_0}(\rho_0) \cap \mathcal{P}$. Then, for all $\tilde{\rho} \in B_{4\nu_0}(\rho) \cap \mathcal{P}$, we find

$$\psi(\rho, \tilde{\rho}) < \psi(\rho, \rho_0) + \psi(\tilde{\rho}, \rho_0) < \eta/2. \quad (\text{A1})$$

Let $\rho' \in \partial B_{\epsilon}(\rho) \cap \mathcal{P}$, $0 < \epsilon < \nu_0$. Furthermore, let $\gamma : [0, \lambda] \rightarrow \mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a geodesic that connects ρ and ρ' with $\gamma(0) = \rho$ and $\frac{d}{dt}\gamma(t)|_{t=0} = v$, $\|v\| = 1$. Since $(\rho, v) \mapsto \frac{d^2}{dt^2}\gamma(t)|_{t=0} = \frac{d^2}{dt^2}\exp(\rho, vt)|_{t=0}$ is a smooth function from the compact set $\mathcal{P} \times S^{\dim \mathcal{P}-1}$ to $H(\mathcal{H})$, there is $k \geq 0$ such that $k := \max_{(\rho, v) \in \mathcal{P} \times S^{\dim \mathcal{P}-1}} \|\frac{d^2}{dt^2}\exp(\rho, vt)|_{t=0}\|$. It follows from the geodesic equation

$$\begin{aligned} \pi_{\gamma(t)}^T \left(\frac{d^2}{dt^2}\gamma(t) \right) &= 0 \\ \pi_{\gamma(0)}^T \left(\frac{d^2}{dt^2}\gamma(t) \right) + (\pi_{\gamma(t)}^T - \pi_{\gamma(0)}^T) \left(\frac{d^2}{dt^2}\gamma(t) \right) &= 0 \\ \pi_{\gamma(0)}^T \left(\frac{d}{dt}\gamma(t) \right) + \int_0^t dt' (\pi_{\gamma(t')}^T - \pi_{\gamma(0)}^T) \left(\frac{d^2}{dt'^2}\gamma(t') \right) &= v \\ \pi_{\gamma(0)}^T (\gamma(t)) + \int_0^t dt' \int_0^{t'} dt'' (\pi_{\gamma(t'')}^T - \pi_{\gamma(0)}^T) \left(\frac{d^2}{dt''^2}\gamma(t'') \right) &= vt + \pi_{\gamma(0)}^T(\rho). \end{aligned}$$

However, for $t \in [0, 4\nu_0]$,

$$\begin{aligned} &\left\| \int_0^t dt' (\pi_{\gamma(t')}^T - \pi_{\gamma(0)}^T) \left(\frac{d^2}{dt'^2}\gamma(t') \right) \right\| \\ &\leq \int_0^t dt' \left\| (\pi_{\gamma(t')}^T - \pi_{\gamma(0)}^T) \left(\frac{d^2}{dt'^2}\gamma(t') \right) \right\| \\ &< \int_0^t dt' k\eta/2 = kt\eta/2 \leq 2\eta k\nu_0. \end{aligned}$$

as well as

$$\begin{aligned} & \left\| \int_0^t dt' \int_0^{t'} dt'' (\pi_{\gamma(t'')}^T - \pi_{\gamma(0)}^T) \left(\frac{d^2}{dt''^2} \gamma(t'') \right) \right\| \\ & \leq \int_0^t dt' \int_0^{t'} dt'' \left\| (\pi_{\gamma(t'')}^T - \pi_{\gamma(0)}^T) \left(\frac{d^2}{dt''^2} \gamma(t'') \right) \right\| \\ & < \int_0^t dt' \int_0^{t'} dt'' k\eta/2 = kt^2\eta/4 \leq 4\eta k(\nu_0)^2. \end{aligned}$$

Thus, we find

$$\left\| \pi_{\gamma(0)}^T (\gamma(t) - (vt + \rho)) \right\| < kt^2\eta/4 < 4\eta k(\nu_0)^2$$

and

$$\begin{aligned} \left\| \frac{d}{dt} \gamma(t) - v \right\| &= \left\| \pi_{\gamma(t)}^T \left(\frac{d}{dt} \gamma(t) \right) - \pi_{\gamma(0)}^T \left(\frac{d}{dt} \gamma(t) \right) + \pi_{\gamma(0)}^T \left(\frac{d}{dt} \gamma(t) \right) - v \right\| \\ &\leq \left\| \pi_{\gamma(t)}^T \left(\frac{d}{dt} \gamma(t) \right) - \pi_{\gamma(0)}^T \left(\frac{d}{dt} \gamma(t) \right) \right\| + \left\| \pi_{\gamma(0)}^T \left(\frac{d}{dt} \gamma(t) \right) - v \right\| \\ &\leq \eta/2 + 2\eta k\nu_0, \end{aligned}$$

Note that γ stays inside $B_{5\nu_0}(\rho_0)$. Furthermore, for η and ν_0 small enough, γ intersects $\partial B_{2\nu_0}(\rho_0)$ and γ intersects $\partial B_\epsilon(\rho)$ close to radial. In particular it follows that $t \mapsto \|\gamma(t) - \rho\|$ is strictly increasing as long as γ stays inside $B_{2\nu_0}(\rho_0)$. Then, each geodesic intersects $\partial B_\epsilon(\rho)$ exactly once before it passes through $\partial B_{2\nu_0}(\rho_0)$ for each ϵ with $0 < \epsilon < \nu_0$. Let K_ρ be the connected component of $\mathcal{P} \cap B_{\nu_0}(\rho)$ containing ρ and let K_{ρ_0} be the connected component of $\mathcal{P} \cap B_{2\nu_0}(\rho_0)$ containing ρ_0 . The above reasoning implies that $K_\rho \cap B_\epsilon(\rho)$ is connected and that $K_{\rho_0} \cap B_{\tilde{\epsilon}}(\rho_0)$ is connected for $0 < \tilde{\epsilon} < 2\nu_0$.

We can assume w.l.o.g. that $B_{\nu_0}(\rho) \cap \mathcal{P}$ is connected for all $\rho \in B_{\nu_0}(\rho_0)$. Because if not, we can shrink ν_0 until $B_{2\nu_0}(\rho_0)$ contains a single connected component. To see this, shrink ν_0 such that $0 < \nu_0 < \max_{\rho \in K_{\rho_0}, \rho' \in \overline{B_{2\nu_0}(\rho_0)} \cap \mathcal{P} - K_{\rho_0}} \|\rho - \rho'\|$.

We then find for $0 < \epsilon < \nu_0$ and $t \in [0, 4\epsilon]$,

$$\begin{aligned} \left\| \pi_{\gamma(0)}^T (\gamma(t) - \rho) \right\| &= \left\| \pi_{\gamma(0)}^T (\gamma(t) - (vt + \rho) + vt) \right\| \\ &> \|vt\| - \left\| \pi_{\gamma(0)}^T (\gamma(t) - (vt + \rho)) \right\| > t - 4\eta k\epsilon^2. \end{aligned}$$

Hence, $\gamma(t) = \rho'$ and $\gamma([0, t]) \subseteq B_{2\nu_0}(\rho_0) \Rightarrow t < \epsilon + 4\eta k\epsilon^2$.

Choosing ν_0 such that $4k\nu_0 < 1$, we find for the component of $\rho' - \rho = \gamma(t) - \gamma(0)$ normal to $T_\rho \mathcal{P}$,

$$\begin{aligned} \left\| \pi_\rho^N (\rho' - \rho) \right\| &= \left\| \int_0^t dt' \frac{d}{dt'} \gamma(t') - \pi_{\gamma(0)}^T \left(\frac{d}{dt'} \gamma(t') \right) \right\| \\ &\leq \int_0^t dt' \left\| \pi_{\gamma(t')}^T \left(\frac{d}{dt'} \gamma(t') \right) - \pi_{\gamma(0)}^T \left(\frac{d}{dt'} \gamma(t') \right) \right\| \\ &< \int_0^t dt' \eta/2 = t\eta/2 < \eta(\epsilon + 4k\epsilon^2)/2 \leq \epsilon\eta(1 + 4k\nu_0)/2 < \eta\epsilon = \eta\|\rho - \rho'\|. \end{aligned}$$

Now, for a given $\eta > 0$, construct such neighbourhoods for all $\rho \in \mathcal{P}$ to obtain a cover of \mathcal{P} by open sets, $\{B_{\nu_\rho}(\rho) \cap \mathcal{P}\}_{\rho \in \mathcal{P}}$. By compactness of \mathcal{P} there is a finite subcover $\{B_{\nu_{\rho_i}}(\rho_i) \cap \mathcal{P}\}_{i \in I}$ and set $\epsilon := \min_{i \in I} \nu_{\rho_i}$. \square

Remark Note that the proof of this lemma shows that for $0 < \tilde{\epsilon} < \epsilon$, $B_{\tilde{\epsilon}}(\rho) \cap \mathcal{P}$ is connected.

3. Proof of Proposition III.10

Proof. To see that i is smooth, choose an orthonormal basis $\{\sigma_i\}_{i \in I}$ of hermitian operators for $H(\mathcal{H})$. Expansion in this basis gives global coordinates on $H(\mathcal{H})$. Expansion in $\{\sigma_{i_1} \otimes \dots \otimes \sigma_{i_k}\}_{i_1, \dots, i_k \in I}$ gives global coordinates on $H(\mathcal{H})^{\otimes k}$. In these coordinates i is just a polynomial and hence smooth.

\mathcal{P} is a smooth submanifold and since $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H}) \subseteq H(\mathcal{H})$, $i|_{\mathcal{P}}$ is smooth. We prove that $i|_{\mathcal{P}}$ is injective. Note that $\rho^{\otimes k} = \sigma^{\otimes k}$ iff $\sigma = a \cdot \rho$, $a^k = 1$. But then $\sigma, \rho \in H(\mathcal{H})$ implies $a \in \{-1, 1\}$ and the positivity of both σ and ρ yields $a = 1$.

Finally, $i|_{\mathcal{P}}$ is an immersion. To see this let $\rho \in \mathcal{P}$ and $v \in T_\rho \mathcal{P}$. Furthermore, let $\gamma : (-1, 1) \rightarrow H(\mathcal{H})$ be a smooth curve with $\gamma(0) = \rho$ and $\frac{d}{dt}\gamma(0) = v$. First, observe that for $k = 2$,

$$\begin{aligned} di_\rho v &= \frac{d}{dt}\Big|_{t=0} (i \circ \gamma) = \lim_{t \rightarrow 0} \frac{\gamma(t)^{\otimes 2} - \gamma(0)^{\otimes 2}}{t} \\ &= \lim_{t \rightarrow 0} \frac{(\gamma(t) - \gamma(0)) \otimes \gamma(t) - (\gamma(0) - \gamma(t)) \otimes \gamma(0)}{t} = \rho \otimes v + v \otimes \rho. \end{aligned}$$

This inductively generalizes to arbitrary $k \in \mathbb{N}$ and we get

$$di_\rho v = \sum_{i=1}^k \rho^{\otimes i-1} \otimes v \otimes \rho^{\otimes k-i}.$$

This is zero if and only if $v = 0$, what can be easily seen by orthogonally decomposing v with respect to ρ .

Finally, since $i|_{\mathcal{P}}$ is smooth, injective and an immersion, it is a smooth embedding by the compactness of $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$. \square

Remark Note that the proof shows that $i|_{H(\mathcal{H}) - \{0\}}$ is an immersion.

4. Proof of Lemma III.13

In this lemma we prove that, for a submanifold $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$, every smooth embedding $\psi : \mathcal{P} \rightarrow \mathbb{R}^m$ can be approximated by a polynomial map $F : \mathbf{1}_{\mathcal{H}} + H_0(\mathcal{H}) \simeq \mathbb{R}^{d^2-1} \rightarrow \mathbb{R}^m$. Let us first state lemma 1.3 of [31].

Lemma A.1. [31] Let $U \subseteq \mathbb{R}^n$ be open and $W \subseteq U$ be open with compact closure $\overline{W} \subseteq U$. Let $f : U \rightarrow \mathbb{R}^n$ be a smooth embedding. There exists $\epsilon > 0$ such that if $g : U \rightarrow \mathbb{R}^n$ is smooth and

$$\|D_\alpha g(x) - D_\alpha f(x)\|_2 < \epsilon \quad \text{and} \quad \|g(x) - f(x)\|_2 < \epsilon$$

for all $x \in W$, $|\alpha| = 1$, then $g|_W$ is an embedding.

Now we give the proof of lemma III.13.

Proof. Note that $\psi' = \tilde{\psi}'|_{\mathcal{P}}$ is smooth because it is a restriction of smooth functions to a smooth submanifold. ψ can be extended to a compactly supported smooth map $\tilde{\psi}$ on $\mathbb{1}_{\mathcal{H}} + H(\mathcal{H})_0 \simeq \mathbb{R}^{n^2-1}$ and let $K \subseteq \mathbb{R}^{n^2-1}$ be a compact set containing $\text{supp } \tilde{\psi}$. In the following we make use of an approximation result given by theorem 1 in [32]. The relevant part for us is that for every $\eta > 0$, there is a $k \in \mathbb{N}$ such that $\tilde{\psi}$ and $d\tilde{\psi}$ can be approximated simultaneously by a map $\tilde{\psi}' = (p_1, \dots, p_i)$, $p_i \in P^{\leq k}(\mathbb{R}^{n \times n-1})$, i.e $\sup_{x \in K} \|\tilde{\psi}(x) - \tilde{\psi}'(x)\|_2 < \eta$ and $\sup_{(x,v) \in TK, \|v\| \leq 1} \|d\tilde{\psi}_x(v) - d\tilde{\psi}'_x(v)\|_2 < \eta$.

Let $\{(\phi_i, W_i)\}_{i \in I}$ be a finite atlas on \mathcal{P} and let $\tilde{B}(r_i) := \phi_i(B(r_i)) \subseteq W_i$ be the image of an open ball of radius r_i around the origin such that $\bigcup_{i \in I} \tilde{B}(r_i/2) = \mathcal{P}$ and $\overline{\tilde{B}(r_i)} \subseteq W_i$. Applying lemma 1.3 of [31] to $\psi \circ \phi_i$, $\phi_i^{-1}(W_i)$ and $B(r_i)$, we obtain for each $i \in I$ an $\epsilon_i > 0$ such that for all ψ' with $\sup_{B(r_i)} \|D_\alpha(\psi \circ \phi_i) - D_\alpha(\psi' \circ \phi_i)\|_2 < \epsilon_i$, $|\alpha| \leq 1$, $\psi'|_{\tilde{B}(r_i)}$ is an embedding.

For $|\alpha| = 1$, we have

$$\begin{aligned} & \sup_{x \in B(r_i)} \|D_\alpha(\psi \circ \phi_i)(x) - D_\alpha(\psi' \circ \phi_i)(x)\|_2 \\ &= \sup_{x \in B(r_i)} \|(d\psi_{\phi(x)} - d\psi'_{\phi(x)}) \circ D_\alpha \phi_i(x)\|_2 \\ &\leq \sup_{(x,v) \in T\tilde{B}(r_i), \|v\| \leq 1} \|(d\psi_x - d\psi'_x)v\|_2 \sup_{B(r_i)} \|D_\alpha \phi_i\|_2 \\ &\leq \sup_{(x,v) \in TK, \|v\| \leq 1} \|(d\tilde{\psi}_x - d\tilde{\psi}'_x)v\|_2 \sup_{B(r_i)} \|D_\alpha \tilde{\phi}_i\|_2. \end{aligned}$$

For $|\alpha| = 1$, let $\kappa_{i,\alpha} := \epsilon_i / \sup_{B(r_i)} \|D_\alpha \phi_i\|_2$ and let $\epsilon := \min_{i, |\alpha|=1} \{\epsilon_i, \kappa_{i,\alpha}\}$. Then, for every $\tilde{\psi}'$ with $\sup_{x \in K} \|\tilde{\psi}(x) - \tilde{\psi}'(x)\|_2 < \epsilon$ and $\sup_{(x,v) \in TK, \|v\| \leq 1} \|d\tilde{\psi}_x(v) - d\tilde{\psi}'_x(v)\|_2 < \epsilon$, $\tilde{\psi}'|_{\tilde{B}(r_i)}$ is an embedding for all $i \in I$ and by theorem 1 of [32] such a $\tilde{\psi}'$ exists for some $k \in \mathbb{N}$.

Finally we show that there is an $\epsilon \geq 0$ such that for every $\tilde{\psi}'$ with $\|D_\alpha \tilde{\psi} - D_\alpha \tilde{\psi}'\| < \epsilon$, $\psi'|_{\mathcal{P}}$ is injective. Then ψ' is both an immersion and injective and thus a smooth embedding by the compactness of \mathcal{P} .

$\tilde{B}(r_i)^c := \mathcal{P} - \tilde{B}(r_i)$ and $\overline{\tilde{B}(r_i/2)}$ are closed and therefore compact as closed subsets of a compact set. Then, by the continuity of the norm, $\eta_i := \min_{q \in \tilde{B}(r_i)^c, p \in \overline{\tilde{B}(r_i/2)}} \|\psi(p) - \psi(q)\|$ exists and it is bigger than 0 because ψ is injective and $\tilde{B}(r_i)^c \cap \tilde{B}(r_i/2) = \emptyset$. By possibly shrinking ϵ , make sure that $\epsilon \leq \frac{1}{4} \min_{i \in I} \eta_i$.

Assume $\psi'(p) = \psi'(q)$, $p, q \in \mathcal{P}$. Since $\psi'|_{\mathcal{P}}$ is an embedding around p , there is an $i \in I$ such that $p \in \tilde{B}(r_i/2)$ and $q \in \tilde{B}(r_i)^c$. Thus,

$$\begin{aligned} \|\psi'(p) - \psi'(q)\|_2 &= \|\psi(p) - \psi(q) + \psi'(p) - \psi(p) + \psi(q) - \psi'(q)\|_2 \\ &\geq \|\psi(p) - \psi(q)\|_2 - \|\tilde{\psi}'(p) - \tilde{\psi}'(q) + \tilde{\psi}(q) - \tilde{\psi}'(q)\|_2 \\ &\geq 4\epsilon - 2\epsilon > 0, \end{aligned}$$

a contradiction. \square

A direct consequence of this proof is the following corollary, which was used in the third section.

Corollary A.2. *Let $\mathcal{P} \subseteq \mathcal{S}(\mathcal{H})$ be a submanifold and let $L : H(\mathcal{H}) \rightarrow \mathbb{R}^m$ be a linear map such that $L|_{\mathcal{P}}$ is a smooth embedding. Then, there is an $\epsilon > 0$ such that for every linear map $L' : H(\mathcal{H}) \rightarrow \mathbb{R}^m$ with $\sup_{v \in H(\mathcal{H}), \|v\| \leq 1} \|(L - L')v\|_2 < \epsilon$, $L'|_{\mathcal{P}}$ is a smooth embedding.*

Proof. Let $K \subseteq H(\mathcal{H})$ be a compact set containing \mathcal{P} . Furthermore let $b : H(\mathcal{H}) \rightarrow H(\mathcal{H})$ be a smooth and compactly supported bump function which equals the identity on K . Then, the proof of III.13 shows that there is an $\eta > 0$ such that for every smooth map $\psi : H(\mathcal{H}) \rightarrow \mathbb{R}^m$ with $\sup_{x \in K} \|\psi(x) - (L \circ b)(x)\|_2 < \epsilon$ and $\sup_{(x,v) \in TK, \|v\| \leq 1} \|d\psi_x(v) - d(L \circ b)_x(v)\|_2 < \eta$, $\psi|_{\mathcal{P}}$ is a smooth embedding. But for ψ linear we find

$$\begin{aligned} &\sup_{(x,v) \in TK, \|v\| \leq 1} \|d\psi_x(v) - d(L \circ b)_x(v)\|_2 \\ &= \sup_{(x,v) \in TK, \|v\| \leq 1} \|\psi(v) - L(v)\|_2 \\ &= \sup_{v \in H(\mathcal{H}), \|v\| \leq 1} \|\psi(v) - L(v)\|_2 \end{aligned}$$

and

$$\begin{aligned} &\sup_{x \in K} \|\psi(x) - (L \circ b)(x)\|_2 \\ &\leq \sup_{x \in K} \|x\| \sup_{v \in H(\mathcal{H}), \|v\| \leq 1} \|\psi(v) - L(v)\|_2. \end{aligned}$$

Thus, the claim holds for $\epsilon := \eta / \sup_{x \in K} \|x\| > 0$. \square

5. Proof of Proposition IV.6

In order to prove IV.6, let us first fix some notation. Let X be an oriented smooth compact manifold and $K(X)$ be the K-ring of X , i.e. the ring of equivalence classes of complex vector bundles on X , where $E \sim E'$ if $E + n \simeq E' + m$ (An introductory text on this topic is e.g. [33]).⁹ Let $p_i(X) := p_i(TX) \in H^{2i}(X, \mathbb{Q})$ be the image of i -th rational Pontryagin class evaluated on the tangent bundle TX . Furthermore, let $\hat{A}(p_1, \dots, p_n)$ be the \hat{A} -genus, i.e. the genus associated to the power series $\frac{\sqrt{z}/2}{\sinh(\sqrt{z}/2)}$ [34].¹⁰

⁹ Here m denotes the m -dimensional trivial bundle.

¹⁰ This means that by construction the identity $p(E \oplus F) = p(E)p(F)$ of the total rational Pontryagin class transfers to the \hat{A} -genus.

Let $\text{ch} : K(X) \rightarrow H^*(X, \mathbb{Q})$ be the Chern class and let $\text{ch}(X) := \text{ch}(K(X)) \subseteq H^*(X, \mathbb{Q})$. For $z := \sum_{i=0}^{\infty} z^{2i} \in H^*(X, \mathbb{Q})$, with $z_{2j} \in H^{2i}(X, \mathbb{Q})$, let $z^{(t)} := \sum_{i=0}^{\infty} z^{2i} t^i$ for $t \in \mathbb{Q}$. Note that $(yz)^{(t)} = y^{(t)} z^{(t)}$.

For $z \in \text{ch}(X)$, $d \in H^2(X, \mathbb{Q})$, $t \in \mathbb{Q}$ we define the Hilbert polynomial in t to be $H_{X,z,d}(t) := (z^t e^{d/2} \hat{A}(p(X))) [X]$, where $[X]$ is the fundamental class of X [35, 36]. Furthermore for $q \in \mathbb{Q}$ let $\nu_2(q) :=$ exponent of 2 as primefactor of q . The result of Walgenbach in [21] is based on the result of Mayer in [37],

Theorem A.3. [37] *Let X be a $2n$ -dimensional compact oriented smooth manifold and H be the Hilbert polynomial associated with $d \in H^2(X; \mathbb{Z})$ and $z \in \text{ch}(X)$. Then X cannot be immersed in Euclidean space of dimension $-2\nu_2(H(\frac{1}{2})) - 1$ and cannot be embedded in Euclidean space of dimension $-2\nu_2(H(\frac{1}{2}))$.*

Walgenbach obtains his results by computing $H(\frac{1}{2})$ for some $d \in H^2(X; \mathbb{Z})$ and $z \in \text{ch}(X)$ using combinatorial methods. With the following lemma, IV.6 follows directly by observing that $\nu_2(a \cdot b) = \nu_2(a) + \nu_2(b)$. Let X_1, X_2 be $2n$ -dimensional compact oriented smooth manifolds and let $\pi_i : X_1 \times X_2 \rightarrow X_i$, $i = 1, 2$, be the canonical projections.

Lemma A.4. *For $z_i \in \text{ch}(X_i)$, $d_i \in H^2(X_i, \mathbb{Q})$ and $[X_i] = (\pi_i)_*[X_1 \times X_2]$, $i = 1, 2$, let $z := \pi_1^*(z_1)\pi_2^*(z_2)$, $d := \pi_1^*(d_1) + \pi_2^*(d_2)$. Then,*

$$H_{X_1 \times X_2, z, d}(t) = H_{X_1, z_1, d_1}(t) H_{X_2, z_2, d_2}(t).$$

Proof. First, note that $z \in K(X_1 \times X_2)$, since

$$\pi_1^*(z_1)\pi_2^*(z_2) = \pi_1^*(\text{ch}(E_1))\pi_2^*(\text{ch}(E_2)) = \text{ch}(\pi_1^*(E_1))\text{ch}(\pi_2^*(E_2)) = \text{ch}(\pi_1^*(E_1) \oplus \pi_2^*(E_2)).$$

Furthermore note that

$$\begin{aligned} p(X_1 \times X_2) &= p(T(X_1 \times X_2)) = p(\pi_1^*TX_1 \oplus \pi_2^*TX_2) \\ &= p(\pi_1^*(TX_1))p(\pi_2^*(TX_2)) = \pi_1^*(p(TX_1))\pi_2^*(p(TX_2)). \end{aligned}$$

This, together with the fact that all cohomology classes involved are even dimensional and hence commute, yields

$$\begin{aligned} H_{X_1 \times X_2, z, d}(t) &= \left(z^{(t)} e^{d/2} \hat{A}(X_1 \times X_2) \right) [X_1 \times X_2] \\ &= \left(\pi_1^*(z_1)^{(t)} \pi_2^*(z_2)^{(t)} e^{\pi_1^*(d_1/2) + \pi_2^*(d_2/2)} \pi_1^*(\hat{A}(TX_1)) \pi_2^*(\hat{A}(TX_2)) \right) [X_1 \times X_2] \\ &= \left(\pi_1^* \left(z_1^{(t)} e^{\pi_1^*(d_1)/2} \hat{A}(X_1) \right) \pi_2^* \left(z_2^{(t)} e^{\pi_2^*(d_2)/2} \hat{A}(X_2) \right) \right) [X_1 \times X_2] \\ &= \left(z_1^{(t)} e^{\pi_1^*(d_1)/2} \hat{A}(X_1) \right) [X_1] \left(z_2^{(t)} e^{\pi_2^*(d_2)/2} \hat{A}(X_2) \right) [X_2] \\ &= H_{X_1, z_1, d_1}(t) H_{X_2, z_2, d_2}(t). \end{aligned}$$

□

6. Proof of Proposition IV.8

Let X be a smooth compact n -manifold with tangent bundle TX . Let $\phi : X \rightarrow \mathbb{R}^{n+k}$ be an immersion and let NX be the normal bundle, i.e. $TX \oplus NX \simeq n + k$. Furthermore let ω be the total Stiefel-Whitney class. Let us state the following well-known result.

Proposition A.5. [25] *Let i be the degree of $\bar{\omega}(X) = \omega(NX) \in H^*(X, \mathbb{Z}_2)$. Then X cannot be immersed in Euclidean space of dimension $n + i$ and cannot be embedded in Euclidean space of dimension $n + i + 1$.*

In order to use this result, we need to compute $\bar{\omega}(PW_{n,k})$. The following is similar to [24], where the dual Stiefel-Whitney class of the real projective Stiefel manifolds is computed. Let L be the complex line bundle associated to the $U(1)$ -principal bundle $W_{n,k} \rightarrow PW_{n,k}$ and let x be the mod 2 Euler class of L^{11} . In [38], the cohomology ring $H^*(PW_{n,k}, \mathbb{Z}_2)$ for $k < n$ is found to be

$$\mathbb{Z}_2[x]/(x^N) \oplus \Lambda(y_{n-k+1}, \dots, y_n),$$

with $y_i \in H^{2i-1}(PW_{n,k}, \mathbb{Z}_2)$. It is shown in [27], that $TPW_{n,k}$ is stably isomorphic to nkL^* , where L^* is regarded as a real vector bundle. Hence $\omega(TPW_{n,k}) = \omega(L^*)^{nk}$ and we obtain

$$\omega(NPW_{n,k}) = \bar{\omega}(TPW_{n,k}) = \omega(L^*)^{-nk}.$$

Since the odd Stiefel-Whitney classes of complex vector bundles (regarded as real vector bundles) vanish [33], and the Euler class is mapped to the top Stiefel Whitney class under the coefficient homomorphism $H^*(PW_{n,k}, \mathbb{Z}) \rightarrow H^*(PW_{n,k}, \mathbb{Z}_2)$ [33], we get $\omega(L) = \omega(L^*) = 1 + x$. Thus,

$$\begin{aligned} \omega(NPW_{n,k}) &= (1 + x)^{-nk} \\ &= \sum_{j=1}^{\infty} (-1)^j \binom{nk + j - 1}{j} x^j. \end{aligned}$$

We now want to find

$$\gamma(m, k) = \text{the biggest } j \text{ such that the coefficient of } x^j \text{ does not vanish in } \mathbb{Z}_2[x]/(x^N)$$

Since we factor over the ideal generated by x^N , we clearly have $j \leq N(n, k)$. Passing to mod2, we get $2\gamma(m, k) = \sigma(n, k)$. This proves IV.8.

[1] H Häffner, W Hänsel, CF Roos, J Benhelm, et al. Scalable multiparticle entanglement of trapped ions. *Nature*, 438(7068):643–646, 2005.

¹¹ I.e. the image of the Euler class of L under the coefficient homomorphism $H^*(PW_{n,k}, \mathbb{Z}) \rightarrow H^*(PW_{n,k}, \mathbb{Z}_2)$

- [2] Teiko Heinosaari, Luca Mazzarella, and Michael M Wolf. Quantum tomography under prior information. *Communications in Mathematical Physics*, 318(2):355–374, 2013.
- [3] Stefan Weigert. Pauli problem for a spin of arbitrary length: A simple method to determine its wave function. *Physical Review A*, 45(11):7688, 1992.
- [4] Jean-Pierre Amiet and Stefan Weigert. Reconstructing the density matrix of a spin s through Stern-Gerlach measurements: II. *Journal of Physics A: Mathematical and General*, 32(25):L269, 1999.
- [5] Jean-Pierre Amiet and Stefan Weigert. Reconstructing the density matrix of a spin s through Stern-Gerlach measurements. *Journal of Physics A: Mathematical and General*, 31(31):L543, 1998.
- [6] J Finkelstein. Pure-state informationally complete and “really” complete measurements. *Physical Review A*, 70(5):052107, 2004.
- [7] V Voroninski D Mondragon. Determination of all pure quantum states from a minimal number of observables. *arXiv:1306.1214*, 2013.
- [8] Claudio Carmeli, Teiko Heinosaari, Jussi Schultz, and Alessandro Toigo. Tasks and premises in quantum state determination. *Journal of Physics A: Mathematical and Theoretical*, 47(7):075302, 2014.
- [9] Claudio Carmeli, Teiko Heinosaari, Jussi Schultz, and Alessandro Toigo. Expanding the principle of local distinguishability. *Phys. Rev. A*, 91:042121, Apr 2015.
- [10] Steven T Flammia, Andrew Silberfarb, and Carlton M Caves. Minimal informationally complete measurements for pure states. *Foundations of Physics*, 35(12):1985–2006, 2005.
- [11] David Gross. Recovering low-rank matrices from few coefficients in any basis. *IEEE Trans. on Information Theory*, 57:1548–1566, 2011.
- [12] Steven T Flammia, David Gross, Yi-Kai Liu, and Jens Eisert. Quantum tomography via compressed sensing: error bounds, sample complexity and efficient estimators. *New Journal of Physics*, 14(9):095022, 2012.
- [13] R. Millane. Phase retrieval in crystallography and optics. *JOSA A*, 7:394–411, 1990.
- [14] David Gross, Felix Kraemer, and Richard Kueng. A partial derandomization of phaselift using spherical designs. *Journal of Fourier Analysis and Applications*, pages 1–38, 2014.
- [15] David Gross, Yi-Kai Liu, Steven T Flammia, Stephen Becker, and Jens Eisert. Quantum state tomography via compressed sensing. *Physical review letters*, 105(15):150401, 2010.
- [16] Nathaniel Johnston Gus Gutoski. Process tomography for unitary quantum channels. *J. Math. Phys.*, 55:032201, 2014.
- [17] Ivan H. Deutsch Charles H. Baldwin, Amir Kalev. Quantum process tomography of unitary and near-unitary maps. *Phys. Rev. A*, 90:012110, 2014.
- [18] Alexander S Holevo. *Probabilistic and statistical aspects of quantum theory*, volume 1. Springer, 2011.
- [19] Paul Busch, Marian Grabowski, and Pekka Johannes Lahti. *Operational quantum physics*, volume 31. Springer, 1995.
- [20] Frank W Warner. *Foundations of differentiable manifolds and Lie groups*, volume 94. Springer, 1971.
- [21] Markus Walgenbach. Lower bounds for the immersion dimension of homogeneous spaces. *Topology and its Applications*, 112(1):71–86, 2001.
- [22] Toby Cubitt, Ashley Montanaro, and Andreas Winter. On the dimension of subspaces with bounded schmidt rank. *Journal of Mathematical Physics*, 49(2):022107, 2008.
- [23] Kenneth R Davidson. *C*-algebras by example*, volume 6. American Mathematical Soc., 1996.
- [24] Nelza Elisabete Barufatti. Obstructions to immersions of projective stiefel manifolds. *CONTEMPORARY MATHEMATICS*, 161:281–281, 1994.
- [25] John Willard Milnor and James D Stasheff. *Characteristic classes*, volume 93. Princeton University Press Princeton, 1974.

- [26] R James Milgram. Immersing projective spaces. *The Annals of Mathematics*, 85(3):473–482, 1967.
- [27] L. Astey, S. Gitler, E. Micha, and G. Pastor. Parallelizability of complex projective stiefel manifolds. *Proceedings of the American Mathematical Society*, 128(5):pp. 1527–1530, 2000.
- [28] Morris W Hirsch. Immersions of manifolds. *Transactions of the American Mathematical Society*, 93(2):242–276, 1959.
- [29] M.F. Atiyah. Immersions and embeddings of manifolds. *Topology*, 1(2):125 – 132, 1962.
- [30] Elmer Rees. Some embeddings of lie groups in euclidean space. *Mathematika*, 18(01):152–156, 1971.
- [31] Morris W Hirsch. *Differential topology*. Springer, 1976.
- [32] Thomas Bagby, Len Bos, and Norman Levenberg. Multivariate simultaneous approximation. *Constructive approximation*, 18(4):569–577, 2002.
- [33] Allen Hatcher. Vector bundles and k-theory. <http://www.math.cornell.edu/~hatcher>, 2003.
- [34] Friedrich Hirzebruch. *Topological methods in algebraic geometry.*, volume 131. Springer, 1995.
- [35] Armand Borel and Friedrich Hirzebruch. Characteristic classes and homogeneous spaces, ii. *American Journal of Mathematics*, 81(2):315–382, 1959.
- [36] Michael F Atiyah and Friedrich Hirzebruch. Quelques théorèmes de non-plongement pour les variétés différentiables. *Bulletin de la Société Mathématique de France*, 87:383–396, 1959.
- [37] Karl Heinz Mayer. Elliptische Differentialoperatoren und Ganzzahligkeitssätze für charakteristische Zahlen. *Topology*, 4(3):295–313, 1965.
- [38] L Astey, S Gitler, E Micha, and G Pastor. Cohomology of complex projective stiefel manifolds. *Canadian Journal of Mathematics*, 51(5):897–914, 1999.