

Bifurcations in models of a society of reasonable contrarians and conformists

Franco Bagnoli*

*Dipartimento di Fisica e Astronomia, Università di Firenze,
Via G. Sansone 1, 50017 Sesto Fiorentino (FI), Italy;
also INFN, sez. Firenze.*

Raúl Rechtman†

*Instituto de Energías Renovables,
Universidad Nacional Autónoma de México,
Apdo. Postal 34, 62580 Temixco Mor., Mexico*

We study models of a society composed of a mixture of conformist and reasonable contrarian agents that at any instant hold one of two opinions. Conformists tend to agree with the average opinion of their neighbors and reasonable contrarians to disagree, but revert to a conformist behavior in the presence of an overwhelming majority, in line with psychological experiments. The model is studied in the mean field approximation and on small-world and scale-free networks. In the mean field approximation, a large fraction of conformists triggers a polarization of the opinions, a pitchfork bifurcation, while a majority of reasonable contrarians leads to coherent oscillations, with an alternation of period-doubling and pitchfork bifurcations up to chaos. Similar scenarios are obtained by changing the fraction of long-range rewiring and the parameter of scale-free networks related to the average connectivity.

PACS numbers: 05.45.Ac,05.50.+q,64.60.aq,64.60.Ht

I. INTRODUCTION

In a previous work [1], we studied the collective behavior of a society of reasonable contrarian agents. The rationale was that in some cases, and in particular in the presence of frustrated situations like in minority games [2], it is not convenient to always follow the majority, since in this case one is always on the “loosing side” of the market. This is one of the main reasons for the emergence of a contrarian attitude. On the other hand, if all or almost all agents in a market take the same decision, it is often wise to follow such a trend. We can denote such a situation with the word “social norm”. Following an overwhelming majority is an ecological strategy since it is probable that this coherent behavior is due to some unknown piece of information, and in any case the competitive loss is minimal since it equally affects the other agents. Indeed, it is well known that in the presence of an overwhelming majority, individuals tend to align with it, even if it is in contrast to evidence [3].

The agents in our model can hold opinion 0 or opinion 1, representing one of two parties, or against/in favor of an option. Each agent gathers the average opinion of his neighbors and changes his opinion according to this average. The time evolution is synchronous, so the model is essentially a cellular automaton.

Contrarians were introduced in a different socio-physical model by S. Galam [4, 5]. In this case they had the effect of destroying consensus in a society mainly

formed by conformists. In this case, when the fraction of contrarians becomes opinion-dependent, chaotic dynamics appear [6].

A society fully composed by reasonable contrarians exhibits interesting behaviors when changing the topology of the connections. On a one-dimensional regular lattice, there is no long-range order, the evolution is disordered and the average opinion is always halfway between the extreme values 0 and 1. However, adding long-range connections or rewiring existing ones, we observe the Watts-Strogatz “small-world” effect, with a transition towards a mean field behavior. But since in this case the mean field equation is, for a suitable choice of parameters, chaotic, we observe the emergence of coherent oscillations, with a bifurcation cascade eventually leading to a chaotic-like behavior of the average opinion. The small-world transition is essentially a synchronization effect. Similar effects with a bifurcation diagram resembling that of the logistic map have been observed in a different model of “adapt if novel - drop if ubiquitous” behavior, upon changing the connectivity [7, 8].

Since a homogeneous society of unreasonable contrarians is not so reasonable, we study here the problem of collective behaviors in the presence of a mixture of conformists and contrarians. To keep things simple, reasonable contrarian and conformist agents have the same behavior in the presence of an overwhelming majority of their neighbors. Conformists become less conformists in the presence of a large majority. We can call them “slightly unreasonable”. The presence of reasonable contrarians and slightly unreasonable contrarians avoids absorbing states which are rather unusual in real societies.

In the presence of a strong fraction of conformists, we have the classical ferromagnetic Ising scenario, with the

*Electronic address: franco.bagnoli@unifi.it

†Electronic address: rrs@cie.unam.mx

appearance of a stationary average opinion different from one half. As this fraction becomes smaller, this polarized opinion vanishes, as expected. What is unexpected is another bifurcation, with the appearance of oscillations and chaos as the fraction of conformists becomes smaller.

The outline of the present paper is as follows: in Sec. II we present the model in detail. Its mean field approximation is discussed in Sec. III. Then the model is studied on small-world networks, Sec. IV and on on scale-free networks, Sec. V. We end the presentation with some conclusions, Sec. VI.

II. THE MODEL

Our model society is formed by N agents with a fraction ξ of conformists and a fraction $1 - \xi$ of contrarians. Agent i , $i = 0, \dots, N - 1$ holds an opinion $s(i, t) \in \{0, 1\}$ at time t . The opinions of all agents change synchronously in time. Agent i gathers the average opinion of his neighbors and changes opinion tending to agree with his neighbors if he is a conformist, or to disagree if he is a contrarian.

The neighborhoods of all agents are defined by the adjacency matrix A with components $a_{ij} \in \{0, 1\}$ in a way that $a_{ij} = 1$ if agent j belongs to i 's neighborhood and $a_{ij} = 0$ if he doesn't. The connectivity k_i of agent i is the number of agents in i 's neighborhood,

$$k_i = \sum_j a_{ij} \quad (1)$$

and the average opinion h_i of his neighbors is

$$h_i = \frac{1}{k_i} \sum_j a_{ij} s(j). \quad (2)$$

The average opinion c of the society is

$$c = \frac{1}{N} \sum_i s_i. \quad (3)$$

Given the average opinion h of the neighbors of agent i at time t , $s(i, t + 1) = 1$ according to the transition probability $\tau(h; J)$ defined by [1]

$$\tau(h; J) = \begin{cases} \varepsilon & \text{if } h < q, \\ \frac{1}{1 + \exp(-2J(2h - 1))} & \text{if } q \leq h \leq 1 - q, \\ 1 - \varepsilon & \text{if } h > 1 - q. \end{cases} \quad (4)$$

In this last expression J represents the agent's conviction, conformists have $J > 0$, reasonable contrarians, $J < 0$. The graphs of τ are shown in Fig. 1. A reasonable conformist or a contrarian can assume opinion 1 with probability ε ($1 - \varepsilon$) when $0 \leq h \leq q$ ($1 - q \leq h \leq 1$). When $q < h < 1 - q$ a reasonable conformist will probably agree with his neighbors, and a contrarian will probably disagree. Both conformists and contrarians share the




FIG. 1: (Color online) The transition probabilities τ given by Eq. (4) with $|J| = 3$. For $q < h < 1 - q$, τ is an increasing function of h (in red) for conformists, $J = 3$, and a decreasing one (in blue), $J = -3$, for reasonable contrarians. In this and the following Figs. $q = 0.1$ and $\varepsilon = 0.2$ unless stated otherwise.

same values of $|J|$, q , and ε . A nonzero value of ε avoids the presence of absorbing states but causes the “slightly unreasonable” behavior of conformists. Unless otherwise stated, we always use the value $q = 0.1$ and $\varepsilon = 0.2$, as in Ref. [1].

In the mean field approximation c changes deterministically and we can characterize the properties of its trajectory by means of the Lyapunov exponent λ . On the other hand, c changes probabilistically on networks, so we use Boltzmann's entropy η [1, 9, 10], which is applicable in both cases. To define η we partition the unit interval in L disjoint equal sized subintervals I_i , $i = 1, \dots, L$ and find the probability q_i that c falls in subinterval I_i . Then

$$\eta = \frac{-1}{\log L} \sum_{i=1}^L q_i \log q_i. \quad (5)$$

The probabilities q_i are found numerically by finding the frequency with which an orbit visits each subinterval I_i after a transient. A fixed point of the trajectory corresponds to $\eta = 0$, and when the orbit visits every subinterval I_i with the same frequency $\eta = 1$. If $L = 2^b$ and the orbit is periodic with period 2^a , $\eta = a/b$. In the limit $L \rightarrow \infty$, $\eta \rightarrow 0$ for periodic orbits.

For deterministic maps we can compare the behavior of the Lyapunov exponent λ and that of Boltzmann's entropy η . Values of $\lambda > 0$ are equivalent to $\eta > 1/2$ when L is sufficiently large. In other words, deterministic chaos corresponds to $\eta > 1/2$ and order to $\eta < 1/2$. By extending this correspondence to the probabilistic network dynamics, we define disorder whenever $\eta \gtrsim 1/2$ and order when $\eta \lesssim 1/2$.

III. MEAN FIELD APPROXIMATION

We start with a model of a society where the neighborhood of each agent i , either conformist or contrarian,




FIG. 2: (Color online.) Return maps of the mean field approximation, Eq. (6), smooth curves (in red), and 40 iterates of the map starting from a random initial c , stepping lines (in blue). (a) $\xi = 0.06$. (b) $\xi = 0.2$. (c) $\xi = 0.4$. (d) $\xi = 0.9$. In this and the following Figs., $|J| = 5$, $k = 20$, $\varepsilon = 0.2$, and $q = 0.1$ unless stated otherwise.

is formed by k random neighbors, i.e., the mean field approximation for a fixed connectivity k . With a fraction ξ of conformists and a fraction $1 - \xi$ of reasonable contrarians, the time evolution of the average opinion c is

$$c' = \sum_{w=0}^k \binom{k}{w} c^w (1-c)^{k-w} [\xi \tau(w/k; J) + (1-\xi) \tau(w/k; -J)], \quad (6)$$

with c and c' the average opinions at times t and $t + 1$ respectively. In the right-hand side term, the first parenthesis is the w combinations from a set of k elements.

In Fig. 2 we show return maps of Eq. (6) for different values of ξ . For small ξ , Fig. 2 (a), the map is chaotic, and for larger values of ξ , we find periodic orbits or fixed points, Figs. 2 (b), (c) and (d).

In Fig. 3 we show the mean field bifurcation diagrams of c given by Eq. (6) as a function of the fraction of conformists ξ for $\varepsilon = 0.2$ and $\varepsilon = 0$. In this second case, there are absorbing states for large ξ but for smaller values the diagram hardly depends on the value of ε . The leftmost vertical line at ξ_c marks the threshold at which the chaotic region ends. In Fig. 4 (a) we show an amplification of the bifurcation diagram for small ξ and in Fig. 4 (b) the corresponding Lyapunov exponent λ and Boltzmann's entropy η . For $\xi > \xi_c$, $\lambda < 0$ and $\eta < 1/2$. In Figs. 3 (a) and (b), the next vertical line at ξ_a corresponds to the period-doubling bifurcation from a period two orbit to a fixed point. For $\xi_c < \xi < \xi_a$ the rather large fraction of contrarians causes symmetric oscillations of c . The rightmost vertical line at ξ_b corresponds to a pitchfork bifurcation from $c = 1/2$ to $c > 1/2$ when




FIG. 3: (Color online.) The mean field average opinion c as a function of the fraction of conformists ξ with $|J| = 5$, $k = 20$, and $q = 0.1$. (a) $\varepsilon = 0.2$, $\xi_c = 0.0665$. (b) $\varepsilon = 0.0$, $\xi_c = 0.0795$. In (a) and (b), $\xi_a = 0.3245$ and $\xi_b = 0.6755$ so that $1 - \xi_b = \xi_a$. For ξ slightly larger than ξ_c the first and third branches (starting from the bottom) correspond to $c_0 = 0.9$ (in red) and the other two to $c_0 = 0.1$ (in blue). For $\xi_b < \xi \leq 1$, the upper branch (in red) corresponds to $c_0 = 0.9$, the lower one (in blue) to $c_0 = 0.1$. For all values of ξ , two consecutive iterations are plotted after a transient of 1,000 time steps for both $c_0 = 0.9$ (in red) and $c_0 = 0.1$ (in blue).

$c_0 > 1/2$ and to $c < 1/2$ when $c_0 < 1/2$ with c_0 the average opinion at $t = 0$, (see also Fig. 2).

We prove in the Appendix that $\xi_a = 1 - \xi_b$ and find that the approximate behavior of ξ_a as a function of J and k is given by

$$\xi_a(J, k) = \frac{1}{2} \left(1 - \frac{1}{J} \sqrt{1 + \frac{2J^2}{k}} \right). \quad (7)$$

In Fig. 5 we show that Eq. (7) agrees with the numerical results for large connectivities k .

In Fig. 6, we show bifurcation diagrams of the average opinion c as $|J|$ changes for small values of ξ . For $\xi = 0$, Fig. 6 (a), there are no conformists and we have the case studied in Ref. [1]. The bifurcation diagram is stretched horizontally as ξ grows. For small $|J|$, $c = 1/2$, and as this parameter grows, there is a first bifurcation to a period 2 orbit. What seems like a second bifurcation to a period-four orbit, corresponds to two period-two bifurcations that depend on the initial opinion c_0 . Starting from below, the first and third branches (in red) correspond to $c_0 = 0.9$, and the other two (in blue) to $c_0 = 0.1$.

IV. SMALL-WORLD NETWORKS

Real societies are not random, nor regular. It is interesting to study what happens when the topology changes, due for instance to advances in the transportation system, or to politics favoring mixing, etc. We studied the effect of rewiring a fraction p of links in a regular, one-dimensional society with connectivity k . This leads to the small-world networks first discussed by Watts and Strogatz [11]. As p , the long range connection probability, grows, small-world networks approach a mean field behavior. As we show in Fig. 7, the bifurcation diagrams




FIG. 4: (Color online.) (a) The mean field average opinion c as a function of the fraction of conformists ξ . For ξ slightly smaller and also larger than ξ_c , we see four branches. Starting from below, the first and third branches correspond to $c_0 = 0.9$ (in red), and the second and fourth to $c_0 = 0.1$ (in blue). After a 1,000 time steps transient, 32 consecutive iterations are plotted for each value of ξ starting with $c_0 = 0.9$ and $c_0 = 0.1$. (b) The Lyapunov exponent λ , bottom curve for small ξ (in blue), and the entropy η , top curve for small ξ (in red), of the mean field average opinion c , Eq. (6), as functions of ξ . For each value of ξ , λ is evaluated during 1,000 time steps and for η , the unit interval is divided in $2^{10} = 1,024$ equal size subintervals and the frequency with which each subinterval is visited is found during $100 \times 2^{10} = 102,400$ time steps after a 300 time steps transient.




FIG. 5: (Color online.) Mean field critical values ξ_a as functions of k for $|J| = 5$, circles (in red), and $|J| = 10$ squares (in blue). The continuous curves are the graphs of Eq. (7), the bottom one (in red) is for $|J| = 5$, and the top one (in blue) corresponds to $|J| = 10$. The horizontal dashed lines show the asymptotic values (derived in the Appendix), $\xi_a(5, \infty) = 0.4$, bottom dashed line (in red), and $\xi_a(10, \infty) = 0.45$ top dashed line (in blue). Numerical data obtained after a transient of 1,000 time steps starting with $c_0 = 0.9$.

of c as functions of p are similar to those obtained by varying J in the mean field approximation.

In Fig. 7 (a) we study a society of reasonable contrarians. For small values of p , c fluctuates around $c = 0.5$. For slightly larger values, there are noisy oscillations around two symmetric values, in a way reminiscent of the period-doubling bifurcations of deterministic systems. For even larger values, $p \simeq 0.4$, we observe the appearance of two different noisy oscillating states of pe-




FIG. 6: (Color online.) Mean field bifurcation diagrams of the average opinion c as functions of $|J|$ for several values of the fraction of conformists ξ . (a) $\xi = 0.0$. (b) $\xi = 0.04$. (c) $\xi = 0.07$. (d) $\xi = 0.08$. Starting from small $|J|$ there is a first bifurcation to a period two orbit. For larger values of $|J|$ there are four branches that lead. Starting from below, the first and third branches are obtained with an initial average opinion $c_0 = 0.9$ (in red), and the second and fourth with $c_0 = 0.1$ (in blue). For each value of $|J|$, after a transient of 1,000 time steps the next 32 iterations are plotted for two initial average opinions c_0 .

riod 2. For $c_0 = 0.1$ the orbit oscillates between the first and third branches, starting from the bottom (blue points), and for $c_0 = 0.9$ the orbit oscillates between the second and fourth branches (red points). This roughly corresponds to what is shown in Fig. 6 (a), although in that case the diagram is drawn as a function of $|J|$ and here as a function of p . For $\xi = 0.05$, Fig. 7 (b) we have a similar scenario and for larger values of ξ , Figs. 7 (c) and (d), we find a great similarity with the mean field behavior.

For $\xi = 0.2$, Fig. 7 (c), c fluctuates around $c = 0.5$ for small values of p and fluctuates around two branches for larger values. These branches agree with the period 2 orbit of the mean field approximation, Eq. (6) shown in Fig. 3. For $\xi = 0.8$, Fig. 7 (d), and $p \gtrsim 0.2$, c fluctuates around one of two values, depending on c_0 , the average opinion at $t = 0$. These values also agree with the mean field approximation shown in Fig. 3.

It is possible to roughly understand these results assuming that the main contributions to the mean field character of the collective behavior come from the fraction of links that are rewired (long-range connections) that depend on p . The actual value of the field $(2h - 1)$ in Eq. (4) is multiplied by a factor p , so that changing p is roughly equivalent to changing J .

In Fig. 8 we show bifurcation diagrams of c on small-world networks as a function of the fraction of conformists ξ for some values of the long range connection




FIG. 7: (Color online.) Bifurcation diagrams of the average opinion c on a small-world network as a function of the long range connection probability p for different values of the fraction of conformists ξ with $N = 20,000$, $|J| = 5$, $k = 20$. The initial opinion of each agent is chosen at random between 0 and 1 with $c_0 = 0.1$ (blue points) and $c_0 = 0.9$ (red points). (a) $\xi = 0.0$. (b) $\xi = 0.05$. (c) $\xi = 0.20$. (d) $\xi = 0.8$. In (c) and (d) the horizontal lines are the values of c of the period two orbit of Eq. (6). For each value of p , 32 points are plotted after a 1,000 time step transient for each c_0 .

probability p . This sequence of plots illustrate an unexpected behavior. For $p = 0$, Fig. 8 (a), there is a pitchfork bifurcation at $\xi \sim 0.9$. For larger values of p , the pitchfork bifurcation occurs at smaller values of ξ and an oscillating “bubble” is formed, Fig. 8 (b) for $0.12 \lesssim \xi \lesssim 0.22$. This oscillating region grows with c and for $p = 0.8$, Fig. 8 (d), the bifurcation diagram is similar to the mean field one, Fig. 3 (a).

We can explain this behavior assuming that the conformist behavior promotes synchronization, as does p . So the system progressively synchronizes starting from high values of ξ , but this synchronization is not visible if the dynamics leads to fixed points. When the synchronization reaches the oscillating phases, it becomes manifest by means of the coherent oscillation of the population. So, this coherent dynamical behavior appears to start first in the vicinity of the bifurcation for $\xi \simeq 0.2$ and then, by increasing p , it propagates to lower values of ξ .

V. SCALE-FREE NETWORKS

In this section we present results of the model on uncorrelated scale-free networks [12]. Starting from a fully connected group of m agents, other $N - m$ agents join sequentially, each one choosing m neighbors among those already in the group. The choice is preferential, the probability that a new member chooses agent i is proportional to its connectivity k_i , the number of neighboring agents




FIG. 8: (Color online.) Bifurcation diagrams of the average opinion c on a small-world network as functions of the fraction of conformists ξ for different values of the long range connection probability p with $N = 50,000$, $|J| = 5$, $k = 20$. (a) $p = 0.0$. (b) $p = 0.05$. (c) $p = 0.20$. (d) $p = 0.8$. For each value of p , at $t = 0$ the opinion of each agent is chosen at random in such a way that the average opinion is $c_0 = 0.1$ or $c_0 = 0.9$. After a 50,000 time step transient, 32 points are plotted. For $0.6 \lesssim \xi \lesssim 0.8$, the orbits that started with $c_0 = 0.99$ have $c > 1/2$, (points in red), and those that started with $c_0 = 0.1$ have $c < 1/2$, (points in blue).

that agent i already has. Another way of building the network is choosing a random edge of a random node and connecting to the other end of the edge, since such an edge arrives to a vertex with probability proportional to $kp(k)$ with $p(k)$ the probability that a randomly selected node has connectivity k [13].

In Ref. [1] we showed that the dynamics of a model of a society whose agents are all reasonable contrarians on a scale-free network with m initially connected agents is comparable to the mean field approximation of Sec. III with connectivity k provided that

$$k = \alpha m \quad (8)$$

with $\alpha \sim 1.7$ for scale-free networks with $p(k) \propto k^{-3}$. In Fig. 9 we show that this result also holds for the model of societies studied here. In this Figure we compare the bifurcation diagrams, as ξ changes, of the dynamics on a scale-free network with the corresponding mean field one, Figs. 9 (a) and (b), and the scale-free entropies with those of the mean field map, Figs. 9 (c) and (d). There is a reasonable agreement in both bifurcation diagrams. Both entropies show a good agreement where there is disorder, that is $\eta > 1/2$ but not when $\eta < 1/2$. This can be understood from the bifurcation diagrams. While the scale-free network dynamics is stochastic and therefore the orbit visits many subintervals, the mean field one visits a smaller number. For example, for $\xi = 0.2$, the

mean field dynamics has period two so $\eta = 2/8$ both in Figs. 9 (c) and (d).




FIG. 9: (a) and (b). Bifurcation diagrams of the average opinion c as functions of the fraction of conformists ξ on a scale-free network, large points (in blue) with $m = 20$ in (a) and $m = 30$ in (b), and the mean field approximation, Eq. (6) with $k = 1.7 \times 20 = 34$ in (a) and $k = 1.7 \times 30 = 51$ in (b), smaller points (in red). (c) and (d) Boltzmann's entropy η of c as functions of ξ on a scale-free network, top curve for $\xi = 1/2$ (in red), and of Eq. (6), bottom curve for $\xi = 1/2$ (in blue). (c) $m = 20$, $k = 34$. (d) $m = 30$, $k = 51$. In (a) and (b), for c on a scale-free network and for each value of ξ , at $t = 0$ the opinion of each agent is chosen at random in such a way that the average opinion is $c_0 = 0.9$ and for the mean field bifurcation diagram $c_0 = 9$ and $c_0 = 0.1$. After a transient of 300 time steps, the next 32 values of c are plotted. For the entropy η and each value of ξ , the unit interval is divided in $2^8 = 256$ equal size subintervals and the frequency with which each subinterval is visited is found during $100 \times 2^8 = 25,600$ time steps after a 300 time steps transient.

VI. CONCLUSIONS

The dynamics of the mean field approximation of the average opinion c , Eq. (6) as a function of ξ is chaotic with periodic windows when $0 < \xi < \xi_c$ and oscillates periodically between two symmetric values when $\xi_c < \xi < \xi_a$. For $\xi_a < \xi < \xi_b$, $c = 1/2$, and for $\xi_b < \xi$, $c > 1/2$ ($c < 1/2$) if $c_0 = 0.9$ ($c_0 = 0.1$). Also $\xi_a \sim 1 - \xi_b$ and ξ_a approaches a limit value value as k grows. Eq. (6) depends on the connectivity k and on the transition probability τ that in turn depends on k , $|J|$, ε , and q . As far as we have explored the parameter space, the above description is generic with the exception that for large values of k the chaotic phase is bounded below by $\xi_d > 0$. For $0 < \xi < \xi_d$ there are two period two orbits that depend on c_0 and for $\xi_d < \xi < \xi_c$ the orbits are chaotic with periodic windows.

In small-world networks with small values of p , Fig. 8 (b), the coherent oscillations appear first for a population with a small fraction of conformist rather than for a pure-contrarian one. This is probably due to the fact that coherent oscillations are a signal of synchronization, and the presence of conformists increases the synchronization. On the other hand, if the synchronized dynamics leads to a stable fixed point, the degree of synchronization is not manifest. As a result, the first synchronized zone is near the bifurcation point $\xi \simeq 0$.

Dynamics on small-world networks approaches that of the mean field approximation as the long range connection probability p grows. Dynamics of c on scale-free networks is similar to that of the mean field approximation provided $k = \alpha m$ with $\alpha \sim 1.7$.

Acknowledgements

Interesting discussions with Héctor D. Cortés González and Maximiliano Valdez González are acknowledged. This work was partially supported by project PAPIIT-DGAPA-UNAM IN109213. F.B. acknowledges partial financial support from European Commission (ICT-20111.1.6) Proposal No. 288021 EINS - Network of Excellence in Internet Science and European Commission (FP7-ICT-2013-10) Proposal No. 611299 SciCafe 2.0.

We thank S. Galam and K.D. Harris for having pointed out their interesting contributions about the effects induced by the presence of contrarians and of chaotic bifurcations upon rewiring.

Appendix

The bifurcation points ξ_a and ξ_b can be found from the mean field evolution, Eq. (6), since at both values, the absolute value of the derivative of this expression must be 1. For large k the mean field approximation for the average opinion, can be approximated by [1]

$$c' = \int dx \sqrt{\frac{k}{2\pi c(1-c)}} \exp\left(-\frac{k(x-c)^2}{2c(1-c)}\right) \cdot [\xi\tau(x; J) + (1-\xi)\tau(x; -J)]. \quad (9)$$

Expanding the right-hand side term of this expression around $c = 1/2$ up to first order

$$\begin{aligned} c' &= \sqrt{\frac{2k}{\pi}} \int dx \exp(-2k(x-c)^2) \cdot [\xi\tau(x; J) + (1-\xi)\tau(x; -J)] \\ &= \sqrt{\frac{2k}{\pi}} \int dy \exp(-2ky^2) \cdot [\xi\tau(y+c; J) + (1-\xi)\tau(y+c; -J)]. \quad (10) \end{aligned}$$

We denote Eq. (10) by $g(\xi; J)$.

To proceed we need the derivative of τ given by Eq. (4) near $c = 1/2$ and small y . From Eq. (4)

$$f(y; J) = \left. \frac{\partial \tau}{\partial c} \right|_{x=1/2+y} = \frac{4J \exp(-4Jy)}{(1 + \exp(-4Jy))^2}.$$

We note that $f(y; -J) = -f(y; J)$. Then

$$g(\xi; J) = \int dy \sqrt{\frac{2k}{\pi}} \exp(-2ky^2) (2\xi - 1) f(y; J).$$

It is now straightforward to check that

$$g(1 - \xi; J) = g(\xi; -J) = -g(\xi; J).$$

If $g(\xi_a; J) = -1$ then $g(\xi_b) = 1$ with $\xi_b = 1 - \xi_a$ and the two bifurcation points are symmetric with respect to $\xi = 1/2$.

By approximating

$$f(y; J) \simeq J(1 - 4J^2y^2) \simeq J \exp(-4J^2y^2)$$

we get for ξ_a

$$(2\xi_a - 1)J \sqrt{\frac{k}{2J^2 + k}} = -1$$

i.e.,

$$\xi_a(J, k) = \frac{1}{2} \left(1 - \frac{1}{J} \sqrt{1 + \frac{2J^2}{k}} \right).$$

The limit of this last expression when $k \rightarrow \infty$ is $\xi_a(J, \infty) = (1/2)(1 - 1/J)$.

-
- [1] F. Bagnoli, R. Rechtman, Phys. Rev. E, **88** 062914 (2013).
- [2] W. B. Arthur, The American Economic Review **84**, 406–411 (1994); D. Challet, Y.-C. Zhang, Physica A **246**, 407–418 (1997) D. Challet, M. Marsili, Y.-C. Zhang, Minority Games (Oxford University Press 2005); A. C. C. Coolen, The Mathematical Theory of Minority Games (Oxford University Press, 2005).
- [3] S.E. Asch, *Effects of group pressure on the modification and distortion of judgments*, in *Groups, leadership and men*, H. Guetzkow editor (Carnegie Press, Pittsburgh PA, 1951) pp. 177–190; S.E. Asch, *Social psychology*, (Prentice Hall, Englewood Cliffs NJ, 1952); S.E. Asch, *Studies of independence and conformity. A minority of one against a unanimous majority*. Psychological Monographs, **70d**, 1–70 (1956).
- [4] S. Galam, *Contrarian deterministic effects on opinion dynamics: “the hung elections scenario”*, Physica A **333**, 453 (2004).
- [5] S. Galam, *Modeling the Forming of Public Opinion: An approach from Sociophysics*, Global Economics and Management Review **18** 11-20 (2013).
- [6] C. Borghesi, S. Galam, *Chaotic, staggered, and polarized dynamics in opinion forming: The contrarian effect*, Phys. Rev. E **73**, 066118 (2006).
- [7] P.S. Dodds, K.D. Harris, C.M. Danforth, *Limited Imitation Contagion on Random Networks: Chaos, Universality, and Unpredictability*, Phys. Rev. Lett. **110**, 158701 (2013).
- [8] K.D. Harris, P.S. Dodds, C.M. Danforth, *Dynamical influence processes on networks: General theory and applications to social contagion*, Phys. Rev. E **88**, 022816 (2013).
- [9] L. Boltzmann, *Vorlesungen über Gastheorie*, Leipzig, J. A. Barth,; Part I, 1896, Part II, 1898. English translation by S. G. Brush, *Lectures on Gas Theory*, University of California Press, 1964, Chapter I, Sec. 6.
- [10] E. Ott, *Chaos in Dynamical Systems*, Cambridge University Press (1993).
- [11] D.J. Watts and S.H. Strogatz, Nature **393**, 409 (1998).
- [12] L. Barabási, R. Albert, Science **286** 509 (1999).
- [13] M. E. J. Newman, S. H. Strogatz, D. J. Watts, Phys. Rev. E **64**, 026118 (2001).