INFINITE MEASURE PRESERVING TRANSFORMATIONS WITH RADON MSJ

Alexandre I. Danilenko

We introduce concepts of Radon MSJ and Radon disjointness for infinite Radon measure preserving homeomorphisms of the locally compact Cantor space. We construct an uncountable family of pairwise Radon disjoint infinite Chacon like transformations. Every such transformation is Radon strictly ergodic, totally ergodic, asymmetric (not isomorphic to its inverse), has Radon MSJ and possesses Radon joinings whose ergodic components are not joinings.

0. INTRODUCTION

This paper (inspired by a recent progress in [JaRoRu]) is about property of minimal self-joinings (MSJ) for infinite measure preserving transformations. The concept of MSJ for probability preserving maps was introduced in [Rud] as a powerful tool to construct systems with certain prescribed dynamical properties (see also [dJRu], [Ru], [Da2] and references therein for further developments). Some subsequent works are devoted to extension of MSJ to infinite measure preserving and nonsingular systems: [RudSi], [SiWi], [dJSi], [JaRoRu], etc. However such extensions are only partial due to principal obstacles such as the following ones:

- given a continuous transformation of a Polish space with a recurrent point, there exist uncountably many ergodic pairwise orthogonal nonsingular (as well as infinite invariant) measures [Gli], [Ef],
- ergodic components of a non-ergodic joining need not be joinings [Aa1],
- given an ergodic probability preserving transformation S and $\lambda \in (0,1)$, there is en ergodic 2-fold self-joining of S of Krieger's type III_λ [RudSi].

To bypass these obstacles the authors of the aforementioned papers have to select special classes of measures invariant or quasi-invariant under Cartesian powers of the transformations under question. For instance, only boundedly finite infinite measures were considered in [JaRoRu] and mostly rational nonsingular joinings were studied in [RudSi]. We also go this way.

We first develop elements of the general theory of joinings for ergodic measure preserving homeomorphisms T of a locally compact non-compact Cantor space X endowed with an infinite Radon measure μ . This is not a restriction because every ergodic infinite (non-atomic) σ -finite measure preserving transformation on a standard Borel space is measure theoretically isomorphic to a Radon strictly ergodic homeomorphism on X [Yu]. Considering only Radon joinings and self-joinings of ergodic Radon dynamical systems, we introduce notions of Radon MSJ and Radon disjointness. Using the topological nature of X, T and μ and Hopf ratio ergodic theorem we can define generic points for the system.

 T and T

In the remaining part of the paper we present examples of transformations with Radon MSJ. First we consider the *infinite Chacon transformation T*. It was introduced in 1997 by Adams, Friedman and Silva (see [AdFrSi, Section 2]) as an infinite measure preserving counterpart of the classical Chacon map. In a recent paper [JaRoRu] Janvresse, Roy and de la Rue showed that this transformation has MSJ in the class of so-called boundedly finite (in fact, Radon) measures¹ and that there exist ergodic infinite $T^{\times d}$ -invariant measures whose marginals are singular with respect to the original T-invariant measure. Such measures are called *weird* in [JaRoRu]. We provide different (shorter and more algebraic) proofs of these results in Theorem 5.2. While in $[AdFrSi]$ and $[JaRoRu]$, T is a Borel bijection (with countably many discontinuities) of a co-countable subset of the infinite interval $[0, \infty)$ furnished with Lebesgue measure, in this paper T appears as a Radon strictly ergodic homeomorphism of X. In addition to what was done in [JaRoRu] we

- describe explicitly the recurrent points for $T \times T$ (Proposition 4.3),
- describe the weird $T^{\times d}$ -invariant measures (for each $d > 1$) as quasi-graph measures (see Section 3.2),
- introduce an uncountable family of infinite Chacon like Radon strictly ergodic transformations T^{ω} , $\omega \in \{0,1\}^{\mathbb{N}}$ (in Section 6),
- show that each T^{ω} is totally ergodic, has Radon MSJ and is not isomorphic to the inverse $(T^{\omega})^{-1}$ (Theorem 6.1, Corollary 6.2),
- prove that T^{ω} and $T^{\omega'}$ are isomorphic if the infinite sequences ω and ω' are tail equivalent and that T^{ω} and $T^{\omega'}$ are Radon disjoint otherwise (Theorem 6.1),
- give an example of nonergodic Radon 2-fold joining of T such that almost all ergodic components of this joining are conservative quasi-graphs whose coordinate projections are singular to the original T-invariant measure (Example 5.4).

The outline of the paper is as follows. In Section 1 we introduce some basic notions for Radon dynamical systems such as recurrent points and generic points and discuss ergodic decompositions. Radon joinings and Radon disjointness for general Radon dynamical systems are studied in Section 2. In Section 3 we recall the (C, F) -construction of rank-one actions for Abelian groups. We introduce their quasi-graph invariant measures and study their properties. The (C, F) -construction is used in Section 4 to define the infinite Chacon transformation T. The recurrent points for the Cartesian square of T are explicitly described there. In Section 5 we show that T has Radon MSJ. As a byproduct we describe all $T^{\times d}$ -invariant measures on X^d . An example of a Radon joining of T whose ergodic components are not joinings is also given in this section. In Section 6 we introduce and study the infinite Chacon like transformations. In the final Section 7 we discuss possible generalizations of the class of dynamical systems to which the main results of this paper extend. Some open problems are also stated there.

Acknowledgements. I thank the anonymous referee for useful remarks.

¹This extends the well known result that the classical Chacon has MSJ [dJRaSw].

1. Basic definitions: recurrent points, generic points, ergodic decomposition

Let X be a locally compact non-compact Cantor space and let T be an aperiodic homeomorphism of X.

Definition 1.1. A point $x \in X$ is called T-recurrent if for each neighborhood U of x there is an integer $n \neq 0$ such that $T^n x \in U$.

We are interested in T-invariant (non-negative) σ -finite Borel measures on X. Since X is non-compact, T -invariant probability measures on X need not exist. On the other hand, it follows from $\lbrack\text{Gli}\rbrack$ and $\lbrack\text{Sc}\rbrack$ that if there exists a T-recurrent point then there are uncountably many non-equivalent infinite T-invariant σ -finite Borel measures on X. In this work our main concern is Radon measures. We recall that a Borel measure λ on X is called Radon if it is finite on every compact subset of X. Of course, every Radon measure is σ -finite. The converse is not true. Denote by $\mathcal{M}_{\text{Ra}}(X)$ the cone of all Radon measures on X. It is a Polish space in the ∗-weak topology. Let $\mathcal{M}_{\text{Ra}}(X,T)$ stand for the subset of T-invariant Radon measures on X and let $\mathcal{M}_{\mathrm{Ra}}^e(X,T)$ denote the subset of ergodic T-invariant Radon measures on X. The two subsets are Borel. Given a pair of measures $\lambda, \lambda' \in \mathcal{M}_{\text{Ra}}^e(X,T)$, we have that either $\lambda = c\lambda'$ for some real $c > 0$ or $\lambda \perp \lambda'$. If $\mathcal{M}_{\text{Ra}}(X,T)$ consists of a single ray we say that T is Radon uniquely ergodic. If, in addition, T is minimal then we say that T is Radon strictly ergodic.

Definition 1.2. Let $\lambda \in \mathcal{M}_{\text{Ra}}^e(X,T)$. A point $x \in X$ is called *generic* for λ if for each pair of compact open subsets $K_1, K_2 \subset X$ such that $\lambda(K_2) > 0$, the following limit exists and is given by

$$
\lim_{n \to \infty} \frac{\sum_{a_n \le j \le b_n} 1_{K_1}(T^j x)}{\sum_{a_n \le j \le b_n} 1_{K_2}(T^j x)} = \frac{\lambda(K_1)}{\lambda(K_2)}
$$

whenever $a_n \leq 0 \leq b_n$ and $b_n - a_n \to +\infty$.

If λ is non-atomic (i.e. the dynamical system (X, λ, T) is conservative) then every generic point is recurrent. Denote by $\mathcal{G}(\lambda)$ the set of all generic points for λ. Then $G(\lambda)$ is a Borel subset of X and $TG(\lambda) = G(\lambda)$ (pointwise, not only almost everywhere). By the Hopf ratio ergodic theorem $[Aa]$, λ -almost every point of X is generic for λ^2 . If λ' is another ergodic T-invariant Radon measure then $\mathcal{G}(\lambda) = \mathcal{G}(\lambda')$ if $\lambda' = c\lambda$ for some $c > 0$ and $\mathcal{G}(\lambda) \cap \mathcal{G}(\lambda') = \emptyset$ otherwise.

Let $\eta \in \mathcal{M}_{\text{Ra}}(X,T)$. Then there is a probability measure κ on $\mathcal{M}_{\text{Ra}}^e(X,T)$ such that $\eta = \int_{\mathcal{M}_{\text{Ra}}^e(X,T)} \lambda \, d\kappa(\lambda)$. This integral is called an *ergodic decomposition* of η . It is not unique.

2. Radon self-joinings and Radon disjointness

Let X be a locally compact Cantor space, G a discrete countable infinite Abelian group, $T = (T_q)_{q \in G}$ a continuous action of G on X and μ a T-invariant Radon measure on X. We call the triple (X, T, μ) Radon dynamical system. From now on we assume that μ is ergodic.

²This theorem was stated there only for conservative systems. It is also true for dissipative (ergodic, Radon) ones with a trivial proof.

Definition 2.1. Let $d > 1$. A Radon $((T_g)^{\times d})_{g \in G}$ -invariant measure on X^d whose coordinate projections (marginals) are all equivalent³ to μ is called a *d-fold Radon* self-joining of T. The set of all Radon d-fold self-joinings of T will be denoted by $J_{d,\text{Ra}}(T)$. The subset of ergodic Radon d-fold self-joinings of T will be denoted by $J_{d,\text{Ra}}^e(T).$

For instance, the product $\mu \otimes \cdots \otimes \mu(d \text{ times})$ belongs to $J_{d, \text{Ra}}(T)$.

Denote by $C(T)$ the centralizer of T, i.e. the group of all invertible μ -nonsingular transformations commuting with T_g for each $g \in G$. If $S_1, \ldots, S_{d-1} \in C(T)$ then there are constants $c_i > 0$ such that $\mu \circ S_i = c_i \mu$ for all $i = 1, \ldots, d^4$. The measure $\mu_{S_1,...,S_{d-1}}$ on X^d , given by

$$
\mu_{S_1,\ldots,S_{d-1}}(A_1 \times \cdots \times A_d) := \mu(A_1 \cap S_1^{-1}A_2 \cap \cdots \cap S_{d-1}^{-1}A_d)
$$

for all Borel subsets $A_1, \ldots, A_d \subset X$, belongs to $J_{d, \text{Ra}}^e(T)$. Since $\lambda_{S_1, \ldots, S_{d-1}}$ is supported on the graph of the map $X \ni x \mapsto (S_1x, \ldots, S_{d-1}x) \in X^{d-1}$, it is called a graph-joining of T.

Definition 2.2. If for each $\lambda \in J_{d,\text{Ra}}^e(T)$ there is a partition of $\{1,\ldots,d\}$ into subsets J_1, \ldots, J_k such that λ splits into direct product of its marginals λ_i on X^{J_i} , each λ_i is (up to multiplicative constant) a graph-joining of T if $\#J_i > 1$ or $\lambda_i = \mu$ if $\#J_i = 1, i = 1,\ldots,k$, and $C(T) = \{T_g | g \in G\}$ then we say that T has the property of Radon MSJ_d . If T has Radon MSJ_d for each $d > 1$ then we say that T has Radon MSJ.

We note that if λ is a non-ergodic Radon self-joining of T then the ergodic components of λ need not be self-joinings of T. For instance, there exist Radon strictly ergodic homeomorphism R of X with a T-invariant Radon measure μ such that the system $(X \times X, \mu \times \mu, R \times R)$ is totally dissipative [Da1, Theorem 0.1(4)], i.e. almost every ergodic component of the Radon 2-fold self-joining $\mu \times \mu$ of R is an $(R \times R)$ -orbit equipped with the "counting" purely atomic measure. Purely atomic measures are not self-joinings of R because their coordinate projections are also atomic and hence singular to μ which is continuous.

We also need the following analogue of the Furstenberg concept of disjointness for probability preserving systems [Fu].

Definition 2.3. Let $(X,(T_g)_{g\in G},\mu)$ and $(Y,(S_g)_{g\in G},\nu)$ be two ergodic Radon dynamical systems. We say that they are Radon disjoint if $\mu \times \nu$ is the only (up to a multiplicative constant) ergodic Radon $(T_q \times S_q)_{q \in G}$ -invariant measure on $X \times Y$ whose coordinate projections are equivalent to μ and ν respectively.

Of course, if T and S are Radon disjoint then they are not isomorphic.

3. (C, F) -CONSTRUCTION OF RANK-ONE ABELIAN ACTIONS

3.1. (C, F) -nomenclature. For a detailed exposition of the (C, F) -construction we refer the reader to [Da1] and [Da3].

Let G be a discrete countable infinite Abelian group. Let $(F_n)_{n>0}$ and $(C_n)_{n\geq 1}$ be two sequences of finite subsets in G such that for each $n > 0$,

(I) $F_0 = \{0\}, \#C_n > 1,$

³We note that these projections are not necessarily σ -finite.

- (II) $F_0 \subset F_1 \subset \cdots$ with $\bigcup_{n \geq 0} F_n = G$,
- (III) $F_n + F_n + C_{n+1} \subset F_{n+1}$,

(IV) $(F_n + c) \cap (F_n + c') = \emptyset$ for all $c, c' \in C_{n+1}$ with $c \neq c'$.

We let $X_n := F_n \times C_{n+1} \times C_{n+2} \times \cdots$ and endow this set of infinite sequences with the infinite product topology. Then X_n is a compact Cantor (i.e. totally disconnected perfect metric) space. The mapping

$$
X_n \ni (f_n, c_{n+1}, c_{n+2}, \dots) \mapsto (f_n + c_{n+1}, c_{n+2}, \dots) \in X_{n+1}
$$

is a topological embedding of X_n into X_{n+1} . From now on we can consider X_n an a subset of X_{n+1} . Let X denote the inductive limit of the sequence $(X_n)_{n>0}$ furnished with these embeddings. Then X is a well defined locally compact Cantor space. We call it the (C, F) -space associated with the sequence $(C_n, F_{n-1})_{n>1}$. Given a subset $A \subset F_n$, we let

$$
[A]_n := \{ x = (f_n, c_{n+1}, \dots) \in X_n \mid f_n \in A \}
$$

and call this set an *n-cylinder* in X. It is open and compact in X. It is easy to verify that

$$
[A]_n \cap [B]_n = [A \cap B]_n, \quad [A]_n \cup [B]_n = [A \cup B]_n \quad \text{and}
$$

$$
[A]_n = [A + C_{n+1}]_{n+1}
$$

for all $A, B \subset F_n$ and $n \geq 0$. For brevity, we will write $[f]_n$ for $[\{f\}]_n, f \in F_n$. We note that the collection of all cylinders coincides with the family of all compact open subset in X.

Let R denote the *tail equivalence relation* on X. This means that the restriction of R to X_n is the tail equivalence relation on X_n for each $n \geq 0$. We note that $\mathcal R$ is minimal, i.e. the $\mathcal R$ -class of every point is dense in X. There exists a unique σ-finite R-invariant Borel measure μ on X such that $\mu(X_0) = 1$. It is Radon. Moreover, μ is strictly positive on every non-empty open subset. We note that the $\mathcal{R}\text{-}invariance^5$ of μ is equivalent to the following property:

$$
\mu([f]_n) = \mu([f']_n) \quad \text{ for all } f, f' \in F_n, n \ge 0.
$$

This and the normalization condition on μ imply that

$$
\mu([A]_n) = \frac{\#A}{\#C_1 \cdots \#C_n} \quad \text{for each subset } A \subset F_n, \ n > 0.
$$

We call μ the (C, F) -measure associated with $(C_n, F_{n-1})_{n \geq 1}$. It is infinite if and only if 6

(3-1)
$$
\lim_{n \to \infty} \frac{\#F_n}{\#C_1 \cdots \#C_n} = \infty.
$$

 ${}^5\mu$ is called R-invariant if μ is invariant under each Borel transformation whose graph is contained in R.

⁶In view of (I)–(IV), the sequence $(\frac{\#F_n}{\#C_1\cdots\#C_n})_{n=1}^{\infty}$ is non-decreasing and bounded by 1 from below.

It is easy to see that μ is \mathcal{R} -ergodic, i.e. each Borel \mathcal{R} -saturated subset of X is either μ -null or μ -conull. We now define an action of G on X. Given $g \in G$, let

$$
X_n^g := \{ (f_n, c_{n+1}, c_{n+2} \dots) \in X_n \mid g + f_n \in F_n \}.
$$

Then X_n^g is a compact open subset of X_n and $X_n^g \subset X_{n+1}^g$. In view of (III) and (II), $X_n \subset X_{n+1}^g$ eventually in n. Hence $\bigcup_{n\geq 0} X_n^g = X$. Therefore, given $x \in X$ and $g \in G$, there is $n > 0$ such that if we consider an expansion $x = (f_n, c_{n+1}, \dots)$ of x in X_n then $g + f_n \in F_n$. We now let

$$
T_g x := (g + f_n, c_{n+1}, \dots) \in X_n \subset X.
$$

It is standard to verify that

- (i) the map $T_q : X \ni x \mapsto T_q x \in X$ is a homeomorphism of X and
- (ii) $T_gT_{g'} = T_{g+g'}$ for all $g, g' \in G$.

Thus $T := (T_q)_{q \in G}$ is a continuous action of G on X. We call it the (C, F) -action of G associated with the sequence $(C_n, F_{n-1})_{n \geq 1}$. It is free. The T-orbit equivalence relation is R. Indeed, if $(x, x') \in \mathcal{R}$ then there is $n > 0$ such that $x, x' \in X_n$. Consider now the expansions $x = (f_n, c_{n+1}, \dots)$ and $x' = (f'_n, c'_{n+1}, \dots)$ of x and x' in X_n . Then $x' = T_g x$, where $g := f'_n - f_n + \sum_{j>n} (c'_j - c_j)$. It follows that T preserves μ . We call (X, μ, T) the dynamical system associated with $(C_n, F_{n-1})_{n \geq 1}$. It is Radon strictly ergodic⁷.

We will need the following standard lemma (see $[Da4, Lemma 1.1(ii)]$ for the proof of a more general fact).

Lemma 3.1. Let $g \in G$. If there is $\delta > 0$ such that for each $n \in \mathbb{N}$ and $v, w \in F_n$, there is a subset $A \subset [v]_n$ and $m \in \mathbb{Z}$ such that $T_{mg}A \subset [w]_n$ and $\mu(A) \geq \delta \mu([v]_n)$ then the transformation T_g is ergodic.

3.2. Quasi-graph invariant Radon measures. In this subsection we introduce an important class of Radon $((T_g)^{\times d})_{g \in G}$ -invariant measures on X^d which are "close" to graph-joinings but whose marginals can be singular to μ . Such measures can appear as ergodic components of non-ergodic Radon d-fold self-joinings of T (see §5 and §6). Throughout this subsection we will assume that the following condition holds:

(V) $F_n + C_{n+1} + F_n - F_n \subset F_{n+1}$ for each $n > 0$.

This condition is stronger than (III). Let J a subset of N. For each $n \notin J$, fix an element $\alpha_n \in C_n$. Thus we obtain a sequence $\alpha := (\alpha_n)_{n \notin J}$. We now define a new (C, F) -sequence $(C_n^{\alpha}, F_{n-1}^{\alpha})_{n \geq 1}$ by setting $F_n^{\alpha} := F_n$ and

$$
C_n^{\alpha} := \begin{cases} C_n, & \text{if } n \in J \\ \{\alpha_n\}, & \text{if } n \notin J. \end{cases}
$$

Then $(C_n^{\alpha}, F_{n-1}^{\alpha})_{n\geq 1}$ satisfies (I)–(V) except for the fact that the cardinality of C_n^{α} can be 1. This happens if and only if $n \notin J$. Nevertheless the associated (C, F) action $T^{\alpha} = (T_g^{\alpha})_{g \in G}$ on a locally compact (C, F) -space X^{α} equipped with an invariant σ -finite measure μ^{α} is well defined (via the same inductive construction

⁷We also note that T has rank one along the sequence $(F_n)_{n=1}^{\infty}$ (in the measure theoretical sense, see [Da1]).

process described in §3.1). The space X^{α} is totally disconnected. It is either perfect (i.e. Cantor) or purely discrete and countable. The latter happens if and only if J is finite. We note that μ^{α} is finite if and only if μ is finite and J is cofinite⁸. Given $A \subset F_n$, the corresponding cylinder in X^{α} will de denoted by $[A]_n^{\alpha}$ (in order to not confuse it with the cylinder $[A]_n$ in X). We now define an embedding ϕ_α of X^α into X by setting for each $n > 0$ and a point $x = (f_n, c_{n+1}, c_{n+2}, \dots) \in X_n^{\alpha} \subset X^{\alpha}$,

$$
\phi_{\alpha}(x) := (f_n, c_{n+1}, c_{n+2}, \dots) \in X_n \subset X.
$$

It is straightforward to verify that ϕ_{α} is a well-defined, one-to-one continuous equivariant⁹ map. Hence the image $\phi^{\alpha}(X^{\alpha})$ of X^{α} is a T-invariant dense F_{σ} -subset of X. Now given $m > 0$ and a subset $A \subset F_m$, take a point $x \in [A]_m \cap \phi_\alpha(X^\alpha)$. If we write the expansion $x = (f_m, c_{m+1}, c_{m+2}, \dots)$ of x in X_m then $f_m \in A$ and $c_l = \alpha_l$ for all $l \notin J$ that are greater than some $n > m$. It follows that

(3-2)
$$
[A]_m \cap \phi_\alpha(X^\alpha) = \bigsqcup_{g \in \sum_{J \not\ni l > m} (C_l - \alpha_l)} T_g \phi_\alpha([A]_m^\alpha).
$$

We have the following trichotomy:

- if J is cofinite then $\mu_\alpha \circ \phi_\alpha^{-1}$ is a multiple of μ and hence ϕ_α is an isomorphism (both topological and measure theoretical) of $(X^{\alpha}, \mu^{\alpha}, T^{\alpha})$ onto (X, μ, T) ;
- if J is finite then $\mu_{\alpha} \circ \phi_{\alpha}^{-1} \perp \mu$ and $(X^{\alpha}, \mu^{\alpha}, T^{\alpha})$ is an ergodic totally dissipative dynamical system;
- if *J* is neither finite nor cofinite then $\mu_\alpha \circ \phi_\alpha^{-1} \perp \mu$ and $(X^\alpha, \mu^\alpha, T^\alpha)$ is an ergodic conservative dynamical system.

Consider now another sequence $\beta = (\beta_n)_{n \notin J}$ with $\beta_n \in C_n$ for each $n \notin J$. Let $(X^{\beta}, \mu^{\beta}, T^{\beta})$ be the dynamical system associated with $(C_n^{\beta}, F_{n-1}^{\beta})_{n \geq 1}$, where C_n^{β} and F_n^{β} n_{n-1}^{β} are defined in a similar way as C_n^{α} and F_{n-1}^{α} . Then there is a natural measure preserving topological equivariant isomorphism $\phi_{\alpha,\beta}$ of $(X^{\alpha}, \mu^{\alpha}, T^{\alpha})$ onto $(X^{\beta}, \mu^{\beta}, T^{\beta})$. To define $\phi_{\alpha,\beta}$ we take a point $x \in X_{n-1}^{\alpha}$ for some $n > 0$ and write the expansion of x in X_n^{α} : $x = (f_n, c_{n+1}, c_{n+2}, ...) \in F_n \times C_{n+1}^{\alpha} \times C_{n+2}^{\alpha} \cdots$. Then $f_n \in F_{n-1} + C_n^{\alpha}$. We now set

$$
\phi_{\beta,\alpha}(x) := (\widetilde{f}_n, \widetilde{c}_{n+1}, \widetilde{c}_{n+2}, \dots),
$$

where $\widetilde{f}_n := f_n + \sum_{J \neq j \leq n} (\beta_j - \alpha_j)$ and for each $l > n$,

$$
\widetilde{c}_l := \begin{cases} c_l, & \text{if } l \in J \\ \beta_l, & \text{if } l \notin J. \end{cases}
$$

We note that if $n \in J$ then $\sum_{J \neq j \leq n} (\beta_j - \alpha_j) \in F_{n-1} - F_{n-1}$ and hence $\tilde{f}_n \in$ $f_n + F_{n-1} - F_{n-1} \subset F_{n-1} + C_n^{\alpha} + F_{n-1} - F_{n-1}$. If $n \notin J$ then $\sum_{J \not\ni j \leq n} (\beta_j - \alpha_j) \in$ $\beta_n - \alpha_n + F_{n-1} - F_{n-1}$ and hence $f_n \in F_{n-1} + C_n^{\alpha} + \beta_n - \alpha_n + F_{n-1} - F_{n-1} =$ $F_{n-1} + C_n^{\beta} + F_{n-1} - F_{n-1}$. Therefore in every of the two possible cases $f_n \in F_n$

⁸I.e. the complement $\mathbb{N} \setminus J$ is finite.

⁹I.e. $\phi^{\alpha} \circ T_g^{\alpha} = T_g \circ \phi^{\alpha}$ for each $g \in G$.

in view of (V). Hence $\phi_{\alpha,\beta}(x) \in X_n^{\beta}$. Thus $\phi_{\alpha,\beta}$ is well defined. Moreover, for each subset $A \subset F_{n-1} + C_n^{\alpha}$,

(3-3)
$$
\phi_{\beta,\alpha}([A]_n^{\alpha}) = [\widetilde{A}]_n^{\beta}, \text{ where } \widetilde{A} := A + \sum_{J \not\ni j \leq n} (\beta_j - \alpha_j).
$$

It is straightforward to verify that $\phi_{\beta,\alpha}$ is as claimed. It is easy to see that the converse $\phi_{\beta,\alpha}^{-1}$ to $\phi_{\beta,\alpha}$ is $\phi_{\alpha,\beta}$. We now define a map $\psi_{\alpha,\beta}$ from X^{α} to $X \times X$ by setting

$$
\psi_{\alpha,\beta}(x) := (\phi_{\alpha}(x), \phi_{\beta}(\phi_{\beta,\alpha}(x))).
$$

It is one-to-one, continuous and $\psi_{\alpha,\beta} \circ T_g^{\alpha} = (T_g \times T_g) \circ \psi_{\alpha,\beta}$ for all $g \in G$.

Theorem 3.2. Suppose that for each $l \notin J$,

(3-4)
$$
(C_l - \alpha_l + F_{l-1}^{\bullet}) \cap (C_l - \beta_l + F_{l-1}^{\bullet}) = F_{l-1}^{\bullet},
$$

where $F_{l-1}^{\bullet} := F_{l-2} + C_{l-1} - F_{l-2} - C_{l-1}$. Then the following are satisfied.

- (i) The subset $\psi_{\alpha,\beta}(X^{\alpha})$ is closed in $X \times X$. Hence $\psi_{\alpha,\beta}(X^{\alpha})$ is locally compact in the induced topology.
- (ii) The topological dynamical system $(\psi_{\alpha,\beta}(X^{\alpha}), (T_{\alpha} \times T_{\alpha})_{\alpha \in G})$ is Radon strictly ergodic.
- (iii) $\mu^{\alpha} \circ \psi_{\alpha,\beta}^{-1}$ is the only (up to scaling) $T_1 \times T_1$ -invariant Radon measure supported on $\psi_{\alpha,\beta}(X^{\alpha})$.
- (iv) Given $a, b \in F_m$, represent the cylinder $[a]_m \times [b]_m$ in $X \times X$ as the infinite product $\{(a, b)\}\times (C_{m+1}\times C_{m+1})\times (C_{m+2}\times C_{m+2})\times \cdots$. Then

$$
\mu^{\alpha} \circ \psi_{\alpha,\beta}^{-1} \upharpoonright ([a]_m \times [b]_m) = \mu^{\alpha}([a]_m^{\alpha}) \cdot \widehat{\delta}_{a',b'} \otimes \bigotimes_{n>m} \gamma_n,
$$

where γ_m is a probability on $C_m \times C_m$ given by

$$
\gamma_n := \begin{cases} \frac{1}{\#C_n} \sum_{c \in C_n} \delta_c \times \delta_c, & \text{if } n \in J \\ \delta_{(\alpha_n, \beta_n)}, & \text{if } n \notin J, \end{cases}
$$

$$
a' = a + \sum_{J \not\ni j \le m} \alpha_j \text{ and } b' = b + \sum_{J \not\ni j \le m} \beta_j.^{10}
$$

Proof. (i) If J is cofinite then it is straightforward to verify that $\psi_{\alpha,\beta}(X_{\alpha})$ is the the graph of T_h , where $h := \sum_{j \notin J} (\beta_j - \alpha_j)$. The graph is closed in $\overline{X} \times \overline{X}$.

Consider now another particular case where J is finite. Then it is straightforward to verify that $\psi_{\alpha,\beta}(X_\alpha)$ is the $(T_g \times T_g)_{g \in G}$ -orbit of a point $(x,y) \in X \times X$ such that $x = (f_m, \alpha_{m+1}, \alpha_{m+2}, \dots) \in X_m$ and $y = (r_m, \beta_{m+1}, \beta_{m+2}, \dots) \in X_m$ for some $m \geq 0$ and elements $f_m, r_m \in F_m$. It suffices to show that given $l > m$ and subsets $A, B \subset F_l$, the intersection of the orbit of (x, y) with the cylinder $[A]_l \times [B]_l$ is finite. Indeed, $(T_gx, T_gy) \in [A]_l \times [B]_l$ if and only if

$$
g \in A - f_m - \alpha_{m+1} - \dots - \alpha_l + \sum_{i>l} (C_i - \alpha_i) \text{ and}
$$

$$
g \in B - r_m - \beta_{m+1} - \dots - \beta_l + \sum_{i>l} (C_i - \beta_i).
$$

¹⁰Here and below δ_c or $\delta_{(a,b)}$ denotes the probability measure supported at a single point $\{c\}$ or $\{(a, b)\}\)$ respectively while $\hat{\delta}_{a', b'}$ denotes the Kronecker delta.

Hence $g \in F_{l+1}^{\bullet} + \sum_{i>l+1} (C_i - \alpha_i)$ and $g \in F_{l+1}^{\bullet} + \sum_{i>l+1} (C_i - \beta_i)$. It follows from (3-4) that $(F_{l+1}^{\bullet} + \sum_{i>l+1} (C_i - \alpha_i)) \cap (F_{l+1}^{\bullet} + \sum_{i>l+1} (C_i - \beta_i)) = F_{l+1}^{\bullet}$. It remains to note that F_{l+1}^{\bullet} is finite.

Suppose now that J is neither finite nor cofinite. Fix $n > 0$ and subsets $A \subset$ $F_{n-1} + C_n$ and $B \subset F_{n-1} + C_n^{\beta}$. Applying (3-2) we obtain that

$$
\psi_{\alpha,\beta}^{-1}([A]_n \times [B]_n) \subset \phi_{\alpha}^{-1}([A]_n) \cap \phi_{\beta,\alpha}^{-1}(\phi_{\beta}^{-1}([B]_n))
$$
\n
$$
= \left(\bigcup_{g \in \sum_{J \not\exists l > n} (C_l - \alpha_l)} T_g^{\alpha}[A]_n^{\alpha}\right) \cap \phi_{\alpha,\beta} \left(\bigcup_{g \in \sum_{J \not\exists l > n} (C_l - \beta_l)} T_g^{\beta}[B]_n^{\beta}\right)
$$
\n
$$
= \left(\bigcup_{J \not\exists l > n} [A + C_{n+1} + \dots + C_l]_l^{\alpha}\right) \cap \phi_{\alpha,\beta} \left(\bigcup_{J \not\exists l > n} [B + C_{n+1} + \dots + C_l]_l^{\beta}\right)
$$
\n
$$
= \bigcup_{J \not\exists l > n} ([A + C_{n+1} + \dots + C_l]_l^{\alpha} \cap \phi_{\alpha,\beta}([B + C_{n+1} + \dots + C_l]_l^{\beta})).
$$

Given $l \notin J$, let l' be the least integer such that $l' > l$ and $l' \notin J$. Then $C_k^{\alpha} = C_k^{\beta} =$ C_k if $l < k < l'$, $C_{l'}^{\alpha} = {\alpha_{l'}}$, $C_{l'}^{\beta}$ $\ell_l^{\beta} = {\beta_{l'}}$ and hence

$$
[A + C_{n+1} + \dots + C_l]_l^{\alpha} = [A' + \alpha_{l'}]_{l'}^{\alpha} \text{ and}
$$

$$
[B + C_{n+1} + \dots + C_l]_l^{\beta} = [B' + \beta_{l'}]_{l'}^{\beta},
$$

where $A' := A + C_{n+1} + \cdots + C_{l'-1} \subset F_{l'-1}$ and $B' := B + C_{n+1} + \cdots + C_{l'-1} \subset F_{l'-1}$. Utilizing this and $(3-3)$ we obtain that 11

$$
\psi_{\alpha,\beta}^{-1}([A]_n \times [B]_n) \subset \bigsqcup_{J \not\ni l > n} \left([A' + \alpha_{l'}]_{l'}^{\alpha} \cap \left[B' + \beta_{l'} + \sum_{J \not\ni j \le l'} (\alpha_j - \beta_j) \right]_{l'}^{\alpha} \right)
$$
\n
$$
= \bigsqcup_{J \not\ni l > n} \left(\left[\left(A' \cap \left(B' + \sum_{J \not\ni j \le l} (\alpha_j - \beta_j) \right) \right) + \alpha_{l'} \right]_{l'}^{\alpha} \right).
$$

It follows from (3-4) that for all subsets $D, D' \subset F_n^{\bullet}$,

$$
\left(D+\sum_{j=n+1}^{l'-1}C_j-\sum_{n
$$

¹¹In the formulas below, $l' = l'(l)$.

Therefore we have

$$
A' \cap \left(B' + \sum_{J \not\ni j \le l} (\alpha_j - \beta_j) \right)
$$

=
$$
\left(\left(A + \sum_{j=n+1}^{l'-1} C_j - \sum_{j
=
$$
\left(\left(\hat{A} + \sum_{j=n+1}^{l'-1} C_j - \sum_{n
=
$$
(\hat{A} \cap \hat{B}) + \sum_{n
=
$$
(A \cap \tilde{B}) + \sum_{n
$$
$$
$$
$$

where $\widehat{A} := A - \sum_{J \not\in j \leq n} \alpha_j \subset F_{n-1} + C_n - F_{n-1} - C_n = F_n^{\bullet}, \widehat{B} := B - \sum_{J \not\in j \leq n} \beta_j \subset F_n$ F_n^{\bullet} and $\widetilde{B} := B + \sum_{J \not\ni j \leq n} (\alpha_j - \beta_j)$. Hence

$$
\psi_{\alpha,\beta}^{-1}([A]_n \times [B]_n) \subset \bigsqcup_{J \not\ni l > n} \left[(A \cap \widetilde{B}) + \sum_{n < j \le l'} C_j^{\alpha} \right]_{l'}^{\alpha} = [A \cap \widetilde{B}]_n^{\alpha}.
$$

On the other hand, it is straightforward to verify that the converse inclusion holds. Therefore,

(3-5)
$$
\psi_{\alpha,\beta}^{-1}([A]_n \times [B]_n) = [A \cap \widetilde{B}]_n^{\alpha}.
$$

It follows that the $\psi_{\alpha,\beta}$ -preimage of $[A]_n \times [B]_n$ is compact in X^{α} . Hence $[A]_n \times [B]_n$ does not intersect the set $\psi_{\alpha,\beta}(X_{\alpha}) \setminus \psi_{\alpha,\beta}(X_{\alpha})$. Since J is not finite then we take $A := B := F_{n-1} + C_n$ for $n \in J$ and obtain that the union

$$
\bigcup_{n \in J} ([F_{n-1} + C_n]_n \times [F_{n-1} + C_n]_n) = \bigcup_{n \in J} (X_{n-1} \times X_{n-1}) = X \times X
$$

does not intersect $\overline{\psi_{\alpha,\beta}(X_\alpha)} \setminus \psi_{\alpha,\beta}(X_\alpha)$. Hence $\psi_{\alpha,\beta}(X_\alpha)$ is closed.

(ii) It follows from (i) that ψ is an equivariant topological isomorphism of the dynamical system (X^{α}, T^{α}) onto $(\psi_{\alpha,\beta}(X^{\alpha}), (T_g \times T_g)_{g \in G})$. Since the former system is Radon strictly ergodic, so is the latter.

(iii) is obvious now.

(iv) Let m, a, b be the same as in the statement of (iv). Suppose that J is infinite. Take $n > m$ with $n \in J$ and elements $c_j, \hat{c}_j \in C_j$ whenever $m < j \leq n$. Then by $(3-5)$,

$$
\psi_{\alpha,\beta}^{-1}\left(\left[a+\sum_{m
=
$$
\left[\left\{a+\sum_{m
$$
$$

Therefore, by (3-4), the intersection in the above formula is non-empty if and only if $a = b + \sum_{J \not\ni j \le m} (\alpha_j - \beta_j)$, $c_j = \widehat{c}_j$ if $J \ni j$ and $c_j = \alpha_j$ and $\widehat{c}_j = \beta_j$ if $J \not\ni j$. Hence

$$
\mu^{\alpha}\left(\psi_{\alpha,\beta}^{-1}\left(\left[a+\sum_{j=m+1}^{n}c_j\right]_n\times\left[b+\sum_{j=m+1}^{n}\widehat{c}_j\right]_n\right)\right)=\mu^{\alpha}([a]_m^{\alpha})\cdot\widehat{\delta}_{a',b'}\prod_{j=m+1}^{n}\gamma_j(c_j,\widehat{c}_j),
$$

where a', b' and γ_j were introduced in the statement of (iv). \Box

Definition 3.3. Given a subset $J \subset \mathbb{N}$, two sequences $\alpha = (\alpha_n)_{n \notin J}$ and $\beta =$ $(\beta_n)_{n\in\mathcal{J}}$ with $\{\alpha_n,\beta_n\}\subset C_n$ for each $n\not\in J$ and an element $h\in G$, we call the measure $(\mu^{\alpha} \circ \psi_{\alpha,\beta}^{-1}) \circ (I \times T_{h}^{-1})$ h^{-1}) the *quasi-graph* measure.

It follows from Theorem 3.2 that if (3-4) is satisfied then $(\mu^{\alpha} \circ \psi_{\alpha,\beta}^{-1}) \circ (I \times$ T_h^{-1} $(h_h^{-1}) \in \mathcal{M}_{\text{Ra}}^e(X \times X, (T_g \times T_g)_{g \in G})$. The projection of this measure to the first coordinate is equivalent to μ^{α} and the projection of this measure to the second coordinate is equivalent to μ^{β} . The conditional measures corresponding to the disintegrations of $(\mu^{\alpha} \circ \psi_{\alpha,\beta}^{-1}) \circ (I \times T_{h}^{-1})$ μ_h^{-1}) over μ^{α} (or over μ^{β}) are delta-measures almost everywhere. The quasi-graph Radon measure is a 2-fold self-joining of T if and only if $\mu^{\alpha} \sim \mu^{\beta} \sim \mu$. This happens if and only if J is cofinite. In this case the quasi-graph measure is a graph-joining of T^{12} .

In a similar way we may define quasi-graph invariant Radon measures for higher Cartesian powers of T. Indeed, given $d > 1$ and sequences $\alpha^{(i)} = (\alpha^{(i)})_{n \notin J}$, $i =$ $1, \ldots, d$, such that $\alpha_n^{(i)} \in C_n$ for all $i = 1, \ldots, d$, we define a map $\psi_{\alpha^{(1)}, \ldots, \alpha^{(d)}}$ from $X^{\alpha^{(1)}}$ to the d-th Cartesian power of X by setting

$$
\psi_{\alpha^{(1)},...,\alpha^{(d)}}(x) := (\phi_{\alpha^{(1)}}(x), \phi_{\alpha^{(2)}}(\phi_{\alpha^{(2)},\alpha^{(1)}}(x)), \ldots, \phi_{\alpha^{(d)}}(\phi_{\alpha^{(d)},\alpha^{(1)}}(x))).
$$

This map is one-to-one, continuous and equivariant, i.e. $\psi_{\alpha^{(1)},...,\alpha^{(d)}} \circ (T^{\alpha^{(1)}})_1 =$ $(T_1)^{\times d} \circ \psi_{\alpha^{(1)},\dots,\alpha^{(d)}}$. The following theorem is a d-fold analogue of Theorem 3.2. It can be proved in a similar way. Therefore we leave the proof to the reader.

Theorem 3.4. Suppose that for each $l \notin J$, there are $u_l, v_l \in \{1, ..., d\}$ such that

(3-6)
$$
(C_l - \alpha_l^{(u_l)} + F_{l-1}^{\bullet}) \cap (C_l - \alpha_l^{(v_l)} + F_{l-1}^{\bullet}) = F_{l-1}^{\bullet},
$$

where F_{l-1}^{\bullet} is same as in the statement of Theorem 3.2. Then the following are satisfied.

- (i) The subset $\psi_{\alpha^{(1)},...,\alpha^{(d)}}(X^{\alpha^{(1)}})$ is closed in X^d . Hence it is locally compact in the induced topology.
- (ii) The topological dynamical system $(\psi_{\alpha^{(1)},...,\alpha^{(d)}}(X^{\alpha^{(1)}}),((T_g)^{\times d})_{g\in G})$ is Radon strictly ergodic.
- (iii) $\mu^{\alpha^{(1)}} \circ \psi_{\alpha^{(1)}}^{-1}$ $\sigma_{\alpha^{(1)},\ldots,\alpha^{(d)}}^{-1}$ is the only (up to scaling) $((T_g)^{\times d})_{g\in G}$ -invariant Radon measure supported on $\psi_{\alpha^{(1)},...,\alpha^{(d)}}(X^{\alpha^{(1)}})$.

 12 The quasi-graph invariant Radon measures which are not graph-joinings are analogues of the so-called weird measures considered in [JaRoRu] in the case when T is an infinite Chacon transformation.

(iv) Given $a^{(1)}, \ldots, a^{(d)} \in F_m$, represent the cylinder $[a^{(1)}]_m \times \cdots \times [a^{(d)}]_m$ in X^d as the infinite product $\{(a^{(1)}, \ldots, a^{(d)})\} \times C_{m+1}^d \times C_{m+2}^d \times \cdots$. Then

$$
\mu^{\alpha^{(1)}} \circ \psi_{\alpha^{(1)},...,\alpha^{(d)}}^{-1} \upharpoonright ([a^{(1)}]_m \times \cdots \times [a^{(d)}]_m)
$$

=
$$
\mu^{\alpha^{(1)}}([a]_m^{\alpha^{(1)}}) \prod_{i=1}^d \widehat{\delta}_{a_{\bullet}^{(1)},a_{\bullet}^{(i)}} \otimes \bigotimes_{n>m} \gamma_n,
$$

where γ_n is a probability on $C_n \times C_n$ given by

$$
\gamma_n := \begin{cases} \frac{1}{\#C_n} \sum_{c \in C_n} \bigotimes_{i=1}^d \delta_c, & \text{if } n \in J \\ \delta_{(c_n^{(1)}, \dots, c_n^{(d)})}, & \text{if } n \notin J, \end{cases}
$$

 $a_{\bullet}^{(i)}:=a^{(i)}+\sum_{J\not\ni j\leq m}\alpha_{j}^{(i)}$ $j_j^{(i)}$ and elements $c_n^{(i)} \in C_n$, $i=1,\ldots,d,$ are such that $c_n^{(u_n)} = \alpha_n^{(u_n)}$ and $c_n^{(v_n)} = \alpha_n^{(v_n)}$ for each $n > m$.

By analogy with Definition 3.3 we now give the following definition.

Definition 3.5. Given $\alpha^{(1)}, \ldots, \alpha^{(d)}$ as above and let h_1, \ldots, h_{d-1} be elements of G. The measure $\mu^{\alpha^{(1)}} \circ \psi_{\alpha^{(1)},...,\alpha^{(d)}}^{-1} \circ (I \times T_{h_1}^{-1})$ $T_{h_1}^{-1} \times \cdots \times T_{h_{d-1}}^{-1}$ $\binom{n-1}{h_{d-1}}$ is called a *quasi-graph* measure.

It follows from Theorem 3.4 that if (3-6) is satisfied for $\alpha^{(1)}, \ldots, \alpha^{(d)}$ then the corresponding quasi-graph measure is Radon, $((T_g)^{\times d})_{g \in G}$ -invariant and ergodic. In this case a quasi-graph Radon measure is purely atomic if and only if J is finite. On the other hand, a quasi-graph Radon measure is a d -fold Radon joining $(d$ -fold graph-joining indeed) if and only if J is cofinite.

4. Infinite Chacon transformation and recurrent points for its Cartesian square

To define the infinite Chacon transformation we will use the (C, F) -construction. From now on $G = \mathbb{Z}$. We first define recurrently a sequence of positive integers $(h_n)_{n=0}^{\infty}$ by setting $h_0 := 1$ and $h_{n+1} := 6h_n + 1$ for $n > 0^{13}$. For each $n \geq 0$, we now let

$$
F_n := \left\{ -\frac{h_n - 1}{2}, -\frac{h_n - 3}{2}, \dots, \frac{h_n - 1}{2} \right\} \quad \text{and} \quad C_{n+1} := \left\{ -h_n, 0, h_n + 1 \right\}.
$$

The sequence $(C_n, F_{n-1})_{n\geq 1}$ satisfies (I) – (IV) from Section 3 and $(3-1)$. We also note that $#F_n = h_n$ and

(4-1)
$$
F_{n-1} + C_n - F_{n-1} - C_n \subset F_n.
$$

Let $(X, \mu, (T_n)_{n \in \mathbb{Z}})$ denote the dynamical system associated with $(C_n, F_{n-1})_{n \geq 1}$.

¹³In [AdFrSi] and [JaRoRu], it was assumed that $h_{n+1} = 6h_n + 2$. To achieve the "symmetric" version of the infinite Chacon we use one spacer less at each step of the inductive construction. See Section 7 for a discussion about other possibilities to add extra spacers.

Definition 4.1. The transformation T_1 (or, rather, the dynamical system (X, μ, T_1)) is called the infinite Chacon transformation.

For a (more common) cutting-and-stacking definition of the infinite Chacon transformation we refer to [AdFrSi], [JaRoRu] and Section 7 of the present paper.

We will need the following simple lemma.

Lemma 4.2. If $x \in X_{n-1} \cap [f]_n$ and $T_gx \in X_{n-1}$ for some $g, f \in F_n$, then $g + f \in F_n$.

Proof. Indeed, consider expansions $x = (f_{n-1}, c_n, c_{n+1}, ...) \in X_{n-1}$ and $T_g x =$ $(f'_{n-1}, c'_n, c'_{n+1}, \dots) \in X_{n-1}$ of x and $T_g x$ respectively. Then there is $m \ge n$ such that $g = (f'_{n-1} - f_{n-1}) + \sum_{l=n}^{m} (c'_l - c_l)$. Moreover, if $m > n$ we can additionally assume that $c_m \neq c'_m$. It follows from (III) and (4-1) that $g \in F_{m-1}+c'_m-c_m$. Since $g \in F_n \subset F_{m-1}$, we get a contradiction with (IV). Hence $m = n$. Since $x \in [f]_n$, it follows that $f = f_{n-1} + c_n$. This implies that $g = f'_{n-1} + c_n - f \in F_n - f$, as desired. \square

We first describe the recurrent points of the Cartesian square of T_1 .

Proposition 4.3. Let $(x, x') \in X_n \times X_n$ for some $n > 0$. Expand x and x' as $x = (f_n, c_{n+1}, \dots)$ and $x' = (f'_n, c'_{n+1}, \dots).$

- (i) If there is $L > n$ such that $\{c_l, c'_l\} = \{-h_{l-1}, h_{l-1} + 1\}$ for each $l > L$ then the $(T_1 \times T_1)$ -orbit of (x, x') intersects each compact subset of $X \times X$ at most finitely many times.
- (ii) Otherwise (x, x') is $(T_1 \times T_1)$ -recurrent.

Proof. (i) Let $J := \{1, ..., L\}$. We now let $\alpha_l := c_l$ and $\beta_l := c'_l$ for all $n \notin J$. It is easy to see that (3-4) holds. Then by Theorem 3.2(i), the subset $\psi_{\alpha,\beta}(X^{\alpha})$ is closed in $X \times X$. However this subset is a single $(T_1 \times T_1)$ -orbit and the point (x, x') belongs to this orbit. This follows directly from the definition of $\psi_{\alpha,\beta}$ (see §3).

(ii) Let $J := \{l > 0 \mid \{c_l, c'_l\} \neq \{-h_{l-1}, h_{h-1} + 1\}\}\.$ We set $\alpha_l := c_l$ and $\beta_l := c'_l$ for each $l \notin J$. Since J is infinite, X^{α} is a perfect space. It follows now from Theorem 3.2(i) that $\psi_{\alpha,\beta}$ is a perfect subset of $X \times X$. Of course, $(x, x') \in \psi_{\alpha,\beta}(X^{\alpha})$. In view of Theorem 3.2(ii), the $(T_1 \times T_1)$ -orbit of (x, x') is dense in $\psi_{\alpha,\beta}(X^{\alpha})$. Hence (x, x') is $(T_1 \times T_1)$ -recurrent. \Box

Remark 4.4. In a similar way one can prove the following generalization of Proposition 4.3(i). Let $d > 1$ and let $(x^{(1)},...,x^{(d)}) \in X^d$. If there is $L > 0$ such that $\{c_l^{(1)}\}$ $\binom{1}{l},\ldots,\binom{c^{(d)}_{l}}{l}$ $\{a^{(d)}\}$ \supset $\{-h_{l-1}, h_{l-1} + 1\}$ for all $l > L$ then the $(T_1)^{\times d}$ -orbit of $(x^{(1)},...,x^{(d)})$ intersects each compact subset of X^d at most finitely many times. Hence the point $(x^{(1)},...,x^{(d)})$ is not $T_1^{\times d}$ $i_1^{\times d}$ -recurrent. All the other points of X^d are $T_1^{\times d}$ $i_1^{\times d}$ -recurrent.

Corollary 4.5. The counting measure¹⁴ on the $(T_1 \times T_1)$ -orbit of a point $(x, x') \in$ $X\times X$ is a Radon measure on $X\times X$ if and only if the condition of Proposition 4.3(i) is satisfied.

We also show that T_1 is totally ergodic.

¹⁴This measure is infinite, σ -finite, $(T_1 \times T_1)$ -invariant, ergodic but non-conservative.

Proposition 4.6. If $m > 1$ then the transformation T_m is ergodic.

Proof. Let $k > 0$. We set $h := h_k + h_{k+1} + \cdots + h_{k+m-1}$. Take $f, f' \in F_k$. There is $j \in \{0, 1, \ldots, m-1\}$ such that $f' - f + h + j$ is divisible by m. Take $f, f' \in F_k$ Then $[f - h_{k+j} - \cdots - h_{k+m-1}]_{k+m} \subset [f]_k$,

$$
T_{h+j+f'-f}[f-h_{k+j}-\cdots-h_{k+m-1}]_{k+m} = [f'+(h_k+1)+\cdots+(h_{k+j-1}+1)]_{k+m}
$$

$$
\subset [f']_k
$$

and $\mu([f-h_{k+j}-\cdots-h_{k+m-1}]_{k+m})=3^{-m}\mu([f]_k)$. It remains to apply Lemma 3.1. \Box

5. Radon invariant measures for Cartesian powers of the infinite Chacon transformation

We will need the following general lemma about splitting of Radon measures into direct products (cf. with [RuSi, Lemma 3.1.1]). It is, in fact, equivalent to [JaRoRu, Lemma A.1]. We provide a short proof of it.

Lemma 5.1. Let T and S be homeomorphisms of locally compact Cantor spaces X and Y respectively. If $\lambda \in \mathcal{M}_{\text{Ra}}^e(X \times Y, T \times S)$ and $\lambda \circ (\text{Id} \times S) = \lambda$ then there exist $\mu \in \mathcal{M}_{\text{Ra}}^e(X,T)$ and $\nu \in \mathcal{M}_{\text{Ra}}^e(Y,S)$ such that $\lambda = \mu \times \nu$.

Proof. Let $\tilde{\mu}$ be a probability measure on (X, \mathfrak{B}) which is equivalent (i.e. it has the same ideal of Borel subsets of 0-measure) to the projection of λ to X^{15} . Then $\tilde{\mu}$ is quasi-invariant and ergodic under T. Consider the disintegration of λ with respect to $\tilde{\mu}$:

$$
\lambda = \int_X \delta_x \times \lambda_x d\widetilde{\mu}(x),
$$

where the map $X \ni x \mapsto \lambda_x$ is the corresponding system of σ -finite measures on Y. Since $\lambda \in \mathcal{M}_{\text{Ra}}(X \times Y)$, it follows that $\lambda_x \in \mathcal{M}_{\text{Ra}}(X)$ for $\tilde{\mu}$ -a.e. x. We have that

$$
\lambda \circ (T \times S) = \int_X \delta_{T^{-1}x} \times \lambda_x \circ S \, d\widetilde{\mu}(x) = \int_X \delta_x \times \frac{d\widetilde{\mu} \circ T}{d\widetilde{\mu}}(x) \cdot \lambda_{Tx} \circ S \, d\widetilde{\mu}(x).
$$

In view of the uniqueness of the disintegration we obtain that λ is invariant under $T \times S$ if and only if $\lambda_x = \frac{d\widetilde{\mu} \circ T}{d\widetilde{\mu}}$ $\frac{\mu \circ T}{d\widetilde{\mu}}(x) \cdot \lambda_{Tx} \circ S$ for a.e. x. In a similar way, λ is invariant under Id×S if and only if $\lambda_x = \lambda_x \circ S$ for a.e. x. Hence $\lambda_x = \frac{d\widetilde{\mu} \circ T}{d\widetilde{\mu}}$ $\frac{\mu \circ T}{d\widetilde{\mu}}(x) \cdot \lambda_{Tx}$ for a.e. x. Since T is ergodic and λ_x is a Radon measure for a.e. $x \in X$, it follows that there is a single Radon measure ν on Y and a measurable map $X \ni x \mapsto a(x) \in \mathbb{R}_+^*$ such that $\lambda_x = a(x)\nu$ for μ -a.e. x. We now obtain that $\nu \circ S = \nu$ and $a(x) = \frac{d\widetilde{\mu} \circ T}{d\widetilde{\mu}}(x)a(Tx)$ for a.e. x. Define a σ -finite measure measure μ on X by setting $\frac{d\mu}{d\tilde{\mu}}(x) = a(x)$ for all $x \in X$. Then μ is invariant under T. Moreover, $\lambda = \mu \times \nu$ and hence $\mu \in \mathcal{M}_{\text{Ra}}^e(X, T)$ and $\nu \in \mathcal{M}_{\text{Ra}}^e(Y, S)$. \Box

We now state one of the main results of the paper.

¹⁵Such a measure can be obtained as the projection (to X) of a probability measure on $X \times Y$ equivalent to λ .

Theorem 5.2. The infinite Chacon transformation has Radon MSJ.

Proof. Let (X, μ, T_1) denote the infinite Chacon transformation as above. We fix $d > 1$ and take $\lambda \in \mathcal{M}_{\text{Ra}}^e(X^d, (T_1)^{\times d})$. Let $z = (x^{(1)}, \ldots, x^{(d)}) \in \mathcal{G}(\lambda)$. We recall that $X = \bigcup_{n\geq 0} X_n$ and $X_n := F_n \times C_{n+1} \times \cdots$. If m is large so that $z \in (X_m)^d$, we consider expansion $x^{(j)} = (f_m^{(j)}, c_{m+1}^{(j)}, c_{m+2}^{(j)}, \dots) \in X_m$ of $x^{(j)}$ for each $j = 1, \ldots, d$. We call z extreme if for each sufficiently large $n > m$, either $\{-h_{n-1}, h_{n-1} + 1\} \subset \{c_n^{(1)}, \ldots, c_n^{(d)}\}$ or $c_n^{(1)} = \cdots = c_n^{(d)}$.

We first show that if z is not extreme then λ splits into direct product of two its marginals. Indeed, there is an infinite subset $\mathcal{N} \subset \mathbb{N}$ and a non-empty subset $J \subset \{1, \ldots, d\}$ such that $c_n^{(j)} = 0$ if $j \in J$ and either $c_n^{(j)} = -h_{n-1}$ for all $j \notin J$ or $c_n^{(j)} = h_{n-1} + 1$ for all $j \in \{1, ..., d\} \setminus J$ for each $n \in \mathcal{N}$. We consider only the former case since the latter one is similar. Let $A^{(1)}, \ldots, A^{(d)}$ be compact open subsets in X. Then there is $r > 0$ such that these subsets are r-cylinders. Hence for each $n \geq r$ and each $j = 1, \ldots, d$, there is a subset $A_n^{(j)} \subset F_n$ such that $A^{(j)} = [A_n^{(j)}]_n$. It follows from Lemma 4.2 that

(5-1)
$$
\sum_{i \in F_n} 1_{A^{(1)} \times \dots \times A^{(d)}} ((T_1^{\times d})^i z) = \# \left(F_n \cap \bigcap_{j=1}^d (A_n^{(j)} - f_n^{(j)}) \right).
$$

Since we may assume without loss of generality that $A^{(j)} \cup \{x^{(j)}\} \subset X_{n-1}$, it follows from $(4-1)$ that

(5-2)
$$
A_n^{(j)} - f_n^{(j)} \subset F_n
$$
 for each $j = 1, ..., d$.

By the definition of generic point, for each $\epsilon > 0$ and each sufficiently large n, we have (in view of $(5-1)$ and $(5-2)$) that

(5-3)
$$
\frac{\# \left(\bigcap_{j=1}^d (A_n^{(j)} - f_n^{(j)}) \right)}{\# \left(\bigcap_{j=1}^d (F_{r,n} - f_n^{(j)}) \right)} = \frac{\lambda(A^{(1)} \times \dots \times A^{(d)})}{\lambda(X_r^d)} \pm \epsilon
$$

where $F_{r,n} := F_r + C_{r+1} + \cdots + C_n$. Now choose n so that $n+1 \in \mathcal{N}$. Then

$$
f_{n+1}^{(j)} = f_n^{(j)} + c_{n+1}^{(j)} = \begin{cases} f_n^{(j)}, & \text{if } j \in J \\ f_n^{(j)} - h_n, & \text{if } j \notin J. \end{cases}
$$

Since $A_{n+1}^{(j)} = A_n^{(j)} + C_{n+1}$, we obtain that

$$
\bigcap_{j=1}^{d} (A_{n+1}^{(j)} - f_{n+1}^{(j)}) = \bigcap_{j \in J} (A_n^{(j)} - f_n^{(j)} + C_{n+1}) \cap \bigcap_{j \notin J} (A_n^{(j)} - f_n^{(j)} + C_{n+1} + h_n)
$$
\n
$$
= \bigcap_{j \in J} (A_n^{(j)} - f_n^{(j)} + \{0, h_n + 1\}) \cap \bigcap_{j \notin J} (A_n^{(j)} - f_n^{(j)} + \{0, h_n\})
$$
\n
$$
= \bigcap_{j=1}^{d} (A_n^{(j)} - f_n^{(j)}) \sqcup \left(h_n + \bigcap_{\substack{j \in J \\ j \in J}} (A_n^{(j)} - f_n^{(j)} + 1) \cap \bigcap_{j \notin J} (A_n^{(j)} - f_n^{(j)}) \right).
$$

We define a map $S: X^d \to X^d$ by setting $S(x^{(1)}, \ldots, x^{(d)}) = (y^{(1)}, \ldots, y^{(d)})$, where $y^{(j)} := x^{(j)}$ if $j \in J$ and $y^{(j)} := T_1 x^{(j)}$ if $j \notin J$. Of course, S is a homeomorphism of X^d . It commutes with $T_1^{\times d}$ $i^{\times d}$. Let

$$
g^{(j)} := \begin{cases} 0, & \text{if } j \in J \\ 1, & \text{if } j \notin J. \end{cases}
$$

Then

$$
\#\bigg(\bigcap_{j=1}^d (A_{n+1}^{(j)} - f_{n+1}^{(j)})\bigg) = \#\bigg(\bigcap_{j=1}^d (A_n^{(j)} - f_n^{(j)})\bigg) + \#\bigg(\bigcap_{j=1}^d (A_n^{(j)} + g^{(j)} - f_n^{(j)})\bigg).
$$

In a similar way we obtain that

$$
\#\bigg(\bigcap_{j=1}^d (F_{r,n+1} - f_{n+1}^{(j)})\bigg) = \#\bigg(\bigcap_{j=1}^d (F_{r,n} - f_n^{(j)})\bigg) + \#\bigg(\bigcap_{j=1}^d (F_{r,n} + g^{(j)} - f_n^{(j)})\bigg).
$$

Thus we obtain that the lefthand side of $(5-3)$ (with $n+1$ in place of n) equals

(5-4)
$$
\frac{\# \left(\bigcap_{j=1}^{d} (A_n^{(j)} + g^{(j)} - f_n^{(j)}) \right) + \# \left(\bigcap_{j=1}^{d} (A_n^{(j)} - f_n^{(j)}) \right)}{\# \left(\bigcap_{j=1}^{d} (F_{r,n} + g^{(j)} - f_n^{(j)}) \right) + \# \left(\bigcap_{j=1}^{d} (F_{r,n} - f_n^{(j)}) \right)}.
$$

It is straightforward to verify that $\lambda(S(X_r)^d)/\lambda((X_r)^d) = 1 \pm \epsilon$ if r is large enough. Therefore it follows from (5-3) and (5-4) that

$$
\frac{\lambda(A^{(1)} \times \cdots \times A^{(d)})}{\lambda(X_r^d)} \pm \epsilon = \frac{\frac{\lambda \circ S(A^{(1)} \times \cdots \times A^{(d)})}{\lambda(X_r^d)} + \frac{\lambda(A^{(1)} \times \cdots \times A^{(d)})}{\lambda(X_r^d)} \pm 2\epsilon}{2 \pm \epsilon}.
$$

Hence $\lambda = (\lambda + \lambda \circ S)/2$, i.e. $\lambda \circ S = \lambda$. Lemma 5.1 yields that $\lambda = \lambda_1 \times \lambda_2$, where $\lambda_1 \in \mathcal{M}_{\text{Ra}}^e(X^J, T_1^J)$ and $\lambda_2 \in \mathcal{M}_{\text{Ra}}^e(X^{\{1,\dots,d\}\setminus J}, T_1^{\{1,\dots,d\}\setminus J})$. Continuing this way several times we obtain finally a splitting of λ into direct product of its marginals whose generic points are all extreme.

Thus to complete the proof of the thorem it suffices to prove the following fact: if every λ -generic point is extreme then either there is $s \in \{1, \ldots, d\}, d > 1$, such that the projection of $\mathcal{G}(\lambda)$ to the s-coordinate is μ -negligible or there exist $n_1, \ldots, n_{d-1} \in \mathbb{Z}$ such that $\lambda = \mu_{T^{n_1}, \ldots, T^{n_{d-1}}}^{16}$. Thus we fix $z \in \mathcal{G}(\lambda)$ as above and find $m > 0$ such that for each $n > m$, either $\{-h_{n-1}, h_{n-1} + 1\} \subset \{c_n^{(1)}, \ldots, c_n^{(d)}\}$ or $c_n^{(1)} = \cdots = c_n^{(d)}$. Put

$$
I := \{ i > m \mid c_i^{(1)} = \dots = c_i^{(d)} \}.
$$

We can now describe explicitly the set $\mathcal{G}(\lambda)$. Namely, we claim that

(5-5)
\n
$$
\mathcal{G}(\lambda) \cap ([f_m^{(1)}]_m \times \cdots \times [f_m^{(d)}]_m)
$$
\n
$$
= \bigcap_{l > m} \bigcap_{j_1, j_2 = 1}^d {\{\widetilde{z} \in (X_m)^d \mid \widetilde{c}_l^{(j_1)} - \widetilde{c}_l^{(j_2)} = c_l^{(j_1)} - c_l^{(j_2)}\}},
$$

¹⁶We omit the case where $d = 1$ because it is trivial. We recall that every (C, F) -action is Radon strictly ergodic.

where the coordinates $\tilde{c}_i^{(j)}$ $i^{(j)}$ are taken from the expansion

$$
\widetilde{z} = ((\widetilde{f}_m^{(1)}, \widetilde{c}_{m+1}^{(1)}, \widetilde{c}_{m+2}^{(1)}, \dots), \dots, (\widetilde{f}_m^{(d)}, \widetilde{c}_{m+1}^{(d)}, \widetilde{c}_{m+2}^{(d)}, \dots))
$$

of \tilde{z} . Indeed, let $A^{(1)}, \ldots, A^{(d)}$ be arbitrary r-cylinders with $r \geq m$. Then, as above, we have

(5-6)
$$
\frac{\sum_{i \in F_{r+1}} 1_{A^{(1)} \times \cdots \times A^{(d)}} ((T^{\times d})^{i} z)}{\sum_{i \in F_{r+1}} 1_{X_r^d} ((T^{\times d})^{i} z)} = \frac{\# \left(\bigcap_{j=1}^d (A_{r+1}^{(j)} - f_{r+1}^{(j)}) \right)}{\# \left(\bigcap_{j=1}^d (F_{r,r+1} - f_{r+1}^{(j)}) \right)}.
$$

Consider now 2 cases. If $r + 2 \in I$ then

$$
\bigcap_{j=1}^{d} (A_{r+2}^{(j)} - f_{r+2}^{(j)}) = \bigcap_{j=1}^{d} (A_{r+1}^{(j)} - f_{r+1}^{(j)} + C_{r+2} - c_{r+2}^{(j)}) = \bigsqcup_{c \in C_{r+2} - c_{r+2}^{(1)}} \bigcap_{j=1}^{d} (A_{r+2}^{(j)} - f_{r+2}^{(j)} + c)
$$

and hence $\#(\bigcap_{j=1}^d (A^{(j)}_{r+2} - f^{(j)}_{r+2})) = 3\#(\bigcap_{j=1}^d (A^{(j)}_{r+1} - f^{(j)}_{r+1}))$. In a similar way, $\#(\bigcap_{j=1}^d (F_{r,r+2} - f_{r+2}^{(j)})) = 3\#(\bigcap_{j=1}^d (F_{r,r+1} - f_{r+1}^{(j)})).$ If $r + 2 \notin I$ then

$$
\bigcap_{j=1}^{d} (A_{r+2}^{(j)} - f_{r+2}^{(j)}) = \bigcap_{j=1}^{d} (A_{r+1}^{(j)} - f_{r+1}^{(j)} + C_{r+2} - c_{r+2}^{(j)}) = \bigcap_{j=1}^{d} (A_{r+1}^{(j)} - f_{r+1}^{(j)})
$$

and hence $\#(\bigcap_{j=1}^d (A^{(j)}_{r+2} - f^{(j)}_{r+2})) = \#(\bigcap_{j=1}^d (A^{(j)}_{r+1} - f^{(j)}_{r+1}))$. In a similar way, $\#(\bigcap_{j=1}^d (F_{r,r+2} - f_{r+2}^{(j)})) = \#(\bigcap_{j=1}^d (F_{r,r+1} - f_{r+1}^{(j)}))$. Thus in every of the two cases we obtain that

(5-7)
$$
\frac{\# \left(\bigcap_{j=1}^{d} (A_{r+1}^{(j)} - f_{r+1}^{(j)}) \right)}{\# \left(\bigcap_{j=1}^{d} (F_{r,r+1} - f_{r+1}^{(j)}) \right)} = \frac{\# \left(\bigcap_{j=1}^{d} (A_{r+2}^{(j)} - f_{r+2}^{(j)}) \right)}{\# \left(\bigcap_{j=1}^{d} (F_{r,r+2} - f_{r+2}^{(j)}) \right)}
$$

for each $r \geq m$. Moreover, if we assume additionally that $A^{(j)} \subset X_{r-1}$ or, equivalently, $A_r^{(j)} \subset F_{r-1} + C_r$ for each $j = 1, ..., d$ then the same argument as above yields that (5-7) holds also if we replace r with $r - 1$. It follows from this, (5-6) and the fact that z is λ -generic that

(5-8)
$$
\frac{\lambda(A^{(1)} \times \cdots \times A^{(d)})}{\lambda(X_r^d)} = \frac{\# \left(\bigcap_{j=1}^d (A_r^{(j)} - f_r^{(j)}) \right)}{\# \left(\bigcap_{j=1}^d (F_r - f_r^{(j)}) \right)}.
$$

If \tilde{z} is another λ -generic point with $\tilde{z} \in [f_m^{(1)}]_m \times \cdots \times [f_m^{(d)}]_m$ then it is extreme and therefore (5-8) holds with $\tilde{f}_r^{(j)}$ in place of $f_r^{(j)}$, $j = 1, ..., d$. This yields

(5-9)
$$
\frac{\# \left(\bigcap_{j=1}^{d} (A_r^{(j)} - f_r^{(j)}) \right)}{\# \left(\bigcap_{j=1}^{d} (F_r - f_r^{(j)}) \right)} = \frac{\# \left(\bigcap_{j=1}^{d} (A_r^{(j)} - \tilde{f}_r^{(j)}) \right)}{\# \left(\bigcap_{j=1}^{d} (F_r - \tilde{f}_r^{(j)}) \right)}.
$$

We obtain that $\bigcap_{j=1}^d (A^{(j)} - f_r^{(j)}) \neq \emptyset$ if and only if $\bigcap_{j=1}^d (A^{(j)} - \widetilde{f}_r^{(j)}) \neq \emptyset$. Substituting $A^{(j)} := \{f_r^{(j)}\}$, we obtain that $f_r^{(j_1)} - f_r^{(j_2)} = \tilde{f}_r^{(j_1)} - \tilde{f}_r^{(j_2)}$ for each $1 \leq j_1 < j_2 \leq d$. Since r is arbitrary, we conclude that $c_l^{(j_1)} - c_l^{(j_2)} = \tilde{c}_l^{(j_1)} - \tilde{c}_l^{(j_2)}$ $\int_l^{(J2)}$ for each $1 \leq j_1 < j_2 \leq d$ and all sufficiently large l. This proves the " \subset "-part of the equality in $(5-5)$. The " \supset "-part of this equality follows easily from $(5-9)$ and $(5-6)$. Thus (5-5) is proved.

Consider now two cases. If $\mathbb{N} \setminus I$ is finite then we have $x^{(j)} = T_{g_j} x^{(1)}$ for $g_j := f_m^{(j)} - f_m^{(1)} + \sum_{n>m} (c_n^{(j)} - c_n^{(1)}), j = 2, \ldots, d$. Hence for each sufficiently large r and each $j \in \{1, ..., d\}$, we have that $f_r^{(j)} = g_j + f_r^{(1)}$. Then (5-7) implies that

$$
\frac{\lambda(A^{(1)} \times \cdots \times A^{(d)})}{\lambda(X_r^d)} = \frac{\# \left(\bigcap_{j=1}^d (A_r^{(j)} - g_j) \right)}{\# \left(\bigcap_{j=1}^d (F_r - g_j) \right)} = \frac{\mu_{T_{g_2}, \dots, T_{g_d}}(A^{(1)} \times \cdots \times A^{(d)})}{\mu_{T_{g_2}, \dots, T_{g_d}}(X_r^d)}.
$$

Hence λ is a multiple of $\mu_{T_{g_2},...,T_{g_d}}$. Thus λ is a graph-joining. Conversely, if λ is a graph-joining then $\mathbb{N} \setminus I$ is finite.

Now consider the second case, where $\mathbb{N} \setminus I$ is infinite. Then it follows from (5-5) that there exists $s \in \{1, ..., d\}$ and an infinite subset $I_0 \subset \{m+1, m+2, ...\} \setminus I$ such that for every $\widetilde{z} \in \mathcal{G}(\lambda) \cap ([f_m^{(1)}]_m \times \cdots \times [f_m^{(d)}]_m)$, we have that $\widetilde{c}_j^{(s)}$ $j_j^{(s)} \neq 0$ whenever $j \in$ I₀. However, $\mu({x = (f_m, c_{m+1}, \dots) \in X_m \mid c_j \neq 0 \text{ for all } j \in I_0}) = 0$. Thus the projection of the subset $\mathcal{G}(\lambda) \cap ([f_m^{(1)}]_m \times \cdots \times [f_m^{(d)}]_m) \subset X^d$ to the s-th coordinate is μ -negligible. Since the projection is equivariant and $\lambda([f_m^{(1)}]_m \times \cdots \times [f_m^{(d)}]_m) > 0$, it follows that the projection of the entire $\mathcal{G}(\lambda)$ to the s-th coordinate is also μ negligible. Hence $\lambda \notin J_{d,\text{Ra}}((T_1)^{\times d})$, as desired. Thus, T_1 has Radon MSJ. \Box

We note that not only the Radon MSJ property for T_1 was established in the proof of Theorem 5.2 but also a complete description of the set of ergodic $(T_1)^{\times d}$ invariant Radon measures on X^d was, in fact, obtained. Namely, the following theorem was proved indeed.

Theorem 5.3. Let $\lambda \in \mathcal{M}_{\text{Ra}}^e(X^d, (T_1)^{\times d})$. Then there is a partition of $\{1, \ldots, d\}$ into subsets J_1, \ldots, J_k such that λ splits into direct product of its marginals λ_i on X^{J_i} , each λ_i is (up to multiplicative constant) either μ if $\#J_i = 1$ or an invariant quasi-graph Radon measure if $\#J_i > 1$.

We now show that there are non-ergodic 2-fold Radon self-joinings ρ of T_1 such that almost every ergodic component of ρ is not a joining of T: the ergodic components of ρ are conservative quasi-graph invariant Radon measures which are not graph-joinings.

Example 5.4. Let $\mathbf{A} := \{-1,0,1\}^{\mathbb{N}}$. Given a sequence $a := (a_n)_{n=1}^{\infty} \in \mathbf{A}$, we define a measure λ_a on X_0^2 considered as the infinite product $\{(0,0)\}\times C_1^2\times C_2^2\times\cdots$ by setting

(5-10)
$$
\lambda_a = \delta_{(0,0)} \otimes \bigotimes_{n>0} \gamma_n^a,
$$

where γ_n is a probability on C_n^2 given by

(5-11)
$$
\gamma_n^a := \begin{cases} \delta_{(-h_{n-1}, h_{n-1}+1)} & \text{if } a_n = -1, \\ \delta_{(h_{n-1}+1, -h_{n-1})} & \text{if } a_n = 1, \\ \frac{1}{3} \sum_{c \in C_n} \delta_{(c, c)} & \text{if } \alpha_n = 0. \end{cases}
$$

We recall that $C_n = \{-h_{n-1}, 0, h_{n-1} + 1\}$. Let $J_a := \{n \mid a_n \neq 0\}$. We now set for each $n \notin J_a$,

$$
\alpha_n^a := \begin{cases}\n-h_{n-1}, & \text{if } a_n = -1 \\
h_{n-1} + 1, & \text{if } a_n = 1\n\end{cases}, \quad \beta_n^a := \begin{cases}\nh_{n-1} + 1, & \text{if } a_n = -1 \\
-h_{n-1}, & \text{if } a_n = 1\n\end{cases}.
$$

It is easy to see that the sequences $\alpha^a := (\alpha^a_n)_{n \notin J}$ and $\beta^a := (\beta^a_n)_{n \notin J}$ satisfy (3-4). Then it follows from Theorem 3.2(iv) that $\lambda_a = \mu^{\alpha^a} \circ \psi_{\alpha^a, \beta^a}^{-1} \upharpoonright X_0^2$ and that $\mu^{\alpha^a} \circ$ $\psi_{\alpha^a,\beta^a}^{-1}$ is a $(T_1 \times T_1)$ -invariant quasi-graph Radon measure¹⁷. For each $n > 0$, let κ_n be a probability on $\{-1,0,1\}$ such that $\kappa_n(-1) = \kappa_n(1) = \frac{1}{2n}$ and $\kappa_n(0) = 1 - \frac{1}{n}$ $\frac{1}{n}$. We now set $\kappa := \bigotimes_{n>0} \kappa_n$ and $\lambda := \int_{\mathbf{A}} \lambda_a d\kappa(a)$. Then

$$
\lambda = \left(\int_{\mathbf{A}} \mu^{\alpha^a} \circ \psi_{\alpha^a, \beta^a}^{-1} d\kappa(a) \right) \restriction X_0^2.
$$

We claim that the Radon measure

(5-12)
$$
\rho := \int_{\mathbf{A}} \mu^{\alpha^a} \circ \psi_{\alpha^a, \beta^a}^{-1} d\kappa(a)
$$

is a 2-fold self-joining of T_1 . For that we have to verify that the two coordinate projections of this measure are equivalent to μ . It suffices to show that the two coordinate projections of λ are equivalent to $\mu \upharpoonright X_0$. A straightforward computation shows that the projection of λ to the first coordinate is the following measure¹⁸:

$$
\delta_0 \otimes \bigotimes_{n>0} \left(\kappa_n(-1)\delta_{-h_{n-1}} + \kappa_n(1)\delta_{h_{n-1}+1} + \frac{\kappa_n(0)}{3} \sum_{c \in C_n} \delta_c \right).
$$

It is equivalent to the measure $\delta_0 \otimes \bigotimes_{n>0} \left(\frac{1}{3}\right)$ $\frac{1}{3}\sum_{c\in C_n} \delta_c$ (i.e. to $\mu \restriction X_0$) by the Kakutani theorem on equivalence of infinite product measures [Ka]. In a similar way one can prove that the projection of ρ to the second coordinate is also equivalent to μ . Therefore $\rho \in J_{2,\text{Ra}}(T_1)$. On the other hand,

$$
0 = \kappa(\{a \mid J_a \text{ is cofinite}\}) = \kappa(\{a \mid \mu^{\alpha^a} \circ \psi_{\alpha^a, \beta^a}^{-1} \text{ is a 2-fold joining of } T_1\}),
$$

$$
0 = \kappa(\{a \mid J_a \text{ is finite}\}) = \kappa(\{a \mid \mu^{\alpha^a} \circ \psi_{\alpha^a, \beta^a}^{-1} \text{ is purely atomic}\}).
$$

We note that (5-12) is the ergodic decomposition of ρ and κ is the corresponding measure on the space of ergodic components of ρ . Hence almost all ergodic components of ρ are conservative Radon dynamical systems whose coordinate projections are singular to μ .

6. Uncountable family of pairwise Radon disjoint infinite Chacon like transfrmations

Take an infinite sequence $\omega \in \{0,1\}^{\mathbb{N}}$. Define a sequence of positive integers $(h_n)_{n=0}^{\infty}$ by setting $h_0 := 1$ and $h_{n+1} := 6h_n + 1$ for $n > 0$. For each $n \geq 0$, we now let

$$
F_n := \left\{ -\frac{h_n - 1}{2}, -\frac{h_n - 3}{2}, \dots, \frac{h_n - 1}{2} \right\} \quad \text{and} \quad
$$

$$
C_{n+1}^{\omega} := \left\{ -h_n - 1 + \omega(n), 0, h_n + \omega(n) \right\}.
$$

¹⁷For the definition of μ^{α} and $\psi_{\alpha,\beta}$ we refer to §3.2.

 18 We use (5-10) and (5-11) to obtain this.

The sequence $(C_n^{\omega}, F_{n-1})_{n\geq 1}$ satisfies (1-1) and (I)–(IV) from Section 1. We also note that $#F_n = h_n$ and

$$
F_n + C_{n+1}^{\omega} - F_n - C_{n+1}^{\omega} \subset F_{n+1}.
$$

Let $(X^{\omega}, \mu^{\omega}, T^{\omega})$ denote the associated (C, F) -dynamical system. It is Radon strictly ergodic. We call it an infinite Chacon like transformation. The infinite Chacon transformation corresponds to the case $\omega(n) = 1$ for all n.

If $\omega, \omega' \in \{0,1\}^{\mathbb{N}},$ we write $\omega \sim \omega'$ if the pair (ω, ω') belongs to the tail equivalence relation on $\{0,1\}^{\mathbb{N}}$, i.e. there is $N > 0$ such that $\omega(n) = \omega'(n)$ for all $n > N$.

Theorem 6.1.

- (i) For each $\omega \in \{0,1\}^{\mathbb{N}}$, the dynamical system $(X^{\omega}, \mu^{\omega}, T_1^{\omega})$ is totally ergodic. It has Radon MSJ. $C(T_1^{\omega}) = \{T_n^{\omega} \mid n \in \mathbb{Z}\}.$
- (ii) If $\omega \sim \omega'$ then $(X^{\omega}, \mu^{\omega}, T_1^{\omega})$ is isomorphic to $(X^{\omega'}, \mu^{\omega'}, T_1^{\omega'})$ $\binom{1}{1}$.
- (iii) If $\omega \nsim \omega'$ then $(X^{\omega}, \mu^{\omega}, T_1^{\omega})$ is Radon disjoint with $(X^{\omega'}, \mu^{\omega'}, T_1^{\omega'})$ $\binom{m'}{1}$.

Sketch of the proof. (i) is proved in the same way as Theorem 5.2 and Proposition 4.6.

(ii) Let $\omega(i) = \omega'(i)$ for all $i > N$. Given $x \in X^{\omega}$, let $x = (f_n, c_{n+1}, \dots) \in (X^{\omega})_n$ for some $n > N$. We now define $\phi: X^{\omega} \to X^{\omega'}$ by setting $\phi(x) := (f_n, c_{n+1}, \dots) \in$ $(X^{\omega'})_n$. Then ϕ is an isomorphism of $(X^{\omega}, \mu^{\omega}, T_1^{\omega})$ onto $(X^{\omega'}, \mu^{\omega'}, T_1^{\omega'}$ $\binom{1}{0}$.

(iii) Let λ be an ergodic $(T_1^{\omega} \times T_1^{\omega'})$ $\int_1^{\omega'}$)-invariant measure on $X^{\omega} \times X^{\omega'}$ whose marginals are equivalent to μ^{ω} and $\mu^{\omega'}$ respectively. Fix a generic point (x, x') of λ . Find m such that $x \in (X^{\omega})_m$ and $x' \in (X^{\omega'})_m$. Consider expansions $x = (f_m, c_{m+1}, \dots)$ and $x' = (f'_m, c'_{m+1}, \dots)$ of x and x' in $(X^{\omega})_m$ and $(X^{\omega})_m$ respectively. Let

$$
J := \{ i \in \mathbb{N} \mid \omega(i) \neq \omega'(i) \}.
$$

This set is infinite. One of the following cases takes place.

- (A) The subset $J_A := \{j \in J \mid \{c_j, c'_j\} = \{h_{j-1} + 1, h_{j-1}\}\}\$ is infinite.
- (B) The subset $J_B := \{j \in J \mid \{c_j, c'_j\} = \{-h_{j-1}, -h_{j-1} 1\}\}\$ is infinite.
- (C) The subset $J_C := \{ j \notin J \mid 0 \in \{c_j, c'_j\} \neq \{0\} \}$ is infinite.
- (D) The subset $J_A \cup J_B \cup J_C$ is finite.

We consider separately every case.

Case (A). Let A and A' be two r-cylinders with $r > m$. Then $A = [A_n]_n$ and $A' = [A'_n]_n$ for some subsets $A_n \subset F_n$ and $A'_n \subset F'_n$ for each $n \geq r$. As in the proof of Theorem 5.2, we obtain that

$$
\sum_{i \in F_n} 1_{A \times A'}((T_1^{\omega} \times T_1^{\omega'})^i(x, x')) = \#((A_n - f_n) \cap (A'_n - f'_n)).
$$

Therefore for each $\epsilon > 0$ and each sufficiently large n, we have that

(6-1)
$$
\frac{\#((A_n - f_n) \cap (A'_n - f'_n))}{\#((F_{r,n} - f_n) \cap (F_{r,n} - f'_n))} = \frac{\lambda(A \times A')}{\lambda(X_r)} \pm \epsilon.
$$

This follows from the fact that $(x, x') \in \mathcal{G}(T_1^{\omega} \times T_1^{\omega'})$ $\binom{n}{1}$. Now choose *n* so that $n+1 \in$ J_A . Then $c_{n+1} = h_n + 1$ and $c'_{n+1} = h_n$ (or $c_{n+1} = h_n$ and $c'_{n+1} = h_n + 1$, which 20

is considered in a similar way) and hence $f_{n+1} = f_n + h_n + 1$ and $f'_{n+1} = f'_n + h_n$. We now have

$$
(A_{n+1}-f_{n+1})\cap (A'_{n+1}-f'_{n+1})=(A_n-1-f_n+C_{n+1}^{\omega}-h_n)\cap (A_n-f_n+C_{n+1}^{\omega'}-h_n).
$$

Since $C_{n+1}^{\omega} = \{-h_n, 0, h_n + 1\}$ and $C_{n+1}^{\omega'} = \{-h_n - 1, 0, h_n\}^{19}$, it follows that

$$
#((A_{n+1}-f_{n+1}) \cap (A'_{n+1}-f'_{n+1})) = #((A_n - f_n) \cap (A'_n - f'_n))
$$

+
$$
#((A_n - 1 - f_n) \cap (A'_n - f'_n)) + #((A_n - f_n) \cap (A'_n - f'_n))
$$

In a similar way we obtain that

$$
#((F_{r,n+1}-f_{n+1})\cap (F_{r,n+1}-f'_{n+1})) = 2\#((F_{r,n}-f_n)\cap (F_{r,n}-f'_n))
$$

+ $\#((F_{r,n}-1-f_n)\cap (F_{r,n}-f'_n)).$

Applying (6-1) we now obtain that

$$
\frac{\lambda(A \times A')}{\lambda(X_r)} \pm \epsilon = \frac{\frac{\lambda(A \times A')}{\lambda(X_r)} + \frac{2\lambda \circ (T_{-1}^{\omega} \times \text{Id})(A \times A')}{\lambda(X_r)} \pm 3\epsilon}{3 \pm \epsilon}.
$$

This yields that $\lambda \circ (T_{-1}^{\omega} \times \text{Id}) = \lambda$. By Lemma 5.1, λ equals $\mu \times \mu'$ (up to a multiplicative constant).

Case (B) is analogous to Case (A).

Case (C) was considered, in fact, in the first part of the proof of Theorem 5.2. In this case we also obtain that $\lambda = q \cdot \mu \times \mu'$ for some $q > 0$.

Case (D). We will show that in this case one of the two marginals of the set $\mathcal{G}(\lambda)$ is of null measure. For that we will argue as in the final part of the proof of Theorem 5.2. Without loss of generality we may assume that $\{c_n, c'_n\} \cap (J_A \cup J_B \cup$ J_C = \emptyset for each $n > m$. We claim that

$$
(6-2) \mathcal{G}(\lambda) \cap ([f_m]_m \times [f']_m) = \bigcap_{l>m} \{ (\widetilde{x}, \widetilde{x}') \in (X^{\omega})_m \times (X^{\omega'})_m \mid \widetilde{c}_l - \widetilde{c}'_l = c_l - c'_l \},
$$

where the coordinates \tilde{c}_l and \tilde{c}'_l are taken from the expansion

$$
\widetilde{x} = (\widetilde{f}_m, \widetilde{c}_{m+1}, \widetilde{c}_{m+2}, \dots), \quad \widetilde{x}' = (\widetilde{f}'_m, \widetilde{c}'_{m+1}, \widetilde{c}'_{m+2}, \dots)
$$

of \tilde{x} and \tilde{x}' . We let $\alpha_n := \#((A_n - f_n) \cap (A'_n - f'_n))$. One can verify (as in the proof of Theorem 5.2) that

(6-3)
$$
\alpha_{n+1} = \begin{cases} 3\alpha_n & \text{if } c_{n+1} = c'_{n+1}, \\ \alpha_n & \text{if } 0 \notin \{c_{n+1}, c'_{n+1}\} \text{ but } c_{n+1} \neq c'_{n+1}, \\ 2\alpha_n & \text{if } 0 \in \{c_{n+1}, c'_{n+1}\} \text{ but } c_{n+1} \neq c'_{n+1}. \end{cases}
$$

This implies

$$
\frac{\#((A_{r+1}-f_{r+1})\cap (A'_{r+1}-f'_{r+1})}{\#((F_{r,r+1}-f_{r+1})\cap (F'_{r,r+1}-f'_{r+1})}=\frac{\#((A_{r+2}-f_{r+2})\cap (A'_{r+2}-f'_{r+2})}{\#((F_{r,r+2}-f_{r+2})\cap (F'_{r,r+2}-f'_{r+2})}.
$$

¹⁹Because $h_n + 1 \in C_{n+1}^{\omega}$ and $h_n \in C_{n+1}^{\omega'}$.

Slightly modifying the argument in the proof of Theorem 5.2 we deduce (6-2). If either the second or the third condition from $(6-3)$ is satisfied for infinitely many n then one can easily deduce from (6-2) and (6-3) that one of the coordinate projection of $\mathcal{G}(\lambda)$ is of 0 measure. If the second and the third condition are satisfied for only finitely many *n* then the first condition in $(6-3)$ is satisfied for all but finitely many n. Hence it is satisfied for infinitely many elements of J. However, if $\tilde{c}_n = \tilde{c}'_n$ for $n \in J$ then $\tilde{c}_n = \tilde{c}'_n = 0$. Therefore the two marginals of $\mathcal{G}(\lambda)$ are of 0 measure. \Box

We also note that one can construct²⁰, for each ω , a non-ergodic Radon 2fold self-joining of T_1^{ω} whose ergodic components are not joinings because their coordinate projections are singular to μ^{ω} .

Given $\omega \in \{0,1\}^{\mathbb{N}},$ we define $\omega^* \in \{0,1\}^{\mathbb{N}}$ by setting $\omega^*(i) = 1 - \omega(i)$. It is easy to see that $T_1^{\omega^*}$ j^{ω^*} is conjugate to the inverse to T_1^{ω} .

Corollary 6.2. For each ω , the transformation T_1^{ω} is not conjugate to its inverse. Moreover, T_1^{ω} and its inverse are Radon disjoint.

7. Further generalizations and some open problems

It is easy to see that the above argument works almost verbally for a more general class of rank-one transformations. To specify this class we first give an alternative (but equivalent) description of the infinite Chacon transformation. For that we will use the classical language of cutting-and-stacking construction [Fr]. The initial 0-th tower consists of a single interval $[0, 1)$. We now describe the inductive procedure of passing from the *n*-th tower consisting of h_n levels of width $\frac{1}{3^n}$ to the $(n+1)$ -th tower. For that we cut the *n*-th tower into 3 subtowers (called copies) of equal width. Then we place the second copy over the first one, add an additional level (called spacer) over the second copy and put the third copy over this spacer. Next we put $[1.5h_n]$ spacers over the top of the third copy and $[1.5h_n] + 1$ spacers under the bottom of the first copy. Here [.] stands for the integer part. We thus obtain the $(n + 1)$ -tower consisting of $h_{n+1} = 6h_n + 1$ levels of width $\frac{1}{3^{n+1}}$. The transformation T moves each (except for the highest one) level of the tower one level up. The transformation is not defined on the highest level of the tower. However in the limit we obtain a well defined transformation on $\mathbb R$ (which is the union of all levels of all towers) endowed with Lebesgue measure. It is measure theoretically isomorphic to the infinite Chacon transformation described above via the (C, F) -construction in Section 4.

We now can obtain a family of transformations using almost the same cuttingand-stacking algorithm but adding "more" spacers. I mean the following. Let $(\alpha_n)_{n\in\mathbb{N}}$ and $(\beta_n)_{n\in\mathbb{N}}$ be two arbitrary sequences of nonnegative integers. When constructing the $(n+1)$ -th tower we first do verbally what we did in the above construction of the infinite Chacon transformation and after that we put α_n additional spacers on the top and β_n spacers under the bottom of the tower. In the limit of the inductive construction we obtain a transformation that has Radon MSJ. The proof is almost the same as for the infinite Chacon transformation.

We conclude the paper with a list of open problems.

— Are there two ergodic Radon dynamical systems which are Radon disjoint but which admit a non-ergodic Radon joining?

 20 Via a slight modification of Example 5.4.

- Does the infinite Chacon transformation have invariant sub- σ -algebras admitting equivalent σ -finite invariant measures?
- More generally, given a (C, F) -transformation T with an infinite invariant Radon measure μ such that all ergodic $T \times T$ -invariant Radon measures are either quasi-graphs or $\mu \times \mu$, does it have factors admitting equivalent σ -finite invariant measures?

REFERENCES

- [Aa1] J. Aaronson, The intrinsic normalizing constants of transformations preserving infinite measures, J. d'Analyse Math. 49 (1987), 239–270.
- [Aa2] J. Aaronson, An introduction to infinite ergodic theory, Mathematical Surveys and Monographs, vol. 50, American Mathematical Society, 1997.
- [AdFrSi] T. Adams, N. Friedman and C. E. Silva, Rank-one weak mixing for nonsingular transformations, Israel J. Math. 102 (1997), 269–281.
- [Da1] A. I. Danilenko, Funny rank-one weak mixing for nonsingular Abelian actions, Isr. J. Math. **121** (2001), 29–54.
- [Da2] , On simplicity concepts for ergodic actions, J. d'Anal. Math. 102 (2007), 77– 118.
- [Da3] $\qquad \qquad \qquad \ldots$, (C, F) -actions in ergodic theory, Geometry and Dynamics of Groups and Spaces, Progr. Math., vol. 265, 2008, pp. 325–351.
- [Da4] , Finite ergodic index and asymmetry for infinite measure preserving actions, Proc. Amer. Math. Soc. 144 (2016), 2521–2532.
- [DaSi] A. I. Danilenko and C. E. Silva, Ergodic theory: nonsingular transformations, Encyclopedia of Complexity and Systems Science, Springer, 2009, pp. 3055–3083.
- [Ef] E. G. Effros, Transformation groups and C∗-algebras, Ann. Math. 81 (1965), 38–55.
- [dJRaSw] A. del Junco, M. Rahe and L. Swanson, *Chacon's automorphism has minimal self*joinings, J. Analyse Math. **37** (1980), 276-284.
- [dJRud] A. del Junco and D. Rudoplh, On ergodic actions whose self-joinings are graphs, Ergodic Theory and Dynamical Systems 7 (1987), 531–557.
- [dJSi] A. del Junco and C. E. Silva, On factors of non-singular Cartesian products, Ergodic Theory and Dynamical Systems 23 (2003), 1445–1465.
- [Fr] N. Friedman, Introduction to ergodic theory, Van Nostrand Reinhold, New York, NY, 1970.
- [Fu] H. Furstenberg, Disjointness in ergodic theory, minimal sets and diophantine approximation, Math. Syst. Theory **1** (1967), 1-49.
- [Gli] J. Glimm, Locally compact transformation groups, Trans. Amer. Math. Soc. 101 (1961), 124–138.
- [Ka] S. Kakutani, On equivalence of infinite product measures, Ann. Math. 49 (1948), 214– 224.
- [JaRoRu] E. Janvresse, E. Roy and T. de la Rue, Invariant measures for Cartesian powers of Chacon infinite transformation, arXiv:1509.07802.
- [RudSi] D. J. Rudolph and S. E. Silva, *Minimal self-joinings for nonsingular transformations*, Ergodic Theory and Dynamical Systems 9 (1989), 759–800.
- [Ru] T. de la Rue, Joinings in ergodic theory, Encyclopedia of Complexity and Systems Science, Springer, 2009, pp. 5037–5051.
- [Sc] K. Schmidt, Infinite invariant measures on the circle, Symposia Mathematica 21 (1977), 37–43.
- [SiWi] C. E. Silva and D. Witte, On quotients of nonsingular actions whose self-joinings are graphs, Internat. J. Math. 5 (1994), 219–237.
- [Yu] H. Yuasa, Uniform sets for infinite measure-preserving systems, J. d'Anal. Math. 120 (2013), 333–356.

Institute for Low Temperature Physics & Engineering of National Academy of Sciences of Ukraine, 47 Lenin Ave., Kharkov, 61164, UKRAINE E-mail address: alexandre.danilenko@gmail.com