MIXING PROPERTIES IN CODED SYSTEMS

JEREMIAS EPPERLEIN, DOMINIK KWIETNIAK, AND PIOTR OPROCHA

ABSTRACT. We show that topological mixing, weak mixing and total transitivity are equivalent for coded systems. We provide an example of a mixing coded system which cannot be approximated by any increasing sequence of mixing shifts of finite type, has only periodic points of even period and each set of its generators consists of blocks of even length. We prove that such an example cannot be a synchronized system. We also show that a mixing coded systems has the strong property P.

1. INTRODUCTION

We consider coded systems and their recurrence properties that are stronger than topological transitivity. We are interested in topological (weak) mixing, and properties like the strong property P, which is a variant of the specification property.

Recall that a shift space is a *coded system* if it can be presented by an irreducible directed graph whose edges are labelled by symbols from a finite alphabet \mathcal{A} . Here, "presented" means that the shift space is the closure in $\mathcal{A}^{\mathbb{Z}}$ of all bi-infinite sequences of symbols which are labels for bi-infinite paths in the graph. Equivalently, X is a coded system if it is a closure in $\mathcal{A}^{\mathbb{Z}}$ of the set of all bi-infinite sequences obtained by freely concatenating the words in a (possibly infinite) list of words over \mathcal{A} . Such a list is the set of generators of X.

Transitive sofic shifts are coded systems which can be presented by a finite irreducible digraph. Coded systems were introduced by Blanchard and Hansel [5], who showed that any factor of a coded system is coded. This is a generalization of a well-known property of sofic shifts [16, Corollary 3.2.2].

It is natural to ask which properties of irreducible sofic shifts extend to coded systems. We address an aspect of this problem which leads to an extension of known results and a quite unexpected example. The counter-intuitive nature of this example is in our opinion the most interesting feature of this paper, but we hope that our other results will fill a gap in the literature. Furthermore, there has been a resurgence in interest in coded systems in general, and their notable subclasses in particular (for example S-gap shifts [2, 8, 10], β -shifts [8], Dyck shifts [17, 18]). Coded systems often provide a testing ground for further extensions (see [9], where the line of investigation initiated in [8] is developed and extended to non-symbolic systems). Therefore understanding the situation for coded systems may lead to a solution of more general problems.

In order to describe our results, note first that any coded system is topologically transitive (irreducible). Recall also that for a non trivial transitive sofic shift X the following stronger variants of transitivity are equivalent:

(a) X has the periodic specification property;

⁽b) X is topologically mixing;

⁽c) X has the strong property P;

Date: July 10, 2018.

²⁰¹⁰ Mathematics Subject Classification. 37B10, 37B20.

Key words and phrases. coded system, synchronized system, shift of finite type, sofic shift, mixing, property P, specification property, Fisher cover.

(d) X is topologically weakly mixing;

(e) X is totally transitive;

(f) X has two periodic points with relatively prime primary periods.

It seems that this is a folklore theorem. It follows from various well-known results, but we could not find it in this form in the literature.

Here we examine the connections between transitivity variants (b)-(f) for non necessarily sofic coded systems. Note that the proof of equivalence of properties (b)-(f) for shifts of finite type can be adapted for irreducible Markov shifts over countable alphabets. Furthermore, every coded system contains a dense subset which is a factor of an irreducible countable Markov shift. This suggests that properties (b)-(f) should remain equivalent for coded systems. And this is indeed the case with one notable exception.

It is easy to see that property (f) implies topological mixing for coded systems. We prove, somewhat surprisingly, that the converse does not hold for all coded systems. We construct a mixing coded systems without a periodic point of odd period and hence without a generator of an odd length. Such a system cannot contain a mixing shift of finite type. Note that Krieger [14] characterized coded systems as those shift spaces which contain an increasing sequence of irreducible shifts of finite type with dense union. Krieger's characterization is the best possible, in the sense that there exists an increasing sequence of sofic shifts whose closure is a shift space which is not a coded system [6]. It follows from our result that there are mixing coded systems, which cannot be approximated from the inside by mixing shifts of finite type as they do not contain any mixing sofic shift.

On the other hand, (b)–(e) are still equivalent for coded systems. We also show that if X is a transitive but not totally transitive coded system, then for some prime p we can write $X = X_1 \cup \ldots X_p$, where X_j 's are closed subsets of X cyclically permuted by the shift map σ , each X_j is σ^p invariant, $X_i \cap X_j$ is nowhere dense for $i \neq j$, and (X_j, σ^p) is topologically mixing. Finally, we note that (a) implies (b) by the definition, but is not implied by any of the conditions (b)–(f), since all β -shifts are mixing but some do not have the specification property (see [19]).

This paper is organized as follows: In the next section we set up the notation and terminology. In Section 3 we prove a structure theorem for topologically mixing coded systems and that total transitivity, weak mixing and mixing are equivalent for coded systems. Section 4 contains an example of a shift space which has the strong property P, but is not topologically mixing. In Section 5 we describe a mixing coded systems without a periodic point of odd period. In Section 6 we establish an equivalence of the strong property P and topological mixing for coded systems. In section 7 we present some complementary results on synchronized systems. They imply that (b)–(f) are equivalent for synchronized systems, thus our example does not have any synchronizing words.

2. NOTATION AND DEFINITIONS

We assume the reader is familiar with elementary symbolic dynamics as in [16]. We fix a finite set \mathcal{A} with at least two elements and call it the *alphabet*. Let $\mathcal{A}^{\mathbb{Z}}$ denote the set of bi-infinite (two-sided) sequences

$$x = (x_i)_{i \in \mathbb{Z}} = \dots x_{-3} x_{-2} x_{-1} x_0 x_1 x_2 \dots,$$

such that $x_i \in \mathcal{A}$ for all *i*. We equip \mathcal{A} with the discrete topology and we consider $\mathcal{A}^{\mathbb{Z}}$ as a compact metric space in the product topology. The shift operator on $\mathcal{A}^{\mathbb{Z}}$ is denoted by σ . A *shift space* over the alphabet \mathcal{A} is a shift-invariant subset of $\mathcal{A}^{\mathbb{Z}}$ which is closed in that topology. The set $\mathcal{A}^{\mathbb{Z}}$ itself is a shift space called the *full shift*. In this paper all shift spaces will be two-sided and transitive.

 $\mathbf{2}$

A block of length k is an element $w = w_1 w_2 \dots w_k$ of \mathcal{A}^k . Throughout this paper "a word" is a synonym for "a block". The length of a block is denoted |w|. The set of all words over \mathcal{A} is denoted by \mathcal{A}^* . Given $x \in \mathcal{A}^{\mathbb{Z}}$ and $i, j \in \mathbb{Z}$ with $i \leq j$ we write $x_{[i,j]}$ to denote the block $x_i x_{i+1} \dots x_j$. We say that a block w occurs in x if $w = x_{[i,j]}$ for some $i, j \in \mathbb{Z}$. A language of a shift space X is the set $\mathcal{B}(X)$ of all blocks that occur in X. The set of blocks of length n in the language of X is denoted $\mathcal{B}_n(X)$. Similarly, the set of all words that occur in a point $x \in \mathcal{A}^{\mathbb{Z}}$ is denoted $\mathcal{B}(x)$. The empty word \perp is the unique word of length 0. We write \mathcal{A}^+ for the set of nonempty words over \mathcal{A} .

A central cylinder set of a word $u \in \mathcal{B}_{2r+1}(X)$, where $r \in \mathbb{N}$, is the set $[u] \subset X$ of points from X in which the block u occurs starting at position -r, that is, $\{y \in X : y_{[-r,r]} = u\}$. Central cylinders (or cylinders for short) are open and closed subsets of X. The family

$$\{[x_{[-r,r]}]:r\in\mathbb{N}\}$$

of cylinder sets determined by a central subblock of x is a neighbourhood basis for a point $x \in X$. We use a multiplicative notation for *concatenation* of words, so that $w^n = w \dots w$ (*n*-times) and $w^{\infty} = www \dots \in \mathcal{A}^{\mathbb{N}}$.

Given a set of words $\Omega \subset \mathcal{A}^+$, we define $\Omega^0 = \{\bot\}$, and $\Omega^n = \Omega^{n-1}\Omega = \{uw : u \in \Omega^{n-1}, w \in \Omega\}$. We also let Ω^+ denote the set of all possible finite concatenations of words from Q, that is, $\Omega^+ = \bigcup_{n=1}^{\infty} \Omega^n$. In particular, $\Omega \subset \Omega^+$.

By $\Omega^{\mathbb{Z}}$ we denote the set containing all possible bi-infinite concatenations of elements of Ω , that is, $x \in \Omega^{\mathbb{Z}}$ if x can be partitioned into elements of Ω .

By a *countable graph* we mean a directed graph with at most countably many vertices and edges. A countable graph G is *labeled* if there is a *labeling* Θ which is simply a function from the set of edges of G to the alphabet \mathcal{A} . A labeling of edges extends, in an obvious way, to a labeling of all finite (respectively, infinite, bi-infinite) paths on G by blocks (respectively, infinite or bi-infinite sequences) over \mathcal{A} . The set Y_G of bi-infinite sequences constructed by reading off labels along a bi-infinite path on a labeled graph (G, Θ) is shift invariant, but usually it is not closed and therefore not a shift space. Nevertheless, its closure $X = \overline{Y_G}$ in $\mathcal{A}^{\mathbb{Z}}$ is a shift space, and we say that X is presented by (G, Θ) . Any shift space admitting such a presentation is a *coded system*. A set of *generators* for a shift space X is a family of words $\mathcal{Q} \subset \mathcal{A}^*$ such that the language of X coincides with the set of all subblocks occurring in elements of Q^+ . Equivalently, X is the closure of $Q^{\mathbb{Z}}$ in $\mathcal{A}^{\mathbb{Z}}$. Every coded systems has a set of generators, and conversely, if a transitive shift space has a set of generators then it is a coded system. A countable graph is *irreducible* if given any pair of its vertices, say (v_i, v_j) , there is a path from v_i to v_j .

Dynamical properties like those mentioned in (a)–(e) above are usually defined for a continuous map acting on a metric space. Here we define them in the language of symbolic dynamics.

A shift space X is:

- (1) *transitive* if for any $u, v \in \mathcal{B}(X)$ there is $w \in \mathcal{B}(X)$ such that $uwv \in \mathcal{B}(X)$;
- (2) totally transitive if for any $u, v \in \mathcal{B}(X)$ and any n > 0 there is $w \in \mathcal{B}(X)$ such that $uwv \in \mathcal{B}(X)$ and n divides |uw|;
- (3) weakly mixing if for any $u_1, v_1, u_2, v_2 \in \mathcal{B}(X)$ there are $w_1, w_2 \in \mathcal{B}(X)$ such that $u_1w_1v_1, u_2w_2v_2 \in \mathcal{B}(X)$ and $|u_1w_1| = |u_2w_2|$;
- (4) mixing if for every $u, v \in \mathcal{B}(X)$ there is N > 0 such that for every n > N there is $w \in \mathcal{B}_n(X)$ such that $uwv \in \mathcal{B}(X)$.

We say that a shift space has:

- (1) the strong property P if for any $k \ge 2$ and any words $u_1, \ldots, u_k \in \mathcal{B}(X)$ with $|u_1| = \ldots = |u_k|$ there is an $n \in \mathbb{N}$ such that for any $N \in \mathbb{N}$ and function $\varphi \colon \{1, \ldots, N\} \to \{1, \ldots, k\}$ there are words $w_1, \ldots, w_{N-1} \in \mathcal{B}_n(X)$ such that $u_{\varphi(1)}w_1u_2\ldots u_{\varphi(N-1)}w_{N-1}u_{\varphi(N)} \in \mathcal{B}(X)$;
- (2) the specification property if there is an integer $N \ge 0$ such that for any $u, v \in \mathcal{B}(X)$ there is $w \in \mathcal{B}_N(X)$ such that $uwv \in \mathcal{B}(X)$.

Blanchard [7] proved that the strong property P implies weak mixing, and does not imply mixing.

It is convenient to rephrase the above definitions using the sets

 $N_{\sigma}([u], [v]) = \{\ell \in \mathbb{N} : uwv \in \mathcal{B}(X) \text{ for some } w \in \mathcal{A}^* \text{ such that } |uw| = \ell\},\$

where $u, v \in \mathcal{B}(X)$. For example a shift space X is weakly mixing if for every $u, v \in \mathcal{B}(X)$ the set $N_{\sigma}([u], [v])$ contains arbitrarily long intervals of consecutive integers (see [13, Theorem 1.11]). A shift space X is transitive for σ^k where $k \in \mathbb{N}$ if and only if the set

 $N_{\sigma^k}([u], [v]) = \{\ell \in \mathbb{N} : uwv \in \mathcal{B}(X) \text{ for some } w \in \mathcal{A}^* \text{ such that } |uw| = k\ell\},$ is non-empty for every $u, v \in \mathcal{B}(X)$.

If a dynamical systems on a compact metric space is transitive, then there is a dense G_{δ} -set of points with dense orbit. In particular, in a transitive shift space X in every cylinder set there is a point x such that every block in $\mathcal{B}(X)$ occurs infinitely many times in x.

A synchronizing word for a shift space X is an element v of $\mathcal{B}(X)$ such that $uv, vw \in \mathcal{B}(X)$ for some blocks u, w over \mathcal{A} imply $uvw \in \mathcal{B}(X)$. A synchronized system is a shift space with a synchronizing word. Synchronized systems were introduced in [5]. Every synchronized system is coded. The uniqueness of the minimal right-resolving presentation known for sofic shifts extends to synchronized systems as outlined in [16, p. 451] (see also [20, p. 1241] and references therein). Synchronized systems and their generalizations were extensively studied in [11].

Let x_1, x_2, \ldots, x_n be positive integers. It is well-known that every sufficiently large integer can be represented as a non-negative integer linear combination of the x_i if and only if $gcd(x_1, x_2, \ldots, x_n) = 1$. For a later reference we formulate an important consequence of this result as a remark.

Remark 2.1. Let x_1, x_2, x_3, \ldots be positive integers. If $gcd(x_1, x_2, x_3, \ldots) = k$, then every sufficiently large multiple of k can be represented as a non-negative integer linear combination of the x_i .

3. TOTAL TRANSITIVITY IMPLIES MIXING

We prove that total transitivity, weak mixing and mixing are equivalent for coded systems. This leads to a structure theorem for coded systems which are not totally transitive.

Theorem 3.1. Suppose that X is a coded system and let $D \subset X$ be a closed set with nonempty interior such that $\sigma^k(D) \subset D$ for some k > 0. If the shift space (D, σ^k) is totally transitive, then it is mixing.

Proof. Let G be an irreducible countable labeled graph presenting X. Since D has nonempty interior, for each $r \in \mathbb{N}$ large enough there is $w \in \mathcal{B}(X)$ of length 2r + 1 such that the cylinder $[w] = \{x \in X : x_{[-r,r]} = w\}$ is contained in the interior of D.

We claim that the periodic points of (D, σ^k) are dense in D. Let V be a nonempty open subset of D. Then there is an open set $U \subset X$ with $V = U \cap D$. Without loss of generality we may assume that U is a cylinder set of some block $u \in \mathcal{B}(X)$. By total transitivity there is $s \in \mathcal{B}(X)$ and a path on G labeled by wsu such that the length

4

of ws is jk for some integer j. We can join the last vertex on the path labeled by wsu with the first vertex of the same path by a path labeled by a word t. Therefore there is a bi-infinite periodic sequence $y \in X$ such that $y_{[-r,\infty)} = (wsut)^{\infty}$ for some words $s, t \in \mathcal{B}(X)$. But then $y \in [w] \subset D$ and $\sigma^{jk}(y) \in [u]$ because |ws| = jk. On the other hand $y \in D$, hence $\sigma^{jk}(y) \in D$ and then $\sigma^{jk}(y) \in U \cap D = V$. This shows that periodic points are dense in D under σ^k . Since every totally transitive dynamical system with dense set of periodic points is weakly mixing (e.g. see [3]), it follows that (D, σ^k) is weakly mixing.

We proceed to a proof of mixing of (D, σ^k) . We recall that by [12, Lemma 3.1] a shift system (D, σ^k) is mixing if for each cylinder v in some neighborhood basis of a point with dense orbit in D the set $N_{\sigma^k}([v], [v])$ is cofinite. Let $\bar{x} \in [w]$ be a point whose orbit is dense in D under σ^k . Given $a \ge r$ set $v_a = \bar{x}_{[-a,a]}$ and

$$[v_a] = \{ x \in D : x_{[-a,a]} = v_a \}.$$

Note that $\bar{x} \in [v_a] \subset [w] \subset D$, hence the cylinder of v_a in X and in D coincide. Furthermore, $([v_a])_{a>r}$ is a neighborhood basis of \bar{x} .

It is now enough to show that the set $N_{\sigma^k}([v_a], [v_a])$ is cofinite. Let u be a word such that $v_a u$ is a labeling of a loop in G and let m be the length of $v_a u$. Without loss of generality we may assume that k divides m (we replace vu by $(v_a u)^k$ if necessary). Denote the loop presenting $v_a u$ on G by ξ . By weak mixing of (D, σ^k) the set $N_{\sigma^k}([v_a], [v_a])$ contains a set of m consecutive integers, hence there is an integer q > 0 such that for each $i = 1, \ldots, m$ the graph G contains a path η_i labeled $v_a v_i v_a$ where $|v_i| = (q+i)k - |v_a|$. Since G is irreducible, for each $i = 0, 1, \ldots, m$ there exists a path γ_i in G such that the following path is a loop on G:

 $\pi = \xi \gamma_0 \eta_1 \gamma_1 \eta_2 \dots \gamma_{m-1} \eta_m \gamma_m.$

Let $p = |\pi|$. We claim that for every $j \ge 1$ and $i = 1, \ldots, m$ we have

 $p + (mj) + (q+i)k \in N_{\sigma}([v_a], [v_a]).$

In order to show this, consider the labeling of the following path:

$$\eta_i \gamma_i \dots \eta_m \gamma_m(\xi)^j \gamma_0 \eta_1 \gamma_1 \dots \gamma_{i-1} \eta_i.$$

It starts and ends with v_a (it is a path on G because π and ξ are loops). This proves that $N_{\sigma^k}([v_a], [v_a])$ is cofinite.

Corollary 3.2. If a coded system X is totally transitive, then it is mixing.

Proof. Take D = X and apply Theorem 3.1.

We will now describe the structure of coded systems which are not totally transitive. Banks proved in [3] that if a dynamical system (X, T) is transitive, but (X, T^k) is not transitive for some k > 1 then there is a regular periodic decomposition of X, that is, one can find a finite cover $\{D_0, \ldots, D_{k-1}\}$ of X by non-empty regular closed sets with pairwise disjoint interiors such that $T(D_{i-1}) \subseteq D_{i \pmod{k}}$ for each $1 \leq i \leq k$. In this case we say that k is the length of the decomposition. Recall that a set is regular closed if it is the closure of its interior. If (D_i, T^n) is topologically mixing for some regular periodic decomposition $\mathcal{D} = \{D_0, \ldots, D_{k-1}\}$ of X then we say that T is relatively mixing with respect to \mathcal{D} . Observe that (X, T) is relatively mixing with respect to \mathcal{D} if and only if (D_0, T^n) is topologically mixing. It was proved in [3] that if there is an upper bound on the possible lengths of periodic decompositions of a transitive dynamical system then there exists a regular periodic decomposition $D_0, D_1, \ldots, D_{n-1}$ such that (D_i, T^n) is totally transitive for every $0 \leq i < n$ (this decomposition is called terminal).

Theorem 3.3. Every coded system is relatively mixing.

Proof. Let X be a coded system. By Theorem 3.1 and the result of Banks mentioned above, it suffices to show that there is an upper bound on the possible lengths of regular periodic decompositions of X.

Let G be an irreducible countable labeled graph presenting X and let k be the length of a cycle η in G. We claim that the length of a periodic decomposition of X can not be greater than k. On the contrary, assume that D_0, \ldots, D_{n-1} is a regular periodic decomposition and n > k. Since D_0 is regularly closed, there is $r \in \mathbb{N}$ and a word w of length 2r+1 such that the cylinder $[w] = \{x : x_{[-r,r]} = w\} \subset D_0$. Since for each i > 0 the interior of a regular closed set D_i is disjoint with the interior of D_0 , we have $D_i \cap [w] = \emptyset$ for each i > 0.

Since G is irreducible, there are paths π, γ with $|\pi| \ge |w|$ such that $\pi\eta\gamma$ is a cycle on G labeled $wu \in \mathcal{B}(X)$ for some word u. Repeating the path $\pi\eta\gamma$ if necessary we may assume that n divides $|\pi\eta\gamma| = |wu|$. Let a word $wu' \in \mathcal{B}(X)$ be the label of the path $\pi\eta\eta\gamma$ on G. We have |wu'| = nj + k for some $j \ge 1$. Note that $wu'w \in \mathcal{B}(X)$ because $\pi\eta\eta\gamma\pi$ is a path on G and hence there is $x \in X$ with $x_{[-r,t]} = wu'w$ for some t > r. But then x and $\sigma^{nj+k}(x)$ both belong to $[w] \subset D_0$. On the other hand

$$\sigma^{nj+k}(x) \in \sigma^{nj+k}(D_0) = \sigma^k(D_0) = D_k.$$

Since k < n, we have $[w] \cap D_k = \emptyset$ which leads to a contradiction.

4. Property P does not imply mixing

We construct a weakly mixing but not mixing spacing shift Y with the strong property P. This shows that the property P and topological mixing are not equivalent in general. Note that Y can not be coded system by Corollary 3.2. A similar example was first given by Blanchard [7], but our construction is much simpler.

Given $R \subset \mathbb{N}$ we define a spacing shift Ω_R as the set of all $x \in \{0, 1\}^{\mathbb{Z}}$ such that the condition $x_i = x_j = 1$ for some $i, j \in \mathbb{Z}$ with $i \neq j$ implies $|i - j| \in R$. Elements of $\mathcal{B}(\Omega_R)$ are called *R*-allowed blocks (see [4, 15] for more details).

Theorem 4.1. There is a non-mixing shift space Y with the strong property P.

Proof. We construct a *spacing shift* with the desired properties. Below we write $\langle n \rangle_2$ for the binary representation of a positive integer n, that is,

$$\langle 1 \rangle_2 = 1, \, \langle 2 \rangle_2 = 10, \, \langle 3 \rangle_2 = 11, \dots$$

Also for $u = u_1 \dots u_n \in \{0, 1\}^n$ we let $\Delta(u) = \{|i-j| : 1 \le j < i \le n, u_i = u_j = 1\}.$

Let $R = \mathbb{N} \setminus \{2^k : k \in \mathbb{N}\}$. Define $Y = \Omega_R$, and note that R is thick, thus Ω_R is nontrivial and weakly mixing (see [15]). We claim that for every $k \in \mathbb{N}$, $L = 2^k$, $w = 0^{2L}$ and any family of R-allowed blocks v_0, v_1, \ldots, v_t of length L, the block $u = v_0 w v_1 w v_2 \ldots v_{t-1} w v_t$ is also R-allowed. This clearly implies that the spacing shift Ω_R has the strong property P. A simple calculation yields that

$$\Delta(u) \subset \left(\{1, \dots, L-1\} \cap R\right) \cup \left(\bigcup_{m=0}^{\infty} \{(3m+2)L+1, \dots, (3m+4)L-1\}\right).$$

It is enough to show that no power of 2 is in $\Delta(u)$.

Note that any

$$q \in \bigcup_{m=0}^{\infty} \{(3m+2)L+1, \dots, (3m+4)L-1\}$$

can be written as q = a + b where $a \in \{2^{k+1} + 1, \ldots, 2^{k+2} - 1\}$ and $b = 3m \cdot 2^k$. If b = 0 then clearly q = a is not a power of 2, hence we may assume that m > 0. Then $\langle b \rangle_2 = \langle 3m \rangle_2 0^k$ and $\langle a \rangle_2 = 1x_k \ldots x_0$, where not all x_i 's are 0. Denote

$$\langle a+b\rangle_2 = y_l y_{l-1} \dots y_k \dots y_0.$$

Note that l > k + 1. If $x_i \neq 0$ for some i = 0, 1, ..., k - 1, then $y_i \neq 0$ and a + b is not a power of 2. If $x_0 = x_1 = ... = x_{k-1} = 0$ and $x_k = 1$, then $a = 3 \cdot 2^k$. In that case, a + b is also divisible by 3 and hence it is not a power of 2. Therefore $\Delta(u) \subset R$ and Ω_R has the strong property P.

On the other hand Ω_R is not topologically mixing because it is easy to see that $R = N([1]_R, [1]_R)$ is not cofinite which is a necessary condition for topological mixing (see [4], cf. [15]). Here $[1]_R = \{x \in \Omega_R : x_0 = 1\}$ is a nonempty open subset of Ω_R .

5. A mixing coded system without a generators of coprime length

We construct a mixing coded system without periodic points of odd period and such that every set of generators for this system contains only words of even length.

Let $t = t_0 t_1 t_2 \ldots = 10010110 \ldots$ be the Prouhet-Thue-Morse sequence (see [1]). Recall that it obeys $t_{2n} = t_n$ and $t_{2n+1} = 1 - t_n$. It is well-known that t is a cube-free sequence, hence neither 000 nor 111 occur in t. Furthermore, $\mathcal{B}(t)$ is a language of a minimal and non-periodic shift space X_{TM} .

We first define auxiliary sets $L_n \subset \{0,1\}^*$ for n = 1, 2, ... and a sequence of words $\{a_k\}_{k=0}^{\infty}$. We begin by setting $a_0 = 01$ and $L_1 := \{a_0\}$. Assume that we have performed n-1 steps of our construction $(n \in \mathbb{N})$. We are given the set L_{n-1} and $\{a_k\}_{k=0}^{\infty}$ is defined for indices $0, 1, ..., s_n - 1$, that is, s_n denotes the number of words in the sequence $\{a_k\}$ constructed up to the step n. In particular, we have $s_1 = 0$ and $s_2 = 1$. At each step $n \ge 2$ we enumerate the blocks in L_{n-1} starting from s_n , that is, we write

$$L_{n-1} = \{w_{s_n}, \dots, w_{s_n+|L_{n-1}|-1}\}$$

We extend the sequence $\{a_k\}$ by adding words

$$a_j = 01110t_{[0,4j-3]}011110w_j011110t_{[0,4j-1]}01110.$$

for $j = s_n, ..., s_n + |L_{n-1}| - 1$. Then we set

$$L_n := \left\{ a_{s_n}, a_{s_n+1}, \dots, a_{s_{n+1}-1} \right\} \cup \bigcup_{k=1}^n L_n^k,$$

where $L_n^k = \{w_1 w_2 \dots w_k : w_j \in L_n \text{ for } j = 1, \dots, k\}$. This completes the step n and our induction. Let $\mathcal{Q} := \{a_i ; i \in \mathbb{N}_0\}$.

We will call the words 01110 and 011110 markers. Note that a_0 is the only element of Q without markers, and since 111 is never a subblock of $\mathcal{B}(t)$ we can identify positions of all markers in a_k and therefore we can identify also positions of blocks $t_{[0,4j-3]}$ and $t_{[0,4j-1]}$. Hence knowing that $w \in Q$ and the length of the longest subblock from $\mathcal{B}(t)$ in w between two markers (when $w \neq a_0$) we can uniquely determine k such that $w = a_k$.

Notice that

(1)
$$L_n \subset L_{n+1}$$
 and $Q^+ = \bigcup_{n=1}^{\infty} L_n$,

thus Ω and $\bigcup_{n=1}^{\infty} L_n$ generate the same coded system denoted by X.

Lemma 5.1. The coded system X is mixing.

Proof. Every block $u \in \mathcal{B}(\mathbb{X})$ is a subword of some concatenation of generators. Therefore it is enough to show that for any $k, \ell \in \mathbb{N}$ and $u_1, \ldots, u_k, v_1 \ldots v_\ell \in \Omega$ there is $M \in \mathbb{N}$ such that for all m > M there is a block $w \in \{0, 1\}^m$ with $u_1 \ldots u_\ell w v_1 \ldots v_k \in \mathcal{B}(\mathbb{X})$. Observe that by (1) there is $n \in \mathbb{N}$ such that $u_1, \ldots, u_k \in L_n$. Clearly, we may also assume that n > k. Then it follows directly from the construction that $u = u_1 \ldots u_k \in L_{n+1}$ and there is $q \in \mathbb{N}$ such that

$$a_q = 01110t_{[0,4q-3]}011110u011110t_{[0,4q-1]}01110 \in L_{n+2}.$$

Define $s_{2q-1} = t_{[0,4q-3]}$ and $s_{2q} = t_{[0,4q-1]}$. Set M = 4q + 11. If m > M is odd, then we have

$$a_{q}a_{0}^{\frac{m-4q-11}{2}}v_{1}\ldots v_{k} = 01110s_{2q-1}011110\underbrace{u_{1}\ldots u_{\ell}}_{u}\underbrace{011110s_{2q}01110(01)^{\frac{m-4q-11}{2}}}_{w\in\{0,1\}^{m}}v_{1}\ldots v_{k}\in \mathbb{Q}^{+}.$$

For even m > M we have

$$ua_0^{\frac{m}{2}}v_1\dots v_k = u(01)^{\frac{m}{2}}v_1\dots v_k \in \mathbb{Q}^+$$

This completes the proof, since $Q^+ \subset \mathcal{B}(\mathbb{X})$.

Lemma 5.2. If X is a non trivial coded system generated by a set Q and there is a word $w \in Q$ with odd length, then X contains a periodic point of odd prime period greater than one.

Proof. Since X is nontrivial, there is a word in $\mathcal{B}(X)$ containing symbol 0 and a word containing symbol 1. Hence there is a non-constant word u in Q^+ , thus $uu \in Q^*$ is a non-constant word of even length. Then the word uuw is a non-constant word of odd length k in Q^+ whose infinite concatenation is a non-constant periodic point with an odd prime period dividing k.

We are going to prove that \mathbb{X} has no periodic points with odd period. We first show that an odd periodic point cannot occur in one of the sofic shifts Y_n generated by L_n .

Lemma 5.3. For every $n \in \mathbb{N}$ the sofic shift Y_n generated by the set L_n does not contain a periodic point with odd prime period.

Proof. Fix any $n \in \mathbb{N}$ and let be a point $x \in Y_n$ with prime period q. Clearly, q > 1 because the lengths of runs of 0's and 1's in $\mathcal{B}(\mathbb{X})$ are bounded. If x does not contain any marker then $x = (01)^{\infty}$ and q is even. Thus we may assume that there are (infinitely many) markers in x. Let ℓ be the length of the longest block w from $\mathcal{B}(t)$ appearing in x between two markers. Then $\ell = 4k$ for some k, and there must be $j \in \mathbb{Z}$ such that $x_{[j,j+|a_k|-1]} = a_k$ (no word a_r with r > k can appear in x, since then we would have $\ell \ge 4k + 4$). Because x has period q, we also have $x_{[j+q,j+q+|a_k|-1]} = a_k$. Let $w \in Q^+$ be a word which contains $x_{[j,j+q+|a_k|-1]}$ as a subblock and which does not contain a block from $\mathcal{B}(t)$ of length greater then 4k. Write $w = v_1 \dots v_t$ where $v_j \in Q$. By the above observation, we have that $x_{[j,j+|a_k|-1]} = x_{[j+q,j+q+|a_k|-1]} = a_k \in Q$ must be among v_i . This immediately implies that $x_{[j,j+q-1]} \in Q^+$ and since all words in Q have even length, we find that $q = |x_{[j,j+q-1]}|$ is even.

Finally we show that taking the closure of $\bigcup_{n=1}^{\infty}Y_n$ does not introduce periodic points.

Lemma 5.4. Any element x of $\mathbb{X} \setminus \bigcup_{n=1}^{\infty} Y_n$ contains arbitrary long blocks from $\mathcal{B}(t)$. In particular, x cannot be periodic.

Proof. Assume on the contrary that there is $x \in \mathbb{X} \setminus \bigcup_{n=1}^{\infty} Y_n$ such that the longest block from $\mathcal{B}(t)$ appearing in x has length at most 4k for some integer

k > 0. Then x must contain infinitely many markers as subwords, as otherwise $x = \ldots a_0 a_0 w a_0 a_0 \ldots$ for some $w \in L_n$ and some n, thus $x \in Y_n$.

There exists an infinite set $J \subset \mathbb{Z}$ and a strictly increasing infinite sequence of integers $(n_i)_{i \in J}$ such that $x_{[n_i,n_i+4]} = 01110$ if and only if there is $i \in J$ such that $j = n_i$. Let m be the least integer such that $a_k \in L_m$.

We claim that every word $w_i := x_{[n_{-i},...,n_i+4]}$ is contained in $\mathcal{B}(Y_m)$. Let j be the smallest positive integer, such that $w_i \in \mathcal{B}(Y_j)$. There must be some word $g \in L_j^+$ such that $g = bw_i c$ with $b, c \in \mathcal{B}(\mathbb{X})$. Either $j \leq m$, in which case we are done, or j > m and g must contain a_ℓ for some $\ell > k$. Since w_i starts and ends with 01110, the two longest blocks of symbols from $\mathcal{B}(t)$ occurring in a_ℓ must be contained in b and c and thus w_i is already contained in the middle word of a_ℓ , which is an element of L_{j-1} . This contradicts the minimality of j. Therefore our claim holds. If J is bi-infinite, then $x \in Y_m$ which is a contradiction. Otherwise, either

$$a = \inf\{n_i : i \in J\} = \min\{n_i : i \in J\} > -\infty, \text{ or} b = \sup\{n_i : i \in J\} = \max\{n_i : i \in J\} < \infty.$$

If $a > -\infty$, then $x_{(-\infty,a-1]}$ does not contain markers and $x_{[a,a+4]} = 011110$ which implies that $x_{(-\infty,a-1]} = \dots a_0 a_0 a_0$. In the second case $b < \infty$ and we obtain that $x_{[b+1,\infty)} = a_0 a_0 a_0 \dots$, hence both cases imply that $x \in Y_m$ completing the proof.

Theorem 5.5. There exists a coded system X which is mixing, but does not have periodic points with odd periods. In particular, every set of generators for X contains only blocks of even length.

Proof. The shift X = X is mixing by Lemma 5.1. Combining Lemma 5.3 and 5.4 we see that X does not contain a periodic point with odd prime period. Then it follows from Theorem 5.2 that every set of generators for X contains only blocks of even length.

6. Strong property P and mixing coded systems

As we have seen before, the strong property P does not imply topological mixing. Clearly, the converse implication is neither true, since Blanchard [7] proved that the property P implies positive topological entropy and there are examples of mixing shifts with zero topological entropy.

The purpose of this section is to show that every mixing coded system has strong property P.

Lemma 6.1. Let X be a mixing coded system, Q be its generator, and $k = gcd\{|u| : u \in Q\}$. Then for each $u \in \mathcal{B}(X)$ with $|u| = 0 \mod k$ there is a word $v \in Q^+$ such that v = aub with $|a| = 0 \mod k$ and $|b| = 0 \mod k$.

Proof. Since X is mixing, there is a word $v \in Q^+$ such that for some $\tilde{a}, \tilde{b}, z_1, \ldots, z_k \in \mathcal{B}(X)$ we have

$$v = \tilde{a}uz_1uz_2uz_3\dots uz_k \tilde{b}$$

with $|z_i| = 1 \mod k$ for each $i = 1, \ldots, k$. Replacing v by v^k if necessary, we may assume that $|v| = 0 \mod k$. There is $\ell \in \{0, \ldots, k-1\}$ such that $|\tilde{a}| = -\ell \mod k$. Let $a = \tilde{a}uz_1uz_2\ldots uz_\ell$ and $b = z_{\ell+1}uz_{\ell+2}\ldots uz_k\tilde{b}$ and observe that $|a| = 0 \mod k$, $|b| = |v| - |a| - |u| = 0 \mod k$ and v = aub.

Theorem 6.2. If X is a mixing coded system, then X has the strong property P.

10

Proof. Blanchard [7, Proposition 4] proved that a shift space X over \mathcal{A} has the property P if for any integer p belonging to some infinite strictly increasing sequence of integers there exists an integer q = q(p) such for any $k \geq 2$ and any words $u_1, \ldots, u_k \in \mathcal{B}_p(X)$ there are words $w_1, \ldots, w_{k-1} \in \mathcal{B}_q(X)$ such that $u_1w_1u_2 \ldots u_{k-1}w_{k-1}u_k \in \mathcal{B}(X)$. Let Q be a set of generators of X and $k = \gcd\{|u| : u \in Q\}$.

We will show that Blanchard's criterion [7, Proposition 4] applies to any $p \in \{\ell \in \mathbb{N} : \ell = 0 \mod k\}$. To this end, fix $p = 0 \mod k$ and enumerate all blocks of length p by v_1, \ldots, v_n . We use Lemma 6.1 to obtain $a_1, \ldots, a_n, b_1, \ldots, b_n \in \mathcal{B}(X)$ such that

$$0 = |a_1| = \dots = |a_n| = |b_1| = \dots = |b_n| \mod k$$

and $a_i v_i b_i \in Q^+$ for $i \in \{1, \ldots, n\}$. By Remark 2.1 there exists N such that for all n > N there is a word $w \in Q^+$ with |w| = kn, thus we can find $c_1, \ldots, c_n \in Q^+$ and $d_1, \ldots, d_n \in Q^+$ such that $q' = |c_1 a_1| = \cdots = |c_n a_n|$ and $q'' = |b_1 d_1| = \cdots = |b_n d_n|$. Now let u_1, \ldots, u_k be any words in $\mathcal{B}_p(X)$. Then there is a function $\varphi \colon \{1, \ldots, k\} \to \{1, \ldots, n\}$ with $u_i = v_{\varphi(i)}$. Therefore

$$a_{\varphi(1)}u_1(\underbrace{b_{\varphi(1)}d_{\varphi(1)}c_{\varphi(2)}a_{\varphi(2)}}_{w_1})u_2(\underbrace{b_{\varphi(2)}d_{\varphi(2)}c_{\varphi(3)}d_{\varphi(3)}}_{w_2})\dots c_{\varphi(n)}a_{\varphi(n)}u_nb_{\varphi(n)} \in Q^+.$$

We set q(p) = q' + q'' and we obtain that X has the strong property P by [7, Proposition 4].

7. Two folklore results

We finish the paper with two results which are probably folklore, but we were unable to find them in the literature so we attach them for completeness. Combining them with Corollary 3.2 we obtain that the stronger forms of transitivity mentioned in the Introduction are equivalent for synchronized systems.

Lemma 7.1. Let X be a coded system presented by a labelled graph G. If there are two cycles on G with relatively prime lengths, then X is mixing.

Proof. Denote by α_1 and α_2 two cycles on G with relatively prime lengths, $k_j = |\alpha_j|, j = 1, 2$. Let e_j be a vertex of some edge belonging to α_j for j = 1, 2. Let u_j be the label of α_j for j = 1, 2 read off traversing α_j from e_j . Take any words $w_1, w_2 \in \mathcal{B}(X)$. Since X is coded, we can find paths γ_1, γ_2 on G labelled, respectively, by w_1, w_2 . Let a_2 be the initial vertex of γ_2 , and b_1 be the terminal vertex of γ_1 . Let ℓ_1 (ℓ_2) be the length of the shortest path π_1 (π_2) on G from b_1 to e_1 (from e_2 to a_2) and m be the length of the the shortest path ρ on G from e_1 to e_2 . Let $v_1, v_2, z \in \mathcal{B}(X)$ be labels of π_1, π_2, ρ , respectively. It follows that for each $p, q \in \mathbb{N}$ the path

$\pi_{pq} = \gamma_1 \pi_1(\alpha_1)^p \rho(\alpha_2)^q \pi_2$

is labeled by $w_1v_1(u_1)^p z(u_2)^q v_2w_2$. Since k_1 and k_2 are relatively prime, the set $\{pk_1 + qk_2 : p, q \in \mathbb{N}\}$ is cofinite. Therefore there is N > 0 such that if we fix any $t \geq N$ then we can find $p = p(t), q = q(t) \in \mathbb{N}$ so that $t = \ell_1 + p_t k_1 + m + q_t k_2 + \ell_2$ and the path $\pi_{p(t)q(t)}$ on G has length t. Therefore for each $n \geq N - \ell_1$ there is a word w of length n such that $w_1ww_2 \in \mathcal{B}(X)$ and hence X is mixing. \Box

Lemma 7.2. A synchronized shift X is topologically mixing if and only if there are two closed paths with relatively prime lengths in its Fisher cover.

Proof. The "if" part follows from Lemma 7.1. For the "only if" part assume that X is topologically mixing. Let w be a synchronizing word for X. Then w is a magic word for the Fisher cover (G, Θ) of X, that is, there is a vertex e of G such that every path labelled by w ends at e. Since X is mixing there is $N \in \mathbb{N}$ and there are

words u_1, u_2 with $|u_1| = N$, $|u_2| = N + 1$ such that $wu_1w, wu_2w \in \mathcal{B}(X)$. Because each path labelled by w ends at e, there are closed paths in G labelled by u_1w and u_2w with relatively prime lengths.

The following theorem summarizes our results on connections between variants of transitivity for coded systems.

Theorem 7.3. Let X be a non trivial coded system. Then the following conditions are equivalent:

- (a) X is topologically mixing;
- (b) X has the strong property P;
- (c) X is topologically weakly mixing;
- (d) X is totally transitive.

Additionally, if X is synchronized, then any of the above conditions is equivalent to

(e) X has two periodic points with relatively prime primary periods.

Moreover, there exists a coded system X fulfilling (a)-(d), but not (e).

Acknowledgements

The research of P. Oprocha was supported by the Polish Ministry of Science and Higher Education from sources for science in the years 2013–2014, Grant No. IP2012 004272. The research of D. Kwietniak was supported by the National Science Centre (NCN) under grant no. DEC-2012/07/E/ST1/00185. The research of J. Epperlein was partly supported by the German Research Foundation (DFG) through the Cluster of Excellence (EXC 1056), Center for Advancing Electronics Dresden (cfaed).

References

- Allouche, Jean-Paul; Shallit, Jeffrey. The ubiquitous Prouhet-Thue-Morse sequence. Sequences and their applications (Singapore, 1998), 1–16, Springer Ser. Discrete Math. Theor. Comput. Sci., Springer, London, 1999. MR1843077 (2002e:11025)
- S. Baker, A. Ghenciu, Dynamical properties of S-gap shifts and other shift spaces, arXiv preprint arXiv:1411.7167 (2014).
- J. Banks, Regular periodic decompositions for topologically transitive maps. Ergodic Theory Dynam. Systems, 17 (1997), no. 3, 505–529.
- [4] J. Banks, T. D. Nguyen, P. Oprocha, B. Stanley and B. Trotta, *Dynamics of spacing shifts*. Discrete Contin. Dyn. Syst., **33** (2013), no. 9, 4207–4232.
- [5] F. Blanchard, G. Hansel, G. Systèmes codés (French) [Coded systems], Theoret. Comput. Sci. 44 (1986), no. 1, 17–49.
- [6] F. Blanchard, G. Hansel, Systèmes codéss et limites de systèmes sofiques. (French) [Coded systems and limits of sofic systems] C. R. Acad. Sci. Paris Sèr. I Math., 303 (1986), 475–477.
- [7] F. Blanchard, Fully positive topological entropy and topological mixing in Symbolic dynamics and its applications (New Haven, CT, 1991), 95–105, Contemp. Math., 135, Amer. Math. Soc., Providence, RI, 1992.
- [8] V. Climenhaga, D. Thompson, Intrinsic ergodicity beyond specification: β-shifts, S-gap shifts, and their factors, Israel J. Math. 192 (2012), no. 2, 785–817.
- [9] V. Climenhaga, D. Thompson, Intrinsic ergodicity via obstruction entropies, Ergodic Theory Dynam. Systems 34 (2014), no. 6, 1816–1831.
- [10] D. Dastjerdi, S. Jangjoo, Dynamics and topology of S-gap shifts Topology and its Applications 159 (2012), no. 10, 2654–2661.
- [11] D. Fiebig, U.-R. Fiebig, Covers for coded systems. Symbolic dynamics and its applications (New Haven, CT, 1991), 139–179, Contemp. Math., 135, Amer. Math. Soc., Providence, RI, 1992.
- [12] G. Harańczyk, D. Kwietniak, P. Oprocha, A note on transitivity, sensitivity and chaos for graph maps. J. Difference Equ. Appl., 17 (2011), 1549–1553.
- [13] E. Glasner, Ergodic Theory via Joinings, Mathematical Surveys and Monographs, 101. American Mathematical Society, Providence, RI, 2003. xii+384 pp.

- [14] W. Krieger. On subshifts and topological Markov chains. Numbers, Information and Complexity. Eds. I. Althöfer et al. Kluwer, 2000, pp. 453–472.
- [15] K. Lau, A. Zame, On weak mixing of cascades. Math. Systems Theory. 6 (1972/73), 307–311.
- [16] D. Lind and B. Marcus, An Introduction to Symbolic Dynamics, Cambridge University Press, New York, 1995.
- [17] T. Meyerovitch, Tail invariant measures of the Dyck shift, Israel Journal of Mathematics 163, (2008), no. 1, 61–83.
- [18] T. Meyerovitch, Gibbs and equilibrium measures for some families of subshifts, Ergodic Theory Dynam. Systems 33 (2013), no. 3, 934–953.
- [19] D.J. Thompson, Irregular sets, the β-transformation and the almost specification property, Trans. Amer. Math. Soc., 364 (2012), 5395–5414.
- [20] K. Thomsen, On the ergodic theory of synchronized systems, Ergodic Theory Dynam. Systems 26 (2006), no. 4, 1235–1256.

(J. Epperlein) INSTITUTE FOR ANALYSIS AND CENTER FOR DYNAMICS, TECHNISCHE UNIVER-SITÄT DRESDEN, ZELLESCHER WEG 12-14, 01069 DRESDEN, GERMANY

E-mail address: jeremias.epperlein@tu-dresden.de

 (D. Kwietniak) Faculty of Mathematics and Computer Science, Jagiellonian University in Krakow, ul. Łojasiewicza 6, 30-348 Kraków, Poland

E-mail address: dominik.kwietniak@uj.edu.pl *URL*: www.im.uj.edu.pl/DominikKwietniak/

(P. Oprocha) AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY, FACULTY OF APPLIED MATHEMATICS, AL. MICKIEWICZA 30, 30-059 KRAKÓW, POLAND

E-mail address: oprocha@agh.edu.pl