# arXiv:1503.05233v3 [quant-ph] 14 Dec 2015

# Quantum Model of Cooling and Force Sensing With an Optically Trapped Nanoparticle

B. Rodenburg<sup>\*</sup>,<sup>1,2</sup> L. P. Neukirch,<sup>2,3</sup> A. N. Vamivakas,<sup>2,4</sup> and M. Bhattacharya<sup>1,2</sup>

<sup>1</sup>School of Physics and Astronomy, Rochester Institute of Technology, Rochester, NY 14623, USA

<sup>2</sup>Center for Coherence and Quantum Optics, University of Rochester, Rochester, NY 14627, USA\*

<sup>3</sup>Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627, USA

<sup>4</sup>Institute of Optics, University of Rochester, Rochester, NY 14627, USA

(Dated: March 1, 2022)

Optically trapped nanoparticles have recently emerged as exciting candidates for tests of quantum mechanics at the macroscale and as versatile platforms for ultrasensitive metrology. Recent experiments have demonstrated parametric feedback cooling, nonequilibrium physics, and temperature detection, all in the classical regime. Here we provide the first quantum model for trapped nanoparticle cooling and force sensing. In contrast to existing theories, our work indicates that the nanomechanical ground state may be prepared without using an optical resonator; that the cooling mechanism corresponds to nonlinear friction; and that the energy loss during cooling is nonexponential in time. Our results show excellent agreement with experimental data in the classical limit, and constitute an underlying theoretical framework for experiments aiming at ground state preparation. Our theory also addresses the optimization of, and the fundamental quantum limit to, force sensing, thus providing theoretical direction to ongoing searches for ultra-weak forces using levitated nanoparticles.

PACS numbers: 42.50.-p, 42.50.Wk, 37.10.Vz, 62.25.-g

### INTRODUCTION

Optically trapped nanoparticles can support explorations of macroscopic quantum mechanics as well as ultrasensitive metrology very well since they can be isolated from the environment in a trap, cooled, and detected all using a single laser beam without the need for an optical cavity [1–5]. Experiments with optically trapped harmonically oscillating subwavelength dielectric particles [6–10] (see Fig. 1) have recently realized feedback cooling [1, 11], nonlinear dynamics [2], non-equilibrium physics [4], coupling to spin degrees of freedom [12] and thermometry [3]. All experiments thus far have been carried out in the classical regime.

While several groups are currently exploring ways to access the nonclassical regime of such systems, we present here the first quantum theory of trapped nanoparticle optical feedback cooling and force sensing. The impetus for investigating cooling comes from the fact that although levitated particles have been successfully cooled in optical resonators (to 10 K [13] and 64 K [14]), cavityless cooling has been able to reach much lower temperatures (50 mK [11]) starting from the same initial (room) temperature. It is however an open theoretical question as to whether the quantum ground state can be prepared without using a cavity. Our model addresses this important question and reveals a viable route to nanoparticle ground state preparation for ongoing cavityless experiments. It also identifies the dissipative mechanism underlying experimental cooling to be nonlinear in nature, in contrast to standard experimental techniques, which depend on linear damping, and standard theory, which relies on linear response analysis [14–20]. Lastly, the model yields analytical results for the ensuing nonexponential decay of phonon number, which shows excellent agreement with our experimental data in the classical regime. The motivation for investigating force detection is provided by the use of levitated cavityless nanoparticles in several ongoing searches for various ultraweak forces [5, 21–23]. Again it is an open question as to whether and to what extent force detection is limited by the effects of quantum backaction in those systems. To clarify this issue we derive in this article expressions for the force sensitivity and the standard quantum limit of force detection. We expect our new results on cooling and force sensing in the quantum limit will be invaluable as this still nascent field matures.

A schematic of the physical system under consideration is shown in Fig. 1. A subwavelength polarizable dielectric sphere is confined at the focus of a Gaussian trapping beam, and its motion is detected using a probe beam, polarized orthogonal to the trap. The detected signal is processed and fed back to the trap beam to cool the particle. We analyze this configuration by dividing it into a 'system' and a 'bath'. The system consists of the nanomechanical oscillator and the optical probe and trap. The bath consists of the optical modes into which light is scattered by the nanosphere, and the background



FIG. 1: a) Image of the trapped nanoparticle. b) Schematic of the experiment modeled in this article.

thermal gas present in the experiment. We proceed to derive a quantum model by identifying the electromagnetic modes relevant to the problem, constructing the system and bath Hamiltonians, and deriving the master equation for the system [24, 25]. All conclusions stated in this article follow from this master equation.

### MODEL

The configuration Hamiltonian can be written as,

$$H = H_m + H_f + H_{\text{int}}.$$
 (1)

In Eq. (1), the first term on the right hand side represents the mechanical kinetic energy  $H_m = |\mathbf{p}|^2 / 2m$ , where  $\mathbf{p}$ is the three dimensional momentum of the nanoparticle, and m its mass. The second term in Eq. (1) is the field energy  $H_f = \epsilon_0 \int |\mathbf{E}(\mathbf{r})|^2 d^3 \mathbf{r}$ , where  $\mathbf{E}(\mathbf{r})$  is the sum of the trap  $\mathbf{E}_t$ , probe  $\mathbf{E}_p$ , and background  $\mathbf{E}_b$  electric fields. We model the trap and probe modes as Gaussian beams, and the background using a plane wave expansion. We find after dropping a constant term,

$$H_f = \hbar \omega_p a^{\dagger} a + \sum_{\mu} \int d^3 \mathbf{k} \hbar \omega_{\mathbf{k}} a^{\dagger}_{\mu}(\mathbf{k}) a_{\mu}(\mathbf{k}), \qquad (2)$$

which is simply the sum of the probe and background field energies with a and  $a_{\mu}(\mathbf{k})$  representing the corresponding standard bosonic field operators. Finally, in Eq. (1) the interaction Hamiltonian is given by  $H_{\text{int}} = -\int_{V} \mathbf{P}(\mathbf{r}) \cdot \mathbf{E}(\mathbf{r}) \, \mathrm{d}^{3}\mathbf{r}/2 = -\alpha_{p} \int_{V} \left|\mathbf{E}(\mathbf{r})\right|^{2} \mathrm{d}^{3}\mathbf{r}/2$ , where we have assumed that the dielectric has a volume V, and that it has a linear polarizability density  $\alpha_{p}$ , i.e. the polarization density is  $\mathbf{P}(\mathbf{r}) = \alpha_{p} \mathbf{E}(\mathbf{r})$ . Using the expressions for the electric fields from the Supplementary Material we can evaluate  $H_{\text{int}}$  for small particle displacements  $\mathbf{r}$ , and rewrite Eq. (1) as

$$H = H_S + H_B + H_{SB},\tag{3}$$

where this system Hamiltonian is

$$H_S = \hbar \omega_p a^{\dagger} a + \sum_j \hbar \omega_j b_j^{\dagger} b_j - \sum_j \hbar g_j a^{\dagger} a (b_j + b_j^{\dagger}), \quad (4)$$

with mechanical trapping frequencies  $\omega_j$ , optomechanical coupling constants  $g_j$ , and mechanical operators which obey the standard commutation relations  $[b_j, b_j^{\dagger}] = 1$  $(j = \{x, y, z\})$ . In Eq. (3), the bath Hamiltonian is  $H_B = \sum_{\mu} \int d^3 \mathbf{k} \hbar \omega_{\mathbf{k}} a_{\mu}^{\dagger}(\mathbf{k}) a_{\mu}(\mathbf{k})$  and the system-bath interaction Hamiltonian is  $H_{SB} = -\epsilon_c \epsilon_0 \int_V d^3 \mathbf{r} [\mathbf{E}_{\mathbf{t}}(\mathbf{r}) + \mathbf{E}_{\mathbf{p}}(\mathbf{r})] \cdot \mathbf{E}_{\mathbf{b}}(\mathbf{r})$ , which represents the scattering of the trap and probe fields into the background (see Supplementary Material for details).

We now trace over the bath modes, applying the standard Born and Markov approximations, since the systembath coupling is weak and the bath correlations decay quickly [17, 24]. We also trace over the x and y degrees of particle motion, since the dynamics along the three axes are independent of each other, and it suffices to analyze a single direction [11]. The net result of our calculation is a master equation for the density matrix  $\rho(t)$  describing the optical probe and the z-motion of the nanoparticle

$$\dot{\rho}(t) = \frac{1}{i\hbar} [H'_S, \rho] - \frac{A_t}{2} \mathcal{D}[Q] \rho + \mathcal{L}_{\rm sc} \rho, \qquad (5)$$

where the first term on the right hand side represents unitary evolution of the system with  $H'_S = \hbar \omega_p a^{\dagger} a + \hbar \omega_z b_z^{\dagger} b_z - \hbar g_z a^{\dagger} a (b_z + b_z^{\dagger})$ . The second term corresponds to the positional decoherence of the nanoparticle due to scattering of trap photons, with the Lindblad superoperator  $\mathcal{D}[Q_z]\rho \equiv Q_z^{\dagger}Q_z\rho + \rho Q_z^{\dagger}Q_z - 2Q_z\rho Q_z^{\dagger}$ , where  $Q_z = b_z^{\dagger} + b_z$ , and  $A_t$  is the heating rate due to trap beam scattering as defined in the Supplementary Material. The third superoperator describes the loss of photons from the probe, also due to scattering by the nanoparticle,  $\mathcal{L}_{\rm sc}[\rho(t)] = -B\left(\mathcal{D}[a] + (7\omega_p^2 \ell_z^2/5c^2)\mathcal{D}[aQ_z]\right)\rho$ , where  $\ell_z$ is the oscillator length. The nanoparticle also experiences collisions with background gas particles at the ambient temperature T. This effect may be accounted for by adding to the right hand of Eq. (5) the superoperator [26]

$$\mathcal{B}[\rho(t)] = -\frac{D_p}{2} \mathcal{D}[Q_z]\rho - \frac{D_q}{2} \mathcal{D}[P_z]\rho - i\frac{\eta_f}{4m} \left[Q_z, \{P_z, \rho\}\right],$$
(6)

where  $P_z = i(b_z^{\dagger} - b_z)$ , and curly braces denote an anticommutator. The first term on the right hand side corresponds to momentum diffusion and  $D_p = 2\eta_f k_B T \ell_z^2/\hbar^2$ , where  $k_B$  is Boltzmann's constant. The second term describes position diffusion with  $D_q = \eta_f \hbar^2/(24k_B T m^2 \ell_z^2)$ . The third term accounts for friction, and by Stokes law we have  $\eta_f = 6\pi \mu r_d$ , where  $r_d$  is the radius of the nanoparticle and  $\mu$  is the dynamic viscosity of the background gas. As shown earlier, internal and center-ofmass heating of the nanoparticle due to optical absorption and blackbody radiation are negligible in systems such as ours, as are particle size and shape effects, as well as trap beam shot noise [11, 27].

We now characterize the measurement of the oscillator displacement using input-output theory from quantum optics [28] applied to the nanoparticle. Specifically, the incoming probe field  $a_{in}$  interacts with the nanoparticle, and the outgoing probe field  $a_{out}$  carries a signature of this interaction (as shown in the Supplementary Material)

$$a_{\rm out} = a_{\rm in} + \frac{\alpha \chi}{2} Q_z(t), \tag{7}$$

where  $\chi = 4g_z \Delta t$  is the scaled optomechanical coupling, with integration time  $\Delta t$  (determined by the detection bandwidth), and we have written the probe beam as a coherent state  $a = -i\alpha + v$ , with  $\alpha$  a classical number and v a bosonic annihilation operator. A homodyne measurement on the output field yields a current [28]

$$I_h = \chi^2 \Phi \langle Q_z \rangle(t) + \sqrt{\chi^2 \Phi} \xi(t), \qquad (8)$$

where  $\Phi = \alpha^2 \Delta \omega$  is the average detected flux of probe photons, and  $\xi(t)$  is a stochastic variable with mean  $\langle \xi(t) \rangle = 0$  and correlation  $\langle \xi(t)\xi(t') \rangle = \delta(t-t')$ .

In the experiment, the detected current  $I_h$  is frequency doubled, phase shifted, and fed back to modulate the power of the trapping beam [11]. This results in a feedback Hamiltonian  $H_{\rm fb} = \hbar G I_{\rm fb} Q_z^3$ , where G is the dimensionless feedback gain related to the trap intensity modulation [5, 29],

$$M \equiv \frac{\Delta I_t}{I_t} \approx \frac{G\chi^2 \Phi \langle b_z^{\dagger} b_z \rangle}{\omega_z}, \tag{9}$$



FIG. 2: a) Shows the diffusive evolution of the trapped nanoparticle's position at atmospheric pressure. b) Harmonic motion of the nanoparticle at a lower pressure of  $4 \times 10^{-3}$  mbar. The reduction of the amplitude of harmonic motion corresponds to the turn-on of feedback, i.e. cooling.

and the feedback current is  $I_{\rm fb} = \chi^2 \Phi \langle P_z \rangle + \sqrt{\chi^2 \Phi} \xi'(t)$ , where  $\xi'(t)$  has the same properties as  $\xi(t)$ . This form of the Hamiltonian implies a feedback force  $F_{\rm fb} = -\partial H_{\rm fb}/\partial Q_z$  which is equivalent to that used in experiments in the classical regime [5]. Taking the Markovian limit where the feedback occurs faster than any system timescale, and applying quantum feedback theory for homodyne detection [30], we find that the following superoperator must be added to Eq. (5)

$$\mathcal{F}[\rho(t)] = -i\chi^2 \Phi G[Q_z^3, \{P_z, \rho\}] - \frac{\chi^2 \Phi}{2} G^2 \mathcal{D}[Q_z^3]\rho, \quad (10)$$

where the first term on the right hand side represents the desired cooling effect of the feedback, and the second term the accompanying backaction. We emphasize that in contrast to standard optomechanics, the feedback and backaction terms are highly nonlinear in the oscillator variables. The presence of this nonlinearity distinguishes our system from conventional cavity optomechanics and results in qualitatively different dynamics, as we show below.

The full master equation, assembled from Eqs. (5), (6)



FIG. 3: a) The y and z phonon cooling dynamics [Eq. (13)]. b) Steady state phonon number versus pressure [Eq. (14a)]. Circles represent experimental data, and the solid curve our theoretical model for a fused silica sphere ( $\epsilon_r = 2.1$  and density = 2200 kg/m<sup>3</sup>) radius  $r_d = 50$  nm, 1064 nm trap (100 mW) and probe (10 mW) beams, a mechanical frequency  $\omega_z/2\pi = 38$  kHz,  $\chi \approx 10^{-7}$ , and trap intensity modulation  $M \lesssim 0.1\%$ . The dotted lines represent the equivalent curves for one of the transverse degrees of freedom ( $\omega_y/2\pi = 138$  kHz). The dashed curve in a) represents the prediction of our theory for a setup placed in a cryostat with the feedback chosen optimally, keeping  $M \leq 10\%$ .

and (10) is then

$$\dot{\rho}(t) = \frac{1}{i\hbar} [\tilde{H}_S, \rho(t)] - (A_t + A_p) \mathcal{D}[Q_z]\rho(t)/2$$

$$-B\mathcal{D}[a]\rho(t) + \mathcal{B}[\rho(t)] + \mathcal{F}[\rho(t)],$$
(11)

where the new system Hamiltonian  $\tilde{H}_S = \hbar \omega_p v^{\dagger} v + \hbar \omega_z b_z^{\dagger} b_z - i \alpha \hbar g_z (v^{\dagger} - v) (b_z + b_z^{\dagger})$ , accounts for the linearization of the probe implemented above, and the master equation now includes a mechanical decoherence term due to scattering from the probe, in addition to the trap beam, with heating rate  $A_p$ .

A sample experimental data set of the measured position of the nanoparticle along the y axis is shown in Fig. 2. At atmospheric pressures, Brownian effects, as given by Eq. (6) dominate, and the position of the particle follows a diffusive evolution, as can be seen in Fig. 2a. At lower pressures the particle's evolution becomes increasingly ballistic. The ensuing harmonic motion is shown in Fig. 2b, both in the absence as well as in presence of feedback [Eq. (10)]. The decrease in amplitude of the harmonic motion is due to the presence of parametric feedback cooling.

### PHONON DYNAMICS

Employing the master equation [Eq. (11)] to consider the question of ground state occupation, tracing out the optical probe field, and using the resulting reduced master equation for the nanoparticle only, we find the equation for the dynamics of the phonon number  $(N \equiv b_z^{\dagger}b_z)$ ,  $\langle \dot{N} \rangle = -J \langle N^2 \rangle - K \langle N \rangle + L$ , where  $J = [12G - 54G^2] \chi^2 \Phi$ ,  $K = \eta_f/2m + J$ , L = D - J/2, the dot denotes a time derivative, and  $D = D'_p + D_q$  with  $D'_p = D_p + A_t + A_p$ accounting for positional decoherence. We assume that the nanoparticle is described by a thermal state [24, 31– 34], for which  $\langle N^2 \rangle = 2 \langle N \rangle^2 + \langle N \rangle$  [35], a relation which simplifies the phonon dynamics to

$$\langle \dot{N} \rangle = -2J \langle N \rangle^2 - (J+K) \langle N \rangle + L.$$
 (12)

The total effect of parametric feedback on the phonon dynamics is contained in the parameter J, which is determined by the difference between the feedback cooling and backaction heating. In the experiments  $J \neq 0$  [1, 11], making the phonon dynamics of cooling nonlinear, and the oscillator energy loss nonexponential, as shown below. We stress that this behavior is *qualitatively* different from standard quantum cavity optomechanical theory, which characterizes cooling as a linear damping process resulting in an exponential decay of energy (see, e.g. [31, 32] and Eq. (82) in [18]). We note that  $G = G_{\text{opt}} = 1/9$  maximizes  $\langle N \rangle$  in Eq. (12) with the maximum nonlinear cooling rate  $J_{\text{max}} = 2\chi^2 \Phi/3$ .

Assuming the initial condition  $\langle N(0) \rangle \equiv N_0 = D'_p 2m/\eta_f \equiv k_B T_{\text{eff}}/\hbar\omega_z$ , where  $T_{\text{eff}}$  is the effective temperature of a bath due to gas and optical scattering combined, the analytical solution to Eq. (12) is

$$\langle N(t) \rangle = -\frac{(J+K)}{4J} + \frac{1}{2J\tau} \tanh\left(\frac{t}{\tau} + \theta\right),$$
 (13)

where  $\theta = \tanh^{-1} \left[ (2JN_0 + J + K)\tau \right]$  and the cooling timescale  $\tau = 2 \left[ (J + K)^2 + 8JL \right]^{-1/2}$ . From Eq. (13)

the steady state phonon number is

$$N_{\rm ss} \equiv \lim_{t \to \infty} \langle N(t) \rangle = \frac{1}{2J\tau} - \frac{(J+K)}{4J} \tag{14a}$$

$$\approx \sqrt{\frac{\eta_f}{2m}} \frac{N_0}{2J} = \sqrt{\frac{D_p + A_t + A_p}{2J}},$$
 (14b)

where the approximation is valid for  $N_0 \gg 1$ . To reach the ground state, we need to maximize the feedback cooling J, which can be done by setting  $G = G_{\text{opt}}$ . We also need to minimize gas heating, which can be accomplished by going to low pressures and cryogenic temperatures, such that  $D_p$  is negligible in Eq. (14). Below we discuss situations involving realistic experimental parameters.

Two plots of the nonlinear phonon dynamics are shown in Fig. 3a for the z and y motion at  $10^{-3}$  mbar along with experimental values measured by us (circles). The solid curve represents  $\langle N(t) \rangle$  as given in Eq. (13), while the dotted curve gives the corresponding equation for motion along one of the transverse directions y (which is nearly degenerate with x i.e.  $\omega_y \approx \omega_x$ ), see Fig. 1b. In Fig. 3b we show three plots of the steady state phonon number, as the vacuum pressure is tuned. The solid and dotted curves represent  $N_{\rm ss}$  [Eq. (14a)] at 300 K for z and y motion respectively, while the circles are experimental data. As can be seen, in all cases there is very good agreement between theory and experiment. The dashed curve in Fig. 3b predicts the steady state phonon number for an identical configuration, but placed in a cryostat at 4K. The ground state can be prepared if starting at high pressures, the particle is cooled while continuously increasing the feedback gain as in [11], and keeping the trap modulation M = 10%. Proceeding in this manner, we find that below  $\lesssim 10^{-5}\,{\rm mbar}$  optimal feedback J =  $J_{\rm max}$  can be achieved, and the ground state occupied.

We note that practical cooling to lower phonon numbers is currently limited by a number of factors. These include high pressures enforced by nanosphere loading technologies, classical errors from the electronic feedback loop and laser noise, measurement uncertainties due to detector bandwidth limitations, and collection inefficiencies of the scattered light [3, 5, 11, 22]. However, these problems are technical rather than fundamental, and efforts are underway to overcome these limitations [36], suggesting strongly the possibility of cooling to the ground state using the parameters presented in this article.

### FORCE SENSING

We now consider force sensing using the nanoparticle model given by the master equation of Eq. (11). Since the state of the nanoparticle is continuously monitored,



FIG. 4: Experimentally measured positional PSDs for all three degrees of freedom with dark lines representing the theoretical fits to the data [Eq. (18)]. Data was taken at a moderate vacuum pressure of 10 mbar and clearly shows the Lorentzian shape of the resonance. These fits were used to extract the values of  $\omega_j$ ,  $\Gamma$ , and  $N_{\rm ss}$ .

the master equation can be unraveled in terms of a set of Langevin equations describing the evolution of the quadratures  $Q_z$  and  $P_z$  plus a stochastic force due to the measurement backaction which gives [37, 38]

$$\dot{Q}_{z} = \mathcal{L}_{0}^{\#}[Q_{z}] = \omega_{z}P_{z}$$
  
$$\dot{P}_{z} = \mathcal{L}_{0}^{\#}[P_{z}] + F/m\omega_{z}\ell_{z}$$
  
$$= -\omega_{z}Q_{z} - \Gamma P_{z} + F/m\omega_{z}\ell_{z},$$
  
(15)

where  $\mathcal{L}_0^{\#}$  is the Liouvillian superoperator (dual to the superoperator  $\mathcal{L}_0$  appearing in the master equation  $\dot{\rho} = \mathcal{L}_0[\rho]$ ) defined by  $\operatorname{Tr}(\rho \mathcal{L}_0^{\#}[A]) \equiv \operatorname{Tr}(A\mathcal{L}_0[\rho])$  for any arbitrary operator A [39]. The parameter  $\Gamma = \Gamma_0 + \delta \Gamma$ , where  $\Gamma_0$  is the gas damping and  $\delta \Gamma \approx 12\chi^2 \Phi G(\langle N \rangle + 1/2)$  is the nonlinear feedback damping [2, 5, 11]. Finally F = $F_T + F_F$  is the sum of the (independent) stochastic forces due to thermal and feedback backaction heating, respectively, with zero mean and correlations  $\langle F_T(t)F_T(t') \rangle =$  $S_T \delta(t - t')$  and  $\langle F_F(t)F_F(t') \rangle = S_F \delta(t - t')$ , with

$$S_T = 2m\Gamma_0 k_B T_{\text{eff}}$$

$$S_F = 27m\hbar\omega_z \chi^2 \Phi G^2 \left( 2\langle N \rangle^2 + 2\langle N \rangle + 1 \right).$$
(16)

The presence of the  $\langle N \rangle$ -dependent factor in Eq. (16) implies that the feedback noise is dependent on the system state, and is therefore non-additive. Furthermore, the dependence is nonlinear in  $\langle N \rangle$ . Both of these features are fundamentally different from the typical additive feedback noise in standard cavity optomechanics, which is independent of the state of the system [33].

We convert Eq. (15) into the second order differential equation for the position  $\ddot{q}_z + \Gamma \dot{q}_z + \omega_z^2 q_z = F/m$ , and take its Fourier transform to find the position spectrum  $\tilde{q}_z(\omega) = \chi_m(\omega)\tilde{F}(\omega)$ , where

$$\chi_m(\omega) = \left\{ m \left[ \left( \omega_z^2 - \omega^2 \right) - i\omega\Gamma \right] \right\}^{-1}, \qquad (17)$$

is the optomechanical susceptibility of our oscillator. Finally the positional power spectral density (PSD) noise spectrum is given by

$$\left\langle \left| \tilde{q}_{z}(\omega) \right|^{2} \right\rangle = \left| \chi_{m} \right|^{2} \left( S_{T} + S_{F} \right) + \frac{\ell_{z}^{2}}{\chi^{2} \Phi}, \qquad (18)$$

where the last term in the equation comes from the shot noise of the measured signal [Eq. (8)]. A typical example data set of the positional PSDs at moderate vacuum is shown in Fig. 4, along with fits to the theoretical expression of Eq. (18).

In view of the fact that trapped nanoparticles offer the possibility of ultrasensitive force measurements [5, 21–23], we express our measurement noise spectrum [Eq. (18)] in terms of the estimator  $\tilde{q}(\omega)/\chi_m$  in order to investigate the fundamental limits of such measurements. The sensitivity of force estimation is set by the force noise PSD

$$\left\langle \left| \tilde{F}(\omega) \right|^2 \right\rangle = S_T + S_F + S_S(\omega),$$
 (19)

where  $S_S(\omega) = S_S(0) \left[ \left( 1 - (\omega/\omega_z)^2 \right)^2 + (\omega\Gamma/\omega_z^2)^2 \right]$  and  $S_S(0) = (m\ell_z\omega_z^2)^2/\chi^2 \Phi$ . Only the last term carries an  $\omega$  dependence in Eq. (19). A plot of  $S_S(\omega)$  is shown in Fig 5 in the high as well as low total damping  $\Gamma$  regimes, both of which are experimentally accessible [11, 12]. The minimum value of  $S_S(\omega)$ , and therefore the optimal force sensitivity, occurs at the response frequency  $\omega_{\text{opt}} = \sqrt{\omega_z^2 - \Gamma^2/2}$ .

The first two terms in Eq. (19) scale linearly with the optical power while the shot noise scales inversely (i.e.  $S_T + S_F \sim \Phi$  and  $S_S \sim 1/\Phi$ ). Therefore there is a power that minimizes the total noise, representing the standard quantum limit for our system. Assuming that the feedback is optimal (i.e.  $J = J_{\text{max}}$ ), the standard quantum limit is reached when  $\chi^2 \Phi_{\text{SQL}} \approx \Gamma/8$  and equals

$$\left\langle \left| \tilde{F} \right|^2 \right\rangle_{\text{SQL}} = 2m\Gamma_0 k_B T + 4m\hbar\omega_z \left( A_t + \sqrt{2}\Gamma \right).$$
 (20)

The first term in (20) represents a thermal contribution from the background gas; the second term is due



FIG. 5: a) Plots of the shot noise force PSD [the last term of Eq. (19)] versus the normalized mechanical frequency  $\omega/\omega_z$  for low and high total damping  $\Gamma$ . The minimum occurs for  $\omega_{\text{opt}} = \sqrt{\omega_z^2 - \Gamma^2/2}$ . b) Plot of the force sensitivity as a function of the normalized optical power  $\Phi/\Phi_{\text{SQL}}$  at high vacuum. The standard quantum limit is reached when the shot noise balances the recoil and backaction noises [Eq. (19)].

to scattering of photons from the trapping beam; and the third term contains the effects of light scattering and shot noise from the probe, as well as the feedback backaction. At the low vacuum pressures currently available (i.e.  $\leq 10^{-7}$  mbar) the gas contribution is negligible, implying a minimum force sensitivity of  $\sqrt{\left<\left|\tilde{F}\right|^2\right>_{\rm SQL}}\approx$  $10^{-21} \,\mathrm{N}/\sqrt{\mathrm{Hz}} = \mathrm{zN}/\sqrt{\mathrm{Hz}}$  and optimal probe power of  $\hbar\omega_p \Phi_{SQL} \approx 2 \,\mathrm{mW}$ , where the remaining system parameters have been taken from the caption of Fig. 3. Even at this limit the system can be readily used to test for violations of Newtonian gravity (~  $10^{-18}$  N) [22] with moderate measurement bandwidths. However, backaction effects will impose long interrogation times on experiments searching for new small scale ( $\sim 10^{-21}$  N) [21], and Casimir forces (~  $10^{-24}$  N) [5, 21]. Conversely for short measurement times, our calculations show that backaction effects, which are of interest in their own right in optomechanics [40], can be observed at moderate laser powers and readily attainable vacuum pressures.

### CONCLUSIONS

To conclude, we have presented a quantum model that describes the cooling and force sensing characteristics of an optically trapped subwavelength dielectric particle. We have shown that the predictions of this model for cooling are in very good agreement with experimentally measured occupation values in the classical regime. Further, we have demonstrated that quantum ground state preparation is challenging, but achievable in anticipated experiments. Finally, we have derived the standard quantum limit to force sensing, indicating experiments where the role of quantum backaction needs to be accounted for. The model presented by us opens the door to the characterization of the quantum behavior of a system important for macroscopic quantum mechanics [1, 11], optical tweezing [2], ultrasensitive metrology [3], and nonequilibrium physics [4]. With the proper identifications, our theory is also applicable to electromechanical systems with parametric feedback [41].

# ACKNOWLEDGEMENTS

We are grateful to C. Stroud, A. Aiello, B. Zwickl, and S. Agarwal for useful discussions. This material is based upon work supported by the Office of Naval Research under Award Nos. N00014-14-1-0803 and N00014-14-1-0442. ANV thanks the Institute of Optics for support. LPN acknowledges support from a University of Rochester Messersmith fellowship.

\* Electronic address: brandon.rodenburg@gmail.com

- Neukirch, L. P., Gieseler, J., Quidant, R., Novotny, L. & Nick Vamivakas, A. Observation of nitrogen vacancy photoluminescence from an optically levitated nanodiamond. *Opt. Lett.* 38, 2976–9 (2013).
- [2] Gieseler, J., Spasenović, M., Novotny, L. & Quidant, R. Nonlinear Mode Coupling and Synchronization of a Vacuum-Trapped Nanoparticle. *Phys. Rev. Lett.* **112**, 103603 (2014).
- [3] Millen, J., Deesuwan, T., Barker, P. & Anders, J. Nanoscale temperature measurements using non-equilibrium Brownian dynamics of a levitated nanosphere. *Nat. Nanotechnol.* 9, 425–9 (2014).
- [4] Gieseler, J., Novotny, L., Moritz, C. & Dellago, C. Nonequilibrium steady state of a driven levitated particle with feedback cooling. New J. Phys. 17, 045011 (2015).
- [5] Neukirch, L. P. & Vamivakas, A. N. Nano-optomechanics with optically levitated nanoparticles. *Contemporary Physics* 56, 48–62 (2015).

- [6] Li, T., Kheifets, S. & Raizen, M. G. Millikelvin cooling of an optically trapped microsphere in vacuum. *Nature Phys.* 7, 527–530 (2011).
- [7] Yin, Z. Q., Geraci, A. A. & Li, T. C. Optomechanics of levitated dielectric particles. *Int. J. Mod. Phys. B* 27, 1330018 (2013).
- [8] Bateman, J., Nimmrichter, S., Hornberger, K. & Ulbricht, H. Near-field interferometry of a free-falling nanoparticle from a point-like source. *Nat. Commun.* 5 (2013).
- [9] Arita, Y., Mazilu, M. & Dholakia, K. Laser-induced rotation and cooling of a trapped microgyroscope in vacuum. *Nat. Commun.* 4, 2374 (2013).
- [10] Scala, M., Kim, M. S., Morley, G. W., Barker, P. F. & Bose, S. Matter-wave interferometry of a levitated thermal nano-oscillator induced and probed by a spin. *Phys. Rev. Lett.* **111**, 180403 (2013).
- [11] Gieseler, J., Deutsch, B., Quidant, R. & Novotny, L. Sub-Kelvin Parametric Feedback Cooling of a Laser-Trapped Nanoparticle. *Phys. Rev. Lett.* **109**, 103603 (2012).
- [12] Neukirch, L. P., von Haartman, E., Rosenholm, J. M. & Nick Vamivakas, A. Multi-dimensional single-spin nanooptomechanics with a levitated nanodiamond. *Nat. Photonics* (2015).
- [13] Millen, J., Fonseca, P. Z. G., Mavrogordatos, T., Monteiro, T. S. & Barker, P. F. Cavity Cooling a Single Charged Levitated Nanosphere. *Phys. Rev. Lett.* **114**, 123602 (2015).
- [14] Kiesel, N. et al. Cavity cooling of an optically levitated submicron particle. PNAS 110, 14180–5 (2013).
- [15] Kippenberg, T. J. & Vahala, K. J. Cavity optomechanics. Opt. Exp. 15, 17172–17205 (2007).
- [16] Marquardt, F. & Girvin, S. M. Optomechanics (a brief review). *Physics* 2, 40 (2009).
- [17] Romero-Isart, O. et al. Large Quantum Superpositions and Interference of Massive Nanometer-Sized Objects. *Phys. Rev. Lett.* **107**, 020405 (2011).
- [18] Aspelmeyer, M., Kippenberg, T. J. & Marquardt, F. Cavity optomechanics. *Rev. Mod. Phys.* 86, 1391–1452 (2014).
- [19] Meystre, P. A short walk through quantum optomechanics. Annalen der Physik 525, 215–233 (2013).
- [20] Asenbaum, P., Kuhn, S., Nimmrichter, S., Sezer, U. & Arndt, M. Cavity cooling of free silicon nanoparticles in high vacuum. *Nat. Nanotechnol.* 4, 2743 (2013).
- [21] Geraci, A. A., Papp, S. B. & Kitching, J. Short-Range Force Detection Using Optically Cooled Levitated Microspheres. *Phys. Rev. Lett.* **105**, 101101 (2010).
- [22] Ranjit, G., Atherton, D. P., Stutz, J. H., Cunningham, M. & Geraci, A. A. Attonewton force detection using microspheres in a dual-beam optical trap in high vacuum. *Phys. Rev. A* **91**, 051805 (2015).
- [23] Moore, D. C., Rider, A. D. & Gratta, G. Search for Millicharged Particles Using Optically Levitated Microspheres. *Phys. Rev. Lett.* **113**, 251801 (2014).
- [24] Pflanzer, A. C., Romero-Isart, O. & Cirac, J. I. Masterequation approach to optomechanics with arbitrary dielectrics. *Phys. Rev. A* 86, 013802 (2012).
- [25] Carmichael, H. J. Statistical Methods in Quantum Op-

tics 1: Master Equations and Fokker-Planck Equations (Springer, 2002).

- [26] Diósi, L. Quantum Master Equation of a Particle in a Gas Environment. EuroPhys. Lett. (EPL) 30, 63–68 (1995).
- [27] Chang, D. E. et al. Cavity opto-mechanics using an optically levitated nanosphere. PNAS 107, 1005–10 (2010).
- [28] Gardiner, C. W. & Zoller, P. Quantum Noise: A Handbook of Markovian and Non-Markovian Quantum Stochastic Methods with Applications to Quantum Optics (Springer, 2004), 3rd edn.
- [29] Mancini, S., Vitali, D. & Tombesi, P. Optomechanical Cooling of a Macroscopic Oscillator by Homodyne Feedback. *Phys. Rev. Lett.* **80**, 688–691 (1998).
- [30] Wiseman, H. M. & Milburn, G. J. Quantum theory of optical feedback via homodyne detection. *Phys. Rev. Lett.* **70**, 548 (1993).
- [31] Wilson-Rae, I., Nooshi, N., Zwerger, W. & Kippenberg, T. J. Theory of Ground State Cooling of a Mechanical Oscillator Using Dynamical Backaction. *Phys. Rev. Lett.* **99**, 093901 (2007).
- [32] Marquardt, F., Chen, J. P., Clerk, A. A. & Girvin, S. M. Quantum Theory of Cavity-Assisted Sideband Cooling of Mechanical Motion. *Physical Review Letters* **99**, 093902 (2007).
- [33] Genes, C., Vitali, D., Tombesi, P., Gigan, S. & Aspelmeyer, M. Ground-state cooling of a micromechanical

oscillator: Comparing cold damping and cavity-assisted cooling schemes. *Phys. Rev. A* **77**, 033804 (2008).

- [34] Romero-Isart, O. et al. Optically levitating dielectrics in the quantum regime: Theory and protocols. Phys. Rev. A 83, 013803 (2011).
- [35] Gerry, C. & Knight, P. Introductory Quantum Optics (Cambridge University Press, Cambridge, 2004).
- [36] Mestres, P. *et al.* Cooling and manipulation of a levitated nanoparticle with an optical fiber trap. *Appl. Phys. Lett.* **107**, 151102 (2015).
- [37] Gisin, N. & Percival, I. C. The quantum-state diffusion model applied to open systems. J. Phys. A. Math. Gen. 25, 5677–5691 (1992).
- [38] Halliwell, J. & Zoupas, A. Quantum state diffusion, density matrix diagonalization, and decoherent histories: A model. *Phys. Rev. D* 52, 7294 (1995).
- [39] Hornberger, K. Entanglement and Decoherence, vol. 768 of Lecture Notes in Physics (Springer Berlin Heidelberg, Berlin, Heidelberg, 2009).
- [40] Purdy, T. P., Peterson, R. W. & Regal, C. A. Observation of radiation pressure shot noise on a macroscopic object. *Science* **339**, 801–4 (2013).
- [41] Villanueva, L. G. et al. A Nanoscale Parametric Feedback Oscillator. Nano Lett. 11, 5054–5059 (2011).

# Supplemental Materials: Quantum Model of Cooling and Force Sensing With an Optically Trapped Nanoparticle

### ELECTRIC FIELDS

The total electric field as defined in the main article is written as

$$\mathbf{E}(\mathbf{r},t) = \mathbf{E}_{\mathbf{t}}(\mathbf{r},t) + \mathbf{E}_{\mathbf{p}}(\mathbf{r},t) + \mathbf{E}_{\mathbf{b}}(\mathbf{r},t).$$
(S1)

The  $\mathbf{E}_{\mathbf{t}}$  and  $\mathbf{E}_{\mathbf{b}}$  fields are both assumed to be Gaussian beams with the trap treated classically and the probe treated as a quantized beam with frequency  $\omega_p$ , linewidth  $\Delta \omega$ , waist  $w_0$ , and canonical bosonic operators  $[a, a^{\dagger}] = 1$  [S1]

$$\mathbf{E}_{\mathbf{p}}(\mathbf{r},t) = i \left(\frac{\hbar\omega_p \Delta\omega}{4\pi\epsilon_0 c}\right)^{1/2} e^{i\omega_p(z/c-t)} \mathbf{e}_{\mathbf{p}} G(\mathbf{r}',\omega_p) a + \text{h.c.},\tag{S2}$$

where  $G(\mathbf{r}, \omega)$  is the spatial mode function of the beam. The background field  $\mathbf{E}_{\mathbf{b}}$  is simply the quantum field of all other modes and can be represented via a typical plane-wave expansion [S2].

### FREE FIELD HAMILTONIAN

Our configuration Hamiltonian as given in the main text is

$$H = H_m + H_f + H_{\text{int}},\tag{S3}$$

where  $H_m = p^2/2m$  is the particle's kinetic energy for the momentum  $p = \sqrt{p_x^2 + p_y^2 + p_z^2}$ , and  $H_f$  and  $H_{int}$  are the free field and interaction Hamiltonians respectively. The energy of the free field Hamiltonian is

$$H_f = \epsilon_0 \int \left| \mathbf{E}(\mathbf{r}, \mathbf{t}) \right|^2 \mathrm{d}^3 \mathbf{r}.$$
 (S4)

The term proportional to  $|\mathbf{E}_{t}(\mathbf{r})|^{2}$  corresponds to the energy of the trap beam, and can be neglected as it represents a constant offset of the Hamiltonian. The energy of the free probe field is given by [S1]

$$\epsilon_0 \int \left| \mathbf{E}_{\mathbf{p}}(\mathbf{r}) \right|^2 \mathrm{d}^3 \mathbf{r} = \hbar \omega_p a^{\dagger} a.$$
(S5)

The energy of the background field, which can be found in several textbooks (e.g. [S2]) can be written in a plane wave expansion as

$$H_B = \sum_{\mu} \int d^3 \mathbf{k} \hbar \omega_{\mathbf{k}} a^{\dagger}_{\mu}(\mathbf{k}) a_{\mu}(\mathbf{k}).$$
(S6)

The cross term between the trap and background fields, given by  $\int (\mathbf{E}_t + \mathbf{E}_p) \cdot \mathbf{E}_b d^3 \mathbf{r}$ , vanishes as their mutual overlap is very small. This cancellation also represents the avoidance of self-interference and mode overcounting in our model. The cross term between the probe and the background fields vanishes for the same reason. The cross term between the trap and probe fields given by  $\mathbf{E}_t \cdot \mathbf{E}_p$ , vanishes due to polarization orthogonality. Finally, combining Eqs. (S5) and (S6), the free field Hamiltonian is

$$H_f = \hbar \omega_p a^{\dagger} a + \sum_{\mu} \int \mathrm{d}^3 \mathbf{k} \hbar \omega_{\mathbf{k}} a^{\dagger}_{\mu}(\mathbf{k}) a_{\mu}(\mathbf{k}), \tag{S7}$$

which is simply the sum of the probe and background field energies.

### INTERACTION HAMILTONIAN

The interaction Hamiltonian between the fields and the nanoparticle is given by

$$H_{\rm int} = -\frac{1}{2} \int \mathbf{P}(\mathbf{r}) \cdot \mathbf{E}(\mathbf{r}) \,\mathrm{d}^3 \mathbf{r}.$$
 (S8)

Assuming that the particle has a linear polarizability described by  $\mathbf{P}(\mathbf{r}) = \alpha_p \mathbf{E}(\mathbf{r})$ , the interaction Hamiltonian is

$$H_{\rm int} = -\frac{\epsilon_0 \epsilon_c}{2} \int_V \left| \mathbf{E}(\mathbf{r}) \right|^2 \mathrm{d}^3 \mathbf{r},\tag{S9}$$

where  $\epsilon_c = 3(\epsilon_r - 1)/(\epsilon_r + 2)$  is the Clausius-Mossotti relation for the effective relative permittivity of a dielectric due to local field effects, and V denotes integration over the volume of the dielectric particle.

Now when we use the total electric field from Eq. (S1) in the above equation, we again get a number of terms. The terms proportional to  $|\mathbf{E}_{\mathbf{b}}|^2$  and  $|\mathbf{E}_{\mathbf{t}} \cdot \mathbf{E}_{\mathbf{p}}|^2$  we neglect as these represent a renormalization of the background modes and we again assume the trap and probe are cross-polarized.

### Trap potential

The first non-negligible term is the effect of the trap beam proportional to  $\int_{V} |\mathbf{E}_{\mathbf{p}}|^{2} d^{3}\mathbf{r}$ . Since the nanoparticle is smaller than the wavelength of any relevant optical field, the integration can be written as  $\int_{V} d^{3}\mathbf{r} = V \int \delta(\mathbf{q}) d^{3}\mathbf{r}$ , where  $\mathbf{q}$  is the center of mass position of the particle. Therefore  $H_{\text{int}}$  for the trap-particle interaction is given by

$$-\frac{\epsilon_c \epsilon_0}{2} V \left| \mathbf{E}_{\mathbf{t}}(\mathbf{q}) \right|^2 = -\frac{\epsilon_c \epsilon_0}{2} V E_0^2 w_0^2 \frac{\pi}{2} \left| G(\mathbf{q}) \right|^2$$
$$= \frac{\epsilon_c \epsilon_0}{2} V E_0^2 \left( -1 + \left(\frac{q_z}{z_R}\right)^2 + 2 \left(\frac{q_x}{w_0}\right)^2 + 2 \left(\frac{q_y}{w_0}\right)^2 \right) + \mathcal{O}(q^4),$$
(S10)

where  $q_z$  is the longitudinal coordinate, and  $q_x$  and  $q_y$  are the transverse coordinates. Ignoring the overall constant in the above equation, we finally have for our trap Hamiltonian

$$H_{\rm trap} = \frac{p^2}{2m} + \frac{1}{2}m\left(\omega_z^2 q_z^2 + \omega_x^2 q_x^2 + \omega_y^2 q_y^2\right),\tag{S11}$$

where  $\omega_z = \sqrt{\epsilon_c \epsilon_0 E_0^2 V/(m z_R^2)}$  and  $\omega_{x,y} = \sqrt{2\epsilon_c \epsilon_0 E_0^2 V/(m w_0^2)}$ . Writing our canonical nanoparticle variables as  $q_j = \sqrt{\hbar/(2m\omega_j)}(b_j^{\dagger} + b_j) = \ell_j (b_j^{\dagger} + b_j) = \ell_j Q_j$  and  $p_j = i\sqrt{\hbar m \omega_j/2}(b^{\dagger} - b)$  allows us to rewrite Eq. (S11) as  $H_{\text{trap}} = \sum_j \hbar \omega_j b_j^{\dagger} b_j$ .

### **Optomechanical coupling**

Now we examine the coupling between the probe beam and the trapped nanoparticle given. The Hamiltonian for this term is

$$H_{\rm OM} = -\frac{\epsilon_c \epsilon_0}{2} \int_V \left| \hat{\mathbf{E}}_{\mathbf{p}}(\mathbf{r}) \right|^2 \mathrm{d}^3 \mathbf{r}$$
  
$$= -V \frac{\epsilon_c \hbar \omega_p \Delta \omega}{4\pi c} \left( a^{\dagger} a + \frac{1}{2} \right) \left| G(\mathbf{q} - \Delta \mathbf{r}) \right|^2$$
  
$$= -V \frac{\epsilon_c \hbar \omega_p \Delta \omega}{2\pi^2 w_0^2 c} \left( a^{\dagger} a + \frac{1}{2} \right) \left( 1 + 2 \frac{\Delta z}{z_R^2} q_z + \frac{4}{w_0^2} (\Delta x q_x + \Delta y q_y) + \mathcal{O}(\mathbf{q}^2) \right),$$
(S12)

where we have assumed that the probe beam is shifted from the trap beam by a small amount  $\Delta \mathbf{r} \equiv (\Delta x, \Delta y, \Delta z)$ . Now the terms proportional to a constant times either  $a^{\dagger}a$  or  $q_j$  (for  $j \in \{x, y, z\}$ ) can be incorporated into shifts in the optical and mechanical frequencies, and the oscillator position. The optomechanical coupling term between the probe and particle is then given by

$$H_{\rm OM} = -V \frac{\epsilon_c \hbar \omega_p \Delta \omega}{\pi c} a^{\dagger} a \left( \frac{\Delta z}{z_R^2} \ell_z Q_z + 2 \frac{\Delta x}{z_R^2} \ell_x Q_x + 2 \frac{\Delta y}{z_R^2} \ell_y Q_y \right) = -\sum_j \hbar g_j a^{\dagger} a Q_j.$$
(S13)

The system Hamiltonian is then a combination of the energies of the probe field, optical trap, and optomechanical coupling and is given by

$$H_S = \hbar \omega_p a^{\dagger} a + \sum_j \hbar \omega_j b_j^{\dagger} b_j - \sum_j \hbar g_j a^{\dagger} a (b_j + b_j^{\dagger}).$$
(S14)

# **Optical scattering**

Computing the cross coupling Hamiltonian between the field and the background due to scattering by the nanoparticle gives

$$-\epsilon_{c}\epsilon_{0}\int_{V}\mathbf{E}_{t}\cdot\mathbf{E}_{b}\,\mathrm{d}^{3}\mathbf{r}\approx-i\frac{\epsilon_{c}V}{2}\left(\frac{\hbar\epsilon_{0}}{16\pi^{3}}\right)^{1/2}\int\mathrm{d}^{3}\mathbf{k}\sqrt{\omega_{k}}\sum_{\mu}\left[1+i\left(\mathbf{k}_{\perp}\cdot\mathbf{q}_{\perp}+(k_{z}-k_{0})q_{z}\right)\right]$$

$$\times\left(\mathbf{e}_{\mu}(\mathbf{k})\cdot\mathbf{E}_{0}^{*}a_{\mu}(\mathbf{k})e^{-i(\omega_{k}-\omega_{t})t}+\mathrm{h.c.}\right),$$
(S15)

for the trap beam and

$$-\epsilon_{c}\epsilon_{0}\int_{V}\mathbf{E}_{\mathbf{p}}\cdot\mathbf{E}_{\mathbf{b}}\,\mathrm{d}^{3}\mathbf{r}\approx-i\frac{\hbar\epsilon_{c}V}{8\pi^{2}w_{0}}\sqrt{\frac{2\omega_{p}\Delta\omega}{\pi c}}\int\mathrm{d}^{3}\mathbf{k}\sqrt{\omega_{k}}\sum_{\mu}\left[1+i\left(\mathbf{k}_{\perp}\cdot\mathbf{q}_{\perp}+(k_{z}-k_{0})q_{z}\right)\right]\times\left(\mathbf{e}_{p}\cdot\mathbf{e}_{\mu}(\mathbf{k})a_{\mu}^{\dagger}(\mathbf{k})a_{p}e^{-i(\omega_{p}-\omega_{k})t)}+\mathrm{h.c.}\right)$$
(S16)

for the probe beam, assuming elastic scattering and amplitude of the particle motion small relative to an optical wavelength [S3]. Applying standard Born-Markov theory and tracing over the background optical modes and particle motion transverse to the z-axis allows us to derive a master equation

$$\dot{\rho}(t) = -\frac{7\omega_t^5}{\hbar c^6} \frac{(E_0^2 \epsilon_0 c/2)\epsilon_c^2 V^2 \ell_z^2}{60\pi} \mathcal{D}[Q_z]\rho(t) + \frac{1}{\hbar^4 c^3} \frac{(\hbar \omega_p)^4 \epsilon_c^2 V^2}{24\pi^3 w_0^2 (c/\Delta \omega)} \left( \mathcal{D}[a_p] + \frac{7\omega_p^2 \ell_z^2}{5c^2} \mathcal{D}[aq_z] \right) \rho(t)$$

$$= -\frac{A_t}{2} \mathcal{D}[Q_z] + \mathcal{L}_{\rm sc}[\rho(t)]$$
(S17)

For reference, if we do not trace over the transverse motion, the three-dimensional analogue of the trap scattering is

$$\mathcal{L}_t^{3D}\left[\rho(t)\right] \equiv -\frac{\omega_t^5}{\hbar c^6} \frac{\left(E_0^2 \epsilon_0 c/2\right) \epsilon_c^2 V^2}{60\pi} \left(2\mathcal{D}[q_{\times}] + \mathcal{D}[q_{\parallel}] + 7\mathcal{D}[q_z]\right) \rho(t),\tag{S18}$$

where  $\parallel$  and  $\times$  indicate transverse motion parallel (or not parallel) to the polarization direction of the trap beam. A similar expression holds for the probe

$$\mathcal{L}_{sc}^{3D}\left[\rho(t)\right] \equiv \frac{1}{\hbar^4 c^3} \frac{(\hbar\omega_p)^4 \epsilon_c^2 V^2}{24\pi^3 w_0^2(c/\Delta\omega)} \left\{ \mathcal{D}[a_p] + \frac{\omega_p^2 \ell_z^2}{5c^2} \left( \mathcal{D}[aq_{\times}] + 2\mathcal{D}[aq_{\parallel}] + 7\mathcal{D}[aq_z] \right) \right\} \rho(t).$$
(S19)

### DERIVATION OF THE DETECTED HOMODYNE CURRENT AND FEEDBACK

In order to find the input-output relations for the probe field, we write the system Hamiltonian (for the single degree of freedom  $Q_z$ ) as  $H_s = \hbar \omega_p a^{\dagger} a + \hbar \omega_z b_z^{\dagger} b_z - \hbar g a^{\dagger} a Q_z$ . If we move into the interaction picture for the probe field where  $a \to a e^{-i\omega_p t}$ , then this becomes  $H_s = \hbar \omega_z b_z^{\dagger} b_z - \hbar g a^{\dagger} a Q_z$ .

We assume the probe field is initially a coherent state which can be written as  $a = -i\alpha + v$ , where  $\alpha$  is a constant and v is a field annihilation operator. In this case the optomechanical coupling in our system Hamiltonian becomes

$$H_{\rm OM} = -\hbar g_z Q_z a^{\dagger} a = \hbar g_z Q_z \left( \left| \alpha \right|^2 + i \alpha v^{\dagger} - i \alpha^* v \right).$$
(S20)

The term proportional to  $|\alpha|^2 Q_z$  is responsible for simply shifting the mean position of the oscillator and can safely be ignored, leaving us with  $H_{\rm OM} = i\hbar g_z Q_z (\alpha v^{\dagger} - \alpha^* v)$ .

The Heisenberg equation of motion for v is given by

$$\dot{v} = \frac{1}{i\hbar}[a, H] = \alpha g_z Q_z, \tag{S21}$$

which can be integrated formally to give

$$v(t) = v(t_0) + \int_{t_0}^t \alpha g_z Q_z \, \mathrm{d}t' \approx v(t_0) + \alpha g_z Q_z \Delta t, \tag{S22}$$

where the integration is taken over a time  $\Delta t$  short compared to  $1/\omega_z$ . If we had picked an initial time  $t_f > t$  then we would have computed  $v(t) = v(t_f) - \alpha g_z Q_z \Delta t$ . By identifying the input state as  $a_{in} = -i\alpha + v(t_0)$ , and the output state as  $a_{out} = -i\alpha + v(t_f)$ , then we can relate the output and input fields by

$$a_{\rm out} = a_{\rm in} + 2\alpha g_z \Delta t \equiv a_{\rm in} + \frac{\alpha \chi}{2} Q_z, \qquad (S23)$$

where we have defined the variable  $\chi \equiv 4g_z \Delta t$ .

Now detected homodyne current is proportional to

$$I_h = \chi^2 \Phi \langle Q_z \rangle (t) + \sqrt{\chi^2 \Phi} \xi(t), \qquad (S24)$$

where we have converted from photon numbers to rates by using the identity for the photon flux  $\Phi \equiv \langle a^{\dagger}a \rangle \Delta \omega = \alpha^2 \Delta \omega$ , and have introduced the stochastic variable  $\xi(t)$  which is due to the shot noise of the detection.

As described in the main text, phase shifting the measured signal (or equivalently adding a short time delay) is equivalent to measuring a different quadrature of motion, i.e. the current that is fed back is  $I_{\rm fb} = \chi^2 \Phi \langle P_z \rangle + \sqrt{\chi^2 \Phi} \xi'(t)$ , where  $\xi'(t)$  has the same properties as  $\xi(t)$ . Now writing  $\sigma \equiv \chi^2 \Phi P_z$  allows us to write the master equation for the feedback in standard notation [S4]

$$\dot{\rho} = \mathcal{K}[\sigma\rho + \rho\sigma^{\dagger}] + \frac{1}{2\chi^2 \Phi} \mathcal{K}^2[\rho] \equiv -i\chi^2 \Phi G\left[Q_z^3, \{P_z, \rho\}\right] - \frac{\chi^2 \Phi}{2} G^2 \mathcal{D}[Q_z^3]\rho \tag{S25}$$

where the Liouvillian superoperator  $\mathcal{K}$  is defined as  $\mathcal{K}[\rho] = [F, \rho]/i\hbar$  and where F is the feedback term that comes from the feedback Hamiltonian  $H_{\rm fb} = I_{\rm fb}F$ , and is chosen to be  $F = \hbar G Q_z^3$  to match the classical physics [S5] of the problem as described in the main text. The gain coefficient G may be related to the experimental trap beam intensity modulation  $M \equiv \Delta I_t/I_t$  by using the fact that  $H_{\rm fb} = \hbar \omega_z M Q_z^2 = I_{\rm fb}F$ , and therefore [S5]

$$M = \frac{G\chi^2 \Phi \langle Q_z \rangle \langle P_z \rangle}{\omega_z} \approx \frac{G\chi^2 \Phi N}{\omega_z}.$$
 (S26)

### FULL MASTER EQUATION

In the main text of this Letter we presented the master equations describing the density matrix of only the single degree of freedom represented by the z motion of the nanoparticle. For convenience, we present the full master equation of the three-dimensional motion of the nanoparticle, as well as of the probe beam itself, which can be used to calculate the statistics of any quantum observable for the nanoparticle or probe beam.

The master equation for the density matrix  $\rho$  describing the full three-dimensional motion of the nanoparticle, as well as the probe field is given by

$$\dot{\rho}(t) = \frac{1}{i\hbar} \left[ H_S, \rho(t) \right] \quad \left\{ \begin{array}{l} \text{Unitary dynamics} \\ + \mathcal{L}_t^{3D} \left[ \rho(t) \right] + \mathcal{L}_{\text{sc}}^{3D} \left[ \rho(t) \right] \end{array} \right\} \text{Photon scattering} \\ - \frac{\eta_f k_B T}{\hbar^2} \left[ \mathbf{q}, \left[ \mathbf{q}, \rho(t) \right] \right] - \frac{\eta_f}{12k_B T m^2} \left[ \mathbf{p}, \left[ \mathbf{p}, \rho(t) \right] \right] - \frac{\eta_f}{2m\hbar} \left[ \mathbf{q}, \left\{ \mathbf{p}, \rho(t) \right\} \right] \end{array} \right\} \text{Gas scattering.}$$
(S27)

This equation can be used for studying full dynamics, light-matter entanglement and photon statistics.

# HOMODYNE PHOTODETECTION: A COMPARISON OF THEORY AND EXPERIMENT

The experimental data in Fig. 2 of the main article was obtained by using a scheme more involved, and of more general applicability, than indicated in the simplified detection model presented in the main article. In the laboratory, the position of the nanoparticle was determined from the optical interference between the unscattered probe and a spherical wave radiated by the induced dipole of the polarizable sphere. The interference signal varies linearly with the particle position for oscillation amplitudes small compared to the optical wavelength. In order to detect the particle position and also to eliminate the large constant background term (equal to the unscattered probe flux  $\approx \Phi$ ) in an experimental setting, we leveraged a balanced homodyne detection scheme, as described in [S3] and [S5].

In this configuration, one of the detectors sampled the entire spatial profile of the probe beam, while the other was apodized so that only the center of the beam was sampled. As the trapped particle moved in the axial direction, the relative amount of scattered light collected by each detector, and thus the AC term of the resulting homodyne current, was modulated. The optical channels were adjusted so that each detector recorded the same average power, and the resulting homodyne signals were subtracted to eliminate the common DC bias.

In our theoretical model, rather than include an additional quantized mode corresponding to the full induced dipole emission pattern, we considered for simplicity only the field scattered back into the probe mode by the nanoparticle. This simplification is admissible as long as the value of the linear optomechanical coupling constant  $\chi$  in the model is taken from experiment [S3]. We emphasize that our experimental method of feedback cooling works even if the trap and probe foci are very close to each other, in which case the optomechanical coupling is essentially quadratic in the particle position, while the position detection (using the dipole wave mode) is still linear. In the theoretical calculation of the phonon number, the linear (or quadratic) optomechanical coupling simply adds a position offset (or frequency shift) to the trap. Both the offset as well as the shift are negligible for weak probe light used in the experiment, rendering the theoretical predictions of ground state cooling identical for linear as well as quadratic optomechanical coupling.

# COMPUTING THE GAS DAMPING RATE AS A FUNCTION OF GAS PRESSURE

The damping rate of the particle motion due to the background gas is given by  $\Gamma_0 \equiv \eta_f/m$ , where *m* is the particle mass and  $\eta_f = 6\pi\mu r_d$  is the coefficient of friction for a spherical particle of radius  $r_d$  in a fluid of dynamic viscosity  $\mu$ . This expression is for a particle much larger than the mean free path of the gas bath  $\lambda_{mfp}$ . For a nanoparticle in

a rarefied gas, as assumed in Eq. (9) in the main text, the damping constant becomes [S6]

$$\Gamma_0 = \frac{6\pi\mu r_d}{m} \times \text{correction} = \frac{6\pi\mu r_d}{m} \frac{0.619}{0.619 + \text{Kn}} \left( 1 + \frac{0.31\text{Kn}}{0.785 + 1.152\text{Kn} + \text{Kn}^2} \right),$$
(S28)

where the correction term is in terms of the Knudsen number  $\text{Kn} \equiv \lambda_{\text{mfp}}/r_d \propto 1/Pr_d$ . Using the fact that  $\lambda_{\text{mfp}} \approx 70 \text{ nm}$  at atmospheric pressure, and matching the experimentally measured rates [S3, S5], we can write the damping for a general pressure and particle size as

$$\Gamma_0 \approx \frac{r_d}{70\,\mathrm{nm}} \frac{2\pi \times 10^6\,\mathrm{Hz}}{0.619 + \mathrm{Kn}} \left( 1 + \frac{0.31\mathrm{Kn}}{0.785 + 1.152\mathrm{Kn} + \mathrm{Kn}^2} \right).$$
(S29)

In order to ensure that this model of damping holds in the deep quantum regime [S7], we consider the effects of individual scattering events as described by the master equation [S8]

$$\dot{\rho} = \sum_{V} V \rho V^{\dagger} - \frac{1}{2} \left\{ V^{\dagger} V, \rho \right\}, \tag{S30}$$

where the Lindblad operators are given by

$$V \sim \exp\left(-i\frac{\Delta pq}{\hbar}\right),$$
 (S31)

where  $\Delta p$  is the momentum transferred to the nanoparticle during a collision with a gas molecule. Now if the momentum kick is sufficiently small, then we can approximate

$$\exp\left(-i\Delta pq/\hbar\right) \approx 1 - i\Delta pq/\hbar,\tag{S32}$$

which ultimately allows us to recover the Brownian contribution, derived in [S8], to our master equation presented in the main Letter. In order for the approximation of Eq. (S32) to hold, it is necessary that  $\Delta p \ell_{\rm coh}/\hbar < 1$ , where  $\ell_{\rm coh}$ is the coherence length of the system. Considering the coherence length of the system in its ground state,  $\ell_{\rm coh} \approx \ell_z$ . Using the parameters listed in the main article for the system at T = 4 K, and using the maximum RMS thermal momentum change  $\Delta p = 2\sqrt{3m_{\rm gas}k_BT}$  assuming a gas made of N<sub>2</sub> gives

$$\frac{\Delta p \ell_z}{\hbar} \approx 0.6 < 1,\tag{S33}$$

which ensures that our linearized treatment of Brownian motion is valid over all considered regimes. This inequality actually overestimates the magnitude of a typical momentum kick as Hydrogen and Helium comprise the majority of the species of gas molecules in an environment at 4 K, and are much lighter than N<sub>2</sub>.

# CONVERTING A MASTER EQUATION TO A LANGEVIN EQUATION

Consider a master equation in standard Linblad form

$$\dot{\rho} = \mathcal{L}_0 \rho = \frac{1}{i\hbar} \left[ H, \rho \right] - \sum_j \frac{\gamma_j}{2} \mathcal{D}[L_j] \rho, \qquad (S34)$$

where  $H_0$  is a Hamiltonian representing the unitary evolution and  $\mathcal{D}[L_j]\rho$  is the standard Lindblad dissipation superoperator for the Lindblad operators  $L_j$  [S9]. Now following the quantum-state diffusion model presented by Gisin and Percival [S10], the density matrix can be considered as the expectation over pure states e.g.  $\rho \equiv E [|\psi\rangle \langle \psi|]$ , whose dynamics are stochastic in nature. The (stochastic) differential increment for the equation of motion for  $|\psi\rangle$ representing the unraveling of the master equation given by the QSD model is [S10, S11]

$$|\mathrm{d}\psi\rangle = \frac{1}{i\hbar}H\,\mathrm{d}t\,|\psi\rangle + \sum_{j} \left[\frac{\gamma_{j}}{2}\left(2\,\langle L_{j}^{\dagger}\rangle\,L_{j} - L_{j}^{\dagger}L_{j} - \langle L_{j}^{\dagger}\rangle\,\langle L_{j}\rangle\right)\mathrm{d}t + \left(L_{j} - \langle L_{j}\rangle\right)\sqrt{\gamma_{j}}\,\mathrm{d}W_{j}\right]|\psi\rangle \tag{S35}$$

where  $dW_j$  are complex differentials representing Wiener processes defined by  $E\left[dW_j\right] = E\left[dW_j^*\right] = E\left[dW_j dW_k\right] = 0$ , and  $dW_j^* dW_k = \delta_{jk} dt$ . Note, this complex Wiener process is equivalent to  $dW = (dR_1 + i dR_2)/\sqrt{2}$ , where  $dR_j$  are real valued Wiener increments. Now such a model is not unique, but represents an open quantum system which is continuously monitored by it's environment [S12], and thus provides an appropriate trajectory or unraveling for a system that we will ultimately wish to observe in terms of a PSD or measurement spectrum (i.e. which we will continuously monitor).

Now we are interested in expressing the random forces acting on our system due to interaction with the environment (as expressed by the Lindblad operators  $L_j$ ). Therefore computing the increment for the momentum p of the system gives

$$dp = \langle d\psi | p | \psi \rangle + \langle \psi | p | d\psi \rangle + \langle d\psi | p | d\psi \rangle + \mathcal{O}(dt^2) = -\frac{1}{i\hbar} [H_0, p] dt - \sum_j \frac{\gamma_j}{2} \mathcal{D} \left[ L_j^{\dagger} \right] p dt + \sum_j \left( dB_j^{\dagger} p + p dB_j \right) = \mathcal{L}_0^{\#}[p] dt + \sum_j \left( dB_j^{\dagger} p + p dB_j \right),$$
(S36)

where  $dB_j \equiv (L_j - \langle L_j \rangle) \sqrt{\gamma_j} dW_j$ . Applying this to our master equation gives us our Langevin equation for the stochastic forces listed in Eq. (15) in the main article.

\* Electronic address: brandon.rodenburg@gmail.com

- [S1] A. Aiello, C. Marquardt, and G. Leuchs, "Transverse angular momentum of photons," Phys. Rev. A 81, 053838 (2010).
- [S2] C. Gerry and P. Knight, Introductory Quantum Optics (Cambridge University Press, Cambridge, 2004).
- [S3] J. Gieseler, B. Deutsch, R. Quidant, and L. Novotny, "Sub-Kelvin Parametric Feedback Cooling of a Laser-Trapped Nanoparticle," Phys. Rev. Lett. 109, 103603 (2012).
- [S4] H. Wiseman and G. Milburn, "Quantum theory of optical feedback via homodyne detection," Phys. Rev. Lett. 70, 548–551 (1993).
- [S5] L. P. Neukirch and A. N. Vamivakas, "Nano-optomechanics with optically levitated nanoparticles," Contemporary Physics 56, 48–62 (2015).
- [S6] S. A. Beresnev, V. G. Chernyak, and G. A. Fomyagin, "Motion of a spherical particle in a rarefied gas. Part 2. Drag and thermal polarization," Journal of Fluid Mechanics 219, 405 (2006).
- [S7] K. Hornberger, J. E. Sipe, and M. Arndt, "Theory of decoherence in a matter wave Talbot-Lau interferometer," Phys. Rev. A 70, 053608 (2004).
- [S8] L. Diósi, "Quantum Master Equation of a Particle in a Gas Environment," Europhys. Lett. **30**, 63–68 (1995).
- [S9] G. Lindblad, "On the generators of quantum dynamical semigroups," Commun. Math. Phys. 48, 119–130 (1976).
- [S10] N. Gisin and I. C. Percival, "The quantum-state diffusion model applied to open systems," J. Phys. A. Math. Gen. 25, 5677–5691 (1992).
- [S11] J. Halliwell and A. Zoupas, "Quantum state diffusion, density matrix diagonalization, and decoherent histories: A model," Phys. Rev. D 52, 7294 (1995).
- [S12] T. A. Brun, "Continuous measurements, quantum trajectories, and decoherent histories," Phys. Rev. A 61, 31 (1997).