

Higher order elicibility and Osband's principle

Tobias Fissler* Johanna F. Ziegel*

December 4, 2018

Abstract

A statistical functional, such as the mean or the median, is called elicitable if there is a scoring function or loss function such that the correct forecast of the functional is the unique minimizer of the expected score. Such scoring functions are called strictly consistent for the functional. The elicibility of a functional opens the possibility to compare competing forecasts and to rank them in terms of their realized scores. In this paper, we explore the notion of elicibility for multi-dimensional functionals and give both necessary and sufficient conditions for strictly consistent scoring functions. We cover the case of functionals with elicitable components, but we also show that one-dimensional functionals that are not elicitable can be a component of a higher order elicitable functional. In the case of the variance this is a known result. However, an important result of this paper is that spectral risk measures with a spectral measure with finite support are jointly elicitable if one adds the 'correct' quantiles. A direct consequence of applied interest is that the pair (Value at Risk, Expected Shortfall) is jointly elicitable under mild conditions that are usually fulfilled in risk management applications.

Keywords: Consistency; Decision theory; Elicibility; Expected Shortfall; Point forecasts; Propriety; Scoring functions; Scoring rules; Spectral risk measures; Value at Risk

AMS 2010 Subject Classification: 62C99; 91B06

1 Introduction

Point forecasts for uncertain future events are issued in a variety of different contexts such as business, government, risk-management or meteorology, and they are often used as the basis for strategic decisions. In all these situations, one has a random quantity Y with unknown distribution F . One is interested in a statistical property of F , that is a functional $T(F)$. Here, Y can be real-valued (GDP growth for next year), vector-valued (wind-speed, income from taxes for all cantons of Switzerland), functional-valued (path of the interchange rate Euro - Swiss franc over one day), or set-valued (area of rain tomorrow,

*University of Bern, Department of Mathematics and Statistics, Institute of Mathematical Statistics and Actuarial Science, Sidlerstrasse 5, 3012 Bern, Switzerland, e-mails: tobias.fissler@stat.unibe.ch and johanna.ziegel@stat.unibe.ch

area of influenza in a country). Likewise, also the functional T can have a variety of different sorts of values, amongst them the real- and vector-valued case (mean, vector of moments, covariance matrix, expectiles), the set-valued case (confidence regions) or also the functional-valued case (distribution functions). This article is concerned with the situation where Y is a d -dimensional random vector and T is a k -dimensional functional, thus also covering the the real-valued case.

It is common to assess and compare competing point forecasts in terms of a loss function or *scoring function*. This is a function S such as the squared error or the absolute error which is negatively oriented in the following sense: If the forecast $x \in \mathbb{R}^k$ is issued and the event $y \in \mathbb{R}^d$ materializes, the forecaster is *penalized* by the real value $S(x, y)$. In the presence of several different forecasters one can compare their performances by ranking their realized scores. Hence, forecasters have an incentive to minimize their Bayes risk or *expected loss* $\mathbb{E}_F[S(x, Y)]$. Gneiting (2011) demonstrated impressively that scoring functions should be incentive compatible in that they should encourage the forecasters to issue truthful reports; see also Murphy and Daan (1985); Engelberg et al. (2009). In other words, the choice of the scoring function S must be consistent with the choice of the functional T . We say a scoring function S is \mathcal{F} -consistent for a functional T if $T(F) \in \arg \min_x \mathbb{E}_F[S(x, Y)]$ for all $F \in \mathcal{F}$ where the class \mathcal{F} of probability distributions is the domain of T . If $T(F)$ is the unique minimizer of the expected score for all $F \in \mathcal{F}$ we say that S is *strictly \mathcal{F} -consistent for T* . Hence, a strictly \mathcal{F} -consistent scoring function for T elicits T . Following Lambert et al. (2008) and Gneiting (2011), we call a functional T with domain \mathcal{F} *elicitable* if there exists a strictly \mathcal{F} -consistent scoring function for T .

The elicibility of a functional allows for regression, such as quantile regression and expectile regression (Koenker, 2005; Newey and Powell, 1987) and for M-estimation (Huber, 1964). Early work on elicibility is due to Osband (1985); Osband and Reichelstein (1985). More recent advances in the one-dimensional case, that is $k = d = 1$ are due to Gneiting (2011); Lambert (2013); Steinwart et al. (2014) with the latter showing the intimate relation between elicibility and identifiability. Under mild conditions, many important functionals are elicitable such as moments, ratios of moments, quantiles and expectiles. However, there are also relevant functionals which are not elicitable such as variance, mode, or Expected Shortfall (Osband, 1985; Weber, 2006; Gneiting, 2011; Heinrich, 2013).

With the so-called *revelation principle* (see Proposition 2.13) Osband (1985) was one of the first to show that a functional, albeit itself not being elicitable, can be a component of an elicitable vector-valued functional. The most prominent example in this direction is that the pair (mean, variance) is elicitable despite the fact that variance itself is not. However, it is crucial for the validity of the revelation principle that there is a bijection between the pair (mean, variance) and the first two moments. Until now, it appeared as an open problem if there are elicitable functionals with non-elicitable components other than those which can be connected to a functional with elicitable components via a bijection. Frongillo and Kash (2014) conjectured that this is generally not possible. We solve this open problem and can reject their conjecture: Corollary 5.5 shows that the pair (Value at Risk, Expected Shortfall) is elicitable, subject to mild regularity assumptions, improving a recent partial result of Acerbi and Szekely (2014). To the best of our knowledge, we

provide the first proof of this result in full generality. In fact, Corollary 5.4 demonstrates more generally that *spectral risk measures* with a spectral measure having finite support in $(0, 1]$ can be a component of an elicitable vector-valued functional. These results may lead to a new direction in the contemporary discussion about what risk measure is best in practice, and in particular about the importance of elicibility in risk measurement contexts (Embrechts and Hofert, 2014; Emmer et al., 2013; Davis, 2013; Acerbi and Szekely, 2014).

Complementing the question whether a functional is elicitable or not, it is interesting to determine the class of strictly consistent scoring functions for a functional, or at least to characterize necessary and sufficient conditions for the strict consistency of a scoring function. Most of the existing literature focuses on real-valued functionals meaning that $k = 1$. For the case $k > 1$, mainly linear functionals, that is, vectors of expectations of certain transformations, are classified where the only strictly consistent scoring functions are *Bregman functions* (Savage, 1971; Osband and Reichelstein, 1985; Dawid and Sebastiani, 1999; Banerjee et al., 2005; Abernethy and Frongillo, 2012); for a general overview of the existing literature, we refer to Gneiting (2011). To the best of our knowledge, only Osband (1985), Lambert et al. (2008) and Frongillo and Kash (2014) investigated more general cases of functionals, the latter also treating vectors of ratios of expectations as the first non-linear functionals. In his doctoral thesis, Osband (1985) established a necessary representation for the first order derivative of a strictly consistent scoring function with respect to the report x which connects it with identification functions. Following Gneiting (2011) we call results in the same flavor *Osband's principle*. Theorem 3.2 in this paper complements and generalizes Osband (1985, Theorem 2.1). Using our techniques, we retrieve the results mentioned above concerning the Bregman representation, however under somewhat stronger regularity assumptions than the one in Frongillo and Kash (2014); see Corollary 4.3. On the other hand, we are able to treat a much broader class of functionals; see Proposition 4.1, Remark 4.4 and Theorem 5.2. In particular, we show that under mild richness assumptions on the class \mathcal{F} , any strictly \mathcal{F} -consistent scoring function for a vector of quantiles and / or expectiles is the sum of strictly \mathcal{F} -consistent one dimensional scoring functions for each quantile / expectile; see Corollary 4.2.

The paper is organized as follows. In Section 2, we introduce notation and derive some basic results concerning the elicibility of k -dimensional functionals. Section 3 is concerned with *Osband's principle*, Theorem 3.2, and its immediate consequences. We investigate the situation where a functional is composed of elicitable components in Section 4, whereas Section 5 is dedicated to the elicibility of spectral risk measures. We end our article with a brief discussion; see Section 6. Most proofs are deferred to Section 7.

2 Properties of higher order elicibility

2.1 Notation and definitions

Following Gneiting (2011), we introduce a decision-theoretic framework for the evaluation of point forecasts. To this end, we introduce an *observation domain* $\mathcal{O} \subseteq \mathbb{R}^d$. We equip

\mathbf{O} with the Borel σ -algebra \mathcal{O} using the induced topology of \mathbb{R}^d . We identify a Borel probability measure P on $(\mathbf{O}, \mathcal{O})$ with its cumulative distribution function (cdf) $F_P: \mathbf{O} \rightarrow [0, 1]$ defined as $F_P(x) := P((-\infty, x] \cap \mathbf{O})$, where $(-\infty, x] = (-\infty, x_1] \times \cdots \times (-\infty, x_d]$ for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$. Let \mathcal{F} be a class of distribution functions on $(\mathbf{O}, \mathcal{O})$. Furthermore, for some integer $k \geq 1$, let $\mathbf{A} \subseteq \mathbb{R}^k$ be an *action domain*. To shorten notation, we usually write $F \in \mathcal{F}$ for a cdf and also omit to mention the σ -algebra \mathcal{O} .

Let $T: \mathcal{F} \rightarrow \mathbf{A}$ be a functional. We introduce the notation $T(\mathcal{F}) := \{x \in \mathbf{A}: x = T(F) \text{ for some } F \in \mathcal{F}\}$. For a set $M \subseteq \mathbb{R}^k$ we will write $\text{int}(M)$ for its interior with respect to \mathbb{R}^k , that is, $\text{int}(M)$ is the biggest open set $U \subseteq \mathbb{R}^k$ such that $U \subseteq M$. The convex hull of M is defined as

$$\text{conv}(M) := \left\{ \sum_{i=1}^n \lambda_i x_i \mid n \in \mathbb{N}, x_1, \dots, x_n \in M, \lambda_1, \dots, \lambda_n > 0, \sum_{i=1}^n \lambda_i = 1 \right\}.$$

We say that a function $a: \mathbf{O} \rightarrow \mathbb{R}$ is \mathcal{F} -integrable if it is F -integrable for each $F \in \mathcal{F}$. A function $g: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ is \mathcal{F} -integrable if $g(x, \cdot)$ is \mathcal{F} -integrable for each $x \in \mathbf{A}$. If g is \mathcal{F} -integrable, we introduce the map

$$\bar{g}: \mathbf{A} \times \mathcal{F} \rightarrow \mathbb{R}, \quad (x, F) \mapsto \bar{g}(x, F) = \int g(x, y) \, dF(y).$$

Consequently, for fixed $F \in \mathcal{F}$ we can consider the function $\bar{g}(\cdot, F): \mathbf{A} \rightarrow \mathbb{R}$, $x \mapsto \bar{g}(x, F)$, and for fixed $x \in \mathbf{A}$ we can consider the (linear) functional $\bar{g}(x, \cdot): \mathcal{F} \rightarrow \mathbb{R}$, $F \mapsto \bar{g}(x, F)$. We introduce the \mathbb{R} -vector space $\hat{\mathcal{F}} := \text{span}(\mathcal{F})$ which consists of all finite linear combinations of elements in \mathcal{F} . For a fixed $x \in \mathbf{A}$ we can use linearity to extend the functional $\bar{g}(x, \cdot)$ to a linear functional on $\hat{\mathcal{F}}$. If we fix $y \in \mathbf{O}$ and g is sufficiently smooth in its first argument, then for $m \in \{1, \dots, k\}$ we denote the m -th partial derivative of the function $g(\cdot, y)$ with $\partial_m g(\cdot, y)$. More formally, we set

$$\partial_m g(\cdot, y): \text{int}(\mathbf{A}) \rightarrow \mathbb{R}, \quad (x_1, \dots, x_k) \mapsto \frac{\partial}{\partial x_m} g(x_1, \dots, x_k, y).$$

We denote by $\nabla g(\cdot, y)$ the gradient of $g(\cdot, y)$ defined as $\nabla g(\cdot, y) := (\partial_1 g(\cdot, y), \dots, \partial_k g(\cdot, y))^\top$; and with $\nabla^2 g(\cdot, y) := (\partial_l \partial_m g(\cdot, y))_{l, m=1, \dots, k}$ the Hessian of $g(\cdot, y)$. *Mutatis mutandis*, we use the same notation for $\bar{g}(\cdot, F)$, $F \in \mathcal{F}$. We call a function on \mathbf{A} differentiable if it is differentiable in $\text{int}(\mathbf{A})$ and use the notation as given above. The restriction of a function f to some subset M of its domain is denoted by $f|_M$.

Definition 2.1 (Consistency). A *scoring function* is an \mathcal{F} -integrable function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$. It is said to be \mathcal{F} -consistent for a functional $T: \mathcal{F} \rightarrow \mathbf{A}$ if

$$\bar{S}(T(F), F) \leq \bar{S}(x, F)$$

for all $F \in \mathcal{F}$ and for all $x \in \mathbf{A}$. Furthermore, S is *strictly \mathcal{F} -consistent* for T if it is \mathcal{F} -consistent for T and if

$$\bar{S}(T(F), F) = \bar{S}(x, F) \implies x = T(F)$$

for all $F \in \mathcal{F}$ and for all $x \in \mathbf{A}$. Wherever it is convenient we assume that $S(x, \cdot)$ is locally bounded for all $x \in \mathbf{A}$.

Definition 2.2 (*k*-Elicitability). A functional $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ is called *k-elicitable*, if there exists a strictly \mathcal{F} -consistent scoring function for T .

Definition 2.3 (Identification function). An *identification function* is an \mathcal{F} -integrable function $V: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}^k$. It is said to be an *\mathcal{F} -identification function* for a functional $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ if

$$\bar{V}(T(F), F) = 0$$

for all $F \in \mathcal{F}$. Furthermore, V is a *strict \mathcal{F} -identification function* for T if

$$\bar{V}(x, F) = 0 \iff x = T(F)$$

for all $F \in \mathcal{F}$ and for all $x \in \mathbf{A}$. Wherever it is convenient we assume that $V(x, \cdot)$ is locally bounded for all $x \in \mathbf{A}$ and that $V(\cdot, y)$ is locally Lebesgue-integrable for all $y \in \mathbf{O}$.

Definition 2.4 (*k*-identifiability). A functional $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ is said to be *k-identifiable*, if there exists a strict \mathcal{F} -identification function for T .

If the dimension k is clear from the context, we say that a functional is elicitable (identifiable) instead of *k-elicitable* (*k-identifiable*).

Remark 2.5. Depending on the class \mathcal{F} , some statistical functionals such as quantiles can be set-valued. In such situations, one can define $T: \mathcal{F} \rightarrow 2^{\mathbf{A}}$. Then, a scoring function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ is called (strictly) \mathcal{F} -consistent for T if $\bar{S}(t, F) \leq \bar{S}(x, F)$ for all $x \in \mathbf{A}$, $F \in \mathcal{F}$ and $t \in T(F)$ (with equality implying $x \in T(F)$). The definition of a (strict) \mathcal{F} -identification function for T can be generalized *mutatis mutandis*. Many of the results of this paper can be extended to the case of set-valued functionals – at the cost of a more involved notation and analysis. To allow for a clear presentation, we confine ourselves to functionals with values in \mathbb{R}^k in this paper.

If $V: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}^k$ is an \mathcal{F} -identification function for a functional $T: \mathcal{F} \rightarrow \mathbf{A}$ and $h: \mathbf{A} \rightarrow \mathbb{R}^{k \times k}$ is a matrix-valued function, then the function

$$hV: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}^k, \quad (x, y) \mapsto hV(x, y) := h(x)V(x, y)$$

is again an \mathcal{F} -identification function for T . If V is a strict \mathcal{F} -identification function for T and $\det h(x) \neq 0$ for all $x \in \mathbf{A}$, then hV is also a strict \mathcal{F} -identification function for T .

Remark 2.6. Steinwart et al. (2014) introduced the notion of an *oriented* strict \mathcal{F} -identification function for the case $k = 1$ (and $d = 1$). They say that $V: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ is an oriented strict \mathcal{F} -identification function for the functional $T: \mathcal{F} \rightarrow \mathbf{A}$ if V is a strict \mathcal{F} -identification function for T and moreover

$$\bar{V}(x, F) > 0 \iff x > T(F) \tag{2.1}$$

for all $F \in \mathcal{F}$ and for all $x \in \mathbf{A}$. They show – under some regularity assumptions such as the continuity of the functional T – that if V is a strict \mathcal{F} -identification function for the functional T then either V or $-V$ is oriented; see Steinwart et al. (2014, Lemma 6). This notion of orientation can also be generalized to the case $k > 1$.

Definition 2.7 (Orientation). Let $T: \mathcal{F} \rightarrow \mathbf{A}$ be a functional with a strict \mathcal{F} -identification function $V: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}^k$. Then V is called an *oriented* strict \mathcal{F} -identification function for T if

$$v^\top \bar{V}(T(F) + sv, F) > 0 \iff s > 0$$

for all $v \in \mathbb{S}^{k-1} := \{x \in \mathbb{R}^k: \|x\| = 1\}$, for all $F \in \mathcal{F}$ with $T(F) \in \mathbf{A}$ and for all $s \in \mathbb{R}$ such that $T(F) + sv \in \mathbf{A}$.

Indeed, the one-dimensional definition of orientation at (2.1) is nested in Definition 2.7 upon recalling that $\mathbb{S}^0 = \{-1, 1\}$. Under some smoothness assumptions, we can give a necessary condition for the orientation of a strict \mathcal{F} -identification function V : Assume that the function $\mathbf{A} \rightarrow \mathbb{R}^k$, $x \mapsto \bar{V}(x, F)$ is partially differentiable. If V is oriented then the matrix $(\partial_l \bar{V}_r(t, F))_{r,l=1,\dots,k}$ is positive semi-definite for all $F \in \mathcal{F}$ and $t = T(F)$. It appears to be an open question under which conditions there exists an oriented identification function for an identifiable functional. In the light of Lemma 2.9 (ii), Remark 2.10 and Proposition 3.5 this would give insight whether the construction of a strictly proper scoring function is possible.

Remark 2.8. Our notion of orientation differs from the one proposed by Frongillo and Kash (2014). In contrast to their definition, our definition is *per se* independent of a (possibly non-existing) strictly consistent scoring function for T . Moreover, with respect to Lemma 2.9 (ii) and Remark 2.10, the orientation of the gradient of a scoring function implies its strict consistency.

2.2 Basic results

The first lemma gives a sufficient condition for strict consistency and connects the notions of scoring functions and identification functions.

Lemma 2.9. (i) A scoring function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ is strictly \mathcal{F} -consistent for $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ if and only if the function

$$\psi: D \rightarrow \mathbb{R}, \quad s \mapsto \bar{S}(t + sv, F)$$

has a global (unique) minimum at $s = 0$ for all $F \in \mathcal{F}$, $t = T(F)$ and $v \in \mathbb{S}^{k-1}$ where $D = \{s \in \mathbb{R}: t + sv \in \mathbf{A}\}$.

(ii) Let $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ be a scoring function that is continuously differentiable in its first argument and let $\mathcal{F}' = T^{-1}(\text{int}(\mathbf{A})) \subseteq \mathcal{F}$. If $\nabla S: \text{int}(\mathbf{A}) \times \mathbf{O} \rightarrow \mathbb{R}^k$ is an oriented strict \mathcal{F}' -identification function for $T|_{\mathcal{F}'}$ then $S|_{\text{int}(\mathbf{A}) \times \mathbf{O}}$ is a strictly \mathcal{F}' -consistent scoring function for $T|_{\mathcal{F}'}$.

Remark 2.10. One can weaken the assumptions of Lemma 2.9 (ii) on the smoothness of S . Let $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ be a scoring function such that $\bar{S}(\cdot, F)$ is continuously differentiable for all $F \in \mathcal{F}$. If \mathcal{F} consists of absolutely continuous distributions, this is a much weaker requirement; see Section 3 for a detailed discussion. Let $\mathcal{F}' = T^{-1}(\text{int}(\mathbf{A})) \subseteq \mathcal{F}$. If for all

$F' \in \mathcal{F}$, $t = T(F) \in \text{int}(\mathbf{A})$, for all $v \in \mathbb{S}^{k-1}$ and for all $s \in \mathbb{R}$ such that $t + sv \in \text{int}(\mathbf{A})$ we have that

$$v^\top \nabla \bar{S}(t + sv, F) \begin{cases} > 0, & \text{if } s > 0 \\ = 0, & \text{if } s = 0 \\ < 0, & \text{if } s < 0 \end{cases}$$

then $S_{|\text{int}(\mathbf{A}) \times \mathbf{O}}$ is a strictly \mathcal{F}' -consistent scoring function for $T_{|\mathcal{F}'}$.

The following result follows directly from the definition of consistency (Definition 2.1). However, it is crucial to understand many of the results of this paper.

Lemma 2.11. *Let $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ be a functional with a strictly \mathcal{F} -consistent scoring function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$. Then the following two assertions hold.*

(i) *Let $\mathcal{F}' \subseteq \mathcal{F}$ and $T_{|\mathcal{F}'}$ be the restriction of T to \mathcal{F}' . Then S is also a strictly \mathcal{F}' -consistent scoring function for $T_{|\mathcal{F}'}$.*

(ii) *Let $\mathbf{A}' \subseteq \mathbf{A}$ such that $T(\mathcal{F}) \subseteq \mathbf{A}'$ and $S_{|\mathbf{A}' \times \mathbf{O}}$ be the restriction of S to $\mathbf{A}' \times \mathbf{O}$. Then $S_{|\mathbf{A}' \times \mathbf{O}}$ is also a strictly \mathcal{F} -consistent scoring function for T .*

The main results of this paper consist of necessary and sufficient conditions for the strict \mathcal{F} -consistency of a scoring function S for some functional T . What are the consequences of Lemma 2.11 for such conditions? Assume that we start with a functional $T': \mathcal{F}' \rightarrow \mathbf{A}' \subseteq \mathbb{R}^k$ and deduce some *necessary* conditions for a scoring function $S': \mathbf{A}' \times \mathbf{O} \rightarrow \mathbb{R}$ to be strictly \mathcal{F}' -consistent for T' . Then Lemma 2.11 (i) implies that these conditions continue to be necessary conditions for the strict \mathcal{F} -consistency of S' for $T: \mathcal{F} \rightarrow \mathbf{A}'$ where $\mathcal{F}' \subseteq \mathcal{F}$, and T is some extension of T' such that $T(\mathcal{F}) \subseteq \mathbf{A}'$. On the other hand, Lemma 2.11 (ii) implies that the necessary conditions for the strict \mathcal{F}' -consistency of a scoring function $S': \mathbf{A}' \times \mathbf{O} \rightarrow \mathbb{R}$ continue to be necessary conditions for the strict \mathcal{F} -consistency of S for T' , where $\mathbf{A}' \subseteq \mathbf{A}$ and S is some extension of S' .

Summarizing, given a functional $T: \mathcal{F} \rightarrow \mathbf{A}$, a collection of necessary conditions for the strict \mathcal{F} -consistency of scoring functions for T is the more restrictive the smaller the class \mathcal{F} and the smaller the set \mathbf{A} is (provided that $T(\mathcal{F}) \subseteq \mathbf{A}$, of course). Hence, in the forthcoming results concerning necessary conditions, it is no loss of generality to just mention which distributions must necessarily be in the class \mathcal{F} to guarantee the validity of the results. Furthermore, it is no loss of generality to make the assumption that T is surjective, so $\mathbf{A} = T(\mathcal{F})$.

Some of the subsequent results also provide *sufficient* conditions for the strict \mathcal{F} -consistency of a scoring function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ for a functional $T: \mathcal{F} \rightarrow \mathbf{A}$. Those results are the stronger the bigger the class \mathcal{F} and the bigger the set \mathbf{A} is. For the notion of elicibility this means that the assertion that a functional $T: \mathcal{F} \rightarrow \mathbf{A}$ is elicitable is also the stronger the bigger the class \mathcal{F} and the bigger the set \mathbf{A} is. To demonstrate this reasoning, be aware that if the functional $T: \mathcal{F} \rightarrow \mathbf{A}$ is degenerate in the sense that it is constant, so $T = t$ for some $t \in \mathbf{A}$ (which covers the particular case that \mathcal{F} contains only one element), then T is automatically elicitable with a strictly \mathcal{F} -consistent scoring function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$, defined as $S(x, y) := \|x - t\|$.

Strictly consistent scoring functions for a given functional T are not unique. In particular, the following result generalizes directly from the one-dimensional case. Let $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ be a strictly \mathcal{F} -consistent scoring function a functional $T: \mathcal{F} \rightarrow \mathbf{A}$. Then, for any $\lambda > 0$ and any \mathcal{F} -integrable function $a: \mathbf{O} \rightarrow \mathbb{R}$, the scoring function

$$\tilde{S}(x, y) := \lambda S(x, y) + a(y) \quad (2.2)$$

is again strictly \mathcal{F} -consistent for T . Gneiting (2011, Theorem 2) shows that in the one-dimensional case under the assumption $S(x, y) \geq 0$, the class of consistent scoring functions is a convex cone. Generally, the assumption of scoring functions being nonnegative is natural if $\delta_y \in \mathcal{F}$ for all $y \in \mathbf{O}$ because for an \mathcal{F} -consistent scoring function S , the scoring function $\tilde{S}(x, y) := S(x, y) - \bar{S}(T(\delta_y), \delta_y) \geq 0$ and it is of the form (2.2) if $y \mapsto \bar{S}(T(\delta_y), \delta_y)$ is \mathcal{F} -integrable. As we are particularly interested in classes \mathcal{F} of absolutely continuous distributions in this manuscript, we do not require scoring functions to be nonnegative. We generalize Gneiting (2011, Theorem 2) as follows showing that the class of strictly \mathcal{F} -consistent scoring functions for T is a convex cone (not including zero). The proof follows easily using Fubini's theorem and is omitted.

Proposition 2.12. *Let $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ be a functional. Let (Z, \mathcal{Z}) be a measurable space with a σ -finite measure ν where $\nu \neq 0$. Let $\{S_z: z \in Z\}$ be a family of strictly \mathcal{F} -consistent scoring functions $S_z: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ for T . If for all $x \in \mathbf{A}$ and for all $F \in \mathcal{F}$ the map*

$$Z \times \mathbf{O} \rightarrow \mathbb{R}, \quad (z, y) \mapsto S_z(x, y)$$

is $\nu \otimes F$ -integrable, then the scoring function

$$S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}, \quad (x, y) \mapsto S(x, y) = \int_Z S_z(x, y) \nu(dz)$$

is strictly \mathcal{F} -consistent for T .

Point forecasts and probabilistic forecasts are closely related. Probabilistic forecasts, issuing a whole probability distribution, can be evaluated in terms of *scoring rules* (Winkler, 1996; Gneiting and Raftery, 2007). A scoring rule is a map $R: \mathcal{F} \times \mathbf{O} \rightarrow \mathbb{R}$ such that for each $G \in \mathcal{F}$, the map $\mathbf{O} \rightarrow \mathbb{R}$, $y \mapsto R(G, y)$ is \mathcal{F} -integrable. A scoring rule is (strictly) \mathcal{F} -proper if $\bar{R}(F, F) \leq \bar{R}(G, F)$ for all $F, G \in \mathcal{F}$ (with equality implying $F = G$). As in the one-dimensional case (Gneiting, 2011, Theorem 3), each \mathcal{F} -consistent scoring function S for a functional $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ induces an \mathcal{F} -proper scoring rule R via

$$R: \mathcal{F} \times \mathbf{O} \rightarrow \mathbb{R}, \quad (F, y) \mapsto R(F, y) = S(T(F), y).$$

However, if we do not impose that the functional T is injective, we cannot conclude that R is a *strictly* \mathcal{F} -proper scoring rule even if the scoring function S is strictly \mathcal{F} -consistent.

Many important statistical functionals are transformations of other statistical functionals, for example variance and first and second moment are related in this manner. The following *revelation principle*, which originates from Osband (1985, p.8) and is also given in Gneiting (2011, Theorem 4) states that if two functionals are related by a bijection, then one of them is elicitable if and only if the other one is elicitable. The assertion also holds upon replacing 'elicitable' with 'identifiable'. We omit the proof which is straight forward.

Proposition 2.13 (Revelation principle). *Let $g: \mathbf{A} \rightarrow \mathbf{A}'$ be a bijection with inverse g^{-1} , where $\mathbf{A}, \mathbf{A}' \subseteq \mathbb{R}^k$. Let $T: \mathcal{F} \rightarrow \mathbf{A}$ be a functional. Then the following two assertions hold.*

- (i) *The functional $T: \mathcal{F} \rightarrow \mathbf{A}$ is identifiable if and only if $T_g = g \circ T: \mathcal{F} \rightarrow \mathbf{A}'$ is identifiable. The function $V: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}^k$ is a strict \mathcal{F} -identification function for T if and only if*

$$V_g: \mathbf{A}' \times \mathbf{O} \rightarrow \mathbb{R}^k, \quad (x', y) \mapsto V_g(x', y) = V(g^{-1}(x'), y)$$

is a strict \mathcal{F} -identification function for T_g .

- (ii) *The functional $T: \mathcal{F} \rightarrow \mathbf{A}$ is elicitable if and only if $T_g = g \circ T: \mathcal{F} \rightarrow \mathbf{A}'$ is elicitable. The function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ is a strictly \mathcal{F} -consistent scoring function for T if and only if*

$$S_g: \mathbf{A}' \times \mathbf{O} \rightarrow \mathbb{R}, \quad (x', y) \mapsto S_g(x', y) = S(g^{-1}(x'), y)$$

is a strictly \mathcal{F} -consistent scoring function for T_g .

We remark that also (Gneiting, 2011, Theorem 5) on weighted scoring functions carries over directly to the higher order case. Furthermore, convexity of level sets continues to be a necessary condition for elicibility. The result is classical in the literature and was first presented in Osband (1985, Proposition 2.5); see also Gneiting (2011, Theorem 6).

Proposition 2.14 (Osband). *Let $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ be an elicitable functional. Then for all $F_0, F_1 \in \mathcal{F}$ with $t := T(F_0) = T(F_1)$ and for all $\lambda \in (0, 1)$ such that $F_\lambda := (1-\lambda)F_0 + \lambda F_1 \in \mathcal{F}$ it holds that $t = T(F_\lambda)$.*

As a last result in this section, we present the intuitive observation that a vector of elicitable functionals itself is elicitable.

Lemma 2.15. *Let $k_1, \dots, k_l \geq 1$ and let $T_m: \mathcal{F} \rightarrow \mathbf{A}_m \subseteq \mathbb{R}^{k_m}$ be a k_m -elicitable functional, $m \in \{1, \dots, l\}$. Then the functional $T = (T_1, \dots, T_l): \mathcal{F} \rightarrow \mathbf{A}$ is k -elicitable where $k = k_1 + \dots + k_l$ and $\mathbf{A} = \mathbf{A}_1 \times \dots \times \mathbf{A}_l \subseteq \mathbb{R}^k$.*

Proof. For $m \in \{1, \dots, l\}$ let $S_m: \mathbf{A}_m \times \mathbf{O} \rightarrow \mathbb{R}$ be a strictly \mathcal{F} -consistent scoring function for T_m . Let $\lambda_1, \dots, \lambda_l > 0$ be positive real numbers. Then

$$S: \mathbf{A}_1 \times \dots \times \mathbf{A}_l \times \mathbf{O} \rightarrow \mathbb{R}, \quad (x_1, \dots, x_l, y) \mapsto S(x_1, \dots, x_l, y) := \sum_{m=1}^l \lambda_m S_m(x_m, y) \quad (2.3)$$

is a strictly \mathcal{F} -consistent scoring function for T . □

A particularly simple and relevant case of Lemma 2.15 is the situation $k_1 = \dots = k_l = 1$ such that $k = l$. It is an interesting question whether the scoring functions of the form (2.3) are the only strictly \mathcal{F} -consistent scoring functions for T , which amounts to the question of *separability* of scoring rules that was posed by Frongillo and Kash (2014). The answer is generally negative. As mentioned in the introduction, it is known that

all Bregman functions elicit T , if the components of T are all expectations of transformations of Y (Savage, 1971; Osband and Reichelstein, 1985; Dawid and Sebastiani, 1999; Banerjee et al., 2005; Abernethy and Frongillo, 2012) or ratios of expectations with the same denominator (Frongillo and Kash, 2014); see also Corollary 4.3. However, for other situations, such as a combination of different quantiles and/or expectiles, the answer is positive; see Corollary 4.2. These results rely on ‘Osband’s principle’ which gives necessary conditions for scoring functions to be strictly \mathcal{F} -consistent for a given functional T ; see Section 3.

There are more involved functionals that are k -elicitable than just the mere combination of k 1-elicitable components. To illustrate this with a first example, recall that the variance does not have convex level sets in the sense of Proposition 2.14, whence it is not elicitable. However, we can easily show that the pair (expectation, variance) is 2-elicitable.

Corollary 2.16. *Let \mathcal{F} be a class of distribution functions on \mathbb{R} with finite second moments. Then, the functional $T = (T_1, T_2): \mathcal{F} \rightarrow \mathbb{R}^2$, defined as $T_1(F) = \int_{\mathbb{R}} y \, dF(y)$, $T_2(F) = \int_{\mathbb{R}} y^2 \, dF(y) - (\int_{\mathbb{R}} y \, dF(y))^2$ is 2-elicitable.*

Proof. Let $\phi: \mathbb{R} \rightarrow \mathbb{R}$, $z \mapsto \phi(z) = z^2/(1 + |z|)$. The scoring function $S_1: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $(x_1, y) \mapsto S_1(x_1, y) = \phi(y) - \phi(x_1) - \phi'(x_1)(y - x_1)$ is a strictly \mathcal{F} -consistent scoring function for the expectation and $S_2: [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$, $(x_2, y) \mapsto S_2(x_2, y) = \phi(y^2) - \phi(x_2) - \phi'(x_2)(y^2 - x_2)$ is a strictly \mathcal{F} -consistent scoring function for the second moment. Hence, invoking Lemma 2.15, the pair (expectation, second moment) is 2-elicitable. Using the revelation principle given in Proposition 2.13 yields the assertion. \square

In Section 5, we show that the concept of k -elicitability is not restricted to functionals that can be obtained by combining Lemma 2.15 and the revelation principle. It is shown in Weber (2006, Example 3.4) and Gneiting (2011, Theorem 11) that the coherent risk measure Expected Shortfall at level α , $\alpha \in (0, 1)$, does not have convex level sets and is therefore not elicitable. In contrast, we show in Corollary 5.5 that the pair (Value at Risk $_{\alpha}$, Expected Shortfall $_{\alpha}$) is 2-elicitable relative to the class of distributions on \mathbb{R} with finite first moment. This refutes both Conjecture 1 of Frongillo and Kash (2014) and Proposition 2.3 of Osband (1985); see Remark 5.7 for a discussion.

3 Osband’s principle

In this section, we give necessary conditions for the strict \mathcal{F} -consistency of a scoring function S for a functional $T: \mathcal{F} \rightarrow \mathbf{A}$. In the light of Lemma 2.11 and the discussion thereafter, we have to impose some richness conditions on the class \mathcal{F} as well as on the ‘variability’ of the functional T . To this end, we establish a link between strictly \mathcal{F} -consistent scoring functions and strict \mathcal{F} -identification functions. We illustrate the idea in the one-dimensional case. Let \mathcal{F} be a class of distribution functions on \mathbb{R} , $T: \mathcal{F} \rightarrow \mathbb{R}$ a functional and $S: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ a strictly \mathcal{F} -consistent scoring function for T . Furthermore, let $V: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be an oriented strict \mathcal{F} -identification function for T . Then, under

certain regularity conditions, there is a non-negative function $h: \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\frac{d}{dx}S(x, y) = h(x)V(x, y). \quad (3.1)$$

If we naïvely swap differentiation and expectation and h does not vanish, the form (3.1) plus the identification property of V are sufficient for the first order condition on $\bar{S}(\cdot, F)$, $F \in \mathcal{F}$, to be satisfied and the orientation of V as well as the fact that h is positive are sufficient for $\bar{S}(\cdot, F)$ to satisfy the second order condition for strict \mathcal{F} -consistency. So the really interesting part is to show that the form given in (3.1) is necessary for the strict \mathcal{F} -consistency of a scoring function for T .

The idea of this characterization originates from Osband (1985). He gives a characterization including \mathbb{R}^k -valued functionals, but for his proof he assumes that \mathcal{F} contains all distributions with finite support. This is not a problem *per se*, but in the light of Lemma 2.11 and the discussion thereafter it would be desirable to weaken this assumption or to complement the result. Gneiting (2011) illustrates Osband's principle in a quite intuitive manner for the one-dimensional case. In Steinwart et al. (2014, Theorem 5) there is a rigorous statement of Osband's principle for the one-dimensional case. We shall give a proof in the setting of an \mathbb{R}^k -valued functional that does not rely on the existence of distributions with finite support in \mathcal{F} .

Let \mathcal{F} be a class of distribution functions on $\mathbf{O} \subseteq \mathbb{R}^d$ and $\hat{\mathcal{F}} = \text{span}(\mathcal{F})$. Fix a functional $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$, an identification function $V: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}^k$ and a scoring function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$. We introduce the following collection of regularity assumptions.

Assumption (V1). For every $x \in \text{int}(\mathbf{A})$ there are $F_1, \dots, F_{k+1} \in \hat{\mathcal{F}}$ such that

$$0 \in \text{int}(\text{conv}(\{\bar{V}(x, F_1), \dots, \bar{V}(x, F_{k+1})\})).$$

Remark 3.1. We remark that assumption (V1) is equivalent to the condition that for every $x \in \text{int}(\mathbf{A})$ there are $F_1, \dots, F_k \in \hat{\mathcal{F}}$ such that the vectors

$$\bar{V}(x, F_1), \dots, \bar{V}(x, F_k)$$

are linearly independent. We will often make use of this equivalent formulation.

Assumption (V1) implies that the class \mathcal{F} is 'rich' enough meaning that the functional T varies sufficiently in order to derive a necessary form of the scoring function S in Theorem 3.2. We emphasize that assumptions like (V1) are classical in the literature. For the case of k -elicitability, Osband (1985) assumes that $0 \in \text{int}(\text{conv}(\{V(x, y): y \in \mathbf{O}\}))$. Steinwart et al. (2014, Definition 8) and Lambert (2013) treat the case $k = 1$ and work under the assumption that the functional is *strictly locally non-constant* which implies assumption (V1) if the functional is identifiable.

Assumption (V2). For every $F \in \mathcal{F}$, the function

$$\bar{V}(\cdot, F): \mathbf{A} \rightarrow \mathbb{R}^k, \quad x \mapsto \bar{V}(x, F)$$

is continuous.

Assumption (V3). For every $F \in \mathcal{F}$, the function $\bar{V}(\cdot, F)$ is continuously differentiable.

If the function $x \mapsto V(x, y)$, $y \in \mathbf{O}$, is continuous (continuously differentiable), assumption (V2) (assumption (V3)) is directly satisfied, and it is even equivalent to (V2) ((V3)) if \mathcal{F} contains all measures with finite support. However, (V2) and (V3) are much weaker requirements if we move away from distributions with finite support. To illustrate this fact, let $k = 1$ and $V(x, y) = \mathbb{1}\{y \leq x\} - \alpha$, $\alpha \in (0, 1)$, which is a strict \mathcal{F} -identification function for the α -quantile. Of course, $V(\cdot, y)$ is not continuous. But if \mathcal{F} contains only probability distributions F that have a continuous derivative $f = F'$, then $\bar{V}(x, F) = F(x) - \alpha$ and $\frac{d}{dx}\bar{V}(x, F) = f(x)$ and V satisfies (V2) and (V3). The following assumptions (S1) and (S2) are similar conditions as (V2) and (V3) but for scoring functions instead of identification functions.

Assumption (S1). For every $F \in \mathcal{F}$, the function

$$\bar{S}(\cdot, F): \mathbf{A} \rightarrow \mathbb{R}, \quad x \mapsto \bar{S}(x, F)$$

is continuously differentiable.

Assumption (S2). For every $F \in \mathcal{F}$, the function $\bar{S}(\cdot, F)$ is continuously differentiable and the gradient is locally Lipschitz continuous. Furthermore, $\bar{S}(\cdot, F)$ is twice continuously differentiable at $t = T(F) \in \text{int}(\mathbf{A})$.

Note that assumption (S2) implies that the gradient of $\bar{S}(\cdot, F)$ is (totally) differentiable for almost all $x \in \mathbf{A}$ by Rademacher's theorem, which in turn indicates that the Hessian of $\bar{S}(\cdot, F)$ exists for almost all $x \in \mathbf{A}$ and is symmetric by Schwarz's theorem; see Grauert and Fischer (1978, p. 57).

Theorem 3.2 (Osband's principle). *Let \mathcal{F} be a class of distribution functions on $\mathbf{O} \subseteq \mathbb{R}^d$. Let $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ be a surjective, elicitable and identifiable functional with a strict \mathcal{F} -identification function $V: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}^k$ and a strictly \mathcal{F} -consistent scoring function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$. If the assumptions (V1) and (S1) hold, then there exists a matrix-valued function $h: \text{int}(\mathbf{A}) \rightarrow \mathbb{R}^{k \times k}$ such that for $l \in \{1, \dots, k\}$*

$$\partial_l \bar{S}(x, F) = \sum_{m=1}^k h_{lm}(x) \bar{V}_m(x, F) \tag{3.2}$$

for all $x \in \text{int}(\mathbf{A})$ and $F \in \hat{\mathcal{F}}$. If in addition, assumption (V2) holds, then h is continuous. Under the additional assumptions (V3) and (S2), the function h is locally Lipschitz continuous.

The proof of Theorem 3.2 follows closely the idea of the proof of Osband (1985, Theorem 2.1). However, the latter proof only works under the condition that the class \mathcal{F} contains all distributions with finite support. He conjectures that the assertion also holds if \mathcal{F} consists only of absolutely continuous distributions, but we do not believe that his approach is feasible for this case. To show Theorem 3.2, we apply a similar technique as in the proof of Osband (1985, Lemma 2.2) which is based on a finite-dimensional argument.

Remark 3.3. Let $\tilde{h}: \mathbf{A} \rightarrow \mathbb{R}^{k \times k}$ be a function such that the restriction $\tilde{h}|_{\text{int}(\mathbf{A})}$ to $\text{int}(\mathbf{A})$ coincides with the function h in (3.2). Then the function

$$\tilde{h}V: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}^k, \quad (x, y) \mapsto \tilde{h}V(x, y) = \tilde{h}(x)V(x, y)$$

is an \mathcal{F} -identification function for T . If $\det(\tilde{h}(x)) \neq 0$ for all $x \in \mathbf{A}$, then $\tilde{h}V$ is even a strict \mathcal{F} -identification function for T . However, even if V is oriented, $\tilde{h}V$ is not necessarily an oriented strict \mathcal{F} -identification function. To see this, note that due to the strict \mathcal{F} -consistency of S we have that for all $v \in \mathbb{S}^{k-1}$ and $F \in \mathcal{F}$ such that $t = T(F)$ the function

$$\psi: D \rightarrow \mathbb{R}, \quad s \mapsto \psi(s) = \bar{S}(t + sv, F)$$

has a global minimum at $s = 0$ where $D = \{s \in \mathbb{R}: t + sv \in \mathbf{A}\}$. Hence, $\psi'(s) \leq 0$ for all $s < 0$ and $\psi'(s) \geq 0$ for all $s > 0$. We have that

$$\psi'(s) = v^\top \nabla \bar{S}(t + sv, F) = v^\top \overline{\tilde{h}V}(t + sv, F).$$

However, despite the fact that $\tilde{h}V$ is a strict \mathcal{F} -identification function for T , $\overline{\tilde{h}V}(t + sv, F)$ may be orthogonal to v for some $s \neq 0$ which implies $\psi'(s) = 0$.

Under the conditions of Theorem 3.2, equation (3.2) gives a characterization of the partial derivatives of the expected score. If we impose more smoothness assumptions on the expected score, we are also able to give a characterization of the *second* order derivatives of the expected score. In particular, one has the following result.

Corollary 3.4. *For a surjective, elicitable and identifiable functional $T: \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ with a strict \mathcal{F} -identification function $V: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}^k$ and a strictly \mathcal{F} -consistent scoring function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ that satisfy assumptions (V1), (V3) and (S2) we have the following identities for the second order derivatives*

$$\begin{aligned} \partial_m \partial_l \bar{S}(x, F) &= \sum_{i=1}^k \partial_m h_{li}(x) \bar{V}_i(x, F) + h_{li}(x) \partial_m \bar{V}_i(x, F) \\ &= \sum_{i=1}^k \partial_l h_{mi}(x) \bar{V}_i(x, F) + h_{mi}(x) \partial_l \bar{V}_i(x, F) = \partial_l \partial_m \bar{S}(x, F), \end{aligned} \tag{3.3}$$

for all $l, m \in \{1, \dots, k\}$, for all $F \in \mathcal{F}$ and almost all $x \in \text{int}(\mathbf{A})$, where h is the matrix-valued function appearing at (3.2). In particular, 3.3 holds for $x = T(F) \in \text{int}(\mathbf{A})$.

Theorem 3.2 and Corollary 3.4 establish necessary conditions for strictly \mathcal{F} -consistent scoring functions on the level of the expected scores. If the class \mathcal{F} is rich enough and the scoring and identification function smooth enough pointwise in the following sense, we can also deduce a necessary condition for S which holds pointwise.

Assumption (F1). For every $y \in \mathbf{O}$ there exists a sequence $(F_n)_{n \in \mathbb{N}}$ of distributions $F_n \in \mathcal{F}$ that converges weakly to the Dirac-measure δ_y such that the support of F_n is contained in a compact set K for all n .

Assumption (VS1). Let

$$C := \{(x, y) \in \mathbf{A} \times \mathbf{O} \mid V(x, \cdot) \text{ and } S(x, \cdot) \text{ are continuous at the point } y\}.$$

Suppose that the complement of C has $(k + d)$ -dimensional Lebesgue measure zero.

Proposition 3.5. *Assume that $\text{int}(A) \subseteq \mathbb{R}^k$ is a star domain and let $T: \mathcal{F} \rightarrow A$ be a surjective, elicitable and identifiable functional with a strict \mathcal{F} -identification function $V: A \times \mathcal{O} \rightarrow \mathbb{R}^k$ and a strictly \mathcal{F} -consistent scoring function $S: A \times \mathcal{O} \rightarrow \mathbb{R}$. Suppose that assumptions (V1), (V3), (S2), (F1) and (VS1) hold. Let h be the matrix valued function appearing at (3.2). Then, the scoring function S is necessarily of the form*

$$S(x, y) = \sum_{r=1}^k \sum_{m=1}^k \int_{z_r}^{x_r} h_{rm}(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k) \times V_m(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k, y) dv + a(y) \quad (3.4)$$

for almost all $(x, y) \in A \times \mathcal{O}$ for some star point $z = (z_1, \dots, z_k) \in \text{int}(A)$ and some \mathcal{F} -integrable function $a: \mathcal{O} \rightarrow \mathbb{R}$. On the level of the expected score $\bar{S}(x, F)$, equation (3.4) holds for all $x \in \text{int}(A)$, $F \in \hat{\mathcal{F}}$.

While Theorem 3.2, Corollary 3.4 and Proposition 3.5 only establish necessary conditions for strictly \mathcal{F} -consistent scoring functions for some functional T , often, they guide a way how to construct strictly \mathcal{F} -consistent scoring functions starting with a strict \mathcal{F} -identification function V for T . For the one-dimensional case, one can use the fact that, subject to some mild regularity conditions, if V is a strict \mathcal{F} -identification function, then either V or $-V$ is oriented; see Remark 2.6. Supposing that V is oriented, we can choose any strictly positive function $h: A \rightarrow \mathbb{R}$ to get the derivative of a strictly \mathcal{F} -consistent scoring function. Then integration yields the desired strictly \mathcal{F} -consistent scoring function.

Establishing *sufficient* conditions for scoring functions to be strictly \mathcal{F} -consistent for T is generally more involved in the case $k > 1$. First of all, working under assumption (S2), the symmetry of the Hessian $\nabla^2 \bar{S}(x, F)$ imposes strong necessary conditions on the functions h_{lm} ; see for example Proposition 4.1 which treats the case where all components of the functional $T = (T_1, \dots, T_k)$ are elicitable and identifiable. The example of spectral risk measures is treated in Section 5. Secondly, (3.2) and (3.3) are necessary conditions for $\bar{S}(x, F)$ having a *local* minimum in $x = T(F)$, $F \in \mathcal{F}$. Even if we additionally suppose that the Hessian $\nabla^2 \bar{S}(x, F)$ is strictly positive definite at $x = T(F)$, this is a sufficient condition only for a *local* minimum at $x = T(F)$, but does not provide any information concerning a *global* minimum.¹ Consequently, even if the functions h_{lm} satisfy (3.3), one must verify the strict consistency of the scoring function on a case by case basis. This can often be done by showing that the one-dimensional functions $\mathbb{R} \rightarrow \mathbb{R}$, $s \mapsto \bar{S}(t + sv, F)$, with $t = T(F)$, have a global minimum in $s = 0$ for all $v \in \mathbb{S}^{k-1}$ and for all $F \in \mathcal{F}$. This holds for example if the function $(x, y) \mapsto h(x)V(x, y)$ is an oriented strict \mathcal{F} -identification function for T ; see Lemma 2.9. In this step, one may have to impose additional conditions

¹We illustrate this phenomenon with the function

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}, \quad (x_1, x_2) \mapsto f(x_1, x_2) := x_1^2 + x_2^2(1 - x_1)^3.$$

Indeed, f has only one critical point, that is $\nabla f(x_1, x_2) = 0$, if and only if $(x_1, x_2) = (0, 0)$. The Hessian $\nabla^2 f(0, 0)$ is strictly positive definite. Therefore, f has a local minimum at $(x_1, x_2) = (0, 0)$. However, this is not a global minimum since $f(2, 4) = -12 < 0 = f(0, 0)$.

Functional	Strict identification function
Ratio $\mathbb{E}_F[p(Y)]/\mathbb{E}_F[q(Y)]$	$V(x, y) = xq(y) - p(y)$
α -Quantile	$V(x, y) = \mathbf{1}\{y \leq x\} - \alpha$
τ -Expectile	$V(x, y) = 2 \mathbf{1}\{y \leq x\} - \tau (x - y)$

Table 1: Possible choices for strict identification functions for one-dimensional functionals taken from Gneiting (2011, Table 9).

on the functions h_{lm} to ensure sufficiency which cannot always be shown to be necessary; see e.g. Theorem 5.2.

We conclude this section with a remark clarifying how the function h in Osband's principle behaves under the revelation principle.

Remark 3.6. Let $g : \mathbf{A} \rightarrow \mathbf{A}'$ be a bijection, $\mathbf{A}, \mathbf{A}' \subseteq \mathbb{R}^k$. Suppose we have an identification function V for a functional $T : \mathcal{F} \rightarrow \mathbf{A}$ and we choose the identification function $V_g(x', y) = V(g^{-1}(x'), y)$ as an identification function for the functional $T_g = g \circ T$. If the functional T (and hence also T_g by Proposition 2.13) is elicitable, then the gradient of the expected scores of T and T_g are of the form (3.2) with functions h and h_g , respectively. The functions h and h_g are connected by the following relation

$$(h_g)_{lm}(x') = \sum_{r=1}^k \partial_l(g^{-1})_r(x') h_{rm}(g^{-1}(x')), \quad x' \in \mathbf{A}'.$$

4 Functionals with elicitable components

Suppose that the functional $T = (T_1, \dots, T_k) : \mathbb{R} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ consists of 1-elicitable components T_m . As prototypical examples of such 1-elicitable components, we consider the functionals given in Table 1 where we implicitly assume that $\mathbf{0} \subseteq \mathbb{R}$ if a quantile or an expectile are a part of T . With the given identification functions, it turns out that usually T (or some subset of its components) fulfills either one of the following two assumptions.

Assumption (V4). Let assumption (V3) hold. For all $r \in \{1, \dots, k\}$ and for all $t \in \text{int}(\mathbf{A}) \cap T(\mathcal{F})$ there are $F_1, F_2 \in T^{-1}(\{t\})$ such that

$$\partial_l \bar{V}_l(t, F_1) = \partial_l \bar{V}_l(t, F_2) \quad \forall l \in \{1, \dots, k\} \setminus \{r\}, \quad \partial_r \bar{V}_r(t, F_1) \neq \partial_r \bar{V}_r(t, F_2).$$

Assumption (V5). Let assumption (V3) hold. For all $F \in \mathcal{F}$ there is a constant $c_F \neq 0$ such that for all $r \in \{1, \dots, k\}$ and for all $x \in \text{int}(\mathbf{A})$ it holds that

$$\partial_r \bar{V}_r(x, F) = c_F.$$

Following Frongillo and Kash (2014), we call a functional that fulfills assumption (V5) with $c_F = 1$ for all $F \in \mathcal{F}$ a *linear functional*.

Prima facie, assumptions (V4) and (V5) are mutually exclusive. Considering the functionals in Table 1 with the associated identification functions, we obtain, for $x =$

$(x_1, \dots, x_k) \in \mathbb{R}^k$, $F \in \mathcal{F}$ with derivative $F' = f$ and $m \in \{1, \dots, k\}$

$$\partial_m \bar{V}_m(x, F) = \begin{cases} \bar{q}_m(F), & \text{if } V_m(x, y) = x_m q_m(y) - p_m(y) \\ f(x_m), & \text{if } V_m(x, y) = \mathbf{1}\{y \leq x_m\} - \alpha_m \\ (2 - 4\tau_m)F(x_m) + 2\tau_m, & \text{if } V_m(x, y) = 2|\mathbf{1}\{y \leq x_m\} - \tau_m|(x_m - y), \end{cases}$$

where $p_m, q_m: \mathcal{O} \rightarrow \mathbb{R}$ are some \mathcal{F} -integrable functions such that $\bar{q}_m(F) \neq 0$ for all $F \in \mathcal{F}$ and $\alpha_m, \tau_m \in (0, 1)$. We see that (V5) is satisfied if e.g. T is a vector of ratios of expectations with *the same denominator* (compare the situation in Frongillo and Kash (2014)). In this situation, we have that $c_F = \bar{q}(F)$. On the other hand, if the components of T are quantiles, expectiles with $\tau_m \neq 1/2$ or ratios of expectations with *different denominators* and additionally the class \mathcal{F} is rich enough, then (V4) might be satisfied.

Proposition 4.1. *Let $T_m: \mathcal{F} \rightarrow \mathbf{A}_m \subseteq \mathbb{R}$ be 1-elicitable and 1-identifiable functionals with oriented strict \mathcal{F} -identification functions $V_m: \mathbf{A}_m \times \mathcal{O} \rightarrow \mathbb{R}$ for $m \in \{1, \dots, k\}$. Let $\mathbf{A} := T(\mathcal{F}) \subseteq \mathbf{A}_1 \times \dots \times \mathbf{A}_k$. Then $V: \mathbf{A} \times \mathcal{O} \rightarrow \mathbb{R}^k$ defined as*

$$V(x_1, \dots, x_k, y) = (V_1(x_1, y), \dots, V_k(x_k, y))^\top \quad (4.1)$$

is an oriented strict \mathcal{F} -identification function for T .

Let $S: \mathbf{A} \times \mathcal{O} \rightarrow \mathbb{R}$ be a strictly \mathcal{F} -consistent scoring function for T . Suppose that assumptions (V1), (V3) and (S2) hold, and let $h: \text{int}(\mathbf{A}) \rightarrow \mathbb{R}^{k \times k}$ be the function given at (3.2). Define $\mathbf{A}'_m := \{x_m: (x_1, \dots, x_k) \in \text{int}(\mathbf{A})\}$.

(i) *If assumption (V4) holds and \mathbf{A} is connected then there are functions $g_m: \mathbf{A}'_m \rightarrow \mathbb{R}$, $m \in \{1, \dots, k\}$, $g_m > 0$, such that*

$$h_{mm}(x_1, \dots, x_k) = g_m(x_m)$$

for all $m \in \{1, \dots, k\}$ and $(x_1, \dots, x_k) \in \text{int}(\mathbf{A})$ and

$$h_{rl}(x) = 0 \quad (4.2)$$

for all $r, l \in \{1, \dots, k\}$, $l \neq r$, and for all $x \in \text{int}(\mathbf{A})$.

(ii) *If assumption (V5) holds then*

$$\partial_l h_{rm}(x) = \partial_r h_{lm}(x), \quad h_{rl}(x) = h_{lr}(x) \quad (4.3)$$

for all $r, l, m \in \{1, \dots, k\}$, $l \neq r$, where the first identity holds for almost all $x \in \text{int}(\mathbf{A})$ and the second identity for all $x \in \text{int}(\mathbf{A})$. Moreover, the matrix $(h_{rl}(x))_{l,r=1,\dots,k}$ is positive definite for all $x \in \text{int}(\mathbf{A})$.

A direct consequence of Proposition 4.1 (i) and Proposition 3.5 is the following characterization of the class of strictly \mathcal{F} -consistent scoring functions for functionals with elicitable components satisfying assumption (V4). In particular, it gives a characterization of the class of strictly \mathcal{F} -consistent scoring functions for a vector of different quantiles and/or different expectiles (with the exception of the 1/2-expectile), thus answering a question raised in Gneiting and Raftery (2007, p. 370).

Corollary 4.2. *Suppose that $T = (T_1, \dots, T_k): \mathcal{F} \rightarrow \mathbf{A}$ is a functional with 1-identifiable components having oriented strict \mathcal{F} -identification functions. Assume that the interior of $\mathbf{A} := T(\mathcal{F}) \subseteq \mathbf{A}_1 \times \dots \times \mathbf{A}_k$ is a star domain and that assumptions (V1), (V3), (S2), (F1) and (VS1) hold for T . If assumption (V4) holds, then a scoring function $S: \mathbf{A} \times \mathbf{O} \rightarrow \mathbb{R}$ is strictly \mathcal{F} -consistent for T if and only if it is of the form*

$$S(x_1, \dots, x_k, y) = \sum_{m=1}^k S_m(x_m, y), \quad (4.4)$$

for almost all $(x, y) \in \mathbf{A} \times \mathbf{O}$, where $S_m: \mathbf{A}_m \times \mathbf{O} \rightarrow \mathbb{R}$, $m \in \{1, \dots, k\}$, are strictly \mathcal{F} -consistent scoring functions for T_m .

If we are in the situation of Proposition 4.1 (ii), that is, T satisfies assumption (V5), it is well-known that a statement analogous to Corollary 4.2 is false. Let $F \in \mathcal{F}$ and $t = T(F)$. Recalling the orientation of the components V_m , we can immediately deduce that there is $c_F > 0$ such that $\bar{V}(t + sv, F) = c_F sv$ for $s \in \mathbb{R}$ and $v \in \mathbb{S}^{k-1}$. Hence, one obtains

$$v^\top h(t + sv) \bar{V}(t + sv, F) = c_F sv^\top h(t + sv)v.$$

Consequently, if \mathbf{A} is open and convex, the positive definiteness of $h(x)$ for all $x \in \mathbf{A}$ is a sufficient condition for the strict \mathcal{F} -consistency of S for T by Lemma 2.9 (i). Moreover, we now assume that T is a ratio of expectations with the same denominator $q: \mathbf{O} \rightarrow \mathbb{R}$ implying that $c_F = \bar{q}(F)$ for all $F \in \mathcal{F}$. Using Proposition 3.5, we obtain that for almost all $(x, y) \in \mathbf{A} \times \mathbf{O}$ strictly \mathcal{F} -consistent scoring functions for T are of the form

$$S(x, y) = -\phi(x)q(y) + \sum_{m=1}^k V_m(x, y)\partial_m\phi(x) + a(y), \quad (4.5)$$

with

$$\phi(x) = \sum_{r=1}^k \int_{z_r}^{x_r} \int_{z_r}^v h_{rr}(x_1, \dots, x_{r-1}, w, z_{r+1}, \dots, z_k) dw dv,$$

where $(z_1, \dots, z_k) \in \mathbf{A}$ and $a: \mathbf{O} \rightarrow \mathbb{R}$ is some \mathcal{F} -integrable function. Using (4.3), it follows that the function ϕ has Hessian h . Therefore, for \mathbf{A} open and convex, ϕ is strictly convex. Hence we have shown the following corollary.

Corollary 4.3. *Let $T = (T_1, \dots, T_k): \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^k$ be a surjective functional with 1-identifiable components with oriented strict identification functions $V_m: \mathbf{A}_m \times \mathbf{O} \rightarrow \mathbb{R}$, $m \in \{1, \dots, k\}$, that fulfills assumption (V5). Suppose that $\mathbf{A} \subseteq \mathbf{A}_1 \times \dots \times \mathbf{A}_k$ is open and convex and that assumptions (V1), (V3), (S2), (F2) and (VS1) hold. Then, a scoring function S is strictly \mathcal{F} -consistent for T if and only if it is of the form (4.5) for almost all $(x, y) \in \mathbf{A} \times \mathbf{O}$ with a twice continuously differentiable strictly convex function $\phi: \mathbf{A} \rightarrow \mathbb{R}$ and \mathcal{F} -integrable functions $q, a: \mathbf{O} \rightarrow \mathbb{R}$, where q is such that $\bar{q}(F) > 0$ for all $F \in \mathcal{F}$.*

This corollary recovers results of Osband and Reichelstein (1985); Banerjee et al. (2005); Abernethy and Frongillo (2012) if T is linear (meaning $q \equiv 1$), which show that all consistent scoring functions for linear functionals are so-called *Bregman functions*, that is,

functions of the form (4.5) with $q \equiv 1$ and a convex function ϕ . Frongillo and Kash (2014, Theorem 5) also treat the case of more general functions q . Comparing these results with Corollary 4.3, on the one hand, they are stronger as they require weaker smoothness assumptions on the scoring function, but on the other hand, they are weaker since they assume that \mathcal{F} contains all one-point distributions δ_y .

Remark 4.4. One might wonder about necessary conditions on the matrix-valued function h in the flavor of Proposition 4.1 if the k components of the functional T can be regrouped into (i) a new functional $T'_1: \mathcal{F} \rightarrow \mathbf{A}'_1 \subset \mathbb{R}^{k'_1}$ with an oriented strict \mathcal{F} -identification function $V'_1: \mathbf{A}'_1 \times \mathcal{O} \rightarrow \mathbb{R}^{k'_1}$ which satisfies assumption (V4), and (ii) several, say l , new functionals $T'_m: \mathcal{F} \rightarrow \mathbf{A}'_{k'_m} \subseteq \mathbb{R}^{k'_m}$, $m \in \{2, \dots, l+1\}$ with oriented strict \mathcal{F} -identification functions $V'_m: \mathbf{A}'_{k'_m} \times \mathcal{O} \rightarrow \mathbb{R}^{k'_m}$ such that each one satisfies assumption (V5), and $k'_1 + \dots + k'_{l+1} = k$. We can apply Proposition 4.1 to obtain necessary conditions for each of the $(k'_m \times k'_m)$ -valued functions h'_m , $m \in \{1, \dots, l+1\}$. Applying Lemma 2.15 we get a possible choice for a strictly \mathcal{F} -consistent scoring function S for T . On the level of the $k \times k$ -valued function h associated to S this means that h is a block diagonal matrix of the form $\text{diag}(h'_1, \dots, h'_{l+1})$. But what about the necessity of this form? Indeed, if we assume that the blocks in (ii) have maximal size (or equivalently that l is minimal) then one can verify that h must be necessarily of the block diagonal form described above.

While Remark 4.4 is consistent with Conjecture 1 of Frongillo and Kash (2014), the next section contains results refuting their conjecture; see Theorem 5.2 and Corollary 5.5.

5 Spectral risk measures

Risk measures are a common tool to measure the risk of a financial position Y . A risk measure is usually defined as a mapping ρ from some space of random variables, for example L^∞ , to the real line. Arguably, the most common risk measure in practice is *Value at Risk* at level α (VaR_α) which is the generalized α -quantile $F^{-1}(\alpha)$, that is,

$$\text{VaR}_\alpha(Y) := F^{-1}(\alpha) := \inf\{x \in \mathbb{R} : F(x) \geq \alpha\},$$

where F is the distribution function of Y . An important alternative to VaR_α is *Expected Shortfall* at level α (ES_α) (also known under the names *Conditional Value at Risk* or *Average Value at Risk*). It is defined as

$$\text{ES}_\alpha(Y) := \frac{1}{\alpha} \int_0^\alpha \text{VaR}_u(Y) \, du, \quad \alpha \in (0, 1], \quad (5.1)$$

and $\text{ES}_0(Y) = \text{ess inf } Y$. Since the influential paper of Artzner et al. (1999) introducing coherent risk measures, there has been a lively debate about which risk measure is best in practice, one of the requirements under discussion being the coherence of a risk measure. We call a functional ρ *coherent* if it is monotone, meaning that $Y \leq X$ a.s. implies that $\rho(Y) \leq \rho(X)$; it is superadditive in the sense that $\rho(X + Y) \geq \rho(X) + \rho(Y)$; it is positively homogeneous which means that $\rho(\lambda Y) = \lambda \rho(Y)$ for all $\lambda \geq 0$; and it is translation invariant which amounts to $\rho(Y + a) = \rho(Y) + a$ for all $a \in \mathbb{R}$. In the literature

on risk measures there are different sign conventions which co-exist. In this paper, a positive value of Y denotes a profit. Moreover, the position Y is considered the more risky the smaller $\rho(Y)$ is. Strictly speaking, we have chosen to work with utility functions instead of risk measures as for example in Delbaen (2012). The risk measure ρ is called *comonotonically additive* if $\rho(X + Y) = \rho(X) + \rho(Y)$ for comonotone random variables X and Y . Coherent and comonotonically additive risk measures are also called *spectral* risk measures (Acerbi, 2002). All risk measures of practical interest are law-invariant, that is, if two random variable X and Y have the same law F , then $\rho(X) = \rho(Y)$. As we are only concerned with law-invariant risk measures in this paper, we will abuse notation and write $\rho(F) := \rho(X)$, if X has distribution F .

One of the main criticisms on VaR_α is its failure to fulfill the superadditivity property in general (Acerbi, 2002). Furthermore, it fails to take the size of losses beyond the level α into account (Daníelsson et al., 2001). In both of these aspects, ES_α is a better alternative as it is a coherent and comonotonically additive, that is, a spectral risk measure. However, with respect to robustness, some authors argue that VaR_α should be preferred over ES_α (Cont et al., 2010; Kou et al., 2013), whereas others argue that the classical statistical notions of robustness are not necessarily appropriate in a risk measurement context (Krättschmer et al., 2012, 2013, 2014). Finally, ES_α fails to be 1-elicitable (Weber, 2006; Gneiting, 2011), whereas VaR_α is 1-elicitable for most classes of distributions \mathcal{F} of practical relevance. In fact, except for the expectation, all spectral risk measures fail to be 1-elicitable (Ziegel, 2015); further recent results on elicitable risk measures include (Kou and Peng, 2014; Wang and Ziegel, 2015) showing that distortion risk measures are rarely elicitable and (Weber, 2006; Bellini and Bigozzi, 2014; Delbaen et al., 2014) demonstrating that convex risk measures are only elicitable if they are shortfall risk measures.

We show in Theorem 5.2 (see also Corollary 5.4 and 5.5) that spectral risk measures having a spectral measure with finite support can be a component of a k -elicitable functional. In particular, the pair $(\text{VaR}_\alpha, \text{ES}_\alpha): \mathcal{F} \rightarrow \mathbb{R}^2$ is 2-elicitable for any $\alpha \in (0, 1)$ subject to mild conditions on the class \mathcal{F} . We remark that our results substantially generalize the result of Acerbi and Székely (2014) as detailed below.

Assumption (F2). \mathcal{F} is a class of distribution functions F on \mathbb{R} having a continuous density $f = F'$. Assume that the distributions F in \mathcal{F} have unique quantiles in the sense that the equation $F(q) = \alpha$ has a unique solution for all $\alpha \in (0, 1)$.

Definition 5.1 (Spectral risk measures). Let μ be a probability measure on $[0, 1]$ (called *spectral measure*) and let \mathcal{F} be a class of distribution functions on \mathbb{R} with finite first moments. Then, the *spectral risk measure* associated to μ is the functional $\nu_\mu: \mathcal{F} \rightarrow \mathbb{R}$ defined as

$$\nu_\mu(F) := \int_{[0,1]} \text{ES}_\alpha(F) \mu(d\alpha).$$

Kusuoka (2001); Jouini et al. (2006) have shown that law-invariant coherent and comonotonically additive risk measures are exactly the spectral risk measures in the sense of Definition 5.1 for distributions with compact support. If $\mu = \delta_\alpha$ for some $\alpha \in [0, 1]$, then $\nu_\mu(F) = \text{ES}_\alpha(F)$. In particular, $\mu_{\delta_1}(F) = \int y dF(y)$ is the expectation of F .

In the following theorem, we show that spectral risk measures whose spectral measure μ has finite support in $(0, 1)$ are k -elicitable for some k . It is possible to extend the result to spectral measures with finite support in $(0, 1]$; see Corollary 5.4. If μ has mass at zero, we believe that ν_μ is not k -elicitable for any k with respect to interesting classes \mathcal{F} . In this case, if the support of F is unbounded below, we have $\nu_\mu(F) = \text{ess inf}(F) = -\infty$.

Theorem 5.2. *Let \mathcal{F} be a class of distribution functions on \mathbb{R} with finite first moments satisfying assumption (F2). Let $\nu_\mu: \mathcal{F} \rightarrow \mathbb{R}$ be a spectral risk measure where μ is given by*

$$\mu = \sum_{m=1}^{k-1} p_m \delta_{q_m},$$

with $p_m \in (0, 1]$, $\sum_{m=1}^{k-1} p_m = 1$, $q_m \in (0, 1)$ and the q_m 's are pairwise distinct. Define the functional

$$T = (T_1, \dots, T_k): \mathcal{F} \rightarrow \mathbb{R}^k,$$

where $T_m(F) := F^{-1}(q_m)$, $m \in \{1, \dots, k-1\}$, and $T_k(F) := \nu_\mu(F)$. Then the following assertions are true:

- (i) *The functional T is k -elicitable with respect to \mathcal{F} .*
- (ii) *Let assumption (F1) be satisfied and let the interior of $\mathbf{A} := T(\mathcal{F}) \subseteq \mathbb{R}^k$ be a star domain. Define the function $V: \mathbf{A} \times \mathbb{R} \rightarrow \mathbb{R}^k$ with components*

$$\begin{aligned} V_m(x_1, \dots, x_k, y) &= \mathbb{1}\{y \leq x_m\} - q_m, \quad m \in \{1, \dots, k-1\}, \\ V_k(x_1, \dots, x_k, y) &= x_k - \sum_{m=1}^{k-1} \frac{p_m}{q_m} y \mathbb{1}\{y \leq x_m\}. \end{aligned} \quad (5.2)$$

Then V is a strict \mathcal{F} -identification for T satisfying assumption (V3).

Assume that (V1) and (F1) are satisfied and that (V_1, \dots, V_{k-1}) satisfies (V4). Then every strictly \mathcal{F} -consistent scoring function $S: \mathbf{A} \times \mathbb{R} \rightarrow \mathbb{R}$ for T satisfying (S2) and (VS1) is necessarily of the form

$$\begin{aligned} S(x_1, \dots, x_k, y) &= \sum_{r=1}^{k-1} \int_{z_r}^{x_r} \left(\frac{p_r}{q_r} G_k(z_k) + g_r(v) \right) (\mathbb{1}\{y \leq v\} - q_r) \, dv \\ &\quad + (G_k(x_k) - G_k(z_k)) \sum_{m=1}^{k-1} \frac{p_m}{q_m} (x_m (\mathbb{1}\{y \leq x_m\} - q_m) - y \mathbb{1}\{y \leq x_m\}) \\ &\quad + G_k(x_k) x_k - \mathcal{G}_k(x_k) + a(y), \end{aligned} \quad (5.3)$$

for almost all (x, y) , where $(z_1, \dots, z_k) \in \text{int}(\mathbf{A})$ is some star point in $\text{int}(\mathbf{A})$, $g_m: \mathbf{A}'_m \rightarrow \mathbb{R}$, $\mathbf{A}'_m := \{x_m: (x_1, \dots, x_k) \in \text{int}(\mathbf{A})\}$, $m \in \{1, \dots, k\}$, are locally Lipschitz continuous functions and $\mathcal{G}_k, G_k: \mathbf{A}'_k \rightarrow \mathbb{R}$ are such that $G'_k = g_k$, $\mathcal{G}'_k = G_k$ with

$$g_k(x_k) > 0, \quad \frac{p_m}{q_m} G_k(x_k) + g_m(x_m) > 0 \quad (5.4)$$

for all $m \in \{1, \dots, k-1\}$, $x = (x_1, \dots, x_k) \in \text{int}(\mathbf{A})$ and $a: \mathbb{R} \rightarrow \mathbb{R}$ is some \mathcal{F} -integrable function.

(iii) If $A \supseteq T(\mathcal{F})$ is open and convex and if a scoring function $S: A \times \mathbb{R} \rightarrow \mathbb{R}$ is of the form given at (5.3) where the functions g_m , $m \in \{1, \dots, k\}$, are continuous, $G'_k = g_k$, $\mathcal{G}'_k = G_k$ and for all $x = (x_1, \dots, x_k) \in A$, $m \in \{1, \dots, k-1\}$ we have (5.4) and additionally

$$G_k(x_k) > 0, \quad (5.5)$$

then S is strictly \mathcal{F} -consistent for T .

Remark 5.3. In part (iii) of Theorem 5.2, it is possible and often convenient to choose z on the boundary of A . This works if the chosen functions g_m , $m \in \{1, \dots, k\}$, G_k have continuous extensions to the closure of A . One can also choose $z_r = -\infty$ for some $r \in \{1, \dots, k\}$ if A is large enough. For $r = k$ one has to ensure that $G_k(z_k) = \lim_{x \rightarrow -\infty} G_k(x)$ is finite, and for $r < k$ we need that the functions $g_r(\cdot) + (p_r/q_r)G_k(z_k)$ are integrable over intervals of the form $(-\infty, x]$.

Using Theorem 5.2 and the revelation principle (Proposition 2.13) we can now state one of the main results of this paper.

Corollary 5.4. *Let \mathcal{F} be a class of distribution functions on \mathbb{R} with finite first moments satisfying assumption (F2). Let $\nu_\mu: \mathcal{F} \rightarrow \mathbb{R}$ be a spectral risk measure. If the support of μ is finite and contained in $(0, 1]$, then ν_μ is a component of a k -elicitable functional where*

- (i) $k = 1$, if μ is concentrated at 1 meaning $\mu(\{1\}) = 1$;
- (ii) $k = 1 + L$, if $\mu(\{1\}) < 1$, where L is the cardinality of the support of μ .

Theorem 5.2 (iii) immediately gives a large class of strictly consistent scoring functions for the pair $(\text{VaR}_\alpha, \text{ES}_\alpha)$.

Corollary 5.5. *Let $\alpha \in (0, 1)$. Let \mathcal{F} be a class of distribution functions on \mathbb{R} with finite first moments. If \mathcal{F} satisfies assumption (F2), then the pair $T = (\text{VaR}_\alpha, \text{ES}_\alpha): \mathcal{F} \rightarrow A$ with $A := \mathbb{R}^2$ is 2-elicitable. Strictly \mathcal{F} -consistent scoring functions for T are given by*

$$\begin{aligned} S(x_1, x_2, y) &= (\mathbb{1}\{y \leq x_1\} - \alpha)(G_1(x_1) - G_1(y)) \\ &\quad + \frac{1}{\alpha}G_2(x_2)\mathbb{1}\{y \leq x_1\}(x_1 - y) + G_2(x_2)(x_2 - x_1) - \mathcal{G}_2(x_2), \end{aligned}$$

where G_1 and G_2 are strictly increasing continuously differentiable functions, G_1 is \mathcal{F} -integrable and $\lim_{x \rightarrow -\infty} G_2(x) = 0$, $\mathcal{G}'_2 = G_2$.

Remark 5.6. Peering into the proof of Theorem 5.2 part (iii), one can see that it is possible to weaken assumption (F2) in Corollary 5.5. We must require that all $F \in \mathcal{F}$ have finite first moments, their α -quantiles must be unique, and they must be continuously differentiable at their α -quantiles. This remark also applies to Corollary 5.8 below.

Remark 5.7. According to Corollary 5.5 (and Theorem 5.2) the functional $(\text{VaR}_\alpha, \text{ES}_\alpha)$ admits non-separable strictly consistent scoring functions. Hence, this refutes both Osband (1985, Proposition 2.3) and Frongillo and Kash (2014, Conjecture 1). Proposition 2.3 in Osband (1985) states that any strictly consistent scoring function for a functional with a

quantile as a component must be separable in the sense that it must be the sum of a strictly consistent scoring function for the quantile and a strictly consistent scoring function for the rest of the functional. However, there is a flaw in the proof of the proposition. Conjecture 1 in Frongillo and Kash (2014) asserts that functionals admitting non-separable strictly consistent scoring functions must necessarily be ratios of expectations (with the same denominator).

Acerbi and Szekely (2014) also give an example of a scoring function for the pair $T = (\text{VaR}_\alpha, \text{ES}_\alpha): \mathcal{F} \rightarrow \mathbf{A} \subseteq \mathbb{R}^2$. They use a different sign convention for VaR_α and ES_α than we do in this paper. Using our sign convention, their proposed scoring function $S^W: \mathbf{A} \times \mathbb{R} \rightarrow \mathbb{R}$ reads

$$S^W(x_1, x_2, y) = \alpha(x_2^2/2 + Wx_1^2/2 - x_1x_2) + \mathbf{1}\{y \leq x_1\}(-x_2(y - x_1) + W(y^2 - x_1^2)/2), \quad (5.6)$$

where $W \in \mathbb{R}$. The authors claim that S^W is a strictly \mathcal{F} -consistent scoring function for $T = (\text{VaR}_\alpha, \text{ES}_\alpha)$ provided that

$$\text{ES}_\alpha(F) > W \text{VaR}_\alpha(F) \quad (5.7)$$

for all $F \in \mathcal{F}$. They assume that \mathcal{F} satisfies assumption (F2) and that $F(x) \in (0, 1)$ implies $f(x) > 0$ for all $F \in \mathcal{F}$ with density f . Furthermore, in order to ensure that $\bar{S}^W(\cdot, F)$ is finite one needs to impose the assumption that $\int_{-\infty}^x y^2 dF(y)$ is finite for all $x \in \mathbb{R}$ and $F \in \mathcal{F}$. This is slightly less than requiring finite second moments. As a matter of fact, they only show that $\nabla \bar{S}^W(t_1, t_2, F) = 0$ for $F \in \mathcal{F}$ and $(t_1, t_2) = T(F)$ and that $\nabla^2 \bar{S}^W(t_1, t_2, F)$ is positive definite. This only shows that $\bar{S}^W(x, F)$ has a *local* minimum at $x = T(F)$ but does not provide a proof concerning a *global* minimum; see also the discussion after Corollary 3.4. However, we can use Theorem 5.2 to verify their claims at least partially. A straightforward calculation shows that S^W given at (5.6) is of the form given at (5.3) with $z_1 = z_2 = 0$, $a(y) = \mathbf{1}\{y \leq 0\} y^2/2$ and

$$g_1(x_1) = -Wx_1, \quad G_2(x_2) = \alpha x_2, \quad g_2(x_2) = G_2'(x_2) = \alpha.$$

Note that (5.4) is satisfied for x_1, x_2 with $Wx_1 < x_2$. However, (5.5) is only satisfied for $x_2 > 0$. Therefore, we can only guarantee that S^W is a strictly \mathcal{F}' -consistent scoring function for T if we restrict S^W to $\mathbf{A}' \times \mathbb{R}$ where

$$\mathbf{A}' := \{(x_1, x_2) \in \mathbb{R}^2: Wx_1 < x_2, x_2 > 0\},$$

and $\mathcal{F}' \subseteq \mathcal{F}$ is such that $T(\mathcal{F}') \subseteq \mathbf{A}'$ is satisfied. This is unfortunate, as our sign convention described in the beginning of this section has the interpretation that risky assets correspond negative values of $\text{VaR}_\alpha, \text{ES}_\alpha$. Therefore, the interesting action domain for risk management applications would be (a subset) of $\{(x_1, x_2) \in \mathbb{R}^2: x_1 < 0, x_2 < 0\}$.

While we believe that it may be possible to construct a distribution F with the property $W \text{VaR}_\alpha(F) < \text{ES}_\alpha(F) < 0$ such that $\bar{S}^W(\cdot, \cdot, F)$ does not have a global minimum at $(\text{VaR}_\alpha(F), \text{ES}_\alpha(F))$, Acerbi and Szekely (2014) have tested their score on a variety of distributions that are relevant in risk management contexts. In these numerical examples the graphs of the expected scores appear to be convex, so there is hope that S^W is

strictly \mathcal{F}' -consistent for some smaller class of distributions \mathcal{F}' that is still large enough to accommodate all practically relevant distributions.

The scoring function S^W has two properties which are interesting and potentially relevant in applications. If x_1, x_2 and y are expressed in the same units of measurement, then $S^W(x_1, x_2, y)$ is a quantity with these units squared. If one insists that we should only add quantities with the same units, then the necessary condition (5.4) enforces a condition of the type (5.7). We are currently not able to provide a scoring function respecting units in this sense, which can be shown to be strictly consistent for $(\text{VaR}_\alpha, \text{ES}_\alpha)$ under condition (5.4), even if we restrict the action domain to a subset of $\{(x_1, x_2) \in \mathbb{R}^2: x_1 < 0, x_2 < 0\}$.

The second property of S^W that may be interesting for risk management applications is the fact that the expected score $\bar{S}^W(x_1, x_2, F)$ only depends on the distribution of the losses, that is, on the distribution F on $(-\infty, x_1]$. Restricting the action domain accordingly, it is possible to provide a rich class of strictly consistent scoring functions with the same property using Theorem 5.2 part (iii).

Corollary 5.8. *Let $\alpha \in (0, 1)$. Let \mathcal{F} be a class of distribution functions on \mathbb{R} with finite first moments such that $\text{VaR}_\alpha(F) < 0$ and $\text{ES}_\alpha(F) < 0$ for all $F \in \mathcal{F}$. If \mathcal{F} satisfies assumption (F2), then the pair $T = (\text{VaR}_\alpha, \text{ES}_\alpha): \mathcal{F} \rightarrow \mathbf{A}$ with $\mathbf{A} := (-\infty, 0) \times (-\infty, 0)$ is 2-elicitable. Strictly \mathcal{F} -consistent scoring functions for T are given by*

$$S(x_1, x_2, y) = -(\alpha G_1(x_1) + x_1 G_2(x_2) + \mathcal{G}_2(x_2)) \\ + \frac{1}{\alpha} \mathbf{1}\{y \leq x_1\}(\alpha G_1(x_1) - y G_1(y) + G_2(x_2)(x_1 - y)),$$

where G_1 and G_2 are strictly increasing continuously differentiable functions on $(-\infty, 0]$, G_1 is \mathcal{F} -integrable and $G_2 > 0$, $\mathcal{G}'_2 = G_2$.

As it is currently unclear, whether S^W given at (5.6) remains strictly consistent for negative values of x_2 , we recommend to use one of the scoring functions given in Corollaries 5.5 or 5.8 for evaluating forecasts for $(\text{VaR}_\alpha, \text{ES}_\alpha)$. Possible choices arise by choosing $G_1(x) = x$ and $G_2(x) = e^x$. We remark that for the scoring functions in Corollaries 5.5 and 5.8 it is also not necessary to impose any conditions such as (5.7), which may be impossible to check in practice.

6 Discussion

We have investigated necessary and sufficient conditions for the elicibility of k -dimensional functionals of d -dimensional distributions. In order to derive necessary conditions we have adapted Osband's principle for the case where the class \mathcal{F} of distributions does not necessarily contain distributions with finite support. This comes at the cost of certain smoothness assumptions on the expected scores $\bar{S}(\cdot, F)$. For particular situations, e.g. when characterizing the class of strictly \mathcal{F} -consistent scoring functions for ratios of expectations, it is possible to weaken the smoothness assumptions; see Frongillo and Kash (2014). However, Frongillo and Kash (2014) assume that the class \mathcal{F} of distributions contains all distributions with finite support, which is not necessary for the validity of our

result. While this is not a great gain in the case of linear functionals or ratios of expectations it comes in handy when considering spectral risk measures. Value at Risk, VaR_α , being defined as the smallest α -quantile, is generally not elicitable for distributions where the α -quantile is not unique. Therefore, we believe that it is also not possible to show joint elicibility of $(\text{VaR}_\alpha, \text{ES}_\alpha)$ for classes \mathcal{F} of distributions with non-unique α -quantiles. Fortunately, the classes \mathcal{F} of distributions that are relevant in risk management usually consist of absolutely continuous distributions.

Emmer et al. (2013) have remarked that ES_α is *conditionally elicitable*. One can slightly generalize their definition of conditional elicibility as follows.

Definition 6.1. Fix an integer $k \geq 1$. A functional $T_k: \mathcal{F} \rightarrow \mathbf{A}_k \subseteq \mathbb{R}$ is called *conditionally elicitable of order k* if there are $k - 1$ elicitable functionals $T_m: \mathcal{F} \rightarrow \mathbf{A}_m \subseteq \mathbb{R}$, $m \in \{1, \dots, k - 1\}$, such that T_k is elicitable restricted to the class

$$\mathcal{F}_{x_1, \dots, x_{k-1}} := \{F \in \mathcal{F} : T_1(F) = x_1, \dots, T_{k-1}(F) = x_{k-1}\}$$

for any $(x_1, \dots, x_{k-1}) \in \mathbf{A}_1 \times \dots \times \mathbf{A}_{k-1}$.

Mutatis mutandis, one can define a notion of *conditional identifiability* by replacing the term ‘elicitable’ with ‘identifiable’ in the above definition. It is not difficult to check that any conditionally identifiable functional T_k of order k is a component of an identifiable functional $T = (T_1, \dots, T_k)$. Spectral risk measures ν_μ with spectral measure μ with finite support in $(0, 1)$ provide an example of a conditionally elicitable functional of order $L + 1$, where L is the cardinality of the support of μ ; see Theorem 5.2. However, we would like to stress that it is generally an open question whether any conditionally elicitable and identifiable functional T_k of order $k \geq 2$ is always a component of a k -elicitable functional.

Slightly modifying Lambert et al. (2008, Definition 11), one could define the *elicibility order* of a real-valued functional T as the smallest number k such that the functional is a component of a k -elicitable functional. It is clear that the elicibility order of the variance is two, and we have shown that the same is true for ES_α for reasonably large classes \mathcal{F} . For spectral risk measures ν_μ , the elicibility order is at most $L + 1$, where L is the cardinality of the support; see Theorem 5.2.

In the one-dimensional case, Steinwart et al. (2014) have shown that having convex level sets in the sense of Proposition 2.14 is a sufficient condition for elicibility of a functional T under continuity assumptions on T . Without such continuity assumptions, the converse of Proposition 2.14 is generally false; see Heinrich (2013) for the example of the mode functional. It is an open (and potentially difficult) question under which conditions a converse of Proposition 2.14 is true for higher order elicibility.

7 Proofs

Proof of Lemma 2.9

The first part is a direct consequence of the definition of strict \mathcal{F} -consistency. For the second part, using part (i), we must prove that any function

$$\psi: D \rightarrow \mathbb{R}, \quad s \mapsto \bar{S}(t + sv, F)$$

has a (unique) global minimum at $s = 0$ for all $F \in \mathcal{F}'$, $t = T(F) \in \text{int}(A)$ and $v \in \mathbb{S}^{k-1}$ where $D = \{s \in \mathbb{R}: t + sv \in \text{int}(A)\}$. This can be easily verified by observing that for $s \in D$ we have

$$\psi'(s) = v^\top \nabla \bar{S}(t + sv, F) \begin{cases} > 0, & \text{if } s > 0 \\ = 0, & \text{if } s = 0 \\ < 0, & \text{if } s < 0 \end{cases}$$

where we used the fact that ∇S is an oriented strict \mathcal{F}' -identification function for $T|_{\mathcal{F}'}$. \square

Proof of Theorem 3.2

Let $x \in \text{int}(A)$. The identifiability property of V plus the first order condition stemming from the strict \mathcal{F} -consistency of S yields the relation

$$\ker(\bar{V}(x, \cdot)) \subseteq \ker(\nabla \bar{S}(x, \cdot)),$$

where we consider $\bar{V}(x, \cdot)$ and $\nabla \bar{S}(x, \cdot)$ as linear functionals $\hat{\mathcal{F}} \rightarrow \mathbb{R}^k$. Let $l \in \{1, \dots, k\}$. To show (3.2), consider the composed linear functional

$$\bar{B}(x, \cdot): \hat{\mathcal{F}} \rightarrow \mathbb{R}^{k+1}, \quad F \mapsto (\partial_l \bar{S}(x, F), \bar{V}(x, F)).$$

By construction, we know that

$$\ker(\bar{V}(x, \cdot)) = \ker(\bar{B}(x, \cdot)). \tag{7.1}$$

Assumption (V1) implies that there are $F_1, \dots, F_{k+1} \in \hat{\mathcal{F}}$ such that the matrix

$$\mathbb{V} = \text{mat}(\bar{V}(x, F_1), \dots, \bar{V}(x, F_{k+1})) \in \mathbb{R}^{k \times (k+1)}$$

has maximal rank, meaning $\text{rank}(\mathbb{V}) = k$.² Let $G \in \hat{\mathcal{F}}$. Then still

$$0 \in \text{int}(\text{conv}(\{\bar{V}(x, G), \bar{V}(x, F_1), \dots, \bar{V}(x, F_{k+1})\})),$$

such that $\text{rank}(\mathbb{V}_G) = k$ where

$$\mathbb{V}_G = \text{mat}(\bar{V}(x, G), \bar{V}(x, F_1), \dots, \bar{V}(x, F_{k+1})) \in \mathbb{R}^{k \times (k+2)}.$$

²If $\text{rank}(\mathbb{V}) < k$, then $\text{span}\{\bar{V}(x, F_1), \dots, \bar{V}(x, F_{k+1})\}$ would be a linear subspace such that $\text{int}(\text{conv}(\{\bar{V}(x, F_1), \dots, \bar{V}(x, F_{k+1})\})) = \emptyset$.

Define the matrix

$$\mathbb{B}_G = \begin{pmatrix} \partial_l \bar{S}(x, G) & \partial_l \bar{S}(x, F_1) & \cdots & \partial_l \bar{S}(x, F_{k+1}) \\ & \mathbb{V}_G & & \end{pmatrix} \in \mathbb{R}^{(k+1) \times (k+2)}.$$

Due to (7.1) we have that $\ker(\mathbb{V}_G) = \ker(\mathbb{B}_G)$. With the rank-nullity theorem, this gives $\text{rank}(\mathbb{B}_G) = \text{rank}(\mathbb{V}_G) = k$. Hence, there is a unique vector $(h_{l1}(x), \dots, h_{lk}(x)) \in \mathbb{R}^k$ such that

$$\partial_l \bar{S}(x, G) = \sum_{m=1}^k h_{lm}(x) \bar{V}_m(x, G).$$

Since $G \in \hat{\mathcal{F}}$ was arbitrary, the assertion at (3.2) follows.

The second part of the claim can be seen as follows. For $x \in \text{int}(\mathbf{A})$ pick $F_1, \dots, F_k \in \hat{\mathcal{F}}$ such that $\bar{V}(x, F_1), \dots, \bar{V}(x, F_k)$ are linearly independent and let $\mathbb{V}(z)$ be the matrix with columns $\bar{V}(z, F_i)$, $i \in \{1, \dots, k\}$ for $z \in \text{int}(\mathbf{A})$. Due to assumption (V2) or (V3), $\mathbb{V}(z)$ has full rank in some neighborhood U of x . Let $r \in \{1, \dots, k\}$ and let e_r be the r th standard unit vector of \mathbb{R}^k . We define $\lambda(z) := \mathbb{V}(z)^{-1} e_r$ for $z \in U$. Taking the inverse of a matrix is a continuously differentiable operation, so it is in particular locally Lipschitz continuous. Therefore, the vector λ inherits the regularity properties of $\bar{V}(z, F_i)$, that is, under (V2) λ is continuous, and under (V3) λ is locally Lipschitz continuous. Therefore, these properties carry over to h because for $l \in \{1, \dots, k\}$, $z \in U$

$$h_{lr}(z) = \sum_{i=1}^k \lambda_i(z) \sum_{m=1}^k h_{lm}(z) \bar{V}_m(z, F_i) = \sum_{i=1}^k \lambda_i(z) \partial_l \bar{S}_m(z, F_i)$$

using the assumptions on S . □

Proof of Proposition 3.5

We denote the right-hand side of (3.4) minus $a(y)$ by $I(x, y)$. Using Fubini's theorem, we obtain

$$\begin{aligned} \bar{I}(x, F) &= \sum_{r=1}^k \sum_{m=1}^k \int_{z_r}^{x_r} h_{rm}(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k) \\ &\quad \times \bar{V}_m(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k, F) dv. \end{aligned}$$

We will show that for any $l \in \{1, \dots, k\}$, $F \in \mathcal{F}$ and almost all $x \in \text{int}(\mathbf{A})$

$$\partial_l \bar{I}(x, F) = \sum_{m=1}^k h_{lm}(x) \bar{V}_m(x, F)$$

implying the necessity of the representation $\bar{S}(x, F) = \bar{I}(x, F) + c(F)$ where $c(F)$ is a constant depending on F . We obtain,

$$\begin{aligned} \partial_l \bar{I}(x, F) &= \sum_{m=1}^k h_{lm}(x_1, \dots, x_l, z_{l+1}, \dots, z_k) \bar{V}_m(x_1, \dots, x_l, z_{l+1}, \dots, z_k, F) \\ &+ \sum_{r=l+1}^k \int_{z_r}^{x_r} \sum_{m=1}^k \partial_l h_{rm}(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k) \bar{V}_m(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k, F) \\ &\quad + h_{rm}(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k) \partial_l \bar{V}_m(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k, F) dv. \end{aligned}$$

Using (3.3) we obtain that

$$\begin{aligned} \partial_l \bar{I}(x, F) &= \sum_{m=1}^k h_{lm}(x_1, \dots, x_l, z_{l+1}, \dots, z_k) \bar{V}_m(x_1, \dots, x_l, z_{l+1}, \dots, z_k, F) \\ &+ \sum_{r=l+1}^k \int_{z_r}^{x_r} \sum_{m=1}^k \left(\partial_r h_{lm}(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k) \bar{V}_m(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k, F) \right. \\ &\quad \left. + h_{lm}(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k) \partial_r \bar{V}_m(x_1, \dots, x_{r-1}, v, z_{r+1}, \dots, z_k, F) \right) dv \\ &= \sum_{m=1}^k h_{lm}(x_1, \dots, x_l, z_{l+1}, \dots, z_k) \bar{V}_m(x_1, \dots, x_l, z_{l+1}, \dots, z_k, F) \\ &\quad + \sum_{r=l+1}^k \sum_{m=1}^k \left(h_{lm}(x_1, \dots, x_r, z_{r+1}, \dots, z_k) \bar{V}_m(x_1, \dots, x_r, z_{r+1}, \dots, z_k, F) \right. \\ &\quad \left. - h_{lm}(x_1, \dots, x_{r-1}, z_r, \dots, z_k) \bar{V}_m(x_1, \dots, x_{r-1}, z_r, \dots, z_k, F) \right) \\ &= \sum_{r=l}^k \sum_{m=1}^k h_{lm}(x_1, \dots, x_r, z_{r+1}, \dots, z_k) \bar{V}_m(x_1, \dots, x_r, z_{r+1}, \dots, z_k, F) \\ &\quad - \sum_{r=l}^{k-1} \sum_{m=1}^k h_{lm}(x_1, \dots, x_r, z_{r+1}, \dots, z_k) \bar{V}_m(x_1, \dots, x_r, z_{r+1}, \dots, z_k, F) \\ &= \sum_{m=1}^k h_{lm}(x_1, \dots, x_k) \bar{V}_m(x_1, \dots, x_k, F). \end{aligned}$$

For almost all $y \in \mathcal{O}$, the set $\{x \in \mathbb{R}^k \mid (x, y) \in C^c\} =: A_y$ has k -dimensional Lebesgue measure zero, where C^c is the complement of the set C defined in assumption (VS1). Let $y \in \mathcal{O}$ be such that A_y has measure zero. Then we obtain that for almost all x the sets $\{x_i \in \mathbb{R} \mid (x, y) \in A_y\} =: N_i$ have one-dimensional Lebesgue-measure zero for all $i \in \{1, \dots, k\}$. Therefore, the summands of $I(x, y)$ are continuous in y for almost all x .

Let $(F_n)_{n \in \mathbb{N}}$ be a sequence as in assumption (F1), that is, $(F_n)_{n \in \mathbb{N}}$ converges weakly to δ_y and the support of all F_n is contained in some compact set K . Let φ be a function on \mathcal{O} which is locally bounded and continuous at y . We claim that then $\int_{\mathcal{O}} \varphi(z) dF_n(z) \rightarrow \varphi(y)$. By Skorohod's theorem there are random variables ξ_n , $n \in \mathbb{N}$, and ξ on some probability

space with distributions F_n , $n \in \mathbb{N}$, and δ_y , respectively, such that $\xi_n \rightarrow y$ almost surely. By the continuous mapping theorem, we obtain that $\varphi(\xi_n) \rightarrow \varphi(y)$ almost surely and since $\varphi \mathbb{1}_K$ is bounded we can apply dominated convergence theorem to obtain $\mathbb{E}\varphi(\xi_n) = \mathbb{E}\varphi(\xi_n)\mathbb{1}_K(\xi_n) = \int_{\mathcal{O}} \varphi dF_n \rightarrow \varphi(y)$.

By this argument (recalling that $S(x, \cdot)$, $V(x, \cdot)$ are assumed to be locally bounded), if $S(x, \cdot)$ and $I(x, \cdot)$ are continuous at y , then $\bar{S}(x, F_n) - \bar{I}(x, F_n) \rightarrow S(x, y) - I(x, y)$. We have shown that $S(x, F_n) - \bar{I}(x, F_n)$ does not depend on x , hence the same is true for the limit. Therefore, we can define $a(y) = S(x, y) - I(x, y)$ for almost all y . The function a is \mathcal{F} -integrable, since S and I are \mathcal{F} -integrable. \square

Proof of Proposition 4.1

It is clear that V given at (4.1) is a strict \mathcal{F} -identification function for T . Also the orientation of V follows directly from its form and the orientation of its components. We have that $\partial_l \bar{V}_r(x, F) = 0$ for all $l, r \in \{1, \dots, k\}$, $l \neq r$, and $x \in \text{int}(\mathbf{A})$, $F \in \mathcal{F}$. Hence, (3.3) takes the form

$$\begin{aligned} \partial_l \partial_r \bar{S}(x, F) &= \sum_{m=1}^k \partial_l h_{rm}(x) \bar{V}_m(x, F) + h_{rl}(x) \partial_l \bar{V}_l(x, F) \\ &= \sum_{m=1}^k \partial_r h_{lm}(x) \bar{V}_m(x, F) + h_{lr}(x) \partial_r \bar{V}_r(x, F) = \partial_r \partial_l \bar{S}(x, F). \end{aligned} \tag{7.2}$$

Equation (7.2) evaluated at $x = t = T(F)$ yields

$$h_{rl}(t) \partial_l \bar{V}_l(t, F) = h_{lr}(t) \partial_r \bar{V}_r(t, F). \tag{7.3}$$

We distinguish the two cases of the proposition.

- (i) Assume that (V4) holds. Then (7.3) implies that $h_{rl}(t) = 0$ for $r \neq l$, hence we obtain (4.2) with the surjectivity of T .
- (ii) Assume that (V5) holds. This means that (7.3) implies that $h_{rl}(t) = h_{lr}(t)$, whence the second part of (4.3) is shown, again using the surjectivity of T .

In both cases, (7.2) is equivalent to

$$\sum_{m=1}^k (\partial_l h_{rm}(x) - \partial_r h_{lm}(x)) \bar{V}_m(x, F) = 0. \tag{7.4}$$

The equation (7.4) continues to hold for all $F \in \hat{\mathcal{F}}$. Hence, using assumption (V1) there are $F_1, \dots, F_k \in \hat{\mathcal{F}}$ such that $\bar{V}(x, F_1), \dots, \bar{V}(x, F_k)$ are linearly independent. This yields that $\partial_l h_{rm}(x) = \partial_r h_{lm}(x)$ for almost all $x \in \mathbf{A}$. Now, we obtain the claims as follows.

- (i) We can conclude that $\partial_l h_{rr}(x) = \partial_l h_{rl}(x) = 0$ for $r \neq l$ for almost all $x \in \mathbf{A}$. Consequently, invoking that \mathbf{A} is connected, the functions h_{mm} only depend on x_m

and we can write $h_{mm}(x) = g_m(x_m)$ for some function $g_m: \mathbb{A}'_m \rightarrow \mathbb{R}$. By Lemma 2.9 (i), for $v \in \mathbb{S}^{k-1}$, $t = T(F) \in \text{int}(\mathbf{A})$, the function $s \mapsto \bar{S}(t + sv, F)$ has a minimum at $s = 0$, hence

$$v^\top \nabla \bar{S}(t + sv, F) = \sum_{m=1}^k g_m(t_m + sv_m) \bar{V}_m(t_m + sv_m, F) v_m$$

vanishes for $s = 0$, is negative for $s < 0$ and positive for $s > 0$, where s is in some neighborhood of zero. Choosing v as the l th standard basis vector of \mathbb{R}^k we obtain that $g_l > 0$ exploiting the orientation of V_l and the surjectivity of T .

- (ii) For the assertion about the definiteness, observe that due to assumption (V5), we have for $v \in \mathbb{S}^{k-1}$, $t = T(F) \in \text{int}(\mathbf{A})$ that $\bar{V}_m(t + sv, F) = c_F s v_m$ where $c_F > 0$ due to assumption (V5) and the orientation of each component of V . Hence,

$$v^\top \nabla \bar{S}(t + sv, F) = c_F s v^\top h(t + sv)v,$$

which implies the claim using again the surjectivity of T . \square

Proof of Corollary 4.2

The sufficiency, meaning that S of the form given at (4.4) is strictly \mathcal{F} -consistent for T , is immediate; see the proof of Lemma 2.15. For the necessity, we choose an identification function $V(x, y) = (V_1(x_1, y), \dots, V_k(x_k, y))$ with oriented strict \mathcal{F} -identification functions V_m for each T_m . We apply Proposition 3.5 and Proposition 4.1 to obtain that there are positive functions g_m and some \mathcal{F} -integrable function a such that

$$S(x_1, \dots, x_k, y) = \sum_{m=1}^k \int_{z_m}^{x_m} g_m(v) V_m(v, y) dv + a(y),$$

for almost all $(x, y) \in \mathbf{A} \times \mathbf{O}$, where $(z_1, \dots, z_k) \in \text{int}(\mathbf{A})$ is a star point of $\text{int}(\mathbf{A})$. Let $F \in \mathcal{F}$, $t = T(F)$ and $x_m \neq t_m$. Then the strict consistency of S implies that $\bar{S}(t, F) < \bar{S}(t_1, \dots, t_{m-1}, x_m, t_{m+1}, \dots, t_m)$. This is equivalent to $\bar{S}_m(t_m, F) < \bar{S}_m(x_m, F)$ with $S_m(x_m, y) := \int_{z_m}^{x_m} g_m(v) V_m(v, y) dv$. \square

Proof of Theorem 5.2

If the class \mathcal{F} satisfies assumption (F2), we can rewrite $\text{ES}_\alpha(F)$ defined at (5.1) in an equivalent form. If f denotes the density of F , it holds that

$$\text{ES}_\alpha(F) = \frac{1}{\alpha} \int_{-\infty}^{F^{-1}(\alpha)} y f(y) dy, \quad \alpha \in (0, 1]. \quad (7.5)$$

We start by proving part (ii). We first show the assertions concerning V given at (5.2). Let $F \in \mathcal{F}$ with density f and let $t = T(F)$. Then we have for $m \in \{1, \dots, k-1\}$, $x \in \mathbf{A}$,

that $\bar{V}_m(x, F) = F(x_m) - q_m$ which is zero if and only if $x_m = t_m$. On the other hand, using the identity at (7.5) (recall that we work under assumption (F2))

$$\bar{V}_k(t_1, \dots, t_{k-1}, x_k, F) = x_k - \sum_{m=1}^{k-1} \frac{p_m}{q_m} \int_{-\infty}^{t_m} y f(y) \, dy = x_k - t_k.$$

Hence, it follows that V is a strict \mathcal{F} -identification function for T . Moreover, V satisfies assumption (V3), and we have for $m \in \{1, \dots, k-1\}$, $l \in \{1, \dots, k\}$ and $x \in \text{int}(\mathbf{A})$ that

$$\begin{aligned} \partial_l \bar{V}_m(x, F) &= \begin{cases} 0, & \text{if } l \neq m \\ f(x_l), & \text{if } l = m, \end{cases} \\ \partial_l \bar{V}_k(x, F) &= \begin{cases} -(p_l/q_l)x_l f(x_l), & \text{if } l \in \{1, \dots, k-1\} \\ 1, & \text{if } l = k. \end{cases} \end{aligned}$$

From now on, we assume that $t = T(F) \in \text{int}(\mathbf{A})$. Let S be a strictly \mathcal{F} -consistent scoring function for T satisfying (S2). Then we can apply Theorem 3.2 and Corollary 3.4 to get that there are locally Lipschitz continuous functions $h_{lm}: \text{int}(\mathbf{A}) \rightarrow \mathbb{R}$ such that (3.2) holds and for all $r, l \in \{1, \dots, k\}$

$$\begin{aligned} \partial_l \partial_r \bar{S}(x, F) &= \sum_{m=1}^k \partial_l h_{rm}(x) \bar{V}_m(x, F) + h_{rm}(x) \partial_l \bar{V}_m(x, F) \\ &= \sum_{m=1}^k \partial_r h_{lm}(x) \bar{V}_m(x, F) + h_{lm}(x) \partial_r \bar{V}_m(x, F) = \partial_r \partial_l \bar{S}(x, F) \end{aligned} \tag{7.6}$$

for $x = t$ and all almost all other $x \in \text{int}(\mathbf{A})$. If we evaluate (7.6) for $r = k, l \in \{1, \dots, k-1\}$ at the point $x = t$ we get

$$h_{kl}(t) \partial_l \bar{V}_l(t, F) + h_{kk}(t) \partial_l \bar{V}_k(t, F) = h_{lk}(t) \partial_k \bar{V}_k(t, F),$$

which takes the form

$$h_{kl}(t) f(t_l) - h_{kk}(t) \frac{p_l}{q_l} t_l f(t_l) = h_{lk}(t).$$

Invoking assumption (V4) for (V_1, \dots, V_{k-1}) we get that $h_{lk}(t) = 0$ and $h_{kl}(t) = (p_l/q_l) t_l h_{kk}(t)$. So with the surjectivity of T we get for $x \in \text{int}(\mathbf{A})$ that

$$h_{lk}(x) = 0, \quad h_{kl}(x) = \frac{p_l}{q_l} x_l h_{kk}(x) \quad \text{for all } l \in \{1, \dots, k-1\}. \tag{7.7}$$

Now, we can evaluate (7.6) for $r, l \in \{1, \dots, k-1\}$, $r \neq l$, at $x = t$ and use the first part of (7.7) to get that

$$h_{rl}(t) f(t_l) = h_{lr}(t) f(t_r).$$

Using again the same argument, we get for $x \in \text{int}(\mathbf{A})$ that

$$h_{rl}(x) = 0 \quad \text{for all } r, l \in \{1, \dots, k-1\}, \quad l \neq r. \tag{7.8}$$

At this stage, we can evaluate (7.6) for $l \in \{1, \dots, k-1\}$, $r \in \{1, \dots, k\}$, $r \neq l$, for some $x \in \text{int}(\mathbf{A})$. Using (7.7) and (7.8) we obtain

$$\sum_{m=1}^k (\partial_l h_{rm}(x) - \partial_r h_{lm}(x)) \bar{V}_m(x_m, F) = 0.$$

Invoking assumption (V1) and using (7.7) and (7.8), we can conclude that for almost all $x \in \mathbf{A}$,

$$\partial_l h_{rr}(x) = 0 \quad \text{for all } l \in \{1, \dots, k-1\}, r \in \{1, \dots, k\}, l \neq r. \quad (7.9)$$

and

$$\partial_l h_{ll}(x) = \frac{p_l}{q_l} h_{kk}(x) \quad \text{for all } l \in \{1, \dots, k-1\}. \quad (7.10)$$

Equation (7.9) for $r = k$ shows that there is a locally Lipschitz continuous function $g_k: \mathbf{A}'_k \rightarrow \mathbb{R}$ such that for all $(x_1, \dots, x_k) \in \text{int}(\mathbf{A})$

$$h_{kk}(x_1, \dots, x_k) = g_k(x_k).$$

Equation (7.10) together with (7.9) gives that for $l \in \{1, \dots, k-1\}$, $(x_1, \dots, x_k) \in \text{int}(\mathbf{A})$

$$h_{ll}(x_1, \dots, x_k) = \frac{p_l}{q_l} G_k(x_k) + g_l(x_l),$$

where $g_l: \mathbf{A}'_l \rightarrow \mathbb{R}$ is locally Lipschitz continuous and $G_k: \mathbf{A}'_k \rightarrow \mathbb{R}$ is such that $G'_k = g_k$.

To show (5.4), we use Lemma 2.9 part (i). Let $D = \{s \in \mathbb{R}: t + sv \in \text{int}(\mathbf{A})\}$, and let $v = (v_1, \dots, v_k) \in \mathbb{S}^{k-1}$ with $v_k \geq 0$. We define $\psi: D \rightarrow \mathbb{R}$ by $\psi(s) := \bar{S}(t + sv, F)$, that is,

$$\begin{aligned} \psi(s) &= \sum_{r=1}^{k-1} \int_{z_r}^{\bar{s}_r} \left(\frac{p_r}{q_r} G_k(z_k) + g_r(v) \right) (F(v) - q_r) dv \\ &\quad + (G_k(\bar{s}_k) - G_k(z_k)) \sum_{m=1}^{k-1} \frac{p_m}{q_m} \left(\bar{s}_m (F(\bar{s}_m) - q_m) - \int_{-\infty}^{\bar{s}_m} y f(y) dy \right) \\ &\quad + \bar{s}_k G_k(\bar{s}_k) - \mathcal{G}_k(\bar{s}_k) + \bar{a}(F), \end{aligned} \quad (7.11)$$

where we use the notation $\bar{s} = t + sv$. The function ψ has a minimum at $s = 0$. Hence, there is $\varepsilon > 0$ such that $\psi'(s) < 0$ for $s \in (-\varepsilon, 0)$ and $\psi'(s) > 0$ for $s \in (0, \varepsilon)$. If $v_k = 0$, then

$$\psi'(s) = \sum_{r=1}^{k-1} (F(\bar{s}_r) - q_r) v_r \left(g_r(\bar{s}_r) + \frac{p_r}{q_r} G_k(\bar{s}_k) \right). \quad (7.12)$$

Choosing v as the m th standard basis vector of \mathbb{R}^k for $m \in \{1, \dots, k-1\}$, the second part of (5.4) follows. On the other hand, if v is the k th standard basis vector, we obtain that $\psi'(s) = g_k(\bar{s}_k)s$, implying the first part of (5.4), in both cases again exploiting the surjectivity of T .

Now, we show part (iii) again using Lemma 2.9 part (i). Let S be a scoring function of the form given at (5.3). For $m \in \{1, \dots, k\}$ let $\mathbf{A}'_m := \{x_m \in \mathbb{R}: (x_1, \dots, x_k) \in \mathbf{A}\}$. Let

$g_m: A'_m \rightarrow \mathbb{R}$ be continuous functions such and $\mathcal{G}_k, G_k: A'_k \rightarrow \mathbb{R}$ with $\mathcal{G}'_k = G_k, G'_k = g_k$. Let (5.4) and (5.5) be satisfied. Fix a distribution $F \in \mathcal{F}$ with continuous density $f = F'$, set $t = (t_1, \dots, t_k) = T(F)$, $D = \{s \in \mathbb{R}: t + sv \in A\}$, and let $v = (v_1, \dots, v_k) \in \mathbb{S}^{k-1}$ with $v_k \geq 0$. We define $\psi: D \rightarrow \mathbb{R}$ as at (7.11). If $v_k = 0$, then it is clear from (7.12) that ψ' is negative for $s < 0$ and positive for $s > 0$ under our assumptions. Therefore ψ has a unique minimum at $s = 0$ as desired. Suppose now that $v_k > 0$. Then, we obtain $\psi'(s) = v_k g_k(\bar{s}_k)(R(s) - L(s))$, where

$$R(s) = \sum_{m=1}^{k-1} \frac{p_m}{q_m} \left(\bar{s}_m (F(\bar{s}_m) - q_m) - \int_{-\infty}^{\bar{s}_m} y f(y) dy \right)$$

and

$$L(s) = -\bar{s}_k - \sum_{l=1}^{k-1} v_l (F(\bar{s}_l) - q_l) \frac{g_l(\bar{s}_l) + \frac{p_l}{q_l} G_k(\bar{s}_k)}{v_k g_k(\bar{s}_k)}.$$

We have $L(0) = R(0) = -t_k$. Furthermore,

$$R'(s) = \sum_{m=1}^{k-1} \frac{p_m}{q_m} v_m (F(\bar{s}_m) - q_m),$$

which is an increasing function with $R'(0) = 0$. Hence the function R is convex, increasing for $s > 0$ and decreasing for $s < 0$. The first summand, $-\bar{s}_k = -t_k - sv_k$, of $L(s)$ is a linear function in s with slope $-v_k < 0$. The second summand vanishes for $s = 0$, it is ≤ 0 for $s > 0$ and ≥ 0 for $s < 0$. This implies that $R(s) > L(s)$ for all $s > 0$, hence $\psi'(s) > 0$ for $s > 0$. The assumptions ensure that $L'(0) \leq -v_k < 0$, so for $s < 0$ close enough to zero, we obtain $\psi'(s) < 0$ indicating that ψ has a local minimum at $s = 0$. It is possible that there exists an $s^* < 0$ such that $\psi'(s^*) = 0$, that is, $R(s^*) = L(s^*)$. We will show that then $\psi(s^*) > \psi(0)$ which leads to the conclusion that ψ has indeed a global minimum at $s = 0$.

Using that $R(s^*) = L(s^*)$, we obtain

$$\begin{aligned} \psi(s^*) &= \sum_{r=1}^{k-1} \int_{z_r}^{\bar{s}_r^*} \left(\frac{p_r}{q_r} G_k(z_k) + g_r(v) \right) (F(v) - q_r) dv \\ &\quad + (G_k(\bar{s}_k^*) - G_k(z_k)) R(s^*) + \bar{s}_k^* G_k(\bar{s}_k^*) - \mathcal{G}_k(\bar{s}_k^*) + \bar{a}(F) \\ &= \sum_{r=1}^{k-1} \int_{z_r}^{\bar{s}_r^*} g_r(v) (F(v) - q_r) dv \\ &\quad - G_k(\bar{s}_k^*) \sum_{l=1}^{k-1} v_l (F(\bar{s}_l^*) - q_l) \frac{g_l(\bar{s}_l^*) + \frac{p_l}{q_l} G_k(\bar{s}_k^*)}{v_k g_k(\bar{s}_k^*)} - \mathcal{G}_k(\bar{s}_k^*) \\ &\quad + G_k(z_k) \sum_{r=1}^{k-1} \frac{p_r}{q_r} \left(\int_{z_r}^{\bar{s}_r^*} (F(v) - q_r) dv - \bar{s}_r^* (F(\bar{s}_r^*) - q_r) + \int_{-\infty}^{\bar{s}_r^*} y f(y) dy \right) + \bar{a}(F). \end{aligned}$$

With partial integration we obtain that

$$\int_{z_r}^{\bar{s}_r^*} (F(v) - q_r) dv - \bar{s}_r^* (F(\bar{s}_r^*) - q_r) + \int_{-\infty}^{\bar{s}_r^*} y f(y) dy = \int_{-\infty}^{z_r} y f(y) dy - z_r (F(z_r) - q_r).$$

Furthermore,

$$-G_k(\bar{s}_k^*) \sum_{l=1}^{k-1} v_l(F(\bar{s}_l^*) - q_l) \frac{g_l(\bar{s}_l^*) + \frac{p_l}{q_l} G_k(\bar{s}_k^*)}{v_k g_k(\bar{s}_k^*)} \geq 0$$

due to (5.4), (5.5) and because $v_l(F(\bar{s}_l^*) - q_l) \leq 0$ for $s^* \leq 0$. As $-G_k(\bar{s}_k)$ is strictly decreasing in s due to assumption (5.5), we obtain

$$\begin{aligned} \psi(s^*) &> \sum_{r=1}^{k-1} \int_{z_r}^{t_r} g_r(v)(F(v) - q_r) dv - G_k(t_k) \\ &+ G_k(z_k) \sum_{r=1}^{k-1} \frac{p_r}{q_r} \left(\int_{-\infty}^{z_r} y f(y) dy - z_r(F(z_r) - q_r) \right) + \bar{a}(F) = \psi(0), \end{aligned}$$

where we used the fact that $\int_{z_r}^{\bar{s}_r} g_r(v)(F(v) - q_r) dv$ is decreasing for $s \leq 0$.

Finally, part (i) follows from part (iii) and Remark 5.3 by the definition of elicibility upon choosing, for example, $\mathbf{A} = \mathbb{R}^k$, $g_m \equiv 1$, $z_m = 0$ for $m \in \{1, \dots, k-1\}$, $g_k(s) = G_k(s) = \mathcal{G}_k(s) = e^s$, $z_k = -\infty$, $a \equiv 0$. \square

Proof of Corollary 5.4

For the first part of the claim, note that if $\mu(\{1\}) = 1$, then ν_μ coincides with the expectation and is thus 1-elicitable. If $\mu(\{1\}) = 0$, the assertion of the corollary is a direct consequence of Theorem 5.2 (i). If $\lambda := \mu(\{1\}) \in (0, 1)$, then we can write

$$\mu = \sum_{m=1}^{k-2} p_m \delta_{q_m} + \lambda \delta_1$$

where $p_m \in (0, 1)$, $\sum_{m=1}^{k-2} p_m = 1 - \lambda$, $q_m \in (0, 1)$ and the q_m 's are pairwise distinct. Define the probability measure

$$\tilde{\mu} := \sum_{m=1}^{k-2} \frac{p_m}{1 - \lambda} \delta_{q_m}.$$

Using Theorem 5.2 (i), the functional $(T'_1, \dots, T'_{k-1}): \mathcal{F} \rightarrow \mathbb{R}^{k-1}$ is $(k-1)$ -elicitable where $T'_m(F) := F^{-1}(q_m)$, $m \in \{1, \dots, k-2\}$, and $T'_{k-1}(F) = \nu_{\tilde{\mu}}(F)$. Using Lemma 2.15 we can deduce that the functional $(T'_1, \dots, T'_{k-1}, \nu_{\delta_1}): \mathcal{F} \rightarrow \mathbb{R}^k$ is k -elicitable. Note that

$$\nu_\mu = (1 - \lambda)\nu_{\tilde{\mu}} + \lambda\nu_{\delta_1}.$$

Hence, we can apply Proposition 2.13 to deduce that the functional $T = (T_1, \dots, T_k): \mathcal{F} \rightarrow \mathbb{R}^k$ is k -elicitable where $T_m = T'_m$, $m \in \{1, \dots, k-2\}$, $T_{k-1} = \nu_{\delta_1}$ and $T_k = \nu_\mu$. \square

Acknowledgements

This project is supported by the Swiss National Science Foundation (SNF) via grant 152609. We would like to thank Carlo Acerbi, Rafael Frongillo and Tilmann Gneiting for fruitful discussions which inspired some of the results of this paper.

References

- J. Abernethy and R. Frongillo. A Characterization of Scoring Rules for Linear Properties. In *Proceedings of the 25th Conference on Learning Theory*, 2012.
- C. Acerbi. Spectral measures of risk: A coherent representation of subjective risk aversion. *Journal of Banking & Finance*, 26:1505–1518, 2002.
- C. Acerbi and B. Szekely. Backtesting Expected Shortfall. *Risk Magazine*, 2014.
- P. Artzner, F. Delbaen, J.-M. Eber, and D. Heath. Coherent measures of risk. *Math. Finance*, 9:203–228, 1999.
- A. Banerjee, X. Guo, and H. Wang. On the Optimality of Conditional Expectation as a Bregman Predictor. *IEEE Trans. Inform. Theory*, 51:2664–2669, 2005.
- F. Bellini and V. Bignozzi. Elicitable Risk Measures. *Quant. Finance*, 2014. To appear.
- R. Cont, R. Deguest, and G. Scandolo. Robustness and sensitivity analysis of risk measurement procedures. *Quant. Finance*, 10:593–606, 2010.
- J. Daniélsson, P. Embrechts, C. Goodhart, C. Keating, F. Muennich, O. Renault, and H. S. Shin. An Academic Response to Basel II. Special paper no. 130, Financial Markets Group, London School of Economics, 2001.
- M. Davis. Consistency of Risk Measure Estimates. *Preprint*, <http://ssrn.com/abstract=2342279>, 2013.
- A. P. Dawid and P. Sebastiani. Coherent dispersion criteria for optimal experimental design. *Ann. Statist.*, 27:65–81, 1999.
- F. Delbaen. *Monetary utility functions*. Osaka University Press, 2012.
- F. Delbaen, F. Bellini, V. Bignozzi, and J. F. Ziegel. Risk Measures with the CxLS property. *Preprint*, <http://arxiv.org/pdf/1411.0426v1.pdf>, 2014.
- P. Embrechts and M. Hofert. Statistics and Quantitative Risk Management for Banking and Insurance. *Ann. Rev. Stat. Appl.*, 1, 2014.
- S. Emmer, M. Kratz, and D. Tasche. What is the best risk measure in practice? A comparison of standard measures. *Preprint*, <http://arxiv.org/abs/1312.1645v3>, 2013.
- J. Engelberg, C. F. Manski, and J. Williams. Comparing the point predictions and subjective probability distributions of professional forecasters. *J. Bus. Econ. Stat.*, 27:30–41, 2009.
- R. Frongillo and I. Kash. Vector-Valued Property Elicitation. *Preprint*, <http://people.seas.harvard.edu/~raf/media/papers/vec-props.pdf>, 2014.

- T. Gneiting. Making and Evaluating Point Forecasts. *J. Amer. Statist. Assoc.*, 106: 746–762, 2011.
- T. Gneiting and A. Raftery. Strictly Proper Scoring Rules, Prediction, and Estimation. *J. Amer. Statist. Assoc.*, 102:359–378, 2007.
- H. Grauert and W. Fischer. *Differential- und Integralrechnung II*. Springer-Verlag, Berlin Heidelberg New York, 1978.
- C. Heinrich. The mode functional is not elicitable. *Biometrika*, 2013.
- P. J. Huber. Robust Estimation of a Location Parameter. *Ann. Math. Statist.*, pages 73–101, 1964.
- E. Jouini, W. Schachermayer, and N. Touzi. Law invariant risk measures have the Fatou property. In *Adv. Math. Econ.*, volume 9, pages 46–71. Springer, Tokyo, 2006.
- R. Koenker. *Quantile Regression*. Cambridge University Press, Cambridge, 2005.
- S. Kou and X. Peng. On the Measurement of Economic Tail Risk. *Preprint*, <http://arxiv.org/pdf/1401.4787v2.pdf>, 2014.
- S. Kou, X. Peng, and C. C. Heyde. External Risk Measures and Basel Accords. *Math. Oper. Res.*, 38:393–417, 2013.
- V. Krätschmer, A. Schied, and H. Zähle. Qualitative and infinitesimal robustness of tail-dependent statistical functionals. *J. Multivariate Anal.*, 103:35–47, 2012.
- V. Krätschmer, A. Schied, and H. Zähle. Quasi-Hadamard differentiability of general risk functionals and its applications. *Statistics & Risk Modeling*, 2013. To appear.
- V. Krätschmer, A. Schied, and H. Zähle. Comparative and qualitative robustness for law-invariant risk measures. *Finance Stoch.*, 18:271–295, 2014.
- S. Kusuoka. On law-invariant coherent risk measures. *Adv. Math. Econ.*, 3:83–95, 2001.
- N. Lambert. Elicitation and Evaluation of Statistical Functionals. *Preprint*, <http://web.stanford.edu/~nlambert/papers/elicitation.pdf>, 2013.
- N. Lambert, D. M. Pennock, and Y. Shoham. Eliciting properties of probability distributions. In *Proceedings of the 9th ACM Conference on Electronic Commerce*, pages 129–138, Chicago, IL, USA, 2008. ACM.
- A. H. Murphy and H. Daan. Forecast Evaluation. In A. H. Murphy and R. W. Katz, editors, *Probability, Statistics and Decision Making in the Atmospheric Sciences*, pages 379–437. Westview Press, Boulder, Colorado, 1985.
- W. K. Newey and J. L. Powell. Asymmetric Least Squares Estimation and Testing. *Econometrica*, 55:819–847, 1987.

- K. Osband and S. Reichelstein. Information-Eliciting Compensation Schemes. *J. Public Econ.*, 27:107–115, 1985.
- K. H. Osband. *Providing Incentives for Better Cost Forecasting*. PhD thesis, University of California, Berkeley, 1985.
- L. J. Savage. Elicitation of Personal Probabilities and Expectations. *J. Amer. Statist. Assoc.*, 66:783–801, 1971.
- I. Steinwart, C. Pasin, R. Williamson, and S. Zhang. Elicitation and Identification of Properties. *JMLR: Workshop and Conference Proceedings*, 35:1–45, 2014.
- R. Wang and J. F. Ziegel. Elicitable distortion risk measures: A concise proof. *Statistics and Probability Letters*, 100:172–175, 2015.
- S. Weber. Distribution-Invariant Risk Measures, Information, and Dynamic Consistency. *Math. Finance*, 16:419–441, 2006.
- R. L. Winkler. Scoring rules and the evaluation of probabilities. *Test*, 5:1–60, 1996. with discussion.
- J. F. Ziegel. Coherence and elicibility. *Math. Finance*, 2015. To appear.