Efficient Online Minimization for Low-Rank Subspace Clustering

Jie Shen

Dept. of Computer Science School of Arts and Sciences Rutgers University Piscataway, NJ 08854, USA js2007@rutgers.edu

Ping Li

Dept. of Statistics & Biostatistics
Department of Computer Science
Rutgers University
Piscataway, NJ 08854, USA
pingli@stat.rutgers.edu

Huan Xu

Industrial & Systems Engineering Georgia Institute of Technology Atlanta, GA 30332, USA huan.xu@isye.gatech.edu

Abstract

Low-rank representation (LRR) has been a significant method for segmenting data that are generated from a union of subspaces. It is, however, known that solving the LRR program is challenging in terms of time complexity and memory footprint, in that the size of the nuclear norm regularized matrix is n-by-n (where n is the number of samples). In this paper, we thereby develop a fast online implementation of LRR that reduces the memory cost from $O(n^2)$ to O(pd), with p being the ambient dimension and p being some estimated rank (p < p < p). The crux for this end is a non-convex reformulation of the LRR program, which pursues the basis dictionary that generates the (uncorrupted) observations. We build the theoretical guarantee that the sequence of the solutions produced by our algorithm converges to a stationary point of the empirical and the expected loss function asymptotically. Extensive experiments on synthetic and realistic datasets further substantiate that our algorithm is fast, robust and memory efficient.

Keywords: Low-Rank Representation, Subspace Clustering, Online Optimization

1 Introduction

In modern scientific computing, data are routinely generated from a union of small subspaces for which, traditional tools such as principal component analysis are not able to identify the group structure. As an alternative, in the past a few years, subspace clustering (Vidal, 2010; Soltanolkotabi and Candès, 2012) has been extensively studied and has established solid applications in, for example, computer vision (Elhamifar and Vidal, 2009) and network topology inference (Eriksson et al., 2011). Among many subspace clustering methods which seek a structured representation to fit the underlying data, two prominent examples are sparse subspace clustering (SSC) (Elhamifar and Vidal, 2009; Soltanolkotabi et al., 2014) and low-rank representation (LRR) (Liu et al., 2013). Both of them utilize the idea of self-expressiveness, i.e., expressing each sample as a linear combination of the remaining. Hence, the coefficients quickly suggest which data points belong to the same group (i.e., subspace). What is of difference is that SSC pursues a sparse solution, i.e., using a small fraction of samples to represent each data point. In this light, not only the clustering structure can be found by SSC, but also the most correlated samples are detected. For LRR, it prefers a (global) low-rank structure. This is motivated by many applications where the union of the subspaces is still of low

rank. An appealing property of LRR is that it comes up with the theoretical guarantee of recovering the true low-rank model in addition to clustering the samples.

In this paper, we are interested in the LRR method, which is shown to achieve the state-of-the-art performance on a broad range of real-world problems (Liu et al., 2013). Recently, Liu and Li (2014) demonstrated that, when equipped with a proper dictionary, LRR can even handle the coherent data – a challenging issue in the literature (Candès and Recht, 2009; Candès et al., 2011) but is ubiquitous in realistic datasets such as the Netflix movie rating problem.

Formally, the LRR program we investigate here is formulated as follows (Liu et al., 2013):

$$\min_{X,E} \frac{\lambda_1}{2} \|Z - YX - E\|_F^2 + \|X\|_* + \lambda_2 \|E\|_1.$$
 (1)

Here, $Z=(z_1,z_2,\cdots,z_n)\in\mathbb{R}^{p\times n}$ is the observation matrix with n samples, each of which is a p-dimensional column vector. The matrix $Y\in\mathbb{R}^{p\times n}$ is a given dictionary, E is some possible sparse corruption and $\lambda_1>0$ and $\lambda_2>0$ are two tunable parameters. Typically, Y is chosen as the dataset Z itself, hence the idea of self-expressiveness. As X is penalized by the nuclear norm which is a convex surrogate of the rank function, the program seeks a low-rank representation among all samples, each of which can be approximated by a linear combination of the atoms in the dictionary Y. Note that a column of X is the coefficients associated with an individual sample. When the optimal solution is obtained, one may perform spectral clustering (Ng et al., 2001) on X since the entries reflect the correlation between the (uncorrupted) data.

To aid intuition and to introduce more background for the LRR program (1), we discuss the ability of subspace clustering of LRR in the noiseless case. That is, the data matrix Z is generated from a union of (low-rank) subspaces. Since each sample can be represented only by those belonging to the same subspace, one may simply solve the linear system

$$Z = ZX$$
.

which certainly produces a block diagonal solution of X as long as the small subspaces are disjoint. That being said, a careful reader may observe that the above linear system does not exclude the trivial identity matrix $X = I_n$. In order to alleviate it and to conform the low-rank structure of the data, LRR looks for the one with the lowest rank, i.e.,

$$\min_{X} \|X\|_*, \text{ s.t. } Z = ZX.$$
(2)

The following lemma, which is due to Liu et al. (2010), justifies that the above program correctly segments the data.

Lemma 1. Suppose that there are k number of small subspaces and without loss of generality, the data matrix is organized as $Z = (Z_1, Z_2, \dots, Z_k)$, where $Z_i \in \mathbb{R}^{p \times n_i}$ is the collection of samples coming from the ith subspace. Further assume that there are sufficient samples for each subspace such that $n_i > \text{rank}(Z_i)$. If the subspaces are mutually disjoint, then there exists an optimal solution to (2),

$$X^* = \begin{pmatrix} X_1^* & 0 & 0 & 0 \\ 0 & X_2^* & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & X_k^* \end{pmatrix}$$

such that $\operatorname{rank}(X_i^*) = \operatorname{rank}(Z_i)$ for all $1 \le i \le k$.

More generally, one may think of the noisy model Z = YX + E + G for some given dictionary Y, a sparse corruption E and a Gaussian noise G. In this case, program (1) is a straightforward extension to achieve the robustness. While a large body of work has shown that LRR is able to segment the data (see, e.g., Liu and Li (2014)), three issues are immediately incurred for LRR in the face of big data:

- (I1) Memory cost of X. In the LRR formulation (1), there is typically no sparsity assumption on X. Hence, the memory footprint of X is proportional to n^2 which precludes most of the recently developed nuclear norm solvers (Lin et al., 2010; Jaggi and Sulovský, 2010; Avron et al., 2012; Hsieh and Olsen, 2014).
- (I2) Computational cost of $||X||_*$. Since the size of the nuclear norm regularized matrix X is $n \times n$, optimizing such problems can be computationally expensive even when n is moderate, say n = 1000 (Recht et al., 2010).
- (13) Memory cost of Y. As the size of the dictionary Y is proportional to sample size n, it is prohibitive to store the entire dictionary Y during optimization when n is large.

To remedy these problems, especially the memory bottleneck, one potential way is solving the LRR program in an online manner. That is, we sequentially reveal the samples z_1, z_2, \cdots, z_n and update the components in X and E. Nevertheless, it turns out that the LRR program (1) is not suitable for online minimization. To see this, we note that each column of X is the coefficients of a sample with respect to the *entire* dictionary Y, for example, $z_1 \approx Yx_1 + e_1$. This indicates that without further technique, we have to load the entire dictionary Y so as to update x_1 and x_2 and x_3 and x_4 and

(I4) Partial realization of Y. We are required to guarantee the optimality of the solution but can only access part of the atoms of Y in each iteration.

1.1 Contribution

In this paper, henceforth, we propose a new algorithm termed online low-rank subspace clustering (OLRSC), which admits a low computational complexity. Compared to existing solvers, OLRSC reduces the memory cost of LRR from $O(n^2)$ to O(pd) where d is an estimated rank of the uncorrupted data (d). This nice property makes OLRSC an appealing solution for large-scale subspace clustering problems. Furthermore, we prove that the sequence of the solutions produced by OLRSC converges to a stationary point of the expected loss function asymptotically even though one atom of <math>Y is available at each iteration. In a nutshell, OLRSC resolves all practical issues of LRR and still promotes global low-rank structure – the merit of LRR. Finally, concerning the robustness of the algorithm, our empirical study suggests accurate recovery of the true subspace even when a large fraction of the entries are corrupted.

The present paper considerably extends our earlier conference version (Shen et al., 2016). In particular, we make extra efforts to remove the assumption that the given dictionary Y has to be full-rank, which indicates that our algorithm can be applied to a broader range of problems. We also propose refined conditions on parameter scaling under which the algorithm converges. Lastly, we draw additional connections to the literature.

1.2 Related Work

Low-rankness has been studied for more than two decades. As one of the earliest work, Fazel et al. (2001) appealed to a rank minimization program for system identification and signal processing. It was further

suggested that the nuclear norm (also known as trace norm) is a good convex surrogate to the rank of a matrix. Such a key observation was utilized in machine learning for recommender systems (Srebro et al., 2004; Rennie and Srebro, 2005). Though empirically effective, it was not clear under which conditions such a nuclear norm based program exactly returns the true low-rank solution. In the seminal work of Candès et al. (2011), it was shown that if the singular vectors of the true low-rank matrix does not concentrate around the canonical basis, formally referred to as the incoherence property, then a convex program with proper parameter setting recovers the underlying matrix even a constant fraction of the entries are corrupted. Following Candès et al. (2011), a considerable number of work extends the low-rank model to accommodate outliers (Xu et al., 2013), high-dimensional case (Xu et al., 2010), graph clustering (Chen et al., 2014), to name a few. This work considers the variant of multiple subspaces, which is due to Liu et al. (2010). In particular, in place of drawing insight on the statistical performance, we mainly focus on an efficient and provable algorithm for subspace clustering. There is a plethora of work attempting to mitigate the memory and computational bottleneck of the nuclear norm regularizer. However, to the best of our knowledge, none of them can handle Issue (13) and Issue (14).

One of the most popular ways to alleviate the huge memory cost is online implementation. Feng et al. (2013) devised an online algorithm for the robust principal component analysis (RPCA) problem, which makes the memory cost independent of the sample size. Yet, compared to RPCA where the size of the nuclear norm regularized matrix is $p \times n$, that of LRR is $n \times n$ – a worse and more challenging case. Moreover, their algorithm cannot address the partial dictionary issue that emerges in our case.

To tackle the computational overhead, Jaggi and Sulovský (2010) utilized a sparse semi-definite programming solver to derive a simple yet efficient algorithm. Unfortunately, the memory requirement of their algorithm is proportional to the number of observed entries, making it impractical when the regularized matrix is large and dense (which is the case of LRR). Avron et al. (2012) combined stochastic subgradient and incremental SVD to boost efficiency. But for the LRR problem, the type of the loss function does not meet the requirements and thus, it is still not practical to use that algorithm in our case.

Another line in the literature explores a structured formulation of LRR beyond the low-rankness. For example, Wang et al. (2013) provably showed that combining LRR and SSC can take advantages of both methods. Whereas Wang et al. (2013) promotes the sparsity on the representation matrix X, Shen and Li (2016) demonstrated how to pursue a sparsity structure on the factors of X, which is known to be more flexible (Haeffele et al., 2014). Due to the non-convexity of matrix factorization, a large body of work is dedicated to characterizing the conditions under which gradient descent ensures global optimum (Jain et al., 2013; Lee et al., 2016; Ge et al., 2016).

Subsequent to the conference version, the authors made use of similar idea to present a structured low-rank representation framework (Shen and Li, 2016). Inspired by the unified framework, Zhao et al. (2017) presents a non-asymptotic algorithm to solve the subspace clustering problem.

1.3 Overview

In Section 2, we reformulate the LRR program (1) so that it is amenable to online optimization. The detailed algorithm is presented in Section 2.2, along with a theoretical justification on its convergence behavior. We illustrate the efficiency and efficacy of the proposed algorithm by performing extensive experiments in Section 4. Finally, we conclude the paper in Section 5. All the proofs can be found in the appendix.

1.4 Notation

We use bold lowercase letters, e.g., v, to denote column vectors. The ℓ_2 norm and ℓ_1 norm of a vector v are denoted by $||v||_2$ and $||v||_1$ respectively. Capital letters such as M are used to denote a matrix, and its transpose is denoted by M^{\top} . For an invertible matrix M, we write its inverse as M^{-1} . The capital letter I

is reserved for the identity matrix. When necessary, we write I_r to emphasize its size. The jth column of a matrix M is denoted by m_j , and we write m_{ij} for the (i,j)-th component of M. Three matrix norms will be used: $\|M\|_*$ for the nuclear norm which is the sum of the singular values of M, $\|M\|_F$ for the Frobenius norm and $\|M\|_1$ for the ℓ_1 norm of a matrix seen as a long vector. The trace of a square matrix M is denoted by $\mathrm{Tr}(M)$.

For an integer n, we use [n] to denote the integer set $\{1, 2, \cdots, n\}$. We follow the standard asymptotic notation. That is, for two functions f(n) and g(n), we use f(n) = O(g(n)) to denote that there exists an absolute constant C such that $f(n) \leq C \cdot g(n)$ as n goes to infinity. We also write $f(n) = \Omega(g(n))$ if $f(n) \geq C \cdot g(n)$. When $C_1 \cdot g(n) \leq f(n) \leq C_2 \cdot g(n)$, we write $f(n) = \Theta(g(n))$.

2 Problem Formulation and Algorithm

Recall the LRR program given in (1). Our goal is to efficiently learn the representation matrix X and the corruption matrix E in an online manner so as to mitigate the issues mentioned in Section 1. To this end, we reformulate it as an empirical risk minimization problem which is amenable for online optimization.

The first technique for our purpose is a *non-convex reformulation* of the nuclear norm. Assume that the rank of X is at most d. Then Fazel et al. (2001) showed that,

$$||X||_* = \min_{U,V,X=U^\top V} \frac{1}{2} \left(||U||_F^2 + ||V||_F^2 \right), \tag{3}$$

where $U \in \mathbb{R}^{d \times n}$ and $V \in \mathbb{R}^{d \times n}$. The minimum can be attained at, for example, $U^{\top} = U_0 S_0^{1/2}$ and $V^{\top} = V_0 S_0^{1/2}$ where $X = U_0 S_0 V_0^{\top}$ is the singular value decomposition. In this way, (1) can be written as follows:

$$\min_{UVE} \frac{\lambda_1}{2} \left\| Z - YU^\top V - E \right\|_F^2 + \frac{1}{2} \left\| U \right\|_F^2 + \frac{1}{2} \left\| V \right\|_F^2 + \lambda_2 \left\| E \right\|_1. \tag{4}$$

Note that by this reformulation, updating the entries in X amounts to sequentially updating the columns of U and V, as shown below:

$$\underbrace{\begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nn} \end{pmatrix}}_{Y \in \mathbb{P}^{n \times n}} = \underbrace{\begin{pmatrix} \boldsymbol{u}_1^\top \\ \boldsymbol{u}_2^\top \\ \vdots \\ \boldsymbol{u}_n^\top \end{pmatrix}}_{U^\top \subset \mathbb{P}^{n \times d}} \times \underbrace{\begin{pmatrix} \boldsymbol{v}_1 & \boldsymbol{v}_2 & \cdots & \boldsymbol{v}_n \end{pmatrix}}_{V \in \mathbb{R}^{d \times n}}.$$

For instance, when we have $\{u_i\}_{i=1}^t$ and $\{v_j\}_{j=1}^t$ on hand, we are able to recover the components $\{x_{ij}\}_{1\leq i,j\leq t}$ since each x_{ij} equals $u_i^{\top}v_j$. It is worth mentioning that this technique is utilized in Feng et al. (2013) for online RPCA. Unfortunately, the size of U and V in our problem are both proportional to the sample size n and the dictionary Y is partially observed in each iteration, making the algorithm in Feng et al. (2013) not applicable to LRR. Related to the online implementation, another challenge is that, all the columns of U are coupled together at this moment as U^{\top} is left multiplied by Y in the first term of (4). Since we do not want to load the entire dictionary Y, this makes it difficult to sequentially compute the columns of U.

For the sake of decoupling the columns of U, as part of the crux of our techniques, we introduce an auxiliary variable $D = YU^{\top}$, whose size is $p \times d$ (i.e., independent of the sample size n). Interestingly, in this way, we are approximating the term Z - E with DV, which provides an intuition on the role of D: namely, D can be seen as a *basis dictionary* of the clean data, with V being the coefficients.

 $^{^{1}}$ The matrix Y is referred to as atom dictionary, and D is referred to as basis dictionary.

These key observations allow us to derive an equivalent reformulation to LRR (1):

$$\min_{D,U,V,E} \frac{\lambda_1}{2} \|Z - DV - E\|_F^2 + \frac{1}{2} \left(\|U\|_F^2 + \|V\|_F^2 \right) + \lambda_2 \|E\|_1, \quad \text{s.t. } D = YU^\top.$$

By penalizing the constraint in the objective, we obtain a *regularized* version of LRR on which our new algorithm is based:

$$\min_{D,U,V,E} \frac{\lambda_1}{2} \|Z - DV - E\|_F^2 + \frac{1}{2} \left(\|U\|_F^2 + \|V\|_F^2 \right) + \lambda_2 \|E\|_1 + \frac{\lambda_3}{2} \|D - YU^\top\|_F^2. \tag{5}$$

We note that there are two advantages of (5) compared to (1). First, it is amenable for online optimization. Second, it is more informative since it explicitly models the basis of the union of subspaces, which yields a better subspace recovery and clustering to be shown in Section 4.

We also point out that due to our explicit modeling of the basis, we unify LRR and RPCA as follows: for LRR, $D \approx YU^{\top}$ (or $D = YU^{\top}$ if λ_3 tends to infinity) while for RPCA, $D = U^{\top}$. That is, ORPCA (Feng et al., 2013) considers a problem of $Y = I_p$ whose size is independent of n, hence can be kept in memory which naturally resolves Issue (I3) and (I4). This is why RPCA can be easily implemented in an online fashion while LRR cannot.

Now we return to the online implementation of LRR. The main idea is optimizing a surrogate of the empirical risk function (to be defined) in each iteration, which is fast and memory efficient. Let z_i , y_i , e_i , u_i , and v_i be the *i*th column of matrices Z, Y, E, U and V respectively and let

$$\tilde{\ell}(D, \boldsymbol{v}, \boldsymbol{e}; \boldsymbol{z}) \stackrel{\text{def}}{=} \frac{\lambda_1}{2} \|\boldsymbol{z} - D\boldsymbol{v} - \boldsymbol{e}\|_2^2 + \frac{1}{2} \|\boldsymbol{v}\|_2^2 + \lambda_2 \|\boldsymbol{e}\|_1,
\ell(D; \boldsymbol{z}) \stackrel{\text{def}}{=} \min_{\boldsymbol{v}, \boldsymbol{e}} \tilde{\ell}(D, \boldsymbol{v}, \boldsymbol{e}; \boldsymbol{z}).$$
(6)

Further, we define

$$\tilde{h}(D, U; Y) \stackrel{\text{def}}{=} \sum_{i=1}^{n} \frac{1}{2} \|\boldsymbol{u}_i\|_2^2 + \frac{\lambda_3}{2} \left\| D - \sum_{i=1}^{n} \boldsymbol{y}_i \boldsymbol{u}_i^{\top} \right\|_F^2,$$

$$h(D; Y) \stackrel{\text{def}}{=} \min_{U} \tilde{h}(D, U; Y). \tag{7}$$

Then (5) can be rewritten as follows:

$$\min_{D} \min_{U,V,E} \sum_{i=1}^{n} \tilde{\ell}(D, \boldsymbol{v}_i, \boldsymbol{e}_i; \boldsymbol{z}_i) + \tilde{h}(D, U; Y), \tag{8}$$

which amounts to minimizing the *empirical loss* function:

$$\min_{D} f_n(D) \stackrel{\text{def}}{=} \frac{1}{n} \sum_{i=1}^n \ell(D; \boldsymbol{z}_i) + \frac{1}{n} h(D; Y). \tag{9}$$

2.1 Expected Loss

In stochastic approximation, we are also interested in analyzing the optimality of the obtained solution with respect to the expected loss function (Bottou and Bousquet, 2007). To this end, we first derive the optimal solutions \tilde{U} , \tilde{V} and \tilde{E} that minimize (8) which renders a concrete form of the empirical loss function $f_n(D)$.

Given D, we need to compute the optimal solutions \tilde{U} , \tilde{V} and \tilde{E} to evaluate the objective value of $f_n(D)$. What is of interest here is that, the optimization procedure of U is quite different from that of V and E. According to (6), when D is given, each \tilde{v}_i and \tilde{e}_i can be solved by only accessing the ith sample z_i . However, the optimal \tilde{u}_i depends on the whole dictionary Y as the second term in $\tilde{h}(Y,D,U)$ couples all the u_i 's. Fortunately, we can obtain a closed form solution for each \tilde{u}_i , as stated below.

Proposition 1. Suppose that Y and D are fixed. Then the optimal solution $\tilde{U} = (\tilde{u}_1, \tilde{u}_2, \dots, \tilde{u}_n)$ that minimizes $\tilde{h}(D, U; Y)$ and Eq. (5) is given by

$$\tilde{\boldsymbol{u}}_i = D^{\top} \left(\frac{1}{\lambda_3} I_p + Y Y^{\top} \right)^{-1} \boldsymbol{y}_i, \ \forall i \in [n].$$
 (10)

Hence,

$$h(D;Y) = \frac{1}{2} \left\| D^{\top} \left(\frac{1}{\lambda_3} I_p + Y Y^{\top} \right)^{-1} Y \right\|_F^2 + \frac{1}{2\lambda_3} \left\| \left(\frac{1}{\lambda_3} I_p + Y Y^{\top} \right)^{-1} D \right\|_F^2.$$
 (11)

The result follows by noting the first order optimality condition and some basic algebra. The proof can be found in Appendix B.2.

Let $\sigma_i(Y)$ be the singular values of the matrix $Y \in \mathbb{R}^{p \times n}$ where $1 \leq i \leq p$. By calculation, it is not hard to see that

$$\sigma_{\max}\left(\left(\frac{1}{\lambda_3}I_p + YY^{\top}\right)^{-1}\right) = \frac{1}{(\sigma_p(Y))^2 + \lambda_3^{-1}} \le \lambda_3,$$

$$\sigma_{\max}\left(\left(\frac{1}{\lambda_3}I_p + YY^{\top}\right)^{-1}Y\right) = \frac{\sigma_i(Y)}{(\sigma_i(Y))^2 + \lambda_3^{-1}} \le \frac{\sqrt{\lambda_3}}{2} \text{ for some } i.$$

Thereby, we obtain the upper bound

$$h(D;Y) \le \frac{\lambda_3}{8} \|D\|_F^2 + \frac{\lambda_3}{2} \|D\|_F^2 \le \frac{5\lambda_3}{8} \|D\|_F^2.$$

Suppose that D is fixed. Also notice that the size of D does not grow with n. It follows that

$$\lim_{n \to \infty} \frac{1}{n} h(D; Y) = 0,$$

if λ_3 is such that $\lim_{n\to\infty}\lambda_3/n=0$. Hence, asymptotically, minimizing the empirical loss $f_n(D)$ amounts to optimizing $n^{-1}\sum_{i=1}^n\ell(D;\mathbf{z})$. Note that when λ_1 and λ_2 are independent of n, we quickly get

$$\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \ell(D; \boldsymbol{z}_i) = \mathbb{E}_{\boldsymbol{z}}[\ell(D; \boldsymbol{z})], \tag{12}$$

assuming that all the samples are drawn i.i.d. from some unknown distribution. This gives the *expected loss* function

$$f(D) \stackrel{\text{def}}{=} \mathbb{E}_{\boldsymbol{z}}[\ell(\boldsymbol{z}, D)] = \lim_{n \to \infty} f_n(D).$$
 (13)

2.2 Algorithm

We are now in the position to elaborate an online optimization algorithm for low-rank subspace clustering. Namely, we show how to solve (9) in an efficient manner. From a high level, we alternatively minimize over the variables D and $\{u_i, v_i, e_i\}_{i=1}^n$. That is, we begin with an initial (and inaccurate) guess of the variable D, and solve the subproblems (6) and (7). Using the obtained solution, we construct a surrogate function of the empirical loss (9), which is further optimized to refine our initial guess of D. Such a paradigm was utilized in Mairal et al. (2010); Mairal (2013); Shen et al. (2014) for different problems. It turns out that the case studied here is more challenging due to the component h(D; Y) of the empirical loss function. To optimize it in an online manner and to analyze the performance, we develop novel techniques that will be described in the following.

2.2.1 Optimize v and e

For exposition, we first investigate the sum of $\ell(D; \mathbf{z}_i)$ where $1 \le i \le n$ in (9). This is where the variables \mathbf{v}_i and \mathbf{e}_i are involved. Suppose that at the t-th iteration, a fresh sample \mathbf{z}_t is drawn and we have an initial guess of D, say D_{t-1} . Then it is easy to see that the optimal solution $\{\mathbf{v}_t^*, \mathbf{e}_t^*\}$ that minimizes $\tilde{\ell}(D_{t-1}, \mathbf{v}, \mathbf{e}; \mathbf{z}_t)$ is given by

$$\{v_t^*, e_t^*\} = \operatorname*{arg\,min}_{v,e} \ rac{\lambda_1}{2} \| oldsymbol{z}_t - D_{t-1} oldsymbol{v} - oldsymbol{e} \|_2^2 + rac{1}{2} \| oldsymbol{v} \|_2^2 + \lambda_2 \| oldsymbol{e} \|_1.$$

Since the objective function is jointly strongly convex over the variables $\{v, e\}$, one may apply coordinate descent to obtain the optimum (Bertsekas, 1999). In particular, we observe that if e is fixed, we can optimize v in closed form:

$$v = (D_{t-1}^{\top} D_{t-1} + I_d / \lambda_1)^{-1} D_{t-1}^{\top} (z_t - e).$$
(14)

Conversely, given v, the variable e is obtained via soft-thresholding (Donoho, 1995):

$$e = \mathcal{S}_{\lambda_2/\lambda_1}[z_t - D_{t-1}v]. \tag{15}$$

See Algorithm 2 in Appendix A for details. It follows that $\tilde{\ell}(D_{t-1}, \boldsymbol{v}_t^*, \boldsymbol{e}_t^*; \boldsymbol{z}_t)$ is a surrogate to $\ell(D; \boldsymbol{z}_t)$, since D_{t-1} is our guess of the optimal D.

2.2.2 Optimize u

We move on to describe how to optimize the second component h(D;Y), which contains the variables $\{u_i\}_{i\geq 1}$. Again, we will assume that an initial guess of D is available, so we only need to minimize a surrogate. However, looking at the solution (10), we notice that even D is given (e.g., some initial guess D_{t-1}), we have to load the *entire* dictionary $Y \in \mathbb{R}^{p \times n}$ so as to obtain the local optimum \tilde{u}_t , and hence a surrogate of h(D;Y). This is quite different from the scheme of optimizing v and e, where optimum only depends on the new sample. Indeed, though the framework of our algorithm follows from Mairal et al. (2010); Feng et al. (2013); Shen et al. (2014), none of them considers the situation as we stated here (the optimum in their work can be computed directly when a new sample arrives).

In order to remedy the issue, our novelty here is constructing a proxy function by minimizing which, we are generating a solution that gradually approximates to the one of $\tilde{h}(D,U;Y)$. The proxy function is given as follows

$$\tilde{\ell}_2(D, \boldsymbol{u}; M, \boldsymbol{y}) \stackrel{\text{def}}{=} \frac{1}{2} \|\boldsymbol{u}\|_2^2 + \frac{\lambda_3}{2} \|D - M - \boldsymbol{y} \boldsymbol{u}^\top\|_F^2.$$
(16)

Algorithm 1 Online Low-Rank Subspace Clustering

Require: $Z \in \mathbb{R}^{p \times n}$ (observed samples), $Y \in \mathbb{R}^{p \times n}$, parameters λ_1 , λ_2 and λ_3 , random matrix $D_0 \in \mathbb{R}^{p \times d}$ (initial basis), zero matrices M_0 , A_0 and B_0 .

Ensure: Optimal basis D_n .

- 1: **for** t = 1 to n **do**
- 2: Access the t-th sample z_t and the t-th atom y_t .
- 3: Compute the coefficient and noise:

$$\{\boldsymbol{v}_{t}^{*}, \boldsymbol{e}_{t}^{*}\} = \underset{\boldsymbol{v}, \boldsymbol{e}}{\operatorname{arg \, min}} \, \tilde{\ell}(D_{t-1}, \boldsymbol{v}, \boldsymbol{e}; \boldsymbol{z}_{t}),$$
$$\boldsymbol{u}_{t}^{*} = \underset{\boldsymbol{u}}{\operatorname{arg \, min}} \, \tilde{\ell}_{2}(D_{t-1}, M_{t-1}, \boldsymbol{u}; \boldsymbol{y}_{t}).$$

4: Update the accumulation matrices:

$$M_{t} \leftarrow M_{t-1} + \boldsymbol{y}_{t}(\boldsymbol{u}_{t}^{*})^{\top},$$

$$A_{t} \leftarrow A_{t-1} + \boldsymbol{v}_{t}^{*}(\boldsymbol{v}_{t}^{*})^{\top},$$

$$B_{t} \leftarrow B_{t-1} + (\boldsymbol{z}_{t} - \boldsymbol{e}_{t}^{*})(\boldsymbol{v}_{t}^{*})^{\top}.$$

5: Update the basis:

$$D_t = \underset{D}{\operatorname{arg\,min}} \frac{1}{t} \left[\frac{1}{2} \operatorname{Tr} \left(D^{\top} D(\lambda_1 A_t + \lambda_3 I_d) \right) - \operatorname{Tr} \left(D^{\top} (\lambda_1 B_t + \lambda_3 M_t) \right) \right].$$

6: end for

Suppose that we store the accumulation matrix

$$M_{t-1} \stackrel{\text{def}}{=} \sum_{i=1}^{t-1} \boldsymbol{y}_i(\boldsymbol{u}_i^*)^\top \in \mathbb{R}^{p \times d}, \tag{17}$$

where we define M_0 as a zero matrix. At the t-th iteration, when a new atom y_t is revealed, we calculate u_t^* as the minimizer of $\tilde{\ell}_2(D_{t-1}, u; M_{t-1}, y_t)$, for which we have the closed form solution

$$\boldsymbol{u}_{t}^{*} = (\|\boldsymbol{y}_{t}\|_{2}^{2} + 1/\lambda_{3})^{-1} (D_{t-1} - M_{t-1})^{\top} \boldsymbol{y}_{t}.$$
(18)

The solution in (18) differs from that of (10) a lot. Though (18) is not an accurate solution that minimizes $\tilde{h}(D,U;Y)$, the computation is efficient. More importantly, \boldsymbol{u}_t^* only depends on \boldsymbol{y}_t rather than the entire atom dictionary. We will also show that (18) suffices to guarantee the convergence of the algorithm to a stationary point of the expected loss function.

To gain intuition on the connection between $\tilde{\ell}_2(D, \boldsymbol{u}; M, \boldsymbol{y})$ and $\tilde{h}(D, U; Y)$, assume we have n atoms in total (i.e., Y has n columns). It turns out that for the function $\tilde{h}(D, U; Y)$, U can be optimized using block coordinate minimization (BCM), with each column of U being a block variable. Initially, we may set all these columns to a zero vector. As BCM proceeds, we update \boldsymbol{u}_t while keeping the other \boldsymbol{u}_i 's for $i \neq t$. Note that in this way, optimizing $\tilde{h}(D, U; Y)$ over \boldsymbol{u}_t amounts to minimizing the function $\tilde{\ell}_2(D, \boldsymbol{u}; M_{t-1}, \boldsymbol{y}_t)$ where M_{t-1} is defined in (17). So after revealing all the atoms, each \boldsymbol{u}_t is sequentially updated only once. Henceforth, our strategy can be seen as a one-pass BCM algorithm for the function $\tilde{h}(D, U; Y)$.

2.2.3 Optimize *D*

As soon as $\{v_i^*, e_i^*, u_i^*\}_{i=1}^t$ are available, we can refine the initial guess D_{t-1} by optimizing the surrogate function

$$g_t(D) \stackrel{\text{def}}{=} \frac{1}{t} \left(\sum_{i=1}^t \tilde{\ell}(D, \boldsymbol{v}_i^*, \boldsymbol{e}_i^*; \boldsymbol{z}_i) + \sum_{i=1}^t \frac{1}{2} \|\boldsymbol{u}_i^*\|_2^2 + \frac{\lambda_3}{2} \|D - M_t\|_F^2 \right). \tag{19}$$

Let us look at the first term:

$$\tilde{g}_t(D) \stackrel{\text{def}}{=} \frac{1}{t} \sum_{i=1}^t \tilde{\ell}(D, \boldsymbol{v}_i^*, \boldsymbol{e}_i^*; \boldsymbol{z}_i), \tag{20}$$

which is a surrogate of $\frac{1}{t} \sum_{i=1}^{t} \ell(D; z_i)$. Though the above objective function involves the past iterates $\{v_i^*, e_i^*\}_{i=1}^t$ whose memory cost is proportional to the sample size, we show that the minimizer of (20) can be computed by accessing only two accumulation matrices whose sizes are independent of the sample size. To see this, we may expand the function $\tilde{\ell}(D, v_i^*, e_i^*; z_i)$ as follows:

$$\tilde{\ell}(D, \boldsymbol{v}_{i}^{*}, \boldsymbol{e}_{i}^{*}; \boldsymbol{z}_{i}) = \frac{\lambda_{1}}{2} \|\boldsymbol{z}_{i} - D\boldsymbol{v}_{i}^{*} - \boldsymbol{e}_{i}^{*}\|_{2}^{2} + \frac{1}{2} \|\boldsymbol{v}_{i}^{*}\|_{2}^{2} + \lambda_{2} \|\boldsymbol{e}_{i}^{*}\|_{1}$$

$$= \frac{\lambda_{1}}{2} \operatorname{Tr} \left(D^{\top} D \boldsymbol{v}_{i}^{*} (\boldsymbol{v}_{i}^{*})^{\top} \right) + \lambda_{1} \operatorname{Tr} \left(D^{\top} (\boldsymbol{e}_{i}^{*} - \boldsymbol{z}_{i}) (\boldsymbol{v}_{i}^{*})^{\top} \right)$$

$$+ \frac{\lambda_{1}}{2} \|\boldsymbol{z}_{i} - \boldsymbol{e}_{i}^{*}\|_{2}^{2} + \frac{1}{2} \|\boldsymbol{v}_{i}^{*}\|_{2}^{2} + \lambda_{2} \|\boldsymbol{e}_{i}^{*}\|_{1}.$$

Since the variable here is D, it holds that the minimizer of (20) is given by the following:

$$\min_{D} \frac{1}{t} \sum_{i=1}^{t} \left[\frac{\lambda_1}{2} \operatorname{Tr} \left(D^{\top} D \boldsymbol{v}_i^* (\boldsymbol{v}_i^*)^{\top} \right) + \lambda_1 \operatorname{Tr} \left(D^{\top} (\boldsymbol{e}_i^* - \boldsymbol{z}_i) (\boldsymbol{v}_i^*)^{\top} \right) \right]. \tag{21}$$

As the trace is a linear mapping, solving the above program only needs to record two accumulation matrices

$$A_t \stackrel{\text{def}}{=} \sum_{i=1}^t \boldsymbol{v}_i^* (\boldsymbol{v}_i^*)^\top \in \mathbb{R}^{d \times d}, \quad B_t \stackrel{\text{def}}{=} \sum_{i=1}^t (\boldsymbol{z}_i - \boldsymbol{e}_i^*) (\boldsymbol{v}_i^*)^\top \in \mathbb{R}^{p \times d}.$$
 (22)

Using the result of (21) and doing some calculation, we derive D_t as follows:

$$D_t = \underset{D}{\operatorname{arg min}} \frac{1}{t} \left[\frac{1}{2} \operatorname{Tr} \left(D^{\top} D(\lambda_1 A_t + \lambda_3 I_d) \right) - \operatorname{Tr} \left(D^{\top} (\lambda_1 B_t + \lambda_3 M_t) \right) \right]$$
$$= (\lambda_1 B_t + \lambda_3 M_t) (\lambda_1 A_t + \lambda_3 I_d)^{-1}, \tag{23}$$

where A_t and B_t are given in (22) and M_t is defined in (17). Numerically, we apply coordinate descent to solve the above program owing to its efficiency. See more details in Appendix A.

2.2.4 Memory Cost

It is remarkable that the memory cost of Algorithm 1 is O(pd). To see this, note that when solving v_t^* and e_t^* , we load D_{t-1} and a sample z_t into the memory, which costs O(pd). To compute the optimal u_t^* , we need to access D_{t-1} and $M_{t-1} \in \mathbb{R}^{p \times d}$. Although we aim to minimize (19), which seems to require all the past information, we actually only need to record A_t , B_t and M_t , whose sizes are at most O(pd) (recall that d < p).

2.2.5 Time Complexity

In addition to the memory efficiency, we further elaborate that the computation in each iteration is cheap. To compute $\{v_t^*, e_t^*\}$, one may utilize the block coordinate method in Richtárik and Takác (2014) which enjoys linear convergence due to strong convexity. One may also apply the stochastic variance reduced algorithms which also ensure a geometric rate of convergence (Xiao and Zhang, 2014; Defazio et al., 2014). The u_t^* is obtained by simple matrix-vector multiplications, which costs O(pd). It is easy to see the complexity of Step 4 is O(pd) and that of Step 5 is $O(pd^2)$.

2.2.6 A Fully Online Subspace Clustering Scheme

Now we have provided a way to learn the low-rank representation matrix X in an online manner. Usually, researchers in the literature will take an optional post-processing step to refine the segmentation accuracy, for example, applying spectral clustering (Ng et al., 2001) on the obtained representation matrix X^* . In this case, one has to collect all the u_i^* 's and v_i^* 's to compute $X^* = (U^*)^T V^*$ which will again increase the memory cost to $O(n^2)$. Here, we suggest an alternative scheme which admits O(kd) memory usage where k is the number of subspaces. The idea is appealing to the well-known k-means clustering in place of spectral clustering. One notable advantage is that updating the k-means model can be implemented in an online manner. In fact, the online k-means algorithm can be easily integrated into Algorithm 1 by observing that v_i^* is the robust feature for the ith sample. Second, we observe that v_i^* is actually a robust feature for the ith sample. On the other hand, updating the k-means model is quite cheap, since the computational cost is O(kd).

2.2.7 An Alternative Online Implementation

Our strategy for solving u_t is based on a carefully designed proxy function which resolves Issue (I4) and has a low complexity. Yet, to tackle Issue (I4), another potential way is to avoid the variable u_t^2 . Recall that we derive the optimal solution \tilde{U} (provided that D is given) to $\tilde{h}(D, U; Y)$ as follows (see Prop 1):

$$\tilde{U} = Y^{\top} \left(\lambda_3^{-1} I_p + Y Y^{\top} \right)^{-1} D.$$

Plugging it back to $\tilde{h}(D, U; Y)$, we obtain

$$\tilde{h}(D, U; Y) = \frac{1}{2} \operatorname{Tr} \left(DD^{\top} \left(Q_n - \lambda_3^{-1} Q_n^2 \right) \right) + \frac{\lambda_3}{2} \left\| D - \lambda_3^{-1} Q_n D \right\|_F^2,$$

where

$$Q_n = \left(\lambda_3^{-1} I_p + Y Y^{\top}\right)^{-1}.$$

Here, the subscript of Q_n denotes the number of atoms in Y. Note that the size of Q_n is $p \times p$. Hence, if we incrementally compute the accumulation matrix $YY^{\top} = \sum_{i=1}^{t} \boldsymbol{y}_i \boldsymbol{y}_i^{\top}$, we can update the variable D in an online fashion. Namely, at t-th iteration, we re-define the surrogate function as follows:

$$g_t(D) \stackrel{\text{def}}{=} \frac{1}{t} \left[\sum_{i=1}^t \tilde{\ell}(\boldsymbol{z}_i, D, \boldsymbol{v}_i, \boldsymbol{e}_i) + \frac{\lambda_3}{2} \left\| D - \frac{1}{\lambda_3} Q_t D \right\|_F^2 + \frac{1}{2} \operatorname{Tr} \left(D D^\top \left(Q_t - \frac{1}{\lambda_3} Q_t^2 \right) \right) \right].$$

²We'd like to thank the anonymous NIPS 2015 reviewer for pointing out this potential solution to the online algorithm.

Again, by noting the fact that $\tilde{\ell}(z_i, D, v_i, e_i)$ only involves recording A_t and B_t , we show that the memory cost is independent of sample size. However, the main shortcoming is that the time complexity of computing the inverse of a $p \times p$ matrix in each iteration is $O(p^3)$ which is not efficient in the high-dimensional regime (the time complexity of ours is proportional to pd^2). Either, it is not clear how to analyze the convergence.

3 Theoretical Analysis

In this section, we present theoretical evidence that our algorithm guarantees that the sequence $\{D_t\}_{t\geq 1}$ converges to a stationary point of the expected loss (9) asymptotically. We note that the problem studied here is non-convex, and hence convergence to a stationary point is the best one can hope in general (Bertsekas, 1999). As we have mentioned in related work, there are elegant results showing that convergence to global optimum is possible for batch alternating minimization under more stringent conditions on the data. See, for example, Jain et al. (2013); Netrapalli et al. (2014); Ge et al. (2016).

We make two assumptions throughout our analysis.

- (A1) The observed data are generated i.i.d. from some (unknown) distribution and there exist constants α_0 and α_1 , such that the conditions $0 < \alpha_0 \le \|\boldsymbol{z}_t\|_2 \le \alpha_1$ and $\alpha_0 \le \|\boldsymbol{y}_t\|_2 \le \alpha_1$ hold almost surely for all $t \ge 1$.
- (A2) The smallest singular value of $\frac{1}{t}A_t$ is bounded away from zero almost surely.

Note that the first assumption is very mild. In fact, in many applications the data points are normalized to unit length, and in this case $\alpha_0 = \alpha_1 = 1$. To understand the second assumption, recall that D_t is computed in (23). If $\lambda_3 = o(t)$, then asymptotically the solution D_t is not unique. Hence, Assumption (A2) ensures that we always have a unique solution that minimizes the surrogate function $g_t(D)$. Geometrically, as we have noted that v_i is the coefficient of z_i in terms of the basis dictionary D, the assumption simply requires that the data points are in general positions.

We also need the following parameter settings.

(P1) λ_1 and λ_2 are independent of the sample size n.

$$(P2) \lim_{n\to\infty} \lambda_3/n = 0.$$

As we have shown in Section 2, these parameter scalings give the expected loss function as in (13). We note that (P1) is also required in previous work (Mairal et al., 2010; Feng et al., 2013; Shen et al., 2014), though not set out explicitly. (P2) is specific to our problem. It facilitates the characterization of the empirical loss function when n tends to infinity. Otherwise, even the convergence of the sequence $\{f_n(D)\}_{n\geq 1}$ is not clear (consider, e.g., $\lambda_3 = n \sin n$).

We present a fundamental result that will be heavily invoked in the subsequent analysis.

Proposition 2. Let $\{u_t^*\}_{t\geq 1}$, $\{v_t^*\}_{t\geq 1}$, $\{e_t^*\}_{t\geq 1}$ and $\{D_t\}_{t\geq 1}$ be the sequence of the optimal solutions produced by Algorithm 1. Assume (A1) and (A2). Further suppose that λ_2 does not grow with t. Then, for all $t\geq 1$,

- 1. v_t^* , e_t^* , $\frac{1}{t}A_t$ and $\frac{1}{t}B_t$ are uniformly bounded from above;
- 2. M_t is uniformly upper bounded;
- 3. D_t is supported on some compact set \mathcal{D} ;
- 4. u_t^* is uniformly upper bounded.

Note that we prove the result by assuming a weaker condition than (P1): the parameter λ_2 does not grow with the sample size t. While it is not surprising to see that u_t^*, v_t^*, e_t^* and D_t are bounded from above due to the regularization, it is interesting to note that $M_t = \sum_{i=1}^t y_i (u_i^*)^{\top}$ is also upper bounded. Intuitively, this holds since the last term in (5) imposes that M_t cannot deviate far from D_t . However, the technical challenge is that D_t itself depends on M_t , as shown in (23), and neither D_t nor M_t is bounded without the boundedness of the other. To remedy this, we propose a novel technique which conducts mathematical induction simultaneously on D_t and M_t . This tremendously simplifies the proof of our earlier version (Shen et al., 2016). See Appendix B.3 for the proof.

The proposition has many implications. For example, the uniform boundedness of the solutions immediately implies that the surrogate function $g_t(D)$ and the empirical loss function $f_t(D)$ are bounded from above for all $D \in \mathcal{D}$, which is necessary for the convergence.

Next, we prove that the sequence of $\{g_t(D_t)\}_{t\geq 1}$ converges almost surely. We will make use of the following lemma, which is due to Bottou (1998).

Lemma 2. Let (Ω, \mathcal{F}, P) be a measurable probability space, ψ_t , for $t \ge 1$, be the realization of a stochastic process and \mathcal{F}_t be the filtration by the past information at time t. Let

$$\delta_t = \begin{cases} 1 & \text{if } \mathbb{E}[\psi_{t+1} - \psi_t \mid \mathcal{F}_t] > 0, \\ 0 & \text{otherwise.} \end{cases}$$

If for all t, $\psi_t \ge 0$ and $\sum_{t=1}^{\infty} \mathbb{E}[\delta_t(\psi_{t+1} - \psi_t)] < \infty$, then ψ_t is a quasi-martingale and converges almost surely. Moreover, it holds almost surely that

$$\sum_{t=1}^{\infty} |\mathbb{E}[\psi_{t+1} - \psi_t \mid \mathcal{F}_t]| < +\infty.$$

Based on the lemma, we show the following result.

Theorem 1. Assume (A1) and (A2). Set the parameters λ_1 , λ_2 and λ_3 such that they satisfy (P1) and (P2). Then the sequence of $\{g_t(D_t)\}_{t\geq 1}$ converges almost surely, where $\{D_t\}_{t\geq 1}$ is the solution produced by Algorithm 1. Moreover, the following holds almost surely:

$$\sum_{t=1}^{\infty} \left| \mathbb{E} \left[\psi_{t+1} - \psi_t \mid \mathcal{F}_t \right] \right| < +\infty, \tag{24}$$

where $\mathcal{F}_t = \{\boldsymbol{z}_i, \boldsymbol{y}_i\}_{i=1}^t$.

Proof. (Sketch) We will view $\{g_t(D_t)\}_{t\geq 1}$ as a non-negative stochastic process, with $\mathcal{F}_t = \{\boldsymbol{z}_i, \boldsymbol{y}_i\}_{i=1}^t$ be the filtration of the past information. In order to apply Lemma 2, we write $\psi_t := g_t(D_t)$ and compute the variation between two consecutive iterations:

$$\psi_{t+1} - \psi_{t} = \underbrace{g_{t+1}(D_{t+1}) - g_{t+1}(D_{t})}_{\zeta_{1}} + \underbrace{\underbrace{\frac{\hat{f}_{t}(D_{t}) - \tilde{g}_{t}(D_{t})}{t+1}}_{\zeta_{2}}}_{+ \underbrace{1}} + \underbrace{\underbrace{\frac{\ell(z_{t+1}, D_{t}) - \tilde{f}_{t}(D_{t})}{t+1}}_{\zeta_{3}}}_{+ \underbrace{1}} + \underbrace{\frac{1}{t+1} \sum_{i=1}^{t+1} \frac{1}{2} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} + \frac{\lambda_{3}}{2(t+1)} \|D_{t} - M_{t+1}\|_{F}^{2}}_{-\frac{1}{t} \sum_{i=1}^{t} \frac{1}{2} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} - \frac{\lambda_{3}}{2t} \|D_{t} - M_{t}\|_{F}^{2}}.$$

Here,

$$\tilde{f}_t(D_t) = \frac{1}{t} \sum_{i=1}^t \ell(D_t; \boldsymbol{z}_i).$$

Since D_{t+1} minimizes $g_{t+1}(D)$, we have $\zeta_1 \leq 0$. Also, recall that $\tilde{g}_t(D)$ is a surrogate of $\tilde{f}_t(D)$, implying $\zeta_2 \leq 0$. For ζ_3 , the P-Donsker lemma (see Prop. 5) gives

$$\mathbb{E}\big[\mathbb{E}[\zeta_3 \mid \mathcal{F}_t]\big] \le \frac{C}{\sqrt{t(t+1)}}$$

for some constant C. Finally, we need to upper bound the terms in the brackets which are specific to the online subspace clustering problem. In light of the bound of ζ_3 , one may look for a way to bound them with $O(1/t^{3/2})$. Surprisingly, it turns out that our algorithm automatically guarantees that the sum of the terms in the brackets is not greater than zero. This follows by noting the closed-form solution of u_i^* (18) and the fact that $M_t = \sum_{i=1}^t y_i(u_i^*)^\top$. Putting all the pieces together completes the proof.

We move on to show that the sequence of $\{f_t(D_t)\}_{t\geq 1}$ converges. In particular, we justify that $g_t(D)$ acts as a good surrogate of $f_t(D)$ in the sense that the sequence of the surrogate converges to the same limit of that of the empirical loss.

Theorem 2. Assume (A1) and (A2). Set the parameters λ_1 , λ_2 and λ_3 such that they satisfy (P1) and (P2). Let $\{D_t\}_{t\geq 1}$ be the solution produced by Algorithm 1. Then, the sequence of $\{f_t(D_t)\}_{t\geq 1}$ converges almost surely to the same limit of $\{g_t(D_t)\}_{t\geq 1}$.

Corollary 1. Assume sames conditions as in Theorem 2. Then the sequence of $\{f(D_t)\}_{t\geq 1}$ converges almost surely to the same limit of $\{f_t(D_t)\}_{t\geq 1}$ or equivalently, $\{g_t(D_t)\}_{t\geq 1}$.

Proof. (Sketch) We need several tools in the literature to prove this result. From a high level, since we have already shown that $\{g_t(D_t)\}_{t\geq 1}$ converges, we only need to deduce that the limit of $g_t(D_t) - f_t(D_t)$ is zero. A useful result for this purpose is stated below, which is borrowed from Mairal et al. (2010).

Lemma 3 (Lemma 8 in Mairal et al. (2010)). Let $\{a_t\}_{t\geq 1}$, $\{b_t\}_{t\geq 1}$ be two real sequences such that for all t, $a_t\geq 0$, $b_t\geq 0$, $\sum_{t=1}^{\infty}a_t=\infty$, $\sum_{t=1}^{\infty}a_tb_t<\infty$, there exists a scalar C>0, such that $|b_{t+1}-b_t|< C\cdot a_t$. Then, $\lim_{t\to\infty}b_t=0$.

In particular, we set $a_t = (t+1)^{-1}$ and $b_t = g_t(D_t) - f_t(D_t) \ge 0$. First, we verify that the sum of the infinite series $\{a_tb_t\}_{t\ge 1}$ is finite. By algebra, we obtain

$$\frac{b_t}{t+1} \le \frac{\tilde{g}_t(D_t) - \tilde{f}_t(D_t)}{t+1} + \frac{q_t(D_t)}{t+1},$$

where

$$q_t(D_t) = \frac{1}{t} \sum_{i=1}^{t} \frac{1}{2} \|\boldsymbol{u}_i^*\|_2^2 + \frac{\lambda_3}{2t} \|D_t - M_t\|_F^2.$$

Then by utilizing the uniform boundedness of D_t and M_t , it is possible to derive

$$\frac{b_t}{t+1} \le \frac{\ell(D_t; \boldsymbol{z}_{t+1}) - \tilde{f}_t(D_t)}{t+1} + \psi_t - \psi_{t+1} + \frac{C}{2t(t+1)}$$

for some absolute constant C. As we have shown in the proof sketch of Theorem 1, the first term scales as $O(1/t^{3/2})$, which combined with (24) imply that $\sum_{t=1}^{\infty} \frac{b_t}{t+1}$ is finite.

Next, we claim that

$$|b_{t+1} - b_t| \le \frac{\mathcal{C}_0}{t+1},$$

for some absolute constant C_0 . The follows heavily from Prop. 2 where we showed that all the variables are uniformly bounded, and hence all involved functions evaluated at these solutions are bounded from above. To be more concrete, using triangle inequality we get

$$|b_{t+1} - b_t| \le |g_{t+1}(D_{t+1}) - g_t(D_{t+1})| + |f_{t+1}(D_{t+1}) - f_t(D_{t+1})| + |g_t(D_{t+1}) - g_t(D_t)| + |f_t(D_{t+1}) - f_t(D_t)|.$$

The first two terms on the right-hand side are upper bounded by O(1/t) due to uniform boundedness. For the last two terms, we utilize the fact that $g_t(D)$ and $f_t(D)$ are both Lipschitz to show that they are bounded by $O(\|D_{t+1} - D_t\|_F)$ from above. The following proposition illustrates that the variation of D_{t+1} and D_t vanishes with the rate O(1/t), whose proof can be found in Appendix B.6.

Proposition 3. Assume (A1) and (A2). Further suppose that λ_1 and λ_2 do not grow with t. Let $\{D_t\}_{t\geq 1}$ be the basis sequence produced by Algorithm 1. Then,

$$||D_{t+1} - D_t||_F = O\left(\frac{1}{t}\right).$$

Thus, in allusion to Lemma 3, we complete the proof of Theorem 2. The corollary follows immediately because the central limit theorem asserts that $\sqrt{t}(f(D_t) - f_t(D_t))$ is upper bounded.

Finally, we show that asymptotically, D_t acts as a stationary point of the expected loss function.

Theorem 3. Assume (A1), (A2), (P1) and (P2). Let $\{D_t\}_{t=1}^{\infty}$ be the sequence of optimal bases produced by Algorithm 1. Then, the sequence converges to a stationary point of the expected loss function f(D) when t goes to infinity.

4 Experiments

This section gives empirical evidence that our algorithm OLRSC is fast and robust. We demonstrate by simulations that the solution is good enough. In fact, we find that after revealing all the data points, OLRSC is able to recover the true subspace even with grossly corrupted entries. Thus, it is interesting to study the statistical performance of OLRSC in the future work. We also illustrate that OLRSC is orders of magnitude faster than batch methods such as LRR and SSC.

4.1 Settings

Before presenting the empirical results, we first introduce the universal settings used throughout the section.

4.1.1 Baselines

For the subspace recovery task, we compare our algorithm with ORPCA (Feng et al., 2013), LRR (Liu et al., 2013) and PCP (Candès et al., 2011). For the subspace clustering task, we choose ORPCA, LRR and SSC (Elhamifar and Vidal, 2009) as the competitive baselines. Recently, Liu and Li (2014) improved the vanilla LRR by utilizing some low-rank matrix for Y. We denote this variant of LRR by LRR2 and accordingly, our algorithm equipped with such a basis dictionary Y is denoted as OLRSC2.

4.1.2 Evaluation Metric

We evaluate the fitness of the recovered subspaces D (with each column being normalized) and the ground truth L by the Expressed Variance (EV) (Xu et al., 2010):

$$EV(D, L) \stackrel{\text{def}}{=} \frac{\text{Tr}(DD^{\top}LL^{\top})}{\text{Tr}(LL^{\top})}.$$
 (25)

The value of EV is between 0 and 1, and a higher value means better recovery (EV = 1 means exact recovery).

The performance of subspace clustering is measured by clustering accuracy which is provided in the SSC toolkit. Its value also ranges in the interval [0, 1], and a higher value indicates a more accurate clustering.

4.1.3 Parameters

We set $\lambda_1=1$, $\lambda_2=1/\sqrt{p}$ and $\lambda_3=\sqrt{t/p}$, where t is the iteration counter. Note that the parameter settings satisfy (P1) and (P2). In particular, $\lim_{t\to\infty}\lambda_3/t=0$. We follow the default parameter setting for the baselines.

4.2 Subspace Recovery

4.2.1 Simulation Data

We use 4 disjoint subspaces $\{S_k\}_{k=1}^4 \subset \mathbb{R}^p$, whose bases are denoted by $\{L_k\}_{k=1}^4 \in \mathbb{R}^{p \times d_k}$. The clean data matrix $\bar{Z}_k \in \mathcal{S}_k$ is then produced by $\bar{Z}_k = L_k R_k^{\top}$, where $R_k \in \mathbb{R}^{n_k \times d_k}$. The entries of L_k 's and R_k 's are sampled i.i.d. from the normal distribution. Finally, the observed data matrix Z is generated by $Z = \bar{Z} + E$, where \bar{Z} is the column-wise concatenation of \bar{Z}_k 's followed by a random permutation, E is the sparse corruption whose ρ fraction entries are non-zero and follow an i.i.d. uniform distribution over [-2,2]. We independently conduct each experiment 10 times and report the averaged results.

4.2.2 Robustness

We illustrate by simulation results that OLRSC can effectively recover the underlying subspaces, confirming that D_t converges to the union of subspaces. For the two online algorithms OLRSC and ORPCA, We compute the EV after revealing all the samples. We examine the performance under different intrinsic dimension d_k 's and corruption ρ . To be more detailed, the d_k 's are varied from 0.01p to 0.1p with a step size 0.01p, and the ρ is from 0 to 0.5, with a step size 0.05.

The results are presented in Figure 1. The most intriguing observation is that OLRSC as an online algorithm outperforms its batch counterpart LRR! Such improvement may come from the explicit modeling for the basis, which makes OLRSC more informative than LRR. To fully understand the rationale behind this phenomenon is an important direction for future research. Notably, OLRSC consistently beats ORPCA (an online version of PCP), in that OLRSC takes into account that the data are produced by a union of small subspaces. While PCP works well for almost all scenarios, OLRSC degrades a little when addressing difficult cases (high rank and corruption). This is not surprising since Theorem 3 is based on asymptotic analysis and hence, we expect that OLRSC will converge to the true subspace after acquiring more samples.

4.2.3 Convergence Rate and Time Complexity

Now we test on a large dataset to show that our algorithm usually converges to the true subspace faster than ORPCA. We plot the EV curve against the number of samples in Figure 2. Firstly, when equipped




Figure 1: Subspace recovery under different intrinsic dimensions and corruptions. Brighter is better. We set p=100, $n_k=1000$ and $d=4d_k$. LRR and PCP are batch methods. OLRSC consistently outperforms ORPCA and even improves the performance of LRR. Compared to PCP, OLRSC is competitive in most cases and degrades a little for highly corrupted data, possibly due to the number of samples is not sufficient for its convergence.

with a proper matrix Y, OLRSC2 and LRR2 can always produce an exact recovery of the subspace as PCP does. When using the dataset itself for Y, OLRSC still converges to a favorable point after revealing all the samples. Compared to ORPCA, OLRSC is more robust and converges much faster for hard cases (see, e.g., $\rho=0.5$). Again, we note that in such hard cases, OLRSC outperforms LRR, which agrees with the observation in Figure 1.

We also illustrate the time complexity of the algorithms in the last panel of Figure 2. In short, our algorithms (OLRSC and OLRSC2) admit the lowest computational complexity for all cases. One may argue that PCP spends slightly less time than ours for a small ρ (0.01 and 0.1). However, we remark here that PCP utilizes a highly optimized C++ toolkit to boost computation while our algorithms are fully written in Matlab. We believe that ours will work more efficiently if properly optimized by, e.g., the blas routine. Another important message conveyed by the figure is that, OLRSC is always being orders of magnitude computationally more efficient than the batch method LRR, as well as producing comparable or even better solution.




Figure 2: Convergence rate and time complexity. A higher EV means better subspace recovery. We set p=1000, $n_k=5000$, $d_k=25$ and d=100. OLRSC always converges to or outperforms the batch counterpart LRR. For hard cases, OLRSC converges much faster than ORPCA. Both PCP and LRR2 achieve the best EV value. When equipped with the same dictionary as LRR2, OLRSC2 also well handles the highly corrupted data ($\rho=0.5$). Our methods are more efficient than the competitors but PCP when ρ is small, possibly because PCP utilizes a highly optimized C++ toolkit while ours are written in Matlab.

4.3 Subspace Clustering

4.3.1 Datasets

We examine the performance of subspace clustering on 5 realistic databases shown in Table 1, which can be downloaded from the LibSVM website. For MNIST, We randomly select 20 thousands samples to form MNIST-20K since we find it time consuming to run the batch methods on the entire database.

4.3.2 Standard Clustering Pipeline

In order to focus on the solution quality of different algorithms, we follow the standard pipeline which feeds X to a spectral clustering algorithm (Ng et al., 2001). To this end, we collect all the u's and v's produced by OLRSC to form the representation matrix $X = UV^{\top}$. For ORPCA, we use $R_0R_0^{\top}$ as the similarity matrix (Liu et al., 2013), where R_0 is the row space of $Z_0 = L_0\Sigma_0R_0^{\top}$ and Z_0 is the clean matrix recovered by ORPCA. We run our algorithm and ORPCA with 2 epochs so as to refine the coefficients (i.e., U and V in ours and R_0 in ORPCA). Note that for subspace clustering, this step is essential because the initial guess of D results in bad solutions of the coefficients at the beginning.

Table 1: Datasets for subspace clustering.

	#classes	#samples	#features
Mushrooms	2	8124	112
DNA	3	3186	180
Protein	3	24,387	357
USPS	10	9298	256
MNIST-20K	10	20,000	784

4.3.3 Fully Online Pipeline

As we discussed in Section 2.2, the (optional) spectral clustering procedure needs the similarity matrix X, making the memory proportional to n^2 . To tackle this issue, we proposed a fully online scheme where the key idea is performing k-means on V. Here, we examine the efficacy of this variant, which is called OLRSC-F.

4.3.4 Results

The results are recorded in Table 2, where the time cost of spectral clustering or k-means is not included so we can focus on comparing the efficiency of the algorithms themselves. Also note that we use the dataset itself as the dictionary Y because we find that an alternative choice of Y does not help too much on this task. For OLRSC and ORPCA, they require an estimation on the true rank. Here, we use 5k as such estimation where k is the number of classes of a dataset. Our algorithm significantly outperforms the two state-of-the-art methods LRR and SSC both in terms of accuracy and efficiency. One may argue that SSC is slightly better than OLRSC on Protein. Yet, it spends 1 hour while OLRSC only costs 25 seconds. Hence, SSC is not practical. Compared to ORPCA, OLRSC always identifies more correct samples as well as consumes comparable running time. For example, on the USPS dataset, OLRSC achieves the accuracy of 65.95% while that of ORPCA is 55.7%. Regarding the running time, OLRSC uses only 7 seconds more than ORPCA – same order of computational complexity, which agrees with the qualitative analysis in Section 2.2 and the one in Feng et al. (2013).

More interestingly, it shows that the k-means alternative (OLRSC-F) usually outperforms the spectral clustering pipeline. This suggests that perhaps for *robust* subspace clustering formulations, the simple k-means paradigm suffices to guarantee an appealing result. On the other hand, we report the running time of spectral clustering and k-means in Table 3. As expected, since spectral clustering computes SVD for an n-by-n similarity matrix, it is quite slow. In fact, it sometimes dominates the running time of the whole pipeline. In contrast, k-means is extremely fast and scalable, as it can be implemented in online fashion.

4.4 Influence of d

A key ingredient of our formulation is a factorization on the nuclear norm regularized matrix, which requires an estimation on the rank of the X (see (3)). Here we examine the influence of the choice of d (which plays as an upper bound of the true rank). We report both EV and clustering accuracy for different d under a range of corruptions. The simulation data are generated as in Section 4.2 and we set p=200, $n_k=1000$ and $d_k=10$. Since the four subspaces are disjoint, the true rank is 40.

From Figure 3, we observe that our algorithm cannot recover the true subspace if d is smaller than the true rank. On the other hand, when d is sufficiently large (at least larger than the true rank), our algorithm can perfectly estimate the subspace. This agrees with the results in Burer and Monteiro (2005) which says as long as d is large enough, any local minima is global optima. We also illustrate the influence of d on

Table 2: Clustering accuracy (%) and computational time (seconds in default). For each dataset, the first row indicates the accuracy and the second row the running time. For all the large-scale datasets, OLRSC (or OLRSC-F) has the highest clustering accuracy. Regarding the running time, our method spends comparable time as ORPCA (the fastest solver) does while dramatically improves the accuracy. Although SSC is slightly better than SSC on Protein, it consumes one hour while OLRSC takes 25 seconds.

	OLRSC	OLRSC-F	ORPCA	LRR	SSC
Mush-	85.09	89.36	65.26	58.44	54.16
rooms	8.78	8.78	8.30	46.82	32 min
DNA	67.11 2.58	83.08 2.58	53.11 2.09	44.01 23.67	52.23 3 min
Protein	43.30 24.66	43.94 24.66	40.22 22.90	40.31 921.58	44.27 65 min
USPS	65.95 33.93	70.29 33.93	55.70 27.01	52.98 257.25	47.58 50 min
MNIST- 20K	57.74 129	55.50 129	54.10 121	55.23 32 min	43.91 7 hours

Table 3: Time cost (seconds) of spectral clustering and k-means.

	Mushrooms	DNA	Protein	USPS	MNIST-20K
Spectral	295	18	7567	482	4402
k-means	2	6	5	19	91

subspace clustering. Generally speaking, OLRSC can consistently identify the cluster of the data points if d is sufficiently large. Interestingly, different from the subspace recovery task, here the requirement for d seems to be slightly relaxed. In particular, we notice that if we pick d as 20 (smaller than the true rank), OLRSC still performs well. Such relaxed requirement of d may benefit from the fact that the spectral clustering step can correct some wrong points as suggested by Soltanolkotabi et al. (2014).

5 Conclusion

In this paper, we have proposed an online algorithm termed OLRSC for subspace clustering, which dramatically reduces the memory cost of LRR from $O(n^2)$ to O(pd). One of the key techniques is an explicit basis modeling, which essentially renders the model more informative than LRR. Another important component is a non-convex reformulation of the nuclear norm. Combining these techniques allows OLRSC to simultaneously recover the union of the subspaces, identify the possible corruptions and perform subspace clustering. We have also established the theoretical guarantee that solutions produced by our algorithm converge to a stationary point of the expected loss function. Moreover, we have analyzed the time complexity and empirically demonstrated that our algorithm is computationally very efficient compared to competing baselines. Our extensive experimental study on synthetic and realistic datasets also illustrates the robustness of OLRSC. In a nutshell, OLRSC is an appealing algorithm in all three worlds: memory cost, computation and robustness.




Figure 3: **Examine the influence of** d. The true rank is 40. We experiment on $d=\{2,20,40,60,80,100,120,140,160,180\}.$

A Algorithm Details

Algorithm 2 Solving $oldsymbol{v}$ and $oldsymbol{e}$

Require: $D \in \mathbb{R}^{p \times d}$, $\boldsymbol{z} \in \mathbb{R}^p$, parameters λ_1 and λ_2

Ensure: Optimal v and e.

1: Set e = 0.

2: repeat

3: Update v:

$$\boldsymbol{v} = (D^{\top}D + \frac{1}{\lambda_1}I)^{-1}D^{\top}(\boldsymbol{z} - \boldsymbol{e}).$$

4: Update *e*:

$$e = S_{\lambda_2/\lambda_1}[z - Dv].$$

5: until convergence

Algorithm 3 Solving *D*

Require: $D \in \mathbb{R}^{p \times d}$ in the previous iteration, accumulation matrix M, A and B, parameters λ_1 and λ_3 . **Ensure:** Optimal D (updated).

1: Denote $\widehat{A} = \lambda_1 A + \lambda_3 I$ and $\widehat{B} = \lambda_1 B + \lambda_3 M$.

2: repeat

3: **for** j = 1 to d **do**

4: Update the jth column of D:

$$oldsymbol{d}_j \leftarrow oldsymbol{d}_j - rac{1}{\widehat{a}_{jj}} \left(D \widehat{oldsymbol{a}}_j - \widehat{oldsymbol{b}}_j
ight)$$

5: end for

6: until convergence

For Algorithm 2, we set a threshold $\epsilon = 10^{-3}$. Let $\{v', e'\}$ and $\{v'', e''\}$ be the two consecutive iterates. If the maximum of $\|v' - v''\|_2 / \|v'\|_2$ and $\|e' - e''\|_2 / \|e'\|_2$ is less than ϵ , then we stop Algorithm 2.

For Algorithm 3, we observe that a one-pass update on the dictionary D is enough for the final convergence of D, as we showed in the experiments. This is also observed in Mairal et al. (2010).

B Proofs

B.1 Technical Lemmas

We need several technical lemmas for our proof.

Lemma 4 (Corollary of Thm. 4.1 in Bonnans and Shapiro (1998)). Let $f : \mathbb{R}^p \times \mathbb{R}^q \to \mathbb{R}$. Suppose that for all $x \in \mathbb{R}^p$ the function $f(x, \cdot)$ is differentiable, and that f and $\nabla_{\boldsymbol{u}} f(x, \boldsymbol{u})$ are continuous on $\mathbb{R}^p \times \mathbb{R}^q$. Let $\boldsymbol{v}(\boldsymbol{u})$ be the optimal value function $\boldsymbol{v}(\boldsymbol{u}) = \min_{\boldsymbol{x} \in \mathcal{C}} f(\boldsymbol{x}, \boldsymbol{u})$, where \mathcal{C} is a compact subset of \mathbb{R}^p . Then $\boldsymbol{v}(\boldsymbol{u})$ is differentiable. Furthermore, if for $\boldsymbol{u}_0 \in \mathbb{R}^q$, $f(\cdot, \boldsymbol{u}_0)$ has unique minimizer \boldsymbol{x}_0 then $\boldsymbol{v}(\boldsymbol{u})$ is differentiable in \boldsymbol{u}_0 and $\nabla_{\boldsymbol{u}} \boldsymbol{v}(\boldsymbol{u}_0) = \nabla_{\boldsymbol{u}} f(\boldsymbol{x}_0, \boldsymbol{u}_0)$.

Lemma 5 (Corollary of Donsker theorem (van der Vaart, 2000)). Let $F = \{f_{\theta} : \mathcal{X} \to \mathbb{R}, \theta \in \Theta\}$ be a set of measurable functions indexed by a bounded subset Θ of \mathbb{R}^d . Suppose that there exists a constant K such that

$$|f_{\theta_1}(x) - f_{\theta_2}(x)| \le K \|\theta_1 - \theta_2\|_2$$

for every θ_1 and θ_2 in Θ and x in X. Then, F is P-Donsker. For any f in F, let us define $\mathbb{P}_n f$, $\mathbb{P} f$ and $\mathbb{G}_n f$ as

$$\mathbb{P}_n f = \frac{1}{n} \sum_{i=1}^n f(X_i), \ \mathbb{P} f = \mathbb{E}[f(X)], \ \mathbb{G}_n f = \sqrt{n}(\mathbb{P}_n f - \mathbb{P} f).$$

Let us also suppose that for all f, $\mathbb{P}f^2 < \delta^2$ and $\|f\|_{\infty} < M$ and that the random elements X_1, X_2, \cdots are Borel-measurable. Then, we have

$$\mathbb{E} \|\mathbb{G}\|_F = O(1),$$

where $\|\mathbb{G}\|_F = \sup_{f \in F} |\mathbb{G}_n f|$.

B.2 Proof of Prop. 1

Proof. The optimal solution U for (7) is given by the first order optimality condition:

$$\frac{\partial \tilde{h}(Y, D, U)}{\partial U} = U + \lambda_3 (UY^{\top} - D^{\top})Y = 0,$$

by which we have

$$\tilde{U} = D^{\top} Y \left(\frac{1}{\lambda_3} I_n + Y^{\top} Y \right)^{-1}$$
$$= D^{\top} \left(\frac{1}{\lambda_3} I_p + Y Y^{\top} \right)^{-1} Y.$$

Note that \tilde{u}_i is the *i*th column of \tilde{U} . So for each $i \in [n]$,

$$\tilde{m{u}}_i = D^{ op} \left(rac{1}{\lambda_3} I_p + Y Y^{ op}
ight)^{-1} m{y}_i.$$

Also, we have

$$Y\tilde{U}^{\top} = YY^{\top} \left(\frac{1}{\lambda_3} I_p + YY^{\top}\right)^{-1} D$$

$$= \left(\frac{1}{\lambda_3} I_p + YY^{\top} - \frac{1}{\lambda_3} I_p\right) \left(\frac{1}{\lambda_3} I_p + YY^{\top}\right)^{-1} D$$

$$= D - \frac{1}{\lambda_3} \left(\frac{1}{\lambda_3} I_p + YY^{\top}\right)^{-1} D.$$

Thus,

$$h(D;Y) = \frac{1}{2} \left\| D^{\top} \left(\frac{1}{\lambda_3} I_p + Y Y^{\top} \right)^{-1} Y \right\|_{F}^{2} + \frac{1}{2\lambda_3} \left\| \left(\frac{1}{\lambda_3} I_p + Y Y^{\top} \right)^{-1} D \right\|_{F}^{2}.$$

B.3 Proof of Prop. 2

Proof. Let us consider the optimization problem of solving v and e. As the trivial solution $\{v'_t, e'_t\} = \{0, 0\}$ is feasible, we have

$$\tilde{\ell}_1(D_{t-1}, \boldsymbol{v}_t', \boldsymbol{e}_t'; \boldsymbol{z}_t) = \lambda_2 \|\boldsymbol{z}_t\|_1.$$

Therefore, the optimal solution should satisfy:

$$\frac{\lambda_1}{2} \| \boldsymbol{z}_t - D_{t-1} \boldsymbol{v}_t^* - \boldsymbol{e}_t^* \|_2^2 + \frac{1}{2} \| \boldsymbol{v}_t^* \|_2^2 + \lambda_2 \| \boldsymbol{e}_t^* \|_1 \le \lambda_2 \| \boldsymbol{z}_t \|_1,$$

which implies

$$\frac{1}{2} \left\| \bm{v}_t^* \right\|_2^2 \leq \lambda_2 \left\| \bm{z}_t \right\|_1, \ \lambda_2 \left\| \bm{e}_t^* \right\|_1 \leq \lambda_2 \left\| \bm{z}_t \right\|_1.$$

Since z_t is uniformly bounded (Assumption (A1)) and λ_2 does not grow with t, v_t^* and e_t^* are both uniformly bounded from above.

To examine the uniform boundedness for $\frac{1}{t}A_t$ and $\frac{1}{t}B_t$, note that

$$egin{aligned} & rac{1}{t}A_t = rac{1}{t}\sum_{i=1}^t oldsymbol{v}_i^*(oldsymbol{v}_i^*)^ op, \ & rac{1}{t}B_t = rac{1}{t}\sum_{i=1}^t \left(oldsymbol{z}_i - oldsymbol{e}_i^*
ight)(oldsymbol{v}_i^*)^ op. \end{aligned}$$

Since for each i, v_i^* , e_i^* and z_i are uniformly bounded from above, $\frac{1}{t}A_t$ and $\frac{1}{t}B_t$ are uniformly upper bounded.

Now we derive the bound for D_t and M_t . We inductively show that both the sequences $\{D_t\}_{t\geq 1}$ and $\{M_t\}_{t\geq 1}$ are uniformly bounded. First, let us denote the upper bound of $\frac{1}{t}B_t$ by C_2 , i.e.,

$$\left\| \frac{1}{t} B_t \right\| \le C_2, \ \forall \ t \ge 1. \tag{26}$$

Also, Assumption (A2) indicates that there exists an absolute constant C_3 , such that

$$\left\| \frac{1}{t} A_t \right\| \ge C_3, \ \forall \ t \ge 1. \tag{27}$$

Now suppose that for all $1 \le i \le t - 1$, it holds for some absolute constant $C_1 > C_2/C_3$ that

$$||D_i|| \le C_1, \quad ||M_i|| \le C_1.$$

Using the closed form solution of u_t^* , we have

$$M_{t} = M_{t-1} + \boldsymbol{y}_{t}(\boldsymbol{u}_{t}^{*})^{\top}$$

$$= M_{t-1} + \|\boldsymbol{y}_{t}\|_{2}^{2} \left(\|\boldsymbol{y}_{t}\|_{2}^{2} + \frac{1}{\lambda_{3}}\right)^{-1} (D_{t-1} - M_{t-1})$$

$$= \frac{\|\boldsymbol{y}_{t}\|_{2}^{2}}{\|\boldsymbol{y}_{t}\|_{2}^{2} + \frac{1}{\lambda_{3}}} D_{t-1} + \frac{\lambda_{3}^{-1}}{\|\boldsymbol{y}_{t}\|_{2}^{2} + \frac{1}{\lambda_{3}}} M_{t-1}.$$

Hence,

$$||M_t|| \le \frac{||\boldsymbol{y}_t||_2^2}{||\boldsymbol{y}_t||_2^2 + \lambda_3^{-1}} ||D_{t-1}|| + \frac{\lambda_3^{-1}}{||\boldsymbol{y}_t||_2^2 + \lambda_3^{-1}} ||M_{t-1}|| \le C_1.$$

Now using the closed form solution of D_t , we have

$$D_t = (\lambda_1 B_t + \lambda_3 M_t)(\lambda_1 A_t + \lambda_3 I_d)^{-1} = \left(\frac{\lambda_1}{t} B_t + \frac{\lambda_3}{t} M_t\right) \left(\frac{\lambda_1}{t} A_t + \frac{\lambda_3}{t} I_d\right)^{-1}.$$

Combining the above with (26) and (27) gives us

$$||D_t|| \le \left(\lambda_1 C_2 + \frac{\lambda_3}{t} C_1\right) \left(\lambda_1 C_3 + \frac{\lambda_3}{t}\right)^{-1} = \frac{\lambda_3}{\lambda_1 C_3} \frac{C_1 - C_2/C_3}{t + \frac{\lambda_3}{\lambda_1 C_2}} + \frac{C_2}{C_3}.$$

It turns out that the maximum of the right hand side is attained at t = 1 due to our earlier choice $C_1 > C_2/C_3$. Hence, we have

$$||D_t|| \leq C_1.$$

The induction is complete.

By examining the closed form of u_t^* , and note that we have proved the uniform boundedness of D_t and M_t , we conclude that u_t^* is uniformly upper bounded.

Corollary 2. Assume same conditions as in Prop. 2. Further suppose that λ_1 does not grow with t. Then, for all $t \geq 1$,

- 1. $\tilde{\ell}(D_t, v_t^*, e_t^*; z_t)$ and $\ell(D_t; z_t)$ are uniformly bounded from above.
- 2. $\frac{1}{t}\tilde{h}(D_t, U_{1:t}^*; Y_{1:t})$ is uniformly upper bounded where $U_{1:t}^* = (\boldsymbol{u}_1^*, \dots, \boldsymbol{u}_t^*)$ and $Y_{1:t} = (\boldsymbol{y}_1, \dots, \boldsymbol{y}_t)$.
- 3. The surrogate function $g_t(D_t)$ defined in (19) is uniformly upper bounded and Lipschitz.

Proof. To show Claim 1, we just need to examine the definition of $\tilde{\ell}(D_t, v_t^*, e_t^*; z_t)$ (see Eq. (6)) and notice that z_t , D_t , v_t^* and e_t^* are all uniformly bounded. This implies that $\tilde{\ell}(D_t, v_t^*, e_t^*; z_t)$ is uniformly bounded and so is $\ell(D_t; z_t)$. Likewise, we prove that $\frac{1}{t}\tilde{h}(D_t, U_{1:t}^*; Y_{1:t})$ is uniformly bounded. The uniform boundedness of $g_t(D_t)$ follows immediately.

To show that $g_t(D)$ is Lipschitz, we show that the gradient of $g_t(D)$ is uniformly bounded for all $D \in \mathcal{D}$.

$$\|\nabla g_{t}(D)\|_{F} = \left\|\lambda_{1} D\left(\frac{1}{t} A_{t} + \frac{\lambda_{3}}{t} I_{d}\right) - \frac{\lambda_{1}}{t} B_{t} - \frac{\lambda_{3}}{t} M_{t}\right\|_{F}$$

$$\leq \lambda_{1} \|D\|_{F} \left(\left\|\frac{1}{t} A_{t}\right\|_{F} + \left\|\frac{\lambda_{3}}{t} I_{d}\right\|_{F}\right) + \lambda_{1} \left\|\frac{1}{t} B_{t}\right\|_{F} + \left\|\frac{\lambda_{3}}{t} M_{t}\right\|_{F}.$$

Notice that each term on the right hand side of the inequality is uniformly bounded from above and λ_1 does not grow with t. Thus the gradient of $g_t(D)$ is uniformly bounded, implying that $g_t(D)$ is Lipschitz. \square

Proposition 4. Let $D \in \mathcal{D}$ and denote the minimizer of $\tilde{\ell}(D, \boldsymbol{v}, \boldsymbol{e}; \boldsymbol{z})$ as:

$$\{v', e'\} = \underset{v,e}{\operatorname{arg min}} \tilde{\ell}(D, v, e; z).$$

Then, the function $\ell(D; z)$ is continuously differentiable and

$$\nabla_D \ell(D; \boldsymbol{z}) = (D\boldsymbol{v}' + \boldsymbol{e}' - \boldsymbol{z})(\boldsymbol{v}')^{\top}.$$

Furthermore, $\ell(D; z)$ is uniformly Lipschitz.

Proof. By fixing z, the function $\tilde{\ell}$ can be seen as a mapping:

$$\mathbb{R}^{d+p} \times \mathcal{D} \to \mathbb{R}$$
$$([\boldsymbol{v}; \, \boldsymbol{e}], D) \mapsto \tilde{\ell}(D, \boldsymbol{v}, \boldsymbol{e}; \boldsymbol{z}).$$

It is easy to show that for all $[v; e] \in \mathbb{R}^{d+p}$, $\tilde{\ell}(D, v, e; z)$ is differentiable with respect to D. Also $\tilde{\ell}(D, v, e; z)$ is continuous on $\mathbb{R}^{d+p} \times \mathcal{D}$ and so is its gradient $\nabla_D \tilde{\ell}(D, v, e; z) = (Dv + e - z)v^{\top}$. For all $D \in \mathcal{D}$, since $\tilde{\ell}(D, v, e; z)$ is strongly convex w.r.t. v and e, it has a unique minimizer $\{v', e'\}$. Thus Lemma 4 applies and we prove that $\ell(D; z)$ is differentiable in D and

$$\nabla_D \ell(D; \boldsymbol{z}) = (D\boldsymbol{v}' + \boldsymbol{e}' - \boldsymbol{z})(\boldsymbol{v}')^{\top}.$$

Since every term in $\nabla_D \ell(D; z)$ is uniformly bounded (Assumption (A1) and Proposition 2), we conclude that the gradient of $\ell(D; z)$ is uniformly bounded, implying that $\ell(z, D)$ is uniformly Lipschitz w.r.t. D. \square

Corollary 3. Let $f_t(D)$ be the empirical loss function defined in (9). Then $f_t(D)$ is uniformly bounded from above and Lipschitz for all $t \ge 1$ and $D \in \mathcal{D}$.

Proof. Since $\ell(D; \mathbf{z})$ can be uniformly bounded (Corollary 2), we only need to show that $\frac{1}{t}h(D; Y)$ is uniformly bounded, where $Y = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_t)$. Note that we have derived the form for h(D; Y) in Prop. 1:

$$\frac{1}{t}h(D;Y) = \frac{1}{2t}\sum_{i=1}^{t} \left\| D^{\top} \left(\frac{1}{\lambda_3} I_p + YY^{\top} \right)^{-1} \boldsymbol{y}_i \right\|_2^2 + \frac{1}{2\lambda_3 t} \left\| \left(\frac{1}{\lambda_3} I_p + YY^{\top} \right)^{-1} D \right\|_F^2.$$

Since every term in the above equation can be uniformly bounded, $\frac{1}{t}h(D; Z_t)$ is uniformly bounded and so is $f_t(D)$.

To show that $f_t(D)$ is uniformly Lipschitz, it amounts to prove that its gradient can be uniformly bounded from above. Using Prop. 4, we have

$$\nabla f_t(D) = \frac{1}{t} \sum_{i=1}^t \nabla \ell(D; \boldsymbol{z}_i) + \frac{1}{t} \nabla h(D; \boldsymbol{Z}_t)$$

$$= \frac{1}{t} \sum_{i=1}^t (D \boldsymbol{v}_i^* + \boldsymbol{e}_i^* - \boldsymbol{z}_i) (\boldsymbol{v}_i^*)^\top$$

$$+ \frac{1}{t} \sum_{i=1}^t \left(\frac{1}{\lambda_3} I_p + Y Y^\top \right)^{-1} \boldsymbol{y}_i \boldsymbol{y}_i^\top \left(\frac{1}{\lambda_3} I_p + Y Y^\top \right)^{-1} D$$

$$+ \frac{\lambda_3}{t} \left(\frac{1}{\lambda_3} I_p + Y Y^\top \right)^{-2} D.$$

Then the Frobenius norm of $\nabla f_t(D)$ can be bounded by:

$$\begin{split} \|\nabla f_t(D)\|_F &\leq \frac{1}{t} \sum_{i=1}^t \|D \boldsymbol{v}_i^* + \boldsymbol{e}_i^* - \boldsymbol{z}_i\|_2 \cdot \|\boldsymbol{v}_i^*\|_2 \\ &+ \frac{1}{t} \sum_{i=1}^t \left\| \left(\frac{1}{\lambda_3} I_p + YY^\top \right)^{-1} \right\|_F^2 \cdot \|\boldsymbol{y}_i\|_2^2 \cdot \|D\|_F \\ &+ \frac{\lambda_3}{t} \left\| \left(\frac{1}{\lambda_3} I_p + YY^\top \right)^{-1} \right\|_F^2 \cdot \|D\|_F \,. \end{split}$$

One can easily check that the right hand side of the inequality is uniformly bounded from above. Thus $\|\nabla f_t(D)\|_F$ is uniformly bounded, implying that $f_t(D)$ is uniformly Lipschitz.

B.4 Proof of P-Donsker

Proposition 5. Let $\tilde{f}_t(D) = \frac{1}{t} \sum_{i=1}^t \ell(D; \boldsymbol{z}_i)$. Then we have

$$\mathbb{E}[\sqrt{t} \left\| \tilde{f}_t - \mathbb{E}_{\boldsymbol{z}}[\ell(D; \boldsymbol{z})] \right\|_{\infty}] = O(1).$$

Proof. Let us consider $\{\ell(D; z)\}$ as a set of measurable functions indexed by $D \in \mathcal{D}$. As we showed in Proposition 2, \mathcal{D} is a compact set. Also, we have proved that $\ell(D; z)$ is uniformly Lipschitz over D (Proposition 4). Thus, $\{\ell(D; z)\}$ is P-Donsker (see the definition in Lemma 5). Furthermore, as $\ell(D; z)$ is non-negative and its magnitude is uniformly upper bounded (Corollary 2), so is $\ell^2(D; z)$. Hence we have $\mathbb{E}_z[\ell^2(D; z)] \leq c$ for some absolute constant c. Note that we have verified all the hypotheses in Lemma 5. Hence the proof is complete.

B.5 Proof of Theorem 1

Proof. Note that $g_t(D_t)$ can be viewed as a stochastic positive process since every term in $g_t(D_t)$ is non-negative and the samples are drawn randomly. We define for all $t \ge 1$

$$\psi_t \stackrel{\text{def}}{=} g_t(D_t).$$

To show the convergence of ψ_t , we need to bound the difference of ψ_{t+1} and ψ_t :

$$\psi_{t+1} - \psi_{t} = g_{t+1}(D_{t+1}) - g_{t}(D_{t})
= g_{t+1}(D_{t+1}) - g_{t+1}(D_{t}) + g_{t+1}(D_{t}) - g_{t}(D_{t})
= g_{t+1}(D_{t+1}) - g_{t+1}(D_{t}) + \frac{1}{t+1}\ell(\mathbf{z}_{t+1}, D_{t}) - \frac{1}{t+1}\tilde{g}_{t}(D_{t})
+ \left[\frac{1}{t+1} \sum_{i=1}^{t+1} \frac{1}{2} \|\mathbf{u}_{i}^{*}\|_{2}^{2} + \frac{\lambda_{3}}{2(t+1)} \|D_{t} - M_{t+1}\|_{F}^{2} \right]
- \frac{1}{t} \sum_{i=1}^{t} \frac{1}{2} \|\mathbf{u}_{i}^{*}\|_{2}^{2} - \frac{\lambda_{3}}{2t} \|D_{t} - M_{t}\|_{F}^{2} \right]
= g_{t+1}(D_{t+1}) - g_{t+1}(D_{t}) + \frac{\tilde{f}_{t}(D_{t}) - \tilde{g}_{t}(D_{t})}{t+1} + \frac{\ell(\mathbf{z}_{t+1}, D_{t}) - \tilde{f}_{t}(D_{t})}{t+1}
+ \left[\frac{1}{t+1} \sum_{i=1}^{t+1} \frac{1}{2} \|\mathbf{u}_{i}^{*}\|_{2}^{2} + \frac{\lambda_{3}}{2(t+1)} \|D_{t} - M_{t+1}\|_{F}^{2} \right]
- \frac{1}{t} \sum_{i=1}^{t} \frac{1}{2} \|\mathbf{u}_{i}^{*}\|_{2}^{2} - \frac{\lambda_{3}}{2t} \|D_{t} - M_{t}\|_{F}^{2} \right].$$
(28)

Here,

$$\tilde{g}_t(D_t) = \frac{1}{t} \sum_{i=1}^t \tilde{\ell}(D, \boldsymbol{v}_i, \boldsymbol{e}_i; \boldsymbol{z}_i),$$

and

$$\tilde{f}_t(D_t) = \frac{1}{t} \sum_{i=1}^t \ell(D_t; \boldsymbol{z}_i).$$

First, we bound the four terms in the brackets of (28). We have

$$\frac{1}{t+1} \sum_{i=1}^{t+1} \frac{1}{2} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} - \frac{1}{t} \sum_{i=1}^{t} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} = \frac{-1}{t(t+1)} \sum_{i=1}^{t} \frac{1}{2} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} + \frac{1}{2(t+1)} \|\boldsymbol{u}_{t+1}^{*}\|_{2}^{2}
\leq \frac{1}{2(t+1)} \|\boldsymbol{u}_{t+1}^{*}\|_{2}^{2},$$
(29)

and

$$\frac{\lambda_{3}}{2(t+1)} \|D_{t} - M_{t+1}\|_{F}^{2} - \frac{\lambda_{3}}{2t} \|D_{t} - M_{t}\|_{F}^{2}
= \frac{-\lambda_{3}}{2t(t+1)} \|D_{t} - M_{t}\|_{F}^{2} + \frac{\lambda_{3}}{2(t+1)} \|\mathbf{z}_{t+1}(\mathbf{u}_{t+1}^{*})^{\top}\|_{F}^{2}
- \frac{\lambda_{3}}{t+1} \operatorname{Tr} \left((D_{t} - M_{t})^{\top} \mathbf{z}_{t+1}(\mathbf{u}_{t+1}^{*})^{\top} \right)
= \frac{-\lambda_{3}}{2t(t+1)} \|D_{t} - M_{t}\|_{F}^{2} + \frac{\lambda_{3}}{2(t+1)} \|\mathbf{z}_{t+1}(\mathbf{u}_{t+1}^{*})^{\top}\|_{F}^{2}
- \frac{\lambda_{3}}{t+1} \left(\|\mathbf{z}_{t+1}\|_{2}^{2} + \frac{1}{\lambda_{3}} \right) \|\mathbf{u}_{t+1}^{*}\|_{2}^{2}
\leq \frac{1}{t+1} \left(\frac{\lambda_{3}}{2} \|\mathbf{z}_{t+1}(\mathbf{u}_{t+1}^{*})^{\top}\|_{F}^{2} - (\lambda_{3} \|\mathbf{z}_{t+1}\|_{2}^{2} + 1) \|\mathbf{u}_{t+1}^{*}\|_{2}^{2} \right)
\leq \frac{1}{t+1} \left(-\frac{\lambda_{3}}{2} \|\mathbf{z}_{t+1}\|_{2}^{2} \|\mathbf{u}_{t+1}^{*}\|_{2}^{2} - \|\mathbf{u}_{t+1}^{*}\|_{2}^{2} \right), \tag{30}$$

where the first equality is derived by the fact that $M_{t+1} = M_t + z_{t+1}(u_{t+1}^*)^{\top}$, and the second equality is derived by the closed form solution of u_{t+1}^* (see (18)).

Combining (29) and (30), we know that

$$\frac{1}{t+1} \sum_{i=1}^{t+1} \frac{1}{2} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} - \frac{1}{t} \sum_{i=1}^{t} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2}
+ \frac{\lambda_{3}}{2(t+1)} \|D_{t} - M_{t+1}\|_{F}^{2} - \frac{\lambda_{3}}{2t} \|D_{t} - M_{t}\|_{F}^{2}
\leq \frac{1}{2(t+1)} \|\boldsymbol{u}_{t+1}^{*}\|_{2}^{2} + \frac{1}{t+1} \left(-\frac{\lambda_{3}}{2} \|\boldsymbol{z}_{t+1}\|_{2}^{2} \|\boldsymbol{u}_{t+1}^{*}\|_{2}^{2} - \|\boldsymbol{u}_{t+1}^{*}\|_{2}^{2} \right)
= \frac{1}{t+1} \left(-\frac{\lambda_{3}}{2} \|\boldsymbol{z}_{t+1}\|_{2}^{2} \|\boldsymbol{u}_{t+1}^{*}\|_{2}^{2} - \frac{1}{2} \|\boldsymbol{u}_{t+1}^{*}\|_{2}^{2} \right) \leq 0.$$

Therefore,

$$\psi_{t+1} - \psi_t \leq g_{t+1}(D_{t+1}) - g_{t+1}(D_t) + \frac{1}{t+1}\ell(\boldsymbol{z}_{t+1}, D_t) - \frac{1}{t+1}\tilde{g}_t(D_t)$$

$$= g_{t+1}(D_{t+1}) - g_{t+1}(D_t) + \frac{\tilde{f}_t(D_t) - \tilde{g}_t(D_t)}{t+1} + \frac{\ell(\boldsymbol{z}_{t+1}, D_t) - \tilde{f}_t(D_t)}{t+1}$$

$$\leq \frac{\ell(\boldsymbol{z}_{t+1}, D_t) - \tilde{f}_t(D_t)}{t+1},$$

where the last inequality holds because D_{t+1} is the minimizer of $g_{t+1}(D)$ and $\tilde{g}_t(D)$ is a surrogate function of $\tilde{f}_t(D)$.

Let \mathcal{F}_t be the filtration of the past information. We take the expectation on the above equation conditional on \mathcal{F}_t :

$$\mathbb{E}[\psi_{t+1} - \psi_t \mid \mathcal{F}_t] \leq \frac{\mathbb{E}[\ell(\boldsymbol{z}_{t+1}, D_t) \mid \mathcal{F}_t] - \tilde{f}_t(D_t)}{t+1}$$
$$\leq \frac{f(D_t) - \tilde{f}_t(D_t)}{t+1}$$
$$\leq \frac{\left\| f - \tilde{f}_t \right\|_{\infty}}{t+1}.$$

From Proposition 5, we know

$$\mathbb{E}\Big[\left\|f - \tilde{f}_t\right\|_{\infty}\Big] = O\left(\frac{1}{\sqrt{t}}\right).$$

Thus,

$$\mathbb{E}\left[\mathbb{E}[\psi_{t+1} - \psi_t \mid \mathcal{F}_t]^+\right] = \mathbb{E}\left[\max\{\mathbb{E}[\psi_{t+1} - \psi_t \mid \mathcal{F}_t], 0\}\right] \le \frac{c}{\sqrt{t(t+1)}},\tag{31}$$

where c is some constant.

Now let us define the index set

$$\mathcal{T} = \Big\{ t : \ t \ge 1, \mathbb{E} \big[\psi_{t+1} - \psi_t \mid \mathcal{F}_t \big] > 0 \Big\},\,$$

and the indicator function

$$\delta_t = \begin{cases} 1, & \text{if } t \in \mathcal{T}, \\ 0, & \text{otherwise.} \end{cases}$$

It follows that

$$\sum_{t=1}^{\infty} \mathbb{E}[\delta_t(\psi_{t+1} - \psi_t)] = \sum_{t \in \mathcal{T}} \mathbb{E}[\psi_{t+1} - \psi_t]$$

$$= \sum_{t \in \mathcal{T}} \mathbb{E}[\mathbb{E}[\psi_{t+1} - \psi_t \mid \mathcal{F}_t]]$$

$$= \sum_{t=1}^{\infty} \mathbb{E}[\mathbb{E}[\psi_{t+1} - \psi_t \mid \mathcal{F}_t]^+]$$

$$\leq + \infty,$$

where the last inequality holds in view of (31).

Thus, Lemma 2 applies. That is, $\{g_t(D_t)\}_{t\geq 1}$ is a quasi-martingale and converges almost surely. In addition,

$$\sum_{t=1}^{\infty} \left| \mathbb{E} \left[\psi_{t+1} - \psi_t \mid \mathcal{F}_t \right] \right| < +\infty, \ a.s.$$

B.6 Proof of Prop. 3

Proof. According the strong convexity of $g_t(D)$ (Assumption (A2)), we have,

$$g_t(D_{t+1}) - g_t(D_t) \ge \frac{\beta_0}{2} \|D_{t+1} - D_t\|_F^2,$$
 (32)

On the other hand,

$$g_{t}(D_{t+1}) - g_{t}(D_{t})$$

$$= g_{t}(D_{t+1}) - g_{t+1}(D_{t+1}) + g_{t+1}(D_{t+1}) - g_{t+1}(D_{t})$$

$$+ g_{t+1}(D_{t}) - g_{t}(D_{t})$$

$$\leq g_{t}(D_{t+1}) - g_{t+1}(D_{t+1}) + g_{t+1}(D_{t}) - g_{t}(D_{t}).$$
(33)

Note that the inequality is derived by the fact that $g_{t+1}(D_{t+1}) - g_{t+1}(D_t) \le 0$, as D_{t+1} is the minimizer of $g_{t+1}(D)$. Let

$$G_t(D) = g_t(D) - g_{t+1}(D).$$
 (34)

By a simple calculation, we obtain the gradient of $G_t(D)$:

$$\nabla G_{t}(D) = \nabla g_{t}(D) - \nabla g_{t+1}(D)$$

$$= \frac{1}{t} \Big[D \left(\lambda_{1} A_{t} + \lambda_{3} I_{d} \right) - \left(\lambda_{1} B_{t} + \lambda_{3} M_{t} \right) \Big]$$

$$- \frac{1}{t+1} \Big[D \left(\lambda_{1} A_{t+1} + \lambda_{3} I_{d} \right) - \left(\lambda_{1} B_{t+1} + \lambda_{3} M_{t+1} \right) \Big]$$

$$= \frac{1}{t} \Big[D \left(\lambda_{1} A_{t} + \lambda_{3} I_{d} - \frac{\lambda_{1} t}{t+1} A_{t+1} - \frac{\lambda_{3} t}{t+1} I_{d} \right)$$

$$+ \frac{\lambda_{1} t}{t+1} B_{t+1} - \lambda_{1} B_{t} + \frac{\lambda_{3} t}{t+1} M_{t+1} - \lambda_{3} M_{t} \Big]$$

$$= \frac{1}{t} \Big[D \left(\frac{\lambda_{1}}{t+1} A_{t+1} - \lambda_{1} v_{t+1} v_{t+1}^{\top} + \frac{\lambda_{3}}{t+1} I_{d} \right)$$

$$+ \lambda_{1} (z_{t+1} - e_{t+1}) v_{t+1}^{\top} - \frac{\lambda_{1}}{t+1} B_{t+1} + \lambda_{3} z_{t+1} u_{t+1}^{\top} - \frac{\lambda_{3}}{t+1} M_{t+1} \Big]$$

So the upper bound of the Frobenius norm of $\nabla G_t(D)$ follows immediately:

$$\begin{split} &\|\nabla G_{t}(D)\|_{F} \\ &\leq \frac{1}{t} \left[\|D\|_{F} \left(\lambda_{1} \left\| \frac{A_{t+1}}{t+1} \right\|_{F} + \lambda_{1} \left\| \boldsymbol{v}_{t+1} \boldsymbol{v}_{t+1}^{\top} \right\|_{F} + \frac{\lambda_{3}}{t+1} \|I_{d}\|_{F} \right) \\ &+ \lambda_{1} \left\| (\boldsymbol{z}_{t+1} - \boldsymbol{e}_{t+1}) \boldsymbol{v}_{t+1}^{\top} \right\|_{F} + \lambda_{1} \left\| \frac{B_{t+1}}{t+1} \right\|_{F} + \lambda_{3} \left\| \boldsymbol{z}_{t+1} \boldsymbol{u}_{t+1}^{\top} \right\|_{F} + \frac{\lambda_{3}}{t+1} \|M_{t+1}\|_{F} \right] \\ &= \frac{1}{t} \left[\|D\|_{F} \left(\lambda_{1} \left\| \frac{A_{t+1}}{t+1} \right\|_{F} + \lambda_{1} \left\| \boldsymbol{v}_{t+1} \boldsymbol{v}_{t+1}^{\top} \right\|_{F} \right) + \lambda_{1} \left\| (\boldsymbol{z}_{t+1} - \boldsymbol{e}_{t+1}) \boldsymbol{v}_{t+1}^{\top} \right\|_{F} \\ &+ \lambda_{1} \left\| \frac{B_{t+1}}{t+1} \right\|_{F} + \lambda_{3} \left\| \boldsymbol{z}_{t+1} \boldsymbol{u}_{t+1}^{\top} \right\|_{F} \right] + \frac{\lambda_{3}}{t(t+1)} \left[\|I_{d}\|_{F} + \|M_{t+1}\|_{F} \right]. \end{split}$$

We know from Proposition 2 that all the terms in the above equation are uniformly bounded from above. Thus, there exist constants c_1 , c_2 and c_3 , such that

$$\|\nabla G_t(D)\|_F \le \frac{1}{t} \left(c_1 \|D\|_F + c_2 \right) + \frac{c_3}{t}. \tag{35}$$

According to the first order Taylor expansion,

$$G_{t}(D_{t+1}) - G_{t}(D_{t})$$

$$= \operatorname{Tr} \left((D_{t+1} - D_{t})^{\top} \nabla G_{t} \left(\rho D_{t} + (1 - \rho) D_{t+1} \right) \right)$$

$$\leq \|D_{t+1} - D_{t}\|_{F} \cdot \|\nabla G_{t} \left(\rho D_{t} + (1 - \rho) D_{t+1} \right)\|_{F},$$

where ρ is some scalar between 0 and 1. According to Proposition 2, D_t and D_{t+1} are uniformly bounded, indicating that $\rho D_t + (1 - \rho) D_{t+1}$ is uniformly bounded. In view of (35), there exists a constant c_4 , such that

$$\|\nabla G_t \left(\alpha D_t + (1 - \alpha) D_{t+1}\right)\|_F \le \frac{c_4}{t},$$

resulting in

$$G_t(D_{t+1}) - G_t(D_t) \le \frac{c_4}{t} \cdot ||D_{t+1} - D_t||_F.$$

Combining (32), (33) and the above equation, we have

$$||D_{t+1} - D_t||_F = \frac{2c_4}{\beta_0 t}.$$

B.7 Proof of Theorem 2

Proof. Recall that $\tilde{f}_t(D) = \frac{1}{t} \sum_{i=1}^t \ell(D; \boldsymbol{z}_i)$ and $\tilde{g}_t(D) = \frac{1}{t} \sum_{i=1}^t \tilde{\ell}(D, \boldsymbol{v}_i^*, \boldsymbol{e}_i^*; \boldsymbol{z}_i)$. Define $b_t \stackrel{\text{def}}{=} g_t(D_t) - f_t(D_t)$

$$= \tilde{g}_{t}(D_{t}) - \tilde{f}_{t}(D_{t}) + \left[\frac{1}{t} \sum_{i=1}^{t} \frac{1}{2} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} + \frac{\lambda_{3}}{2t} \|D_{t} - M_{t}\|_{F}^{2} - \frac{1}{t} \sum_{i=1}^{t} \frac{1}{2} \|D_{t}^{\top} \left(\frac{1}{\lambda_{3}} I_{p} + YY^{\top}\right)^{-1} \boldsymbol{z}_{i} \|_{2}^{2} - \frac{1}{2\lambda_{3}} \left\| \left(\frac{1}{\lambda_{3}} I_{p} + YY^{\top}\right)^{-1} D_{t} \right\|_{F}^{2} \right]$$

$$= \tilde{g}_{t}(D_{t}) - \tilde{f}_{t}(D_{t}) + g_{t}(D_{t}),$$

where $q_t(D_t)$ denotes the four terms in the brackets. Using (28), we have

$$\frac{b_t}{t+1} = \frac{\tilde{g}_t(D_t) - \tilde{f}_t(D_t)}{t+1} + \frac{q_t(D_t)}{t+1}$$

$$= g_{t+1}(D_{t+1}) - g_{t+1}(D_t) + \frac{\ell(D_t; \mathbf{z}_{t+1}) - \tilde{f}_t(D_t)}{t+1} + \psi_t - \psi_{t+1}$$

$$+ \left[\frac{q_t(D_t)}{t+1} + \frac{1}{t+1} \sum_{i=1}^{t+1} \frac{1}{2} \|\mathbf{u}_i^*\|_2^2 + \frac{\lambda_3}{2(t+1)} \|D_t - M_{t+1}\|_F^2 \right]$$

$$- \frac{1}{t} \sum_{i=1}^{t} \frac{1}{2} \|\mathbf{u}_i^*\|_2^2 - \frac{\lambda_3}{2t} \|D_t - M_t\|_F^2 \right].$$

Note that it always holds that for some constant c,

$$\frac{q_t(D_t)}{t+1} \le \frac{1}{t(t+1)} \sum_{i=1}^t \frac{1}{2} \|\boldsymbol{u}_i^*\|_2^2 + \frac{\lambda_3}{2t(t+1)} \|D_t - M_t\|_F^2
\le \frac{1}{t(t+1)} \sum_{i=1}^t \frac{1}{2} \|\boldsymbol{u}_i^*\|_2^2 + \frac{c}{2t(t+1)},$$

where the second inequality is due to the fact that D_t and M_t are both uniformly bounded (see Proposition 2). On the other hand, from (29) we know that

$$\frac{1}{t+1} \sum_{i=1}^{t+1} \frac{1}{2} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} - \frac{1}{t} \sum_{i=1}^{t} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} = \frac{-1}{t(t+1)} \sum_{i=1}^{t} \frac{1}{2} \|\boldsymbol{u}_{i}^{*}\|_{2}^{2} + \frac{1}{2(t+1)} \|\boldsymbol{u}_{t+1}^{*}\|_{2}^{2},$$

while (30) implies

$$\frac{\lambda_3}{2(t+1)} \|D_t - M_{t+1}\|_F^2 - \frac{\lambda_3}{2t} \|D_t - M_t\|_F^2 \le \frac{1}{t+1} \left(-\frac{\lambda_3}{2} \|\boldsymbol{z}_{t+1}\|_2^2 \|\boldsymbol{u}_{t+1}^*\|_2^2 - \|\boldsymbol{u}_{t+1}^*\|_2^2 \right).$$

Combining these pieces, we have

$$\frac{q_{t}(D_{t})}{t+1} + \frac{1}{t+1} \sum_{i=1}^{t+1} \frac{1}{2} \| \boldsymbol{u}_{i}^{*} \|_{2}^{2} + \frac{\lambda_{3}}{2(t+1)} \| D_{t} - M_{t+1} \|_{F}^{2}
- \frac{1}{t} \sum_{i=1}^{t} \frac{1}{2} \| \boldsymbol{u}_{i}^{*} \|_{2}^{2} - \frac{\lambda_{3}}{2t} \| D_{t} - M_{t} \|_{F}^{2}
\leq \frac{c}{2t(t+1)} + \frac{1}{2(t+1)} \| \boldsymbol{u}_{t+1}^{*} \|_{2}^{2} + \frac{1}{t+1} \left(-\frac{\lambda_{3}}{2} \| \boldsymbol{z}_{t+1} \|_{2}^{2} \| \boldsymbol{u}_{t+1}^{*} \|_{2}^{2} - \| \boldsymbol{u}_{t+1}^{*} \|_{2}^{2} \right)
= \frac{c}{2t(t+1)} - \frac{1}{2(t+1)} \| \boldsymbol{u}_{t+1}^{*} \|_{2}^{2} - \frac{\lambda_{3}}{2(t+1)} \| \boldsymbol{z}_{t+1} \|_{2}^{2} \| \boldsymbol{u}_{t+1}^{*} \|_{2}^{2}
\leq \frac{c}{2t(t+1)}.$$

Therefore,

$$\frac{b_t}{t+1} \le g_{t+1}(D_{t+1}) - g_{t+1}(D_t) + \frac{\ell(D_t; \mathbf{z}_{t+1}) - \tilde{f}_t(D_t)}{t+1}
+ \psi_t - \psi_{t+1} + \frac{c}{2t(t+1)}
\le \frac{\ell(D_t; \mathbf{z}_{t+1}) - \tilde{f}_t(D_t)}{t+1} + \psi_t - \psi_{t+1} + \frac{c}{2t(t+1)},$$

where we use the fact that D_{t+1} minimizes $g_{t+1}(D)$ in the second inequality and we denote $\psi_t = g_t(D_t)$. By taking the expectation over z conditional on the past filtration \mathcal{F}_t , we have

$$\frac{b_t}{t+1} \le \frac{c_1}{\sqrt{t}(t+1)} + |\mathbb{E}[\psi_t - \psi_{t+1} \mid \mathcal{F}_t]| + \frac{c}{2t(t+1)},$$

which is an immediate result from Prop. 5. Thereby,

$$\sum_{t=1}^{\infty} \frac{b_t}{t+1} \le \sum_{t=1}^{\infty} \frac{c_1}{\sqrt{t(t+1)}} + \sum_{t=1}^{\infty} |\mathbb{E}[\psi_t - \psi_{t+1} \mid \mathcal{F}_t]| + \sum_{t=1}^{\infty} \frac{c}{2t(t+1)} < +\infty.$$

Here, the last inequality is derived by applying (24).

Next, we examine the difference between b_{t+1} and b_t :

$$|b_{t+1} - b_t| = |g_{t+1}(D_{t+1}) - f_{t+1}(D_{t+1}) - g_t(D_t) + f_t(D_t)|$$

$$\leq |g_{t+1}(D_{t+1}) - g_t(D_{t+1})| + |g_t(D_{t+1}) - g_t(D_t)|$$

$$+ |f_{t+1}(D_{t+1}) - f_t(D_{t+1})| + |f_t(D_{t+1}) - f_t(D_t)|.$$
(36)

For the first term on the right hand side, we have

$$\begin{aligned} &|g_{t+1}(D_{t+1}) - g_{t}(D_{t+1})| \\ &= \left| \tilde{g}_{t+1}(D_{t+1}) - \tilde{g}_{t}(D_{t+1}) + \frac{1}{t+1} \sum_{i=1}^{t+1} \frac{1}{2} \| \boldsymbol{u}_{i}^{*} \|_{2}^{2} - \frac{1}{t} \sum_{i=1}^{t} \frac{1}{2} \| \boldsymbol{u}_{i}^{*} \|_{2}^{2} \right. \\ &+ \frac{\lambda_{3}}{2(t+1)} \| D_{t+1} - M_{t+1} \|_{F}^{2} - \frac{\lambda_{3}}{2t} \| D_{t+1} - M_{t} \|_{F}^{2} \right| \\ &= \left| \tilde{g}_{t+1}(D_{t+1}) - \tilde{g}_{t}(D_{t+1}) - \frac{1}{t(t+1)} \sum_{i=1}^{t} \frac{1}{2} \| \boldsymbol{u}_{i}^{*} \|_{2}^{2} - \frac{1}{2(t+1)} \| \boldsymbol{u}_{t+1}^{*} \|_{2}^{2} \right. \\ &- \frac{\lambda_{3}}{2t(t+1)} \| D_{t+1} - M_{t} \|_{F}^{2} - \frac{\lambda_{3}}{2(t+1)} \| \boldsymbol{z}_{t+1}(\boldsymbol{u}_{t+1}^{*})^{\top} \|_{F}^{2} \right| \\ &\leq \left| \tilde{g}_{t+1}(D_{t+1}) - \tilde{g}_{t}(D_{t+1}) \right| + \frac{1}{t(t+1)} \sum_{i=1}^{t} \frac{1}{2} \| \boldsymbol{u}_{i}^{*} \|_{2}^{2} + \frac{1}{2(t+1)} \| \boldsymbol{u}_{t+1}^{*} \|_{2}^{2} \\ &+ \frac{\lambda_{3}}{2t(t+1)} \| D_{t+1} - M_{t} \|_{F}^{2} + \frac{\lambda_{3}}{2(t+1)} \| \boldsymbol{z}_{t+1}(\boldsymbol{u}_{t+1}^{*})^{\top} \|_{F}^{2} \\ &\stackrel{\zeta_{1}}{\leq} \left| \tilde{g}_{t+1}(D_{t+1}) - \tilde{g}_{t}(D_{t+1}) \right| + \frac{c_{1}}{t+1} \\ &= \left| \frac{1}{t+1} \ell(D_{t+1}; \boldsymbol{z}_{t+1}) - \frac{1}{t+1} \tilde{g}_{t}(D_{t+1}) \right| + \frac{c_{1}}{t+1} \\ &\stackrel{\zeta_{2}}{\leq} \frac{c_{2}}{t+1}, \end{aligned}$$

where c_1 and c_2 are some uniform constants. Note that ζ_1 holds because all the \boldsymbol{u}_t^* , D_{t+1} , M_t and \boldsymbol{z}_{t+1} are uniformly bounded (see Proposition 2), and ζ_2 holds because $\ell(\boldsymbol{z}_{t+1}, D_{t+1})$ and $\tilde{g}_t(D_{t+1})$ are uniformly bounded (see Corollary 2).

For the third term on the right hand side of (36), we can similarly show that

$$|f_{t+1}(D_{t+1}) - f_t(D_{t+1})| \le \left| \tilde{f}_{t+1}(D_{t+1}) - \tilde{f}_t(D_{t+1}) \right| + \frac{c_3}{t+1}$$

$$= \left| \frac{1}{t+1} \ell(D_{t+1}; \mathbf{z}_{t+1}) - \frac{1}{t+1} \tilde{f}_t(D_{t+1}) \right| + \frac{c_3}{t+1}$$

$$\stackrel{\zeta_3}{\le} \frac{c_4}{t+1},$$

where c_3 and c_4 are some uniform constants, and ζ_3 holds as $\ell(D_{t+1}; \boldsymbol{z}_{t+1})$ and $\tilde{f}_t(D_{t+1})$ are both uniformly bounded (see Corollary 3).

Using Corollary 2 and Corollary 3, we know that both $g_t(D)$ and $f_t(D)$ are uniformly Lipschitz. That

is, there exist uniform constants κ_1 , κ_2 , such that

$$|g_t(D_{t+1}) - g_t(D_t)| \le \kappa_1 \|D_{t+1} - D_t\|_F \le \frac{\zeta_4}{t+1},$$

$$|f_t(D_{t+1}) - f_t(D_t)| \le \kappa_2 \|D_{t+1} - D_t\|_F \le \frac{\zeta_5}{t+1}.$$

Here, ζ_4 and ζ_5 are derived by applying Prop. 3 and κ_3 and κ_4 are some uniform constants. Finally, we have a bound for (36):

$$|b_{t+1} - b_t| \le \frac{\kappa_0}{t+1},$$

where κ_0 is some uniform constant.

By applying Lemma 3, we conclude that $\{b_t\}_{t>1}$ converges to zero. That is,

$$\lim_{t \to +\infty} g_t(D_t) - f_t(D_t) = 0.$$

Since we have proved in Theorem 1 that $g_t(D_t)$ converges almost surely, we conclude that $f_t(D_t)$ converges almost surely to the same limit of $g_t(D_t)$.

B.8 Proof of Theorem 3

We need a technical result to prove Theorem 3.

Proposition 6. Let f(D) be the expected loss function which is defined in (13). Then, f(D) is continuously differentiable and $\nabla f(D) = \mathbb{E}_{\mathbf{z}}[\nabla_D \ell(D; \mathbf{z})]$. Moreover, $\nabla f(D)$ is uniformly Lipschitz on \mathcal{D} .

Proof. We have shown in Proposition 4 that $\ell(D; z)$ is continuously differentiable, f(D) is also continuously differentiable and we have $\nabla f(D) = \mathbb{E}_{z}[\nabla_{D}\ell(D; z)]$.

Next, we prove the Lipschitz of $\nabla f(D)$. Let v'(D'; z') and e'(D'; z') be the minimizer of $\tilde{\ell}(D', v, e; z')$. Since $\tilde{\ell}(D, v, e; z)$ has a unique minimum and is continuous in D, v, e and z, v'(D'; z') and e'(D'; z') is continuous in D and z.

Let $\Lambda = \{j \mid e'_j \neq 0\}$. According the first order optimality condition, we know that

$$\frac{\partial \tilde{\ell}(D, \boldsymbol{v}, \boldsymbol{e}; \boldsymbol{z})}{\partial \boldsymbol{e}} = 0,$$

which implies

$$\lambda_1(\boldsymbol{z}' - D'\boldsymbol{v}' - \boldsymbol{e}') \in \lambda_2 \partial \|\boldsymbol{e}'\|_1$$

Hence,

$$\left|(\boldsymbol{z}'-D'\boldsymbol{v}'-\boldsymbol{e}')_{j}\right|=rac{\lambda_{2}}{\lambda_{1}},\ \forall j\in \Lambda.$$

Since z-Dv-e is continuous in z and D, there exists an open neighborhood $\mathcal V$, such that for all $(z'',D'')\in \mathcal V$, if $j\notin \Lambda$, then $|(z''-D''v''-e'')_j|<\frac{\lambda_2}{\lambda_1}$ and $e''_j=0$. That is, the support set of e' will not change.

Let us denote $H = [D\ I_p], \, m{r} = [m{v}^{ op}\ m{e}^{ op}]^{ op}$ and define the function

$$ilde{\ell}(H_{A},oldsymbol{r}_{A};oldsymbol{z}) = rac{\lambda_{1}}{2}\left\|oldsymbol{z}-H_{A}oldsymbol{r}_{A}
ight\|_{2}^{2} + rac{1}{2}\left\|[I_{d}\ 0]oldsymbol{r}_{A}
ight\|_{2}^{2} + \lambda_{2}\left\|[0\ I_{|A|}]oldsymbol{r}_{A}
ight\|_{1}.$$

Above, $r_{\Lambda} = [\boldsymbol{v}^{\top} \ \boldsymbol{e}_{\Lambda}^{\top}]^{\top}$, and accordingly for H_{Λ} . Since $\tilde{\ell}(D_{\Lambda}, \boldsymbol{r}_{\Lambda}; \boldsymbol{z})$ is strongly convex with respect to \boldsymbol{r}_{Λ} , there exists a uniform constant κ_1 , such that for all $\boldsymbol{r}''_{\Lambda}$,

$$\tilde{\ell}(H'_{\Lambda}, \mathbf{r}''_{\Lambda}; \mathbf{z}') - \tilde{\ell}(H'_{\Lambda}, \mathbf{r}'_{\Lambda}; \mathbf{z}') \ge \kappa_1 \|\mathbf{r}''_{\Lambda} - \mathbf{r}'_{\Lambda}\|_2^2 = \kappa_1 (\|\mathbf{v}'' - \mathbf{v}'\|_2^2 + \|\mathbf{e}''_{\Lambda} - \mathbf{e}'_{\Lambda}\|_2^2).$$
(37)

On the other hand,

$$\tilde{\ell}(H'_{\Lambda}, \mathbf{r}''_{\Lambda}; \mathbf{z}') - \tilde{\ell}(H'_{\Lambda}, \mathbf{r}'_{\Lambda}; \mathbf{z}')$$

$$= \tilde{\ell}(H'_{\Lambda}, \mathbf{r}''_{\Lambda}; \mathbf{z}') - \tilde{\ell}(H''_{\Lambda}, \mathbf{r}''_{\Lambda}; \mathbf{z}'') + \tilde{\ell}(H''_{\Lambda}, \mathbf{r}''_{\Lambda}; \mathbf{z}'') - \tilde{\ell}(D'_{\Lambda}, \mathbf{r}'_{\Lambda}; \mathbf{z}')$$

$$\leq \tilde{\ell}(H'_{\Lambda}, \mathbf{r}''_{\Lambda}; \mathbf{z}') - \tilde{\ell}(H''_{\Lambda}, \mathbf{r}''_{\Lambda}; \mathbf{z}'') + \tilde{\ell}(H''_{\Lambda}, \mathbf{r}'_{\Lambda}; \mathbf{z}'') - \tilde{\ell}(H'_{\Lambda}, \mathbf{r}'_{\Lambda}; \mathbf{z}'), \tag{38}$$

where the last inequality holds because r'' is the minimizer of $\tilde{\ell}(H'', r; z'')$.

We shall prove that $\tilde{\ell}(H'_{\Lambda}, r_{\Lambda}; z') - \tilde{\ell}(H''_{\Lambda}, r_{\Lambda}; z'')$ is Lipschitz w.r.t. r, which implies the Lipschitz of v'(D; z) and e'(D; z). By algebra, we have

$$\nabla_{\boldsymbol{r}} \left(\tilde{\ell}(H'_{\Lambda}, \boldsymbol{r}_{\Lambda}; \boldsymbol{z}') - \tilde{\ell}(H''_{\Lambda}, \boldsymbol{r}_{\Lambda}; \boldsymbol{z}'') \right) = \lambda_1 \left[H'^{\top}_{\Lambda} (H'_{\Lambda} - H''_{\Lambda}) + (H'_{\Lambda} - H''_{\Lambda})^{\top} H''_{\Lambda} + H'^{\top}_{\Lambda} (\boldsymbol{z}'' - \boldsymbol{z}') + (H''_{\Lambda} - H'_{\Lambda})^{\top} \boldsymbol{z}'' \right].$$

Note that $||H'_A||_F$, $||H''_A||_F$ and z'' are all uniformly bounded by Assumption (A1) and Prop. 2. Hence, there exists uniform constants c_1 and c_2 , such that

$$\left\|
abla_{oldsymbol{r}} \left(ilde{\ell}(oldsymbol{z}', H_{A}', oldsymbol{r}_{A}) - ilde{\ell}(oldsymbol{z}'', H_{A}'', oldsymbol{r}_{A})
ight)
ight\|_{2} \leq c_{1} \left\| H_{A}' - H_{A}''
ight\|_{F} + c_{2} \left\| oldsymbol{z}' - oldsymbol{z}''
ight\|_{2},$$

which implies that $\tilde{\ell}(H'_{\Lambda}, \boldsymbol{r}_{\Lambda}; \boldsymbol{z}') - \tilde{\ell}(H''_{\Lambda}, \boldsymbol{r}_{\Lambda}; \boldsymbol{z}'')$ is Lipschitz w.r.t \boldsymbol{r}_{Λ} where the Lipschitz coefficient $c(H'_{\Lambda}, H''_{\Lambda}, \boldsymbol{z}', \boldsymbol{z}'') = c_1 \|H'_{\Lambda} - H''_{\Lambda}\|_F + c_2 \|\boldsymbol{z}' - \boldsymbol{z}''\|_2$. Combining this fact with (37) and (38), we obtain

$$\kappa_1 \| \boldsymbol{r}_{\Lambda}'' - \boldsymbol{r}_{\Lambda}' \|_2^2 \le c(H_{\Lambda}', H_{\Lambda}'', \boldsymbol{z}', \boldsymbol{z}'') \| \boldsymbol{r}_{\Lambda}'' - \boldsymbol{r}_{\Lambda}' \|_2.$$

Therefore, r(D; z) is Lipschitz and so are v(D; z) and e(D; z). Note that according to Proposition 4,

$$\nabla f(D') - \nabla f(D'') = \mathbb{E}_{\boldsymbol{z}} \Big[(H'\boldsymbol{r}' - \boldsymbol{z})(\boldsymbol{v}')^{\top} - (H''\boldsymbol{r}'' - \boldsymbol{z})(\boldsymbol{v}'')^{\top} \Big]$$
$$= \mathbb{E}_{\boldsymbol{z}} \Big[H'\boldsymbol{r}'(\boldsymbol{v}' - \boldsymbol{v}'')^{\top} + (H' - H'')\boldsymbol{r}'(\boldsymbol{v}'')^{\top}$$
$$+ H''(\boldsymbol{r}' - \boldsymbol{r}'')(\boldsymbol{v}'')^{\top} + \boldsymbol{z}(\boldsymbol{v}'' - \boldsymbol{v}')^{\top} \Big].$$

Thus,

$$\begin{split} & \left\| \nabla f(D') - \nabla f(D'') \right\|_F \\ & \stackrel{\zeta_1}{\leq} \mathbb{E}_{\boldsymbol{z}} \Big[\left\| H' \boldsymbol{r}' \right\|_2 \left\| \boldsymbol{v}' - \boldsymbol{v}'' \right\|_2 + \left\| H' - H'' \right\|_F \left\| \boldsymbol{r}' \boldsymbol{v}''^\top \right\|_F \\ & + \left\| H'' \right\|_F \left\| \boldsymbol{r}' - \boldsymbol{r}'' \right\|_2 \left\| \boldsymbol{v}'' \right\|_2 + \left\| \boldsymbol{z} \right\|_2 \left\| \boldsymbol{v}' - \boldsymbol{v}'' \right\|_2 \Big] \\ & \stackrel{\zeta_2}{\leq} \mathbb{E}_{\boldsymbol{z}} \Big[\left(\gamma_1 + \gamma_2 \left\| \boldsymbol{z} \right\|_2 \right) \left\| H' - H'' \right\|_F \Big] \\ & \stackrel{\zeta_3}{\leq} \gamma_0 \left\| D' - D'' \right\|_F, \end{split}$$

where γ_0, γ_1 and γ_2 are all uniform constants. Here, ζ_1 holds due to the convexity of $\|\cdot\|_F$. ζ_2 is derived by using the result that r(H; z) and v(H; z) are both Lipschitz and H', H'', r', r'', v' and v'' are all uniformly bounded. ζ_3 holds because z is uniformly bounded and $\|H' - H''\|_F = \|D' - D''\|_F$. Thus, we complete the proof.

Proof. (Proof of Theorem 3) Since $\frac{1}{t}A_t$ and $\frac{1}{t}B_t$ are uniformly bounded (Proposition 2), there exist subsequences of $\{\frac{1}{t}A_t\}$ and $\{\frac{1}{t}B_t\}$ that converge to A_{∞} and B_{∞} respectively. Then D_t will converge to D_{∞} . Let W be an arbitrary matrix in $\mathbb{R}^{p\times d}$ and $\{h_k\}_{k>1}$ be any positive sequence that converges to zero.

As $g_t(D)$ is a surrogate function of $f_t(D)$, for all t and k, we have

$$g_t(D_t + h_k W) \ge f_t(D_t + h_k W).$$

Let t tend to infinity, and note that $f(D) = \lim_{t \to \infty} f_t(D)$, we have

$$q_{\infty}(D_{\infty} + h_k W) > f(D_{\infty} + h_k W).$$

Note that the Lipschitz property of $\nabla f(D)$ indicates that the second derivative of f(D) is uniformly bounded. By a simple calculation, we can also show that it also holds for $g_t(D)$. This fact implies that we can take the first order Taylor expansion for both $g_t(D)$ and f(D) even when t tends to infinity (because the second order derivatives of them always exist). That is,

$$\operatorname{Tr}(h_k W^{\top} \nabla g_{\infty}(D_{\infty})) + o(h_k \|W\|_F) \ge \operatorname{Tr}(h_k W^{\top} \nabla f(D_{\infty})) + o(h_k \|W\|_F).$$

Above, we use the fact that $\lim_{t\to\infty} g_t(D_t) - f(D_t) = 0$ as implied by Corollary 1. By multiplying $\frac{1}{h_k \|W\|_F}$ on both sides and note that $\{h_k\}_{k\geq 1}$ is a positive sequence, it follows that

$$\operatorname{Tr}\left(\frac{1}{\|W\|_{F}}W^{\top}\nabla g_{\infty}(D_{\infty})\right) + \frac{o(h_{k}\|W\|_{F})}{h_{k}\|W\|_{F}} \ge \operatorname{Tr}\left(\frac{1}{\|W\|_{F}}W^{\top}\nabla f(D_{\infty})\right) + \frac{o(h_{k}\|W\|_{F})}{h_{k}\|W\|_{F}}.$$

Now let k go to infinity, we obtain

$$\operatorname{Tr}\left(\frac{1}{\|W\|_F}W^{\top}\nabla g_{\infty}(D_{\infty})\right) \ge \operatorname{Tr}\left(\frac{1}{\|W\|_F}W^{\top}\nabla f(D_{\infty})\right).$$

Note that this inequality holds for any matrix $W \in \mathbb{R}^{p \times d}$, so we actually have

$$\nabla g_{\infty}(D_{\infty}) = \nabla f(D_{\infty}).$$

As D_{∞} is the minimizer of $g_{\infty}(D)$, we have

$$\nabla f(D_{\infty}) = \nabla g_{\infty}(D_{\infty}) = 0.$$

The proof is complete.

References

- H. Avron, S. Kale, S. P. Kasiviswanathan, and V. Sindhwani. Efficient and practical stochastic subgradient descent for nuclear norm regularization. In *Proceedings of the 29th International Conference on Machine Learning*, 2012.
- D. P. Bertsekas. Nonlinear programming. Massachusetts: Athena Scientific, 1999.
- J. F. Bonnans and A. Shapiro. Optimization problems with perturbations: A guided tour. *SIAM Review*, 40 (2):228–264, 1998.
- L. Bottou. Online learning and stochastic approximations. On-line learning in neural networks, 17(9), 1998.
- L. Bottou and O. Bousquet. The tradeoffs of large scale learning. In *Proceedings of the 21st Annual Conference on Neural Information Processing Systems*, pages 161–168, 2007.
- S. Burer and R. D. C. Monteiro. Local minima and convergence in low-rank semidefinite programming. *Mathematical Programming*, 103(3):427–444, 2005.
- E. J. Candès and B. Recht. Exact matrix completion via convex optimization. *Foundations of Computational Mathematics*, 9(6):717–772, 2009.
- E. J. Candès, X. Li, Y. Ma, and J. Wright. Robust principal component analysis? *Journal of the ACM*, 58 (3):11, 2011.
- Y. Chen, A. Jalali, S. Sanghavi, and H. Xu. Clustering partially observed graphs via convex optimization. *Journal of Machine Learning Research*, 15(1):2213–2238, 2014.
- A. Defazio, F. R. Bach, and S. Lacoste-Julien. SAGA: A fast incremental gradient method with support for non-strongly convex composite objectives. In *Proceedings of 28th Annual Conference on Neural Information Processing Systems*, pages 1646–1654, 2014.
- D. L. Donoho. De-noising by soft-thresholding. IEEE Trans. Information Theory, 41(3):613–627, 1995.
- E. Elhamifar and R. Vidal. Sparse subspace clustering. In *IEEE Conference on Computer Vision and Pattern Recognition*, pages 2790–2797, 2009.
- B. Eriksson, L. Balzano, and R. D. Nowak. High-rank matrix completion and subspace clustering with missing data. *CoRR*, abs/1112.5629, 2011.
- M. Fazel, H. Hindi, and S. P. Boyd. A rank minimization heuristic with application to minimum order system approximation. In *Proceedings of the American Control Conference*, volume 6, pages 4734–4739, 2001.
- J. Feng, H. Xu, and S. Yan. Online robust PCA via stochastic optimization. In *Proceedings of the 27th Annual Conference on Neural Information Processing Systems*, pages 404–412, 2013.
- R. Ge, J. D. Lee, and T. Ma. Matrix completion has no spurious local minimum. In *Proceedings of the 30th Annual Conference on Neural Information Processing Systems*, pages 2973–2981, 2016.
- B. D. Haeffele, E. Young, and R. Vidal. Structured low-rank matrix factorization: Optimality, algorithm, and applications to image processing. In *Proceedings of the 31st International Conference on Machine Learning*, pages 2007–2015, 2014.

- C. Hsieh and P. A. Olsen. Nuclear norm minimization via active subspace selection. In *Proceedings of the* 31st International Conference on Machine Learning, pages 575–583, 2014.
- M. Jaggi and M. Sulovský. A simple algorithm for nuclear norm regularized problems. In *Proceedings of the 27th International Conference on Machine Learning*, pages 471–478, 2010.
- P. Jain, P. Netrapalli, and S. Sanghavi. Low-rank matrix completion using alternating minimization. In *Proceedings of the 45th ACM Symposium on Theory of Computing*, pages 665–674, 2013.
- J. D. Lee, M. Simchowitz, M. I. Jordan, and B. Recht. Gradient descent only converges to minimizers. In *Proceedings of the 29th Conference on Learning Theory*, pages 1246–1257, 2016.
- Z. Lin, M. Chen, and Y. Ma. The augmented lagrange multiplier method for exact recovery of corrupted low-rank matrices. *arXiv preprint arXiv:1009.5055*, 2010.
- G. Liu and P. Li. Recovery of coherent data via low-rank dictionary pursuit. In *Proceedings of the 28th Annual Conference on Neural Information Processing Systems*, pages 1206–1214, 2014.
- G. Liu, Z. Lin, and Y. Yu. Robust subspace segmentation by low-rank representation. In *Proceedings of the 27th International Conference on Machine Learning*, pages 663–670, 2010.
- G. Liu, Z. Lin, S. Yan, J. Sun, Y. Yu, and Y. Ma. Robust recovery of subspace structures by low-rank representation. *IEEE Trans. Pattern Analysis and Machine Intelligence*, 35(1):171–184, 2013.
- J. Mairal. Stochastic majorization-minimization algorithms for large-scale optimization. In *Proceedings of the 27th Annual Conference on Neural Information Processing Systems*, pages 2283–2291, 2013.
- J. Mairal, F. R. Bach, J. Ponce, and G. Sapiro. Online learning for matrix factorization and sparse coding. *Journal of Machine Learning Research*, 11:19–60, 2010.
- P. Netrapalli, N. U. N, S. Sanghavi, A. Anandkumar, and P. Jain. Non-convex robust PCA. In *Proceedings of the 28th Annual Conference on Neural Information Processing Systems*, pages 1107–1115, 2014.
- A. Y. Ng, M. I. Jordan, and Y. Weiss. On spectral clustering: Analysis and an algorithm. In *Proceedings of the 15th Annual Conference on Neural Information Processing Systems*, pages 849–856, 2001.
- B. Recht, M. Fazel, and P. A. Parrilo. Guaranteed minimum-rank solutions of linear matrix equations via nuclear norm minimization. *SIAM Review*, 52(3):471–501, 2010.
- J. D. M. Rennie and N. Srebro. Fast maximum margin matrix factorization for collaborative prediction. In *Proceedings of the 22nd International Conference on Machine Learning*, pages 713–719, 2005.
- P. Richtárik and M. Takác. Iteration complexity of randomized block-coordinate descent methods for minimizing a composite function. *Mathematical Programming*, 144(1-2):1–38, 2014.
- J. Shen and P. Li. Learning structured low-rank representation via matrix factorization. In *Proceedings of the 19th International Conference on Artificial Intelligence and Statistics*, pages 500–509, 2016.
- J. Shen, H. Xu, and P. Li. Online optimization for max-norm regularization. In *Proceedings of the 28th Annual Conference on Neural Information Processing Systems*, pages 1718–1726, 2014.
- J. Shen, P. Li, and H. Xu. Online low-rank subspace clustering by basis dictionary pursuit. In *Proceedings* of the 33rd International Conference on Machine Learning, pages 622–631, 2016.

- M. Soltanolkotabi and E. J. Candès. A geometric analysis of subspace clustering with outliers. *The Annals of Statistics*, 40:2195–2238, 2012.
- M. Soltanolkotabi, E. Elhamifar, and E. J. Candès. Robust subspace clustering. *The Annals of Statistics*, 42 (2):669–699, 2014.
- N. Srebro, J. D. M. Rennie, and T. S. Jaakkola. Maximum-margin matrix factorization. In *Proceedings of the 18th Annual Conference on Neural Information Processing Systems*, pages 1329–1336, 2004.
- A. van der Vaart. Asymptotic statistics. Cambridge University Press, 2000.
- R. Vidal. A tutorial on subspace clustering. IEEE Signal Processing Magazine, 28(2):52-68, 2010.
- Y. Wang, H. Xu, and C. Leng. Provable subspace clustering: When LRR meets SSC. In *Proceedings of 27th Annual Conference on Neural Information Processing Systems*, pages 64–72, 2013.
- L. Xiao and T. Zhang. A proximal stochastic gradient method with progressive variance reduction. *SIAM Journal on Optimization*, 24(4):2057–2075, 2014.
- H. Xu, C. Caramanis, and S. Mannor. Principal component analysis with contaminated data: The high dimensional case. In *Proceedings of the 23rd Conference on Learning Theory*, pages 490–502, 2010.
- H. Xu, C. Caramanis, and S. Mannor. Outlier-robust PCA: the high-dimensional case. *IEEE Trans. Information Theory*, 59(1):546–572, 2013.
- R. Zhao, W. B. Haskell, and J. Feng. A unified framework for stochastic matrix factorization via variance reduction. *CoRR*, abs/1705.06884, 2017.