H. Jardón-Kojakhmetov

June 28, 2018

#### Abstract

An  $A_k$  slow fast system is a particular type of singularly perturbed ODE. The corresponding slow manifold is defined by the critical points of a universal unfolding of an  $A_k$  singularity. In this note we propose a formal normal form of  $A_k$  slow fast systems.

### 1 Introduction

In this note we propose a formal normal form of a particular class of slow fast systems. A slow fast system (SFS) is a singularly perturbed ODE usually written as

$$\begin{aligned} \dot{x} &= f(x, z, \varepsilon) \\ \varepsilon \dot{z} &= g(x, z, \varepsilon) \end{aligned} \tag{1}$$

where  $x \in \mathbb{R}^m$ ,  $z \in \mathbb{R}^n$  and  $0 < \varepsilon \ll 1$  is a small parameter, and where the over-dot denotes the derivative with respect to a time parameter t. Slow fast systems are often used as mathematical models of phenomena which occur in two time scales. Observe that as  $\varepsilon$  decreases, the time scale difference between x and z increases. A couple of classical examples of real life phenomena that were modeled by a SFS are the Zeeman's heartbeat and nerve-impulse models [15]. For  $\varepsilon \neq 0$ , we can define a new time parameter  $\tau$  by  $t = \varepsilon \tau$ . With this new time  $\tau$  we can write (1) as

$$\begin{aligned} x' &= \varepsilon f(x, z, \varepsilon) \\ z' &= g(x, z, \varepsilon), \end{aligned} \tag{2}$$

where the prime denotes derivative with respect to  $\tau$ . An important geometric object in the study of SFSs is the *slow manifold* which is defined by

$$S = \{ (x, z) \in \mathbb{R}^m \times \mathbb{R}^n \, | \, g(x, z, 0) = 0 \} \,.$$
(3)

When  $\varepsilon = 0$ , the manifold S serves as the phase space of (1) and as the set of equilibrium points of (2). In the rest of the document, we prefer to work with a SFS written as (2). Furthermore, to avoid working with an  $\varepsilon$ -parameter family of vector fields as in (2), we plug-in into (2) the trivial equation  $\varepsilon' = 0$ . To be more precise, we treat a  $C^{\infty}$ -smooth vector field defined as follows.

**Definition 1.1** ( $A_k$  slow fast system). Let  $k \in \mathbb{N}$  with  $k \geq 2$ . An  $A_k$  slow fast system (for short  $A_k$ -SFS) is a vector field X of the form

$$X = \varepsilon (1 + \varepsilon f_1) \frac{\partial}{\partial x_1} + \sum_{i=1}^{k-1} \varepsilon^2 f_i \frac{\partial}{\partial x_i} - (G_k - \varepsilon f_k) \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}.$$
 (4)

where  $G_k = z^k + \sum_{i=1}^{k-1} x_i z^{i-1}$  and where each  $f_i = f_i(x_1, \ldots, x_{k-1}, z, \varepsilon)$  is a  $C^{\infty}$ -smooth function vanishing at the origin.

**Remark 1.1.** The slow manifold associated to an  $A_k$ -SFS is defined by

$$S = \left\{ (x, z) \in \mathbb{R}^k \, | \, z^k + \sum_{i=1}^{k-1} x_i z^{i-1} = 0 \right\}.$$
 (5)

The manifold S can be regarded as the critical set of the universal unfolding of a smooth function with an  $A_k$  singularity at the origin [1, 3]. Hence the name  $A_k$ -SFS.

Observe that the origin is a non-hyperbolic equilibrium point of X and thus, it is not possible to study its local dynamics with the classical Geometric Singular Perturbation Theory [6]. In this case, a technique called blow-up [4, 5, 9] is usually applied to desingularize the SFS. This methodology has been successfully used in many cases, e.g. [2, 8, 10, 11, 13, 14], where many of these deal with an  $A_k$ -SFS with fixed k = 2 or k = 3. Briefly speaking, the blow-up technique consists in an appropriate change of coordinates under which the induced vector field is regular or has simpler singularities (hyperbolic or partially-hyperbolic). However, in this work we propose a normal form of  $A_k$ -SFS to be performed prior to the blow-up, see theorem 2.2. This normalization greatly simplifies the local analysis of  $A_k$ -SFSs as shown in [7, 8].

## 2 Formal normal form of an $A_k$ -slow fast system

We regard the vector field X of definition 1.1 as X = F + P, where F and P are smooth vector fields called "the principal part" and "the perturbation" respectively. That is

$$F = \varepsilon \frac{\partial}{\partial x_1} + \sum_{i=2}^{k-1} 0 \frac{\partial}{\partial x_i} - G_k \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}, \qquad P = \sum_{i=1}^{k-1} \varepsilon^2 f_i \frac{\partial}{\partial x_i} + \varepsilon f_k \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}.$$
 (6)

The idea of the rest of the document is motivated by [12]. In short, we want to formally simplify the expression of X by eliminating the perturbation P. The terminology used below is that of [12].

The vector field F is quasihomogeneous of type r = (k, k-1, ..., 1, 2k-1) and quasidegree k-1 [1, 12]. From now on, we fix the type of quasihomogeneity r. A quasihomogeneous object of type r will be called r-quasihomogeneous.

**Definition 2.1** (Good perturbation). Let F be an r-quasihomogeneous vector field of quasidegree k - 1. A good perturbation X of F is a smooth vector field X = F + P, where  $P = P(x_1, \ldots, x_{k-1}, z, \varepsilon)$  satisfies the following conditions

- P is a smooth vector field of quasiorder greater than k-1,
- $P = \sum_{i=1}^{k-1} P_i \frac{\partial}{\partial x_i} + P_k \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}$ , with  $P|_{\varepsilon=0} = 0$ .

**Notation** By  $\mathcal{P}_{\delta}$  we denote the space of *r*-quasihomogeneous polynomials (in k+1 variables) of quasidegree  $\delta$ . By  $\mathcal{H}_{\gamma}$  we denote the space of *r*-quasihomogeneous vector fields (in  $\mathbb{R}^{k+1}$ ) of quasidegree  $\gamma$  and such that for all  $U \in \mathcal{H}_{\delta}$  we have  $U = \sum_{i=1}^{k} U_k \frac{\partial}{\partial x_i} + 0 \frac{\partial}{\partial x_{k+1}}$ . The formal series expansion of a function f is be denoted by  $\hat{f}$ .

**Definition 2.2** (The inner product  $\langle \cdot, \cdot \rangle_{r,\delta}$  [12]). Let  $x = (x_1, \ldots, x_n)$ , and  $s, q \in \mathbb{N}^n$ . Let  $f, g \in \mathcal{P}_{\delta}$ , that is

$$f = \sum_{(r,s)=\delta} f_s x^s,\tag{7}$$

where  $f_s \in \mathbb{R}$ ,  $x^s = x_1^{s_1} \cdots x_n^{s_n}$ ; and similarly for g. Then the inner product  $\langle \cdot, \cdot \rangle_{r,\delta}$  is defined as

$$\langle f,g\rangle_{r,\delta} = \sum_{(r,s)=\delta} f_s g_s \frac{(s!)^r}{\delta !},\tag{8}$$

where  $(s!)^r = (s_1!)^{r_1} \cdots (s_n!)^{r_n}$ , and where (r,s) denotes the dot product  $r \cdot s$ . So for monomials one has

$$\langle x^s, x^q \rangle_{r,\delta} = \begin{cases} \frac{(s_1!)^{r_1} \cdots (s_n!)^{r_n}}{\delta!} & \text{if } s = q \quad \text{with} \quad (s,r) = \delta, \\ 0 & \text{otherwise.} \end{cases}$$
(9)

Accordingly, for vector fields: let  $X = \sum_{i=1}^{n} X_i \frac{\partial}{\partial x_i} \in \mathcal{H}_{\delta}$ , and  $Y = \sum_{i=1}^{n} Y_i \frac{\partial}{\partial x_i} \in \mathcal{H}_{\delta}$ . Then

$$\langle X, Y \rangle_{r,\delta} = \sum_{i=1}^{n} \langle X_i, Y_i \rangle_{r,\delta+r_i}.$$
 (10)

**Definition 2.3** (The operators  $d, d^*$  and  $\Box$  [12]). The operator  $d : \mathcal{H}_{\gamma} \to \mathcal{H}_{\gamma+k-1}$  (associated to F) is defined by d(U) = [F, U] for any  $U \in \mathcal{H}_{\gamma}$ , where  $[\cdot, \cdot]$  denotes the Lie bracket. The operator  $d^*$  is the adjoint operator of d with respect to the inner product of definition 2.2. This is, given  $U \in \mathcal{H}_{\gamma}, V \in \mathcal{H}_{\gamma+k-1}$  we have

$$\langle d(U), V \rangle_{r,\gamma+k-1} = \langle U, d^*(V) \rangle_{r,\gamma}$$
(11)

For any quasidegree  $\beta > k - 1$ , the self adjoint operator  $\Box_{\beta} : \mathcal{H}_{\beta} \to \mathcal{H}_{\beta}$  is defined by  $\Box_{\beta}(U) = dd^*(U)$  for all  $U \in \mathcal{H}_{\beta}$ .

**Definition 2.4** (Resonant vector field [12]).

- We say that a vector field  $U \in \mathcal{H}_{\beta}$  is resonant if  $U \in \ker \Box_{\beta}$ .
- A formal vector field is called resonant if all its quasihomogeneous components are resonant.

**Definition 2.5** (Normal Form [12]). A good perturbation X = F + R of F is a normal form with respect to F if R is resonant.

It is important to note the following.

Lemma 2.1. ker  $\Box_{\beta} = \ker d^* |_{\mathcal{H}_{\beta}}$ .

*Proof.* Let  $\alpha = k - 1$ , then  $d : \mathcal{H}_{\gamma} \to \mathcal{H}_{\gamma+\alpha}$  and  $d^* : \mathcal{H}_{\gamma+\alpha} \to \mathcal{H}_{\gamma}$ . Due to the fact that  $d^*$  is the adjoint of d, we have the decomposition  $\mathcal{H}_{\gamma} = \operatorname{Im} d^*|_{\mathcal{H}_{\gamma+\alpha}} \oplus \ker d|_{\mathcal{H}_{\gamma}}$ . Now let  $U \in \mathcal{H}_{\gamma+\alpha} = \mathcal{H}_{\beta}$ , then  $\Box_{\beta}(U) = dd^*(U) = 0$  if and only if  $d^*U \in \ker d$ . Furthermore,  $d^*U \in \operatorname{Im} d^*$ . That is  $d^*U \in \operatorname{Im} d^* \cap \ker d$ . However  $\operatorname{Im} d^*$  and  $\ker d$  are orthogonal. Then  $\Box_{\beta}(U) = 0$  if and only if  $d^*U = 0$ .

We now recall a result of [12] (Proposition 4.4), we only adapt it for the present context.

**Theorem 2.1** (Formal normal form [12]). Let X = F + P be a good perturbation of F as in definition 2.1. Then there exists a formal diffeomorphism  $\hat{\Phi}$  such that  $\hat{\Phi}$  conjugates  $\hat{X}$  to a vector field F + R, where R is a resonant formal vector field in the sense of definition 2.4.

Finally, we present our result. In short, we prove that the resonant vector field R in theorem 2.1 associated to F given by (6) is R = 0.

**Theorem 2.2.** Let X = F + P be a good perturbation of the vector field

$$F = \varepsilon \frac{\partial}{\partial x_1} + \sum_{i=2}^{k-1} 0 \frac{\partial}{\partial x_i} - \left( z^k + \sum_{j=1}^{k-1} x_j z^{j-1} \right) \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}.$$
 (12)

Then, there exists a formal diffeomorphism  $\hat{\Phi}$  that conjugates  $\hat{X}$  with F, this is  $\hat{\Phi}_* \hat{X} = F$ .

*Proof.* From theorem 2.1 and lemma 2.1 we will show that if  $P \in \ker d^*|_{\mathcal{H}_{\geq k}}$  then P = 0. Let us start by rewriting  $d^*(P)$  in a more workable format.

To simplify the notation, let  $\alpha \geq k$ ,  $P \in \mathcal{H}_{\alpha}$ ,  $\beta = \alpha - k + 1$ , and  $Q \in \mathcal{H}_{\beta}$ ; and let  $x = (x_1, \ldots, x_{k-1}, z, \varepsilon) = (x_1, \ldots, x_{k-1}, x_k, x_{k+1})$ . If D is an operator, its adjoint with respect to the inner product definition 2.2 is always denoted as  $D^*$ .

We start with the inner product (definition 2.2)

$$\langle d(Q), P \rangle_{r,\alpha} = \langle Q, d^*(P) \rangle_{r,\beta}.$$
(13)

We can write  $d(Q) = \sum_{i=1}^{k+1} F(Q_i) - Q(F_i)$ , where  $F(Q_i) = \sum_{j=1}^{k+1} F_j \frac{\partial Q_i}{\partial x_j}$  and similarly for  $Q(F_i)$ , then

$$\langle d(Q), P \rangle_{r,\alpha} = \sum_{i=1}^{k+1} \langle F(Q_i) - Q(F_i), P_i \rangle_{r,\beta} = \sum_{i=1}^{k+1} \langle F(Q_i), P_i \rangle_{r,\alpha+r_i} - \langle Q(F_i), P_i \rangle_{r,\alpha+r_i} = \sum_{i=1}^{k+1} \langle Q_i, F^*(P_i) \rangle_{r,\beta+r_i} - \sum_{j=1}^{k+1} \langle Q_j, \left(\frac{\partial F_i}{\partial x_j}\right)^* (P_i) \rangle_{\beta+r_j} = \sum_{i=1}^{k+1} \langle Q_i, F^*(P_i) - \sum_{j=1}^{k+1} \left(\frac{\partial F_j}{\partial x_i}\right)^* (P_j) \rangle_{\beta+r_i}$$

$$(14)$$

Comparing (14) to  $\langle Q, d^*(P) \rangle_{r,\beta}$  we can write

$$d^{*}(P) = \begin{bmatrix} F^{*} - \left(\frac{\partial F_{1}}{\partial x_{1}}\right)^{*} & - \left(\frac{\partial F_{2}}{\partial x_{1}}\right)^{*} & \cdots & - \left(\frac{\partial F_{k+1}}{\partial x_{1}}\right)^{*} \\ - \left(\frac{\partial F_{1}}{\partial x_{2}}\right)^{*} & F^{*} - \left(\frac{\partial F_{2}}{\partial x_{2}}\right)^{*} & \cdots & - \left(\frac{\partial F_{k+1}}{\partial x_{2}}\right)^{*} \\ \vdots & \vdots & \ddots & \vdots \\ - \left(\frac{\partial F_{1}}{\partial x_{k+1}}\right)^{*} & - \left(\frac{\partial F_{2}}{\partial x_{k+1}}\right)^{*} & \cdots & F^{*} - \left(\frac{\partial F_{k+1}}{\partial x_{k+1}}\right)^{*} \end{bmatrix} \begin{bmatrix} P_{1} \\ P_{2} \\ \vdots \\ P_{k+1} \end{bmatrix}.$$
(15)

Plugging in the expressions of F and P into (15) we get

$$d^{*}(P) = \begin{bmatrix} F^{*} & 0 & \cdots & 0 & 1 & 0 \\ 0 & F^{*} & \cdots & 0 & z^{*} & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \\ 0 & 0 & \cdots & F^{*} & (z^{k-1})^{*} & 0 \\ 0 & 0 & \cdots & 0 & F^{*} + Z^{*} & 0 \\ -1 & 0 & \cdots & 0 & 0 & F^{*} \end{bmatrix} \begin{bmatrix} P_{1} \\ P_{2} \\ \vdots \\ P_{k-1} \\ P_{k} \\ 0 \end{bmatrix} = 0.$$
(16)

where  $Z^* = \left(kz^{k-1} + \sum_{i=2}^{k-1} (i-1)x_i z^{i-2}\right)^*$ . Now note that (16) implies  $F^*(P_j) = 0$  for all  $j = 2, \dots, k-1$  and  $P_1 = P_k = 0$ .

**Remark 2.1.** For k = 2, the result is trivial: we have  $F = \varepsilon \frac{\partial}{\partial x_1} - (z^2 + x_1) \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}$ , and therefore  $d^*(P) = 0$  is written as

$$d^{*}(P) = \begin{bmatrix} F^{*} & 1 & 0\\ 0 & F^{*} + 2z^{*} & 0\\ -1 & 0 & F^{*} \end{bmatrix} \begin{bmatrix} P_{1}\\ P_{2}\\ 0 \end{bmatrix} = 0,$$
(17)

which immediately implies  $P_1 = P_2 = 0$ .

Now, we study  $F^*(P_j) = 0$ . Recall that  $P = P(x_1, \ldots, x_{k-1}, z, \varepsilon)$  is not any vector field, but it has the property that  $P(x_1, \ldots, x_{k-1}, z, 0) = 0$ . That is, we can write

$$P = \sum_{i=1}^{k-1} \varepsilon \bar{P}_i \frac{\partial}{\partial x_i} + \varepsilon \bar{P}_k \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}, \tag{18}$$

where  $\bar{P}_j \in \mathcal{P}_{\alpha+r_j-2k+1}$ . This is because the (quasihomogeneous) weight of  $\varepsilon$  is 2k-1. Now, since it is complicated to work with the adjoint, we first rewrite the problem  $F^*(\varepsilon \bar{P}_j) = 0$ . We then prove that  $F^*(\varepsilon \bar{P}_j) = 0$  implies that  $\bar{P}_j = 0$ .

Note that  $F^*(\varepsilon \bar{P}_j) = 0$  is equivalent to  $\langle Q, F^*(\varepsilon \bar{P}_j) \rangle_{\alpha+r_j-k+1} = 0$  for all  $Q \in \mathcal{P}_{\beta+r_j}$ . Next, we use the definition of  $F^*$  that is

$$\langle Q, F^*(\varepsilon \bar{P}_j) \rangle_{r,\beta+r_j} = \langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j} = 0.$$
 (19)

We will now show that if  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j} = 0$  for all  $Q \in \mathcal{P}_{\beta+r_j}$ , then  $\bar{P}_j = 0$ . Note that by (19), this is the same as proving that  $F^*(\varepsilon \bar{P}_j) = 0$  implies  $\bar{P}_j = 0$ .

Start by choosing an element  $x^q$  of the basis of  $\mathcal{P}_{\beta+r_i}$ , this is

$$x^{q} = x_{1}^{q_{1}} \cdots x_{k-1}^{q_{k-1}} z^{q_{k}} \varepsilon^{q_{k+1}}, \qquad (r,q) = \beta + r_{j}.$$
 (20)

Then we have

$$F(x^{q}) = q_{1}x_{1}^{q_{1}-1} \cdots x_{k-1}^{q_{k-1}} z^{q_{k}} \varepsilon^{q_{k+1}+1} - \left(z^{k} + \sum_{i=1}^{k-1} x_{i} z^{i-1}\right) q_{k}x_{1}^{q_{1}} \cdots x_{k-1}^{q_{k-1}} z^{q_{k}-1} \varepsilon^{q_{k+1}}.$$
 (21)

Let us write  $\varepsilon \bar{P}_j \in \mathcal{P}_{\alpha+r_j}$  as

$$\varepsilon \bar{P}_j = \varepsilon \sum_{(r,p)=\alpha+r_j-2k+1} a_p x_1^{p_1} \cdots x_{k-1}^{p_{k-1}} z^{p_k} \varepsilon^{p_{k+1}}, \qquad (22)$$

where  $a_p \in \mathbb{R}$ . We now proceed by recursion on the exponent of  $\varepsilon$ . Let  $q_{k+1} = 0$ , then the inner product  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{\alpha+r_j}$  has only one term since F(Q) has only one monomial containing  $\varepsilon$ . That is

$$\langle F(Q), \varepsilon \bar{P}_j \rangle_{\alpha+r_j} |_{q_{k+1}=0} = \langle q_1 x_1^{q_1-1} \cdots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon, \varepsilon a_p x_1^{p_1} \cdots x_{k-1}^{p_{k-1}} z^{p_k} \rangle_{r,\alpha+r_j} = 0.$$
(23)

We naturally consider  $q_1 > 0$ . If  $q_1 = 0$ , then the equality is automatically satisfied. Recalling the definition 2.2 of the inner product, the equality (23) means that

$$\langle q_1 x_1^{q_1-1} \cdots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon, \varepsilon a_p x_1^{p_1} \cdots x_{k-1}^{p_{k-1}} z^{p_k} \rangle_{r,\alpha+r_j} = q_1 a_p \frac{(q!)^r}{(\alpha+r_j)!} = 0,$$
(24)

and therefore from (23) we have

$$a_p = a_{q_1-1, p_2, \dots, p_k, 1} = 0, (25)$$

for all  $q_1 > 0$ ,  $p_2, \ldots, p_k \ge 0$  (naturally, also satisfying the degree condition  $(r, p) = \alpha + r_j$ ). Next, let  $q_{k+1} = 1$ . Then

$$F(x^{q}) = q_{1}x_{1}^{q_{1}-1} \cdots x_{k-1}^{q_{k-1}} z^{q_{k}} \varepsilon^{2} - \left(z^{k} + \sum_{i=1}^{k-1} x_{i} z^{i-1}\right) q_{k}x_{1}^{q_{1}} \cdots x_{k-1}^{q_{k-1}} z^{q_{k}-1} \varepsilon.$$
(26)

Once again, the inner product  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j}$  has only one term, now this is due to the fact that all coefficients  $a_p$  of monomials containing  $\varepsilon$  are zero due to (25). Then

$$\langle F(Q), \varepsilon \bar{P}_j \rangle_{\alpha+r_j} |_{q_{k+1}=1} = \langle q_1 x_1^{q_1-1} \cdots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon^2, \varepsilon a_p x_1^{p_1} \cdots x_{k-1}^{p_{k-1}} z^{p_k} \varepsilon \rangle_{r,\alpha+r_j} = 0.$$
(27)

Therefore, similarly as above, we have the condition

$$a_p = a_{q_1-1, p_2, \dots, p_k, 2} = 0, (28)$$

for all  $q_1 > 0, p_2, \ldots, p_k \ge 0$  (naturally, also satisfying the degree condition  $(r, p) = \alpha + r_j$ ). By recursion arguments, assume  $q_{k+1} = n$  and that all the coefficients

$$a_p = a_{p_1, p_2, \dots, p_k, m} = 0, \qquad \forall m \le n.$$
 (29)

Then again the inner product  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j}$  has only one term, namely

$$\langle F(Q), \varepsilon \bar{P}_j \rangle_{\alpha+r_j} |_{q_{k+1}=n} = \langle q_1 x_1^{q_1-1} \cdots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon^{n+1}, \varepsilon a_p x_1^{p_1} \cdots x_{k-1}^{p_{k-1}} z^{p_k} \varepsilon^n \rangle_{r,\alpha+r_j} = 0.$$
(30)

The latter then implies

$$a_p = a_{q_1-1,p_2,\dots,p_k,n+1} = 0. (31)$$

This finishes the proof of  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j} = 0$  implies  $\bar{P}_j = 0$ .

**Remark 2.2.** Theorem 2.2 together with Borel's lemma [3], imply that an  $A_k$ -SFS X = F + P is smoothly conjugate to a smooth vector field Y = F + H where H is flat at the origin. The benefits of this normal form are exploited in [7, 8].

# Acknowledgments

The author gratefully acknowledges Henk Broer, Robert Roussarie, and Laurent Stolovitch for fruitful discussions and valuable comments and suggestions. This work is partially supported by a CONACyT postgraduate grant.

# References

- V.I. Arnold, S.M. Gusein-Zade, and A.N. Varchenko. Singularities of Differentiable Maps, Volume I, volume 17. Birkhäuser, 1985.
- [2] H. W. Broer, T. J. Kaper, and M. Krupa. Geometric Desingularization of a Cusp Singularity in Slow–Fast Systems with Applications to Zeeman's Examples. J. Dyn. Diff. Equat., 2013.

- [3] Th. Bröcker. Differentiable Germs and Catastrophes, volume 17 of Lecture Note Series. Cambridge University Press, 1975.
- [4] F. Dumortier and R. Roussarie. Geometric singular perturbation theory beyond normal hyperbolicity. In C.K.R.T. Jones and A. Khibnik, editors, *Multiple-Time-Scale Dynamical* Systems, volume 122, pages 29–63. Springer, 2001.
- [5] Freddy Dumortier and Robert Roussarie. Canard Cycles and Center Manifolds, volume 121. American Mathematical Society, 1996.
- [6] N. Fenichel. Geometric singular perturbation theory. JDE, pages 53–98, 1979.
- [7] H. Jardón-Kojakhmetov. Geometric desingularization of constrained differential equations in terms of slow fast systems. PhD Thesis, University of Groningen, 2015.
- [8] H. Jardón-Kojakhmetov, Henk W. Broer, and R. Roussarie. Analysis of a slow fast system near a cusp singularity. *in preparation*.
- [9] M. Krupa and P. Szmolyan. Extending geometric singular perturbation theory to non hyperbolic points: fold and canard points in two dimensions. SIAM J. Math. Anal., 33:286–314, 2001.
- [10] M. Krupa and P. Szmolyan. Relaxation oscillation and canard explosion. J. Diff. Eqns., 174:312–368, 2001.
- [11] Martin Krupa and Martin Wechselberger. Local analysis near a folded saddle-node singularity. Journal of Differential Equations, 248(12):2841 – 2888, 2010.
- [12] Eric Lombardi and Laurent Stolovitch. Normal forms of analytic perturbations of quasihomogeneous vector fields: rigidity, invariant analytic sets and exponentially small approximation. Ann. Sci. Éc. Norm. Supér., 43(4), 2010.
- [13] Peter Szmolyan and Martin Wechselberger. Canards in r3. Journal of Differential Equations, 177(2):419 - 453, 2001.
- [14] S. van Gils, M. Krupa, and P. Szmolyan. Asymptotic expansions using blow-up. Z. angew. Math. Phys., 56(8):369–397, 2005.
- [15] E.C. Zeeman. Differential equations for the heart beat and nerve impulse. In Towards a theoretical biology, volume 4, pages 8–67. Edinburgh University Press.