New infinite families of directed strongly regular graphs via equitable partitions

Štefan Gyürki*

Matej Bel University, 974 11 Banská Bystrica, Slovak Republic

April 2, 2015

Abstract

In this paper we introduce a construction of directed strongly regular graphs from smaller ones using equitable partitions. Each equitable partition of a single DSRG satisfying several conditions leads to an infinite family of directed strongly regular graphs. We construct in this way dozens of infinite families. For order at most 110, we confirm the existence of DSRGs for 30 previously open parameter sets.

1 Introduction

The main subject of this paper is directed strongly regular graph (briefly DSRG), a possible generalisation of the well-known (undirected) strongly regular graphs for the directed case introduced by Duval in [10]. The undirected version plays a central role in Algebraic Graph Theory, while the directed version received less attention.

We show how to construct larger DSRGs from smaller ones with the aid of suitably defined equitable partitions for directed graphs. According to our best knowledge, equitable partitions for directed graphs have not been defined in exact terms, though similar concept can be found for association schemes in [15] with an application for (undirected) strongly regular graphs in [19].

The main results of the introduced idea involve constructions of many infinite families of DSRGs. Several families have new parameter sets, others generalise and connect constructions which have been previously known from various ideas.

For the small parameter sets there is an evidence of the current state-of-the-art on the webpage of A.Brouwer and S.Hobart (see [4]). According to their catalogue of parameter sets with order at most 110, we have confirmed the existence of DSRGs for 30 open parameter sets.

^{*}e-mail address: gyurki@savbb.sk

The paper is organised as follows: In Section 2 the necessary and most significant terms are repeated. In Section 3 a brief summary is given on the used computer algebra packages. In Section 4 we define a new construction called π -join. It serves as a key concept for our project. We derive the necessary and sufficient conditions, when this construction gives us a DSRG if the basic graph is a smaller DSRG. It turns out that the crucial moment is finding a certain partition in the basic graph. In Section 5 we present good partitions for several well-known families of DSRGs and in Section 6 for DSRGs with some sporadic parameter sets.

2 Preliminaries

In this section we present the basic and most significant concepts used in this paper.

2.1 General concepts in algebraic graph theory

All graphs considered in this paper are finite, simple, directed or undirected graphs, thus no loops or multiple edges are allowed. If there is a dart from vertex x to vertex y in a directed graph (digraph), then we say that y is an *outneighbour* of x, and x is an *inneighbour* of y. The number of outneighbours (inneighbours) of a vertex x is called *outdegree (indegree)* of x. If the in- and outdegrees of all vertices in a digraph are equal to k, then we say that the digraph is k-regular, or regular of degree (valency) k.

Let *H* be a group. Suppose that $X \subseteq H$ and $e \notin X$, where *e* is the identity element of the group *H*. Then the digraph $\Gamma = \operatorname{Cay}(H, X)$ with vertex set *H* and dart set $\{(x, y) : x, y \in H, yx^{-1} \in X\}$ is called the *Cayley digraph over H with respect to X*.

Let $\Gamma = (V, E)$ be a simple undirected graph. Assume that $\pi = \{C_1, \ldots, C_r\}$ is a partition of V. It is called *equitable* if for all $1 \leq i, j \leq r$ the amount of neighbours of a vertex $u \in C_i$ in the set C_j is equal to a constant q_{ij} which does not depend on the choice of $u \in C_i$. Thus, to an equitable partition we can associate a matrix $Q = (q_{i,j})_{1 \leq i,j \leq r}$ which is called *quotient matrix*. An equitable partition π is called *homogeneous of degree* d, if each cell has size d. Equitable partitions of graphs can be defined also using their adjacency matrix. We say that a partition π of a matrix A is *column equitable*, if there exists such an $r \times r$ matrix Q, usually called *quotient matrix*, for which $A^T H = HQ$, where H is an $n \times r$ matrix, the *characteristic matrix* of π , whose columns are the characteristic

vectors of
$$\pi$$
. In other words, $h_{ij} = \begin{cases} 1 & \text{if } v_i \in C_j; \\ 0 & \text{otherwise.} \end{cases}$

Similarly, a partition π of a matrix A is row equitable, if π is column equitable for A^T . Clearly, if π is both row and column equitable for an adjacency matrix A, then π defines an equitable partition of the corresponding graph.

Following this idea one can define a kind of equitable partitions also for directed graphs. We will say that π is a *column equitable partition* of a directed graph Γ , if it is a column equitable partition of its adjacency matrix $A(\Gamma)$.

Let A be an adjacency matrix of a (di)graph Γ of order n, and π a partition of $V(\Gamma)$ into a cells C_1, C_2, \ldots, C_a . We say that adjacency matrix A respects the partition π , if the corresponding ordering (v_1, v_2, \ldots, v_n) of the vertices satisfies: $i \leq j \iff x \leq y$ for all vertices $v_i \in C_x$, $v_j \in C_y$.

We assume that the reader is familiar with the terms like *adjacency matrix*, *Kronecker product of matrices*, group of automorphisms of a graph, 1-factor (perfect matching), clique, coclique and line graphs. Otherwise, we refer the reader to [8].

We will use also terms like *blocks of imprimitivity, rank of a permutation group* or *transitivity* from permutation group theory. For definitions see [9].

2.2 Strongly regular graphs

An undirected graph Γ is called *regular* of valency k, or k-regular, if each vertex is incident to the same number k of edges. In the terms of adjacency matrix of a graph we can equivalently say that Γ is k-regular if and only if for its adjacency matrix $A = A(\Gamma)$ the equation AJ = JA = kJ holds, where J is the all-one matrix. A simple regular graph with valency k is said to be *strongly regular* (SRG, for short) if there exist integers λ and μ such that for each edge $\{u, v\}$ the number of common neighbours of u and v is exactly λ ; while for each non-edge $\{u, v\}$ the number of common neighbours of uand v is equal to μ . Previous condition can be equivalently rewritten into the equation $A^2 = kI + \lambda A + \mu (J - I - A)$ using the adjacency matrix of Γ . The quadruple (n, k, λ, μ) is called the *parameter set* of an SRG Γ .

Strongly regular graphs play a central importance in Algebraic Graph Theory. There are many dozens of papers with interesting constructions and results. The most significant ones altogether with crucial information on their properties are collected on the webpage of A. Brouwer [4]. A database of small SRGs is located on the webpage of E. Spence [39].

2.3 Directed strongly regular graphs

A possible generalisation of the notion of SRGs for directed graphs was given by Duval [10]. While the family of SRGs has been well-studied, the directed version has received less attention.

A directed strongly regular graph (DSRG) with parameters (n, k, t, λ, μ) is a regular directed graph on n vertices with valency k, such that every vertex is incident with tundirected edges, and the number of directed paths of length 2 directed from a vertex xto another vertex y is λ , if there is an arc from x to y, and μ otherwise. In particular, a DSRG with t = k is an SRG, and a DSRG with t = 0 is a doubly regular tournament.

Remark. In this paper we prefer to use the ordering (n, k, t, λ, μ) for the 5-tuple of parameters of a DSRG, however in several other papers the ordering (n, k, μ, λ, t) is used.

The adjacency matrix $A = A(\Gamma)$ of a DSRG with parameters (n, k, t, λ, μ) , satisfies

$$AJ = JA = kJ \quad \text{and} \quad A^2 = tI + \lambda A + \mu(J - I - A).$$
(1)

Oppositely, if an $n \times n$ zero-one matrix A with zeros on the diagonal satisfies (1) for some k, t, λ and μ , then it is an adjacency matrix of a DSRG with parameters (n, k, t, λ, μ) . **Proposition 1 ([10])** If Γ is a DSRG with parameter set (n, k, t, λ, μ) and adjacency matrix A, then the complementary graph $\overline{\Gamma}$ is a DSRG with parameter set $(n, \overline{k}, \overline{t}, \overline{\lambda}, \overline{\mu})$ with adjacency matrix $\overline{A} = J - I - A$, where

$$\begin{split} \bar{k} &= n-k-1\\ \bar{t} &= n-2k+t-1\\ \bar{\lambda} &= n-2k+\mu-2\\ \bar{\mu} &= n-2k+\lambda. \end{split}$$

For a directed graph Γ let Γ^T denote the digraph obtained by reversing all the darts in Γ . Then Γ^T is called the *reverse* of Γ . In other words, if A is the adjacency matrix of Γ , then A^T is the adjacency matrix of Γ^T .

The following proposition was observed by Pech, and presented in [25]:

Proposition 2 ([25]) Let Γ be a DSRG. Then the graph Γ^T is a DSRG with the same parameter set.

We say that two DSRGs Γ_1 and Γ_2 are *equivalent*, if $\Gamma_1 \cong \Gamma_2$, or $\Gamma_1 \cong \Gamma_2^T$, or $\Gamma_1 \cong \bar{\Gamma}_2$, or $\Gamma_1 \cong \bar{\Gamma}_2^T$; otherwise they are called *non-equivalent*. (In other words, Γ_1 is equivalent to Γ_2 if and only if Γ_1 is isomorphic to Γ_2 or to a graph, obtained from Γ_2 via reverse and complementation.) From our point of view the interesting DSRGs are those which are non-equivalent.

Under *eigenvalues* of a DSRG we mean the eigenvalues of its adjacency matrix. It is known, that there are precisely three different eigenvalues, all of them are integers, and their values together with their multiplicities are uniquely determined by the parameter set. More explicitly, a DSRG with parameter set (n, k, t, λ, μ) has eigenvalues:

$$\theta_0 = k, \quad \theta_{1,2} = \frac{1}{2} \left(\lambda - \mu \pm \sqrt{(\mu - \lambda)^2 + 4(t - \mu)} \right),$$

with respective multiplicities:

$$m_0 = 1$$
, $m_1 = \frac{k + \theta_1(n-1)}{\theta_2 - \theta_1}$, $m_2 = \frac{k + \theta_2(n-1)}{\theta_1 - \theta_2}$.

The parameters n, k, t, λ, μ are not independent. Relations, which have to be satisfied for such parameter sets are usually called *feasibility conditions*. Most important, and, in a sense, basic conditions are the following (for their proof see [10]):

$$k(k + \mu - \lambda) = t + (n - 1)\mu.$$
 (2)

There exists a positive integer d such that:

$$d^{2} = (\mu - \lambda)^{2} + 4(t - \mu)$$
(3)

$$d \mid (2k - (\mu - \lambda)(n - 1)) \tag{4}$$

$$n-1 \equiv \frac{2k - (\mu - \lambda)(n-1)}{d} \pmod{2} \tag{5}$$

$$n-1 \geq \left| \frac{2k - (\mu - \lambda)(n-1)}{d} \right|.$$
(6)

Further:

We have to mention that for a feasible parameter set it is not guaranteed that a DSRG does exist. A feasible parameter set for which at least one DSRG Γ exists is called *realizable*, otherwise *non-realizable*. The smallest example of a feasible, but non-realizable parameter set is (14, 5, 4, 1, 2). The non-existence of such graph was shown in [25].

There are known several general constructions of DSRGs. The most common constructions are based on algebraic or combinatorial approaches, or even on their combination. They include approaches using block matrices [1, 10], Kronecker product of matrices [10], finite geometries [12, 13, 16, 26], combinatorial block designs [13, 17], finite incidence structures [5, 30], coherent algebras [13, 25], association schemes [17], regular tournaments [23], partial sum families [2, 27], Cayley graphs [11, 17, 20, 23, 25] or computer algebra experimentation [17, 24]. In this paper we extend this list by a construction which is on the border of matrix construction, equitable partitions and computer algebra experimentation.

3 Computer algebra tools

This project has been supported by an essential amount of results which have been obtained with the aid of a computer. All such computations have been run in GAP (Groups, Algorithms, Programming [14]) jointly with using its share packages: GRAPE – a package for computation with graphs [35] together with nauty for testing automorphisms of graphs [28]; COCO II – an unreleased package for computation with coherent configurations and association schemes, written by Reichard [34]; and the package SetOrbit for finding representatives of orbits of group actions on sets of various size, written by Pech and Reichard [32, 33].

4 π -join construction

In this section we introduce a new construction, which serves as a basic idea for constructing new graphs from old ones. We will investigate when this construction gives us a larger DSRG from smaller ones and derive its most important properties. Starting from this section it often happens that we do not need to distinguish between directed and undirected strongly regular graphs. In such a case we use abbreviation (D)SRG.

4.1 Description of the construction

Let us consider a directed or an undirected strongly regular graph Γ of order n. Let us call it *basic graph*. Suppose that $\pi = \{C_1, C_2, \ldots, C_a\}$ is a homogeneous partition of its vertex set into a cells of size b, where ab = n. Consider a positive integer j and define a digraph Γ^j_{π} as following. Create ja + 1 disjoint copies $\Gamma_0, \Gamma_1, \ldots, \Gamma_{ja}$ of the graph Γ , i.e.

for its vertex set we have

$$V(\Gamma^{j}_{\pi}) = V(\Gamma) \times \{0, 1, \dots, ja\}$$

In each of the copies we can distinguish the images of the cells of the partition π . Let us denote C_s^r the image of the cell C_s in the r-th copy of Γ . We define the adjacency in Γ_{π}^j between vertices by giving the set of outneighbours of vertices. For fixed r there is a dart from vertex (u, r) to vertex (v, r) whenever there is a dart from u to v in Γ . Further, let $(u, r) \in C_s^r$ for some s and r. There is a dart from (u, r) to all of the vertices in cells $C_1^{r+1}, C_1^{r+2}, \ldots, C_1^{r+j}, C_2^{r+j+1}, C_2^{r+j+2}, \ldots, C_2^{r+2j}, \ldots, C_a^{r+(a-1)j+1}, \ldots, C_a^{r+aj}$, where the superscripts are taken modulo aj+1. Let us call this construction π -join of Γ in power j, or just shortly π -join when Γ and/or j are either clear from the context or irrelevant.

According to our best knowledge this construction is new. However, it can be regarded as a mixture of the lexicographic product of graphs [36] and the X-join of graphs [37]. In particular, if a = 1 then our construction coincides with the lexicographic product of Γ with the complete graph K_{j+1} . The similarity with X-joins can be seen after collapsing each cell in Γ^{j}_{π} into a single vertex. From this follows that the only difference between our construction and X-join is, that we define differently adjacencies between vertices in the cells which do belong to the same copy of Γ , and those which do not.

Now, we translate the π -join construction into the language of matrices, since it will be easier to prove statements using adjacency matrices of graphs.

Let Γ be a (di)graph and $\pi = \{C_1, C_2, \ldots, C_a\}$ be a homogeneous partition of its vertices into a cells of size b, for suitable positive integers a and b. Reorder the vertices of Γ according to π and create adjacency matrix A of Γ which respects π . Then we may consider A as a block matrix with blocks of size $b \times b$. Let us define for $i \in \{1, \ldots, a\}$ matrix U_i as a matrix whose all entries are ones in the columns with indices $(i-1)a+1, (i-1)a+2, \ldots, i \cdot a$, and all other entries are zeros, i.e. $U_i = (0, \ldots, 0, 1, 0, \ldots, 0) \otimes J$, where J is the all-ones matrix of size $n \times b$, and \otimes stands for the Kronecker product. Further, for an arbitrary integer j let us create a circulant block matrix $M_j(A)$ whose first row of blocks is

$$(A, \underbrace{U_1, \ldots, U_1}_{j}, \underbrace{U_2, \ldots, U_2}_{j}, \ldots, \underbrace{U_a, \ldots, U_a}_{j}),$$

and all other rows of blocks are obtained by shifting the blocks to right by one position.

The matrix $M_i(A)$ is the adjacency matrix of the π -join of Γ in power j, defined above.

Example 1 For n = 6, a = 3 and b = 2 then we have the following matrices:

$$U_{1} = \begin{pmatrix} 1 & 1 & | & 0 & 0 & | & 0 & 0 \\ 1 & 1 & | & 0 & 0 & 0 & 0 \\ 1 & 1 & | & 0 & 0 & 0 & 0 \\ 1 & 1 & | & 0 & 0 & 0 & 0 \\ 1 & 1 & | & 0 & 0 & | & 0 & 0 \\ 1 & 1 & | & 0 & 0 & | & 0 & 0 \\ \end{pmatrix}, U_{2} = \begin{pmatrix} 0 & 0 & | & 1 & 1 & | & 0 & 0 \\ 0 & 0 & 1 & 1 & | & 0 & 0 \\ 0 & 0 & 1 & 1 & | & 0 & 0 \\ 0 & 0 & 1 & 1 & | & 0 & 0 \\ 0 & 0 & 1 & 1 & | & 0 & 0 \\ \end{pmatrix}, U_{3} = \begin{pmatrix} 0 & 0 & | & 0 & 0 & | & 1 & 1 \\ 0 & 0 & 0 & 0 & | & 1 & 1 \\ 0 & 0 & 0 & 0 & | & 1 & 1 \\ 0 & 0 & 0 & 0 & | & 1 & 1 \\ 0 & 0 & 0 & 0 & | & 1 & 1 \\ \end{pmatrix},$$

while for j = 1 and j = 2 we have:

$$M_{1}(A) = \begin{pmatrix} A & U_{1} & U_{2} & U_{3} \\ U_{3} & A & U_{1} & U_{2} \\ U_{2} & U_{3} & A & U_{1} \\ U_{1} & U_{2} & U_{3} & A \end{pmatrix}, \qquad M_{2}(A) = \begin{pmatrix} A & U_{1} & U_{1} & U_{2} & U_{2} & U_{3} & U_{3} \\ U_{3} & A & U_{1} & U_{1} & U_{2} & U_{2} & U_{3} \\ U_{2} & U_{3} & A & U_{1} & U_{1} & U_{2} & U_{2} \\ U_{2} & U_{3} & U_{3} & A & U_{1} & U_{1} & U_{2} \\ U_{2} & U_{2} & U_{3} & U_{3} & A & U_{1} & U_{1} \\ U_{1} & U_{2} & U_{2} & U_{3} & U_{3} & A & U_{1} \end{pmatrix}$$

4.2 Motivation

The motivation behind the π -join construction defined in the previous subsection comes from an analysis of previously known DSRGs. It showed that a significant amount of them can be constructed as a π -join of certain smaller DSRG Γ , its partition π and a suitable integer j. Inspired by this observation we started to investigate the following problem:

Problem: For a given $(D)SRG \Gamma$, what are the necessary and sufficient conditions for a homogeneous partition π in order to get a DSRG from the *j*-th power of its π -join?

We will solve this problem in two steps. First, we will consider the smallest case j = 1 to derive some necessary conditions on Γ and π . After that we will show that if these conditions are satisfied, then for arbitrary integer j the j-th power of the π -join produces again a DSRG. If a π -join of a (D)SRG Γ produces a larger DSRG, then we say that π is a good partition for Γ .

4.3 Necessary conditions

Here we derive the necessary conditions for a homogeneous partition π in a (D)SRG Γ in order to be good for the π -join construction. Let us start with an easy observation about matrices U_i . Computational arguments lead to the following properties:

Observation 1: Let a, b, k, A, U_i are as in the definition of the π -join construction. Then for any $i \in \{1, \ldots, a\}$ and $l \in \{1, \ldots, b\}$ we have:

- (i) $\sum_{i=1}^{a} U_i = J$,
- (ii) $U_i \cdot U_l = b \cdot U_l$,
- (iii) $A \cdot U_l = k \cdot U_l$,
- (iv) $U_i \cdot A$ has constant columns, and this constant in an arbitrary column represents the number of darts starting from cell C_i and terminating in the vertex corresponding to the given column.

The following theorem is very important, since it has many consequences, which significantly restricts our search space for possibly good partitions. **Theorem 3** Let A be an adjacency matrix of a (D)SRG Γ with parameters (n, k, t, λ, μ) , which respects the homogeneous partition $\pi = \{C_1, \ldots, C_a\}$ of degree b. Suppose that the π -join Γ^1_{π} for j = 1 is a DSRG with parameters $(\tilde{n}, \tilde{k}, \tilde{t}, \tilde{\lambda}, \tilde{\mu})$. Then

- (a) $\tilde{n} = (a+1)n$, $\tilde{k} = n+k$, $\tilde{t} = b+t$, $\tilde{\lambda} = b+\lambda$, and $\tilde{\mu} = b+\mu$.
- (b) For arbitrary $i, l \in \{1, ..., a\}$ the number $q_{i,l}$ of darts starting in C_i and terminating in $v \in C_l$ does not depend on the concrete choice of v, just on i and l. Moreover,

$$q_{i,l} = \begin{cases} \lambda + b - k & \text{if } i = l, \\ \mu & \text{if } i \neq l. \end{cases}$$

Proof. Denote $M_1(A)$ the adjacency matrix of Γ^1_{π} coming from the construction. Clearly, $M_1^2(A)$ will be a circulant block matrix which is completely determined by its first row of blocks, so let us say that its first row is (B_0, B_1, \ldots, B_a) .

(a) The equations for \tilde{n} and \tilde{k} follow immediately from the method of construction. For other parameters let us compute:

$$B_0 = A^2 + U_1 U_a + U_2 U_{a-1} + \ldots + U_a U_1 = A^2 + b \cdot (U_a + U_{a-1} + \ldots + U_1) =$$

= $A^2 + b \cdot J = \lambda \cdot A + \mu (J - I - A) + t \cdot I + b \cdot J = (\lambda + b)A + (\mu + b)(J - I - A) + (t + b)I,$
and the claim follows.

(b) Again compute:

$$B_1 = AU_1 + U_1A + U_2U_a + \ldots + U_aU_2 = U_1A + kU_1 + b(J - U_1),$$

but from strong regularity we have also $B_1 = \lambda U_1 + \mu (J - U_1)$. Altogether this means that $U_1A = (\lambda + b - k)U_1 + \mu (J - U_1)$. Now using part (*iv*) from Observation 1 we get, that the number of darts starting in C_1 and terminating in a concrete vertex in C_1 is precisely $\lambda + b - k$; while the number of those starting from C_1 and terminating in a concrete vertex in C_i , where i > 1, is precisely μ . Analogous computations with B_2, B_3, \ldots, B_a prove the claim.

Corollary 1. Let $M_1(A)$ and A are as above. Then if $M_1(A)$ is an adjacency matrix of a DSRG then necessarily

$$2k + \mu - \lambda = a\mu + b. \tag{eq1}$$

Proof. Counting the number of incoming darts to a fixed vertex in C_l in Γ we get

$$k = \sum_{i=1}^{a} q_{i,l} = (\lambda + b - k) + (a - 1)\mu.$$

Corollary 2. Necessarily:

$$\lambda + b - k \ge 0. \tag{eq2}$$

Corollary 3. If $M_1(A)$ is an adjacency matrix of a DSRG, then π is a homogeneous column-equitable partition with quotient matrix $Q = (\lambda + b - k)I + \mu(J - I)$.

Proof. Immediately follows from Corollary 1.

Corollary 4. If A is an adjacency matrix with respect to a partition π of an undirected strongly regular graph, then necessarily π is an equitable partition and the cells are regular graphs of valency $\lambda + b - k$.

Proof. In an undirected graph indegree and outdegree of a vertex are equal, therefore the partition π satisfies all conditions for being equitable.

Corollary 5. For given (D)SRG with parameter set (n, k, t, λ, μ) the number of solutions of the equation (eq1) is at most two.

Proof. Since ab = n, replacing a by n/b results in a quadratic equation in b.

Corollary 6. If a = 1, then b = n, and the potential parameter set of the π -join is $(2n, n+k, n+t, n+\lambda, n+\mu)$. Feasibility conditions for parameter sets imply that n = 1, which is not an interesting case.

Proof. If $(2n, n + k, n + t, n + \lambda, n + \mu)$ is a parameter set of a DSRG, then from (2) we have

$$(n+k)(n+k+n+\mu-n-\lambda) = (t+n) + (2n-1)(\mu+n).$$

After simple algebraic manipulations, using equations $k(k + \mu - \lambda) = t + (n - 1)\mu$ and (eq1) we obtain that n = 1.

Corollary 7. If b = 1, then the initial graph Γ is necessarily complete.

Proof. The cells are single vertices, therefore part (b) of Lemma 1 implies that the initial graph is necessarily complete.

4.4 Sufficient conditions

In this section we state that the conditions derived in the previous subsection are, in fact, sufficient.

Theorem 4 Let Γ be a (D)SRG with parameter set (n, k, t, λ, μ) . Let a and b are positive integers such that ab = n and there exists a homogeneous column equitable partition $\pi = \{C_1, \ldots, C_a\}$ of vertices with quotient matrix $Q = (\lambda + b - k)I + \mu(J - I)$. Let A be an adjacency matrix respecting π , and let us define matrix $M_j(A)$ in accordance with our π -join construction for an arbitrary positive integer j. Then $M_j(A)$ is an adjacency matrix of a DSRG with parameter set

$$((ja+1)n, jn+k, jb+t, jb+\lambda, jb+\mu).$$

Proof. The necessity of conditions has been shown above. By direct computations one can show that if A satisfies all the posed conditions, then they are also sufficient, therefore $M_j(A)$ corresponds to a DSRG for any positive integer j.

This theorem provides an algorithm how to proceed when we are looking for a π -join of a DSRG:

- 1. Take a directed or an undirected SRG Γ ;
- 2. Solve (eq1) for the parameter set of Γ ;
- 3. For the solutions of (eq1) satisfying (eq2) count the quotient matrix and find a column equitable partition π with this quotient matrix.
- 4. For π , Γ and an arbitrary integer j create the π -join in power j for Γ according to the construction.

Of course, the key moment of the construction is the finding of a good column equitable partition. In the last sections we try to provide such partitions (if they do exist) at least for the small cases of DSRGs and for some general families. As we will see, in some cases the method of construction of the initial DSRG immediately provides us such a partition. In other cases some combinatorial tricks help to find it; and it happens sometimes that we are forced to leave this job to a computer.

4.5 Properties of the construction

Here we investigate two properties of the π -joins: behaviour under the complementation, and the set of eigenvalues.

4.5.1 Complementation

In this subsection we investigate when does there exist a partition π for which both (D)SRG and its complementary (di)graph can be π -joined into a new DSRG, simultaneously. After that we derive the sufficient condition when we can get two DSRGs with complementary parameter sets as a result of the π -join construction.

Theorem 5 Suppose that Γ is a (D)SRG with parameters (n, k, t, λ, μ) , $\overline{\Gamma}$ its complementary digraph with parameters $(n, \overline{k}, \overline{t}, \overline{\lambda}, \overline{\mu})$ and π a homogeneous column equitable partition for both Γ and $\overline{\Gamma}$. Then $\mu + \overline{\mu} = n/2$ and π has precisely two cells.

Proof. From the necessary conditions on the quotient matrix follow that $\tilde{\mu} + \mu = b$ and $(\lambda + b - k) + (\tilde{\lambda} + b - \tilde{k}) = b - 1$. Using the identities for the parameters of the complementary digraph of a DSRG we get: $n - 2k + \lambda + \mu = b$ from the first equation. After this the second equation gives us:

$$b - 1 = (\lambda + \tilde{\lambda}) + 2b - (k + \tilde{k}) = \lambda + n - 2k + \mu - 2 + 2b - (n - 1) =$$
$$= b - 2 + 2b - n + 1 = 3b - 1 - n,$$

hence n = 2b and the claim follows.

From the proof and the assumptions of the previous theorem we can derive the following corollary:

Corollary 8. Let Γ be a DSRG with parameters (n, k, t, λ, μ) , $\overline{\Gamma}$ its complement with parameters $(n, \overline{k}, \overline{t}, \overline{\lambda}, \overline{\mu})$. Suppose that $\mu + \overline{\mu} = n/2$ and π is a good homogeneous column-equitable partition with two cells for Γ . Then it is a good partition for $\overline{\Gamma}$ as well.

Theorem 6 Let Γ be a DSRG with parameters (n, k, t, λ, μ) , $\overline{\Gamma}$ its complement with parameters $(n, \overline{k}, \overline{t}, \overline{\lambda}, \overline{\mu})$. Suppose that they can be simultaneously π -joined via partition π to a DSRG. Then Γ^{π} and $\overline{\Gamma}^{\pi}$ have complementary parameter sets.

Proof. We already know from Theorem 5 that under the posed assumptions the partition π has two cells. Now putting a = 2 into the formulas for parameter sets of the π -join and using relations between parameter sets of mutually complementary digraphs we get that the parameter sets for $\overline{\Gamma}^{\pi}$ and $\overline{\Gamma}^{\pi}$ coincide.

4.5.2 Eigenvalues

It is well-known that a DSRG has three different eigenvalues and all of them are integers. These eigenvalues are uniquely determined by the parameter set. It is not hard to see that if the basic graph Γ with parameter set (n, k, t, λ, μ) has eigenvalues $\{k, \theta_1, \theta_2\}$ then its π -join has eigenvalues $\{jn + k, \theta_1, \theta_2\}$.

5 Good partitions for general families of (D)SRGs

In this section we try to describe some good equitable partitions for well-known families of (D)SRGs. Consequently, the π -join construction gives us many infinite families of DSRGs, which are in many cases new. For fixed parameter set we consider just one graph and we focus on finding at least one good equitable partition.

5.1 Complete graphs

The complete graph K_n can be regarded as an SRG with parameters $(n, n - 1, n - 2, \mu)$, where the parameter μ is undefined. Thus, we cannot apply our previous theory. However, if we apply the π -join construction for the complete graph K_{ab} using the homogeneous equitable partition with a cells of size b, then it is easy to show that the result is again a DSRG. Its parameter set is

$$(ja^{2}b + ab, jab + ab - 1, jb + ab - 1, jb + ab - 2, jb + b).$$

This parameter set is known for example from [30], Section 4.

5.2 Duval's construction for $\lambda = 0, t = \mu$

Duval (in [10], Lemma 8.2) constructed an infinite family of DSRGs with parameter set $(k^2 + k, k, 1, 0, 1)$ by giving its adjacency matrix.

It is an easy exercise to check that his matrix defines a good partition for both feasible integer solutions of (eq1): (a, b) = (k, k+1) and (a, b) = (k+1, k). More precisely, if the vertices v_1, v_2, \ldots, v_n of the graph are in accordance with the order of rows/columns of the adjacency matrix given by Duval, then

 $\pi_1 = \{\{v_1, v_2, \dots, v_{k+1}\}, \{v_{k+2}, \dots, v_{2k+2}\}, \dots, \{v_{k^2}, v_{k^2+1}, \dots, v_{k^2+k}\}\},\$

and

$$\pi_2 = \{\{v_1, v_2, \dots, v_k\}, \{v_{k+1}, \dots, v_{2k}\}, \dots, \{v_{k^2+1}, \dots, v_{k^2+k}\}\}$$

are the corresponding good partitions.

This proves the existence of DSRGs with the following parameter sets:

$$(k(k+1)(jk+1), jk(k+1)+k, jk+j+1, j(k+1), jk+j+1),$$

and

$$(k(k+1)(jk+j+1), jk(k+1)+k, jk+1, jk, jk+1),$$

where j and k are arbitrary positive integers. Among these parameter sets there are several new: (60,15,4,3,4), (60,20,7,6,7), (78,26,9,8,9), (96,32,11,10,11), and (108,27,7,6,7) for $n \leq 110$.

In similar manner one can show that from Duval's generalized construction ([10], Theorem 8.3), in conjunction with our construction it is possible to obtain DSRGs with parameter sets

$$(mk(k+1)(jk+1), mjk(k+1) + mk, m(jk+j+1), mj(k+1), m(jk+j+1)), \\ (mk(k+1)(jk+j+1), mjk(k+1) + mk, m(jk+1), mjk, m(jk+1)),$$

for arbitrary positive integers j, m, k.

5.3 Jørgensen's construction

For any positive integers k and μ such that $\mu \mid k-1$, Jørgensen in [23] constructed a DSRG Γ with parameter set $((k^2 - 1)/\mu, k, \mu + 1, \mu, \mu)$. The vertices of his graphs are integers modulo $n = (k^2 - 1)/\mu$, and there is a dart from x to y if and only if $x + ky \in \{1, 2, \ldots, k\}$. (Operations are also taken modulo n.) In other words, a fixed vertex x has outneighbours of form k(s - x) and inneighbours of form s - ky for $s = 1, 2, \ldots, k$.

For this parameter set (eq1) has always a solution in integers: $(a, b) = ((k-1)/\mu, k+1)$. The possibly second solution is $(a, b) = ((k+1)/\mu, k-1)$, but since $\mu \mid k-1$ an easy number theoretical argument shows that $(k+1)/\mu$ is integer if and only if $\mu \in \{1, 2\}$.

5.3.1 Case
$$a = \frac{k-1}{\mu}, b = k+1$$

Let us define for $i \in \{0, 1, ..., a - 1\}$ the following sets:

$$C_i = \{bi\} \cup \{k(s - bi) \mid s \in \{1, 2, \dots, k\}\},\$$

i.e. C_i contains the number bi and its outneighbours. It is not hard to show that if from vertex bi there is a dart to some y, then there is no dart from bj to y for $j \neq i$, and there is no dart between bi and bj. Therefore the sets C_i are cells of a partition of $\{0, 1, 2, \ldots, n-1\}$ which was identified with the vertex set of Γ .

Pick a vertex $r \neq b \cdot i$ in C_i . Then, clearly, $bi \rightarrow r$ is a dart in Γ , and we know that there are precisely λ directed two-paths from bi to r in Γ . However, all these two-paths are inside the cell C_i , because all the outneighbours of bi belong to C_i . Hence, the number of inneighbours of r starting from C_i is precisely $\lambda + 1$, and the number of inneighbours of bi in C_i is also $\lambda + 1$, which follows from the property that Γ is a DSRG with parameter $t = \lambda + 1$. Altogether this means that for each vertex in C_i there are precisely $\lambda + 1$ inneighbours in C_i , thus if our partition is column equitable, then $q_{i,i} = \lambda + 1 = \mu + 1$.

Now take $i \neq j$, an arbitrary vertex $r \in C_j$, and count the number of inneighbours of r in C_i . Clearly, bi can not be an inneighbour of r, since $r \notin C_i$. Therefore there are precisely μ oriented paths of length two from bi to r, and we know that the midvertices on these paths lie in C_i . Hence, $q_{i,j} = \mu$.

In fact, we proved the following lemma:

Lemma 1 For arbitrary k and μ satisfying $\mu \mid k-1$ let $a = (k-1)/\mu$, b = k+1 and Γ be the corresponding DSRG from Jørgensen's construction. For $i \in \{0, 1, ..., a-1\}$ the sets

$$C_i = \{ib\} \cup \{k(s - ib) \mid s \in \{1, 2, \dots, k\}\}$$

form a column equitable partition of Γ with quotient matrix $Q = (\mu + 1)I + \mu(J - I)$. Therefore for an arbitrary positive integer j there exists a DSRG with parameter set $(N, K, \Lambda + 1, \Lambda, \Lambda)$, where:

$$N = \left(\frac{j(k-1)}{\mu} + 1\right) \cdot \frac{k^2 - 1}{\mu}, \quad K = \frac{j(k^2 - 1)}{\mu} + k, \quad \Lambda = j(k+1) + \mu.$$

Calculations show that $\Lambda \mid K - 1$ does hold, therefore this parameter set is not new, since it is covered by the construction given by Jørgensen. However, the resulting graphs from different constructions are not isomorphic, in general. For example, when j = 1, k = 7 and $\mu = 3$ then K = 23 and $\Lambda = 11$, but the resulting DSRGs from our and Jørgensen's construction with parameters (48, 23, 12, 11, 11) are non-equivalent.

5.3.2 Case
$$a = \frac{k+1}{\mu}, b = k-1$$

Recall that in this case the assumption $\mu \mid k-1$ on the initial parameters imply that the only possibilities are $\mu \in \{1, 2\}$, otherwise *a* cannot be an integer. Now let us consider a partition of the set $\{0, 1, 2, ..., n-1\}$ into cells C_i as equivalence classes modulo *a*, i.e.

$$C_i = \{ s \mid s \equiv i \pmod{a} \}.$$

We show that $\{C_0, C_1, \ldots, C_{a-1}\}$ is a good equitable partition.

Let us select any $y \in C_i$ and count the number of $x \in C_i$ such that $x \to y$ is a dart. We repeat that if x is an inneighbour of y, then for some $s \in \{1, 2, ..., k\}$ we have x = s - ky, therefore necessarily $s - ky \equiv y \pmod{a}$. Equivalently, $s \equiv (k+1)y \equiv 0 \pmod{a}$, because k + 1 is a multiple of a.

- If $\mu = 1$, then $s \equiv 0 \pmod{k+1}$ has no solution in $s \in \{1, 2, \dots, k\}$.
- If $\mu = 2$, then $s \equiv 0 \pmod{(k+1)/2}$ has a unique solution s = (k+1)/2.

Now let us consider the number of inneighbours in C_j of a fixed vertex $y \in C_i$, where $i \neq j$. Similarly as above, for an inneighbour $x \in C_j$ there is $s \in \{1, 2, ..., k\}$ such that x = s - ky, and $y - x \equiv j - i \pmod{a}$. After simplification: $s \equiv i - j \pmod{a}$.

- If $\mu = 1$, then $s \equiv i j \pmod{k+1}$ has a unique solution s = i j (taken modulo a).
- If $\mu = 2$, then $s \equiv i j \pmod{(k+1)/2}$ has exactly two solutions.

We proved the following:

Lemma 2 For $\mu \in \{1, 2\}$ and arbitrary k satisfying $\mu \mid k-1$ let $a = (k+1)/\mu$, b = k-1and Γ be the corresponding DSRG from Jørgensen's construction. For $i \in \{0, 1, ..., a-1\}$ the sets:

$$C_i = \{s \mid s \equiv i \pmod{a}\}$$

form a column equitable partition of Γ with quotient matrix $(\mu - 1)I + \mu(J - I)$. Therefore for arbitrary positive integer j there exists a DSRG with parameter set $(N, K, \Lambda + 1, \Lambda, \Lambda)$, where:

$$N = \left(\frac{j(k+1)}{\mu} + 1\right) \cdot \frac{k^2 - 1}{\mu}, \quad K = \frac{j(k^2 - 1)}{\mu} + k, \quad \Lambda = j(k-1) + \mu.$$

In the case, when k = 1 the resulting parameter set can be also obtained from Jørgensen's construction. Also this is the case, when k = 2, $\mu = j = 1$. In all other cases Λ does not divide K - 1. According to [4] we constructed first DSRG with parameters (40, 11, 4, 3, 3), (72, 19, 6, 5, 5), (90, 19, 5, 4, 4) and (104, 27, 8, 7, 7), if we consider just $N \leq 110$.

5.4 Triangular graphs

The triangular graph T(n) is defined as the line graph of the complete graph K_n . For $n \ge 4$ it is an SRG with parameters $\binom{n}{2}, 2(n-2), n-2, 4$. Moreover, graphs with these parameter set are unique, when $n \ne 8$. For n = 8 there are three more graphs, the so-called *Chang graphs* (see [6, 7]), apart from T(8).

By solving (eq1) we get that for $n \equiv 1 \pmod{2}$ there is a unique solution $(a, b) = (\frac{n-1}{2}, n)$; for $n \equiv 0 \pmod{4}$ also a unique solution (a, b) = (n/4, 2n - 2), while for $n \equiv 2 \pmod{4}$ there is no integer solution.

5.4.1 Case $n \equiv 1 \pmod{2}$

For odd *n* we have $a = \frac{n-1}{2}$ and b = n. So the cells of the potential equitable partitions are regular of valency $\lambda + b - k = 2$. In other words, each cell is a collection of cycles.

Let us consider the vertex set of the triangular graph as 2-subsets of the set $\mathbb{Z}_n = \{0, 1, 2, \ldots, n-1\}$ and there is an edge between two different 2-subsets if and only if they are not disjoint. For $i = 1, 2, 3, \ldots, (n-1)/2$ let us define sets $C_i = \{\{\ell, \ell+i\} | \ell \in \mathbb{Z}_n\}$. The partition $\pi = \{C_1, C_2, \ldots, C_{(n-1)/2}\}$ is equitable with quotient matrix Q = 2I + 4(J - I), thus the existence of DSRGs with parameter set (N, K, T, Λ, M) is proved,

where N = m(2m+1)(jm+1), K = jm(2m+1) + 4m - 2, T = j(2m+1) + 4m - 2, $\Lambda = j(2m+1) + 2m - 1$, M = j(2m+1) + 4 and j, m are arbitrary integers. In all the cases when $N \leq 110$ the parameter set has been known previously.

5.4.2 Case $n \equiv 0 \pmod{4}$

Here the cells are of order 2n - 2 and valency $\lambda + b - k = n$. Again, equivalently we can consider the graph K_n and correspondingly an edge partition in it. The partition determines an edge-coloring (not proper) in the sense, that the same color is assigned to edges from the same cell and different colors to edges from different cells. Each color defines a subgraph of K_n . We can assume that the edge u_1u_2 is black and we further consider the "black subgraph" of K_n . This edge is incident to n black edges, therefore $\deg_B(u_1) + \deg_B(u_2) = n+2$, where $\deg_B(u)$ stands for "black-degree", i.e. the number of black edges incident to u. By pigeonhole principle there exists a vertex u_3 such that u_1u_3 and u_2u_3 are black edges. Thus, $\deg_B(u_1) + \deg_B(u_3) = \deg_B(u_2) + \deg_B(u_3) = n+2$, hence $\deg_B(u_1) = \deg_B(u_2) = \deg_B(u_3) = 1 + \frac{n}{2}$, and necessarily all the n/2+1 neighbours of u_1 have black-degree n/2+1. So the number of black edges is at least $\frac{1}{2}(1+\frac{n}{2})(2+\frac{n}{2})$. However, for n > 6 this number is greater than 2n - 2, thus for $n \ge 8$ there is no good partition. For n = 4 the partition is trivial.

For the three exceptional cases, Chang graphs, we executed an exhaustive computer search for finding induced regular subgraphs of size 14 and valency 8. Since the number of subsets of size 14 in a set of size 28 is quite large, we used the **SetOrbit** package written by Pech and Reichard [34]. With its aid we can generate just one representative for each orbit of the group of automorphisms on subsets of size 14. This tool reduced the amount of subsets being considered for a few hundreds of thousands, and therefore it took just a few hours of computation. However, this exhaustive search showed that there is no such induced subgraph, and therefore there is no good partition for our construction.

5.5 Complements to triangular graphs

Triangular graph T(n) has been described above. Let us now consider its complement $\overline{T}(n)$. Again, to avoid trivial cases we suppose that n > 4. The parameter set of $\overline{T}(n)$ is $\binom{n}{2}, \binom{n-2}{2}, \binom{n-4}{2}, \binom{n-3}{2}$. Equation (eq1) has integer solution just for n = 5, 6, 8. For n = 5 see the section about the Petersen graph. For n = 6 exhaustive computer search showed that there is no good partition for the only solution (a, b) = (3, 5). Finally, for n = 8 we have just one solution (a, b) = (2, 14), but $\mu + \overline{\mu} = 14$ and Theorem 5 applies.

5.6 Lattice square graphs

Another famous family of strongly regular graphs is the so-called *lattice square graphs*. The lattice square graph $L_2(n)$ has vertex set $V = \{1, 2, ..., n\} \times \{1, 2, ..., n\}$ and two vertices are adjacent if their coordinates agree in exactly one position. The parameter set of $L_2(n)$ is $(n^2, 2n - 2, n - 2, 2)$. For each positive integer n the equation (eq1) has at least one solution a = b = n. For odd n this is the only solution, but for even n there is a second solution (a, b) = (n/2, 2n).

Conditions posed on the quotient matrix Q of a good partition say that in the first case Q = 2(J - I), and in the second Q = nI + 2(J - I).

First, consider partition $\pi = \{C_1, C_2, \ldots, C_n\}$ of vertices of $L_2(n)$ such that

$$C_1 = \{(1,1), (2,2), \dots, (n-1,n-1), (n,n)\}, C_2 = \{(1,2), (2,3), \dots, (n-1,n), (n,1)\}, \vdots C_n = \{(1,n), (2,1), \dots, (n-1,n-2), (n,n-1)\}.$$

It is not hard to see that this partition is equitable with quotient matrix Q = 2(J-I), and it guarantees the existence of a DSRG with parameters

$$(n^2(jn+1), jn^2+2n-2, jn+2n-2, jn+n-2, jn+2)$$

for arbitrary positive integers j and n. The resulting parameter set is new at least in the case when j = 1 and n = 4.

When n is even, we can consider the partition $\pi = \{C_1, C_2, \ldots, C_{n/2}\}$ such that for all $m = 1, 2, \ldots, n/2$ the cell C_m is:

$$C_m = \{(x, y) \mid x \in \{1, 2, \dots, n\}, y \in \{2m - 1, 2m\}\}.$$

It is an equitable partition with quotient matrix Q = nI + 2(J - I). The corresponding π -join of the lattice square graph $L_2(n)$ is a DSRG for each integer j with parameter set

$$\left(n^2(1+\frac{jn}{2}), jn^2+2n-2, 2jn+2n-2, 2jn+n-2, 2jn+2\right).$$

5.7 Construction by Klin et al.

Klin et al. in [25] constructed a family of DSRGs with parameter set $(2n, n-1, \frac{n-1}{2}, \frac{n-3}{2}, \frac{n-1}{2})$, where *n* is odd, as Cayley graphs over dihedral group of order 2*n*. So let $D_n = \langle r, s \mid r^n = s^2 = 1 \rangle$ be the dihedral group of order 2*n*. Let $X = \{r, r^2, \ldots, r^{(n-1)/2}, sr, sr^2, \ldots, sr^{(n-1)/2}\}$. The Cayley graph with connection set X over D_n is the requested DSRG.

For this parameter set (eq1) has always a unique solution, when n > 3. This is (a,b) = (2,n). For a good partition the quotient matrix is $(n-1)/2 \cdot J$, and it is not hard to see that the partition

$$\pi = \{\{1, r, r^2, \dots, r^{n-1}\}, \{s, sr, sr^2, \dots, sr^{n-1}\}\}\$$

satisfies this condition, therefore π is a good partition. Hence the existence of DSRGs with parameter set

$$\left((4j+2)n, 2jn+n-1, jn+\frac{n-1}{2}, jn+\frac{n-3}{2}, jn+\frac{n-1}{2}\right)$$

is proved for arbitrary j and odd n.

5.8 Construction by Duval and Iourinski

Duval and Iourinski in [11] gave a sufficient condition for construction DSRGs as Cayley graphs over metacirculant groups. For our construction it is sufficient to choose their orbit partition satisfying the q-orbit condition in order to get a good partition for (a, b) = (q, m).

For their graphs equation (eq1) has a second solution just in the case, when m = q+1. This case coincides with the construction leading to DSRGs with $\mu = 0$, $t = \lambda = 1$.

6 Good partitions for small cases of (D)SRGs

In this section we discuss the π -joins of small (D)SRGs with parameter sets which have not been included in the previous section. We show just a single good partition for each parameter set, usually. In fact, in the smallest cases ($n \leq 16$) we run an exhaustive search with the aid of a computer for the values j = 1, 2, 3. In many cases we have found dozens of good partitions and non-isomorphic π -joins. These are available upon request from the author.

Exceptionally, we will consider more graphs sharing the same parameter set or a graph of larger order. It happens since we want to include into consideration some famous graphs, as well. For example, the Petersen graph, the Shrikhande graph, the Schläfli graph or the Hoffman-Singleton graph. On the other hand, we will not consider parameter sets (n, k, t, λ, μ) , where $k \ge n/2$ and the only solution of the (eq1) is the pair (a, b) = (2, n/2), since in these cases we can apply Theorem 5 and Theorem 6.

A summary of the all new parameter sets of DSRGs is given in Appendix.

$6.1 \quad SRG(4,2,0,2)$

The smallest undirected SRG is a 4-cycle. The only good solution of (eq1) is a = b = 2. Though there are two equitable partitions into two cells of size 2, just one of them, the pairs of non-adjacent vertices, satisfies the condition posed on the quotient matrix. Our construction creates DSRGs with parameter set (8j + 4, 4j + 2, 2j + 2, 2j, 2j + 2). For this parameter set and its complementary parameter set there are known several constructions, see [10], [20], [23] and [30].

6.2 SRG(10,3,0,1) – Petersen graph

One of the most famous graphs in graph theory, Petersen graph (see Fig. 1), is SRG with parameter set (10, 3, 0, 1). It is the complementary graph to the triangular graph T(5) and it is known to be unique with this parameter set. All of its equitable partitions have been enumerated in [40]. Equation (eq1) has two solutions: (a, b) = (2, 5) and (a, b) = (5, 2), but the second is ruled out by (eq2). For (a, b) = (2, 5), a good equitable partition is determined by a partition into two disjoint 5-cycles. The corresponding DSRGs from the π -join have parameter set (20j + 10, 10j + 3, 5j + 3, 5j, 5j + 1). For j = 1 the parameter set (30, 13, 8, 5, 6) is not new, since in [27] the authors constructed a DSRG with this parameter set. Their digraph is not vertex-transitive, while ours is, thus these graphs are non-isomorphic. The first DSRG with parameter set (50, 23, 13, 10, 11) has been just recently constructed in [17]. For $j \ge 3$ the resulting parameter set is new, according to our best knowledge.



Figure 1: SRG(10,3,0,1).

6.3 DSRG(15,5,2,1,2)

There are 1292 non-isomorphic DSRGs sharing this parameter set, all with very small group of automorphisms (orders less than 6). We investigated the DSRG corresponding to the adjacency matrix $A_{15,5}$. Equation (eq1) has unique solution: (a, b) = (3, 5). A good partition is

 $\{\{1, 2, 7, 8, 10\}, \{3, 4, 9, 11, 15\}, \{5, 6, 12, 13, 14\}\}.$

The parameter set of the π -join is (45j+15, 15j+5, 5j+2, 5j+1, 5j+2), which is new.

6.4 SRG(16,5,0,2) – Clebsch graph

There is a unique SRG with parameters (16, 5, 0, 2), it is the Clebsch graph (see Fig. 2). Equation (eq1) has two solutions: (a, b) = (2, 8) and (a, b) = (4, 4). The latter solution is ruled out by (eq2). A good partition is $\{\{1, 2, \ldots, 8\}, \{9, 10, \ldots, 16\}\}$. The parameter set of the π -join is (32j + 16, 16j + 5, 8j + 5, 8j, 8j + 2) which also comes from [16].



Figure 2: Clebsch graph.



Figure 3: Shrikhande graph.

6.5 SRG(16, 6, 2, 2)

There are precisely two non-isomorphic graphs sharing this parameter set: the Shkrikhande graph (see Fig. 3) and the lattice square graph $L_2(4)$ (considered in Section 5.6). Using the labeling from the figure, the partitions

$$\{\{1, 2, 5, 6, 9, 10, 13, 14\}, \{3, 4, 7, 8, 11, 12, 15, 16\}\},\$$

and

$$\{\{1, 2, 3, 4\}, \{5, 6, 7, 8\}, \{9, 10, 11, 12\}, \{13, 14, 15, 16\}\}$$

are good partitions of the Shrikhande graph for (a, b) = (2, 8) and (a, b) = (4, 4), respectively. First case lead to the parameter set (32j + 16, 16j + 6, 8j + 2, 8j + 2), which is known from [16], and the second case to the parameter set (64j + 16, 16j + 6, 4j + 6, 4j + 2, 4j + 2), which is new.

6.6 DSRG(16,7,5,4,2)

There is a unique DSRG on this parameter set, its adjacency matrix is A_{16} . Equation (eq1) has two solutions: (a, b) = (2, 8) and (a, b) = (4, 4). In the case when (a, b) = (2, 8) a good partition is:

$$\{\{1, 2, 5, 6, 9, 10, 13, 14\}, \{3, 4, 7, 8, 11, 12, 15, 16\}\}.$$

The resulting parameter sets is (32j + 16, 16j + 7, 8j + 5, 8j + 4, 8j + 2), which is known from [10]. In the case when (a, b) = (4, 4) a good partition

$$\{\{1, 2, 3, 4\}, \{5, 6, 7, 8\}, \{9, 10, 11, 12\}, \{13, 14, 15, 16\}\}$$

The resulting parameter set (64j + 16, 16j + 7, 4j + 5, 4j + 4, 4j + 2) is new.

6.7 DSRG(18,4,3,0,1)

Duval in his seminal paper presented also a DSRG with parameter set (18, 4, 3, 0, 1). This graph has been known already from other problems in graph theory, and often is referred to as *Bosák graph* [3]. It is known to be unique on this parameter set.

Though (eq1) has two solutions, one of them is ruled out by condition (eq2). For (a, b) = (3, 6) a good partition is given by the partition into three vertex-disjoint induced copies of the subgraph isomorphic to the unique DSRG(6,2,1,0,1). Hence the existence of a DSRG with parameters (54j + 18, 18j + 4, 6j + 3, 6j, 6j + 1) is proven. For j = 1 a DSRG with parameter set (72, 22, 9, 6, 7) has been recently found in [17].

6.8 DSRG(18,7,5,2,3)

For a DSRG(18,7,5,2,3) with adjacency matrix $A_{18,7}$ we have to consider two possibilities: (*a*, *b*) = (2, 9) and (*a*, *b*) = (3, 6). A good partition is {{1, 2, ..., 9}, {10, ..., 18}} in the first case, and {{1, 4, 7, 10, 13, 16}, {2, 5, 8, 11, 14, 17}, {3, 6, 9, 12, 15, 18}} in the second case. The resulting parameter sets (36j + 18, 18j + 7, 9j + 5, 9j + 2, 9j + 3) and (54j + 18, 18j + 7, 6j + 5, 6j + 2, 6j + 3) are both new, in general.

6.9 Jørgensen's sporadic examples

Jørgensen in [22] besides a remarkable amount of non-existence results also constructed a few DSRGs as Cayley graphs.

6.9.1 DSRG(21, 6, 2, 1, 2)

Let us consider his DSRG(21,6,2,1,2), which was constructed as a Cayley graph over the group $G = \langle x, y | x^3 = y^7 = 1, xy^2 = yx \rangle$ with connection set $S = \{y, y^3, x, xy^2, x^2, x^2y^5\}$. After solving the equation (eq1) and counting the quotient matrix we figure out that (a, b) = (3, 7) is a solution and $\lambda + b - k = \mu = 2$. Therefore the partition $\pi = \{C_0, C_1, C_2\}$ where $C_i = \{x^i y^j | j = 0, 1, 2, \dots, 6\}$ is a good partition leading to the existence of a family of DSRGs with parameter set (63j + 21, 21j + 6, 7j + 2, 7j + 1, 7j + 2).

6.9.2 DSRGs of order 24

Jørgensen ([22], Theorem 14) constructed three DSRGs on 24 vertices as Cayley graphs of S_4 . Their connection sets are

$$X_{1} = \{(3,4), (2,3), (2,3,4), (1,2)(3,4), (1,2,3,4), (1,3,2), (1,3,4,2), (1,3,4)\},$$

$$X_{2} = \{(3,4), (2,3), (1,2,3), (1,2,4,3), (1,3,2), (1,3,4), (1,3)(2,4), (1,4,3), (1,4,2,3)\},$$

and

$$X_3 = \{(3,4), (2,3), (2,3,4), (2,4,3), (1,2), (1,2,3), (1,2,3,4), (1,4,3,2), (1,4,3), (1,4)\}.$$

The corresponding Cayley graphs are DSRGs with parameters (24, 8, 3, 2, 3), (24, 9, 7, 2, 4), and (24, 10, 8, 4, 4), respectively.

For DSRG(24,8,3,2,3) the equation (eq1) has only one solution (a, b) = (3, 8), while for (24,9,7,2,4) and (24,10,8,4,4) there are two solutions: (2, 12) and (3,8).

In order to get a good partition for the DSRG(24,8,3,2,3) we run an exhaustive computer search. It returned us hundreds of such partitions. One of them has cells: $C_1 = \{id, (3, 4), (1, 2, 3), (2, 3, 4), (1, 2, 3, 4), (1, 2, 4), (2, 4, 3), (2, 3)\},\$ $C_2 = \{(1, 3)(2, 4), (2, 4), (1, 3, 2, 4), (1, 3, 2), (1, 3, 4, 2), (1, 2), (1, 4, 3, 2), (1, 2)(3, 4)\},\$ $C_3 = \{(1, 4, 2), (1, 4, 3), (1, 2, 4, 3), (1, 3), (1, 4), (1, 3, 4), (1, 4, 2, 3), (1, 4)(2, 3)\}.$

Thus the existence of DSRGs with parameter set (72j+24, 24j+8, 8j+3, 8j+2, 8j+3) is proved.

For the remaining two DSRGs we have found good partitions combinatorially.

Using conditions on the quotient matrix we immediately get that the partition of S_4 into even an odd permutations gives us a good partition in the case of DSRG(24,9,7,2,4). Thus the existence of a DSRG with parameters (48j+24,24j+9,12j+7,12j+2,12j+4)is proved. For (a,b) = (3,8) the situation is a little bit more complicated. After some computer algebra experimentation we concluded, that considering the index 3 subgroup $H = \langle (1,2)(3,4), (1,2,3,4) \rangle \leq S_4$ the partition $\pi = \{H, H \cdot (2,3), H \cdot (3,4)\}$ is good. Hence, the existence of DSRGs with parameter set (72j+24,24j+9,8j+7,8j+2,8j+4)is proved.

Let us put permutation $p \in S_4$ into C_1 if $1^p \in \{1, 2\}$, otherwise put p into C_2 . Partition $\pi = \{C_1, C_2\}$ is a good partition for DSRG(24,10,8,4,4) with (a, b) = (2, 12). This proves the existence of a DSRG with parameter set (48j + 24, 24j + 10, 12j + 8, 12j + 4, 12j + 4). Considering the index 3 subgroup $H = \langle (1, 2, 3, 4), (1, 3) \rangle \leq S_4$ the partition $\pi = \{H, H \cdot (1, 2), H \cdot (1, 4)\}$ is good for DSRG(24,10,8,4,4) for (a, b) = (3, 8), therefore the existence of DSRG with parameters (72j + 24, 24j + 10, 8j + 8, 8j + 4, 8j + 4) is proved.

In all the cases, the resulting parameter sets for j = 1 has been displayed in [4] as open.

6.9.3 DSRG(26,11,7,4,5)

Jørgensen ([22], Theorem 15) constructed a DSRG(26,11,7,4,5). He prescribed adjacencies between elements of two sets of size 13. For this case (eq1) says that (a, b) = (2, 13)and from the quotient matrix it is easy to check that the two original sets define a good partition. This confirms the existence of DSRGs with parameters (52j+26, 26j+11, 13j+7, 13j+4, 13j+5). For j = 1 the parameter set (78, 37, 20, 17, 18) is new according to [4].

6.10 SRG(26,10,3,4)

According to Paulus [31] there are precisely 10 SRGs with parameters (26,10,3,4). The second most symmetric of them has group of automorphisms of order 39, which has two orbits of size 13 on the vertex set. This orbit-partition is good for the unique solution (a, b) = (2, 13) of the equation (eq1), hence the existence of DSRGs with parameters

$$(52j + 26, 26j + 10, 13j + 10, 13j + 3, 13j + 4)$$

is proved for all positive integer j. For j = 1 the resulting parameter set (78,36,23,16,17) is new according to [4].

6.11 SRG(27,10,1,5) – Complement to the Schläfli graph

The equation (eq1) gives us a unique solution (a, b) = (3, 9). For diagonal elements of the quotient matrix of a good partition we get $\lambda + b - k = 0$, what means that in the basic graph we have to find a coclique of size 9, or equivalently, a clique of size 9 in the Schläfli graph. This graph has been well-studied, and we know that the maximal clique is of size 6. Therefore there is no good partition for our construction.

6.12 Sporadic examples by Martinez and Araluze

Martinez and Araluze [27] constructed a handful of DSRGs with previously unknown parameter sets with order at most 45 using partial sum families over cyclic groups. From their construction immediately follows that the equivalence classes modulo a (where a is the number of blocks) form a column equitable partition, and these partitions are not excluded neither by condition (eq1) nor by (eq2). Moreover, from the blocks it is easy to count the quotient matrix, since it is enough to count the amount of numbers modulo ain each block for each possibility.

In this way we checked all their 14 graphs constructed from cyclic groups, and figured out that six of them have quotient matrix satisfying part (b) in Theorem 1. These correspond to parameter sets: (27,10,6,3,4), (30,13,8,5,6), (34,14,12,5,6), (34,15,9,6,7), (39,10,6,1,3) and (40,17,11,6,8). Consequently, using our π -join construction we can create for any positive integer j a DSRG with parameter set: (81j+27,27j+10,9j+6,9j+3,9j+4), (60j+30,30j+13,15j+8,15j+5,15j+6), (68j+34,34j+14,17j+12,17j+5,17j+6), (68j+34,34j+15,17j+9,17j+6,17j+7), (117j+39,39j+10,13j+6,13j+1,13j+3) and (80j+40,40j+17,20j+11,20j+6,20j+8).

According to [4] we constructed the first DSRGs with parameters (102, 48, 29, 22, 23), (102, 49, 26, 23, 24), (108, 37, 15, 12, 13), while the parameter set (90, 43, 23, 20, 21) appears only in this paper, but here already for the third time.

6.13 DSRG(36,13,11,2,6)

The first DSRG with parameter set (36, 13, 11, 2, 6) has been recently constructed in [17] as a union of classes in a non-commutative association scheme. The equation (eq1) has two solutions (a, b) = (2, 18) and (3, 12). The full group of automorphisms of the graph is transitive of order 144, therefore we can consider its system of imprimitivity. One can obtain a good partition for the first solution as a union of a block of imprimitivity of size 12, and a suitable block of size 6. Hence the existence of DSRGs with parameter sets

$$(72j + 36, 36j + 13, 18j + 11, 18j + 2, 18j + 6)$$

is confirmed for all positive integer j. For j = 1 the parameter set (108, 49, 29, 20, 24) is new according to [4].

For the case (a, b) = (3, 12) a good partition is given by the system of imprimivity with 3 cells of size 12, where each cell is inducing a regular graph of valency 1. This confirms the existence of DSRGs with parameters

(108j + 36, 36j + 13, 12j + 11, 12j + 2, 12j + 6)

for all positive integer j.

6.14 SRG(36, 14, 4, 6)

There are precisely 180 SRGs (for more information see [29, 38]) with parameters (36,14,4,6). For this parameter set (eq1) has two solutions: (a, b) = (2, 18) and (3, 12).

Let us consider the second most symmetric graph of them, i.e. which has group of automorphisms of order 432. The imprimitivity system of the group of automorphisms has blocks of size 18. The corresponding partition fits the conditions for being good for our construction, therefore we can construct for every positive integer j a DSRG with parameter set

$$(72j + 36, 36j + 14, 18j + 14, 18j + 4, 18j + 6).$$

For j = 1 the resulting parameter set (108, 50, 32, 22, 24) is new according to our best knowledge.

If we consider the only imprimitivity system of the group of automorphisms with blocks of size 6, then after some experimentation we can obtain 3 pairs of blocks whose unions results in a 3-cell partition with cells of size 12 and quotient matrix 2J + 6(J - I). This proves the existence of a DSRG with parameter set

$$(108j + 36, 36j + 14, 12j + 14, 12j + 4, 12j + 6)$$

for arbitrary positive integer j.

6.15 SRG(36,15,6,6)

It is known that the exact number of SRG sharing the parameter set (36,15,6,6) is 32548 (for more information see [29, 38]). We considered a highly symmetric one of them with automorphism group of order 648. (In the catalogue of E. Spence it is the last graph on this parameter set.) It has a unique system of imprimitivity with blocks of size 9. Union of any two of these blocks and the remaining two blocks gives us a suitable partition for (a, b) = (2, 18). Hence, the existence of a DSRG with parameter set

$$(72j + 36, 36j + 15, 18j + 15, 18j + 6, 18j + 6)$$

is proved for every integer j. For j = 1 the parameter set (108, 51, 33, 24, 24) is new according to [4].

In a similar fashion, using unions of 4 blocks of imprimitivity of size 3 we have found a good partition into 3 cells of size 12, therefore the existence of a DSRG with parameter set

$$(108j + 36, 36j + 15, 12j + 15, 12j + 6, 12j + 6)$$

is proved for all integer j.

6.16 SRG(50,7,0,1) – Hoffman-Singleton graph

Just for curiosity we considered one of the most famous SRGs, the Hoffman-Singleton graph. Solutions of (eq1) are (5, 10) and (10, 5), however the latter is excluded by (eq2). The first solution challenges us to consider the partition consisting from 5 vertex-disjoint copies of the Petersen graph. By counting the possible quotient matrix it is not hard to see that this is really a good partition, therefore there exists a DSRG (250j + 50, 50j + 7, 10j + 7, 10j, 10j + 1) for each positive integer j.

7 Summary

Using our π -join construction introduced in Section 4 we constructed dozens of infinite families of DSRGs from smaller ones. Among the constructed graphs with order less than 110 there are 30 with new parameter sets according to [4]. The new parameter sets are displayed in the Appendix.

Acknowledgement

The author gratefully acknowledges the contribution of the Scientific Grant Agency of the Slovak Republic under the grant 1/0151/15, as well as the constribution of the Slovak Research and Development Agency under the project APVV 0136-12. This research was also supported by the Project: Mobility - enhancing research, science and education at the Matej Bel University, ITMS code: 26110230082, under the Operational Program Education cofinanced by the European Social Fund.

References

- F. Adams, A. Gendreau, O. Olmez, S.Y. Song, Construction of directed strongly regular graphs using block matrices, preprint, arXiv: 1311.0494.
- [2] A. Araluze, I. Kovács, K. Kutnar, L. Martinez, D. Marušič, Partial sum quadruples and bi-Abelian digraphs, J. Combin. Th. A 119(2012), 1811–1831.
- [3] J. Bosák, Partially directed Moore graphs, Math. Slovaca 29(1979), 181–196.
- [4] A.E. Brouwer, S. Hobart, Tables of directed strongly regular graphs, (November 2014), http://homepages.cwi.nl/~aeb/.
- [5] A.E. Brouwer, O. Olmez, S.Y. Song, Directed strongly regular graphs from 1¹/₂-designs, Europ. J. Combin. 33(2012), 1174–1177.
- [6] L.C. Chang, The uniqueness and nonuniqueness of triangular association schemes, Sci. Record 3(1959) 604–613.
- [7] L.C. Chang, Association schemes of partially balanced block designs with parameters v = 28, $n_1 = 12$, $n_2 = 15$ and $p_{1,1}^2 = 4$, Sci. Record 4(1960), 12–18.
- [8] G. Chartrand, L. Lesniak, Graphs & Digraphs (4th edition), Chapman & Hall/CRC, Boca Raton, 2005.
- [9] J.D. Dixon, B. Mortimer, *Permutation groups*. Graduate Texts in Mathematics, 163. Springer-Verlag, New York, 1996. xii+346 pp.
- [10] A.M. Duval, A directed graph version of strongly regular graphs, J. Combin. Th. A 47(1988), 71–100.

- [11] A.M. Duval, D. Iourinski, Semidirect product construction of directed strongly regular graphs, J. Combin. Th. A 104(2003), 157–167.
- [12] F. Fiedler, M.H. Klin, M. Muzychuk, Small vertex-transitive directed strongly regular graphs, Discrete Math. 255(2002), 87–115.
- [13] F. Fiedler, M. Klin, Ch. Pech, Directed strongly regular graphs as elements of coherent algebras. In: K. Denecke, H.-J. Vogel, (eds.) General Algebra and Discrete Mathematics: Proc. Conf. on General Algebra and Discrete Mathematics, Potsdam 1998, pp. 69–87. Shaker Verlag, Aachen (1999).
- [14] GAP Groups, Algorithms, Programming a System for Computational Discrete Algebra, www.gap-system.org.
- [15] C.D. Godsil, Algebraic Combinatorics, Chapman and Hall Mathematics Series. Chapman & Hall, New York, 1993. xvi+362 pp.
- [16] C.D. Godsil, S.A. Hobart, W.J. Martin, Representations of directed strongly regular graphs, Europ. J. Combin. 28(2007), 1980–1993.
- [17] S. Gyürki, M. Klin, Sporadic examples of directed strongly regular graphs obtained by computer algebra experimentation, in Gerdt, V.P. (ed.) et al., Computer algebra in scientific computing. CASC 2014, Proceedings. Berlin: Springer. Lecture Notes in Computer Science 8660 (2014), 155–170.
- [18] J.M. Hammersley, The friendship theorem and the love problem, pp. 31–54 in: Surveys in Combinatorics (E. Keith Lloyd, ed.), Cambridge University Press 1983.
- [19] M. Hirasaka, H. Kang, K. Kim, Characterization of association schemes by equitable partitions, Europ. J. Combin. 27(2006), 139–152.
- [20] S.A. Hobart, T.J. Shaw, A note on a family of directed strongly regular graphs, Europ. J. Combin. 20(1999), 819–820.
- [21] L.K. Jørgensen, Search for directed strongly regular graphs, Aalborg University preprint R-99-2016, October 1999.
- [22] L.K. Jørgensen, Non-existence of directed strongly regular graphs, Discrete Math. 264 (2003), 111–126.
- [23] L.K. Jørgensen, Directed strongly regular graphs with $\mu = \lambda$, Discrete Math. 231 (2001), 289–293.
- [24] L.K. Jørgensen, New mixed Moore graphs and directed strongly regular graphs, Discrete Math. 338 (2015), 1011–1016.
- [25] M. Klin, A. Munemasa, M. Muzychuk, P.H. Zieschang, Directed strongly regular graphs obtained from coherent algebras, Lin. Alg. Appl., 377 (2004), 83–109.

- [26] M. Klin, Ch. Pech, P.H. Zieschang, Flag algebras of block designs: I. Initial notions, Steiner 2-designs and generalized quadrangles. Preprint, MATH-AL-10-1998, Technische Universität Dresden (1998)
- [27] L. Martinez, A. Araluze, New tools for construction of directed strongly regular digraphs: Difference digraphs and partial sum families, J. Combin. Th. B, 100(2010), 720–728.
- [28] B. McKay, nauty user's guide, ver. 1.5, Technical Report TR-CS-90-02. Computer Science Department, Australian National Univ. (1990).
- [29] B. McKay, E. Spence, Classification of regular two-graphs on 36 and 38 vertices, Australasian J. of Combin., 24(2001), p.293.
- [30] O. Olmez, S.Y. Song, Some Families of Directed Strongly Regular Graphs Obtained from Certain Finite Incidence Structures, Graphs and Combinatorics, 30 (2014), 1529–1549.
- [31] A.J.L. Paulus, Conference matrices and graphs of order 26, Technische Hogeschool Eindhoven, report WSK 73/06, Eindhoven, 1983, 89 pp.
- [32] Ch. Pech, S. Reichard, *The* SetOrbit *package for* GAP, http://www.math.tu-dresden.de/~pech.
- [33] Ch. Pech, S. Reichard, *Enumerating set orbits*. in: M. Klin et al., Algorithmic Algebraic Combinatorics and Gröbner Bases, pp. 137–150. Springer, Berlin, Heidelberg (2009).
- [34] S. Reichard, COCO II, (personal communication).
- [35] L.H. Soicher, GRAPE: A system for computing with graphs and groups, Groups and computation (New Brunswick, 1991), DIMACS Ser. Discrete Math. Theoret. Comput. Sci. 11, pp. 287–291, Amer. Math. Soc., Providence, RI (1993)
- [36] G. Sabidussi, *Graph derivatives*, Math. Z. **76**(1961), 385–401.
- [37] G. Sabidussi, The lexicographic product of graphs, Duke Math. J. 28(1961), 573–578.
- [38] E. Spence, Regular two-graphs on 36 vertices, Lin. Alg. Appl. 226–228(1995), 459-497.
- [39] E. Spence, Strongly regular graphs on at most 64 vertices, http://www.maths.gla.ac.uk/~es/srgraphs.php
- [40] M. Ziv-Av, Interactions between coherent configurations and some classes of objects in extremal combinatorics, PhD thesis, Ben-Gurion University of the Negev, 2014, http://my.svgalib.org/phdfiles/thesis.pdf

Appendix

Parameter set	Parameter set			
of the π -join	of the basic graph	a	b	j
(40, 11, 4, 3, 3)	(8,3,2,1,1)	4	2	1
(50, 23, 13, 10, 11)	(10,3,0,1)	2	5	2
(60, 15, 4, 3, 4)	(12, 3, 1, 0, 1)	4	3	1
(60, 20, 7, 6, 7)	(6,2,1,0,1)	3	2	3
(60, 20, 7, 6, 7)	(15, 5, 2, 1, 2)	3	5	1
(70, 33, 18, 15, 16)	(10,3,0,1)	2	5	3
$(72,\!19,\!6,\!5,\!5)$	(8,3,2,1,1)	4	2	2
(72, 25, 11, 8, 9)	(18, 7, 5, 2, 3)	3	6	1
(72, 33, 19, 14, 16)	(24, 9, 7, 2, 4)	2	12	1
(72, 34, 20, 16, 16)	(24, 10, 8, 4, 4)	2	12	1
$(78,\!26,\!9,\!8,\!9)$	(6,2,1,0,1)	3	2	4
(78, 36, 23, 16, 17)	(26, 10, 3, 4)	2	13	1
(78, 37, 20, 17, 18)	(26, 11, 7, 4, 5)	2	13	1
(80, 22, 10, 6, 6)	(16, 6, 2, 2)	4	4	1
$(80,\!23,\!9,\!8,\!6)$	(16, 7, 5, 4, 2)	4	4	1
(90, 19, 5, 4, 4)	(15, 4, 2, 1, 1)	5	3	1
(90, 31, 13, 10, 11)	(9,4,1,2)	3	3	3
(90, 43, 23, 20, 21)	(10,3,0,1)	2	5	4
(90, 43, 23, 20, 21)	(18, 7, 5, 2, 3)	2	9	2
(90, 43, 23, 20, 21)	$(30,\!13,\!8,\!5,\!6)$	2	15	1
(96, 32, 11, 10, 11)	(6,2,1,0,1)	3	2	5
(96, 32, 11, 10, 11)	(24, 8, 3, 2, 3)	3	8	1
(96, 33, 15, 10, 12)	(24, 9, 7, 2, 4)	3	8	1
(96, 34, 16, 12, 12)	(24, 10, 8, 4, 4)	3	8	1
(102, 48, 29, 22, 23)	(34, 14, 12, 5, 6)	2	17	1
(102, 49, 26, 23, 24)	(34, 15, 9, 6, 7)	2	17	1
(104, 27, 8, 7, 7)	(8,3,2,1,1)	4	2	3
(105, 35, 12, 11, 12)	(15, 5, 2, 1, 2)	3	5	2
(108, 27, 7, 6, 7)	(12, 3, 1, 0, 1)	4	3	2
(108, 37, 15, 12, 13)	(27, 10, 6, 3, 4)	3	9	1
(108, 49, 29, 20, 24)	(36, 13, 11, 2, 6)	2	18	1
(108, 50, 32, 22, 24)	(36, 14, 4, 6)	2	18	1
(108, 51, 33, 24, 24)	(36, 15, 6, 6)	2	18	1
(110, 53, 28, 25, 26)	(10,3,0,1)	2	5	5

Table 1: New parameter sets up to $n \leq 110$ appearing in this paper.