PRIME VALUES OF $a^2 + p^4$

D.R. HEATH-BROWN AND XIANNAN LI

ABSTRACT. We prove an asymptotic formula for the number of primes of the shape $a^2 + p^4$, thereby refining the well known work of Friedlander and Iwaniec [4]. Along the way, we prove a result on equidistribution of primes up to x, in which the moduli may be almost as large as x^2 .

1. INTRODUCTION

Many remarkably difficult conjectures in prime number theory take the form that there are infinitely many primes in some set of natural numbers S. In many interesting examples, we even have conjectured asymptotic formulas for the number of primes in S. Thus, we think that there are infinitely many primes of the form p + 2, $a^2 + 1$, $a^2 + b^6$, and so on. Here, and everywhere in the paper, p shall always denote a prime.

The generality of our belief is in stark contrast with the paucity of examples for which we can prove our conjectures. In this paper, we are interested in the problem of finding primes in sequences which occur as the special values of a polynomial in two variables. For polynomials in one variable, only the linear case is understood, from the work of Dirichlet.

A classical result is that there are infinitely many primes of the form $a^2 + b^2$. Indeed, by a result of Fermat, primes of that form are essentially the same as primes of the form 4n + 1, so that this reduces to a special case of Dirichlet's theorem on primes in arithmetic progressions. Let us define the exponential density of the sequence of values of the polynomial P(a, b) as the infimum of those real λ for which

$$#\{P(a,b) \le x\} \ll x^{\lambda}.$$

Then the density of the sequence defined by $a^2 + b^2$ is 1, the same as the set of all natural numbers.

It is much more challenging to prove a similar result when the sequence given by P(a, b) has density less than 1. The first result in this direction was the breakthrough of Friedlander and Iwaniec [4] on the prime values of $a^2 + b^4$, which was followed by the result of Heath-Brown [8] on primes values of $a^3 + 2b^3$. It is worth mentioning that the density of the sequence is not the only measure of difficulty, as no results are available for prime values of $a^2 + b^3$ due to its lack of structure.

Aside from generalizations of Heath-Brown's result to more general cubic polynomials by Heath-Brown and Moroz [9], the theorems of Friedlander and Iwaniec [4] and Heath-Brown [8] remain the only results of this type. In this paper, we add the following example on prime values of $a^2 + p^4$.

Theorem 1.

$$\#\{a^2 + p^4 \le x : a > 0 \text{ and } a^2 + p^4 \text{ is prime}\} = \nu \frac{4Jx^{3/4}}{\log^2 x} \left(1 + O_\epsilon\left(\frac{1}{(\log x)^{1-\epsilon}}\right)\right),$$

²⁰¹⁰ Mathematics Subject Classification. Primary: 11M06, Secondary: 11M26.

where

$$J = \int_0^1 \sqrt{1 - t^4} dt,$$

and

$$\nu = \prod_{p} \left(1 - \frac{\chi_4(p)}{p-1} \right),$$

for χ_4 the non-principal character modulo 4.

In the statement of the Theorem above, and in the rest of the paper, ϵ denotes any sufficiently small, positive constant, not necessarily the same in each occurrence.

Remark 1. The reader may check that the number of elements $a^2 + p^4 \leq x$ is well approximated by $\frac{4Jx^{3/4}}{\log x}$.

Our method also applies to more general sequences $a^2 + y^4$ where y is restricted to a set Y which is regularly distributed and is not too sparse.

As in the works [4] and [8], the proof rests on establishing a level of distribution for the sequence, and the estimation of special bilinear sums. The first appears in classical sieve theory, and simply asks for good estimates for the remainder term in counting numbers of the form $a^2 + p^4$ divisible by a given integer d, averaged over d. The bilinear sums estimate is the ingredient which allows us to overcome the parity barrier, and as in previous works is the most significant part of the proof.

One ingredient used in Friedlander and Iwaniec's work on prime values of $a^2 + b^4$ is the regularity of the distribution of the squares of integers, which allows them to use a delicate harmonic analysis argument to extract certain main terms in their bilinear sum, and prove that the error terms are small on average (see Sections 4-9 in [4]). This is the portion of their work which overcomes the sparsity of their sequence.

This regularity does not exist in the case of squares of primes, and we need to develop a method which applies for more general sequences. In particular, we prove a result about the distribution of sequences in arithmetic progressions, which is similar in spirit to a general form of the Barban–Davenport–Halberstam theorem. In our case, for $a^2 + p^4 \leq x$, we have $p \leq x^{1/4}$, while the modulus appearing in our bilinear sum can be as large as $x^{1/2-\delta}$. We thus need an equidistribution result which holds when the modulus goes up to nearly the square of the length of the sum, in contrast to the Barban–Davenport–Halberstam theorem. Since this result is of independent interest, we first illustrate our result in the case of primes below.

Corollary 1. For (a,q) = 1, let

$$S(x; a, q) = \sum_{\substack{m,n \le x \\ n \equiv am \pmod{q}}} \Lambda(m) \Lambda(n).$$

Then for any A > 0, there exists B = B(A) such that

$$\sum_{q \le Q} \sum_{a \pmod{q}}^{*} \left| S(x;a,q) - \frac{x^2}{\phi(q)} \right|^2 \ll \frac{x^4}{\log^A x}$$

for $Q \le x^2 (\log x)^{-B}$.

At first sight it seems remarkable that the primes up to x should be well distributed for moduli as large as $x^2(\log x)^{-B}$, but this is essentially the conclusion of the Corollary. The

reader may verify however that the analogous result does not hold without the average over $a \pmod{q}$.

For our application, we will require a more general result which applies for sequences satisfying the Siegel–Walfisz condition, which is essentially saying that the sequence is equidistributed for small moduli. Remark 2 below makes this precise.

Remark 2. We say that an arithmetic function c(n) satisfies a Siegel–Walfisz condition if for any constant κ , we have that for any $q < (\log x)^{\kappa}$, and any nonprincipal character $\chi \pmod{q}$, we have

(1)
$$\sum_{n \le x} \chi(n) c(n) \ll_{\kappa} x^{1/2} ||c|| (\log x)^{-\kappa}.$$

Here $||c||^2 = \sum_n |c(n)|^2$ as usual.

This is known in the case $c(n) = \Lambda(n)$ by the Siegel–Walfisz Theorem so Corollary 1 follows immediately from Corollary 2 below.

Corollary 2. Let $c_1(n)$ and $c_2(n)$ be arithmetic functions supported on $n \le x$ with $c_1(n)$ satisfying the Siegel-Walfisz condition given by (1). For (a,q) = 1, let

$$S(x; a, q) = \sum_{\substack{m,n \le x \\ (mn,q)=1 \\ m \equiv an \pmod{q}}} c_1(m) c_2(n),$$

and let

$$S(x;q) = \frac{1}{\phi(q)} \left(\sum_{\substack{m \le x \\ (m,q)=1}} c_1(m) \right) \left(\sum_{\substack{n \le x \\ (n,q)=1}} c_2(n) \right).$$

Then for any A > 0, there exists B = B(A) such that for $Q \le x^2 (\log x)^{-B}$

$$\sum_{q \le Q} \sum_{a \pmod{q}}^{*} |S(x;a,q) - S(x;q)|^2 \ll \frac{x^2}{\log^A x} ||c_1\tau||^2 ||c_2\tau||^2.$$

Corollary 2 follows from the more technical Theorem 2 stated in Section 12 which applies to general sequences not necessarily satisfying the Siegel–Walfisz condition.

After using Corollary 2 to extract main terms in our bilinear sum, we still need to estimate the sum of these main terms. In Friedlander and Iwaniec's treatment, this involves a difficult direct estimation (see Sections 10-26 in [4]). We avail ourselves of their work in our estimates as well.

In our treatment, we compare our $a^2 + p^4$ with the sequence given by $a^2 + p^2$, previously studied by Fouvry and Iwaniec [2], which helps to streamline our arguments. For our result, we may also use the sequence $a^2 + b^2$ where b has no small prime factors below $(\log x)^A$, but using the result of Fouvry and Iwaniec is convenient and elegant. We now move to more precise definitions.

Since the number of $n = a^2 + p^4 \le x$ with $p < x^{1/4} / \log^2 x$ is bounded by

$$\sum_{p < \frac{x^{1/4}}{\log^2 x}} \sum_{a \le \sqrt{x-p^4}} 1 \ll \frac{x^{3/4}}{\log^3 x},$$

we may assume that $p \ge x^{1/4}/\log^2 x$. Let $x^{1/2}/\log^4 x \le X \le x^{1/2}$ and $\eta = (\log x)^{-1}$, and fix an interval $I = (X, X(1+\eta)]$. In our treatment of the bilinear sum, we need the two

sequences to behave alike even when restricted to small sets, so that it becomes necessary to introduce proper weights. Define the sequences $\mathcal{A} = \{a(n)\}$ and $\mathcal{B} = \{b(n)\}$ by

(2)
$$a(n) = \begin{cases} \sum_{\substack{n=a^2+p^4 \\ p^2 \in I \\ (a,p)=1 \\ 0 \\ \end{cases}} \log p & \text{if } n \le x \\ a(n) = \begin{cases} \sum_{\substack{n=a^2+p^4 \\ p^2 \in I \\ (a,p)=1 \\ 0 \\ \end{array}} \log p & \text{otherwise,} \end{cases}$$

and

$$b(n) = \begin{cases} \sum_{\substack{n=a^2+p^2 \ \text{log } p \\ p \in I \\ (a,p)=1 \\ 0 & \text{otherwise}} \end{cases}$$

Note that for any fixed natural number d and any real A > 0, we have

$$\sum_{d|n} a(n) = \sum_{d|n} b(n) + O_{d,A} \left(x (\log x)^{-A} \right).$$

It is this property and the choice of weights which makes the sequence \mathcal{B} suitable for our method.

Further, define

$$\pi(\mathcal{A}) = \sum_{p} a(p)$$

and

$$\pi(\mathcal{B}) = \sum_{p} b(p).$$

We claim that it suffices to show

Proposition 1. With notation as above,

$$\pi(\mathcal{A}) - \pi(\mathcal{B}) \ll \frac{1}{(\log x)^{2-\epsilon}} \mu(I),$$

where

$$\mu(I) = \int_{I} \sqrt{x - t^2} dt.$$

Note that $\mu(I) \leq \eta X \sqrt{x} \leq \eta x$ for all values of X, and $\mu(I) \gg \eta X \sqrt{x}$ for $X \leq \sqrt{x}/2$ and $\mu(I) \gg \eta^{3/2} x$ for $\sqrt{x}/2 \leq X \leq \sqrt{x}$.

We verify that our main Theorem follows from Proposition 1 in the next section. Most of this paper will be devoted to proving the Proposition.

We conclude this introduction by recording one convention of notation. We shall use the familiar convention that the positive number ε may vary between occurrences. This allows us to write $x^{\epsilon} \log x \ll x^{\epsilon}$, for example.

Acknowledgement. This work was supported by EPSRC grant EP/K021132X/1. We also thank Pierre Le Boudec for a careful reading of an earlier version of this paper. This resulted in the detection of a significant oversight, which has now been corrected.

2. Setting up the sieve

Let us first verify that Theorem 1 follows from Proposition 1. Note that the condition (a, p) = 1 in the definition of b(n) may be removed since

$$\sum_{p \in I} \log p \sum_{\substack{a \le \sqrt{x-p^2} \\ p \mid a}} 1 \ll x^{1/2+\epsilon}$$

Then, by the work of Fouvry and Iwaniec [2] and summation by parts, we have that

$$\pi(\mathcal{B}) = \frac{\nu\mu(I)}{\log x} \left(1 + O\left(\frac{1}{\log^{1-\epsilon} x}\right) \right),\,$$

for ν as given in Theorem 1. Thus Proposition 1 gives us that

$$\pi(\mathcal{A}) = \frac{\nu\mu(I)}{\log x} \left(1 + O\left(\frac{1}{\log^{1-\epsilon} x}\right) \right).$$

Our main result then follows by partial summation, possible since the length of I is short. To be precise, let $I_j = (X_j, X_j(1 + \eta)]$ be disjoint intervals for $1 \le j \le m$ such that

$$\cup_j I_j = (Y, x^{1/2}],$$

where $Y \gg \frac{x^{1/2}}{\log^2 x}$. Further, let \mathcal{A}_j be defined as in (2) with $I = I_j$. Recall that $\eta = \frac{1}{\log x}$, and note that the number of elements $a^2 + p^4 \leq x$ where p|a with $p^2 \in I$ is bounded by

$$\sum_{p^2 \in I} \frac{\sqrt{x}}{p} \ll x^{1/2 + \epsilon}.$$

Hence,

$$\begin{split} &\#\{a^2 + p^4 \le x : a^2 + p^4 \text{ is prime and } a > 0\} \\ &= \sum_j \frac{1}{\sqrt{X_j} \log X_j} \pi(\mathcal{A}_j) \left(1 + O\left(\frac{1}{\log x}\right) \right) + O(x^{1/2 + \epsilon}) \\ &= \frac{2\nu + O\left(\frac{1}{\log^{1 - \epsilon} x}\right)}{\log^2 x} \sum_j \frac{\mu(I_j)}{\sqrt{X_j}} + O(x^{1/2 + \epsilon}) \\ &= \frac{2\nu + O\left(\frac{1}{\log^{1 - \epsilon} x}\right)}{\log^2 x} \int_Y^{\sqrt{x}} \sqrt{x - t^2} \frac{dt}{\sqrt{t}} + O(x^{1/2 + \epsilon}) \\ &= \frac{4\nu x^{3/4}}{\log^2 x} \int_0^1 \sqrt{1 - t^4} dt \left(1 + O\left(\frac{1}{\log^{1 - \epsilon} x}\right) \right). \end{split}$$

Thus Theorem 1 follows from Proposition 1.

We now fix some basic notation. For $\mathcal{C} = \{c(n)\}\$ any sequence supported on $(\sqrt{x}, x]$, let

$$\pi(\mathcal{C}) = \sum_{p \le x} c(p),$$
$$\mathcal{C}_d = \{c(dn) : n \in \mathbb{N}\},$$

$$#\mathcal{C}_d = \sum_{d|n} c(n),$$

and

$$R_d(\mathcal{C}) = |\#\mathcal{C}_d - M_d(\mathcal{C})|,$$

for some $M_d(\mathcal{C})$ depending on d and \mathcal{C} . Note that the use of $\#\mathcal{C}_d$ here denotes the sum of the elements in \mathcal{C}_d rather than the number of elements in \mathcal{C}_d .

We shall prove Proposition 1 by applying the same sieving procedure to both sequences \mathcal{A} and \mathcal{B} . As mentioned in the Introduction, this requires a level of distribution result and an understanding of certain bilinear forms. We refer the reader to Friedlander and Iwaniec's asymptotic sieve for primes [6], Harman's alternative sieve [7] and Heath-Brown's proof of primes of the form $x^3 + 2y^3$ [8] for several perspectives on this. Here, we develop what we need from scratch along the lines of [8]. For our application, this eases some technical details involving the bilinear sum. We begin by stating a level of distribution result for \mathcal{A} and \mathcal{B} .

Proposition 2. Let

$$g(d) = \frac{\rho(d)}{d},$$

where $\rho(d)$ denotes the number of solutions to

$$a^2 + 1 \equiv 0 \pmod{d}.$$

Now define

$$M_d(\mathcal{C}) = g(d)\mu(I).$$

Then for any constants $A \ge 0$ and $k \ge 0$, there exists a constant B = B(A, k) such that for $D = \frac{x^{3/4}}{(\log x)^B}$, we have

$$\sum_{d \le D} \tau^k(d) R_d(\mathcal{C}) \ll \frac{x}{(\log x)^A};$$

for both C = A and C = B.

For $C = \mathcal{B}$, the Proposition holds in the larger range $D \leq \frac{x}{(\log x)^B}$, but we do not require this. Proposition 2 is a direct consequence of a result of Friedlander and Iwaniec [5] to allow for the weights we need and to include the $\tau(d)^k$ factor. The derivation of Proposition 2 is given in Section 3.

Fix for the rest of the paper $\delta = (\log x)^{\varpi^{-1}}$ for some small constant $\varpi > 0$ (ϖ must be smaller than the ϵ appearing in the statement of Theorem 1), and $Y = x^{1/3+1/48}$. In fact, the $\frac{1}{48}$ in the previous definition can be replaced with any positive number less than $\frac{1}{24}$. The following Lemma begins our sieving procedure.

Lemma 1. For any $x^{\delta} < Y < x^{1/2-\delta}$ and C = A and C = B, we have that

$$\pi(\mathcal{C}) = S_1(\mathcal{C}) - S_2(\mathcal{C}) - S_3(\mathcal{C}) + O\left(\frac{\delta}{\log x}\mu(I)\right),$$

where

$$S_1(\mathcal{C}) = S(\mathcal{C}, x^{\delta})$$

$$S_2(\mathcal{C}) = \sum_{x^{\delta} \le p < Y} S(\mathcal{C}_p, p)$$

$$S_3(\mathcal{C}) = \sum_{Y \le p < x^{1/2 - \delta}} S(\mathcal{C}_p, p).$$

Proof. By Buchstab's identity, we have

$$\pi(\mathcal{C}) = S(\mathcal{C}, x^{1/2}) = S_1(\mathcal{C}) - S_2(\mathcal{C}) - S_3(\mathcal{C}) - \sum_{x^{1/2-\delta} \le p \le x^{1/2}} S(\mathcal{C}_p, p).$$

Using Selberg's upper bound sieve, we have that

$$\sum_{x^{1/2-\delta} \le p \le x^{1/2}} S(\mathcal{C}_p, p) \le \sum_{x^{1/2-\delta} \le p \le x^{1/2}} S(\mathcal{C}_p, x^{1/10}) \\ \ll \sum_{x^{1/2-\delta} \le p \le x^{1/2}} \frac{\mu(I)}{p \log x} + \sum_{x^{1/2-\delta} \le p \le x^{1/2}} \sum_{d \le x^{1/5}} \tau_3(d) R_{dp}(\mathcal{C}) \\ \ll \frac{\delta\mu(I)}{\log x} + \frac{x}{\log^A x},$$

for any $A \ge 0$ by Proposition 2.

While $S_1(\mathcal{C})$ may be handled via the Fundamental Lemma, and $S_3(\mathcal{C})$ can be readily written in terms of a bilinear form in the right range, $S_2(\mathcal{C})$ requires more attention.

Lemma 2. With notation as in Lemma 1 and $n_0 = \left[\frac{\log Y}{\delta \log x}\right]$, we have

(3)
$$S_2(\mathcal{C}) = \sum_{1 \le n \le n_0} (-1)^{n-1} (T^{(n)}(\mathcal{C}) - U^{(n)}(\mathcal{C})),$$

where

$$T^{(n)}(\mathcal{C}) = \sum_{\substack{x^{\delta} \le p_n < \dots < p_1 < Y \\ p_1 \dots p_n < Y}} S(\mathcal{C}_{p_1 \dots p_n}, x^{\delta}),$$

and

$$U^{(n)}(\mathcal{C}) = \sum_{\substack{x^{\delta} \le p_{n+1} < \dots < p_1 < Y \\ p_1 \dots p_n < Y \le p_1 \dots p_{n+1}}} S(\mathcal{C}_{p_1 \dots p_{n+1}}, p_{n+1}).$$

Proof. Let

$$V^{(n)}(\mathcal{C}) = \sum_{\substack{x^{\delta} \le p_n < \dots < p_1 < Y \\ p_1 \dots p_n < Y}} S(\mathcal{C}_{p_1 \dots p_n}, p_n).$$

The proof of (3) follows immediately upon observing that $S_2(\mathcal{C}) = V^{(1)}(\mathcal{C})$ and from the identity

$$V^{(n)}(\mathcal{C}) = T^{(n)}(\mathcal{C}) - U^{(n)}(\mathcal{C}) - V^{(n+1)}(\mathcal{C}).$$

By Lemmas 1 and 2, in order to prove Proposition 1, it suffices to prove the following two propositions.

Proposition 3. Let Q be a set of squarefree numbers not exceeding Y. Then for any A > 0,

$$\left|\sum_{q\in Q} S(\mathcal{A}_q, x^{\delta}) - \sum_{q\in Q} S(\mathcal{B}_q, x^{\delta})\right| \ll_A \frac{x}{\log^A x}.$$

Note that Proposition 3 immediately implies that

$$|S_1(\mathcal{A}) - S_1(\mathcal{B})| \ll_A \frac{x}{\log^A x}.$$

Since $n_0 \simeq 1/\delta \ll \log x$, Proposition 3 also implies that

$$\sum_{1 \le n \le n_0} |T^{(n)}(\mathcal{A}) - T^{(n)}(\mathcal{B})| \ll_A \frac{x}{\log^A x}.$$

The rest of the pieces from the decompositions in Lemmas 1 and 2 are handled below.

Proposition 4. For any A > 0 and $n \ge 3$,

$$|S_3(\mathcal{A}) - S_3(\mathcal{B})| \ll \frac{x}{\log^A x}$$
, and
 $|U^{(n)}(\mathcal{A}) - U^{(n)}(\mathcal{B})| \ll \frac{x}{\log^A x}$.

For $n \leq 2$,

$$|U^{(n)}(\mathcal{A}) - U^{(n)}(\mathcal{B})| \ll \frac{\delta}{\log x} \mu(I).$$

Propositions 2 and 3 are proven in Sections 3 and 4 respectively. We reduce the proof of Proposition 4 to a related statement about bilinear sums in Section 5, which is in turn proven in a number of stages in Sections 6 to 11.

3. Level of distribution

Here, we prove Proposition 2. Define the sequences $\tilde{\mathcal{A}}$ and $\tilde{\mathcal{B}}$ by

$$\tilde{a}(n) = \begin{cases} \sum_{\substack{n=a^2+p^4 \ x^{1/4}\log x \\ p^2 \in I \\ p \nmid a \\ 0 \\ \end{cases}} & \text{if } \sqrt{x} < n \le x \\ x \le 1 \\ p < n \le 1 \\ 0 \\ 0 \\ \text{otherwise,} \end{cases}$$

and

$$\tilde{b}(n) = \begin{cases} \sum_{\substack{n=a^2+p^2 \ \log p \\ p \in I \\ p \nmid a \\ 0 \\ \end{cases}} & \text{if } \sqrt{x} < n \le x \\ p \neq a \\ 0 \\ \text{otherwise.} \end{cases}$$

Friedlander and Iwaniec's main result in [5] gives that for

$$M_d(\tilde{\mathcal{A}}) = g(d) \sum_{p^2 \in I} \frac{p \log p}{x^{1/4} \log x} \frac{p-1}{p} \sqrt{x-p^4},$$

and

$$M_d(\tilde{\mathcal{B}}) = g(d) \sum_{p \in I} \frac{\log p}{\log x} \frac{p-1}{p} \sqrt{x-p^2},$$

and any A > 0, there exists B = B(A) such that

$$\sum_{d < D} R_d(\tilde{\mathcal{A}}) \ll_A \frac{x^{3/4}}{\log^A x}$$

and

$$\sum_{d < D} R_d(\tilde{\mathcal{B}}) \ll_A \frac{x}{\log^A x},$$

for $D = \frac{x^{3/4}}{\log^B x}$. This immediately implies the corresponding results for \mathcal{A} and \mathcal{B} . Specifically, the following Lemma holds.

Lemma 3. For any A > 0, there exists B = B(A) such that

$$\sum_{d < D} R_d(\mathcal{C}) \ll \frac{x}{(\log x)^A}$$

for both $\mathcal{C} = \mathcal{A}$ and $\mathcal{C} = \mathcal{B}$, where $D = \frac{x^{3/4}}{\log^B x}$.

We want to prove a version of the above lemma which includes a $\tau(d)^k$ term. In order to develop this, we first state an elementary Lemma.

Lemma 4. For any $n, k \ge 1$, there exists a divisor d|n such that $d \le n^{1/2^k}$ and such that $\tau(n) \le 2^{2^k - 1} \tau(d)^{2^k}.$

Proof. Assume n > 1. We prove the result for k = 1, the rest following by induction on k. Let d|n with $d \leq \sqrt{n}$ be such that $\tau(d)$ is maximal. Write n = dd', and note that d' > 1. By maximality, any prime divisor p|d' satisfies $pd > \sqrt{n}$ so that $d'/p \le \sqrt{n}$. Again, by maximality, $\tau(d'/p) \leq \tau(d)$. Thus we have that $\tau(n) \leq \tau(d)\tau(d'/p)\tau(p) \leq 2\tau(d)^2$.

Now we prove the following trivial bound.

Lemma 5. Recall that
$$D = \frac{x^{3/4}}{\log^B x}$$
. For $C = \mathcal{A}$ and $C = \mathcal{B}$, we have

$$\sum_{d < D} \tau(d)^k \# C_d \ll x (\log x)^{2^{2k+3}}$$

Proof. We prove the result for $\mathcal{C} = \mathcal{A}$, the proof for $\mathcal{C} = \mathcal{B}$ being essentially the same. By Lemma 4,

$$\sum_{d < D} \tau(d)^k \# \mathcal{A}_d \le \sum_n \tau(n)^{k+1} a(n)$$
$$\ll \sum_{d \le \sqrt{x}} \tau(d)^{2k+2} \# \mathcal{A}_d$$

For $d \leq \sqrt{x}$,

$$#\mathcal{A}_d \ll \sum_{p^2 \in I} p \log p \sum_{\substack{a < \sqrt{x} \\ a^2 \equiv -p^4 \pmod{d} \\ (a,p)=1}} 1$$
$$\ll g(d)\sqrt{x} \sum_{p^2 \in I} p \log p$$
$$\ll g(d)x.$$

Using the bound $g(d) \leq \frac{\tau(d)}{d}$, we have

$$\sum_{d \le \sqrt{x}} \tau(d)^{2k+2} \# \mathcal{A}_d \ll x \sum_{d \le \sqrt{x}} \frac{\tau(d)^{2k+3}}{d} \ll x (\log x)^{2^{2k+3}}.$$

Now, we are ready to prove our Proposition 2. By Cauchy–Schwarz, we have

$$\sum_{d < D} \tau(d)^k R_d(\mathcal{C}) \ll \left(\sum_{d < D} \tau(d)^{2k} R_d(\mathcal{C}) \right)^{1/2} \left(\sum_{d < D} R_d(\mathcal{C}) \right)^{1/2} \\ \ll \left(\sum_{d < D} \tau(d)^{2k} (\#\mathcal{C}_d + M_d(\mathcal{C})) \right)^{1/2} \left(\sum_{d < D} R_d(\mathcal{C}) \right)^{1/2} \\ \ll \left(x (\log x)^{2^{4k+3}} \right)^{1/2} \left(\frac{x}{(\log x)^{\tilde{A}}} \right)^{1/2},$$

by Lemmas 3 and 5. This concludes the proof since we may make \tilde{A} as large as we like by choosing B suitably large.

4. Application of the Fundamental Lemma

We now prove Proposition 3. Recall that we are interested in studying

$$\sum_{q \in Q} S(\mathcal{C}_q, x^{\delta}),$$

for $\mathcal{C} = \mathcal{A}$ or $\mathcal{C} = \mathcal{B}$, $\delta = (\log x)^{\varpi^{-1}}$, and Q a set of square-free numbers not exceeding Y. The level of distribution provided by Proposition 2 is sufficient to derive the correct asymptotic for this quantity, with a small error term. To be precise, we apply an upper and lower bound sieve of level of distribution $x^{\frac{1}{4}}$ so that the sifting variable $s = \frac{1}{4\delta}$. For $z \geq 1$, we use the usual notation

$$V(z) = \prod_{p < z} \left(1 - g(p) \right)$$

By the Fundamental Lemma (see e.g. Corollary 6.10 in [4]) and Proposition 2, we have

1

$$\sum_{q \in Q} S(\mathcal{C}_q, x^{\delta}) = V(x^{\delta}) \sum_{q \in Q} g(q)\mu(I) \left(1 + O\left(\exp\left(-(4\delta)^{-1}\right)\right)\right) + O\left(\sum_{q \in Q} \sum_{d < x^{1/4}} R_{dq}(\mathcal{C})\right)$$
$$= V(x^{\delta}) \sum_{q \in Q} g(q)\mu(I) \left(1 + O\left(\frac{1}{\log^A x}\right)\right) + O\left(\sum_{d < x^{3/4-1/8}} \tau(d)R_d(\mathcal{C})\right)$$
$$= V(x^{\delta}) \sum_{q \in Q} g(q)\mu(I) \left(1 + O\left(\frac{1}{\log^A x}\right)\right) + O(x\log^{-A} x)$$

for any A > 0. Note that the last line is independent of whether $\mathcal{C} = \mathcal{A}$ or $\mathcal{C} = \mathcal{B}$. Thus

$$\sum_{q \in Q} S(\mathcal{A}_q, x^{\delta}) - \sum_{q \in Q} S(\mathcal{B}_q, x^{\delta}) \ll \frac{1}{\log^A x} \mu(I) \sum_{q \in Q} g(q) + x \log^{-A} x$$
$$\ll x \log^{-A+2} x,$$

upon noting that $g(q) \ll \frac{\tau(q)}{q}$.

5. Reduction of Proposition 4 to a bilinear form bound

We first rewrite $U^{(1)}$ and $U^{(2)}$ into a more convenient form.

Lemma 6. For C = A and C = B, and $U^{(j)}$ as defined in Lemma 2,

(4)

$$U^{(1)}(\mathcal{C}) = \sum_{\substack{x^{\delta} \le p_2 < p_1 < Y \\ Y \le p_1 p_2 < x^{1/2 - \delta}}} S(\mathcal{C}_{p_1 p_2}, p_2) + \sum_{\substack{x^{\delta} \le p_2 < p_1 < Y \\ p_1 p_2 \ge x^{1/2 + \delta}}} S(\mathcal{C}_{p_1 p_2}, p_2) + O\left(\frac{\delta\mu(I)}{\log x}\right)$$

$$=: U_1^{(1)}(\mathcal{C}) + U_2^{(1)}(\mathcal{C}) + O\left(\frac{\delta}{\log x}\mu(I)\right),$$

and

$$U^{(2)}(\mathcal{C}) = \sum_{\substack{x^{\delta} \le p_3 < \dots < p_1 < Y \\ p_1 p_2 < Y \le p_1 p_2 p_3 < x^{1/2 - \delta}}} S(\mathcal{C}_{p_1 p_2 p_3}, p_3) + \sum_{\substack{x^{\delta} \le p_3 < \dots < p_1 < Y \\ p_1 p_2 < Y \le p_1 p_2 p_3 \\ p_1 p_2 q_3 \ge x^{1/2 + \delta}}} S(\mathcal{C}_{p_1 p_2 p_3}, p_3) + O\left(\frac{\delta\mu(I)}{\log x}\right)$$
(5)
$$=: U_1^{(2)}(\mathcal{C}) + U_2^{(2)}(\mathcal{C}) + O\left(\frac{\delta\mu(I)}{\log x}\right).$$

Proof. To prove (4), it suffices to show that

(6)
$$\sum_{\substack{x^{\delta} \le p_2 < p_1 < Y\\ x^{1/2-\delta} \le p_1 p_2 < x^{1/2+\delta}}} S(\mathcal{C}_{p_1 p_2}, p_2) \ll \frac{\delta \mu(I)}{\log x}$$

In the sum above, $p_2 \ge x^{1/2-\delta}/p_1 > x^{1/2-\delta}/Y > x^{1/10}$, so that by Selberg's upper bound sieve, and Proposition 2, the left hand side of (6) is bounded by

$$\sum_{\substack{x^{\delta} \le p_2 < p_1 < Y \\ x^{1/2 - \delta} \le p_1 p_2 < x^{1/2 + \delta}}} S(\mathcal{C}_{p_1 p_2}, x^{1/10}) \ll \frac{\mu(I)}{\log x} \sum_{\substack{x^{1/10} < p_2 < p_1 < Y \\ x^{1/2 - \delta} \le p_1 p_2 < x^{1/2 + \delta}}} \frac{1}{p_1 p_2}$$
$$\ll \frac{\delta}{\log x} \mu(I).$$

Similarly, to prove (5), it suffices to show that

(7)
$$\sum_{\substack{x^{\delta} \le p_3 < p_2 < p_1 < Y \\ p_1 p_2 < Y \\ x^{1/2 - \delta} \le p_1 p_2 p_3 < x^{1/2 + \delta}}} S(\mathcal{C}_{p_1 p_2 p_3}, p_3) \ll \frac{\delta}{\log x} \mu(I).$$

In the sum above, $p_3 > x^{1/2-\delta}/Y > x^{1/10}$, so by Selberg's upper bound sieve, and Proposition 2, the quantity on the left hand side of (7) is bounded by

$$\sum_{\substack{x^{\delta} \le p_{3} < p_{2} < p_{1} < Y \\ p_{1}p_{2} < Y \\ x^{1/2-\delta} \le p_{1}p_{2}p_{3} < x^{1/2+\delta}}} S(\mathcal{C}_{p_{1}p_{2}p_{3}}, x^{1/10}) \ll \frac{\mu(I)}{\log x} \sum_{\substack{x^{1/10} < p_{3} < p_{2} < p_{1} < Y \\ x^{1/2-\delta} \le p_{1}p_{2}p_{3} < x^{1/2+\delta}}} \frac{1}{p_{1}p_{2}p_{3}} \ll \frac{\delta}{\log x} \mu(I).$$

(-> >

In order to simplify the conditions on the primes in our sieving functions the following lemma will be useful.

Lemma 7. Let $x^{-\delta} \leq \kappa \leq 1$. Then for any $P_1, P_2 \in [x^{\delta}, x^{1/3}]$ we have

$$\sum_{P_1 \le p_1 \le (1+\kappa)P_1} \sum_{P_2 \le p_2 \le (1+\kappa)P_2} \sum_{n \equiv 0 \pmod{p_1 p_2}} c(n)\tau(n) \ll \kappa^2 x (\log x)^{2^{17}} + \frac{x}{(\log x)^A}$$

for any A > 0.

Proof. By Lemma 4, n has at least one divisor $d \leq n^{1/16}$ such that $\tau(n) \ll \tau(d)^{16}$. Thus according to Proposition 2 we obtain

$$\sum_{P_1 \le p_1 \le (1+\kappa)P_1} \sum_{P_2 \le p_2 \le (1+\kappa)P_2} \sum_{n \equiv 0 \pmod{p_1 p_2}} c(n)\tau(n)$$

$$\ll \sum_{P_1 \le p_1 \le (1+\kappa)P_1} \sum_{P_2 \le p_2 \le (1+\kappa)P_2} \sum_{d \le x^{1/16}} \tau(d)^{16} \sum_{n \equiv 0 \pmod{dp_1 p_2}} c(n)$$

$$\ll \sum_{P_1 \le p_1 \le (1+\kappa)P_1} \sum_{P_2 \le p_2 \le (1+\kappa)P_2} \sum_{d \le x^{1/16}} \tau(d)^{16} (M_{dp_1 p_2}(\mathcal{C}) + R_{dp_1 p_2}(\mathcal{C}))$$

$$\ll \sum_{P_1 \le p_1 \le (1+\kappa)P_1} \sum_{P_2 \le p_2 \le (1+\kappa)P_2} \sum_{d \le x^{1/16}} \tau(d)^{17} \frac{x}{dp_1 p_2} + \frac{x}{(\log x)^A}.$$

To complete the proof we merely observe that if $P = P_1$ or P_2 then

$$\sum_{P \le p \le (1+\kappa)P} \frac{1}{p} \ll \kappa.$$

For $k \geq 3$, in the sum in $U^{(k)}(\mathcal{C})$, we have

$$Y \le p_1 \dots p_{k+1} < (p_1 \dots p_k)^{\frac{k+1}{k}} \le Y^{4/3} < x^{1/2-\delta}.$$

Thus if we define

$$U_*^{(k)}(\mathcal{C}) = \sum_{\substack{x^{\delta} \le p_{k+1} < \dots < p_1 < Y \\ p_1 \dots p_k < Y \le p_1 \dots p_{k+1} < x^{1/2-\delta}}} S(\mathcal{C}_{p_1 \dots p_{k+1}}, p_{k+1}),$$

then we will have

$$S_3(\mathcal{C}) = U_*^{(0)}(\mathcal{C}), \ U_1^{(1)}(\mathcal{C}) = U_*^{(1)}(\mathcal{C}), \ U_1^{(2)}(\mathcal{C}) = U_*^{(2)}(\mathcal{C}),$$

and

$$U^{(k)}(\mathcal{C}) = U^{(k)}_*(\mathcal{C}) \text{ for } k \ge 3.$$

If $p \in J = [V, (1 + \kappa)V)$ and an integer *n* is counted by $S(\mathcal{C}_{pq}, V)$ but not by $S(\mathcal{C}_{pq}, p)$, then *n* has at least two prime factors in *J*. In our application we have $V \leq x^{1/2-\delta}$ and $n \geq x(\log x)^{-8}$. Thus *n* will have at least one further prime factor. Thus $V^3 \leq n \leq x$ in this situation. A given integer *n* may be counted in many ways by $U_*^{(k)}(\mathcal{C})$. However the number of ways is at most the number of choices for $p_{k+1} < \ldots < p_1$ all dividing *n*. This will be

$$\left(\begin{array}{c}\omega(n)\\k+1\end{array}\right) \le 2^{\omega(n)} \le \tau(n).$$

The total contribution from such integers n is therefore bounded as in Lemma 7. Now, let

$$J(r) = [V_r, V_{r+1}) = [x^{\delta}(1+\kappa)^r, x^{\delta}(1+\kappa)^{r+1}), \ (r \ge 0)$$

Г		٦.
L		L
L		1

and let $R \ll \kappa^{-1} \log x$ be such that $x^{\delta}(1+\kappa)^R > x$. We then see that

$$U_*^{(k)}(\mathcal{C}) = \sum_{0 \le r \le R} \sum_{p \in J(r)} \sum_{\substack{p < p_k < \dots < p_1 < Y \\ p_1 \dots p_k < Y \le p_1 \dots p_k p < x^{1/2-\delta}}} S(\mathcal{C}_{p_1 \dots p_k p}, V_r) + O\left(\kappa x (\log x)^{1+2^{17}}\right) + O\left(\kappa^{-1} \frac{x}{\log^{A-1} x}\right).$$

This procedure enables us to reduce considerations to a bilinear sum. Indeed we have

(8)
$$\sum_{p \in J(r)} \sum_{\substack{p < p_k < \dots < p_1 < Y \\ p_1 \dots p_k < Y \le p_1 \dots p_k p < x^{1/2-\delta}}} S(\mathcal{C}_{p_1 \dots p_k p}, V_r) = \sum_{m,n} \alpha_m^{(r)} \beta_n^{(r)} c(mn)$$

where $\alpha_m^{(r)}$ is the characteristic function for the integers *m* all of whose prime factors are at least V_r , and $\beta_n^{(r)}$ is the characteristic function for integers $n = p_1 \dots p_k p$ with

$$p \in J(r), \quad p < p_k < \ldots < p_1 < Y \quad \text{and} \quad p_1 \ldots p_k < Y \le p_1 \ldots p_k p < x^{1/2-\delta}.$$

Note that $\beta_n^{(r)}$ is supported on integers $n \in [Y, x^{1/2-\delta})$.

The procedure for $U_2^{(1)}(\mathcal{C})$ and $U_2^{(2)}(\mathcal{C})$ will be somewhat different. As before we may use Lemma 7 to replace $S(\mathcal{C}_{p_1p_2}, p_2)$ in $U_2^{(1)}(\mathcal{C})$ by $S(\mathcal{C}_{p_1p_2}, V_r)$, when $p_2 \in J(r)$. For example, this yields

$$U_2^{(1)}(\mathcal{C}) = \sum_{0 \le r \le R} \sum_{\substack{p_2 \in J(r) \\ p_1 \ge x^{1/2+\delta}/p_2 \\ p_2 < p_1 < Y}} S(\mathcal{C}_{p_1 p_2}, V_r) + O(\kappa x (\log x)^{1+2^{17}}) + O\left(\kappa^{-1} \frac{x}{\log^{A-1} x}\right).$$

The sum on the right can be expressed as

$$\sum_{0 \le r \le R} \sum_{m,n} \alpha_m^{(r)} \beta_n^{(r)} c(mn),$$

where we now take $\alpha_m^{(r)}$ as the characteristic function for numbers $m = p_1 p_2$ with $p_2 \in J(r)$, $p_2 < p_1 < Y$, and $p_1 p_2 \ge x^{1/2+\delta}$, and $\beta_n^{(r)}$ as the characteristic function for numbers n all of whose prime factors are at least V_r . Since c(n) is supported in

$$(X^2, x] \subseteq (x(\log x)^{-8}, x]$$

we may assume that $\beta_n^{(r)}$ is supported in

$$\left(x(\log x)^{-8}Y^{-2}, x^{1/2-\delta}\right] \subseteq \left(x^{1/4+1/48}, x^{1/2-\delta}\right]$$

say. This is satisfactory for our purposes.

We may handle $U_2^{(2)}(\mathcal{C})$ in a precisely analogous fashion. On choosing $\kappa = (\log x)^{-A/2}$ we find that each of $S_3(\mathcal{C})$, $U_1^{(1)}(\mathcal{C})$, $U_2^{(1)}(\mathcal{C})$, $U_1^{(2)}(\mathcal{C})$, $U_2^{(2)}(\mathcal{C})$ and $U^{(k)}(\mathcal{C})$ (for $k \ge 3$), can be expressed as a sum of O(R) bilinear sums as in (8), together with an error term $O(x(\log x)^{1+2^{17}-A/2})$. Thus it will suffice to prove the following result.

Proposition 5. Fix any $\xi > 0$ and suppose $x^{1/4+\xi} \leq N \leq x^{1/2-\delta}$. Then, for coefficients α_m and β_n as above, we have

$$\sum_{N < n \le 2N} \sum_{m < x/N} \alpha_m \beta_n \left(a(mn) - b(mn) \right) \ll_{A,\xi} \frac{x}{\log^A x},$$

for any A > 0.

We note for future reference that α_m and β_n are supported on integers all of whose prime factors are at least x^{δ} . In particular they vanish unless m and n are odd. We also note that $|\alpha_m|, |\beta_n| \leq 1$ for all m, n.

6. The bilinear form over Gaussian integers

Our purpose is to prove Proposition 5. The expression $a^2 + p^4$ is a special value of the norm form of the Gaussian integers, and we now take advantage of that structure.

For $w, z \in \mathbb{Z}[i]$, let N(w) denote the usual Gaussian norm and

$$S_1(z,w) = \sum_{\substack{p^2 \in I \\ \operatorname{Re} \, \bar{w}z = p^2}} 2p \log p,$$

and

$$S_2(z,w) = \sum_{\substack{p \in I \\ \operatorname{Re} \, \bar{w}z = p}} \log p.$$

Note that both sums are either empty or contain only one term. We would now like to convert the sum over m and n present in Proposition 5 to a sum over Gaussian integers. We shall call $\gamma \in \mathbb{Z}[i]$ primitive if γ is not divisible by any rational prime.

Lemma 8. Let $\gamma \in \mathbb{Z}[i]$ be primitive and coprime to 2, and let m be a positive integer such that $m|N(\gamma)$. Then there exist exactly four associate choices for $\lambda \in \mathbb{Z}[i]$ such that $\lambda|\gamma$ and $N(\lambda) = m$. Of these exactly one has $\operatorname{Re}(\lambda)$ positive and odd.

Proof. Suppose the ideal (γ) factors as

$$(\gamma) = P_1^{e_1} \dots P_k^{e_k}.$$

Since γ is primitive and coprime to 2, we have $P_i \neq \overline{P_j}$ for every pair i, j. Moreover $N(P_i)$ will be a rational prime p_i , and we will have

$$m = p_1^{f_1} \dots p_k^{f_k},$$

with exponents $f_i \leq e_i$. It is then clear that (λ) must be

$$(\lambda) = P_1^{f_1} \dots P_k^{f_k},$$

and the result follows.

By Lemma 8,

$$a(mn) = \frac{1}{2} \sum_{\substack{N(w)=m \\ \bar{w}z \text{ primitive}}} S_1(z, w),$$

and

$$b(mn) = \frac{1}{2} \sum_{\substack{N(w)=m \\ \bar{w}z \text{ primitive}}} \sum_{\substack{N(z)=n \\ \bar{w}z \text{ primitive}}} S_2(z, w),$$

where we restrict z to have $\operatorname{Re}(z)$ positive and odd in both sums. Note that the double sum counts pairs with $\overline{w}z = p^2 + ia$ (or p + ia) with no restriction on the sign of a. In our original definition of a(n) and b(n), we have the condition a > 0, and this is accounted for by the factor of $\frac{1}{2}$.

We let $\beta_z = \beta_{N(z)}$ and $\alpha_w = \alpha_{N(w)}$. It now suffices to show that

(9)
$$\sum_{z} \sum_{\substack{w \\ \bar{w}z \text{ primitive}}} \beta_z \alpha_w \left(S_1(z, w) - S_2(z, w) \right) \ll_A \frac{x}{\log^A x},$$

for any A > 0, and for coefficients β_z and α_w satisfying $|\beta_z| \leq 1$ and $|\alpha_w| \leq 1$. Further, we may assume that β_z is supported on primitive z satisfying $2 \nmid \operatorname{Re}(z) > 0$ and $N \leq N(z) < 2N$, while α_w is supported on primitive w such that $N(w) \leq M := x/N$. Note that N < M since $N \leq x^{1/2-\delta}$. We also remark that β_z is supported on values with N(z)free of small prime factors. Hence N(z) is odd, and since $\operatorname{Re}(z)$ is also odd we must have $z \equiv 1 \pmod{2}$.

We first remove the primitivity condition on $\bar{w}z$ with negligible error. Indeed, the contribution of $S_1(z, w)$ for imprimitive $\bar{w}z$ of the form

$$\bar{w}z = p^2 + ia$$

must have p|a so that

$$\bar{w}z = p^2 + ibp,$$

for $p^2 \in I$, whence $b \leq \frac{x^{1/2}}{p} \leq x^{1/4} \log^2 x$. Hence, there are at most $x^{1/4+\epsilon}$ choices for b, and thus at most $x^{1/4+\epsilon}$ choices for w and z, given p. These are counted with weight $2p \log p \ll x^{1/4+\epsilon}$. Thus, the total contribution is bounded by

$$\sum_{p < x^{1/4}} x^{1/2 + \epsilon} \ll x^{3/4 + \epsilon} \ll \frac{x}{\log^A x}$$

The contribution from $S_2(z, w)$ for imprimitive $\bar{w}z$ is bounded similarly.

Let $\theta(z) = \arg z \in [0, 2\pi)$ and

$$\mathcal{R} = \mathcal{R}(A) = \{ z \in \mathbb{Z}[i] : N \le N(z) < 2N, |\theta(z) - k\pi/2| \le (\log x)^{-A} \text{ for } k \in \mathbb{Z} \}.$$

We now note that we may discard the part of the sum (9) with $z \in \mathcal{R}$.

Lemma 9. Suppose that both z and q are fixed. Then the number of possible w with $q = \operatorname{Re} \bar{w}z$ is

$$\ll \frac{\sqrt{M}}{\sqrt{N}}.$$

Proof. Let

$$z = s + it$$
$$w = u + iv,$$

so that

(10)
$$q := \operatorname{Re} \bar{w}z = us + vt$$

We have either $|s| \gg \sqrt{N}$ or $|t| \gg \sqrt{N}$. We deal with the case $|s| \gg \sqrt{N}$, the other case being similar. Since z is primitive, (s,t) = 1 so we may write

$$v \equiv \bar{t}q \pmod{s}.$$

Thus, there are $\ll \sqrt{\frac{M}{N}}$ choices for v. Once v is fixed, u is uniquely determined by (10).

Lemma 10.

$$\sum_{z \in \mathcal{R}} \sum_{w} \beta_z \alpha_w S_j(z, w) \ll x (\log x)^{-A}$$

for j = 1, 2.

Proof. We apply Lemma 9 to get

$$\sum_{z \in \mathcal{R}} \sum_{w} \beta_{z} \alpha_{w} S_{1}(z, w) \ll \sum_{\substack{z \in \mathcal{R} \\ z \text{ primitive}}} \sum_{p^{2} \in I} p \log p \sum_{\text{Re } \bar{w} z = p^{2}} 1$$
$$\ll \sqrt{\frac{M}{N}} \sum_{p^{2} \in I} p \log p \sum_{z \in \mathcal{R}} 1$$
$$\ll \sqrt{\frac{M}{N}} N(\log x)^{-A} \sum_{p^{2} \in I} p \log p$$
$$\ll x (\log x)^{-A}.$$

In the case of $S_2(z, w)$, the sum is simpler and we get

$$\sum_{z \in \mathcal{R}} \sum_{w} \beta_{z} \alpha_{w} S_{2}(z, w) \ll \sqrt{\frac{M}{N}} \sum_{p \in I} \log p \sum_{z \in \mathcal{R}} 1$$
$$\ll x (\log x)^{-A}.$$

In the sequel, let $\sum_{i=1}^{b}$ denote a sum over primitive $z \notin \mathcal{R}$ for which $\operatorname{Re}(z)$ is positive and $z \equiv 1 \pmod{2}$. Then Cauchy–Schwarz gives that

$$\left(\sum_{w} \alpha_{w} \sum_{z} \beta_{z} (S_{1}(z,w) - S_{2}(z,w))\right)^{2} \leq \sum_{w} \alpha_{w}^{2} \sum_{w} \left(\sum_{z} \beta_{z} (S_{1}(z,w) - S_{2}(z,w))\right)^{2},$$

where we now extend the sum over w over all Gaussian integers w satisfying $N(w) \leq x/N$, possible by positivity. We then see that it suffices to show that

(11)
$$\sum_{z_1, z_2}^{\flat} \beta_{z_1} \beta_{z_2} \sum_{w} \left(S_1(z_1, w) - S_2(z_1, w) \right) \left(S_1(z_2, w) - S_2(z_2, w) \right) \ll \frac{xN}{\log^A x}$$

for any A > 0.

Lemma 11. The contribution of the diagonal term $z_1 = z_2$ in (11) is at most

$$\sum_{z} \beta_{z}^{2} \sum_{w} \left(S_{1}(z,w)^{2} - 2S_{1}(z,w)S_{2}(z,w) + S_{2}(z,w)^{2} \right) \ll x^{1-\xi/2} N.$$

Proof. Since it is impossible for Re $\bar{w}z$ to be both a prime and the square of a prime, $S_1(z,w)S_2(z,w) = 0$. Let us record the trivial bounds

(12)
$$\sum_{z} \sum_{w} S_{1}(z, w) \ll \sum_{n \le x} \left(\sum_{\substack{a^{2} + p^{4} = n \\ p^{2} \in I \\ 16}} 2p \log p \right) \tau(n) \ll x^{1+\epsilon}$$

and similarly

(13)
$$\sum_{z} \sum_{w} S_2(z,w) \ll x^{1+\epsilon}$$

Now

$$\sum_{z} \sum_{w} S_1(z,w)^2 + S_2(z,w)^2 \ll \sqrt{X} \log X \sum_{z} \sum_{w} S_1(z,w) + \log X \sum_{z} \sum_{w} S_2(z,w) \\ \ll x^{5/4+\epsilon},$$

by (12) and (13). Since $N > x^{1/4+\xi}$ this suffices on choosing ϵ sufficiently small.

Thus, in considering (11), we will assume that $z_1 \neq z_2$. For any pair z_1, z_2 , we let $\theta = \theta(z_1, z_2) = \arg z_2 - \arg z_1$ denote the angle between z_1 and z_2 . Moreover, we define $\Delta = \Delta(z_1, z_2) = \operatorname{Im} \bar{z}_1 z_2 = |z_1 z_2| \sin \theta(z_1, z_2)$. Note that z_1 and z_2 being primitive and $z_1 \neq z_2$ implies that $\theta \neq 0$. Further, Re $z_i > 0$ implies that $\theta \neq \pi$. Hence $\Delta \neq 0$. Since $z_1 \equiv z_2 \equiv 1 \pmod{2}$ we will have $2 \mid \Delta$.

Remark 3. For ease of notation, we restrict our attention to those z_1, z_2 satisfying $\Delta > 0$, and henceforth assume this condition to be included in \sum_{z_1, z_2}^{\flat} .

If we write

Re $\bar{w}z_i = q_i$

for i = 1, 2, then

(14)
$$w = -i\Delta(z_1, z_2)^{-1}(q_1z_2 - q_2z_1).$$

Of course, we must have

(15)
$$q_1 z_2 \equiv q_2 z_1 \pmod{\Delta}.$$

Let $C(q_1, q_2, z_1, z_2)$ be the statement that q_1, q_2, z_1 and z_2 satisfy (15). From (14) and since $N(w) \leq x/N$, we have the additional condition

(16)
$$|q_1 z_2 - q_2 z_1| \le \Delta(z_1, z_2) \sqrt{\frac{x}{N}}$$

We also wish to dispose of the case in which Δ is small. In particular, we wish to only consider those z_1, z_2 such that

(17)
$$\Delta(z_1, z_2) > \mathfrak{D}_0 := N(\log x)^{-A-6}.$$

For brevity, let

$$f(q) = \begin{cases} 2p \log p & \text{if } q = p^2 \in I \\ 0 & \text{otherwise,} \end{cases}$$

and

$$g(q) = \begin{cases} \log p & \text{if } q = p \in I \\ 0 & \text{otherwise.} \end{cases}$$

 Set

$$h(q) = f(q) - g(q).$$
¹⁷

For any $J \subset I$, we have by the Prime Number Theorem that

$$\sum_{q \in J} h(q) = O\left(\frac{\sqrt{x}}{\log^C x}\right),$$

for any C > 0. This is a result of our choice of weights.

The conditions (16) and (17) are quite awkward, so we shall remove them by dissecting our sum in (11) into smaller pieces. To be precise, for some constant L to be determined, let

$$\omega_1 \asymp \omega_2 \asymp \omega := (\log x)^{-L}$$

and let $I = (X, X(1 + \eta)]$ be a disjoint union of intervals J of length $\simeq X\omega_1$. We need $\ll 1/\omega_1$ such intervals to cover I. Further, split the sum over z_1 and z_2 into regions \mathcal{U} , where each \mathcal{U} is of the form

$$\mathcal{U}(c,\theta_0) = \mathcal{U} := \{ z : c\sqrt{N} < |z| \le c(1+\omega_1)\sqrt{N}, \theta_0 < \arg(z) < \theta_0 + \omega_2 \},$$

for fixed $1 \leq c < \sqrt{2}$ and θ_0 . Note that we may chose ω_1 and ω_2 so that our regions \mathcal{U} form a partition of the region $\{z : N \leq N(z) < 2N, \operatorname{Re}(z) > 0\} - \mathcal{R}$. The number of regions needed for the sum over z_1 and z_2 is $O(\log^{4L} x)$. Here, we have allowed ω_1 to possibly be distinct from ω_2 in order to cover our region perfectly. They are the same size and can frequently be replaced by ω in our estimates.

Now, write $\mathfrak{C}_1(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ as the condition that all $(z_1, z_2, q_1, q_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \times J_1 \times J_2$ satisfy (16) and (17). Also, let $\mathfrak{C}_2(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ be the condition that there exists some (z_1, z_2, q_1, q_2) in $\mathcal{U}_1 \times \mathcal{U}_2 \times J_1 \times J_2$ which satisfies (16), and there exists some (z'_1, z'_2, q'_1, q'_2) in $\mathcal{U}_1 \times \mathcal{U}_2 \times J_1 \times J_2$ which does not satisfy (16). Finally, let $\mathfrak{C}_3(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ be the condition that all (z_1, z_2, q_1, q_2) in $\mathcal{U}_1 \times \mathcal{U}_2 \times J_1 \times J_2$ satisfy (16) but there exists some (z_1, z_2, q_1, q_2) in $\mathcal{U}_1 \times \mathcal{U}_2 \times J_1 \times J_2$ which does not satisfy (17).

For $\mathcal{U}_1, \mathcal{U}_2, J_1, J_2$ satisfying $\mathfrak{C}_1(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$, set

$$T(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2) = \sum_{\substack{z_1 \in \mathcal{U}_1 \\ z_2 \in \mathcal{U}_2}}^{\flat} \beta_{z_1} \beta_{z_2} \sum_{\substack{q_1 \in J_1 \\ q_2 \in J_2 \\ C(q_1, q_2, z_1, z_2)}} h(q_1) h(q_2),$$

and otherwise set $T(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2) = 0.$

Further, let

$$T'(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2) = \sum_{\substack{z_1 \in \mathcal{U}_1 \\ z_2 \in \mathcal{U}_2}}^{\flat} \sum_{\substack{q_1 \in J_1 \\ q_2 \in J_2 \\ C(q_1, q_2, z_1, z_2)}} |h(q_1)h(q_2)|.$$

Then (11) reduces to proving that for any constant A > 0, (18)

$$\sum_{\substack{\mathcal{U}_1,\mathcal{U}_2,J_1,J_2\\\mathfrak{C}_1(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2)}} T(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2) + \sum_{\substack{\mathcal{U}_1,\mathcal{U}_2,J_1,J_2\\\mathfrak{C}_2(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2) \text{ or } \mathfrak{C}_3(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2)}} T'(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2) \ll_A \frac{xN}{\log^A x}$$

Since L may be freely chosen, it suffices to prove the following Propositions.

Proposition 6. With notation as above and for $L \ge A + 6$, we have

$$\sum_{\substack{\mathcal{U}_1,\mathcal{U}_2,J_1,J_2\\\mathfrak{C}_2(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2) \text{ or } \mathfrak{C}_3(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2)}} T'(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2) \ll \frac{xN}{\log^A x}$$

Proposition 7. With notation as above and for fixed J_1 , J_2 and for L = 6A + 52 we have that

$$\sum_{\substack{\mathcal{U}_1,\mathcal{U}_2\\(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2)}} T(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2) \ll \frac{xN}{\log^{A+2L}x}.$$

Note that (18) follows from the Propositions above.

 \mathfrak{C}_1

Remark 4. When $\mathfrak{C}_1(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ holds and $z_i \in \mathcal{U}_i$, we automatically have $\operatorname{Re}(z_i) > 0$, $z_i \notin \mathcal{R}$ and $\Delta(z_1, z_2) > 0$.

7. PROOF OF PROPOSITIONS 6 AND 7; PRELIMINARY STEPS

We first note that we may essentially assume that q_1q_2 is coprime with Δ in the sums defining $T(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ and $T'(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$.

Lemma 12. We have that

$$\sum_{z_1, z_2}^{\flat} \sum_{\substack{q_1 \in J_1, q_2 \in J_2\\C(q_1, q_2, z_1, z_2)\\(q_1 q_2, \Delta) > 1}} |h(q_1)h(q_2)| \ll N^2 \sqrt{x} \log^3 x.$$

Proof. Note the number of z_1, z_2 appearing in the sum is $O(N^2)$, and that $h(q_1)h(q_2) \ll \sqrt{x \log^2 x}$. Hence, it suffices to show that for fixed z_1 and z_2 , the number of choices for q_1 and q_2 is bounded by $O(\log x)$.

Suppose $(q_1, \Delta) > 1$. We have that either $q_1 = p$ or $q_1 = p^2$ for some prime p, so $p|\Delta$. The congruence

$$q_1 z_2 \equiv q_2 z_1 \pmod{\Delta}$$

implies that $p|q_2$ as well, since z_1 is primitive. Thus $q_2 = p$ or $q_2 = p^2$ as well. Since the number of prime factors of Δ is $O(\log x)$, we have that the number of choices for p is $O(\log x)$, which suffices.

If $(q_i, \Delta) = 1$, (15) is equivalent to there existing $a \pmod{\Delta}$ with $(a, \Delta) = 1$ such that

$$q_1 \equiv aq_2 \pmod{\Delta}$$
$$az_2 \equiv z_1 \pmod{\Delta}.$$

Then, by Lemma 12, we may rewrite $T(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ as

(19)
$$T(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2) = \sum_{D \le 2N} \sum_{a \pmod{D}}^* Y(a, D; h, h) Z(a, D) + O(N^2 \sqrt{x} \log^3 x)$$

where

$$Z(a, D) = \sum_{\substack{(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ \Delta = D \\ az_2 \equiv z_1 \pmod{D}}}^{\flat} \beta_{z_1} \beta_{z_2}$$

and

$$Y(a, D; h_1, h_2) = Y(a, D) = \sum_{\substack{q_1 \in J_1, q_2 \in J_2 \\ q_1 \equiv aq_2 \pmod{D} \\ (q_1q_2, D) = 1}} h_1(q_1)h_2(q_2)$$

The rewriting of $T(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ in (19) separates the sum Z(a, D) containing the coefficients β_z from the congruence sum Y(a, D) involving the primes. This key procedure

has transformed the sum into the right form for us to extract the main terms from $T(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ using Corollary 2. Of course, we also need some understanding of the behaviour of Z(a, D) for which the following bounds will suffice for the moment.

Lemma 13. Let

$$\tilde{Z}(a,D) = \sum_{\substack{(z_1,z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ \Delta = D \\ az_2 \equiv z_1 \pmod{D}}}^{\flat} 1.$$

We have

(20)
$$\sum_{D} \tau(D) \sum_{a \pmod{D}}^{*} \tilde{Z}(a, D) \ll \omega^{4} N^{2} (\log x)^{16},$$

(21)
$$\sum_{\mathcal{U}_1,\mathcal{U}_2} \sum_{a \pmod{D}}^* \tilde{Z}(a,D) \ll N,$$

and

(22)
$$\sum_{a \pmod{D}}^{*} \tilde{Z}(a, D)^{2} \ll (\log x)^{3} \frac{N^{2}}{D} \tau(D)^{6}.$$

Proof. We write $z_k = x_k + iy_k$ for k = 1, 2, and assume without loss of generality that $|x_2|$ is maximal among $|x_1|, |x_2|, |y_1|$ and $|y_2|$. For (20) we apply Lemma 4 with k = 2 to deduce that

$$\sum_{D} \tau(D) \sum_{\substack{a \pmod{D} \\ a \pmod{D}}} \tilde{Z}(a, D) \\ \ll \sum_{\substack{d \le (2N)^{1/4}}} \tau(d)^4 \# \{ (z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 : (x_1, y_1) = 1, d \mid x_1 y_2 - x_2 y_1 \}.$$

Since the regions \mathcal{U}_i are contained in squares of side $O(\omega\sqrt{N})$, and $N^{1/4} \ll \omega\sqrt{N}$ we deduce that

$$\sum_{D} \tau(D) \sum_{a \pmod{D}}^{*} \tilde{Z}(a, D)$$

$$\ll \sum_{d \le (2N)^{1/4}} \tau(d)^4 \frac{\omega^4 N^2}{d^4} \#\{(z_1, z_2) \pmod{d} : (x_1, y_1, d) = 1, d \mid x_1 y_2 - x_2 y_1\}.$$

One can easily show that if d is a prime power p^e , then

$$#\{(z_1, z_2) \pmod{d} : (x_1, y_1, d) = 1, d \mid x_1 y_2 - x_2 y_1\} \le d^3,$$

whence the same bound holds for all d, and we obtain

$$\sum_{D} \tau(D) \sum_{a \pmod{D}}^{*} \tilde{Z}(a, D) \ll \omega^4 N^2 \sum_{d \le (2N)^{1/4}} \frac{\tau(d)^4}{d} \ll \omega^4 N^2 (\log x)^{16}$$

as required.

For (21) we note that the condition $\Delta = D$ implies $x_1y_2 - x_2y_1 = D$, and since $(x_2, y_2) = 1$ we have $x_1 \equiv \overline{y}_2 D \pmod{|x_2|}$. However we arranged that $|x_1| \leq |x_2|$, so that there are at most 2 possibilities for x_1 once x_2 and y_2 are given. Since x_1, x_2 and y_2 determine y_1 via the equation $x_1y_2 - x_2y_1 = D$, the required bound O(N) follows.

Finally, to prove (22) we decompose $\tilde{Z}(a, D)$ into 4 parts according to which of $|x_1|$, $|x_2|$, $|y_1|$ and $|y_2|$ is maximal. Using Cauchy's inequality it then suffices to handle the

analogue of $\tilde{Z}(a, D)$ in which $|x_2|$, say, is largest. If $(x_1, x_2, D) = k$, say, we see that the congruence $x_1y_2 \equiv D \pmod{|x_2|}$ determines at most 2k values of y_2 with $|y_2| \leq |x_2|$. Now, note that if we have two solutions $x_1 \equiv ax_2 \pmod{D}$ and $x'_1 \equiv ax'_2 \pmod{D}$, then $x_1x'_2 \equiv x_2x'_1 \pmod{D}$. Thus, we have

$$\sum_{a \pmod{D}}^{*} \tilde{Z}(a, D)^{2} \ll \sum_{\substack{x_{1}, x_{2}, x_{1}', x_{2}'\\x_{1}x_{2}' \equiv x_{2}x_{1}' \pmod{D}}} (x_{1}, x_{2}, D)(x_{1}', x_{2}', D).$$

However $(x_1, x_2, D)(x'_1, x'_2, D)$ divides $(x_1x'_2, x_2x'_1, D^2)$. Thus,

$$\sum_{a \pmod{D}} \tilde{Z}(a, D)^2 \ll \sum_{\substack{m,n \le 2N \\ m \equiv n \pmod{D}}} \tau(m)\tau(n)(m, n, D^2)$$
$$\leq \frac{1}{2} \sum_{\substack{m,n \le 2N \\ m \equiv n \pmod{D}}} (\tau(m)^2(m, D^2) + \tau(n)^2(n, D^2))$$
$$\ll \frac{N}{D} \sum_{m \le 2N} \tau(m)^2(m, D^2).$$

Finally

$$\begin{split} \sum_{m \leq 2N} \tau(m)^2(m, D^2) &\leq \sum_{d \mid D^2} d \sum_{\substack{m \leq 2N \\ d \mid m}} \tau(m)^2 \\ &\leq \sum_{d \mid D^2} d \sum_{v \leq 2N/d} \tau(v)^2 \tau(d)^2 \\ &\ll \sum_{d \mid D^2} d\{Nd^{-1}\log^3 N\} \tau(d)^2 \\ &\ll N\tau(D^2)^3\log^3 N \\ &\ll N\tau(D)^6\log^3 N \end{split}$$

which suffices to prove (22).

Remark 5. Trivially $|Z(a, D)| \leq \tilde{Z}(a, D)$ so that Lemma 13 applies with $\tilde{Z}(a, D)$ replaced by Z(a, D).

Now for an interval J and any function \tilde{h} , let

$$Y(J,\tilde{h};D) = \sum_{\substack{q \in J \\ (q,D)=1}} \tilde{h}(q)$$

and

$$Y_{h_1,h_2}(D) = Y(D) = \frac{1}{\phi(D)}Y(J_1,h_1;D)Y(J_2,h_2;D).$$

Recall that q_1 and q_2 appearing in Y(a, D) satisfy $(q_1q_2, D) = 1$. If h_1 or h_2 is g, then Y(D) is the expected value of Y(a, D).

If $h_1 = h_2 = f$, note that $p_1^2 \equiv ap_2^2 \pmod{D}$ implies that $p_1 \equiv bp_2 \pmod{D}$ for some b such that $a \equiv b^2 \pmod{D}$. Here, Y(a, D) = 0 if a is not a square modulo D so

$$\sum_{a \pmod{D}}^{*} Y(a, D) Z(a, D) = \sum_{b \pmod{D}}^{*} Y_f(b, D) Z(b^2, D),$$

where

$$Y_f(b, D) = \sum_{\substack{p_1^2 \in J_1 \\ p_2^2 \in J_2 \\ p_1 \equiv bp_2 \pmod{D} \\ (p_1p_2, D) = 1}} f(p_1^2) f(p_2^2).$$

When $h_1 = h_2 = f$, Y(D) is the expected value of $Y_f(b, D)$.

The following proposition makes the above discussion precise.

Proposition 8. If either $h_1 = g$ or $h_2 = g$, let

$$\mathcal{E}(N) = \sum_{D \le 2N} \sum_{a \pmod{D}}^{*} |Y(a, D; h_1, h_2) - Y_{h_1, h_2}(D)| \tilde{Z}(a, D)$$

Then for any constant C > 0,

(23)
$$\mathcal{E}(N) \ll_C \frac{xN}{\log^C x}.$$

For $h_1 = h_2 = f$, let

$$\mathcal{E}_f(N) = \sum_{D \le 2N} \sum_{b \pmod{D}}^* |Y_f(b, D) - Y_{f,f}(D)| \tilde{Z}(b^2, D).$$

Then for any constant C > 0,

$$\mathcal{E}_f(N) \ll_C \frac{xN}{\log^C x}.$$

Proof. We first prove the bound (23) for \mathcal{E} . We prove the result for $h_1 = g$ and $h_2 = f$. The proof is similar for the cases $h_1 = g$ and $h_2 = f$ and $h_1 = h_2 = g$. We write

$$\mathcal{E}(N) \le \sum_{D \le 2N} \sum_{\substack{q_2 \in J_2 \\ (q_2,D)=1}} f(q_2) \sum_{\substack{a \pmod{D}}}^* \left| \sum_{\substack{q_1 \in J_1 \\ q_1 \equiv aq_2 \pmod{D}}} g(q_1) - \frac{Y(J_1,g;D)}{\phi(D)} \right| \tilde{Z}(a,D).$$

By Cauchy–Schwarz and Lemma 13, we have that $\mathcal{E}(N) \leq \mathcal{E}_1^{1/2} \mathcal{E}_2^{1/2}$, where

$$\mathcal{E}_{1} = \sum_{D \leq 2N} Y(J_{2}, f; D) \sum_{a \pmod{D}}^{*} \tilde{Z}(a, D)^{2}$$

$$\ll \sum_{D \leq 2N} Y(J_{2}, f; D) (\log x)^{3} \frac{N^{2}}{D} \tau(D)^{6},$$

and

$$\mathcal{E}_{2} = \sum_{D \leq 2N} \sum_{\substack{q_{2} \in J_{2} \\ (q_{2},D)=1}} f(q_{2}) \sum_{a \pmod{D}}^{*} \left(\sum_{\substack{q_{1} \in J_{1} \\ q_{1} \equiv aq_{2} \pmod{D}}} g(q_{1}) - \frac{Y(J_{1},g;D)}{\phi(D)} \right)^{2}$$
$$= \sum_{D \leq 2N} \sum_{\substack{q_{2} \in J_{2} \\ (q_{2},D)=1}} f(q_{2}) \sum_{b \pmod{D}}^{*} \left(\sum_{\substack{q_{1} \in J_{1} \\ q_{1} \equiv b \pmod{D}}} g(q_{1}) - \frac{Y(J_{1},g;D)}{\phi(D)} \right)^{2}.$$

We have $Y(J_2, f, D) \ll X \ll x^{1/2}$ and

$$\sum_{D \le 2N} D^{-1} \tau(D)^6 \ll (\log x)^{64}.$$

whence $\mathcal{E}_1 \ll N^2 x^{1/2} (\log x)^{67}$. Moreover the bound $Y(J_2, f, D) \ll X \ll x^{1/2}$ shows that

$$\mathcal{E}_{2} \ll x^{1/2} \sum_{D \leq 2N} \sum_{b \pmod{D}}^{*} \left(\sum_{\substack{q_{1} \in J_{1} \\ q_{1} \equiv b \pmod{D}}} g(q_{1}) - \frac{Y(J_{1}, g; D)}{\phi(D)} \right)^{2},$$

whence it suffices to show that

$$\sum_{D \le 2N} \sum_{b \pmod{D}}^{*} \left(\sum_{\substack{q_1 \in J_1 \\ q_1 \equiv b \pmod{D}}} g(q_1) - \frac{Y(J_1, g; D)}{\phi(D)} \right)^2 \ll_C \frac{x}{\log^C x};$$

for any C > 0. This follows by the Barban–Davenport–Halberstam theorem (see, e.g. Theorem 9.14 in [3]), and noting that $\sqrt{x} \ge X \ge \sqrt{x}/\log^4 x$ while $N \le x^{1/2-\delta}$.

It is necessary to use Corollary 2 in the bound for \mathcal{E}_f . Here, Cauchy–Schwarz on \mathcal{E}_f produces $\mathcal{E}_f \leq \mathcal{E}_1^{1/2} \mathcal{E}_2^{1/2}$, where now

$$\mathcal{E}_1 = \sum_{D \le N} \sum_{b}^* \tilde{Z}(b^2, D)^2$$
; and $\mathcal{E}_2 = \sum_{D \le 2N} \sum_{b \pmod{D}}^* (Y_f(b, D) - Y_{f,f}(D))^2$.

For \mathcal{E}_1 we note that any residue class *a* coprime to *D* arises $O(\tau(D))$ times as a square, whence

$$\mathcal{E}_1 \ll \sum_{D \le N} \tau(D) \sum_a^* \tilde{Z}(a, D)^2$$
$$\ll \sum_{D \le N} \tau(D)^7 \frac{N^2}{D} \log^3 x$$
$$\ll N^2 \log^{131} x,$$

by Lemma 13. Thus, it remains to show that $\mathcal{E}_2 \ll x^2 (\log x)^{-C}$ for any constant C. But \mathcal{E}_2 is simply

$$\sum_{D \le 2N} \sum_{b \pmod{D}}^{*} \left| \sum_{\substack{m,n \\ m \equiv bn \pmod{D} \\ (mn,D)=1}} c_1(m)c_2(n) - \frac{1}{\phi(D)} Y(J_1, f; D) Y(J_2, f; D) \right|^2$$

where

$$c_1(m) = \begin{cases} 2p \log p & \text{if } m = p \text{ is prime and } p^2 \in J_1, \\ 0 & \text{otherwise,} \end{cases}$$

and similarly

$$c_2(n) = \begin{cases} 2p \log p & \text{if } n = p \text{ is prime and } p^2 \in J_2, \\ 0 & \text{otherwise.} \end{cases}$$

Since c_1 satisfies the Siegel–Walfisz condition, we now apply Corollary 2 to complete the proof. Note that

$$||c_i \tau||^2 \ll x^{3/4} \log^2 x,$$

for i = 1 or 2, and that $c_i(n)$ is supported on $n \leq x^{1/4}$ (so that the value of x appearing in the statement of Corollary 2 is $x^{1/4}$ in this application).

8. Proof of Proposition 6

We begin by observing that Lemma 12 yields

$$T'(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2) \ll \sum_{D \le 2N} \sum_{a \pmod{D}}^* Y(a, D; f_1, f_2) \tilde{Z}(a, D) + N^2 \sqrt{x} (\log x)^3$$

for some pair of functions $f_1, f_2 = f$ or g. We consider the case in which $f_1 = f_2 = f$, the others being similar, or easier. Since

$$\sum_{D \le 2N} \sum_{a \pmod{D}}^* Y(a, D; f, f) \tilde{Z}(a, D) = \sum_{D \le 2N} \sum_{b \pmod{D}}^* Y_f(b, D) \tilde{Z}(b^2, D)$$

it follows from Proposition 8 that

(24)
$$T'(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2) \ll \sum_{D \le 2N} Y_{f,f}(D) \sum_{b \pmod{D}} \tilde{Z}(b^2, D) + \frac{xN}{\log^C x}.$$

This holds for any C > 0, and since there are $O((\log x)^{6L})$ possible regions $\mathcal{U}_1 \times \mathcal{U}_2 \times J_1 \times J_2$, we see that the final term contributes $O(xN(\log x)^{6L-C})$ in Proposition 6. This is satisfactory on taking $C \ge A + 6L$.

To handle the main terms we note

(25)
$$Y(J_i, f) = |J_i| + O(\sqrt{x} \exp(-\sqrt{\log x}))$$

by the Prime Number Theorem. It follows that $Y(J_i, f) \ll |J_i| \ll \omega X$, and hence that $Y_{f,f}(D) \ll \omega^2 X^2 / \phi(D)$. The main terms in (24) are thus

$$\ll \omega^2 X^2 \sum_{D \le 2N} \frac{\tau(D)}{\phi(D)} \sum_{a \pmod{D}} \tilde{Z}(a, D),$$

since any residue class a coprime to D arises $O(\tau(D))$ times as a square.

Hence, to establish Proposition 6 it will suffice to show that

(26)
$$\mathcal{\mathcal{E}} := \omega^2 X^2 \sum_{\substack{\mathcal{U}_1, \mathcal{U}_2, J_1, J_2 \\ \mathfrak{C}_2(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2) \text{ or } \mathfrak{C}_3(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)} \sum_{D \le 2N} \frac{\tau(D)}{\phi(D)} \sum_{a \pmod{D}} \tilde{Z}(a, D)$$
$$\ll \frac{xN}{\log^A x}$$

when $L \ge A + 6$. For this we use the following two lemmas.

Lemma 14. For fixed $\mathcal{U}_1, \mathcal{U}_2$, D and J_1 , the number of choices for J_2 subject to the condition $\mathfrak{C}_2(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ is $\ll 1$.

Proof. Let $X_i = \inf J_i$, and fix $Z_i \in \mathcal{U}_i$. Then $\mathfrak{C}_2(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ implies that there exists $q_i \in J_i$ and $z_i \in \mathcal{U}_i$ such that

$$D\sqrt{\frac{x}{N}} \ge |q_1 z_2 - q_2 z_1| = |X_1 Z_2 - X_2 Z_1| + O(\omega X \sqrt{N}),$$

and that there exists $q'_i \in J_i$ and $z'_i \in \mathcal{U}_i$ such that

$$D\sqrt{\frac{x}{N}} < |q_1'z_2' - q_2'z_1'| = |X_1Z_2 - X_2Z_1| + O(\omega X\sqrt{N}).$$

Then

$$|X_1Z_2 - X_2Z_1| = D\sqrt{\frac{x}{N}} + O(\omega X\sqrt{N}),$$

and since $|Z_1| \gg \sqrt{N}$ we have

$$|X_2 - \frac{X_1 Z_2}{Z_1}| = \frac{D}{|Z_1|} \sqrt{\frac{x}{N}} + O(\omega X).$$

Since the different values for X_2 increase in steps of length $\simeq \omega X$ it follows that $\mathcal{U}_1, \mathcal{U}_2, D$ and J_1 determine O(1) choices for J_2 .

Lemma 15. For fixed $\mathcal{U}_1, \mathcal{U}_2$, D and J_1 , the number of choices for J_2 subject to the condition $\mathfrak{C}_3(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)$ is $\ll \omega^{-1}(\log x)^{-A-2}$.

Proof. Let $X_i = \inf J_i$, and fix $Z_i \in \mathcal{U}_i$. For all $(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2$, we have that $\Delta(z_1, z_2) = \Delta(Z_1, Z_2) + O(\omega N)$. Then

$$|X_1Z_2 - X_2Z_1| \ll (\mathfrak{D}_0 + \omega N)\sqrt{\frac{x}{N}},$$

and since $|Z_1| \gg \sqrt{N}$ and $\omega N \ll \mathfrak{D}_0$ for $L \ge A + 6$ this yields

$$X_2 = X_1 \frac{Z_2}{Z_1} + O\left(\frac{\mathfrak{D}_0}{N}\sqrt{x}\right).$$

This tells us that X_2 is restricted to be in an interval of length $\ll \sqrt{x}(\log x)^{-A-6}$. Since J_2 is of length $\omega_1 X \gg \omega \sqrt{x}(\log x)^{-4}$, this gives $O(\omega^{-1}(\log x)^{-A-2})$ choices for J_2 . \Box

By Lemma 14, Lemma 15 and (26),

$$\mathcal{E} \ll \left(\omega + (\log x)^{-A-2}\right) X^2 \sum_{\mathcal{U}_1, \mathcal{U}_2} \sum_{D \le 2N} \frac{\tau(D)}{\phi(D)} \sum_{a \pmod{D}} \tilde{Z}(a, D)$$
$$\ll \left(\omega + (\log x)^{-A-2}\right) X^2 N \sum_{D \le 2N} \frac{\tau(D)}{\phi(D)},$$

where we have used Lemma 13 for the last line. Hence if $L \ge A + 6$ we obtain

$$\mathcal{E} \ll \left(\omega + (\log x)^{-A-2}\right) X^2 N (\log x)^2 \ll \frac{xN}{\log^{L-2} x} + \frac{xN}{\log^A x} \ll \frac{xN}{\log^A x},$$

as required.

9. PROOF OF PROPOSITION 7; FURTHER MANOEUVRES

Supposing that one of the functions h_1 and h_2 is g, we have according to Proposition 8 that

$$\sum_{D \le 2N} \sum_{a \pmod{D}}^{*} Y(a, D; h_1, h_2) Z(a, D) = \sum_{D \le 2N} \sum_{a \pmod{D}}^{*} Y_{h_1, h_2}(D) Z(a, D) + O_C\left(\frac{xN}{\log^C x}\right),$$

for any C > 0. In the remaining case $h_1 = h_2 = f$, we have

$$\sum_{D \le 2N} \sum_{a \pmod{D}}^{*} Y(a, D; f, f) Z(a, D) = \sum_{D \le 2N} \sum_{b \pmod{D}}^{*} Y_{f, f}(D) Z(b^{2}, D) + O_{C}\left(\frac{xN}{\log^{C} x}\right).$$

From (25) we see that we may replace $Y_{h_1,h_2}(D)$ by $|J_1||J_2|/\phi(D)$ in each case, with a total error

$$\ll x \exp(-\sqrt{\log x}) \sum_{\mathcal{U}_{1},\mathcal{U}_{2}} \sum_{D \leq 2N} \frac{1}{\phi(D)} \left(\sum_{a \pmod{D}}^{*} |Z(a,D)| + \sum_{b \pmod{D}}^{*} |Z(b^{2},D)| \right)$$
$$\ll x \exp(-\sqrt{\log x}) \sum_{D \leq 2N} \frac{\tau(D)}{\phi(D)} \sum_{\mathcal{U}_{1},\mathcal{U}_{2}} \sum_{a \pmod{D}}^{*} \tilde{Z}(a,D)$$
$$\ll x \exp(-\sqrt{\log x}) \sum_{D \leq 2N} \frac{\tau(D)}{\phi(D)} N$$
$$\ll x \exp(-\sqrt{\log x}) N \log x,$$

by Lemma 13. This is satisfactory for the proposition.

It therefore remains to show that

$$|J_1||J_2| \sum_{\substack{\mathcal{U}_1,\mathcal{U}_2\\\mathfrak{C}_1(\mathcal{U}_1,\mathcal{U}_2,J_1,J_2)}} \sum_{D \le 2N} \frac{1}{\phi(D)} \left(\sum_{b \pmod{D}}^* Z(b^2, D) - \sum_{a \pmod{D}}^* Z(a, D) \right) \ll \frac{xN}{\log^{A+2L} x}$$

or indeed that

$$\mathcal{E} := \sum_{\substack{\mathcal{U}_1, \mathcal{U}_2\\\mathfrak{C}_1(\mathcal{U}_1, \mathcal{U}_2, J_1, J_2)}} \sum_D \frac{1}{\phi(D)} \left(\sum_{b \pmod{D}}^* Z(b^2, D) - \sum_{a \pmod{D}}^* Z(a, D) \right) \ll \frac{N}{\log^A x}.$$

、

Here we have dropped the condition $D \leq 2N$, which follows automatically since β_z is supported on $N(z) \leq 2N$.

It turns out that it is easier to estimate the corresponding sum in which the factor $1/\phi(D)$ is replaced by $D/\phi(D)$. To make this transition we will use summation by parts on D to see that

$$\mathcal{E} \ll \sum_{\mathcal{U}_1, \mathcal{U}_2} \frac{|\mathcal{E}_0(t_0)|}{\mathfrak{D}_0},$$

where $t_0 = t_0(\mathcal{U}_1, \mathcal{U}_2) > \mathfrak{D}_0$ is such that $|\mathcal{E}_0(t)|$ is maximal, with

,

$$\mathcal{E}_0(t) = \mathcal{E}_0(t, \mathcal{U}_1, \mathcal{U}_2) := \sum_{D < t} \frac{D}{\phi(D)} \left(\sum_{\substack{b \pmod{D} \\ 26}}^* Z(b^2, D) - \sum_{\substack{a \pmod{D} \\ a \pmod{D}}}^* Z(a, D) \right)$$

We now wish to remove the condition D < t by arguing that the contribution of those $\mathcal{U}_1, \mathcal{U}_2$ such that there exists $z_1, z'_1 \in \mathcal{U}_1, z_2, z'_2 \in \mathcal{U}_2$ with $\Delta(z_1, z_2) < t$ and $\Delta(z'_1, z'_2) \geq t$ is negligible. Indeed, for such $\mathcal{U}_1, \mathcal{U}_2$, we must have

$$\Delta(z_1, z_2) = t + O(N\omega) = t \left(1 + O((\log x)^{A+6-L}) \right)$$

for all $z_1 \in \mathcal{U}_1, z_2 \in \mathcal{U}_2$. If $\mathcal{U}_1 = \mathcal{U}(c_1, \theta_1)$ and $\mathcal{U}_2 = \mathcal{U}(c_2, \theta_2)$, then

$$\sin(\theta_1 - \theta_2) = \frac{t}{c_1 c_2 N} \left(1 + O((\log x)^{A+6-L}) \right).$$

In general that if one restricts $\sin(\phi)$ to an interval of length μ , then ϕ will be confined to a set of measure $O(\sqrt{\mu})$ modulo $\pi/2$ and hence modulo 2π as well. Indeed, suppose $\phi_1, \phi_2 \in [0, \pi/2 - \sqrt{\mu}]$, and that $|\sin \theta_1 - \sin \theta_2| \ll \mu$. We want to show that $|\theta_1 - \theta_2| \ll \sqrt{\mu}$. This follows from the elementary formula

$$\sin \phi_1 - \sin \phi_2 = 2 \cos \left(\frac{\phi_1 + \phi_2}{2}\right) \sin \left(\frac{\phi_1 - \phi_2}{2}\right),$$

provided that $\cos\left(\frac{\phi_1+\phi_2}{2}\right) \gg \sqrt{\mu}$. The latter bound follows from $\pi/2 - (\phi_1+\phi_2)/2 \gg \sqrt{\mu}$.

Hence if we fix c_1 , θ_1 and c_2 then the total number of choices for θ_2 is $\ll (\log x)^{(A+6+L)/2}$. This gives $\ll (\log x)^{(A+6+7L)/2}$ total possibilities for $(\mathcal{U}_1, \mathcal{U}_2)$. A given residue class a occurs $O(\tau(D))$ times as b^2 , and $D/\phi(D) \ll \log x$. Thus each pair $(\mathcal{U}_1, \mathcal{U}_2)$ contributes $\ll \omega^4 N^2 (\log x)^{17}$ to the sum, by Lemma 13, giving a total contribution of

$$\ll \frac{N^2 (\log x)^{(A+40-L)/2}}{\mathfrak{D}_0} \ll \frac{N}{(\log x)^{(L-3A-52)/2}}$$

to \mathcal{E} , which suffices when L = 6A + 52.

It now suffices to show that

(27)
$$\mathcal{E}_1 = \mathcal{E}_1(\mathcal{U}_1, \mathcal{U}_2) = \sum_D \frac{D}{\phi(D)} \left(\sum_{b \pmod{D}}^* Z(b^2, D) - \sum_{a \pmod{D}}^* Z(a, D) \right) \ll \frac{N^2}{\log^{C_1} x},$$

for any $C_1 > 0$ and for fixed $\mathcal{U}_1, \mathcal{U}_2$. Since we have $\Delta > 0$ whenever $(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2$ we see that if $D \mid \Delta$ then

$$\sum_{k|\Delta/D} \mu(k) = \begin{cases} 1, & D = \Delta, \\ 0, & D \neq \Delta. \end{cases}$$

We therefore have

$$\mathcal{E}_1 = \sum_{D=1}^{\infty} \sum_{k=1}^{\infty} \frac{D\mu(k)}{\phi(D)} \left(\sum_{b \pmod{D}}^* W(b^2, k, D) - \sum_{a \pmod{D}}^* W(a, k, D) \right)$$

where

$$W(a,k,D) := \sum_{\substack{(z_1,z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ kD \mid \Delta \\ az_2 \equiv z_1 \pmod{D}}}^{\flat} \beta_{z_1} \beta_{z_2}.$$

When $kD \mid \Delta$, there is a unique integer $c = c(z_1, z_2; kD)$ modulo kD such that $cz_2 \equiv z_1 \pmod{kD}$, and conversely this congruence implies that $kD \mid \Delta$. For this integer c we have (c, kD) = 1 and

$$\#\{b \pmod{D} : b^2 z_2 \equiv z_1 \pmod{D}\} = \#\{b \pmod{D} : b^2 \equiv c \pmod{D}\} = \sum_{\substack{\chi \pmod{D} \\ \chi^2 = \chi_0}} \chi(c).$$

It now follows that

$$\sum_{\substack{b \pmod{D} \\ b \pmod{D}}}^{*} W(b^{2}, k, D) - \sum_{\substack{a \pmod{D} \\ a \pmod{D}}}^{*} W(a, k, D) = \sum_{\substack{\chi \pmod{D} \\ \chi^{2} = \chi_{0} \\ \chi \neq \chi_{0}}} \sum_{\substack{c \pmod{kD} \\ c = z_{1} \pmod{kD}}}^{*} \sum_{\substack{(z_{1}, z_{2}) \in \mathcal{U}_{1} \times \mathcal{U}_{2} \\ (z_{2} \equiv z_{1} \pmod{kD})}}^{\flat} \beta_{z_{1}} \beta_{z_{2}} \chi(c),$$

and hence that

$$\mathcal{E}_1 = \sum_{D=1}^{\infty} \sum_{k=1}^{\infty} \frac{D\mu(k)}{\phi(D)} \sum_{\substack{\chi \pmod{D} \\ \chi^2 = \chi_0 \\ \chi \neq \chi_0}} \sum_{\substack{c \pmod{kD} \\ cz_2 \equiv z_1 \pmod{kD}}}^* \sum_{\substack{(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ cz_2 \equiv z_1 \pmod{kD}}}^{\flat} \beta_{z_1} \beta_{z_2} \chi(c),$$

Let $d = d(\chi)$ be the conductor of χ and write D = de and ek = f, giving

$$\mathcal{E}_1 = \sum_{d>1} \sum_f C(d, f) \sum_{\substack{\chi \pmod{d} \\ \chi^2 = \chi_0}}^* \sum_{\substack{c \pmod{df} \\ cz_2 \equiv z_1 \pmod{df}}}^* \sum_{\substack{(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ cz_2 \equiv z_1 \pmod{df}}}^{\rho} \beta_{z_1} \beta_{z_2} \chi(c),$$

where

$$C(d,f) = \sum_{ek=f} \frac{de\mu(k)}{\phi(de)} = \frac{d}{\phi(d)} \sum_{ek=f} \frac{\phi(d)e\mu(k)}{\phi(de)}.$$

Note that the sum for $\chi \pmod{d}$ is empty unless $d = d_1$ or $4d_1$ or $8d_1$, with d_1 odd and square-free, in which cases there are at most two possible characters χ . When d is given, the function $\kappa(e) := \phi(d)e/\phi(de)$ is multiplicative in e. Moreover if $v \ge 1$ then

$$(\kappa * \mu)(p^{v}) = \begin{cases} (p-1)^{-1}, & \text{if } v = 1 \text{ and } p \nmid d, \\ 0, & \text{otherwise.} \end{cases}$$

We then see that

$$C(d,f) = \frac{d\mu^2(f)}{\phi(df)}$$

if (d, f) = 1 and C(d, f) = 0 otherwise. This leads to the expression

(28)
$$\mathcal{E}_1 = \sum_{\substack{f,d \\ (d,f)=1}} \frac{d\mu^2(f)}{\phi(df)} \sum_{\substack{\chi \pmod{d} \\ \chi^2 = \chi_0 \\ \chi \neq \chi_0}}^* \left(\sum_{\substack{c \pmod{df} \ (z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ cz_2 \equiv z_1 \pmod{df}}} \beta_{z_1} \beta_{z_2} \chi(c) \right).$$

We proceed to show that large values of f make a negligible contribution. Since $df \mid \Delta(z_1, z_2)$ we will have $df \leq 2N$. On recalling that $0 \leq \beta_z \leq 1$ we find that

Τ

$$\begin{split} \sum_{f>F} \sum_{\substack{d \\ (d,f)=1}} \frac{d\mu^2(f)}{\phi(df)} \sum_{\substack{\chi \pmod{d} \\ \chi^2 = \chi_0}}^* \left| \sum_{\substack{c \pmod{d} \\ (\log x)}}^* \sum_{\substack{(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ cz_2 \equiv z_1 \pmod{d}f}}^{\flat} \beta_{z_1} \beta_{z_2} \chi(c) \right| \\ \ll (\log x) \sum_{f>F} f^{-1} \sum_{\substack{d \le 2N/f}} \sum_{\substack{df \mid D \\ D \le 2N}} \sum_{a \pmod{D}}}^* \tilde{Z}(a, D) \\ \ll (\log x) \sum_{f>F} f^{-1} \sum_{\substack{d \le 2N/f}} \sum_{\substack{df \mid D \\ D \le 2N}} N \\ \ll \frac{N^2 (\log x)^2}{F}, \end{split}$$

by Lemma 13. An inspection of (27) and (28) reveals that this is satisfactory if we take

$$F = (\log x)^{C_1 + 2}.$$

Next, we split our sum into three different ranges for d, namely $d \leq D_1$, $D_1 < d \leq D_2$, $d > D_2$ where

$$D_1 = F^{10} (\log x)^{2C_1 + 14}$$
 and $D_2 = \frac{N}{F^{15} (\log x)^{3C_1 + 21}}$.

We deal with the middle range for d in Section 10, and the remaining ranges in Section 11.

10. Proof of Proposition 7; Middle d

We proceed to dispose of the middle range of values for d, making use of a major intermediate result from the work of Friedlander and Iwaniec [4]. Thus we will consider the case in which $D < d \leq 2D$ say. (We should make it clear to the reader that there is no longer any connection between D and Δ .) We will set

$$\mathcal{E}_{1}(D) := \sum_{f \leq F} f^{-1} \mu^{2}(f) \sum_{\substack{D < d \leq 2D \\ (d,f)=1}} \sum_{\substack{\chi \pmod{d} \\ \chi^{2} = \chi_{0}}}^{*} \left| \sum_{\substack{c \pmod{df} \\ cz_{2} \equiv z_{1} \pmod{df}}}^{*} \sum_{\substack{\beta_{z_{1}} \beta_{z_{2}} \chi(c) \\ cz_{2} \equiv z_{1} \pmod{df}}}^{\flat} \beta_{z_{1}} \beta_{z_{2}} \chi(c) \right|.$$

Let $d = d_1 d_2$ where d_1 is odd and square-free, and d_2 is 1, 4, or 8, and write $\chi = \chi_1 \chi_2$ accordingly, so that $\chi_1(n)$ is the Jacobi symbol $(\frac{n}{d_1})$. If we set $g = f d_2$ then g and d_1 will be coprime. We now split the variables z_j into congruence classes $z_j \equiv w_j \pmod{g}$ and find that

$$\mathcal{E}_1(D) \ll \sum_{f \leq F} \sum_{d_2=1,4,8} \sum_{w_1,w_2 \pmod{g}} \mathcal{E}_2(D;g,w_1,w_2),$$

where

$$\mathcal{E}_{2}(D; g, w_{1}, w_{2}) := \sum_{\substack{D/8 < d_{1} \leq 2D \\ (d_{1}, g) = 1}} \left| \sum_{\substack{b \pmod{d_{1}} (z_{1}, z_{2}) \in \mathcal{U}_{1} \times \mathcal{U}_{2} \\ z_{j} \equiv w_{j} \pmod{g} \\ bz_{2} \equiv z_{1} \pmod{d_{1}}}^{\flat} \beta_{z_{1}} \beta_{z_{2}} \left(\frac{b}{d_{1}} \right) \right|$$

We now write $\gamma_{z_j} = \beta_{z_j}$ if $z_j \equiv w_j \pmod{g}$, and $\gamma_{z_j} = 0$ otherwise, so that $0 \leq \gamma_1, \gamma_2 \leq 1$. It follows that there is some choice of f, d_2 and w_1, w_2 such that

$$\mathcal{E}_1(D) \ll F^5 \mathcal{E}_3(D)$$

ī

with

$$\mathcal{E}_{3}(D) = \sum_{D/8 < d \le 2D} \mu^{2}(2d) \left| \sum_{\substack{b \pmod{d} \\ bz_{2} \equiv z_{1} \pmod{d}}}^{*} \sum_{\substack{(z_{1}, z_{2}) \in \mathcal{U}_{1} \times \mathcal{U}_{2} \\ bz_{2} \equiv z_{1} \pmod{d}}}^{\flat} \gamma_{z_{1}} \gamma_{z_{2}} \left(\frac{b}{d}\right) \right|.$$

(Here we have replaced the dummy variable d_1 by d, for notational convenience.)

1

Writing $z_j = x_j + iy_j$ for j = 1, 2 we classify the numbers z_j according to the value of $(x_1, d) = r$, say. Since $bz_2 \equiv z_1 \pmod{d}$ with (b, d) = 1 we will have $(x_2, d) = r$ also. We now write d = rs and $x_j = ru_j$ for j = 1, 2, so that $(u_j, s) = 1$. Since z_j is primitive and $r \mid x_j$ it follows that $(y_j, r) = 1$. Then $d \mid x_1y_2 - x_2y_1$ if and only if there is a b such that $bz_2 \equiv z_1 \pmod{d}$. Moreover $d \mid x_1y_2 - x_2y_1$ if and only if $u_1y_2 - u_2y_1$. Thus there is an

integer c such that $y_1 u_1^{-1} \equiv y_2 u_2^{-1} \equiv c \pmod{s}$. We will also have $bu_2 \equiv u_1 \pmod{s}$ and $by_2 \equiv y_1 \pmod{r}$, so that

$$\left(\frac{b}{d}\right) = \left(\frac{b}{r}\right)\left(\frac{b}{s}\right) = \left(\frac{y_1y_2}{r}\right)\left(\frac{u_1u_2}{s}\right).$$

These considerations show that

$$\mathcal{E}_1(D) \ll F^3 \sum_{D/8 < rs \le 2D \ c \ (\text{mod } s)} \mu^2(2rs) |\Sigma_1(r, s, c) \Sigma_2(r, s, c)|,$$

with

$$\Sigma_j(r, s, c) := \sum_{\substack{ru_j + iy_j \in \mathcal{U}_j \\ y_j u_j^{-1} \equiv c \pmod{s}}}^{\flat} \gamma_{ru_j + iy_j} \left(\frac{y_j}{r}\right) \left(\frac{u_j}{s}\right).$$

We have dropped the condition $(u_j, s) = 1$ since the Legendre symbol vanishes when this fails to hold.

By Cauchy's inequality we deduce that for either j = 1 or j = 2,

(29)
$$\mathcal{E}_1(D) \ll F^5 \sum_{r \le 2D} \sum_{D/8r < s \le 2D/r} \mu^2(2rs) \sum_{c \pmod{s}} |\Sigma_j(r, s, c)|^2.$$

Suppressing the dependence on r and j we may write

$$\Sigma_j(r, s, c) := \sum_{yu^{-1} \equiv c \pmod{s}} \alpha_{u,y}\left(\frac{u}{s}\right)$$

where

$$\alpha_{u,y} = \gamma_{ru+iy} \left(\frac{y}{r}\right)$$

when $z = ru + iy \in \mathcal{U}_j$ and the conditions for $\sum_{i=1}^{b}$ hold, and $\alpha_{u,y} = 0$ otherwise. The coefficients $\alpha_{u,y}$ are therefore supported in $|u| \leq \sqrt{2N}/r$ and $|y| \leq \sqrt{2N}$. Since β_z is supported on values with z primitive and $\operatorname{Re}(z)$ odd, we may assume that (u, 2y) = 1.

We are now ready to apply Proposition 14.1 of Friedlander and Iwaniec [4]. This shows that if $\alpha_{u,y}$ is supported in U < u < 2U and Y < y < 2Y, then

$$\sum_{S < s \le 2S} \mu^2(2s) \sum_{c \pmod{s}} \left| \sum_{yu^{-1} \equiv c \pmod{s}} \alpha_{u,y} \left(\frac{u}{s}\right) \right|^2 \le N(S, U, Y) \sum_{u,y} \tau(u) |\alpha_{u,y}|^2$$

with

$$N(S, U, Y) \ll_{\epsilon} S + S^{-1/2}UY + S^{1/3}(UY)^{2/3}(\log UY)^4 + (U+Y)^{1/12}(UY)^{11/12+\epsilon}$$

for any fixed $\epsilon > 0$.

In our case we have

$$\sum_{u,y} \tau(u) |\alpha_{u,y}|^2 \ll UY(\log x)$$

Thus, if we sum over dyadic ranges for u, y, and s, we obtain

$$\sum_{D/8r < s \le 2D/r} \mu^2(2rs) \sum_{c \pmod{s}} |\Sigma_j(r, s, c)|^2 \\ \ll \left\{ \frac{D}{r} + \left(\frac{D}{r}\right)^{-1/2} \frac{N}{r} + \left(\frac{D}{r}\right)^{1/3} \left(\frac{N}{r}\right)^{2/3} (\log x)^4 + N^{1/24} \left(\frac{N}{r}\right)^{11/12} N^\epsilon \right\} \frac{N}{r} (\log x) \\ \ll \left\{ D + D^{-1/2}N + D^{1/3}N^{2/3} + N^{23/24+\epsilon} \right\} \frac{N}{r} (\log x)^5.$$

Summing over r, we see that from (29) that

$$\mathcal{E}_1(D) \ll F^5(\log x)^6 \{D + D^{-1/2}N + D^{1/3}N^{2/3} + N^{23/24+\epsilon}\}N.$$

Thus, on summing over dyadic ranges for D, we see that values of d with $D_1 \leq d \leq D_2$ make a satisfactory contribution given our choices of D_1 and D_2 .

11. Proof of Proposition 7 : large d and small d

11.1. Large d. Now, we will show for fixed $f \leq F$, and any C > 0 that

$$\sum_{\substack{d>D_2\\(d,f)=1}} \frac{d}{\phi(d)} \sum_{\substack{\chi \pmod{d}\\\chi^2=\chi_0}}^* \left(\sum_{\substack{c \pmod{df} \ (z_1,z_2)\in\mathcal{U}_1\times\mathcal{U}_2\\cz_2\equiv z_1 \pmod{df}}}^* \beta_{z_1}\beta_{z_2}\chi(c) \right) \ll_C \frac{N^2}{\log^C x}.$$

In the sequel, we shall use the convention that C denotes a large positive constant, not necessarily the same from line to line.

As in the previous section we decompose d as d_1d_2 and χ as $\chi_1\chi_2$. Writing $\Delta = \Delta(z_1, z_2)$ for short, we must have $df \mid \Delta$, and so we may set $\Delta = d_1et$ where e is odd and t is a power of 2. Our conditions on \mathcal{U}_1 and \mathcal{U}_2 ensure that $0 < \Delta \leq 2N$, whence $1 \leq et \leq 16N/D_2 \ll (\log x)^{18C_1+51}$. We split the sums over z_j into congruence classes $z_j \equiv w_j \pmod{8et}$, and fix the parameters

(30)
$$f, d_2, \chi_2, e, w_1, w_2 \text{ and } t.$$

Each admissible pair z_1, z_2 corresponds to a unique integer $k \pmod{\Delta}$ with the property that $kz_2 \equiv z_1 \pmod{\Delta}$, and then

$$\chi(c) = \chi(k) = \chi_2(k) \left(\frac{k}{d_1}\right)$$

where $\chi_2(k)$ is determined by the parameters (30). The number of choices for the parameters (30) is bounded by a fixed power of log x so that it suffices to show that

$$\sum_{\substack{d_1 > D_2/d_2 \\ (d_1, 2f) = 1}} \frac{d_1 \mu^2(d_1)}{\phi(d_1)} \left(\sum_{k \pmod{\Delta}} \sum_{z_1, z_2} \beta_{z_1} \beta_{z_2} \left(\frac{k}{d_1} \right) \right) \ll_C \frac{N^2}{\log^C x}$$

for every C > 0, where the sum over z_1, z_2 satisfies the conditions

 $(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2, \quad kz_2 \equiv z_1 \pmod{\Delta} \quad z_j \equiv w_j \pmod{8et} \quad (j = 1, 2) \text{ and } \Delta = d_1 et.$

We proceed to investigate the Jacobi symbol which occurs here. Our analysis is very close to that given by Friedlander and Iwaniec [4, Lemma 17.1]. However our situation is not quite the same as theirs. In what follows we will make repeated use of the fact that $z_j = x_j + iy_j$ with $x_j > 0$ and $(x_j, 2y_j) = 1$. If we set $r = (x_1, \Delta)$ then, as in the previous section, we have $r = (x_2, \Delta)$, allowing us to write $x_j = ru_j$ for j = 1 and 2. Since the x_j

are odd we see that $r \mid d_1 e$, and setting $s = d_1 e/r$ we find that $ku_2 \equiv u_1 \pmod{s}$ and $ky_2 \equiv y_1 \pmod{r}$. Then (k, e) = 1 since $(k, \Delta) = 1$, and $(y_i, r) = 1$, since z_i is primitive. We therefore find that

$$\left(\frac{k}{d_1}\right) = \left(\frac{k}{e}\right)\left(\frac{k}{r}\right)\left(\frac{k}{s}\right) = \left(\frac{k}{e}\right)\left(\frac{y_1y_2}{r}\right)\left(\frac{u_1u_2}{s}\right).$$

Recalling that $z_i \equiv w_i \pmod{e}$, we see that $\binom{k}{e}$ is determined by w_1 and w_2 . By quadratic reciprocity we have

$$\left(\frac{u_1u_2}{s}\right) = \left(\frac{s}{u_1u_2}\right) < s, u_1u_2 >,$$

where

$$< a, b >:= (-1)^{(a-1)(b-1)/4}$$

for odd integers a, b. However $\Delta = x_1y_2 - x_2y_1$, whence $st = u_1y_2 - u_2y_1$. It follows that

$$\left(\frac{s}{u_1}\right) = \left(\frac{-tu_2y_1}{u_1}\right)$$

and

$$\left(\frac{s}{u_2}\right) = \left(\frac{tu_1y_2}{u_2}\right),\,$$

so that

$$\begin{pmatrix} \frac{s}{u_1 u_2} \end{pmatrix} = \begin{pmatrix} \frac{t}{u_1 u_2} \end{pmatrix} \begin{pmatrix} \frac{-1}{u_1} \end{pmatrix} \begin{pmatrix} \frac{y_1}{u_1} \end{pmatrix} \begin{pmatrix} \frac{y_2}{u_2} \end{pmatrix} < u_1, u_2 >$$

$$= \begin{pmatrix} \frac{t}{x_1 x_2} \end{pmatrix} \begin{pmatrix} \frac{-1}{x_1} \end{pmatrix} \begin{pmatrix} \frac{-1}{r} \end{pmatrix} \begin{pmatrix} \frac{y_1}{u_1} \end{pmatrix} \begin{pmatrix} \frac{y_2}{u_2} \end{pmatrix} < u_1, u_2 >$$

The first two factors on the right are determined by the parameters (30), since $z_j \equiv$ $w_i \pmod{8}$, whence

$$\left(\frac{k}{d_1}\right) = \eta\left(\frac{y_1y_2}{r}\right) < s, u_1u_2 > \left(\frac{-1}{r}\right)\left(\frac{y_1}{u_1}\right)\left(\frac{y_2}{u_2}\right) < u_1, u_2 >,$$

for some value of $\eta = \pm 1$ determined by the parameters (30). We then deduce that

$$\left(\frac{k}{d_1}\right) = \eta\left(\frac{y_1}{x_1}\right)\left(\frac{y_2}{x_2}\right) < r, r > < s, u_1u_2 > < u_1, u_2 > < u_2 > < u_1, u_2 > < u_2 > < u_1, u_2 > < u$$

We can now use the relations $\langle ab, c \rangle = \langle a, c \rangle \langle b, c \rangle$ and $\langle a, bc \rangle = \langle a, b \rangle \langle a, c \rangle$ to conclude that

$$\begin{pmatrix} \frac{k}{d_1} \end{pmatrix} = \eta \begin{pmatrix} \frac{y_1}{x_1} \end{pmatrix} \begin{pmatrix} \frac{y_2}{x_2} \end{pmatrix} < sr, u_1 u_2 > < x_1, x_2 >$$
$$= \eta \begin{pmatrix} \frac{y_1}{x_1} \end{pmatrix} \begin{pmatrix} \frac{y_2}{x_2} \end{pmatrix} < d_1 e, x_1 x_2 > < x_1, x_2 >$$

The parameters (30) determine $x_1, x_2 \pmod{8et}$, and it follows that the factor $\langle x_1, x_2 \rangle$ is uniquely determined by the parameters (30). Moreover,

$$d_1et = \Delta = \operatorname{Im}(\bar{z}_1z_2) \equiv \operatorname{Im}(\bar{w}_1w_2) \pmod{8et}.$$

Writing Im $(\bar{w}_1 w_2) = etn$, we see that $d_1 \equiv n \pmod{8}$ so that $d_1 \pmod{8}$ is determined by the parameters (30), and so $\langle d_1e, x_1x_2 \rangle$ is also.

We therefore see that it will be enough to show that

$$\sum_{z_1, z_2} \stackrel{\flat}{=} \frac{d_1 \mu^2(d_1)}{\phi(d_1)} \beta_{z_1}\left(\frac{y_1}{x_1}\right) \beta_{z_2}\left(\frac{y_2}{x_2}\right) \ll_C \frac{N^2}{\log^C x}$$
32

for each given set of parameters (30), where the sum over z_1, z_2 is restricted by the conditions

$$(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2, \ z_j \equiv w_j \pmod{8et} \ (j = 1, 2), \ et \mid \Delta, \ d_1 > D_2/d_2, \ and \ (d_1, 2f) = 1.$$

Moreover d_1 is then defined by $d_1 := \Delta/et$. We may remove the conditions for $\sum_{i=1}^{\flat}$, since β_z is supported on primitive $z \equiv 1 \pmod{2}$ with positive real part. Moreover the condition $et \mid \Delta$ depends only on the choice of w_1 and w_2 . Following the notation of Friedlander and Iwaniec [4, Section 17] we write

$$\beta'_z = \beta_z[z] = \beta_z i^{(x-1)/2} \left(\frac{y}{x}\right)$$

for z = x + iy. Since the factors corresponding to $i^{(x-1)/2}$ are determined by w_1 and w_2 we then see that it suffices to show that

$$\mathcal{E}_4 := \sum_{z_1, z_2} \frac{d_1 \mu^2(d_1)}{\phi(d_1)} \beta'_{z_1} \beta'_{z_2} \ll_C \frac{N^2}{\log^C x},$$

where the sum is subject to

$$(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2, \ z_j \equiv w_j \pmod{8et} \ (j = 1, 2), \ d_1 > D_2/d_2, \ \text{and} \ (d_1, 2f) = 1,$$

with $d_1 := \Delta/et$ as before.

The multiplicative function

$$\kappa(n) := \begin{cases} \frac{n\mu^2(n)}{\phi(n)}, & (n, 2f) = 1, \\ 0, & (n, 2f) > 1, \end{cases}$$

may be written as $\kappa = \kappa_0 * 1$, where $\kappa_0(n) = \mu(n)$ for $n \mid 2f$, and for $p \nmid 2f$,

$$\kappa_0(p) = \frac{1}{p-1}, \quad \kappa_0(p^2) = \frac{-p}{p-1}, \text{ and } \kappa_0(p^r) = 0, \ (r \ge 3).$$

It follows that we may write

$$\mathcal{E}_{4} = \sum_{n=1}^{\infty} \kappa_{0}(n) \sum_{z_{1}, z_{2}} \beta_{z_{1}}' \beta_{z_{2}}'$$

where the inner sum is subject to

(31) $(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2, \ z_j \equiv w_j \pmod{8et} \ (j = 1, 2), \ \Delta > etD_2/d_2, \ \text{and} \ etn \mid \Delta.$

Since

$$\sum_{\substack{(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ n \mid \Delta}}^{\flat} \leq \sum_{\substack{D \leq 2N \\ n \mid D}} \sum_{\substack{a \pmod{D}}}^* \tilde{Z}(a, D) \ll N^2/n$$

by (21), the contribution to \mathcal{E}_4 from integers $n \ge (\log x)^B$ say is

$$\ll \sum_{n \ge (\log x)^B} \frac{N^2}{n} |\kappa_0(n)|$$

$$\ll \sum_{n_1|2f} \sum_{n_2 n_3^2 \ge (\log x)^B/n_1} \frac{N^2 \log(n_2 n_3)}{n_2^2 n_3^2}$$

$$\ll \sum_{n_1|2f} \sum_{m^2 \ge (\log x)^B/n_1} \frac{N^2 \tau(m) \log m}{m^2}$$

$$\ll \sum_{n_1|2f} N^2 (\log x)^{1-B/2} n_1^{1/2}$$

$$\ll F N^2 (\log x)^{1-B/2}.$$

In order to achieve a bound $O(N^2(\log x)^{-C})$ for \mathcal{E}_4 we choose B so that $F(\log x)^{1-B/2} = (\log x)^{-C}$. Thus it will now be enough to show that

$$\sum_{z_1, z_2} \beta'_{z_1} \beta'_{z_2} \ll_C \frac{N^2}{\log^{C+B} x}$$

for every fixed C > 0, for each individual value of $n \leq (\log x)^B$. The sum is again subject to (31), and we can allow for the constraint $etn \mid \Delta$ by subdividing the sum according to the values of z_j modulo 8etn. Thus there are Gaussian integers w_1, w_2 for which it will suffice to show that

$$\sum_{\substack{(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ z_j \equiv w_j \pmod{8etn} \\ \Delta > etD_2/d_2}} \beta'_{z_1} \beta'_{z_2} \ll_C \frac{N^2}{\log^C x}$$

for every fixed C > 0, and for each choice of $e, t, n \leq (\log x)^C$ and of w_1 and w_2 .

Our next task is to remove the condition $\Delta > etD_2/d_2$. This may be done by further subdividing the regions \mathcal{U}_i and following the methods of Section 8, used to prove Proposition 6. We do not repeat the details.

We now have to handle

 $z \equiv$

$$\sum_{\substack{(z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ z_j \equiv w_j \pmod{\det n}}} \beta'_{z_1} \beta'_{z_2} = \left\{ \sum_{\substack{z \in \mathcal{U}_1 \\ z \equiv w_1 \pmod{\det n}}} \beta'_z \right\} \left\{ \sum_{\substack{z \in \mathcal{U}_2 \\ z \equiv w_2 \pmod{\det n}}} \beta'_z \right\}$$

we note that 8etn is at most a power of $\log x$, while β'_z is supported on values for which every prime factor of N(z) is at least x^{δ} . Thus z is automatically coprime to 8etn, and so we may assume that w_1 and w_2 are coprime to 8etn. This allows us to pick out the congruence conditions $z \equiv w \pmod{8etn}$ using multiplicative characters $\chi \pmod{8etn}$ over $\mathbb{Z}[i]$. We therefore deduce that

$$\sum_{\substack{z \in \mathcal{U}_j \\ \equiv w \pmod{8etn}}} \beta'_z = \frac{1}{\phi_{\mathbb{Q}[i]}(8etn)} \sum_{\chi \pmod{8etn}} \overline{\chi}(w) S(\chi, \mathcal{U}_j),$$

where $\phi_{\mathbb{Q}(i)}(q)$ is the Euler ϕ -function for the Gaussian integers, and

$$S(\chi, \mathcal{U}) = \sum_{\substack{z \in \mathcal{U} \\ 34}} \beta'_z \chi(z).$$

We have then reduced our problem to one of showing that for all C > 0,

$$S(\chi, \mathcal{U}) \ll_C N(\log x)^{-C},$$

for all $\chi \pmod{8etn}$ and $\mathcal{U} = \mathcal{U}_1$ or $\mathcal{U} = \mathcal{U}_2$. From now on it will be convenient to set f = 8etn, so that χ is a character to modulus f. The reader should note that we are recycling some of our previous notation — the symbol f no longer has its former meaning!

We recall here that \mathcal{U} is of the form

(32)
$$\mathcal{U} = \{z : \sqrt{N'} < |z| \le (1+\omega_1)\sqrt{N'}, \theta_0 < \arg z < \theta_0 + \omega_2\},$$

where $N' \simeq N$. We pick out the condition $\arg z \in (\theta_0, \theta_0 + \omega_2)$ using a twice continuously differentiable periodic function $w(\theta)$, where

$$w(\theta) \begin{cases} = 1 & \text{if } \theta \in (\theta_0, \theta_0 + \omega_2) \pmod{2\pi} \\ = 0 & \text{if } \theta \notin [\theta_0 - \log^{-C} x, \theta_0 + \omega_2 + \log^{-C} x)] \pmod{2\pi}, \end{cases}$$

and where $|w''(\theta)| \ll \log^{2C} x$. Then

$$S(\chi, \mathcal{U}) = \sum_{N' < N(z) \le N'(1+\omega)} \beta'_z \chi(z) w(\arg z) + O\left(\frac{N}{\log^C x}\right).$$

The Fourier coefficients of w satisfy $c_k \ll k^{-2} \log^{2C} x$ for $k \neq 0$, whence

$$w(\arg z) = \sum_{k} c_k \left(\frac{z}{|z|}\right)^k = \sum_{|k| \le \log^{3C} x} c_k \left(\frac{z}{|z|}\right)^k + O(\log^{-C} x).$$

It then suffices to show that

$$S(\chi, N', k) := \sum_{N' < N(z) \le N'(1+\omega)} \beta'_z \chi(z) \left(\frac{z}{|z|}\right)^k \ll_C N(\log x)^{-4C}$$

for any C > 0, and for $|k| \le \log^{3C} x$. Indeed we will do rather better, and show that one can achieve a small power saving in N.

Recall that $\beta_z = \beta_{N(z)}$, where β_n is the indicator function of a set of one of the shapes

$$Q_j := \{ p_1 \dots p_{j+1} \in (N', N'(1+\omega)] : p_{j+1} \in J, \ p_{j+1} < \dots < p_1 < Y, \\ p_1 \dots p_j < Y \le p_1 \dots p_{j+1} < x^{1/2-\delta} \}$$

or

 $R:=\{n\in (N',N'(1+\omega)]:(n,P(V))=1\}.$

Here we will have $0 \leq j \leq n_0 = \left[\frac{\log Y}{\delta \log x}\right]$, and $J = [V, V(1 + \kappa)) \subseteq [x^{\delta}, Y)$. In particular we interpret Q_0 to be $\{p : p \in J \cap (N', N'(1 + \omega))\}$.

It will be convenient to write

(33)
$$\lambda(n) = \sum_{N(z)=n}^{P} \chi(z) \left(\frac{z}{|z|}\right)^{k} [z],$$

where $\sum_{i=1}^{P} denotes a sum over primitive z.$ Then

$$S(\chi, N', k) = \sum_{n} \lambda(n),$$

where n runs over R, or one of the sets Q_j . We discuss the procedure in the case of Q_j , the situation for the set R being similar, or indeed easier. One small difference is that

elements of R need not be square-free. However integers $n \in R$ which are not square-free make a negligible contribution, since any prime factor must be at least x^{δ} .

We begin by handling the terms in which the largest prime factor of n, which we write as P(n) say, exceeds $N^{99/100}$. The contribution from such integers is

$$\sum_{\substack{m \le 2N^{1/100} \ p > \max(P(m), N^{99/100}) \\ mp \in Q_j}} \lambda(mp).$$

Since p is the largest prime factor of mp one sees from the definition of the set Q_j that one may rewrite the conditions p > P(m) and $mp \in Q_j$ to say that p runs over a certain interval $I_j(m) \subseteq [N/m, 2N/m)$. We may then apply the following result of Iwaniec and Friedlander [4, Theorem 2^{ψ}]

Lemma 16. For any $m \ge 1$ we have

$$\sum_{n \le U} \Lambda(n)\lambda(mn) \ll f(|k|+1)mU^{76/77},$$

where λ is given by (33) and χ is a character modulo f.

Note that in our situation fk is at most a power of $\log x$. The above result shows that

$$\sum_{\substack{m \le 2N^{1/100} \ p > \max(P(m), N^{99/100}) \\ mp \in Q_j}} \sum_{\substack{\lambda(mp) \\ mp \in Q_j}} \lambda(mp) \ll f(|k|+1) \sum_{\substack{m \le 2N^{1/100} \\ m \le 2N^{1/100}}} m(N/m)^{76/77} \\ \ll f(|k|+1)N^{76/77 + (78/77)/100}.$$

Since 76/77 + (78/77)/100 < 1 this is satisfactory.

We now examine the terms in which every prime factor of n is at most $N^{99/100}$. To do this, we first rewrite our sum in terms of bilinear sums. Let $n = p_1 \dots p_{j+1}$, as in the description of the set Q_j , and divide the range for each prime p_i into intervals of the form $(P_i, 2P_i]$. This will give us at most $(2 \log N)^{1+n_0}$ sets of dyadic ranges, and since $n_0 \ll \delta^{-1} = (\log x)^{1-\varpi}$ there will be $O(N^{\epsilon})$ sets of ranges. Moreover we may suppose that

$$\prod_{i=1}^{j+1} P_i \ll N \ll 2^{j+1} \prod_{i=1}^{j+1} P_i.$$

Since we may now assume that $P_1 \leq N^{99/100}$ there will be an index u such that

$$N^{1/100} \le \prod_{i=1}^{u} P_i \le N^{99/100}.$$

Fixing such an index we split n as n = ab with

$$a = \prod_{i=1}^{u} p_i$$
, and $b = \prod_{i=u+1}^{j+1} p_i$,

so that $a \leq N_1$ and $b \leq N_2$ with

$$N_1 := 2^{1+n_0} \prod_{i=1}^u P_i$$
 and $N_2 := 2^{1+n_0} \prod_{i=u+1}^{j+1} P_i$,

and hence

(34)
$$N_1 N_2 \ll N^{1+\epsilon} \text{ and } N_1, N_2 \ll N^{99/100+\epsilon}$$

We will then have $N_1 N^{-\epsilon} \ll a \leq N_1$, and similarly for *b*. Our description of Q_j may now be expressed by requiring that $a \in Q_{j,u}$ and $b \in Q'_{j,u}$ for appropriate sets $Q_{j,u}$ and $Q'_{j,u}$, together with the conditions that $ab \in I = (N', N'(1+\omega)] \cap [Y, x^{1/2-\delta})$, that $p_{j+1}^{-1}ab < Y$, and that $p_{u+1} < p_u$. Specifically, we take

$$Q_{j,u} = \{ a = p_1 \dots p_u : p_i \in (P_i, 2P_i], p_u < \dots < p_1 < Y \},\$$

and

$$Q'_{j,u} = \{b = p_{u+1} \dots p_{j+1} : p_i \in (P_i, 2P_i], p_{j+1} \in J, p_{j+1} < \dots < p_{u+1} < Y\}$$

In order to separate the variables a and b completely we subdivide the available ranges for a, b, p_{j+1}, p_u and p_{u+1} into intervals of the shape (A, A + A/L], (B, B + B/L], $(P'_{j+1}, P'_{j+1} + P'_{j+1}/L]$, $(P'_u, P'_u + P'_u/L]$ and $(P'_{u+1}, P'_{u+1} + P'_{u+1}/L]$. Here the parameter L will be a small power of N. The reader should note that we do not insist that each of these intervals should have length 1 or more. Indeed such an interval may not contain any integers at all. There will be $O(L^5(\log x)^2)$ collections of such intervals. There will be some for which the conditions $ab \in I$, $p_{j+1}^{-1}ab < Y$, and $p_{u+1} < p_u$ hold for every choice of p_1, \ldots, p_{j+1} satisfying

 $a \in (A, A + A/L], \quad b \in (B, B + B/L]$

 $p_{j+1} \in (P'_{j+1}, P'_{j+1} + P'_{j+1}/L], \quad p_u \in (P'_u, P'_u + P'_u/L], \quad p_{u+1} \in (P'_{u+1}, P'_{u+1} + P'_{u+1}/L],$ and

$$p_i \in I_i \quad (i \neq 1, u, u+1)$$

In this case the corresponding subsum is

$$\sum_{\substack{a \in Q_{j,u} \cap (A,A+N_1/L] \\ p_u \in (P'_u, P'_u + P'_u/L]}} \sum_{\substack{b \in Q'_{j,u} \cap (B,B+B/L] \\ p_{j+1} \in (P'_{j+1}, P'_{j+1} + P'_{j+1}/L] \\ p_{u+1} \in (P'_{u+1}, P'_{u+1} + P'_{u+1}/L]}} \lambda(ab)$$

so that we have separated the variables a and b. For such sums we can apply the following consequence of Friedlander and Iwaniec [4, Proposition 23.1].

Lemma 17. Let $\mathfrak{d}_1(m)$ and $\mathfrak{d}_2(n)$ be bounded arithmetic functions supported on $1 \le m \le N_1$ and $1 \le n \le N_2$ respectively. Then

$$\sum_{m,n} \mathfrak{d}_1(m) \mathfrak{d}_2(n) \lambda(mn) \ll (N_1 + N_2)^{\frac{1}{12}} (N_1 N_2)^{\frac{11}{12} + \epsilon}.$$

Since $A \leq N_1$ and $B \leq N_2$ this gives us a bound

$$\ll (N^{99/100})^{1/12} N^{11/12+\epsilon} = N^{1-1/1200+\epsilon}$$

in view of (34). Since there are $O(L^5 N^{\epsilon})$ such subsums the overall contribution is $O(L^5 N^{1-1/1200+\epsilon})$.

It remains to consider the contribution from "bad" sets of ranges which are not excusively contained in the region given by $ab \in I$, $p_{j+1}^{-1}ab < Y$, and $p_{u+1} < p_u$. Suppose that the interval I is given by $I = [e_1, e_2]$, for example, and that there are integers $a, a' \in (A, A + A/L]$ and $b, b' \in (B + B/L]$ for which $ab \in I$ but $a'b' \notin I$. Then we must have $ab = (1 + O(L^{-1}))e_1$ or $ab = (1 + O(L^{-1}))e_2$. We now consider the total contribution from integers $n \in Q_j$ for all such "bad" choices of intervals (A, A + A/L], (B, B + B/L], $(P'_{j+1}, P'_{j+1} + P'_{j+1}/L]$, $(P'_u, P'_u + P'_u/L]$ and $(P'_{u+1}, P'_{u+1} + P'_{u+1}/L]$. Since each integer n occurs at most once, and $\lambda(n) = O(\tau(n))$, the contribution will be

$$\ll \sum_{\substack{n = (1 + O(L^{-1}))e_i \\ 37}} \tau(n) \ll N^{1 + \epsilon} L^{-1}.$$

Similarly, if we have $p_{j+1}^{-1}ab < Y$ but $p'_{j+1}^{-1}a'b' \ge Y$, then $p_{j+1}^{-1}ab = (1 + O(L^{-1}))Y$. Now $P_{j+1}Y \simeq AB \le N_1N_2 \ll N^{1+\epsilon}$, so any *n* which is to be counted will have a prime factor $p \ll N^{1+\epsilon}/Y$ such that $p^{-1}n = (1 + O(L^{-1}))Y$. Thus, writing n = pm, we see that the total contribution in this case is

$$\ll \sum_{p \ll N^{1+\epsilon}/Y} \sum_{m=(1+O(L^{-1}))Y} \tau(pm) \ll N^{1+\epsilon}Y^{-1}(1+L^{-1}Y) \ll N^{1+\epsilon}L^{-1},$$

for $L \leq Y$.

Lastly, if $P_u = P_{u+1}$, then it may happen that the condition $p_{u+1} < p_u$ is satisfied by some, but not all, pairs of primes (p_u, p_{u+1}) from the intervals $(P'_u, P'_u + P'_u/L)$ and $(P'_{u+1}, P'_{u+1} + P'_{u+1}/L)$. Clearly this problem cannot arise when $L \ge 2P_u$ since then the intervals $(P'_u, P'_u + P'_u/L)$ and $(P'_{u+1}, P'_{u+1} + P'_{u+1}/L)$ contain at most one prime each. It follows that any *n* to be counted in this case will have two prime factors $p' > p \ge P_u \ge L/2$ with $p' = (1 + O(L^{-1}))p$. Hence the corresponding contribution is

$$\ll \sum_{\substack{p' > p \ge L/2 \\ p' = (1+O(L^{-1}))p}} \sum_{\substack{n \ll N \\ p'p|n}} \tau(n) \ll \sum_{\substack{p' > p \ge L/2 \\ p' = (1+O(L^{-1}))p}} \frac{N^{1+\epsilon}}{p'p} \ll N^{1+\epsilon}L^{-1}$$

We therefore find that our sum is

$$\ll L^5 N^{1-1/1200+\epsilon} + N^{1+\epsilon} L^{-1}$$

if $L \leq Y$. We may then choose $L = N^{1/10000}$ for example, to achieve the claimed power saving.

11.2. Small d. To handle small d it will be enough to show for any $f \leq F$, $d \leq D_1$, and any non-principal $\chi \pmod{d}$, that

$$\sum_{\substack{c \pmod{df} \ (z_1, z_2) \in \mathcal{U}_1 \times \mathcal{U}_2 \\ cz_2 \equiv z_1 \pmod{df}}}^* \beta_{z_1} \beta_{z_2} \chi(c) \ll_C \frac{N^2}{\log^C x}$$

for every C > 0. Since

$$\sum_{c \pmod{df}}^* \chi(c) = 0,$$

it suffices to prove that if $\mathcal{U} = \mathcal{U}_1$ or \mathcal{U}_2 then there is an $\mathfrak{M} = \mathfrak{M}(\mathcal{U}, df)$ such that

(35)
$$\sum_{\substack{z \in \mathcal{U} \\ z \equiv \alpha \pmod{df}}} \beta_z = \mathfrak{M} + O\left(\frac{N}{\log^C x}\right),$$

for any $(\alpha, df) = 1$, and any C > 0.

As in Section 11.1, we may assume that $\beta_z = \beta_{N(z)}$, where β_n is the indicator function of either Q_j or R. We describe the procedure for Q_j , the method for R being similar. We decompose z as z_1z_2 with $N(z_1)$ being the largest prime factor of $N(z_1z_2)$. The requirement that $n \in Q_j$ is then equivalent to a condition of the form $N(z_2) \in Q'_j$ together with a restriction of the type $N(z_1) \in I(z_2)$ for some real interval $I(z_2)$. Specifically we have

$$Q'_{j} = \{ p_{2} \dots p_{j+1} : p_{j+1} \in J, \ p_{j+1} < \dots < p_{2} \}$$

and

$$I(z_2) = (p_2, Y) \cap \left(\frac{N'}{N(z_2)}, \frac{N'(1+\omega)}{N(z_2)}\right] \cap \left[\frac{Y}{N(z_2)}, \frac{x^{1/2-\delta}}{N(z_2)}\right),$$

where p_2 is the largest prime factor of $N(z_2)$. When \mathcal{U} is given by (32) the condition on the size of $N(z_1z_2)$ is exactly the condition

$$N(z_1) \in \left(\frac{N'}{N(z_2)}, \frac{N'(1+\omega)}{N(z_2)}\right],$$

and we have $\theta_0 < \arg z < \theta_0 + \omega_2$ exactly when

$$\theta_1(z_2) < \arg z_1 < \theta_1(z_2) + \omega_2,$$

with $\theta_1(z_2) = \theta_1 - \arg z_2$.

It follows that

(36)
$$\sum_{\substack{z \in \mathcal{U} \\ z \equiv \alpha \pmod{df}}} \beta_z = \sum_{\substack{z_2 \in Q'_j \\ (z_2, df) = 1}} \mathcal{N}(z_2, \alpha),$$

where $\mathcal{N}(z_2, \alpha)$ is the number of Gaussian integers z_1 satisfying

$$z_1 \equiv \alpha \overline{z_2} \pmod{df}, \quad N(z_1) \in I(z_2), \quad \text{and} \quad \theta_1(z_2) < \arg z_1 < \theta_1(z_2) + \omega_2,$$

and for which $N(z_1)$ is prime. Here $\overline{z_2}$ is the inverse of z_2 modulo df.

We can estimate $\mathcal{N}(z_2, \alpha)$ using a form of the Prime Number Theorem for arithmetic progressions, over number fields. Given $q \in \mathbb{N}$, any Gaussian integer α coprime to q, and any $\theta \in [0, 2\pi]$ write $\pi(x; q, \alpha, \theta)$ for the number of Gaussian primes $\mu \equiv \alpha \pmod{q}$ of norm at most x and with $0 \leq \arg(\mu) \leq \theta$. The principal result of Mitsui [10] tells us that there is an absolute constant c such that

(37)
$$\pi(x;q,\alpha,\theta) = \frac{4}{\phi_{\mathbb{Q}(i)}(q)} \frac{\theta}{2\pi} \mathrm{Li}(x) + O_A(x \exp(-c\sqrt{\log x}))$$

uniformly for all $\theta \in [0, 2\pi]$ and all $q \leq (\log x)^A$. Here $\phi_{\mathbb{Q}(i)}(q)$ is the Euler ϕ -function for the Gaussian integers.

We now apply (37) to estimate $\mathcal{N}(z_2, \alpha)$. We have $I(z_2) \subseteq (0, 2N/N(z_2)]$, and so we will need to know that $df \leq (\log 2N/N(z_2))^A$ for some constant A. However we recall that if p divides an element of Q_j then one has $p \geq x^{\delta}$ with $\delta = (\log x)^{\varpi-1}$. Thus we will have $2N/N(z_2) \geq x^{\delta}$ so that $df \leq (\log 2N/N(z_2))^{C/\varpi}$ whenever $df \leq (\log x)^C$. The required condition is therefore satisfied when $f \leq F$ and $d \leq D_1$.

We therefore find that

$$\mathcal{N}(z_2, \alpha) = \mathfrak{M}(z_2, df, j, \mathcal{U}) + O\left(\frac{N}{N(z_2)} \exp(-c(\log x)^{\varpi/2})\right),$$

where the main term $\mathfrak{M}(z_2, df, j, \mathcal{U})$ is, crucially, independent of α . If we feed this into (36) we then obtain the desired estimate (35). This completes our treatment of small d.

12. DISTRIBUTION OF SEQUENCES IN ARITHMETIC PROGRESSIONS

The purpose of this final section is to prove Corollary 2 and the more general Theorem 2 below. Theorem 2 is motivated by the possibility that an interesting arithmetic function can be biased for certain small moduli — that is, the sequences does not satisfy the Siegel–Walfisz condition. In this case, we can still prove a result about the distribution of such a sequence in arithmetic progressions by adjusting the main term.

We first fix some notation for the rest of the section: for any arithmetic function a(n) with finite support, we let

$$||a\tau||^2 = \sum_{\substack{n \\ 39}} |a(n)|^2 \tau(n)^2.$$

For arithmetic functions c_1 and c_2 , we shall see that $||c_1\tau||^2 ||c_2\tau||^2$ is an upper bound for $||c_1 * c_2||^2$.

Let $\gamma(n)$ and $\delta(n)$ be arithmetic functions supported on $n \leq N_1$ and $n \leq N_2$ respectively and let $Q_0 \geq 1$. For a character χ , let $Q(\chi)$ be the conductor of the unique primitive character which induces χ . Now, let

$$S(a,q) = \sum_{\substack{n_1 \equiv an_2 \pmod{q} \\ (n_1n_2,q)=1}} \gamma(n_1)\delta(n_2),$$

$$\mathcal{M}(a,q) = \frac{1}{\phi(q)} \sum_{n_1,n_2} \gamma(n_1)\delta(n_2) \sum_{\substack{\chi \pmod{q} \\ Q(\chi) \le Q_0}} \chi(n_1) \overline{\chi(an_2)}$$

and

$$\mathcal{E}(a,q) = S(a,q) - \mathcal{M}(a,q).$$

Note that $\mathcal{M}(a,q)$ is the expected main term for S(a,q). The main result of this section is below.

Theorem 2. For any $Q \in \mathbb{N}$, let

$$\mathcal{E} = \sum_{q \le Q} \sum_{a \pmod{q}}^{*} |\mathcal{E}(a,q)|^2.$$

Then,

$$\mathcal{E} \ll \left(Q + \frac{N_1 N_2}{Q_0}\right) (\log Q) \|\gamma \tau\|^2 \|\delta \tau\|^2.$$

12.1. **Proof of Theorem 2.** We first prove a consequence of the large sieve which lies at the heart of our result.

Proposition 9. Let a(n) be any arithmetic function supported on $n \leq N$, and for any character χ , let

$$A(\chi) = \sum_{n} a(n)\chi(n).$$

Further, let $Q(\chi)$ denote the conductor of the unique primitive character that induces χ . Then

$$S := \sum_{h \le H} \frac{1}{\phi(h)} \sum_{\substack{\chi \pmod{h} \\ Q(\chi) > h_0}} |A(\chi)|^2 \ll \left(H + \frac{N}{h_0}\right) (\log H) ||a||^2.$$

Proof. For each non-principal character $\chi \pmod{h}$, let $\psi \pmod{h_1}$ be the unique primitive character which induces χ , where we may write $h = h_1h_2$ for $h_1 > 1$. Then for any n, $\chi(n) = \psi(n)$ if $(n, h_2) = 1$ and $\chi(n) = 0$ otherwise. Let us write

$$A(\psi, h_2) = \sum_{\substack{n \ (n,h_2)=1}}^{n} a(n)\psi(n),$$

whence

$$S \leq \sum_{\substack{h_1h_2 \leq H \\ h_1 \geq h_0}} \frac{1}{\phi(h_2)} \frac{1}{\phi(h_1)} \sum_{\psi \pmod{h_1}}^* |A(\psi, h_2)|^2,$$

where $\sum_{\psi \pmod{h_1}}^{*}$ denotes a sum over all primitive characters modulo h_1 . Applying the multiplicative large sieve (see e.g. (9.52) in [3]) in dyadic ranges, we see that

$$\sum_{h_0 \le h_1 \le H/h_2} \frac{1}{\phi(h_1)} \sum_{\psi \pmod{h_1}}^* |A(\psi, h_2)|^2 \ll \left(\frac{H}{h_2} + \frac{N}{h_0}\right) \|a\|^2,$$

whence

$$S \ll \sum_{h_2 \le H} \frac{1}{\phi(h_2)} \left(\frac{H}{h_2} + \frac{N}{h_0} \right) \|a\|^2$$
$$\ll \left(H + \frac{N \log H}{h_0} \right) \|a\|^2.$$

The Proposition then follows.

We now proceed to the proof of Theorem 2. *Proof.* We have

$$\mathcal{E}(a,q) = \frac{1}{\phi(q)} \sum_{\substack{\chi \pmod{q} \\ Q(\chi) > Q_0}} G(\chi) \overline{D(\chi)\chi(a)},$$

where

$$G(\chi) = \sum_{n} \gamma(n) \chi(n),$$

and

$$D(\chi) = \sum_{n} \overline{\delta(n)} \chi(n).$$

Then

$$\begin{aligned} \mathcal{E} &= \sum_{q \leq Q} \frac{1}{\phi(q)^2} \sum_{\substack{a \pmod{q} \\ a \pmod{q}}} \left| \sum_{\substack{\chi \pmod{q} \\ Q(\chi) > Q_0}} G(\chi) \overline{D(\chi)\chi(a)} \right|^2 \\ &= \sum_{q \leq Q} \frac{1}{\phi(q)} \sum_{\substack{\chi \pmod{q} \\ Q(\chi) > Q_0}} \left| G(\chi) \overline{D(\chi)} \right|^2 \\ &\ll \left(Q + \frac{N_1 N_2}{Q_0} \right) (\log Q) ||a||^2, \end{aligned}$$

where

$$a(n) = \sum_{n=n_1n_2} \gamma(n_1) \overline{\delta(n_2)},$$

and where we have used Proposition 9 with a(n) supported on $n \leq N_1 N_2$.

However Cauchy's inequality yields

$$|a(n)|^{2} \leq \tau(n) \sum_{n=n_{1}n_{2}} |\gamma(n_{1})\delta(n_{2})|^{2} \leq \sum_{n=n_{1}n_{2}} |\gamma(n_{1})\delta(n_{2})|^{2} \tau(n_{1})\tau(n_{2}),$$

whence

$$\|a\|^{2} \leq \sum_{n_{1},n_{2}} |\gamma(n_{1})\delta(n_{2})|^{2} \tau(n_{1})\tau(n_{2}) \leq \|\gamma\tau\|^{2} \|\delta\tau\|^{2}.$$

It follows that

$$\mathcal{E} \ll \left(Q + \frac{N_1 N_2}{Q_0}\right) (\log Q) \|\gamma \tau\|^2 \|\delta \tau\|^2,$$

as desired.

12.2. Proof of Corollary 2. We have

$$S(x; a, q) = \frac{1}{\phi(q)} \sum_{\chi \pmod{q}} \sum_{n_1, n_2} \chi(n_1) \overline{\chi(an_2)} c_1(n_1) c_2(n_2)$$

Then, taking $Q_0 = (\log x)^{A+1}$, we see that

$$|S(x; a, q) - S(x; q)|^{2}$$

$$= \left| \frac{1}{\phi(q)} \sum_{\substack{\chi \pmod{q} \\ \chi \neq \chi_{0}}} \sum_{n_{1}, n_{2}} \chi(n_{1}) \overline{\chi(an_{2})} c_{1}(n_{1}) c_{2}(n_{2}) \right|^{2}$$

$$= |\mathcal{E}(a, q) + E(a, q, Q_{0})|^{2}$$

$$\ll |\mathcal{E}(a, q)|^{2} + |E(a, q, Q_{0})|^{2},$$

where

$$E(a,q,Q_0) = \frac{1}{\phi(q)} \sum_{\substack{\chi \pmod{q} \\ 1 < Q(\chi) \le Q_0}} \left\{ \sum_n c_1(n)\chi(n) \right\} \left\{ \sum_n c_2(n)\overline{\chi(an)} \right\}$$

and $\mathcal{E}(a,q)$ is as defined in Theorem 2 with $\gamma = c_1$ and $\delta = c_2$. We let B(A) = A + 1 and apply Theorem 2 to see that

$$\sum_{q \le Q} \sum_{a \pmod{q}}^{*} |S(x;a,q) - S(x;q)|^2 \ll \frac{x^2}{\log^A x} ||c_1\tau||^2 ||c_2\tau||^2 + \sum_{q \le Q} \sum_{a \pmod{q}}^{*} |E(a,q,Q_0)|^2.$$

It now suffices to bound the last term. We apply the Siegel–Walfisz condition (1) with $Q_0 \leq (\log x)^{\kappa}$, for some constant $\kappa \geq A + 1$ to be determined, and deduce that

$$\sum_{n} c_1(n)\chi(n) \ll x^{1/2} \|c_1\| (\log x)^{-\kappa}.$$

The sum involving c_2 may be bounded trivially as

$$\sum_{n} |c_2(n)| \le x^{1/2} ||c_2||,$$

and since the number of characters with modulus less than Q_0 is at most Q_0^2 , we have

$$E(a, q, Q_0) \ll \frac{Q_0^2 x \|c_1\| \|c_2\|}{\phi(q) (\log x)^{\kappa}}$$

Thus

$$\sum_{q \le Q} \sum_{a \pmod{q}}^{*} |E(a, q, Q_0)|^2 \ll \frac{Q_0^4 x^2 ||c_1||^2 ||c_2||^2}{(\log x)^{2\kappa}} \sum_{q \le Q} \frac{1}{\phi(q)}$$
$$\ll \frac{Q_0^4 x^2 ||c_1||^2 ||c_2||^2}{(\log x)^{2\kappa - 1}}.$$

We then see that the Corollary follows upon choosing

$$2\kappa = 5A + 5.$$

References

- [1] H. Davenport, Multiplicative Number Theory, GTM vol.74, Springer-Verlag, New York, 2000.
- [2] E. Fouvry and H. Iwaniec, Gaussian primes. Acta Arith. 79 (1997), no. 3, 249–287.
- [3] J. Friedlander and H. Iwaniec, Opera de cribro. American Mathematical Society Colloquium Publications, 57. American Mathematical Society, Providence, RI, 2010.
- [4] J. Friedlander and H. Iwaniec, The polynomial $X^2 + Y^4$ captures its primes. Ann. of Math. (2) 148 (1998), no. 3, 945–1040.
- [5] J. Friedlander and H. Iwaniec, Gaussian sequences in arithmetic progressions. Funct. Approx. Comment. Math. 37 (2007), no. 1, 149–157.
- [6] J. Friedlander and H. Iwaniec, Asymptotic sieve for primes. Ann. of Math. (2) 148 (1998), no. 3, 1041–1065.
- [7] G. Harman, *Prime-detecting sieves*. London Mathematical Society Monographs Series, 33. Princeton University Press, Princeton, NJ, 2007.
- [8] D.R. Heath-Brown, Primes represented by $x^3 + 2y^3$. Acta Math. 186 (2001), no. 1, 1–84.
- [9] D.R. Heath-Brown and B.Z. Moroz, On the representation of primes by cubic polynomials in two variables. Proc. London Math. Soc. (3) 88 (2004), no. 2, 289–312.
- [10] T. Mitsui, Generalized prime number theorem. Jap. J. Math. 26 (1956), 1–42.
- [11] E.C. Titchmarsh, The theory of the Riemann zeta-function. Second edition. The Clarendon Press, Oxford University Press, New York, 1986.

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING, RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD, UK, OX2 6GG

E-mail address: Roger.Heath-Brown@maths.ox.ac.uk

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING, RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD, UK, OX2 6GG *E-mail address*: lix1@maths.ox.ac.uk