Quartic isospin asymmetry energy of nuclear matter from chiral pion-nucleon dynamics^{[1](#page-0-0)}

N. Kaiser

Physik Department T39, Technische Universität München, D-85747 Garching, Germany

email: nkaiser@ph.tum.de

Abstract

Based on a chiral approach to nuclear matter, we calculate the quartic term in the expansion of the equation of state of isospin-asymmetric nuclear matter. The contributions to the quartic isospin asymmetry energy $A_4(k_f)$ arising from 1 π -exchange and chiral 2 π -exchange in nuclear matter are calculated analytically together with three-body terms involving virtual $\Delta(1232)$ isobars. From these interaction terms one obtains at saturation density $\rho_0 = 0.16 \,\text{fm}^{-3}$ the value $A_4(k_{f0}) = 1.5 \,\text{MeV}$, more than three times as large as the kinetic energy part. Moreover, iterated 1 π-exchange exhibits components for which the fourth derivative with the respect to the isospin asymmetry parameter δ becomes singular at $\delta = 0$. The genuine presence of a non-analytical term $\delta^4 \ln |\delta|$ in the expansion of the energy per particle of isospin-asymmetric nuclear matter is demonstrated by evaluating a s-wave contact interaction at second order.

PACS: 21.60.Jz, 21.65.+f, 24.10.Cn.

1 Introduction and summary

The determination of the equation of state of isospin-asymmetric nuclear matter has been a longstanding goal shared by both nuclear physics and astrophysics [\[1\]](#page-5-0). Usually one assumes a parabolic form for the energy per nucleon at zero temperature, $\bar{E}_{as}(\rho_p, \rho_n) = \bar{E}(\rho) + A_2(\rho) \delta^2 + \mathcal{O}(\delta^4)$, where $\rho = \rho_p + \rho_n$ is the total nucleon density and $\delta = (\rho_n - \rho_p)/\rho$ is the isospin asymmetry related to unequal proton and neutron densities $\rho_p \neq \rho_n$. The validity of the quadratic approximation has been verified with good numerical accuracy from isospin-symmetric nuclear matter $(\delta = 0)$ up to pure neutron matter $(\delta = 1)$ by most of the existing nuclear many-body theories using various interactions [\[2\]](#page-5-1). Nonetheless, it has been shown consistently in numerous studies [\[3\]](#page-5-2) that for some properties of neutron stars, such as the proton fraction at beta-equilibrium, the core-crust transition density and the critical density for the direct URCA process to occur, even a very small quartic isospin asymmetry energy $A_4(\rho)$ (multiplied with δ^4 in the expansion of the energy per nucleon) can make a big difference.

Given the fact that all the available numerical solutions of the nuclear many-body problem confirm the validity of the quadratic approximation, the quartic isospin asymmetry $A_4(\rho)$ should be rather small. However, in the recent work by Cai and Li [\[4\]](#page-5-3), which employs an empirically constrained isospin-dependent single-nucleon momentum-distribution and the equation of state of pure neutron matter near the unitary limit, a significant quartic isospin asymmetry energy of $A_4(\rho_0) = (7.2 \pm 2.5)$ MeV has been found. This value amounts to about 16 times the free Fermi gas prediction, see eq.(2). On the other hand, the calculations by the Darmstadt group [\[5\]](#page-5-4) based on chiral low-momentum interactions and many-body perturbation theory lead to a small value of $A_4(\rho_0) = (1.0 \pm 0.2)$ MeV.

The purpose of the present paper it to give a prediction for the density-dependent quartic isospin asymmetry energy $A_4(k_f)$ in the chiral approach to nuclear matter developed in refs.[\[6,](#page-5-5) [7\]](#page-5-6).

¹This work has been supported in part by DFG and NSFC (CRC 110).

In this approach the long- and medium-range NN-interactions arising from multi-pion exchange are treated explicitly and few parameters encoding the relevant short-distance dynamics are adjusted to bulk properties of nuclear matter. Single-particle potentials [\[7\]](#page-5-6), quasi-particle interactions [\[8\]](#page-5-7), the thermodynamic behavior of nuclear matter at finite temperatures [\[9\]](#page-5-8) and the density-dependence of the in-medium quark condensate [\[10\]](#page-5-9) follow then as predictions in that framework (see also the recent review article [\[11\]](#page-5-10)).

The present paper is organized as follows. In section 2, analytical expressions are given for the contributions to the quartic isospin asymmetry energy $A_4(k_f)$ as they arise from 1π -exchange and chiral 2π -exchange. The three-nucleon interaction generated by 2π -exchange and excitation of a virtual ∆(1232)-isobar is considered as well. These interaction contributions lead at saturation density $\rho_0 = 0.16 \,\text{fm}^{-3}$ (or $k_{f0} = 263 \,\text{MeV}$) to the (small) value $A_4(k_{f0}) = 1.5 \,\text{MeV}$, which amounts to about three times the kinetic energy part. Moreover, in the course of the calculation one encounters components of the second-order 1π -exchange whose representation of the fourth derivative with the respect to δ at $\delta = 0$ is singular. In section 3, the generic presence of a non-analytical term $\delta^4 \ln |\delta|$ in the expansion of the energy per particle of isospin-asymmetric nuclear matter is demonstrated by calculating in closed form the second-order contribution from a s-wave contact-interaction. Clearly, after having established its existence, the non-analytical term δ^4 ln $|\delta|$ should be included in future fits of the equation of state of (zero-temperature) isospin-asymmetric nuclear matter.

2 One-pion and two-pion exchange contributions

In this section we collect the expressions for the quartic isospin asymmetry $A_4(k_f)$ as they arise from one-pion and two-pion exchange diagrams following refs.[\[6,](#page-5-5) [7\]](#page-5-6). Isospin-asymmetric (spin-saturated) nuclear matter is characterized by different proton and neutron Fermi momenta, $k_{p,n} = k_f (1 \mp \delta)^{1/3}$. Expanding the energy per particle at fixed nucleon density $\rho = 2k_f^3/3\pi^2$ in the isospin asymmetry parameter δ up to fourth order gives:

$$
\bar{E}_{\rm as}(k_p, k_n) = \bar{E}(k_f) + \delta^2 A_2(k_f) + \delta^4 A_4(k_f) + \mathcal{O}(\delta^6) \,, \tag{1}
$$

with $A_2(k_f)$ the (usual) quadratic isospin asymmetry energy. We view the density-dependent expansion coefficients $E(k_f)$, $A_2(k_f)$ and $A_4(k_f)$ as functions of the Fermi momentum k_f , since they emerge in this form directly from the calculation. The first contribution to $A_4(k_f)$ comes from the relativistically improved kinetic energy $T_{\text{kin}}(p) = p^2/2M - p^4/8M^3$, and it reads:

$$
A_4(k_f)^{(\text{kin})} = \frac{k_f^2}{162M} \left(1 + \frac{k_f^2}{4M^2} \right),\tag{2}
$$

with $M = 939$ MeV the average nucleon mass. The corresponding value at nuclear matter saturation density $\rho_0 = 0.16 \,\text{fm}^{-3}$ (or $k_{f0} = 263 \,\text{MeV}$) is $A_4 (k_{f0})^{\text{(kin)}} = 0.464 \,\text{MeV}$.

For the treatment of two-body interactions that depend on the momentum transfer $|\vec{p}_1-\vec{p}_2|$ the following expansion formulas for integrals over two Fermi spheres are most helpful:

$$
\int \frac{d^3 p_1 d^3 p_2}{(2\pi)^6} F(|\vec{p_1} - \vec{p_2}|) \left[\theta(k_p - |\vec{p_1}|) \theta(k_p - |\vec{p_2}|) + \theta(k_n - |\vec{p_1}|) \theta(k_n - |\vec{p_2}|) \right]
$$

=
$$
\frac{2k_f^6}{3\pi^4} \int_0^1 dz \left\{ \left[z^2 (1-z)^2 (2+z) + \frac{\delta^2 z^3}{3} \right] F(2zk_f) + \frac{\delta^4 k_f}{162} \left[F'(2k_f) - 7z^4 F'(2zk_f) \right] \right\},
$$
 (3)

$$
\int \frac{d^3 p_1 d^3 p_2}{(2\pi)^6} F(|\vec{p_1} - \vec{p_2}|) \theta(k_p - |\vec{p_1}|) \theta(k_n - |\vec{p_2}|)
$$

=
$$
\frac{k_f^6}{3\pi^4} \int_0^1 dz \left\{ \left[z^2 (1-z)^2 (2+z) + \frac{\delta^2 z}{3} (z^2 - 1) \right] F(2zk_f) + \frac{\delta^4 k_f}{162} (8z^2 - 1 - 7z^4) F'(2zk_f) \right\}.
$$
 (4)

The z-dependent weighting functions at order δ^2 and δ^4 have been obtained by applying several partial integrations. The contribution of the 1π -exchange Fock diagram to the quartic isospin asymmetry energy reads:

$$
A_4(k_f)^{(1\pi)} = \frac{g_A^2 m_\pi^3}{(36\pi f_\pi)^2} \left\{ \left(4u + \frac{21}{8u} \right) \ln(1 + 4u^2) - 2u^3 - \frac{33u}{4} - \frac{u(9 + 44u^2)}{4(1 + 4u^2)^2} + \frac{m_\pi^2}{M^2} \left[2u^5 + 2u^3 + \frac{3u}{8} - u^3 \ln(1 + 4u^2) - \frac{u(3 + 16u^2)}{8(1 + 4u^2)^2} - \frac{3u^2}{2} \arctan 2u \right] \right\}, \quad (5)
$$

with the dimensionless variable $u = k_f/m_\pi$. The second line in eq.(5) gives the relativistic $1/M^2$ correction. The occurring physical parameters are: nucleon axial-vector coupling constant $g_A = 1.3$, (neutral) pion mass $m_{\pi} = 135 \text{ MeV}$, and pion decay constant $f_{\pi} = 92.4 \text{ MeV}$.

Next in the chiral expansion comes the iterated (second-order) 1π -exchange. With two medium insertions $\frac{1}{2}(1+\tau_3)\theta(k_p-|\vec{p}_i|)+\frac{1}{2}(1-\tau_3)\theta(k_n-|\vec{p}_i|)$ one gets a Hartree contribution of the form:

$$
A_4(k_f)^{(\text{H2})} = \frac{g_A^4 M m_\pi^4}{(24\pi)^3 f_\pi^4} \left\{ 10u^3 - \frac{61u}{2} + \frac{200u^2 + 49}{6u} \ln(1 + 4u^2) - \frac{u(13 + 60u^2)}{6(1 + 4u^2)^2} - \frac{128u^2}{3} \arctan 2u \right\}, \tag{6}
$$

and the corresponding Fock exchange-term reads:

$$
A_4(k_f)^{(\text{F2})} = \frac{g_A^4 M m_\pi^4}{(12\pi)^3 f_\pi^4} \left\{ \frac{u}{8} - \frac{u^3}{3} - \frac{u}{12(1+2u^2)} - \frac{u}{24(1+u^2)} \right. \\
\left. + u^4 \arctan u + \frac{u^2 (2+11u^2+16u^4)}{6(1+2u^2)^2} \left[\arctan u - \arctan 2u \right] \\
+ \int_0^u dx \frac{21x^2 - 16u^2}{6u(1+2x^2)} \left[(1+8x^2+8x^4) \arctan x - (1+4x^2) \arctan 2x \right] \right\}.
$$
 (7)

Pauli-blocking effects at second order are included through diagrams with three (isospin-asymmetric) medium insertions [\[6\]](#page-5-5). We consider here only the factorizable Fock contribution for which the energy denominator gets canceled by factors from the momentum-dependent πN -vertices (see eqs.(11,26) in ref.[\[6\]](#page-5-5)). Its contribution to the quartic isospin asymmetry energy can be represented as a oneparameter integral, $A_4(k_f)^{(\text{fac})} = g_A^4 M m_{\pi}^4 (12 \pi f_{\pi})^{-4} \int_0^u dx I(x, u)$, where the lengthy integrand $I(x, u)$ involves the function $\ln[1 + (u+x)^2] - \ln[1 + (u-x)^2]$ and its square. The corresponding value at saturation density is $A_4(k_{f0})^{\text{(fac)}} = -1.35 \text{ MeV}$, thus counterbalancing most of the Fock term $A_4(k_{f0})^{(F2)} = 1.70 \text{ MeV}$ without Pauli-blocking written in eq.(7). For the non-factorizable pieces the representation of the fourth derivative with respect to δ at $\delta = 0$ includes singularities of the form $(u-x)^{-\nu}, \nu = 1, 2$. When subtracting these singular terms from the integrand only very small numerical values are obtained for the non-factorizable Hartree contribution. In the case of the quadratic isospin asymmetry energy $A_2(k_{f0})$ one finds that the non-factorizable pieces (see eqs.(24,26) in ref.[\[6\]](#page-5-5)) tend to cancel each other almost completely as: (−11.6+12.0) MeV. Therefore one can expect that the omission of the non-factorizable pieces does not change much the final result for the quartic isospin asymmetry energy $A_4(k_f)$. However, the observation that the iterated 1 π exchange has components with a singular representation of their fourth derivative with respect to δ at $\delta = 0$, indicates that the expansion in eq.(1) becomes non-analytic beyond the quadratic order δ^2 . This feature is demonstrated in section 3 by calculating in closed form the second-order contribution from a s-wave contact-interaction.

We continue with the contribution of the irreducible 2π -exchange to the quartic isospin asymmetry energy. Using a twice-subtracted dispersion relation for the 2π -exchange NN-potential in momentum-space and the master formulas in eqs. $(3,4)$, one obtains:

$$
A_4(k_f)^{(2\pi)} = \frac{1}{81\pi^3} \int_{2m\pi}^{\infty} d\mu \left\{ Im(V_C + 2\mu^2 V_T) \left[\frac{7\mu k_f}{4} - \frac{2k_f^5}{3\mu^3} - \frac{\mu k_f^3 (7\mu^2 + 36k_f^2)}{2(\mu^2 + 4k_f^2)^2} \right] \right\}
$$

Figure 1: Quartic isospin asymmetry $A_4(k_f)$ as a function of the nucleon density $\rho = 2k_f^3/3\pi^2$.

$$
-\frac{7\mu^3}{16k_f} \ln\left(1+\frac{4k_f^2}{\mu^2}\right) + \text{Im}(W_C + 2\mu^2 W_T) \left[\frac{2k_f^5}{\mu^3} + \frac{k_f^3}{\mu} + \frac{21\mu k_f}{4} - \frac{\mu k_f^3 (7\mu^2 + 36k_f^2)}{2(\mu^2 + 4k_f^2)^2} - \frac{\mu}{16k_f} (21\mu^2 + 32k_f^2) \ln\left(1 + \frac{4k_f^2}{\mu^2}\right) \right],
$$
 (8)

where $\text{Im}V_{C,T}$ and $\text{Im}W_{C,T}$ are the spectral functions of the isoscalar and isovector central and tensor NN-amplitudes, respectively. These imaginary parts are composed of the functions $\sqrt{\mu^2 - 4m_\pi^2}$ and $arctan(\sqrt{\mu^2-4m_{\pi}^2}/2\Delta)$, with $\Delta = 293 \text{ MeV}$ the delta-nucleon mass splitting. Note that due to the implemented subtractions the k_f -expansion of $A_4(k_f)^{(2\pi)}$ in eq.(8) starts with the power k_f^7 . A short-distance contribution proportional to k_f^5 is supplemented by the subtraction constants:

$$
A_4(k_f)^{(\text{sc})} = \frac{10k_f^5}{(3M)^4} \left(\frac{2B_5}{3} - B_{n,5}\right),\tag{9}
$$

with the parameters $B_5 = 0$ and $B_{n,5} = -3.58$ adjusted in ref.[\[7\]](#page-5-6) to the empirical nuclear matter saturation point and quadratic isospin asymmetry energy $A_2(k_{f0})^{\text{(emp)}} = 34 \text{ MeV}.$

Finally, we consider the long-range three-nucleon interaction generated by 2π -exchange and virtual excitation of a $\Delta(1232)$ -isobar [\[7\]](#page-5-6). The corresponding three-body Hartree contribution reads:

$$
A_4(k_f)^{(\Delta)} = \frac{g_A^4 m_\pi^6 u^2}{\Delta (6\pi f_\pi)^4} \left\{ \left(\frac{16u^2}{3} + \frac{21}{4} \right) \ln(1 + 4u^2) - \frac{4u^4}{3} - \frac{41u^2}{3} - \frac{2u^2(11 + 99u^2 + 236u^4)}{3(1 + 4u^2)^3} \right\}, \tag{10}
$$

while the associated three-body Fock term can be represented as $g_A^4 m_\pi^6 (12\pi f_\pi)^{-4} \Delta^{-1} \int_0^u dx J(x, u)$, where the lengthy integrand $J(x, u)$ involves the functions $\arctan(u + x) + \arctan(u - x)$ and $\ln(1 +$ $(u+x)^2$ – ln[1 + $(u-x)^2$]. Note that the three-body contact-term proportional to ζ introduced additionally in ref. [\[7\]](#page-5-6) does not contribute to the quartic isospin asymmetry energy $A_4(k_f)$.

Summing up all the calculated contributions, one obtains the result for the density-dependent quartic isospin asymmetry energy $A_4(k_f)$ of nuclear matter as shown in Fig. 1 in the density region $0 < \rho < 2\rho_0 = 0.32 \,\text{fm}^{-3}$. The predicted value at saturation density $\rho_0 = 0.16 \,\text{fm}^{-3}$ is $A_4(k_{f0}) =$ 1.49 MeV. It amounts to 3.2 times the free Fermi-gas part $A_4(k_{f0})^{(kin)} = 0.464 \text{ MeV}$. Note that

interaction contributions to $A_4(k_f)$ start (at least) with the power k_f^5 . The density-dependence of the full line in Fig. 1 is to a good approximation $\rho^{5/4}$. Actually, it should be noted that the present calculation of the quartic isospin asymmetry energy $A_4(k_f)$ is performed in a framework where empirical constraints from bulk properties of nuclear matter are satisfied. This does not apply to the recent work in ref. [\[4\]](#page-5-3), where a large value of $A_4(k_{f0}) = (7.2 \pm 2.5)$ MeV has been found.

3 S-wave contact interaction to second order

The analysis of the Pauli-blocking corrections to the second-order (iterated) 1π -exchange has indicated that non-analytical terms may occur in the δ -expansion of the energy per particle of isospinasymmetric nuclear matter beyond the quadratic order. In the extreme case there could be a cubic term $|\delta|^3$, which is after all even under the exchange of protons and neutrons: $\delta \to -\delta$. In order to clarify the situation, we consider a s-wave contact interaction:

$$
V_{\rm ct} = \frac{\pi}{M} \Big[a_s + 3a_t + (a_t - a_s) \vec{\sigma}_1 \cdot \vec{\sigma}_2 \Big], \qquad (11)
$$

and examine it in second-order many-body perturbation theory. For this simple interaction, the occurring integrals over three Fermi spheres with (at most) two different radii k_p, k_n can be solved in closed analytical form. The pertinent function to express the result in the isospin-asymmetric configuration of interest is:

$$
35 \int_0^1 dz \left(z - z^4\right) \left\{ 2xz + \left(x^2 - z^2\right) \ln \frac{x + z}{|x - z|} \right\}
$$

= $\frac{x}{2} (15 + 33x^2 - 4x^4) + \frac{1}{4} (42x^2 - 15 - 35x^4) \ln \frac{x + 1}{|x - 1|} + 2x^7 \ln \frac{x^2}{|x^2 - 1|},$ (12)

where the variable $x > 0$ is set to a ratio of Fermi momenta, $[(1+\delta)/(1-\delta)]^{\pm 1/3}$ or 1. Note that the function defined in eq.(12) has at $x = 1$ the value $22 - 4 \ln 2$. Combining the second-order Hartree and Fock diagrams generated by V_{ct} according to their spin- and isospin-factors and performing the expansion in powers of δ , one obtains the following result for the energy per particle:

$$
\bar{E}_{\rm as}(k_p, k_n)^{(\rm 2nd)} = \frac{k_f^4}{5\pi^2 M} \left\{ \frac{3}{7} (a_s^2 + a_t^2)(11 - 2\ln 2) + \frac{4\delta^2}{3} \left[a_s^2 (3 - \ln 2) - a_t^2 (2 + \ln 2) \right] + \frac{\delta^4}{81} \left[a_s^2 \left(10 \ln \frac{|\delta|}{3} + 2 \ln 2 - \frac{41}{6} \right) + a_t^2 \left(30 \ln \frac{|\delta|}{3} + 2 \ln 2 + \frac{3}{2} \right) \right] + \mathcal{O}(\delta^6) \right\} . (13)
$$

The crucial and novel feature which becomes evident from this expression is the presence of the nonanalytical logarithmic term $\delta^4 \ln(|\delta|/3)$. Interestingly, the corresponding coefficient is three times as large in the spin-triplet channel as in the spin-singlet channel. For comparison the first-order contribution of the s-wave contact interaction V_{ct} reads, $\bar{E}_{\text{as}}(k_p, k_n)^{(\text{1st})} = k_f^3[-a_s - a_t + \delta^2(a_t$ $a_s/3$) $/2\pi M$, without any higher powers of δ . Note that the sign-convention for the scattering lengths $a_{s,t}$ is chosen here such that positive values correspond to attraction. As a check we have rederived the same results at first and second order by using the alternative (and equivalent) form of the s-wave contact interaction $V'_{\rm ct} = \pi [3a_s + a_t + (a_s - a_t)\vec{\tau}_1 \cdot \vec{\tau}_2]/M$.

In Fig. 2 the dependence of the second-order energy per particle $\bar{E}_{\text{as}}(k_p, k_n)^{(\text{2nd})}$ on the isospin asymmetry parameter δ is shown for three different choices of the s-wave scattering lengths: $a_s = a_t$, $a_t = 0$ and $a_s = 0$. In each case the full line shows the exact result and the (nearby) dashed line gives the expansion in powers of δ truncated at fourth order according to eq.(13). One observes that these expansions reproduce the full δ-dependence very well over the whole range $-1 \le \delta \le 1$. Note also that the prefactor $k_f^4 a_{s,t}^2/5\pi^2 M$ of dimension energy has been scaled out in Fig. 2.

Figure 2: Dependence of the 2nd order energy per particle $\bar{E}_{\text{as}}(k_p, k_n)^{(\text{2nd})}$ on the isospin asymmetry δ. Three different choices for the scattering lengths, $a_s = a_t$, $a_t = 0$ and $a_s = 0$, are considered.

If one performs for the second-order energy density the fourth derivative with respect to δ at $\delta = 0$ under the integral, then one encounters integrands with singularities of the form $(1-z)^{-\nu}$, $\nu = 1, 2$. The origin of these singularities, or in the proper treatment the non-analytical term $\delta^4 \ln(|\delta|/3)$, lies in the energy denominator of second-order diagrams. For an infinite (normal) many-fermion system the energy spectrum has a vanishing gap between bound states in the Fermi sea and excited states in the continuum. Such a gap-less energy spectrum causes a singularity, respectively a non-analyticity, if small asymmetries of the Fermi levels of two components are analyzed with too high resolution.

In summary, we have demonstrated that the non-analytical term $\delta^4 \ln(|\delta|/3)$ will be generically present in calculations of isospin-asymmetric nuclear matter that go beyond the mean-field Hartree-Fock level. Therefore, a term $\delta^4 \ln |\delta|$ should be included in future fits of the equation of state of (zero-temperature) isospin-asymmetric nuclear matter and its role should be further examined.

References

- [1] "Topical issue on nuclear symmetry energy", eds., B.A. Li, A. Ramos, G. Verde and I. Vidana, Eur. Phys. J. A50, No. 2 (2014).
- [2] I. Bombaci and U. Lombardo, Phys. Rev. C44, 1892 (1991).
- [3] O. Sjöberg, *Nucl. Phys.* **A222**, 161 (1974); A.W. Steiner, *Phys. Rev.* **C74**, 0452808 (2006); C. Ducoin, J. Margueron and P. Chomaz, *Nucl. Phys.* **A809**, 30 (2008); B.J. Cai and L.W. Chen, Phys. Rev. C85, 024302 (2012); D.N. Basu, Phys. Rev. C89, 028801 (2014).
- [4] Bao-Jun Cai and Bao-An Li, arXiv, nucl-th/1503.01167.
- [5] C. Drischler, master thesis, TU-Darmstadt, 2014.
- [6] N. Kaiser, S. Fritsch and W. Weise, *Nucl. Phys.* **A697**, 255 (2002).
- [7] S. Fritsch, N. Kaiser and W. Weise, Nucl. Phys. A750, 259 (2005).
- [8] N. Kaiser, *Nucl. Phys.* **A768**, 99 (2006).
- [9] S. Fiorilla, N. Kaiser and W. Weise, Nucl. Phys. A880, 65 (2012).
- [10] S. Fiorilla, N. Kaiser and W. Weise, Phys. Lett. B714, 251 (2012).
- [11] J.W. Holt, N. Kaiser and W. Weise, Prog. Part. Nucl. Phys. 73, 35 (2013).