

# Rapid host switching in generalist *Campylobacter* strains erodes the signal for tracing human infections

Bethany L. Dearlove<sup>1§</sup>, Alison J. Cody<sup>2</sup>, Ben Pascoe<sup>3,4</sup>, Guillaume Méric<sup>3,4</sup>, Daniel J. Wilson,<sup>1,5\*</sup>, Samuel K. Sheppard<sup>2,3,4\*</sup>

<sup>1</sup>Nuffield Department of Medicine, Experimental Medicine Division, University of Oxford, Oxford, UK

<sup>2</sup>Department of Zoology, University of Oxford, Oxford, UK

<sup>3</sup>College of Medicine, Institute of Life Science, Swansea University, Swansea, UK

<sup>4</sup>MRC CLIMB Consortium, Institute of Life Science, Swansea University, Swansea, UK

<sup>5</sup>Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK

\*These authors contributed equally to this work.

§Corresponding author. Present address: Department of Veterinary Medicine, University of Cambridge, Cambridge, United Kingdom. Email: [bd357@cam.ac.uk](mailto:bd357@cam.ac.uk)

**Running Title:** Rapid zoonosis in generalist *Campylobacter* strains

**Subject Category:** Evolutionary genetics

## Abstract

*Campylobacter jejuni* and *Campylobacter coli* are the biggest causes of bacterial gastroenteritis in the developed world, with human infections typically arising from zoonotic transmission associated with infected meat, especially poultry. Because this organism is not thought to survive well outside of the gut, host associated populations are genetically isolated to varying degrees. Therefore the likely origin of most *Campylobacter* strains can be determined by host-associated variation in the genome. This is instructive for characterizing the source of human infection at the population level. However, some common strains appear to have broad host ranges, hindering source attribution. Whole genome sequencing has the potential to reveal fine-scale genetic structure associated with host specificity within each of these strains.

We found that rates of zoonotic transmission among animal host species in ST-21, ST-45 and ST-828 clonal complexes were so high that the signal of host association is all but obliterated. We attributed 89% of clinical cases to a chicken source, 10% to cattle and 1% to pig. Our results reveal that common strains of *C. jejuni* and *C. coli* infectious to humans are adapted to a generalist lifestyle, permitting rapid transmission between different hosts. Furthermore, they show that the weak signal of host association within these complexes presents a challenge for pinpointing the source of clinical infections, underlining the view that whole genome sequencing, powerful though it is, cannot substitute for intensive sampling of suspected transmission reservoirs.

**Keywords:** attribution / *Campylobacter* / multilocus sequence typing / transmission / zoonosis

## Introduction

*Campylobacter jejuni* and *Campylobacter coli* are zoonotic pathogens with broad host ranges, carried, apparently asymptotically, as part of the gut microbiota of a range of wild and domesticated mammal and bird species. Common carriage in the gut of animals and poultry farmed for meat leads to numerous opportunities for contamination of food products, which may take place at various points from farm to fork (Neimann *et al.*, 2003). Human-to-human transmission is rare (Allos, 2001), hence *Campylobacter* infection in humans tends to be sporadic, and seldom manifests as outbreaks except when a single point source results in direct transmission to many people, for example via contaminated drinking water (Palmer *et al.*, 1983; Pebody *et al.*, 1997; Engberg *et al.*, 1998; Clark *et al.*, 2003).

The sources of human *Campylobacter* infection have been well characterized at the population level. Genetic analysis, particularly using seven-locus multilocus sequence typing (MLST) data, has helped to attribute the sources of clinical infections by exploiting differences in the frequency of *Campylobacter* strains that live in different animal and environmental reservoirs (McCarthy *et al.*, 2007). For example, isolates belonging to related sequence types (STs) from the ST-257 and ST-61 clonal complexes are strongly associated with chicken and ruminants respectively, while wild bird species are often colonised by phylogenetically divergent *Campylobacter* lineages (Sheppard *et al.*, 2011; Griekspoor *et al.*, 2013). Independent MLST-based studies in England, Scotland and New Zealand found that over 95% of human infections are attributable to animals farmed for meat and poultry, with 56-76% of cases attributable to poultry in particular (Wilson *et al.*, 2008; Mullner *et al.*, 2009; Sheppard *et al.*, 2009).

However, at the individual isolate level, there is often considerable uncertainty in source attribution because some of the most common disease causing strains, notably those belonging to the ST-21, ST-45 and ST-828 clonal complexes, are regularly isolated from multiple animal species. This means that quantitative attribution to a single host reservoir is difficult using MLST data alone. The reason for the apparently broader host range of these lineages is not currently understood, and may reflect true host generalism (Gripp *et al.*, 2011; Sheppard *et al.*, 2014) or the existence of host restricted sublineages within these clonal complexes.

The increasing availability of whole genome sequence data provides a potential solution to the challenge of attributing the origin of generalist strains in clinical samples, as a host-specific signal might be gleaned from the remaining 99.8% of the genome outside the seven loci sequenced by standard MLST. Genomics has been used to investigate direct transmission events in a range of bacterial species including *Staphylococcus aureus* (Harris *et al.*, 2010; Price *et al.*, 2014), *Clostridium difficile* (Eyre *et al.*, 2013) and *Mycobacterium tuberculosis* (Walker *et al.*, 2013). This raises the prospect of enhancing understanding of *Campylobacter* epidemiology, for example by identifying individual retailers or producers of *Campylobacter*-contaminated food in cases of human infection, and detecting cryptic point source outbreaks (Cody *et al.*, 2013). Ultimately this information can help to inform targeted interventions and reduce the incidence of human campylobacteriosis (Sears *et al.*, 2011).

In this study we investigated the utility of whole genome sequence data for improving the accuracy with which human cases can be attributed to particular animal reservoirs, by focusing on common strains that are difficult to attribute with MLST. Specifically, we sought to detect fine-scale host-specific genetic structuring within *C. jejuni* ST-21 and ST-45

complex and *C. coli* ST-828 complex isolates sampled from different animal species and to exploit that signal to improve human source attribution. We observed extraordinarily rapid rates of zoonotic transfer within these strains leading to little or no phylogenetic association with host species, limiting the ability of whole genome sequencing to improve source attribution, but revealing new insight into the transmission dynamics within these common *Campylobacter* strains.

## Materials and Methods

### Sequences

The ST-21 and ST-45 (*C. jejuni*) and ST-828 (*C. coli*) clonal complexes were chosen as the focus of the study because they are among the most common lineages causing human disease, but are difficult to attribute to source populations using seven-locus MLST data (Wilson *et al.*, 2008; Sheppard *et al.*, 2009). Isolates from these three clonal complexes were chosen from MLST collections ([www.pubmlst.org/campylobacter](http://www.pubmlst.org/campylobacter), Jolley and Maiden (2010)), sequenced and assembled according to the protocols described in Sheppard et al. (2013) and Cody et al. (2013). In total, 348, 87 and 158 sequences were obtained for the ST-21, ST-45 and ST-828 complexes, respectively (Lefébure *et al.*, 2010; Cody *et al.*, 2013; Sheppard, Didelot, Jolley, *et al.*, 2013; Sheppard, Didelot, Meric, *et al.*, 2013; Sheppard *et al.*, 2014) (see Supplementary Table 1). Isolates belonging these complexes are predominantly associated with chicken and cattle, from where the majority of animal samples were obtained (13 chicken isolates from each of clonal complexes, and 7, 9 and 10 cattle isolates for the ST-21, ST-45 and ST-828 clonal complexes respectively). In addition, three ST-45 complex isolates were included from wild bird species, and six ST-828 complex isolates from pigs. Clinical disease isolates (328, 68 and 128 for the ST-21, ST-45 and ST-828 clonal complexes

respectively) were obtained from a surveillance study of samples submitted to the microbiology laboratory of the John Radcliffe Hospital, Oxfordshire, United Kingdom (UK).

After Illumina sequencing, the high coverage short reads were *de novo* assembled using Velvet (Zerbino and Birney, 2008), and the resulting contiguous sequences ('contigs') stored using BIGSdb (Jolley and Maiden, 2010). These contigs were then compared to the NCTC11168 reference sequence (Genbank accession number: AL111168) to identify genes using a BLAST search (Parkhill *et al.*, 2000; Gundogdu *et al.*, 2007). Orthologous genes were defined as homologous genes that had 70% or greater nucleotide identity, and less than 50% difference in alignment length. Genes for all isolates were then aligned using MUSCLE (Edgar, 2004), and concatenated into a single sequence per sample.

### **Accounting for ancestral recombination**

There is good evidence that novel diversity in *Campylobacter* is generated frequently by the continued movement of genes between lineages, more so even than by the evolution of new variants through mutation (Wilson *et al.*, 2009; Sheppard *et al.*, 2010). High levels of recombination lead to mosaic genomes with differing ancestral histories, and the full evolution of samples cannot be represented by a coalescent genealogy.

During preliminary analyses using BEAST (Drummond *et al.*, 2012), a relaxed clock model and gamma site heterogeneity were used to try and account for the effect of recombination. However, these analyses diagnosed difficulties in the mixing of the MCMC algorithm underlying BEAST, with multiple runs of the same analysis frequently converging to different topologies. To overcome this, and to account for the effect that recombination has in skewing the branch lengths of the dominant phylogenetic tree (Schierup and Hein, 2000),

homoplasious sites incompatible with the maximum likelihood tree were identified and removed (Pupko *et al.*, 2000; Guindon *et al.*, 2010). For the final dataset, only the biallelic sites compatible with the inferred phylogeny were included in the alignment, alongside all of the non-variable sites. The removal of homoplasies in this way resolved the mixing issues with BEAST. In addition, to improve computation time for later analyses, missing alleles were imputed using ClonalFrameML (Pupko *et al.*, 2000; Didelot and Wilson, 2015).

## Phylogenetic analysis

The analysis was implemented in BEAST v1.7.5 (Drummond *et al.*, 2012), assuming a constant population size. All parameters were scaled in terms of the effective population size,  $N_e$ , by fixing it to 1.0. The HKY85 model of nucleotide substitution (Hasegawa *et al.*, 1985) was used with an uncorrelated log-normal relaxed clock and gamma rate heterogeneity with four categories (Drummond *et al.*, 2006). A log-normal prior with a mean of 1.0 and standard deviation of 1.25 on the logarithmic-scale was assumed for the transition:tranversion ratio  $\kappa$ , and an exponential prior with a mean of 1.0 was utilized for the gamma shape parameter  $\alpha$ . For the log-normal relaxed clock parameters, a uniform prior between 0.0 and 10.0 was assumed for the mean, and an exponential with mean 1.0 for the standard deviation. A uniform (Dirichlet) prior was used for the nucleotide frequencies.

We used the phylogeographic model by Lemey *et al.* (Lemey *et al.*, 2009) to reconstruct zoonotic transmission and infer the source of the infection (host species) along the branches. This allows us to report the posterior probability of any node in the tree being a particular state (i.e. the most likely host of the ancestral lineage), and also to obtain relative rates of transition between states (i.e. switching between host species). Given that human to human

transmission is rare (Allos, 2001), an ambiguity code was set up in the zoonotic model to allow the human isolates to have an ‘unknown’ source population. For each human isolate, the other host species in the analysis were given equal prior probability and thus most likely source of the human isolates could be inferred. A uniform prior from 0.0 to 100.0 was assumed for the host migration rate, a gamma prior with mean 1.0 and scale 1.0 for the relative rates of migration, and a uniform (Dirichlet) prior for the host population equilibrium frequencies. To calculate the number of discrete zoonotic transmissions across the branches, Markov jumping was performed using BEAST (Minin and Suchard, 2008a, 2008b; Talbi *et al.*, 2010).

The MCMC was run for 500 million iterations, with samples taken every 10,000 iterations. For each ST, the analysis was repeated twice (three times for ST-21) with different initial values to check convergence and mixing, and these runs were combined for the final results. Unless otherwise stated, the posterior median was used for point estimates and the (2.5%, 97.5%) quantiles for credible intervals. The inferred ancestral host type for each branch in the phylogeny was taken to be the one with the highest posterior probability. Sample dates were not available for all isolates, so the analysis is given in coalescent time (denoted  $\tau$ ) and calibrated to years using an independent estimate of the mutation rate in *Campylobacter* of  $3.23 \times 10^{-5}$  substitutions per site per year from Wilson *et al.* (2009).

## Results

### Fine-scale Phylogenetic Structure within Sequence Types

Within the ST-21, ST-45 and ST-828 complexes, isolates from different host species were often more closely related than those isolated from the same host species (Figure 1). If the strains harboured previously undetected sub-ST lineages that were strongly host associated,

one would expect to see distinct clusters of the same coloured branches together. However, this was not the case, with branches of the same colour – representing the reconstructed reservoir population of that lineage – scattered throughout the tree in all STs. Furthermore, one could expect that isolates from mammalian hosts would be more closely related than those from avian species reflecting enhanced transfer potential between physiologically similar hosts. Again, this was not observed, with bird isolates in the ST-45 complex and pig isolates in ST-828 complex being closely related to both chicken and cattle isolates. This may suggest that the ability to colonise a specific host has either evolved several times throughout the tree, most plausibly through horizontal gene transfer given that mutation is rare, or that the isolates are adapted to infect all species of host in the sample.

Much of the ancestral history of the lineages was inferred to have occurred within the chicken host population, shown by the dominance of yellow branches. The most recent common ancestor (MRCA) of all three clonal complexes was inferred to have colonized chicken, with posterior probabilities of 0.612, 0.498 and 0.500 for the ST-21, ST-45 and ST-828 complexes respectively.

### **Rates of Zoonotic Transmission in *Campylobacter***

Under the host restricted hypothesis, isolates sampled from the same source reservoir will cluster into a single clade within the phylogeny. Since we sampled ST-21 complex isolates from two host species (chicken and cattle), there must have been exactly one zoonotic transfer in the tree under the restricted host hypothesis, or more under the generalist model. We sampled ST-45 and ST-828 complex isolates from three host species (chicken, cattle and wild birds in ST-45 or swine in ST-828), necessitating exactly two zoonotic transfer events in the tree under the restricted host hypothesis. In fact, the total number of migration events for all

three clonal complexes was estimated to be much higher than these minimum values, with 588.9 (95% credibility interval: 109.8, 1325.9), 468.7 (105.7, 1264.5) and 117.7 (36.6, 456.3) cross-species transmission events for ST-21, ST-45 and ST-828 complexes respectively. In all three cases, the number of zoonotic transfers under the restricted host hypothesis was outside the range of the 95% credible intervals, demonstrating that isolates from these clonal complexes display host generalism and excluding the possibility of host restricted sub-lineages.

The overall estimated migration rate was much more frequent in the *C. jejuni* clonal complexes compared to the *C. coli* (ST-828) complex. Using a mutation rate of  $3.23 \times 10^{-5}$  substitutions per site per year for calibration (Wilson *et al.*, 2009), this corresponds to approximately one host jump every twelve years for ST-828 compared to one every 1.6 or 1.8 years in the ST-21 and ST-45 complexes. Although the credible intervals are wide, particularly for the *C. jejuni* sequence types, the finding of extraordinarily rapid rates of zoonotic transfer were robust to alternative prior beliefs. Thus, based upon the isolate collection in this study, the estimated rate of zoonosis is sufficiently rapid within ST-21 and ST-45 clonal complexes that there is no association between genetic structure and host species. This is consistent with a generalist lifestyle in which the isolates are equally adapted to transmission between versus within species.

### Tracing the Source of Human Infection

In Figure 1, the human cases at the tips of the tree are represented with black circles, and the posterior probability of the sources for the human isolates are illustrated by the bar plots. The majority of clinical cases (462 out of 519 cases, or 89%) were attributed to a chicken source, with 53 cases (10%) attributed to cattle and 4 (1%) to pig (Figure 2). This is consistent with

results found by MLST (Wilson *et al.*, 2008; Mullner *et al.*, 2009; Sheppard *et al.*, 2009), but the difficulty of attributing individual human cases to specific source species with high confidence, even using whole genomes, reflects the remarkable transmission rates of lineages among host species. Even when human isolates were most closely related to clades sampled only from chicken, there remained appreciable probabilities (in the range 30-40%) that the direct source of transmission was non-chicken. This degree of uncertainty is comparable to the 67% accuracy of source attribution that Wilson *et al.* (2008) achieve with MLST alone.

## Discussion

We investigated the source of *Campylobacter* infection in humans using whole genome sequencing, focusing on STs that are frequently isolated from multiple host species and are therefore weakly host-associated on the basis of MLST (Wilson *et al.*, 2008; Sheppard *et al.*, 2009, 2014; Gripp *et al.*, 2011). We tested whether these STs were aggregations of strongly host-restricted sub-lineages, or whether they represented genuine generalists (Gripp *et al.*, 2011). We estimated rates of zoonotic transfer between *Campylobacter* reservoir species, and attributed individual clinical cases to animal sources. Consistent with previous studies (Harris *et al.*, 1986; Wingstrand *et al.*, 2006; Wilson *et al.*, 2008; Mullner *et al.*, 2009; Sheppard *et al.*, 2009), we identified the chicken reservoir as accounting for the majority of human *Campylobacter* infections, emphasizing the importance of measures aimed at controlling food-borne disease in agriculture and the food industry.

We found that fine-scale population structure across the genome within the ST-21, ST-45 and ST-828 clonal complex isolates was not host-associated. This is consistent with the existence of genuine generalist strains, adapted to transmit between and live within multiple host species. There are clear advantages of a generalist lifestyle in agricultural animal species

where multiple mammalian and avian species routinely live in close proximity, giving rise to frequent opportunity for zoonotic transmission.

Some differences were observed in the relative rate of host switching between *C. coli* and *C. jejuni*. Specifically, for the *C. coli* ST-828 complex isolates, there was evidence for slower rates of zoonotic transfer, compared to the *C. jejuni* ST-21 and ST-45 complex isolates. We estimated that there were 588 migration events across the tree in the ST-21 complex and 468 in the ST-45 complex, compared to only 117 in ST-828. In all cases we were able to reject the hypothesis of a unique host jump founding the population in each new species, with significantly more migration events than would be expected if isolates were host restricted (Table 2). This is evidenced by the scattering of isolates sampled from different sources throughout the phylogeny.

The overall rates of migration between different host species provides information about the ecology of the generalist *Campylobacter* strains. Within the ST-45 complex, there was very little difference in the relative rates of transmission between host species (Table 2). However, in ST-828 complex isolates, host switches were twice as frequent between cattle and swine compared with the rates of migration between these species and chicken. This disparity between mammal-mammal and mammal-bird transmission rates suggests that the efficacy of zoonotic transmission may be lower in the ST-828 complex compared to ST-21 or ST-45 complexes. Mammals and birds exhibit many physiological differences including a difference in core body temperature of about 38°C in cattle and pigs versus 42°C in chickens. Variation in rates of zoonosis may be explained by differences in the route of transmission between the mammalian species, either through opportunity – for example if cattle and pigs have more commonly shared the same physical environment, or through affinity – involving factors

associated with the ability to colonize different species. Suggestively, the average genome size of ST-828 complex isolates is smaller than that of ST-21 and ST-45 complex isolates, potentially reflecting more limited phenotypic plasticity that would limit the ability to occupy multiple divergent niches.

By comparing an approach using relatively few isolates characterized at many loci, by whole genome sequencing, with methods using hundreds or thousands of isolates characterized at few loci, such as MLST studies (Wilson *et al.*, 2008; Mullner *et al.*, 2009; Sheppard *et al.*, 2009), we investigated the added benefit of whole genome sequencing for investigating lineages with rapid host-switching within clonal complexes. There was little additional information regarding source of infection provided by the whole genome sequence in the majority of human cases. This suggests an inherent trade-off between the number of samples and the number of loci sequenced, and that the approximately thousand-fold increase in sequence information afforded by whole genomes over 7-locus MLST does not, on its own, eliminate the need for detailed sampling. While there is evidence of the potential for whole genome sequencing to provide greater resolution in detecting recent transmission of human *Campylobacter* infection from non-chicken sources, this study highlights the importance of intensive sampling of potential source populations and an understanding of host transmission ecology for effective source attribution.

## Acknowledgements

This study was supported by the Oxford NIHR Biomedical Research Centre and the UKCRC Modernising Medical Microbiology Consortium, the latter funded under the UKCRC Translational Infection Research Initiative supported by the Medical Research Council, the Biotechnology and Biological Sciences Research Council and the National Institute for

Health Research on behalf of the UK Department of Health (Grant G0800778) and the Wellcome Trust (Grant 087646/Z/08/Z). BLD is supported by a Medical Research Council Methodology Research Programme grant (grant number MR/J013862/1). AJC was supported by the United Kingdom Department for Environment, Food, and Rural Affairs and Food Standards Agency (grant number OZ0624). DJW is a Sir Henry Dale Fellow, jointly funded by the Wellcome Trust and the Royal Society (Grant 101237/Z/13/Z). SKS is funded by the Biotechnology and Biological Sciences Research Council (BBSRC), the Medical Research Council (MR/L015080/1) and the Wellcome Trust. This publication made use of the *Campylobacter* Multi Locus Sequence Typing website (<http://pubmlst.org/campylobacter/>) developed by Keith Jolley and sited at the University of Oxford (24). The development of this site has been funded by the Wellcome Trust.

Supplementary information is available at The ISME Journal's website.

## **Conflicts of Interest**

The authors have no conflicts of interest to declare.

## **References**

- Allos BM. (2001). *Campylobacter jejuni* infections: update on emerging issues and trends. *Clin Infect Dis* 32:1201–6.
- Clark CG, Price L, Ahmed R, Woodward DL, Melito PL, Rodgers FG, et al. (2003). Characterization of waterborne outbreak-associated *Campylobacter jejuni*, Walkerton, Ontario. *Emerg Infect Dis* 9:1232–1241.

- Cody AJ, McCarthy ND, Jansen van Rensburg M, Isinkaye T, Bentley SD, Parkhill J, et al. (2013). Real-time genomic epidemiological evaluation of human *Campylobacter* isolates by use of whole-genome multilocus sequence typing. *J Clin Microbiol* 51:2526–34.
- Didelot X, Wilson DJ. (2015). ClonalFrameML: Efficient Inference of Recombination in Whole Bacterial Genomes. *PLOS Comput Biol* 11:e1004041.
- Drummond AJ, Ho SYW, Phillips MJ, Rambaut A. (2006). Relaxed phylogenetics and dating with confidence. *PLoS Biol* 4:e88.
- Drummond AJ, Suchard MA, Xie D, Rambaut A. (2012). Bayesian phylogenetics with BEAUTi and the BEAST 1.7. *Mol Biol Evol* 29:1969–73.
- Edgar RC. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–7.
- Engberg J, Gerner-Smidt P, Scheutz F, Møller Nielson E, On SLW, Mølbak K. (1998). Water-borne *Campylobacter jejuni* infection in a Danish town - a 6-week continuous source outbreak. *Clin Microbiol Infect* 4:648–656.
- Eyre DW, Cule ML, Wilson DJ, Griffiths D, Vaughan A, O'Connor L, et al. (2013). Diverse sources of *C. difficile* infection identified on whole-genome sequencing. *N Engl J Med* 369:1195–205.
- Griekspoor P, Colles FM, McCarthy ND, Hansbro PM, Ashurst-Smith C, Olsen B, et al. (2013). Marked host specificity and lack of phylogeographic population structure of *Campylobacter jejuni* in wild birds. *Mol Ecol* 22:1463–72.
- Gripp E, Hlahla D, Didelot X, Kops F, Maurischat S, Tedin K, et al. (2011). Closely related *Campylobacter jejuni* strains from different sources reveal a generalist rather than a specialist lifestyle. *BMC Genomics* 12:584.

- Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 59:307–21.
- Gundogdu O, Bentley SD, Holden MT, Parkhill J, Dorrell N, Wren BW. (2007). Re-annotation and re-analysis of the *Campylobacter jejuni* NCTC11168 genome sequence. *BMC Genomics* 8. doi:10.1186/1471-2164-8-162.
- Harris N V, Weiss NS, Nolan CM. (1986). The role of poultry and meats in the etiology of *Campylobacter jejuni/coli* enteritis. *Am J Public Health* 76:407–11.
- Harris SR, Feil EJ, Holden MTG, Quail MA, Nickerson EK, Chantratita N, et al. (2010). Evolution of MRSA during hospital transmission and intercontinental spread. *Science* 327:469–74.
- Hasegawa M, Kishino H, Yano T. (1985). Dating of the Human-Ape Splitting by a Molecular Clock of Mitochondrial DNA. *J Mol Evol* 22:160–174.
- Jolley KA, Maiden MCJ. (2010). BIGSdb: Scalable analysis of bacterial genome variation at the population level. *BMC Bioinformatics* 11:595.
- Lefébure T, Pavinski Bitar PD, Suzuki H, Stanhope MJ. (2010). Evolutionary dynamics of complete *Campylobacter* pan-genomes and the bacterial species concept. *Genome Biol Evol* 2:646–55.
- Lemey P, Rambaut A, Drummond AJ, Suchard MA. (2009). Bayesian phylogeography finds its roots. *PLoS Comput Biol* 5:e1000520.
- McCarthy ND, Colles FM, Dingle KE, Bagnall MC, Manning G, Maiden MCJ, et al. (2007). Host-associated genetic import in *Campylobacter jejuni*. *Emerg Infect Dis* 13:267–72.
- Minin VN, Suchard MA. (2008a). Counting labeled transitions in continuous-time Markov models of evolution. *J Math Biol* 56:391–412.

- Minin VN, Suchard MA. (2008b). Fast, accurate and simulation-free stochastic mapping. *Phil Trans R Soc B* 363:3985–95.
- Muellner P, Spencer SEF, Wilson DJ, Jones G, Noble AD, Midwinter AC, et al. (2009). Assigning the source of human campylobacteriosis in New Zealand: a comparative genetic and epidemiological approach. *Infect Genet Evol* 9:1311–9.
- Neimann J, Engberg J, Mølbak K, Wegener HC. (2003). A case-control study of risk factors for sporadic campylobacter infections in Denmark. *Epidemiol Infect* 130:353–66.
- Palmer SR, Gully PR, White JM, Person AD, Suckling WG, Jones DM, et al. (1983). Waterborne outbreak of *Campylobacter* gastroenteritis. *Lancet* 321:287–290.
- Parkhill J, Wren BW, Mungall K, Ketley JM, Churcher C, Basham D, et al. (2000). The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hypervariable sequences. *Nature* 403:665–8.
- Pebody RG, Ryan MJ, Wall PG. (1997). Outbreaks of campylobacter infection: rare events for a common pathogen. *Commun Dis Rep Rev* 7:R33–7.
- Price JR, Golubchik T, Cole K, Wilson DJ, Crook DW, Thwaites GE, et al. (2014). Whole-genome sequencing shows that patient-to-patient transmission rarely accounts for acquisition of *Staphylococcus aureus* in an intensive care unit. *Clin Infect Dis* 58:609–18.
- Pupko T, Pe'er I, Shamir R, Graur D. (2000). A fast algorithm for joint reconstruction of ancestral amino acid sequences. *Mol Biol Evol* 17:890–6.
- Schierup MH, Hein J. (2000). Consequences of recombination on traditional phylogenetic analysis. *Genetics* 156:879–91.
- Sears A, Baker MG, Wilson N, Marshall J, Muellner P, Campbell DM, et al. (2011). Marked Campylobacteriosis decline after interventions aimed at poultry, New Zealand. *Emerg Infect Dis* 17:1007–1015.

- Sheppard SK, Cheng L, Méric G, de Haan CPA, Llarena A-K, Marttinen P, et al. (2014). Cryptic ecology among host generalist *Campylobacter jejuni* in domestic animals. *Mol Ecol* 23:2442–51.
- Sheppard SK, Colles FM, McCarthy ND, Strachan NJC, Ogden ID, Forbes KJ, et al. (2011). Niche segregation and genetic structure of *Campylobacter jejuni* populations from wild and agricultural host species. *Mol Ecol* 20:3484–90.
- Sheppard SK, Dallas JF, Strachan NJC, MacRae M, McCarthy ND, Wilson DJ, et al. (2009). *Campylobacter* genotyping to determine the source of human infection. *Clin Infect Dis* 48:1072–8.
- Sheppard SK, Didelot X, Jolley KA, Darling AE, Pascoe B, Meric G, et al. (2013). Progressive genome-wide introgression in agricultural *Campylobacter coli*. *Mol Ecol* 22:1051–64.
- Sheppard SK, Didelot X, Meric G, Torralbo A, Jolley KA, Kelly DJ, et al. (2013). Genome-wide association study identifies vitamin B5 biosynthesis as a host specificity factor in *Campylobacter*. *Proc Natl Acad Sci USA* 110:11923–11927.
- Sheppard SK, Maiden MCJ, Falush D. (2010). Population Genetics of *Campylobacter*. In: *Bacterial populations in infectious disease*, Robinson, DA, Falush, D, & Feil, EJ (eds), Wiley-Blackwell: Hoboken, NJ, USA, p. 184.
- Talbi C, Lemey P, Suchard MA, Abdelatif E, Elharrak M, Nourlil J, et al. (2010). Phylodynamics and human-mediated dispersal of a zoonotic virus. *PLoS Pathog* 6:e1001166.
- Walker TM, Ip CLC, Harrell RH, Evans JT, Kapatai G, Dedicoat MJ, et al. (2013). Whole-genome sequencing to delineate *Mycobacterium tuberculosis* outbreaks: a retrospective observational study. *Lancet Infect Dis* 13:137–46.

Wilson DJ, Gabriel E, Leatherbarrow AJH, Cheesbrough J, Gee S, Bolton E, et al. (2009).

Rapid evolution and the importance of recombination to the gastroenteric pathogen

*Campylobacter jejuni*. Mol Biol Evol 26:385–97.

Wilson DJ, Gabriel E, Leatherbarrow AJH, Cheesbrough J, Gee S, Bolton E, et al. (2008).

Tracing the source of campylobacteriosis. PLoS Genet 4:e1000203.


Wingstrand A, Neumann J, Engberg J, Møller Nielsen E, Gerner-Smidt P, Wegener HC, et al. (2006). Fresh chicken as main risk factor for campylobacteriosis, Denmark. Emerg Infect Dis 12:280–284.

Zerbino DR, Birney E. (2008). Velvet: algorithms for de novo short read assembly using de Bruijn graphs. Genome Res 18:821–9.

## Figures and Tables


### Figure 1: Maximum clade credibility trees for the ST-21, ST-45 ST-828 complexes.

Tips are coloured by host from which the sample was isolated: chicken (yellow), cattle (red), pig (pink), wild bird (green) and human (black). Branches are coloured according to the ancestral source population inferred using the maximum posterior probability. Pie charts show the posterior probability for the root of the tree. For each human case, the posterior probability of source is shown as a stacked bar plot. Scale is given in units of coalescent time. Note that a change in host may have occurred at any point on the branch, not necessarily at the node, and it is also possible to have a number of host switches occurring along a branch.



**Figure 2: Probability of source for clinical cases for a) ST-21, b) ST-45 and c) ST-828.**

The posterior probability of each human isolate (vertical bars) broken down by source population: chicken (yellow), cattle (red), pig (pink) and wild bird (green). The isolates have been reordered along the x-axis for visualisation purposes.



**Table 1: Summary of sequence data for the three STs.**

	ST-21 complex	ST-45 complex	ST-828 complex
<b>Genome alignment length:</b>	<b>1544595</b>	<b>1465323</b>	<b>1421603</b>
Non-polymorphic sites	1447536	1386349	1272113
Polymorphic sites	97059	78929	149490
<b>Biallelic sites:</b>	<b>90239</b>	<b>722233</b>	<b>136375</b>
Compatible with ML tree	41895	34960	60692
<b>Total sites used in analysis</b>	<b>1489431</b>	<b>1421309</b>	<b>1332805</b>

**Table 2: Parameter estimates for each ST.**

Parameter	ST-21 complex	ST-45 complex	ST-828 complex
<b>Substitution rate</b>			
( $10^{-3}$ /site / $\tau$ )	2.815 (1.950, 3.560)	3.715 (2.315, 5.158)	9.582 (8.019, 11.534)
<b>TMRCA</b>			
Coalescent time, $\tau$	1.022 (0.442, 3.068)	0.709 (0.288, 2.583)	0.159 (0.110, 0.232)
Years <sup>1</sup>	89.069 (38.521, 267.381)	81.545 (33.124, 297.085)	47.168 (32.632, 68.824)
<b>Host switching rate</b>			
/ $\tau$	54.078 (10.089, 97.519)	61.634 (14.540, 98.069)	23.491 (6.802, 85.357)
/year <sup>1</sup>	0.621 (0.116, 1.119)	0.536 (0.126, 0.853)	0.079 (0.023, 0.287)
<b>Number of migrations</b>			
	588.89 (109.80, 1325.90)	468.67 (105.70, 1264.48)	117.691 (36.558, 456.318)
<b>Estimated host frequencies</b>			
Chicken	0.614 (0.388, 0.809)	0.498 (0.311, 0.685)	0.425 (0.136, 0.691)
Cattle	0.386 (0.191, 0.612)	0.348 (0.185, 0.540)	0.282 (0.095, 0.562)
Wild bird	-	0.140 (0.0429, 0.313)	-
Pig	-	-	0.270 (0.103, 0.545)
<b>Relative migration rates between host species</b>			
Chicken-Cattle	0.695 (0.023, 3.705)	0.820 (0.051, 3.779)	0.303 (0.010, 2.139)
Chicken-Wild Bird	-	0.592 (0.021, 3.282)	-
Chicken-Pig	-	-	0.581 (0.024, 3.129)
Cattle-Wild Bird	-	0.728 (0.031, 3.643)	-
Cattle-Pig	-	-	1.367 (0.192, 4.783)

<sup>1</sup>Using the mutation rate  $3.23 \times 10^{-5}$  substitutions per site per year [21] for calibration with the median evolutionary rate from the BEAST analysis, one unit of coalescent time ( $\tau$ ) is equal to: 87.151 years in the ST-21 complex, 115.015 years in ST-45 complex, and 296.656 years in ST-828 complex. The median value was used for estimating all other parameters in years.

**Supplementary Table 1: Isolates included in the study.**

All isolates are available from <http://pubmlst.org/campylobacter/>.

\*Sequence type was derived from the allelic profile of seven housekeeping genes by multilocus sequence typing (MLST).

\*\*Clonal complex is defined as including any ST that matches a previously defined central genotype (<http://pubmlst.org/campylobacter/>) at three or more loci.

\*\*\*References refer to:

1. Sheppard SK, Didelot X, Jolley KA, Darling AE, Pascoe B, Meric G, Kelly DJ, Cody A, Colles FM, Strachan NJC, Ogden ID, Forbes K, French NP, Carter P, Miller WG, McCarthy ND, Owen R, Litrup E, Egholm M, Affourtit JP, Bentley SD, Parkhill J, Maiden MCJ, Falush D (2013) Progressive genome-wide introgression in agricultural *Campylobacter coli*. *Mol Ecol*, **22**:1051–64.
2. Sheppard SK, Didelot X, Meric G, Torralbo A, Jolley KA, Kelly DJ, Bentley SD, Maiden MCJ, Parkhill J, Falush D (2013) Genome-wide association study identifies vitamin B5 biosynthesis as a host specificity factor in *Campylobacter*. *Proc Natl Acad Sci USA*, **110**:11923–11927.
3. Sheppard SK, Cheng L, Méric G, de Haan CPA, Llarena A-K, Marttinen P, Vidal A, Ridley A, Clifton-Hadley F, Connor TR, Strachan NJC, Forbes K, Colles FM, Jolley KA, Bentley SD, Maiden MCJ, Hänninen M-L, Parkhill J, Hanage WP, Corander J (2014) Cryptic ecology among host generalist *Campylobacter jejuni* in domestic animals. *Mol Ecol*, **23**:2442–51.
4. Lefébure T, Pavinski Bitar PD, Suzuki H, Stanhope MJ (2010) Evolutionary dynamics of complete *Campylobacter* pan-genomes and the bacterial species concept. *Genome Biol Evol*, **2**:646–55.
5. Cody AJ, McCarthy ND, Jansen van Rensburg M, Isinkaye T, Bentley SD, Parkhill J, Dingle KE, Bowler ICJW, Jolley KA, Maiden MCJ (2013) Real-time genomic epidemiological evaluation of human *Campylobacter* isolates by use of whole-genome multilocus sequence typing. *J Clin Microbiol*, **51**:2526–34.

ID	Isolate	source	species	aspA	glnA	gltA	glyA	pgm	tkt	uncA	Sequence Type*	Clonal Complex**	Reference***
1	NC_002163	human	<i>Campylobacter jejuni</i>	2	1	5	3	4	1	5	43	ST-21 complex	2
2	CampsClin262	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	3	262	ST-21 complex	2
3	CampsClin266	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	54	5	266	ST-21 complex	2
4	CampsClin883	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	2
5	chicka21	chicken	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	2
6	cow518	cattle	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	2
7	CampsClin53	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	2
8	cowa21	cattle	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	2
9	chickc21	chicken	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	2
10	chick104	chicken	<i>Campylobacter jejuni</i>	2	1	1	3	7	1	5	104	ST-21 complex	2
11	chick19	chicken	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	2
12	chick50	chicken	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	2
13	chick53	chicken	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	2
14	chick262	chicken	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	3	262	ST-21 complex	2
15	chick266	chicken	<i>Campylobacter jejuni</i>	2	1	5	3	2	54	5	266	ST-21 complex	2
16	chick1086	chicken	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	2
17	chick1360	chicken	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	2
18	cowb21	cattle	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	2
19	cow104	cattle	<i>Campylobacter jejuni</i>	2	1	1	3	7	1	5	104	ST-21 complex	2
20	cow3201	cattle	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	2
21	chickb21	chicken	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	3
22	chick883	chicken	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	3
23	CampsClin21	human	<i>Campylobacter jejuni</i>	2	1	1	462	2	1	5		ST-21 complex	3
24	OxCrina21	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	3
25	CjLMG9879	human	<i>Campylobacter jejuni</i>	2	1	1	5	2	1	5	47	ST-21 complex	4
26	Cj2008-1025	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	4
27	Cj2008-831	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	4
28	Cj110-21	cattle	<i>Campylobacter jejuni</i>	2	1	2	3	2	1	5	982	ST-21 complex	4

29	Cj87330	chicken	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	4
30	Cj1928	cattle	<i>Campylobacter jejuni</i>	2	1	1	3	140	3	5	806	ST-21 complex	4
31	OXC5333	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
32	OXC5335	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
33	OXC5344	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
34	OXC5349	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
35	OXC5350	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
36	OXC5364	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
37	OXC5372	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
38	OXC5378	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
39	OXC5393	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
40	OXC5397	human	<i>Campylobacter jejuni</i>	2	1	1	5	11	343	5	5242	ST-21 complex	5
41	OXC5474	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
42	OXC5476	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
43	OXC5664	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
44	OXC5671	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	5
45	OXC5691	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
46	OXC5696	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
47	OXC5708	human	<i>Campylobacter jejuni</i>	2	1	42	3	148	1	5	861	ST-21 complex	5
48	OXC5710	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
49	OXC5713	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
50	OXC5720	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
51	OXC5495	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
52	OXC5617	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
53	OXC5618	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
54	OXC5625	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
55	OXC5628	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
56	OXC5639	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	5
57	OXC5647	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
58	OXC5649	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5

59	OXC5651	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
60	OXC5652	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
61	OXC5835	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
62	OXC5498	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
63	OXC5840	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
64	OXC5841	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
65	OXC5843	human	<i>Campylobacter jejuni</i>	2	1	42	3	148	1	5	861	ST-21 complex	5
66	OXC5846	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
67	OXC5848	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
68	OXC5862	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
69	OXC5871	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
70	OXC5879	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
71	OXC5880	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
72	OXC5413	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
73	OXC5414	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
74	OXC5416	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	23	3574	ST-21 complex	5
75	OXC5431	human	<i>Campylobacter jejuni</i>	2	1	1	3	492	1	5	5018	ST-21 complex	5
76	OXC5435	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
77	OXC5437	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	5
78	OXC5438	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
79	OXC5444	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
80	OXC5445	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
81	OXC5451	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
82	OXC5462	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
83	OXC5463	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
84	OXC5465	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
85	OXC5470	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	1	44	ST-21 complex	5
86	OXC5724	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
87	OXC5725	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
88	OXC5728	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	61	5	2355	ST-21 complex	5

89	OXC5731	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
90	OXC5737	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	54	5	266	ST-21 complex	5
91	OXC5739	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
92	OXC5754	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
93	OXC5757	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
94	OXC5766	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
95	OXC5767	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
96	OXC5771	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
97	OXC5772	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
98	OXC5774	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	3	5	190	ST-21 complex	5
99	OXC5791	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
100	OXC5799	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
101	OXC5804	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
102	OXC5808	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
103	OXC5811	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
104	OXC5818	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
105	OXC5819	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
106	OXC5821	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
107	OXC5822	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
108	OXC5823	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
109	OXC5825	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
110	OXC5826	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
111	OXC5829	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
112	OXC5832	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
113	OXC5887	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
114	OXC5888	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
115	OXC5895	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
116	OXC5899	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
117	OXC5902	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
118	OXC5905	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5

119	OXC5906	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
120	OXC5909	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
121	OXC5918	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
122	OXC5937	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
123	OXC5924	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
124	OXC5925	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
125	OXC5926	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
126	OXC5929	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
127	OXC6548	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
128	OXC6552	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	12	3769	ST-21 complex	5
129	OXC6558	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
130	OXC6562	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
131	OXC6563	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
132	OXC6564	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
133	OXC6565	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
134	OXC6571	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
135	OXC6573	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
136	OXC6251	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
137	OXC6257	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
138	OXC6266	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
139	OXC6270	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
140	OXC6275	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
141	OXC6277	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
142	OXC6282	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
143	OXC6285	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
144	OXC6286	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
145	OXC6289	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
146	OXC6292	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
147	OXC6300	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
148	OXC6301	human	<i>Campylobacter jejuni</i>	2	1	42	3	148	1	5	861	ST-21 complex	5

149	OXC6303	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
150	OXC6305	human	<i>Campylobacter jejuni</i>	2	1	42	3	148	1	5	861	ST-21 complex	5
151	OXC6310	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
152	OXC6317	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
153	OXC6324	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
154	OXC6325	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
155	OXC6326	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
156	OXC6329	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
157	OXC6331	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
158	OXC6334	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
159	OXC6335	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
160	OXC6340	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	3	262	ST-21 complex	5
161	OXC6347	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
162	OXC6355	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
163	OXC6367	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
164	OXC6370	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
165	OXC6379	human	<i>Campylobacter jejuni</i>	2	1	5	462	2	1	5	5726	ST-21 complex	5
166	OXC6383	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
167	OXC6384	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
168	OXC6393	human	<i>Campylobacter jejuni</i>	2	1	12	462	2	1	5	5727	ST-21 complex	5
169	OXC6394	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
170	OXC6405	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
171	OXC6420	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
172	OXC6429	human	<i>Campylobacter jejuni</i>	2	1	1	3	492	1	5	5018	ST-21 complex	5
173	OXC6449	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
174	OXC6457	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
175	OXC6459	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
176	OXC6461	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
177	OXC6464	human	<i>Campylobacter jejuni</i>	2	1	1	3	492	1	5	5018	ST-21 complex	5
178	OXC6479	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	5

179	OXC6483	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
180	OXC6489	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
181	OXC6493	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
182	OXC6496	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
183	OXC6500	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
184	OXC6502	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
185	OXC6505	human	<i>Campylobacter jejuni</i>	2	1	10	3	2	1	5	141	ST-21 complex	5
186	OXC6508	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
187	OXC6514	human	<i>Campylobacter jejuni</i>	2	84	12	3	11	1	5	3102	ST-21 complex	5
188	OXC6516	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
189	OXC6519	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
190	OXC6521	human	<i>Campylobacter jejuni</i>	2	1	1	5	2	1	5	47	ST-21 complex	5
191	OXC6524	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
192	OXC6527	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
193	OXC6530	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
194	OXC6531	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
195	OXC6538	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
196	OXC6539	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
197	OXC6543	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
198	OXC6255	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
199	OXC6590	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
200	OXC6596	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
201	OXC6598	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
202	OXC6600	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
203	OXC6602	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
204	OXC6603	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
205	OXC6604	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
206	OXC6613	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
207	OXC6615	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
208	OXC6616	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5

209	OXC6619	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
210	OXC6625	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
211	OXC6627	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
212	OXC6629	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
213	OXC6633	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
214	OXC6636	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
215	OXC6637	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
216	OXC6642	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
217	OXC6643	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
218	OXC6656	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
219	OXC6664	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
220	OXC6665	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
221	OXC6666	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
222	OXC6667	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	1	44	ST-21 complex	5
223	OXC6671	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
224	OXC6672	human	<i>Campylobacter jejuni</i>	2	1	1	5	2	1	5	47	ST-21 complex	5
225	OXC6673	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
226	OXC6677	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
227	OXC6678	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
228	OXC6681	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
229	OXC6682	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
230	OXC6689	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
231	OXC6690	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
232	OXC6691	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	3	5	190	ST-21 complex	5
233	OXC6702	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
234	OXC6703	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
235	OXC6706	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
236	OXC6711	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
237	OXC6712	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
238	OXC6713	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5

239	OXC6714	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
240	OXC6727	human	<i>Campylobacter jejuni</i>	2	2	52	3	2	100	5	5811	ST-21 complex	5
241	OXC6728	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
242	OXC6730	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
243	OXC6731	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
244	OXC6734	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
245	OXC6741	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
246	OXC6745	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
247	OXC6751	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
248	OXC6752	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
249	OXC6763	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
250	OXC6764	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
251	OXC6766	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	3	262	ST-21 complex	5
252	OXC6779	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
253	OXC6781	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
254	OXC6786	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	12	2135	ST-21 complex	5
255	OXC6789	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
256	OXC6791	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
257	OXC6798	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	301	4526	ST-21 complex	5
258	OXC6804	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
259	OXC6812	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
260	OXC6815	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
261	OXC6820	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
262	OXC6823	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	5
263	OXC6824	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	3	262	ST-21 complex	5
264	OXC6833	human	<i>Campylobacter jejuni</i>	2	1	1	3	7	1	5	104	ST-21 complex	5
265	OXC6932	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
266	OXC6941	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
267	OXC6945	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
268	OXC6946	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5

269	OXC6948	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
270	OXC6949	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
271	OXC6953	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
272	OXC6956	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
273	OXC6959	human	<i>Campylobacter jejuni</i>	2	1	12	462	2	1	5	5727	ST-21 complex	5
274	OXC6961	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
275	OXC6964	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
276	OXC6979	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
277	OXC6981	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	1	44	ST-21 complex	5
278	OXC6985	human	<i>Campylobacter jejuni</i>	2	17	12	462	2	1	5	6135	ST-21 complex	5
279	OXC6988	human	<i>Campylobacter jejuni</i>	2	1	5	462	2	1	5	5726	ST-21 complex	5
280	OXC6994	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
281	OXC6998	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
282	OXC7000	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
283	OXC7104	human	<i>Campylobacter jejuni</i>	2	1	5	10	2	1	6	6137	ST-21 complex	5
284	OXC7105	human	<i>Campylobacter jejuni</i>	2	1	5	10	2	1	6	6137	ST-21 complex	5
285	OXC7118	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	54	5	266	ST-21 complex	5
286	OXC7132	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
287	OXC7134	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
288	OXC7136	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
289	OXC7137	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
290	OXC7140	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
291	OXC7141	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	1	44	ST-21 complex	5
292	OXC7142	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
293	OXC7147	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
294	OXC7148	human	<i>Campylobacter jejuni</i>	2	1	79	3	2	1	5	822	ST-21 complex	5
295	OXC7150	human	<i>Campylobacter jejuni</i>	8	1	6	3	2	1	1	44	ST-21 complex	5
296	OXC7153	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
297	OXC7160	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
298	OXC7163	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5

299	OXC7168	human	<i>Campylobacter jejuni</i>	2	1	5	3	2	1	5	19	ST-21 complex	5
300	OXC6836	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
301	OXC6845	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
302	OXC6850	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
303	OXC6852	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	3	262	ST-21 complex	5
304	OXC6854	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
305	OXC6856	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
306	OXC6857	human	<i>Campylobacter jejuni</i>	2	1	12	88	2	1	5	520	ST-21 complex	5
307	OXC6860	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
308	OXC6861	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
309	OXC6862	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
310	OXC6865	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
311	OXC6867	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
312	OXC6868	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
313	OXC6872	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
314	OXC6876	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
315	OXC6879	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
316	OXC6886	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
317	OXC6892	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
318	OXC6900	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
319	OXC6909	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
320	OXC6922	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
321	OXC6924	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	5
322	OXC6927	human	<i>Campylobacter jejuni</i>	2	1	21	3	2	1	5	53	ST-21 complex	5
323	OXC7002	human	<i>Campylobacter jejuni</i>	2	1	1	3	2	1	5	21	ST-21 complex	5
324	OXC7015	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
325	OXC7020	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
326	OXC7029	human	<i>Campylobacter jejuni</i>	2	1	12	3	2	1	5	50	ST-21 complex	5
327	OXC7031	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	5
328	OXC7035	human	<i>Campylobacter jejuni</i>	2	17	2	3	2	1	5	883	ST-21 complex	5

					2	1	1	3	2	1	5	21	ST-21 complex	5	
329	OXC7037	human	<i>Campylobacter jejuni</i>		2	1	12	3	2	1	5	50	ST-21 complex	5	
330	OXC7041	human	<i>Campylobacter jejuni</i>		2	1	6	3	2	1	12	2135	ST-21 complex	5	
331	OXC7046	human	<i>Campylobacter jejuni</i>		8	1	5	3	2	1	54	5	266	ST-21 complex	5
332	OXC7049	human	<i>Campylobacter jejuni</i>		2	1	1	3	2	1	5	21	ST-21 complex	5	
333	OXC7058	human	<i>Campylobacter jejuni</i>		2	1	12	3	2	1	5	50	ST-21 complex	5	
334	OXC7065	human	<i>Campylobacter jejuni</i>		2	1	5	3	2	1	5	50	ST-21 complex	5	
335	OXC7068	human	<i>Campylobacter jejuni</i>		2	1	12	3	2	1	5	19	ST-21 complex	5	
336	OXC7071	human	<i>Campylobacter jejuni</i>		2	1	1	3	2	1	5	50	ST-21 complex	5	
337	OXC7073	human	<i>Campylobacter jejuni</i>		2	1	5	3	2	1	5	21	ST-21 complex	5	
338	OXC7078	human	<i>Campylobacter jejuni</i>		2	1	1	3	2	25	5	1949	ST-21 complex	5	
339	OXC7080	human	<i>Campylobacter jejuni</i>		2	1	12	3	2	1	5	50	ST-21 complex	5	
340	OXC7084	human	<i>Campylobacter jejuni</i>		2	1	1	3	2	1	5	21	ST-21 complex	5	
341	OXC7091	human	<i>Campylobacter jejuni</i>		2	1	1	3	2	1	3	262	ST-21 complex	5	
342	OXC7089	human	<i>Campylobacter jejuni</i>		2	1	21	3	2	1	5	53	ST-21 complex	5	
343	OXC7173	human	<i>Campylobacter jejuni</i>		8	1	6	3	2	1	1	44	ST-21 complex	5	
344	OXC7187	human	<i>Campylobacter jejuni</i>		2	1	1	3	2	1	5	21	ST-21 complex	5	
345	OXC7188	human	<i>Campylobacter jejuni</i>		2	1	12	3	2	1	5	50	ST-21 complex	5	
346	OXC7190	human	<i>Campylobacter jejuni</i>		2	1	1	3	2	1	5	21	ST-21 complex	5	
347	OXC7195	human	<i>Campylobacter jejuni</i>		2	1	1	3	2	1	5	21	ST-21 complex	5	
348	OXC6605	human	<i>Campylobacter jejuni</i>		2	1	1	3	2	1	5	21	ST-21 complex	5	
349	CAMP45	chicken	<i>Campylobacter jejuni</i>		4	7	10	4	1	7	1	45	ST-45 complex	1	
350	CampsClin11	human	<i>Campylobacter jejuni</i>		48	7	10	4	1	7	1	11	ST-45 complex	2	
351	chick2219	chicken	<i>Campylobacter jejuni</i>		10	7	10	4	1	7	1	2219	ST-45 complex	2	
352	chick594	chicken	<i>Campylobacter jejuni</i>		4	7	10	4	42	51	1	583	ST-45 complex	2	
353	cow334	cattle	<i>Campylobacter jejuni</i>		4	7	40	4	42	7	1	334	ST-45 complex	2	
354	CampsClin230	human	<i>Campylobacter jejuni</i>		4	7	41	4	42	7	1	230	ST-45 complex	2	
355	cowa45	cattle	<i>Campylobacter jejuni</i>		4	7	10	4	1	7	1	45	ST-45 complex	2	
356	chick2213	chicken	<i>Campylobacter jejuni</i>		4	7	40	4	42	7	1	334	ST-45 complex	2	
357	chickc45	chicken	<i>Campylobacter jejuni</i>		4	7	10	4	1	7	1	45	ST-45 complex	2	
358	chick11	chicken	<i>Campylobacter jejuni</i>		48	7	10	4	1	7	1	11	ST-45 complex	2	

359	chick1003	chicken	<i>Campylobacter jejuni</i>	8	7	4	4	125	7	1	1003	ST-45 complex	2
360	chick2048	chicken	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	2
361	chick2223	chicken	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	2
362	cowb45	cattle	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	2
363	cowc45	cattle	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	2
364	cowd45	cattle	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	2
365	cow137	cattle	<i>Campylobacter jejuni</i>	4	7	10	4	42	7	1	137	ST-45 complex	2
366	cow583	cattle	<i>Campylobacter jejuni</i>	4	7	10	4	42	51	1	583	ST-45 complex	2
367	cow3207	cattle	<i>Campylobacter jejuni</i>	4	7	40	4	42	7	1	334	ST-45 complex	3
368	cow3214	cattle	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	3
369	chickb45	chicken	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	3
370	chickd45	chicken	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	3
371	chick230	chicken	<i>Campylobacter jejuni</i>	4	7	41	4	42	7	1	230	ST-45 complex	3
372	OxCrina45	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	3
373	starling45	wild bird	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	3
374	goose137	wild bird	<i>Campylobacter jejuni</i>	4	7	10	4	42	7	1	137	ST-45 complex	3
375	duck45	wild bird	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	3
376	Cj55037	chicken	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	4
377	OXC5330	human	<i>Campylobacter jejuni</i>	8	7	4	4	125	7	1	1003	ST-45 complex	5
378	OXC5331	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	7	1	137	ST-45 complex	5
379	OXC5332	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
380	OXC5343	human	<i>Campylobacter jejuni</i>	4	7	10	4	10	7	1	2109	ST-45 complex	5
381	OXC5850	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
382	OXC5420	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
383	OXC5434	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	7	1	137	ST-45 complex	5
384	OXC5744	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	7	1	137	ST-45 complex	5
385	OXC5816	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5

386	OXC5941	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
387	OXC6259	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
388	OXC6278	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
389	OXC6293	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
390	OXC6294	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	51	1	583	ST-45 complex	5
391	OXC6313	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
392	OXC6314	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
393	OXC6321	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
394	OXC6330	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	51	1	583	ST-45 complex	5
395	OXC6339	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
396	OXC6351	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
397	OXC6358	human	<i>Campylobacter jejuni</i>	2	7	10	4	1	7	1	233	ST-45 complex	5
398	OXC6365	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
399	OXC6419	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
400	OXC6437	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
401	OXC6448	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	7	1	137	ST-45 complex	5
402	OXC6515	human	<i>Campylobacter jejuni</i>	4	7	73	4	1	7	1	845	ST-45 complex	5
403	OXC6536	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
404	OXC6592	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	7	1	137	ST-45 complex	5
405	OXC6614	human	<i>Campylobacter jejuni</i>	2	7	10	4	1	7	1	233	ST-45 complex	5
406	OXC6624	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
407	OXC6797	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
408	OXC6803	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
409	OXC6819	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	51	1	583	ST-45 complex	5
410	OXC6938	human	<i>Campylobacter jejuni</i>	4	7	10	4	10	7	1	2109	ST-45 complex	5
411	OXC6970	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	51	1	583	ST-45 complex	5
412	OXC7103	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
413	OXC7113	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	25	1	538	ST-45 complex	5
414	OXC7129	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
415	OXC7167	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	532	1	6886	ST-45 complex	5

416	OXC7169	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
417	OXC6894	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
418	OXC6895	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
419	OXC6896	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
420	OXC6916	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
421	OXC7014	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
422	OXC7018	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
423	OXC7025	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
424	OXC7026	human	<i>Campylobacter jejuni</i>	10	7	10	4	1	7	1	2219	ST-45 complex	5
425	OXC7032	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	51	1	583	ST-45 complex	5
426	OXC7061	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
427	OXC7067	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	7	1	137	ST-45 complex	5
428	OXC7074	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
429	OXC7090	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
430	OXC7093	human	<i>Campylobacter jejuni</i>	48	7	10	4	1	7	1	11	ST-45 complex	5
431	OXC7099	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
432	OXC7170	human	<i>Campylobacter jejuni</i>	4	7	10	4	42	7	1	137	ST-45 complex	5
433	OXC7179	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
434	OXC7185	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
435	OXC7186	human	<i>Campylobacter jejuni</i>	4	7	10	4	1	7	1	45	ST-45 complex	5
436	171	human	<i>Campylobacter coli</i>	33	38	30	115	104	85	17	867	ST-828 complex	1
437	CAMP886	pig	<i>Campylobacter coli</i>	33	38	30	82	104	85	68	887	ST-828 complex	1
438	CAMP2588	chicken	<i>Campylobacter coli</i>	33	38	30	115	104	85	17	867	ST-828 complex	1
439	CAMP3129	human	<i>Campylobacter coli</i>	157	39	30	79	668	35	17		ST-828 complex	1
440	CAMP1090	chicken	<i>Campylobacter coli</i>	33	39	30	82	104	43	17	828	ST-828 complex	2
441	CAMP828	chicken	<i>Campylobacter coli</i>	33	39	30	82	104	43	17	828	ST-828 complex	2
442	cow3202	cattle	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	2
443	Cc111-3	pig	<i>Campylobacter coli</i>	33	38	30	167	104	43	17	1467	ST-828 complex	4
444	CcZ163	chicken	<i>Campylobacter coli</i>	33	38	30	82	104	332	17	3336	ST-828 complex	4
445	Cc2553	chicken	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	4

446	Cc2680	chicken	<i>Campylobacter coli</i>	33	39	103	82	104	324	41	3872	ST-828 complex	4
447	Cc2685	chicken	<i>Campylobacter coli</i>	33	39	30	82	211	85	17	1082	ST-828 complex	4
448	Cc2688	chicken	<i>Campylobacter coli</i>	33	39	30	82	104	43	41	1017	ST-828 complex	4
449	Cc2692	chicken	<i>Campylobacter coli</i>	33	39	30	79	113	47	17	860	ST-828 complex	4
450	Cc2698	chicken	<i>Campylobacter coli</i>	33	39	30	82	113	43	17	829	ST-828 complex	4
451	Cc84-2	pig	<i>Campylobacter coli</i>	33	38	30	78	104	35	17	1113	ST-828 complex	4
452	Cc80352	chicken	<i>Campylobacter coli</i>	33	39	30	82	104	43	41	1017	ST-828 complex	4
453	Cc86119	chicken	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	4
454	Cc1091	cattle	<i>Campylobacter coli</i>	33	39	30	78	104	43	17	1068	ST-828 complex	4
455	Cc1098	cattle	<i>Campylobacter coli</i>	33	39	30	82	104	85	68	1104	ST-828 complex	4
456	Cc1148	cattle	<i>Campylobacter coli</i>	33	39	30	78	104	43	17	1068	ST-828 complex	4
457	Cc1417	cattle	<i>Campylobacter coli</i>	33	153	44	82	104	44	17	3221	ST-828 complex	4
458	Cc132-6	pig	<i>Campylobacter coli</i>	33	153	44	82	104	44	17	3221	ST-828 complex	4
459	Cc1891	cattle	<i>Campylobacter coli</i>	33	39	30	78	104	43	17	1068	ST-828 complex	4
460	Cc1909	cattle	<i>Campylobacter coli</i>	33	39	30	82	104	85	68	1104	ST-828 complex	4
461	Cc59-2	pig	<i>Campylobacter coli</i>	33	38	30	82	104	35	36	890	ST-828 complex	4
462	Cc1948	cattle	<i>Campylobacter coli</i>	33	39	30	82	104	85	68	1104	ST-828 complex	4
463	Cc1957	cattle	<i>Campylobacter coli</i>	33	153	30	82	104	85	68	2698	ST-828 complex	4
464	Cc1961	cattle	<i>Campylobacter coli</i>	33	39	30	82	104	85	68	1104	ST-828 complex	4
465	Cc202/04	human	<i>Campylobacter coli</i>	33	39	30	82	189	43	17	1585	ST-828 complex	4
466	Cc67-8	pig	<i>Campylobacter coli</i>	32	39	44	82	104	43	36	1061	ST-828 complex	4
467	CcLMG9854	human	<i>Campylobacter coli</i>	33	39	30	78	104	43	17	1068	ST-828 complex	4
468	CcLMG23336	human	<i>Campylobacter coli</i>	33	176	30	82	451	43	17	3868	ST-828 complex	4
469	CcLMG23341	human	<i>Campylobacter coli</i>	33	39	30	79	104	35	17	855	ST-828 complex	4
470	CcLMG23342	human	<i>Campylobacter coli</i>	33	39	30	79	104	35	17	855	ST-828 complex	4
471	CcLMG23344	human	<i>Campylobacter coli</i>	33	176	30	82	113	43	17	1586	ST-828 complex	4
472	CcLMG9853	human	<i>Campylobacter coli</i>	33	39	261	79	104	64	17	3869	ST-828 complex	4
473	CcH6	human	<i>Campylobacter coli</i>	33	176	30	79	113	43	17	3020	ST-828 complex	4
474	CcH8	human	<i>Campylobacter coli</i>	33	39	30	79	104	43	41	901	ST-828 complex	4
475	CCH9	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	4

476	CcH56	human	<i>Campylobacter coli</i>	33	38	30	82	104	35	17	1096	ST-828 complex	4
477	CcZ156	chicken	<i>Campylobacter coli</i>	33	38	30	82	104	43	17	854	ST-828 complex	4
478	OXC5348	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
479	OXC5353	human	<i>Campylobacter coli</i>	33	39	30	79	530	43	17	5149	ST-828 complex	5
480	OXC5363	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
481	OXC5370	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
482	OXC5386	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5
483	OXC5677	human	<i>Campylobacter coli</i>	33	66	30	174	188	43	17	6132	ST-828 complex	5
484	OXC5685	human	<i>Campylobacter coli</i>	33	39	30	82	112	56	17	1578	ST-828 complex	5
485	OXC5688	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
486	OXC5705	human	<i>Campylobacter coli</i>	33	39	30	82	113	44	17	872	ST-828 complex	5
487	OXC5482	human	<i>Campylobacter coli</i>	33	39	65	79	104	85	17	2273	ST-828 complex	5
488	OXC5605	human	<i>Campylobacter coli</i>	33	39	30	79	113	43	17	832	ST-828 complex	5
489	OXC5629	human	<i>Campylobacter coli</i>	33	39	30	79	104	35	17	855	ST-828 complex	5
490	OXC5635	human	<i>Campylobacter coli</i>	124	39	30	79	104	47	17	1541	ST-828 complex	5
491	OXC5837	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
492	OXC5842	human	<i>Campylobacter coli</i>	33	39	66	82	104	44	174	2178	ST-828 complex	5
493	OXC5849	human	<i>Campylobacter coli</i>	33	39	30	79	104	47	17	830	ST-828 complex	5
494	OXC5853	human	<i>Campylobacter coli</i>	33	39	30	79	104	47	17	830	ST-828 complex	5
495	OXC5855	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5
496	OXC5856	human	<i>Campylobacter coli</i>	33	39	30	79	104	47	17	830	ST-828 complex	5
497	OXC5864	human	<i>Campylobacter coli</i>	33	39	30	82	189	47	17	1191	ST-828 complex	5
498	OXC5473	human	<i>Campylobacter coli</i>	33	39	30	82	23	43	17	6131	ST-828 complex	5
499	OXC5723	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
500	OXC5742	human	<i>Campylobacter coli</i>	33	39	30	82	113	44	17	872	ST-828 complex	5
501	OXC5763	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5
502	OXC5773	human	<i>Campylobacter coli</i>	33	39	30	82	104	43	17	828	ST-828 complex	5
503	OXC5796	human	<i>Campylobacter coli</i>	33	39	30	82	113	44	17	872	ST-828 complex	5
504	OXC5810	human	<i>Campylobacter coli</i>	33	39	30	82	189	44	17	5165	ST-828 complex	5
505	OXC5827	human	<i>Campylobacter coli</i>	33	39	30	79	113	47	17	860	ST-828 complex	5

506	OXC5831	human	<i>Campylobacter coli</i>	33	39	30	82	189	47	17	1191	ST-828 complex	5
507	OXC5922	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
508	OXC5923	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
509	OXC6559	human	<i>Campylobacter coli</i>	33	38	30	466	104	43	17	5757	ST-828 complex	5
510	OXC6568	human	<i>Campylobacter coli</i>	33	39	30	82	113	44	17	872	ST-828 complex	5
511	OXC6576	human	<i>Campylobacter coli</i>	33	39	30	79	104	35	17	855	ST-828 complex	5
512	OXC6577	human	<i>Campylobacter coli</i>	33	39	30	82	113	44	17	872	ST-828 complex	5
513	OXC6253	human	<i>Campylobacter coli</i>	33	39	30	82	113	43	17	829	ST-828 complex	5
514	OXC6258	human	<i>Campylobacter coli</i>	33	39	30	79	113	47	17	860	ST-828 complex	5
515	OXC6263	human	<i>Campylobacter coli</i>	33	39	30	79	104	47	17	830	ST-828 complex	5
516	OXC6267	human	<i>Campylobacter coli</i>	33	39	30	79	247	43	17	5642	ST-828 complex	5
517	OXC6276	human	<i>Campylobacter coli</i>	33	39	65	82	596	47	17	5733	ST-828 complex	5
518	OXC6296	human	<i>Campylobacter coli</i>	87	39	30	79	243	43	17	5737	ST-828 complex	5
519	OXC6297	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5
520	OXC6308	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
521	OXC6309	human	<i>Campylobacter coli</i>	33	110	30	115	104	43	17	5734	ST-828 complex	5
522	OXC6312	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
523	OXC6337	human	<i>Campylobacter coli</i>	33	39	65	140	104	43	17	1600	ST-828 complex	5
524	OXC6338	human	<i>Campylobacter coli</i>	33	39	30	79	104	206	17	1628	ST-828 complex	5
525	OXC6343	human	<i>Campylobacter coli</i>	33	39	30	82	104	47	36	962	ST-828 complex	5
526	OXC6371	human	<i>Campylobacter coli</i>	33	110	30	115	104	43	17	5734	ST-828 complex	5
527	OXC6372	human	<i>Campylobacter coli</i>	33	39	30	139	113	47	17	4425	ST-828 complex	5
528	OXC6376	human	<i>Campylobacter coli</i>	33	110	30	115	104	43	17	5734	ST-828 complex	5
529	OXC6378	human	<i>Campylobacter coli</i>	33	39	30	139	113	47	17	4425	ST-828 complex	5
530	OXC6380	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	353	5735	ST-828 complex	5
531	OXC6385	human	<i>Campylobacter coli</i>	33	39	30	79	104	206	17	1628	ST-828 complex	5
532	OXC6386	human	<i>Campylobacter coli</i>	33	39	30	82	113	43	17	829	ST-828 complex	5
533	OXC6400	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5
534	OXC6416	human	<i>Campylobacter coli</i>	33	39	30	82	104	44	17	1145	ST-828 complex	5
535	OXC6424	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5

536	OXC6426	human	<i>Campylobacter coli</i>	32	66	66	82	104	43	17	2464	ST-828 complex	5
537	OXC6428	human	<i>Campylobacter coli</i>	33	39	30	79	113	43	17	832	ST-828 complex	5
538	OXC6447	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
539	OXC6460	human	<i>Campylobacter coli</i>	124	39	30	79	104	47	17	1541	ST-828 complex	5
540	OXC6471	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
541	OXC6472	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
542	OXC6474	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
543	OXC6476	human	<i>Campylobacter coli</i>	33	39	30	79	113	47	17	860	ST-828 complex	5
544	OXC6504	human	<i>Campylobacter coli</i>	33	39	30	79	113	43	17	832	ST-828 complex	5
545	OXC6513	human	<i>Campylobacter coli</i>	33	39	30	82	104	43	17	828	ST-828 complex	5
546	OXC6523	human	<i>Campylobacter coli</i>	33	66	30	82	113	206	17	5659	ST-828 complex	5
547	OXC6537	human	<i>Campylobacter coli</i>	124	39	30	79	104	47	17	1541	ST-828 complex	5
548	OXC6597	human	<i>Campylobacter coli</i>	33	39	30	79	104	35	17	855	ST-828 complex	5
549	OXC6601	human	<i>Campylobacter coli</i>	33	39	30	79	104	85	17	1614	ST-828 complex	5
550	OXC6630	human	<i>Campylobacter coli</i>	33	39	30	82	104	44	17	1145	ST-828 complex	5
551	OXC6651	human	<i>Campylobacter coli</i>	33	39	66	174	104	43	41	1181	ST-828 complex	5
552	OXC6684	human	<i>Campylobacter coli</i>	33	39	30	82	104	47	36	962	ST-828 complex	5
553	OXC6685	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
554	OXC6705	human	<i>Campylobacter coli</i>	33	39	65	79	104	35	17	1957	ST-828 complex	5
555	OXC6710	human	<i>Campylobacter coli</i>	2	39	30	82	113	47	17	5810	ST-828 complex	5
556	OXC6725	human	<i>Campylobacter coli</i>	33	38	30	82	104	35	17	1096	ST-828 complex	5
557	OXC6735	human	<i>Campylobacter coli</i>	33	39	30	82	104	44	17	1145	ST-828 complex	5
558	OXC6738	human	<i>Campylobacter coli</i>	33	39	65	79	104	85	17	2273	ST-828 complex	5
559	OXC6744	human	<i>Campylobacter coli</i>	53	39	30	82	603	43	17	5813	ST-828 complex	5
560	OXC6761	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
561	OXC6765		<i>Campylobacter coli</i>	33	39	30	79	113	43	17	832	ST-828 complex	5
562	OXC6785	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
563	OXC6810	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5
564	OXC6817	human	<i>Campylobacter coli</i>	33	39	30	82	104	47	36	962	ST-828 complex	5
565	OXC6825	human	<i>Campylobacter coli</i>	33	39	30	79	113	43	41	2183	ST-828 complex	5

566	OXC6933	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
567	OXC6937	human	<i>Campylobacter coli</i>	33	176	30	82	104	43	17	3753	ST-828 complex	5
568	OXC6962	human	<i>Campylobacter coli</i>	33	39	65	79	104	85	17	2273	ST-828 complex	5
569	OXC6984	human	<i>Campylobacter coli</i>	33	39	30	82	104	47	17	1055	ST-828 complex	5
570	OXC6987	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5
571	OXC6996	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
572	OXC7102	human	<i>Campylobacter coli</i>	33	39	30	82	189	47	17	1191	ST-828 complex	5
573	OXC7110	human	<i>Campylobacter coli</i>	33	39	30	82	189	47	17	1191	ST-828 complex	5
574	OXC7124	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
575	OXC7130	human	<i>Campylobacter coli</i>	33	39	30	79	113	43	41	2183	ST-828 complex	5
576	OXC7131	human	<i>Campylobacter coli</i>	33	39	30	82	104	43	17	828	ST-828 complex	5
577	OXC7154	human	<i>Campylobacter coli</i>	33	39	30	82	104	43	17	828	ST-828 complex	5
578	OXC7164	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
579	OXC6847	human	<i>Campylobacter coli</i>	33	39	30	82	113	43	17	829	ST-828 complex	5
580	OXC6864	human	<i>Campylobacter coli</i>	33	39	30	82	113	43	17	829	ST-828 complex	5
581	OXC6873	human	<i>Campylobacter coli</i>	33	39	30	81	561	47	17	5349	ST-828 complex	5
582	OXC6901	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
583	OXC6920	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
584	OXC7027	human	<i>Campylobacter coli</i>	33	38	30	82	118	43	17	1016	ST-828 complex	5
585	OXC7051	human	<i>Campylobacter coli</i>	33	38	30	82	118	43	17	1016	ST-828 complex	5
586	OXC7054	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5
587	OXC7070	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
588	OXC7082	human	<i>Campylobacter coli</i>	33	39	30	82	113	47	17	825	ST-828 complex	5
589	OXC7083	human	<i>Campylobacter coli</i>	33	39	30	79	104	35	17	855	ST-828 complex	5
590	OXC7097	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
591	OXC7177	human	<i>Campylobacter coli</i>	33	39	30	139	113	47	17	4425	ST-828 complex	5
592	OXC7199	human	<i>Campylobacter coli</i>	33	39	30	82	104	56	17	827	ST-828 complex	5
593	OXC7200	human	<i>Campylobacter coli</i>	33	39	30	82	113	43	17	829	ST-828 complex	5