An inexact Noda iteration for computing the smallest eigenpair of a large irreducible monotone matrix

Ching-Sung Liu

SUMMARY

In this paper, we present an inexact Noda iteration with inner-outer iterations for finding the smallest eigenvalue and the associated eigenvector of an irreducible monotone matrix. The proposed inexact Noda iteration contains two main relaxation steps for computing the smallest eigenvalue and the associated eigenvector, respectively. These relaxation steps depend on the relaxation factors, and we analyze how the relaxation factors in the relaxation steps affect the convergence of the outer iterations. By considering two different relaxation factors for solving the inner linear systems involved, we prove that the convergence is globally linear or superlinear, depending on the relaxation factor, and that the relaxation factor also influences the convergence rate. The proposed inexact Noda iterations are structure preserving and maintain the positivity of approximate eigenvectors. Numerical examples are provided to illustrate that the proposed inexact Noda iterations are practical, and they always preserve the positivity of approximate eigenvectors. Copyright © 2010 John Wiley & Sons, Ltd.

Received ...

KEY WORDS: Inexact Noda iteration, Modified inexact Noda iteration, *M*-matrix, non-negative matrix, monotone matrix, smallest eigenpair, singular value, Perron vector, Perron root.

Department of Applied Mathematics, National Chiao Tung University, Taiwan.

Ching-Sung Liu, Department of Applied Mathematics, National Chiao Tung University, Taiwan. E-mail: chingsungliu@nctu.edu.tw

1. INTRODUCTION

Monotone matrices arise in many areas of mathematics, such as stability analysis [19], and bounds for eigenvalues and singular values [3, 4]. In many applications, one is interested in finding the smallest eigenvalue λ and the associated eigenvector **x** of an irreducible nonsingular monotone matrix $A \in \mathbb{R}^{n \times n}$. The smallest eigenvalue λ of a monotone matrix A is defined as $\sigma_{\min}(A) =$ $\min\{|\lambda| \mid \lambda \in \sigma(A)\}$, where $\sigma(A)$ denotes the set of eigenvalues of A. In [23, 25], a real matrix A is called monotone if and only if A^{-1} is a non-negative matrix. The irreducible nonsingular Mmatrices are one of the most important classes of matrices for applications such as discretized PDEs, Markov chains [2] and electric circuits [24], and they have been studied extensively in the literature [5, Chapter 6]. It is well known that there exist some monotone matrices that are not M-matrices, such as matrices that can be written as a product of M-matrices.

There are some differences between an *M*-matrix and a monotone matrix. For example, an *M*-matrix can be expressed in the form $\sigma I - B$ with a non-negative matrix *B* and some constant $\sigma > \rho(B)$, where $\rho(\cdot)$ denotes the spectral radius, see [5]. Thus, the smallest eigenvalue λ of an irreducible nonsingular *M*-matrix *A* is equal to $\sigma - \rho(B) > 0$. In contrast, the smallest eigenvalue of a monotone matrix *A* can only be expressed as $\sigma_{\min}(A) = \rho(A^{-1})^{-1}$. However, the smallest eigenvalue retains the same properties [12, p. 487], that is, the largest eigenvalue of an irreducible non-negative matrix A^{-1} is the Perron root, which is simple and equal to the spectral radius of A^{-1} with a positive associated eigenvector.

For the computation of the Perron vector of a non-negative matrix B, many methods exist [21, 22, 28, 20, 14, 13, 17, 6, 26, 16] but the power methods are not structure preserving and cannot guarantee the desired positivity of approximations when the Perron vector \mathbf{x} has very small components. Therefore, a central concern is how to preserve strict positivity of approximations to the Perron vector. In 1971, Noda introduced an inverse iteration method with shifted Rayleigh quotientlike approximations [18] for non-negative matrix eigenvalue problems. This iteration method is called Noda iteration (NI), and it has also been adapted to the computation of the smallest eigenvalue and the eigenvector of an irreducible nonsingular M-matrix [29, 1]. The major advantages of Noda iteration are structure preservation and global convergence. More precisely, it generates a monotonically decreasing sequence of approximate eigenvalues that is guaranteed to converge to $\rho(B)$, and maintains the positivity of approximate eigenvectors. Furthermore, the convergence has been proven to be superlinear [18] and asymptotically quadratic [9]. In [15], the authors introduced two inexact strategies for Noda iteration, which are called inexact Noda iteration (INI) to find the Perron vector of a non-negative matrix (or *M*-matrix). The proposed INI algorithms are practical, and they always preserve the positivity of approximate eigenvectors. Moreover, the convergence of INI with these two strategies is globally linear and superlinear with convergence order $\frac{1+\sqrt{5}}{2}$, respectively.

In this paper, we propose an inexact Noda iteration (INI) to find the smallest eigenvalue and the associated eigenvector of an irreducible monotone matrix *A*. The major contribution of this paper is to provide two main relaxation steps for computing the smallest eigenvalue λ and the associated eigenvector **x**, respectively. The first step is to use $O(\gamma_k \min(\mathbf{x}_k))$ as a stopping criterion for inner iterations, with $0 < \gamma_k < 1$, where \mathbf{x}_k is the current positive approximate eigenvector. The second step is to update the approximate eigenvalues by using the recurrence relations $\overline{\lambda}_{k+1} = \overline{\lambda}_k - (1 - \gamma_k) \min(\frac{\mathbf{x}_k}{\mathbf{y}_{k+1}})$, where y_{k+1} is the next normalized positive approximate eigenvector, so resulting INI algorithms are structure preserving and globally convergent. The above parameter γ_k is called the "relaxation factor". We then establish a rigorous convergence theory of INI with two different relaxation factors γ_k , and prove that the convergence of the resulting INI algorithms is globally linear, and superlinear with the relaxation factor γ_k as the convergence rate, respectively.

In fact, the inner iterations of INI (or NI) require the solution of ill-conditioned linear systems when the sequence of approximate eigenvalues converges to $\rho(A^{-1})$ (or $\rho(B)$). In order to reduce the condition number of the inner linear system, we propose a modified Noda iteration (MNI) by using rank one update for the inner iterations, and we show that MNI and NI are mathematically equivalent. For monotone matrix eigenvalue problems, we also develop an integrated algorithm that combines INI with MNI, and we call this modified inexact Noda iteration (MINI). This hybrid iterative method can significantly improve the condition number of inner linear systems of INI.

The paper is organized as follows. In Section 2, we introduce the Noda iteration and some preliminaries. Section 3 contains the new strategy for inexact Noda iteration, and proves some basic properties for it. In Section 4, we establish its convergence theory, and derive the asymptotic convergence factor precisely. In Section 5, we present the integrated algorithm that combines INI with MNI. Finally, in Section 6 we present some numerical examples illustrating the convergence theory and the effectiveness of INI, and we make some concluding remarks in Section 7.

2. PRELIMINARIES AND NOTATION

For any real matrix $B = [b_{ij}] \in \mathbb{R}^{n \times n}$, we write $B \ge 0$ (> 0) if $b_{ij} \ge 0$ (> 0) for all $1 \le i, j \le n$. We define $|B| = [|b_{ij}|]$. If $B \ge 0$, we say *B* is a non-negative matrix, and if B > 0, we say *B* is a positive matrix. For real matrices *B* and *C* of the same size, if B - C is a non-negative matrix, we write $B \ge C$. A non-negative (positive) vector is similarly defined. A non-negative matrix *B* is said to be reducible if it can be placed into block upper-triangular form by simultaneous row/column permutations; otherwise it is irreducible. If μ is not an eigenvalue of *B*, the function $sep(\mu, B)$ is defined as

$$\operatorname{sep}(\mu, B) = \|(\mu I - B)^{-1}\|^{-1}.$$
(1)

 \angle (**w**,**z**) denotes the acute angle of any two nonzero vectors **w** and **z**. Throughout the paper, we use a 2-norm for vectors and matrices, and the superscript *T* denotes its transpose.

We review some fundamental properties of non-negative matrices, monotone matrices and *M*-matrices.

Definition 1

A matrix *A* is said to be "monotone" if $Ax \ge 0$ implies $x \ge 0$ for any positive vector.

Another characterization of monotone matrices is given by the following well known theorem.

Theorem 1 ([7]) A is monotone if and only if A is non-singular and $A^{-1} > 0$.

Definition 2

A monotone matrix *M* is an *M*-matrix if $M = (m_{ij}), m_{ij} \leq 0$ for $i \neq j$.

Lemma 1 ([5])

Let M is a nonsingular M-matrix. Then the following statements are equivalent:

1. $M = (a_{ij}), a_{ij} \leq 0$ for $i \neq j$, and $M^{-1} \geq 0$; 2. $M = \sigma I - B$ with some $B \geq 0$ and $\sigma > \rho(B)$.

For a pair of positive vectors v and w, define

$$\max\left(\frac{\mathbf{w}}{\mathbf{v}}\right) = \max_{i}\left(\frac{\mathbf{w}^{(i)}}{\mathbf{v}^{(i)}}\right), \quad \min\left(\frac{\mathbf{w}}{\mathbf{v}}\right) = \min_{i}\left(\frac{\mathbf{w}^{(i)}}{\mathbf{v}^{(i)}}\right)$$

where $\mathbf{v} = [\mathbf{v}^{(1)}, \mathbf{v}^{(2)}, \dots, \mathbf{v}^{(n)}]^T$ and $\mathbf{w} = [\mathbf{w}^{(1)}, \mathbf{w}^{(2)}, \dots, \mathbf{w}^{(n)}]^T$. The following lemma gives bounds for the spectral radius of a non-negative matrix *B*.

Lemma 2 ([12, p. 493])

Let *B* be an irreducible non-negative matrix. If $\mathbf{v} > \mathbf{0}$ is not an eigenvector of *B*, then

$$\min\left(\frac{B\mathbf{v}}{\mathbf{v}}\right) < \rho(B) < \max\left(\frac{B\mathbf{v}}{\mathbf{v}}\right). \tag{2}$$

2.1. The Noda iteration

The Noda iteration [18] is an inverse iteration shifted by a Rayleigh quotient-like approximation of the Perron root of an irreducible non-negative matrix B.

Given an initial vector $\mathbf{x}_0 > \mathbf{0}$ with $\|\mathbf{x}_0\| = 1$, the Noda iteration (NI) consists of three steps:

$$(\lambda_k I - B) \mathbf{y}_{k+1} = \mathbf{x}_k, \tag{3}$$

$$\mathbf{x}_{k+1} = \mathbf{y}_{k+1} / \|\mathbf{y}_{k+1}\|, \tag{4}$$

$$\widehat{\lambda}_{k+1} = \max\left(\frac{B\mathbf{x}_{k+1}}{\mathbf{x}_{k+1}}\right). \tag{5}$$

The main task is to compute a new approximation \mathbf{x}_{k+1} to \mathbf{x} by solving the inner linear system (3). From Lemma 2, we know that $\hat{\lambda}_k > \rho(B)$ if \mathbf{x}_k is not a scalar multiple of the eigenvector \mathbf{x} . This result shows that $\hat{\lambda}_k I - B$ is an irreducible nonsingular *M*-matrix, and its inverse is an irreducible non-negative matrix. Therefore, we have $\mathbf{y}_{k+1} > \mathbf{0}$ and $\mathbf{x}_{k+1} > \mathbf{0}$, i.e., \mathbf{x}_{k+1} is always a positive vector if \mathbf{x}_k is. After variable transformation, we get $\hat{\lambda}_{k+1}$ from the following relation:

$$\widehat{\lambda}_{k+1} = \widehat{\lambda}_k - \min\left(\frac{\mathbf{x}_k}{\mathbf{y}_{k+1}}\right),$$

so $\{\widehat{\lambda}_k\}$ is monotonically decreasing.

Copyright © 2010 John Wiley & Sons, Ltd. Prepared using nlaauth.cls

2.2. The inexact Noda iteration

The inexact Noda iteration. Based on the Noda iteration, in [15] the authors propose an inexact Noda iteration (INI) for the computation of the spectral radius of a non-negative irreducible matrix *B*. In this paper, since *A* is a monotone matrix, i.e., A^{-1} is a non-negative matrix, we replace *B* by A^{-1} in (3), i.e.,

$$(\widehat{\lambda}_k I - A^{-1}) \mathbf{y}_{k+1} = \mathbf{x}_k.$$
(6)

When A is large and sparse, we see that we must resort to an iterative linear solver to get an approximate solution. In order to reduce the computational cost of (6), we solve y_{k+1} in (6) by inexactly satisfying

$$(\widehat{\lambda}_k I - A^{-1}) \mathbf{y}_{k+1} = \mathbf{x}_k + A^{-1} \mathbf{f}_k, \tag{7}$$

which is equivalent to

$$(\lambda_k A - I) \mathbf{y}_{k+1} = A \mathbf{x}_k + \mathbf{f}_k, \tag{8}$$

$$\mathbf{x}_{k+1} = \mathbf{y}_{k+1} / \|\mathbf{y}_{k+1}\|, \tag{9}$$

where \mathbf{f}_k is the residual vector between $(\widehat{\lambda}_k A - I) \mathbf{y}_{k+1}$ and $A\mathbf{x}_k$. Here, the residual norm (inner tolerance) $\xi_k := \|\mathbf{f}_k\|$ can be changed at each iterative step *k*.

Theorem 2 ([15])

Let *A* be an irreducible monotone matrix and $0 \le \gamma < 1$ be a fixed constant. For the unit length $\mathbf{x}_k > \mathbf{0}$, if $\mathbf{x}_k \ne \mathbf{x}$ and \mathbf{f}_k in (8) satisfies

$$\left\|A^{-1}\mathbf{f}_k\right\| \le \gamma \min\left(\mathbf{x}_k\right),\tag{10}$$

then $\{\hat{\lambda}_k\}$ is monotonically decreasing and $\lim_{k\to\infty} \hat{\lambda}_k = \rho(A^{-1})$. Moreover, the convergence of INI is at least globally linear.

Based on (8)-(10), we describe INI as Algorithm 1.

Algorithm 1 Inexact Noda Iteration (INI)

- 1. Given $\widehat{\lambda}_0$, $\mathbf{x}_0 > \mathbf{0}$ with $\|\mathbf{x}_0\| = 1$, $0 \le \gamma < 1$ and tol > 0.
- 2. for $k = 0, 1, 2, \dots$
- 3. Solve $(\widehat{\lambda}_k A I) \mathbf{y}_{k+1} = A \mathbf{x}_k$ inexactly such that the inner tolerance ξ_k satisfies condition (10)
- 4. Normalize the vector $\mathbf{x}_{k+1} = \mathbf{y}_{k+1} / ||\mathbf{y}_{k+1}||$.

5. Compute
$$\lambda_{k+1} = \max\left(\frac{A - \mathbf{x}_{k+1}}{\mathbf{x}_{k+1}}\right)$$

6. **until** convergence: Resi= $||A\mathbf{x}_{k+1} - \widehat{\lambda}_k^{-1}\mathbf{x}_{k+1}|| < \text{tol.}$

Using the relation (7), step 5 in Algorithm 1 can be rewritten as

$$\widehat{\lambda}_{k+1} = \widehat{\lambda}_k - \min\left(\frac{\mathbf{x}_k + A^{-1}\mathbf{f}_k}{\mathbf{y}_{k+1}}\right).$$

Unfortunately, A^{-1} is not explicitly available; in other words, we need to compute " $A^{-1}\mathbf{f}_k$ " exactly for the required approximate eigenvalue $\hat{\lambda}_{k+1}$. Hence, in the next section, we propose a new strategy to estimate the approximate eigenvalues without increasing the computational cost. This strategy is practical and preserves the strictly decreasing property of the approximate eigenvalue sequence.

3. THE RELAXATION STRATEGY FOR INI AND SOME BASIC PROPERTIES

In order to ensure that INI is correctly implemented, we now propose two main relaxation steps to define Algorithm 2:

• The residual norm satisfies

$$\boldsymbol{\xi}_{k} = \|\mathbf{f}_{k}\| \le \gamma_{k} \operatorname{sep}(0, A) \min(\mathbf{x}_{k}), \tag{11}$$

where $0 \le \gamma_k \le \gamma < 1$ with a constant upper bound γ .

• The update of the approximate eigenvalue satisfies

$$\overline{\lambda}_{k+1} = \overline{\lambda}_k - (1 - \gamma_k) \min\left(\frac{\mathbf{x}_k}{\mathbf{y}_{k+1}}\right).$$
(12)

Algorithm 2 Inexact Noda Iteration for monotone matrices (INI)

- 1. Given $\overline{\lambda}_0$, $\mathbf{x}_0 > \mathbf{0}$ with $\|\mathbf{x}_0\| = 1$, $0 \le \gamma < 1$ and tol > 0.
- 2. for $k = 0, \underline{1}, 2, \ldots$
- 3. Solve $(\overline{\lambda}_k A I) \mathbf{y}_{k+1} = A \mathbf{x}_k$ inexactly such that the inner tolerance ξ_k satisfies condition (11)
- 4. Normalize the vector $\mathbf{x}_{k+1} = \mathbf{y}_{k+1} / ||\mathbf{y}_{k+1}||$.
- 5. Compute $\overline{\lambda}_{k+1}$ that satisfies condition (12).
- 6. **until** convergence: $||A\mathbf{x}_{k+1} \overline{\lambda}_k^{-1}\mathbf{x}_{k+1}|| < \text{tol.}$

In step 3 of Algorithm 2, it leads to two equivalent inexact relation satisfying

$$(\overline{\lambda}_k I - A^{-1}) \mathbf{y}_{k+1} = \mathbf{x}_k + A^{-1} \mathbf{f}_k$$
(13)

$$(\overline{\lambda}_k A - I) \mathbf{y}_{k+1} = A \mathbf{x}_k + \mathbf{f}_k, \tag{14}$$

We remark that $\overline{\lambda}_{k+1}$ in (12) is no longer equal to $\max\left(\frac{A^{-1}\mathbf{x}_{k+1}}{\mathbf{x}_{k+1}}\right)$, and therefore that $\overline{\lambda}_{k+1}$ cannot be clearly demonstrated to be greater than its lower bound $\rho(A^{-1})$. The following lemma ensures that $\rho(A^{-1})$ is still the lower bound of $\overline{\lambda}_k$.

Lemma 3

Let *A* be an irreducible monotone matrix. For the unit length $\mathbf{x}_k \neq \mathbf{x} > \mathbf{0}$ and the relaxation factor $\gamma_k \in [0, 1)$, if $\overline{\lambda}_k > \rho(A^{-1})$, \mathbf{f}_k in (14) satisfies condition (11) and the approximate eigenvalue satisfies (12), then the new approximation $\mathbf{x}_{k+1} > \mathbf{0}$ and the sequence $\{\overline{\lambda}_k\}$ is monotonically decreasing and bounded below by $\rho(A^{-1})$, i.e.,

$$\overline{\lambda}_k > \overline{\lambda}_{k+1} \ge \rho(A^{-1}).$$

Proof

From (12) and $\gamma_k \in [0, 1)$, it is easy to know that $\{\overline{\lambda}_k\}$ is monotonically decreasing, i.e., $\overline{\lambda}_k > \overline{\lambda}_{k+1}$. From (11),

$$\left\|A^{-1}\mathbf{f}_{k}\right\| \leq \left\|A^{-1}\right\| \left\|\mathbf{f}_{k}\right\| \leq \gamma_{k} \min(\mathbf{x}_{k}),\tag{15}$$

which implies $|A^{-1}\mathbf{f}_k| \leq \gamma_k \mathbf{x}_k$, then $\mathbf{x}_k + A^{-1}\mathbf{f}_k > 0$. Consequently, $\overline{\lambda}_k I - A^{-1}$ is a nonsingular *M*-matrix, and the vector \mathbf{y}_{k+1} satisfies

$$\mathbf{y}_{k+1} = (\overline{\lambda}_k I - A^{-1})^{-1} \left(\mathbf{x}_k + A^{-1} \mathbf{f}_k \right) > \mathbf{0}.$$

This implies $\mathbf{x}_{k+1} = \mathbf{y}_{k+1} / ||\mathbf{y}_{k+1}|| > \mathbf{0}$.

Copyright © 2010 John Wiley & Sons, Ltd. Prepared using nlaauth.cls We now prove $\overline{\lambda}_k$ is bounded below by $\rho(A^{-1})$. From (15), we have

$$(1-\gamma_k)\mathbf{x}_k \le \mathbf{x}_k + A^{-1}\mathbf{f}_k \le (1+\gamma_k)\mathbf{x}_k, \tag{16}$$

and

$$(1-\gamma_k)\frac{\mathbf{x}_k}{\mathbf{y}_{k+1}} \le \frac{\mathbf{x}_k + A^{-1}\mathbf{f}_k}{\mathbf{y}_{k+1}} \le (1+\gamma_k)\frac{\mathbf{x}_k}{\mathbf{y}_{k+1}}$$

Hence, we obtain

$$(1-\gamma_k)\min\left(\frac{\mathbf{x}_k}{\mathbf{y}_{k+1}}\right) \leq \min\left(\frac{\mathbf{x}_k+A^{-1}\mathbf{f}_k}{\mathbf{y}_{k+1}}\right) \leq (1+\gamma_k)\min\left(\frac{\mathbf{x}_k}{\mathbf{y}_{k+1}}\right),$$

then

$$\left|\min\left(\frac{\mathbf{x}_{k}+A^{-1}\mathbf{f}_{k}}{\mathbf{y}_{k+1}}\right)-\min\left(\frac{\mathbf{x}_{k}}{\mathbf{y}_{k+1}}\right)\right| \leq \gamma_{k}\min\left(\frac{\mathbf{x}_{k}}{\mathbf{y}_{k+1}}\right).$$
(17)

From (13), it follows that

$$\rho(A^{-1}) \le \widehat{\lambda}_{k+1} = \max\left(\frac{A^{-1}\mathbf{x}_{k+1}}{\mathbf{x}_{k+1}}\right) = \overline{\lambda}_k - \min\left(\frac{\mathbf{x}_k + A^{-1}\mathbf{f}_k}{\mathbf{y}_{k+1}}\right).$$
(18)

Combine (12), (18) and (17), then

$$\overline{\lambda}_{k+1} = \overline{\lambda}_{k} - (1 - \gamma_{k}) \min\left(\frac{\mathbf{x}_{k}}{\mathbf{y}_{k+1}}\right)$$

$$= \widehat{\lambda}_{k+1} + \min\left(\frac{\mathbf{x}_{k} + A^{-1}\mathbf{f}_{k}}{\mathbf{y}_{k+1}}\right) - (1 - \gamma_{k}) \min\left(\frac{\mathbf{x}_{k}}{\mathbf{y}_{k+1}}\right)$$

$$\geq \widehat{\lambda}_{k+1} + (1 - \gamma_{k} - (1 - \gamma_{k})) \min\left(\frac{\mathbf{x}_{k}}{\mathbf{y}_{k+1}}\right)$$

$$> \rho(A^{-1}).$$
(19)

By induction, $\{\overline{\lambda}_k\}$ is bounded below by $\rho(A^{-1})$, i.e.,

$$\overline{\lambda}_k > \rho(A^{-1})$$
 for all k .

From Lemma 3, since $\{\overline{\lambda}_k\}$ is a monotonically decreasing and bounded sequence, we must have $\lim_{k\to\infty} \overline{\lambda}_k = \alpha \ge \rho(A^{-1})$, where $\alpha = \rho(A^{-1})$ or $\alpha > \rho(A^{-1})$. We next investigate the case $\alpha > \rho(A^{-1})$, and present some basic results; this plays an important role later in proving the convergence of INI.

Lemma 4

For Algorithm 2, if $\overline{\lambda}_k$ is converge to $\alpha > \rho(A^{-1})$, then (i) $\|\mathbf{y}_k\|$ is bounded;(ii) $\lim_{k \to \infty} \min(\mathbf{x}_k) = 0$;(iii) $\sin \angle (\mathbf{x}_k, \mathbf{x}) \ge m > 0$ for some constant m > 0, where $\angle (\mathbf{x}_k, \mathbf{x})$ the acute angle of \mathbf{x}_k and \mathbf{x} .

Proof (i). From (16), we get

$$\|\mathbf{y}_{k+1}\| = \|\left(\overline{\lambda}_{k}I - A^{-1}\right)^{-1} \left(\mathbf{x}_{k} + A^{-1}\mathbf{f}_{k}\right)\| \le (1 + \gamma_{k}) \|(\overline{\lambda}_{k}I - A^{-1})^{-1}\|$$

= $2 \operatorname{sep}(\overline{\lambda}_{k}, A^{-1})^{-1} \le 2 \operatorname{sep}(\alpha, A^{-1})^{-1} < \infty.$ (20)

Copyright © 2010 John Wiley & Sons, Ltd. Prepared using nlaauth.cls

(ii). From (12) it follows that

$$\lim_{k \to \infty} \min\left(\frac{\mathbf{x}_k}{\mathbf{y}_{k+1}}\right) = \lim_{k \to \infty} \left(\frac{\overline{\lambda}_k - \overline{\lambda}_{k+1}}{1 - \gamma_k}\right) = 0.$$
(21)

On the other hand, from (20) and (21) we have

$$\min\left(\frac{\mathbf{x}_k}{\mathbf{y}_{k+1}}\right) \geq \frac{\min(\mathbf{x}_k)}{\max\left(\mathbf{y}_{k+1}\right)} \geq \frac{\min(\mathbf{x}_k)\operatorname{sep}(\alpha, A^{-1})}{2} > 0.$$

Thus, it is holds that

$$\lim_{k \to \infty} \min(\mathbf{x}_k) = 0. \tag{22}$$

(iii) Suppose there is a subsequence $\{\sin \angle (\mathbf{x}_{k_i}, \mathbf{x})\}$ which converges to zero, then

$$\lim_{j\to\infty}\widehat{\lambda}_{k_j} = \lim_{j\to\infty} \max\left(\frac{A^{-1}\mathbf{x}_{k_j}}{\mathbf{x}_{k_j}}\right) = \max\left(\lim_{j\to\infty}\frac{A^{-1}\mathbf{x}_{k_j}}{\mathbf{x}_{k_j}}\right) = \rho(A^{-1}).$$

By the definition of $\hat{\lambda}_k$, from (19) and (17), we have

$$\begin{aligned} \left| \widehat{\lambda}_{k} - \overline{\lambda}_{k} \right| &= \left| \min \left(\frac{\mathbf{x}_{k-1} + A^{-1} \mathbf{f}_{k-1}}{\mathbf{y}_{k}} \right) - (1 - \gamma_{k}) \min \left(\frac{\mathbf{x}_{k-1}}{\mathbf{y}_{k}} \right) \right| \\ &\leq \left| (1 + \gamma_{k} - (1 - \gamma_{k})) \min \left(\frac{\mathbf{x}_{k-1}}{\mathbf{y}_{k}} \right) \right| \text{ for any } k. \end{aligned}$$
(23)

From (21) and (23),

$$\lim_{i\to\infty}\overline{\lambda}_{k_j} = \lim_{j\to\infty} \left(\overline{\lambda}_{k_j} - \widehat{\lambda}_{k_j} + \widehat{\lambda}_{k_j}\right) = \rho(A^{-1}).$$

This is a contradiction.

Lemma 5 ([15])

Let $\mathbf{x} > \mathbf{0}$ be the unit length eigenvector of A associated with $\sigma_{\min}(A)$. For any vector $\mathbf{z} > \mathbf{0}$ with $\|\mathbf{z}\| = 1$, it holds that $\cos \angle (\mathbf{z}, \mathbf{x}) > \min(\mathbf{x})$ and

$$\inf_{\|\mathbf{z}\|=1,\mathbf{z}>\mathbf{0}} \cos \angle(\mathbf{z},\mathbf{x}) = \min(\mathbf{x}). \tag{24}$$

4. CONVERGENCE ANALYSIS FOR INI

In Sections 4.1–4.2, we will prove the global convergence and the convergence rate of INI. Furthermore, we will derive the explicit linear convergence factor and the superlinear convergence order with different γ_k .

4.1. Convergence Analysis

For an irreducible non-negative matrix A^{-1} , recall that the largest eigenvalue $\rho(A^{-1})$ of A^{-1} is simple. Let **x** be the unit length positive eigenvector corresponding to $\rho(A^{-1})$. Then for any orthogonal matrix $\begin{bmatrix} \mathbf{x} & V \end{bmatrix}$ it holds (cf. [10]) that

$$\begin{bmatrix} \mathbf{x}^T \\ V^T \end{bmatrix} A^{-1} \begin{bmatrix} \mathbf{x} & V \end{bmatrix} = \begin{bmatrix} \rho(A^{-1}) & \mathbf{c}^T \\ 0 & L \end{bmatrix}$$
(25)

with $L = V^T A^{-1}V$ whose eigenvalues constitute the other eigenvalues of A^{-1} . Therefore, we now define

$$\varepsilon_k = \overline{\lambda}_k - \rho(A^{-1}), \quad A_k = \overline{\lambda}_k I - A^{-1}.$$
 (26)

Copyright © 2010 John Wiley & Sons, Ltd. Prepared using nlaauth.cls

Similar to (25) we also have the spectral decomposition

$$\begin{bmatrix} \mathbf{x}^T \\ V^T \end{bmatrix} A_k \begin{bmatrix} \mathbf{x} & V \end{bmatrix} = \begin{bmatrix} \boldsymbol{\varepsilon}_k & 0 \\ 0 & L_k \end{bmatrix},$$
(27)

where $L_k = \overline{\lambda}_k I - L$. For $\overline{\lambda}_k \neq \rho(A^{-1})$, it is easy to verify that

$$\begin{bmatrix} \mathbf{x}^{T} \\ V^{T} \end{bmatrix} A_{k}^{-1} \begin{bmatrix} \mathbf{x} & V \end{bmatrix} = \begin{bmatrix} \boldsymbol{\varepsilon}_{k}^{-1} & \mathbf{b}_{k}^{T} \\ 0 & L_{k}^{-1} \end{bmatrix} \text{ with } \mathbf{b}_{k}^{T} = -\frac{\mathbf{c}^{T} L_{k}^{-1}}{\boldsymbol{\varepsilon}_{k}},$$
(28)

from which we get

$$A_{k}^{-1}V = \mathbf{x}\mathbf{b}_{k}^{T} + VL_{k}^{-1} = -\mathbf{x}\frac{\mathbf{c}^{T}L_{k}^{-1}}{\varepsilon_{k}} + VL_{k}^{-1},$$
(29)

$$A_k^{-1} = \varepsilon_k^{-1} \mathbf{x} \mathbf{x}^T - \varepsilon_k^{-1} \mathbf{x} \mathbf{c}^T L_k^{-1} V^T + V L_k^{-1} V^T, \qquad (30)$$

and

$$\mathbf{x}^T A_k^{-1} = \boldsymbol{\varepsilon}_k^{-1} \mathbf{x}^T - \boldsymbol{\varepsilon}_k^{-1} \mathbf{c}^T L_k^{-1} V^T.$$

Let $\{\mathbf{x}_k\}$ be generated by Algorithm 2. We decompose \mathbf{x}_k into the orthogonal direct sum by

$$\mathbf{x}_k = \mathbf{x}\cos(\boldsymbol{\varphi}_k) + \mathbf{p}_k\sin(\boldsymbol{\varphi}_k), \quad \mathbf{p}_k \in \operatorname{span}(V) \perp \mathbf{x}$$
(31)

with $\|\mathbf{p}_k\| = 1$ and $\varphi_k = \angle (\mathbf{x}_k, \mathbf{x})$ being the acute angle between \mathbf{x}_k and \mathbf{x} . So by definition, we have $\cos \varphi_k = \mathbf{x}^T \mathbf{x}_k$ and $\sin \varphi_k = \|V^T \mathbf{x}_k\|$. Evidently, $\mathbf{x}_k \to \mathbf{x}$ if and only if $\tan \varphi_k \to 0$, i.e., $\sin \varphi_k \to 0$. Since $\xi_k = \|\mathbf{f}_k\| \le \gamma_k \operatorname{sep}(0, A) \min(\mathbf{x}_k)$ in INI, it holds that $|A^{-1}\mathbf{f}_k| \le \gamma_k \mathbf{x}_k$. Therefore, we have

$$(1 - \gamma_k) \mathbf{x}_k \le \mathbf{x}_k + A^{-1} \mathbf{f}_k \le (1 + \gamma_k) \mathbf{x}_k.$$
(32)

As $A_k^{-1} \ge 0$, it follows from the above relation that

$$(1-\gamma_k)A_k^{-1}\mathbf{x}_k \le \mathbf{y}_{k+1} \le (1+\gamma_k)A_k^{-1}\mathbf{x}_k.$$
(33)

Using the above relation, we obtain

$$\tan \varphi_{k+1} = \frac{\sin \varphi_{k+1}}{\cos \varphi_{k+1}} = \frac{\|V^T \mathbf{x}_{k+1}\|}{\mathbf{x}^T \mathbf{x}_{k+1}} = \frac{\|V^T \mathbf{y}_{k+1}\|}{\mathbf{x}^T \mathbf{y}_{k+1}}$$

$$= \frac{\|V^T A_k^{-1} (\mathbf{x}_k + A^{-1} \mathbf{f}_k)\|}{\mathbf{x}^T A_k^{-1} (\mathbf{x}_k + A^{-1} \mathbf{f}_k)}$$

$$= \frac{\|L_k^{-1} V^T (\mathbf{x}_k + A^{-1} \mathbf{f}_k)\|}{(\varepsilon_k^{-1} \mathbf{x}^T - \varepsilon_k^{-1} \mathbf{c}^T L_k^{-1} V^T) (\mathbf{x}_k + A^{-1} \mathbf{f}_k)}$$

$$= \frac{\|L_k^{-1} V^T (\mathbf{x}_k + A^{-1} \mathbf{f}_k)\|}{\varepsilon_k^{-1} \mathbf{x}^T \mathbf{x}_k - \varepsilon_k^{-1} \mathbf{c}^T L_k^{-1} V^T \mathbf{x}_k + \varepsilon_k^{-1} \mathbf{x}^T A^{-1} \mathbf{f}_k - \varepsilon_k^{-1} \mathbf{c}^T L_k^{-1} V^T A^{-1} \mathbf{f}_k}$$

$$\leq \frac{\|L_k^{-1}\| \varepsilon_k (\sin \varphi_k + \|A^{-1} \mathbf{f}_k\|)}{\cos \varphi_k - \mathbf{c}^T L_k^{-1} V^T \mathbf{x}_k - \|A^{-1} \mathbf{f}_k\| - \|\mathbf{c}\|\|L_k^{-1}\|\|A^{-1} \mathbf{f}_k\|}.$$
(34)

Note that if we solve the inner linear system exactly, i.e., $f_k = 0$, we recover NI and get 1

$$\tan \varphi_{k+1} \le \frac{\|L_k^{-1}\| \varepsilon_k}{1 - \mathbf{c}^T L_k^{-1} V^T \mathbf{x}_k / \cos \varphi_k} \tan \varphi_k := \beta_k \tan \varphi_k.$$
(35)

Since L. Elsner [9] proved the quadratic convergence of the proposed Noda iteration, for k large enough we must have

$$\beta_k = O(\tan \varphi_k) \to 0.$$

Copyright © 2010 John Wiley & Sons, Ltd. Prepared using nlaauth.cls

It follows that

$$\beta_k < \beta < 1 \tag{36}$$

for any given positive constant $\beta < 1$. Therefore, we have

 $\tan \varphi_{k+1} < \beta \tan \varphi_k$

for $k \ge N$ with *N* large enough.

Theorem 3 (Main Theorem)

Let *A* be an irreducible monotone matrix. If the sequence $\{\overline{\lambda}_k\}$ is generated by INI with the relaxation strategies (11) and (12), then $\{\overline{\lambda}_k\}$ is monotonically decreasing and $\lim_{k\to\infty} \overline{\lambda}_k^{-1} = \sigma_{\min}(A)$.

Proof

From Lemma 3, the sequence $\{\overline{\lambda}_k\}$ is bounded and monotonically decreasing, and we must have either $\lim_{k\to\infty} \overline{\lambda}_k = \rho(A^{-1})$ or $\lim_{k\to\infty} \overline{\lambda}_k = \alpha > \rho(A^{-1})$. Next we prove by contradiction that, for INI, $\lim_{k\to\infty} \overline{\lambda}_k = \rho(A^{-1})$ must hold.

Suppose that $\lim_{k\to\infty} \overline{\lambda}_k = \alpha > \rho(A^{-1})$. It follows (iii) of Lemma 4 show that

$$\frac{1}{\cos\varphi_k} \le \frac{1}{\cos\varphi_k} \frac{\sin\varphi_k}{m} = \frac{\tan\varphi_k}{m}.$$
(37)

From (ii) of Lemma 4, we have

$$\lim_{k\to\infty}\min(\mathbf{x}_k)=0$$

This implies the inner tolerance $||f_k|| \to 0$, i.e., $||A^{-1}\mathbf{f}_k||$ is suitably small. In addition, by Lemma 5, it holds that $\cos \varphi_k$ is uniformly bounded below by $\min(\mathbf{x})$, therefore,

$$(1 + ||c|| ||L_k^{-1}) ||A^{-1}\mathbf{f}_k|| / \cos\varphi_k < 1 - \mathbf{c}^T L_k^{-1} V^T \mathbf{x}_k / \cos\varphi_k$$
(38)

for *k* large enough.

Using (34), (37) and (38), we obtain

$$\tan \varphi_{k+1} \leq \frac{\|L_k^{-1}\| \varepsilon_k \left(\tan \varphi_k + \|A^{-1}\mathbf{f}_k\| / \cos \varphi_k \right)}{1 - \mathbf{c}^T L_k^{-1} V^T \mathbf{x}_k / \cos \varphi_k - (1 + \|\mathbf{c}\| \|L_k^{-1}\|) \|A^{-1}\mathbf{f}_k\| / \cos \varphi_k} \\ \leq \frac{\|L_k^{-1}\| \varepsilon_k \left(\tan \varphi_k + \gamma_k \min(\mathbf{x}_k) \tan \varphi_k / m \right)}{1 - \mathbf{c}^T L_k^{-1} V^T \mathbf{x}_k / \cos \varphi_k - (1 + \|\mathbf{c}\| \|L_k^{-1}\|) \gamma_k \min(\mathbf{x}_k) \tan \varphi_k / m} \\ \leq \frac{\|L_k^{-1}\| \varepsilon_k \left(1 + \gamma_k \min(\mathbf{x}_k) / m \right)}{\left(1 - \mathbf{c}^T L_k^{-1} V^T \mathbf{x}_k / \cos \varphi_k \right) - (1 + \|\mathbf{c}\| \|L_k^{-1}\|) \gamma_k \min(\mathbf{x}_k) \tan \varphi_k / m} \tan \varphi_k$$

Define

$$\beta_k' = \frac{\|L_k^{-1}\|\boldsymbol{\varepsilon}_k(1+\gamma_k\min(\mathbf{x}_k)/m)}{\left(1-\mathbf{c}^T L_k^{-1} V^T \mathbf{x}_k/\cos\varphi_k\right) - (1+\|\mathbf{c}\|\|L_k^{-1}\|)\gamma_k\min(\mathbf{x}_k)\tan\varphi_k/m}$$

Note that β'_k is a continuous function with respect to $\min(\mathbf{x}_k)$ for $0 < \gamma_k < 1$. Then it holds that $\beta'_k \rightarrow \beta_k$ defined by (35) as $\min(\mathbf{x}_k) \rightarrow 0$. Therefore, from (36), for *k* large enough we can choose a sufficiently small δ such that

 $\beta_k' \leq (1+\delta)\beta_k < \beta < 1$

for $min(\mathbf{x}_k)$ sufficiently small. As a result, we have

$$an arphi_{k+1} \leq eta$$
 $an arphi_k$

for $k \ge N$ with N large enough and min(\mathbf{x}_k) sufficiently small. This means that $\tan \varphi_k \to 0$, i.e., $\sin \varphi_k \to \mathbf{0}$. From (iii) of Lemma 4, $\sin \varphi_k$ is uniformly bounded below by a positive constant. So $\sin \varphi_k \to \mathbf{0}$ and $\sin \varphi_k \ge m$, a contradiction. Therefore the initial assumption " $\lim_{k\to\infty} \overline{\lambda}_k = \alpha > \rho(A^{-1})$ "must be false.

Copyright © 2010 John Wiley & Sons, Ltd. Prepared using nlaauth.cls

4.2. Convergence Rates

Theorem 3 has proved the global convergence of INI, but the results are only qualitative and do not tell us anything about how fast the INI method converges. In this subsection, we will show the convergence rate of INI with different relaxation factors γ_k . More precisely, we prove that INI converges at least linearly with an asymptotic convergence factor bounded by $\frac{2\gamma}{1+\gamma}$ for $0 \le \gamma_k \le \gamma < 1$ and superlinearly for decreasing $\gamma_k = \frac{\overline{\lambda}_{k-1} - \overline{\lambda}_k}{\overline{\lambda}_{k-1}}$, respectively.

From (12) we have

$$\varepsilon_{k+1} = \varepsilon_k \left(1 - (1 - \gamma_k) \min\left(\frac{\mathbf{x}_k}{\varepsilon_k \mathbf{y}_{k+1}}\right) \right)$$

=: $\varepsilon_k \rho_k.$ (39)

Since $\overline{\lambda}_k - \overline{\lambda}_{k+1} < \overline{\lambda}_k - \rho(A^{-1})$, from (39) and (12), we have

$$\rho_k = 1 - (1 - \gamma_k) \min\left(\frac{\mathbf{x}_k}{\varepsilon_k \mathbf{y}_{k+1}}\right) = \left(1 - \frac{\overline{\lambda}_k - \overline{\lambda}_{k+1}}{\overline{\lambda}_k - \rho(A^{-1})}\right) < 1.$$

Theorem 4

For INI, we have $\rho_k \leq \frac{2\gamma_k}{1+\gamma_k} < 1$. Moreover, if $\gamma_k \leq \gamma < 1$, then $\lim_{k \to \infty} \rho_k \leq \frac{2\gamma}{1+\gamma} < 1$, i.e., the convergence of INI is at least globally linear. If $\lim_{k \to \infty} \gamma_k = 0$, then $\lim_{k \to \infty} \rho_k = 0$, that is, the convergence of INI is globally superlinear.

Proof From (32) and (33), we have

$$\min\left(\frac{\mathbf{x}_k}{\varepsilon_k \mathbf{y}_{k+1}}\right) \geq \min\left(\frac{\mathbf{x}_k}{(1+\gamma_k)\varepsilon_k A_k^{-1} \mathbf{x}_k}\right) = \frac{1}{1+\gamma_k} \min\left(\frac{\mathbf{x}_k}{\varepsilon_k A_k^{-1} \mathbf{x}_k}\right).$$

From (30), we get

$$\varepsilon_k A_k^{-1} \mathbf{x}_k = \mathbf{x} \mathbf{x}^T \mathbf{x}_k - \mathbf{x} \mathbf{c}^T L_k^{-1} \mathbf{V}^T \mathbf{x}_k + \varepsilon_k V L_k^{-1} V^T \mathbf{x}_k.$$
(40)

From Theorem 3, we know that $\lim_{k\to\infty} \mathbf{x}_k = \mathbf{x}$ and $\lim_{k\to\infty} \overline{\lambda}_k = \rho(A^{-1})$, from which it follows that $\varepsilon_k \to 0$ and $L_k^{-1} \to (\rho(A^{-1})I - L)^{-1}$. On the other hand, since $L_k^{-1} \to (\rho(A^{-1})I - L)^{-1}$ and $\lim_{k\to\infty} V^T \mathbf{x}_k = V^T \mathbf{x} = 0$, from (40) we get

$$\lim_{k\to\infty}\varepsilon_k A_k^{-1}\mathbf{x}_k = \mathbf{x}$$

Consequently, we obtain

$$egin{aligned} &\lim_{k o\infty}\min\left(rac{\mathbf{x}_k}{arepsilon_k\mathbf{y}_{k+1}}
ight) \geq rac{1}{1+\gamma_k}\min\left(\lim_{k o\infty}rac{\mathbf{x}_k}{arepsilon_kB_k^{-1}\mathbf{x}_k}
ight) \ &= rac{1}{1+\gamma_k}\min\left(rac{\mathbf{x}}{\mathbf{x}}
ight) = rac{1}{1+\gamma_k} > 0, \end{aligned}$$

leading to

$$\rho_k \le 1 - \frac{1 - \gamma_k}{1 + \gamma_k} = \frac{2\gamma_k}{1 + \gamma_k} < 1.$$

$$\tag{41}$$

		۰.
L		
L		

Copyright © 2010 John Wiley & Sons, Ltd. Prepared using nlaauth.cls

It can be seen from (41) that if γ_k is small then INI must ultimately converge quickly. Although Theorem 4 has established the superlinear convergence of INI, it does not reveal the convergence order. Our next concern is to derive the precise convergence order of INI. This is more informative and instructive because it lets us understand how fast INI converges.

Theorem 5

If the inner tolerance ξ_k in INI satisfies condition (11) with the relaxation factors

$$\gamma_k = \frac{\overline{\lambda}_{k-1} - \overline{\lambda}_k}{\overline{\lambda}_{k-1}},\tag{42}$$

then INI converges quadratically (asymptotically) in the form of

$$\overline{\varepsilon}_k \le 2\overline{\varepsilon}_{k-1}^2 \tag{43}$$

for *k* large enough, where the relative error $\overline{\varepsilon}_{k+1} = \varepsilon_k / \rho(A^{-1})$.

 $\frac{Proof}{\text{Since }\overline{\lambda}_{k-1}} > \overline{\lambda}_k > \rho(A^{-1}), \text{ we have }$

$$\gamma_{k} = \frac{\overline{\lambda}_{k-1} - \overline{\lambda}_{k}}{\overline{\lambda}_{k-1}} \le \frac{\overline{\lambda}_{k-1} - \rho(A^{-1})}{\rho(A^{-1})} = \frac{\varepsilon_{k-1}}{\rho(A^{-1})}.$$
(44)

From (39), (41) and (44), we have

$$\begin{split} \varepsilon_k &= \varepsilon_{k-1}\rho_{k-1} \leq \varepsilon_{k-1}\frac{2\gamma_k}{1+\gamma_k} = \varepsilon_{k-1}\frac{2}{1+\frac{1}{\gamma_k}} \\ &\leq \varepsilon_{k-1}\frac{2}{1+\frac{\rho(A^{-1})}{\varepsilon_{k-1}}} = \varepsilon_{k-1}^2\frac{2}{\varepsilon_{k-1}+\rho(A^{-1})} \\ &\leq \frac{2}{\rho(A^{-1})}\varepsilon_{k-1}^2. \end{split}$$

Dividing both sides of the above inequality by $\rho(A^{-1})$, we get

$$\overline{\varepsilon}_k = \frac{\varepsilon_k}{\rho(A^{-1})} \le \frac{2}{\rho(A^{-1})^2} \varepsilon_{k-1}^2 = 2\overline{\varepsilon}_{k-1}^2.$$

5. THE MODIFIED INEXACT NODA ITERATION

In this section, we propose a modified Noda iteration (MNI) for a non-negative matrix, and show that MNI and NI are equivalent. Thus, by combining INI (Algorithm 2) with MNI we can propose a modified inexact Noda iteration for a monotone matrix

5.1. The modified Noda iteration

When $\hat{\lambda}_k I - B$ tends to a singular matrix, the Noda iteration requires us to solve a possibly illconditioned linear system (3). Hence, we propose a rank one update technique for the ill-conditioned linear system (3), i.e.,

$$\begin{pmatrix} B - \hat{\lambda}_k I & -\mathbf{x}_k \\ -\mathbf{x}_k^T & 0 \end{pmatrix} \begin{pmatrix} \Delta \mathbf{y}_k \\ \delta_k \end{pmatrix} = \begin{bmatrix} (\hat{\lambda}_k I - B) \mathbf{x}_k \\ 0 \end{bmatrix},$$
(45)

Copyright © 2010 John Wiley & Sons, Ltd. Prepared using nlaauth.cls

where $\Delta \mathbf{y}_k = \mathbf{x}_{k+1} - \mathbf{x}_k$. Let $\mathbf{r}_k = (\hat{\lambda}_k I - B) \mathbf{x}_k$. In general, the linear system (45) is a well-conditioned linear system, unless *B* has the Jordan form corresponding to the largest eigenvalue, which contradicts the Perron–Frobenius theorem.

From (45),

$$0 = (B - \widehat{\lambda}_k I) (\mathbf{x}_{k+1} - \mathbf{x}_k) - \delta_k \mathbf{x}_k - \mathbf{r}_k$$

= $(B - \widehat{\lambda}_k I) \mathbf{x}_{k+1} - (B - \widehat{\lambda}_k I) \mathbf{x}_k - \delta_k \mathbf{x}_k - \mathbf{r}_k$
= $(B - \widehat{\lambda}_k I) \mathbf{x}_{k+1} - \delta_k \mathbf{x}_k.$

Hence, we have the following linear system

$$\left(\widehat{\lambda}_k I - B\right)\left(\frac{\mathbf{x}_{k+1}}{-\delta_k}\right) = \mathbf{x}_k,$$

or

 $\left[\widehat{\lambda}_k I - B\right] \mathbf{y}_{k+1} = \mathbf{x}_k,$

with $\mathbf{y}_k = \frac{-1}{\delta_k} \mathbf{x}_{k+1}$. Thus, from (3) and (45), we have the new iterative vector

$$\mathbf{x}_{k+1} = \frac{\mathbf{y}_{k+1}}{\|\mathbf{y}_{k+1}\|} = \frac{\mathbf{x}_k + \Delta \mathbf{y}_k}{\|\mathbf{x}_k + \Delta \mathbf{y}_k\|}.$$
(46)

This means the Noda iteration and the modified Noda iteration are mathematically equivalent. Based on (45) and (46), we state our algorithm as follows.

Algorithm 3 Modified Noda Iteration (MNI)

1. Given $\widehat{\lambda}_0$, $\mathbf{x}_0 > \mathbf{0}$ with $\|\mathbf{x}_0\| = 1$ and tol > 0. 2. for $k = 0, 1, 2, \ldots$ if $||B\mathbf{x}_{k+1} - \widehat{\lambda}_k \mathbf{x}_{k+1}|| > \sqrt{\mathsf{tol}}$ 3. 4. Solve $(\widehat{\lambda}_k I - B) \mathbf{y}_{k+1} = \mathbf{x}_k$. 5. Normalize the vector $\mathbf{x}_{k+1} = \mathbf{y}_{k+1} / ||\mathbf{y}_{k+1}||$. 6. else if Solve $\begin{pmatrix} B - \widehat{\lambda}_k & -\mathbf{x}_k \\ -\mathbf{x}_k^T & 0 \end{pmatrix} \begin{pmatrix} \Delta \mathbf{y}_k \\ \delta_k \end{pmatrix} = \begin{bmatrix} \widehat{\lambda}_k \mathbf{x}_k - B\mathbf{x}_k \\ 0 \end{bmatrix}.$ 7. 8. Normalize the vector $\mathbf{x}_{k+1} = (\mathbf{x}_k + \Delta \mathbf{y}_k) / \|\mathbf{x}_k + \Delta \mathbf{y}_k\|$ 9. end Compute $\widehat{\lambda}_{k+1} = \max\left(\frac{B\mathbf{x}_{k+1}}{\mathbf{x}_{k+1}}\right)$. 10. 11. **until** convergence: $||B\mathbf{x}_{k+1} - \lambda_k \mathbf{x}_{k+1}|| < \text{tol}.$

Note that the sequence $\{\hat{\lambda}_k I - B\}$ tends to a singular matrix, meaning that $\{\hat{\lambda}_k\}$ converges to an eigenvalue of *B*, and (3) becomes an ill-conditioned linear system. Based on practical experiments, we propose taking $\|B\mathbf{x}_{k+1} - \hat{\lambda}_k \mathbf{x}_{k+1}\| \le \sqrt{\text{tol}}$ and switching from (3) to (45).

5.2. The modified inexact Noda iteration

For a monotone matrix A, we replaced B by A^{-1} in (45). The linear system (45) can be rewritten as

$$\begin{pmatrix} I - \widehat{\lambda}_k A & -A\mathbf{x}_k \\ -\mathbf{x}_k^T & 0 \end{pmatrix} \begin{pmatrix} \Delta \mathbf{y}_k \\ \delta_k \end{pmatrix} = \begin{bmatrix} \widehat{\lambda}_k A \mathbf{x}_k - \mathbf{x}_k \\ 0 \end{bmatrix}.$$
 (47)

Based on MNI, by combining INI (Algorithm 2) with equation (47), we can propose a modified inexact Noda iteration for a monotone matrix, which is described as Algorithm 4.

Algorithm 4 Modified Inexact Noda Iteration (MINI)

1. Given $\widehat{\lambda}_0$, $\mathbf{x}_0 > \mathbf{0}$ with $\|\mathbf{x}_0\| = 1$ and tol > 0. 2. for $k = 0, 1, 2, \ldots$ $\mathbf{if} = \|A\mathbf{x}_{k+1} - \overline{\lambda}_k^{-1}\mathbf{x}_{k+1}\| > \sqrt{\mathsf{tol}}$ 3. Run INI for monotone matrix A (Algorithm 2). 4. 5. else if Solve $\begin{pmatrix} I - \overline{\lambda}_k A & -A \mathbf{x}_k \\ -\mathbf{x}_k^T & 0 \end{pmatrix} \begin{pmatrix} \Delta \mathbf{y}_k \\ \delta_k \end{pmatrix} = \begin{bmatrix} \overline{\lambda}_k A \mathbf{x}_k - \mathbf{x}_k \\ 0 \end{bmatrix}$ exactly. 6. 7. Normalize the vector $\mathbf{x}_{k+1} = (\mathbf{x}_k + \Delta \mathbf{y}_k) / \|\mathbf{x}_k + \Delta \mathbf{y}_k\|$. Compute $\overline{\lambda}_{k+1}$ that satisfies condition (12). 8. 9. end 10. **until** convergence: $||A\mathbf{x}_{k+1} - \overline{\lambda}_k^{-1}\mathbf{x}_{k+1}|| < \text{tol.}$

6. NUMERICAL EXPERIMENTS

In this section we present numerical experiments to support our theoretical results for INI, and to illustrate the effectiveness of the proposed MINI algorithms. All numerical tests were performed on an Intel (R) Core (TM) i7 CPU 4770@ 3.4GHz with 16 GB memory using Matlab R2013*a* with the machine precision $\varepsilon = 2.22 \times 10^{-16}$ under the Microsoft Windows 7 64-bit.

 I_{outer} denotes the number of outer iterations to achieve the convergence, and I_{inner} denotes the total number of inner iterations, which measures the overall efficiency of MNI and MINI. In view of the above, we have the average number $I_{ave} = I_{inner}/I_{outer}$ at each outer iteration for our test algorithms. In the tables, "Positivity"illustrates whether the converged Perron vector preserves the strict positivity property. If "No", then the percentage in the brace indicates the proportion that the converged Perron vector has the positive components. We also report the CPU time of each algorithm, which measures the overall efficiency too.

6.1. INI for computing the smallest eigenvalue of a monotone matrix

We present an example to illustrate the numerical behavior of NI, INI_1 and INI_2 for monotone matrices. The approximate solution y_{k+1} of (14) satisfies

$$(\overline{\lambda}_k A - I) \mathbf{y}_{k+1} = A \mathbf{x}_k + \mathbf{f}_k$$

by requiring the following inner tolerances:

- for NI: $\|\mathbf{f}_k\| \le 10^{-14}$;
- for INL1: $\|\mathbf{f}_k\| \leq \gamma_k \operatorname{sep}(0, A) \min(\mathbf{x}_k)$ with some $0 < \gamma_k < 1$;

• for INI_2:
$$\|\mathbf{f}_k\| \leq \frac{\overline{\lambda}_{k-1} - \overline{\lambda}_k}{\overline{\lambda}_{k-1}} \operatorname{sep}(0, A) \min(\mathbf{x}_k)$$
 for $k \geq 1$ and $\overline{\lambda}_0 > \rho(A^{-1})$.

We use the minimal residual method to solve the inner linear systems. For the implementations, we use the standard Matlab function minres. The outer iteration starts with the normalized vector of $(1, ..., 1)^T$, and the stopping criterion for outer iterations is

$$\frac{\|A\mathbf{x}_k - \overline{\lambda}_k^{-1}\mathbf{x}_k\|}{(\|A\|_1 \|A\|_{\infty})^{1/2}} \le 10^{-10},$$

where $\|\cdot\|_1$ and $\|\cdot\|_{\infty}$ are the one norm and the infinity norm of a matrix, respectively.

Condition (11) ensures that the eigenvector in Lemma 3 does indeed preserve the strict positivity property. However, the formula in (11) is not applicable in practice, because it uses sep(0,A), which is unknown at the time it needs to be computed. Therefore, for practical implementations, we suggest a relaxation strategy to replace sep(0,A) by $\overline{\lambda_k}^{-1}$. The quantity $\overline{\lambda_k}^{-1}$ is related to the lower



Figure 1. The outer residual norms versus outer iterations in Examples 1.

Table I. The total outer and inner iterations in Example 1

Method	Iouter	I _{inner}	Iave	CPU time	Positivity
INI_1 with $\gamma = 0.8$	51	19622	384	76	Yes
INI_1 with $\gamma = 0.5$	18	11233	624	38	Yes
NI	5	3621	724	25	Yes
INI_2	5	3591	718	19	Yes

bound of the smallest eigenvalue of A, i.e., $\sigma_{min}(A) \ge \overline{\lambda}_k^{-1}$. For all examples, the stopping criterion for the inner iteration is set at

$$\|\mathbf{f}_k\| \leq \max\{\gamma_k \min(\mathbf{x}_k)/\overline{\lambda}_k, 10^{-13}\}$$
 for INI_1

and

$$\|\mathbf{f}_k\| \le \max\{\frac{\overline{\lambda}_{k-1} - \overline{\lambda}_k}{\overline{\lambda}_{k-1}\overline{\lambda}_k}\min(\mathbf{x}_k), 10^{-13}\} \text{ for INI_2}.$$

Example 1

We consider the finite-element discretization of the boundary value problem in [3, Example 4.2.4]

$$\begin{aligned} -u_{xx} - u_{yy} &= g(x, y) \quad \text{in } \Omega = [0, a] \times [0, b], \\ a, b &> 0, \qquad u = f(x, y) \text{ on } \partial \Omega, \end{aligned}$$

using piecewise quadratic basis functions on the uniform mesh of $p \times m$ isosceles right-angled triangles. This is a matrix of order n = (2p-1)(2m-1) = 127,041 with p = 400 and m = 80.

For Example 1, we see that, for this monotone matrix eigenproblem, INI_1, with two different $\gamma_k = 0.5$ and 0.8 exhibits distinct convergence behaviors and uses 51 and 18 outer iterations to achieve the desired accuracy, respectively. As Figure 1 indicates, NI and INI_2 typically converge superlinearly, and INI_1 with $\gamma_k = 0.5, 0.8$ typically converge linearly. This confirms our theory and demonstrates that the results of our theorem can be realistic and pronounced.

We observe from Table I that all the converged eigenvectors are positive, and that INI_2 improves the overall efficiency of NI. As we see, the INI_1 algorithm converges linearly and slowly. To be precise, INI_1 needs between twice and three times the CPU time of INI_2, but I_{ave} for INI_1 is only half I_{ave} of INI_2. There are two reasons for this. First, since the approximate eigenvalues are obtained from the relation (12), then the parameter γ_k will lead to a difference in the convergence rates, as is seen from Figure 1. Second, from (11), INI_2 solves the inner linear systems more and more accurately as k increases . In contrast, the inner tolerance used by INI_1 is fixed except for the factor min(\mathbf{x}_k), which also makes the average number of the iterations of INI_1 only about half of those for INI_2.

6.2. MINI for computing the smallest singular value of an M-matrix

In the above section, INI_2 was considerably better than NI and INI_1 for overall efficiency. Therefore, in this subsection, we use MINI (INI_2 combined with MNI) to find the smallest singular value and the associated eigenvector of an *M*-matrix, and confirm the effectiveness of MINI and the theory we presented in Sections 3 and 4. For MINI, the stopping criteria for inner and outer iterations are the same as those for monotone matrices. In the meantime , we compare MINI with the algorithms JDQR [27] and JDSVD [11] and the Matlab function svds; none of these are positivity preserving for approximate eigenvectors. We show that the MINI algorithm always reliably computes positive eigenvectors, while the other algorithms generally fail to do so.

Since JDQR and JDSVD use the absolute residual norms to decide the convergence, then we set the stopping criteria "TOL= $10^{-10}(||A||_1 ||A||_{\infty})^{1/2}$ " for outer iterations, and then we will get the same stopping criteria as used for MINI. We set the parameters "sigma=SM" for JDQR, "opts.target=0" for JDSVD, and the inner solver "OPTIONS.LSolver=minres". All the other options use defaults. We do not use any preconditioning for inner linear systems. For svds, we set the stopping criteria "OPTS.tol= $10^{-10}(||A||_1 ||A||_{\infty})^{1/2}$, and take the maximum and minimum subspace dimensions as 20 and 2 at each restart, respectively.

Suppose that we want to compute the smallest singular value, and the corresponding singular vector, of a real $n \times n$ M-matrix M. This partial SVD can be computed by using equivalent eigenvalue decomposition, that is, the augmented matrix

$$A = \left[\begin{array}{cc} 0 & M \\ M^T & 0 \end{array} \right]$$

Obviously, such a matrix A is no longer an M-matrix but will indeed be monotone.

Example 2

We consider a symmetric *M*-matrix of the form $M = \sigma I - B$, where *B* is the non-negative matrix rgg_n_2_19_s0 from the DIMACS10 test set [8]. This matrix is a random geometric graph with 2^{19} vertices. Each vertex is a random point in the unit square and edges connect vertices whose Euclidean distance is below 0.55 $(\log(n)/n)$. This threshold is chosen in order to ensure that the graph is almost connected. This matrix is a binary matrix with $n = 2^{19} = 524,288$ and 6,539,532 nonzero entries.

For this problem, MINI works very well and uses only six outer iterations to attain the desired accuracy. Furthermore, it is reliable and positivity preserving. In contrast, JDQR, JDSVD, and svds compute the desired eigenvalue, but the converged eigenvectors are not positive. More precisely, Table II indicates that for these algorithms roughly 50% of the components of each converged eigenvector are negative.

As far as overall efficiency is concerned, MINI is the most efficient in terms of I_{inner} , I_{outer} and the CPU time. JDQR and svds require at least five times the CPU time of MINI; they are also more expensive than JDSVD in terms of the CPU time.

Method	Iouter	I _{inner}	Iave	CPU time	Positivity
MINI	6	331	55	30	Yes
JDQR	25	4068	162	243	No (52%)
JDSVD	34	1432	42	58	No (51%)
svds	140		140	144	No (57%)

Table II. The total outer and inner iterations in Example 2

7. CONCLUSIONS

We have proposed an inexact Noda iteration method for computing the smallest eigenpair of a large irreducible monotone matrix, and have considered the convergence of the modified inexact

Noda iteration with two relaxation factors. We have proved that the convergence of INI is globally linear and superlinear, with the asymptotic convergence factor bounded by $\frac{2\gamma_k}{1+\gamma_k}$. More precisely, the modified inexact Noda iteration with inner tolerance $\xi_k = ||\mathbf{f}_k|| \le \gamma_k \operatorname{sep}(0, A) \min(\mathbf{x}_k)$ converges at least linearly if the relaxation factors meet the condition $\gamma_k \le \gamma < 1$, and superlinearly if the relaxation factors meet the condition $\gamma_k \le \gamma < 1$, and superlinearly if the relaxation factors meet the condition $\gamma_k = \frac{\overline{\lambda_{k-1} - \overline{\lambda_k}}}{\overline{\lambda_{k-1}}}$, respectively. The results for INI clearly show how the accuracy of the inner iterations affects the convergence of the outer iterations.

In the experiments, we also compared MINI with Jacobi–Davidson type methods (JDQR, JDSVD) and the implicitly restarted Arnoldi method (svds). The contribution of this paper is twofold. First, MINI always preserves the positivity of approximate eigenvectors, while the other three methods often fail to do so. Second, the proposed MINI algorithms have been shown to be practical and effective for large monotone matrix eigenvalue problems and *M*-matrix singular value problems.

REFERENCES

- 1. A. S. ALFA, J. XUE AND Q. YE, Accurate computation of the smallest eigenvalue of a diagonally dominant *M*-matrix, Math. Comp., 71, pp. 217–236 (2002).
- J. ABATE, L. CHOUDHURY AND W. WHITT, Asymptotics for steady-state tail probabilities in structured Markov queueing models, Commun. Statist-Stochastic Models, 1, pp. 99–143 (1994)
- O. AXELSSON, L. KOLOTILINA, Monotonicity and discretization error estimates, SIAM J.Numer.Anal., 27 (1990), 1591-1611.
- 4. O.AXELSSON, S.V.GOLOLOBOV, Monotonicity and discretization error estimates for convection-diffusion problems, Technical Report, Department of mathematic university of Nijmegen (2003).
- A. BERMAN AND R. J. PLEMMONS, Non-negative Matrices in the Mathematical Sciences, Vol. 9 of Classics in Applied Mathematics, SIAM, Philadelphia, PA (1994)
- 6. J. BERNS-MÜLLER, I. G. GRAHAM AND A. SPENCE, *Inexact inverse iteration for symmetric matrices*, Linear Algebra Appl., 416, pp. 389–413 (2006)
- 7. L. COLLATZ, Numerical Treatment of Differential Equations, 3rd ed., Springer, berlin, (1960)
- DIMACS10 test set and the University of Florida Sparse Matrix Collection. Available at www.cise.ufl.edu/research/sparse/dimacs10/.
- 9. L. ELSNER, Inverse iteration for calculating the spectral radius of a non-negative irreducible matrix, Linear Algebra Appl., 15, pp. 235–242 (1976)
- G. H. GOLUB AND C. F. VAN LOAN, *Matrix Computations*, The John Hopkins University Press, Baltimore, London, 4th ed. (2012)
- 11. M.E. HOCHSTENBACH, A Jacobi-Davidson type SVD method, SIAM J. Sci. Comp. 23(2), pp. 606-628 (2001)
- 12. R. A. HORN AND C. R. JOHNSON, *Matrix Analysis*, The Cambridge University Press, Cambridge, UK (1985)
- Z. JIA, On convergence of the inexact Rayleigh quotient iteration with MINRES, J. Comput. Appl. Math., 236, pp. 4276–4295 (2012)
- 14. ____, On convergence of the inexact Rayleigh quotient iteration with the Lanczos method used for solving linear systems, technical report, September 2011. http://arxiv.org/pdf/0906.2239v4.pdf.
- Ž. JIA, W.-W. LIN, AND C.-S. LIU, A positivity preserving inexact Noda iteration for computing the smallest eigenpair of a large irreducible M-matrix, Numer. Math., accepted, 2014.
- C. LEE, Residual Arnoldi method: theory, package and experiments, Ph D thesis, TR-4515, Department of Computer Science, University of Maryland at College Park, (2007)
- Y.-L. LAI, K.-Y. LIN, AND W.-W. LIN, An inexact inverse iteration for large sparse eigenvalue problems, Numer. Linear Algebra Appl., 4 (1997), pp. 425–437.
- T. NODA, Note on the computation of the maximal eigenvalue of a non-negative irreducible matrix, Numer. Math., 17 (1971), pp. 382–386.
- D. D. OLESKY AND P. VAN DEN DRIESSCHE, Monotone positive stable matrices, Linear Algebra Appl., Volume 48, pp. 381–401 (1983).
- A. M. OSTROWSKI, On the convergence of the Rayleigh quotient iteration for the computation of the characteristic roots and vectors. V. (Usual Rayleigh quotient for non-Hermitian matrices and linear elementary divisors), Arch. Rational Mech. Anal., 3 (1959), pp. 472–481.
- 21. B. N. PARLETT, *The Symmetric Eigenvalue Problem*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1998.
- Y. SAAD, Numerical Methods for Large Eigenvalue Problems, Manchester University Press, Manchester, UK, 1992.
- J. SCHROEDER, M-Matrices and generalizations using an operator theory approach, SIAM Rev. 20, 213-244 (1978).
- P. SHIVAKUMAR, J. WILLIAMS, Q. YE AND C. MARINOV, On two-sided bounds related to weakly diagonally dominant M-matrices with applications to digital circuit dynamics, SIAM J. Matrix Anal. Appl., 17, pp. 298–312 (1996)
- K.C. SIVAKUMAR, A new characterization of nonnegativity of Moore–Penrose inverses of Gram operators, Positivity 13 (1) (2009), 277–286.

- G. L. G. SLEIJPEN AND H. A. VAN DER VORST, A Jacobi-Davidson iteration method for linear eigenvalue problems, SIAM J. Matrix Anal. Appl., 17 (1996), pp. 401-425.
 G. L. G. SLEIJPEN, JDQR and JDQZ codes. Available at http://www.math.uu.nl/people/sleijpen
 G. W. STEWART, Matrix Algorithms. Vol. II, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PM. 2001 PA, 2001.
- 29. J. XUE, Computing the smallest eigenvalue of an M-matrix, SIAM J. Matrix Anal. Appl., 17 (1996), pp. 748–762.