# On the number of edges in a graph with no (k + 1)-connected subgraphs

Anton Bernshteyn \* Alexandr Kostochka †

#### Abstract

Mader proved that for  $k \geq 2$  and  $n \geq 2k$ , every n-vertex graph with no (k+1)-connected subgraphs has at most  $(1+\frac{1}{\sqrt{2}})k(n-k)$  edges. He also conjectured that for n large with respect to k, every such graph has at most  $\frac{3}{2}\left(k-\frac{1}{3}\right)(n-k)$  edges. Yuster improved Mader's upper bound to  $\frac{193}{120}k(n-k)$  for  $n \geq \frac{9k}{4}$ . In this note, we make the next step towards Mader's Conjecture: we improve Yuster's bound to  $\frac{19}{12}k(n-k)$  for  $n \geq \frac{5k}{2}$ .

Mathematics Subject Classification: 05C35, 05C40 Key words and phrases: Average degree, connectivity, k-connected subgraphs.

## 1 Introduction

All graphs considered here are finite, undirected, and simple. For a graph G, V(G) and E(G) denote its vertex set and edge set respectively. If  $U \subseteq V(G)$ , then G[U] denotes the induced subgraph of G whose vertex set is U, and  $G - U := G[V(G) \setminus U]$ . For  $v \in V(G)$ ,  $N(v) := \{u \in V(G) : uv \in E(G)\}$  denotes the neighborhood of v in G.

Let  $k \in \mathbb{N}$ . Recall that a graph G is (k+1)-connected if, for every set  $S \subset V(G)$  of size k, the graph  $G[V(G) \setminus S]$  is connected and contains at least two vertices (so  $|V(G)| \ge k+2$ ). Mader [1] posed the following question:

What is the maximum possible number of edges in an n-vertex graph that does not contain a (k + 1)-connected subgraph?

It is easy to see that for k = 1 the answer is n - 1: every tree on n vertices contains n - 1 edges and no 2-connected subgraphs, whereas every graph on n vertices with at least n edges contains a cycle, and cycles are 2-connected. Thus for the rest of the note we will assume  $k \ge 2$ .

The following construction due to Mader [2] gives an example of a graph with no (k+1)connected subgraphs and a large number of edges. Fix k and n, and suppose that n = kq + r,
where  $1 \le r \le k$ . The graph  $G_{n,k}$  has vertex set  $\bigcup_{i=0}^q V_i$ , where the sets  $V_0, \ldots, V_q$  are pairwise
disjoint and satisfy the following contitions.

<sup>\*</sup>Department of Mathematics, University of Illinois at Urbana-Champaign, IL, USA, bernsht2@illinois.edu. Research of this author is supported by the Illinois Distinguished Fellowship.

<sup>&</sup>lt;sup>†</sup>Department of Mathematics, University of Illinois at Urbana–Champaign, IL, USA and Sobolev Institute of Mathematics, Novosibirsk 630090, Russia, kostochk@math.uiuc.edu. Research of this author is supported in part by NSF grant DMS-1266016 and by grant 15-01-05867 of the Russian Foundation for Basic Research.

- 1.  $|V_0| = \ldots = |V_{q-1}| = k$ , while  $|V_q| = r$ .
- 2.  $V_0$  is an independent set in  $G_{n,k}$ .
- 3. For  $1 \leq i \leq q$ ,  $V_i$  is a clique in  $G_{n,k}$ .
- 4. Every vertex in  $V_0$  is adjacent to every vertex in  $\bigcup_{i=1}^q V_i$ .
- 5.  $G_{n,k}$  has no other edges.

Note that  $V_0$  is a separating set of size k and every component of  $G_{n,k} - V_0$  has at most k vertices. It follows that  $G_{n,k}$  has no (k+1)-connected subgraphs. A direct calculation shows that  $G_{n,k}$  has at most  $\frac{3}{2}\left(k-\frac{1}{3}\right)(n-k)$  edges, where the equality holds if n is a multiple of k. Mader [2] conjectured that this example is, in fact, best possible.

**Conjecture 1** (Mader [2]). Let  $k \geq 2$ . Then for n sufficiently large, the number of edges in an n-vertex graph without a (k+1)-connected subgraph cannot exceed  $\frac{3}{2}\left(k-\frac{1}{3}\right)(n-k)$ .

Mader himself proved Conjecture 1 for  $k \leq 6$ . Moreover, he showed that for all k, the weaker version of the conjecture, where the coefficient  $\frac{3}{2}$  is replaced by  $1 + \frac{1}{\sqrt{2}}$ , holds. Yuster [4] improved this result by showing that the coefficient can be taken to be  $\frac{193}{120}$ .

**Theorem 2** (Yuster [4]). Let  $k \geq 2$  and  $n \geq \frac{9k}{4}$ . Then every n-vertex graph G with  $|E(G)| > \frac{193}{120}k(n-k)$  contains a (k+1)-connected subgraph.

Here we improve Yuster's bound, obtaining the value  $\frac{19}{12}$  for the coefficient.

It turns out that for this problem, computations work out nicer if we "normalize" vertex and edge counts by assigning a weight  $\frac{1}{k}$  to each vertex and a weight  $\frac{1}{k^2}$  to each edge in a graph. Using this terminology, we can restate Conjecture 1 in the following way.

**Conjecture 1'.** Let  $k \geq 2$ . Then for  $\gamma$  sufficiently large, every graph G with  $\frac{1}{k}|V(G)| = \gamma$  and  $\frac{1}{k^2}|E(G)| > \frac{3}{2}(\gamma - 1)$  contains a (k + 1)-connected subgraph.

Our main result in these terms is as follows.

**Theorem 3.** Let  $k \geq 2$ . Then every graph G with  $\frac{1}{k}|V(G)| = \gamma \geq \frac{5}{2}$  and  $\frac{1}{k^2}|E(G)| > \frac{19}{12}(\gamma - 1)$  contains a (k+1)-connected subgraph.

We follow the ideas of Mader and Yuster: Use induction on the number of vertices for graphs with at least  $\frac{5}{2}k$  vertices. The hardest part is to prove the case when after deleting a separating set of size k, exactly one of the components of the remaining graph has fewer than  $\frac{3}{2}k$  vertices, since the induction assuption does not hold for  $n < \frac{5}{2}k$ . New ideas in the proof are in Lemmas 8, 9, and 10 below.

## 2 Proof of Theorem 3

We want to derive a linear in (n-k) bound on the number of edges in a graph that does not contain (k+1)-connected subgraphs. But the bound becomes linear only for graphs with large number of vertices; while for small graphs the dependency is quadratic in n-k. The main difficulties we encounter are around the transition between the quadratic and linear regimes. To deal with small n, we use the following lemma due to Matula [3], whose bound is asymptotically exact for n < 2k.

**Lemma 4** (Matula [3]). Let  $k \ge 2$ . Then every graph G with  $|V(G)| = n \ge k+1$  and  $|E(G)| > \binom{n}{2} - \frac{1}{3}((n-k)^2 - 1)$  contains a (k+1)-connected subgraph.

We will use the following "normalized" version of this lemma.

**Lemma 4'.** Let  $k \geq 2$ . Then every graph G with  $\frac{1}{k}|V(G)| = \gamma > 1$ , and

$$\frac{1}{k^2}|E(G)| > \frac{1}{6}\left(\gamma^2 + 4\gamma - 2\right) \tag{1}$$

contains a (k+1)-connected subgraph.

*Proof.* Indeed, (1) yields

$$|E(G)| > \frac{k^2}{6} \left( \gamma^2 + 4\gamma - 2 \right)$$

$$= \binom{\gamma k}{2} - \frac{1}{3} ((\gamma k - k)^2 - 1) + \frac{\gamma k}{2} - \frac{1}{3}$$

$$> \binom{\gamma k}{2} - \frac{1}{3} ((\gamma k - k)^2 - 1),$$

and we are done by original Matula's lemma.

From now on, fix a graph G with  $\frac{1}{k}|V(G)| = \gamma \ge \frac{5}{2}$  and  $\frac{1}{k^2}|E(G)| > \frac{19}{12}(\gamma - 1)$ , and suppose for contradiction that G does not contain a (k+1)-connected subgraph. Choose G to have the least possible number of vertices (so we can apply induction hypothesis for subgraphs of G). Since G itself is not (k+1)-connected, it contains a separating set  $S \subset V(G)$  of size K. Let  $K \subset V(G) \setminus S$  be such that  $K \subset K$  is a smallest connected component of  $K \subset K$ , and let  $K \subset K$  is a smallest connected component of  $K \subset K$  and let  $K \subset K$  is a smallest connected component of  $K \subset K$ .

We start by showing that the graph G cannot be too small, using Matula's Lemma.

#### Lemma 5. $\gamma > 3$ .

*Proof.* Suppose that  $\gamma \leq 3$ . Then, by Lemma 4',

$$0 \le \frac{1}{k^2} |E(G)| - \frac{19}{12} (\gamma - 1) \le \frac{1}{6} \left( \gamma^2 + 4\gamma - 2 \right) - \frac{19}{12} (\gamma - 1) = \frac{1}{12} (2\gamma^2 - 11\gamma + 15). \tag{2}$$

The function  $g(\gamma) = 2\gamma^2 - 11\gamma + 15$  on the right-hand side of (2) is convex in  $\gamma$ . Hence it is maximized on the boundary of the interval  $[\frac{5}{2}; 3]$ . But it is easy to check that  $g(\frac{5}{2}) = g(3) = 0$ , hence it is nonpositive on the whole interval. Therefore,  $\gamma > 3$ .

All the edges in G either belong to the graph  $G[S \cup B]$ , or are incident to the vertices in A. The number of edges in  $G[S \cup B]$  can be bounded either using Matula's lemma (which is efficient for  $\beta \leq \frac{3}{2}$ ) or using the induction hypothesis (which can be applied if  $\beta > \frac{3}{2}$ ). Hence the difficulty is in bounding the number of edges incident to the vertices in A.

The first step is to show that A cannot be too large, because otherwise we can use induction.

## Lemma 6. $\alpha < \frac{3}{2}$ .

*Proof.* If  $\alpha \geq \frac{3}{2}$ , then we can apply the induction hypothesis both for  $G[S \cup A]$  and for  $G[S \cup B]$ , and thus obtain

 $\frac{1}{k^2}|E(G)| \le \frac{19}{12}\alpha + \frac{19}{12}\beta = \frac{19}{12}(\alpha + \beta) = \frac{19}{12}(\gamma - 1).$ 

The next lemma shows that A cannot be too small either, since otherwise the total number of edges between the vertices in A and the vertices in  $S \cup A$  is small.

#### Lemma 7. $\alpha > 1$ .

*Proof.* Suppose that  $\alpha \leq 1$ . Then  $\beta > 1$ , since  $\alpha + \beta + 1 = \gamma > 3$ . If  $\beta \geq \frac{3}{2}$ , then using the induction hypothesis for  $G[S \cup B]$ , we get

$$\frac{1}{k^2}|E(G)| \le \frac{1}{2}\alpha^2 + \alpha + \frac{19}{12}\beta \le \frac{3}{2}\alpha + \frac{19}{12}\beta < \frac{19}{12}(\alpha + \beta) = \frac{19}{12}(\gamma - 1).$$

Thus  $\beta < \frac{3}{2}$ . Therefore,  $\alpha > \frac{1}{2}$ . In this case, applying Lemma 4' to  $G[S \cup B]$  reduces the problem to proving the inequality

$$\frac{1}{2}\alpha^2 + \alpha + \frac{1}{6}\left((\beta+1)^2 + 4(\beta+1) - 2\right) \le \frac{19}{12}(\alpha+\beta),$$

which is equivalent to

$$6\alpha^2 + 2\beta^2 - 7\alpha - 7\beta + 6 \le 0. (3)$$

For  $\alpha$  fixed, the left-hand side of (3) is monotone decreasing in  $\beta$  when  $\beta < \frac{7}{4}$ , so its maximum is attained at  $\beta = 1$ . Thus (3) will hold if the function  $g_1(\alpha) = 6\alpha^2 - 7\alpha + 1$  is nonpositive. Since  $g_1(\alpha)$  is a convex function, its maximum on the interval  $\left[\frac{1}{2};1\right]$  is attained at one of the boundary points. We have

$$g_1\left(\frac{1}{2}\right) = 6 \cdot \left(\frac{1}{2}\right)^2 - 7 \cdot \frac{1}{2} + 1 = -1 < 0,$$
 and  $g_1(1) = 6 \cdot 1^2 - 7 \cdot 1 + 1 = 0.$ 

So we know that  $1 < \alpha < \frac{3}{2}$ . How can we bound the number of edges incident to the vertices in A? The tricks of Lemma 7 and of Lemma 4' are not sufficient here. The idea is to combine them by applying Lemma 4' only to the graph  $G[A \cup (S \setminus S')]$ , where S' is a subset of S with relatively few edges between A and S'. To obtain such set S', we will use Lemma 8 below, which asserts that there are many vertices in S that have not too many neighbors in A.

**Lemma 8.** Let 
$$S_1 := \{v \in S : \frac{1}{k} | N(v) \cap A| \le \frac{1}{2} (\alpha + 1) \}$$
. Then  $\frac{1}{k} |S_1| > \frac{1}{3}$ .

*Proof.* Suppose that  $\frac{1}{k}|S_1| = \sigma \leq \frac{1}{3}$ . Let  $G_1 := G[A \cup (S \setminus S_1)]$ . Since  $G_1$  is not (k+1)-connected, it has a separating set  $T \subset V(G_1)$  of size k. Let X and Y form a partition of  $V(G_1) \setminus T$  and be separated by T in  $G_1$ . Without loss of generality assume that  $|X \cap A| \geq |Y \cap A|$ . Then

$$\frac{1}{k}|X\cap A| \ge \frac{1}{2} \cdot \frac{1}{k}|A\setminus T| \ge \frac{1}{2}(\alpha - 1).$$

Hence if  $v \in Y \cap S$ , then

$$\frac{1}{k}|N(v) \cap A| \le \frac{1}{k}(|A| - |X \cap A|) \le \frac{1}{2}(\alpha + 1),$$

which means that  $v \in S_1$ . But that is impossible, since  $S_1 \cap V(G_1) = \emptyset$ . Thus  $Y \cap S = \emptyset$ , i.e.  $Y \subset A$ . In particular, since  $|X \cap A| \ge |Y \cap A| = |Y|$ , we have  $\frac{1}{k}|Y| \le \frac{1}{2}\alpha$ . Then

$$\frac{1}{k}|V(G)\setminus Y| \,=\, \alpha+\beta+1-\frac{1}{k}|Y| \,\geq\, \frac{1}{2}\alpha+\beta+1 \,\geq\, \frac{5}{2},$$

so the induction hypothesis holds for G - Y, and

$$\frac{1}{k^2}|E(G-Y)| \le \frac{19}{12} \left(\frac{1}{k}|V(G-Y)|-1\right).$$

Hence we are done if

$$\frac{1}{k^2}(|E(G)| - |E(G - Y)|) \le \frac{19}{12} \cdot \frac{1}{k}|Y|,$$

so assume that is not the case. Let  $\mu := \frac{1}{k}|Y|$ . Then

$$\frac{1}{2}\mu^2 + \mu(1+\sigma) > \frac{19}{12}\mu,$$

SO

$$\mu > \frac{7}{6} - 2\sigma.$$

We consider two cases.

CASE 1:  $X \cap S \neq \emptyset$ . Let  $v \in X \cap S$ . Then v has more than  $k \cdot \frac{1}{2}(\alpha + 1)$  neighbors in A, none of which belong to Y. Hence  $\mu < \frac{1}{2}(\alpha - 1)$ , and so

$$\frac{1}{2}(\alpha - 1) > \frac{7}{6} - 2\sigma.$$

Therefore,

$$\alpha > \frac{10}{3} - 4\sigma \ge \frac{10}{3} - 4 \cdot \frac{1}{3} = 2;$$

a contradiction.

CASE 2:  $X \cap S = \emptyset$ . Then  $S \setminus S_1 \subset T$ , and the set  $T \cap A$  separates X and Y in G[A] and satisfies  $\frac{1}{k}|T \cap A| = \frac{1}{k}(|T| - |T \cap S|) = 1 - (1 - \sigma) = \sigma$ . Note that since  $|Y| \leq |X|$ , we have

$$\frac{7}{6} - 2\sigma < \mu \le \frac{1}{2}(\alpha - \sigma),$$

SO

$$\sigma > \frac{7}{9} - \frac{1}{3}\alpha > \frac{7}{9} - \frac{1}{3} \cdot \frac{3}{2} = \frac{5}{18}$$

Now observe that

$$\frac{1}{k^2}|E(G[A])| \le \frac{1}{2}\alpha^2 - \mu(\alpha - \sigma - \mu).$$

Since  $\frac{7}{6} - 2\sigma < \mu \le \frac{1}{2}(\alpha - \sigma)$ , the latter expression is less than

$$\frac{1}{2}\alpha^2 - \left(\frac{7}{6} - 2\sigma\right) \cdot \left(\alpha + \sigma - \frac{7}{6}\right).$$

Hence  $\frac{1}{k^2}(|E(G)| - |E(G - A)|)$  is less than

$$\frac{1}{2}(\alpha+1)\sigma + \alpha(1-\sigma) + \frac{1}{2}\alpha^2 - \left(\frac{7}{6} - 2\sigma\right) \cdot \left(\alpha + \sigma - \frac{7}{6}\right).$$

Case 2.1.  $\beta \leq \frac{3}{2}$ . Then, after adding Matula's estimate for the number of edges in  $G[S \cup B]$  and subtracting  $\frac{19}{12}(\alpha + \beta)$ , it is enough to prove that the following quantity is nonpositive:

$$\frac{1}{2}(\alpha+1)\sigma + \alpha(1-\sigma) + \frac{1}{2}\alpha^2 - \left(\frac{7}{6} - 2\sigma\right) \cdot \left(\alpha + \sigma - \frac{7}{6}\right) + \frac{1}{6}\left((\beta+1)^2 + 4(\beta+1) - 2\right) - \frac{19}{12}(\alpha+\beta),$$

which in equal to

$$\frac{1}{36}(18\alpha^2 + 54\alpha\sigma - 63\alpha + 6\beta^2 - 21\beta + 72\sigma^2 - 108\sigma + 67).$$

Note that for  $\alpha$  and  $\sigma$  fixed, the last expression is monotone decreasing in  $\beta$  (recall that  $\beta \leq \frac{3}{2}$ , while the minimum is attained at  $\beta = \frac{7}{4}$ ), so its maximum is attained when  $\beta = \alpha$ , where it turns

$$\varphi_1(\alpha, \sigma) = \frac{1}{36}(24\alpha^2 + 54\alpha\sigma - 84\alpha + 72\sigma^2 - 108\sigma + 67).$$

Since  $\varphi_1(\alpha, \sigma)$  is convex in both  $\alpha$  and  $\sigma$ , it attains its maximum at some point  $(\alpha_0, \sigma_0)$ , where  $\alpha_0 \in \{1, \frac{3}{2}\}$  and  $\sigma_0 \in \{\frac{5}{18}, \frac{1}{3}\}$ . It remains to check the four possibilities:

$$\varphi_1\left(1, \frac{5}{18}\right) = -\frac{11}{162} < 0, \quad \varphi_1\left(1, \frac{1}{3}\right) = -\frac{1}{12} < 0,$$

$$\left(\frac{3}{18}, \frac{5}{18}\right) = -\frac{125}{12} < 0, \quad \text{and} \quad \varphi_1\left(\frac{3}{18}, \frac{1}{18}\right) = -\frac{1}{12} < 0,$$

$$\varphi_1\left(\frac{3}{2}, \frac{5}{18}\right) = -\frac{125}{648} < 0, \text{ and } \varphi_1\left(\frac{3}{2}, \frac{1}{3}\right) = -\frac{1}{6} < 0.$$

CASE 2.2.  $\beta > \frac{3}{2}$ . Then, instead of using Matula's bound for  $G[S \cup B]$ , we can apply the induction hypothesis, so it is enough to prove that the function

$$\varphi_2(\alpha,\sigma) = \frac{1}{2}(\alpha+1)\sigma + \alpha(1-\sigma) + \frac{1}{2}\alpha^2 - \left(\frac{7}{6} - 2\sigma\right) \cdot \left(\alpha + \sigma - \frac{7}{6}\right) - \frac{19}{12}\alpha$$

is nonpositive. Again, we only have to check the boundary values:

$$\varphi_2\left(1, \frac{5}{18}\right) = -\frac{49}{324} < 0, \quad \varphi_2\left(1, \frac{1}{3}\right) = -\frac{1}{6} < 0,$$

$$\varphi_2\left(\frac{3}{2}, \frac{5}{18}\right) = -\frac{125}{648} < 0, \text{ and } \varphi_2\left(\frac{3}{2}, \frac{1}{3}\right) = -\frac{1}{6} < 0.$$

This finishes the proof.

Now we can simply try to use as the set S' the set  $S_1$  itself. This choice indeed gives a good bound if A is large, as the next lemma shows.

Lemma 9.  $\alpha < \frac{4}{3}$ .

*Proof.* Suppose that  $\alpha \geq \frac{4}{3}$ . Recall that  $\sigma = |S_1| > \frac{1}{3}$ . Using Lemma 4' for  $G[A \cup (S \setminus S_1)]$ , we get that

$$\frac{1}{k^2}(|E(G)| - |E(G - A)|)$$

$$\leq \frac{1}{6} \left( (\alpha + 1 - \sigma)^2 + 4(\alpha + 1 - \sigma) - 2 \right) + \frac{1}{2}(\alpha + 1)\sigma$$

$$= \frac{1}{6} (\alpha^2 + \alpha\sigma + 6\alpha + \sigma^2 - 3\sigma + 3).$$

Case 1:  $\beta \leq \frac{3}{2}$ . Adding Matula's estimate for  $G[B \cup S]$  and subtracting  $\frac{19}{12}(\alpha + \beta)$ , we get

$$\frac{1}{k^2}|E(G)| - \frac{19}{12}(\alpha + \beta)$$

$$\leq \frac{1}{6}(\alpha^2 + \alpha\sigma + 6\alpha + \sigma^2 - 3\sigma + 3) + \frac{1}{6}((\beta + 1)^2 + 4(\beta + 1) - 2) - \frac{19}{12}(\alpha + \beta)$$

$$= \frac{1}{12}(2\alpha^2 + 2\alpha\sigma - 7\alpha + 2\beta^2 - 7\beta + 2\sigma^2 - 6\sigma + 12).$$

Again, the maximum is attained when  $\beta = \alpha$ , so we should consider the expression

$$\varphi_3(\alpha,\sigma) = \frac{1}{6}(2\alpha^2 + \alpha\sigma - 7\alpha + \sigma^2 - 3\sigma + 6).$$

It is convex in both  $\alpha$  and  $\sigma$ , so again it is enough to check the boundary points:

$$\varphi_3\left(\frac{4}{3}, \frac{1}{3}\right) = -\frac{1}{27} < 0, \quad \varphi_3\left(\frac{4}{3}, 1\right) = -\frac{2}{27} < 0,$$

$$\varphi_3\left(\frac{3}{2}, \frac{1}{3}\right) = -\frac{7}{108} < 0, \quad \text{and} \quad \varphi_3\left(\frac{3}{2}, 1\right) = -\frac{1}{12} < 0.$$

CASE 2:  $\beta > \frac{3}{2}$ . Then, instead of using Matula's bound for  $G[S \cup B]$ , we can apply the induction hypothesis, so it is enough to prove that the function

$$\varphi_4(\alpha, \sigma) = \frac{1}{6}(\alpha^2 + \alpha\sigma + 6\alpha + \sigma^2 - 3\sigma + 3) - \frac{19}{12}\alpha = \frac{1}{12}(2\alpha^2 + 2\alpha\sigma - 7\alpha + 2\sigma^2 - 6\sigma + 6)$$

is nonpositive. The function is convex in both  $\alpha$  and  $\sigma$ , so we check the boundary points:

$$\varphi_4\left(\frac{4}{3}, \frac{1}{3}\right) = -\frac{1}{18} < 0, \quad \varphi_4\left(\frac{4}{3}, 1\right) = -\frac{5}{54} < 0,$$

$$\varphi_4\left(\frac{3}{2}, \frac{1}{3}\right) = -\frac{7}{108} < 0, \quad \text{and} \quad \varphi_4\left(\frac{3}{2}, 1\right) = -\frac{1}{12} < 0.$$

This finishes the proof.

The next lemma is the final piece of the jigsaw. It shows that if A is small, we can still obtain the desired bound if we take the set S' to be slightly bigger than  $S_1$ .

**Lemma 10.**  $\alpha > \frac{4}{3}$ .

*Proof.* Suppose that  $\alpha \leq \frac{4}{3}$ . Then  $1-2(\alpha-1) \geq \frac{1}{3}$ . Let S' be a subset of S with  $\frac{1}{k}|S'| = 1-2(\alpha-1)$  such that  $\frac{1}{k}|S' \cap S_1| \geq \frac{1}{3}$ . Observe that the normalized number of edges between A and S' is at most

$$\frac{1}{k^2}|A|\cdot|S'| - \frac{1}{2}(\alpha-1)\cdot\frac{1}{3},$$

by the definition of  $S_1$ . Hence, using Lemma 4' for  $G[A \cup (S \setminus S')]$ , we get that

$$\frac{1}{k^2}(|E(G)| - |E(G - A)|) \le \frac{1}{6}\left((3\alpha - 2)^2 + 4(3\alpha - 2) - 2\right) + \alpha(3 - 2\alpha) - \frac{1}{6}(\alpha - 1)$$

$$= \frac{1}{6}(-3\alpha^2 + 17\alpha - 5).$$

CASE 1:  $\beta \leq \frac{3}{2}$ . Adding Matula's estimate for  $G[B \cup S]$  and subtracting  $\frac{19}{12}(\alpha + \beta)$ , we get

$$\frac{1}{k^2}|E(G)| - \frac{19}{12}(\alpha + \beta) \le \frac{1}{6}(-3\alpha^2 + 17\alpha - 5) + \frac{1}{6}\left((\beta + 1)^2 + 4(\beta + 1) - 2\right) - \frac{19}{12}(\alpha + \beta)$$

$$= \frac{1}{12}(-6\alpha^2 + 15\alpha + 2\beta^2 - 7\beta - 4).$$

Since  $\alpha \leq \beta \leq \frac{3}{2}$ , the maximum is attained when  $\beta = \alpha$ , in which case the last expression turns into  $-\frac{1}{3}(\alpha - 1)^2 \leq 0$ .

CASE 2:  $\beta > \frac{3}{2}$ . Then, instead of using Matula's bound for  $G[S \cup B]$ , we can apply the induction hypothesis, so it is enough to prove that

$$\frac{1}{6}(-3\alpha^2 + 17\alpha - 5) - \frac{19}{12}\alpha = -\frac{1}{12}(6\alpha^2 - 15\alpha + 10) \le 0.$$
 (4)

Since the discriminant of the quadratic  $6\alpha^2 - 15\alpha + 10$  is negative, (4) holds for all  $\alpha$ , and we are done.

Since Lemmas 9 and 10 contradict each other, we conclude that such graph G does not exist. This completes the proof of the theorem.

### References

- [1] W. Mader. Existenz n-fach zusammenhängender Teilgraphen in Graphen genügend großen Kantendichte, Abh. Math. Sem. Univ. Hamburg Volume 37, 1972. Pages 86–97.
- [2] W. Mader. Connectivity and edge-connectivity in finite graphs. Surveys in Combinatorics, B. Bollobás, Ed., Cambridge University Press, London, 1979. Pages 66–95.
- [3] D.W. Matula. Ramsey theory for graph connectivity. J. Graph Theory, Volume 7, 1983. Pages 95–105.
- [4] R. Yuster. A note on graphs without k-connected subgraphs. Ars Combinatoria, Volume 67, 2003. Pages 231–235.