

On the second largest distance eigenvalue of a graph

Ruifang Liu^{a*} Jie Xue^a Litao Guo^{b†}

^a School of Mathematics and Statistics, Zhengzhou University, Zhengzhou, Henan 450001, China

^b School of Applied Mathematics, Xiamen University of Technology, Xiamen, Fujian 361024, China

Abstract

Let G be a simple connected graph of order n and $D(G)$ be the distance matrix of G . Suppose that $\lambda_1(D(G)) \geq \lambda_2(D(G)) \geq \dots \geq \lambda_n(D(G))$ are the distance spectrum of G . A graph G is said to be determined by its D -spectrum if with respect to the distance matrix $D(G)$, any graph with the same spectrum as G is isomorphic to G . In this paper, we consider spectral characterization on the second largest distance eigenvalue $\lambda_2(D(G))$ of graphs, and prove that the graphs with $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2} \approx -0.5692$ are determined by their D -spectra.

AMS Classification: 05C50, 05C12

Key words: Distance matrix; The second largest distance eigenvalue; D -spectrum determined

1 Introduction

All graphs considered here are simple, undirected and connected. Let G be a graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G)$. Two vertices u and v are called adjacent if they are connected by an edge. Let $d_G(v)$ and $N_G(v)$ denote the degree and the neighbor set of a vertex v in G , respectively. The distance between vertices u and v of a graph G is denoted by $d_G(u, v)$. The diameter of G , denoted by d or $d(G)$, is the maximum distance between any pair of vertices of G . Let X and Y be subsets of vertices of G . The induced subgraph $G[X]$ is the subgraph of G whose vertex set is X and whose edge set consists of all edges of G which have both ends in X . For any $v \in V(G)$, denote by $G - v$ the induced subgraph $G[V \setminus \{v\}]$. The complete product $G_1 \nabla G_2$ of graphs

*Supported by NSFC (No. 11201432) and NSF-Henan (Nos. 15A110003 and 15IRTSTHN006). E-mail address: rfliu@zzu.edu.cn (R.Liu).

†Supported by NSFC (No. 11301440) and NSF-Fujian (No. JA13240).

G_1 and G_2 is the graph obtained from $G_1 \cup G_2$ by joining every vertex of G_1 with every vertex of G_2 .

The distance matrix $D(G) = (d_{ij})_{n \times n}$ of a connected graph G is the matrix indexed by the vertices of G , where d_{ij} denotes the distance between the vertices v_i and v_j . Let $\lambda_1(D(G)) \geq \lambda_2(D(G)) \geq \dots \geq \lambda_n(D(G))$ be the distance spectrum of G , where $\lambda_2(D(G))$ is called the second largest distance eigenvalue of G . The polynomial $P_D(\lambda) = \det(\lambda I - D(G))$ is defined as the distance characteristic polynomial of the graph G . Two graphs are said to be D -cospectral if they have the same distance spectrum. A graph G is said to be determined by the D -spectra if there is no other nonisomorphic graph D -cospectral to G .

Which graphs are determined by their spectrum seems to be a difficult and interesting problem in the theory of graph spectra. This question was proposed by Dam and Haemers in [2]. In this paper, Dam and Haemers investigated the cospectrality of graphs up to order 11. Later, Dam et al. [3, 4] provided two excellent surveys on this topic. Up to now, only a few families of graphs were shown to be determined by their spectra. In particular, there are much fewer results on which graphs are determined by their D -spectra. In [6], Lin et al. proved that the complete bipartite graph K_{n_1, n_2} and the complete split graph $K_a \nabla K_b^c$ are determined by D -spectra, and conjecture that the complete k -partite graph K_{n_1, n_2, \dots, n_k} is determined by its D -spectrum. Recently, Jin and Zhang [5] have confirmed the conjecture. Xue and Liu [8] showed that some special graphs K_n^h , K_n^{s+t} and $K_n^{s,t}$ are determined by their D -spectra. Lin, Zhai and Gong [7] characterized all connected graphs with $\lambda_{n-1}(D(G)) = -1$, and showed that these graphs are determined by their D -spectra. Moreover, in this paper, they also proved that the graphs with $\lambda_{n-2}(D(G)) > -1$ are determined by their distance spectra. In this paper, we prove that the graphs with $\lambda_2(D(G)) \leq \frac{17 - \sqrt{329}}{2} \approx -0.5692$ are determined by their D -spectra.

Next, we introduce two kinds of graphs K_s^t and $K_n^{n_1, n_2, \dots, n_k}$ as shown in Fig. 1.

- K_s^t : the graph obtained by adding a pendant edge to t vertices of K_s , where $2 \leq t \leq s$ and $n = s + t$.
- $K_n^{n_1, n_2, \dots, n_k}$: the graph G with $d_G(v) = n - 1$ and $G - v$ is the disjoint union of some complete graphs, where $n = \sum_{i=1}^k n_i + 1$ and $k \geq 2$.




Fig. 1. Graphs K_s^t and $K_n^{n_1, n_2, \dots, n_k}$.

2 Preliminaries

Before presenting the proof of the main result, we give some important lemmas and theorems. The following lemma is well-known Cauchy Interlace Theorem.

Lemma 2.1 ([1]) *Let A be a Hermitian matrix of order n with eigenvalues $\lambda_1(A) \geq \lambda_2(A) \geq \dots \geq \lambda_n(A)$, and B be a principal submatrix of A of order m with eigenvalues $\mu_1(B) \geq \mu_2(B) \geq \dots \geq \mu_m(B)$. Then $\lambda_{n-m+i}(A) \leq \mu_i(B) \leq \lambda_i(A)$ for $i = 1, 2, \dots, m$.*

Applying Lemma 2.1 to the distance matrix D of a graph, we have

Lemma 2.2 *Let G be a graph of order n with distance spectrum $\lambda_1(D(G)) \geq \lambda_2(D(G)) \geq \dots \geq \lambda_n(D(G))$, and H be an induced subgraph of G on m vertices with the distance spectrum $\mu_1(D(H)) \geq \mu_2(D(H)) \geq \dots \geq \mu_m(D(H))$. Moreover, if $D(H)$ is a principal submatrix of $D(G)$, then $\lambda_{n-m+i}(D(G)) \leq \mu_i(D(H)) \leq \lambda_i(D(G))$ for $i = 1, 2, \dots, m$.*

Theorem 2.3 *Let G be a connected graph and $D(G)$ be the distance matrix of G . Then $\lambda_2(D(G)) \geq -1$, with the equality if and only if $G \cong K_n$.*

Proof. Let G be a connected graph with order $n \geq 2$, then P_2 is an induced subgraph of G , and $D(P_2)$ is a principal matrix of $D(G)$. Note that $\lambda_2(D(P_2)) = -1$, then by Lemma 2.2, $\lambda_2(D(G)) \geq \lambda_2(D(P_2)) = -1$.

For the equality, we only need to prove the necessity. Suppose that G is not complete graph, then $d(G) \geq 2$. As a result, $D(P_3)$ is a principal submatrix of $D(G)$. Note that $\lambda_2(D(P_3)) = 1 - \sqrt{3}$, by Lemma 2.2, $\lambda_2(D(G)) \geq \lambda_2(D(P_3)) = 1 - \sqrt{3} \approx -0.7321$, a contradiction. Hence $G \cong K_n$. \square

Remark 2.4 *From the proof of the above theorem, there is no graph of order n with $-1 < \lambda_2(D(G)) < 1 - \sqrt{3}$.*

First we will investigate which graphs satisfy $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2} \approx -0.5692$. Let G be a graph with $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2}$. We call H a forbidden subgraph of G if G contains no H as an induced subgraph.

For any $S \subseteq V(G)$, let $D_G(S)$ denote the principal submatrix of $D(G)$ obtained by S .

Lemma 2.5 *Let G be a connected graph and $D(G)$ be the distance matrix of G . If $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2}$, then C_4 , C_5 , P_5 and H_i ($i \in \{1, 2, 3\}$) are forbidden subgraphs of G .*




Fig. 2. Graphs C_4 , C_5 , P_5 and $H_1 - H_3$.

Proof. Note that the diameters of graphs C_4 , C_5 and H_1 are all 2. Their D -spectra are shown as follows.

	λ_1	λ_2	λ_3	λ_4	λ_5
C_4	4.0000	0.0000	-2.0000	-2.0000	
C_5	6.0000	-0.3820	-0.3820	-2.6180	-2.6180
H_1	3.5616	-0.5616	-1.0000	-2.0000	

If C_4 is an induced subgraph of G , then $D(C_4)$ is a principal submatrix of $D(G)$. By Lemma 2.2, then $\lambda_2(D(G)) \geq \lambda_2(D(C_4)) > \frac{17-\sqrt{329}}{2} \approx -0.5692$, a contradiction. Thus C_4 is a forbidden subgraph of G . Similarly, we can prove that C_5 and H_1 are also forbidden subgraphs of G .

Consider P_5 . Suppose that P_5 is an induced subgraph of G , then $d_G(v_1, v_5) \in \{2, 3, 4\}$. If $d_G(v_1, v_5) = 4$, then $D_G(\{v_1, v_2, v_3, v_4, v_5\}) = D(P_5)$, thus $D(P_5)$ is a principal submatrix of $D(G)$. By Lemma 2.2, we have $\lambda_2(D(G)) \geq \lambda_2(D(P_5)) = -0.5578 > \frac{17-\sqrt{329}}{2}$, a contradiction. If $d_G(v_1, v_5) \in \{2, 3\}$, let $d_G(v_1, v_4) = a$, $d_G(v_1, v_5) = b$ and $d_G(v_2, v_5) = c$, then $a, b, c \in \{2, 3\}$. Hence the principal submatrix of $D(G)$

$$D_G(\{v_1, v_2, v_3, v_4, v_5\}) = \begin{pmatrix} 0 & 1 & 2 & a & b \\ 1 & 0 & 1 & 2 & c \\ 2 & 1 & 0 & 1 & 2 \\ a & 2 & 1 & 0 & 1 \\ b & c & 2 & 1 & 0 \end{pmatrix}.$$

By a simple calculation, we have

(a, b, c)	(3, 3, 3)	(3, 2, 2)	(3, 2, 3)	(3, 3, 2)	(2, 3, 3)	(2, 3, 2)	(2, 2, 2)	(2, 2, 3)
λ_2	-0.4348	-0.3260	0	-0.3713	-0.3713	-0.1646	-0.2909	-0.3260

From Lemma 2.2, this contradicts $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2}$. Hence P_5 is a forbidden subgraph of G .

Consider H_2 . Suppose that H_2 is an induced subgraph of G . Let $d_G(v_1, v_4) = a$ and $d_G(v_4, v_5) = b$, then $a, b \in \{2, 3\}$. We get the principal submatrix of $D(G)$:

$$D_G(\{v_1, v_2, v_3, v_4, v_5\}) = \begin{pmatrix} 0 & 1 & 2 & a & 2 \\ 1 & 0 & 1 & 2 & 1 \\ 2 & 1 & 0 & 1 & 2 \\ a & 2 & 1 & 0 & b \\ 2 & 1 & 2 & b & 0 \end{pmatrix}.$$

By a simple calculation, we have

(a, b)	$(3, 3)$	$(2, 3)$	$(3, 2)$	$(2, 2)$
λ_2	-0.5120	-0.3583	-0.3583	-0.2245

By Lemma 2.2, $\lambda_2(D(G)) \geq \lambda_2(D_G(\{v_1, v_2, v_3, v_4, v_5\})) > \frac{17-\sqrt{329}}{2}$, a contradiction. Hence H_2 is a forbidden subgraph of G .

Consider H_3 . Suppose that H_3 is an induced subgraph of G . Let $d_G(v_1, v_4) = a$ and $d_G(v_4, v_5) = b$, then $a, b \in \{2, 3\}$. Then the principal submatrix of $D(G)$

$$D_G(\{v_1, v_2, v_3, v_4, v_5\}) = \begin{pmatrix} 0 & 1 & 2 & a & 1 \\ 1 & 0 & 1 & 2 & 1 \\ 2 & 1 & 0 & 1 & 2 \\ a & 2 & 1 & 0 & b \\ 1 & 1 & 2 & b & 0 \end{pmatrix}.$$

By a simple calculation, we have

(a, b)	$(3, 3)$	$(2, 3)$	$(3, 2)$	$(2, 2)$
λ_2	-0.5686	-0.3626	-0.3626	-0.3311

From Lemma 2.2, we also get a contradiction since $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2}$. Therefore H_3 is a forbidden subgraph of G . \square

Let $\mathbb{K}_s^t = \{K_s^t | 2 \leq t \leq s, s+t = n\}$ and $\mathbb{K}_n^{n_1, n_2, \dots, n_k} = \{K_n^{n_1, n_2, \dots, n_k} | \sum_{i=1}^k n_i + 1 = n, k \geq 2\}$.

Lemma 2.6 *Let G be a connected graph and $D(G)$ be the distance matrix of G . If $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2}$, then $G \in K_n \cup \mathbb{K}_s^t \cup \mathbb{K}_n^{n_1, n_2, \dots, n_k}$.*

Proof. Let $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2}$. Note that P_5 is a forbidden subgraph of G , then $d(G) \leq 3$. $d(G) = 1$ indicates that $G \cong K_n$. Next we consider $d(G) \geq 2$, then there exist non-adjacent vertices in G . For any two non-adjacent vertices, we claim that they have at most one common neighbor. Otherwise there exists forbidden subgraph C_4 or H_1 . Next we distinguish the following two cases:

Case 1. There exist two non-adjacent vertices u and v such that $|N_G(u) \cap N_G(v)| = 0$.

Then $d_G(u, v) = 3$. Let $P = uxyv$ be the shortest path between u and v . Thus we have $d_G(u) = d_G(v) = 1$. Otherwise there exists forbidden subgraph P_5, H_3, C_4 or H_1 .

Let $S = (N_G(x) \cup N_G(y)) \setminus \{u, v\}$ and $T = V(G) \setminus (S \cup \{u, v, x, y\})$.

We claim that $G[S \cup \{x, y\}]$ is a clique, otherwise there exists forbidden subgraph H_2 or H_1 . If $T = \emptyset$, then $G \cong K_s^2$. If $T \neq \emptyset$, then we claim that each vertex in T is adjacent to some vertices in S . Otherwise the distance from the vertex to u or v is at least 4, a contradiction. Furthermore, we claim that each vertex in T is adjacent to exactly one vertex in S and the vertices in T have no common neighbor in S . Otherwise there exists forbidden subgraph H_1 , H_2 or H_3 . Moreover, by forbidden subgraph C_4 , T is an independent set. Therefore $G \cong K_s^t$.

Case 2. For any two non-adjacent vertices, they have exactly one common neighbor.

Let $S = \{v_1, v_2, \dots, v_k\}$ be the maximum independent set of G , where $k \geq 2$. Let $\bar{S} = V(G) \setminus S$. Then by the definition of the maximum independent set, each vertex in \bar{S} is adjacent to some vertices in S .

Let v be the only common neighbor of v_1 and v_2 . We claim that v is adjacent to each vertex in S . Otherwise there exists forbidden subgraph H_2, C_5, C_4 or H_1 . Since any two non-adjacent vertices of G have exactly one common neighbor, then each vertex in $\bar{S} \setminus \{v\}$ is adjacent to exactly one vertex in S .

Let V_1, V_2, \dots, V_k be a partition of $\bar{S} \setminus \{v\}$, where V_i (possibly empty) is the vertex subset whose vertices are all adjacent to v_i . Next we show that each vertex in $\bar{S} \setminus \{v\}$ is adjacent to v . Without loss of generality, we consider the vertex subset V_1 . Let $v_1^* \in V_1$. Suppose that $v_1^*v \notin E(G)$. Note that v_1^* is not adjacent to v_2 , hence we can suppose that w is the only common neighbor of v_1^* and v_2 . Then $G[vv_1v_1^*wv_2] = C_5$ or $G[vv_1v_1^*w] = C_4$, a contradiction.

Moreover, we claim that $G[V_i]$ ($i = 1, 2, \dots, k$) is a clique and $E[V_i, V_j] = \emptyset$ ($i \neq j$). Otherwise there exists forbidden subgraph H_1 . Thus we have $G \cong K_n^{n_1, n_2, \dots, n_k}$. \square

3 Main results

In this section, we will show that the graphs with $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2}$ are determined by their D -spectra. First, we give the distance characteristic polynomials of K_s^t and $K_n^{n_1, n_2, \dots, n_k}$.

Lemma 3.1 *Let $G = K_s^t$, where $2 \leq t \leq s$ and $n = s+t$. Then the distance characteristic polynomial of G is as follows.*

For $s \geq t + 1$,

$$P_D(\lambda) = (\lambda+1)^{s-t-1}(\lambda+2-\sqrt{2})^{t-1}(\lambda+2+\sqrt{2})^{t-1}[\lambda^3+(5-s-3t)\lambda^2+(6-4s-2t-st)\lambda+2-2s-st].$$

For $s = t$,

$$P_D(\lambda) = (\lambda+2-\sqrt{2})^{t-1}(\lambda+2+\sqrt{2})^{t-1}[\lambda^2+(4-4t)\lambda+2-2t-t^2].$$

Proof. For $s \geq t + 1$. The distance matrix of K_s^t can be written as

$$D = \begin{pmatrix} \sigma & \alpha & \cdots & \alpha & \beta \\ \alpha & \sigma & \cdots & \alpha & \beta \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \alpha & \alpha & \cdots & \sigma & \beta \\ \beta^\top & \beta^\top & \cdots & \beta^\top & B \end{pmatrix},$$

where $\sigma = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\alpha = \begin{pmatrix} 3 & 2 \\ 2 & 1 \end{pmatrix}$, $\beta = \begin{pmatrix} 2 & 2 & \cdots & 2 \\ 1 & 1 & \cdots & 1 \end{pmatrix}_{2 \times (s-t)}$, and

$$B = \begin{pmatrix} 0 & 1 & \cdots & 1 \\ 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 0 \end{pmatrix}_{(s-t) \times (s-t)}.$$

Thus

$$\begin{aligned} \det(\lambda I - D) &= \begin{vmatrix} \lambda I_{2 \times 2} - \sigma & -\alpha & \cdots & -\alpha & -\beta \\ -\alpha & \lambda I_{2 \times 2} - \sigma & \cdots & -\alpha & -\beta \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -\alpha & -\alpha & \cdots & \lambda I_{2 \times 2} - \sigma & -\beta \\ -\beta^\top & -\beta^\top & \cdots & -\beta^\top & \lambda I_{(s-t) \times (s-t)} - B \end{vmatrix} \\ &= \begin{vmatrix} \lambda I_{2 \times 2} - \sigma - (t-1)\alpha & -\alpha & \cdots & -\alpha & -\beta \\ 0 & \lambda I_{2 \times 2} - \sigma + \alpha & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \lambda I_{2 \times 2} - \sigma + \alpha & 0 \\ -t\beta^\top & -\beta^\top & \cdots & -\beta^\top & \lambda I_{(s-t) \times (s-t)} - B \end{vmatrix} \\ &= |\lambda I_{2 \times 2} - \sigma + \alpha|^{t-1} \begin{vmatrix} \lambda I_{2 \times 2} - \sigma - (t-1)\alpha & -\beta \\ -t\beta^\top & \lambda I_{(s-t) \times (s-t)} - B \end{vmatrix} \\ &= (\lambda+1)^{s-t-1} (\lambda+2-\sqrt{2})^{t-1} (\lambda+2+\sqrt{2})^{t-1} [\lambda^3 + (5-s-3t)\lambda^2 + (6-4s-2t-st)\lambda + 2-2s-st]. \end{aligned}$$

For $s = t$. The distance matrix of K_s^t can be written as

$$D = \begin{pmatrix} \sigma & \alpha & \cdots & \alpha \\ \alpha & \sigma & \cdots & \alpha \\ \vdots & \vdots & \ddots & \vdots \\ \alpha & \alpha & \cdots & \sigma \end{pmatrix}.$$

Thus

$$\begin{aligned}
\det(\lambda I - D) &= \begin{vmatrix} \lambda I_{2 \times 2} - \sigma & -\alpha & \cdots & -\alpha \\ -\alpha & \lambda I_{2 \times 2} - \sigma & \cdots & -\alpha \\ \vdots & \vdots & \ddots & \vdots \\ -\alpha & -\alpha & \cdots & \lambda I_{2 \times 2} - \sigma \end{vmatrix} \\
&= \begin{vmatrix} \lambda I_{2 \times 2} - \sigma - (t-1)\alpha & -\alpha & \cdots & -\alpha \\ 0 & \lambda I_{2 \times 2} - \sigma + \alpha & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda I_{2 \times 2} - \sigma + \alpha \end{vmatrix} \\
&= |\lambda I_{2 \times 2} - \sigma + \alpha|^{t-1} |\lambda I_{2 \times 2} - \sigma - (t-1)\alpha| \\
&= (\lambda + 2 - \sqrt{2})^{t-1} (\lambda + 2 + \sqrt{2})^{t-1} [\lambda^2 + (4-4t)\lambda + 2 - 2t - t^2].
\end{aligned}$$

□

Corollary 3.2 *Let $G = K_s^t$, where $3 \leq t+1 \leq s$ and $n = s+t$. Then $\lambda_2(D(G)) = \sqrt{2} - 2$.*

Proof. By Lemma 3.1,

$$P_D(\lambda) = (\lambda+1)^{s-t-1} (\lambda+2-\sqrt{2})^{t-1} (\lambda+2+\sqrt{2})^{t-1} [\lambda^3 + (5-s-3t)\lambda^2 + (6-4s-2t-st)\lambda + 2-2s-st].$$

Let $f(\lambda) = \lambda^3 + (5-s-3t)\lambda^2 + (6-4s-2t-st)\lambda + 2-2s-st$. By a simple calculation,

$$\begin{cases} f(0) = 2 - 2s - st < 0, \\ f(-\frac{2}{3}) = -\frac{2}{27} + \frac{2}{9}s - \frac{1}{3}st < 0, \\ f(-1) = s - t > 0. \end{cases}$$

Note that $f(\lambda) \rightarrow +\infty$ ($\lambda \rightarrow +\infty$) and $f(0) < 0$, so there is at least one root in $(0, +\infty)$. Since $f(-\frac{2}{3}) < 0$ and $f(-1) > 0$, then there is at least one root in $(-1, -\frac{2}{3})$. By $f(\lambda) \rightarrow -\infty$ ($\lambda \rightarrow -\infty$) and $f(-1) > 0$, so there is at least one root in $(-\infty, -1)$. Thus there is exactly one root in each interval. So $\lambda_2(D(G)) = \sqrt{2} - 2$. □

Theorem 3.3 *No two non-isomorphic graphs in \mathbb{K}_s^t are D -cospectral.*

Proof. For any $K_{s_1}^{t_1}, K_{s_2}^{t_2} \in \mathbb{K}_s^t$. Suppose that they are D -cospectral, by Lemma 3.1, then $t_1 = t_2$. Note that $s_1 + t_1 = s_2 + t_2 = n$, thus $s_1 = s_2$, that is, $K_{s_1}^{t_1} \cong K_{s_2}^{t_2}$. □

Lemma 3.4 *Let $G = K_n^{n_1, n_2, \dots, n_k}$, where $n = \sum_{i=1}^k n_i + 1$ and $k \geq 2$. Then the distance characteristic polynomial of G is*

$$P_D(\lambda) = (\lambda + 1)^{n-k-1} \left(\lambda - \sum_{i=1}^k \frac{n_i(2\lambda + 1)}{\lambda + n_i + 1} \right) \prod_{i=1}^k (\lambda + n_i + 1).$$

Proof. The distance matrix of G has the form

$$D = \begin{pmatrix} J_{n_1} - I_{n_1} & 2J_{n_1 \times n_2} & \cdots & 2J_{n_1 \times n_k} & J_{n_1 \times 1} \\ 2J_{n_2 \times n_1} & J_{n_2} - I_{n_2} & \cdots & 2J_{n_2 \times n_k} & J_{n_2 \times 1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 2J_{n_k \times n_1} & 2J_{n_k \times n_2} & \cdots & J_{n_k} - I_{n_k} & J_{n_k \times 1} \\ J_{1 \times n_1} & J_{1 \times n_2} & \cdots & J_{1 \times n_k} & 0 \end{pmatrix},$$

where J_{n_i} is the all-one square matrix with order n_i , I_{n_i} is the identity square matrix with order n_i , and $J_{n_i \times n_j}$ ($i \neq j$) is the all-one matrix with n_i rows and n_j columns.

Then

$$\begin{aligned} \det(\lambda I - D) &= (\lambda+1)^{n_1-1} (\lambda+1)^{n_2-1} \cdots (\lambda+1)^{n_k-1} \begin{vmatrix} \lambda - (n_1 - 1) & -2n_2 & \cdots & -2n_k & -1 \\ -2n_1 & \lambda - (n_2 - 1) & \cdots & -2n_k & -1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -2n_1 & -2n_2 & \cdots & \lambda - (n_k - 1) & -1 \\ -n_1 & -n_2 & \cdots & -n_k & \lambda \end{vmatrix} \\ &= (\lambda+1)^{\sum_{i=1}^k n_i - k} \begin{vmatrix} \lambda + n_1 + 1 & 0 & \cdots & 0 & -1 - 2\lambda \\ 0 & \lambda + n_2 + 1 & \cdots & 0 & -1 - 2\lambda \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \lambda + n_k + 1 & -1 - 2\lambda \\ -n_1 & -n_2 & \cdots & -n_k & \lambda \end{vmatrix} \\ &= (\lambda+1)^{n-k-1} \left(\lambda - \sum_{i=1}^k \frac{n_i(2\lambda+1)}{\lambda+n_i+1} \right) \prod_{i=1}^k (\lambda+n_i+1). \end{aligned}$$

□

Theorem 3.5 *No two non-isomorphic graphs in $\mathbb{K}_n^{n_1, n_2, \dots, n_k}$ are D -cospectral.*

Proof. Let $G = K_n^{n_1, n_2, \dots, n_k}$. By Lemma 3.4, -1 is the eigenvalue of G with multiplicity $n-k-1$. Suppose that $G^* \in \mathbb{K}_n^{n_1, n_2, \dots, n_k}$ is D -cospectral to G , then -1 is also the eigenvalue of G^* with multiplicity $n-k-1$. Hence we may assume that $G^* = K_n^{n_1^*, n_2^*, \dots, n_k^*}$ (k parts) with $\sum_{i=1}^k n_i^* = n-1$. Let us denote the rest of common distance eigenvalues of G and G^* by $\lambda_1, \lambda_2, \dots, \lambda_{k+1}$. Let

$$\begin{aligned} g(\lambda, n_1, n_2, \dots, n_k) &= \left(\lambda - \sum_{i=1}^k \frac{n_i(2\lambda+1)}{\lambda+n_i+1} \right) \prod_{i=1}^k (\lambda+n_i+1) \\ &= \lambda \prod_{i=1}^k (\lambda+n_i+1) - (2\lambda+1) \sum_{i=1}^k n_i \prod_{j=1, j \neq i}^k (\lambda+n_j+1) \\ &= \sum_{i=0}^{k+1} \mu_i \lambda^{k+1-i}, \end{aligned}$$

then $\lambda_1, \lambda_2, \dots, \lambda_{k+1}$ are the roots of $g(\lambda, n_1, n_2, \dots, n_k)$.

Denote $s_1 = \sum_{j=1}^k n_j$, $s_2 = \sum_{1 \leq j_1 < j_2 \leq k} n_{j_1} n_{j_2}, \dots, s_i = \sum_{1 \leq j_1 < j_2 < \dots < j_i \leq k} n_{j_1} n_{j_2} \dots n_{j_i}, \dots, s_k = n_1 n_2 \dots n_k$. Obviously, $\mu_0 = 1$ and $\mu_1 = -\sum_{i=1}^k n_i + k = -s_1 + k$. For $2 \leq i \leq k$, by the calculation, we have

$$\begin{aligned} \mu_i &= \sum_{1 \leq j_1 < j_2 < \dots < j_i \leq k} (n_{j_1} + 1) \dots (n_{j_i} + 1) - 2 \sum_{l=1}^k n_l \sum_{1 \leq j_1 < j_2 < \dots < j_{i-1} \leq k, l \notin \{j_1, \dots, j_{i-1}\}} (n_{j_1} + 1) \dots (n_{j_{i-1}} + 1) \\ &\quad - \sum_{l=1}^k n_l \sum_{1 \leq j_1 < j_2 < \dots < j_{i-2} \leq k, l \notin \{j_1, \dots, j_{i-2}\}} (n_{j_1} + 1) \dots (n_{j_{i-2}} + 1) \\ &= t_i s_i + t_{i-1} s_{i-1} + t_{i-2} s_{i-2} + \dots + t_1 s_1 + t_0, \end{aligned}$$

where $t_i = 1 - 2i$, and $\mu_{k+1} = -k s_k - (k-1) s_{k-1} - \dots - 2 s_2 - s_1$.

Define $s_1^* = \sum_{j=1}^k n_j^*$, $s_2^* = \sum_{1 \leq j_1 < j_2 \leq k} n_{j_1}^* n_{j_2}^*, \dots, s_i^* = \sum_{1 \leq j_1 < j_2 < \dots < j_i \leq k} n_{j_1}^* n_{j_2}^* \dots n_{j_i}^*, \dots, s_k^* = n_1^* n_2^* \dots n_k^*$. Let

$$\begin{aligned} g(\lambda, n_1^*, n_2^*, \dots, n_k^*) &= \left(\lambda - \sum_{i=1}^k \frac{n_i^* (2\lambda + 1)}{\lambda + n_i^* + 1} \right) \prod_{i=1}^k (\lambda + n_i^* + 1) \\ &= \lambda \prod_{i=1}^k (\lambda + n_i^* + 1) - (2\lambda + 1) \sum_{i=1}^k n_i^* \prod_{j=1, j \neq i}^k (\lambda + n_j^* + 1) \\ &= \sum_{i=0}^{k+1} \mu_i^* \lambda^{k+1-i}. \end{aligned}$$

Similar to the calculation of μ_i , then

$$\mu_i^* = t_i s_i^* + t_{i-1} s_{i-1}^* + t_{i-2} s_{i-2}^* + \dots + t_1 s_1^* + t_0, \quad i = 1, 2, \dots, k.$$

Note that $\lambda_1, \lambda_2, \dots, \lambda_{k+1}$ are also the roots of $g(\lambda, n_1^*, n_2^*, \dots, n_k^*)$, hence $\mu_i = \mu_i^*$. Recursively, we can obtain that $s_1 = s_1^*, s_2 = s_2^*, \dots, s_k = s_k^*$. Since n_1, n_2, \dots, n_k are the roots of $x^k - s_1 x^{k-1} + s_2 x^{k-2} + \dots + (-1)^k s_k = 0$, then $n_1^*, n_2^*, \dots, n_k^*$ are also the roots of the same polynomial, and hence $n_1^*, n_2^*, \dots, n_k^*$ must be a permutation of n_1, n_2, \dots, n_k . So $G^* \cong G$. \square

Theorem 3.6 For any $G \in \mathbb{K}_s^t$ and $\tilde{G} \in \mathbb{K}_n^{n_1, n_2, \dots, n_k}$, then they are not D -cospectral.

Proof. We may assume that $G = K_s^t$ and $\tilde{G} = K_n^{n_1, n_2, \dots, n_k}$ with m and \tilde{m} edges, respectively. Suppose that G and \tilde{G} have the same distance spectrum $\lambda_1(D) \geq \lambda_2(D) \geq \dots \geq \lambda_n(D)$. Let $D(G) = (d_{ij})$ and $D(\tilde{G}) = (\tilde{d}_{ij})$ be the distance matrices of G and \tilde{G} , respectively. Then we have

$$\sum_{l=1}^n \lambda_l^2 = \sum_{i=1}^n \sum_{j=1}^n d_{ij}^2 = 2[m + \left(\frac{n(n-1)}{2} - m\right) \times 4 + \frac{t(t-1)}{2} \times 5] = 4n(n-1) - 6m + 5t(t-1)$$

and

$$\sum_{l=1}^n \lambda_l^2 = \sum_{i=1}^n \sum_{j=1}^n \tilde{d}_{ij}^2 = 2[\tilde{m} + (\frac{n(n-1)}{2} - \tilde{m}) \times 4] = 4n(n-1) - 6\tilde{m}.$$

Hence $6m - 5t(t-1) = 6\tilde{m}$, i.e., $6(m - \tilde{m}) = 5t(t-1)$. Note that $t \geq 2$, then $m - \tilde{m}$ is a nonzero positive integer, and hence $6|t(t-1)$.

If $s \geq t + 1$. By Lemmas 3.1 and 3.4, we know that -1 is the eigenvalue of G and \tilde{G} with multiplicity $s - t - 1$ and $n - k - 1$, respectively. Hence $s - t - 1 = n - k - 1$, then $k = 2t$, since $s = n - t$. By Corollary 3.2, $\lambda_2(D(G)) = \sqrt{2} - 2 \approx -0.5858$, then $\lambda_2(D(\tilde{G})) = \sqrt{2} - 2$. We claim that $k < 6$. Otherwise, the star $K_{1,6}$ is an induced subgraph of \tilde{G} , by Lemma 2.2, $\lambda_2(D(\tilde{G})) \geq \lambda_2(D(K_{1,6})) = -0.5678$, a contradiction. Note that $t \geq 2$, then $k = 2t = 4$, contradicting $6|t(t-1)$.

If $s = t$. By Lemma 3.1, -1 is the eigenvalue of G with multiplicity 0, hence -1 is also the eigenvalue of \tilde{G} with multiplicity 0. By Lemma 3.4, $k = n - 1$. Then $n_1 = n_2 = \dots = n_k = 1$, that is $\tilde{G} = K_{1,n-1}$. Note that -2 is the eigenvalue of \tilde{G} , then -2 is also the eigenvalue of G , we can deduce that $t^2 - 6t + 2 = 0$, a contradiction. \square

Theorem 3.7 *The connected graphs with $\lambda_2(D(G)) \leq \frac{17-\sqrt{329}}{2} \approx -0.5692$ are determined by their D -spectra.*

Proof. By Lemma 2.6, $G \in K_n \cup \mathbb{K}_s^t \cup \mathbb{K}_n^{n_1, n_2, \dots, n_k}$. Obviously, K_n is determined by its D -spectrum. By Theorems 3.3, 3.5 and 3.6, the result follows immediately.

References

- [1] D.M. Cvetković, M. Doob, H. Sachs, Spectra of Graphs, third edition, Johann Ambrosius Barth Verlag, 1995.
- [2] E. R. van Dam, W. H. Haemers, Which graphs are determined by their spectrum?, Linear Algebra Appl. 373 (2003) 241-272.
- [3] E. R. van Dam, W. H. Haemers, J. H. Koolen, Cospectral graphs and the generalized adjacency matrix, Linear Algebra Appl. 423 (2007) 33-41.
- [4] E. R. van Dam, W. H. Haemers, Developments on spectral characterizations of graphs, Discrete Math. 309 (2009) 576-586.
- [5] Y.-L. Jin and X.-D. Zhang, Complete multipartite graphs are determined by their distance spectra, Linear Algebra Appl. 448 (2014) 285-291.
- [6] H.Q. Lin, Y. Hong, J.F. Wang, J.L. Shu, On the distance spectrum of graphs, Linear Algebra Appl. 439 (2013) 1662-1669.
- [7] H.Q. Lin, M.Q. Zhai, S.C. Gong, On graphs with at least three distance eigenvalues less than -1 , Linear Algebra Appl. 458 (2014) 548-558.
- [8] J. Xue, R.F. Liu, Some graphs determined by their D -spectra, in submission.