Detecting genomic signatures of natural selection with principal component analysis: application to the 1000 Genomes data

Nicolas Duforet-Frebourg^{1,2,3}, Guillaume Laval^{4,5}, Eric Bazin⁶, Michael G.B. Blum^{1,2,*}

1 Univ. Grenoble Alpes, TIMC-IMAG UMR 5525, F-38000 Grenoble, France

2 CNRS, TIMC-IMAG, F-38000 Grenoble, France

3 Department of Integrative Biology, University of California, Berkeley, California 94720-3140, USA

4 Institut Pasteur, Human Evolutionary Genetics, Department of Genomes and Genetics, Paris, France

5 Centre National de la Recherche Scientifique, URA3012, Paris, France

6 Univ. Grenoble Alpes, CNRS, Laboratoire d'Ecologie Alpine UMR 5553, F-38000 Grenoble, France

* michael.blum@imag.fr.edu

Abstract

Large-scale genomic data offers the perspective to decipher the genetic architecture of natural selection. To characterize natural selection, various analytical methods for detecting candidate genomic regions have been developed. We propose to perform genome-wide scans of natural selection using principal component analysis. We show that the common F_{ST} index of genetic differentiation between populations can be viewed as a proportion of variance explained by the principal components. Looking at the correlations between genetic variants and each principal component provides a conceptual framework to detect genetic variants involved in local adaptation without any prior definition of populations. To validate the PCA-based approach, we consider the 1000 Genomes data (phase 1) after removal of recently admixed individuals resulting in 850 individuals coming from Africa, Asia, and Europe. The number of genetic variants is of the order of 36 millions obtained with a low-coverage sequencing depth (3X). The correlations between genetic variation and each principal component provide well-known targets for positive selection (EDAR, SLC24A5, SLC45A2, DARC), and also new candidate genes (APPBPP2, TP1A1, RTTN, KCNMA, MYO5C) and non-coding RNAs. In addition to identifying genes involved in biological adaptation, we identify two biological pathways involved in polygenic adaptation that are related to the innate immune system (beta defensins) and to lipid metabolism (fatty acid omega oxidation). PCA-based statistics retrieve well-known signals of human adaptation, which is encouraging for future whole-genome sequencing project, especially in non-model species for which defining populations can be difficult. Genome scan based on PCA is implemented in the open-source and freely available PCAdapt software.

Introduction

Because of the flood of genomic data, the ability to understand the genetic architecture of natural selection has dramatically increased. Of particular interest is the study of local positive selection which explains why individuals are adapted to their local environment. In humans, the availability of genomic data fostered the identification of loci involved in positive selection (Sabeti et al. 2007; Barreiro et al. 2008; Pickrell et al. 2009; Grossman et al. 2013). Local positive selection tends to increase genetic differentiation, which can be measured by difference of allele frequencies between populations (Sabeti et al. 2006; Nielsen 2005; Colonna et al. 2014). For instance, a mutation in the DARC gene that confers resistance to malaria is fixed in Sub-Saharan African populations whereas it is absent elsewhere (Hamblin et al. 2002). In addition to the variants that confer resistance to pathogens, genome scans also identify other genetic variants, and many of these are involved in human metabolic phenotypes and morphological traits (Barreiro et al. 2008; Hancock et al. 2010).

In order to provide a list of variants potentially involved in natural selection, genome scans compute measures of genetic differentiation between populations and consider that extreme values correspond to candidate regions (Luikart et al. 2003). The most widely used index of genetic differentiation is the F_{ST} index which measures the amount of genetic variation that is explained by variation between populations (Excoffier et al. 1992). However, the F_{ST} statistic is not optimal in many settings especially when there is hierarchical population structure, which can arise in models of divergence between populations (Bonhomme et al. 2010; Bierne et al. 2013). When there is hierarchical population structure, other related statistics can provide less false discoveries than F_{ST} although they generally require to group individuals into populations (Bonhomme et al. 2010; Fariello et al. 2013; Günther and Coop 2013). Grouping individuals into population can be subjective, and important signals of selection may be missed with an inadequate choice of populations (Yang et al. 2012). We have previously developed an indivial-based approach for selection scan based on a Bayesian factor model but the MCMC algorithm required for model fitting does not scale well to large data sets containing a million of variants or more (Duforet-Frebourg et al. 2014).

In this paper, we propose to detect candidates for natural selection using principal component analysis (PCA). PCA is technique of multivariate analysis used to ascertain population structure (Patterson et al. 2006). PCA decomposes the total genetic variation into K axes of genetic variation called principal components. In population genomics, the principal components correspond to evolutionary processes such as evolutionary divergence between populations (McVean 2009). In this paper, we show that, with an appropriate choice of K, the common F_{ST} statistic can be viewed as the proportion of variation explained by the principal components (see also Laloë and Gautier 2011). With this point of view, the F_{ST} for a given variant is obtained as a sum of squared correlations with the different principal components opening the door to new statistics for genome scans. At a genome-wide level, it is known that there is a relationship between F_{ST} and PCA (McVean 2009), and we show that the relationship also applies at the level of

a single variant. Using the 1000 Genomes data, we show that PCA can provide useful insights for genome scans. Looking at the correlations between SNPs and each of the principal component provides a conceptual framework to detect genomic regions that are candidates for local adaptation at different levels of population structure.

Methods

New statistics for genome scan

We denote by \mathbf{Y} the $(n \times p)$ centered and scaled genotype matrix where *n* is the number of individuals and *p* is the number of loci. The new statistics for genome scan are based on principal component analysis. The objective of PCA is to find a new set of orthogonal variables called the principal components, which are linear combinations of (centered and standardized) allele counts, such that the projections of the data onto these axes lead to an optimal summary of the data. To present the method, we introduce the truncated singular value decomposition (SVD), which approximates the data matrix \mathbf{Y} as

$$\mathbf{Y} \approx \mathbf{U} \mathbf{\Sigma} \mathbf{V}^T,\tag{1}$$

where U is a $(n \times K)$ orthonormal matrix, V is a $(p \times K)$ orthonormal matrix, Σ is a diagonal $(K \times K)$ matrix and K corresponds to the number of principal components. The solution of PCA with K components can be obtained using the truncated SVD of equation (1) : the K columns of V contain the coefficients of the linear combinations used to form the new orthogonal variables, the K columns of U contain the projections (called *scores*) of the original variables onto the principal components and capture population structure (S1 Fig.), and the squares of the elements of Σ are proportional to the proportion of variance explained by each principal component (Jolliffe 2005). We denote the diagonal elements of Σ by $\sqrt{\lambda_k}$, $k = 1, \ldots, K$ where the λ_k 's are the ranked eigenvalues of the covariance matrix $\mathbf{Y}\mathbf{Y}^T$. Denoting by \mathbf{V}_{jk} , the entry of V at the j^{th} line and k^{th} column, then the correlation ρ_{jk} between the j^{th} SNP and the k^{th} principal component is given by $\rho_{jk} = \sqrt{\lambda_k}V_{jk}$ (Cadima and Jolliffe 1995). In the following, the correlations ρ_{jk} are referred as *loadings* and will be used for detecting selection.

The second statistic we consider for genome scan corresponds to the proportion of variance of a SNP that is explained by the first K PCs. It is called the communality in exploratory factor analysis because it is the variance of observed variables accounted for by the common factors (Suhr 2009). Denoting by h_j the communality of the j^{th} SNP, we have

$$h_j^2 = \sum_{k=1}^K \lambda_k V_{jk}^2.$$
 (2)

The last statistic we consider for genome scans sums the squared of normalized loadings. It is defined as $h_j^{\prime 2} = \sum_{k=1}^{K} V_{jk}^2$. Compared to the communality h^2 , the statistic h'^2 should theoretically give the same importance to each

PC because the normalized loadings are on the same scale as we have $\sum_{j=1}^{p} V_{jk}^2 = 1$, for $k = 1 \dots K$.

Numerical computations

The method of selection scan should be able to handle a large number p of genetic variants. In order to compute truncated SVD with large values of p, we form the $n \times n$ covariance matrix $\Omega = \mathbf{Y}\mathbf{Y}^T/(p-1)$. The covariance matrix Ω is typically of much smaller dimension than the $p \times p$ covariance matrix. Considering the $n \times n$ covariance matrix Ω speeds up matrix operations. Computation of the covariance matrix is the most costly operation and it requires a number of arithmetic operations proportional to pn^2 . After computing the covariance matrix Ω , we compute its first K eigenvalues and eigenvectors to find $\Sigma^2/(p-1)$ and U. Eigenanalysis is performed with the *dsyevr* routine of the linear algebra package LAPACK (Anderson et al. 1999). The matrix V, which captures the relationship between each SNPs and population structure, is obtained by the matrix operation $\mathbf{V}^T = \boldsymbol{\Sigma}^{-1} \mathbf{U}^T \mathbf{Y}$ that arises from equation (1). In the software *PCAdapt*, data are processed as a stream and never stored in order to have a very low memory access whatever the size of the data.

Results

To compare the performance of the different PCA-based summary statistics, we simulate genetic variation in models of population divergence. The divergence models assume that there are three populations, A, B_1 and B_2 with B_1 and B_2 being the most related populations (Figs. 1 and 2). The first simulation scheme assumes that local adaptation took place in the lineages corresponding to the environments of populations A and B_1 (Fig. 1). The SNPs, which are are assumed to be independent, are divided into 3 groups: 9,500 SNPs evolve neutrally, 250 SNPs confer a selective advantage in the environment of A, and 250 other SNPs confer a selective advantage in the environment of B_1 . Performing principal component analysis with K = 2 shows that the first component mainly separates the A population from the B populations whereas the second component separates B_1 from B_2 (S1 Fig.). The choice of K = 2 is evident when looking at the scree plot because the eigenvalues, which are proportional to the proportion of variance explained by each PC, drop beyond K = 2 and stay almost constant as K further increases (S2 Fig.).

We investigate the relationship between the communality statistic h^2 , which measures the proportion of variance explained by the first two PCs, and the F_{ST} statistic. We find a squared Pearson correlation between the two statistics larger than 98.8% in the simulations corresponding to Figs. 1 and 2 (S3 Fig.). For these two simulations, we look at the SNPs in the top 5% (respectively 1%) of the ranked lists based on h^2 and F_{ST} , and we find an overlap coefficient always larger than 95% for the lists provided by the two different statistics (respectively 93%). Providing a ranking of the SNPs similar to the ranking provided by F_{ST} is therefore possible without considering that individuals originate from predefined populations.




Figure 1: Repartition of the 5% top-ranked SNPs for each PCA-based statistic under a divergence model with two types of adaptive constraints. Thicker and colored lineages correspond to lineages where adaptation took place. The squared loadings with PC1 ρ_{j1}^2 pick a large proportion of SNPs involved in selection in the environment of population A whereas the squared loadings with PC2 ρ_{j2}^2 pick SNPs involved in selection in the environment of population B_1 . This difference is reflected in the different repartition of the top-ranked SNPs for the communality h^2 and the statistic h'^2 .

We then compare the performance of the different statistics based on PCA by investigating if the top-ranked SNPs (top 5%) manage to pick the SNPs involved in local adaptation (Fig. 1). The squared loadings ρ_{j1}^2 with the first PC pick SNPs involved in selection in the environment of population A (26% of the top 5%), just a few SNPs involved in selection in the B_1 environment (5%), and many false positive SNPs (69%). The squared loadings with the second PC ρ_{j2}^2 pick also false positives (63%) but also many SNPs involved in selection in the B_1 environment (36%) with just a few involved in selection in A (1%). When adaptation took place in two different evolutionary lineages of a divergence tree between populations, a genome scan based on PCA has the nice property that outlier loci correlated with PC1 or with PC2 correspond to adaptive constraints that occurred in different parts of the tree.

Because the communality h^2 gives more importance to the first PC (equation(2)), it picks preferentially the SNPs that are the most correlated with PC1. There is a large overlap of 84% between the 5% top-ranked lists provided by h^2 and ρ_{j1}^2 . Therefore, the communality statistic h^2 is more sensitive to ancient adaptation events that occurred in the environment of population A. By contrast, the alternative statistics h'^2 is more sensitive to recent adaptation events that occurred in the environment of population B_1 . When considering the top-ranked 5% of the SNPs, h'^2 captures few SNPs involved in selection in A (13% of the top 5%) and three times more SNPs related to adaptation in B_1 (37% of the top 5%). The h'^2 statistic is mostly influenced by the second principal component because the distribution of squared loadings corresponding to the second PC has a heavier tail, and this result holds for all simulations and data




Figure 2: Repartition of the 5% top-ranked SNPs of each PCA-based statistic under a divergence model with four types of adaptive constraints. Thicker and colored lineages correspond to lineages where adaptation occurred. The different types of SNPs picked by the squared loadings ρ_{j1}^2 and ρ_{j2}^2 is also found when comparing the communality h^2 and the statistic h'^2 .

we have investigated (S4 Fig.). To summarize, the h^2 and h'^2 statistics give too much importance to either PC1 or PC2 and they fail to capture in an equal manner both types of adaptive events occurring in the environment of populations A and B_1 .

We also investigate a more complex simulation in which adaptation occurs in the four branches of the divergence tree (Fig. 2). Among the 10,000 simulated SNPs, we assume that there are four sets of 125 adaptive SNPs with each set being related to adaptation in one of the four branches of the divergence tree. Compared to the simulation of Fig. 1, we find the same pattern of population structure (S1 Fig.). The squared loadings ρ_{j1}^2 with the first PC pick SNPs involved in selection in the branch that predates the split between B_1 and B_2 (19% of the top 5%), SNPs involved in selection in the environment of population A (9%), and false positive SNPs (67% of the top-ranked 5%). Except for false positives (63%), the squared loadings ρ_{j2}^2 with the second PC rather pick SNPs involved in selection in the environment of the B_1 and B_2 populations (19% for B_1 and 18% for B_2). Because the first PC discriminates the A population from the two B subpopulations (S1 Fig.), the SNPs most correlated with PC1 pick SNPs related to adaptation in the (red and green) branches that separate A from the two B populations, whereas the SNPs most correlated to PC2 pick SNPs related to adaptation in the two (blue and yellow) branches that separate B_1 from B_2 (Fig. 2).

In the models of Figs. 1 and 2, we also compare the false discovery rates obtained with the communality h^2 , with

the statistic h'^2 and with a Bayesian factor model implemented in the software *PCAdapt* (Duforet-Frebourg et al. 2014). For the optimal choice of K = 2, the statistic h'^2 and the Bayesian factor model provide the smallest false discovery rates (FDR, S5 Fig.). However, it is not a general result because the communality h^2 can provide the smallest FDR in another divergence model (S6 Fig.). When varying the value of K from K = 1 to K = 6, we find that the communality h^2 and the Bayesian approach are robust to over-specification of K (K > 3) whereas the false discovery rate obtained with h'^2 increases importantly as K increases beyond K = 2 (S5 Fig.).

The main results obtained under the divergence models can be described as follows. The principal components correspond to different evolutionary lineages of the divergence tree. The communality statistic h^2 provides similar list of candidate SNPs than F_{ST} and it is mostly influenced by the first principal component which can be problematic if other PCs also convey adaptive events. To counteract this limitation, which can potentially lead to the loss of important signals of selection, we show that looking at the squared loadings with each of the principal components provide adaptive SNPs that are related to different evolutionary events.

1000 Genomes data

Since we are interested in selective pressures that occur during the human diaspora out of Africa, we decide to exclude individuals whose genetic makeup is the result of recent admixture events (African Americans, Columbians, Puerto Ricans and Mexicans). The first three principal components capture population structure whereas the following components separate individuals within populations (Figs. 3 and S7). The first and second PCs ascertains population structure between Africa, Asia and Europe (Fig. 3) and the third principal component separates the Yoruba from the Luhya population (S7 Fig.). The decay of eigenvalues suggest to use K = 2 because the eigenvalues drop between K = 2 and K = 3 where a plateau of eigenvalue is reached (S2 Fig.).

When performing a genome scan with PCA, there are different choices of statistics. The first choice is the h^2 communality statistic. Using the three continents as labels, there is a squared correlation between h^2 and F_{ST} of $R^2 = 0.989$ (Fig. S3). To investigate if h^2 is mostly influenced by the first PC, we determine if the outliers for the h^2 statistics are related with PC1 or with PC2. Among the top 0.1% of SNPs with the largest values of h^2 , we find that 74% are in the top 0.1% of the squared loadings ρ_{j1}^2 corresponding to PC1 and 20% are in the top 0.1% of the squared loadings ρ_{j2}^2 corresponding to PC2. The repartition of the outliers for the communality h^2 confirms the results of the simulations, which showed that h^2 is mostly influenced by the first PC. The second possible choice of summary statistics is the h'^2 statistic. Investigating the repartition of the 0.1% outliers for h', we find that 0.005% are in the top 0.1% of the squared loadings ρ_{j1}^2 corresponding to PC1 and 85% are in the top 0.1% of the squared loadings ρ_{j2}^2 corresponding to PC2. The h'^2 statistic is mostly influenced by the second PC because the distribution of the V_{2j}^2 (normalized squared loadings) has a longer tail than the corresponding distribution for PC1 (S4 Fig.). Because the h^2




Figure 3: Principal component analysis with K = 2 applied to the 1000 Genomes data. The sampled populations are the following: British in England and Scotland (GBR), Utah residents with Northern and Western European ancestry (CEU), Finnish in Finland (FIN), Iberian populations in Spain (IBS), Toscani in Italy (TSI), Han Chinese in Bejing (CHB), Southern Han Chinese (CHS), Japanese in Tokyo (JPT), Luhya in Kenya (LWK), Yoruba in Nigeria (YRI).

statistic is mostly influenced by PC1 and h'^2 is mostly influenced by PC2 — confirming the results obtained under the divergence models — we rather decide to perform two separate genome scans based on the squared loadings ρ_{j1}^2 and ρ_{j2}^2 .

The two Manhattan plots based on the squared loadings for PC1 and PC2 are displayed in Figs. 4 and 5 (S1 table contains the loadings for all variants). Because of Linkage Disequilibrium, Manhattan plots generally produce clustered outliers. To investigate if the top 0.1% outliers are clustered in the genome, we count—for various window sizes—the proportion of contiguous windows containing at least one outlier. We find that the outlier SNPs correlated with PC1 or with PC2 are more clustered than expected if they would have been uniformly distributed among the 36, 536, 154 variants (S8 Fig.). Additionally, the clustering is larger for the outliers related to the second PC as they

cluster in fewer windows (S8 Fig.). As the genome scan for PC2 captures more recent adaptive events, it reveals larger genomic windows that experienced fewer recombination events.




Figure 4: Manhattan plot for the 1000 Genomes data of the squared loadings ρ_{j1}^2 with the first principal component. For sake of presentation, only the top-ranked SNPs (top 0.1%) are displayed and the 100 top-ranked SNPs are colored in red.




Figure 5: Manhattan plot for the 1000 Genomes data of the squared loadings ρ_{j2}^2 with the second principal component. For sake of presentation, only the top-ranked SNPs (top 0.1%) are displayed and the 100 top-ranked SNPs are colored in red.

The 100 SNPs that are the most correlated with the first PC are located in 24 genomic regions (S2 Table). Most of the regions contain just one or a few SNPs except a peak in the gene APPBP2 that contains 33 out of the 100 top SNPs, a peak encompassing the RTTN and CD226 genes containing 17 SNPS and a peak in the ATP1A1 gene containing 7 SNPs (Fig. 4). Confirming a larger clustering for PC2 outliers, the 100 SNPs that are the most correlated with PC2 cluster in fewer genomic regions (S3 Table). They are located in 14 genomic regions including a region overlapping with EDAR contains 44 top hits, two regions containing 8 SNPs and located in the pigmentation genes SLC24A5 and SLC45A2, and two regions with 7 top hit SNPs, one in the gene KCNMA1 and another one encompassing the RGLA/MYO5C genes (Fig. 5).

We perform Gene Ontology enrichment analyses using *Gowinda* for the SNPs that are the most correlated with PC1 and PC2. For PC1, we find, among others, enrichment ($FDR \le 5\%$) for ontologies related to the regulation of arterial blood pressure, the endocrine system and the immunity response (interleukin production, response to viruses) (S4 Table). For PC2, we find enrichment ($FDR \le 5\%$) related to olfactory receptors, keratinocyte and epidermal cell differentiation, and ethanol metabolism (S5 Table). We also search for polygenic adaptation by looking for biological pathways enriched with outlier genes (Daub et al. 2013). For PC1, we find one enriched ($FDR \le 5\%$) pathway consisting of the beta defensin pathway (S6 Table). The beta defensin pathway contains mainly genes involved in the innate immune system consisting of 36 defensin genes and of 2 Toll-Like receptors (TLR1 and TLR2). There are additionally 2 chemokine receptors (CCR2 and CCR6) involved in the beta defensin pathway. For PC2, we also find one enriched pathway consisting of fatty acid omega oxidation ($FDR \le 5\%$, S7 Table). This pathway consists of genes involved in alcohol oxidation (CYP, ALD and ALDH genes). Performing a less stringent enrichment analysis which can find pathways containing overlapping genes, we find more enriched pathways: the beta defensin and the defensin pathways for PC1 and ethanol oxidation, glycolysis/gluconeogenesis and fatty acid omega oxidation for PC2 (S8 Table).

To further validate the proposed list of candidate SNPs involved in local adaptation, we test for an enrichment of genic or non-synonymous SNP among the SNPs that are the most correlated with the PC. We measure the enrichment among outliers by computing odds ratio (Kudaravalli et al. 2009; Fagny et al. 2014). For PC1, we do not find significant enrichments (Table 1) except when measuring the enrichment of genic regions compared to non-genic regions (OR = 10.18 for the 100 most correlated SNPS, P < 5% using a permutation procedure). For PC2, we find an enrichment of genic regions among outliers as well as an enrichment of non-synonymous SNPs (Table 1). By contrast with the enrichment of genic regions for SNPs extremely correlated with the first PC, the enrichment for the variants extremely correlated with PC2 outliers is significant when using different thresholds to define outliers (Table 1).

Discussion

The promise of a fine characterization of natural selection in humans fostered the development of new analytical methods for detecting candidate genomic regions (Vitti et al. 2013). Population-differentiation based methods such as genome scans based on F_{ST} look for marked differences in allele frequencies between population (Holsinger and Weir 2009). Here, we show that the communality statistic h^2 , which measures the proportion of variance of a SNP that is explained by the first K principal components, provides a similar list of outliers than the F_{ST} statistic when there are K + 1 well-defined populations. In addition, the communality statistic h^2 based on PCA can be viewed as an extension of F_{ST} because it does not require to define populations in advance and can even be applied in the absence of well-defined populations.

Although it is possible to reproduce genome scans based on F_{ST} with PCA, the problems related to F_{ST} approaches remain pregnant. The drawbacks of F_{ST} arise because F_{ST} does not account for correlations of allele frequencies among subpopulations (Bonhomme et al. 2010; Bierne et al. 2013). An alternative presentation of the issues arising with F_{ST} is that it implicitly assumes either a model of instantaneous divergence between populations or an islandmodel (Bonhomme et al. 2010). Deviations from these models severely impact false discovery rates (Duforet-Frebourg et al. 2014). Viewing F_{ST} from the point of view of PCA provides new keys to understand the limits of genome scans based on F_{ST} or on the h^2 communality statistic. It is mostly influenced by the first principal component and the relative importance of the first PC increases with the difference between the first and second eigenvalues of the covariance matrix of the data. Because the first PC can represent ancient adaptive events, especially under population divergence models (McVean 2009), it explains why F_{ST} and the communality h^2 are biased toward ancient evolutionary events. We therefore proposed an alternative statistic h'^2 , which should give equal weights to the different PCs. However, the analysis of the 1000 Genomes data shows that outliers identified by h'^2 are almost always related to the last PC kept in the analysis. The difficulty of constructing a comprehensive summary statistic which captures all axes of genetic variation in an adequate manner leads us to perform a genome scan for each principal component separately.

The analysis of the phase 1 release of the 1000 Genomes data demonstrates the suitability of a genome scan based on PCA to detect signals of positive selection. We search for variants extremely correlated with the first PC, which corresponds to differentiation between Africa and Eurasia and with the second PC, which corresponds to differentiation between Europe and Asia. For the variants most correlated with the second PC, there is a significant enrichment of genic and non-synonymous SNPs whereas the enrichment is less detectable for variants related to the first PC. The enrichment analysis confirms that positive selection may favor local adaptation of human population by increasing differentiation in genic regions especially in non synonymous variants (Barreiro et al. 2008). Consistent with LD, we find that the candidate variants are clustered along the genome with a larger clustering for the variants correlated with the Europe-Asia axis of differentiation (PC2). The difference of clustering illustrates that statistical methods based on LD for detecting selection will perform differently depending on the time frame under which adaptation had the opportunity to occur (Sabeti et al. 2006). The fact that population divergence, and its concomitant adaptive events, between Europe and Asia is more recent that the out-of-Africa event is a putative explanation of the difference of clustering between PC1 and PC2 outliers. Explaining the difference of enrichment between PC1 and PC2 outliers is more difficult. The weaker enrichment for PC1 outliers can be attributed either to a larger number of false discoveries or to a larger importance of other forms of natural selection such as background selection (Hernandez et al. 2011).

When looking at the 100 SNPs most correlated with PC1 or PC2, we find genes for which selection in humans was already documented (9/24 for PC1 and 5/14 for PC2, Table S9). Known targets for selection include genes involved in pigmentation (MATP, OCA2 for PC1 and SLC45A2, SLC24A5, and MYO5C for PC2), in the regulation of sweating (EDAR for PC2), and in adaptation to pathogens (DARC, SLC39A4, and VAV2 for PC1). A 100 kb region in the vicinity of the APPBPP2 gene contains one third of the 100 SNPs most correlated with PC1. This APPBPP2 region is a known candidate for selection and has been identified by looking for miRNA binding sites with extreme population differentiation (Li et al. 2012). APPBPP2 is a nervous system gene that has been associated with Alzheimer disease, and it may have experienced a selective sweep (Williamson et al. 2007). For some SNPs in APPBPP2, the differences of allele frequencies between Eurasiatic population and SubSaharan populations from Africa are of the order of 90% (http://www.popgen.uchicago.edu/ggv) calling for a further functional analysis. Moreover, looking at the 100 SNPs most correlated with PC1 and PC2 confirms the importance of non-coding RNA (FAM230B, D21S2088E, LOC100133461, LINC00290, LINC01347, LINC00681), such as miRNA (MIR429), as a substrate for human adaptation (Li et al. 2012; Grossman et al. 2013). Among the other regions with a large number of candidate SNPs, we also found the RTTN/CD226 regions, which contain many SNPs correlated with PC1. In different selection scans, the RTTN genes has been detected (Carlson et al. 2005; Barreiro et al. 2008), and it is involved in the development of the human skeletal system (Wu and Zhang 2010). An other region with many SNPs correlated with PC1 contains the ATP1A1 gene involved in osmoregulation and associated with hypertension (Gurdasani et al. 2015). The regions containing the largest number of SNPs correlated with PC2 are well-documented instances of adaptation in humans and includes the EDAR, SLC24A5 and SLC45A2 genes. The KCNMA1gene contains 7 SNPs correlated with PC2 and is involved in breast cancer and obesity (Oeggerli et al. 2012; Jiao et al. 2011) and deserves additional population genomic and functional studies to better understand how adaptation took place. As for KCNMA1, the MYO5C has already been reported in selection scans although no mechanism of biological adaption has been proposed yet (Chen et al. 2010; Fumagalli et al. 2010). To summarize, the list of most correlated SNPs with the PCs identifies well-known genes related to biological adaptation in humans (EDAR, SLC24A5, SLC45A2, DARC), but also provides candidate genes that deserve further studies such as the APPBPP2, TP1A1, RTTN, KCNMA and MYO5C genes, as well as the ncRNAs listed above.

We also show that the scan based on PCA can also be used to detect more subtle footprints of positive selection. We conduct an enrichment analysis that detects polygenic adaptation at the level of biological pathways (Daub et al. 2013). We find that the genes in the beta-defensin pathway are enriched in SNPs correlated with PC1. The beta-defensin genes are key components of the innate immune system and have evolved through positive selection in the catarrhine primate lineages (Hollox and Armour 2008). As for the HLA complex, some beta-defensing genes (DEFB1, DEFB127) show evidence of long-term balancing selection with major haplotypic clades coexisting since millions of years (Cagliani et al. 2008; Hollox and Armour 2008). We also find that the genes in the omega fatty acid oxidation pathways are enriched in SNPs correlated with PC2. This pathway was also found when investigating polygenic adaptation to altitude in humans (Foll et al. 2014). The proposed explanation was that omega oxidation becomes a more important metabolic pathway when beta oxidation is defective, which can occur in case of hypoxia (Foll et al. 2014). However, this explanation is not valid in the context of the 1000 Genomes data when there are no populations living in hypoxic environments. Proposing phenotypes on which selection operates is complicated by the fact that the omega fatty acid oxidation pathway strongly overlaps with two other pathways: ethanol oxidation and glycolysis. Evidence of selection on the alcohol dehydrogenase locus have already been provided (Han et al. 2007) with some authors proposing that a lower risk for alcoholism might have been beneficial after rice domestication in Asia (Peng et al. 2010). This hypothesis is only speculative and we lack a confirmed biological mechanism explaining the enrichment of the fatty acid oxidation pathway. More generally, the enrichment of the beta-defensin and of the omega fatty acid oxidation pathways confirms the importance of pathogenic pressure and of metabolism in human adaptation to different environments (Hancock et al. 2008; Barreiro and Quintana-Murci 2009; Fumagalli et al. 2011; Daub et al. 2013).

In conclusion, we propose a new approach to scan genomes for local adaptation that works with individual genotype data. Because the method is efficiently implemented in the software *PCAdapt*, analyzing 36, 536, 154 SNPs took only 502 minutes using a single core of an Intel(R) Xeon(R) (E5-2650, 2.00GHz, 64 bits). Even with low-coverage sequence data (3x), PCA-based statistics retrieve well-known examples of biological adaptation which is encouraging for future whole-genome sequencing project, especially for non-model species, aiming at sampling many individuals with limited cost.

Materials and Methods

Simulations of the divergence models

We assume that each population has a constant effective population size of Ne = 1,000 diploid individuals, with 50 individuals sampled in each population. The genotypes consist of 10,000 independent SNPs. The simulations were performed in two steps. In the first step, we used the software *ms* to simulate genetic diversity (Hudson 2002) in the

ancestral population. We kept only variants with a minor allele frequency larger than 5% at the end of the first step. The second step was performed with *SimuPOP* (Peng and Kimmel 2005) and simulations were started using the allele frequencies generated with *ms* in the ancestral population. Looking forward in time, we consider that there are 100 generations between the initial split and the following split between the two *B* subpopulations, and 200 generations following the split between the two *B* subpopulations. In the simulation of Fig. 1, we assume that 250 SNPs confer a selective advantage in the branch leading to population A and 250 other SNPs confer a selective advantage in the branch leading to population B_1 . We consider an additive model for selection with a selection coefficient of s = 1.025 for heterozygotes. For the simulation of Fig. 2, we assume that there are four non-overlapping sets of 125 adaptive SNPs with each set being related to adaptation in one of the four branches of the divergence tree. A SNP can confer a selective advantage in a single branch only. To estimate the F_{ST} statistic, we consider the estimator of Weir and Cockerham(Weir and Cockerham 1984).

1000 Genomes data

We downloaded the 1000 Genomes data (phase 1 v3) at ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/ phasel/analysis_results/integrated_call_sets/ (Altshuler et al. 2012). We kept low-coverage genome data and excluded exomes and triome data to minimize variation in read depth. Filtering the data resulted in a total of 36, 536, 154 SNPs that have been typed on 1, 092 individuals. Because the analysis focuses on biological adaptation that took place during the human diaspora out of Africa, we removed recently admixed populations (Mexican, Columbian, PortoRican, and AfroAmerican individuals from the Southwest of the USA). The resulting dataset contains 850 individuals coming from Asia (two Han Chinese and one Japanese populations), Africa (Yoruba and Luhya) and Europe (Finish, British in England and Scotland, Iberian, Toscan, and Utah residents with Northern and Western European ancestry).

Enrichment analyses

We used *Gowinda* (Kofler and Schlötterer 2012) to test for enrichment of Gene Ontology (GO). A gene is considered as a candidate if there is at least one of the most correlated SNPs (top 1%) that is mapped to the gene (within an interval of 50Kb upstream and downstream of the gene). Enrichment was computed as the proportion of genes containing at least one outlier SNPs among the genes of the given GO category that are present in the dataset. In order to sample a null distribution for enrichment, *Gowinda* performs resampling without replacement of the SNPs. We used the *-gene* option of *Gowinda* that assumes complete linkage within genes.

We performed a second enrichment analysis to determine if outlier SNPs are enriched for genic regions. We

computed odds ratio (Kudaravalli et al. 2009)

$$OR = \frac{Pr(genic|outlier)}{Pr(not genic|outlier)} \frac{Pr(not genic|not outlier)}{Pr(genic|not outlier)}.$$

We implemented a permutation procedure to test if an odds ratio is significantly larger than 1 (Fagny et al. 2014). The same procedure was applied when testing for enrichment of UTR regions and of non-synonymous SNPs.

Polygenic adaptation

To test for polygenic adaptation, we determined whether genes in a given biological pathway show a shift in the distribution of the loadings (Daub et al. 2013). We computed the SUMSTAT statistic for testing if there is an excess of selection signal in each pathway (Daub et al. 2013). We downloaded 63,693 known genes from the UCSC website and we mapped SNPs to a gene if a SNP is located within a gene transcript or within 50kb of a gene. A total of 18,267 genes were mapped with this approach. We downloaded 2,681 gene sets from the NCBI Biosystems database. After discarding genes that were not part of the aforementioned gene list, removing gene sets with less than 10 genes and pooling nearly identical gene sets, we kept 1,532 sets for which we test if there was a shift of the distribution of loadings. To account for redundancy of genes within pathways, we applied a pruning method, and we further considered the Gaussian approximation to test for significant shifts (Daub et al. 2013).

Acknowledgments

This work has been supported by the LabEx PERSYVAL-Lab (ANR-11-LABX-0025-01) and the ANR AGRHUM project (ANR-14-CE02-0003).

Supplementary Material

Available at http://membres-timc.imag.fr/Michael.Blum/SuppMat_PCA.zip

Literature Cited

Altshuler, D. et al., 2012. An integrated map of genetic variation from 1,092 human genomes. Nature 491:56–65.

Anderson, E., Z. Bai, C. Bischof, S. Blackford, J. Demmel, J. Dongarra, J. Du Croz, A. Greenbaum, S. Hammarling,A. McKenney, and D. Sorensen, 1999. LAPACK Users' Guide. Third ed. Society for Industrial and Applied Mathematics, Philadelphia, PA.

- Barreiro, L. B., G. Laval, H. Quach, E. Patin, and L. Quintana-Murci, 2008. Natural selection has driven population differentiation in modern humans. Nature genetics 40:340–345.
- Barreiro, L. B. and L. Quintana-Murci, 2009. From evolutionary genetics to human immunology: how selection shapes host defence genes. Nature Reviews Genetics 11:17–30.
- Bierne, N., D. Roze, and J. J. Welch, 2013. Pervasive selection or is it...? why are F_{ST} outliers sometimes so frequent? Molecular ecology 22:2061–2064.
- Bonhomme, M., C. Chevalet, B. Servin, S. Boitard, J. Abdallah, S. Blott, and M. SanCristobal, 2010. Detecting selection in population trees: the Lewontin and Krakauer test extended. Genetics 186:241–262.
- Cadima, J. and I. T. Jolliffe, 1995. Loading and correlations in the interpretation of principle compenents. Journal of Applied Statistics 22:203–214.
- Cagliani, R., M. Fumagalli, S. Riva, U. Pozzoli, G. P. Comi, G. Menozzi, N. Bresolin, and M. Sironi, 2008. The signature of long-standing balancing selection at the human defensin beta-1 promoter. Genome Biol 9:R143.
- Carlson, C. S., D. J. Thomas, M. A. Eberle, J. E. Swanson, R. J. Livingston, M. J. Rieder, and D. A. Nickerson, 2005. Genomic regions exhibiting positive selection identified from dense genotype data. Genome research 15:1553–1565.
- Chen, H., N. Patterson, and D. Reich, 2010. Population differentiation as a test for selective sweeps. Genome research 20:393–402.
- Colonna, V., Q. Ayub, Y. Chen, L. Pagani, P. Luisi, M. Pybus, E. Garrison, Y. Xue, C. Tyler-Smith, et al., 2014. Human genomic regions with exceptionally high levels of population differentiation identified from 911 whole-genome sequences. Genome biology 15:R88.
- Daub, J. T., T. Hofer, E. Cutivet, I. Dupanloup, L. Quintana-Murci, M. Robinson-Rechavi, and L. Excoffier, 2013. Evidence for polygenic adaptation to pathogens in the human genome. Molecular biology and evolution 30:1544– 1558.
- Duforet-Frebourg, N., E. Bazin, and M. G. B. Blum, 2014. Genome scans for detecting footprints of local adaptation using a Bayesian factor model. Molecular Biology and Evolution 31:2483–2495.
- Excoffier, L., P. E. Smouse, and J. M. Quattro, 1992. Analysis of molecular variance inferred from metric distances among dna haplotypes: application to human mitochondrial DNA restriction data. Genetics 131:479–491.
- Fagny, M., E. Patin, D. Enard, L. B. Barreiro, L. Quintana-Murci, and G. Laval, 2014. Exploring the occurrence of classic selective sweeps in humans using whole-genome sequencing data sets. Molecular biology and evolution 31:1850–1868.

- Fariello, M. I., S. Boitard, H. Naya, M. SanCristobal, and B. Servin, 2013. Detecting signatures of selection through haplotype differentiation among hierarchically structured populations. Genetics 193:929–941.
- Foll, M., O. E. Gaggiotti, J. T. Daub, A. Vatsiou, and L. Excoffier, 2014. Widespread signals of convergent adaptation to high altitude in Asia and America. The American Journal of Human Genetics 95:394–407.
- Fumagalli, M., U. Pozzoli, R. Cagliani, G. P. Comi, N. Bresolin, M. Clerici, and M. Sironi, 2010. Genome-wide identification of susceptibility alleles for viral infections through a population genetics approach. PLoS genetics 6:e1000849.
- Fumagalli, M., M. Sironi, U. Pozzoli, A. Ferrer-Admettla, L. Pattini, and R. Nielsen, 2011. Signatures of environmental genetic adaptation pinpoint pathogens as the main selective pressure through human evolution. PLoS genetics 7:e1002355.
- Grossman, S. R., K. G. Andersen, I. Shlyakhter, S. Tabrizi, S. Winnicki, A. Yen, D. J. Park, D. Griesemer, E. K. Karlsson, S. H. Wong, et al., 2013. Identifying recent adaptations in large-scale genomic data. Cell 152:703–713.
- Günther, T. and G. Coop, 2013. Robust identification of local adaptation from allele frequencies. Genetics 195:205–220.
- Gurdasani, D., T. Carstensen, F. Tekola-Ayele, L. Pagani, I. Tachmazidou, K. Hatzikotoulas, S. Karthikeyan, L. Iles, M. O. Pollard, A. Choudhury, et al., 2015. The African genome variation project shapes medical genetics in Africa. Nature 517:327–332.
- Hamblin, M. T., E. E. Thompson, and A. Di Rienzo, 2002. Complex signatures of natural selection at the duffy blood group locus. The American Journal of Human Genetics 70:369–383.
- Han, Y., S. Gu, H. Oota, M. V. Osier, A. J. Pakstis, W. C. Speed, J. R. Kidd, and K. K. Kidd, 2007. Evidence of positive selection on a class I *adh* locus. The American Journal of Human Genetics 80:441–456.
- Hancock, A. M., D. B. Witonsky, E. Ehler, G. Alkorta-Aranburu, C. Beall, A. Gebremedhin, R. Sukernik, G. Utermann, J. Pritchard, G. Coop, et al., 2010. Human adaptations to diet, subsistence, and ecoregion are due to subtle shifts in allele frequency. Proceedings of the National Academy of Sciences 107:8924–8930.
- Hancock, A. M., D. B. Witonsky, A. S. Gordon, G. Eshel, J. K. Pritchard, G. Coop, and A. Di Rienzo, 2008. Adaptations to climate in candidate genes for common metabolic disorders. PLoS Genetics 4:e32.
- Hernandez, R. D., J. L. Kelley, E. Elyashiv, S. C. Melton, A. Auton, G. McVean, G. Sella, M. Przeworski, et al., 2011. Classic selective sweeps were rare in recent human evolution. science 331:920–924.

- Hollox, E. J. and J. A. Armour, 2008. Directional and balancing selection in human beta-defensins. BMC evolutionary biology 8:113.
- Holsinger, K. E. and B. S. Weir, 2009. Genetics in geographically structured populations: defining, estimating and interpreting FST. Nature Reviews Genetics 10:639–650.
- Hudson, R. R., 2002. Generating samples under a Wright-Fisher neutral model of genetic variation. Bioinformatics 18:337–338.
- Jiao, H., P. Arner, J. Hoffstedt, D. Brodin, B. Dubern, S. Czernichow, F. van't Hooft, T. Axelsson, O. Pedersen, T. Hansen, et al., 2011. Genome wide association study identifies KCNMA1 contributing to human obesity. BMC medical genomics 4:51.
- Jolliffe, I., 2005. Principal component analysis. Wiley Online Library.
- Kofler, R. and C. Schlötterer, 2012. Gowinda: unbiased analysis of gene set enrichment for genome-wide association studies. Bioinformatics 28:2084–2085.
- Kudaravalli, S., J.-B. Veyrieras, B. E. Stranger, E. T. Dermitzakis, and J. K. Pritchard, 2009. Gene expression levels are a target of recent natural selection in the human genome. Molecular biology and evolution 26:649–658.
- Laloë, D. and M. Gautier, 2011. On the genetic interpretation of between-group PCA on SNP data. HAL hal-00661214.
- Li, J., Y. Liu, X. Xin, T. S. Kim, E. A. Cabeza, J. Ren, R. Nielsen, J. L. Wrana, and Z. Zhang, 2012. Evidence for positive selection on a number of MicroRNA regulatory interactions during recent human evolution. PLoS genetics 8:e1002578.
- Luikart, G., P. R. England, D. Tallmon, S. Jordan, and P. Taberlet, 2003. The power and promise of population genomics: from genotyping to genome typing. Nature reviews genetics 4:981–994.
- McVean, G., 2009. A genealogical interpretation of principal components analysis. PLoS Genetics 5:e1000686.
- Nielsen, R., 2005. Molecular signatures of natural selection. Annu. Rev. Genet. 39:197-218.
- Oeggerli, M., Y. Tian, C. Ruiz, B. Wijker, G. Sauter, E. Obermann, U. Güth, I. Zlobec, M. Sausbier, K. Kunzelmann, et al., 2012. Role of KCNMA1 in breast cancer. PloS one 7:e41664.
- Patterson, N., A. L. Price, and D. Reich, 2006. Population structure and eigenanalysis. PLoS Genetics 2:e190.
- Peng, B. and M. Kimmel, 2005. simuPOP: a forward-time population genetics simulation environment. Bioinformatics 21:3686–3687.

- Peng, Y., H. Shi, X.-b. Qi, C.-j. Xiao, H. Zhong, Z. M. Run-lin, and B. Su, 2010. The ADH1B Arg47His polymorphism in East Asian populations and expansion of rice domestication in history. BMC evolutionary biology 10:15.
- Pickrell, J. K., G. Coop, J. Novembre, S. Kudaravalli, J. Z. Li, D. Absher, B. S. Srinivasan, G. S. Barsh, R. M. Myers, M. W. Feldman, et al., 2009. Signals of recent positive selection in a worldwide sample of human populations. Genome research 19:826–837.
- Sabeti, P., S. Schaffner, B. Fry, J. Lohmueller, P. Varilly, O. Shamovsky, A. Palma, T. Mikkelsen, D. Altshuler, and E. Lander, 2006. Positive natural selection in the human lineage. science 312:1614–1620.
- Sabeti, P. C., P. Varilly, B. Fry, J. Lohmueller, E. Hostetter, C. Cotsapas, X. Xie, E. H. Byrne, S. A. McCarroll, R. Gaudet, et al., 2007. Genome-wide detection and characterization of positive selection in human populations. Nature 449:913–918.
- Suhr, D. D., 2009. Principal component analysis vs. exploratory factor analysis. SUGI 30 Proceedings Pp. 203–230.
- Vitti, J. J., S. R. Grossman, and P. C. Sabeti, 2013. Detecting natural selection in genomic data. Annual review of genetics 47:97–120.
- Weir, B. S. and C. C. Cockerham, 1984. Estimating F-statistics for the analysis of population structure. evolution Pp. 1358–1370.
- Williamson, S. H., M. J. Hubisz, A. G. Clark, B. A. Payseur, C. D. Bustamante, and R. Nielsen, 2007. Localizing recent adaptive evolution in the human genome. PLoS genetics 3:e90.
- Wu, D.-D. and Y.-P. Zhang, 2010. Positive selection drives population differentiation in the skeletal genes in modern humans. Human molecular genetics 19:2341–2346.
- Yang, W.-Y., J. Novembre, E. Eskin, and E. Halperin, 2012. A model-based approach for analysis of spatial structure in genetic data. Nature genetics 44:725–731.

	$\mathrm{top} 0.1\%$	$\mathrm{top} 0.01\%$	$\mathrm{top} 0.005\%$	top 100 SNPs
pc1 - genic/nogenic	$1,60^{*}$	1,24	1,09	1,93
pc1 - nonsyn/all	1,70	1,18	2,42	$10,07^{*}$
pc1 - UTR/all	1,37	0,80	1,65	3,44
pc2 - genic/nogenic	1,51*	2,27	$4,73^{**}$	4,44*
pc2 - nonsyn/all	1,72	$4,66^{*}$	7,40	$12, 18^{*}$
pc2 - UTR/all	1,68	$4,01^{*}$	3,36	2,73

Table 1: Enrichment measured with Odds ratio (OR) of the variants most correlated with the principal components. Enrichment significant at the 1% (resp. 5%) level are indicated with ** (resp. *).