

A SHORT PROOF OF SMOOTH IMPLIES FLAT

JESÚS CONDE-LAGO

ABSTRACT. Proofs that a smooth morphism is flat available in the literature are long and difficult. We give a short proof of this fact.

Let $(A, \mathfrak{m}) \rightarrow (B, \mathfrak{n})$ be a local homomorphism of noetherian local rings. It is said [EGA IV, 0_{IV}.19.3.1] that B is formally smooth over A (for the \mathfrak{n} -adic topology) if for any A -algebra C and any nilpotent ideal N of C , each A -algebra homomorphism $u : B \rightarrow C/N$ satisfying $u(\mathfrak{n}^k) = 0$ for some k , factorizes as $B \xrightarrow{v} C \xrightarrow{p} C/N$ where p is the canonical map.

A fundamental result, proved by Grothendieck in [EGA IV, 0_{IV}.19.7.1], says that if B is formally smooth over A then B is flat over A . His proof is long. In the following few years shorter proofs appeared, but based on results whose proofs were long.

In many situations, one is concerned exclusively with *smooth* algebras, i.e., formally smooth algebras essentially of finite type [EGA IV, 17.3.1, 17.1.2] (which are then formally smooth for the discrete topology). In this case, a shorter proof of this result can be seen in [Bo, §7 n.10, Lemme 5], but it uses non-trivial results on smoothness and regularity. There are also two short papers deducing flatness from the Jacobian criterion of smoothness in some particular cases (for an extension of polynomial algebras over a field in [MRW] and in characteristic zero in [MR]).

In this paper we deduce flatness from the very definition of smoothness in a short way.

Theorem. *Let A be a noetherian ring and B a smooth A -algebra. Then B is flat as A -module.*

Proof. Let R be a localization of a polynomial A -algebra of finite type such that we have a surjective homomorphism $\varphi : R \rightarrow B$. Let $I = \ker \varphi$. Then $\hat{R}/\hat{I} = R/I = B$ and so $\hat{I} = \ker(\hat{\varphi} : \hat{R} \rightarrow B)$. As in [EGA IV, 0_{IV}.19.3.11] we are going to see that we have a section $B \rightarrow \hat{R}$. By induction on $i \geq 2$, we construct A -algebra homomorphisms $f_i : B \rightarrow \hat{R}/\hat{I}^i$

$$\begin{array}{ccccccc}
 & & & & B & & \\
 & & & & \downarrow f_{i-1} & & \\
 & & & \swarrow f_i & & & \\
 0 & \longrightarrow & \hat{I}^{i-1}/\hat{I}^i & \longrightarrow & \hat{R}/\hat{I}^i & \longrightarrow & \hat{R}/\hat{I}^{i-1} \longrightarrow 0
 \end{array}$$

2000 *Mathematics Subject Classification.* 14B25 (primary); 13B40 (secondary).
Key words and phrases. Formal smoothness, flat algebra.

making commutative the triangles. Since \hat{R} is complete for the \hat{I} -adic topology, we have a limit homomorphism

$$f = \varprojlim_i f_i : B \longrightarrow \varprojlim_i \hat{R}/\hat{I}^i = \hat{R}$$

such that $\hat{\varphi}f = \text{id}_B$. Therefore B is a direct summand of the flat A -module \hat{R} and thus also flat. \square

REFERENCES

- [Bo] Bourbaki, N. *Algèbre Commutative*, Chapitre X. Masson, Paris, 1998.
- [EGA IV] Grothendieck, A. *Éléments de Géométrie Algébrique*, Chap. IV. Publ. Math. IHES 20, 24, 28, 32, 1964-1967.
- [MR] Majadas, J.; Rodicio, A.G. *A short proof of Jacobian hypothesis implies flatness*. Comm. Algebra, 28 (2000), 4995-4996.
- [MRW] Miyanishi, M.; Robbiano, L.; Wang, S.S.-S. *A brief proof of Jacobian hypothesis implies flatness*. Proc. Amer. Math. Soc., 109 (1990), 327-330.

DEPARTAMENTO DE ÁLXEBRA, FACULDADE DE MATEMÁTICAS, UNIVERSIDADE DE SANTIAGO DE COMPOSTELA, E-15782 SANTIAGO DE COMPOSTELA, SPAIN
E-mail address: `jesus.conde@usc.es`