

GLOBAL DYNAMICS BELOW EXCITED SOLITONS FOR THE NONLINEAR SCHRÖDINGER EQUATION WITH A POTENTIAL

KENJI NAKANISHI

ABSTRACT. Consider the nonlinear Schrödinger equation (NLS) with a potential with a single negative eigenvalue. It has solitons with negative small energy, which are asymptotically stable, and, if the nonlinearity is focusing, then also solitons with positive large energy, which are unstable. In this paper we classify the global dynamics below the second lowest energy of solitons under small mass and radial symmetry constraints.

CONTENTS

1. Introduction	2
1.1. Background and motivation	2
1.2. Setting and the main result	3
1.3. Difficulties and ideas in the proof	5
1.4. Notation	7
1.5. Assumptions on V	7
Acknowledgements	7
2. Standing waves	8
2.1. Energy functionals	8
2.2. Small solitons	9
2.3. Focusing case $\mathfrak{s} = +$	10
2.4. Defocusing case $\mathfrak{s} = -$	12
3. Blow-up below the excited energy	14
4. Modulation and linearized equations around the ground state	15
5. Linearized profile decomposition	20
6. Nonlinear perturbation estimates	24
7. Nonlinear profile decomposition	29
8. Scattering below the excited energy	34
Appendix A. Decay of the potential	37
Appendix B. Table of Notation	37
References	38

1. INTRODUCTION

1.1. Background and motivation. Nonlinear dispersive equations have solutions with various types of behavior in time, typically *scattering* (globally dispersive), *blow-up*, and solitary waves, i.e., *solitons*. In the recent years, especially since the work of Kenig and Merle [12], global dynamics leading to those different types have been revealed among large general solutions, so that one can partially predict evolution of each solution from the initial data. Kenig and Merle [12] studied the energy-critical NLS

$$i\dot{u} - \Delta u = |u|^4 u, \quad u(t, x) : \mathbb{R}^{1+3} \rightarrow \mathbb{C}, \quad (1.1)$$

and proved that all solutions with energy less than the ground state W

$$E(u) = \int_{\mathbb{R}^3} \frac{|\nabla u|^2}{2} - \frac{|u|^6}{6} dx < E(W), \quad (1.2)$$

$$W(x) := (1 + |x|^2/3)^{-1/2}, \quad -\Delta W = W^5,$$

either scatter or blow-up, and that the two types of behavior are distinguished by some explicit functionals of the initial data. For example,

$$K(u(0)) = \int_{\mathbb{R}^3} |\nabla u(0)|^2 - |u(0)|^6 dx \begin{cases} \geq 0 & \implies \text{scattering,} \\ < 0 & \implies \text{blow-up.} \end{cases} \quad (1.3)$$

The distinction is essentially the same as that in the classical result for the nonlinear Klein-Gordon equation by Payne and Sattinger [18] into global existence vs. blow-up, but the crucial aspect of Kenig-Merle's work is to reveal and exploit the global dispersion in the scattering part. It was extended to the threshold energy $E(u) \leq E(W)$ by Duyckaerts and Merle [6], and then slightly above the ground state by Schlag and the author [15], for the nonlinear Klein-Gordon equation

$$\ddot{u} - \Delta u + u = u^3, \quad u(t, x) : \mathbb{R}^{1+3} \rightarrow \mathbb{R},$$

$$E(u) = \int_{\mathbb{R}^3} \frac{|\dot{u}|^2 + |\nabla u|^2 + |u|^2}{2} - \frac{|u|^4}{4} dx < E(Q) + \varepsilon^2, \quad (1.4)$$

where $Q \in H^2(\mathbb{R}^3)$ is the unique positive radial solution or the ground state of

$$-\Delta Q + Q = Q^3. \quad (1.5)$$

The types of behavior in that case are separated into 9 sets of solutions by center-stable and center-unstable manifolds of the ground state, and the mechanism of transition between scattering and blow-up is revealed. Furthermore, Duyckaerts, Kenig and Merle [5] established a complete classification of asymptotic behavior of solutions, for the energy-critical wave equation

$$\ddot{u} - \Delta u = u^5, \quad u(t, x) : \mathbb{R}^{1+3} \rightarrow \mathbb{R}, \quad (1.6)$$

in terms of resolution into solitons (i.e. rescaled W), without any size restriction on the initial data. All of these works have been extended to several equations and settings including the above examples, except the soliton resolution which is yet limited to variants of energy-critical wave equations.

General dynamics are, however, far more complicated for more general or physical equations. In particular, many equations, especially of the NLS type, have many

solitons, differing in shape, energy, stability, etc. Heuristically, unstable solitons are expected to collapse into stable solitons, radiating dispersive waves. For small solitons of the NLS with a decaying potential $V(x)$, Tsai and Yau [21–24] first proved such a phenomenon, as well as asymptotic stability of the ground soliton, in the case $-\Delta + V$ has two well-positioned negative eigenvalues. Since then, there have been intensive studies (cf. [3, 8, 14, 19, 20]) on global behavior of small solutions including many solitons, but very little is rigorously known about dynamical relation between solitons which are neither close nor similar to each other. It seems hard in such cases to construct or control solutions in a precise way along some anticipated evolution. A more natural strategy is to deal altogether with general solutions including or at least close to those solitons, with less precise information on individual trajectories.

1.2. Setting and the main result. As a first step toward the above problem, we consider the NLS with a potential

$$\begin{aligned} i\ddot{u} + Hu &= \mathfrak{s}|u|^2u, \quad H := -\Delta + V, \quad \mathfrak{s} = \pm, \\ u(t, x) &: \mathbb{R}^{1+3} \rightarrow \mathbb{C}, \quad V(x) : \mathbb{R}^3 \rightarrow \mathbb{R}, \end{aligned} \quad (1.7)$$

in the simplest non-trivial setting, namely the case with the unique eigenvalue

$$H\phi_0 = e_0\phi_0, \quad e_0 < 0, \quad 0 < \phi_0 \in H^2(\mathbb{R}^3), \quad \|\phi_0\|_{L^2(\mathbb{R}^3)} = 1, \quad (1.8)$$

with $\text{spec}(H|_{\phi_0^\perp}) = [0, \infty)$ absolutely continuous, and the radial symmetry restriction

$$u(t, x) = u(t, |x|), \quad V(x) = V(|x|). \quad (1.9)$$

Hence the initial data set is the radial subspace of the Sobolev space

$$H_r^1(\mathbb{R}^3) := \{u \in L^2(\mathbb{R}^3) \mid \nabla u \in L^2(\mathbb{R}^3), u(x) = u(|x|)\}. \quad (1.10)$$

The nonlinearity can be either defocusing $\mathfrak{s} = -$ or focusing $\mathfrak{s} = +$. In the focusing case $\mathfrak{s} = +$, the above equation is one of the simplest equations with both stable and unstable solitons, where the former is small and the latter is large. The goal of this study is a complete description of global dynamics in a fairly large solution space, containing both the stable and the unstable solitons. In this paper, we consider the region of small mass and an upper energy constraint which eliminates the unstable solitons. An implication of the main result is that if an unstable (large) soliton with small mass and the second largest energy is perturbed to decrease its energy and the mass, then it either blows up or collapses into a (small) ground state soliton, radiating most of the energy (which is large) into a dispersive wave. The two types of behavior is distinguished by a functional of the initial data, similarly to Kenig-Merle or Payne-Sattinger.

In order to state the main result, we need a few more assumptions on V . A simple sufficient condition is that V is in the Schwartz class and H has no resonance:

$$\dot{H}^1(\mathbb{R}^3) \ni \varphi, \quad (-\Delta + V)\varphi = 0 \implies \varphi = 0. \quad (1.11)$$

The existence of small solitons is well known in the above setting. The function $u(t, x) = e^{-it\omega}\varphi(x)$ is a solution of (1.7) iff

$$(H + \omega)\varphi = \mathfrak{s}|\varphi|^2\varphi. \quad (1.12)$$

In this paper we call a solution of (1.12) a soliton, denoting the set of solitons by

$$\mathcal{S} := \{\varphi \in H_r^1(\mathbb{R}^3) \mid \exists \omega \in \mathbb{R} \text{ s.t. (1.12)}\} \quad (1.13)$$

and the energy (Hamiltonian) and the mass (charge) by

$$\mathbb{E}(u) := \int_{\mathbb{R}^3} \frac{|\nabla u|^2 + V|u|^2}{2} - \frac{\mathfrak{s}|u|^4}{4} dx, \quad \mathbb{M}(u) := \int_{\mathbb{R}^3} \frac{|u|^2}{2} dx, \quad (1.14)$$

which are continuous on $H^1(\mathbb{R}^3)$ and conserved for (1.7). For each fixed mass $\mathbb{M}(\varphi) = \mu > 0$, we can define the energy levels of solitons by induction on $j = 0, 1, 2, \dots$,

$$\mathcal{E}_j(\mu) := \inf\{\mathbb{E}(\varphi) \mid \varphi \in \mathcal{S}, \mathbb{M}(\varphi) = \mu, \mathbb{E}(\varphi) > \mathcal{E}_{j-1}(\mu)\}, \quad (1.15)$$

where $\mathcal{E}_{-1}(\mu) := -\infty$ and $\inf \emptyset := \infty$, then classify the solitons

$$\mathcal{S}_j := \{\varphi \in \mathcal{S} \mid \mathbb{E}(\varphi) = \mathcal{E}_j(\mathbb{M}(\varphi))\}. \quad (1.16)$$

\mathcal{S}_0 is the set of least energy solitons, namely *the ground states*, while \mathcal{S}_j is the j -th *excited state* for $j \geq 1$. In this paper, we are concerned only with \mathcal{S}_0 and \mathcal{S}_1 .

It is easy to observe that the ground states for small mass are bifurcation from 0 generated by the linear ground state ϕ_0 in (1.8). More precisely, there exists $0 < z_b \ll 1$ and a C^1 map

$$(\Phi, \Omega) : D_b := \{z \in \mathbb{C} \mid |z|^2 < 2z_b\} \rightarrow H_r^1(\mathbb{R}^3) \times \mathbb{R} \quad (1.17)$$

such that $(\varphi, \omega) = (\Phi[z], \Omega[z])$ solves (1.12) for each $z \in D_b$ and

$$\Phi[z] = z\phi_0 + \gamma, \quad \gamma \perp \phi_0, \quad \|\gamma\|_{H^1} \lesssim |z|^3. \quad (1.18)$$

See [9] for a proof in a more general setting. We can prove that $\Phi(D_b) = \mathcal{S}_0$ under the small mass constraint $\mathbb{M} < z_b^2$, while the first excited energy satisfies

$$\mathcal{E}_1(\mu) = \begin{cases} \mathbb{E}^0(Q)\mathbb{M}(Q)\mu^{-1}(1 + o(1)) & (\mathfrak{s} = +) \\ \infty & (\mathfrak{s} = -), \end{cases} \quad (1.19)$$

as $\mu \rightarrow 0$, where \mathbb{E}^0 denotes the energy without the potential, namely

$$\mathbb{E}^0(\varphi) := \int_{\mathbb{R}^3} \frac{|\nabla u|^2}{2} - \mathfrak{s} \frac{|u|^4}{4} dx. \quad (1.20)$$

In fact, in the defocusing case $\mathfrak{s} = -$, the soliton (1.12) is unique for each fixed $\mathbb{M}(\varphi) = \mu > 0$ modulo the gauge symmetry $e^{i\theta}$. In the focusing case $\mathfrak{s} = +$, the first excited states are generated by scaling of Q

$$\mathcal{S}_1 \ni \varphi = \omega^{1/2}(Q + o(1))(\omega^{1/2}x) \quad (\mathbb{M}(\varphi) \rightarrow 0) \quad (1.21)$$

We do not need the above characterizations of \mathcal{S}_1 , but the variational property with respect to the virial-type functional

$$\mathbb{K}_2(u) := \int_{\mathbb{R}^3} |\nabla u|^2 - \frac{rV_r|u|^2}{2} - \mathfrak{s} \frac{3|u|^4}{4} dx = \partial_{\alpha=1} \mathbb{E}(\alpha^{3/2}u(\alpha x)) \quad (1.22)$$

plays a crucial role as in the case $V = 0$. The following is the main result of this paper.

Theorem 1.1. *There exists $0 < \mu_* \ll 1$ such that for any $u(0) \in H_r^1(\mathbb{R}^3)$ satisfying $\mathbb{M}(u(0)) \leq \mu_*$ and $\mathbb{E}(u(0)) < \mathcal{E}_1(\mathbb{M}(u(0)))$, the corresponding solution of (1.7) either blows up in finite time both in $t > 0$ and in $t < 0$, or scatters as $t \rightarrow \pm\infty$ to the ground states \mathcal{S}_0 . More precisely, in the former case, there are $T_\pm \in (0, \infty)$ such that the unique solution $u \in C((-T_-, T_+); H_r^1)$ exists and*

$$\lim_{t \rightarrow \pm(T_\pm - 0)} \|\nabla u(t)\|_{L^2(\mathbb{R}^3)} = \infty = \limsup_{t \rightarrow \pm(T_\pm - 0)} \|u(t)\|_{L^\infty(\mathbb{R}^3)}. \quad (1.23)$$

In the latter case, there are a C^1 function $z : \mathbb{R} \rightarrow D_b \subset \mathbb{C}$ and $u_\pm \in H_r^1(\mathbb{R}^3)$ such that $|z(t)|$ converges as $t \rightarrow \pm\infty$ and

$$\lim_{t \rightarrow \pm\infty} \|u(t) - \Phi[z(t)] - e^{-it\Delta} u_\pm\|_{H^1(\mathbb{R}^3)} = 0. \quad (1.24)$$

Moreover, the blow up occurs if and only if

$$\mathfrak{s} = +, \quad \|\nabla u(0)\|_{L^2(\mathbb{R}^3)} > 1, \quad \text{and} \quad \mathbb{K}_2(u(0)) < 0, \quad (1.25)$$

which persists in t as long as the solution u exists.

The above theorem contains the asymptotic stability of the ground state \mathcal{S}_0 for small H^1 radial solutions. This part is contained in the asymptotic stability for small solutions in [9] by Gustafson, Tsai and the author, which does not need the radial symmetry restriction.

If the potential $V = 0$, then there is no small soliton such as \mathcal{S}_0 , but the ground state Q as in (1.4) exists and unstable. In that case, the above result regarding $\mathcal{S}_0 = \{0\}$ and $\mathcal{S}_1 = \{\alpha Q(\alpha x)\}_{\alpha > 0}$ was obtained by Holmer and Roudenko [10], extended to the non-radial case by Duyckaerts, Holmer and Roudenko [4], to the threshold energy by Duyckaerts and Roudenko [7], and slightly above the threshold (in the radial case) by Schlag and the author [16]. In these works there is no small-mass constraint as above, but it is not an essential difference, because the scale invariance in the case $V = 0$ allows one freely to add or remove such a restriction.

1.3. Difficulties and ideas in the proof. The proof follows the strategy of Kenig and Merle [12], which consists of a stationary part based on the classical variational argument for the elliptic equation (1.12), and a dynamical (or scattering) part based on the variational argument in space-time: the profile decomposition by Bahouri and Gérard [1].

The problem caused by the potential in the stationary variational argument can be read immediately from the virial identity

$$\partial_t \langle i\dot{u} | x \cdot \nabla u \rangle = -2\mathbb{K}_2(u). \quad (1.26)$$

In the absence of V , the functional \mathbb{K}_2 can not vanish under the energy constraint except at 0, and so sign-definite along each trajectory. This leads to monotonicity in the virial identity, which has been the crucial starting point for $V = 0$, including the case slightly above the ground state [15], where possible change of sign \mathbb{K}_2 was controlled by using the linearized operator around Q .

In the presence of V , the functional \mathbb{K}_2 changes the sign around the ground solitons \mathcal{S}_0 . Note that this problem does not arise in the elliptic equation (1.12) using the

Nehari functional

$$\partial_{\alpha=1}(\mathbb{E} + \omega\mathbb{M})(\alpha u) = \int_{\mathbb{R}^3} |\nabla u|^2 + (V + \omega)|u|^2 - |u|^4 dx, \quad (1.27)$$

because the excited states \mathcal{S}_1 can be distinguished from the ground states \mathcal{S}_0 by the time frequency ω . Indeed, $\omega \rightarrow -e_0$ on \mathcal{S}_0 while $\omega \rightarrow \infty$ on \mathcal{S}_1 as $\mathbb{M} \rightarrow 0$. In contrast, the virial functional \mathbb{K}_2 is independent of ω , since it is derived by the L^2 -preserving dilation.

The above problem in the virial identity is however easily solved using the fact that the disturbance of sign \mathbb{K}_2 occurs only in a small neighborhood of $H^1(\mathbb{R}^3)$, where we have the asymptotic stability of \mathcal{S}_0 from [9]. In fact, the region $\mathbb{K}_2 \lesssim \mathbb{M} \ll 1$ splits into two sets far from each other in $H^1(\mathbb{R}^3)$: one around 0 satisfying

$$\|\nabla\varphi\|_{L^2(\mathbb{R}^3)}^2 + \|\varphi\|_{L^4(\mathbb{R}^3)}^2 \lesssim \mathbb{M}(\varphi) \ll 1, \quad (1.28)$$

and the other with large energy satisfying

$$\min(\|\nabla\varphi\|_{L^2(\mathbb{R}^3)}^2, \|\varphi\|_{L^4(\mathbb{R}^3)}^4) \gtrsim \mathbb{M}(\varphi)^{-1} \gg 1. \quad (1.29)$$

See Lemma 2.3 for a more general statement with a proof. In (1.25), the condition $\|\nabla u\|_{L^2} > 1$ is imposed only to distinguish the above two cases, so there are many alternative conditions, such as $\|u\|_{L^4} > 1$.

The problems in the space-time variational argument, caused by the potential, or more precisely by the stable solitons \mathcal{S}_0 , appear more fundamental. First, we should obviously remove the stable soliton part from the solution to apply the profile decomposition, as it aims at global dispersion or space-time integrability of the solutions. Second, the linear terms of the dispersive part, namely the interaction with the small soliton, can not be treated as part of the nonlinear perturbation, since it would require smallness in L_t^2 of the remainder of the profile decomposition, which is impossible as long as we take the initial data from the L_x^2 Sobolev space.

Therefore, we have to consider the linearized equation around the small soliton as the reference equation in the profile decomposition for the dispersive component. Since the modulation in time, namely $z(t)$ in (1.24), depends on the solution, it means that we have to consider a sequence of equations corresponding to the sequence of solutions to which we apply the concentration compactness.

Another problem is that we have very poor control on the global or asymptotic behavior of $z(t)$. For example, the convergence of $|z(t)|$ as $t \rightarrow \infty$ becomes arbitrarily slow by choosing small H^1 data, see [9, Theorem 1.9]. This causes difficulties at least in the following two places.

First, the nonlinear profile decomposition is a method to approximate solutions globally in time, but we can not do it for the soliton part $z(t)$. Therefore we have to distinguish time into two regimes: around and away from the profiles, approximating z only in the former, while relying on the smallness of the dispersive component in the latter.

Second, the nonlinear profiles moving to $t \rightarrow \pm\infty$ were defined in Kenig-Merle [12] by the wave operator, i.e., solving the final state problem with the linear profile as the scattering state. The existence of solution to the final state problem in the current setting, namely around the ground states \mathcal{S}_0 , was proved in [9], but we do not even know the uniqueness, while we would need some continuity estimate. Hence

we have to define the nonlinear profiles in another way, that is the weak limit along a time sequence, proving afterward that the linear profile is the scattering state. The drawback of this definition is that we can not construct global approximation at one stroke as in Kenig-Merle, but have to proceed step by step over each profile. The approach in this paper can be roughly regarded as a hybrid between Bahouri-Gérard [1] and Kenig-Merle [12]. The former used the scattering to describe the limit of sequence of solutions, while the latter used the limit of sequence to obtain the scattering. We need to proceed from both the sides.

Yet another complication in the estimates is due to the quadratic nonlinearity in the equation after linearization, to which we can not directly apply the Strichartz estimate to obtain Lipschitz estimate in the energy space for global perturbation, together with the smallness of the remainder in the profile decomposition. To solve this problem, we follow the idea in [13], using non-admissible Strichartz norms and measuring the initial data by the Strichartz norm.

1.4. Notation. L_x^p , $B_{p,q}^s$, and $H^s = B_{2,2}^s$ denote respectively the standard Lebesgue, inhomogeneous Besov, and inhomogeneous Sobolev spaces on \mathbb{R}^3 . The L_x^p norm is often denoted by $\|\cdot\|_p$. For any function space X on \mathbb{R}^3 , the subspace of radial functions is denoted by X_r , and the L^p space in $t \in \mathbb{R}$ with values in X is denoted by $L_t^p X$. For any function space Z on \mathbb{R}^{1+3} and $I \subset \mathbb{R}$, $Z(I)$ denotes the restriction onto $I \times \mathbb{R}^3$. The L^2 inner products on \mathbb{R}^3 are denoted by

$$(f|g) := \int_{\mathbb{R}^3} f(x)\overline{g(x)}dx, \quad \langle f|g \rangle := \operatorname{Re}(f|g). \quad (1.30)$$

1.5. Assumptions on V . Let $L_0^\infty(\mathbb{R}^3) = \{\varphi \in L^\infty(\mathbb{R}^3) \mid \lim_{R \rightarrow \infty} \|\varphi\|_{L^\infty(|x|>R)} = 0\}$.

The precise assumption on V is as follows.

- (i) $V : \mathbb{R}^3 \rightarrow \mathbb{R}$ is radially symmetric.
- (ii) $V, x\nabla V, x^2\nabla^2 V \in (L^2 + L_0^\infty)(\mathbb{R}^3)$ and $V/|x| \in L^1(\mathbb{R}^3)$.
- (iii) $-\Delta + V$ on $L_r^2(\mathbb{R}^3)$ has a unique and negative eigenvalue.
- (iv) The wave operator $W = \lim_{t \rightarrow \infty} e^{itH} e^{it\Delta}$ and its adjoint W^* are bounded on the Sobolev space $W^{k,p}(\mathbb{R}^3)$ for some $\mathfrak{p} > 6$ and $k = 0, 1$.

The above assumption (ii) implies that

$$\lim_{|x| \rightarrow \infty} |V(x)| + |x\nabla V(x)| = 0 \quad (1.31)$$

by the radial Sobolev inequalities, cf. Appendix A. By Beceanu [2, Corollary 1.5], the assumption (iv) is fulfilled if 0 is neither an eigenvalue nor resonance of H , and $V \in (L^p \cap \mathcal{F}B_{2,1}^{1/2})(\mathbb{R}^3)$.

For example, if $\psi \in \mathcal{S}(\mathbb{R}^3)$ is a radial positive function, the above assumptions (i)-(iv) are satisfied by $V = -a\psi$ for $a \in (1/a_1, 1/a_2)$, where $a_1 > a_2 > 0$ are the largest and the second largest eigenvalues of the compact self-adjoint operator $(-\Delta)^{-1/2}\psi(-\Delta)^{-1/2}$ on $L_r^2(\mathbb{R}^3)$.

ACKNOWLEDGEMENTS

This work was originally started from intensive discussions with Stephen Gustafson and Tai-Peng Tsai. The author would also like to thank them for useful comments on the manuscript.

2. STANDING WAVES

This section collects some properties of the solutions of (1.12), namely solitons. It is easy to see $\omega > 0$ for $\varphi \in H_r^1(\mathbb{R}^3)$, using the asymptotic behavior of the ODE as $r = |x| \rightarrow \infty$. We will see that in the defocusing case $\mathfrak{s} = -$, there is a unique soliton φ for each $\omega \in (0, -e_0)$ and nothing else. In the focusing case $\mathfrak{s} = +$, there is a soliton for each $\omega \in (-e_0, \infty)$, among which we can specify the ground state and the first excited state for each fixed small mass under the radial constraint.

2.1. Energy functionals. For any $\varphi \in H^1(\mathbb{R}^3)$, and $V : \mathbb{R}^3 \rightarrow \mathbb{R}$, we define the following functionals on $H^1(\mathbb{R}^3)$.

$$\begin{aligned} [V](\varphi) &:= \langle V\varphi|\varphi \rangle / 2, \quad \mathbb{M}(\varphi) := [1](\varphi) = \|\varphi\|_2^2 / 2, \quad \mathbb{G}(\varphi) := \mathfrak{s}\|\varphi\|_4^4 / 4, \\ \mathbb{H}(\varphi) &:= \langle H\varphi|\varphi \rangle / 2 = \|\nabla\varphi\|_2^2 / 2 + [V](\varphi), \quad \mathbb{E} := \mathbb{H} - \mathbb{G}. \end{aligned} \quad (2.1)$$

The energy \mathbb{E} and the mass \mathbb{M} are conserved in time for (1.7). The corresponding quantities without the potential V are denoted by $\mathbb{H}^0, \mathbb{E}^0$, etc.

$$\mathbb{H}^0(\varphi) := \|\nabla\varphi\|_2^2 / 2, \quad \mathbb{E}^0(\varphi) := \|\nabla\varphi\|_2^2 / 2 - \mathfrak{s}\|\varphi\|_4^4 / 4. \quad (2.2)$$

For the variational property in the focusing case, we need the dilation operator

$$\mathcal{S}_p^t \varphi(x) := e^{3t/p} \varphi(e^t x), \quad \mathcal{S}_p' \varphi(x) = (x \cdot \nabla + 3/p) \varphi(x), \quad (2.3)$$

which preserves the L_x^p norm. The same notation is used for the functional

$$(\mathcal{S}_p' F)(\varphi) := \partial_{t=0} F(\mathcal{S}_p^t \varphi). \quad (2.4)$$

Then we have

$$\begin{aligned} \mathcal{S}_p' \mathbb{M} &= (6/p - 3)\mathbb{M}, \quad \mathcal{S}_p' \mathbb{H}^0 = (6/p - 1)\mathbb{H}^0, \\ \mathcal{S}_p' \mathbb{G} &= (12/p - 3)\mathbb{G}, \quad \mathcal{S}_p' [V] = -[\mathcal{S}_{p/(p-2)}' V]. \end{aligned} \quad (2.5)$$

The L^2 -scaling derivative plays a crucial role via the virial identity

$$\mathbb{K}_2 := \mathcal{S}_2' \mathbb{E} = 2\mathbb{H}^0 - 3\mathbb{G} - [\mathcal{S}_\infty' V]. \quad (2.6)$$

The following functional is used for convexity of \mathbb{E} in \mathcal{S}_p^t :

$$\mathbb{I} := \mathbb{E} - \mathbb{K}_2 / 2 = \mathbb{G} / 2 + [\mathcal{S}_{3/2}' V] / 2. \quad (2.7)$$

If there is a family of solitons $\omega \mapsto \varphi_\omega \in H^1$ differentiable in $\omega \in \mathbb{R}$, then we have

$$\partial_\omega \mathbb{E}(\varphi_\omega) = \langle \mathbb{E}'(\varphi_\omega) | \varphi_\omega' \rangle = \langle -\omega \varphi_\omega | \varphi_\omega' \rangle = -\omega \partial_\omega \mathbb{M}(\varphi_\omega). \quad (2.8)$$

For the potential part, we will frequently use the following bound

Lemma 2.1. *Let $V \in (L^2 + L_0^\infty)(\mathbb{R}^3)$. Then for any $\varepsilon > 0$, there is $C > 0$ such that*

$$H^1(\mathbb{R}^3) \ni \forall \varphi, \quad |[V](\varphi)| \leq \min(\varepsilon \|\varphi\|_4^2 + C \|\varphi\|_2^2, \varepsilon \|\varphi\|_2^2 + C \|\varphi\|_4^2). \quad (2.9)$$

Proof. Let $V = V_2 + V_\infty$ where $V_2 \in L^2$ and $V_\infty \in L_0^\infty$. For any $h > 0$, we have

$$|[V_2](\varphi)| \leq \|V_2\|_{L^2(V>h)} \|\varphi\|_4^2 + h \|\varphi\|_2^2, \quad (2.10)$$

where $\|V_2\|_{L^2(V>h)} \rightarrow 0$ as $h \rightarrow \infty$. We have $|[V_\infty](\varphi)| \lesssim \|\varphi\|_2^2$ and for any $R > 0$,

$$|[V_\infty](\varphi)| \lesssim \|V_\infty\|_{L^\infty(|x|<R)} \|\varphi\|_4^2 R^{3/2} + \|V_\infty\|_{L^\infty(|x|>R)} \|\varphi\|_2^2, \quad (2.11)$$

where $\|V_\infty\|_{L^\infty(|x|>R)} \rightarrow 0$ as $R \rightarrow \infty$. Adding these yields the conclusion. \square

2.2. Small solitons. For small mass, the ground state is the bifurcation from zero, generated by the ground state ϕ_0 of H . The following precise statement can be extracted from [9, Lemma 2.1]

Lemma 2.2. *There exists $0 < z_b \ll 1$ and a C^1 map*

$$(\Phi, \Omega) : D_b := \{z \in \mathbb{C} \mid |z|^2 < 2z_b\} \rightarrow H_r^1(\mathbb{R}^3) \times \mathbb{R}, \quad (2.12)$$

such that $(\varphi, \omega) = (\Phi[z], \Omega[z])$ is a soliton for each $z \in D_b$, with a decomposition

$$\Phi[z] = z\phi_0 + \gamma[z], \quad \Omega[z] = -e_0 + o(z), \quad \gamma[z] \perp \phi_0, \quad \|\gamma[z]\|_{H^1} = o(|z|^2), \quad (2.13)$$

satisfying the gauge covariance $(\Phi[e^{i\theta}z], \Omega[e^{i\theta}z]) = (e^{i\theta}\Phi[z], \Omega[z])$, and $\mathbb{M}(\Phi[z]) = |z|^2/2 + o(|z|^4)$ is an increasing function of $|z|$. Moreover, there is an open set in $H_r^1(\mathbb{R}^3)$ which contains $\Phi[D_b]$ but no other soliton.

Let $\mu_b > 0$ be the maximal mass among those solitons:

$$\mu_b := \sup \mathbb{M}(\Phi[D_b]). \quad (2.14)$$

Then the monotonicity implies that

$$[0, z_b) \ni z \mapsto \mathbb{M}(\Phi[z]) \in [0, \mu_b) \quad (2.15)$$

is an increasing bijection. Let $z_0 : [0, \mu_b) \rightarrow [0, z_b)$ be the inverse function, so that

$$\mathbb{M}(\Phi[z_0(\mu)]) = \mu. \quad (2.16)$$

The following lemma shows that the above solitons are the ground states for small mass. It will be crucial also for identifying the first excited state.

Lemma 2.3 (Small mass dichotomy). *For any $\varphi \in H^1(\mathbb{R}^3)$ satisfying $\mathbb{K}_2(\varphi) \ll \mathbb{M}(\varphi)^{-1}$ and $\mathbb{M}(\varphi) \ll 1$, we have one of the following (i)-(iii)*

- (i) $\mathbb{H}^0(\varphi) \lesssim \mathbb{M}(\varphi)$,
- (ii) $\mathbb{M}(\varphi) \lesssim \mathbb{H}^0(\varphi) \sim \mathbb{E}(\varphi) \sim \mathbb{K}_2(\varphi)$,
- (iii) $\mathfrak{s} = +$ and $\mathbb{H}^0(\varphi) \sim \mathbb{G}(\varphi) \gtrsim \mathbb{M}(\varphi)^{-1}$,

Moreover, in the case (iii), we have

$$|[\mathbb{V}](\varphi)| + |[\mathcal{S}'_p \mathbb{V}](\varphi)| + |[\mathcal{S}'_p \mathcal{S}'_q \mathbb{V}](\varphi)| \ll \mathbb{G}(\varphi) \quad (2.17)$$

for any $p, q > 0$. In particular, \mathbb{G} in (iii) can be replaced with $2\mathbb{I}$.

Note that the first two regions overlap each other, but the last one is separated. Thus the above lemma gives a dichotomy into (i)-(ii) and (iii). The case (ii) can be removed if $\mathbb{K}_2(\varphi) \lesssim \mathbb{M}(\varphi)$, which is mostly satisfied when the above lemma is used.

Proof. Let $\mu := \mathbb{M}(\varphi)$ and $h := \mathbb{H}^0(\varphi)$. Using Gagliardo-Nirenberg, we obtain

$$\begin{aligned} |2h - \mathbb{K}_2(\varphi)| &= | - [\mathcal{S}'_\infty \mathbb{V}](\varphi) + 3\mathbb{G}(\varphi) | \\ &\leq \|\mathcal{S}'_\infty \mathbb{V}\|_{L^2+L^\infty} (\|\varphi\|_2^2 + \|\varphi\|_4^2) + \|\varphi\|_4^4 \\ &\lesssim \mu + h^{3/4} \mu^{1/4} + h^{3/2} \mu^{1/2}. \end{aligned} \quad (2.18)$$

Hence either $h \leq \mathbb{K}_2(\varphi) \ll \mu^{-1}$, $h \lesssim \mu$, $h \lesssim h^{3/4} \mu^{1/4}$ or $h \lesssim h^{3/2} \mu^{1/2}$. The third case is the same as the second, up to the constants, leading to (i). If $\mu \ll h \ll \mu^{-1}$, then (2.18) $\ll h$, and the same estimate works for $|\mathbb{H}^0(\varphi) - \mathbb{E}(\varphi)|$, leading to (ii).

In the last case, we obtain $h \gtrsim \mu^{-1} \gg 1$ hence \mathbb{G} is dominating, thus $\mathfrak{s} = +$ and $\mathbb{I}(\varphi) \sim \mathbb{G}(\varphi) \sim h \gtrsim \mu^{-1}$, leading to (iii). \square

The above lemma enables us to identify the solitons in Lemma 2.2 with \mathcal{S}_0 :

Proposition 2.4. *There exists $0 < \mu_d \leq \mu_b$ and $0 < z_d \leq z_b$ such that*

$$\begin{aligned} \{\varphi \in \mathcal{S}_0 \mid \mathbb{M}(\varphi) < \mu_d\} &= \{\Phi[z] \mid |z| < z_d\}, \\ 0 < \mu < \mu_d &\implies \mathcal{E}_0(\mu) = \mathbb{E}(\Phi[z_0(\mu)]) \sim e_0\mu < 0. \end{aligned} \quad (2.19)$$

Proof. Since $\mathbb{K}_2 = 0$ on \mathcal{S} , we can apply the above lemma to any $\varphi \in \mathcal{S}$ with $\mathbb{M}(\varphi) < \mu_d \ll 1$, leading to either (i) or (iii). Taking $\mu_d > 0$ small ensures that the region (i) is in the uniqueness region of $\Phi[D_b]$ in Lemma 2.2, as well as that the region (iii) is far away. Then every $\varphi \in \mathcal{S}$ with $\mathbb{M}(\varphi) < \mu_d$ is either in $\Phi[D_b]$ or in the region (iii). In the latter case, φ is an excited state, as Φ gives a soliton with the same mass and negative energy. Thus we obtain the first identity in (2.19). The second one is its obvious consequence. The behavior of \mathcal{E}_0 follows from (2.8) together with the differentiability of Φ from Lemma 2.2. \square

2.3. Focusing case $\mathfrak{s} = +$. Next we investigate the first excited state of small mass in the focusing case. The small mass dichotomy Lemma 2.3 allows us to ignore the potential effect, leading to the same variational characterization as $V = 0$ in the higher energy region.

Proposition 2.5. *Let $\mathfrak{s} = +$. There exists $0 < \mu_e \leq \mu_d$ such that for $0 < \mu < \mu_e$*

$$\begin{aligned} \mathcal{E}_1(\mu) &= \inf\{\mathbb{E}(\varphi) \mid \varphi \in H_r^1, \mathbb{M}(\varphi) = \mu, \mathbb{K}_2(\varphi) = 0, \mathbb{G}(\varphi) \geq 1\} \\ &= \inf\{\mathbb{I}(\varphi) \mid \varphi \in H_r^1, \mathbb{M}(\varphi) \leq \mu, \mathbb{K}_2(\varphi) \leq 0, \mathbb{G}(\varphi) \geq 1\} \\ &= \mu^{-1}\mathbb{M}(Q)\mathbb{E}^0(Q)(1 + o(1)) \gg 1 \quad (\mu \rightarrow 0), \end{aligned} \quad (2.20)$$

where \inf is attained by some $\varphi \in \mathcal{S}_1$, and $\mathcal{E}_1(\mu)$ is decreasing in μ . Moreover, there is a continuous function $\kappa : (0, \mu_e) \times (0, \infty) \rightarrow (0, 1/2)$ such that for any $\delta > 0$ and any $\varphi \in H_r^1$ satisfying $\mathbb{M}(\varphi) < \mu_e$, $\mathbb{E}(\varphi) \leq \mathcal{E}_1(\mu) - \delta$ and $\|\nabla\varphi\|_2 \geq 1$, we have

$$|\mathbb{K}_2(\varphi)| \geq \kappa(\mathbb{M}(\varphi), \delta). \quad (2.21)$$

The above minimization is well known in the case $V = 0$ without the restriction on \mathbb{G} . Some restriction to higher energy is needed in the case $V \neq 0$, since those \inf in (2.20) would become $\mathcal{E}_0(\mu)$ without $\mathbb{G} \geq 1$.

Proof. First, the \geq part of (2.20) is obvious from $\mathbb{K}_2 = 0$ on \mathcal{S} , the dichotomy Lemma 2.3, and $\mathbb{I} = \mathbb{E} - \mathbb{K}_2/2$. The second infimum is obviously decreasing in μ .

To show the \leq part of the second equality, let $\varphi \in H_r^1$, $\mathbb{M}(\varphi) < \mu$, $\mathbb{K}_2(\varphi) \leq 0$ and $\mathbb{G}(\varphi) \geq 1$. Consider the one-parameter scaling $v(t) := \mathcal{S}'_{3.5} \varphi$ for $t \leq 0$. The dichotomy implies $\mathbb{G}(\varphi) \gtrsim \mu^{-1} \gg 1$. Since $\mathcal{S}'_{3.5}\mathbb{M} = -9\mathbb{M}/7 < 0$, there exists $T < 0$ such that $\mathbb{M}(v(T)) = \mu$. Moreover, since

$$\mathcal{S}'_{3.5}\mathbb{I} = \frac{3}{14}\mathbb{G} - \frac{1}{2}[\mathcal{S}'_{7/3}\mathcal{S}'_{3/2}V], \quad \mathcal{S}'_{3.5}\mathbb{K}_2 = \frac{5}{7}\mathbb{K}_2 + \frac{6}{7}\mathbb{G} + [\mathcal{S}'_{3/2}V], \quad (2.22)$$

we have, using Lemma 2.1,

$$\mathcal{S}'_{3.5}\mathbb{I}(v) \gtrsim \mu^{-1}, \quad \mathcal{S}'_{3.5}\mathbb{K}_2(v) > 5(\mathbb{K}_2(v) + 1)/7 \quad (2.23)$$

as long as $\mathbb{K}_2(v) \leq \mathbb{M}(v) \leq \mu$. The second inequality of (2.23) implies that if $\mathbb{K}_2(v) \geq -1$, $\mathbb{K}_2(v)$ is decreasing as t decreases. Therefore $\mathbb{K}_2(v) < 0$ and (2.23) are preserved for $0 > t > T$, hence the infimum is reduced to the case $\mathbb{M}(\varphi) = \mu$.

Next consider the one-parameter scaling $v(t) := \mathcal{S}_2^t \varphi$ for $t \leq 0$. Since

$$\mathcal{S}_2' \mathbb{I} = 3\mathbb{G}/2 - [\mathcal{S}'_\infty \mathcal{S}'_{3/2} V]/2, \quad \mathcal{S}'_2 \mathbb{K}_2 = 2\mathbb{K}_2 - 2\mathcal{S}'_2 \mathbb{I}, \quad (2.24)$$

a similar argument as above implies that $\mathbb{I}(v)$ is decreasing and $\mathbb{K}_2(v)$ is increasing as t decreases, as long as $\mathbb{K}_2(v) < 0$, while $\mathbb{G}(v) \geq 1$ is preserved by the dichotomy. Thus the infimum is further reduced to the case $\mathbb{K}_2(\varphi) = 0$, which means the second equality in (2.20).

To prove the existence of minimizer as well as the lower bound (2.21) on $|\mathbb{K}_2|$, take any sequence $\varphi_n \in H_r^1$ satisfying $\mathbb{M}(\varphi_n) \rightarrow \mu \in (0, \mu_e)$, $\mathbb{K}_2(\varphi_n) \rightarrow 0$, $\mathbb{G}(\varphi_n) + \|\nabla \varphi_n\|_2 \geq 1$ and $\mathbb{E}(\varphi_n) \rightarrow E_\infty \leq E_*(\mu)$, where E_* is the infimum in (2.20). Using Lemma 2.1 with Gagliardo-Nirenberg, we have

$$\mathbb{H}^0 = 3\mathbb{E} - \mathbb{K}_2 - [\mathcal{S}'_1 V] \leq 3\mathbb{E} - \mathbb{K}_2 + C\mathbb{M} + \mathbb{H}^0/2. \quad (2.25)$$

Hence φ_n is bounded in H_r^1 , and so, we may assume, passing to a subsequence, that $\varphi_n \rightarrow \exists \varphi \in H_r^1$ weakly. Since $\mathbb{K}_2 = 2\mathbb{H}^0 - 3\mathbb{G} - [\mathcal{S}'_\infty V]$, disposing of the potential part as above, we deduce that $\mathbb{G}(\varphi_n) \gtrsim 1$ and $\|\nabla \varphi_n\|_2 \gtrsim 1$ are equivalent for large n , and then the dichotomy implies $\mathbb{H}^0(\varphi_n) \sim \mathbb{G}(\varphi_n) \gtrsim \mu^{-1}$.

Since \mathbb{G} and the potential functionals are weakly continuous on H_r^1 , we have

$$\mathbb{I}(\varphi) = E_\infty, \quad \mathbb{G}(\varphi) \geq 1, \quad \mathbb{M}(\varphi) \leq \mu, \quad \mathbb{K}_2(\varphi) \leq 0, \quad (2.26)$$

hence $E_\infty = E_*$ and φ is a minimizer of (2.20). Moreover, the above argument implies that $\mathbb{M}(\varphi) = \mu$ and $\mathbb{K}_2(\varphi) = 0$. Since the dichotomy implies $\mathbb{G}(\varphi) \gtrsim \mu^{-1} \gg 1$, we have Lagrange multipliers $\omega, \alpha \in \mathbb{R}$ such that

$$\mathbb{E}'(\varphi) + \omega \mathbb{M}'(\varphi) = \alpha \mathbb{K}_2(\varphi). \quad (2.27)$$

Differentiation along the curve $\mathcal{S}_2^t \varphi$ at $t = 0$ yields

$$0 = \mathbb{K}_2(\varphi) = \mathcal{S}'_2 \mathbb{E}(\varphi) = \alpha \mathcal{S}'_2 \mathbb{K}_2(\varphi) = -2\alpha \mathcal{S}'_2 \mathbb{I}(\varphi), \quad (2.28)$$

where $\mathcal{S}'_2 \mathbb{I}(\varphi) \neq 0$ by (2.24) with the dichotomy, hence $\alpha = 0$. This means that $\varphi \in \mathcal{S}$ and so $\mathcal{E}_1(\mu) = E_*(\mu)$, as well as the lower bound (2.21) on $|\mathbb{K}_2|$.

Finally, we prove the asymptotic formula. In the second infimum in (2.20), put $\psi(x) := \mu \varphi(\mu x)$ and $V_\mu(x) := \mu^2 V(\mu x)$. Then

$$\mu \mathcal{E}_1(\mu) = \min_{\psi \in A_\mu} \mathbb{G}(\psi)/2 + [(\mathcal{S}'_{3/2} V)_\mu](\psi), \quad (2.29)$$

$$A_\mu := \{\psi \in H_r^1 \mid \mathbb{M}(\psi) \leq 1, \mathbb{K}_2^0(\psi) \leq [(\mathcal{S}'_\infty V)_\mu](\psi), \mathbb{G}(\psi) \geq \mu\}.$$

Since $\|V_\mu\|_{L^2+L^\infty} \leq \mu^{1/2} \|V\|_{L^2+L^\infty}$, we have for $p > 0$

$$|[(\mathcal{S}'_p V)_\mu](\psi)| \lesssim \mu^{1/2} (\|\psi\|_4^2 + \|\psi\|_2^2). \quad (2.30)$$

Hence if $\psi \in H_r^1$ satisfies $\mathbb{M}(\psi) \leq 1$ and $\mathbb{K}_2^0(\psi) \leq -1$, then $\psi \in A_\mu$ for $0 < \mu \ll 1$. Therefore as $\mu \rightarrow 0$, the minimizer ψ is bounded in L_x^4 . Since $\mathbb{G}(\varphi) \gtrsim \mu^{-1}$, we obtain $\mathbb{G}(\psi) \sim 1$ and $|[(\mathcal{S}'_p V)_\mu](\psi)| \leq O(\mu^{1/2})$. On the other hand, for any $\psi \in H_x^1$

satisfying $\mathbb{M}(\psi) \leq 1 \sim \mathbb{G}(\psi)$ and $\mathbb{K}_2^0(\psi) = 0$, we deduce from (2.24) that for $0 < \mu \ll 1$ there exists $t = O(\mu^{1/2})$ such that $\mathcal{S}_2^t \psi \in A_\mu$. Therefore

$$\lim_{\mu \rightarrow 0} \mu \mathcal{E}_1(\mu) = \inf\{\mathbb{G}(\psi)/2 \mid \psi \in H_r^1, \mathbb{M}(\psi) \leq 1, \mathbb{K}_2^0(\psi) \leq 0\}, \quad (2.31)$$

and by the well-known characterization of Q , the right hand equals $\mathbb{M}(Q)\mathbb{E}^0(Q)$. \square

2.4. Defocusing case $\mathfrak{s} = -$. In the defocusing case, the variational structure is much simpler, and so we can determine the entire set of solitons \mathcal{S} without the mass constraint. In this subsection, we prove the following

Proposition 2.6. *Let $\mathfrak{s} = -$. Under the assumptions on V in Section 1.5, the equation (1.12) has a unique positive solution φ_ω for each $\omega \in (0, -e_0)$, and*

$$\mathcal{S} = \{e^{i\theta} \varphi_\omega \mid 0 < \omega < -e_0, \theta \in \mathbb{R}\}. \quad (2.32)$$

The function $(0, -e_0) \ni \omega \mapsto \mathbb{M}(\varphi_\omega) \in (0, \infty)$ is C^1 , decreasing and bijective. Let $\omega_0(\mu)$ be its inverse function. Then for all $\mu > 0$,

$$e_0 \mu < \mathcal{E}_0(\mu) = \mathbb{E}(\varphi_{\omega_0(\mu)}) < 0 < \infty = \mathcal{E}_1(\mu), \quad \omega_0'(\mu) < 0. \quad (2.33)$$

Since $H \geq e_0$, multiplying (1.12) with φ

$$0 = \langle (H + \omega)\varphi | \varphi \rangle + \|\varphi\|_4^4 = \|\nabla \varphi\|_2^2 + \omega \|\varphi\|_2^2 + \langle V\varphi | \varphi \rangle + \|\varphi\|_4^4 \quad (2.34)$$

implies that $\omega < -e_0$ is necessary for existence of a non-trivial solution.

If $\omega \leq 0$, then putting $\psi := r\varphi$ we have from (1.12)

$$\psi_{rr} + \omega\psi = (V + |\varphi|^2)\psi, \quad \liminf_{r \rightarrow \infty} \{|\psi(r)| + |\psi_r(r)|\} = 0. \quad (2.35)$$

Rewriting the above into an integral equation from $r = \infty$, we obtain

$$|\psi(s)| \leq \int_s^\infty (r-s) |(V + |\varphi|^2)\psi| dr \leq \|\psi\|_{L^\infty(r>s)} \|r(V + |\varphi|^2)\|_{L^1(r>s)}, \quad (2.36)$$

where the last norm is vanishing as $s \rightarrow \infty$ by the assumption $V/|x| \in L^1(\mathbb{R}^3)$ and $\varphi \in L^2(\mathbb{R}^3)$. Hence taking $s \rightarrow \infty$ and then solving the ODE, we deduce that $\varphi = 0$.

Therefore $0 < \omega < -e_0$ for every non-trivial $\varphi \in \mathcal{S}$. Moreover, using Lemma 2.1 with $\varepsilon = \omega/2$, we deduce that $\varphi \in \mathcal{S}$ is uniformly bounded in $H^1(\mathbb{R}^3)$ on any interval of ω away from 0, while $\varphi_\omega \rightarrow 0$ in $H^1(\mathbb{R}^3)$ as $\omega \rightarrow -e_0 - 0$.

For each $\omega \in (0, -e_0)$, we have a solution φ_ω of (1.12) which is a global minimizer

$$(\mathbb{E} + \omega\mathbb{M})(\varphi_\omega) = \inf\{(\mathbb{E} + \omega\mathbb{M})(\varphi) \mid \varphi \in H^1\} < 0. \quad (2.37)$$

The proof is easy and omitted. The positivity is also standard. The uniqueness of the solution φ_ω for each ω , modulo the phase $e^{i\theta}$, follows from a general argument:

Lemma 2.7. *Let H be a self-adjoint operator on L^2 with non-degenerate eigenvalue e_0 , and assume the rest of the spectrum of H is contained in $[e_1, \infty)$ for some $e_1 > e_0$. Let $f : [0, \infty) \rightarrow [0, \infty)$ be a strictly monotone function such that $f(a)a$ is non-decreasing. Then the nonlinear eigenvalue problem*

$$(H + f(|\varphi|))\varphi = \omega\varphi$$

can have at most one non-trivial solution (up to the phase symmetry) for each $\omega < e_1$. The same conclusion holds for $\omega = e_1$ if $f(a)a$ is strictly increasing.

The above lemma may be known, but a proof is given below for the sake of completeness.

Proof. Let $f(z) := f(|z|)$ for $z \in \mathbb{C}$. Let φ and ψ be two non-zero solutions, and let ϕ_0 be an eigenfunction of H for e_0 . We must have $(\phi_0|\varphi) \neq 0$, or else

$$(e_1 - \omega)\|\varphi\|_2^2 \leq (\varphi|(H - \omega)\varphi) = -(|\varphi|^2|f(\varphi)), \quad (2.38)$$

which contradicts either $\omega < e_1$ or $\omega = e_1$ with strictly increasing $f(a)a$. Thus we can find $\beta \in \mathbb{C} \setminus \{0\}$ so that $(\phi_0|\beta\varphi + \psi) = 0$. Using the invariance of the equation for $\varphi \mapsto e^{i\theta}\varphi$, we may take $\beta > 0$ by appropriate complex rotation of φ . Then

$$\begin{aligned} (e_1 - \omega)\|\beta\varphi + \psi\|_2^2 &\leq \langle \beta\varphi + \psi | (H - \omega)(\beta\varphi + \psi) \rangle \\ &= -\beta^2 \langle f(\varphi) ||\varphi|^2 \rangle - \langle f(\psi) ||\psi|^2 \rangle + 2\beta \langle \varphi | (H - \omega)\psi \rangle \\ &\leq 2\beta \left[|\langle \varphi | (H - \omega)\psi \rangle| - \sqrt{\langle f(\varphi) ||\varphi|^2 \rangle \langle f(\psi) ||\psi|^2 \rangle} \right]. \end{aligned} \quad (2.39)$$

First consider the case where f is non-decreasing. Then using Schwarz,

$$|\langle \varphi | (H - \omega)\psi \rangle| = \begin{cases} |\langle \varphi | f(\psi)\psi \rangle| \leq \sqrt{\langle f(\psi) ||\varphi|^2 \rangle \langle f(\psi) ||\psi|^2 \rangle}, \\ |\langle f(\varphi)\varphi | \psi \rangle| \leq \sqrt{\langle f(\varphi) ||\varphi|^2 \rangle \langle f(\varphi) ||\psi|^2 \rangle}. \end{cases} \quad (2.40)$$

So we arrive at

$$\langle f(\psi) ||\varphi|^2 \rangle \geq \langle f(\varphi) ||\varphi|^2 \rangle, \quad \langle f(\varphi) ||\psi|^2 \rangle \geq \langle f(\psi) ||\psi|^2 \rangle, \quad (2.41)$$

and hence $\langle f(\varphi) - f(\psi) ||\varphi|^2 - |\psi|^2 \rangle \leq 0$. Since f is non-decreasing, this implies that $f(\varphi) = f(\psi)$ (a.e.).

Next consider the case where f is non-increasing. By Schwarz, we have

$$\begin{aligned} |\langle \varphi | f(\psi)\psi \rangle| &\leq \sqrt{\langle f(\varphi) ||\varphi|^2 \rangle \langle f(\psi)^2 |\psi|^2 |1/f(\varphi)\rangle}, \\ |\langle f(\varphi)\varphi | \psi \rangle| &\leq \sqrt{\langle f(\varphi)^2 |\varphi|^2 |1/f(\psi)\rangle \langle f(\psi) ||\psi|^2 \rangle}. \end{aligned} \quad (2.42)$$

Hence (2.39) implies that

$$\begin{aligned} \langle f(\psi)^2 |\psi|^2 |1/f(\varphi)\rangle &\geq \langle f(\psi)^2 |\psi|^2 |1/f(\psi)\rangle, \\ \langle f(\varphi)^2 |\varphi|^2 |1/f(\psi)\rangle &\geq \langle f(\varphi)^2 |\varphi|^2 |1/f(\varphi)\rangle, \end{aligned} \quad (2.43)$$

and so, $\langle f(\varphi)^2 |\varphi|^2 - f(\psi)^2 |\psi|^2 |1/f(\varphi) - 1/f(\psi)\rangle \leq 0$. Since $1/f(a)$ and $f(a)a$ are both non-decreasing, we have $f(\varphi) = f(\psi)$ or $f(\varphi)|\varphi| = f(\psi)|\psi|$ (a.e.). If $e_1 > \omega$, then $\psi = -\beta\varphi$, otherwise the above must be a strict inequality, contradicting the monotonicity of $f(\varphi)$ and $f(\varphi)|\varphi|$. If $e_1 = \omega$, then $f(z)|z|$ is strictly increasing, so we get $f(\varphi) = f(\psi)$, and then going back to (2.39),

$$\begin{aligned} 0 &= (e_1 - \omega)\|\beta\varphi + \psi\|_2^2 \leq \langle \beta\varphi + \psi | (H - \omega)(\beta\varphi + \psi) \rangle \\ &= -\langle f(\varphi) ||\beta\varphi + \psi|^2 \rangle \leq 0. \end{aligned} \quad (2.44)$$

The strict monotonicity of $f(a)a$ implies that at each x , $f(\varphi(x)) = 0$ implies $f(\psi(x)) = 0$ and $\varphi(x) = 0 = \psi(x)$. Hence $\psi = -\beta\varphi$ (a.e.).

Thus we obtain $\psi = -\beta\varphi$ anyway. Then the equation for φ and ψ implies that

$$(\omega - H)\varphi = f(\varphi)\varphi = -f(\psi)\psi/\beta = f(\beta\varphi)\varphi. \quad (2.45)$$

Since f is strictly monotone, this implies $\beta = 1$ or $\varphi = 0$ a.e. \square

Once we have the uniqueness of φ_ω for ω , it is easy to prove continuity and then differentiability in ω . Differentiating the equation

$$(H + \omega + 3\varphi_\omega^2)\varphi'_\omega + \varphi_\omega = 0, \quad \varphi'_\omega := \partial_\omega \varphi_\omega \quad (2.46)$$

and multiplying it with φ'_ω yield

$$\partial_\omega \mathbb{M}(\varphi_\omega) = \langle \varphi_\omega | \varphi'_\omega \rangle = -\langle (H + \omega + 3\varphi_\omega^2)\varphi'_\omega | \varphi'_\omega \rangle \leq -2\|\varphi_\omega \varphi'_\omega\|_2^2 < 0, \quad (2.47)$$

where we used $H + \omega + \varphi_\omega^2 \geq 0$, because $\varphi_\omega > 0$ is the ground state in the kernel of this Schrödinger operator. Hence $\mathbb{M}(\varphi_\omega)$ is decreasing in ω . Moreover, $\mathbb{M}(\varphi_\omega) \rightarrow \infty$ as $\omega \rightarrow +0$, since otherwise the weak limit yields $0 \neq \varphi_0 \in H_r^1$ satisfying $H\varphi_0 + \varphi_0^3 = 0$, which is impossible. Using (2.8) as well, we conclude the proof of Proposition 2.6.

3. BLOW-UP BELOW THE EXCITED ENERGY

We are now ready to prove the blow-up part of Theorem 1.1, using the above characterization of \mathcal{S}_1 together with the estimate on \mathbb{K}_2 , namely Proposition 2.5.

Let u be a solution of (1.7) with $\mathfrak{s} = +$, satisfying (1.25) as well as $\mathbb{M}(u) =: \mu < \mu_e$ and $\mathbb{E}(u) < \mathcal{E}_1(\mu)$, where μ_e is the small mass condition of Proposition 2.5. Fix $\delta > 0$ such that $\mathbb{E}(u) \leq \mathcal{E}_1(\mu) - \delta$. Suppose for contradiction that u exists on $0 < t < \infty$. Then Proposition 2.5 and Lemma 2.3 together with the continuity of $u(t)$ in H_x^1 imply that (1.25) is preserved for all $t > 0$, and also from (2.21)

$$\mathbb{K}_2(u(t)) \leq -\kappa(\mu, \delta). \quad (3.1)$$

We have the saturated virial identity from [17]

$$\begin{aligned} \partial_t \langle R f_R u | i u_r \rangle &= 2\mathbb{K}_2(u) - \int [2|u_r|^2 f_{0,R} + R^{-2}|u|^2 f_{1,R} + |u|^4 f_{2,R}] dx \\ &\quad - \int (1 - f_R R/r) |u|^2 x \cdot \nabla V dx, \end{aligned} \quad (3.2)$$

where $f_R(x) = f(x/R)$ with $R \gg 1$, and $f_{j,R}(x) = f_j(x/R)$ are derived from f by

$$f_0 = 1 - f_r \geq 0, \quad f_1 = \Delta(\partial_r/2 + 1/r)f, \quad f_2 = -3/2 + (\partial_r/2 + 1/r)f, \quad (3.3)$$

while $f(x) = f(|x|)$ is chosen to be smooth radial satisfying

$$f(r) = \begin{cases} r & (r \leq 1), \\ 3/2 & (r \geq 2). \end{cases} \quad (3.4)$$

The $|u|^4$ integral is bounded by the radial Sobolev inequality

$$\|u\|_{L^4(|x|>R)}^4 \lesssim R^{-2} \|u\|_{L^2(|x|>R)}^3 \|u_r\|_{L^2(|x|>R)}. \quad (3.5)$$

Then we obtain

$$\begin{aligned} & - \int [2|u_r|^2 f_{0,R} + R^{-2}|u|^2 f_{1,R} + |u|^4 f_{2,R} + (1 - f_R R/r) |u|^2 x \cdot \nabla V] dx \\ & \lesssim R^{-4} \|u\|_{L^2(|x|>R)}^6 + o(1) \|u\|_{L^2(|x|>R)}^2, \end{aligned} \quad (3.6)$$

as $R \rightarrow \infty$. See [17], [16, §4.1], for the detail. Note that the potential part was treated by (1.31), using $1 - f_R R/r = 0$ for $|x| < R$. Hence, for large R , we have

$$\partial_t \langle R f_R u | i u_r \rangle < -\kappa(\mu, \delta) < 0. \quad (3.7)$$

Since $\|u\|_{L_x^2}$ is conserved, it implies that $\|u_r\|_2 \rightarrow \infty$ as $t \rightarrow \infty$. Then as $t \rightarrow \infty$,

$$\mathbb{K}_2(u) = 3\mathbb{E}(u) - \mathbb{H}^0(u) - [\mathcal{S}'_1 V](u) \sim -\|\nabla u\|_2^2. \quad (3.8)$$

The rest of the proof is the same as in the case without the potential, see [17]. Thus we obtain the “if”-part for (1.25) of the blow-up in Theorem 1.1.

Next we show the “only if”-part of (1.25), namely the global existence when it is not satisfied. If $\mathfrak{s} = -$, then we have a priori H^1 bound by conservation of the energy and mass, disposing of the potential part by Lemma 2.1, which leads to the global well-posedness in $H^1(\mathbb{R}^3)$.

Hence we may restrict to the case $\mathfrak{s} = +$, $\mathbb{M}(u) = \mu < \mu_e$ and $\mathbb{E}(u) < \mathcal{E}_1(\mu)$. By the persistence of (1.25) proved above, if (1.25) is initially not satisfied, neither is it at any other time. If $\mathbb{K}_2(u(t)) < 0$ and $\|\nabla u(t)\|_2 \leq 1$, then Lemma 2.3 implies that $\|u(t)\|_{H_x^1}^2 \lesssim \mu \ll 1$. If $\mathbb{K}_2(u(t)) \geq 0$, then (2.25) yields a priori bound on H_x^1 by the mass-energy conservation. Hence the solution u is global and bounded in H_x^1 for all $t \in \mathbb{R}$. Moreover, we have the scattering to the ground states by [9] if $\mathbb{H}^0(u(t)) \lesssim \mu$, and it is preserved for all $t \in \mathbb{R}$.

Thus we have obtain the global existence part of Theorem 1.1.

Lemma 3.1. *For any $u(0) \in H_r^1$ satisfying $\mathbb{M}(u(0)) = \mu \leq \mu_e$ and $\mathbb{E}(u(0)) = \varepsilon < \mathcal{E}_1(\mu)$, the corresponding solution u of (1.7) is global in time iff (1.25) fails. Then it is never satisfied at any $t \in \mathbb{R}$. Moreover, the global solution u satisfies one of the following*

- (i) $\|u(t)\|_{H_x^1}^2 \lesssim \mu$ for all $t \in \mathbb{R}$, scattering to \mathcal{S}_0 .
- (ii) $\mu \lesssim \|u(t)\|_{H_x^1}^2 \lesssim \varepsilon + \mu$ and $\mathbb{K}_2(u(t)) \geq \kappa(\mu, \mathcal{E}_1(\mu) - \varepsilon)$ for all $t \in \mathbb{R}$,

where $\kappa > 0$ is the same as in (2.21).

The rest of this paper is devoted to the scattering in (ii).

4. MODULATION AND LINEARIZED EQUATIONS AROUND THE GROUND STATE

Here we recall the coordinate in [9] around the small ground state, and observe that it can be applied to large solutions as long as the mass $\mathbb{M}(u)$ is small, including the excited solitons. For any $\mu > 0$, denote

$$H^1[\mu] := \{\varphi \in H_r^1(\mathbb{R}^3) \mid \mathbb{M}(\varphi) < \mu\}. \quad (4.1)$$

Let $\Phi : D_b \rightarrow H_r^1$ be the small ground states as in Lemma 2.2. We have the following nonlinear projection to them.

Lemma 4.1. *There exist $0 < \mu_p \leq \mu_b$ and a unique mapping $H^1[\mu_p] \ni \varphi \mapsto (z, \eta) \in D_p \times H^1[\mu_p]$, where $D_p := \{z \in \mathbb{C} \mid |z|^2 < 2\mu_p\}$, such that*

$$\begin{aligned} \varphi &= \Phi[z] + \eta, \\ \eta &\in \mathcal{H}_c[z] := \{\psi \in H^1(\mathbb{R}^3) \mid \langle i\psi \mid \partial_j \Phi[z] \rangle = 0 \ (j = 1, 2)\}, \end{aligned} \quad (4.2)$$

where ∂_j denotes the derivative with respect to the real and imaginary parts of $z = z_1 + iz_2$. Moreover, the map $\varphi \mapsto (z, \eta)$ is smooth and injective from $H^1[\mu_p]$ to $\mathbb{C} \times H_r^1$. Furthermore, the orthogonal projection P_c to the continuous spectrum subspace

$$P_c \varphi := 1 - \phi_0(\varphi \mid \phi_0) \quad (4.3)$$

is bijective from $\mathcal{H}_c[z]$ onto

$$\mathcal{H}_c[0] = P_c H^1(\mathbb{R}^3) = \{\varphi \in H^1(\mathbb{R}^3) \mid \langle i\varphi | \phi_0 \rangle = 0 = \langle i\varphi | i\phi_0 \rangle\}, \quad (4.4)$$

for any $z \in D_p$, and

$$R[z] := (P_c|_{\mathcal{H}_c[z]})^{-1} \quad (4.5)$$

is a compact and continuous perturbation of identity in the operator norm on any space between $H^2 \cap W^{1,1}$ and $H^{-2} + L^\infty$.

In [9], the above coordinate was defined on a small ball of $H^1(\mathbb{R}^3)$. However, it is easy to see that the smallness in L^2 suffices, since z is determined by solving the orthogonality conditions

$$\langle \varphi - \Phi[z] | i\partial_j \Phi[z] \rangle = 0 \quad (j = 1, 2), \quad (4.6)$$

by the implicit function theorem. The derivative of the left hand side equals

$$\langle \varphi - \Phi[z] | i\partial_k \partial_j \Phi[z] \rangle - \langle \partial_k \Phi[z] | i\partial_j \Phi[z] \rangle, \quad (j, k \in \{1, 2\}). \quad (4.7)$$

The second term is a non-degenerate 2×2 matrix of $O(1)$, while the first term is bounded by $\|\varphi - \Phi[z]\|_2 \lesssim \sqrt{\mu_p} \ll 1$, thereby the implicit function theorem works, leading to the conclusion. The L^2 bound follows from the orthogonality

$$\mathbb{M}(\varphi) = \mathbb{M}(\Phi[z]) + \mathbb{M}(\eta). \quad (4.8)$$

The operator $R[z]$ is linear, so it does not need any smallness condition. Actually, the above lemma holds without even assuming that the function is in H^1 . Hence every solution u in $H^1[\mu_p]$ can be written as

$$u(t) = \Phi[z(t)] + R[z(t)]\xi(t) = \Phi[z(t)] + \eta(t), \quad (4.9)$$

uniquely, and the equation for u can be rewritten for (z, ξ) as, regarding $\mathbb{C} \simeq \mathbb{R}^2$,

$$\begin{cases} \dot{z} - i\Omega[z]z = \underline{N}(z, R[z]\xi) := M(z, R[z]\xi)^{-1} \langle N(z, R[z]\eta) | D\Phi[z] \rangle, \\ i\dot{\xi} + H\xi = B[z]\xi + \tilde{N}(z, \xi), \end{cases} \quad (4.10)$$

where $M(\cdot)$ is a 2×2 matrix and $N(\cdot, \cdot)$ is a scalar defined by

$$\begin{aligned} M_{j,k}(z, \eta) &:= \langle i\partial_j \Phi[z] | \partial_k \Phi[z] \rangle - \langle i\eta | \partial_j \partial_k \Phi[z] \rangle, \\ N(z, \eta) &:= \mathfrak{s} \left\{ 2\Phi[z]|\eta|^2 + \overline{\Phi[z]}\eta^2 + |\eta|^2\eta \right\}, \end{aligned} \quad (4.11)$$

$B[z]$ is the ‘‘potential part’’ by the small soliton, namely

$$B[z]\xi := \mathfrak{s}P_c\{2|\Phi[z]|^2 R[z]\xi + \Phi[z]^2 \overline{R[z]\xi}\}, \quad (4.12)$$

which is \mathbb{R} -linear but not \mathbb{C} -linear, and $\tilde{N}(\cdot, \cdot)$ is the quadratic part

$$\tilde{N}(z, \xi) := P_c\{N(z, R[z]\xi) - iD\Phi[z]\underline{N}(z, R[z]\xi)\}. \quad (4.13)$$

We introduce some notation for the linearized solutions. For any $s \in \mathbb{R}$ and any set X , the set of X -valued functions defined around s is denoted by

$$X\{s\} := \{u : I \rightarrow X \mid s \in \exists I \subset \mathbb{R}\}. \quad (4.14)$$

For any interval $I \subset \mathbb{R}$, $z \in C(I; \mathbb{C})$, $s_0 \in I$ and $u \in H^1\{s_0\}$, the linear solution v of

$$i\dot{v} + Hv = B[z]v, \quad v(s_0) = u(s_0) \quad (4.15)$$

is denoted by

$$u[z, s_0] := v \in C(I; H^1). \quad (4.16)$$

Note that this depends on $u(s_0)$ but not on $u(t)$ at the other time $t \neq s_0$. Indeed, there is no point for u to depend on t in the definition, but this convention avoids writing the same time s_0 twice. We can apply it to time-independent u as well. Obviously

$$\forall s_1 \in I, \quad u[z, s_0][z, s_1] = u[z, s_0], \quad (4.17)$$

while the solution without the potential $B[z]$ is given by

$$u[0, s_0] = e^{i(t-s_0)H}u(s_0). \quad (4.18)$$

The associated Duhamel integral is denoted by

$$\mathcal{D}f[z, s_0](t) := \int_{s_0}^t f[z, s](t)ds, \quad (4.19)$$

so that $v := \mathcal{D}f[z, s_0]$ satisfies

$$i\dot{v} + Hv = B[z]v + f, \quad v(s_0) = 0. \quad (4.20)$$

Hence for any $\varphi \in H^1$, $z \in C\{s_0\}$ and $f \in H^1\{s_0\}$, the solution of

$$i\dot{\xi} + H\xi = B[z]\xi + f, \quad \xi(s_0) = \varphi \quad (4.21)$$

is uniquely given by

$$\xi = \varphi[z, s_0] + \mathcal{D}f[z, s_0]. \quad (4.22)$$

Another notation

$$u[z, s_0]_> := \begin{cases} u(t) & (t < s_0) \\ u[z, s_0] & (t > s_0) \end{cases} \quad (4.23)$$

is convenient to “turn off the nonlinearity” after some time. Indeed if u solves

$$i\dot{u} + Hu = B[z]u + f \quad (4.24)$$

and $s_0 < s_1$, then we have

$$\begin{cases} u = u[z, s_0] + \mathcal{D}f[z, s_0], \\ u[z, s_1]_> = u[z, s_0] + \mathcal{D}1_{t < s_1}f[z, s_0]. \end{cases} \quad (4.25)$$

Next a few (semi)norms are introduced for space-time functions. For $s \in \mathbb{R}$, put

$$\text{Stz}^s := L_t^\infty H_x^s \cap L_t^2 B_{6,2}^s, \quad \text{Stz}^{**s} := L_t^1 H_x^s + L_t^2 B_{6/5,2}^s, \quad \mathfrak{st} := L_t^4 L_x^6. \quad (4.26)$$

Stz^s is the full Strichartz norm for H^s solutions, and $\text{Stz}^{1/2} \subset L_t^4 B_{3,2}^{1/2} \subset \mathfrak{st}$ by interpolation and Sobolev.

The next semi-norm is a bit more involved. It is needed for long-time perturbation argument for the radiation part ξ , whose equation contains quadratic terms. For $T_0 < T_1 \leq T_2$, $z \in C((T_0, T_2); \mathbb{C})$ and $u \in C((T_0, T_1); H^1)$,

$$\|u\|_{[z; T_0, T_1; T_2]} := \sup_{T_0 < T < T_1} \|u[z, T]_{>} - u[z, T_0]\|_{\text{st}(T_0, T_2)} \quad (4.27)$$

is a semi-norm which vanishes if and only if $u = u[z, T_0]$ solves $iu + Hu = B[z]u$ on (T_0, T_1) in H^1 . It measures how much u deviates from the linear evolution between T_0 and T_1 and its influence until $t < T_2$. If u solves (4.24) on (T_0, T_1) , then

$$\|u\|_{[z; T_0, T_1; T_2]} = \sup_{T_0 < T < T_1} \|\mathcal{D}1_{t < T} f[z, T_0]\|_{\text{st}(T_0, T_2)}. \quad (4.28)$$

Since we use only the Strichartz type estimates, i.e. L_t^p norms, the right hand side will be estimated in the same way as $\|\mathcal{D}f[z, T_0]\|_{\text{st}(T_0, T_1)}$. This semi-norm will be used mostly to bound

$$\|u\|_{[z; T_0, T_1; T_2]} \geq \max(\|u - u[z, T_0]\|_{\text{st}(T_0, T_1)}, \|u[z, T_1] - u[z, T_0]\|_{\text{st}(T_1, T_2)}). \quad (4.29)$$

The idea of long-time perturbation in this type of norms, together with the use of non-admissible Strichartz (as in Lemma 4.4 below), was introduced in [13] to treat quadratic and sub-quadratic nonlinearity.

The above semi-norm is subadditive for gluing intervals:

Lemma 4.2. *For $T_0 < T_1 < T_2 < T_3$,*

$$\|u\|_{[z; T_0, T_2; T_3]} \leq \|u\|_{[z; T_0, T_1; T_3]} + \|u\|_{[z; T_1, T_2; T_3]}. \quad (4.30)$$

Proof. The left is the supremum of $\|u[z, T]_{>} - u[z, T_0]\|_{\text{st}(T_0, T_3)}$ over $T_0 < T < T_2$. For $T_0 < T < T_1$, it is trivially bounded by $\|u\|_{[z; T_0, T_1; T_3]}$. For $T_1 < T < T_2$, we have

$$\begin{aligned} \|u[z, T]_{>} - u[z, T_0]\|_{\text{st}(T_0, T_3)} &\leq \|u[z, T]_{>} - u[z, T_1]_{>}\|_{\text{st}(T_0, T_3)} \\ &\quad + \|u[z, T_1]_{>} - u[z, T_0]\|_{\text{st}(T_0, T_3)}. \end{aligned} \quad (4.31)$$

The first term on the right equals

$$\|u[z, T]_{>} - u[z, T_1]_{>}\|_{\text{st}(T_1, T_3)} \leq \|u\|_{[z; T_1, T_2; T_3]}, \quad (4.32)$$

and the second term is bounded by $\|u\|_{[z; T_0, T_1; T_3]}$. \square

Now we derive the standard Strichartz estimates for the linearized equation, with uniformly small z .

Lemma 4.3. *Let $I = (T_0, T_1)$ be an interval and $z \in C(I; \mathbb{C})$ with $\|z\|_{L^\infty(I)} \ll 1$. Then for any $s_0 \in I$, $\varphi \in P_c H^1(\mathbb{R}^3)$, $f \in C(I; \mathcal{S}')$, $T \in (T_0, T_1)$ and $\theta \in [0, 1]$,*

$$\|\varphi[z, s_0]\|_{\text{Stz}^\theta(I)} \lesssim \|\varphi(s_0)\|_{H^\theta} \sim \inf_{t \in I} \|\varphi[z, s_0](t)\|_{H^\theta} \sim \sup_{t \in I} \|\varphi[z, s_0](t)\|_{H^\theta}, \quad (4.33)$$

$$\|\mathcal{D}P_c f[z, s_0]\|_{\text{Stz}^\theta(I)} \lesssim \|f\|_{\text{Stz}^{*\theta}(I)}, \quad \|\mathcal{D}P_c f[z, s_0]\|_{[z; T_0, T; T_1]} \lesssim \|f\|_{\text{Stz}^{*1/2}(T_0, T)}.$$

Moreover, if $\sup I = \infty$ then we have the scattering of $u := \varphi[z, s_0] + \mathcal{D}P_c f[z, s_0]$ as $t \rightarrow \infty$, namely the strong convergence of $e^{-itH}u(t)$ in H^1 , and furthermore, $u(t) \rightarrow 0$ in L_x^6 . The same holds for $t \rightarrow -\infty$.

Proof. Let $\xi := \varphi[z, s_0] + \mathcal{D}P_c f[z, s_0]$. Then the Strichartz estimate for $e^{itH} P_c$ yields

$$\|\xi\|_{\text{Stz}^\theta(I)} \lesssim \|\varphi(s_0)\|_{H^\theta} + \|B[z]\xi + P_c f\|_{\text{Stz}^{*\theta}(I)}, \quad (4.34)$$

where the term $B[z]\xi$ is bounded by

$$\|B[z]\xi\|_{L_t^2 B_{6/5,2}^\theta(I)} \lesssim \|z\|_{L_t^\infty(I)}^2 \|\xi\|_{L_t^2 B_{6,2}^\theta(I)} \ll \|\xi\|_{L_t^2 B_{6,2}^\theta(I)}, \quad (4.35)$$

and absorbed by the left. This yields the latter three estimates in the lemma.

If $\sup I = \infty$, for any increasing sequence $t_n \rightarrow \infty$, we have

$$[e^{-itH} \xi(t)]_{t=t_m}^{t=t_n} = \int_{t_m}^{t_n} e^{-isH} (B[z]\xi + P_c f) dt, \quad (4.36)$$

where the last term tends to 0 in H_x^1 by the above estimate, together with the property of $L_t^1(I)$ and $L_t^2(I)$. Hence $e^{-itH} \xi(t)$ converges as $t \rightarrow \infty$ strongly in H^1 . Then the decay in L_x^6 follows from the $L^{6/5} \rightarrow L^6$ decay estimate, the Sobolev embedding $H_x^1 \subset L_x^6$ and the density argument.

For the norm equivalence, let $u := \varphi[z, s_0]$ and $u_0 := \varphi[0, s_0]$. Then by the same Strichartz as above,

$$\|u - u_0\|_{\text{Stz}^\theta(I)} \lesssim \|B[z]u\|_{L_t^2 B_{6/5,2}^\theta(I)} \ll \|u\|_{L_t^2 B_{6,2}^\theta(I)}, \quad (4.37)$$

and so the right hand side is equivalent to (since $\varphi \in P_c H^1$)

$$\|u_0\|_{L_t^2 B_{6,2}^\theta(I)} \lesssim \|u_0(s_0)\|_{H^\theta} \sim \|\langle HP_c \rangle^{\theta/2} u_0(s_0)\|_{L^2} \sim \|u_0\|_{L_t^\infty H_x^\theta(I)}, \quad (4.38)$$

which implies the first estimate in the lemma. \square

We also need non-admissible Strichartz estimates.

Lemma 4.4. *Let $(p_0, q_0), (p_1, q_1) \in (1, \infty) \times (2, 6]$ and $\sigma_j := 2/p_j + 3(1/q_j - 1/2)$ satisfy*

$$\sigma_0 + \sigma_1 = 0 < \sigma_j - 1/p_j, \quad |\sigma_j| \leq 2/3. \quad (4.39)$$

Then there exists $C > 0$ such that under the assumptions of the above lemma,

$$\|\mathcal{D}P_c f[z, s_0]\|_{L_t^{p_0} L_x^{q_0}(I)} \leq C \|f\|_{L_t^{p'_1} L_x^{q'_1}(I)}. \quad (4.40)$$

If $(p_0, q_0) = (4, 6)$, then for $T_0 < T \leq T_1$,

$$\|\mathcal{D}P_c f[z, s_0]\|_{[z; T_0, T; T_1]} \leq C \|f\|_{L_t^{p'_1} L_x^{q'_1}(T_0, T)}. \quad (4.41)$$

Proof. This set of estimates for the free Schrödinger equation was proved by Kato [11] for $q_j < 2^*$, and by Vilela [25] for $q_j = 2^*$. It is transferred to the time independent equation $e^{itH} P_c$ by Yajima's argument of bounded wave operators [26]. More precisely, the condition $|\sigma_j| \leq 2/3$ is not needed in Kato, but only in the double endpoint case $q_0 = q_1 = 6$ by Vilela, in the form $p_0 = p_1 \in [6/5, 6]$. The above lemma is infected by this condition including non-endpoint cases, because we use the double endpoint estimate to treat the time-dependent potential part as a small perturbation. Let $u := \mathcal{D}P_c f[z, s_0]$. Then by the above estimates on $e^{itH} P_c$,

$$\|u\|_{(L_t^{p_0} L_x^{q_0} \cap L_t^{p_2} L_x^6)(I)} \lesssim \|B[z]u\|_{L_t^{p_2} L_x^{6/5}(I)} + \|f\|_{L_t^{p'_1} L_x^{q'_1}(I)}, \quad (4.42)$$

where $p_2 := \frac{2}{\sigma_0+1} \in [6/5, 6]$. The potential term is bounded by

$$\|B[z]u\|_{L_t^{p_2} L_x^{6/5}(I)} \lesssim \|z\|_{L_t^\infty(I)}^2 \|u\|_{L_t^{p_2} L_x^6(I)} \ll \|u\|_{L_t^{p_2} L_x^6(I)}, \quad (4.43)$$

and thus we obtain the desired result as in the previous lemma. \square

5. LINEARIZED PROFILE DECOMPOSITION

Now we develop a profile decomposition for the linearized equation of the radiation part in (4.10). For that purpose, we need a similar notation to the above for sequences. For any sequences a, b, c, \dots , the sequence in the form

$$\mathbb{N} \ni n \mapsto \mathcal{X}(a_n, b_n, c_n, \dots) \quad (5.1)$$

for *any expression* \mathcal{X} (as long as it is well defined), is denoted by

$$\mathcal{X}(a, b, c, \dots) := \{\mathcal{X}(a_n, b_n, c_n, \dots)\}_n. \quad (5.2)$$

The same convention applies when the sequence is defined only for large $n \in \mathbb{N}$. When $\mathcal{X} = \{\mathcal{X}_n\}_n$ is a sequence of sets, then it is regarded as the product set:

$$x \in \mathcal{X} \iff \forall n \in \mathbb{N}, x_n \in X_n. \quad (5.3)$$

The same convention is applied to \lim, \sup, \inf , etc., for any sequence $X = \{X_n\}_n$:

$$X \rightarrow \lim X := \lim_{n \rightarrow \infty} X_n, \quad \sup X := \sup_n X_n, \quad \text{etc.} \quad (5.4)$$

unless the limit is explicitly associated with another parameter.

The set of open intervals is denoted by

$$\mathcal{I} := \{(a, b) \subset \mathbb{R} \mid a < b\}. \quad (5.5)$$

For any $I \in \mathcal{I}^{\mathbb{N}}$, the set $\text{SBC}(I)$ of sequences of uniformly small, bounded and continuous functions is defined by the following. For any $z \in C(I; \mathbb{C})$

$$z \in \text{SBC}(I) \quad (5.6)$$

iff $\sup_{n \in \mathbb{N}} \sup_{t \in I_n} |z_n(t)| \ll 1$ and

$$\forall \varepsilon > 0, \exists \delta > 0, \forall s, \forall t \in I, \sup |s - t| < \delta \implies \sup |z(s) - z(t)| < \varepsilon. \quad (5.7)$$

The smallness requirement can be determined by V . Since it can be fixed throughout the paper but does not play any role, we leave it unspecified.

For any $I \in \mathcal{I}^{\mathbb{N}}$, any $z \in C(I; \mathbb{C})$ and any $z_\infty \in C(\mathbb{R}; \mathbb{C})$, we say that $z \rightarrow z_\infty$ locally uniformly on I iff for all $0 < T < \infty$

$$\lim_{n \rightarrow \infty} \sup_{t \in [-T, T] \cap I_n} |z_n(t) - z_\infty(t)| = 0. \quad (5.8)$$

Let $\mathcal{I}^{\mathbb{N}} \ni I \ni s, C(I; \mathbb{C}) \ni z$ and $H^1\{s\} \ni u$. Note that they are all sequences by the above convention. Suppose that $\lim P_c u(s)$ is weakly convergent in H^1 . Then the sequence $v \in C(I; H^1)$ solving

$$\forall n \in \mathbb{N}, \quad i\dot{v}_n + H v_n = B[z_n]v_n \quad (\text{on } I_n), \quad v_n(s_n) = \lim_{k \rightarrow \infty} P_c u_k(s_k) \quad (5.9)$$

is denoted by

$$u[z, s]_{(\infty)} := \{v_n\}_n = \left\{ \lim_{k \rightarrow \infty} P_c u_k(s_k) \right\} [z, s]. \quad (5.10)$$

Let $s' \in I$ be another sequence of times, then

$$u[z, s]_{(\infty)}[z, s'] = u[z, s]_{(\infty)}. \quad (5.11)$$

In the autonomous case $z \equiv 0$, the above object can be defined by translation:

$$\forall t \in \mathbb{R}, \quad u[0, s]_{(\infty)}(t) = e^{i(t-s)H} \lim_{k \rightarrow \infty} P_c u_k(s_k), \quad (5.12)$$

which trivializes the limiting behavior as $n \rightarrow \infty$. The presence of z disables such a precise description. However, we do not need so much to prove the scattering result, but uniform integrability in the Strichartz norms will suffice, which is given by the following lemma.

Lemma 5.1. *Let $s \in I \in \mathcal{I}^{\mathbb{N}}$, $z \in \text{SBC}(I)$ and $u \in P_c H^1\{s\}$ be sequences such that $\sup \|u(s)\|_{H^1} < \infty$. Then after extracting a subsequence, there exist $\varphi \in P_c H^1$ and $z_\infty \in C(\mathbb{R}; \mathbb{C})$ such that $u(s) \rightharpoonup \varphi$ weakly in H^1 , $z(s+t) \rightarrow z_\infty$ locally uniformly on $I - s$, and*

$$\|u[z, s]_{(\infty)} - \varphi[z_\infty, 0](t - s)\|_{\text{Stz}^1(I)} \rightarrow 0. \quad (5.13)$$

Moreover, for any $0 < T < \infty$,

$$\|u[z, s] - u[z, s]_{(\infty)}\|_{L_t^\infty(|t-s| < T; L_x^4)} \rightarrow 0. \quad (5.14)$$

If the convergence $u(s) \rightarrow \varphi$ is strong in H^1 , then

$$\|u[z, s] - u[z, s]_{(\infty)}\|_{\text{Stz}^1(I)} \rightarrow 0. \quad (5.15)$$

Proof. First, the uniform continuity of SBC allows us to extend z_n to \mathbb{R} so that we may assume $I = \mathbb{R}^{\mathbb{N}}$ without losing generality. The uniform boundedness in H^1 allows us to pass to a subsequence such that $u(s) \rightharpoonup \exists \varphi$. Let

$$\begin{aligned} t \mapsto \zeta(t) &:= z(t + s) \in \text{SBC}(\mathbb{R}^{\mathbb{N}}), \\ t \mapsto v(t) &:= \varphi[\zeta, 0](t) = u[z, s]_{(\infty)}(t + s) \in C(\mathbb{R}; P_c H^1)^{\mathbb{N}}. \end{aligned} \quad (5.16)$$

Since $\zeta \in \text{SBC}$, Ascoli-Arzelà implies that, after extracting a subsequence, $\zeta \rightarrow \exists \zeta_\infty$ in $C(\mathbb{R}; \mathbb{C})$ with $\|\zeta_\infty\|_{L^\infty(\mathbb{R})} \ll 1$. Let $v_\infty = \varphi[\zeta_\infty, 0] \in C(\mathbb{R}; P_c H^1)$. Then we have

$$i\dot{v} + Hv = B[\zeta]v, \quad i\dot{v}_\infty + Hv_\infty = B[\zeta_\infty]v_\infty, \quad \forall n, \quad v_n(0) = \varphi = v_\infty(0). \quad (5.17)$$

We have the full Strichartz estimates on v_∞ by Lemma 4.3, and

$$\lim_{T \rightarrow \infty} \|v_\infty\|_{L^2(|t| > T; B_{6,2}^1)} = 0. \quad (5.18)$$

Since $\zeta \rightarrow \zeta_\infty$ in $L^\infty(|t| \leq T)$ for any $T < \infty$, $B[\zeta] \rightarrow B[\zeta_\infty]$ in the operator norm of $B_{6,2}^1 \rightarrow B_{6/5,2}^1$, uniformly on $|t| \leq T$. Hence by the Strichartz estimate on $e^{itH} P_c$, and using $H_{6/5}^1 \subset B_{6/5,2}^1$,

$$\begin{aligned} \|v - v_\infty\|_{\text{Stz}^1(|t| < T)} &\lesssim \|B[\zeta]v - B[\zeta_\infty]v_\infty\|_{L_t^2(|t| < T; H_{6/5}^1)} \\ &\lesssim \|\zeta - \zeta_\infty\|_{L^\infty(|t| < T)} \|v_\infty\|_{L_t^2(|t| < T; H_6^1)} \\ &\quad + \|(\zeta, \zeta_\infty)\|_{L^\infty(|t| < T)} \|v - v_\infty\|_{L_t^2(|t| < T; H_6^1)}. \end{aligned} \quad (5.19)$$

Thus, the last term being absorbed by the left, we obtain

$$\|v - v_\infty\|_{\text{Stz}^1(|t| < T)} \rightarrow 0. \quad (5.20)$$

Applying the same estimate to the Duhamel with $e^{itH}P_c$ from $t = \pm T$, we obtain

$$\begin{aligned} & \|v - v_\infty\|_{\text{Stz}^1(|t|>T)} \\ & \lesssim \|v(\pm T) - v_\infty(\pm T)\|_{H_x^1} + \|B[\zeta]v - B[\zeta_\infty]v_\infty\|_{L_t^2(|t|>T; H_{6/s}^1)} \\ & \lesssim o(1) + \|(\zeta, \zeta_\infty)\|_{L^\infty(|t|>T)} (\|v_\infty\|_{L_t^2(|t|>T; H_6^1)} + \|v - v_\infty\|_{L_t^2(|t|>T; H_6^1)}), \end{aligned} \quad (5.21)$$

where the last term is absorbed by the left. Thus we obtain

$$\limsup \|v - v_\infty\|_{\text{Stz}^1(\mathbb{R})} \ll \|v_\infty\|_{L_t^2(|t|>T; H_6^1)}. \quad (5.22)$$

Sending $T \rightarrow \infty$, we see that the right side is zero, namely (5.13).

To prove (5.14), let $w(t) := u[z, s](t + s)$ and $\gamma := (v - w)[0, 0]$. Then by the Strichartz estimate on $e^{itH}P_c$,

$$\begin{aligned} & \|(v - w) - \gamma\|_{(L_t^\infty L_x^2 \cap L_t^2 L_x^6)(|t|<T)} \\ & \lesssim \|B[\zeta](v - w)\|_{L_t^2 L_x^{6/5}(|t|<T)} \ll \|v - w\|_{L_t^2 L_x^6(|t|<T)}, \end{aligned} \quad (5.23)$$

so it is bounded by $\|\gamma\|_{L_t^2 L_x^6(|t|<T)}$, which tends to 0, because $\gamma \rightarrow 0$ in $L_t^\infty L_x^4(|t| < T)$ and bounded in $L_t^2 H_6^1$. Interpolating with the uniform bounds on v and w , we get $v - w - \gamma \rightarrow 0$ also in $L_t^\infty L_x^4(|t| < T)$, hence for $v - w$ as well.

The proof of (5.15) is similar but easier. We add one derivative to the Strichartz norms and extend to the real line, such as $L_t^2 H_6^1(\mathbb{R})$. Then $\gamma = e^{itH}\gamma(0) \rightarrow 0$ in this norm, since $\gamma(0) \rightarrow 0$ strongly in H^1 . So $v - w$ is also vanishing. \square

The linearized equation does not preserve either the mass or the energy, because $B[z]$ is not even \mathbb{C} -linear, but the next lemma suffices for the profile decomposition. Since $H > 0$ on $P_c H^1$, its fractional power is defined. For any $\theta \in [0, 1]$, the inner product is defined by

$$\mathbb{H}_\theta(\varphi, \psi) := \frac{1}{2} \langle H^\theta P_c \varphi | \psi \rangle, \quad \mathbb{H}_\theta(\varphi) := \mathbb{H}_\theta(\varphi, \varphi), \quad (5.24)$$

such that $\mathbb{M}(\varphi) = \mathbb{H}_0(\varphi)$ and $\mathbb{H}(\varphi) = \mathbb{H}_1(\varphi)$ for all $\varphi \in P_c H^1$.

Lemma 5.2. *Let $s \in I \in \mathcal{I}^\mathbb{N}$ and $z \in \text{SBC}(I)$. Let $v^0, v^1 \in C(I; P_c H^1)$ be two sequences of linearized solutions, i.e. $v^j = v^j[z, s]$ for $j = 0, 1$. Suppose that $v^0(s)$ strongly converges in H^1 and that $v^1(s) \rightarrow 0$ weakly in H^1 . Then*

$$\forall \theta \in [0, 1], \quad \|\mathbb{H}_\theta(v^0, v^1)\|_{L_t^\infty(I)} \rightarrow 0. \quad (5.25)$$

Proof. It suffices to show $\mathbb{H}_\theta(v^0, v^1)(s') \rightarrow 0$ for any $s' \in I$, along a subsequence. We use the unitarity

$$\tilde{v}^j := e^{i(s-t)H} v^j \implies \mathbb{H}_\theta(v^0, v^1) = \mathbb{H}_\theta(\tilde{v}^0, \tilde{v}^1) \quad (5.26)$$

and the Duhamel formula

$$\begin{aligned} \tilde{v}^j(t) &= v^j(s) + \mathcal{D}^j(t), \\ \mathcal{D}^j(t) &:= \int_0^{t-s} e^{-it'H} B[z(s+t')] v^j(s+t') dt'. \end{aligned} \quad (5.27)$$

$v^0(s) \rightarrow \exists \varphi$ strongly in H^1 by the assumption. Extracting a subsequence, we may assume $s' - s \rightarrow \exists s'_\infty \in [-\infty, \infty]$. Then

$$\mathcal{D}^0(s') \rightarrow \int_0^{s'_\infty} e^{-itH} B[z_\infty] \varphi[z_\infty, 0] dt \quad (5.28)$$

strongly in H^1 , by Lemma 5.1, (5.13), after passing to a subsequence. For \mathcal{D}^1 , we use the L^p decay estimate on $e^{itH} P_c$. Fix $0 < \delta \ll 1$ such that $1/q_\pm := 1/6 \pm \delta \in [1/p, 1/2)$. Then for any $0 < T < \infty$,

$$\begin{aligned} \|\mathcal{D}^1\|_{L_t^\infty(I; L^{q_+ + L^{q_-}})_x} &\lesssim \int_{|t-s|>T} |t-s|^{-1-3\delta} \|B[z]v^1(t)\|_{L_x^{q_-}} dt \\ &\quad + \int_{|t-s|<T} |t-s|^{-1+3\delta} \|B[z]v^1(t)\|_{L_x^{q_+}} dt \\ &\lesssim T^{-3\delta} \|v^1\|_{L_t^\infty L_x^2} + \|v^1\|_{L_t^\infty(|t-s|<T; L_x^4)}, \end{aligned} \quad (5.29)$$

and the last term is vanishing by Lemma 5.1, (5.14). Since $T > 0$ is arbitrary, we deduce that $\mathcal{D}^1(s') \rightarrow 0$. Hence $\tilde{v}^1(s') \rightarrow 0$, while $\tilde{v}^0(s')$ is strongly convergent. Thus we obtain $\mathbb{H}_\theta(v^0, v^1)(s') \rightarrow 0$ as desired. \square

Using the above lemmas, we are ready to prove the profile decomposition for the linearized equation for ξ .

Lemma 5.3. *Let $s \in I \in \mathcal{S}^\mathbb{N}$ and $z \in \text{SBC}(I)$. Let $\psi \in (P_c H^1)^\mathbb{N}$ be a bounded sequence. Then passing to a subsequence, there exist $J^* \in \mathbb{N} \cup \{\infty\}$ and $s^j \in I$ for each $\mathbb{N}_0 \ni j < J^*$ with the following properties. Let $\nu := \psi[z, s] \in C(I; P_c H^1)$.*

- (i) $s^0 = s$ and $|s^j - s^k| \rightarrow \infty$ for each $j \neq k < J^*$.
- (ii) For $j < J^*$, $\nu(s^j) \rightarrow \exists \varphi_\infty^j \in H^1$ weakly. Put $\lambda^j := \nu[z, s^j]_{(\infty)} = \varphi_\infty^j[z, s^j]$. Then $\lambda^j(s^k) \rightarrow 0$ weakly in H^1 for $j \neq k$, and $\varphi_\infty^j \neq 0$ for $j > 0$.
- (iii) For each finite $J \leq J^*$, put $\gamma^J := \nu - \sum_{0 \leq j < J} \lambda^j$. For $j < J$, $\gamma^J(s^j) \rightarrow 0$ weakly in H^1 , and for all $\theta \in [0, 1]$,

$$\sum_{0 \leq j < J} \|\lambda^j\|_{L_t^\infty(I; \dot{H}_x^\theta)}^2 + \|\gamma^J\|_{L_t^\infty(I; \dot{H}_x^\theta)}^2 \sim \|\psi\|_{\dot{H}^\theta}^2 + o(1). \quad (5.30)$$

- (iv) For $0 \leq \theta < 1$,

$$\lim_{J \rightarrow J^*} \limsup \|\gamma^J\|_{[L_t^\infty L_x^4, \text{Stz}^1]_\theta(I)} = 0. \quad (5.31)$$

We call the decomposition given by the above lemma

$$\psi[z, s] = \sum_{0 \leq j < J} \lambda^j + \gamma^J, \quad \lambda^j := \psi[z, s][z, s^j]_{(\infty)}, \quad (5.32)$$

the linearized profile decomposition.

Proof. The sequence $s^j \in I$ is defined inductively as follows. First, let $s^0 := s$. Passing to a subsequence, we may assume $\nu(s^0) \rightarrow \exists \varphi_\infty^0$. Then $\lambda^0 := u[z, s^0]_{(\infty)}$ and $\gamma^1 := \nu - \lambda^0$ are defined with $\gamma^1(s^0) \rightarrow 0$. The boundedness (in H^1) of ψ implies that ν and λ^0 are uniformly bounded, so is γ^1 .

Let $J \in \mathbb{N}$ and suppose that uniformly bounded $\gamma^J \in C(I; H^1)$ and $s^j \in I$ have been defined for $j < J$ such that $\gamma^J(s^j) \rightarrow 0$. Since $\|\gamma^J\|_{L_t^\infty(I; L_x^4)}$ is bounded, we can define $s^J \in I$ such that

$$\|\gamma^J\|_{L_t^\infty(I; L_x^4)} = \|\gamma^J(s^J)\|_{L_x^4} + o(1). \quad (5.33)$$

If the left hand sequence tends to 0, put $J^* = J$ and the definition is terminated. Otherwise, put $\lambda^J := \gamma^J[z, s^J]_{(\infty)}$ and $\gamma^{J+1} := \gamma^J - \lambda^J$. Passing to a subsequence, we may assume $\gamma^J(s^J) \rightarrow \exists \varphi_\infty^J \neq 0$ in H^1 and $z(t + s^J) \rightarrow \exists z_\infty^J(t) \in C(\mathbb{R}; \mathbb{C})$ locally uniformly on $I - s^J$. Then $\gamma^{J+1}(s^J) \rightarrow 0$ is obvious. Since $\gamma^J(s^j) \rightarrow 0$ for $j < J$, we have $\gamma^J \rightarrow 0$ in $L_t^\infty(|t - s^j| < T; L_x^4)$ for any $T < \infty$, by Lemma 5.1, (5.14). Hence $|s^J - s^j| \rightarrow \infty$. Then from Lemma 5.1, (5.13), together with the Strichartz bound on $\varphi_\infty^J[z_\infty^J, 0]$, we deduce that $\lambda^J(s^j) \rightarrow 0$, and so $\gamma^{J+1}(s^j) \rightarrow 0$. The same argument implies that $\lambda^j(s^J) \rightarrow 0$ as well.

Hence we can iterate the same procedure. In this way, after the diagonalization argument, we obtain the sequences s^j with the properties that $|s^j - s^k| \rightarrow \infty$ for $j \neq k$, $\gamma^J(s^j) \rightarrow 0$ for $j < J$, $\lambda^j(s^k) \rightarrow 0$ for $j \neq k$, $z(t + s^j) \rightarrow z_\infty^j(t)$, and the decomposition (5.32).

Since $\lambda^j(s^j) = \varphi_\infty^j$ while $\lambda^k(s^j) \rightarrow 0$ for $k \neq j$, Lemma 5.2 implies $\mathbb{H}_\theta(\lambda^j, \lambda^k) \rightarrow 0$, $\mathbb{H}_\theta(\lambda^j, \gamma^k) \rightarrow 0$ for $j < k$ and $\theta \in [0, 1]$. Hence

$$\mathbb{H}_\theta(\psi) = \mathbb{H}_\theta\left(\sum_{0 \leq j < J} \lambda^j(s) + \gamma^J(s)\right) = \sum_{0 \leq j < J} \mathbb{H}_\theta(\lambda^j(s)) + \mathbb{H}_\theta(\gamma^J(s)) + o(1). \quad (5.34)$$

The equivalence $\mathbb{H}_\theta(\varphi) \sim \|\varphi\|_{H^\theta}^2$ on $P_c H^1$ and Lemma 4.3 imply (5.30).

It remains to prove (5.31). By the definition of s^j , cf. (5.33), we have

$$\|\gamma^J\|_{L_t^\infty L_x^4} = \|\varphi_\infty^J\|_{L_x^4} + o(1) \lesssim \|\lambda^J\|_{L_t^\infty H_x^1} + o(1). \quad (5.35)$$

Since the right hand side is vanishing by (5.30),

$$\lim_{J \rightarrow J^*} \limsup \|\gamma^J\|_{L_t^\infty L_x^4} = 0, \quad (5.36)$$

and then by interpolation with the uniform Strichartz bound, we obtain (5.31). \square

6. NONLINEAR PERTURBATION ESTIMATES

In order to use the linearized profile decomposition to approximate the nonlinear solutions, we need a few perturbation lemmas for the nonlinear equation of ξ

$$i\dot{\xi} + H\xi = B[z]\xi + \tilde{N}(z, \xi) \quad (6.1)$$

regarding z as a given time-dependent function. Since our global knowledge on z is very poor (cf. Section 1.3), we should avoid perturbing z for long time. It leads us to prepare the following two lemmas for perturbation: Lemma 6.3 for long time intervals where ξ is small, and Lemma 6.4 for bounded time intervals where ξ may be large.

The first lemma is a perturbation of 0, or construction of dispersed solutions.

Lemma 6.1. *Let $-\infty < T_0 < T_1 \leq \infty$, $z \in C([T_0, T_1]; \mathbb{C})$ and $\varphi \in H^1(\mathbb{R}^3)$. Put*

$$\mathcal{N}_\theta := \|z\|_{L^\infty(T_0, T_1)} + \|\varphi\|_{H^\theta} \quad (6.2)$$

for $\theta \in [0, 1]$ and assume $\mathcal{N}_0 \ll 1$.

(I) If $\|\varphi[z, T_0]\|_{\text{st}(T_0, T_1)} \mathcal{N}_{1/2}^3 \ll 1$, then (6.1) has a unique solution ξ satisfying

$$\xi \in C([T_0, T_1]; H^1), \quad \xi(T_0) = \varphi, \quad \|\xi\|_{\text{st}(T_0, T_1)} \lesssim \|\varphi[z, T_0]\|_{\text{st}(T_0, T_1)}. \quad (6.3)$$

(II) Let $\xi \in C([T_0, T_1]; H^1)$ be a solution of (6.1) with $\xi(T_0) = \varphi$ satisfying

$$\|\xi\|_{\text{st}(T_0, T_1)} \mathcal{N}_{1/2}^3 \ll 1. \quad (6.4)$$

Then for any $T \in (T_0, T_1)$ and all $\theta \in [0, 1]$,

$$\|\xi[z, T]_{>} - \varphi[z, T_0]\|_{\text{Stz}^\theta(T_0, T_1)} \lesssim \mathcal{N}_\theta \|\varphi[z, T_0]\|_{\text{st}(T_0, T)} (\mathcal{N}_0 + \|\varphi[z, T_0]\|_{\text{st}(T_0, T)}), \quad (6.5)$$

and $\|\xi\|_{[z; T_0, T; T_1]} \ll \|\xi[z, T_0]\|_{\text{st}(T_0, T)} \sim \|\xi\|_{\text{st}(T_0, T)}$.

Proof. Let $\xi_0 := \varphi[z, T_0]$ and $\|\xi_0\|_{\text{st}(T_0, T_1)} =: \delta_0$. The solution ξ is obtained by the iteration argument. If ξ is a solution, then by the Strichartz estimate: Lemma 4.3,

$$\|\xi - \xi_0\|_{\text{Stz}^\theta} \lesssim \|\xi\|_{\text{Stz}^\theta} \|\xi\|_{\text{st}} (\|\xi\|_{\text{st}} + \|z\|_{L^\infty}) \quad (6.6)$$

for $\theta \in [0, 1]$. Using the non-admissible Strichartz: Lemma 4.4,

$$\begin{aligned} \|\xi\|_{\text{st}} &\leq \|\xi_0\|_{\text{st}} + C \|\tilde{N}(z, \xi)\|_{L_t^{8/3} L_x^{4/3}} \\ &\lesssim \delta_0 + \|\xi\|_{\text{st}} \|\xi\|_{L_t^8 L_x^4} (\|\xi\|_{L_t^\infty L_x^3} + \|\Phi[z]\|_{L_t^\infty L_x^3}) \\ &\lesssim \delta_0 + \|\xi\|_{\text{st}}^{3/2} \|\xi\|_{\text{Stz}^{1/2}}^{1/2} (\|\xi\|_{\text{Stz}^{1/2}} + \mathcal{N}_0). \end{aligned} \quad (6.7)$$

Suppose that $\|\xi\|_{\text{Stz}^{1/2}} \leq C \mathcal{N}_{1/2}$ and $\|\xi\|_{\text{st}} \leq C \delta_0$ for some constant $C \gg 1$ on some shorter interval. Then

$$\begin{aligned} \|\xi\|_{\text{Stz}^{1/2}} &\lesssim \mathcal{N}_{1/2} + C^2 \delta_0 (\delta_0 + \mathcal{N}_0) \|\xi\|_{\text{Stz}^{1/2}}, \\ \|\xi\|_{\text{st}} &\lesssim \delta_0 + C^2 (\delta_0 \mathcal{N}_{1/2}^3)^{1/2} \|\xi\|_{\text{st}}. \end{aligned} \quad (6.8)$$

Since $\delta_0 + \mathcal{N}_0 \lesssim \mathcal{N}_{1/2}$, we have $\delta_0 (\delta_0 + \mathcal{N}_0) \lesssim (\delta_0 \mathcal{N}_{1/2}^3)^{1/2}$. Hence, if $\delta_0 \mathcal{N}_{1/2}^3 \ll 1$ then $\|\xi\|_{\text{st}} \lesssim \delta_0$ and $\|\xi\|_{\text{Stz}^\theta} \lesssim \mathcal{N}_\theta$ for all $\theta \in [0, 1]$. Then by the continuity for extending the interval, these bounds holds on the whole $(0, T)$.

If we assume $\|\xi\|_{\text{st}(T_0, T_1)} \mathcal{N}_{1/2}^3 \ll 1$ instead of ξ_0 , then we obtain $\|\xi_0\|_{\text{st}(T_0, T)} \lesssim \|\xi\|_{\text{st}(T_0, T)}$ in the same way, starting from $T = T_0 + 0$. In both cases, repeating the Strichartz estimate on H^θ as above, we obtain (II). \square

Remark 1. Since $\dot{H}^{1/2}(\mathbb{R}^3)$ is the scaling invariant norm for the NLS without the potential, $\mathcal{N}_{1/2}$ is in general large, when we use the above lemma. In the focusing case, $\mathcal{N}_{1/2} \sim 1$ on \mathcal{S}_1 , while there is no upper bound on $\mathcal{N}_{1/2}$ in the defocusing case. However, we can expect that st is small for dispersive solutions, by which the assumptions in the above lemma can be satisfied. Note also that the estimate cannot be closed if we use only the admissible Strichartz $\text{Stz}^{1/2}$ as in (6.6), because of the quadratic terms.

If the above solution is obtained for $t \rightarrow \infty$, then it scatters.

Lemma 6.2. Let $T_0 \in \mathbb{R}$, $(z, \xi) \in C([T_0, \infty); \mathbb{C} \times P_c H^1)$ solve (6.1) on (T_0, ∞) satisfying

$$\|z\|_{L^\infty(T_0, \infty)} + \|\xi\|_{L_t^\infty(T_0, \infty; L_x^2)} \ll 1, \quad \|\xi\|_{L_t^\infty(T_0, \infty; H_x^1)} < \infty. \quad (6.9)$$

Then the following (i) and (ii) are equivalent.

- (i) $\exists \varphi \in H^1(\mathbb{R}^3)$ such that $\|\varphi[z, T_0] - \xi\|_{H_x^1} \rightarrow 0$ as $t \rightarrow \infty$.
(ii) $\|\xi\|_{\text{st}(T_0, \infty)} < \infty$.

Then we say that ξ scatters with z as $t \rightarrow \infty$. Moreover, as $T \rightarrow \infty$,

$$\|\xi[z, T] - \varphi[z, T_0]\|_{\text{Stz}^1(T_0, \infty)} + \|\xi - \xi[z, T]\|_{\text{Stz}^1(T, \infty)} \rightarrow 0, \quad (6.10)$$

and for any $\tilde{z} \in C([T_0, \infty); \mathbb{C})$ satisfying $\|\tilde{z}\|_{L_t^\infty} \ll 1$,

$$\|\xi\|_{[\tilde{z}; T, \infty; \infty]} + \|\xi[\tilde{z}, T]\|_{L_t^2 B_{6,2}^1(T, \infty)} + \|\xi\|_{L_t^2 B_{6,2}^1(T, \infty)} \rightarrow 0, \quad (6.11)$$

uniformly with respect to \tilde{z} . A sufficient condition of scattering is

$$\|\xi[z, T_0]\|_{\text{st}(T_0, \infty)} \left\{ \|z\|_{L^\infty(T_0, \infty)} + \|\xi(T_0)\|_{H_x^{1/2}} \right\}^3 \ll 1. \quad (6.12)$$

The scattering with z as $t \rightarrow -\infty$ is defined in the same way, which has the same property as above.

Proof. Assume (i) and let $\xi_+ := \varphi[z, T_0]$. Then by Lemma 4.3, $\|\xi_+\|_{\text{Stz}^1(T_0, \infty)} < \infty$ and so in particular $\|\xi_+\|_{\text{st}(T, \infty)} \rightarrow 0$ as $T \rightarrow \infty$. By (i) and the Strichartz estimate,

$$\|\xi_+ - \xi[z, T]\|_{\text{st}(T, \infty)} \lesssim \|\xi_+(T) - \xi(T)\|_{H^{1/2}} \rightarrow 0. \quad (6.13)$$

Hence for sufficiently large T , the previous lemma implies $\|\xi\|_{\text{st}(T, \infty)} \lesssim \|\xi[z, T]\|_{\text{st}(T, \infty)}$.

Assume (ii) and let $T > T_0$ so large that we can apply the previous lemma on (T, ∞) to get $\|\xi\|_{\text{Stz}^1(T, \infty)} \lesssim \|\xi[z, T]\|_{\text{Stz}^1(T, \infty)}$. Then for any $T_2 > T_1 > T$, by the Strichartz estimate: Lemma 4.3,

$$\begin{aligned} \|\xi[z, T_2] - \xi[z, T_1]\|_{\text{Stz}^1(T_0, \infty)} &\lesssim \|\tilde{N}(z, \xi)\|_{\text{Stz}^{1*}(T_1, T_2)} \\ &\lesssim \|\xi\|_{\text{Stz}^1(T_1, \infty)} [\|\xi\|_{\text{st}(T_1, \infty)} + \|\xi\|_{\text{st}(T_1, \infty)}^2] \rightarrow 0 \end{aligned} \quad (6.14)$$

as $T_1 \rightarrow \infty$. In particular $\xi[z, T](T_0)$ is Cauchy in H_x^1 as $T \rightarrow \infty$, hence convergent to some $\xi_+ \in H_x^1$. Then

$$\begin{aligned} \|\xi_+[z, T_0](T) - \xi(T)\|_{H_x^1} &\leq \|\xi_+[z, T_0] - \xi[z, T]\|_{\text{Stz}^1(T_0, \infty)} \\ &\lesssim \|\xi_+ - \xi[z, T](T_0)\|_{H_x^1} \rightarrow 0 \quad (T \rightarrow \infty) \end{aligned} \quad (6.15)$$

and so (i). Thus in either case, we have $\|\xi[z, T]\|_{\text{st}(T, \infty)} \rightarrow 0$ as $T \rightarrow \infty$, hence the previous lemma implies

$$\|\xi - \xi[z, T]\|_{\text{Stz}^1(T, \infty)} \lesssim \left\{ \|z\|_{L_t^\infty(T_0, \infty)} + \|\xi\|_{L_t^\infty(T_0, \infty; H_x^1)} \right\} \|\xi[z, T]\|_{\text{st}(T, \infty)} \rightarrow 0 \quad (6.16)$$

as $T \rightarrow \infty$. Hence $\|\xi\|_{L_t^2 B_{6,2}^1(T, \infty)} \rightarrow 0$. Let $\xi_0 := \xi[z, T]$ and $\xi_1 := \xi[\tilde{z}, T]$. Then

$$\xi_1 = \xi_0 + \mathcal{D}(B[\tilde{z}] - B[z])\xi_1[z, T] \quad (6.17)$$

and by the Strichartz estimate: Lemma 4.3,

$$\|\xi_1\|_{\text{Stz}^1(T, \infty)} \leq \|\xi_0\|_{\text{Stz}^1(T, \infty)} + C\|z\|_{L_t^\infty(T_0, \infty)}^2 \|\xi_1\|_{L_t^2 B_{6,2}^1(T, \infty)}. \quad (6.18)$$

After the last term is absorbed by the left, we obtain, as $T \rightarrow \infty$,

$$\|\xi[\tilde{z}, T]\|_{L_t^2 B_{6,2}^1(T, \infty)} \leq 2\|\xi[z, T]\|_{L_t^2 B_{6,2}^1(T, \infty)} \rightarrow 0, \quad (6.19)$$

which is uniform with respect to \tilde{z} . By the previous lemma, we also have

$$\|\xi\|_{[\tilde{z}; T; \infty; \infty]} \ll \|\xi[\tilde{z}, T]\|_{\text{st}(T, \infty)} \sim \|\xi\|_{\text{st}(T, \infty)} \rightarrow 0. \quad (6.20)$$

The sufficiency of (6.12) for scattering is now obvious by Lemma 6.1. \square

The next two lemmas are concerned with difference of two solutions. For the sake of brevity, the following notation is introduced for differences. For any expressions \mathcal{X} and a, b, c, \dots ,

$$\triangleleft \mathcal{X}(a_{\triangleright}, b_{\triangleright}, c_{\triangleright}, \dots) := \mathcal{X}(a_0, b_0, c_0, \dots) - \mathcal{X}(a_1, b_1, c_1, \dots). \quad (6.21)$$

The first lemma of difference estimates treats perturbation of dispersed solutions. It will be used either with $z_0 = z_1$ or on a short interval. We need the non-admissible Strichartz for difference of quadratic terms.

Lemma 6.3. *Let $-\infty < T_0 < T_1 < T_2 \leq \infty$, $z_0 \in C([T_0, T_2]; \mathbb{C})$, $z_1 \in C([T_0, T_1]; \mathbb{C})$, and $\xi_0, \xi_1 \in C([T_0, T_1]; H^1)$ solve*

$$\forall t \in (T_0, T_1), \quad i\dot{\xi}_j + H\xi_j = B[z_j]\xi_j + \tilde{N}(z_j, \xi_j). \quad (6.22)$$

Put $\mathcal{N}_\theta := \|z_j\|_{L_t^\infty(T_0, T_1)} + \|\xi_j\|_{L_t^\infty(T_0, T_1; H_x^\theta)}$ for $\theta \in [0, 1]$, and suppose that for some $0 < \delta \lesssim \tilde{\delta}$ satisfying $\mathcal{N}_0 + \tilde{\delta}\mathcal{N}_{1/2}^3 \ll 1$,

$$\|\xi_0[z_0, T_0]\|_{\text{st}(T_0, T_1)} \leq \tilde{\delta}, \quad \|\triangleleft \xi_{\triangleright}[z_0, T_0]\|_{\text{st}(T_0, T_1)} + \|\triangleleft z_{\triangleright}\|_{L^4(T_0, T_1)} \leq \delta. \quad (6.23)$$

Then we have

$$\|\triangleleft \xi_{\triangleright}\|_{[z; T_0, T_1; T_2]} \lesssim (\mathcal{N}_0 \mathcal{N}_1)^{3/7} \tilde{\delta}^{4/7} \delta^{1/7}. \quad (6.24)$$

Proof. The previous lemma 6.1 applies to both (z_j, ξ_j) , which implies

$$\|\xi_j\|_{\text{st}z^\theta(T_0, T_1)} \lesssim \mathcal{N}_\theta, \quad \|\xi_j\|_{\text{st}(T_0, T_1)} \lesssim \tilde{\delta}. \quad (6.25)$$

Now apply the non-admissible Strichartz estimate, Lemma 4.4 to Duhamel

$$\triangleleft \xi_{\triangleright} = \triangleleft \xi_{\triangleright}[z_0, T_0] + \mathcal{D}\{\triangleleft \tilde{N}(z_{\triangleright}, \xi_{\triangleright}) + \triangleleft B[z_{\triangleright}]\xi_1\}[z_0, T_0]. \quad (6.26)$$

Choose

$$(p_0, q_0) = (4, 24/7), \quad (p_1, q_1) = (2, 24/5), \quad (p_2, q_2) = (4, 24/9) \quad (6.27)$$

so that $\sigma_0 = -1/8$, $\sigma_1 = \sigma_2 = 1/8$ and we can apply the lemma. Then

$$\begin{aligned} \|\triangleleft \xi_{\triangleright} - \triangleleft \xi_{\triangleright}[z_0, T_0]\|_{L_t^4 L_x^{24/7}} &\lesssim \|\triangleleft \tilde{N}(z_{\triangleright}, \xi_{\triangleright}) + \triangleleft B[z_{\triangleright}]\xi_1\|_{L_t^2 L_x^{24/19} + L_t^{4/3} L_x^{24/15}} \\ &\lesssim \{\|\triangleleft \xi_{\triangleright}\|_{L_t^4 L_x^{24/7}} + \|\triangleleft z_{\triangleright}\|_{L_t^4}\} (\|\xi_0\|_{\text{st}} + \|\xi_1\|_{\text{st}}) \{1 + \|\xi_0\|_{\text{st}} + \|\xi_1\|_{\text{st}}\}, \end{aligned} \quad (6.28)$$

where the linear and quadratic (in ξ_0, ξ_1) terms are estimated in $L_t^2 L_x^{24/19}$, while the cubic terms are in $L_t^{4/3} L_x^{24/15}$. The factor 1 comes from the term $\triangleleft B[z]\xi_1$, and it also includes the smallness factor $\mathcal{N}_0 \ll 1$. For the cubic term with $\triangleleft z_{\triangleright}$, we used

$$\|\triangleleft R[z_{\triangleright}]\xi_j\|_{H_x^1} \lesssim |\triangleleft z_{\triangleright}| \|\xi_j\|_{L_x^2} \leq |\triangleleft z_{\triangleright}| \mathcal{N}_0. \quad (6.29)$$

Thus, using the smallness of $\|\xi_j\|_{\text{st}(T_0, T_1)}$ and $\|\triangleleft z_{\triangleright}\|_{L^4(T_0, T_1)}$, we obtain

$$\|\triangleleft \xi_{\triangleright}\|_{L_t^4 L_x^{24/7}(T_0, T_1)} \lesssim \|\triangleleft \xi_{\triangleright}[z_0, T_0]\|_{L_t^4 L_x^{24/7}(T_0, T_1)} + \tilde{\delta}. \quad (6.30)$$

Applying the same estimate to the Duhamel formula

$$\triangleleft \xi_{\triangleright}[z_0, T_1]_{>} = \triangleleft \xi_{\triangleright}[z_0, T_0] + \mathcal{D}1_{T_0 < t < T_1} \{\triangleleft \tilde{N}(z_{\triangleright}, \xi_{\triangleright}) + \triangleleft B[z_{\triangleright}]\xi_1\}[z_0, T_0], \quad (6.31)$$

we also obtain

$$\|\langle \xi_{\triangleright} [z_0, T_1] \rangle - \langle \xi_{\triangleright} [z_0, T_0] \rangle\|_{L_t^4 L_x^{24/7}(T_0, T_2)} \lesssim \tilde{\delta} \|\langle \xi_{\triangleright} [z_0, T_0] \rangle\|_{L_t^4 L_x^{24/7}(T_0, T_1)} + \tilde{\delta} \delta, \quad (6.32)$$

and this norm is related to $\mathfrak{st} = L_t^4 L_x^6$ by interpolation and Sobolev as

$$\begin{aligned} \|f\|_{\mathfrak{st}} &\lesssim \|f\|_{L_t^4 L_x^{24/7}}^{4/7} \|f\|_{L^4 B_{\infty,2}^0}^{3/7} \lesssim \|f\|_{L_t^4 L_x^{24/7}}^{4/7} \|f\|_{\text{Stz}^1}^{3/7}, \\ \|f\|_{L_t^4 L_x^{24/7}} &\leq \|f\|_{\mathfrak{st}}^{1/4} \|f\|_{L_t^4 L_x^3}^{3/4} \lesssim \|f\|_{\mathfrak{st}}^{1/4} \|f\|_{\text{Stz}^0}^{3/4}. \end{aligned} \quad (6.33)$$

Injecting these to the above and using the Strichartz bound, we obtain

$$\begin{aligned} \|\langle \xi_{\triangleright} [z_0, T_1] \rangle - \langle \xi_{\triangleright} [z_0, T_0] \rangle\|_{\mathfrak{st}(T_0, T_2)} &\lesssim [\tilde{\delta} \delta^{1/4} \mathcal{N}_0^{3/4}]^{4/7} \mathcal{N}_1^{3/7} \\ &= (\mathcal{N}_0 \mathcal{N}_1)^{3/7} \tilde{\delta}^{4/7} \delta^{1/7}, \end{aligned} \quad (6.34)$$

as desired. \square

The second lemma of difference estimates treats perturbation of large solutions with finite Strichartz. It will be used only on a bounded interval of time.

Lemma 6.4. *Let $-\infty < T_0 < T_1 < T_2 \leq \infty$, $z_0 \in C([T_0, T_2]; \mathbb{C})$, $z_1 \in C([T_0, T_1]; \mathbb{C})$, and $\xi_0, \xi_1 \in C([T_0, T_1]; H^1)$ solve*

$$\forall t \in (T_0, T_1), \quad i\dot{\xi}_j + H\xi_j = B[z_j]\xi_j + \tilde{N}(z_j, \xi_j). \quad (6.35)$$

Put for $\theta \in [0, 1]$

$$\mathcal{N}_{\theta} := \|z_j\|_{L_t^{\infty}(T_0, T_1)} + \|\xi_j\|_{L_t^{\infty}(T_0, T_1; H_x^{\theta})}, \quad \mathcal{N}_2 := \|\xi_0\|_{\mathfrak{st}(T_0, T_1)}, \quad (6.36)$$

and assume $\mathcal{N}_0 \ll 1$. For any $\varepsilon > 0$, there is $\delta_*(\mathcal{N}_1, \mathcal{N}_2, \varepsilon) > 0$, continuous and decreasing for each \mathcal{N}_j , such that if

$$\|\langle \xi_{\triangleright} [z_0, T_0] \rangle\|_{\mathfrak{st}(T_0, T_1)} + \|\langle z_{\triangleright} \rangle\|_{L^4(T_0, T_1)} \leq \delta_* \quad (6.37)$$

then we have

$$\|\langle \xi_{\triangleright} \rangle\|_{[z_0; T_0, T_1; T_2]} \leq \varepsilon. \quad (6.38)$$

Proof. For any $N \in \mathbb{N}$, (T_0, T_1) is decomposed into subintervals I_0, \dots, I_N such that

$$\forall j = 0, \dots, N, \quad \|\xi_0\|_{\mathfrak{st}(I_j)} \leq 2N^{-1/4} \mathcal{N}_2 =: \tilde{\delta}. \quad (6.39)$$

Let $I_j =: (S_j, S_{j+1})$. If $\tilde{\delta} \mathcal{N}_{1/2}^3 \ll 1$, then we can apply Lemma 6.1 starting from S_j , and we obtain

$$\|\xi_0[z_0, S_j]\|_{\mathfrak{st}(I_j)} \sim \|\xi_0\|_{\mathfrak{st}(I_j)} \leq \tilde{\delta}. \quad (6.40)$$

Suppose that for some $\delta_0 > 0$,

$$\|\langle \xi_{\triangleright} [z_0, S_0] \rangle\|_{\mathfrak{st}(S_0, T_1)} + \|\langle z_{\triangleright} \rangle\|_{L^4(S_0, T_1)} \leq \delta_0 \leq \tilde{\delta}, \quad \tilde{\delta} \mathcal{N}_{1/2}^3 \ll 1. \quad (6.41)$$

Then $\|\xi_1[z_0, T_0]\|_{\mathfrak{st}(S_0, S_1)} \lesssim \tilde{\delta}$ and we can apply Lemma 6.3 on I_0 . Then

$$\|\langle \xi_{\triangleright} \rangle\|_{[z_0; S_0, S_1; T_2]} \leq C(\mathcal{N}_0 \mathcal{N}_1)^{3/7} \tilde{\delta}^{4/7} \delta_0^{1/7}. \quad (6.42)$$

Let $\delta_1 := \delta_0 + C(\mathcal{N}_0 \mathcal{N}_1)^{3/7} \tilde{\delta}^{4/7} \delta_0^{1/7}$, then

$$\|\langle \xi_{\triangleright} [z_0, S_1] \rangle\|_{\mathfrak{st}(S_1, T_1)} + \|\langle z_{\triangleright} \rangle\|_{L^4(S_1, T_1)} \leq \delta_1. \quad (6.43)$$

Hence if $\delta_1 \leq \tilde{\delta}$ then we can repeat the same thing on I_1 . Define the sequence δ_j for $j = 0, \dots, N$ inductively from δ_0 by

$$\delta_{j+1} := \delta_j + C(\mathcal{N}_0 \mathcal{N}_1)^{3/7} \tilde{\delta}^{4/7} \delta_j^{1/7}. \quad (6.44)$$

Given \mathcal{N}_1 and \mathcal{N}_2 , we can determine $\tilde{\delta}$ and N such that

$$2N^{-1/4} \mathcal{N}_2 \leq \tilde{\delta} \ll \mathcal{N}_1^{-3} \leq \mathcal{N}_{1/2}^{-3}. \quad (6.45)$$

Then there is $\delta_* = \delta_*(\mathcal{N}_1, \mathcal{N}_2, \varepsilon) > 0$ such that

$$\delta_0 \leq \delta_* \implies \delta_{N+1} < \min(\tilde{\delta}, \varepsilon). \quad (6.46)$$

Then for $\delta_0 \leq \delta_*$, we can iterate the above estimate for all j to get

$$\begin{aligned} \|\langle \xi \rangle\|_{[z_0; T_0, T_1; T_2]} &\leq \sum_{j=0}^N \|\langle \xi \rangle\|_{[z_0; S_j, S_{j+1}; T_2]} \\ &\leq \sum_{j=0}^N C(\mathcal{N}_0 \mathcal{N}_1)^{3/7} \tilde{\delta}^{4/7} \delta_j^{1/7} = \delta_{N+1} - \delta_0 < \varepsilon, \end{aligned} \quad (6.47)$$

where we used the subadditivity for consecutive intervals by Lemma 4.2. \square

7. NONLINEAR PROFILE DECOMPOSITION

We are now ready to develop a profile decomposition for NLS (1.7) in the (z, ξ) coordinate, i.e. the equation (4.10). Let $I \in \mathcal{I}^{\mathbb{N}}$ and $u \in C(I; H^1[\mu_p])$ be a sequence of solutions of (1.7) in the mass region of the (z, ξ) -coordinate. Put

$$I_n =: (\underline{T}_n, \overline{T}_n) \quad (7.1)$$

for each $n \in \mathbb{N}$. We can uniquely write $u = \Phi[z] + R[z]\xi$, where $(z, \xi) \in C(I; \mathbb{C} \times H^1)$ is a sequence of solutions for (4.10). Suppose

$$\mathcal{N}_1 := \sup \{ \|z\|_{L_t^\infty(I)} + \|u\|_{L_t^\infty(I; H^1)} \} < \infty, \quad (7.2)$$

where the z part is uniformly bounded by $\sqrt{\mu_p}$, so that we can omit it. Then (4.10) implies that $z \in \text{SBC}(I)$, where the smallness comes from $\mathbb{M}(u_n) < \mu_p$, but we have not used the second condition of (7.2). By the L^2 conservation

$$\mathcal{N}_0 := \sup \{ \|z\|_{L_t^\infty(I)} + \|u\|_{L_t^\infty(I; L^2)} \} \lesssim \sqrt{\mu_p} \ll 1. \quad (7.3)$$

Similarly for $\theta \in [0, 1]$, put

$$\mathcal{N}_\theta := \sup \{ \|z\|_{L_t^\infty(I)} + \|u\|_{L_t^\infty(I; H^\theta)} \} \leq \mathcal{N}_0^{1-\theta} \mathcal{N}_1^\theta. \quad (7.4)$$

For any $s \in I$, we can apply the linearized profile decomposition: Lemma 5.3 to the sequence $\xi(s)$. Passing to a subsequence, we have for each $J < J^*$,

$$\nu := \xi[z, s] = \lambda^{[0 \dots J]} + \gamma^J, \quad \lambda^{[0 \dots J]} := \sum_{0 \leq j < J} \lambda^j, \quad \lambda^j := \nu[z, s^j]_{(\infty)}. \quad (7.5)$$

Extracting a subsequence if necessary, we may assume

$$s^j - s \rightarrow \sigma^j \infty, \quad \exists \sigma^j \in \{+, -\} \quad (7.6)$$

for each $0 < j < J^*$, and, since (4.10) implies that (z, ξ) is weakly equicontinuous,

$$(z, \xi)(s^j + t) \rightarrow \exists(z_\infty^j, \xi_\infty^j)(t) \text{ in } \mathbb{C} \times \text{w-}H^1, \quad (7.7)$$

locally uniformly on $I - s^j$. Put

$$I_\infty^j := \liminf(I - s^j), \quad I^j := I_\infty^j + s^j. \quad (7.8)$$

(7.6) implies $I_\infty^j \supset \{-\infty < \sigma^j t \leq 0\}$ for $j > 0$. For $j = 0$, we have $I_\infty^0 \supset [0, \infty)$ if $\overline{T} - s \rightarrow \infty$ and $I_\infty^0 \supset (-\infty, 0]$ if $\underline{T} - s \rightarrow -\infty$. Note that if $|I_n|$ is bounded then the decomposition is trivial, i.e. $J^* = 1$. In either case, $I^j \ni s^j, s^0 = s$. By the property of SBC, we can extend z_∞^j to $\text{SBC}(\mathbb{R})$. By the subcritical nature of NLS, it is easy to see that the weak limit $(z_\infty^j, \xi_\infty^j)$ is a solution of (4.10) in $C(I_\infty^j; \mathbb{C} \times P_c H^1)$.

The nonlinear profile $\Lambda^j \in C(I^j; P_c H^1)$ is defined by

$$\Lambda^j(t) := \xi_\infty^j(t - s^j). \quad (7.9)$$

Also put

$$\begin{aligned} \varphi_\infty^j &:= \lim \nu(s^j) = \lambda^j(s^j) \in P_c H^1, \quad \lambda_\infty^j := \varphi_\infty^j[z_\infty^j, 0] \in C(\mathbb{R}; P_c H^1), \\ z_j(t) &:= z(s^j + t), \quad z_\infty^j(t) := z_\infty^j(t - s^j), \quad \lambda_\infty^j(t) := \lambda_\infty^j(t - s^j), \end{aligned} \quad (7.10)$$

such that $z = z_\infty^j + o(1)$ and $z_j = z_\infty^j + o(1)$ in $C(\mathbb{R})$, and using Lemma 5.1, (5.13),

$$\lambda^j = \varphi_\infty^j[z, s^j] = \lambda_\infty^j + o(1) \text{ in } \text{Stz}^1(I), \quad (7.11)$$

while (z_∞^j, Λ^j) is a sequence of solutions of (4.10) on I^j .

The nonlinear remainder $\Gamma^J \in C(I; H^1)$ is defined for each J by the same sequence of equations as ξ , with the same initial data as γ^J :

$$i\dot{\Gamma}^J + H\Gamma^J = B[z]\Gamma^J + \tilde{N}(z, \Gamma^J), \quad \Gamma^J(s) = \gamma^J(s). \quad (7.12)$$

Since $\lim_{J \rightarrow J^*} \limsup \|\gamma^J\|_{\text{st}(I)} = 0$, Lemma 6.1 ensures the unique existence of Γ_n^J for large J and large n , satisfying

$$\|\Gamma_n^J\|_{\text{Stz}^\theta(I_n)} \lesssim \mathcal{N}\theta, \quad \|\Gamma_n^J - \gamma_n^J\|_{\text{st}(I_n)} + \|\Gamma_n^J\|_{[z_n; s_n, \overline{T}_n; \overline{T}_n]} \ll \|\gamma_n^J\|_{\text{st}(I_n)}. \quad (7.13)$$

We have $\gamma^J(s^j) \rightarrow 0$ for each $j < J$. Moreover, for large J and for $0 \leq j < J$,

$$\Gamma^J(s^j) \rightarrow 0, \quad 0 < \forall \tau < \infty, \quad \|\Gamma^J\|_{\text{st}(|t-s^j| < \tau)} + \|\gamma^J\|_{\text{st}(|t-s^j| < \tau)} \rightarrow 0. \quad (7.14)$$

Proof. Let X^j be a sequence of weighted norms on I defined by

$$\|f\|_{X^j} := \sup_{t \in I} \langle t - s^j \rangle^{-\delta} \|f(t)\|_{(L^4 + L^p)_x(\mathbb{R}^3)}, \quad (7.15)$$

where $\delta > 0$ is fixed such that $0 < \delta < 1/2 - 3/p$. By Lemma 5.1, (5.14), we have $\|\gamma^j(t)\|_{L_x^4} \rightarrow 0$ locally uniformly around $t = s^j$, which implies $\|\gamma^j\|_{X^j} \rightarrow 0$, thanks to the decaying weight. Suppose that $\sigma^j = +$, namely $s^j - s^0 \rightarrow \infty$. Put $F^J := B[z]\Gamma^J + \tilde{N}(z, \Gamma^J)$. Then by the L^p decay estimate on $e^{itH}P_c$, we have at any $t \in I_n$ satisfying $t > s_n$,

$$\begin{aligned} \|\Gamma_n^J(t) - \gamma_n^J(t)\|_{L_x^4 + L_x^p} &\lesssim \int_{s_n^0}^t f(t-t') \|F_n^J(t')\|_{L_x^{4/3} \cap L_x^{p'}} dt', \\ f(t) &:= \min(|t|^{-3(1/2-1/p)}, |t|^{-3/4}), \end{aligned} \quad (7.16)$$

and, by Hölder and Sobolev,

$$\begin{aligned} \|F_n^J\|_{L_x^{4/3} \cap L_x^p} &\lesssim \|\Gamma_n^J\|_{L_x^4 + L_x^p} (|z_n|^2 + \|\Gamma_n^J\|_{L_x^4 \cap L_x^2}^2) \\ &\lesssim \{\mathcal{N}_0^2 + \|\Gamma_n^J\|_{L_t^\infty L_x^4(I_n)}^2\} \|\Gamma_n^J\|_{L_x^4 + L_x^p}. \end{aligned} \quad (7.17)$$

By Lemma 6.1, we have

$$\|\Gamma_n^J\|_{L_t^\infty L_x^4(I_n)} \leq \|\gamma_n^J\|_{L_t^\infty L_x^4(I_n)} + C\mathcal{N}_{3/4} \|\gamma_n^J\|_{\text{st}(I_n)} (\mathcal{N}_0 + \|\gamma_n^J\|_{\text{st}(I_n)}) \quad (7.18)$$

Taking J and n larger if necessary, we may assume that the right hand side is bounded by $\mathcal{N}_0 \ll 1$. Inserting this to the above estimate and then to (7.16) yields

$$\begin{aligned} \|\Gamma_n^J(t) - \gamma_n^J(t)\|_{L_x^4 + L_x^p} &\lesssim \mathcal{N}_0^2 \|\Gamma_n^J\|_{X_n^j} \int_{\mathbb{R}} f(t-t') \langle t' - s_n^j \rangle^\delta dt' \\ &\lesssim \mathcal{N}_0^2 \|\Gamma_n^J\|_{X_n^j} \langle t - s_n^j \rangle^\delta, \end{aligned} \quad (7.19)$$

where $-3(1/2 - 1/p) + \delta < -1$ ensures the integrability for $|t'| \rightarrow \mathbb{R}$. The estimate in the cases $t < s_n$ and $\sigma^j = -$ is the same. Thus we obtain

$$\lim \|\Gamma^J\|_{X^j} \lesssim \lim \|\gamma^J\|_{X^j} = 0, \quad (7.20)$$

for large J . By the uniform H^1 bound, this is equivalent to $\Gamma^J(t) \rightarrow 0$ weakly in H^1 , locally uniformly around s^j . Interpolation with (7.13) yields the other part. \square

Henceforth, we concentrate on the estimate on the time interval $t > s_n$, assuming

$$\overline{T}_n - s_n \rightarrow \infty, \quad (n \rightarrow \infty) \quad (7.21)$$

since otherwise uniform Strichartz bound for ξ_n on $t > s_n$ is trivial. The restriction to $t > s_n$ allows us to ignore the profiles with $s^j - s^0 \rightarrow -\infty$.

Fix a finite $J \leq J^*$, so large that (7.14) holds. After neglecting those profiles with $s^j - s^0 \rightarrow -\infty$, and reordering the profiles¹, we may assume for $0 < j < J$

$$s^j - s^{j-1} \rightarrow \infty. \quad (7.22)$$

Since J is now fixed, we can no longer gain a small factor by sending $J \rightarrow \infty$. Instead another parameter $0 < \tau \rightarrow \infty$ is introduced, decomposing the time intervals

$$(s, \overline{T}) = \bigcup_{0 \leq j < J} (s_-^j, s_+^j) \cup (s_+^j, s_-^{j+1}), \quad (7.23)$$

where $s_\pm^j \in \mathbb{R}^{\mathbb{N}}$ are defined for each j by

$$\begin{aligned} s_-^0 &:= s^0, \quad s_-^j := s^j - \tau \quad (0 < j < J), \quad s_-^J := \overline{T}, \\ s_+^j &:= s^j + \tau \quad (0 \leq j < J). \end{aligned} \quad (7.24)$$

Henceforth, o_τ denotes any sequence of real numbers satisfying

$$X(\tau) = o_\tau \iff \lim_{\tau \rightarrow \infty} \limsup_{n \rightarrow \infty} X_n(\tau) = 0. \quad (7.25)$$

¹This reordering can not be performed before fixing $J < \infty$, since more and more linear profiles may well appear between the previous profiles as $J \rightarrow \infty$, which is a typical dispersive behavior of the remainder γ^J .

By the uniform integrability (7.11) of the linearized profiles, and their separation $s^j - s^{j-1} \rightarrow \infty$, we have for $0 \leq j < J$

$$\begin{aligned} & \|\lambda^{[j+1\dots J]}\|_{\text{st}(s^0, s^j_+)} + \|\lambda^{[0\dots j]}\|_{\text{st}(s^j_-, \bar{T})} = o(1), \\ & \|\lambda^j\|_{\text{st}(s^0, s^j_-)} + \|\lambda^j\|_{\text{st}(s^j_+, \bar{T})} = o_\tau. \end{aligned} \quad (7.26)$$

The following is the main property of the nonlinear profile decomposition.

Lemma 7.1. *In the above setting, suppose that ξ_∞^j is scattering with z_∞^j as $t \rightarrow \infty$ for $0 \leq j < J$. Then we have for $0 \leq j < J$,*

- (i) $\|(\xi - \Gamma^J)[z, s^j_-] - \lambda^{[j\dots J]}\|_{\text{st}(s^j_-, \bar{T})} + \|\Lambda^j[z, s^j_-]_> - \lambda^j\|_{\text{Stz}^1(I)} = o_\tau.$
- (ii) $\|\xi - \Lambda^j\|_{[z; s^j_-, s^j_+; \bar{T}]} + \|\Lambda^j\|_{[z; s^0, s^j_+; \bar{T}]} + \|\Lambda^j\|_{[z; s^j_+, \bar{T}; \bar{T}]} = o_\tau.$
- (iii) $\|\xi - \Gamma^J\|_{[z; s^j_+, s^{j+1}_+; \bar{T}]} + \|\Gamma^J\|_{[z; s^j_+, s^j_+; \bar{T}]} = o_\tau.$

For $0 < j < J$, $\|\xi_\infty^j - \lambda_\infty^j\|_{\text{Stz}^1(-\infty, -\tau)} \rightarrow 0$ as $\tau \rightarrow \infty$ (scattering as $t \rightarrow -\infty$).

Proof. The second term of (ii) is bounded by

$$\|\Lambda^j\|_{[z; s^0, s^j_+; \bar{T}]} \leq \|\Lambda^j - \lambda^j\|_{\text{Stz}^1(s^0, s^j_-)} \leq \text{(i)}, \quad (7.27)$$

and the third term is bounded using Lemma 6.2, (6.11) with the scattering of ξ_∞^j

$$\|\Lambda^j\|_{[z; s^j_+, \bar{T}; \bar{T}]} \leq \|\xi_\infty^j\|_{[z; \tau, \infty; \infty]} = o_\tau. \quad (7.28)$$

For the second term of (iii), using (4.29), Lemma 6.1 and (7.14), we obtain

$$\|\Gamma^J[z, s^j_-] - \Gamma^J[z, s^j_+]\|_{\text{st}(s^j_+, \bar{T})} \leq \|\Gamma^J\|_{[z; s^j_-, s^j_+; \bar{T}]} \ll \|\Gamma^J\|_{\text{st}(s^j_-, s^j_+)} = o(1). \quad (7.29)$$

The remaining estimates are proved by induction on j . For $j = 0$, (i) = 0 by the definition and $s_-^0 = s^0$. Assume (i) for some j as an induction hypothesis. Combining (i) with (7.26) and (7.14) yields

$$\|(\xi - \Lambda^j)[z, s^j_-]\|_{\text{st}(s^j_-, s^j_+)} + \|z - z_\infty^j\|_{L_t^4(s^j_-, s^j_+)} = o_\tau. \quad (7.30)$$

This allows us to use Lemma 6.4 leading to the first term of (ii):

$$\|\xi - \Lambda^j\|_{[z; s^j_-, s^j_+; \bar{T}]} = o_\tau, \quad (7.31)$$

By the scattering of ξ_∞^j for $t \rightarrow \infty$, Lemma 6.2 implies

$$\|\Lambda^j[z, s^j_+]\|_{\text{st}(s^j_+, \bar{T})} = \|\xi_\infty^j[z(t + s^j), \tau]\|_{\text{st}(\tau, \bar{T} - s^j)} = o_\tau. \quad (7.32)$$

Combining it with (i), (7.29) and (7.31), using (4.29), we obtain

$$\begin{aligned} & \|(\xi - \Gamma^J)[z, s^j_+] - \lambda^{[j+1\dots J]}\|_{\text{st}(s^j_+, \bar{T})} \\ & \leq \|(\xi - \Lambda^j)[z, s^j_+] - (\xi - \Lambda^j)[z, s^j_-]\|_{\text{st}(s^j_+, \bar{T})} + \|\xi^j[z, s^j_+]\|_{\text{st}(s^j_+, \bar{T})} + o_\tau \\ & \leq \|\xi - \Lambda^j\|_{[z; s^j_-, s^j_+; \bar{T}]} + o_\tau \leq o_\tau. \end{aligned} \quad (7.33)$$

Restricting it and using (7.26), we obtain

$$\|(\xi - \Gamma^J)[z, s^j_+]\|_{\text{st}(s^j_+, s^{j+1}_+)} \leq o_\tau. \quad (7.34)$$

This and the smallness of Γ_n^J in (7.14) allow us to apply Lemma 6.3 to the difference of ξ_n and Γ_n^J for large τ and large n , with the same soliton part z_n . Then the above decay of the linearized solutions leads to the estimate on the first term of (iii):

$$\|\xi - \Gamma^J\|_{[z; s_+^j, s_-^{j+1}; \bar{T}]} = o_\tau. \quad (7.35)$$

If $k := j + 1 < J$, then combining the above with (7.33), using (4.29), we obtain

$$\begin{aligned} & \|(\xi - \Gamma^J)[z, s_-^k] - \lambda^{[k \dots J]}\|_{\mathbf{st}(s_-^k, \bar{T})} \\ & \leq \|(\xi - \Gamma^J)[z, s_+^j] - \lambda^{[k \dots J]}\|_{\mathbf{st}(s_-^k, \bar{T})} + \|\xi - \Gamma^J\|_{[z; s_+^j, s_-^k; \bar{T}]} \leq o_\tau, \end{aligned} \quad (7.36)$$

which is the first term of (i) for k . Using this with (7.26) and (7.14), we obtain

$$\|\xi[z, s_-^k] - \lambda^k\|_{\mathbf{st}(s_-^k, s^k)} = o_\tau. \quad (7.37)$$

Since $\xi(s_-^k) - \lambda^k(s_-^k) \rightarrow \xi_\infty^k(-\tau) - \lambda_\infty^k(-\tau)$, by Lemma 5.1, (5.14), we obtain

$$\begin{aligned} o_\tau & = \|(\xi_\infty^k(-\tau) - \lambda_\infty^k(-\tau))[z, s^k - \tau]\|_{\mathbf{st}(s^k - \tau, s^k)} \\ & = \|(\xi_\infty^k - \lambda_\infty^k)[z(t + s^k), -\tau]\|_{\mathbf{st}(-\tau, 0)}. \end{aligned} \quad (7.38)$$

Taking the limit and using Lemma 5.1, (5.13) with $z(t + s^k) \rightarrow z_\infty^k(t)$,

$$\lim_{\tau \rightarrow \infty} \|\xi_\infty^k[z_\infty^k, -\tau] - \lambda_\infty^k\|_{\mathbf{st}(-\tau, 0)} = 0. \quad (7.39)$$

Since $\lambda_\infty^k \in \mathbf{st}(-\infty, 0)$ by Lemma 4.3, there is $\tau_* > 0$ such that

$$\|\lambda_\infty^k\|_{\mathbf{st}(-\infty, -\tau_*)} + \sup_{\tau > \tau_*} \|\xi_\infty^k[z_\infty^k, -\tau] - \lambda_\infty^k\|_{\mathbf{st}(-\tau, 0)} \ll \mathcal{N}_{1/2}^{-3}. \quad (7.40)$$

For $\tau > \tau_*$, we can apply Lemma 6.1 to ξ_∞^k from $t = -\tau$, thereby obtain

$$\|\xi_\infty^k\|_{\mathbf{st}(-\tau, -\tau_*)} \leq 2\|\xi_\infty^k[z_\infty^k, -\tau]\|_{\mathbf{st}(-\tau, -\tau_*)} \ll \mathcal{N}_{1/2}^{-3}. \quad (7.41)$$

Sending $\tau \rightarrow \infty$ implies $\|\xi_\infty^k\|_{\mathbf{st}(-\infty, -\tau_*)} < \infty$, so by Lemma 6.2, ξ_∞^k scatters with z_∞^k as $t \rightarrow -\infty$. Hence there exists $\varphi_-^k \in H^1$ such that

$$\lim_{\tau \rightarrow \infty} \|\xi_\infty^k[z_\infty^k, -\tau] - \varphi_-^k[z_\infty^k, 0]\|_{\text{Stz}^1(-\infty, 0)} = 0. \quad (7.42)$$

Adding this and (7.39) yields

$$\|(\varphi_-^k - \lambda_\infty^k)[z_\infty^k, 0]\|_{\mathbf{st}(-\infty, 0)} = 0, \quad (7.43)$$

which implies $\varphi_-^k = \lambda_\infty^k(0)$, hence $\|\xi_\infty^k - \lambda_\infty^k\|_{\text{Stz}^1(-\infty, -\tau)} \rightarrow 0$ as $\tau \rightarrow \infty$. Since $\xi_\infty^k = \Lambda^k(t + s^k)$ and $\lambda_\infty^k = \lambda^k(t + s^k) + o(1)$ in Stz^1 , we obtain

$$\|\Lambda^k[z, s_-^k]_{>} - \lambda^k\|_{\text{Stz}^1(I)} \lesssim \|\Lambda^k - \lambda^k\|_{\text{Stz}^1(s^0, s_-^k)} = o_\tau, \quad (7.44)$$

which is the second term of (i) for $k = j + 1$, hence the induction is complete. \square

Suppose that ξ_∞^j is scattering with z_∞^j as $t \rightarrow \infty$ for $0 \leq j < J$ as in the above lemma. Since the profiles Λ^j and the remainder Γ^J are vanishing o_τ in each other intervals, we obtain, using the subadditivity: Lemma 4.2,

$$\begin{aligned} & \|\xi - \Lambda^{[0 \dots J]} - \Gamma^J\|_{[z; s, \bar{T}; \bar{T}]} \\ & \leq \sum_{0 \leq j < J} \|\xi - \Lambda^j\|_{[z; s_-^j, s_+^j; \bar{T}]} + \sum_{0 \leq j < J} \|\xi - \Gamma^J\|_{[z; s_+^j, s_-^{j+1}; \bar{T}]} + o_\tau \leq o_\tau. \end{aligned} \quad (7.45)$$

Since the left hand side is independent of τ , we deduce that

$$\|\xi - \Lambda^{[0\dots J]} - \Gamma^J\|_{[z;s,\overline{T};\overline{T}]} = o(1). \quad (7.46)$$

Thus we have obtained a sequence of nonlinear approximation:

$$\tilde{\xi}^J := \Lambda^{[0\dots J]} + \Gamma^J. \quad (7.47)$$

Since the initial data $\tilde{\xi}^J(s) = \xi_\infty^0(0) + \Lambda^{[1\dots J]}(s) + \gamma^J(s) = \xi(s) + o(1)$ in H^1 , (7.45) with (4.29) implies

$$\|\xi - \tilde{\xi}^J\|_{\text{st}(s,\overline{T})} \leq \|\xi - \tilde{\xi}^J\|_{[z;s,\overline{T};\overline{T}]} + o(1) = o(1). \quad (7.48)$$

Hence we obtain

$$\begin{aligned} \|\xi\|_{\text{st}(s,\overline{T})} &\leq \|\tilde{\xi}^J\|_{\text{st}(s,\overline{T})} + o(1) \\ &\leq \|\xi_\infty^0\|_{\text{st}(0,\infty)} + \sum_{1 \leq j < J} \|\xi_\infty^j\|_{\text{st}(\mathbb{R})} + \|\Gamma^J\|_{\text{st}(s,\overline{T})} + o(1) \\ &\leq \|\xi_\infty^0\|_{\text{st}(0,\infty)} + \sum_{1 \leq j < J} \|\xi_\infty^j\|_{\text{st}(\mathbb{R})} + 2\|\gamma^J\|_{\text{st}(s,\overline{T})} + o(1) \\ &\leq \|\xi_\infty^0\|_{\text{st}(0,\infty)} + \sum_{1 \leq j < J} \|\xi_\infty^j\|_{\text{st}(\mathbb{R})} + 1, \end{aligned} \quad (7.49)$$

where the last number 1 can be made arbitrarily small, taking J bigger if necessary.

The same argument works on the other time direction (\underline{T}, s) , under the scattering assumption of ξ_j^∞ with z_j^∞ as $t \rightarrow -\infty$ for $\sigma^j = -$ and $j = 0$. In order to consider the whole interval $(\underline{T}, \overline{T})$, we should assume the scattering of ξ_j^∞ as $t \rightarrow \sigma^j \infty$ for $\sigma^j = \pm$, and of ξ_0^∞ as $t \rightarrow \pm\infty$. Then we obtain

$$\|\xi\|_{\text{st}(I)} \leq \sum_{0 \leq j < J} \|\xi_\infty^j\|_{\text{st}(\mathbb{R})} + 1, \quad (7.50)$$

if J is close enough to J^* .

8. SCATTERING BELOW THE EXCITED ENERGY

We are now ready to prove the scattering to the ground states. For each $\mu > 0$ and $A \in \mathbb{R}$, let $\text{GS}(\mu, A)$ be the totality of global solution u of (1.7) satisfying

$$\mathbb{M}(u) \leq \mu, \quad \mathbb{E}(u) \leq A. \quad (8.1)$$

Let

$$\begin{aligned} ST(\mu, A) &:= \sup\{\|\xi\|_{\text{st}(0,\infty)} \mid \Phi[z] + R[z]\xi \in \text{GS}(\mu, A)\}, \\ \mathcal{X} &:= \{(\mu, A) \mid ST(\mu, A) < \infty\}. \end{aligned} \quad (8.2)$$

Introduce the following partial orders in \mathbb{R}^2

$$\begin{aligned} (\mu_1, A_1) \leq (\mu_2, A_2) &\iff \mu_1 \leq \mu_2 \text{ and } A_1 \leq A_2, \\ (\mu_1, A_1) \ll (\mu_2, A_2) &\iff \mu_1 < \mu_2 \text{ and } A_1 < A_2. \end{aligned} \quad (8.3)$$

The definition of \mathcal{X} implies that for any $(\mu_j, A_j) \in (0, \infty) \times \mathbb{R}$,

$$(\mu_1, A_1) \leq (\mu_2, A_2) \text{ and } (\mu_2, A_2) \in \mathcal{X} \implies (\mu_1, A_1) \in \mathcal{X}. \quad (8.4)$$

The goal of this section is to prove that for $0 < \mu \ll 1$ and $A \in \mathbb{R}$,

$$(\mu, A) \in \mathcal{X} \iff A < \mathcal{E}_1(\mu). \quad (8.5)$$

\implies is trivial in the defocusing case, obvious by the excited states \mathcal{S}_1 in the focusing case. So the question is the \impliedby part.

For small H^1 data, we have the scattering to \mathcal{S}_0 by [9], together with a uniform bound on the Strichartz norms of ξ in terms of $\|u(0)\|_{H_x^1}$. In fact, Lemma 6.2 implies that $H_x^{1/2}$ smallness is enough. In particular, using Lemma 3.1 and interpolation, we deduce that $(\mu, A) \in \mathcal{X}$ for sufficiently small A for each fixed μ , and for sufficiently small μ for each fixed A . Hence \mathcal{X} contains a neighborhood of both $\{\mu = 0\}$ and $\{A = 0\}$.

Suppose that there exists $(\mu_0, A_0) \in (0, \infty)^2 \setminus \mathcal{X}$ satisfying $A_0 < \mathcal{E}_1(\mu_0)$ and $\mu_0 \ll 1$. Put

$$E_* := \sup\{A < A_0 \mid (\mu_0, A) \in \mathcal{X}\}, \quad M_* := \sup\{\mu < \mu_0 \mid (\mu, E_*) \in \mathcal{X}\}. \quad (8.6)$$

Then

$$0 < M_* \leq \mu_0, \quad 0 < E_* \leq A_0 < \mathcal{E}_1(\mu_0) \leq \mathcal{E}_1(M_*), \quad (8.7)$$

and (M_*, E_*) is minimal on $\partial\mathcal{X}$ in the sense that

$$(\mu_1, A_1) < (M_*, E_*) \ll (\mu_2, A_2) \implies (\mu_1, A_1) \in \mathcal{X}, \quad (\mu_2, A_2) \notin \mathcal{X}. \quad (8.8)$$

In particular, there is a sequence $(\mathbb{R}^2)^\mathbb{N} \ni (M, E) \rightarrow (M_*, E_*)$ and a sequence of solutions $u = \Phi[z] + R[z]\xi \in \text{GS}(M, E)$ such that

$$M \leq \mu_0 + o(1), \quad E < \mathcal{E}_1(M), \quad \|\xi\|_{\text{st}(0, \infty)} \rightarrow \infty. \quad (8.9)$$

See (5.2) for the notation of sequences without index. The mass-energy constraint together with Lemma 3.1 implies that $u(0)$ is bounded in H_x^1 , so is $\xi(0)$, while $|z(t)| \lesssim \mu_0 \ll 1$. Hence we can apply the linearized profile decomposition: Lemma 5.3 on $I = [0, \infty)^\mathbb{N}$, which yields

$$\nu := \xi[z, 0] = \sum_{0 \leq j < J} \lambda^j + \gamma^J, \quad \lambda^j = \nu[z, s^j]_{(\infty)}, \quad (8.10)$$

for each $J < J^*$. $I = [0, \infty)^\mathbb{N}$ implies $s^j \rightarrow \infty$ for all $0 < j < J^*$. Let $\Lambda^j = \xi_\infty^j(t - s^j)$ and Γ^J be the associated nonlinear profiles and the nonlinear remainder, namely

$$(z(t + s^j), \xi(t + s^j)) \rightharpoonup (z_\infty^j, \xi_\infty^j) \quad (8.11)$$

and $(z_\infty^j, \xi_\infty^j)$ is a solution of (4.10), Γ^J is a sequence of solutions to (7.12), and

$$u_\infty^j := \Phi[z_\infty^j] + R[z_\infty^j]\xi_\infty^j \in C(\mathbb{R}; H^1) \quad (C([0, \infty); H^1) \text{ for } j = 0) \quad (8.12)$$

is a solution of NLS.

Fix $J < J^*$ large enough for Lemma 7.1. By the conclusion of the previous section, if all $\xi_\infty^0, \dots, \xi_\infty^J$ are scattering as $t \rightarrow \infty$, then $\|\xi\|_{\text{st}(0, \infty)}$ is bounded, contradicting (8.9). Hence some of the profiles does not scatter as $t \rightarrow \infty$.

Suppose that ξ_∞^k does not scatter as $t \rightarrow \infty$. Since $u(s^k) \rightharpoonup u_\infty^k(0)$ weakly in H_τ^1 ,

$$\mathbb{M}(u) \geq \mathbb{M}(u_\infty^k) + o(1), \quad \mathbb{E}(u) \geq \mathbb{E}(u_\infty^k) + o(1), \quad (8.13)$$

as the negative part of \mathbb{E} is compact on H_r^1 . Hence

$$(\mathbb{M}(u_\infty^k), \mathbb{E}(u_\infty^k)) \leq (M_*, E_*). \quad (8.14)$$

Then the minimality of (M_*, E_*) implies that $(\mathbb{M}(u_\infty^k), \mathbb{E}(u_\infty^k)) = (M_*, E_*)$ and that u_∞^k is a minimal solution which does not scatter to \mathcal{S}_0 as $t \rightarrow \infty$.

In the case $k = 0$, namely u_∞^0 is a minimal solution, then by the orthogonality in the linear profile decomposition, we have

$$\begin{aligned} \mathbb{M}(u) &= \mathbb{M}(u_\infty^0) + \sum_{1 \leq j < J} \mathbb{M}(\lambda_\infty^j(-s^j)) + \mathbb{M}(\gamma^J(0)) + o(1), \\ \mathbb{E}(u) &= \mathbb{E}(u_\infty^0) + \sum_{1 \leq j < J} \mathbb{H}(\lambda_\infty^j(-s^j)) + \mathbb{H}(\gamma^J(0)) + o(1). \end{aligned} \quad (8.15)$$

Positivity of \mathbb{M} and \mathbb{H} on $P_c H^1$ implies $\mathbb{H}_\theta(\lambda_\infty^j(-s^j)) = o(1)$ and $\mathbb{H}_\theta(\gamma^J(0)) = o(1)$. Hence by the norm equivalence in time, $\lambda_\infty^j = 0$ for all j and $\gamma^J \rightarrow 0$ in $L_t^\infty H_x^1(0, \infty)$. This means that $\xi(0) \rightarrow \xi_\infty^0(0)$ in H_x^1 .

Even if ξ_∞^0 is scattering, there can be at most one profile which is not scattering. Suppose for contradiction, ξ_∞^l is another minimal solution. Since u_∞^0 is scattering, $\mathbb{E}(u_\infty^0)$ can be negative only by the soliton component, hence $\mathbb{E}(u_\infty^0) \gtrsim -\mu_0$. Then (8.15) together with the scattering of ξ_∞^j as $t \rightarrow -\infty$ for $j > 0$ (see Lemma 7.1) implies

$$\mathbb{H}(\xi_\infty^k(-s^k)) + \mathbb{H}(\xi_\infty^l(-s^l)) \leq E_* + O(\mu_0). \quad (8.16)$$

Since $E_* \gtrsim 1 \gg \mu_0$, this implies, taking account of the soliton components of u_∞^k, u_∞^l at $t \rightarrow -\infty$,

$$\min(\mathbb{E}(u_\infty^k), \mathbb{E}(u_\infty^l)) \leq E_*/2 + O(\mu_0) < E_*, \quad (8.17)$$

contradicting the minimality of both u_∞^k and u_∞^l .

Next we prove the precompactness of the orbit of a minimal solution. Henceforth, the index n of sequences is made explicit in order to avoid confusion. Let $u = \Phi[z] + R[z]\xi \in \text{GS}(M_*, E_*)$ be a global solution satisfying

$$(\mathbb{M}(u), \mathbb{E}(u)) = (M_*, E_*), \quad \|\xi\|_{\text{st}(0, \infty)} = \infty. \quad (8.18)$$

Then for any sequence $0 < t_n \rightarrow \infty$, the above argument applies to $u_n := u(t - t_n)$ with $I_n = (-t_n, \infty) \rightarrow \mathbb{R}$. Then $s_n^j \rightarrow \sigma^j \infty$ with $\exists \sigma^j \in \{+, -\}$ for each $j > 0$.

If ξ_∞^0 is not scattering in both the time directions, then $\xi_n(0) = \xi(t_n)$ is strongly convergent, so is $u(t_n)$. If ξ_∞^0 is scattering in both the time directions, then there is a unique $k > 0$ such that ξ_∞^k does not scatter as $t \rightarrow \sigma^k \infty$. Hence those profiles ξ_∞^j with $\sigma^j = -\sigma^k$ are all scattering as $t \rightarrow -\sigma^k \infty$, which implies, by the conclusion of the previous section, ξ_n is bounded in $\text{st}(\{t \in I_n \mid \sigma^k t < 0\})$, namely

$$\sup_n \|\xi\|_{\text{st}(0, t_n)} < \infty \quad \text{or} \quad \sup_n \|\xi\|_{\text{st}(t_n, \infty)} < \infty, \quad (8.19)$$

depending on $\sigma^k = +$ or $-$, but both are contradicting $\|\xi\|_{\text{st}(0, \infty)} = \infty$. Therefore

$$\{u(t) \mid t \geq 0\} \subset H_r^1(\mathbb{R}^3) \quad (8.20)$$

is precompact for such a minimal solution u . By Lemma 3.1, we have a lower bound $\mathbb{K}_2(u(t)) \geq \kappa_* := \kappa(M_*, \mathcal{E}_1(M_*) - E_*) > 0$. The precompactness implies that there is $R \gg 1$ such that

$$\sup_{t>0} \int_{|x|>R} [|\nabla u|^2 + |u|^2 + |u|^4] dx \ll \kappa_*. \quad (8.21)$$

Then the saturated virial identity as in Section 3 implies

$$\partial_t \langle Rf_R u | iu_r \rangle > \kappa_* > 0, \quad (8.22)$$

for all $t > 0$, which obviously contradicts the boundedness of $\langle Rf_R u | iu_r \rangle$ in $t > 0$. This concludes the scattering to the ground states \mathcal{S}_0 in (ii) of Lemma 3.1, and so the proof of Theorem 1.1.

APPENDIX A. DECAY OF THE POTENTIAL

Here we prove that $V, x \cdot \nabla V, (x \cdot \nabla)^2 V \in L^2 + L_0^\infty$ for the radial function $V(x) = V(|x|)$ implies $|V(r)| + |rV_r(r)| \rightarrow 0$ as $r \rightarrow \infty$.

Decompose $V = V_2 + V_\infty$ such that

$$\|V\|_{L^2+L^\infty} \sim \|V_2\|_{L^2} + \|V_\infty\|_{L^\infty}, \quad \lim_{R \rightarrow \infty} \|V_\infty\|_{L^\infty(|x|>R)} = 0. \quad (A.1)$$

For any $\varepsilon > 0$, there is $R > 0$ such that $\|V_\infty\|_{L^\infty(|x|>R)} < \varepsilon$. Let $\chi : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function satisfying $\chi(t) = t$ for $|t| \geq 3$, $|\chi(t)| \leq |t|$ and $0 \leq \chi'(t) \leq 10$ for all t , and $\chi(t) = 0$ for $|t| \leq 2$. Put

$$V_{(\varepsilon)} := \varepsilon \chi(V/\varepsilon), \quad (A.2)$$

Then $V_{(\varepsilon)}(r) \neq 0$ implies $|V(r)| > 2\varepsilon$ and so $|V_{(\varepsilon)}(r)| \leq |V(r)| \sim |V_2(r)|$ for $r > R$. Hence for $r > R$,

$$\begin{aligned} |V_{(\varepsilon)}(r)|^2 &\leq 2 \int_r^\infty |V_{(\varepsilon)}(r)| \cdot |r \partial_r V_{(\varepsilon)}(r)| \frac{r^2 dr}{r^3} \\ &\lesssim \|V_{(\varepsilon)}\|_{L^2(|x|>R)} \| |x|^{-3} \|_{L^2 \cap L^\infty(|x|>R)} \|r \partial_r V_{(\varepsilon)}\|_{(L^2+L^\infty)(|x|>R)} \\ &\lesssim \|V_2\|_{L^2(|x|>R)} R^{-3/2} \|r \partial_r V\|_{L^2+L^\infty} \rightarrow 0 \end{aligned} \quad (A.3)$$

as $R \rightarrow \infty$. Since $|V| \leq |V_{(\varepsilon)}| + 3\varepsilon$, we deduce that $V(x) \rightarrow 0$ as $|x| \rightarrow \infty$. The same argument implies that $r \partial_r V(x) \rightarrow 0$.

APPENDIX B. TABLE OF NOTATION

Besides the following list, see (5.2)–(5.4) for notation of sequences without index.

symbols	description	defined in
\mathfrak{s}	sign of nonlinearity	(1.7)
H, V	Schrödinger operator and the potential	(1.7), Section 1.5
e_0, ϕ_0	unique eigenvalue and ground state of H	(1.8)
\mathfrak{p}	L_x^p -exponent where the wave operator is bounded	Section 1.5 (iv)
Q	ground state without V	(1.5)
$\mathbb{E}, \mathbb{M}, \mathbb{K}_2$	energy, mass and virial	(1.14), (1.22)
$[\cdot], \mathbb{G}, \mathbb{H}, \mathbb{I}$	some functionals	(2.1), (2.7)
$\mathbb{E}^0, \mathbb{H}^0$	energy without V	(2.2)
$\mathcal{S}, \mathcal{S}_j, \mathcal{E}_j$	all solitons, j -th bound states and their energy	(1.13), (1.16), (1.15)

(Φ, Ω)	coordinates of ground states	Lemma 2.2
z_b, D_b, μ_b	size of the above coordinates	Lemma 2.2, (2.14)
μ_d, z_d	size of parameterized ground states	(2.19)
μ_p, D_p	size of ground state projection	Lemma 4.1
μ_e	small mass to characterize \mathcal{E}_1	Proposition 2.5
$L_x^p, H^s, B_{p,q}^s$	Lebesgue, Sobolev and Besov spaces on \mathbb{R}^3	Section 1.4, (1.10)
$H_r^1, H^1[\mu]$	radial Sobolev and its subset with small mass	(4.1)
$L_t^p X(I)$	B-space valued L^p space on interval	Section 1.4
$(\cdot \cdot), \langle \cdot \cdot \rangle$	inner products on $L^2(\mathbb{R}^3)$	(1.30)
$\mathcal{S}_p^t, \mathcal{S}_p'$	L^p -preserving scaling and its generator	(2.3)-(2.4)
κ	lower bound on $ \mathbb{K}_2 $	Proposition 2.5
$\mathcal{H}_c[z], R[z]$	normal subspace of \mathcal{S}_0 and its coordinate	(4.2), (4.5)
$B[z], \tilde{N}(z, \xi)$	terms in the equation around \mathcal{S}_0	(4.11), (4.12), (4.13)
$\cdot\{s\}$	functions defined around s	(4.14)
$u[z, s]$	solution of the linearized equation	(4.16)
$\mathcal{D}f[z, s]$	Duhamel form of the linearized equation	(4.19)
$u[z, s]_>$	solution with nonlinearity turnoff	(4.23)
$\text{Stz}^s, \text{Stz}^{*s}, \mathbf{st}$	Strichartz norms	(4.26)
$[z; T_0, T_1; T_2]$	semi-norm to measure deviation from $u[z, T_0]$	(4.27)
\mathcal{I}	totality of intervals	(5.5)
SBC	uniformly small, bounded and continuous functions	(5.6)–(5.7)
$u[z, s]_{(\infty)}$	linearized solution with limit initial data	(5.10)
\mathbb{H}_θ	fractional power of HP_c	(5.24)
J^*, s^j	the number and centers of profiles	Lemma 5.3
$\lambda^j, \lambda^{[0\dots J]}, \gamma^J$	linear profiles and their sum, and remainder	Lemma 5.3, (7.5)
$\triangleleft \dots \triangleright$	difference	(6.21)
Λ^j, Γ^J	nonlinear profiles and remainder	(7.9), (7.12)
s_\pm^j	times around profiles for decomposition	(7.24)
o_τ	vanishing terms as $n \rightarrow \infty$ and $\tau \rightarrow \infty$	(7.25)
$\text{GS}(\mu, A)$	global solutions below some mass and energy	(8.1)
\mathcal{X}	mass-energy region with uniform Strichartz bound	(8.2)
$(\cdot, \cdot) \leq, \ll$	product orders	(8.3)

REFERENCES

- [1] H. Bahouri and P. Gérard, *High frequency approximation of solutions to critical nonlinear wave equations*. Amer. J. Math. **121** (1999), no. 1, 131–175.
- [2] M. Beceanu, *Structure of wave operators for a scaling-critical class of potentials*, Amer. J. Math. **136** (2014), no. 2, 255–308.
- [3] S. Cuccagna and M. Masaya, *On small energy stabilization in the NLS with a trapping potential*, preprint, arXiv:1309.0655.
- [4] T. Duyckaerts, J. Holmer and S. Roudenko, *Scattering for the non-radial 3D cubic nonlinear Schrödinger equation*, Math. Res. Lett. **15** (2008), no. 6, 1233–1250.
- [5] T. Duyckaerts, C. Kenig and F. Merle, *Classification of radial solutions of the focusing, energy-critical wave equation*, preprint, arXiv:1204.0031.
- [6] T. Duyckaerts and F. Merle, *Dynamic of threshold solutions for energy-critical NLS*. Geom. Funct. Anal. **18** (2009), no. 6, 1787–1840.
- [7] T. Duyckaerts and S. Roudenko, *Threshold solutions for the focusing 3D cubic Schrödinger equation*, Rev. Mat. Iberoam. **26** (2010), no. 1, 1–56.
- [8] Z. Gang and I. M. Sigal, *Relaxation of solitons in nonlinear Schrödinger equations with potential*, Adv. Math. **216** (2007), no. 2, 443–490.

- [9] S. Gustafson, K. Nakanishi and T. Tsai, *Asymptotic stability and completeness in the energy space for nonlinear Schrödinger equations with small solitary waves*, Int. Math. Res. Not. (2004) no. 66, 3559–3584.
- [10] J. Holmer and S. Roudenko, *A sharp condition for scattering of the radial 3D cubic nonlinear Schrödinger equation*, Comm. Math. Phys. **282** (2008), no. 2, 435–467.
- [11] T. Kato, *An $L^{q,r}$ -theory for nonlinear Schrödinger equations*, Spectral and scattering theory and applications, Adv. Stud. Pure Math., vol. 23, Math. Soc. Japan, Tokyo, 1994, pp. 223–238.
- [12] C. Kenig and F. Merle, *Global well-posedness, scattering, and blow-up for the energy-critical focusing nonlinear Schrödinger equation in the radial case*, Invent. Math. **166** (2006), no. 3, pp. 645–675.
- [13] K. Nakanishi, *Scattering theory for the nonlinear Klein-Gordon equation with Sobolev critical power*. Internat. Math. Res. Notices **1999**, no. 1, 31–60.
- [14] K. Nakanishi, T. V. Phan, and T.-P. Tsai, *Small solutions of nonlinear Schrödinger equations near first excited states*, J. Funct. Analysis **263** (2012), 703–781.
- [15] K. Nakanishi and W. Schlag, *Global dynamics above the ground state energy for the focusing nonlinear Klein-Gordon equation*, J. Diff. Equations **250** (2011), 2299–2333.
- [16] K. Nakanishi and W. Schlag, *Global dynamics above the ground state energy for the cubic NLS equation in 3D*, Calc. Var. and PDE, **44** (2012), no. 1-2, 1–45.
- [17] T. Ogawa, and Y. Tsutsumi, *Blow-up of H^1 solution for the nonlinear Schrödinger equation*, J. Diff. Eq. **92** (1991), 317–330.
- [18] L. E. Payne and D. H. Sattinger, *Saddle points and instability of nonlinear hyperbolic equations*. Israel J. Math. **22** (1975), no. 3-4, 273–303.
- [19] A. Soffer and M. I. Weinstein, *Selection of the ground state for nonlinear Schrödinger equations*. Rev. Math. Phys. **16** (2004), no. 8, 977–1071.
- [20] T.-P. Tsai, *Asymptotic dynamics of nonlinear Schrödinger equations with many bound states*, J. Diff. Equations **192** (2003) 225–282.
- [21] T.-P. Tsai and H.-T. Yau, *Asymptotic dynamics of nonlinear Schrödinger equations: resonance dominated and radiation dominated solutions*, Comm. Pure Appl. Math. **55** (2002), 153–216.
- [22] T.-P. Tsai and H.-T. Yau, *Relaxation of excited states in nonlinear Schrödinger equations*, Int. Math. Res. Not. **31** (2002), 1629–1673.
- [23] T.-P. Tsai and H.-T. Yau, *Stable directions for excited states of nonlinear Schrödinger equations*, Comm. Partial Differential Equations **27** (2002), 2363–2402.
- [24] T.-P. Tsai and H.-T. Yau, *Classification of asymptotic profiles for nonlinear Schrödinger equations with small initial data*, Adv. Theor. Math. Phys. **6** (2002), 107–139.
- [25] M. C. Vilela. *Inhomogeneous Strichartz estimates for the Schrödinger equation*. Trans. Amer. Math. Soc. **359** (2007), 2123–2136. 35
- [26] K. Yajima, *The $W^{k,p}$ -continuity of wave operators for Schrödinger operators*, J. Math. Soc. Japan **47** (1995), 551–581.

DEPARTMENT OF PURE AND APPLIED MATHEMATICS GRADUATE SCHOOL OF INFORMATION
SCIENCE AND TECHNOLOGY OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN
E-mail address: nakanishi@ist.osaka-u.ac.jp