

A Note on the Rainbow Connectivity of Tournaments

Jesús Alva-Samos¹ Juan José Montellano-Ballesteros²

April 27, 2015

Abstract

An arc-coloured digraph D is said to be *rainbow connected* if for every two vertices u and v there is an uv -path all whose arcs have different colours. The minimum number of colours required to make the digraph rainbow connected is called the *rainbow connection number* of D , denoted $\vec{rc}(D)$. In [4] it was showed that if T is a strong tournament with $n \geq 5$ vertices, then $2 \leq \vec{rc}(T) \leq n - 1$; and that for every n and k such that $3 \leq k \leq n - 1$, there exists a tournament T on n vertices such that $\vec{rc}(T) = k$.

In this note it is showed that for any $n \geq 6$, there is a tournament T of n vertices such that $\vec{rc}(T) = 2$.

1 Introduction

Chartrand, Johns, McKeon and Zhang introduced in [3] the concept of rainbow connection in graphs, and since then such topic has been broadly studied, see [5] for a survey on the matter. The rainbow connection for digraphs was first presented by Dorbec, Schiermeyer, Sidorowicz and Sopena in [4].

A *tournament* is a digraph where every two vertices has exactly one arc joining them. Let ρ be an arc-colouring on a digraph D , a (directed) path in D is called a *rainbow* if it has no two arcs sharing the same colour. If there is a rainbow between every pair of vertices, then the digraph is *rainbow connected* and ρ is called a *rainbow colouring*. The *rainbow connection number* of a digraph D , denoted by $\vec{rc}(D)$, is defined as the smallest number of colours such that D admits a rainbow colouring.

In [4] the following two theorems were proven:

¹ Instituto de Matemáticas, UNAM

²Instituto de Matemáticas, UNAM. juancho@matem.unam.mx

Theorem 1 (Dorbec et al. [4]). *If T is a strong tournament with $n \geq 5$ vertices, then $2 \leq \vec{\text{rc}}(T) \leq n - 1$.*

Theorem 2 (Dorbec et al. [4]). *For every n and k such that $3 \leq k \leq n - 1$, there exists a tournament T on n vertices such that $\vec{\text{rc}}(T) = k$.*

Also, in [4], it is showed that for each $n \cong 8 \pmod{12}$ there is a tournament T of order n and $\vec{\text{rc}}(T) = 2$.

In this note it is showed that for every $n \geq 6$ there is a tournament T of order n and $\vec{\text{rc}}(T) = 2$.

2 The tournaments

When $n = 4, 5$ it is easy to verified that $\vec{\text{rc}}(T) \geq 3$ for each tournament T on n vertices. For an integer $n \geq 2$ and a set $S \subseteq \{1, 2, \dots, n - 1\}$, the *circulant digraph* $C_n(S)$ is defined as the digraph with vertex set $V(C_n(S)) = \{v_0, v_1, \dots, v_{n-1}\}$ and arc set $A(C_n(S)) = \{(v_i, v_j) \mid j - i \equiv s \pmod{n}, s \in S\}$.

Theorem 3. *For every $n \geq 6$, there is a tournament T of order n with $\vec{\text{rc}}(T) = 2$.*

Proof. For $n = 6$, let T be the first tournament in Figure 1. Now, for $n = 2k + 1$, with $k \geq 3$, let $T = C_{2k+1}(1, 2, 4, \dots, 2(k - 1))$ and consider the partition of $A(T)$ into the sets

$$\begin{aligned} A_0 &= \{(u_0, u_1), (u_0, u_2), (u_1, u_{2k-1})\} \\ &\cup (\{(u_r, u_s) \mid r \equiv 0 \pmod{2}, r \geq 2\} \setminus \{(u_2, u_{2k})\}) \end{aligned}$$

and $A_1 = A(T) \setminus A_0$. Now, let ρ be the coloring where $\rho(a) = i$ if $a \in A_i$ for $i = 0, 1$. We will show that there is a rainbow $u_i u_j$ -path for $0 \leq i \leq n - 1$ and $j = i + 3, i + 5, \dots, i + 2k - 1, i + 2k$, and the result will follow.




Figure 1: Tournaments on n vertices with $\vec{rc}(T_n) = 2$ where $n = 6, 7, 8$.

For each i with $3 \leq i \leq 2k - 1$ let consider the paths

$$\begin{cases} u_i u_{i+1} u_r & \text{for } r = 3, 5, \dots, 2k - 1; \\ u_i u_{i+2k-2} u_{i+2k}; \end{cases}$$

and for the vertex u_{2k} consider the paths $u_{2k} u_1 u_2$, $u_{2k} u_{2k-3} u_{2k-1}$ and for $r = 5, 7, \dots, 2k - 1$ let $u_{2k} u_0 u_r$.

For $i \in \{1, 2\}$ consider the paths $u_i u_{i+1} u_{i+r}$ for $r = 3, 5, \dots, 2k - 3$; and the paths $u_1 u_{2k-1} u_{2k}$, $u_1 u_{2k-1} u_0$, $u_2 u_{2k} u_0$, $u_2 u_{2k} u_1$.

Finally, for the vertex u_0 let consider the paths

$$\begin{cases} u_0u_1u_3 \\ u_0u_{r-1}u_r & \text{for } r = 5, 7, \dots, 2k - 1 ; \\ u_0u_{2k-2}u_{2k}. \end{cases}$$

By the definition of ρ , it is not hard to see that all these paths are rainbow paths and therefore $\vec{rc}(T) = 2$.

For the case $n = 2k \geq 8$, let consider the tournament T obtained by adding a new vertex v to $C_{2k-1}(1, 2, 4, \dots, 2(k-2))$ such that $(v, u_i) \in A(T)$ if and only if i is even. Let colouring the arcs of T by extending the coloring ρ of $C_{2k-1}(1, 2, 4, \dots, 2(k-2))$ described above and given color 1 to all the new arcs. Consider the paths vu_iu_{i+1} for $i+1$ odd; $u_iu_{i+1}v$ for i even, $i \neq 2k$; and $u_{2k}u_1v$. By construction all these paths are rainbow paths, therefore $\vec{rc}(T) = 2$. \square

Finally, combining Theorems 2 and 3 we have

Theorem 4. *For every $n \geq 6$ and every k such that $2 \leq k \leq n - 1$, there exists a tournament T on n vertices such that $\vec{rc}(T) = k$.* \square

References

- [1] J. Bang-Jensen, G. Gutin. Digraphs. Theory, Algorithms and Applications. Springer-Verlag, 2002.
- [2] J.A. Bondy, U.S.R. Murty. Graph Theory. Springer. 2008.
- [3] G. Chartrand, G.L. Johns, K.A. McKeon, P. Zhang. Rainbow connection in graphs. Math. Bohem. 133 (2008) 85-98.
- [4] P. Dorbec, I. Schiermeyer, E. Sidorowicz, E. Sopena. Rainbow connection in oriented graphs. Discrete Appl. Math. 179 (2014) 69-78.

- [5] X. Li, Y. Sun, Rainbow Connections of Graphs, in: Springer Briefs in Mathematics, Springer, 2012.
- [6] J.W. Moon, Topics on Tournaments, Holt, Rinehart and Winston Eds., New York, 1968.