

Spectral MLE: Top- K Rank Aggregation from Pairwise Comparisons

Yuxin Chen

*Department of Statistics
Stanford University
Stanford, CA 94305, USA*

YXCHEN@STANFORD.EDU

Changho Suh

*Department of Electrical Engineering
Korea Advanced Institute of Science and Technology
Daejeon, Korea*

CHSUH@KAIST.AC.KR

Editor:

Abstract

This paper explores the preference-based top- K rank aggregation problem. Suppose that a collection of items is repeatedly compared in pairs, and one wishes to recover a consistent ordering that emphasizes the top- K ranked items, based on partially revealed preferences. For concreteness, this work focuses on the popular Bradley-Terry-Luce model that postulates a set of latent preference scores underlying all items, where the odds of paired comparisons depend only on the relative scores of the items involved.

We characterize the minimax limits on the identifiability of top- K ranked items, in the presence of random and non-adaptive sampling. Our findings highlight a separation measure that quantifies the gap of preference scores between the K^{th} and $(K+1)^{\text{th}}$ ranked items. In an attempt to approach this minimax limit, we propose a nearly linear-time ranking scheme, called *Spectral MLE*, that returns the indices of the top- K items in accordance to a careful score estimate. In a nutshell, Spectral MLE starts with an initial score estimate with minimal squared loss (obtained via a spectral method), and then successively refines each component with the assistance of coordinate-wise MLEs. Encouragingly, Spectral MLE achieves perfect top- K item identification with minimal sample complexity. The practical applicability of Spectral MLE is further corroborated by numerical experiments.

Keywords: Bradley-Terry-Luce model, top- K ranking, linear-time algorithm, minimax limits, coordinate-wise MLE

1. Introduction and Motivation

The task of rank aggregation is encountered in a wide spectrum of contexts like social choice (Caplin and Nalebuff, 1991; Soufiani et al., 2014b), web search and information retrieval (Brin and Page, 1998; Dwork et al., 2001), crowd sourcing (Chen et al., 2013), recommendation systems (Baltrunas et al., 2010), to name just a few. Given partial preference information over a collection of items, the aim is to identify a consistent ordering that best respects the revealed preference. In the high-dimensional regime, one is often faced with two challenges: (1) the number of items to be ranked is ever growing, which makes it increasingly harder to recover a consistent total ordering over all items; (2) the observed

data is highly incomplete and inconsistent: only a small number of noisy pairwise / listwise preferences can be acquired.

In an effort to address such challenges, this paper explores a popular pairwise preference-based model, which postulates the existence of a ground-truth ranking. Specifically, consider a parametric model involving n items, each assigned a preference score that determines its rank. Concrete examples of preference scores include the overall rating of an athlete, the academic performance and competitiveness of a university, the dining quality of a restaurant, etc. Each item is then repeatedly compared against a few others in pairs, yielding a set of *noisy* binary comparisons generated based on the relative preference scores. In many situations, the number of repeated comparisons essentially reflects the signal-to-noise ratio (SNR) or the quality of the information revealed for each pair of items. The goal is then to develop a “denoising” procedure that recovers the ground-truth ranking, ideally from a minimal number of noisy observations.

There has been a proliferation of ranking schemes that suggest partial solutions. While the ranking that we are seeking is better treated as a function of the preference parameters, most of the aforementioned schemes adopt the natural “plug-in” procedure, that is, start by inferring the preference scores, and then return a ranking in accordance to the parametric estimates. The most popular paradigm is arguably the maximum likelihood estimation (MLE) (Ford, 1957), where the main appeal of MLE is its inherent convexity under several comparison models, e.g. the Bradley-Terry-Luce (BTL) model (Bradley and Terry, 1952; Luce, 1959). Encouragingly, MLE often achieves low ℓ_2 estimation loss while retaining efficient finite-sample complexity. Another prominent alternative concerns a family of spectral ranking algorithms (e.g. *PageRank* (Brin and Page, 1998)). A provably efficient choice within this family is *Rank Centrality* (Negahban et al., 2012), which produces an estimate with nearly minimax mean squared error (MSE). While both MLE and Rank Centrality allow intriguing guarantees towards finding faithful parametric estimates, the squared loss metric considered therein does not necessarily imply optimality of the ranking accuracy. In fact, there is no shortage of high-dimensional situations that admit parametric estimates with low squared loss while precluding reliable ranking. Furthermore, many realistic scenarios emphasize only a few items that receive the highest ranks. Unfortunately, the above MSE results fall short of ensuring recovery of the top-ranked items.

In this work, we consider accurate identification of top- K ranked items under the popular BTL pairwise comparison model, assuming that the item pairs we can compare are selected in a random and non-adaptive fashion (termed *passive ranking*). In particular, we aim to explore the following questions: (a) what is the minimum number of repeated comparisons necessary for reliable ranking? (b) how is the ranking accuracy affected by the underlying preference score distributions? We will address these two questions from both statistical and algorithmic perspectives.

1.1 Main Contributions

This paper investigates minimax optimal procedures for top- K rank aggregation. Our contributions are two-fold.

To begin with, we characterize the fundamental three-way tradeoff between the number of repeated comparisons, the sparsity of the comparison graph, and the preference score

distribution, from a minimax perspective. In particular, we emphasize a separation measure that quantifies the gap of preference scores between the K^{th} and $(K + 1)^{\text{th}}$ ranked items. Our results demonstrate that the minimal sample complexity or quality of paired evaluation (reflected by the number of repeated comparisons per an observed pair) scales inversely with the separation measure at a quadratic rate.

Algorithmically, we propose a nearly linear-time two-stage procedure, called *Spectral MLE*, which allows perfect top- K identification as soon as the sample complexity exceeds the minimax limits (modulo some constant). Specifically, Spectral MLE starts by obtaining careful initial scores that are faithful in the ℓ_2 sense (e.g. via a spectral ranking method), and then iteratively sharpens the pointwise estimates via a careful comparison between the running iterates and the coordinate-wise MLEs. This algorithm is designed primarily in an attempt to seek a score estimate with minimal pointwise loss which, in turn, guarantees optimal ranking accuracy. Furthermore, numerical experiments demonstrate that Spectral MLE outperforms Rank Centrality by achieving lower ℓ_∞ estimation error and higher ranking accuracy.

1.2 Prior Art

There are two distinct families of observation models that have received considerable interest: (1) *value-based model*, where the observation on each item is drawn only from the distribution underlying this individual; (2) *preference-based model*, where one observes the relative order among a few items instead of revealing their individual values. Best- K identification in the value-based model with adaptive sampling (termed *active ranking*) is closely related to the multi-armed bandit problem, where the fundamental identification complexity has been characterized (Gabillon et al., 2011; Bubeck et al., 2013; Jamieson et al., 2014). In addition, the value-based and preference-based models have been compared in terms of minimax error rates in estimating the latent quantities; see (Shah et al., 2014).

In the realm of pairwise preference settings, many *active ranking* schemes (Busa-Fekete and Hüllermeier, 2014) have been proposed in an attempt to optimize the exploration-exploitation trade-off. For instance, in the noise-free case, (Jamieson and Nowak, 2011) considered perfect total ranking and characterized the query complexity gain of adaptive sampling relative to random queries, provided that the items under study admit a low-dimensional Euclidean embedding. Furthermore, several works (Ailon, 2012; Jamieson and Nowak, 2011; Braverman and Mossel, 2008; Wauthier et al., 2013) explored the query complexity in the presence of noise, but were basically designed to recover “approximately correct” total rankings—a solution with loss at most a factor $(1 + \epsilon)$ from optimal—rather than accurate ordering. Another path-based approach has been proposed to accommodate accurate top- K queries from noisy pairwise data (Eriksson, 2013), where the observation error is assumed to be i.i.d. instead of being item-dependent. Motivated by the success of value-based racing algorithms, (Busa-Fekete et al., 2013; Busa-Fekete and Hüllermeier, 2014) came up with a generalized racing algorithm that often led to efficient sample complexity. In contrast, the current paper concentrates on top- K identification in a *passive* setting, assuming that partial preferences are collected in a noisy, random, and non-adaptive manner. This was previously out of reach.

Apart from Rank Centrality and MLE, the most relevant work is (Rajkumar and Agarwal, 2014). For a variety of rank aggregation methods, they developed intriguing sufficient statistical hypotheses that guarantee the convergence to an optimal ranking, which in turn leads to sample complexity bounds for Rank Centrality and MLE. Nevertheless, they focused on perfect total ordering instead of top- K selection, and their results fall short of a rigorous justification as to whether or not the derived sample complexity bounds are statistically optimal.

Finally, there are many related yet different problem settings considered in the prior literature. For instance, the work (Ammar and Shah, 2012) approached top- K ranking using a maximum entropy principle, assuming the existence of a distribution μ over all possible permutations. Recent work (Soufiani et al., 2013, 2014a) investigated consistent *rank breaking* under more generalized models involving full rankings. A family of distance measures on rankings has been studied and justified based on an axiomatic approach (Farnoud and Milenkovic, 2014). Another line of works considered the popular distance-based Mallows model (Lu and Boutilier, 2011; Busa-Fekete et al., 2014; Awasthi et al., 2014). An online ranking setting has been studied as well (Harrington, 2003; Farnoud et al., 2014). More broadly, the minimax recovery limits under general pairwise measurements have recently been determined by (Chen et al., 2015b). These are beyond the scope of the present work.

1.3 Organization and Notation

The remainder of the paper is organized as follows. Section 2 introduces the pairwise comparison model as well as the key performance metrics for the top- K ranking task. The main results, including a fundamental minimax lower limit and an achievability result by nearly linear-time algorithms, are summarized and discussed in Section 3. Section 4 presents the detailed procedure and performance guarantees of the proposed Spectral MLE algorithm, and provides a heuristic treatment as to why it is expected to control the ℓ_∞ estimation error. We conclude the paper with a summary of our findings and a discussion of about future research directions in Section 5. The proofs of the ranking performance of Spectral MLE (i.e. Theorem 7) and the minimax lower bound (i.e. Theorem 3) are deferred to Appendix A and Appendix B, respectively.

Before continuing, we provide a brief summary of the notations used throughout the paper. Let $[n]$ represent $\{1, 2, \dots, n\}$. We denote by $\|\mathbf{w}\|$, $\|\mathbf{w}\|_1$, $\|\mathbf{w}\|_\infty$ the ℓ_2 norm, ℓ_1 norm, and ℓ_∞ norm of \mathbf{w} , respectively. A graph \mathcal{G} is said to be an Erdős–Rényi random graph, denoted by $\mathcal{G}_{n,p_{\text{obs}}}$, if each pair (i, j) is connected by an edge independently with probability p_{obs} . Besides, we use $\text{deg}(i)$ to represent the degree of vertex i in \mathcal{G} .

Additionally, for any two sequences $f(n)$ and $g(n)$, $f(n) \gtrsim g(n)$ or $f(n) = \Omega(g(n))$ mean that there exists a constant c such that $f(n) \geq cg(n)$; $f(n) \lesssim g(n)$ or $f(n) = O(g(n))$ mean that there exists a constant c such that $f(n) \leq cg(n)$; and $f(n) \asymp g(n)$ or $f(n) = \Theta(g(n))$ mean that there exist constants c_1 and c_2 such that $c_1g(n) \leq f(n) \leq c_2g(n)$.

2. Problem Setup

To formalize matters, we present mathematical setups and key performance metrics in this section.

Comparison Model and Assumptions. Suppose that we observe a few pairwise evaluations over n items. To pursue a statistical understanding towards the ranking limits, we assume that the pairwise comparison outcomes are generated according to the BTL model (Bradley and Terry, 1952; Luce, 1959), a long-standing model that has been applied in numerous applications (Agresti, 2014; Hunter, 2004).

- *Preference Scores.* The BTL model hypothesizes on the existence of some hidden preference vector $\mathbf{w} = [w_i]_{1 \leq i \leq n}$, where w_i represents the underlying preference score / weight of item i . The outcome of each paired comparison depends only on the scores of the items involved. Without loss of generality, we will assume throughout that

$$w_1 \geq w_2 \geq \dots \geq w_n > 0 \quad (1)$$

unless otherwise specified.

- *Comparison Graph.* Denote by $\mathcal{G} = ([n], \mathcal{E})$ the comparison graph such that items i and j are compared if and only if (i, j) belongs to the edge set \mathcal{E} . We will mostly assume that \mathcal{G} is drawn from the Erdős–Rényi model $\mathcal{G} \sim \mathcal{G}_{n, p_{\text{obs}}}$ for some observation ratio p_{obs} .
- *(Repeated) Pairwise Comparisons.* For each $(i, j) \in \mathcal{E}$, we observe L independent paired comparisons between items i and j . The outcome of the l^{th} comparison between them, denoted by $y_{i,j}^{(l)}$, is generated as per the BTL model:

$$y_{i,j}^{(l)} = \begin{cases} 1, & \text{with probability } \frac{w_i}{w_i + w_j}, \\ 0, & \text{else,} \end{cases} \quad (2)$$

where $y_{i,j}^{(l)} = 1$ indicates a win by i over j . We adopt the convention that $y_{j,i}^{(l)} = 1 - y_{i,j}^{(l)}$. It is assumed throughout that conditional on \mathcal{G} , the $y_{i,j}^{(l)}$'s are jointly independent across all l and $i > j$. For ease of presentation, we introduce the collection of sufficient statistics as

$$\mathbf{y}_i := \{y_{i,j} \mid j : (i, j) \in \mathcal{E}\}; \quad y_{i,j} := \frac{1}{L} \sum_{l=1}^L y_{i,j}^{(l)}. \quad (3)$$

- *Signal to Noise Ratio (SNR) / Quality of Comparisons.* The overall faithfulness of the acquired evaluation between items i and j is captured by the sufficient statistic $y_{i,j}$. Its SNR can be captured by

$$\text{SNR} := \frac{\mathbb{E}^2[y_{i,j}]}{\text{Var}[y_{i,j}]} \asymp L. \quad (4)$$

As a result, the number L of repeated comparisons measures the SNR or the *quality* of comparisons over any observed pair of items.

- *Dynamic Range of Preference Scores.* It is assumed throughout that the dynamic range of the preference scores is fixed irrespective of n , namely,

$$w_i \in [w_{\min}, w_{\max}], \quad 1 \leq i \leq n \quad (5)$$

for some positive constants w_{\min} and w_{\max} bounded away from 0, which amounts to the most challenging regime (Negahban et al., 2012). In fact, the case in which the range w_{\max}/w_{\min} grows with n can be readily translated into the above fixed-range regime by first separating out those items with vanishing scores (e.g. via a simple voting method like Borda count (Ammar and Shah, 2011)).

Performance Metric. Given these pairwise observations, one wishes to see whether or not the top- K ranked items are identifiable. To this end, we consider the probability of error P_e in isolating the *set* of top- K ranked items, i.e.

$$P_e(\psi) := \mathbb{P}\left\{\psi(\mathbf{y}) \neq [K]\right\}, \quad (6)$$

where ψ is any ranking scheme that returns a set of K indices. Here, $[K]$ denotes the (unordered) set of the first K indices. We aim to characterize the fundamental *admissible region* of (L, p_{obs}) where reliable top- K ranking is feasible, i.e. P_e can be vanishingly small as n grows.

3. Minimax Ranking Limits

We explore the fundamental ranking limits from a minimax perspective, which centers on the design of *robust* ranking schemes that guard against the worst case in probability of error. The most challenging component of top- K rank aggregation often hinges upon distinguishing the two items near the decision boundary, i.e. the K^{th} and $(K+1)^{\text{th}}$ ranked items. Due to the random nature of the acquired finite-bit comparisons, the information concerning their relative preference could be obliterated by noise, unless their latent preference scores are sufficiently separated. In light of this, we single out a preference separation measure as follows

$$\Delta_K := \frac{w_K - w_{K+1}}{w_{\max}}. \quad (7)$$

As will be seen, this measure plays a crucial role in determining information integrity for top- K identification.

To model random sampling and partial observations, we employ the Erdős–Rényi random graph $\mathcal{G} \sim \mathcal{G}_{n, p_{\text{obs}}}$. As already noted by (Ford, 1957), if the comparison graph \mathcal{G} is not connected, then there is absolutely no basis to determine relative preferences between two disconnected components. Therefore, a reasonable necessary condition that one would expect is the connectivity of \mathcal{G} , which requires

$$p_{\text{obs}} > \frac{\log n}{n}. \quad (8)$$

All results presented in this paper will operate under this assumption.

A main finding of this paper is an order-wise tight sufficient condition for top- K identifiability, as stated in the theorem below.

Theorem 1 (Identifiability) *Suppose that $\mathcal{G} \sim \mathcal{G}_{n, p_{\text{obs}}}$ with $p_{\text{obs}} \geq c_0 \log n/n$. Assume that $L = O(\text{poly}(n))$ and $w_{\max}/w_{\min} = \Theta(1)$. With probability exceeding $1 - c_1 n^{-2}$,*

the set of top- K ranked items can be identified exactly by an algorithm that runs in time $O(|\mathcal{E}|\log^2 n)$, provided that

$$L \geq \frac{c_2 \log n}{np_{\text{obs}}\Delta_K^2}. \quad (9)$$

Here, $c_0, c_1, c_2 > 0$ are some universal constants.

Remark 2 We assume throughout that the input fed to each ranking algorithm is the sufficient statistic $\{y_{i,j} \mid (i,j) \in \mathcal{E}\}$ rather than the entire collection of $y_{i,j}^{(l)}$, otherwise even reading all data takes at least $O(L \cdot |\mathcal{E}|)$ flops / time.

Theorem 1 characterizes an *identifiable region* within which exact identification of top- K items is plausible by nearly linear-time algorithms. The algorithm we propose, as detailed in Section 4, attempts recovery by computing a score estimate whose errors can be uniformly controlled across all entries. Afterwards, the algorithm reports the K items that receive the highest estimated scores.

Encouragingly, the above identifiable region is minimax optimal. Consider a given separation condition Δ_K , and suppose that nature behaves in an adversarial manner by choosing the worst-case scores \mathbf{w} compatible with Δ_K . This imposes a minimax lower bound on the quality of comparisons necessary for reliable ranking, as given below.

Theorem 3 (Minimax Lower Bounds) Fix $\epsilon \in (0, \frac{1}{2})$, and let $\mathcal{G} \sim \mathcal{G}_{n,p_{\text{obs}}}$. If

$$L \leq c \frac{(1-\epsilon) \log n - 2}{np_{\text{obs}}\Delta_K^2} \quad (10)$$

holds for some constant¹ $c > 0$, then for any ranking scheme ψ , there exists a preference vector \mathbf{w} with separation Δ_K such that $P_e(\psi) \geq \epsilon$.

Theorem 3 taken collectively with Theorem 1 determines the scaling of the fundamental ranking boundary on L . Since the sample size sharply concentrates around $n^2 p_{\text{obs}} L$ in our model, this implies that the required sample complexity for top- K ranking scales inversely with the preference separation at a quadratic rate. Put another way, Theorem 3 justifies the need for a minimum separation criterion that applies to any ranking scheme:

$$\Delta_K \gtrsim \sqrt{\frac{\log n}{np_{\text{obs}}L}}. \quad (11)$$

Somewhat unexpectedly, there is no computational barrier away from this statistical limit (at least in an order-wise sense). Several other remarks of Theorem 1 and Theorem 3 are in order.

- **ℓ_2 Loss vs. ℓ_∞ Loss.** A dominant fraction of prior methods focus on the mean squared error in estimating the latent scores \mathbf{w} . It was established by (Negahban et al., 2012) that the minimax ℓ_2 regret is squeezed between

$$\frac{1}{\sqrt{np_{\text{obs}}L}} \lesssim \inf_{\hat{\mathbf{w}}} \sup_{\mathbf{w}} \frac{\mathbb{E}[\|\hat{\mathbf{w}} - \mathbf{w}\|]}{\|\mathbf{w}\|} \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}},$$

1. More precisely, one can take $c = w_{\min}^4 / (2w_{\max}^4)$.

where the infimum is taken over all score estimators $\hat{\mathbf{w}}$. This limit is almost identical to the minimax separation criterion (11) we derive for top- K identification, except for a potential logarithmic factor. In fact, if the pointwise error of $\hat{\mathbf{w}}$ is *uniformly* bounded by $\sqrt{\log n / (np_{\text{obs}}L)}$, then $\hat{\mathbf{w}}$ necessarily achieves the minimax ℓ_2 error. Moreover, the pointwise error bound presents a fundamental bottleneck for top- K ranking — it will be impossible to differentiate the K^{th} and $(K + 1)^{\text{th}}$ ranked items unless their score separation exceeds the aggregate error of the corresponding score estimates for these two items. Based on this observation, our algorithm is mainly designed to control the elementwise estimation error. As will be seen in Section 4, the resulting estimation error will be uniformly spread over all entries, which is optimal in both ℓ_2 and ℓ_∞ sense.

- **From Coarse to Detailed Ranking.** The identifiable region we present depends only on the preference separation between items K and $K + 1$. This arises since we only intend to identify the group of top- K items without specifying the fine details within this group—we term it “coarse top- K ranking”. In fact, our results readily uncover the minimax separation requirements for the case where one further expects fine ordering among these K items. Specifically, this task is feasible—in the minimax sense—if and only if

$$\Delta_i \gtrsim \sqrt{\frac{\log n}{np_{\text{obs}}L}}, \quad 1 \leq i \leq K. \quad (12)$$

In words, the feasibility of detailed top- K ranking relies on sufficient score separation between any consecutive pair of the top- K ranked items.

- **High SNR Requirement for Total Ordering.** In many situations, the separation criterion (12) immediately suggests the hardness (or even impossibility) of recovering the ordering over all items. In fact, to figure out the total order, one expects sufficient score separation between all pairs of consecutive items, namely,

$$\Delta_i \gtrsim \sqrt{\frac{\log n}{np_{\text{obs}}L}}, \quad 1 \leq i < n.$$

Since the Δ_i 's are defined in a normalized way (7), they necessarily satisfy

$$\sum_{i=1}^{n-1} \Delta_i = \frac{w_1 - w_n}{w_{\max}} \leq 1.$$

As can be easily verified, the preceding two conditions would be incompatible unless

$$L \gtrsim \frac{n \log n}{p_{\text{obs}}},$$

which imposes a fairly stringent SNR requirement. For instance, under a sparse graph where $p_{\text{obs}} \asymp \frac{\log n}{n}$, the number of repeated comparisons (and hence the SNR) needs to be at least on the order of n^2 , regardless of the method employed. Such a high SNR requirement could be increasingly more difficult to guarantee as n grows.

- **Passive Ranking vs. Active Ranking.** In our passive ranking model, the sample complexity requirement $n^2 p_{\text{obs}} L$ for reliable top- K identification is given by

$$n^2 p_{\text{obs}} L \gtrsim \frac{n \log n}{\Delta_K^2}.$$

In comparison, when adaptive sampling is employed for the preference-based model, the most recent upper bound on the sample complexity (e.g. Theorem 1 of (Busa-Fekete et al., 2013)) is on the order of

$$\sum_{i=1}^{n-1} \frac{1}{\Delta_i^2} \log n.$$

In the challenging regime where a dominant fraction of consecutive pairs are minimally separated (e.g. $\Delta_1 = \dots = \Delta_{n-1}$), the above results seem to suggest that active ranking may not outperform passive ranking, since the sample complexity reads $(n \log n)/\Delta_1^2$. For the other extreme case where only a single pair is minimally separated (e.g. $\Delta_1 \ll \Delta_i$ ($i \geq 2$)), active ranking is more appealing, because it will adaptively acquire more paired evaluation over the minimally separated items instead of wasting samples on those pairs that are easy to differentiate.

4. Ranking Scheme: Spectral Method Meets MLE

This section presents a nearly linear-time algorithm that attempts recovery of the top- K ranked items. The algorithm proceeds in two stages: (1) an appropriate initialization that concentrates around the ground truth in an ℓ_2 sense, which can be obtained via a spectral ranking method; (2) a sequence of iterative updates sharpening the estimates in a pointwise manner, which consists in computing coordinate-wise MLE solutions. The two stages operate upon different sets of samples, while *no further sample splitting* is needed within each stage. The combination of these two stages will be referred to as *Spectral MLE*.

Before continuing to describe the details of our algorithm, we introduce a few notations that will be used throughout.

- $\mathcal{L}(\mathbf{w}; \mathbf{y}_i)$: the likelihood function of a latent preference vector \mathbf{w} , given the part of comparisons \mathbf{y}_i that have bearing on item i .
- $\mathbf{w}_{\setminus i}$: for any preference vector \mathbf{w} , let $\mathbf{w}_{\setminus i}$ represent $[w_1, \dots, w_{i-1}, w_{i+1}, \dots, w_n]$ excluding w_i .
- $\mathcal{L}(\tau, \mathbf{w}_{\setminus i}; \mathbf{y}_i)$: with a slight abuse of notation, denote by $\mathcal{L}(\tau, \mathbf{w}_{\setminus i}; \mathbf{y}_i)$ the likelihood of the preference vector $[w_1, \dots, w_{i-1}, \tau, w_{i+1}, \dots, w_n]$.

4.1 Algorithm: Spectral MLE

It has been established that the spectral ranking method, particularly *Rank Centrality*, is able to discover a preference vector $\hat{\mathbf{w}}$ that incurs minimal ℓ_2 loss. To enable reliable ranking, however, it is more desirable to obtain an estimate that is faithful in an elementwise sense. Fortunately, the solution returned by the spectral method will serve as an ideal initial guess to seed our algorithm. The two components of the proposed Spectral MLE are described below.

Algorithm 1 Spectral MLE.

Input: The average comparison outcome $y_{i,j}$ for all $(i,j) \in \mathcal{E}$; the score range $[w_{\min}, w_{\max}]$.

Partition \mathcal{E} randomly into two sets $\mathcal{E}^{\text{init}}$ and $\mathcal{E}^{\text{iter}}$ each containing $\frac{1}{2}|\mathcal{E}|$ edges. Denote by $\mathbf{y}_i^{\text{init}}$ (resp. $\mathbf{y}_i^{\text{iter}}$) the components of \mathbf{y}_i obtained over $\mathcal{E}^{\text{init}}$ (resp. $\mathcal{E}^{\text{iter}}$).

Initialize $\mathbf{w}^{(0)}$ to be the estimate computed by *Rank Centrality* on $\mathbf{y}_i^{\text{init}}$ ($1 \leq i \leq n$).

Successive Refinement: for $t = 0 : T$ do

1) Compute the coordinate-wise MLE

$$w_i^{\text{mle}} \leftarrow \arg \max_{\tau \in [w_{\min}, w_{\max}]} \mathcal{L} \left(\tau, \mathbf{w}_{\setminus i}^{(t)}; \mathbf{y}_i^{\text{iter}} \right). \quad (13)$$

2) For each $1 \leq i \leq n$, set

$$w_i^{(t+1)} \leftarrow \begin{cases} w_i^{\text{mle}}, & \text{if } |w_i^{\text{mle}} - w_i^{(t)}| > \xi_t, \\ w_i^{(t)}, & \text{else.} \end{cases} \quad (14)$$

Output the indices of the K largest components of $\mathbf{w}^{(T)}$.

Algorithm 2 Rank Centrality (Negahban et al., 2012)

Input: The average comparison outcome $y_{i,j}$ for all $(i,j) \in \mathcal{E}^{\text{iter}}$.

Compute the transition matrix $\mathbf{P} = [P_{i,j}]_{1 \leq i,j \leq n}$ such that

$$P_{i,j} = \begin{cases} \frac{1}{d_{\max}} \frac{y_{j,i}}{y_{i,j} + y_{j,i}}, & \text{if } (i,j) \in \mathcal{E}^{\text{iter}}; \\ 0, & \text{if } i \neq j \text{ and } (i,j) \notin \mathcal{E}^{\text{iter}}; \\ 1 - \frac{1}{d_{\max}} \sum_{k:(i,k) \in \mathcal{E}^{\text{iter}}} \frac{y_{k,i}}{y_{i,k} + y_{k,i}}, & \text{if } i = j. \end{cases}$$

where d_{\max} is the maximum out-degrees of vertices in $\mathcal{E}^{\text{iter}}$.

Output the stationary distribution of \mathbf{P}

1. **Initialization via Spectral Ranking.** We generate an initialization $\mathbf{w}^{(0)}$ via Rank Centrality. In words, Rank Centrality proceeds by constructing a Markov chain based on the pairwise observations, and then returning its stationary distribution by computing the leading eigenvector of the associated probability transition matrix. For the sake of completeness, we provide the detailed procedure of Rank Centrality in Algorithm 2. Under the Erdős–Rényi model, the estimate $\mathbf{w}^{(0)}$ is known to be reasonably faithful in terms of the mean squared loss (Negahban et al., 2012), that is, with high probability,

$$\frac{\|\mathbf{w}^{(0)} - \mathbf{w}\|}{\|\mathbf{w}\|} \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}}.$$

2. **Successive Refinement via Coordinate-wise MLE.** Note that the state-of-the-art finite-sample analyses for MLE (e.g. (Negahban et al., 2012)) involve only the ℓ_2 accuracy of the global MLE when the locations of all samples are i.i.d. (rather than the

graph-based model considered herein). Instead of seeking a global MLE solution, we propose to carefully utilize the coordinate-wise MLE. Specifically, we cyclically iterate through each component, one at a time, maximizing the log-likelihood function with respect to that component. In contrast to the coordinate-descent method for solving the global MLE, we replace the preceding estimate with the new coordinate-wise MLE only when they are far apart. Theorem 8 (to be stated in Section 4.2) guarantees the contraction of the pointwise error for each cycle, which leads to a geometric convergence rate.

The algorithm then returns the indices of top- K items in accordance to the score estimate. A formal and detailed description of the procedure is summarized in Algorithm 1.

Remark 4 *We split \mathcal{E} into $\mathcal{E}^{\text{init}}$ and $\mathcal{E}^{\text{iter}}$ for analytical convenience. Empirically, if we keep $\mathcal{E}^{\text{init}} = \mathcal{E}^{\text{iter}} = \mathcal{E}$ and reuse all samples, then it seems to slightly outperform the sample-splitting procedure. Thus, we recommend the sample-reusing procedure for practical use, and leave the theoretical justification for future work.*

Remark 5 *Spectral MLE is inspired by recent advances in solving non-convex programs by means of iterative methods (Keshavan et al., 2010, 2009; Jain et al., 2013; Candes et al., 2015; Netrapalli et al., 2013; Balakrishnan et al., 2014). A key message conveyed from these works is: once we arrive at an appropriate initialization (often via a spectral method), the iterative estimates will be rapidly attracted towards the global optimum.*

Remark 6 *While our analysis is restricted to the Erdős–Rényi model, Spectral MLE is applicable to general graphs. We caution, however, that spectral ranking is not guaranteed to achieve minimal ℓ_2 loss for general graphs and, in particular, the kind of graphs exhibiting small spectral gaps. Therefore, Spectral MLE is not necessarily minimax optimal under general graph patterns.*

Notably, the successive refinement stage is developed based on the observation that we are able to characterize the confidence intervals of the coordinate-wise MLEs at each iteration. The role of such confidence intervals is to help detect outlier components that incur large pointwise loss. Since the initial guess is optimal in an overall ℓ_2 sense, a large fraction of its entries are already faithful relative to the ground truth. As a consequence, it suffices to disentangle the “sparse” set of outliers.

One appealing feature of Spectral MLE is its low computational complexity. Recall that the initialization step by Rank Centrality can be solved for ϵ accuracy—i.e. identifying an estimate $\hat{\mathbf{w}}$ such that $\|\hat{\mathbf{w}} - \mathbf{w}\|/\|\mathbf{w}\| \leq \epsilon$ —within $O(|\mathcal{E}| \log \frac{1}{\epsilon})$ time instances by means of a power method. In addition, for each component i , the coordinate-wise likelihood function involves a sum of $\deg(i)$ terms. Since finding the coordinate-wise MLE (13) can be cast as a one-dimensional convex program, one can get ϵ accuracy via a bisection method within $O(\deg(i) \cdot \log \frac{1}{\epsilon})$ time. Therefore, each iteration cycle of the successive refinement stage can be accomplished in time $O(|\mathcal{E}| \cdot \log \frac{1}{\epsilon})$.

The following theorem establishes the ranking accuracy of Spectral MLE under the BTL model.

Theorem 7 *Let $c_0, c_1, c_2, c_3 > 0$ be some universal constants. Suppose that $L = O(\text{poly}(n))$, the comparison graph $\mathcal{G} \sim \mathcal{G}_{n, p_{\text{obs}}}$ with $p_{\text{obs}} > c_0 \log n/n$, and the separation measure (7) satisfies*

$$\Delta_K > c_1 \sqrt{\frac{\log n}{np_{\text{obs}}L}}. \quad (15)$$

Then with probability exceeding $1 - 1/n^2$, Spectral MLE perfectly identifies the set of top- K ranked items, provided that the algorithmic parameters obey $T \geq c_2 \log n$ and

$$\xi_t := c_3 \left\{ \xi_{\min} + \frac{1}{2t} (\xi_{\max} - \xi_{\min}) \right\}, \quad (16)$$

where $\xi_{\min} := \sqrt{\frac{\log n}{np_{\text{obs}}L}}$ and $\xi_{\max} := \sqrt{\frac{\log n}{p_{\text{obs}}L}}$.

Theorem 7 basically implies that the proposed algorithm succeeds in separating out the high-ranking objects with probability approaching one, as long as the preference score satisfies the separation condition

$$\Delta_K \gtrsim \sqrt{\frac{\log n}{np_{\text{obs}}L}}.$$

Additionally, Theorem 7 asserts that the number of iteration cycles required in the second stage scales at most logarithmically, revealing that Spectral MLE achieves the desired ranking precision with nearly linear-time computational complexity.

4.2 Successive Refinement: Convergence and Contraction of ℓ_∞ Error

In the sequel, we would like to provide some interpretation as to why we expect the pointwise error of the score estimates to be controllable. The argument is heuristic in nature, since we will assume for simplicity that each iteration employs a fresh set of samples \mathbf{y} independent of the present estimate $\mathbf{w}^{(t)}$.

Denote by $\ell^*(\tau)$ the true log-likelihood function

$$\ell^*(\tau) := \frac{1}{L} \log \mathcal{L}(\tau, \mathbf{w}_{\setminus i}; \mathbf{y}_i). \quad (17)$$

Straightforward calculation suggests that its expectation around w_i can be controlled through a locally strongly-concave function, due to the existence of a second-order lower bound

$$\begin{aligned} \mathbb{E}_{\mathbf{w}} [\ell^*(w_i) - \ell^*(\tau)] &= \sum_{j:(i,j) \in \mathcal{E}} \text{KL} \left(\frac{w_i}{w_i + w_j} \parallel \frac{\tau}{\tau + w_j} \right) \\ &\gtrsim |\tau - w_i|^2 np_{\text{obs}}, \end{aligned} \quad (18)$$

where $\text{KL}(p \parallel q)$ represents the KullbackLeibler (KL) divergence between $\text{Bernoulli}(p)$ and $\text{Bernoulli}(q)$. These calculations will be made precise in Appendix A.1 (and in particular Eqn. (43) and (44)).

This measures the penalty when τ deviates from the ground truth. Note, however, that we don't have direct access to $\ell^*(\cdot)$ since it relies on the latent scores \mathbf{w} . To obtain a computable surrogate, we replace \mathbf{w} with the present estimate $\mathbf{w}^{(t)}$, resulting in the plug-in likelihood function

$$\hat{\ell}_i(\tau) := \frac{1}{L} \log \mathcal{L}(\tau, \mathbf{w}_{\setminus i}^{(t)}; \mathbf{y}_i).$$

Fortunately, the surrogate loss incurred by employing $\hat{\ell}_i(\tau)$ is locally stable in the sense that,

$$\left| \mathbb{E}_{\mathbf{w}} \left[\hat{\ell}_i(\tau) - \hat{\ell}_i(w_i) - (\ell^*(\tau) - \ell^*(w_i)) \right] \right| \lesssim np_{\text{obs}} |\tau - w_i| \frac{\|\hat{\mathbf{w}} - \mathbf{w}\|}{\|\mathbf{w}\|}, \quad (19)$$

which will be made clear in Appendix A.1. This essentially means that the surrogate loss $\hat{\ell}_i(\tau) - \hat{\ell}_i(w_i)$ is a reasonably good approximation of the true loss $\ell^*(\tau) - \ell^*(w_i)$, as long as τ (resp. $\hat{\mathbf{w}}$) is sufficiently close to w_i (resp. \mathbf{w}). As a result, any candidate $\tau \neq w_i$ will be viewed as *less likely* than and hence distinguishable from the ground truth w_i (i.e. $\hat{\ell}(w_i) > \hat{\ell}(\tau)$) in the mean sense, provided that its deviation penalty (18) dominates the surrogate loss (19). This would hold as long as the pointwise loss exceeds the normalized ℓ_2 loss:

$$|\tau - w_i| \gtrsim \frac{\|\hat{\mathbf{w}} - \mathbf{w}\|}{\|\mathbf{w}\|}.$$

Thus, our procedure is expected to be able to converge to a solution whose pointwise error is as low as the normalized ℓ_2 error of the initial guess.

Encouragingly, the ℓ_∞ estimation error not only converges, but converges at a geometric rate as well. This rapid convergence property does not rely on the ‘‘fresh-sample’’ assumption imposed in the above heuristic argument, as formally stated in the following theorem.

Theorem 8 *Suppose that $\mathcal{G} \sim \mathcal{G}_{n, p_{\text{obs}}}$ with $p_{\text{obs}} > c_0 \log n/n$ for some large constant c_0 , and there exists a score vector $\hat{\mathbf{w}}^{\text{ub}} \in [w_{\min}, w_{\max}]^n$ independent of \mathcal{G} satisfying*

$$\left| \hat{w}_i^{\text{ub}} - w_i \right| \leq \xi w_{\max}, \quad 1 \leq i \leq n; \quad (20)$$

$$\|\hat{\mathbf{w}}^{\text{ub}} - \mathbf{w}\| \leq \delta \|\mathbf{w}\|. \quad (21)$$

Then with probability at least $1 - c_1 n^{-4}$ for some constant $c_1 > 0$, the coordinate-wise MLE

$$w_i^{\text{mle}} := \arg \max_{\tau \in [w_{\min}, w_{\max}]} \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) \quad (22)$$

satisfies

$$\left| w_i - w_i^{\text{mle}} \right| < \frac{20 \left(6 + \frac{\log L}{\log n} \right) w_{\max}^5}{w_{\min}^4} \cdot \max \left\{ \delta + \frac{\log n}{np_{\text{obs}}} \cdot \xi, \sqrt{\frac{\log n}{np_{\text{obs}} L}} \right\} \quad (23)$$

simultaneously for all scores $\hat{\mathbf{w}} \in [w_{\min}, w_{\max}]^n$ obeying $|\hat{w}_i - w_i| \leq |\hat{w}_i^{\text{ub}} - w_i|$, $1 \leq i \leq n$.

In the regime where $L = O(\text{poly}(n))$ and $\frac{\|\mathbf{w}^{(t)} - \mathbf{w}\|}{\|\mathbf{w}\|} \asymp \delta \asymp \sqrt{\frac{\log n}{np_{\text{obs}}L}}$, Theorem 8 asserts that under appropriate conditions, the coordinate-wise MLE \mathbf{w}^{mle} is expected to achieve a lower pointwise error than $\mathbf{w}^{(t)}$ such that

$$\|\mathbf{w}^{\text{mle}} - \mathbf{w}\|_{\infty} \lesssim \frac{\|\mathbf{w}^{(t)} - \mathbf{w}\|}{\|\mathbf{w}\|} + \frac{\log n}{np_{\text{obs}}} \|\mathbf{w}^{(t)} - \mathbf{w}\|_{\infty}. \quad (24)$$

When the replacement threshold ξ_t is chosen to be on the same order as $\|\mathbf{w}^{\text{mle}} - \mathbf{w}\|_{\infty}$, one can detect outliers and drag down the elementwise estimation error at a rate

$$\|\mathbf{w}^{(t+1)} - \mathbf{w}\|_{\infty} \lesssim \frac{\|\mathbf{w}^{(t)} - \mathbf{w}\|}{\|\mathbf{w}\|} + \frac{\log n}{np_{\text{obs}}} \|\mathbf{w}^{(t)} - \mathbf{w}\|_{\infty}. \quad (25)$$

One important feature is that the same collection of samples can be *reused* across all iterations at the successive refinement stage, provided that we can identify in each cycle another slightly looser estimate that is independent of the samples. Another property that will be made clear in the analysis is that the ℓ_2 estimation error obeys

$$\frac{\|\mathbf{w}^{(t)} - \mathbf{w}\|}{\|\mathbf{w}\|} \lesssim \frac{\|\mathbf{w}^{(0)} - \mathbf{w}\|}{\|\mathbf{w}\|} \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}}, \quad (26)$$

which further gives

$$\|\mathbf{w}^{(t+1)} - \mathbf{w}\|_{\infty} \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}} + \frac{\log n}{np_{\text{obs}}} \|\mathbf{w}^{(t)} - \mathbf{w}\|_{\infty}. \quad (27)$$

We recognize that the non-negative recursive sequence $\{f_n\}$ obeying the recurrence equation $f_n = a + bf_{n-1}$ ($0 < b < 1$) must converge to a point² $f_{\infty} = \frac{a}{1-b}$. When specialized to (27), this fact implies that the output of Spectral MLE obeys

$$\|\mathbf{w}^{(T)} - \mathbf{w}\|_{\infty} \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}},$$

as long as $\log n/(np_{\text{obs}})$ is sufficiently small and T is sufficiently large. This is minimally apart from the ground truth.

4.3 Discussion

Choice of Initialization. Careful readers will remark that the success of Spectral MLE can be guaranteed by a broader selection of initialization procedures beyond Rank Centrality. Indeed, Theorem 8 and subsequent analyses lead to the following assertion: as long as the initialization method is able to produce an initial estimate $\mathbf{w}^{(0)}$ that is reasonably faithful in the ℓ_2 sense

$$\frac{\|\mathbf{w}^{(0)} - \mathbf{w}\|}{\|\mathbf{w}\|} \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}}, \quad (28)$$

2. To see this, one can rewrite the recurrence inequality as $f_n - \frac{a}{1-b} = b(f_{n-1} - \frac{a}{1-b})$, which gives $f_n - \frac{a}{1-b} = b^n(f_0 - \frac{a}{1-b})$. When n tends to infinity, this gives $f_n - \frac{a}{1-b} = 0$.




Figure 1: (a) Empirical ℓ_∞ loss v.s. L ; (b) ℓ_∞ loss v.s. p_{obs} ; (c) Rate of success in top- K identification ($n = 100, 200$).

then Spectral MLE will converge to a pointwise optimal preference $\mathbf{w}^{(T)}$ obeying

$$\|\mathbf{w}^{(T)} - \mathbf{w}\|_\infty \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}}.$$

Initialization via Global MLE. One would naturally wonder whether we can employ the global MLE (computed over \mathbf{y}^{init}) to seed the iterative refinement stage (applied over \mathbf{y}^{iter}). In fact, the state-of-the-art analysis (with a different but order-wise equivalent model) (Negahban et al., 2012) asserts that the global MLE satisfies the desired ℓ_2 property (28) for at least two cases: (a) complete graphs, i.e. $p_{\text{obs}} = 1$, and (b) Erdős–Rényi graphs with no repeated comparisons, i.e. $L = 1$. In these two cases, the proposed algorithm achieves minimal ℓ_∞ errors if we initialize it via the global MLE.

Nevertheless, whether the global MLE achieves minimal ℓ_2 loss for other configurations (L, p_{obs}) has not been established. The analytical bottleneck seems to stem from an underlying bias-variance tradeoff when accounting for two successive randomness mechanisms: the random graph \mathcal{G} and the repeated comparisons generated over \mathcal{G} . In general, $y_{i,j}^{(l)}$'s are not jointly independent unless we condition on \mathcal{G} . In contrast, the above two special cases amount to two extreme situations: (a) the randomness of \mathcal{G} goes away when $p_{\text{obs}} = 1$; (b) the condition $L = 1$ avoids repeated sampling. Nevertheless, these two cases alone (as well as the model in Theorem 4 of (Negahban et al., 2012)) are not sufficient in characterizing the complete tradeoff between graph sparsity and the quality of the acquired comparisons.

4.4 Numerical Experiments

A series of synthetic experiments is conducted to demonstrate the practical applicability of Spectral MLE. The important implementation parameters in our approach is the choice of c_2 and c_3 given in Theorem 7, which specify T and ξ_t , respectively. In all numerical simulations performed here, we pick $c_2 = 5$ and $c_3 = 1$, and do not split samples. We focus on the case where $n = 100$, where each reported result is calculated by averaging over 200 Monte Carlo trials.

We first examine the ℓ_∞ error of the score estimates. The latent scores are generated uniformly over $[0.5, 1]$. For each (p_{obs}, L) , the paired comparisons are randomly generated as per the BTL model, and we perform score inference by means of both Rank Centrality and Spectral MLE. Fig. 1(a) (resp. Fig. 1(b)) illustrates the empirical tradeoff between the pointwise score estimation accuracy and the number L of repeated comparisons (resp. graph sparsity p_{obs}). It can be seen from these plots that the proposed Spectral MLE outperforms Rank Centrality uniformly over all configurations, corroborating our theoretical results. Interestingly, the performance gain is the most significant under sparse graphs in the presence of low-resolution comparisons (i.e. when p_{obs} and L are small).

Next, we study the success rate of top- K identification as the number n of items varies. We generate the latent scores randomly over $[0.5, 1]$, except that a separation Δ_K is imposed between items K and $K+1$. The results are shown in Fig. 1(c) for the case where $p_{\text{obs}} = 0.2$, and $L = 5$. As can be seen, Spectral MLE achieves higher ranking accuracy compared to Rank Centrality for all these situations. Interestingly, the benefit of Spectral MLE relative to Rank Centrality is more apparent in the regime where the score separation is small. In addition, it seems that Rank Centrality is capable of achieving good ranking accuracy in the randomized model we simulate, and we leave the theoretical analysis for future work.

5. Conclusion

This paper investigates rank aggregation from pairwise data that emphasizes the top- K items. We developed a nearly linear-time algorithm that performs as well as the best model aware paradigm, from a minimax perspective. The proposed algorithm returns the indices of the best- K items in accordance to a carefully tuned preference score estimate, which is obtained by combining a spectral method and a coordinate-wise MLE. Our results uncover the identifiability limit of top- K ranking, which is dictated by the preference separation between the K^{th} and $(K+1)^{\text{th}}$ items.

This paper comes with some limitations in developing tight sample complexity bounds under general graphs. The performances of Spectral MLE under other sampling models are worth investigating (Osting et al., 2015). In addition, it remains to characterize both statistical and computational ranking limits for other choice models (e.g. the Plackett-Luce model (Hajek et al., 2014)). It would also be interesting to consider the case where the paired comparisons are drawn from a mixture of BTL models (e.g. (Oh and Shah, 2014)), as well as the collaborative ranking setting where one aggregates the item preferences from a pool of different users in order to infer rankings for each individual user (e.g. (Lu and Negahban, 2014; Park et al., 2015)).

Acknowledgments

C. Suh was partly supported by the ICT R&D program of MSIP/IITP [B0101-15-1272]. Y. Chen would like to thank Yuejie Chi for helpful discussion.

Appendix A. Performance Guarantees for Spectral MLE

In this section, we establish the theoretical guarantees of Spectral MLE in controlling the ranking accuracy and ℓ_∞ estimation errors, which are the subjects of Theorem 7 and Theorem 8. The proof of Theorem 7 relies heavily on the claim of Theorem 8; for this reason, we present the proofs of Theorem 7 and Theorem 8 in a reverse order. Before proceeding, we recall that the coordinate-wise log-likelihood of τ is given by

$$\frac{1}{L} \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) := \sum_{j:(i,j) \in \mathcal{E}} y_{i,j} \log \frac{\tau}{\tau + \hat{w}_j} + (1 - y_{i,j}) \log \frac{\hat{w}_j}{\tau + \hat{w}_j}, \quad (29)$$

and we shall use $\mathbf{w}_{\setminus i}$ (resp. $\hat{\mathbf{w}}_{\setminus i}$) to denote the vector $\mathbf{w} = [w_1, \dots, w_n]$ (resp. $\hat{\mathbf{w}} = [\hat{w}_1, \dots, \hat{w}_n]$) excluding the entry w_i (resp. \hat{w}_i).

A.1 Proof of Theorem 8

To prove Theorem 8, we need to demonstrate that for every $\tau \in [w_{\min}, w_{\max}]$ that is sufficiently separated from w_i (or, more formally, $|\tau - w_i| \gtrsim \max \left\{ \delta + \frac{\xi \log n}{np_{\text{obs}}}, \sqrt{\frac{\log n}{np_{\text{obs}}L}} \right\}$), the coordinate-wise likelihood satisfies

$$\log \mathcal{L}(w_i, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) > \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) \quad (30)$$

and, therefore, τ cannot be the coordinate-wise MLE.

To begin with, we provide a lemma (which will be proved later) that concerns (30) for *any single* τ that is well separated from w_i .

Lemma 9 *Fix any $\gamma \geq 3$. Under the conditions of Theorem 8, for any $\tau \in [w_{\min}, w_{\max}]$ obeying*

$$|w_i - \tau| > \gamma \cdot \frac{w_{\max}^5}{w_{\min}^4} \max \left\{ \frac{25}{4} \left(\delta + \frac{\xi \log n}{np_{\text{obs}}} \right), 20 \sqrt{\frac{\log n}{np_{\text{obs}}L}} \right\}, \quad (31)$$

one has

$$\frac{1}{L} \log \mathcal{L}(w_i, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) - \frac{1}{L} \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) > \frac{w_{\max}^6}{100w_{\min}^6} \frac{\log n}{L}. \quad (32)$$

with probability exceeding $1 - 6n^{-\gamma}$; this holds simultaneously for all $\hat{\mathbf{w}}_i \in [w_{\min}, w_{\max}]^n$ satisfying $|\hat{w}_i - w_i| \leq |\hat{w}_i^{\text{ub}} - w_i|$, $1 \leq i \leq n$.

To establish Theorem 8, we still need to derive a uniform control over all τ satisfying (31). This will be accomplished via a standard covering argument. Specifically, for any small quantity $\epsilon > 0$, we construct a set \mathcal{N}_ϵ (called an ϵ -cover) within the interval $[w_{\min}, w_{\max}]$ such that for any $\tau \in [w_{\min}, w_{\max}]$, there exists an $\tau_0 \in \mathcal{N}_\epsilon$ obeying

$$|\tau - \tau_0| \leq \epsilon \quad \text{and} \quad |\tau_0 - w_i| \geq |\tau - w_i|. \quad (33)$$

It is self-evident that one can produce such a cover \mathcal{N}_ϵ with cardinality $\lceil \frac{w_{\max}}{\epsilon} \rceil + 1$. If we set $\gamma = 6 + \frac{\log L}{\log n}$ in Lemma 9 (which obeys $\gamma = \Theta(1)$ since $L = O(\text{poly}(n))$), taking the union bound over \mathcal{N}_ϵ gives

$$\frac{1}{L} \log \mathcal{L}(w_i, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) - \frac{1}{L} \log \mathcal{L}(\tau_0, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) > \frac{w_{\max}^6}{100w_{\min}^6} \frac{\log n}{L} \quad (34)$$

simultaneously over all $\tau_0 \in \mathcal{N}_\epsilon$ obeying

$$|w_i - \tau_0| > \frac{\left(6 + \frac{\log L}{\log n}\right) w_{\max}^5}{w_{\min}^4} \max \left\{ \frac{25}{4} \left(\delta + \frac{\xi \log n}{np_{\text{obs}}} \right), 20 \sqrt{\frac{\log n}{np_{\text{obs}} L}} \right\};$$

this occurs with probability at least $1 - 6 |\mathcal{N}_\epsilon| n^{-6 - \frac{\log L}{\log n}}$.

We proceed by bounding the difference between $\log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i)$ and $\log \mathcal{L}(\tau_0, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i)$ for any $|\tau - \tau_0| \leq \epsilon$. To achieve this, we first recognize that the Lipschitz constant of $\frac{1}{L} \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i)$ (cf. (29)) w.r.t. τ is bounded above by the following inequality:

$$\begin{aligned} \frac{1}{L} \cdot \left| \frac{\partial \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i)}{\partial \tau} \right| &= \left| \sum_{j:(i,j) \in \mathcal{E}} y_{i,j} \left(\frac{1}{\tau} - \frac{1}{\tau + \hat{w}_j} \right) - (1 - y_{i,j}) \frac{1}{\tau + \hat{w}_j} \right| \\ &\stackrel{(a)}{\leq} \deg(i) \cdot \frac{2}{w_{\min}} \stackrel{(b)}{\leq} \frac{12 np_{\text{obs}}}{5 w_{\min}}, \end{aligned}$$

where (a) follows since

$$\begin{aligned} \left| y_{i,j} \left(\frac{1}{\tau} - \frac{1}{\tau + \hat{w}_j} \right) - (1 - y_{i,j}) \frac{1}{\tau + \hat{w}_j} \right| &= \left| \frac{y_{i,j}}{\tau} - \frac{1}{\tau + \hat{w}_j} \right| \\ &\leq \left| \frac{y_{i,j}}{\tau} \right| + \left| \frac{1}{\tau + \hat{w}_j} \right| < \frac{2}{w_{\min}}, \end{aligned}$$

and (b) holds since $\deg(i) \leq 2.4np_{\text{obs}}$ with probability $1 - O(n^{-10})$ as long as $\frac{\log n}{np_{\text{obs}}}$ is sufficiently small. As a result, by picking

$$\epsilon = \frac{\frac{w_{\max}^6 \log n}{100 w_{\min}^6 L}}{\frac{12 np_{\text{obs}}}{5 w_{\min}}} = \frac{w_{\max}^6 \log n}{240 w_{\min}^5 np_{\text{obs}} L}, \quad (35)$$

one can make sure that for any $|\tau - \tau_0| \leq \epsilon$,

$$\frac{1}{L} \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) - \frac{1}{L} \log \mathcal{L}(\tau_0, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) \leq \epsilon \cdot \frac{12 np_{\text{obs}}}{5 w_{\min}}, \quad (36)$$

$$\Rightarrow \frac{1}{L} \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) < \frac{1}{L} \log \mathcal{L}(\tau_0, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) + \frac{w_{\max}^6 \log n}{100 w_{\min}^6 L}. \quad (37)$$

In addition, with the above choice of ϵ , the cardinality of the ϵ -cover is bounded above by

$$|\mathcal{N}_\epsilon| \leq \left\lceil \frac{w_{\max}}{\epsilon} \right\rceil + 1 = \left\lceil \frac{240 np_{\text{obs}} L}{\log n} \cdot \frac{w_{\min}^5}{w_{\max}^5} \right\rceil + 1 \ll n^2 L$$

for any sufficiently large n .

Putting (34) and (37) together suggests that for *all* $\tau \in [w_{\min}, w_{\max}]$ sufficiently apart from the ground truth w_i , namely,

$$\forall \tau \in [w_{\min}, w_{\max}]: \quad |\tau - w_i| \geq \frac{\left(6 + \frac{\log L}{\log n}\right) w_{\max}^5}{w_{\min}^4} \max \left\{ \frac{25}{4} \left(\delta + \frac{\xi \log n}{np_{\text{obs}}} \right), 20 \sqrt{\frac{\log n}{np_{\text{obs}} L}} \right\}, \quad (38)$$

one necessarily has

$$\begin{aligned}
 & \frac{1}{L} \log \mathcal{L}(w_i, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) - \frac{1}{L} \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) \\
 &= \left\{ \frac{1}{L} \log \mathcal{L}(w_i, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) - \frac{1}{L} \log \mathcal{L}(\tau_0, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) \right\} + \left\{ \frac{1}{L} \log \mathcal{L}(\tau_0, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) - \frac{1}{L} \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) \right\} \\
 &> 0,
 \end{aligned} \tag{39}$$

with probability at least $1 - 6|\mathcal{N}_\epsilon|n^{-6 - \frac{\log L}{\log n}} - O(n^{-4}) \geq 1 - 6n^2Ln^{-6 - \frac{\log L}{\log n}} - O(n^{-4}) = 1 - O(n^{-4})$. Consequently, any $\tau \in [w_{\min}, w_{\max}]$ that obeys (38) cannot be the coordinate-wise MLE, which in turn justifies the claim (23) of Theorem 8 (we present Theorem 8 using slightly looser constants for notational simplicity).

Proof [of Lemma 9] We start by evaluating the true coordinate-wise likelihood gap

$$\log \mathcal{L}(w_i, \mathbf{w}_{\setminus i}; \mathbf{y}_i) - \log \mathcal{L}(\tau, \mathbf{w}_{\setminus i}; \mathbf{y}_i) \tag{40}$$

for any fixed $\tau \neq w_i$ independent of \mathbf{y}_i . Here, $\mathbf{y}_i := \{y_{i,j} \mid i : (i,j) \in \mathcal{E}\}$ is assumed to be generated under the BTL model parameterized by \mathbf{w} , which clearly obeys

$$\mathbb{E}[y_{i,j}] = \frac{w_i}{w_i + w_j} \quad \text{and} \quad \mathbf{Var}[y_{i,j}] = \frac{1}{L} \frac{w_i w_j}{(w_i + w_j)^2}.$$

In order to calculate the mean of (40), we rewrite the likelihood function as

$$\frac{1}{L} \log \mathcal{L}(\tau, \mathbf{w}_{\setminus i}; \mathbf{y}_i) = \sum_{j:(i,j) \in \mathcal{E}} \left\{ y_{i,j} \log \left(\frac{\tau}{\tau + w_j} \right) + (1 - y_{i,j}) \log \left(\frac{w_j}{\tau + w_j} \right) \right\} \tag{41}$$

$$= \sum_{j:(i,j) \in \mathcal{E}} y_{i,j} \log \left(\frac{\tau}{w_j} \right) + \sum_{j:(i,j) \in \mathcal{E}} \log \left(\frac{w_j}{\tau + w_j} \right). \tag{42}$$

Taking expectation w.r.t. \mathbf{y}_i using the form (41) reveals that

$$\begin{aligned}
 & \mathbb{E} \left[\frac{1}{L} \log \mathcal{L}(w_i, \mathbf{w}_{\setminus i}; \mathbf{y}_i) - \frac{1}{L} \log \mathcal{L}(\tau, \mathbf{w}_{\setminus i}; \mathbf{y}_i) \middle| \mathcal{G} \right] \\
 &= \sum_{j:(i,j) \in \mathcal{E}} \left\{ \frac{w_i}{w_i + w_j} \log \left(\frac{\frac{w_i}{w_i + w_j}}{\frac{\tau}{\tau + w_j}} \right) + \frac{w_j}{w_i + w_j} \log \left(\frac{\frac{w_j}{w_i + w_j}}{\frac{w_j}{\tau + w_j}} \right) \right\} \\
 &= \sum_{j:(i,j) \in \mathcal{E}} \text{KL} \left(\frac{w_i}{w_i + w_j} \parallel \frac{\tau}{\tau + w_j} \right),
 \end{aligned} \tag{43}$$

where $\text{KL}(p||q)$ stands for the KL divergence of Bernoulli(q) from Bernoulli(p). Using Pinsker's inequality (e.g. (Yeung, 2008, Theorem 2.33)), that is, $\text{KL}(p||q) \geq 2(p - q)^2$, we arrive at the following lower bound

$$\begin{aligned}
 & \mathbb{E} \left[\frac{1}{L} \log \mathcal{L}(w_i, \mathbf{w}_{\setminus i}; \mathbf{y}_i) - \frac{1}{L} \log \mathcal{L}(\tau, \mathbf{w}_{\setminus i}; \mathbf{y}_i) \middle| \mathcal{G} \right] \geq 2 \sum_{j:(i,j) \in \mathcal{E}} \left(\frac{w_i}{w_i + w_j} - \frac{\tau}{\tau + w_j} \right)^2 \\
 &= 2(w_i - \tau)^2 \sum_{j:(i,j) \in \mathcal{E}} \frac{w_j^2}{(w_i + w_j)^2 (\tau + w_j)^2}.
 \end{aligned} \tag{44}$$

That being said, the true coordinate-wise likelihood of w_i strictly dominates that of τ in the mean sense.

Nevertheless, when running Spectral MLE, we do not have access to the ground truth scores $\mathbf{w}_{\setminus i}$; what we actually compute is $\mathcal{L}(w_i, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i)$ (resp. $\mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i)$) rather than $\mathcal{L}(\mathbf{w}; \mathbf{y}_i)$ (resp. $\mathcal{L}(\tau, \mathbf{w}_{\setminus i}; \mathbf{y}_i)$). Fortunately, such surrogate likelihoods are sufficiently close to the true coordinate-wise likelihoods, which we will show in the rest of the proof. For brevity, we shall denote respectively the heuristic and true log-likelihood functions by

$$\begin{cases} \hat{\ell}_i(\tau) & := \frac{1}{L} \log \mathcal{L}(\tau, \hat{\mathbf{w}}_{\setminus i}; \mathbf{y}_i) \\ \ell^*(\tau) & := \frac{1}{L} \log \mathcal{L}(\tau, \mathbf{w}_{\setminus i}; \mathbf{y}_i) \end{cases} \quad (45)$$

whenever it is clear from context. Note that $\hat{\mathbf{w}}_{\setminus i}$ could depend on \mathbf{y}_i .

As seen from (42), for any candidate $\tau \in [w_{\min}, w_{\max}]$, we can quantify the difference between $\hat{\ell}_i(\tau)$ and $\ell^*(\tau)$ as

$$\hat{\ell}_i(\tau) - \ell^*(\tau) = \sum_{j:(i,j) \in \mathcal{E}} y_{i,j} \log \left(\frac{w_j}{\hat{w}_j} \right) + \sum_{j:(i,j) \in \mathcal{E}} \left\{ \log \left(\frac{\hat{w}_j}{\tau + \hat{w}_j} \right) - \log \left(\frac{w_j}{\tau + w_j} \right) \right\}. \quad (46)$$

As a consequence, the gap between the true loss $\ell^*(w_i) - \ell^*(\tau)$ and the surrogate loss $\hat{\ell}_i(w_i) - \hat{\ell}_i(\tau)$ is given by

$$\begin{aligned} \hat{\ell}_i(w_i) - \hat{\ell}_i(\tau) - (\ell^*(w_i) - \ell^*(\tau)) &= \hat{\ell}_i(w_i) - \ell^*(w_i) - (\hat{\ell}_i(\tau) - \ell^*(\tau)) \\ &= \sum_{j:(i,j) \in \mathcal{E}} \left\{ \log \left(\frac{\hat{w}_j}{w_i + \hat{w}_j} \right) - \log \left(\frac{w_j}{w_i + w_j} \right) - \left(\log \left(\frac{\hat{w}_j}{\tau + \hat{w}_j} \right) - \log \left(\frac{w_j}{\tau + w_j} \right) \right) \right\} \end{aligned} \quad (47)$$

$$= \sum_{j:(i,j) \in \mathcal{E}} \left\{ \log \left(\frac{\tau + \hat{w}_j}{w_i + \hat{w}_j} \right) - \log \left(\frac{\tau + w_j}{w_i + w_j} \right) \right\}. \quad (48)$$

This gap thus relies on the function

$$g(t) := \log \left(\frac{\tau + t}{w_i + t} \right) - \log \left(\frac{\tau + w_j}{w_i + w_j} \right), \quad t \in [w_{\min}, w_{\max}],$$

which apparently obeys the following two properties: (i) $g(w_j) = 0$; (ii)

$$\left| \frac{\partial g(t)}{\partial t} \right| = \left| \frac{1}{\tau + t} - \frac{1}{w_i + t} \right| = \frac{|\tau - w_i|}{(w_i + t)(\tau + t)} \leq \frac{|\tau - w_i|}{4w_{\min}^2}, \quad \forall t \in [w_{\min}, w_{\max}].$$

Taken together these two properties demonstrate that

$$\begin{aligned} |g(t)| &\leq |g(w_j)| + |t - w_j| \cdot \sup_{t \in [w_{\min}, w_{\max}]} \left| \frac{\partial g(t)}{\partial t} \right| \\ &\leq \frac{1}{4w_{\min}^2} |\tau - w_i| |t - w_j|, \quad \forall t \in [w_{\min}, w_{\max}]. \end{aligned} \quad (49)$$

Substitution into (48) gives

$$\begin{aligned} \left| \hat{\ell}_i(w_i) - \hat{\ell}_i(\tau) - (\ell^*(w_i) - \ell^*(\tau)) \right| &\leq \frac{1}{4w_{\min}^2} |\tau - w_i| \sum_{j:(i,j) \in \mathcal{E}} |\hat{w}_j - w_j| \\ &\leq \frac{1}{4w_{\min}^2} |\tau - w_i| \sum_{j:(i,j) \in \mathcal{E}} \left| \hat{w}_j^{\text{ub}} - w_j \right|. \end{aligned} \quad (50)$$

Notably, this is a deterministic inequality which holds for all \hat{w}_j obeying $|\hat{w}_j - w_j| \leq |\hat{w}_j^{\text{ub}} - w_j|$, $1 \leq j \leq n$. A desired property of the upper bound (50) is that it is independent of \mathcal{G} and the data \mathbf{y}_i , due to our assumption on $\hat{\mathbf{w}}^{\text{ub}}$.

We now move on to develop an upper bound on (50). From our assumptions on the initial estimate, we have

$$\|\hat{\mathbf{w}} - \mathbf{w}\|^2 \leq \|\hat{\mathbf{w}}^{\text{ub}} - \mathbf{w}\|^2 \leq \delta^2 \|\mathbf{w}\|^2 \leq n w_{\max}^2 \delta^2.$$

Since \mathcal{G} and $\hat{\mathbf{w}}^{\text{ub}}$ are statistically independent, this inequality immediately gives rise to the following two consequences:

$$\begin{aligned} \mathbb{E} \left[\sum_{j:(i,j) \in \mathcal{E}} \left| \hat{w}_j^{\text{ub}} - w_j \right| \right] &= p_{\text{obs}} \|\hat{\mathbf{w}}^{\text{ub}} - \mathbf{w}\|_1 \leq p_{\text{obs}} \sqrt{n} \|\hat{\mathbf{w}}^{\text{ub}} - \mathbf{w}\| \\ &\leq n p_{\text{obs}} w_{\max} \delta \end{aligned} \quad (51)$$

and

$$\mathbb{E} \left[\sum_{j:(i,j) \in \mathcal{E}} \left| \hat{w}_j^{\text{ub}} - w_j \right|^2 \right] = p_{\text{obs}} \|\hat{\mathbf{w}}^{\text{ub}} - \mathbf{w}\|_2^2 \leq n p_{\text{obs}} w_{\max}^2 \delta^2. \quad (52)$$

Recall our assumption that $\max_j |\hat{w}_j^{\text{ub}} - w_j| \leq \xi w_{\max}$. For any fixed $\gamma \geq 3$, if $p_{\text{obs}} > \frac{2 \log n}{n}$, then with probability at least $1 - 2n^{-\gamma}$,

$$\begin{aligned} \sum_{j:(i,j) \in \mathcal{E}} \left| \hat{w}_j^{\text{ub}} - w_j \right| &\stackrel{\text{(i)}}{\leq} \mathbb{E} \left[\sum_{j:(i,j) \in \mathcal{E}} \left| \hat{w}_j^{\text{ub}} - w_j \right| \right] + \sqrt{2\gamma \log n \cdot \mathbb{E} \left[\sum_{j:(i,j) \in \mathcal{E}} \left| \hat{w}_j^{\text{ub}} - w_j \right|^2 \right]} \\ &\quad + \frac{2\gamma}{3} \xi w_{\max} \log n \\ &\leq n p_{\text{obs}} w_{\max} \delta + \sqrt{2\gamma \cdot n p_{\text{obs}} \log n} w_{\max} \delta + \frac{2\gamma}{3} \xi w_{\max} \log n \\ &\stackrel{\text{(ii)}}{\leq} (1 + \sqrt{\gamma}) n p_{\text{obs}} w_{\max} \delta + \frac{2\gamma}{3} \xi w_{\max} \log n \\ &\stackrel{\text{(iii)}}{\leq} \gamma n p_{\text{obs}} w_{\max} \delta + \gamma \xi w_{\max} \log n, \end{aligned}$$

where (i) comes from the Bernstein inequality as given in Lemma 11, (ii) follows since $\log n < \frac{p_{\text{obs}} n}{2}$ by assumption, and (iii) arises since $1 + \sqrt{\gamma} \leq \gamma$ whenever $\gamma \geq 3$. This combined with (50) allows us to control

$$\left| \hat{\ell}_i(w_i) - \hat{\ell}_i(\tau) - (\ell^*(w_i) - \ell^*(\tau)) \right| \leq \frac{|\tau - w_i| \gamma w_{\max}}{4w_{\min}^2} (n p_{\text{obs}} \delta + \xi \log n) \quad (53)$$

with high probability.

The above arguments basically reveal that $\hat{\ell}_i(w_i) - \hat{\ell}_i(\tau)$ is reasonably close to $\ell^*(w_i) - \ell^*(\tau)$. Thus, to show that $\hat{\ell}_i(w_i) - \hat{\ell}_i(\tau) > 0$, it is sufficient to develop a lower bound on $\ell^*(w_i) - \ell^*(\tau)$ that exceeds the gap (53). In expectation, the preceding inequality (44) gives

$$\begin{aligned} \mathbb{E}[\ell^*(w_i) - \ell^*(\tau) \mid \mathcal{G}] &\geq 2(w_i - \tau)^2 \sum_{j:(i,j) \in \mathcal{E}} \frac{w_j^2}{(w_i + w_j)^2 (\tau + w_j)^2} \\ &\geq \frac{w_{\min}^2}{8w_{\max}^4} (w_i - \tau)^2 \deg(i). \end{aligned} \quad (54)$$

Recognizing that $y_{i,j} = \frac{1}{L} \sum_{l=1}^L y_{i,j}^{(l)}$ is a sum of independent random variables $y_{i,j}^{(l)} \sim \text{Bernoulli}\left(\frac{w_i}{w_i + w_j}\right)$, we can control the conditional variance as

$$\begin{aligned} \mathbf{Var}[\ell^*(w_i) - \ell^*(\tau) \mid \mathcal{G}] &\stackrel{(a)}{=} \mathbf{Var} \left[\sum_{j:(i,j) \in \mathcal{E}} y_{i,j} \log\left(\frac{w_i}{\tau}\right) \mid \mathcal{G} \right] \\ &= \log^2\left(\frac{w_i}{\tau}\right) \sum_{j:(i,j) \in \mathcal{E}} \frac{1}{L} \frac{w_i w_j}{(w_i + w_j)^2} \stackrel{(b)}{\leq} \frac{1}{L} \frac{(w_i - \tau)^2}{\min\{w_i^2, \tau^2\}} \sum_{j:(i,j) \in \mathcal{E}} \frac{w_{\max}^2}{4w_{\min}^2} \\ &\leq \frac{w_{\max}^2}{4w_{\min}^4} \cdot \frac{1}{L} (w_i - \tau)^2 \deg(i), \end{aligned} \quad (55)$$

where (a) is an immediate consequence of (42), and (b) follows since $\left| \log \frac{\beta}{\alpha} \right| \leq \frac{\beta - \alpha}{\alpha}$ for any $\beta > \alpha > 0$. Note that $0 \leq \frac{1}{L} y_{i,j}^{(l)} \leq \frac{1}{L}$, and hence each summand of $\ell^*(w_i) - \ell^*(\tau)$ (written in terms of a weighted sum of $y_{i,j}^{(l)}$) is bounded in magnitude by

$$\max_{i,j,l} \frac{1}{L} y_{i,j}^{(l)} \left| \log \frac{w_i}{\tau} \right| \leq \frac{1}{L} \left| \log \frac{w_i}{\tau} \right| \leq \frac{1}{L} \frac{|w_i - \tau|}{w_{\min}}, \quad (56)$$

where the last inequality follows again from the inequality $\left| \log \left(\frac{\beta}{\alpha} \right) \right| \leq \frac{\beta - \alpha}{\alpha}$ for any $\beta \geq \alpha > 0$. Making use of the Bernstein inequality together with (54)-(56) suggests that: conditional on \mathcal{G} ,

$$\begin{aligned} &\ell^*(w_i) - \ell^*(\tau) \\ &\geq \mathbb{E}[\ell^*(w_i) - \ell^*(\tau) \mid \mathcal{G}] - \sqrt{2\gamma \mathbf{Var}[\ell^*(w_i) - \ell^*(\tau) \mid \mathcal{G}] \log n} - \frac{2\gamma \log n}{3} \cdot \frac{\left| \log \left(\frac{w_i}{\tau} \right) \right|}{L} \\ &\geq \frac{w_{\min}^2}{8w_{\max}^4} (w_i - \tau)^2 \deg(i) - \frac{\sqrt{2\gamma} w_{\max} |w_i - \tau|}{2w_{\min}^2} \sqrt{\frac{\deg(i) \log n}{L}} - \frac{2\gamma |w_i - \tau| \log n}{3Lw_{\min}} \end{aligned} \quad (57)$$

holds with probability at least $1 - 2n^{-\gamma}$. The above bound relies on $\deg(i)$, which is on the order of np_{obs} with high probability. More precisely, taking the Chernoff bound (Mitzenmacher and Upfal, 2005, Corollary 4.6) as well as the union bound reveals that: if $\frac{\log n}{np_{\text{obs}}}$ is sufficiently large, then

$$\frac{4}{5} np_{\text{obs}} < \deg(i) < \frac{6}{5} np_{\text{obs}}, \quad 1 \leq i \leq n \quad (58)$$

with probability at least $1 - 2n^{-\gamma}$. This taken collectively with (57) and the assumption $np_{\text{obs}} > 2 \log n$ implies that

$$\begin{aligned}
 \ell^*(w_i) - \ell^*(\tau) &\geq \frac{w_{\min}^2}{8w_{\max}^4} (w_i - \tau)^2 \cdot \frac{4}{5} np_{\text{obs}} - \sqrt{\frac{\gamma}{2}} \frac{w_{\max} |w_i - \tau|}{w_{\min}^2} \sqrt{\frac{6np_{\text{obs}} \log n}{5L}} - \frac{2\gamma |w_i - \tau| \log n}{3Lw_{\min}} \\
 &\geq \frac{w_{\min}^2}{10w_{\max}^4} (w_i - \tau)^2 np_{\text{obs}} - \left(\sqrt{\frac{3\gamma}{5}} + \frac{2\gamma}{3} \frac{1}{\sqrt{2}} \right) \frac{w_{\max} |w_i - \tau|}{w_{\min}^2} \sqrt{\frac{np_{\text{obs}} \log n}{L}} \\
 &\geq \frac{w_{\min}^2}{10w_{\max}^4} (w_i - \tau)^2 np_{\text{obs}} - \gamma \frac{w_{\max} |w_i - \tau|}{w_{\min}^2} \sqrt{\frac{np_{\text{obs}} \log n}{L}} \tag{59} \\
 &\geq \frac{w_{\min}^2}{20w_{\max}^4} (w_i - \tau)^2 np_{\text{obs}} \tag{60}
 \end{aligned}$$

with probability at least $1 - 4n^{-\gamma}$, as long as

$$\gamma \cdot \frac{w_{\max} |w_i - \tau|}{w_{\min}^2} \sqrt{\frac{np_{\text{obs}} \log n}{L}} \leq \frac{w_{\min}^2}{20w_{\max}^4} (w_i - \tau)^2 np_{\text{obs}}$$

or, equivalently,

$$|w_i - \tau| \geq \frac{20\gamma \cdot w_{\max}^5}{w_{\min}^4} \sqrt{\frac{\log n}{np_{\text{obs}} L}}. \tag{61}$$

Finally, we are ready to control $\hat{\ell}_i(w_i) - \hat{\ell}_i(\tau)$ from below. Putting (53) and (60) together, we see that with high probability,

$$\begin{aligned}
 \hat{\ell}_i(w_i) - \hat{\ell}_i(\tau) &\geq \ell^*(w_i) - \ell^*(\tau) - \frac{|\tau - w_i| \gamma w_{\max} (np_{\text{obs}} \delta + \xi \log n)}{4w_{\min}^2} \\
 &\geq \frac{w_{\min}^2}{20w_{\max}^4} (w_i - \tau)^2 np_{\text{obs}} - \frac{|\tau - w_i| \gamma w_{\max}}{4w_{\min}^2} (np_{\text{obs}} \delta + \xi \log n) \\
 &> \frac{w_{\min}^2}{100w_{\max}^4} (w_i - \tau)^2 np_{\text{obs}} \tag{62}
 \end{aligned}$$

$$> \frac{w_{\max}^6}{100w_{\min}^6} \frac{\log n}{L}, \tag{63}$$

where (62) holds under the condition

$$|\tau - w_i| > \frac{25\gamma w_{\max}^5}{4w_{\min}^4} \left(\delta + \frac{\xi \log n}{np_{\text{obs}}} \right), \tag{64}$$

and (63) follows from the assumption (61). The claim (32) is then established under the conditions (61) and (64). \blacksquare

A.2 Proof of Theorem 7

The accuracy of top- K identification is closely related to the ℓ_∞ error of the score estimate. In the sequel, we shall assume that $w_{\max} = 1$ to simplify presentation. Our goal is to demonstrate that

$$\|\mathbf{w}^{(t)} - \mathbf{w}\|_\infty \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}} + \frac{1}{2^t} \sqrt{\frac{\log n}{p_{\text{obs}}L}} \asymp \xi_t, \quad \forall t \in \mathbb{N}, \quad (65)$$

where

$$\xi_t := c_3 \left\{ \xi_{\min} + \frac{1}{2^t} (\xi_{\max} - \xi_{\min}) \right\}, \quad \forall t \geq -1 \quad (66)$$

with $\xi_{\min} = \sqrt{\frac{\log n}{np_{\text{obs}}L}}$ and $\xi_{\max} = \sqrt{\frac{\log n}{p_{\text{obs}}L}}$. If $T \geq c_2 \log n$ for some sufficiently large $c_2 > 0$, then this gives

$$\|\mathbf{w}^{(T)} - \mathbf{w}\|_\infty \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}} = \xi_{\min}.$$

The key implication is the following: if $w_K - w_{K-1} \geq c_1 \sqrt{\frac{\log n}{np_{\text{obs}}L}}$ for some sufficiently large $c_1 > 0$, then

$$\begin{aligned} w_i^{(T)} - w_j^{(T)} &\geq w_i - w_j - |w_i^{(T)} - w_i| - |w_j^{(T)} - w_j| \\ &\geq w_K - w_{K+1} - 2\|\mathbf{w}^{(T)} - \mathbf{w}\|_\infty > 0 \end{aligned}$$

for all $1 \leq i \leq K$ and $j \geq K+1$, indicating that Spectral MLE will output the first K items as desired. The remaining proof then comes down to showing (65).

We start from $t = 0$. When the initial estimate $\mathbf{w}^{(0)}$ is computed by Rank Centrality, the ℓ_2 estimation error satisfies (Negahban et al., 2012)

$$\frac{\|\mathbf{w}^{(0)} - \mathbf{w}\|}{\|\mathbf{w}\|} \leq c_4 \sqrt{\frac{\log n}{np_{\text{obs}}L}} = c_4 \xi_{\min} := \delta \quad (67)$$

with high probability, where $c_4 > 0$ is some universal constant independent of n, p_{obs}, L and Δ_K . A by-product of this result is an upper bound

$$\|\mathbf{w}^{(0)} - \mathbf{w}\|_\infty \leq \|\mathbf{w}^{(0)} - \mathbf{w}\| \leq \delta \|\mathbf{w}\| \leq \delta \sqrt{n} = c_4 \sqrt{\frac{\log n}{p_{\text{obs}}L}}, \quad (68)$$

which together with the fact $\|\mathbf{w}^{(0)} - \mathbf{w}\|_\infty \leq w_{\max} - w_{\min} \leq 1$ gives

$$\|\mathbf{w}^{(0)} - \mathbf{w}\|_\infty \leq \min \left\{ c_4 \sqrt{\frac{\log n}{p_{\text{obs}}L}}, 1 \right\} = \min \{c_4 \xi_{\max}, 1\}. \quad (69)$$

This justifies that $\mathbf{w}^{(0)}$ satisfies the claim (65). Notably, $\mathbf{w}^{(0)}$ is independent of $\mathcal{E}^{\text{iter}}$ and \mathbf{y}^{iter} and, therefore, independent of the iterative steps.

In what follows, we divide the iterative stage into two phases: (1) $t \leq T_0$ and (2) $t > T_0$, where T_0 is a threshold such that

$$\xi_t \geq c_{10}\xi_{\min} = c_{10}\sqrt{\frac{\log n}{np_{\text{obs}}L}}, \quad \text{iff } t \leq T_0, \quad (70)$$

for some large constant $c_{10} > 0$. As is seen from the definition of ξ_t , $T_0 \lesssim \log n$ holds as long as $L = O(\text{poly}(n))$.

For the case where $t \leq T_0$, we proceed by induction on t w.r.t. the following hypotheses:

- \mathcal{M}_t : $\|\mathbf{w}^{\text{mle}} - \mathbf{w}\|_{\infty} < \frac{1}{2}\xi_t$ holds at the t^{th} iteration (the iteration where we compute $\mathbf{w}^{(t+1)}$);
- \mathcal{B}_t : all entries $w_i^{(\tau)}$ of $\mathbf{w}^{(\tau)}$ ($\tau \leq t-1$) satisfying $|w_i^{(\tau)} - w_i| \geq 1.5\xi_t$ have been replaced by time t ;
- \mathcal{H}_t : none of the entries $w_i^{(\tau)}$ ($\tau \leq t-1$) satisfying $|w_i^{(\tau)} - w_i| \leq \frac{1}{2}\xi_t$ have been replaced by time t .

We first note that \mathcal{B}_t is an immediate consequence of \mathcal{M}_t and \mathcal{B}_{t-1} . In fact, given \mathcal{B}_{t-1} , it suffices to examine those entries $w_i^{(\tau)}$ that have *not* been replaced by time $t-1$. To this end, we recall that Spectral MLE replaces $w_i^{(\tau)}$ if $|w_i^{(\tau)} - w_i^{\text{mle}}| > \xi_t$. With \mathcal{M}_t in place, for each i obeying $|w_i^{(\tau)} - w_i| \geq 1.5\xi_t$, one has

$$|w_i^{(\tau)} - w_i^{\text{mle}}| \geq |w_i^{(\tau)} - w_i| - |w_i^{\text{mle}} - w_i| > 1.5\xi_t - \frac{1}{2}\xi_t = \xi_t$$

and hence it will necessarily be replaced by w_i^{mle} at time t . Similarly, \mathcal{H}_t is an immediate consequence of \mathcal{M}_t and \mathcal{H}_{t-1} .³ As a consequence, it boils down to verifying \mathcal{M}_t .

When $t = 0$, applying Theorem 8 and setting $\hat{\mathbf{w}}^{\text{ub}} = \mathbf{w}^{(0)}$, we see that

$$\|\mathbf{w}^{\text{mle}} - \mathbf{w}\|_{\infty} \leq c_7\xi_{\min} + c_9\frac{\log n}{np_{\text{obs}}}\xi_{\max}$$

for some universal constants $c_7, c_9 > 0$, where we have made use of the properties (67) and (69). When c_{10} is sufficiently large, the definition of T_0 (cf. (70)) gives $\xi_0 \gg c_7\sqrt{\frac{\log n}{np_{\text{obs}}L}}$; additionally, $c_9\frac{\log n}{np_{\text{obs}}}\xi_{\max} \ll \xi_{\max} \leq \xi_0$ holds as long as $\frac{\log n}{np_{\text{obs}}}$ is sufficiently small. Putting these conditions together gives

$$\|\mathbf{w}^{\text{mle}} - \mathbf{w}\|_{\infty} \leq c_7\xi_{\min} + c_9c_4\frac{\log n}{np_{\text{obs}}}\xi_{\max} < \frac{1}{2}\xi_0,$$

which verifies the property \mathcal{M}_0 .

3. Given \mathcal{M}_t and \mathcal{H}_{t-1} , for any i obeying $|w_i^{(\tau)} - w_i| \leq 0.5\xi_t$, one has

$$|w_i^{(\tau)} - w_i^{\text{mle}}| \leq |w_i^{(\tau)} - w_i| + |w_i^{\text{mle}} - w_i| < \frac{1}{2}\xi_t + \frac{1}{2}\xi_t = \xi_t$$

and, therefore, it cannot be replaced by time t , which establishes \mathcal{H}_t .

We now turn to extending these inductive hypotheses to the t^{th} iteration, assuming that all of them hold up to time $t - 1$. Taken together \mathcal{M}_{t-1} and \mathcal{B}_{t-1} immediately reveal that

$$\left\| \mathbf{w}^{(t)} - \mathbf{w} \right\|_{\infty} \leq 1.5\xi_{t-1}. \quad (71)$$

In order to invoke Theorem 8 for the coordinate-wise MLEs, we need to construct a looser auxiliary score estimate $\hat{\mathbf{w}}^{\text{ub}}$. With \mathcal{B}_{t-1} , \mathcal{H}_{t-1} and (71) in mind, we propose a candidate for the t^{th} iteration as follows⁴

$$\hat{w}_i^{\text{ub}} = \begin{cases} w_i + 1.5\xi_{t-1}, & \text{if } |w_i^{(0)} - w_i| > \frac{1}{2}\xi_{t-1}, \\ w_i^{(0)} & \text{else.} \end{cases} \quad (72)$$

which is clearly independent of $\mathcal{E}^{\text{iter}}$ and \mathbf{y}^{iter} . According to \mathcal{B}_{t-1} and \mathcal{H}_{t-1} , (i) none of the entries $w_i^{(0)}$ with $|w_i^{(0)} - w_i| \leq \frac{1}{2}\xi_{t-1}$ have been replaced so far; (ii) if an entry $w_i^{(0)}$ has ever been replaced, then the error of the new iterate cannot exceed $1.5\xi_{t-1}$ (otherwise it'll be replaced by the MLE in time $t - 1$ which gives an error below $0.5\xi_{t-1}$). As a result, $\hat{\mathbf{w}}^{\text{ub}}$ satisfies

$$\left| w_i^{(t)} - w_i \right| \leq \left| \hat{w}_i^{\text{ub}} - w_i \right| \leq 1.5\xi_{t-1}, \quad (73)$$

$$\text{and } \left\| \mathbf{w}^{(t)} - \mathbf{w} \right\| \leq \left\| \mathbf{w}^{(\text{ub})} - \mathbf{w} \right\| \stackrel{\text{(a)}}{\leq} 3 \left\| \mathbf{w}^{(0)} - \mathbf{w} \right\| \leq 3\delta \|\mathbf{w}\|. \quad (74)$$

Here, (a) arises since: (1) due to \mathcal{H}_{t-1} , if $w_i^{(0)}$ is ever replaced, then $|w_i^{(0)} - w_i|$ is at least $0.5\xi_{t-1}$; (2) by construction, the pointwise error of $\hat{\mathbf{w}}^{\text{ub}}$ is at most $1.5\xi_{t-1}$, and hence the replacement cannot inflate the original error $|w_i^{(0)} - w_i|$ by more than $\frac{1.5\xi_{t-1}}{0.5\xi_{t-1}} = 3$ times. With these in place, applying Theorem 8 gives

$$\left\| \mathbf{w}^{\text{mle}} - \mathbf{w} \right\|_{\infty} \leq c_8\xi_{\min} + 1.5c_9 \frac{\log n}{np_{\text{obs}}} \xi_{t-1},$$

which relies on the fact $\delta \lesssim \sqrt{\frac{\log n}{np_{\text{obs}}L}}$. Recognize that

$$\xi_t \gg c_8\xi_{\min} \quad \text{and} \quad 1.5c_9 \frac{\log n}{np_{\text{obs}}} \xi_{t-1} \ll \xi_t$$

hold in the regime where $t \leq T_0$ and $\frac{\log n}{np_{\text{obs}}} \ll 1$, which taken together give

$$\left\| \mathbf{w}^{\text{mle}} - \mathbf{w} \right\|_{\infty} < \frac{1}{2}\xi_t$$

as claimed in \mathcal{M}_t . Having verified these inductive hypotheses, we see from the above argument that in any event, the ℓ_{∞} error bound at the t^{th} iteration is at most $1.5\xi_t$, which in turn leads to the claim (65) for any $t \leq T_0$.

4. Careful readers will note that when $|w_i^{(0)} - w_i| \geq \frac{1}{2}\xi_{t-1}$, the resulting \hat{w}_i^{ub} might exceed the range $[w_{\min}, w_{\max}]$. This can be easily addressed if we do the following: (1) change \hat{w}_i^{ub} to $w_i - 1.5\xi_{t-1}$ instead if $w_i - 1.5\xi_{t-1} \in [w_{\min}, w_{\max}]$; (2) if it is still infeasible, set \hat{w}_i^{ub} to be w_{\max} if $|w_i - w_{\max}| > |w_i - w_{\min}|$ and w_{\min} otherwise. For simplicity of presentation, however, we omit these boundary situations and assume $w_i + 1.5\xi_{t-1} \leq w_{\max}$ throughout, which will not change the results anyway.

Starting from $t = T_0 + 1$, we fix the auxiliary score as follows

$$\hat{w}_i^{\text{ub}} = \begin{cases} w_i + 1.5\xi_{T_0}, & \text{if } |w_i^{(0)} - w_i| > \frac{1}{2}\xi_\infty, \\ w_i^{(0)} & \text{else,} \end{cases} \quad (75)$$

where we recall that $\xi_\infty = c_3\xi_{\min}$ and $\xi_{T_0} = c_{10}\xi_{\min}$. This satisfies

$$|w_i^{(t)} - w_i| \leq |\hat{w}_i^{\text{ub}} - w_i| \leq 1.5\xi_{T_0}$$

for $t = T_0 + 1$, due to \mathcal{M}_{T_0} and \mathcal{B}_{T_0} . Moreover, the number of indices that satisfy $|w_i^{(0)} - w_i| > \frac{1}{2}\xi_\infty$, denoted by k , obeys

$$k \cdot \left(\frac{1}{2}\xi_\infty\right)^2 \leq \|\mathbf{w} - \mathbf{w}^{(0)}\|^2 \leq \delta^2\|\mathbf{w}\|^2 \iff k \leq \frac{4\delta^2\|\mathbf{w}\|^2}{\xi_\infty^2},$$

which further gives

$$\begin{aligned} \|\mathbf{w}^{\text{ub}} - \mathbf{w}\|^2 &\leq \|\mathbf{w}^{(0)} - \mathbf{w}\|^2 + \sum_{i: |w_i^{(0)} - w_i| > \frac{1}{2}\xi_\infty} (1.5\xi_{T_0})^2 \leq \delta^2\|\mathbf{w}\|^2 + 2.25k\xi_{T_0}^2 \\ &\leq \delta^2\|\mathbf{w}\|^2 \left(1 + \frac{9\xi_{T_0}^2}{\xi_\infty^2}\right). \end{aligned}$$

Note that the preceding analysis does not depend on the ratio $\frac{c_{10}}{c_3}$ as long as both c_3 and c_{10} are large. If we pick $\frac{c_{10}}{c_3} = \frac{\xi_{T_0}}{\xi_\infty} \leq \sqrt{2}$, then the above inequality gives rise to

$$\|\mathbf{w}^{\text{ub}} - \mathbf{w}\| \leq \sqrt{19}\delta\|\mathbf{w}\|.$$

Applying Theorem 8 we deduce

$$\|\mathbf{w}^{\text{mle}} - \mathbf{w}\|_\infty \lesssim \delta + \frac{\log n}{np_{\text{obs}}}\xi_{T_0} + \sqrt{\frac{\log n}{np_{\text{obs}}L}} \asymp \sqrt{\frac{\log n}{np_{\text{obs}}L}} < \frac{1}{2}\xi_\infty,$$

as long as $\frac{\log n}{p_{\text{obs}}n}$ is small and c_{10}, c_3 are sufficiently large.

The main point of the above calculation is that: for any entry $w_i^{(0)}$ satisfying $|w_i^{(0)} - w_i| < \frac{1}{2}\xi_\infty$, one must have

$$|w_i^{(0)} - w_i^{\text{mle}}| \leq |w_i^{(0)} - w_i| + |w_i^{(\text{mle})} - w_i| < \xi_\infty < \xi_t,$$

and hence it will not be replaced. As a result, the auxiliary score (75) remains valid for the iteration that follows. In fact, these properties continue to hold for all $t > T_0$ if we repeat the same argument as t increases. To finish up, put together the above arguments to obtain

$$\|\mathbf{w}^{(t)} - \mathbf{w}\|_\infty \leq \frac{1}{2}\xi_\infty \asymp \sqrt{\frac{\log n}{np_{\text{obs}}L}}, \quad t > T_0,$$

which establishes the claim (65) for $t > T_0$ and, in turn, Theorem 7.

Appendix B. Proof of the Minimax Lower Bound (Theorem 3)

This section establishes the minimax lower limit given in Theorem 3. To bound the minimax probability of error, we proceed by constructing a finite set of hypotheses, followed by an analysis based on classical Fano-type arguments. For notational simplicity, each hypothesis is represented by a permutation σ over $[n]$, and we denote by $\sigma(i)$ and $\sigma([K])$ the corresponding index of the i^{th} ranked item and the index set of all top- K items, respectively.

We now single out a set of hypotheses and some prior to be imposed on them. Suppose that the values of \mathbf{w} are fixed up to permutation in such a way that

$$w_{\sigma(i)} = \begin{cases} w_K, & 1 \leq i \leq K, \\ w_{K+1}, & K < i \leq n, \end{cases}$$

where we abuse the notation w_K, w_{K+1} to represent any two values satisfying

$$\frac{w_K - w_{K+1}}{w_{\max}} = \Delta_K > 0.$$

Below we suppose that the ranking scheme is informed of the values w_K, w_{K+1} , which only makes the ranking task easier. In addition, we impose a uniform prior over a collection \mathcal{M} of $M := \max\{K, n - K\} + 1$ hypotheses regarding the permutation: if $K < n/2$, then

$$\mathbb{P}\{\sigma([K]) = \mathcal{S}\} = \frac{1}{M}, \quad \text{for } \mathcal{S} = \{2, \dots, K\} \cup \{i\}, \quad (i = 1, K + 1, \dots, n); \quad (76)$$

if $K \geq n/2$, then

$$\mathbb{P}\{\sigma([K]) = \mathcal{S}\} = \frac{1}{M}, \quad \text{for } \mathcal{S} = \{1, \dots, K + 1\} \setminus \{i\}, \quad (i = 1, \dots, K + 1). \quad (77)$$

In words, each alternative hypothesis is generated by swapping two indices of the hypothesis obeying $\sigma([K]) = [K]$. Denoting by $P_{e,M}$ the average probability of error with respect to the prior we construct, one can easily verify that the minimax probability of error is at least $P_{e,M}$.

This Bayesian probability of error will be bounded using classical Fano-type bounds. To accommodate partial observation, we introduce an erased version of $\mathbf{y}_{i,j} := (y_{i,j}^{(1)}, \dots, y_{i,j}^{(L)})$ such that

$$\mathbf{z}_{i,j} = \begin{cases} \mathbf{y}_{i,j}, & \text{with probability } p_{\text{obs}}, \\ \text{erasure}, & \text{else,} \end{cases}$$

and set $\mathbf{Z} := \{z_{i,j}\}_{1 \leq i < j \leq n}$. Applying the generalized Fano inequality (Han and Verdu, 1994, Theorem 7) gives

$$\begin{aligned}
 P_{e,M} &\geq 1 - \frac{1}{\log M} \left\{ \frac{1}{M^2} \sum_{\sigma_1, \sigma_2 \in \mathcal{M}} \text{KL} \left(\mathbb{P}_{\mathbf{Z}|\sigma=\sigma_1} \parallel \mathbb{P}_{\mathbf{Z}|\sigma=\sigma_2} \right) + \log 2 \right\} \\
 &\stackrel{(a)}{=} 1 - \frac{1}{\log M} \left\{ \frac{1}{M^2} \sum_{\sigma_1, \sigma_2 \in \mathcal{M}} \sum_{i \neq j} \text{KL} \left(\mathbb{P}_{z_{i,j}|\sigma=\sigma_1} \parallel \mathbb{P}_{z_{i,j}|\sigma=\sigma_2} \right) + \log 2 \right\} \\
 &\stackrel{(b)}{=} 1 - \frac{1}{\log M} \left\{ \frac{p_{\text{obs}}}{M^2} \sum_{\sigma_1, \sigma_2 \in \mathcal{M}} \sum_{i \neq j} \text{KL} \left(\mathbb{P}_{\mathbf{y}_{i,j}|\sigma=\sigma_1} \parallel \mathbb{P}_{\mathbf{y}_{i,j}|\sigma=\sigma_2} \right) + \log 2 \right\} \\
 &\stackrel{(c)}{=} 1 - \frac{1}{\log M} \left\{ \frac{p_{\text{obs}}L}{M^2} \sum_{\sigma_1, \sigma_2 \in \mathcal{M}} \sum_{i \neq j} \text{KL} \left(\mathbb{P}_{y_{i,j}^{(1)}|\sigma=\sigma_1} \parallel \mathbb{P}_{y_{i,j}^{(1)}|\sigma=\sigma_2} \right) + \log 2 \right\} \\
 &\stackrel{(d)}{\geq} 1 - \frac{1}{\log M} \left\{ \frac{2w_{\max}^4}{w_{\min}^4} np_{\text{obs}}L\Delta_K^2 + \log 2 \right\},
 \end{aligned}$$

where $\text{KL}(P \parallel Q)$ denotes the KL divergence of Q from P . Here, (a) comes from the independence assumption of the $z_{i,j}$'s; (b) arises since $z_{i,j}$ is an erased version of $\mathbf{y}_{i,j}$; (c) follows since $y_{i,j}^{(\ell)}$ ($1 \leq \ell \leq L$) are i.i.d.; and (d) arises from Lemma 10 (see below).

Consequently, one would have $P_e \geq P_{e,M} \geq \epsilon$ if

$$\frac{2w_{\max}^4}{w_{\min}^4} np_{\text{obs}}L\Delta_K^2 \leq (1 - \epsilon) \log M - \log 2.$$

Since $|\mathcal{M}| = M \geq \frac{n}{2}$, the above condition is necessarily satisfied when

$$\frac{2w_{\max}^4}{w_{\min}^4} np_{\text{obs}}L\Delta_K^2 \leq (1 - \epsilon) \log n - 2 \quad \iff \quad L \leq \frac{w_{\min}^4}{2w_{\max}^4} \cdot \frac{(1 - \epsilon) \log n - 2}{np_{\text{obs}}\Delta_K^2},$$

which finishes the proof.

Lemma 10 *If $w_K, w_{K+1} \in [w_{\min}, w_{\max}]$, then for any $\sigma_1, \sigma_2 \in \mathcal{M}$:*

$$\sum_{i \neq j} \text{KL} \left(\mathbb{P}_{y_{i,j}^{(1)}|\sigma=\sigma_1} \parallel \mathbb{P}_{y_{i,j}^{(1)}|\sigma=\sigma_2} \right) \leq \frac{2w_{\max}^4}{w_{\min}^4} n\Delta_K^2. \quad (78)$$

Proof To start with, for any two measures $P \sim \text{Bernoulli}(p)$ and $Q \sim \text{Bernoulli}(q)$, one has (van Erven and Harremoës, 2014, Eqn. (7))

$$\text{KL}(P \parallel Q) \leq \chi^2(P \parallel Q) = \frac{(p-q)^2}{q} + \frac{(p-q)^2}{1-q} = \frac{(p-q)^2}{q(1-q)}. \quad (79)$$

where $\chi^2(P \parallel Q)$ denotes the χ^2 divergence between P and Q .

Recall that given $\sigma = \sigma_1$ (resp. $\sigma = \sigma_2$), $y_{i,j}^{(1)}$ is Bernoulli distributed with mean $r_1 := \frac{w_{\sigma_1(i)}}{w_{\sigma_1(i)} + w_{\sigma_1(j)}}$ (resp. $r_2 := \frac{w_{\sigma_2(i)}}{w_{\sigma_2(i)} + w_{\sigma_2(j)}}$). If we set $\delta = r_1 - r_2$, then (79) yields

$$\text{KL} \left(\mathbb{P}_{y_{i,j}^{(1)} | \sigma = \sigma_1} \parallel \mathbb{P}_{y_{i,j}^{(1)} | \sigma = \sigma_2} \right) \leq \frac{\delta^2}{r_2(1-r_2)} \leq \frac{4w_{\max}^2 \delta^2}{w_{\min}^2},$$

where the last inequality follows since

$$r_2(1-r_2) = \frac{w_{\sigma_2(i)}w_{\sigma_2(j)}}{(w_{\sigma_2(i)} + w_{\sigma_2(j)})^2} \geq \frac{w_{\min}^2}{4w_{\max}^2}.$$

By construction, conditional on any hypotheses $\sigma_1, \sigma_2 \in \mathcal{M}$, the distributions of $\mathbf{y}_{i,j}$ are different over at most $2n$ locations. For each of these $O(n)$ locations, our construction of \mathcal{M} ensures that

$$|\delta| = |r_2 - r_1| \leq \frac{w_K}{w_K + w_{K+1}} - \frac{w_{K+1}}{w_K + w_{K+1}} = \frac{w_K - w_{K+1}}{w_K + w_{K+1}} \leq \frac{w_{\max}}{2w_{\min}} \Delta_K.$$

As a result, the total contribution is bounded above by

$$\sum_{i \neq j} \text{KL} \left(\mathbb{P}_{y_{i,j}^{(1)} | \sigma = \sigma_1} \parallel \mathbb{P}_{y_{i,j}^{(1)} | \sigma = \sigma_2} \right) \leq 2n \cdot \left(\max_{i,j} \delta^2 \right) \frac{4w_{\max}^2}{w_{\min}^2} \leq \frac{2w_{\max}^4}{w_{\min}^4} n \Delta_K^2.$$

■

Appendix C. Bernstein Inequality

Our analysis relies on the Bernstein inequality. To simplify presentation, we state below a user-friendly version of Bernstein inequality.

Lemma 11 *Consider n independent random variables z_l ($1 \leq l \leq n$), each satisfying $|z_l| \leq B$. For any $a \geq 2$, one has*

$$\left| \sum_{l=1}^n z_l - \mathbb{E} \left[\sum_{l=1}^n z_l \right] \right| \leq \sqrt{2a \log n \sum_{l=1}^n \mathbb{E} [z_l^2]} + \frac{2a}{3} B \log n \quad (80)$$

with probability at least $1 - \frac{2}{n^a}$.

This is an immediate consequence of the well-known Bernstein inequality

$$\mathbb{P} \left\{ \left| \sum_{l=1}^n z_l - \mathbb{E} \left[\sum_{l=1}^n z_l \right] \right| > t \right\} \leq 2 \exp \left(- \frac{\frac{1}{2} t^2}{\sum_{l=1}^n \mathbb{E} [z_l^2] + \frac{1}{3} B t} \right). \quad (81)$$

References

- A. Agresti. *Categorical data analysis*. John Wiley & Sons, 2014.
- N. Ailon. Active learning ranking from pairwise preferences with almost optimal query complexity. *Journal of Machine Learning Research*, 13:137–164, 2012.
- A. Ammar and D. Shah. Ranking: Compare, don’t score. In *Allerton Conference*, pages 776–783. IEEE, 2011.
- A. Ammar and D. Shah. Efficient rank aggregation using partial data. In *SIGMETRICS*, volume 40, pages 355–366. ACM, 2012.
- P. Awasthi, A. Blum, O. Sheffet, and A. Vijayaraghavan. Learning mixtures of ranking models. In *Neural Information Processing Systems*, pages 2609–2617, 2014.
- S. Balakrishnan, M. J. Wainwright, and B. Yu. Statistical guarantees for the EM algorithm: From population to sample-based analysis. *arXiv:1408.2156*, 2014.
- L. Baltrunas, T. Makcinskas, and F. Ricci. Group recommendations with rank aggregation and collaborative filtering. In *ACM conference on Recommender systems*, pages 119–126. ACM, 2010.
- R. A. Bradley and M. E. Terry. Rank analysis of incomplete block designs: I. the method of paired comparisons. *Biometrika*, pages 324–345, 1952.
- M. Braverman and E. Mossel. Noisy sorting without resampling. In *ACM-SIAM SODA*, pages 268–276, 2008.
- S. Brin and L. Page. The anatomy of a large-scale hypertextual web search engine. *Computer networks and ISDN systems*, 30(1):107–117, 1998.
- S. Bubeck, T. Wang, and N. Viswanathan. Multiple identifications in multi-armed bandits. *ICML*, 2013.
- R. Busa-Fekete and E. Hüllermeier. A survey of preference-based online learning with bandit algorithms. In *Algorithmic Learning Theory*, pages 18–39, 2014.
- R. Busa-Fekete, B. Szörényi, P. Weng, W. Cheng, and E. Hüllermeier. Top-k selection based on adaptive sampling of noisy preferences. In *ICML*, 2013.
- R. Busa-Fekete, E. Hüllermeier, and B. Szörényi. Preference-based rank elicitation using statistical models: The case of Mallows. In *ICML*, pages 1071–1079, 2014.
- E. Candes, X. Li, and M. Soltanolkotabi. Phase retrieval via Wirtinger flow: Theory and algorithms. *IEEE Trans. on Inf. Theory*, 61(4):1985 – 2007, March 2015.
- A. Caplin and B. Nalebuff. Aggregation and social choice: a mean voter theorem. *Econometrica*, pages 1–23, 1991.
- X. Chen, P. N. Bennett, K. Collins-Thompson, and E. Horvitz. Pairwise ranking aggregation in a crowdsourced setting. In *WSDM*, pages 193–202, 2013.

- Y. Chen, C. Suh, and A. J. Goldsmith. Information recovery from pairwise measurements: A Shannon-theoretic approach. *arXiv preprint arXiv:1504.01369*, 2015b.
- C. Dwork, R. Kumar, M. Naor, and D. Sivakumar. Rank aggregation methods for the web. In *Proceedings of the Tenth International World Wide Web Conference*, 2001.
- B. Eriksson. Learning to top- k search using pairwise comparisons. In *AISTATS*, pages 265–273, 2013.
- F. Farnoud and O. Milenkovic. An axiomatic approach to constructing distances for rank comparison and aggregation. *IEEE Trans on Inf Theory*, 60(10):6417–6439, 2014.
- F. Farnoud, E. Yaakobi, and J. Bruck. Approximate sorting of data streams with limited storage. In *Computing and Combinatorics*, pages 465–476. Springer, 2014.
- L. R. Ford. Solution of a ranking problem from binary comparisons. *American Mathematical Monthly*, 1957.
- V. Gabillon, M. Ghavamzadeh, A. Lazaric, and S. Bubeck. Multi-bandit best arm identification. In *Neural Information Processing Systems*, pages 2222–2230, 2011.
- B. Hajek, S. Oh, and J. Xu. Minimax-optimal inference from partial rankings. In *NIPS*, pages 1475–1483, 2014.
- T. Han and S. Verdú. Generalizing the Fano inequality. *IEEE Transactions on Information Theory*, 40(4):1247–1251, 1994.
- E. F. Harrington. Online ranking/collaborative filtering using the perceptron algorithm. In *ICML*, volume 20, pages 250–257, 2003.
- D. R. Hunter. MM algorithms for generalized Bradley-Terry models. *Annals of Statistics*, pages 384–406, 2004.
- P. Jain, P. Netrapalli, and S. Sanghavi. Low-rank matrix completion using alternating minimization. In *Symposium on Theory of computing*, pages 665–674. ACM, 2013.
- K. Jamieson, M. Malloy, R. Nowak, and S. Bubeck. lil’UCB: An optimal exploration algorithm for multi-armed bandits. *Conference on Learning Theory*, 2014.
- K. G. Jamieson and R. Nowak. Active ranking using pairwise comparisons. In *Advances in Neural Information Processing Systems*, pages 2240–2248, 2011.
- R. Keshavan, A. Montanari, and S. Oh. Matrix completion from noisy entries. In *Advances in Neural Information Processing Systems*, pages 952–960, 2009.
- R. H. Keshavan, A. Montanari, and S. Oh. Matrix completion from a few entries. 56(6): 2980–2998, June 2010.
- T. Lu and C. Boutilier. Learning Mallows models with pairwise preferences. In *ICML*, pages 145–152, 2011.

- Y. Lu and S. N. Negahban. Individualized rank aggregation using nuclear norm regularization. *arXiv preprint arXiv:1410.0860*, 2014.
- R. D. Luce. *Individual choice behavior: A theoretical analysis*. Wiley, 1959.
- M. Mitzenmacher and E. Upfal. *Probability and computing: Randomized algorithms and probabilistic analysis*. Cambridge University Press, 2005.
- S. Negahban, S. Oh, and D. Shah. Rank centrality: Ranking from pair-wise comparisons. 2012. URL <http://arxiv.org/abs/1209.1688>.
- P. Netrapalli, P. Jain, and S. Sanghavi. Phase retrieval using alternating minimization. In *Advances in Neural Information Processing Systems*, pages 2796–2804, 2013.
- S. Oh and D. Shah. Learning mixed multinomial logit model from ordinal data. In *Neural Information Processing Systems*, pages 595–603, 2014.
- B. Osting, J. Xiong, Q. Xu, and Y. Yao. Analysis of crowdsourced sampling strategies for hodgeRank with sparse random graphs. *arXiv:1503.00164*, 2015.
- D. Park, J. Neeman, J. Zhang, S. Sanghavi, and I. S. Dhillon. Preference completion: Large-scale collaborative ranking from pairwise comparisons. 2015.
- A. Rajkumar and S. Agarwal. A statistical convergence perspective of algorithms for rank aggregation from pairwise data. In *ICML*, pages 118–126, 2014.
- N. B. Shah, S. Balakrishnan, J. Bradley, A. Parekh, K. Ramchandran, and M. Wainwright. When is it better to compare than to score? *arXiv:1406.6618*, 2014.
- H. A. Soufiani, W. Z. Chen, D. C. Parkes, and L. Xia. Generalized method-of-moments for rank aggregation. In *Neural Information Processing Systems*, 2013.
- H. A. Soufiani, D. Parkes, and L. Xia. Computing parametric ranking models via rank-breaking. In *International Conference on Machine Learning (ICML)*, 2014a.
- H. A. Soufiani, D. C. Parkes, and L. Xia. A statistical decision-theoretic framework for social choice. In *Neural Information Processing Systems*, 2014b.
- T. van Erven and P. Harremoës. Renyi divergence and Kullback-Leibler divergence. *IEEE Transactions on Information Theory*, 60(7):3797–3820, July 2014. ISSN 0018-9448. doi: 10.1109/TIT.2014.2320500.
- F. Wauthier, M. Jordan, and N. Jovic. Efficient ranking from pairwise comparisons. In *ICML*, pages 109–117, 2013.
- R. W. Yeung. *Information theory and network coding*. Springer, 2008.