

Intrinsic pseudolocal modes in monatomic chain and in graphene

V. Hizhnyakov¹, M. Klopov², A. Shelkan¹

¹*Institute of Physics, University of Tartu, Ravila 14c, 50411 Tartu, Estonia*

²*Department of Physics, Faculty of Science, Tallinn University of Technology,
Ehitajate tee 5, 19086 Tallinn, Estonia
hizh@ut.ee*

Abstract

In this paper an analytical and numerical study of anharmonic vibrations of monatomic chain and graphene in normal to the chain/plane direction is presented. Due to the lack of odd anharmonicities and the presence of hard quartic anharmonicity for these vibrations, there may exist localized anharmonic modes of this type with the frequency above the spectrum of out of-chain/plane phonons. Although this frequency is in resonance with longitudinal (chain) or in-plane (graphene) phonons, the decay of the modes may be very slow. E.g. in chain their lifetime may exceed 10^9 periods. We call these vibrations as intrinsic pseudolocal modes.

1. Introduction

It is well known that point defects in crystals may cause an appearance of spatially localized vibrations, called local modes (see, e.g. [1]). The frequency of such a mode lies outside the phonon spectrum, which prevents the spreading of the vibration to the bulk. Besides, in crystals may exist also resonant or pseudo-local modes (PLMs) with the frequency inside the phonon spectrum. The latter appear when the frequency of the leading vibration associated with the defect gets to a region of small density of states (DOS) of phonons. Unlike local modes, which are stable in harmonic approximation, these modes can live only a finite time due to emission of resonant phonons. However, because of small DOS of the resonant phonons the interaction of the latter with PLM is weak, due to what the lifetime of the mode becomes long. A well-known case of long-living PLMs in 3D lattices is the one by the defect of large mass [1].

Spatially localized modes may exist also in perfect nonlinear lattices. Such modes in chains with cubic and quartic anharmonicity were first described by A. M. Kosevich and A. S. Kovalev [2]. They have found that for sufficiently strong quartic anharmonicity there exist vibrational states localized in space and periodic in time; the frequency of these vibrations exceeds the maximum frequency of phonons. The authors [2] restricted their consideration with localized vibrations of large size and frequency just above the top of the phonon spectrum. Localized vibrations in anharmonic lattices with small size were introduced by Dolgov [3] and then by A. J. Sievers and S. Takeno [4] and were called “intrinsic localized modes” (ILMs).

In numerical studies of ILMs different two-body potential models (Lennard-Jones, Born-Mayer-Coulomb, Toda, and Morse potentials and their combinations) have been used (see, e.g. [5-7]). All these potentials have strong odd anharmonicities and show a strong softening with the increasing of vibrational amplitude. ILMs found in these simulations always drop down from optical band(s) into the phonon gap, if there is one. In this connection see Ref. [8], where gap ILMs in NaI were calculated without taking long-range interactions

into account and Ref. [9], where the calculations ILMs in NaI were made taking these interactions into account (long-range interactions have strong effect to phonons in NaI).

Usually in crystal lattices odd anharmonicities are strong and pair potentials show a strong softening with increase of vibrational amplitude. Therefore, the dropping down of the frequency of ILMs from optical bands is quite common in 3D lattices. Still, as it was shown in Refs. [10-13], in some crystals odd anharmonicities are reduced due to multiparticle or covalent interactions. Examples have been given with germanium [10], diamond [11], metallic Ni, Nb [12], iron [13] and copper [11]; in all these crystals ILMs with frequencies above the top of the phonon band were found in numerical simulations.

However, there are systems in which the odd anharmonicities disappear due to symmetry arguments; at the same time, the quartic anharmonicity is non-zero and is hard (positive). The examples of such systems are given by linear atomic chains and planar atomic structures (e.g. graphene); in these systems odd anharmonicities disappear for vibrations in normal direction (transverse in chain and out-of-plane in graphene). Therefore, one can expect that in these systems there can exist out-of-chain/plane intrinsic localized modes with the frequencies above the maximum frequency of corresponding phonons. The latter frequencies are usually smaller than the maximum frequency of longitudinal/in-plane phonons. These modes fall in resonance with these phonons and as a result can decay. Therefore we call these modes as intrinsic pseudolocal modes (IPLMs). However, unlikely to PLMs the interaction causing the decay of IPLMs is anharmonic, due to what double frequency phonons are created. This interaction in case of a IPLM with small vibrational amplitudes of atoms is very weak. Hence, one can expect that the lifetime of such IPLM may be rather long. Below we will consider IPLMs in a monatomic chain and in graphene, both analytically and numerically. Our considerations confirm the aforesaid expectation.

2. Anharmonic chain

First we consider the anharmonic monatomic chain and examine the vibrations of its atoms in longitudinal (x) and transverse (y) directions. We suppose that the potential energy U of the chain is given by the sum of pair-potentials $V(R_{n-n'})$. The latter potentials can be expanded into the series of atomic displacements. Denoting $a' = (n - n')a$, $x = x_n - x_{n'}$, $y = y_n - y_{n'}$, where a is the atom spacing, x_n and y_n are the longitudinal and transverse displacements of the atom number n from its equilibrium position, we get $R = a' \left(\sqrt{1 + 2x/a' + (x^2 + y^2)/a'^2} - 1 \right)$. Here the term quadratic with respect to y has the same numerical factor as the term linear with respect to x . The same holds also for an arbitrary power of R . In the equilibrium state all linear terms with respect to the coordinates $x \equiv x_n - x_{n'}$ in the potential energy U are cancelled. Therefore, all quadratic terms with respect to $y \equiv y_{n-n'}$ are also cancelled. This means that the frequencies of transverse vibrations tend to zero. Consequently, in pair potentials approximation the chains in the equilibrium position are unstable with respect to small transverse distortions.

To get the chain stable one needs to stretch it [14]. In this case the atom spacing a is replaced by $a + s$ with $s > 0$. Then the terms in U linear with respect to x and y^2 are not cancelled any more. Therefore, the elastic springs for transverse vibrations are also nonzero and positive [14]. This results in the appearance of the transverse phonons with finite,

although small for small stretching maximum frequency. As these phonons do not have any cubic anharmonicity, but have nonzero positive quartic anharmonicity, low-frequency IPLMs with the frequency above the spectrum of transverse phonons should exist here.

Let us consider the IPLM in a monatomic chain with the Morse pair potential $V = D(1 - e^{\alpha(a-r)})^2$. Here D is the energy of dissociation, α is the parameter. We are using dimensionless coordinates with the units corresponding to $a=1$ and the value $\alpha=4$ of the Morse pair potentials of atoms in monatomic metals. For this potential only nearest-neighbor interactions are essential and only these potentials will be taken into account here. We also take for the mass units the mass of the atoms of the chain ($m=1$). The dissociation energy is chosen so that the unit frequency will correspond to the maximum frequency of longitudinal phonons. In this case the pair potential of the stretched chain reads:

$$V = (1 - e^{-4(r-1)})^2 / 128 - x(1 - e^{-4s})e^{-4s} / 16, \quad (1)$$

where $r = \sqrt{(x+1+s)^2 + y^2}$, s is the stretching of the atom spacing. The last term in Eq. (1) stands for the stretching energy. Let us expand the potential into the series of x and y and take into account up to the second order terms with respect to x and forth-order terms with respect to y . We get (up to a constant)

$$V(x, y) \approx \frac{V_{1s}}{8} x^2 + \frac{V_{2s}}{8} sy^2 + \frac{V_{3s}}{8} xy^2 + \frac{V_{4s}}{32} y^4, \quad (2)$$

where $v_{is} \equiv v_i$ are dependent on s parameters. In the small $s \lesssim 10^{-2}$ limit $v_{is} \approx 1$. If $s = 0.05$ then $v_1 = 0.522$, $v_2 = 0.707$, $v_3 = 0.463$, $v_4 = 0.441$. The transverse vibrations alone are described by the potential $V(0, y) = v_2 sy^2 / 8 + v_4 y^4 / 32$. For this pair potential the anharmonic modes exist (called here IPLM) with the frequencies ω_0 close to the maximum frequency of the transverse phonons $\omega_{tm} = \sqrt{v_2 s}$ and the shape [2]

$$y_n(t) \approx (-1)^n \frac{2\omega_{tm}}{\sqrt{3v_4}} \frac{\varepsilon \cos(\omega_0 t)}{\cosh(\varepsilon n)}, \quad \varepsilon = \sqrt{\omega_0^2 - \omega_{tm}^2} \ll 1 \quad (3)$$

These modes interact resonantly with longitudinal acoustic phonons with the frequencies $\omega_{ac} = 2\omega_0 = 2\sqrt{\omega_{tm}^2 + \varepsilon^2}$ and, therefore, they should decay. However, this interaction in the case under consideration, $y_n \ll 1$, is weak and this decay is slow.

In presence of IPLM the pair-potential for the longitudinal motion gets the form

$$V(x, t) \approx \frac{V_1}{8} x^2 + \frac{V_3}{16} xy_0^2 + \frac{V_3}{16} xy_0^2 \cos(2\omega_0 t). \quad (4)$$

The second term stands for a small local compression, while the third term describes periodical force with the frequency $2\omega_0$. For all longitudinal phonons, except those with the resonant frequency $\omega_k = 2\omega_0$, this force causes forced vibrations of atoms with the frequency $2\omega_0$. The resonant term causes the increase of the energy of phonons in time and, consequently, the decay of the transverse anharmonic mode under consideration.

3. Decay of transverse IPLM in chain

To find this decay we consider the equation of motion of the longitudinal phonon with the coordinate $x_k = N^{-1/2} \sum_n x_n e^{ikn}$, where $N \gg 1$ is the number of atoms in the chain, $k = \pi m/N$ is the wave number, $m = 0, \pm 1, \pm 2, \dots$ (we use periodic boundary conditions). This equation reads

$$\ddot{x}_k + \omega_k^2 x_k = \left(N^{-1/2} v_3 / 8 \right) \sum_{n=0}^{\infty} \cos(kn) \left((y_{0,n+1} - y_{0,n})^2 - (y_{0,n} - y_{0,n-1})^2 \right) \cos(2\omega_0 t), \quad (5)$$

where $\omega_k = \sqrt{v_1(1 - \cos(k))}/2$ is the frequency of the phonon, $y_{0,n} \equiv y_n(0)$. In the small ω_k limit $\omega_k \approx k\sqrt{v_1}/2$ the transverse amplitude $y_{0,n}$ slowly changes with n . Therefore, $y_{0,n+1} - y_{0,n} \approx y'_{0,n}$ and therefore the force acting to the atom n in x direction from the IPLM depends on

$$y'_{0,n}(t) - y'_{0,n-1}(t) \approx (y'_{0,n}(t) + y'_{0,n-1}(t))(y'_{0,n}(t) - y'_{0,n-1}(t)) \approx \partial y'_{0,n}(t) / \partial n. \quad (6)$$

This gives

$$\ddot{x}_k + \omega_k^2 x_k \cong \varepsilon^3 s v_3 C_k N^{-1/2} \cos(2\omega_0 t), \quad (7)$$

where

$$C_k \cong \frac{\varepsilon v_2}{6v_4} \sum_{n=0}^{\infty} \cos(kn) \frac{\partial}{\partial n} \left(\frac{\sinh(\varepsilon n)}{\cosh^2(\varepsilon n)} \right)^2 \cong \frac{\varepsilon v_2 k}{6v_4} \sum_{n=0}^{\infty} \sin(kn) \left(\frac{\sinh(\varepsilon n)}{\cosh^2(\varepsilon n)} \right)^2. \quad (8)$$

By using the Green's function of harmonic oscillator $G_k(t) = \omega_k^{-1} \sin(\omega_k t)$, the solution of Eq. (7) can be presented in the following form: $x_k(t) = x_{0,k} \cos(\omega_k t) + x'_k(t)$, where

$$x'_k(t) = \frac{\varepsilon^3 s v_3 C_k}{\sqrt{N} \omega_k} \int_0^t \sin(\omega_k(t-t')) \cos(2\omega_0 t') dt'. \quad (9)$$

The first ($\propto x_{0,k}$) term stands for free oscillations, while the second term ($x'_k(t)$) describes the excitation of vibrations by the transverse mode. The energy of the excited vibrations is given by the sum of the terms $\omega_k^2 x_k'^2$ over all phonons. This gives

$$E(t) = \frac{\varepsilon^6 s^2 v_3^2}{N} \sum_k C_k^2 \left(\int_0^t (\sin(\omega_k t) \cos(\omega_k t') + \cos(\omega_k t) \sin(\omega_k t') / 2) \cos(2\omega_0 t') dt' \right)^2. \quad (10)$$

This energy is taken from the IPLM under consideration. In the large time t limit one can average the energy over the period of vibrations. Then fast oscillating terms average out and we get

$$dE(t)/dt = \left(\varepsilon^6 v_3^2 s^2 / N \right) \sum_k C_k^2 \left[\left(\int_0^t \cos((\omega_k - 2\omega_0)t') dt' \right) + \left(\int_0^t \sin((\omega_k - 2\omega_0)t') dt' \right) \right]. \quad (11)$$

The second term is odd with respect to $\omega_k - 2\omega_0$. Therefore, its contribution is inessential. As a result, in the $t \rightarrow \infty$ limit the rate of increasing the energy of longitudinal phonons, followed by a subsequent decrease of the energy of the IPLM under consideration, equals

$$dE(t)/dt = \left(\pi \varepsilon^6 v_3^2 s^2 / N \right) \sum_k C_k^2 \delta(\omega_k - 2\omega_0), \quad (12)$$

Replacing here the sum over k by the sum over $m = Nk/\pi$ and the latter sum by the integral and taking into account that $dk = Ndm/\pi$, we get

$$dE/dt \approx (\varepsilon^6 k_0^2 s^2 \nu_2^2 \nu_3^2 / 36) \Phi^2(\Omega), \quad (13)$$

where $k_0 = \arccos(\sqrt{1 - 8\omega_0^2/\nu_1})$, $\omega_0 = \sqrt{\omega_{tm}^2 + \varepsilon^2}$, $\Omega = k_0/\varepsilon$,

$$\Phi(\Omega) = \int_0^\infty \frac{\sin(\Omega x) \sinh^2(x)}{\cosh^4(x)} dx, \quad (14)$$

The dependence of $\ln \Phi$ on Ω is given in Fig. 1. In the small ε limit (large size of the IPLM) $\Omega \gg 1$ and $\Phi(\Omega)$ is very small.




Fig. 1. Function $\log_{10} \Phi$ in dependence of $\Omega = k_0/\varepsilon$.

The decay constant Γ of the IPLM is determined by the equation $\Gamma = E^{-1} dE/dt$. Taking into account the relation

$$E = \omega_0^2 \sum_n y_n^2(0) = (4/3\nu_4) \varepsilon \omega_0^2 \omega_{tm}^2, \quad (15)$$

we get

$$\Gamma = (\varepsilon^5 \nu_3^2 k_0^2 / 48 \nu_4 \omega_0^2) \Phi^2(\Omega). \quad (16)$$

Here $\varepsilon = y_0(0) \sqrt{3\nu_4/4s\nu_2}$. Consequently Γ decreases very fast (faster than fifth power) with decreasing of the maximum amplitude $y_0(0)$ of the IPLM. To get an estimation of Γ we consider the IPLM with amplitude $y_0 = 0.03$ in the 5% stretched chain ($s = 0.05$). In this chain $\varepsilon \approx 0.11$, which gives $k_0 = 0.755$, $\omega_0 = 0.23$ and $\Omega = 6.86$. This gives $\Gamma = 1.4 \cdot 10^{-10}$ which corresponds to very long lifetime. Note a specific property of the decay of the IPLMs under consideration: the existence of stable state for definite amplitude and energy (see the sharp minimum going to $-\infty$ in Fig. 1.).

In our numerical study of transverse IPLMs we considered a 5% stretched chain with 40000 atoms and with fixed ends. The small compression of entire chain due to IPLM was taken into account. The potential energy was described by the interactions of nearest neighbors with the Morse pair-potential given by Eq. (1). We have found that transverse IPLMs indeed exist in this chain and they have long lifetime (see Figs. 2, 3; one period of the IPLM with the frequency $\omega_0 = 0.189$ corresponds to 33 time units). E.g. we have found that the IPLM with amplitude $y_0 = 0.03$ indeed decays very slowly: its amplitude (and energy) diminishes on $1.7 \cdot 10^{-8}$ for 1000 periods of vibrations. This corresponds to $\Gamma \approx 1.7 \cdot 10^{-10}$.

Taking into account that used here theory [2] is approximate and the estimation of decay in numerical study has limited accuracy one can conclude that theoretical and numerical results agree very well. The IPLMs with amplitudes ≥ 0.04 decay faster (see Figs. 2b and 3), at that their rate of decay rapidly increases with increasing of the amplitude which is also in agreement with the theory.




Fig. 2a. Time dependence of the shifts (in the units of atom spacing) of the central atom of a transverse IPLM in 5% stretched monatomic chain; initial shift of central atom is 0.05, first 10 periods and 10 periods after 1000 periods of vibration are presented.




Fig. 2b. The same as in Fig. 2a but for another (0.15) initial shift of central atom.




Fig. 3. Time dependence of the shifts of the central atom of transverse IPLMs in 5% stretched monatomic chain for different initial amplitudes of the central atom. The IPLM with the amplitudes < 0.05 do not decay for $40000\omega_0/2\pi \approx 1200$ periods.

4. Graphene: out-of-plane vibrations

The arguments presented in the beginning of Section 2 can be easily extended to planar atomic lattices. To be specific, we are considering here a graphene sheet. It consists of carbon atoms constituting the honeycomb structure. Every carbon atom in this structure is connected with three neighboring atoms by chemical bonds formed by the sp_2 hybrid orbitals.

We suppose that in equilibrium positions the atoms of graphene are situated in (x, y) plane. The distances between an atom and three neighbouring atoms can be presented in the form

$$r_n = \sqrt{(b_n a + x_n)^2 + (c_n a + y_n)^2 + z_n^2} \quad (17)$$

($n=1,2,3$), where x_n , y_n and z_n are the x -, y - and z -components of the displacements of three neighboring atoms with respect to the central atom, $b_1=0, b_2=\sqrt{3}/2, b_3=-\sqrt{3}/2$, $c_1=-1, c_2=1/2, c_3=1/2$. In the pair-potential approximation the potential energy of the vibrations of atoms depends on the distances $R_n = r_n - a$, $n=1,2,3$ (explicit form of this energy in harmonic approximation, see e.g. in [15]). The expansion of these distances over the displacements x_k , y_k and z_k depends on the powers $R^m = (bx + cy + (x^2 + y^2 + z^2)/a)^m$. (here subscript n is omitted for simplicity). In the equilibrium position the expansion of these potential energies with respect to in-plane displacements starts with the terms quadratic with respect to x and y . Therefore, the expansion over z at $x=y=0$ starts with the positive quartic term $\propto z^4$ (in full analogy with expansion over y in monatomic chain). In this case also in the pair-potential approximation the out-of-plane phonons are absent and the plane is unstable with respect to out-of plane distortions. However, in graphene the atomic interactions are determined by covalent bonds causing the angular dependence of pair potentials. This dependence results in appearance of the stiffness of the planes with respect to the transverse short-range displacements of atoms [15]. (In accordance with the Mermin-Wagner theorem [16] the plane of graphene remains unstable with respect to long-range transverse distortions resulting in the appearance of ripples [17, 18]). A stretching of the graphene sheet removes the ripples.

The elementary cell of graphene (honeycomb) lattice includes two atoms. Therefore, there are two branches of out-of-plane vibrations: acoustic and optic. The maximum frequencies of out-of-plane acoustic (≈ 19 THz) and optic phonons (≈ 27 THz) are smaller than the maximum frequency of the in-plane transverse acoustic phonons ≈ 41 THz and in-plane longitudinal acoustic phonons ≈ 51 THz [15]. Due to the existence of two branches of out-of-plane phonons, acoustic and optic, one can expect the existence of two types of out-of plane IPLM – with the frequency above the top frequency of acoustic out-of-plane phonons (the acoustic-like IPLM) and with the frequency above the top frequency of optic out-of-plane phonons (the optic like IPLM). The acoustic-like out-of-plane IPLMs can decay due to a relatively weak anharmonic interaction with the in-plane phonons and also due to in general stronger harmonic resonant interaction with out-of-plane optic phonons. However, the optic like out-of-plane IPLMs can decay only due to a relatively weak anharmonic interaction with in-plane phonons. Therefore the optic-type out-of-plane IPLMs should have longer lifetime.

This conclusion is in agreement with our numerical modelling of out-of-plane IPLMs in weakly (1.5 %) stretched graphene. In our calculations we have used AIREBO potential and taken into account the cluster 246×210 Å (100×50 periods; 20000 atoms) with the periodical boundary conditions. Initially we displaced out-of-plane six atoms of C_6 ring according to the optical mode. We found that out-of-plane optic IPLMs indeed exist (see Fig. 4, where the time-dependences of the displacements of the one of three equivalent atoms of the central C_6 ring are presented; the other three atoms in this ring vibrate out of phase with the mentioned atom).




Fig. 4. Optical-type out-of-plane IPLMs in graphene. The out-of-plane shifts (independent on time) of one of three equivalent atoms of the central C_6 ring is presented for initial amplitudes 0.4 Å (upper) and 0.2 Å (bottom); the other three atoms in this ring vibrate out of phase with the atom mentioned.




Fig. 5. In-plane and out-of-plane phonons DOS for AIREBO potential and the spectrum of out-of-plane IPLM for different initial amplitudes of carbon atoms.

Note that in strongly stretched graphene can exist also anharmonic localized in-plane vibrations (in-plane discrete breathers) [19, 20]. It appears (see Ref. [19]) that strong uniaxial stretching (along the zigzag or armchair direction) results in the opening of the gap in the middle of the phonon spectrum of graphene. This makes possible the existence of in-plane soft ILMs (discrete breathers) with the frequency in the gap of the phonon spectrum [19,20]. The authors [19, 20] have found that these modes remain relatively stable also when, due to large stretching their frequencies get into resonance with out-of-plane phonons. Thus, the physical situation discussed in Refs. [19, 20] is reversed to the one considered in this study: the anharmonic in-plane modes with soft anharmonicity are in resonance with the out-of-plane phonons, while considered here IPLMs are in resonance with in-plane phonons. However, in both cases the lifetime of the anharmonic localized modes is rather long. In our opinion the physical mechanism of decay –emission of phonons of double frequency due to anharmonic interaction, should be common in both cases.

5. Concluding remarks

In this communication it was shown that due to symmetry arguments in linear monatomic chains and in planar monatomic structures (e.g. graphene) odd anharmonicities disappear for normal with respect to the chain/plane motion of atoms. However, even (quartic) anharmonicity exists and is hard (positive). As a result, the anharmonic localized modes of transverse (chain) or out-of-plane (graphene) vibrations (called IPLMs) can exist in these systems with frequencies above the transversal/ out-of-plane phonons. In the case of monatomic chain the theory is developed allowing one to calculate the decay rate of IPLMs. The theory predicts that the lifetime of the IPLMs under consideration may be extremely long (longer than 10^9 periods of vibration). Numerical modelling of these IPLMs fully confirms this result. The normal (out-of-plane) IPLMs with the frequency above the phonon spectrum of out-of-plane phonons were also found for weakly stretched graphene.

The consideration presented here holds generally for linear atomic chains and planar monolayer atomic structures. Therefore one can expect the existence of transverse IPLMs in any linear atomic chain and any planar monolayer atomic structure. Besides, one can expect that analogous situation may exist in 3D lattices with dominating chain and layer structural elements, e.g. in graphite. In such lattices odd anharmonicities exist, but they may be essentially reduced for out of chain/plane vibrations. Analogous IPLMs can be expected to exist also for surface vibrations in crystals with planar atomic surfaces. The change of potential energy of a surface atom for the shift in outer direction also does not have any odd terms with respect to the shift. This leads to decrease of the entire odd-anharmonicity effects for out-of-plane surface vibrations, which may result in the appearance of out-of-plane IPLMs with frequencies above the maximum frequency of transverse surface phonons and with a long lifetime.

Acknowledgements. The research was supported by Estonian research projects SF0180013s07, IUT2-27 and by European Union through the European Regional Development Fund (project 3.2.01.11-0029).

References

1. A. Maradudin, Theory of lattice dynamics in harmonic approximation, Academic Press, 1971.

2. A. M. Kosevich and A. S. Kovalev, Sov. Phys. JETP **40**, 891 (1974).
3. A. S. Dolgov, Sov. Phys. Solid State **28**, 907 (1986).
4. J. Sievers and S. Takeno, Phys. Rev. Lett. **61**, 970 (1988).
5. J. B. Page, Phys. Rev. B **41**, 7835 (1990).
6. J. Sievers and J. B. Page, in Dynamical Properties of Solids: Phonon Physics The Cutting Edge, edited by G. K. Norton and A. A. Maradudin (North Holland, Amsterdam, 1995), Vol. VII, p. 137.
7. S. Flach and C. R. Willis, Phys. Repts. **295**, 182 (1998).
8. S. A. Kiselev and A. J. Sievers, Phys. Rev. B **55**, 5755 (1997).
9. M. Haas, V. Hizhnyakov, M. Klopov, A. Shelkan, IOP Conf. Series: Materials Science and Engineering, **15**, 12045 (2010).
10. N. K. Voulgarakis, S. Hadjisavvas, P. C. Kelires, and G. P. Tsironis, Phys. Rev. B **69**, 113201 (2010).
11. V. Hizhnyakov, M. Haas, A. Shelkan, M. Klopov, Standing and moving discrete breathers with frequencies above phonon spectrum. Springer (forthcoming).
12. V. Haas, V. Hizhnyakov, A. Shelkan, M. Klopov, A. J. Sievers, PRB **84**, 144303 (2011).
13. V. Hizhnyakov, M. Haas, A. Shelkan and M. Klopov, Phys. Scr. **84**, 044003 (2014).
14. S. Cadet, Phys. Lett. A **121**, 77 (1987).
15. V. Adamyan and V. Zavalniuk, J. Phys.: Condens. Matter **23**, 015402 (2011).
16. N. D. Mermin, Phys. Rev. **176**, 250 (1968).
17. J. C. Meyer, A. K. Geim, M. I. Katsnelson, K. S. Novoselov, T. J. Booth and S. Roth, Nature **446**, 60 (2007).
18. A. Fasolino, J. H. Los, M. I. Katsenelson, Nature Materials, **6**, 858 (2007).
19. L. Z. Khadeeva, S. V. Dmitriev, Yu.S. Kivshar, JETP Lett. **94**, 539 (2011).
20. J. A. Baimova, S. V. Dmitriev, K. Zhou, EPL **100**, 36005 (2012).