

A uniformly accurate (UA) multiscale time integrator Fourier pseudospectral method for the Klein-Gordon-Schrödinger equations in the nonrelativistic limit regime

Weizhu Bao · Xiaofei Zhao

Received: date / Accepted: date

Abstract A multiscale time integrator Fourier pseudospectral (MTI-FP) method is proposed and analyzed for solving the Klein-Gordon-Schrödinger (KGS) equations in the nonrelativistic limit regime with a dimensionless parameter $0 < \varepsilon \leq 1$ which is inversely proportional to the speed of light. In fact, the solution to the KGS equations propagates waves with wavelength at $O(\varepsilon^2)$ and $O(1)$ in time and space, respectively, when $0 < \varepsilon \ll 1$, which brings significantly numerical burdens in practical computation. The MTI-FP method is designed by adapting a multiscale decomposition by frequency to the solution at each time step and applying the Fourier pseudospectral discretization and exponential wave integrators for spatial and temporal derivatives, respectively. We rigorously establish two independent error bounds for the MTI-FP at $O(\tau^2/\varepsilon^2 + h^{m_0})$ and $O(\varepsilon^2 + h^{m_0})$ for $\varepsilon \in (0, 1]$ with τ time step size, h mesh size and $m_0 \geq 4$ an integer depending on the regularity of the solution, which imply that the MTI-FP converges uniformly and optimally in space with exponential convergence rate if the solution is smooth, and uniformly in time with linear convergence rate at $O(\tau)$ for $\varepsilon \in (0, 1]$ and optimally with quadratic convergence rate at $O(\tau^2)$ in the regime when either $\varepsilon = O(1)$ or $0 < \varepsilon \leq \tau$. Thus the meshing strategy requirement (or ε -scalability) of the MTI-FP is $\tau = O(1)$ and $h = O(1)$ for $0 < \varepsilon \ll 1$, which is significantly better than classical methods. Numerical results demonstrate that our error bounds are optimal and sharp. Finally, the MTI-FP method is applied to study numerically convergence rates of the KGS equations to the limiting models in the nonrelativistic limit regime.

W. Bao

Department of Mathematics and Center for Computational Science and Engineering, National University of Singapore, Singapore 119076

Fax: +65-6779-5452, Tel.: +65-6516-2765,

URL: <http://www.math.nus.edu.sg/~bao/>

E-mail: matbaowz@nus.edu.sg

X. Zhao

Department of Mathematics, National University of Singapore, Singapore 119076

E-mail: zhxfnus@gmail.com

Keywords Klein-Gordon-Schrödinger equations · nonrelativistic limit regime · multiscale time integrator · uniformly accurate · multiscale decomposition · exponential wave integrator

Mathematics Subject Classification (2000) 65L05 · 65L20 · 65L70

1 Introduction

Consider the Klein-Gordon-Schrödinger (KGS) equations in d -dimensions ($d = 3, 2, 1$) [2, 22, 42]:

$$i\hbar\partial_t\psi(\mathbf{x},t) + \frac{\hbar^2}{2m_1}\Delta\psi(\mathbf{x},t) + g\phi(\mathbf{x},t)\psi(\mathbf{x},t) = 0, \quad \mathbf{x} \in \mathbb{R}^d, \quad (1.1a)$$

$$\frac{1}{c^2}\partial_{tt}\phi(\mathbf{x},t) - \Delta\phi(\mathbf{x},t) + \frac{m_2^2c^2}{\hbar^2}\phi(\mathbf{x},t) - g|\psi(\mathbf{x},t)|^2 = 0, \quad (1.1b)$$

which represent a classical model for describing the dynamics of a complex-valued scalar nucleon field $\psi := \psi(\mathbf{x},t)$ interacting with a neutral real-valued scalar meson field $\phi := \phi(\mathbf{x},t)$ through the Yukawa coupling with $0 \neq g \in \mathbb{R}$ the coupling constant. Here t is time, $\mathbf{x} \in \mathbb{R}^d$ is the spatial coordinate, \hbar is the Planck constant, c is speed of light, $m_1 > 0$ is the mass of a nucleon and $m_2 > 0$ is the mass of a meson.

In order to scale the KGS (1.1), we introduce

$$\tilde{t} = \frac{t}{t_s}, \quad \tilde{\mathbf{x}} = \frac{\mathbf{x}}{x_s}, \quad \tilde{\psi}(\tilde{\mathbf{x}},\tilde{t}) = x_s^{d/2}\psi(\mathbf{x},t), \quad \tilde{\phi}(\tilde{\mathbf{x}},\tilde{t}) = \frac{\phi(\mathbf{x},t)}{\phi_s}, \quad (1.2)$$

where x_s , t_s and ϕ_s are the dimensionless length unit, time unit and meson field unit, respectively, satisfying $t_s = \frac{2m_1x_s^2}{\hbar}$ and $\phi_s = \frac{\hbar x_s^{-d/2}}{\sqrt{2m_1}}$ with $v = \frac{x_s}{t_s} = \frac{\hbar}{2m_1x_s}$ being the wave speed. Plugging (1.2) into (1.1), after a simple computation and then removing all $\tilde{\cdot}$, we obtain the following dimensionless KGS equations in d -dimensions ($d = 3, 2, 1$):

$$i\partial_t\psi(\mathbf{x},t) + \Delta\psi(\mathbf{x},t) + \lambda\phi(\mathbf{x},t)\psi(\mathbf{x},t) = 0, \quad \mathbf{x} \in \mathbb{R}^d, \quad (1.3a)$$

$$\varepsilon^2\partial_{tt}\phi(\mathbf{x},t) - \Delta\phi(\mathbf{x},t) + \frac{\mu^2}{\varepsilon^2}\phi(\mathbf{x},t) - \lambda|\psi(\mathbf{x},t)|^2 = 0, \quad (1.3b)$$

where ε is a dimensionless parameter inversely proportional to the speed of light given by

$$0 < \varepsilon := \frac{v}{c} = \frac{x_s}{t_s c} = \frac{\hbar}{2cm_1x_s} \leq 1, \quad (1.4)$$

and $\mu = \frac{m_2}{2m_1} > 0$ and $\lambda = \frac{g\sqrt{2m_1x_s^{2-d/2}}}{\hbar} \in \mathbb{R}$ are two dimensionless constants which are independent of ε .

We remark here that if one chooses the dimensionless length unit $x_s = \frac{\hbar}{2cm_1}$, $t_s = \frac{x_s}{c} = \frac{\hbar}{2c^2m_1}$ and $\phi_s = \frac{c^{d/2}\hbar^{1-d/2}}{(2m_1)^{(1-d)/2}}$ in (1.2), then $\varepsilon = 1$ in (1.4) and Eqs. (1.3) with $\varepsilon = 1$ take the form often appearing in the literature [2, 22, 42]. This choice of x_s is appropriate when the wave speed is at the same order of the speed of light. However, when the wave speed is much smaller than the speed of light, a different choice of

x_s is more appropriate. Note that the choice of x_s determines the observation scale of the time evolution of the particles and decides: (i) which phenomena are ‘visible’ by asymptotic analysis, and (ii) which phenomena can be resolved by discretization by specified spatial/temporal grids. In fact, there are two important parameter regimes: One is when $\varepsilon = 1$ ($\iff x_s = \frac{\hbar}{2cm_1}$, $t_s = \frac{\hbar}{2c^2m_1}$ and $\phi_s = \frac{e^{d/2}\hbar^{1-d/2}}{(2m_1)^{(1-d)/2}}$), then Eqs. (1.3) describe the case that wave speed is at the same order of the speed of light; the other one is when $0 < \varepsilon \ll 1$, then Eqs. (1.3) are in the nonrelativistic limit regime.

To study the dynamics of the KGS (1.3), the initial data is usually given as

$$\psi(\mathbf{x}, 0) = \psi_0(\mathbf{x}), \quad \phi(\mathbf{x}, 0) = \phi_0(\mathbf{x}), \quad \partial_t \phi(\mathbf{x}, 0) = \frac{1}{\varepsilon^2} \phi_1(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^d, \quad (1.5)$$

where the complex-valued function ψ_0 and the real-valued functions ϕ_0 and ϕ_1 are independent of ε . The KGS equations (1.3) are dispersive and time symmetric. They conserve the mass of the nucleon field

$$\|\psi(\cdot, t)\|_{L^2}^2 := \int_{\mathbb{R}^d} |\psi(\mathbf{x}, t)|^2 d\mathbf{x} \equiv \int_{\mathbb{R}^d} |\psi(\mathbf{x}, 0)|^2 d\mathbf{x} = \int_{\mathbb{R}^d} |\psi_0(\mathbf{x})|^2 d\mathbf{x}, \quad t \geq 0, \quad (1.6)$$

and the *Hamiltonian* or total energy

$$\begin{aligned} E(t) &:= \int_{\mathbb{R}^d} \left[\frac{1}{2} \left(\varepsilon^2 |\partial_t \phi|^2 + |\nabla \phi|^2 + \frac{\mu^2}{\varepsilon^2} |\phi|^2 \right) + |\nabla \psi|^2 - \lambda |\psi|^2 \phi \right] d\mathbf{x} \\ &\equiv \int_{\mathbb{R}^d} \left[\frac{1}{2} \left(\frac{1}{\varepsilon^2} |\phi_1|^2 + |\nabla \phi_0|^2 + \frac{\mu^2}{\varepsilon^2} |\phi_0|^2 \right) + |\nabla \psi_0|^2 - \lambda |\psi_0|^2 \phi_0 \right] d\mathbf{x} \\ &= E(0), \quad t \geq 0. \end{aligned} \quad (1.7)$$

For the KGS equations (1.3) with $\varepsilon = 1$, i.e. O(1)-speed of light regime, there are extensive analytical and numerical results in the literatures. For the existence and uniqueness as well as regularity, we refer to [3, 22–27, 42, 45] and references therein. For the numerical methods and comparison such as the finite difference time domain (FDTD) methods and Crank-Nicolson Fourier pseudospectral method, we refer to [13, 46, 50, 51] and references therein. However, for the KGS equations (1.3) with $0 < \varepsilon \ll 1$, i.e. nonrelativistic limit regime (or the scaled speed of light goes to infinity), the analysis and efficient computation of the KGS equations (1.3) are mathematically and numerically rather complicated issues. The main difficulty is due to that the solution is highly oscillatory in time and the corresponding energy functional $E(t) = O(\varepsilon^{-2})$ in (1.7) becomes unbounded when $\varepsilon \rightarrow 0$.

Formally, in the nonrelativistic limit regime, i.e. $0 < \varepsilon \ll 1$, similarly to the analysis of the nonrelativistic limit of the Klein-Gordon (KG) equation [38–41], taking the ansatz

$$\phi(\mathbf{x}, t) = e^{i\mu t/\varepsilon^2} z(\mathbf{x}, t) + e^{-i\mu t/\varepsilon^2} \bar{z}(\mathbf{x}, t) + o(\varepsilon), \quad \mathbf{x} \in \mathbb{R}^d, \quad t \geq 0, \quad (1.8)$$

where \bar{z} denotes the complex conjugate of a complex-valued function z , and plugging it into (1.3) and (1.5), we obtain the Schrödinger equations with wave operator as a limiting model

$$i\partial_t \psi(\mathbf{x}, t) + \Delta \psi(\mathbf{x}, t) = 0, \quad \mathbf{x} \in \mathbb{R}^d, \quad t > 0, \quad (1.9a)$$

$$2i\mu \partial_t z(\mathbf{x}, t) + \varepsilon^2 \partial_{tt} z(\mathbf{x}, t) - \Delta z(\mathbf{x}, t) = 0, \quad (1.9b)$$




Fig. 1.1: Plots of the solution of the KGS (1.3)-(1.5) with $d = 1$ for different ϵ , where $\text{Im}\{\psi\}$ denotes the imaginary part of ψ .

with the well-prepared initial data [6, 7]

$$z(\mathbf{x}, 0) = \frac{1}{2} \left[\phi_0(\mathbf{x}) - \frac{i}{\mu} \phi_1(\mathbf{x}) \right], \quad \partial_t z(\mathbf{x}, 0) = -\frac{i}{2\mu} \Delta z(\mathbf{x}, 0), \quad \psi(\mathbf{x}, 0) = \psi_0(\mathbf{x}). \quad (1.10)$$

In addition, after dropping the second term in (1.9b), formally we get the Schrödinger equations as another limiting model

$$i\partial_t \psi(\mathbf{x}, t) + \Delta \psi(\mathbf{x}, t) = 0, \quad (1.11a)$$

$$2i\mu \partial_t z(\mathbf{x}, t) - \Delta z(\mathbf{x}, t) = 0, \quad \mathbf{x} \in \mathbb{R}^d, \quad t > 0, \quad (1.11b)$$

with the initial data

$$z(\mathbf{x}, 0) = \frac{1}{2} \left[\phi_0(\mathbf{x}) - \frac{i}{\mu} \phi_1(\mathbf{x}) \right], \quad \psi(\mathbf{x}, 0) = \psi_0(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^d. \quad (1.12)$$

This formally suggests that the solution of (1.3) propagates highly oscillatory waves with amplitude at $O(1)$ and wavelength at $O(\varepsilon^2)$ and $O(1)$ in time and space, respectively, when $0 < \varepsilon \ll 1$. To illustrate this, Fig. 1.1 shows the solution of the KGS equations (1.3)-(1.5) with $d = 1$, $\mu = \lambda = 1$, $\psi_0(x) = \frac{1+i}{2} \operatorname{sech}(x^2)$, $\phi_0(x) = \frac{1}{2} e^{-x^2}$ and $\phi_1(x) = \frac{1}{\sqrt{2}} e^{-x^2}$ for different $0 < \varepsilon \leq 1$.

The highly temporal oscillatory nature of the solution to the KGS equations (1.3) causes severe burdens in practical computation, making the numerical approximation extremely challenging and costly in the regime of $0 < \varepsilon \ll 1$. Recently, different numerical methods have been proposed and/or analyzed for the nonlinear Klein-Gordon equation in the nonrelativistic limit regime in which the solution shares similar oscillatory behavior as that of the KGS equations (1.3), including the FDTD methods [9], exponential wave integrator Fourier pseudospectral (EWI-FP) method [9, 11], asymptotic preserving (AP) method [21], stroboscopic average method (SAM) [18, 19] and multiscale time integrator Fourier pseudospectral (MTI-FP) method [8], etc. Among them, the SAM and MTI-FP methods are uniformly convergent (UA) for $\varepsilon \in (0, 1]$, while the FDTD, EWI-FP and AP methods are not. From the practical computation point of view, the MTI-FP is much simpler and thus more efficient than the SAM method. The main aim of this paper is to propose and analyze a MTI-FP method for the KGS equations (1.3) in the nonrelativistic limit regime by adapting a multiscale decomposition by frequency to the solution at each time step and applying the Fourier pseudospectral discretization and exponential wave integrators for spatial and temporal derivatives, respectively. Two independent error bounds will be established for the MTI-FP, which imply that the MTI-FP converges uniformly and/or optimally for $0 < \varepsilon \leq 1$. The MTI-FP method is also applied to study numerical convergence rates of the KGS equations (1.3) to its limiting models (1.9)-(1.10) and (1.11)-(1.12).

The paper is organized as follows. In section 2, we introduce a multiscale decomposition for the KGS equations (1.3) based on frequency. A MTI-FP method is proposed in section 3, and its rigorous error bounds in energy space are established in section 4. Numerical results are reported in section 5. Finally, some conclusions are drawn in section 6. Throughout this paper, we adopt the standard Sobolev spaces [1] and use the notation $A \lesssim B$ to represent that there exists a generic constant $C > 0$, which is independent of time step τ (or n), mesh size h and ε , such that $|A| \leq CB$.

2 Multiscale decomposition

Let $\tau = \Delta t > 0$ be the time step size, and denote time steps by $t_n = n\tau$ for $n = 0, 1, \dots$. In this section, we present a multiscale decomposition for the solution of (1.3) on the time interval $[t_n, t_{n+1}]$ with given initial data at $t = t_n$ as

$$\phi(\mathbf{x}, t_n) = \phi_0^n(\mathbf{x}) = O(1), \quad \partial_t \phi(\mathbf{x}, t_n) = \frac{1}{\varepsilon^2} \phi_1^n(\mathbf{x}) = O\left(\frac{1}{\varepsilon^2}\right), \quad (2.1a)$$

$$\psi(\mathbf{x}, t_n) = \psi_0^n(\mathbf{x}) = O(1). \quad (2.1b)$$

Similarly to the analytical study of the nonlinear Klein-Gordon equation in the non-relativistic limit regime in [8, 40], we take an ansatz to the variable $\phi(\mathbf{x}, t) := \phi(\mathbf{x}, t_n + s)$ of (1.3b) on the time interval $[t_n, t_{n+1}]$ with (2.1) as [8, 12]

$$\phi(\mathbf{x}, t_n + s) = e^{i\mu s/\varepsilon^2} z^n(\mathbf{x}, s) + e^{-i\mu s/\varepsilon^2} \bar{z}^n(\mathbf{x}, s) + r^n(\mathbf{x}, s), \quad \mathbf{x} \in \mathbb{R}^d, \quad 0 \leq s \leq \tau. \quad (2.2)$$

Differentiating (2.2) with respect to s , we have

$$\begin{aligned} \partial_s \phi(\mathbf{x}, t_n + s) = & e^{i\mu s/\varepsilon^2} \left[\partial_s z^n(\mathbf{x}, s) + \frac{i\mu}{\varepsilon^2} z^n(\mathbf{x}, s) \right] + \partial_s r^n(\mathbf{x}, s) \\ & + e^{-i\mu s/\varepsilon^2} \left[\partial_s \bar{z}^n(\mathbf{x}, s) - \frac{i\mu}{\varepsilon^2} \bar{z}^n(\mathbf{x}, s) \right], \quad \mathbf{x} \in \mathbb{R}^d, \quad 0 \leq s \leq \tau. \end{aligned} \quad (2.3)$$

Plugging (2.2) into (1.3), we get for $\mathbf{x} \in \mathbb{R}^d$, $0 \leq s \leq \tau$ and $\psi(\mathbf{x}, t_n + s) =: \psi^n(\mathbf{x}, s)$

$$\begin{aligned} & e^{i\mu s/\varepsilon^2} \left[\varepsilon^2 \partial_{ss} z^n(\mathbf{x}, s) + 2i\mu \partial_s z^n(\mathbf{x}, s) - \Delta z^n(\mathbf{x}, s) \right] \\ & + e^{-i\mu s/\varepsilon^2} \left[\varepsilon^2 \partial_{ss} \bar{z}^n(\mathbf{x}, s) - 2i\mu \partial_s \bar{z}^n(\mathbf{x}, s) - \Delta \bar{z}^n(\mathbf{x}, s) \right] \\ & + \varepsilon^2 \partial_{ss} r^n(\mathbf{x}, s) - \Delta r^n(\mathbf{x}, s) + \frac{\mu^2}{\varepsilon^2} r^n(\mathbf{x}, s) = \lambda |\psi^n(\mathbf{x}, s)|^2. \end{aligned}$$

Substituting (2.2) into (1.3a) and multiplying the above equation by $e^{-i\mu s/\varepsilon^2}$ and $e^{i\mu s/\varepsilon^2}$, respectively, we can re-formulate the KGS equations (1.3) for $\psi^n := \psi^n(\mathbf{x}, s)$, $z^n := z^n(\mathbf{x}, s)$ and $r^n := r^n(\mathbf{x}, s)$ as

$$\begin{cases} i\partial_s \psi^n + \Delta \psi^n + \lambda \left(e^{\frac{i\mu s}{\varepsilon^2}} z^n \psi^n + e^{-\frac{i\mu s}{\varepsilon^2}} \bar{z}^n \psi^n + r^n \psi^n \right) = 0, \\ 2i\mu \partial_s z^n + \varepsilon^2 \partial_{ss} z^n - \Delta z^n = 0, & \mathbf{x} \in \mathbb{R}^d, \quad 0 \leq s \leq \tau, \\ \varepsilon^2 \partial_{ss} r^n - \Delta r^n + \frac{\mu^2}{\varepsilon^2} r^n = \lambda |\psi^n|^2. \end{cases} \quad (2.4)$$

In order to find proper initial conditions for the system (2.4), setting $s = 0$ in (2.2) and (2.3), noticing (2.1a), we obtain

$$\begin{cases} z^n(\mathbf{x}, 0) + \bar{z}^n(\mathbf{x}, 0) + r^n(\mathbf{x}, 0) = \phi_0^n(\mathbf{x}), & \mathbf{x} \in \mathbb{R}^d, \\ \frac{i\mu}{\varepsilon^2} [z^n(\mathbf{x}, 0) - \bar{z}^n(\mathbf{x}, 0)] + \partial_s z^n(\mathbf{x}, 0) + \partial_s \bar{z}^n(\mathbf{x}, 0) + \partial_s r^n(\mathbf{x}, 0) = \frac{\phi_1^n(\mathbf{x})}{\varepsilon^2}. \end{cases} \quad (2.5)$$

Now we decompose the above initial data so as to: (i) equate $O\left(\frac{1}{\varepsilon^2}\right)$ and $O(1)$ terms in the second equation of (2.5), respectively, and (ii) be well-prepared for the second equation in (2.4) when $0 < \varepsilon \ll 1$, i.e. $\partial_s z^n(\mathbf{x}, 0)$ is determined from the second equation in (2.4), by setting $\varepsilon = 0$ and $s = 0$ [6–8]:

$$\begin{cases} z^n(\mathbf{x}, 0) + \overline{z^n}(\mathbf{x}, 0) = \phi_0^n(\mathbf{x}), & i\mu [z^n(\mathbf{x}, 0) - \overline{z^n}(\mathbf{x}, 0)] = \phi_1^n(\mathbf{x}), \\ 2i\mu \partial_s z^n(\mathbf{x}, 0) - \Delta z^n(\mathbf{x}, 0) = 0, & \mathbf{x} \in \mathbb{R}^d, \\ r^n(\mathbf{x}, 0) = 0, & \partial_s r^n(\mathbf{x}, 0) + \partial_s z^n(\mathbf{x}, 0) + \partial_s \overline{z^n}(\mathbf{x}, 0) = 0. \end{cases} \quad (2.6)$$

Solving (2.6) and noticing (2.1b), we get the initial data for (2.4) as

$$\begin{cases} z^n(\mathbf{x}, 0) = \frac{1}{2} \left[\phi_0^n(\mathbf{x}) - \frac{i}{\mu} \phi_1^n(\mathbf{x}) \right], & \partial_s z^n(\mathbf{x}, 0) = -\frac{i}{2\mu} \Delta z^n(\mathbf{x}, 0), & \mathbf{x} \in \mathbb{R}^d, \\ \psi^n(\mathbf{x}, 0) = \psi_0^n(\mathbf{x}), & r^n(\mathbf{x}, 0) = 0, & \partial_s r^n(\mathbf{x}, 0) = -\partial_s z^n(\mathbf{x}, 0) - \partial_s \overline{z^n}(\mathbf{x}, 0). \end{cases} \quad (2.7)$$

The above decomposition can be called as multiscale decomposition by frequency (MDF). In fact, it can also be regarded as to decompose slow waves at ε^2 -wavelength and fast waves at other wavelengths, thus it can also be called as fast-slow frequency decomposition. After solving the decomposed system (2.4) with the initial data (2.7), we get $\psi^n(\mathbf{x}, \tau)$, $z^n(\mathbf{x}, \tau)$, $\partial_s z^n(\mathbf{x}, \tau)$, $r^n(\mathbf{x}, \tau)$ and $\partial_s r^n(\mathbf{x}, \tau)$. Then we can reconstruct the solution to the KGS equations (1.3) at $t = t_{n+1}$ by setting $s = \tau$ in (2.2) and (2.3), i.e.

$$\begin{cases} \psi(\mathbf{x}, t_{n+1}) = \psi^n(\mathbf{x}, \tau) =: \psi_0^{n+1}(\mathbf{x}), \\ \phi(\mathbf{x}, t_{n+1}) = e^{i\mu\tau/\varepsilon^2} z^n(\mathbf{x}, \tau) + e^{-i\mu\tau/\varepsilon^2} \overline{z^n}(\mathbf{x}, \tau) + r^n(\mathbf{x}, \tau) =: \phi_0^{n+1}(\mathbf{x}), \\ \partial_t \phi(\mathbf{x}, t_{n+1}) = \frac{1}{\varepsilon^2} \phi_1^{n+1}(\mathbf{x}), & \mathbf{x} \in \mathbb{R}^d, \end{cases} \quad (2.8)$$

with

$$\begin{aligned} \phi_1^{n+1}(\mathbf{x}) := & e^{i\mu\tau/\varepsilon^2} [\varepsilon^2 \partial_s z^n(\mathbf{x}, \tau) + i\mu z^n(\mathbf{x}, \tau)] + e^{-i\mu\tau/\varepsilon^2} [\varepsilon^2 \partial_s \overline{z^n}(\mathbf{x}, \tau) - i\mu \overline{z^n}(\mathbf{x}, \tau)] \\ & + \varepsilon^2 \partial_s r^n(\mathbf{x}, \tau). \end{aligned}$$

In summary, the MDF proceeds as a decomposition–solution–reconstruction flow at each time interval, and this makes it essentially different from the classical modulated Fourier expansion [14–17, 34–36] which only carries out the decomposition at the initial time $t = 0$.

3 A MTI-FP method

For the simplicity of notations and without loss of generality, we take $\mu = \lambda = 1$ in (1.3) and present our numerical method in one space dimension (1D). Generalizations to higher dimensions are straightforward and the results remain valid. We truncate the whole-space problem (1.3) into a finite interval $\Omega = (a, b)$ with periodic boundary conditions, where due to the fast decay of the solution at far field, the truncation error

can be negligible by choosing a, b sufficiently large. In 1D, the problem (1.3) with $\mu = \lambda = 1$ collapses to

$$\begin{cases} i\partial_t \psi(x, t) + \partial_{xx} \psi(x, t) + \phi(x, t) \psi(x, t) = 0, & x \in \Omega, \quad t > 0, \\ \varepsilon^2 \partial_{tt} \phi(x, t) - \partial_{xx} \phi(x, t) + \frac{1}{\varepsilon^2} \phi(x, t) = |\psi(x, t)|^2, & x \in \Omega, \quad t > 0, \\ \phi(a, t) = \phi(b, t), \quad \partial_x \phi(a, t) = \partial_x \phi(b, t), \quad \psi(a, t) = \psi(b, t) & t \geq 0, \\ \phi(x, 0) = \phi_0(x), \quad \partial_t \phi(x, 0) = \frac{\phi_1(x)}{\varepsilon^2}, \quad \psi(x, 0) = \psi_0(x), & x \in \overline{\Omega} = [a, b]. \end{cases} \quad (3.1)$$

We remark that the boundary conditions considered here are inspired by the inherent physical nature of the system and they have been widely used in the literatures for the simulation of the KGS equations [10, 13, 50, 51].

Consequently, for $n \geq 0$, the decomposed system MDF (2.4) in 1D collapses to

$$\begin{cases} i\partial_s \psi^n + \partial_{xx} \psi^n + e^{is/\varepsilon^2} z^n \psi^n + e^{-is/\varepsilon^2} \bar{z}^n \psi^n + r^n \psi^n = 0, \\ 2i\partial_s z^n + \varepsilon^2 \partial_{ss} z^n - \partial_{xx} z^n = 0, & a < x < b, \quad 0 < s \leq \tau, \\ \varepsilon^2 \partial_{ss} r^n - \partial_{xx} r^n + \frac{1}{\varepsilon^2} r^n = |\psi^n|^2, \end{cases} \quad (3.2)$$

The initial and boundary conditions for the above system are

$$\begin{cases} z^n(a, s) = z^n(b, s), \quad \partial_x z^n(a, s) = \partial_x z^n(b, s), \\ r^n(a, s) = r^n(b, s), \quad \partial_x r^n(a, s) = \partial_x r^n(b, s), \\ \psi^n(a, s) = \psi^n(b, s), \quad \partial_x \psi^n(a, s) = \partial_x \psi^n(b, s), & 0 \leq s \leq \tau; \\ z^n(x, 0) = \frac{1}{2} [\phi_0^n(x) - i\phi_1^n(x)], \quad \partial_s z^n(x, 0) = -\frac{i}{2} \partial_{xx} z^n(x, 0), \\ r^n(x, 0) = 0, \quad \partial_s r^n(x, 0) = -\partial_s z^n(x, 0) - \partial_s \bar{z}^n(x, 0), \\ \psi^n(x, 0) = \psi_0^n(x), & a \leq x \leq b. \end{cases} \quad (3.3)$$

In what follows, we present a numerical integrator Fourier pseudospectral discretization for the MDF (3.2) with (3.3), which applies the Fourier pseudospectral discretization to spatial derivatives followed by using some proper exponential wave integrators (EWI) for temporal discretizations in phase (Fourier) space.

Choose the mesh size $h := \Delta x = (b - a)/N$ with N a positive integer and denote grid points as $x_j := a + jh$ for $j = 0, 1, \dots, N$. Define

$$X_N := \text{span} \left\{ e^{i\mu_l(x-a)} : x \in \overline{\Omega}, \mu_l = \frac{2\pi l}{b-a}, l = -\frac{N}{2}, \dots, \frac{N}{2} - 1 \right\},$$

$$Y_N := \left\{ \mathbf{v} = (v_0, v_1, \dots, v_N) \in \mathbb{C}^{N+1} : v_0 = v_N \right\}, \quad \|\mathbf{v}\|_{l^2} = h \sum_{j=0}^{N-1} |v_j|^2.$$

For a periodic function $v(x)$ on $\overline{\Omega}$ and a vector $\mathbf{v} \in Y_N$, let $P_N : L^2(\Omega) \rightarrow X_N$ be the standard L^2 -projection operator, and $I_N : C(\Omega) \rightarrow X_N$ or $Y_N \rightarrow X_N$ be the trigonometric interpolation operator [31, 48], i.e.

$$(P_N v)(x) = \sum_{l=-N/2}^{N/2-1} \widehat{v}_l e^{i\mu_l(x-a)}, \quad (I_N \mathbf{v})(x) = \sum_{l=-N/2}^{N/2-1} \widetilde{v}_l e^{i\mu_l(x-a)}, \quad (3.4)$$

where \widehat{v}_l and \widetilde{v}_l are the Fourier and discrete Fourier transform coefficients of the periodic function $v(x)$ and vector \mathbf{v} , respectively, defined as

$$\widehat{v}_l = \frac{1}{b-a} \int_a^b v(x) e^{-i\mu_l(x-a)} dx, \quad \widetilde{v}_l = \frac{1}{N} \sum_{j=0}^{N-1} v_j e^{-i\mu_l(x_j-a)}. \quad (3.5)$$

Then a Fourier spectral method for discretizing (3.2) reads:

Find $z_N^n := z_N^n(x, s)$, $r_N^n := r_N^n(x, s)$, $\psi_N^n := \psi_N^n(x, s) \in X_N$ for $0 \leq s \leq \tau$, i.e.

$$\begin{cases} z_N^n(x, s) = \sum_{l=-N/2}^{N/2-1} (\widehat{z_N^n})_l(s) e^{i\mu_l(x-a)}, & r_N^n(x, s) = \sum_{l=-N/2}^{N/2-1} (\widehat{r_N^n})_l(s) e^{i\mu_l(x-a)}, \\ \psi_N^n(x, s) = \sum_{l=-N/2}^{N/2-1} (\widehat{\psi_N^n})_l(s) e^{i\mu_l(x-a)}, \end{cases} \quad (3.6)$$

such that

$$\begin{cases} i\partial_s \psi_N^n + \partial_{xx} \psi_N^n + e^{is/\varepsilon^2} P_N(z_N^n \psi_N^n) + e^{-is/\varepsilon^2} P_N(\overline{z_N^n} \overline{\psi_N^n}) + P_N(r_N^n \psi_N^n) = 0, \\ 2i\partial_s z_N^n + \varepsilon^2 \partial_{ss} z_N^n - \partial_{xx} z_N^n = 0, & 0 < s \leq \tau, \quad a < x < b, \\ \varepsilon^2 \partial_{ss} r_N^n - \partial_{xx} r_N^n + \frac{1}{\varepsilon^2} r_N^n = P_N(|\psi_N^n|^2). \end{cases} \quad (3.7)$$

Substituting (3.6) into (3.7) and noticing the orthogonality of the Fourier basis, we get

$$\begin{cases} i(\widehat{\psi_N^n})_l'(s) - \mu_l^2 (\widehat{\phi_N^n})_l(s) + e^{\frac{is}{\varepsilon^2}} (\widehat{z_N^n \psi_N^n})_l(s) + e^{-\frac{is}{\varepsilon^2}} (\widehat{\overline{z_N^n} \overline{\psi_N^n}})_l(s) + (\widehat{r_N^n \psi_N^n})_l(s) = 0, \\ 2i(\widehat{z_N^n})_l'(s) + \varepsilon^2 (\widehat{z_N^n})_l''(s) + \mu_l^2 (\widehat{z_N^n})_l(s) = 0, & 0 < s \leq \tau, \\ \varepsilon^2 (\widehat{r_N^n})_l''(s) + \left(\mu_l^2 + \frac{1}{\varepsilon^2}\right) (\widehat{r_N^n})_l(s) = (\widehat{|\psi_N^n|^2})_l(s), & -\frac{N}{2} \leq l \leq \frac{N}{2} - 1. \end{cases} \quad (3.8)$$

For each $-N/2 \leq l \leq N/2 - 1$, we can rewrite (3.8) by using the variation-of-constant formula as

$$(\widehat{z_N^n})_l(s) = a_l(s) (\widehat{z_N^n})_l(0) + \varepsilon^2 b_l(s) (\widehat{z_N^n})_l'(0), \quad 0 \leq s \leq \tau, \quad (3.9a)$$

$$(\widehat{r_N^n})_l(s) = \frac{\sin(\omega_l s)}{\omega_l} (\widehat{r_N^n})_l'(0) + \int_0^s \frac{\sin(\omega_l(s-\theta))}{\varepsilon^2 \omega_l} (\widehat{|\psi_N^n|^2})_l(\theta) d\theta, \quad (3.9b)$$

$$\begin{aligned} (\widehat{\psi_N^n})_l(s) &= e^{-i\mu_l^2 s} (\widehat{\psi_N^n})_l(0) + i e^{-i\mu_l^2 s} \int_0^s e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} (\widehat{z_N^n \psi_N^n})_l(\theta) d\theta \\ &\quad + i e^{-i\mu_l^2 s} \int_0^s e^{i(\mu_l^2 - \frac{1}{\varepsilon^2})\theta} (\widehat{\overline{z_N^n} \overline{\psi_N^n}})_l(\theta) d\theta + i \int_0^s e^{i\mu_l^2(\theta-s)} (\widehat{r_N^n \psi_N^n})_l(\theta) d\theta, \end{aligned} \quad (3.9c)$$

where

$$\begin{cases} a_l(s) := \frac{\lambda_l^+ e^{is\lambda_l^-} - \lambda_l^- e^{is\lambda_l^+}}{\lambda_l^+ - \lambda_l^-}, & b_l(s) := i \frac{e^{is\lambda_l^+} - e^{is\lambda_l^-}}{\varepsilon^2 (\lambda_l^- - \lambda_l^+)}, & 0 \leq s \leq \tau, \\ \lambda_l^\pm = -\frac{1}{\varepsilon^2} \left(1 \pm \sqrt{1 + \mu_l^2 \varepsilon^2}\right), & \omega_l = \frac{1}{\varepsilon^2} \sqrt{1 + \mu_l^2 \varepsilon^2}. \end{cases} \quad (3.10)$$

Differentiating (3.9a) and (3.9b) with respect to s , we obtain

$$\widehat{(z_N^n)}'_l(s) = a'_l(s)\widehat{(z_N^n)}_l(0) + \varepsilon^2 b'_l(s)\widehat{(z_N^n)}'_l(0), \quad 0 \leq s \leq \tau, \quad (3.11a)$$

$$\widehat{(r_N^n)}'_l(s) = \cos(\omega_l s)\widehat{(r_N^n)}'_l(0) + \int_0^s \frac{\cos(\omega_l(s-\theta))}{\varepsilon^2} (\widehat{|\psi_N^n|^2})_l(\theta) d\theta, \quad (3.11b)$$

where

$$a'_l(s) = i\lambda_l^+ \lambda_l^- \frac{e^{is\lambda_l^-} - e^{is\lambda_l^+}}{\lambda_l^+ - \lambda_l^-}, \quad b'_l(s) = \frac{\lambda_l^+ e^{is\lambda_l^+} - \lambda_l^- e^{is\lambda_l^-}}{\varepsilon^2(\lambda_l^+ - \lambda_l^-)}, \quad 0 \leq s \leq \tau.$$

Taking $s = \tau$ in (3.9) and (3.11), we immediately get

$$\begin{aligned} \widehat{(z_N^n)}_l(\tau) &= a_l(\tau)\widehat{(z_N^n)}_l(0) + \varepsilon^2 b_l(\tau)\widehat{(z_N^n)}'_l(0), \\ \widehat{(z_N^n)}'_l(\tau) &= a'_l(\tau)\widehat{(z_N^n)}_l(0) + \varepsilon^2 b'_l(\tau)\widehat{(z_N^n)}'_l(0). \end{aligned} \quad (3.12)$$

In order to approximate the definite integrals in (3.9b), (3.9c) and (3.11b) with $s = \tau$, we adapt the Gautschi's type quadrature [8, 12, 28–30, 33, 33]

$$\int_0^\tau e^{i\delta\theta} f(\theta) d\theta \approx \int_0^\tau e^{i\delta\theta} [f(0) + \theta f'(0)] d\theta = \frac{i - ie^{i\delta\tau}}{\delta} f(0) + \frac{(1 - i\delta\tau)e^{i\delta\tau} - 1}{\delta^2} f'(0)$$

except the last term in (3.9c) which is approximated via a combination of the Gautschi's and Deuffhard's quadrature as [12, 20, 36]

$$\begin{aligned} \int_0^\tau e^{i\mu_l^2(\theta-\tau)} \widehat{(r_N^n \psi_N^n)}_l(\theta) d\theta &\approx \int_0^\tau e^{i\mu_l^2(\theta-\tau)} \left[\widehat{(A_1)}_l(\theta) + \theta \widehat{(A_2)}_l(\theta) \right] d\theta \\ &\approx \frac{\tau}{2} \left[\widehat{(A_1)}_l(\tau) + \tau \widehat{(A_2)}_l(\tau) \right], \end{aligned} \quad (3.13)$$

where $A_1(x, \theta) := r_N^n(x, \theta)\psi_N^n(x, 0)$ and $A_2(x, \theta) := r_N^n(x, \theta)\partial_s \psi_N^n(x, 0)$ for $0 \leq \theta \leq \tau$ and thus $A_1(x, 0) = A_2(x, 0) \equiv 0$ since $r_N^n(x, 0) \equiv 0$. Thus (3.9b), (3.9c) and (3.11b) with $s = \tau$ can be approximated as

$$\left\{ \begin{aligned} \widehat{(r_N^n)}_l(\tau) &\approx \frac{\sin(\omega_l \tau)}{\omega_l} \widehat{(r_N^n)}'_l(0) + p_l(\tau) \widehat{(|\psi_N^n|^2)}_l(0) + q_l(\tau) \widehat{(|\psi_N^n|^2)}'_l(0), \\ \widehat{(r_N^n)}'_l(\tau) &\approx \cos(\omega_l \tau) \widehat{(r_N^n)}'_l(0) + p'_l(\tau) \widehat{(|\psi_N^n|^2)}_l(0) + q'_l(\tau) \widehat{(|\psi_N^n|^2)}'_l(0), \\ \widehat{(\psi_N^n)}_l(\tau) &\approx e^{-i\mu_l^2 \tau} \widehat{(\psi_N^n)}_l(0) + c_l^+(\tau) \widehat{(z_N^n \psi_N^n)}_l(0) + d_l^+(\tau) \widehat{(z_N^n \psi_N^n)}'_l(0) \\ &\quad + c_l^-(\tau) \widehat{(\overline{z_N^n} \overline{\psi_N^n})}_l(0) + d_l^-(\tau) \widehat{(\overline{z_N^n} \overline{\psi_N^n})}'_l(0) + \frac{i\tau}{2} \left[\widehat{(A_1)}_l(\tau) + \tau \widehat{(A_2)}_l(\tau) \right], \end{aligned} \right. \quad (3.14)$$

where $p_l(\tau)$, $q_l(\tau)$, $p'_l(\tau)$, $q'_l(\tau)$, $c_l^\pm(\tau)$ and $d_l^\pm(\tau)$ are given in Appendix A. For the derivatives, we can compute them as

$$\widehat{(|\psi_N^n|^2)}'_l(0) = 2(\operatorname{Re}\{\widehat{\psi_N^n} \partial_s \overline{\psi_N^n}\})_l(0), \quad \widehat{(z_N^n \psi_N^n)}'_l(0) = (\partial_s z_N^n \psi_N^n)_l(0) + (z_N^n \partial_s \psi_N^n)_l(0),$$

where $\text{Re}\{z\}$ denotes the real part of a complex number z and for $-\frac{N}{2} \leq l \leq \frac{N}{2} - 1$,

$$\widehat{(z_N^n)}'_l(0) = \frac{i \sin(\mu_l^2 \tau)}{2\tau} \widehat{(z_N^n)}_l(0), \quad \widehat{(\psi_N^n)}'_l(0) = -i \frac{\sin(\mu_l^2 \tau)}{\tau} \widehat{(\psi_N^n)}_l(0) + i \widehat{(\phi_N^n \psi_N^n)}_l(0),$$

which are approximations of $\partial_s z^n(x, 0) = -\frac{i}{2} \partial_{xx} z^n(x, 0)$ and $\partial_s \psi^n(x, 0) = i \partial_{xx} \psi^n(x, 0) + i \phi^n(x, 0) \psi^n(x, 0)$, respectively [8]. Inserting (3.12) and (3.14) into (3.6) with setting $s = \tau$, and noticing (2.8), we immediately obtain a multiscale time integrator Fourier spectral method based on the MDF (3.2) for the problem (3.1).

In practice, the integrals for computing the Fourier transform coefficients in (3.5), (3.9) and (3.11) are usually approximated by the numerical quadratures [9, 11, 48] given in (3.11). Let Φ_j^n , $\dot{\Phi}_j^n$ and Ψ_j^n be approximations of $\phi(x_j, t_n)$, $\partial_t \phi(x_j, t_n)$ and $\psi(x_j, t_n)$, respectively; Z_j^{n+1} , \dot{Z}_j^{n+1} , R_j^{n+1} and \dot{R}_j^{n+1} be approximations of $z^n(x_j, \tau)$, $\partial_s z^n(x_j, \tau)$, $r^n(x_j, \tau)$ and $\partial_s r^n(x_j, \tau)$, respectively, for $j = 0, 1, \dots, N$. Choosing $\Phi_j^0 = \phi_0(x_j)$, $\dot{\Phi}_j^0 = \phi_1(x_j)/\varepsilon^2$ and $\Psi_j^0 = \psi_0(x_j)$ for $0 \leq j \leq N$, a multiscale time integrator Fourier pseudospectral (MTI-FP) discretization for the problem (3.1) reads for $n \geq 0$,

$$\begin{cases} \Phi_j^{n+1} = e^{i\tau/\varepsilon^2} Z_j^{n+1} + e^{-i\tau/\varepsilon^2} \overline{Z_j^{n+1}} + R_j^{n+1}, & j = 0, 1, \dots, N, \\ \dot{\Phi}_j^{n+1} = e^{i\tau/\varepsilon^2} \left(\dot{Z}_j^{n+1} + \frac{i}{\varepsilon^2} Z_j^{n+1} \right) + e^{-i\tau/\varepsilon^2} \left(\overline{\dot{Z}_j^{n+1}} - \frac{i}{\varepsilon^2} \overline{Z_j^{n+1}} \right) + \dot{R}_j^{n+1}, \\ \Psi_j^{n+1} = \sum_{l=-\frac{N}{2}}^{\frac{N}{2}-1} \widehat{(\Psi^{n+1})}_l e^{i\mu_l(x_j-a)}, \end{cases} \quad (3.15)$$

where

$$\begin{cases} Z_j^{n+1} = \sum_{l=-\frac{N}{2}}^{\frac{N}{2}-1} \widehat{(Z^{n+1})}_l e^{i\mu_l(x_j-a)}, & R_j^{n+1} = \sum_{l=-\frac{N}{2}}^{\frac{N}{2}-1} \widehat{(R^{n+1})}_l e^{i\mu_l(x_j-a)}, \\ \dot{Z}_j^{n+1} = \sum_{l=-\frac{N}{2}}^{\frac{N}{2}-1} \widehat{(\dot{Z}^{n+1})}_l e^{i\mu_l(x_j-a)}, & \dot{R}_j^{n+1} = \sum_{l=-\frac{N}{2}}^{\frac{N}{2}-1} \widehat{(\dot{R}^{n+1})}_l e^{i\mu_l(x_j-a)}, \end{cases}$$

and for $-N/2 \leq l \leq N/2 - 1$,

$$\widehat{(Z^{n+1})}_l = a_l(\tau) \widehat{(Z^0)}_l + \varepsilon^2 b_l(\tau) \widehat{(\dot{Z}^0)}_l, \quad \widehat{(\dot{Z}^{n+1})}_l = a'_l(\tau) \widehat{(Z^0)}_l + \varepsilon^2 b'_l(\tau) \widehat{(\dot{Z}^0)}_l, \quad (3.16a)$$

$$\widehat{(R^{n+1})}_l = \frac{\sin(\omega_l \tau)}{\omega_l} \widehat{(\dot{R}^0)}_l + p_l(\tau) (|\widehat{(\Psi^n)}|^2)_l + 2q_l(\tau) (\text{Re}\{\widehat{(\Psi^n \dot{\Psi}^n)}\})_l, \quad (3.16b)$$

$$\widehat{(\dot{R}^{n+1})}_l = \cos(\omega_l \tau) \widehat{(\dot{R}^0)}_l + p'_l(\tau) (|\widehat{(\Psi^n)}|^2)_l + 2q'_l(\tau) (\text{Re}\{\widehat{(\Psi^n \dot{\Psi}^n)}\})_l, \quad (3.16c)$$

$$\begin{aligned} \widehat{(\Psi^{n+1})}_l &= e^{-i\mu_l^2 \tau} \widehat{(\Psi^n)}_l + c_l^+(\tau) \widehat{(Z^0 \Psi^n)}_l + d_l^+(\tau) \left[\widehat{(\dot{Z}^0 \Psi^n)}_l + \widehat{(Z^0 \dot{\Psi}^n)}_l \right] \\ &\quad + c_l^-(\tau) \widehat{(\overline{Z^0 \Psi^n})}_l + d_l^-(\tau) \left[\widehat{(\overline{\dot{Z}^0 \Psi^n})}_l + \widehat{(\overline{Z^0 \dot{\Psi}^n})}_l \right] + \frac{i\tau}{2} \widehat{(R^{n+1} \Psi^n)}_l \\ &\quad + \frac{i\tau^2}{2} \widehat{(R^{n+1} \dot{\Psi}^n)}_l, \end{aligned} \quad (3.16d)$$

with

$$(\widetilde{Z^0})_l = \frac{1}{2} \left[(\widetilde{\Phi^n})_l - i\varepsilon^2 (\widetilde{\dot{\Phi}^n})_l \right], \quad (\widetilde{Z^0})_l = \frac{i \sin(\mu_l^2 \tau)}{2\tau} (\widetilde{Z^0})_l, \quad (3.17a)$$

$$(\widetilde{R^0})_l = -(\widetilde{Z^0})_l - (\widetilde{\dot{Z}^0})_l, \quad (\widetilde{\Psi^n})_l = -\frac{i \sin(\mu_l^2 \tau)}{\tau} (\widetilde{\Psi^n})_l + i(\widetilde{\Phi^n \Psi^n})_l. \quad (3.17b)$$

This MTI-FP method for the KGS equations (3.1) (or (1.3)) is explicit, accurate, easy to implement and very efficient due to the discrete fast Fourier transform. The memory cost is $O(N)$ and the computational cost per time step is $O(N \log N)$.

4 Uniform convergence of MTI-FP

In this section, we establish an error bound for the MTI-FP method (3.15) of the KGS (3.1), which is uniformly for $\varepsilon \in (0, 1]$. Let $0 < T < T^*$ with T^* the maximum existence time of the solution of the problem (3.1), motivated by the formal asymptotic results (1.8) and (1.9) as well as the numerical results (cf. Fig. 1.1), we make the following assumptions on the solution of the problem (3.1) – there exists an integer $m_0 \geq 4$ such that $\phi \in C^1([0, T]; H_p^{m_0+4}(\Omega))$, $\psi \in C([0, T]; H_p^{m_0+2}(\Omega)) \cap C^1([0, T]; H_p^{m_0}(\Omega)) \cap C^2([0, T]; H_p^{m_0-2}(\Omega))$ and

$$\begin{aligned} \|\phi\|_{L^\infty([0, T]; H^{m_0+4})} + \varepsilon^2 \|\partial_t \phi\|_{L^\infty([0, T]; H^{m_0+4})} &\lesssim 1, \\ \|\psi\|_{L^\infty([0, T]; H^{m_0+2})} + \|\partial_t \psi\|_{L^\infty([0, T]; H^{m_0})} + \varepsilon^2 \|\partial_{tt} \psi\|_{L^\infty([0, T]; H^{m_0-2})} &\lesssim 1, \end{aligned} \quad (4.1)$$

where $H_p^m(\Omega) = \{f(x) \in H^m(\Omega) \mid f^{(k)}(a) = f^{(k)}(b), k = 0, 1, \dots, m-1\} \subset H^m(\Omega)$. From the first equation in (3.1), i.e. the Schrödinger equation, it is easy to see that

$$i\partial_t \rho(x, t) + \partial_x [\overline{\psi} \partial_x \psi(x, t) - \psi \partial_x \overline{\psi}(x, t)] = 0, \quad x \in \Omega, t > 0,$$

where $\rho(x, t) := |\psi(x, t)|^2$. Then under the assumption (4.1), we have

$$\begin{aligned} \rho \in C([0, T]; H_p^{m_0+2}(\Omega)) \cap C^1([0, T]; H_p^{m_0}(\Omega)) \cap C^2([0, T]; H_p^{m_0-2}(\Omega)), \\ \|\partial_t^k \rho\|_{L^\infty([0, T]; H^{m_0+2-2k}(\Omega))} \lesssim 1, \quad k = 0, 1, 2. \end{aligned} \quad (4.2)$$

Denote $C_\phi = \max_{0 < \varepsilon \leq 1} \left\{ \|\phi\|_{L^\infty([0, T]; H^{m_0+4}(\Omega))}, \varepsilon^2 \|\partial_t \phi\|_{L^\infty([0, T]; H^{m_0+4}(\Omega))} \right\}$ and $C_\psi = \max_{0 < \varepsilon \leq 1} \left\{ \|\psi\|_{L^\infty([0, T]; H^{m_0+2}(\Omega))} \right\}$.

Let $\Phi^n = (\Phi_0^n, \Phi_1^n, \dots, \Phi_N^n) \in Y_N$, $\Psi^n = (\Psi_0^n, \Psi_1^n, \dots, \Psi_N^n) \in Y_N$ and $\dot{\Phi}^n = (\dot{\Phi}_0^n, \dot{\Phi}_1^n, \dots, \dot{\Phi}_N^n) \in Y_N$ ($n \geq 0$) are the numerical solution obtained from the MTI-FP method (3.15)-(3.17), denote their interpolations as

$$\phi_I^n(x) := (I_N \Phi^n)(x), \quad \psi_I^n(x) := (I_N \Psi^n)(x), \quad \dot{\phi}_I^n(x) := (I_N \dot{\Phi}^n)(x), \quad x \in \overline{\Omega}, \quad (4.3)$$

and define the error functions as

$$\begin{aligned} e_\phi^n(x) &:= \phi(x, t_n) - \phi_I^n(x), \quad e_\psi^n(x) := \psi(x, t_n) - \psi_I^n(x), \\ \dot{e}_\phi^n(x) &:= \partial_t \phi(x, t_n) - \dot{\phi}_I^n(x), \quad x \in \overline{\Omega}, \quad 0 \leq n \leq \frac{T}{\tau}; \end{aligned} \quad (4.4)$$

then we have the following error estimates for the MTI-FP method (3.15)-(3.17).

Theorem 4.1 (Error bounds of MTI-FP) *Under the assumption (4.1), there exist two constants $0 < h_0 \leq 1$ and $0 < \tau_0 \leq 1$ sufficiently small and independent of ε such that for any $0 < \varepsilon \leq 1$, when $0 < h \leq h_0$ and $0 < \tau \leq \tau_0$, we have*

$$\|e_\phi^n\|_{H^2} + \|e_\psi^n\|_{H^2} + \varepsilon^2 \|\dot{e}_\phi^n\|_{H^2} \lesssim h^{m_0} + \frac{\tau^2}{\varepsilon^2}, \quad (4.5)$$

$$\|e_\phi^n\|_{H^2} + \|e_\psi^n\|_{H^2} + \varepsilon^2 \|\dot{e}_\phi^n\|_{H^2} \lesssim h^{m_0} + \varepsilon^2, \quad 0 \leq n \leq \frac{T}{\tau}, \quad (4.6)$$

$$\|\phi_I^n\|_{H^2} \leq C_\phi + 1, \quad \|\psi_I^n\|_{H^2} \leq C_\psi + 1, \quad \|\dot{\phi}_I^n\|_{H^2} \leq \frac{C_\phi + 1}{\varepsilon^2}. \quad (4.7)$$

Thus, by taking the minimum of the two error bounds (4.5) and (4.6) for $\varepsilon \in (0, 1]$, we obtain a uniform error bound with respect to $\varepsilon \in (0, 1]$ for $0 \leq n \leq \frac{T}{\tau}$

$$\|e_\phi^n\|_{H^2} + \|e_\psi^n\|_{H^2} + \varepsilon^2 \|\dot{e}_\phi^n\|_{H^2} \lesssim h^{m_0} + \min_{0 < \varepsilon \leq 1} \left\{ \frac{\tau^2}{\varepsilon^2}, \varepsilon^2 \right\} \lesssim h^{m_0} + \tau. \quad (4.8)$$

In order to prove the above theorem, we introduce

$$\begin{aligned} e_{\phi,N}^n(x) &:= (P_N \phi)(x, t_n) - \phi_I^n(x), \quad \dot{e}_{\phi,N}^n(x) := P_N(\partial_t \phi)(x, t_n) - \dot{\phi}_I^n(x), \\ e_{\psi,N}^n(x) &:= (P_N \psi)(x, t_n) - \psi_I^n(x), \quad x \in \overline{\Omega}, \quad 0 \leq n \leq \frac{T}{\tau}. \end{aligned} \quad (4.9)$$

Using the triangle inequality and noticing the assumption (4.1), we have

$$\begin{aligned} \|e_\phi^n\|_{H^2} &\leq \|\phi(x, t_n) - (P_N \phi)(x, t_n)\|_{H^2} + \|e_{\phi,N}^n\|_{H^2} \lesssim h^{m_0+2} + \|e_{\phi,N}^n\|_{H^2}, \\ \|e_\psi^n\|_{H^2} &\leq \|\psi(x, t_n) - (P_N \psi)(x, t_n)\|_{H^2} + \|e_{\psi,N}^n\|_{H^2} \lesssim h^{m_0} + \|e_{\psi,N}^n\|_{H^2}, \\ \|\dot{e}_\phi^n\|_{H^2} &\leq \|\dot{\phi}(x, t_n) - (P_N \dot{\phi})(x, t_n)\|_{H^2} + \|\dot{e}_{\phi,N}^n\|_{H^2} \lesssim \frac{h^{m_0+2}}{\varepsilon^2} + \|\dot{e}_{\phi,N}^n\|_{H^2}. \end{aligned} \quad (4.10)$$

Thus we need only obtain estimates for $\|e_{\phi,N}^n\|_{H^2}$, $\|e_{\psi,N}^n\|_{H^2}$ and $\|\dot{e}_{\phi,N}^n\|_{H^2}$, which will be done by introducing the error energy functional

$$\mathcal{E} \left(e_{\phi,N}^n, e_{\psi,N}^n, \dot{e}_{\phi,N}^n \right) := \varepsilon^2 \left\| \dot{e}_{\phi,N}^n \right\|_{H^2}^2 + \left\| \partial_x e_{\phi,N}^n \right\|_{H^2}^2 + \frac{1}{\varepsilon^2} \left(\left\| e_{\phi,N}^n \right\|_{H^2}^2 + \left\| e_{\psi,N}^n \right\|_{H^2}^2 \right), \quad (4.11)$$

and establishing the following several lemmas.

For any $\mathbf{v} \in Y_N$, denote $v_{-1} = v_{N-1}$ and $v_{N+1} = v_1$, and define the difference operators $\delta_x^+ \mathbf{v} \in Y_N$ and $\delta_x^2 \mathbf{v} \in Y_N$ as

$$\delta_x^+ v_j = \frac{v_{j+1} - v_j}{h}, \quad \delta_x^2 v_j = \frac{v_{j+1} - 2v_j + v_{j-1}}{h^2}, \quad j = 0, 1, \dots, N,$$

and the norms as

$$\|\mathbf{v}\|_{Y,1}^2 = \|\mathbf{v}\|_{l^2}^2 + \|\delta_x^+ \mathbf{v}\|_{l^2}^2, \quad \|\mathbf{v}\|_{Y,2}^2 = \|\mathbf{v}\|_{l^2}^2 + \|\delta_x^+ \mathbf{v}\|_{l^2}^2 + \|\delta_x^2 \mathbf{v}\|_{l^2}^2, \quad (4.12)$$

with

$$\|\mathbf{v}\|_{l^2}^2 = h \sum_{j=0}^{N-1} |v_j|^2, \quad \|\delta_x^+ \mathbf{v}\|_{l^2}^2 = h \sum_{j=0}^{N-1} |\delta_x^+ v_j|^2, \quad \|\delta_x^2 \mathbf{v}\|_{l^2}^2 = h \sum_{j=0}^{N-1} |\delta_x^2 v_j|^2,$$

then we can have the following estimates [7, 8]

$$\|I_N \mathbf{v}\|_{H^1} \lesssim \|\mathbf{v}\|_{Y,1} \lesssim \|I_N \mathbf{v}\|_{H^1}, \quad \|I_N \mathbf{v}\|_{H^2} \lesssim \|\mathbf{v}\|_{Y,2} \lesssim \|I_N \mathbf{v}\|_{H^2}, \quad \forall \mathbf{v} \in Y_N. \quad (4.13)$$

Lemma 4.1 (A new formulation of MTI-FP) For the numerical solution ϕ_l^{n+1} and $\dot{\phi}_l^{n+1}$ ($n \geq 0$) obtained from the MTI-FP method (3.15)-(3.17), we have

$$\phi_l^{n+1}(x) = \sum_{l=-N/2}^{N/2-1} (\widetilde{\Phi^{n+1}})_l e^{i\mu_l(x-a)}, \quad \dot{\phi}_l^{n+1}(x) = \sum_{l=-N/2}^{N/2-1} (\dot{\widetilde{\Phi^{n+1}}})_l e^{i\mu_l(x-a)}, \quad (4.14)$$

where

$$\begin{aligned} (\widetilde{\Phi^{n+1}})_l &= \cos(\omega_l \tau) (\widetilde{\Phi^n})_l + \frac{\sin(\omega_l \tau)}{\omega_l} (\dot{\widetilde{\Phi^n}})_l + p_l(\tau) (|\widetilde{\Psi^n}|^2)_l + 2q_l(\tau) (\operatorname{Re}\{\widetilde{\Psi^n} \dot{\widetilde{\Psi^n}}\})_l, \\ (\dot{\widetilde{\Phi^{n+1}}})_l &= \cos(\omega_l \tau) (\dot{\widetilde{\Phi^n}})_l - \omega_l \sin(\omega_l \tau) (\widetilde{\Phi^n})_l + p'_l(\tau) (|\widetilde{\Psi^n}|^2)_l + 2q'_l(\tau) (\operatorname{Re}\{\widetilde{\Psi^n} \dot{\widetilde{\Psi^n}}\})_l. \end{aligned}$$

Proof For $l = -\frac{N}{2}, \dots, \frac{N}{2} - 1$, from (3.15), we get

$$\begin{cases} (\widetilde{\Phi^{n+1}})_l = e^{\frac{i\tau}{\varepsilon^2}} (\widetilde{Z^{n+1}})_l + e^{-\frac{i\tau}{\varepsilon^2}} (\widetilde{Z^{n+1}})_l + (\widetilde{R^{n+1}})_l, \\ (\dot{\widetilde{\Phi^{n+1}}})_l = e^{\frac{i\tau}{\varepsilon^2}} [(\dot{\widetilde{Z^{n+1}}})_l + \frac{i}{\varepsilon^2} (\widetilde{Z^{n+1}})_l] + e^{-\frac{i\tau}{\varepsilon^2}} [(\dot{\widetilde{Z^{n+1}}})_l - \frac{i}{\varepsilon^2} (\widetilde{Z^{n+1}})_l] + (\dot{\widetilde{R^{n+1}}})_l, \end{cases} \quad (4.15)$$

and from (3.17), we obtain

$$(\widetilde{Z^0})_l = \frac{1}{2} [(\widetilde{\Phi^n})_l - i\varepsilon^2 (\dot{\widetilde{\Phi^n}})_l], \quad (\dot{\widetilde{R^0}})_l = -(\dot{\widetilde{Z^0}})_l - (\widetilde{Z^0})_l. \quad (4.16)$$

Noticing (3.10) and (3.4), for $\mathbf{v} \in Y_N$, we have

$$a_l(\tau) = a_{-l}(\tau), \quad b_l(\tau) = b_{-l}(\tau), \quad l = -\frac{N}{2}, \dots, \frac{N}{2} - 1, \quad (\widetilde{\mathbf{v}})_l = \begin{cases} \overline{\widetilde{\mathbf{v}}_{-l}}, & |l| \leq \frac{N}{2} - 1, \\ \widetilde{\mathbf{v}}_l, & l = -\frac{N}{2}. \end{cases}$$

Plugging (4.16) into (3.16a)-(3.16c), and (3.16) into (4.15), we get

$$\begin{aligned} (\widetilde{\Phi^{n+1}})_l &= \operatorname{Re} \left\{ e^{\frac{i\tau}{\varepsilon^2}} a_l(\tau) \right\} (\widetilde{\Phi^n})_l + \varepsilon^2 \operatorname{Im} \left\{ e^{\frac{i\tau}{\varepsilon^2}} a_l(\tau) \right\} (\dot{\widetilde{\Phi^n}})_l + \varepsilon^2 e^{\frac{i\tau}{\varepsilon^2}} b_l(\tau) (\dot{\widetilde{Z^0}})_l \\ &\quad + \varepsilon^2 e^{-\frac{i\tau}{\varepsilon^2}} \overline{b_l(\tau)} (\widetilde{Z^0})_l + \frac{\sin(\omega_l \tau)}{\omega_l} (\dot{\widetilde{R^0}})_l + p_l(\tau) (|\widetilde{\Psi^n}|^2)_l \\ &\quad + 2q_l(\tau) (\operatorname{Re}\{\widetilde{\Psi^n} \dot{\widetilde{\Psi^n}}\})_l, \\ (\dot{\widetilde{\Phi^{n+1}}})_l &= \varepsilon^2 \operatorname{Im} \left\{ e^{\frac{i\tau}{\varepsilon^2}} a'_l(\tau) + \frac{i}{\varepsilon^2} e^{\frac{i\tau}{\varepsilon^2}} a_l(\tau) \right\} (\widetilde{\Phi^n})_l + \varepsilon^2 e^{-\frac{i\tau}{\varepsilon^2}} \left[\overline{b'_l(\tau)} - \frac{i}{\varepsilon^2} \overline{b_l(\tau)} \right] (\widetilde{Z^0})_l \\ &\quad + \operatorname{Re} \left\{ e^{\frac{i\tau}{\varepsilon^2}} a'_l(\tau) + \frac{i}{\varepsilon^2} e^{\frac{i\tau}{\varepsilon^2}} a_l(\tau) \right\} (\dot{\widetilde{\Phi^n}})_l + \varepsilon^2 e^{\frac{i\tau}{\varepsilon^2}} \left[b'_l(\tau) + \frac{i}{\varepsilon^2} b_l(\tau) \right] (\dot{\widetilde{Z^0}})_l \\ &\quad + \cos(\omega_l \tau) (\dot{\widetilde{R^0}})_l + p'_l(\tau) (|\widetilde{\Psi^n}|^2)_l + 2q'_l(\tau) (\operatorname{Re}\{\widetilde{\Psi^n} \dot{\widetilde{\Psi^n}}\})_l, \end{aligned}$$

where $\operatorname{Im}\{z\}$ denotes the imaginary part of a complex number z . Then (4.14) can be obtained by combining the above equalities and (3.10). \square

For the solution $\phi^n(x, s) := \phi(x, t_n + s)$, $\psi^n(x, s) := \psi(x, t_n + s)$ of (3.1), we have

$$\varepsilon^2 \widehat{\phi}_l''(t_n + s) + \mu_l^2 \widehat{\phi}_l(t_n + s) + \frac{1}{\varepsilon^2} \widehat{\phi}_l(t_n + s) = \widehat{(|\psi|^2)}_l(t_n + s), \quad s > 0 \quad (4.17a)$$

$$i \widehat{\psi}_l'(t_n + s) - \mu_l^2 \widehat{\phi}_l(t_n + s) + \widehat{(\phi \psi)}_l(t_n + s) = 0, \quad l \in \mathbb{Z}. \quad (4.17b)$$

By applying the variation-of-constant formula to (4.17) and noticing the ansatz (2.2) with $\rho^n(x, s) := |\psi(x, t_n + s)|^2$, we get

$$\widehat{\phi}_l(t_{n+1}) = \cos(\omega_l \tau) \widehat{\phi}_l(t_n) + \frac{\sin(\omega_l \tau)}{\omega_l} \widehat{\phi}_l'(t_n) + \int_0^\tau \frac{\sin(\omega_l(\tau - \theta))}{\varepsilon^2 \omega_l} \widehat{(\rho^n)}_l(\theta) d\theta, \quad (4.18a)$$

$$\widehat{\phi}_l'(t_{n+1}) = \cos(\omega_l \tau) \widehat{\phi}_l'(t_n) - \omega_l \sin(\omega_l \tau) \widehat{\phi}_l(t_n) + \int_0^\tau \frac{\cos(\omega_l(\tau - \theta))}{\varepsilon^2} \widehat{(\rho^n)}_l(\theta) d\theta, \quad (4.18b)$$

$$\begin{aligned} \widehat{\psi}_l(t_{n+1}) &= e^{-i\mu_l^2 \tau} \widehat{\psi}_l(t_n) + i e^{-i\mu_l^2 \tau} \int_0^\tau e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} \widehat{(z^n \psi^n)}_l(\theta) d\theta \\ &\quad + i e^{-i\mu_l^2 \tau} \int_0^\tau e^{i(\mu_l^2 - \frac{1}{\varepsilon^2})\theta} \widehat{(\overline{z^n \psi^n})}_l(\theta) d\theta + i \int_0^\tau e^{i\mu_l^2(\theta - \tau)} \widehat{(r^n \psi^n)}_l(\theta) d\theta. \end{aligned} \quad (4.18c)$$

Combining the above results, we define local truncation error functions as

$$\begin{aligned} \xi_\phi^n(x) &= \sum_{l=-N/2}^{N/2-1} \widehat{(\xi_\phi^n)}_l e^{i\mu_l(x-a)}, & \xi_\psi^n(x) &= \sum_{l=-N/2}^{N/2-1} \widehat{(\xi_\psi^n)}_l e^{i\mu_l(x-a)}, \\ \xi_\phi^n(x) &= \sum_{l=-N/2}^{N/2-1} \widehat{(\xi_\phi^n)}_l e^{i\mu_l(x-a)}, & x \in \Omega, \end{aligned} \quad (4.19)$$

where

$$\begin{aligned} \widehat{(\xi_\phi^n)}_l &:= \widehat{\phi}_l(t_{n+1}) - \left[\cos(\omega_l \tau) \widehat{\phi}_l(t_n) + \frac{\sin(\omega_l \tau)}{\omega_l} \widehat{\phi}_l'(t_n) + p_l(\tau) \widehat{(|\psi^n|^2)}_l(0) \right. \\ &\quad \left. + 2q_l(\tau) \widehat{(\operatorname{Re}\{\psi^n \phi^n\})}_l(0) \right], \end{aligned} \quad (4.20a)$$

$$\begin{aligned} \widehat{(\xi_\psi^n)}_l &:= \widehat{\psi}_l(t_{n+1}) - \left[-\omega_l \sin(\omega_l \tau) \widehat{\phi}_l(t_n) + \cos(\omega_l \tau) \widehat{\phi}_l'(t_n) + p_l'(\tau) \widehat{(|\psi^n|^2)}_l(0) \right. \\ &\quad \left. + 2q_l'(\tau) \widehat{(\operatorname{Re}\{\psi^n \phi^n\})}_l(0) \right], \end{aligned} \quad (4.20b)$$

$$\begin{aligned} \widehat{(\xi_\psi^n)}_l &:= \widehat{\psi}_l(t_{n+1}) - e^{-i\mu_l^2 \tau} \widehat{\psi}_l(t_n) - c_l^+(\tau) \widehat{(z^n \psi^n)}_l(0) - c_l^-(\tau) \widehat{(\overline{z^n \psi^n})}_l(0) \\ &\quad - d_l^+(\tau) \left[\widehat{(z^n \psi^n)}_l(0) + \widehat{(z^n \phi^n)}_l(0) \right] - d_l^-(\tau) \left[\widehat{(\overline{z^n \psi^n})}_l(0) + \widehat{(\overline{z^n \phi^n})}_l(0) \right] \\ &\quad - \frac{i\tau}{2} \widehat{(\gamma^{n+1} \psi^n)}_l(0) - \frac{i\tau^2}{2} \widehat{(\gamma^{n+1} \phi^n)}_l. \end{aligned} \quad (4.20c)$$

Here we introduce three auxiliary functions as

$$\varphi^n(x) = \sum_{l \in \mathbb{Z}} \widehat{(\varphi^n)}_l e^{i\mu_l(x-a)}, \quad \mathfrak{z}^n(x) = \sum_{l \in \mathbb{Z}} \widehat{(\mathfrak{z}^n)}_l e^{i\mu_l(x-a)}, \quad \gamma^{n+1}(x) = \sum_{l \in \mathbb{Z}} \widehat{(\gamma^{n+1})}_l e^{i\mu_l(x-a)},$$

with

$$\begin{cases} (\widehat{\varphi}^n)_l = -\frac{i}{\tau} \sin(\mu_l^2 \tau) (\widehat{\psi}^n)_l(0) + i (\widehat{\phi}^n \widehat{\psi}^n)_l(0), & (\widehat{z}^n)_l = \frac{i}{2\tau} \sin(\mu_l^2 \tau) (\widehat{z}^n)_l(0), \\ (\widehat{\gamma}^{n+1})_l = p_l(\tau) (|\widehat{\psi}^n|^2)_l(0) + 2q_l(\tau) (\operatorname{Re}\{\widehat{\psi}^n \widehat{\varphi}^n\})_l(0) - \frac{\sin(\omega_l \tau)}{\omega_l} \left((\widehat{z}^n)_l + (\widehat{\beta}^n)_l \right). \end{cases} \quad (4.21)$$

Similarly to the case of the KG equation [8] with details omitted here for brevity, we can prove a priori estimates for the solution of the problem (3.2) with (3.3).

Lemma 4.2 (A priori estimate of MDF) *Let $z^n(x, s)$ and $r^n(x, s)$ be the solution of the MDF (3.2) with initial boundary conditions (3.3). Under the assumption (4.1), there exists a constant $\tau_1 > 0$ independent of ε , h and τ (or n), such that for $0 < \tau \leq \tau_1$,*

$$\|z^n\|_{L^\infty([0, \tau]; H^{m_0+2})} + \|\partial_s z^n\|_{L^\infty([0, \tau]; H^{m_0+1})} + \|\partial_s^2 z_\pm^n\|_{L^\infty([0, \tau]; H^{m_0})} \lesssim 1, \quad (4.22)$$

$$\|\partial_s^k r^n\|_{L^\infty([0, \tau]; H^{6-k})} \lesssim \varepsilon^{2-2k}, \quad k = 0, 1, 2. \quad (4.23)$$

Based on this result, when $0 < \tau \leq \tau_1$, it is easy to see that $\varphi^n, \beta^n \in H_p^{m_0}(\Omega)$, $\gamma^{n+1} \in H_p^4(\Omega)$ and we have the prior estimates

$$\|\varphi^n\|_{H^4} \lesssim 1, \quad \|\beta^n\|_{H^4} \lesssim 1, \quad \|\gamma^{n+1}\|_{H^4} \lesssim \varepsilon^2.$$

With the above prior estimates, we can estimate the local truncation error functions.

Lemma 4.3 (Estimates on local errors) *Under the assumption (4.1), for any $0 < \tau \leq \tau_1$ with τ_1 given in Lemma 4.2, we have two independent bounds as*

$$\mathcal{E} \left(\xi_\phi^n, \xi_\psi^n, \xi_\beta^n \right) \lesssim \frac{\tau^6}{\varepsilon^6}, \quad \mathcal{E} \left(\xi_\phi^n, \xi_\psi^n, \xi_\beta^n \right) \lesssim \tau^2 \varepsilon^2, \quad 0 \leq n \leq \frac{T}{\tau} - 1. \quad (4.24)$$

Proof Subtracting (4.20) from (4.18) and by using the Taylor's expansion, from (4.18a) and (4.20a), we get

$$(\widehat{\xi}_\phi^n)_l = \int_0^\tau \frac{\sin(\omega_l(\tau - \theta))}{\varepsilon^2 \omega_l} \theta^2 \left[\int_0^1 (1-s) (|\widehat{\psi}^n|^2)_l''(\theta s) ds \right] d\theta + 2q_l(\tau) (\widehat{B}_0)_l, \quad (4.25)$$

where

$$B_0(x) = \operatorname{Re}\{\overline{\widehat{\psi}^n(x, 0)} (\partial_s \widehat{\psi}^n(x, 0) - \varphi^n)\}, \quad x \in \Omega.$$

Noticing (A.1), we obtain

$$|(\widehat{\xi}_\phi^n)_l| \lesssim \frac{\tau^2}{\varepsilon^2 \omega_l} \int_0^\tau \int_0^1 \left| (|\widehat{\psi}^n|^2)_l''(\theta s) \right| ds d\theta + \frac{\tau^2}{\varepsilon^2 \omega_l} |(\widehat{B}_0)_l|.$$

Noting $\frac{1}{\varepsilon^2 \omega_l} = \frac{1}{\sqrt{1+\varepsilon^2 \mu_l^2}} \leq 1$ for $l = -\frac{N}{2}, \dots, \frac{N}{2} - 1$ and (4.1), (4.2), we get

$$\begin{aligned} \|\xi_\phi^n\|_{H^2}^2 &\lesssim \tau^6 \|\partial_{ss} \rho\|_{L^\infty([0, T]; H^2)}^2 + \tau^4 \|\psi^n(\cdot, 0) (\partial_s \psi^n(\cdot, 0) - \varphi^n)\|_{H^2}^2 \\ &\lesssim \tau^6 + \tau^4 \|\partial_s \psi^n(\cdot, 0) - \varphi^n\|_{H^2}^2, \quad 0 < \tau \leq \tau_1. \end{aligned} \quad (4.26)$$

By Parseval's identity, we have

$$\begin{aligned}
\|\partial_s \psi^n(\cdot, 0) - \varphi^n\|_{H^2}^2 &= \sum_{l=-N/2}^{N/2-1} (1 + \mu_l^2 + \mu_l^4) \left| (\widehat{\partial_s \psi^n})_l(0) - (\widehat{\varphi^n})_l \right|^2 \\
&\lesssim \sum_{l=-N/2}^{N/2-1} \mu_l^4 \left| \frac{\sin(\mu_l^2 \tau)}{\tau} - \mu_l^2 \right|^2 |(\widehat{\psi^n})_l(0)|^2 \lesssim \tau^2 \sum_{l=-N/2}^{N/2-1} \mu_l^{12} |(\widehat{\psi^n})_l(0)|^2 \\
&\lesssim \tau^2 \|\psi\|_{L^\infty([0, T]; H^6)}^2 \lesssim \tau^2.
\end{aligned} \tag{4.27}$$

This together with (4.26), we get

$$\|\xi_\phi^n\|_{H^2}^2 \lesssim \tau^6, \quad 0 < \tau \leq \tau_1. \tag{4.28}$$

Similarly, noting $\frac{|\mu_l|}{\varepsilon^2 \omega_l} = \frac{|\mu_l|}{\sqrt{1 + \varepsilon^2 \mu_l^2}} \leq \frac{1}{\varepsilon}$ for $l = -\frac{N}{2}, \dots, \frac{N}{2} - 1$, we obtain

$$\|\partial_x \xi_\phi^n\|_{H^2}^2 \lesssim \frac{\tau^6}{\varepsilon^2} \quad \text{and} \quad \|\xi_\phi^n\|_{H^2}^2 \lesssim \frac{\tau^6}{\varepsilon^4}, \quad 0 < \tau \leq \tau_1. \tag{4.29}$$

For ξ_ψ^n , from (4.18c) and (4.20c) and by the Taylor's expansion, we have

$$\begin{aligned}
(\widehat{\xi_\psi^n})_l &= i e^{-i\mu_l^2 \tau} \int_0^\tau e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} \theta^2 \left[\int_0^1 (1-s) (\widehat{z^n \psi^n})_l''(\theta s) ds \right] d\theta \\
&\quad + i e^{-i\mu_l^2 \tau} \int_0^\tau e^{i(\mu_l^2 - \frac{1}{\varepsilon^2})\theta} \theta^2 \left[\int_0^1 (1-s) (\widehat{\bar{z}^n \psi^n})_l''(\theta s) ds \right] d\theta \\
&\quad + d_l^+(\tau) \left[(\widehat{B_1})_l + (\widehat{B_2})_l \right] + d_l^-(\tau) \left[(\widehat{B_3})_l + (\widehat{B_4})_l \right] + \frac{i\tau^3}{4} (\widehat{B_5})_l + \frac{i\tau}{2} (\widehat{B_6})_l \\
&\quad + \frac{i\tau^2}{2} \left[(\widehat{B_7})_l + (\widehat{B_8})_l \right] - \frac{i\tau}{2} (\widehat{r^n \psi^n})_l(\tau) + i \int_0^\tau e^{i\mu_l^2(\theta-\tau)} (\widehat{r^n \psi^n})_l(\theta) d\theta,
\end{aligned} \tag{4.30}$$

where

$$\begin{aligned}
B_1(x) &= [\partial_s z^n(x, 0) - \mathfrak{z}^n(x)] \psi^n(x, 0), & B_2(x) &= z^n(x, 0) [\partial_s \psi^n(x, 0) - \varphi^n(x)], \\
B_3(x) &= [\partial_s \bar{z}^n(x, 0) - \bar{\mathfrak{z}}^n(x)] \psi^n(x, 0), & B_4(x) &= \bar{z}^n(x, 0) [\partial_s \psi^n(x, 0) - \varphi^n(x)], \\
B_5(x) &= r^n(x, \tau) \partial_{ss} \psi^n(x, \theta_0), & B_6(x) &= [r^n(x, \tau) - \gamma^{n+1}(x)] \psi^n(x, 0), \\
B_7(x) &= r^n(x, \tau) [\partial_s \psi^n(x, 0) - \varphi^n(x)], & B_8(x) &= [r^n(x, \tau) - \gamma^{n+1}(x)] \varphi^n(x),
\end{aligned}$$

with $\theta_0 \in [0, \tau]$. For the last two terms in (4.30), we have

$$\begin{aligned}
&\left| i \int_0^\tau e^{i\mu_l^2(\theta-\tau)} (\widehat{r^n \psi^n})_l(\theta) d\theta - \frac{i\tau}{2} (\widehat{r^n \psi^n})_l(\tau) \right| \\
&\lesssim \int_0^\tau \theta(\tau - \theta) \left| \frac{d^2}{d\theta^2} \left[e^{i\mu_l^2(\theta-\tau)} (\widehat{r^n \psi^n})_l(\theta) \right] \right| d\theta.
\end{aligned}$$

Then by using the triangle inequality, we obtain

$$\begin{aligned} |(\widehat{\xi_\psi^n})_l| &\lesssim \tau^2 \int_0^\tau \int_0^1 \left[\left| (\widehat{z^n \psi^n})''_l(\theta s) \right| + \left| (\widehat{r^n \psi^n})''_l(\theta s) \right| \right] ds d\theta \\ &\quad + \tau^2 \left[\left| (\widehat{B_1})_l \right| + \left| (\widehat{B_2})_l \right| \right] + \tau^2 \left[\left| (\widehat{B_3})_l \right| + \left| (\widehat{B_4})_l \right| \right] + \tau^3 \left| (\widehat{B_5})_l \right| + \tau \left| (\widehat{B_6})_l \right| \\ &\quad + \tau^2 \left| (\widehat{B_7})_l \right| + \tau^2 \left| (\widehat{B_8})_l \right| + \tau^2 \int_0^\tau \left| \frac{d^2}{d\theta^2} \left[e^{i\mu_l^2(\theta-\tau)} (\widehat{r^n \psi^n})_l(\theta) \right] \right| d\theta. \end{aligned}$$

Similarly as (4.27), we find from (4.21)

$$\|\partial_s z^n - \mathfrak{z}^n\|_{H^2} \lesssim \tau \|z^n(\cdot, 0)\|_{H^6} \lesssim \tau \left(\|\phi\|_{L^\infty([0, T]; H^6)} + \varepsilon^2 \|\partial_t \phi\|_{L^\infty([0, T]; H^6)} \right) \lesssim \tau.$$

Again, similarly as (4.27) and noticing (3.9b), we have

$$\begin{aligned} \|\gamma^{n+1} - r^n(\cdot, \tau)\|_{H^2}^2 &\lesssim \|\xi_\phi^n\|_{H^2}^2 + \sum_{l \in \mathbb{Z}} (1 + \mu_l^2 + \mu_l^4) \left| \frac{\sin(\omega_l \tau)}{\omega_l} \right|^2 \cdot \left| (\widehat{\mathfrak{z}^n})_l - (\widehat{\partial_s z^n})_l \right|^2 \\ &\lesssim \tau^6 + \tau^2 \|\partial_s z^n - \mathfrak{z}^n\|_{H^2}^2 \lesssim \tau^4. \end{aligned}$$

By the assumption (4.1) and Lemma 4.2, we get

$$\begin{aligned} \|\xi_\psi^n\|_{H^2}^2 &\lesssim \tau^6 \|\partial_{ss}(z^n \psi^n)\|_{L^\infty([0, \tau]; H^2)}^2 + \tau^6 \|z^n\|_{L^\infty([0, \tau]; H^6)}^2 + \tau^6 \|\psi^n\|_{L^\infty([0, \tau]; H^6)}^2 \\ &\quad + \tau^6 \|r^n \partial_{ss} \psi^n\|_{L^\infty([0, \tau]; H^2)}^2 + \tau^2 \|\gamma^{n+1} - r^n(\cdot, \tau)\|_{H^2}^2 + \tau^6 \|r^n \psi^n\|_{L^\infty([0, \tau]; H^6)}^2 \\ &\quad + \tau^6 \|\partial_s(r^n \psi^n)\|_{L^\infty([0, \tau]; H^4)}^2 + \tau^6 \|\partial_{ss}(r^n \psi^n)\|_{L^\infty([0, \tau]; H^2)}^2 \lesssim \frac{\tau^6}{\varepsilon^4}. \end{aligned} \quad (4.31)$$

Plugging (4.28), (4.29) and (4.31) into (4.11) with $e_{\phi, N}^n = \xi_\phi^n$, $e_{\psi, N}^n = \xi_\psi^n$ and $e_{\phi, N}^n = \dot{\xi}_\phi^n$, we immediately get the first inequality in (4.24).

Next we show how to get the second estimate in (4.24). By using Taylor's expansion, truncating at the first order term and integrating by parts, we have

$$\begin{aligned} (\widehat{\xi_\phi^n})_l &= \int_0^\tau \frac{\sin(\omega_l(\tau - \theta))}{\varepsilon^2 \omega_l} \theta \left[\int_0^1 (\widehat{|\psi^n|^2})'_l(\theta s) ds \right] d\theta - 2q_l(\tau) (\text{Re}\{\widehat{\psi^n \phi^n}\})_l(0), \\ &= q_l(\tau) \int_0^1 (\widehat{|\psi^n|^2})'_l(\tau s) ds - \int_0^\tau q_l(\theta) \left[\int_0^1 (\widehat{|\psi^n|^2})''_l(\theta s) ds \right] d\theta \\ &\quad - q_l(0) (\widehat{|\psi^n|^2})'_l(0) - 2q_l(\tau) (\text{Re}\{\widehat{\psi^n \phi^n}\})_l(0). \end{aligned}$$

From (A.1) and noticing $q_l(0) = 0$ and $|q_l(\tau)| \lesssim \tau \varepsilon^2$, we find

$$\begin{aligned} \|\xi_\phi^n\|_{H^2}^2 &\lesssim \tau^2 \varepsilon^4 \left[\|\partial_t \rho\|_{L^\infty([0, T]; H^2)}^2 + \|\partial_{tt} \rho\|_{L^\infty([0, T]; H^2)}^2 + \|\psi \phi^n\|_{L^\infty([0, T]; H^2)}^2 \right] \\ &\lesssim \tau^2 \varepsilon^4, \quad 0 < \tau \leq \tau_1, \quad 0 \leq \varepsilon \leq 1. \end{aligned} \quad (4.32)$$

Similarly, we can get

$$\|\partial_x \xi_\phi^n\|_{H^2}^2 \lesssim \tau^2 \varepsilon^2 \quad \text{and} \quad \|\dot{\xi}_\phi^n\|_{H^2}^2 \lesssim \tau^2, \quad 0 < \tau \leq \tau_1. \quad (4.33)$$

In order to estimate ξ_{ψ}^n , we introduce

$$I^n(x) = \sum_{l=-N/2}^{N/2-1} \widehat{I}_l^n e^{i\mu_l(x-a)}, \quad J^n(x) = \sum_{l=-N/2}^{N/2-1} \widehat{J}_l^n e^{i\mu_l(x-a)}, \quad (4.34)$$

where

$$\begin{aligned} \widehat{I}_l^n &= G_{1,l}^n - d_l^+(\tau) \left[\widehat{(z^n \psi^n)}_l(0) + \widehat{(z^n \phi^n)}_l(0) \right], \\ \widehat{J}_l^n &= G_{2,l}^n - d_l^-(\tau) \left[\widehat{(z^n \psi^n)}_l(0) + \widehat{(z^n \phi^n)}_l(0) \right], \end{aligned} \quad (4.35)$$

with

$$\begin{aligned} G_{1,l}^n &= ie^{-i\mu_l^2 \tau} \int_0^\tau e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} \theta \left[\int_0^1 \widehat{(z^n \psi^n)}_l'(\theta s) ds \right] d\theta, \\ G_{2,l}^n &= ie^{-i\mu_l^2 \tau} \int_0^\tau e^{i(\mu_l^2 - \frac{1}{\varepsilon^2})\theta} \theta \left[\int_0^1 \widehat{(z^n \psi^n)}_l'(\theta s) ds \right] d\theta. \end{aligned} \quad (4.36)$$

Applying the Taylor's expansion in (4.18c) and (4.20c) and truncating at the first order term, we obtain

$$\widehat{(\xi_{\psi}^n)}_l = \widehat{I}_l^n + \widehat{J}_l^n + i \int_0^\tau e^{i\mu_l^2(\theta-\tau)} \widehat{(r^n \psi^n)}_l(\theta) d\theta - \frac{i\tau}{2} \left[\widehat{(\gamma^{n+1} \psi^n)}_l(0) + \tau \widehat{(\gamma^{n+1} \phi^n)}_l \right]. \quad (4.37)$$

From (4.36), integration by parts, we get

$$\begin{aligned} G_{1,l}^n &= ie^{-i\mu_l^2 \tau} \int_0^\tau e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} \theta \int_0^1 \left[\widehat{(\partial_s z^n \psi^n)}_l(\theta s) + \widehat{(z^n \partial_s \psi^n)}_l(\theta s) \right] ds d\theta \\ &= d_l^+(\tau) \int_0^1 \widehat{(\partial_s z^n \psi^n)}_l(\tau s) ds - \int_0^\tau d_l^+(\theta) \int_0^1 \widehat{(\partial_s z^n \psi^n)}_l'(\theta s) ds d\theta \\ &\quad - e^{-i\mu_l^2 \tau} \int_0^\tau e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} \theta \int_0^1 \left[\widehat{(z^n \partial_{xx} \psi^n)}_l(\theta s) + \widehat{(z^n \phi^n \psi^n)}_l(\theta s) \right] ds d\theta \\ &= d_l^+(\tau) \int_0^1 \left[\widehat{(\partial_s z^n \psi^n)}_l(\tau s) + i \widehat{(z^n \partial_{xx} \psi^n)}_l(\tau s) \right] ds \\ &\quad - \int_0^\tau d_l^+(\theta) \int_0^1 \left[\widehat{(\partial_s z^n \psi^n)}_l'(\theta s) + i \widehat{(z^n \partial_{xx} \psi^n)}_l'(\theta s) \right] ds d\theta - G_{3,l}^n, \end{aligned} \quad (4.38)$$

where

$$G_{3,l}^n = e^{-i\mu_l^2 \tau} \int_0^\tau e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} \theta \int_0^1 \widehat{(z^n \phi^n \psi^n)}_l(\theta s) ds d\theta. \quad (4.39)$$

Noticing the ansatz (2.2), we get

$$\begin{aligned} G_{3,l}^n &= e^{-i\mu_l^2 \tau} \int_0^1 \left[\int_0^\tau e^{i(\mu_l^2 + \frac{1+s}{\varepsilon^2})\theta} \theta \widehat{((z^n)^2 \psi^n)}_l(\theta s) d\theta \right. \\ &\quad \left. + \int_0^\tau e^{i(\mu_l^2 + \frac{1-s}{\varepsilon^2})\theta} \theta \widehat{(|z^n|^2 \psi^n)}_l(\theta s) d\theta + \int_0^\tau e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} \theta \widehat{(z^n r^n \psi^n)}_l(\theta s) d\theta \right] ds. \end{aligned}$$

Using integration by parts and the triangle inequality, we have

$$\begin{aligned} |G_{3,l}^n| &\lesssim \int_0^1 \left[\tau \varepsilon^2 \left| (\widehat{(z^n)^2 \psi^n})_l(\tau s) \right| + \tau \varepsilon^2 \left| (\widehat{|z^n|^2 \psi^n})_l(\tau s) \right| + \varepsilon^2 \int_0^\tau \left| (\widehat{(z^n)^2 \psi^n})'_l(\theta s) \right| d\theta \right. \\ &\quad \left. + \varepsilon^2 \int_0^\tau \left| (\widehat{|z^n|^2 \psi^n})'_l(\theta s) \right| d\theta + \int_0^\tau \left| (\widehat{z^n r^n \psi^n})_l(\theta s) \right| d\theta \right] ds. \end{aligned} \quad (4.40)$$

Combining (4.34)-(4.40) and noting $|d_l^+(\tau)| \lesssim \tau \varepsilon^2$ and Lemma 4.2, we get

$$\begin{aligned} \|I^n\|_{H^2} &\lesssim \tau \varepsilon^2 \left[\|\partial_s z^n \psi^n\|_{L^\infty([0,\tau];H^2)} + \|z^n \partial_{xx} \psi^n\|_{L^\infty([0,\tau];H^2)} + \|\partial_{ss} z^n \psi^n\|_{L^\infty([0,\tau];H^2)} \right. \\ &\quad \left. + \|\partial_s z^n \partial_s \psi^n\|_{L^\infty([0,\tau];H^2)} + \|\partial_s z^n \partial_{xx} \psi^n\|_{L^\infty([0,\tau];H^2)} + \|z^n \partial_{sxx} \psi^n\|_{L^\infty([0,\tau];H^2)} \right. \\ &\quad \left. + \|(z^n)^2 \psi^n\|_{L^\infty([0,\tau];H^2)} \right] + \tau \varepsilon^2 \left[\|\mathfrak{z}^n \psi^n\|_{L^\infty([0,\tau];H^2)} + \|z^n \phi^n\|_{L^\infty([0,\tau];H^2)} \right] \\ &\quad + \tau \|z^n r^n \psi^n\|_{L^\infty([0,\tau];H^2)} \lesssim \tau \varepsilon^2. \end{aligned} \quad (4.41)$$

In order to estimate J^n , denote $N_\varepsilon := \frac{b-a}{2\sqrt{2}\pi\varepsilon}$, when $|l| \leq N_\varepsilon$, we have $|\mu_l| \leq \frac{1}{\sqrt{2}\varepsilon}$ which indicates that $|d_l^-(\tau)| \lesssim \tau \varepsilon^2$. Similarly to the estimate of I^n , we have

$$\sum_{|l| \leq N_\varepsilon} (1 + \mu_l^2 + \mu_l^4) |\widehat{J}_l^n|^2 \lesssim \tau^2 \varepsilon^4. \quad (4.42)$$

On the other hand, when $|\mu_l| > \frac{1}{\sqrt{2}\varepsilon}$, we have $|d_l^-(\tau)| \lesssim \tau^2$. Due to the regularity of the solution, i.e. the assumption (4.1) and Lemma 4.2, we can get

$$\begin{aligned} &\sum_{|l| > N_\varepsilon} (1 + \mu_l^2 + \mu_l^4) |\widehat{J}_l^n|^2 \\ &\lesssim \sum_{|l| > N_\varepsilon} \tau^3 \mu_l^4 \left[\int_0^\tau \int_0^1 \left| (\widehat{(z^n \psi^n)})'_l(\theta s) \right|^2 ds d\theta + \tau \left(\left| (\widehat{(z^n \psi^n)})_l(0) \right|^2 + \left| (\widehat{(z^n \phi^n)})_l(0) \right|^2 \right) \right] \\ &\lesssim \tau^4 \varepsilon^4 \left[\|\partial_s(z^n \psi^n)\|_{L^\infty([0,\tau];H^4)}^2 + \|\mathfrak{z}^n \psi^n(\cdot, 0)\|_{H^4}^2 + \|z^n(\cdot, 0) \phi^n\|_{H^4}^2 \right] \\ &\lesssim \tau^4 \varepsilon^4 \left[\|\partial_s(z^n \psi^n)\|_{L^\infty([0,\tau];H^4)}^2 + \frac{1}{\tau^2} \|z^n \psi^n\|_{L^\infty([0,\tau];H^4)}^2 + \|z^n \psi^n \phi^n\|_{L^\infty([0,\tau];H^4)}^2 \right] \\ &\lesssim \tau^2 \varepsilon^4. \end{aligned} \quad (4.43)$$

From (4.42) and (4.43), we obtain

$$\|J^n\|_{H^2} \lesssim \tau \varepsilon^2, \quad 0 < \tau \leq \tau_1, \quad 0 < \varepsilon \leq 1. \quad (4.44)$$

Combining the above estimates and noting (4.37), we have

$$\begin{aligned} \|\xi_\psi^n\|_{H^2}^2 &\lesssim \|I^n\|_{H^2}^2 + \|J^n\|_{H^2}^2 + \tau^2 \|r^n \psi^n\|_{L^\infty([0,\tau];H^2)}^2 + \tau^4 \|\gamma^{n+1} \psi^n\|_{L^\infty([0,\tau];H^2)}^2 \\ &\quad + \tau^4 \|\gamma^{n+1} \phi^n\|_{H^2}^2 \lesssim \tau^2 \varepsilon^4. \end{aligned} \quad (4.45)$$

Plugging (4.32), (4.33) and (4.45) into (4.11), we get the second estimate. \square

Defining the errors from the nonlinear terms as

$$\begin{cases} \eta_\phi^n(x) := \sum_{l=-N/2}^{N/2-1} (\widehat{\eta}_\phi^n)_l e^{i\mu_l(x-a)}, & \dot{\eta}_\phi^n(x) := \sum_{l=-N/2}^{N/2-1} (\dot{\widehat{\eta}}_\phi^n)_l e^{i\mu_l(x-a)}, \\ \eta_\psi^n(x) := \sum_{l=-N/2}^{N/2-1} (\widehat{\eta}_\psi^n)_l e^{i\mu_l(x-a)}, & x \in \overline{\Omega}, \quad n \geq 0, \end{cases} \quad (4.46)$$

where

$$(\widehat{\eta}_\phi^n)_l = p_l(\tau) \left[(|\widehat{\psi^n}|^2)_l(0) - (\widetilde{|\Psi^n|^2})_l \right] + 2q_l(\tau) \left[(\operatorname{Re}\{\widehat{\psi^n}\phi^n\})_l(0) - (\operatorname{Re}\{\widetilde{\Psi^n}\dot{\Psi}^n\})_l \right], \quad (4.47a)$$

$$(\dot{\widehat{\eta}}_\phi^n)_l = p'_l(\tau) \left[(|\widehat{\psi^n}|^2)_l(0) - (\widetilde{|\Psi^n|^2})_l \right] + 2q'_l(\tau) \left[(\operatorname{Re}\{\widehat{\psi^n}\phi^n\})_l(0) - (\operatorname{Re}\{\widetilde{\Psi^n}\dot{\Psi}^n\})_l \right], \quad (4.47b)$$

$$\begin{aligned} (\widehat{\eta}_\psi^n)_l &= c_l^+(\tau) \left[(\widehat{z^n\psi^n})_l(0) - (\widetilde{Z^0\Psi^n})_l \right] + c_l^-(\tau) \left[(\widetilde{z^n\psi^n})_l(0) - (\widetilde{Z^0\Psi^n})_l \right] \\ &\quad + d_l^+(\tau) \left[(\widehat{z^n\psi^n})_l(0) - (\widetilde{Z^0\Psi^n})_l + (\widehat{z^n\phi^n})_l(0) - (\widetilde{Z^0\dot{\Psi}^n})_l \right] \\ &\quad + d_l^-(\tau) \left[(\widetilde{z^n\psi^n})_l(0) - (\widetilde{Z^0\Psi^n})_l + (\widetilde{z^n\phi^n})_l(0) - (\widetilde{Z^0\dot{\Psi}^n})_l \right] \\ &\quad + \frac{i\tau}{2} \left[(\gamma^{n+1}\widehat{\psi^n})_l(0) - (\widetilde{R^{n+1}\Psi^n})_l + \tau \left((\gamma^{n+1}\widehat{\phi^n})_l - (\widetilde{R^{n+1}\dot{\Psi}^n})_l \right) \right], \end{aligned} \quad (4.47c)$$

then we have

Lemma 4.4 (Estimates on η_ϕ^n , η_ψ^n and $\dot{\eta}_\phi^n$) *Under the assumption (4.1) and assume (4.7) holds (which will be proved by induction later), we have for any $0 < \tau \leq \tau_1$,*

$$\mathcal{E} \left(\eta_\phi^n, \eta_\psi^n, \dot{\eta}_\phi^n \right) \lesssim \tau^2 \mathcal{E} \left(e_{N,\phi}^n, e_{N,\psi}^n, e_{N,\phi}^n \right) + \frac{\tau^2 h^{2m_0}}{\varepsilon^2}, \quad 0 \leq n \leq \frac{T}{\tau} - 1. \quad (4.48)$$

Proof From (4.46) and (4.47a), we have

$$\begin{aligned} \|\eta_\phi^n\|_{H^2} &\lesssim \tau \| |\psi^n(\cdot, 0)|^2 - I_N |\Psi^n|^2 \|_{H^2} + \tau^2 \| \overline{\psi^n}(\cdot, 0) \phi^n - I_N (\overline{\Psi^n} \dot{\Psi}^n) \|_{H^2} \\ &\lesssim \tau \| I_N |\psi^n(\cdot, 0)|^2 - I_N |\Psi^n|^2 \|_{H^2} + \tau^2 \| I_N (\overline{\psi^n}(\cdot, 0) \phi^n) - I_N (\overline{\Psi^n} \dot{\Psi}^n) \|_{H^2} + \tau h^{m_0} \\ &\lesssim \tau \| |\psi^n(\cdot, 0)|^2 - |\Psi^n|^2 \|_{Y,2} + \tau^2 \| \overline{\psi^n}(\cdot, 0) \phi^n - \overline{\Psi^n} \dot{\Psi}^n \|_{Y,2} + \tau h^{m_0} \\ &\lesssim \tau \| e_\psi^n \|_{Y,2} + \tau^2 \| \phi^n - \dot{\Psi}^n \|_{Y,2} + \tau h^{m_0} \\ &\lesssim \tau \| e_\psi^n \|_{Y,2} + \tau^2 \| \partial_{xx}^S \psi^n(\cdot, 0) - \partial_{xx}^S \Psi^n \|_{Y,2} + \tau^2 \| \phi^n \psi^n(\cdot, 0) - \Phi^n \Psi^n \|_{Y,2} + \tau h^{m_0} \\ &\lesssim \tau \| e_\psi^n \|_{H^2} + \tau \| e_\phi^n \|_{H^2} + \tau^2 \| \partial_{xx}^S \psi^n(\cdot, 0) - \partial_{xx}^S \Psi^n \|_{H^2} + \tau h^{m_0} \\ &\lesssim \tau \| e_\psi^n \|_{H^2} + \tau \| e_\phi^n \|_{H^2} + \tau h^{m_0}, \end{aligned} \quad (4.49)$$

where the operator ∂_{xx}^S on a function $f(x)$ is defined as

$$\partial_{xx}^S f(x) := - \sum_{l \in \mathbb{Z}} \frac{\sin(\mu_l^2 \tau)}{\tau} \widehat{f}_l e^{i\mu_l(x-a)}.$$

Combining (4.49) and (4.10), we get

$$\|\eta_\phi^n\|_{H^2} \lesssim \tau \|e_{\psi,N}^n\|_{H^2} + \tau \|e_{\phi,N}^n\|_{H^2} + \tau h^{m_0}. \quad (4.50)$$

Similarly, we can obtain

$$\varepsilon^2 \|\widehat{\eta}_\phi^n\|_{H^2}^2 + \|\partial_x \eta_\phi^n\|_{H^2}^2 \lesssim \frac{\tau^2}{\varepsilon^2} \left[\|e_{\psi,N}^n\|_{H^2}^2 + \|e_{\phi,N}^n\|_{H^2}^2 + h^{2m_0} \right]. \quad (4.51)$$

Similarly, we can estimate all the terms of η_ψ^n in (4.47c) except the last term. For the last term, we have

$$\begin{aligned} & \sum_{l=-N/2}^{N/2-1} (1 + \mu_l^2 + \mu_l^4) \left| (\widehat{\gamma^{n+1}\psi^n})_l(0) - (\widetilde{R^{n+1}\Psi^n})_l \right|^2 \\ & \lesssim \|I_N(\gamma^{n+1}\psi^n(\cdot, 0)) - I_N(R^{n+1}\psi^n)\|_{H^2}^2 + h^{2m_0} \\ & \lesssim \|e_\psi^n\|_{H^2}^2 + \|\gamma^{n+1} - I_N R^{n+1}\|_{H^2}^2 + h^{2m_0}. \end{aligned} \quad (4.52)$$

Noting (4.21) and (3.16b), we get

$$\begin{aligned} \left| (\widehat{\gamma^{n+1}})_l - (\widetilde{R^{n+1}})_l \right| & \lesssim \left| (\widehat{z^n})_l(0) - (\widetilde{Z^0})_l \right| + |p_l(\tau)| \cdot \left| (\widehat{|\psi^n|^2})_l(0) - (\widetilde{|\Psi^n|^2})_l \right| \\ & \quad + |q_l(\tau)| \cdot \left| (\widehat{\psi^n \phi^n})_l(0) - (\widetilde{\Psi^n \Psi^n})_l \right|. \end{aligned} \quad (4.53)$$

Combining the above two inequalities, we obtain

$$\|\gamma^{n+1} - I_N R^{n+1}\|_{H^2} \lesssim \|e_\phi^n\|_{H^2} + \varepsilon^2 \|\dot{e}_\phi^n\|_{H^2} + \|e_\psi^n\|_{H^2} + h^{2m_0}. \quad (4.54)$$

Plugging (4.54) into (4.52), we get

$$\begin{aligned} & \sum_{l=-N/2}^{N/2-1} (1 + \mu_l^2 + \mu_l^4) \left| (\widehat{\gamma^{n+1}\psi^n})_l(0) - (\widetilde{R^{n+1}\Psi^n})_l \right|^2 \\ & \lesssim \|e_\psi^n\|_{H^2}^2 + \|e_\phi^n\|_{H^2}^2 + \varepsilon^4 \|\dot{e}_\phi^n\|_{H^2}^2 + h^{2m_0}. \end{aligned} \quad (4.55)$$

Similarly, we can obtain

$$\begin{aligned} & \sum_{l=-N/2+1}^{N/2} (1 + \mu_l^2 + \mu_l^4) \left| (\widehat{\gamma^{n+1}\phi^n})_l - (\widetilde{R^{n+1}\Psi^n})_l \right|^2 \\ & \lesssim \|e_\psi^n\|_{H^2}^2 + \|e_\phi^n\|_{H^2}^2 + \varepsilon^4 \|\dot{e}_\phi^n\|_{H^2}^2 + h^{2m_0}. \end{aligned} \quad (4.56)$$

Combining (4.55), (4.56) and (4.47c), we have

$$\begin{aligned} \|\eta_\psi^n\|_{H^2} & \lesssim \tau \left(\|e_\phi^n\|_{H^2} + \varepsilon^2 \|\dot{e}_\phi^n\|_{H^2} + \|e_\psi^n\|_{H^2} + h^{m_0} \right) \\ & \lesssim \tau \left(\|e_{\phi,N}^n\|_{H^2} + \varepsilon^2 \|e_{\phi,N}^n\|_{H^2} + \|e_{\psi,N}^n\|_{H^2} + h^{m_0} \right). \end{aligned} \quad (4.57)$$

Plugging (4.50), (4.51) and (4.57) into (4.11), we get the estimate (4.48). \square

Proof of Theorem 4.1. The proof will be proceeded by the energy method and the method of mathematical induction [6, 7, 9, 11, 12]. For $n = 0$, from the initial data in the MTI-FP method (3.15)-(3.17) and noticing the assumption (4.1), we have

$$\begin{aligned} & \|e_\phi^0\|_{H^2} + \|e_\psi^0\|_{H^2} + \varepsilon^2 \|\dot{e}_\phi^0\|_{H^2} \\ &= \|\phi_0 - I_N \phi_0\|_{H^2} + \|\psi_0 - I_N \psi_0\|_{H^2} + \|\phi_1 - I_N \phi_1\|_{H^2} \lesssim h^{m_0+2} \lesssim h^{m_0}. \end{aligned}$$

In addition, using the triangle inequality, we know that there exists $h_1 > 0$ independent of ε such that for $0 < h \leq h_1$

$$\begin{aligned} \|\phi_l^0\|_{H^2} &\leq \|\phi_0\|_{H^2} + \|e_\phi^0\|_{H^2} \leq C_\phi + 1, \quad \|\dot{\phi}_l^0\|_{H^2} \leq \frac{\|\phi_1\|_{H^2}}{\varepsilon^2} + \|\dot{e}_\phi^0\|_{H^2} \leq \frac{C_\phi + 1}{\varepsilon^2}, \\ \|\psi_l^0\|_{H^2} &\leq \|\psi_0\|_{H^2} + \|e_\psi^0\|_{H^2} \leq C_\psi + 1. \end{aligned}$$

Thus (4.5)-(4.7) are valid for $n = 0$. Now we assume that (4.5)-(4.7) are valid for $0 \leq n \leq m-1 \leq T/\tau - 1$. Adding (4.20) and (4.14), we have

$$\widehat{(e_\phi^{n+1})}_l = \cos(\omega_l \tau) \widehat{(e_\phi^n)}_l + \frac{\sin(\omega_l \tau)}{\omega_l} \widehat{(\dot{e}_\phi^n)}_l + \widehat{(\xi_\phi^n)}_l + \widehat{(\eta_\phi^n)}_l, \quad (4.58a)$$

$$\widehat{(\dot{e}_\phi^{n+1})}_l = -\omega_l \sin(\omega_l \tau) \widehat{(e_\phi^n)}_l + \cos(\omega_l \tau) \widehat{(\dot{e}_\phi^n)}_l + \widehat{(\dot{\xi}_\phi^n)}_l + \widehat{(\dot{\eta}_\phi^n)}_l, \quad (4.58b)$$

$$\widehat{(e_\psi^{n+1})}_l = e^{-i\mu_l^2 \tau} \widehat{(e_\psi^n)}_l + \widehat{(\xi_\psi^n)}_l + \widehat{(\eta_\psi^n)}_l. \quad (4.58c)$$

Using the Cauchy's inequality, we obtain

$$\left| \widehat{(e_\phi^{n+1})}_l \right|^2 \leq (1 + \tau) \left| \cos(\omega_l \tau) \widehat{(e_\phi^n)}_l + \frac{\sin(\omega_l \tau)}{\omega_l} \widehat{(\dot{e}_\phi^n)}_l \right|^2 + \frac{1 + \tau}{\tau} \left| \widehat{(\xi_\phi^n)}_l + \widehat{(\eta_\phi^n)}_l \right|^2, \quad (4.59a)$$

$$\left| \widehat{(\dot{e}_\phi^{n+1})}_l \right|^2 \leq (1 + \tau) \left| \cos(\omega_l \tau) \widehat{(\dot{e}_\phi^n)}_l - \omega_l \sin(\omega_l \tau) \widehat{(e_\phi^n)}_l \right|^2 + \frac{1 + \tau}{\tau} \left| \widehat{(\dot{\xi}_\phi^n)}_l + \widehat{(\dot{\eta}_\phi^n)}_l \right|^2, \quad (4.59b)$$

$$\left| \widehat{(e_\psi^{n+1})}_l \right|^2 \leq (1 + \tau) \left| \widehat{(e_\psi^n)}_l \right|^2 + \frac{1 + \tau}{\tau} \left| \widehat{(\xi_\psi^n)}_l + \widehat{(\eta_\psi^n)}_l \right|^2. \quad (4.59c)$$

Multiplying (4.59a)-(4.59c) by $(\mu_l^2 + \frac{1}{\varepsilon^2})(1 + \mu_l^2 + \mu_l^4)$, $\varepsilon^2(1 + \mu_l^2 + \mu_l^4)$ and $\frac{1}{\varepsilon^2}(1 + \mu_l^2 + \mu_l^4)$, respectively, then adding them together and summing up for $l = -N/2, \dots, N/2 - 1$, we obtain

$$\mathcal{E}(e_{\phi,N}^{n+1}, e_{\psi,N}^{n+1}, \dot{e}_{\phi,N}^{n+1}) \leq (1 + \tau) \mathcal{E}(e_{\phi,N}^n, e_{\psi,N}^n, \dot{e}_{\phi,N}^n) + \frac{1 + \tau}{\tau} \mathcal{E}(\xi_\phi^n + \eta_\phi^n, \xi_\psi^n + \eta_\psi^n, \dot{\xi}_\phi^n + \dot{\eta}_\phi^n).$$

Using the Cauchy's inequality, we get

$$\begin{aligned} & \mathcal{E}(e_{\phi,N}^{n+1}, e_{\psi,N}^{n+1}, \dot{e}_{\phi,N}^{n+1}) - \mathcal{E}(e_{\phi,N}^n, e_{\psi,N}^n, \dot{e}_{\phi,N}^n) \\ & \lesssim \tau \mathcal{E}(e_{\phi,N}^n, e_{\psi,N}^n, \dot{e}_{\phi,N}^n) + \frac{1 + \tau}{\tau} \left[\mathcal{E}(\xi_\phi^n, \xi_\psi^n, \dot{\xi}_\phi^n) + \mathcal{E}(\eta_\phi^n, \eta_\psi^n, \dot{\eta}_\phi^n) \right]. \end{aligned} \quad (4.60)$$

Inserting (4.48) and the second inequality in (4.24) into (4.60), we get

$$\mathcal{E}(e_{\phi,N}^{n+1}, e_{\psi,N}^{n+1}, \dot{e}_{\phi,N}^{n+1}) - \mathcal{E}(e_{\phi,N}^n, e_{\psi,N}^n, \dot{e}_{\phi,N}^n) \lesssim \tau \mathcal{E}(e_{\phi,N}^n, e_{\psi,N}^n, \dot{e}_{\phi,N}^n) + \frac{\tau h^{2m_0}}{\varepsilon^2} + \frac{\tau^5}{\varepsilon^6}. \quad (4.61)$$

Summing the above inequality for $0 \leq n \leq m-1$ and then applying the discrete Gronwall's inequality, we have

$$\mathcal{E}(e_{\phi,N}^n, e_{\psi,N}^n, \dot{e}_{\phi,N}^n) \lesssim \frac{h^{2m_0}}{\varepsilon^2} + \frac{\tau^4}{\varepsilon^6}. \quad (4.62)$$

Similarly, by using the first inequality in (4.24), we obtain

$$\mathcal{E}(e_{\phi,N}^n, e_{\psi,N}^n, \dot{e}_{\phi,N}^n) \lesssim \frac{h^{2m_0}}{\varepsilon^2} + \varepsilon^2. \quad (4.63)$$

Combining (4.11), (4.62) and (4.63), we get that (4.5) is valid for $n = m$, which implies

$$\|e_{\phi}^m\|_{H^2} + \|e_{\psi}^m\|_{H^2} + \varepsilon^2 \|\dot{e}_{\phi}^m\|_{H^2} \leq h^{m_0} + \tau.$$

Using the triangle inequality, we obtain that there exist $h_2 > 0$ and $\tau_2 > 0$ independent of ε such that

$$\begin{aligned} \|\phi_I^m\|_{H^2} &\leq \|\phi(\cdot, t_m)\|_{H^2} + \|e_{\phi}^m\|_{H^2} \leq C_{\phi} + 1, \\ \|\psi_I^m\|_{H^2} &\leq \|\psi(\cdot, t_m)\|_{H^2} + \|e_{\psi}^m\|_{H^2} \leq C_{\psi} + 1, \quad 0 < h \leq h_2, \quad 0 < \tau \leq \tau_2, \\ \|\dot{\phi}_I^m\|_{H^2} &\leq \|\partial_t \phi(\cdot, t_m)\|_{H^2} + \|\dot{e}_{\phi}^m\|_{H^2} \leq \frac{C_{\phi} + 1}{\varepsilon^2}. \end{aligned}$$

Thus (4.7) is also valid for $n = m$. Then the proof is completed by choosing $\tau_0 = \min\{\tau_1, \tau_2\}$ and $h_0 = \min\{h_1, h_2\}$. \square

Remark 4.1 Here we emphasize that Theorem 4.1 holds in two/three dimensions (2D/3D) and the above approach can be directly extended to 2D/3D without extra effort. The only thing needs to be taken care of is the Sobolev inequality used in Lemma 4.4 in 2D/3D

$$\|u\|_{L^\infty(\Omega)} \lesssim \|u\|_{H^2(\Omega)}, \quad \|u\|_{W^{1,p}(\Omega)} \lesssim \|u\|_{H^2(\Omega)}, \quad 1 < p < 6, \quad (4.64)$$

where Ω is a bounded domain in 2D/3D. By using the assumption (4.7), Lemma 4.4 still holds in 2D/3D. Thus (4.7) and the error bounds can be proved by the method of mathematical induction since our scheme is explicit.

Remark 4.2 Under a weaker assumption on the regularity of the solution, i.e. there exists an integer $m_0 \geq 4$ such that $\phi \in C^1([0, T]; H_p^{m_0+3}(\Omega))$, $\psi \in C([0, T]; H_p^{m_0+1}(\Omega)) \cap C^1([0, T]; H_p^{m_0-1}(\Omega)) \cap C^2([0, T]; H_p^{m_0-3}(\Omega))$ and

$$\begin{aligned} \|\phi\|_{L^\infty([0, T]; H^{m_0+3})} + \varepsilon^2 \|\partial_t \phi\|_{L^\infty([0, T]; H^{m_0+3})} &\lesssim 1, \\ \|\psi\|_{L^\infty([0, T]; H^{m_0+1})} + \|\partial_t \psi\|_{L^\infty([0, T]; H^{m_0-1})} + \varepsilon^2 \|\partial_{tt} \psi\|_{L^\infty([0, T]; H^{m_0-3})} &\lesssim 1, \end{aligned} \quad (4.65)$$

we can establish an H^1 -norm estimate of the MTI-FP method in Theorem 4.1 by a very similar proof with all the H^2 -norm in the proof being changed into H^1 -norm.

In 1D case, the H^1 -norm estimate of the MTI-FP method holds without any stability (or CFL) condition. However in 2D/3D, due to the use of the inverse inequality to provide the bound in l^∞ -norm of the numerical solution [6–8], we need impose the technical condition

$$\tau \lesssim \rho_d(h), \quad \text{with} \quad \rho_d(h) = \begin{cases} 1/|\ln h|, & d = 2, \\ \sqrt{h}, & d = 3. \end{cases} \quad (4.66)$$

Of course, if the solution of the KGS is smooth enough, we can always adapt the H^2 -norm estimate in Theorem 4.1 and thus there is no need to assume the stability (or CFL) condition (4.66).

5 Numerical results

In this section, we report numerical results to demonstrate the uniform convergence of the MTI-FP method for $\varepsilon \in (0, 1]$ and apply it to numerically study convergence rates of the KGS equations to its limiting models (1.9) and (1.11) in the nonrelativistic limit regime. In order to do so, we take $d = 1$ and $\mu = \lambda = 1$ in (1.3) and choose the initial data in (1.5) with $d = 1$ as

$$\psi_0(x) = \frac{1+i}{2} \operatorname{sech}\left(\frac{x^2}{2}\right), \quad \phi_0(x) = \frac{1}{2}e^{-x^2}, \quad \phi_1(x) = \frac{1}{\sqrt{2}}e^{-x^2}, \quad x \in \mathbb{R}. \quad (5.1)$$

5.1 Accuracy test

The problem (1.3) with $d = 1$ and (5.1) is solved on a bounded interval $\Omega = [-32, 32]$, i.e. $b = -a = 32$, which is large enough to guarantee that the periodic boundary condition does not introduce a significant aliasing error relative to the original problem. To quantify the error, we introduce two error functions:

$$\begin{aligned} e_{\phi, \varepsilon}^{\tau, h}(t = t_n) &:= \|\phi(\cdot, t_n) - \phi_I^n\|_{H^2}, & e_{\phi, \infty}^{\tau, h}(t) &:= \max_{0 < \varepsilon \leq 1} \left\{ e_{\phi, \varepsilon}^{\tau, h}(t) \right\}, \\ e_{\psi, \varepsilon}^{\tau, h}(t = t_n) &:= \|\psi(\cdot, t_n) - \psi_I^n\|_{H^2}, & e_{\psi, \infty}^{\tau, h}(t) &:= \max_{0 < \varepsilon \leq 1} \left\{ e_{\psi, \varepsilon}^{\tau, h}(t) \right\}. \end{aligned}$$

Since the analytical solution to this problem is not available, so the ‘reference’ solution here is obtained numerically by the MTI-FP method (3.15)-(3.17) with very fine mesh $h = 1/32$ and time step $\tau = 5 \times 10^{-6}$. Tab. 5.1 and Tab. 5.2 show the spatial error of the MTI-FP method for ψ and ϕ , respectively, at $t = 1$ under different ε and h with a very small time step $\tau = 5 \times 10^{-6}$ such that the discretization error in time is negligible. Tab. 5.3 and Tab. 5.4 show the temporal error of the MTI-FP method for ψ and ϕ , respectively, at $t = 1$ under different ε and τ with a small mesh size $h = 1/16$ such that the discretization error in space is negligible.

From Tabs. 5.1-5.4 and extensive additional results not shown here for brevity, we can draw the following observations:

(i) The MTI-FP method is spectrally accurate in space, which is uniformly for $0 < \varepsilon \leq 1$ (cf. Tabs. 5.1&5.2). The approximation in ϕ is more accurate than ψ .

Table 5.1: Spatial error analysis: $e_{\psi,\varepsilon}^{\tau,h}(t=1)$ with $\tau = 5 \times 10^{-6}$ for different ε and h .

$e_{\psi,\varepsilon}^{\tau,h}(t=1)$	$h_0=1$	$h_0/2$	$h_0/4$	$h_0/8$	$h_0/16$
$\varepsilon_0=1$	6.90E-1	1.39E-1	1.70E-3	2.44E-7	8.96E-11
$\varepsilon_0/2$	7.57E-1	1.40E-1	1.70E-3	2.40E-7	8.37E-11
$\varepsilon_0/2^2$	6.37E-1	1.45E-1	1.70E-3	2.40E-7	9.19E-11
$\varepsilon_0/2^3$	6.46E-1	1.30E-1	1.90E-3	2.40E-7	8.89E-11
$\varepsilon_0/2^4$	6.47E-1	1.31E-1	1.80E-3	2.38E-7	9.26E-11
$\varepsilon_0/2^5$	6.47E-1	1.31E-1	1.80E-3	2.38E-7	9.66E-11
$\varepsilon_0/2^6$	6.46E-1	1.27E-1	1.70E-3	2.52E-7	8.76E-11
$\varepsilon_0/2^7$	6.46E-1	1.31E-1	1.80E-3	2.38E-7	9.08E-11
$\varepsilon_0/2^8$	6.46E-1	1.27E-1	1.70E-3	2.52E-7	9.08E-11
$\varepsilon_0/2^9$	6.46E-1	1.31E-1	1.80E-3	2.38E-7	1.02E-10
$\varepsilon_0/2^{11}$	6.46E-1	1.31E-1	1.80E-3	2.38E-7	9.07E-11
$\varepsilon_0/2^{13}$	6.46E-1	1.31E-1	1.80E-3	2.38E-7	8.61E-11

Table 5.2: Spatial error analysis: $e_{\phi,\varepsilon}^{\tau,h}(t=1)$ with $\tau = 5 \times 10^{-6}$ for different ε and h .

$e_{\phi,\varepsilon}^{\tau,h}(t=1)$	$h_0=1$	$h_0/2$	$h_0/4$	$h_0/8$	$h_0/16$
$\varepsilon_0=1$	4.74E-2	1.60E-3	3.64E-6	3.57E-10	9.23E-11
$\varepsilon_0/2$	8.29E-2	1.90E-3	1.23E-5	3.92E-10	6.18E-11
$\varepsilon_0/2^2$	2.32E-1	1.12E-2	4.61E-5	1.46E-9	5.24E-11
$\varepsilon_0/2^3$	3.49E-1	1.30E-3	9.31E-5	5.33E-10	4.58E-11
$\varepsilon_0/2^4$	2.64E-1	1.20E-3	3.35E-5	5.16E-9	4.49E-11
$\varepsilon_0/2^5$	2.71E-1	2.09E-4	6.92E-6	8.76E-10	5.49E-11
$\varepsilon_0/2^6$	1.13E-1	5.27E-4	2.56E-6	1.73E-10	6.04E-11
$\varepsilon_0/2^7$	3.99E-1	4.29E-4	6.28E-7	4.23E-11	6.20E-11
$\varepsilon_0/2^8$	6.74E-2	5.90E-4	7.20E-8	2.38E-11	6.20E-11
$\varepsilon_0/2^9$	3.37E-1	5.13E-5	2.78E-8	1.86E-11	6.51E-11
$\varepsilon_0/2^{11}$	3.09E-1	7.75E-4	2.98E-9	1.73E-11	6.19E-11
$\varepsilon_0/2^{13}$	1.87E-1	3.64E-4	3.33E-9	1.72E-11	6.13E-11

(ii) The MTI-FP method converges uniformly in time for $\varepsilon \in (0, 1]$ with linear/quadratical convergence rate for ψ/ϕ , respectively (cf. last row in Tabs. 5.3&5.4). In addition, for each fixed $\varepsilon = \varepsilon_0 > 0$, when τ is small enough, it converges quadratically in time (cf. each row in the upper triangle of Tabs. 5.3&5.4); for fixed τ , when $0 < \varepsilon \ll 1$, it converges quadratically in terms of ε (cf. each column in the lower triangle of Tabs. 5.3&5.4).

(iii) The practical temporal uniform convergence order of the MTI-FP in ϕ is better than the theoretical error estimate result. In fact, the temporal local truncation error of the MTI-FP in ϕ is at $O(\tau^3)$, but due to the nonlinear coupling with ψ , the theoretical error bound drops down to uniform linear convergence rate. The rigorous mathematical analysis is still on-going.

(iv) In the nonrelativistic limit regime, i.e. $0 < \varepsilon \ll 1$, the mesh strategy (or ε -scalability) of the MTI-FP method is $\tau = O(1)$ and $h = O(1)$.

Table 5.3: Temporal error analysis: $e_{\psi,\varepsilon}^{\tau,h}(t=1)$ and $e_{\psi,\infty}^{\tau,h}(t=1)$ with $h = 1/16$ for different ε and τ .

$e_{\psi,\varepsilon}^{\tau,h}(t=1)$	$\tau_0 = 0.2$	$\tau_0/2^2$	$\tau_0/2^4$	$\tau_0/2^6$	$\tau_0/2^8$	$\tau_0/2^{10}$	$\tau_0/2^{12}$
$\varepsilon_0 = 1$	1.31E-1	1.60E-2	1.00E-3	6.20E-5	3.84E-6	2.35E-7	1.77E-8
rate	—	1.52	2.00	2.01	2.01	2.02	1.87
$\varepsilon_0/2$	1.59E-1	2.05E-2	1.30E-3	7.70E-5	4.76E-6	2.92E-7	1.37E-8
rate	—	1.48	1.99	2.04	2.01	2.01	2.20
$\varepsilon_0/2^2$	1.20E-1	1.48E-2	1.20E-3	7.65E-5	4.79E-6	2.95E-7	1.38E-8
rate	—	1.35	1.81	1.99	2.00	2.01	2.20
$\varepsilon_0/2^3$	2.23E-2	5.30E-3	5.19E-4	3.20E-5	2.00E-6	1.23E-7	5.77E-9
rate	—	1.04	1.68	2.00	2.00	2.01	2.20
$\varepsilon_0/2^4$	9.50E-3	3.50E-3	2.20E-3	1.32E-4	8.06E-6	4.94E-7	2.31E-8
rate	—	0.72	0.33	2.03	2.02	2.01	2.20
$\varepsilon_0/2^5$	1.40E-3	1.00E-3	6.34E-4	4.67E-4	3.35E-5	2.02E-6	9.41E-8
rate	—	0.24	0.33	0.22	1.91	2.02	2.20
$\varepsilon_0/2^6$	3.82E-4	5.65E-5	1.43E-4	1.51E-4	1.16E-4	8.29E-6	3.80E-7
rate	—	1.38	-0.67	-0.18	0.19	1.90	2.22
$\varepsilon_0/2^7$	5.64E-5	2.30E-5	1.28E-5	3.61E-5	3.83E-5	2.89E-5	1.58E-6
rate	—	0.65	0.42	-0.75	-0.04	0.20	2.10
$\varepsilon_0/2^8$	2.10E-5	5.60E-6	6.41E-6	2.61E-6	1.10E-5	1.12E-5	5.70E-6
rate	—	0.95	-0.10	0.65	-1.00	-0.01	0.49
$\varepsilon_0/2^9$	5.82E-6	4.05E-6	2.40E-6	2.00E-6	2.41E-6	8.49E-7	2.50E-7
rate	—	0.26	0.38	0.13	-0.13	0.75	0.88
$\varepsilon_0/2^{11}$	2.09E-7	2.19E-7	1.67E-7	1.58E-7	8.48E-8	1.43E-7	4.63E-8
rate	—	-0.04	0.20	0.04	0.45	-0.38	0.81
$\varepsilon_0/2^{13}$	1.78E-8	8.93E-9	5.82E-9	5.68E-9	7.48E-9	1.92E-9	9.46E-9
rate	—	0.50	0.31	0.02	-0.20	0.98	-1.15
$e_{\psi,\infty}^{\tau,h}(t=1)$	1.59E-1	2.05E-2	2.20E-3	4.67E-4	1.16E-4	2.89E-5	5.70E-6
rate	—	1.48	1.61	1.11	1.00	1.00	1.17

5.2 Convergence rates of KGS to its limiting models

Let ψ and ϕ be the solution of the problem (1.3) with $d = \mu = \lambda = 1$ and (5.1), which is obtained numerically on a bounded interval $\Omega = [-512, 512]$ by the MTI-FP method (3.15)-(3.17) with very fine mesh $h = 1/16$ and time step $\tau = 10^{-4}$. Similarly, let ψ_{sw} and z_{sw} be the solution of the problem (1.9) with $d = \mu = \lambda = 1$ and (1.10) and (5.1), which is obtained numerically on the bounded interval Ω by the EWI-FP method [7] with very fine mesh $h = 1/16$ and time step $\tau = 10^{-4}$; and let ψ_s and z_s be the solution of the problem (1.11) with $d = \mu = \lambda = 1$ and (1.12) and (5.1), which is obtained numerically on the bounded interval Ω by the TSFP method [5] with very fine mesh $h = 1/16$ and time step $\tau = 10^{-4}$. Denote

$$\phi_{\text{sw}}(x, t) = e^{it/\varepsilon^2} z_{\text{sw}}(x, t) + e^{-it/\varepsilon^2} \bar{z}_{\text{sw}}(x, t), \quad \phi_s(x, t) = e^{it/\varepsilon^2} z_s(x, t) + e^{-it/\varepsilon^2} \bar{z}_s(x, t),$$

and define the error functions as

$$\begin{aligned} \eta_{\text{sw}}(t) &:= \|\phi(\cdot, t) - \phi_{\text{sw}}(\cdot, t)\|_{H^1} + \|\psi(\cdot, t) - \psi_{\text{sw}}(\cdot, t)\|_{H^1}, \\ \eta_s(t) &:= \|\phi(\cdot, t) - \phi_s(\cdot, t)\|_{H^1} + \|\psi(\cdot, t) - \psi_s(\cdot, t)\|_{H^1}. \end{aligned} \quad (5.2)$$

Table 5.4: Temporal error analysis: $e_{\phi,\varepsilon}^{\tau,h}(t=1)$ and $e_{\phi,\infty}^{\tau,h}(t=1)$ with $h = 1/16$ for different ε and τ .

$e_{\phi,\varepsilon}^{\tau,h}(t=1)$	$\tau_0 = 0.2$	$\tau_0/2^2$	$\tau_0/2^4$	$\tau_0/2^6$	$\tau_0/2^8$	$\tau_0/2^{10}$	$\tau_0/2^{12}$
$\varepsilon_0 = 1$	3.74E-2	1.50E-3	3.82E-5	1.78E-6	1.09E-7	6.73E-9	3.32E-10
rate	—	2.32	2.65	2.21	2.01	2.01	2.17
$\varepsilon_0/2$	9.32E-2	4.80E-3	2.29E-4	1.35E-5	8.32E-7	5.11E-8	2.39E-9
rate	—	2.14	2.19	2.04	2.01	2.01	2.20
$\varepsilon_0/2^2$	9.65E-2	1.26E-2	7.09E-4	4.23E-5	2.61E-6	1.60E-7	7.51E-9
rate	—	1.47	2.07	2.03	2.01	2.01	2.20
$\varepsilon_0/2^3$	2.33E-2	4.10E-3	3.02E-4	1.95E-5	1.21E-6	7.47E-8	3.50E-9
rate	—	1.25	1.88	1.98	2.00	2.01	2.20
$\varepsilon_0/2^4$	8.70E-3	3.30E-3	9.09E-5	6.00E-6	3.82E-7	2.36E-8	1.11E-9
rate	—	0.70	2.59	1.96	1.99	2.00	2.20
$\varepsilon_0/2^5$	9.59E-4	8.29E-4	2.42E-4	3.73E-6	2.89E-7	1.81E-8	8.53E-10
rate	—	0.11	0.89	3.01	1.85	2.00	2.20
$\varepsilon_0/2^6$	4.71E-4	1.19E-4	1.33E-5	3.80E-6	2.13E-7	1.67E-8	8.01E-10
rate	—	0.99	1.58	0.90	2.07	1.84	2.19
$\varepsilon_0/2^7$	8.27E-5	2.88E-5	2.49E-6	1.76E-7	4.95E-8	1.30E-8	7.86E-10
rate	—	0.76	1.76	1.91	0.92	0.96	2.02
$\varepsilon_0/2^8$	3.33E-5	5.69E-6	3.28E-7	2.18E-8	3.37E-9	1.70E-9	6.57E-10
rate	—	1.28	2.06	1.95	1.35	0.49	0.69
$\varepsilon_0/2^9$	8.92E-6	3.98E-6	1.04E-7	5.46E-9	3.66E-10	5.75E-11	2.04E-11
rate	—	0.58	2.63	2.12	1.95	1.34	0.75
$\varepsilon_0/2^{11}$	3.05E-7	1.59E-7	5.64E-9	1.32E-9	7.21E-11	7.09E-11	7.43E-11
rate	—	0.47	2.41	1.05	2.10	0.01	-0.03
$\varepsilon_0/2^{13}$	2.30E-8	7.26E-9	2.21E-9	7.77E-10	7.74E-10	7.74E-10	7.74E-10
rate	—	0.83	0.86	0.76	0.01	0.00	0.00
$e_{\phi,\infty}^{\tau,h}(t=1)$	9.65E-2	1.26E-2	7.09E-4	4.23E-5	2.61E-6	1.60E-7	7.51E-9
rate	—	1.46	2.00	2.00	2.00	2.00	2.00

Fig. 5.1 depicts $\eta_{\text{sw}}(t)$ and $\eta_s(t)$ for different ε . From Fig. 5.1 and additional similar results not shown here for brevity, when $\varepsilon \rightarrow 0$, we can observe numerically the solution of the KGS equations (1.3) converges quadratically and uniformly in time to that of its limiting model (1.9) as

$$\|\Psi(\cdot, t) - \Psi_{\text{sw}}(\cdot, t)\|_{H^1(\Omega)} + \|\phi(\cdot, t) - \phi_{\text{sw}}(\cdot, t)\|_{H^1(\Omega)} \leq C_0 \varepsilon^2, \quad t \geq 0, \quad (5.3)$$

and respectively, to its limiting model (1.11) as

$$\|\Psi(\cdot, t) - \Psi_s(\cdot, t)\|_{H^1(\Omega)} + \|\phi(\cdot, t) - \phi_s(\cdot, t)\|_{H^1(\Omega)} \leq (C_1 + C_2 T) \varepsilon^2, \quad 0 \leq t \leq T, \quad (5.4)$$

where C_0, C_1 and $C_2 > 0$ are positive constants which are independent of ε and T . Rigorous mathematical justification of these numerical observations is on-going.

6 Conclusion

A multiscale time integrator Fourier pseudospectral (MTI-FP) method was proposed and analyzed for solving the Klein-Gordon-Schrödinger (KGS) equations with a di-




Fig. 5.1: Time evolution of $\eta_{sw}(t)$ and $\eta_s(t)$ for different ε .

dimensionless parameter $0 < \varepsilon \leq 1$ which is inversely proportional to the speed of light. The key ideas for designing the MTI-FP method are based on (i) carrying out a multiscale decomposition by frequency at each time step with proper choice of

transmission conditions between adjacent time intervals, and (ii) adapting the Fourier spectral method for spatial discretization and the EWI for integrating second-order highly oscillating ODEs. Rigorous error bounds for the MTI-FP method were established, which imply that the MTI-FP method converges uniformly and optimally in space with spectral convergence rate, and uniformly in time with linear convergence rate for $\varepsilon \in (0, 1]$ and optimally with quadratic convergence rate in the regimes when either $\varepsilon = O(1)$ or $0 < \varepsilon \leq \tau$. Numerical results confirmed these error bounds and showed that the MTI-FP method offers compelling advantages over classical methods in the nonrelativistic regime $0 < \varepsilon \ll 1$. Numerical results suggest that the solution of the KGS equations converges quadratically to that of its limiting models when $\varepsilon \rightarrow 0$.

Acknowledgements This work was supported by the Ministry of Education of Singapore grant R-146-000-196-112. Part of this work was done when the authors were visiting the Institute for Mathematical Science at the National University of Singapore in 2015.

Appendix A. Explicit formulas for the coefficients used in (3.14)

Define the functions

$$\left\{ \begin{array}{l} p_l(s) := \int_0^s \frac{\sin(\omega_l(s-\theta))}{\varepsilon^2 \omega_l} d\theta, \quad p'_l(s) := \int_0^s \frac{\cos(\omega_l(s-\theta))}{\varepsilon^2} d\theta, \\ q_l(s) := \int_0^s \frac{\sin(\omega_l(s-\theta))}{\varepsilon^2 \omega_l} \theta d\theta, \quad q'_l(s) := \int_0^s \frac{\cos(\omega_l(s-\theta))}{\varepsilon^2} \theta d\theta, \\ c_l^+(s) := i e^{-i\mu_l^2 s} \int_0^s e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} d\theta, \quad c_l^-(s) := i e^{-i\mu_l^2 s} \int_0^s e^{i(\mu_l^2 - \frac{1}{\varepsilon^2})\theta} d\theta, \\ d_l^+(s) := i e^{-i\mu_l^2 s} \int_0^s e^{i(\mu_l^2 + \frac{1}{\varepsilon^2})\theta} \theta d\theta, \quad d_l^-(s) := i e^{-i\mu_l^2 s} \int_0^s e^{i(\mu_l^2 - \frac{1}{\varepsilon^2})\theta} \theta d\theta. \end{array} \right. \quad (\text{A.1})$$

Taking $s = \tau$ in (A.1), after a detailed computation, we have

$$\begin{aligned} p_l(\tau) &= \frac{1 - \cos(\omega_l \tau)}{\varepsilon^2 \omega_l^2}, & p'_l(\tau) &= \frac{\sin(\omega_l \tau)}{\varepsilon^2 \omega_l}, \\ q_l(\tau) &= \frac{\tau \omega_l - \sin(\omega_l \tau)}{\varepsilon^2 \omega_l^3}, & q'_l(\tau) &= \frac{1 - \cos(\omega_l \tau)}{\varepsilon^2 \omega_l^2}, \\ c_l^+(\tau) &= \frac{\varepsilon^2 e^{-i\mu_l^2 \tau} \left(e^{i\tau(\mu_l^2 + \frac{1}{\varepsilon^2})} - 1 \right)}{1 + \varepsilon^2 \mu_l^2}, & c_l^-(\tau) &= \frac{\varepsilon^2 \left(e^{-i\tau \mu_l^2} - e^{-i\frac{\tau}{\varepsilon^2}} \right)}{1 - \varepsilon^2 \mu_l^2}, \\ d_l^+(\tau) &= \frac{i\varepsilon^2}{(1 + \varepsilon^2 \mu_l^2)^2} \left[e^{\frac{i\tau}{\varepsilon^2}} (\varepsilon^2 - i\tau(1 + \varepsilon^2 \mu_l^2)) - \varepsilon^2 e^{-i\mu_l^2 \tau} \right], \\ d_l^-(\tau) &= \frac{i\varepsilon^2}{(1 - \varepsilon^2 \mu_l^2)^2} \left[e^{-\frac{i\tau}{\varepsilon^2}} (\varepsilon^2 - i\tau(\varepsilon^2 \mu_l^2 - 1)) - \varepsilon^2 e^{-i\mu_l^2 \tau} \right]. \end{aligned}$$

References

1. Adams, R.: Sobolev Spaces. Boston, MA: Academic Press, 1975.
2. Added, H., Added, S.: Equations of Langmuir turbulence and nonlinear Schrödinger equation: Smoothness and approximation. *J. Funct. Anal.* 79, 183-210 (1988)
3. Biler, P.: Attractors for the system of Schrödinger and Klein-Gordon equations with Yukawa coupling. *SIAM J. Math. Anal.* 21, 1190-1212 (1990)
4. Banquet, C., Ferreira, L.C.F., Villamizar-Roa, E.J. : On existence and scattering theory for the Klein-Gordon-Schrödinger system in an infinite L^2 -norm setting. *Ann. Mat. Pura Appl.* DOI 10.1007/s10231-013-0398-7 (2014)
5. Bao, W., Cai, Y.: Mathematical theory and numerical methods for Bose-Einstein condensation *Kinet. Relat. Models*, 61-135 (2013)
6. Bao, W., Cai, Y.: Uniform error estimates of finite difference methods for the nonlinear Schrödinger equation with wave operator. *SIAM J. Numer. Anal.* 50, 492-521 (2012)
7. Bao, W., Cai, Y.: Uniform and optimal error estimates of an exponential wave integrator sine pseudospectral method for the nonlinear Schrödinger equation with wave operator. *SIAM J. Numer. Anal.* 52, 1103-1127 (2014)
8. Bao, W., Cai, Y., Zhao, X.: A uniformly accurate multiscale time integrator pseudospectral method for the Klein-Gordon equation in the nonrelativistic limit regime. *SIAM J. Numer. Anal.* 52, 2488-2511 (2014)
9. Bao, W., Dong, X.: Analysis and comparison of numerical methods for the Klein-Gordon equation in the nonrelativistic limit regime. *Numer. Math.* 120, 189-229 (2012)
10. Bao, W., Dong, X., Wang, S.: Singular limits of Klein-Gordon-Schrödinger equations to Schrödinger-Yukawa equations. *Multiscale Model. Simul.* 8, 1742-1769 (2010)
11. Bao, W., Dong, X., Zhao, X.: An exponential wave integrator sine pseudospectral method for the Klein-Gordon-Zakharov system. *SIAM J. Sci. Comput.* 25, A2903-A2927 (2013)
12. Bao, W., Dong, X., Zhao, X.: Uniformly accurate multiscale time integrators for highly oscillatory second order differential equations. *J. Math. Study* 47, 111-150 (2014)
13. Bao, W., Yang, L.: Efficient and accurate numerical methods for the Klein-Gordon-Schrödinger equations. *J. Comput. Phys.* 225, 1863-1893 (2007)
14. Cohen, D.: Analysis and numerical treatment of highly oscillatory differential equations. PhD thesis, Université de Genève (2004)
15. Cohen, D., Hairer, E., Lubich, Ch.: Modulated Fourier expansions of highly oscillatory differential equations. *Found. Comput. Math.* 3, 327-345 (2003)
16. Cohen, D.: Conservation properties of numerical integrators for highly oscillatory Hamiltonian systems. *IMA J. Numer. Anal.* 26, 34-59 (2005)
17. Cohen, D., Hairer, E., Lubich, Ch.: Conservation of energy, momentum and actions in numerical discretizations of non-linear wave equations. *Numer. Math.* 110, 113-143 (2008)
18. Chartier, Ph., Crouseilles, N., Lemou, M., Méhats, F.: Uniformly accurate numerical schemes for highly oscillatory Klein-Gordon and nonlinear Schrödinger equations. *Numer. Math.* 129, 211-250 (2015)
19. Chartier, Ph., Mauser, N.J., Méhats, F., Zhang, Y.: Solving highly-oscillatory NLS with SAM: numerical efficiency and geometric properties. arXiv:1308.1217 [math.NA]
20. Deuffhard, P.: A study of extrapolation methods based on multistep schemes without parasitic solutions. *ZAMP* 30, 177-189 (1979)
21. Faou, E., Schratz, K.: Asymptotic preserving schemes for the Klein-Gordon equation in the nonrelativistic limit regime. *Numer. Math.* 126, 441-469 (2014)
22. Fukuda, I., Tsutsumi, M.: On the Yukawa-coupled Klein-Gordon-Schrödinger equations in three space dimensions. *Proc. Jpn. Acad.* 51, 402-405 (1975)
23. Fukuda, I., Tsutsumi, M.: On coupled Klein-Gordon-Schrödinger equations II. *J. Math. Anal. Appl.* 66, 358-378 (1978)
24. Fukuda, I., Tsutsumi, M.: On coupled Klein-Gordon-Schrödinger equations III. *Math. Jpn.* 24, 307-321 (1979)
25. Guo, B.L.: Global solution for some problem of a class of equations in interaction of complex Schrödinger field and real Klein-Gordon field. *Sci. China Ser. A* 25, 97-107 (1982)
26. Guo, B.L., Li, Y.S.: Attractor for dissipative Klein-Gordon-Schrödinger equations in \mathbb{R}^3 . *J. Differential Equations* 136, 356-377 (1997)

27. Guo, B.L., Miao, C.X.: Asymptotic behavior of coupled Klein-Gordon-Schrödinger equations. *Sci. China Ser. A* 25, 705-714 (1995)
28. Gautschi, W.: Numerical integration of ordinary differential equations based on trigonometric polynomials. *Numer. Math.* 3, 381-397 (1961)
29. Grimm, V.: A note on the Gautschi-type method for oscillatory second-order differential equations. *Numer. Math.* 102, 61-66 (2005)
30. Grimm, V., Hochbruck, M.: Error analysis of exponential integrators for oscillatory second-order differential equations. *J. Phys.A: Math. Gen.* 39 (2006)
31. Gottlieb, D., Orszag, S.A.: *Numerical Analysis of Spectral Methods: Theory and Applications*. Society for Industrial and Applied Mathematics, Philadelphia, 1993.
32. Garcia-Archilla, B., Sanz-Serna, J. M., Skeel, R.D.: Long-time-step methods for oscillatory differential equations. *SIAM J. Sci. Comput.* 20, 930-963 (1998)
33. Hochbruck, M., Lubich, Ch.: A Gautschi-type method for oscillatory second-order differential equations. *Numer. Math.* 83, 403-426 (1999)
34. Hairer, E., Lubich, Ch.: Long-time energy conservation of numerical methods for oscillatory differential equations. *SIAM J. Numer. Anal.* 38, 414-441 (2000).
35. Hairer, E., Lubich, Ch.: On the energy distribution in Fermi-Pasta-Ulam lattices. *Arch. Ration. Mech. Anal.* 205, 993-1029 (2012)
36. Hairer, E., Lubich, Ch., Wanner, G.: *Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations*. Springer, Berlin (2006)
37. Hayashi, N., von Wahl, W.: On the global strong solutions of coupled Klein-Gordon-Schrödinger equations, *J. Math. Soc. Jpn.* 39, 489-497 (1987)
38. Lu, Y., Zhang, Z.: Long time behavior of the quadratic Klein-Gordon equation in the nonrelativistic limit regime, arXiv: 1412.2787 (math.AP)
39. Machihara, S., Nakanishi, K., Ozawa, T.: Nonrelativistic limit in the energy space for nonlinear Klein-Gordon equations. *Math. Ann.* 322, 603-621 (2002)
40. Masmoudi, N., Nakanishi, K.: From nonlinear Klein-Gordon equation to a system of coupled nonlinear Schrödinger equations. *Math. Ann.* 324, 359-389 (2002)
41. Masmoudi, N., Nakanishi, K.: From the Klein-Gordon-Zakharov system to the nonlinear Schrödinger equation. *J. Hyperbolic Differ. Equ.* 2, 975-1008 (2005)
42. Makhankov, V.G.: Dynamics of classical solitons (in non-integrable systems), *Phys. Lett. C* 35, 1-128 (1978)
43. Missaoui, S., Zahrouni, E.: Regularity of the attractor for a coupled Klein-Gordon-Schrödinger system with cubic nonlinearities in R^2 . *Commun. Pure Appl. Anal.* 14, 695-716 (2015)
44. Sanz-Serna, J.M.: Mollified impulse methods for highly oscillatory differential equations. *SIAM J. Numer. Anal.* 46, 1040-1059 (1998)
45. Wang, M.L., Zhou, Y.B.: The periodic wave solutions for the Klein-Gordon-Schrödinger equations. *Phys. Lett. A* 318, 84-92 (2003)
46. Wang, T.C.: Optimal point-wise error estimate of a compact difference scheme for the Klein-Gordon-Schrödinger equation. *J. Math. Anal. Appl.* 412, 155-167 (2014)
47. Shi, Q.H., Li, W.T., Wang, S.: Wellposedness in energy space for the nonlinear Klein-Gordon-Schrödinger system. *Appl. Math. Comput.* 251, 55-64 (2015)
48. Shen, J., Tang, T.: *Spectral and High-Order Methods with Applications*. Science Press, Beijing, 2006.
49. Shi, Q.H., Wang, S., Li, Y.: Existence and uniqueness of energy solution to Klein-Gordon-Schrödinger equations. *J. Differential Equations* 252, 168-180 (2012)
50. Xiang, X.M.: Spectral method for solving the system of equations of Schrödinger-Klein-Gordon field. *J. Comput. Appl. Math.* 21, 161-171 (1988)
51. Zhang, L.M.: Convergence of a conservative difference scheme for a class of Klein-Gordon-Schrödinger equations in one space dimension. *Appl. Math. Comput.* 163, 343-355 (2005)