ON GLOBAL SOLUTIONS FOR QUASILINEAR ONE-DIMENSIONAL PARABOLIC PROBLEMS WITH DYNAMICAL BOUNDARY CONDITIONS

SIMON GVELESIANI, FRIEDRICH LIPPOTH, AND CHRISTOPH WALKER

ABSTRACT. We provide sufficient and almost optimal conditions for global existence of classical solutions in parabolic Hölder spaces to quasilinear one-dimensional parabolic problems with dynamical boundary conditions.

1. INTRODUCTION

The focus of the present research is on the one-dimensional quasilinear parabolic equation

$$u_t - a(t, x, u, u_x)u_{xx} = f(t, x, u, u_x), \qquad (t, x) \in (0, T) \times (-\ell, \ell), \tag{1.1}$$

supplemented with a nonlinear dynamical boundary condition

$$u_t \pm b(t, x, u, u_x)u_x = g(t, x, u, u_x), \qquad t \in (0, T), \quad x = \pm \ell,$$
(1.2)

and the initial condition $u(0, \cdot) = u^0$ on $[-\ell, \ell]$, where T > 0 and $\ell > 0$ are given fixed numbers.¹ Equations of the form (1.1), (1.2) occur in various fields of natural sciences, we refer e.g. to [1, 7, 10, 17] and the references therein. In the past decades many different aspects of problems with dynamical boundary conditions (also in higher space dimensions) have been investigated by means of different techniques (e.g. [2, 5–8, 10, 16, 18, 25, 31]) for well-posedness issues, also related to the sign of the function *b* in (1.2) (e.g. [3, 5, 28, 30]) or for possibly degenerate equations (see [16] and the references therein). Research has, of course, also focused on questions regarding global existence and related *a priori* estimates and blow up phenomena [3, 4, 7, 9–11, 13–15, 25, 26]. None of these lists of references is complete though.

The starting point of our investigations are, on the one hand, the results of [7] for equation (1.1) subject to the gradient-independent dynamical boundary condition

$$u_t \pm b(t, x, u)u_x = g(t, x, u), \qquad t \in (0, T), \quad x = \pm \ell,$$
(1.3)

and, on the other hand, the results of [29] related to (1.1) but subject to more standard boundary conditions of Dirichlet, Neuman, or Robin type. In [7] criteria were found for the existence of global (i.e. existing on the whole time interval [0, T)) classical solutions for the boundary value problem (1.1), (1.3), though not restricted to one dimension. Roughly speaking, it was shown that if a = b and if the growth of the right-hand side f(t, x, u, p) in the gradient variable p is not faster than $1 + |p|^{1+\alpha}$, then bounded solutions are global provided that $\alpha = 0$ in the general quasilinear case and $\alpha \in [0, 1)$ in the semilinear case (i.e. if a = b is also independent of u). Furthermore, it was shown in [7] that if $\alpha > 1$, then gradients of bounded solutions may blow up in finite time and thus, solutions do not exist globally in general. At least for the case of Dirichlet or Neuman boundary conditions it is well known that a quadratic growth of the quotient f(t, x, u, p)/a(t, x, u, p) as $p \to \infty$ is an almost optimal condition for global a priori estimates

Date: September 23, 2021.

Key words and phrases. Dynamical boundary conditions, global solutions, Bernstein-Nagumo condition, doubling of variables.

¹Here and in the following, equations at the boundary points $x = \pm \ell$ as (1.2) involving \pm signs (or \mp signs) are to be understood as two equations with a + sign (or - sign) at $x = +\ell$ and a - sign (or + sign) at $x = -\ell$. The sign convention is chosen such that $\pm u_x(\pm \ell)$ represents the "outward" normal derivative.

on the gradient of solutions to (1.1), (1.3). Actually, global *a priori* estimates can be derived under the Bernstein-Nagumo condition

$$\left|\frac{f(t,x,u,p)}{a(t,x,u,p)}\right| \le \psi(|p|) \tag{1.4}$$

with a positive function ψ obeying

$$\int_{-\infty}^{\infty} \frac{\rho \, \mathrm{d}\rho}{\psi(\rho)} = \infty \,, \tag{1.5}$$

in which case ψ can grow even faster than quadratic. This last conditions is sharp in the sense that its violation leads in several situation to a gradient blow up, see [7], [27], the introduction of [29], and the references given therein. Nevertheless, in the one-dimensional case with Dirichlet or Neuman boundary conditions, the Bernstein-Nagumo condition was improved in [29] by means of the doubling of variables technique to cope with right-hand sides $f = f_1 + f_2$, where only f_1 satisfies (1.4) and f_2 enjoys some monotonicity properties. The long time behavior of solutions also depends on the boundary conditions [27].

The aim of the present paper is to prove similar results as just described on existence and *a priori* estimates for the case of (1.1) subject to the nonlinear dynamical boundary condition (1.2) or (1.3). More precisely, we first prove with Theorem 2.2 an existence result for (1.1) subject to (1.2) in parabolic Hölder spaces (recalled in the Appendix 4) which includes a general global existence criterion that can be simplified when restricting to the gradient independent boundary condition (1.3). In Section 3 we then show that under the Bernstein-Nagumo condition one may derive L_{∞} -estimates on the gradient of bounded classical solutions to (1.1) subject to (1.2), see Theorem 3.1. A similar result is obtained in Theorem 3.2 for a weaker version of the Bernstein-Nagumo condition and in Theorem 3.3 for mixed boundary conditions, that is, when a dynamic boundary condition is imposed on one boundary point and e.g. a Dirichlet condition on the other. These gradient estimates can be used to derive Hölder estimates on the gradient (see Corollary 3.4) which, for the special case of (1.1) subject to (1.3), imply that bounded solutions exist globally in time, see Corollary 3.5. Finally, in Proposition 3.7 we provide conditions under which solutions to (1.1) subject to (1.3) are bounded and thus exist globally in time as stated in Corollary 3.8.

2. LOCAL SOLUTIONS

In this section we show a local existence result in the parabolic Hölder space $C^{1+\frac{\alpha}{2},2+\alpha}(\Omega_{\tau})$ (see the Appendix for a definition and properties) with $\tau > 0$ and $\Omega_{\tau} := (0,\tau) \times (-\ell,\ell)$ for equation (1.1) subject to the dynamical boundary condition (1.2). To do so, we first consider the corresponding linear problem

$$u_t - a(t, x)u_{xx} = f(t, x), \qquad (t, x) \in (0, T) \times (-\ell, \ell),$$
(2.1)

$$u_t \pm b(t, x)u_x = g(t, x), \qquad t \in (0, T), \quad x = \pm \ell,$$
(2.2)

$$u(0,x) = u^0(x), \qquad x \in [-\ell,\ell]$$
 (2.3)

and recall the following result:

Proposition 2.1 ([5, Theorem 1.1]). Let $\alpha \in (0,1)$. Let $a \in C^{\frac{\alpha}{2},\alpha}(\Omega_T)$ and $b(\cdot,\pm \ell) \in C^{\frac{1+\alpha}{2}}([0,T])$ satisfy

$$m^{-1} \le a(t,x) \le m, \quad (t,x) \in \Omega_T, \qquad m^{-1} \le b(t,\pm\ell) \le m, \quad t \in (0,T),$$
(2.4)

for some m > 0 and let $f \in C^{\frac{\alpha}{2},\alpha}(\Omega_T)$ and $g(\cdot, \pm \ell) \in C^{\frac{1+\alpha}{2}}([0,T])$. If $u^0 \in C^{2+\alpha}([-\ell,\ell])$ satisfies the compatibility condition

$$(au_{xx}^{0} + f)(0, \pm \ell) = (\mp bu_{x}^{0} + g)(0, \pm \ell),$$
(2.5)

then there is a unique solution $u \in C^{1+\frac{\alpha}{2},2+\alpha}(\Omega_T)$ to (2.1)–(2.3) such that $u_t(\cdot,\pm \ell) \in C^{\frac{1+\alpha}{2}}([0,T])$. Moreover, there is a constant

$$c_l := c_l \left(T, m, |a|_{\Omega_T}^{\left(\frac{\alpha}{2}, \alpha\right)}, |b(\cdot, \pm \ell)|_{[0,T]}^{\left(\frac{1+\alpha}{2}\right)} \right),$$

such that

$$|u|_{\Omega_T}^{(1+\frac{\alpha}{2},2+\alpha)} + |u_t(\cdot,\pm\ell)|_{[0,T]}^{(\frac{1+\alpha}{2})} \le c_l \left(|f|_{\Omega_T}^{(\frac{\alpha}{2},\alpha)} + |u^0|_{[-\ell,\ell]}^{(2+\alpha)} + |g(\cdot,\pm\ell)_{[0,T]}^{(\frac{1+\alpha}{2})} \right).$$
(2.6)

Proposition 2.1 is stated and proved in [5, Theorem 1.1] in dimensions higher than one, a detailed proof for the one-dimensional case may also be found in [19]. A more general situation is considered in [8] in an L_p -setting. The constant c_l is increasing with respect to T. Also note that the compatibility condition for the initial value is natural as we consider classical solutions.

Based on the previous result we next prove an existence result for classical solutions to the nonlinear problem (1.1), (1.2) by means of the contraction mapping principle. This naturally yields a global existence criterion that we shall exploit further in the subsequent section. Let us point out that other methods and other solution spaces are possible as well, of course, see e.g. [7] (and the references therein) where a in (1.1) is gradient-independent.

Theorem 2.2. Let $\alpha \in (0,1)$ and

$$a, f \in C^{2}([0,T] \times [-\ell,\ell] \times \mathbb{R} \times \mathbb{R}), \qquad b(\cdot, \pm \ell, \cdot, \cdot), g(\cdot, \pm \ell, \cdot, \cdot) \in C^{2}([0,T] \times \mathbb{R} \times \mathbb{R})$$
(2.7)

satisfy the uniform parabolicity condition

$$a(t, x, z, p) > 0$$
, $\partial_4 b(t, \pm l, z, p) p + b(t, \pm l, z, p) \mp \partial_4 g(t, \pm l, z, p) > 0$ (2.8)

for $(t, x, z, p) \in [0, T] \times [-\ell, \ell] \times \mathbb{R} \times \mathbb{R}$. Let $u^0 \in C^{2+\alpha}([-\ell, \ell])$ satisfy the compatibility condition

$$\mp b(0, x, u^0, u^0_x)u^0_x + g(0, x, u^0, u^0_x) = a(0, x, u^0, u^0_x)u^0_{xx} + f(0, x, u^0, u^0_x) \quad at \ x = \pm \ell.$$
(2.9)

Then the problem (1.1), (1.2) subject to $u(0, \cdot) = u^0$ admits a unique solution u on a maximal interval of existence $[0, \tau_{\infty}) \subset [0, T]$ such that $u \in \mathcal{C}^{1+\frac{\alpha}{2}, 2+\alpha}(\Omega_{\tau})$ for any $\tau < \tau_{\infty}$ and either

$$\lim_{\tau \nearrow \tau_{\infty}} |u|_{\Omega_{\tau}}^{(1+\frac{\alpha}{2},2+\alpha)} = \infty$$
(2.10)

or $u \in \mathcal{C}^{1+\frac{\alpha}{2},2+\alpha}(\Omega_T)$.

Moreover, in the special case (1.3) *when* b *and* g *are gradient-independent and* $b(\cdot, \pm \ell, \cdot) > 0$ *, then either*

$$\lim_{\tau \nearrow \tau_{\infty}} \left(|u|_{\Omega_{\tau}} + |u_{x}|_{\Omega_{\tau}} + \langle u_{x} \rangle_{\Omega_{\tau}}^{\left(\frac{\alpha}{2}, \alpha\right)} \right) = \infty$$

or $u \in \mathcal{C}^{1+\frac{\alpha}{2},2+\alpha}(\Omega_T)$.

Proof. To prove existence of a local solution we formulate the problem as a fixed point equation to which we may apply the contraction mapping principle. To do so, we set

$$\begin{split} h(t, \pm \ell, z, p) &:= b(t, \pm \ell, z, p) p \mp g(t, \pm \ell, z, p) , & (t, z, p) \in [0, T] \times \mathbb{R} \times \mathbb{R} , \\ h_0(t, \pm \ell) &:= \partial_4 h(t, \pm \ell, u^0(\pm \ell), u^0_x(\pm \ell)) , & t \in [0, T] , \\ a_0(t, x) &:= a(t, x, u^0(x), u^0_x(x)) , & (t, x) \in [0, T] \times [-\ell, \ell] . \end{split}$$

Observe that $h_0(t, \pm \ell) > 0$ thanks to (2.8). Given $\tau \in (0, T]$ define the Banach spaces

$$\mathbb{E}_{0}(\tau) := \left\{ (v, w, z) \, ; \, v \in \mathcal{C}^{\frac{\alpha}{2}, \alpha}(\Omega_{\tau}), w(\cdot, \pm \ell) \in C^{\frac{1+\alpha}{2}}([0, \tau]), \, z \in C^{2+\alpha}([-\ell, \ell]), \\ (v + a_{0}z_{xx})(0, \pm \ell) = \left(w \mp h_{0}z_{x}\right)(0, \pm \ell) \right\}$$

and

$$\mathbb{E}_1(\tau) := \left\{ u \in \mathcal{C}^{1+\frac{\alpha}{2},2+\alpha}(\Omega_\tau); \ u_t(\cdot,\pm\ell) \in C^{\frac{1+\alpha}{2}}([0,\tau]) \right\},\$$

equipped with the norms

$$\|(v,w,z)\|_{\mathbb{E}_0(\tau)} := |v|_{\Omega_\tau}^{(\frac{\alpha}{2},\alpha)} + |w(\cdot,\ell)|_{[0,\tau]}^{(\frac{1+\alpha}{2})} + |w(\cdot,-\ell)|_{[0,\tau]}^{(\frac{1+\alpha}{2})} + |z|_{[-\ell,\ell]}^{(2+\alpha)}$$

and, respectively,

$$\|u\|_{\mathbb{E}_{1}(\tau)} := |u|_{\Omega_{\tau}}^{(1+\frac{\alpha}{2},2+\alpha)} + |u_{t}(\cdot,\ell)|_{[0,\tau]}^{(\frac{1+\alpha}{2})} + |u_{t}(\cdot,-\ell)|_{[0,\tau]}^{(\frac{1+\alpha}{2})}$$

Then, introducing the operator L_0 by setting

$$(L_0 u)(t,x) := \left(u_t(t,x) - a_0(t,x) u_{xx}(t,x), u_t(t,\pm \ell) \pm h_0(t,\pm \ell) u_x(t,\pm \ell), u(0,x) \right), \quad (t,x) \in \overline{\Omega}_{\tau} ,$$

for $u \in \mathbb{E}_1(\tau)$, we derive from (2.8) and Proposition 2.1 that $L_0 \in \mathcal{L}(\mathbb{E}_1(\tau), \mathbb{E}_0(\tau))$ is a topological isomorphism for any $\tau \in (0, T]$ with

$$\sup_{\tau \in (0,T]} \|L_0^{-1}\|_{\mathcal{L}(\mathbb{E}_0(\tau),\mathbb{E}_1(\tau))} < \infty.$$
(2.11)

In order to solve problem (1.1), (1.2), we shall seek for a fixed point of the mapping Φ given by

$$\Phi(u) := L_0^{-1} (F(u), H(u), u^0),$$

where

$$F(u) := (a(\cdot, \cdot, u, u_x) - a_0)u_{xx} + f(\cdot, \cdot, u, u_x),$$

$$H(u)(\cdot, \pm \ell) := \pm h_0(\cdot, \pm \ell)u_x(\cdot, \pm \ell) \mp h(\cdot, \pm \ell, u(\cdot, \pm \ell), u_x(\cdot, \pm \ell)),$$

for $u \in \mathbb{E}_1(\tau)$. For this we show that Φ is a contraction on the set

$$\mathcal{V}(\tau) := \{ u \in \mathbb{E}_1(\tau); \ u(0) = u^0, \ \|u\|_{\mathbb{E}_1(\tau)} \le M \}$$

when τ is chosen sufficiently small, where

$$M > 2 \|\Phi(u_0)\|_{\mathbb{E}_1(T)} + \|u_0\|_{\mathbb{E}_1(T)} .$$
(2.12)

First observe that $(F(u), H(u), u(0, \cdot)) \in \mathbb{E}_0(\tau)$ for $u \in \mathcal{V}(\tau)$ due to (2.9) and $u(0, \cdot) = u^0$. We then claim that, given $\varepsilon > 0$, there is $\tau^* \in (0, T]$ such that

$$\left\| \left(F(u), H(u), u^0 \right) - \left(F(v), H(v), u^0 \right) \right\|_{\mathbb{E}_0(\tau)} \le \varepsilon \|u - v\|_{\mathbb{E}_1(\tau)}$$
(2.13)

for all $u, v \in \mathcal{V}(\tau)$ and all $0 < \tau < \tau^*$. Indeed, for $u, v \in \mathcal{V}(\tau)$ we have

$$H(u) - H(v) = \int_0^1 \left[\pm h_0 \mp \partial_4 h(\cdot, \cdot, u, \sigma u_x + (1 - \sigma) v_x) \right] (u_x - v_x) \, \mathrm{d}\sigma$$

and thus

$$\begin{split} |(H(u) - H(v))(\cdot, \pm \ell)|_{[0,\tau]}^{(\frac{1+\alpha}{2})} &\leq \sup_{\sigma \in (0,1)} |(h_0 - \partial_4 h(\cdot, \cdot, u, \sigma u_x + (1-\sigma)v_x)(\cdot, \pm \ell)|_{[0,\tau]}^{(\frac{1+\alpha}{2})} \\ &\times |(u_x - v_x)(\cdot, \pm \ell)|_{[0,\tau]} \\ &+ \sup_{\sigma \in (0,1)} |(h_0 - \partial_4 h(\cdot, \cdot, u, \sigma u_x + (1-\sigma)v_x))(\cdot, \pm \ell)|_{[0,\tau]} \\ &\times |(u_x - v_x)(\cdot, \pm \ell)|_{[0,\tau]}^{(\frac{1+\alpha}{2})}. \end{split}$$

Note that

$$\begin{aligned} \left| u_x(\cdot, \pm \ell) - v_x(\cdot, \pm \ell) \right|_{[0,\tau]} &\leq \tau^{\frac{1+\alpha}{2}} \left| u_x(\cdot, \pm \ell) - v_x(\cdot, \pm \ell) \right|_{[0,\tau]}^{(\frac{1+\alpha}{2})} \leq \tau^{\frac{1+\alpha}{2}} \|u - v\|_{\mathbb{E}_1(\tau)} \end{aligned}$$

and, since $h_0 - \partial_4 h(\cdot, \cdot, u, \sigma u_x + (1 - \sigma) v_x) = 0$ at $(t, x) = (0, \pm \ell)$, we may estimate
 $\left| \left(h_0 - \partial_4 h(\cdot, \cdot, u, \sigma u_x + (1 - \sigma) v_x) \right)(\cdot, \pm \ell) \right|_{[0,\tau]} \end{aligned}$

$$\begin{aligned} h_0 &- \partial_4 n(\cdot, \cdot, u, \sigma u_x + (1 - \sigma) v_x))(\cdot, \pm \ell)|_{[0,\tau]} \\ &\leq \tau^{\frac{1+\alpha}{2}} \left| \left(h_0 - \partial_4 h(\cdot, \cdot, u, \sigma u_x + (1 - \sigma) v_x) \right)(\cdot, \pm \ell) \right|_{[0,\tau]}^{(\frac{1+\alpha}{2})} \\ &\leq c_1(M) \, \tau^{\frac{1+\alpha}{2}} \end{aligned}$$

with a constant $c_1(M)$ depending on second order derivatives of h (here and in the following we ignore possible dependence of constants on the number α). Consequently,

$$\left| \left(H(u) - H(v) \right)(\cdot, \pm \ell) \right|_{[0,\tau]}^{\left(\frac{1+\alpha}{2}\right)} \le c(M) \, \tau^{\frac{1+\alpha}{2}} \, \|u - v\|_{\mathbb{E}_1(\tau)} \,. \tag{2.14}$$

Writing

$$F(u) - F(v) = f(\cdot, \cdot, u, u_x) - f(\cdot, \cdot, v, v_x) + (a(\cdot, \cdot, u, u_x) - a_0)(u_{xx} - v_{xx}) + (a(\cdot, \cdot, u, u_x) - a(\cdot, \cdot, v, v_x))v_{xx}$$

and observing that $a(\cdot, \cdot, u, u_x) - a_0$, $a(\cdot, \cdot, u, u_x) - a(\cdot, \cdot, v, v_x)$, and $f(\cdot, \cdot, u, u_x) - f(\cdot, \cdot, v, v_x)$ are of lower order with $a(\cdot, \cdot, u, u_x) - a_0 = 0$ at t = 0, standard interpolation inequalities for parabolic Hölder spaces (see [22] and Proposition 4.1) together with similar arguments as above imply that

$$\begin{aligned} \left| \left(a(\cdot, \cdot, u, u_x) - a_0 \right) \left(u_{xx} - v_{xx} \right) \right|_{\Omega_{\tau}}^{\left(\frac{\alpha}{2}, \alpha\right)} \\ & \leq \tau^{1/2} \sup_{x \in (-l,l)} \left| a(\cdot, x, u, u_x) - a_0(\cdot, x) \right|_{[0,\tau]}^{\left(\frac{1+\alpha}{2}\right)} \|u - v\|_{\mathbb{E}_1(\tau)} \\ & + c(l) \sup_{t \in (0,\tau)} \|a(t, \cdot, u, u_x) - a_0(t, \cdot)\|_{C^1([-l,l])} \|u - v\|_{\mathbb{E}_1(\tau)} \\ & \leq c(M, l) \left(\tau^{1/2} + \tau^{\alpha/2} \right) \|u - v\|_{\mathbb{E}_1(\tau)} \end{aligned}$$

as well as

$$\left| \left(a(\cdot, \cdot, u, u_x) - a(\cdot, \cdot, v, v_x) \right) v_{xx} \right|_{\Omega_{\tau}}^{\left(\frac{\alpha}{2}, \alpha\right)} \le c(M) \left(\tau^{\frac{1}{2+\alpha}} + \tau^{\frac{1+\alpha}{2}} \right) \|u - v\|_{\mathbb{E}_1(\tau)}$$

and

$$\left|f(\cdot,\cdot,u,u_x) - f(\cdot,\cdot,v,v_x)\right|_{\Omega_{\tau}}^{\left(\frac{\alpha}{2},\alpha\right)} \le c(M)\left(\tau^{\frac{1}{2+\alpha}} + \tau^{\frac{1+\alpha}{2}}\right) \|u - v\|_{\mathbb{E}_1(\tau)}$$

with constants depending on derivatives of a and f up to second order. These estimates combined with (2.14) yield (2.13).

Next note that (2.11) and (2.13) imply

$$\|\Phi(u) - \Phi(v)\|_{\mathbb{E}_{1}(\tau)} \leq \frac{1}{2} \|u - v\|_{\mathbb{E}_{1}(\tau)}, \qquad u, v \in \mathcal{V}(\tau),$$
(2.15)

for $\tau > 0$ small enough. Also note that (2.15) and (2.12) entail that

$$\|\Phi(u)\|_{\mathbb{E}_{1}(\tau)} \leq \|\Phi(u_{0})\|_{\mathbb{E}_{1}(T)} + \frac{1}{2} \left(\|u\|_{\mathbb{E}_{1}(\tau)} + \|u_{0}\|_{\mathbb{E}_{1}(T)}\right) < \frac{M}{2} + \frac{M}{2} = M$$

for $u \in \mathcal{V}(\tau)$, that is, Φ maps $\mathcal{V}(\tau)$ into itself. Consequently, the contraction mapping principle yields a unique fixed point $u \in \mathcal{V}(\tau)$ for the mapping Φ which solves problem (1.1), (1.2) subject to $u(0, \cdot) = u^0$. Clearly, this local solution can be extended to a solution u on a maximal interval of existence $[0, \tau_{\infty}) \subset [0, T]$ such that $u \in \mathcal{C}^{1+\frac{\alpha}{2}, 2+\alpha}(\Omega_{\tau})$ for any $\tau < \tau_{\infty}$ and either

$$\lim_{\tau \nearrow \tau_{\infty}} |u|_{\Omega_{\tau}}^{(1+\frac{\alpha}{2},2+\alpha)} = \infty$$

or $u \in \mathcal{C}^{1+\frac{\alpha}{2},2+\alpha}(\Omega_T)$.

Finally, consider the special case (1.3) when b and g are gradient-independent with $b(\cdot, \pm \ell, \cdot) > 0$. Suppose that there are a constant R > 0 and a sequence $\tau_i \nearrow \tau_{\infty}$ such that

$$\sup_{i\in\mathbb{N}} \left(|u|_{\Omega_{\tau_i}} + |u_x|_{\Omega_{\tau_i}} + \langle u_x \rangle_{\Omega_{\tau_i}}^{\left(\frac{\alpha}{2},\alpha\right)} \right) \le R.$$

This clearly implies that in fact

$$\sup_{\tau < \tau_{\infty}} \left(|u|_{\Omega_{\tau}} + |u_x|_{\Omega_{\tau}} + \langle u_x \rangle_{\Omega_{\tau}}^{\left(\frac{\alpha}{2}, \alpha\right)} \right) \le R.$$
(2.16)

It then follows from equation (1.3) that

$$\sup_{\tau < \tau_{\infty}} |u_t(\cdot, \pm \ell)|_{[0,\tau]} \le c(R) \,. \tag{2.17}$$

Using this and the fact that

$$|u(t,y) - u(s,y)| \le \int_{-\ell}^{y} |u_x(t,z) - u_x(s,z)| \, \mathrm{d}z + |u(t,-\ell) - u(s,-\ell)|$$

for $(t, s, y) \in [0, \tau_{\infty})^2 \times [-\ell, \ell]$, we find

$$\sup_{\tau < \tau_{\infty}} \langle u \rangle_{\Omega_{\tau}}^{\left(\frac{\alpha}{2}, \alpha\right)} \le c(R, \ell) \left(1 + T^{1 - \alpha/2} \right).$$
(2.18)

Now, (2.16), (2.17), and (2.18) imply

$$\max\left\{ |\widetilde{a}|_{\Omega_{\tau}}^{\left(\frac{\alpha}{2},\alpha\right)}, |\widetilde{f}|_{\Omega_{\tau}}^{\left(\frac{\alpha}{2},\alpha\right)}, |\widetilde{b}(\cdot,\pm\ell)|_{[0,\tau]}^{\left(\frac{1+\alpha}{2}\right)}, |\widetilde{g}(\cdot,\pm\ell)|_{[0,\tau]}^{\left(\frac{1+\alpha}{2}\right)} \right\} \le c(R,\ell,T)$$
(2.19)

for each $\tau < \tau_{\infty}$, where $\widetilde{a}(t,x) := a(t,x,u(t,x),u_x(t,x))$ and $\widetilde{f}, \widetilde{b}, \widetilde{g}$ are defined analogously.

As a and b are strictly positive on $[0, T] \times [-\ell, \ell] \times [-R, R]^2$, respectively, on $[0, T] \times \{\pm \ell\} \times [-R, R]$, Proposition 2.1 ensures the existence of a constant $c = c(R, \ell, T, |u^0|_{\Omega}^{(2+\alpha)})$ such that

$$|u|_{\Omega_{\tau}}^{(1+\frac{\alpha}{2},2+\alpha)} \le c, \quad \tau < \tau_{\infty},$$

hence $u \in \mathcal{C}^{1+\frac{\alpha}{2},2+\alpha}(\Omega_T)$ by our previous findings. This proves the theorem.

Actually, a closer look at the proof shows that the global existence criterion (2.10) for the general (i.e. gradient-dependent) case can be weakened. Indeed, it suffices to control

$$|u|_{\Omega_{\tau}}^{(\frac{1+\alpha}{2},\alpha)} + \left|u(\cdot,\pm\ell)\right|_{[0,\tau]}^{(\frac{1+\alpha}{2})}$$

uniformly in $\tau < \tau_{\infty}$ in this case. Moreover, since the maximal regularity result stated in Theorem 2.1 holds also true in higher space dimensions (see [5, Theorem 1.1]), one easily verifies that Theorem 2.2 is true in this case, too.

3. A PRIORI ESTIMATES AND GLOBAL SOLUTIONS

Theorem 2.2 reduces the question of global existence to (1.1) with gradient-independent boundary conditions (1.3) to finding *a priori* estimates for

$$|u|_{\Omega_{\tau}} + |u_x|_{\Omega_{\tau}} + \langle u_x \rangle_{\Omega_{\tau}}^{(\frac{\alpha}{2},\alpha)}$$
(3.1)

independent of $\tau < \tau_{\infty}$ for the solution u on the maximal interval of existence $[0, \tau_{\infty}) \subset [0, T]$. The aim of this section is to further reduce this condition. More precisely, we shall show L_{∞} - and Hölderbounds on the gradient u_x solely based on bounds on $|u|_{\Omega_{\tau}} = \sup_{(t,x)\in\Omega_{\tau}} |u(t,x)|$, that is, we show that a bound on the first term of (3.1) implies bounds on the second and third terms. It is worthwhile to point out that we can obtain such estimates even for equation (1.1) combined with the gradient-dependent boundary condition (1.2). While Hölder-estimates on the gradient, i.e. estimates on $\langle u_x \rangle_{\Omega_{\tau}}^{(\frac{\alpha}{2},\alpha)}$, are rather easy to obtain from the existing theory (see Subsection 3.2), more effort has to be invested in Subsection 3.1 to derive estimates on $|u_x|_{\Omega_{\tau}}$.

3.1. A priori estimates on the gradient. We shall find a priori L_{∞} -bounds on the gradient u_x of a solution to (1.1), (1.2) presupposing a bound on its L_{∞} -norm. The proof is in the spirit of [29] and uses Kruzhkov's idea of introducing a new variable [20, 21]. In the next subsection we derive gradient estimates when imposing the Bernstein-Nagumo condition. In Subsection 3.1.2 we then indicate how to weaken this condition for a right hand side $f + f_1$ in (1.1) where only f satisfies the Bernstein-Nagumo condition and f_1 is allowed to be unbounded in the gradient variable. We also consider the case of mixed boundary conditions in Subsection 3.1.3.

In the following, a *classical solution* to (1.1), (1.2) [resp. to (1.1), (1.3)] on Ω_{τ} with $\tau \leq T$ is a function $u \in C^{1,2}(\Omega_{\tau}) \cap C(\overline{\Omega}_{\tau})$ with derivatives $u_t(\cdot, \pm \ell)$ and $u_x(\cdot, \pm \ell)$ being defined on $(0, \tau)$ and satisfying (1.1), (1.2) [resp. (1.1), (1.3)] pointwise in Ω_{τ} . Note that the existence and uniqueness of such a solution (with higher regularity) is guaranteed by Theorem 2.2 when imposing the assumptions stated there.

Throughout we assume continuity of the data, that is,

 $a, f \in C([0, T] \times [-\ell, \ell] \times \mathbb{R} \times \mathbb{R}), \qquad b(\cdot, \pm \ell, \cdot, \cdot), g(\cdot, \pm \ell, \cdot, \cdot) \in C([0, T] \times \mathbb{R} \times \mathbb{R})$

and the parabolicity condition a > 0 and $b(\cdot, \pm \ell, \cdot, \cdot) > 0$. Stronger assumptions will be indicated explicitly.

3.1.1. Gradient estimates under the Berstein-Nagumo condition. The next theorem provides a priori estimates on the gradient for any classical solution u to (1.1), (1.2) on Ω_{τ} with $\tau \leq T$ in dependence on its L_{∞} -bound.

Theorem 3.1. Let u^0 be Lipschitz continuous on $[-\ell, \ell]$, that is,

$$|u^{0}(x) - u^{0}(y)| \le K|x - y|, \quad x, y \in [-\ell, \ell].$$
(3.2)

Let M > 0 and suppose that there are $q_0 \ge K$ and $\psi \in C^1([0,\infty), [1,\infty))$ with

$$\int_{q_0}^{\infty} \frac{\rho \,\mathrm{d}\rho}{\psi(\rho)} > 2M,\tag{3.3}$$

such that

$$|f(t, x, z, p)| \le a(t, x, z, p)\psi(|p|), \quad (t, x, z, p) \in \overline{\Omega}_T \times [-M, M] \times \mathbb{R}$$
(3.4)

and

$$\pm g(t, \ell, z, \pm p) \le b(t, \ell, z, \pm p)p, \qquad \mp g(t, -\ell, z, \pm p) \le b(t, -\ell, z, \pm p)p, \tag{3.5}$$

for $(t, z) \in (0, T] \times [-M, M]$ and $p \ge q_0$. Then there exists a constant $M_1 := M_1(M, \psi, K)$ such that if u is any classical solution to (1.1), (1.2) on Ω_{τ} with $\tau \le T$ subject to $u(0, \cdot) = u^0$ satisfying $|u|_{\Omega_{\tau}} \le M$, then $|u_x|_{\Omega_{\tau}} \le M_1$.

Proof. We adapt the proof of [29] to the case of the dynamic boundary condition (1.2). Due to (3.3) there is $q_1 > q_0$ with

$$\int_{q_0}^{q_1} \frac{\rho \,\mathrm{d}\rho}{\psi(\rho)} = 2M. \tag{3.6}$$

Let κ be defined by

$$\kappa(\xi) := \int_{\xi}^{q_1} \frac{\mathrm{d}\rho}{\psi(\rho)}, \quad \xi \in [q_0, q_1].$$

Since ψ is positive, κ is strictly decreasing on $[q_0, q_1]$ with $\kappa(q_1) = 0$. Thus, its inverse $q := \kappa^{-1}$ is decreasing on the interval $[0, \kappa_0]$, where $\kappa_0 := \kappa(q_0)$, with $q(0) = q_1$ and $q(\kappa_0) = q_0$. Define

$$h(\xi) := \int_{q(\xi)}^{q_1} \frac{\rho \,\mathrm{d}\rho}{\psi(\rho)}, \quad \xi \in [0, \kappa_0].$$

Noticing that

$$h'(\xi) = q(\xi) \ge q_0, \qquad h''(\xi) = -\psi(q(\xi)),$$
(3.7)

we see that the function h solves

$$h''(\xi) + \psi(|h'(\xi)|) = 0, \quad \xi \in (0, \kappa_0), h(\kappa_0) = 2M, \quad h(0) = 0.$$
(3.8)

Since $h' \ge q_0 \ge K$ and h(0) = 0, there is for $x, y \in [-\ell, \ell]$ with $|x - y| \le \kappa_0$ some $\xi \in [0, |x - y|]$ such that

$$h(|x - y|) = h'(\xi)|x - y| \ge K|x - y|.$$

Hence, by (3.2), we have

$$|u^{0}(x) - u^{0}(y)| \le h(|x - y|), \quad x, y \in [-\ell, \ell], \ |x - y| \le \kappa_{0}.$$
(3.9)

Let us define the sets

$$S_{1} := \{(t, x, y) \in P : -\ell < x < -\ell + \kappa_{0}, y = -\ell, t \in (0, \tau]\},$$

$$S_{2} := \{(t, x, y) \in \overline{P} : x = \ell, \ell - \kappa_{0} < y < \ell, t \in (0, \tau]\},$$

$$S_{3} := \{(t, x, y) \in \overline{P} : x = y, t \in (0, \tau]\},$$

$$S_{4} := \{(t, x, y) \in \overline{P} : x - y = \kappa_{0}, t \in (0, \tau]\},$$
(3.10)

where

$$P := \{ (x, y) \in (-\ell, \ell)^2 : 0 < x - y < \kappa_0 \}, \qquad P_\tau := (0, \tau) \times P.$$

We put

$$S := \bigcup_{i=1}^{4} S_i, \qquad B := \{0\} \times \overline{P}, \qquad \Gamma := S \cup B$$

and define the auxiliary functions v, w, and \tilde{w} by

$$v(t, x, y) := u(t, x) - u(t, y),$$

$$w(t, x, y) := v(t, x, y) - h(x - y),$$

$$\widetilde{w}(t, x, y) := e^{-t}w(t, x, y)$$
(3.11)

for $(t, x, y) \in \overline{P}_{\tau}$. Derivatives with respect to the second and third variable we denote as derivatives with respect to x and y, respectively. To keep notation as simple as possible we also use $u_x := u_y := \partial_2 u$. We shall show that \widetilde{w} does not attain a positive maximum in \overline{P}_{τ} . Let $(t, x, y) \in P_{\tau} \setminus \Gamma$ and note that (t, x) and (t, y) belong to Ω_{τ} . Thus

$$u_t(t,x) + a(t,x,u,u_x)u_{xx} + f(t,x,u,u_x) = -u_t(t,y) + a(t,y,u,u_y)u_{yy} + f(t,y,u,u_y) = 0.$$

by (1.1), where we use here and in the following the notation

$$a(t, y, u, u_y)u_{yy} := a(t, y, u(t, y), u_y(t, y))u_{yy}(t, y)$$

Subtracting the two equations and using the definition of v we obtain

$$-v_t(t, x, y) + a(t, x, u, v_x)v_{xx} + f(t, x, u, v_x) + a(t, y, u, -v_y)v_{xx} - f(t, y, u, -v_y) = 0.$$
(3.12)

Recalling from (3.4) that

$$f(t, x, u, v_x) \le a(t, x, u, v_x)\psi(|v_x|), \qquad -f(t, y, u, -v_y) \le a(t, y, u, -v_y)\psi(|v_y|)$$
(3.13)

and using the notation

$$A(t,x) := a(t,x,u,v_x), \qquad A(t,y) := a(t,y,u,-v_y).$$

we derive from (3.12) and (3.13) the inequality

$$-v_t(t, x, y) + A(t, x)(v_{xx} + \psi(|v_x|)) + A(t, y)(v_{yy} + \psi(|v_y|)) \ge 0$$

Note that (3.8) implies

$$A(t,x)(h''(x-y) + \psi(|h'(x-y)|)) + A(t,y)(h''(x-y) + \psi(|h'(x-y)|)) = 0.$$

Subtracting the previous (in-)equalities yields

$$-w_t(t, x, y) + A(t, x)w_{xx}(t, x, y) + A(t, y)w_{yy}(t, x, y) + r(t, x, y) \ge 0,$$

where

$$r(t,x,y) := \psi(|v_x(t,x,y)|) - \psi(|h'(x-y)|) + \psi(|v_y(t,x,y)|) - \psi(|h'(x-y)|),$$

that is,

$$-\widetilde{w}_t - \widetilde{w} + A(t, x)\widetilde{w}_{xx} + A(t, y)\widetilde{w}_{yy} + re^{-t} \ge 0, \quad (t, x, y) \in \overline{P}_\tau \setminus \Gamma.$$
(3.14)

Assume for contradiction that the function \widetilde{w} attains its positive maximum at $(t_0, x_0, y_0) \in \overline{P}_{\tau} \setminus \Gamma$. At this point of maximum there holds

$$\widetilde{w}(t_0, x_0, y_0) > 0, \quad \widetilde{w}_t(t_0, x_0, y_0) \ge 0, \quad \widetilde{w}_{xx}(t_0, x_0, y_0) \le 0, \quad \widetilde{w}_{yy}(t_0, x_0, y_0) \le 0.$$
 (3.15)

Moreover,

$$\widetilde{w}_x(t_0, x_0, y_0) = \widetilde{w}_y(t_0, x_0, y_0) = 0$$

from which, by definition of \widetilde{w} , we get

$$e^{-t}(v_x(t_0, x_0, y_0) - h'(x_0 - y_0)) = e^{-t}(v_y(t_0, x_0, y_0) + h'(x_0 - y_0)) = 0.$$

Thus

$$v_x(t_0, x_0, y_0) = -v_y(t_0, x_0, y_0) = h'(x_0 - y_0)$$
 and $r(t_0, x_0, y_0) = 0.$ (3.16)

Since both $A(t_0, x_0)$ and $A(t_0, y_0)$ are non-negative it follows from (3.15) and (3.16) that

$$-\widetilde{w}_t - \widetilde{w} + A(t, x)\widetilde{w}_{xx} + A(t, y)\widetilde{w}_{yy} + re^{-t} < 0 \quad \text{at} \ (t, x, y) = (t_0, x_0, y_0)$$
(3.17)

which contradicts (3.14). Therefore, \widetilde{w} does not attain a positive maximum in $\overline{P}_T \setminus \Gamma$.

We next show that \tilde{w} does also not attain a positive maximum on Γ for which we distinguish the two cases $\kappa_0 < 2\ell$ and $\kappa_0 \ge 2\ell$.

Case 1: $\kappa_0 < 2\ell$. In this case we have $-\ell < y < \ell$ for $(t, \ell, y) \in S_2$. Thus, the function u satisfies equations (1.1) and (1.2) at the points (t, y) and (t, ℓ) , that is,

$$\begin{aligned} -u_t(t,y) + a(t,y,u,u_y)u_{yy} + f(t,y,u,u_y) &= 0, \\ -u_t(t,\ell) - b(t,\ell,u,u_x)u_x + g(t,\ell,u,u_x) &= 0. \end{aligned}$$

Subtracting the two equations and recalling (3.4) and the definitions of v and A(t, y), we derive the inequality

$$-v_t(t,\ell,y) + A(t,y)(v_{yy} + \psi(|v_y|)) + r_1(t,\ell,y) \ge 0,$$
(3.18)

where

 $r_1(t, \ell, y) := -b(t, \ell, u, v_x)v_x + g(t, \ell, u, v_x).$

Since $(t, \ell, y) \in S_2 P_T$ and hence $0 \le \ell - y \le \kappa_0$, it follows from (3.8) that

$$A(t,y)(h''(\ell - y) + \psi(|h'(\ell - y)|)) = 0.$$

Subtracting this from (3.18) and using the definition of \tilde{w} , we derive

$$-\widetilde{w}_t - \widetilde{w} + A(t, y)\widetilde{w}_{yy} + r_2 e^{-t} + r_1 e^{-t} \ge 0, \quad (t, \ell, y) \in S_2,$$
(3.19)

where

$$r_2(t,\ell,y) := A(t,y)(\psi(|v_y(t,\ell,y)|) - \psi(|h'(\ell-y)|)).$$

Assume now for contradiction that \tilde{w} attains its maximum at some point $(t_0, \ell, y_0) \in S_2$. At this maximum point there holds

$$\widetilde{w}(t_0,\ell,y_0) > 0, \qquad \widetilde{w}_t(t_0,\ell,y_0) \ge 0, \qquad \widetilde{w}_{yy}(t_0,\ell,y_0) \le 0$$

Furthermore, (t_0, ℓ, y_0) is an inner point of S_2 , hence $\widetilde{w}_y(t_0, \ell, y_0) = 0$. Since, by definition of \widetilde{w} ,

$$e^{-t_0}(v_y(t_0,\ell,y_0) + h'(\ell - y_0)) = 0$$

we have $r_2(t_0, \ell, y_0) = 0$. Also note that

$$\widetilde{w}_x(t_0, \ell, y_0) = e^{-t_0}(v_x(t_0, \ell, y_0) - h'(\ell - y_0)) \ge 0,$$

that is, using (3.7),

$$v_x(t_0, \ell, y_0) \ge h'(\ell - y_0) \ge q_0 > 0 \tag{3.20}$$

and consequently, due to (3.5),

$$r_1(t_0, \ell, y_0) = -b(t_0, \ell, u, v_x)v_x + g(t_0, \ell, u, v_x) \le 0.$$

In summary, we have

$$-\widetilde{w}_t - \widetilde{w} + A(t, y)\widetilde{w}_{yy} + r_2 e^{-t_0} + r_1 e^{-t_0} < 0 \quad \text{at} \ (t, x, y) = (t_0, \ell, y_0), \tag{3.21}$$

in contradiction to (3.19). Therefore, \widetilde{w} does not attain a positive maximum in S_2 . In the same way we consider \widetilde{w} on S_1 . Given $(t, x, -\ell) \in S_1$, we derive from

$$-u_t(t,x) + a(t,x,u,u_x)u_{xx} + f(t,x,u,u_x) = -u_t(t,-\ell) + b(t,-\ell,u,u_x)u_x + g(t,-\ell,u,u_x) = 0$$

by subtraction and by using (3.4) the inequality

$$-v_t(t, x, -\ell) + A(t, x)(v_{xx} + \psi(|v_x|)) + r_3(t, x, -\ell) \ge 0$$

where

$$r_3(t, x, -\ell) := b(t, -\ell, u, -v_y)v_y - g(t, -\ell, u, -v_y)v_y - g(t$$

Since, due to (3.8),

$$A(t,x)(h''(x+\ell) + \psi(|h'(x+\ell)|)) = 0$$

we further obtain

$$-\widetilde{w}_t - \widetilde{w} + A(t, x)\widetilde{w}_{xx} + r_4 e^{-t} + r_3 e^{-t} \ge 0, \quad (t, x, -\ell) \in S_1$$
(3.22)

with

$$r_4(t, x, -\ell) := A(t, x)(\psi(|v_x(t, x, -\ell)|) - \psi(|h'(x+\ell)|)).$$

Assume for contradiction that \widetilde{w} attains its positive maximum at a point $(t_0, x_0, -\ell) \in S_1$ with $x_0 \neq \ell$. From $\widetilde{w}_x(t_0, x_0, -\ell) = 0$ we deduce $r_4(t_0, \ell, y_0) = 0$. Furthermore, since

$$0 \le -\widetilde{w}_y(t_0, x_0, -\ell) = e^{-t_0}(-v_y(t_0, x_0, -\ell) - h'(x_0 + \ell))$$

it follows from (3.7) that

$$-v_y(t_0, x_0, -\ell) \ge h'(x_0 + \ell) \ge q_0 > 0,$$

and thus $r_3(t_0, x_0, -\ell) \leq 0$ due to (3.5). Consequently,

$$-\widetilde{w}_t - \widetilde{w} + A(t, x)\widetilde{w}_{xx} + r_4 e^{-t} + r_3 e^{-t} < 0 \quad \text{at} \ (t, x, y) = (t_0, x_0, -\ell)$$

in contradiction to(3.22). Thus, \tilde{w} does not attain a positive maximum on S_1 .

Next, we consider \widetilde{w} on S_3 and S_4 . Since x = y for $(t, x, y) \in S_3$ and h(0) = 0, we clearly have $\widetilde{w} = 0$ on S_3 so that \widetilde{w} does not attain a positive maximum on S_3 . Given $(t, x, y) \in S_4$ we have $x - y = \kappa_0$. Now (3.8) implies $h(\kappa_0) = 2M = 2|u(t, x)|_{\Omega_\tau}$, hence

$$\widetilde{w}(t, x, y) = u(t, x) - u(t, y) - h(\kappa_0) \le 0,$$

and \widetilde{w} thus does not attain a positive maximum on S_4 either.

Finally, since for $(t, x, y) \in B$ we have t = 0, (3.9) yields

$$\widetilde{w}(t, x, y) = (u^0(x) - u^0(y) - h(x - y)) \le 0,$$

and \widetilde{w} does not attain a positive maximum on Γ if $\kappa_0 < 2\ell$.

Case 2: $\kappa_0 \ge 2\ell$. In this case, the set S_4 is empty while the sets S_1 and S_2 also contain the set $\{(t, l, -\ell) : t \in (0, \tau]\}$. We have to show therefore that \widetilde{w} does not attain a positive maximum on this set. For, note that at the points (t, ℓ) and $(t, -\ell)$ the function u satisfies according to (1.2)

$$-u_t(t,\ell) - b(t,\ell,u,u_x)u_x + g(t,\ell,u,u_x) = -u_t(t,-\ell) + b(t,-\ell,u,u_x)u_x + g(t,-\ell,u,u_x) = 0$$

from which we get

$$-w_t(t,\ell,-\ell) - b(t,\ell,u,u_x)u_x + g(t,\ell,u,u_x) - b(t,-\ell,u,u_x)u_x - g(t,-\ell,u,u_x) = 0.$$
(3.23)

Assume for contradiction that \widetilde{w} attains a positive maximum at some point $(t_0, \ell, -\ell)$ with $t_0 \in (0, \tau]$. Then

$$\widetilde{w}_x(t_0, \ell, -\ell) = e^{-t}(u_x(t_0, \ell) - h'(2\ell)) \ge 0, -\widetilde{w}_y(t_0, \ell, -\ell) = e^{-t}(u_y(t_0, -\ell) - h'(2\ell)) \ge 0$$

and (3.8) implies

$$u_x(t_0, \ell) \ge h'(2\ell) \ge q_0 > 0,$$

 $u_y(t_0, -\ell) \ge h'(2\ell) \ge q_0 > 0.$

From this and (3.5) it follows that

$$-b(t,\ell,u,u_x)u_x + g(t,\ell,u,u_x) \le 0, \qquad -b(t,-\ell,u,u_x)u_x - g(t,-\ell,u,u_x) \le 0$$

and hence, from (3.23), that

$$-w_t(t_0, \ell, -\ell) \ge 0. \tag{3.24}$$

On the other hand, since \widetilde{w} attains on $\{(t,\ell,-\ell):\ t\in (0,\tau]\}$ a positive maximum, we have

 $\widetilde{w}(t_0, \ell, -\ell) > 0, \quad \widetilde{w}_t(t_0, \ell, -\ell) \ge 0$

and further

$$-w_t(t_0, \ell, -\ell) = -e^t(\widetilde{w}(t_0, \ell, -\ell) + \widetilde{w}_t(t_0, \ell, -\ell)) < 0$$

contradicting (3.24). So, \widetilde{w} does also not attain a positive maximum on Γ if $\kappa_0 \ge 2\ell$.

 \widetilde{w}

Consequently, we have shown that \widetilde{w} does not attain a positive maximum in \overline{P}_{τ} and thus

$$(t, x, y) \le 0, \quad (t, x, y) \in \overline{P}_{\tau}.$$

In the same way one shows that the function \widetilde{w}_1 , defined by

$$\widetilde{w}_1(t,x,y) := u(t,y) - u(t,x) - h(x-y), \quad (t,x,y) \in \overline{P}_{\tau_2}$$

satisfies

$$\widetilde{w}_1(t,x,y) \le 0, \quad (t,x,y) \in \overline{P}_{\tau}.$$

Together we deduce

$$|u(t,x) - u(t,y)| \le h(x-y), \quad (t,x,y) \in \overline{P}_{\tau},$$

and putting

$$Q_{\tau} := \{ (t, x, y) : (t, y, x) \in P_{\tau} \},\$$

and using symmetry with respect to the variables we conclude

$$|u(t,x) - u(t,y)| \le h(|x-y|), \quad (t,x,y) \in \overline{Q}_{\tau} \cup \overline{P}_{\tau},$$

with

$$\overline{Q}_{\tau} \cup \overline{P}_{\tau} = \{(t, x, y) \in \mathbb{R}^3 : 0 \le |x - y| \le \kappa_0, |x| \le \ell, |y| \le \ell, t \in [0, \tau]\}.$$

Since h(0) = 0 we obtain

$$|u_x(t,x)| \le h'(0) = q_1, \quad (t,x) \in \Omega_{\tau}.$$
(3.25)

According to (3.6) the number q_1 depends only on K, M, q_0 , and ψ . This proves Theorem 3.1

It may be worthwhile to note that (3.6) yields more information on the gradient bound $M_1 = q_1$. Condition (3.5) is needed because of the dynamical boundary condition (1.2). A simple situation for which (3.5) holds is obtained by strengthening condition (3.3) to

$$\int \int \frac{\rho \,\mathrm{d}\rho}{\psi(\rho)} = \infty. \tag{3.26}$$

Indeed, in this case there exists for each $q_0 > 0$ some $q_1 > 0$ such that (3.6) is satisfied. Then (3.5) holds e.g. if $b \ge \delta > 0$ and g is bounded.

3.1.2. Gradient estimates under a weakened Bernstein-Nagumo condition. In [29] it was shown for classical (i.e. Dirichlet, Neuman, or Robin) boundary conditions that gradient estimates hold true for the more general situation that a term $f_1(t, x, u, u_x)$ is added on the right hand side of (1.1) which may arbitrarily increase in the gradient variable provided it satisfies a homogeneity condition (see (3.29) below). We similarly extend Theorem 3.1 for dynamical boundary conditions.

More precisely, we may consider

$$u_t - a(t, x, u, u_x)u_{xx} = f(t, x, u, u_x) + f_1(t, x, u, u_x), \quad (t, x) \in (0, T) \times (-\ell, \ell),$$
(3.27)

together with the boundary condition

$$u_t \pm b(t, x, u, u_x)u_x = g(t, x, u, u_x) + g_1(t, x, u, u_x), \quad t \in (0, T), \quad x = \pm \ell,$$
(3.28)

where we impose on f_1 and g_1 the following conditions:

$$f_1(t, y, z_1, \pm p) - f_1(t, x, z_2, \pm p) \ge 0$$
(3.29)

for $-\ell \le x \le y \le \ell$, $-M \le z_1 \le z_2 \le M$, $p \ge 0$,

$$f_1(t, x, z_1, \pm p_1) - g_1(t, \ell, z_2, \pm p_2) \ge 0$$
(3.30)

for $-\ell \leq x \leq \ell$, $-M \leq z_1 \leq z_2 \leq M$, $q_0 \leq p_1 \leq p_2$, and

$$g_1(t,\ell,z_1,-p_1) - f_1(t,x,z_2,-p_2) \ge 0, \qquad g_1(t,-\ell,z_1,p_1) - f_1(t,x,z_2,p_2) \ge 0$$
(3.31)

for $-\ell \le x \le \ell$, $-M \le z_1 \le z_2 \le M$, $q_0 \le p_2 \le p_1$. Then we can prove:

Theorem 3.2. Let M > 0 and suppose (3.2)-(3.5) and (3.29)–(3.31). Then there exists a constant $M_1 := M_1(q_0, M, \psi, K)$ such that if u is any classical solution to (3.27), (3.28) on Ω_{τ} with $\tau \leq T$ subject to $u(0, \cdot) = u^0$ satisfying $|u|_{\Omega_{\tau}} \leq M$, then $|u_x|_{\Omega_{\tau}} \leq M_1$.

Proof. Except for small changes the proof is the same as for Theorem 3.1. Indeed, using (3.27) instead of (1.1), inequality (3.14) has to be replaced by

$$-\widetilde{w}_t - \widetilde{w} + A(t,x)\widetilde{w}_{xx} + A(t,y)\widetilde{w}_{yy} + re^{-t} \ge f_1(t,y,u,-v_y) - f_1(t,x,u,v_x), \quad (t,x,y) \in \overline{P}_\tau \setminus \Gamma.$$
(3.32)

Assuming that \widetilde{w} attains a positive maximum at some point $(t_0, \ell, y_0) \in \overline{P}_{\tau} \setminus \Gamma$, we deduce (see (3.16))

$$v_x(t_0, \ell, y_0) = -v_y(t_0, \ell, y_0)$$

and so it follows from (3.32) and (3.29) that

$$-\widetilde{w}_t - \widetilde{w} + A(t, x)\widetilde{w}_{xx} + A(t, y)\widetilde{w}_{yy} + re^{-t} \ge 0 \quad \text{at} \ (t, x, y) = (t_0, x_0, y_0)$$

in contradiction to (3.17), the latter being derived exactly as in Theorem 3.1. Following the proof of Theorem 3.1 one then considers \tilde{w} on S_2 . Replacing f by $f + f_1$ and g by $g + g_1$, we obtain instead of (3.19) the inequality

$$-\widetilde{w}_t - \widetilde{w} + A(t, y)\widetilde{w}_{yy} + r_2 e^{-t} + r_1 e^{-t} \geq e^{-t} \left[f_1(t, y, u(t, y), -v_y(t, \ell, y)) - g_1(t, \ell, u(t, y), -v_y(t, \ell, y)) \right]$$

$$(3.33)$$

for $(t, \ell, y) \in S_2$. Assuming that \widetilde{w} attains a positive maximum at $(t_0, \ell, y_0) \in S_2$ then (see (3.20))

$$v_x(t_0, \ell, y_0) \ge -v_y(t_0, \ell, y_0) \ge q_0$$

and (3.30), (3.33) entail that

$$-\tilde{w}_t - \tilde{w} + A(t, y)\tilde{w}_{yy} + r_2 e^{-t} + r_1 e^{-t} \ge 0$$
 at $(t_0, \ell, y_0) \in S_2$

contradicting (3.21). The rest follows analogously to the proof of Theorem 3.1.

3.1.3. Gradient estimates for mixed boundary conditions. In [29, Lemma 1-Lemma 3] gradient estimates were derived for solutions to (1.1) subject to Dirichlet, Neuman, or Robin type boundary conditions. Inspection of the proofs therein and the one of Theorem 3.1 shows that such estimates can be obtained under the assumptions of Theorem 3.1 when a Dirichlet, Neuman, or Robin type boundary conditions is imposed on one boundary part and the dynamical boundary condition (1.2) on the other. We formulate the precise result in the exemplary constellation that u satisfies at $x = -\ell$ the dynamical boundary condition

$$u_t - b(t, -\ell, u, u_x)u_x = g(t, -\ell, u, u_x), \quad t \in (0, T),$$
(3.34)

and at $x = \ell$ the homogeneous Dirichlet condition

$$u(t,\ell) = 0, \quad t \in (0,T).$$
 (3.35)

All other cases can be considered as well.

Theorem 3.3. *Let* M > 0 *and suppose* (3.2)-(3.4) *and*

 $-g(t,-\ell,z,p) \leq b(t,-\ell,z,p)p, \qquad -g(t,\ell,z,-p) \leq b(t,\ell,z,-p)p,$

for $(t, z) \in (0, T] \times [-M, M]$ and $p \ge q_0$. Then there exists a constant $M_1 := M_1(q_0, M, \psi, K)$ such that if u is any classical solution to (1.1), (3.34), (3.35) on Ω_{τ} with $\tau \le T$ subject to $u(0, \cdot) = u^0$ satisfying $|u|_{\Omega_{\tau}} \le M$, then $|u_x|_{\Omega_{\tau}} \le M_1$.

Proof. The proof is the same as for Theorem 3.1 except when considering the behavior of \tilde{w} and \tilde{w}_1 on S_2 . That both functions do not attain a positive maximum on S_2 can be shown as in [29, Lemma 3].

Theorem 3.3 seems to be interesting in connection with the blow up result proven in [7, Proposition 2.14]. There the special case

$$u_t - u_{xx} = \psi(u_x), \qquad t > 0, \ 0 < x < \ell,$$

$$u(t,0) = 0, \qquad t \ge 0,$$

$$u_t(t,\ell) + u_x(t,\ell) = 0, \qquad t > 0,$$

$$u(0,x) = u^0(x), \qquad t \ge 0,$$

(3.36)

of (1.1), (3.34), (3.35) was considered and shown that if $\psi \in C^2(\mathbb{R}, (0, \infty))$ with

$$\liminf \psi(s) > 0$$

violates (3.26), then there is $\ell > 0$ such that for each $u^0 \in C^2([0, \ell])$ with $u^0(0) = 0$, the corresponding classical solution to (3.36) evolves a gradient singularity in finite time, even if the solution itself stays bounded (see also [7, Remark 2.15]).

Note that assumption (3.5) is satisfied for (3.36). If the function ψ satisfies (3.26), then (3.3) holds for any $q_0 > 0$ and any bounded solution u to (3.36). Thus, if u is any bounded solution to (3.36) subject to

some globally Lipschitz continuous initial value u^0 , then $|u_x|$ does not blow up in finite time according to Theorem 3.3. Consequently, Theorem 3.3 sharpens the statement of [7, Proposition 2.14] in the following way: If (3.26) is violated and if u is a maximal solution to (3.36) on $[0, t^+) \times [-\ell, \ell]$ with $|u_x|_{\Omega_t} \to \infty$ for $t \to t^+$ as constructed in [7, Proposition 2.14], then

$$\frac{1}{2}\int_{0}^{\infty}\frac{\rho\,\mathrm{d}\rho}{\psi(\rho)}\leq |u|_{\Omega_{t^+}}$$

3.2. **Hölder estimates on the gradient and global existence.** We next state a result on Hölder estimates on the gradient of a solution to (1.1), (1.2). It turns out that one can apply the existing theory [23, Chapter VI] rather easily for which the assumptions of Theorem 3.1 have to be strengthened just slightly.

Corollary 3.4. Let $u^0 \in C^{1+\alpha}([-\ell, \ell])$ for some $\alpha \in (0, 1)$ and let (3.2)-(3.5) be satisfied with M > 0. There are numbers $\delta := \delta(M, \alpha) \in (0, \alpha]$ and $C_* := C_*(M, \psi, \langle u_x^0 \rangle_{\Omega}^{\alpha})$ such that if $u \in C^{1,2}(\overline{\Omega}_{\tau})$ is any classical solution to (1.1), (1.2) on Ω_{τ} with $\tau \leq T$ subject to $u(0, \cdot) = u^0$ satisfying $|u|_{\Omega_{\tau}} \leq M$, then $\langle u_x \rangle_{\Omega_{\tau}}^{(\frac{\delta}{2}, \delta)} \leq C_*$.

Proof. Owing to Theorem 3.1 there is a constant $M_1 = M_1(M, \psi, K)$ independent of τ such that $|u_x|_{\Omega_{\tau}} \leq M_1$. Since a and f are continuous and a is positive, it follows that there is $\mu > 0$ such that

$$\mu^{-1} \le a(t, x, z, p) \le \mu, \quad |f(t, x, z, p)| \le \mu, \qquad (t, x, z, p) \in \overline{\Omega}_T \times [-M, M] \times [-M_1, M_1].$$

Moreover, setting

$$c_1 := \max\left\{ |b(t, \pm \ell, z, p)p| + |g(t, \pm \ell, z, p)| ; (t, z, p) \in [0, T] \times [-M, M] \times [-M_1, M_1] \right\}$$

we have $\max_{t \in [0,\tau]} |u_t(t, \pm \ell)| \le c_1$ due to (1.2). The assertion is now a consequence of [23, Chapter VI, Theorem 5.1].

Summarizing our findings we can simplify the criterion for global existence to problem (1.1), (1.3) from Theorem 2.2. While the latter requires a uniform bound on the Hölder norms of solutions, the following corollary states that bounds on the supremum norm are sufficient.

Corollary 3.5. Let $\alpha \in (0, 1)$ and suppose that (2.7) holds, where *b* and *g* are gradient-independent and a > 0 and b > 0. Consider $u^0 \in C^{2+\alpha}([-\ell, \ell])$ satisfying the compatibility condition (2.9). Suppose there is a constant M > 0 such that (3.2)-(3.5) are satisfied. If the unique solution *u* to (1.1), (1.3) subject to $u(0, \cdot) = u^0$ on the maximal interval of existence $[0, \tau_{\infty})$ provided by Theorem 2.2 satisfies $|u|_{\Omega_{\tau}} \leq M$ for $\tau < \tau_{\infty}$, then $u \in C^{1+\frac{\alpha}{2}, 2+\alpha}(\Omega_T)$.

Proof. Let u be the unique solution to (1.1), (1.3) subject to $u(0, \cdot) = u^0$ on the maximal interval of existence $[0, \tau_{\infty})$ provided by Theorem 2.2 and suppose that $|u|_{\Omega_{\tau}} \leq M$ for $\tau < \tau_{\infty}$. Then, by Theorem 3.1, Corollary 3.4 and arguments parallel to those at the end of the proof Theorem 2.2 (cf. (2.16)-(2.19)) we have

$$|u|_{\Omega_\tau}^{(\frac{2+\delta}{2},2+\delta)} \le c(M,\ell,T)$$

for some $\delta \in (0, \alpha]$ and some constant $c(M, \ell, T)$ independent of τ . By embedding we obtain

$$\sup_{\tau < \tau_{\infty}} \left(|u|_{\Omega_{\tau}} + |u_x|_{\Omega_{\tau}} + \langle u_x \rangle_{\Omega_{\tau}}^{\left(\frac{\alpha}{2}, \alpha\right)} \right) < \infty \,.$$

The assertion follows now from a further application of Theorem 2.2.

Remark 3.6. For simplicity we stated Corollary 3.5 only for the case of the Bernstein-Nagumo condition considered in Theorem 3.1. The result is also true in the case of the weakened Bernstein-Nagumo condition considered in Theorem 3.2.

3.3. A priori estimate on the solution. In the previous subsections we have derived *a priori* estimates on the gradient and its Hölder norm of a solution u to (1.1), (1.2) in dependence on $|u|_{\Omega_{\tau}}$. As in [23, Chapt. I, Thm. 2.9] we shall now also consider a simple situation for which one can derive an *a priori* bound on $|u|_{\Omega_{\tau}}$ itself. This yields the bound (3.1) required for global existence.

Let there be a number B > 0 such that f and g satisfy the growth conditions

 $zf(t, x, z, 0) \le \Phi(|z|)|z| + B$, $zg(t, \pm \ell, z, p) \le \Phi(|z|)|z| + B$, $(t, x, z, p) \in \Omega_T \times \mathbb{R} \times \mathbb{R}$, (3.37) where Φ is a non-decreasing positive function on $[0, \infty)$ with

$$\int_0^\infty \frac{\mathrm{d}r}{\Phi(r)} = \infty. \tag{3.38}$$

Let $\phi(\xi)$ be defined for $\xi \in (0,\infty)$ by

$$\int_0^{\phi(\xi)} \frac{\mathrm{d}r}{\Phi(r)} = \ln \xi$$

and note that ϕ is monotonically increasing from $-\infty$ to ∞ with $\phi(1) = 0$ and satisfies $\Phi(\phi(\xi)) = \xi \phi'(\xi)$ for $\xi > 0$. Then solutions to (1.1), (1.2) are bounded:

Proposition 3.7. Suppose (3.37) with (3.38) and set

$$M := \inf \left\{ \phi(\xi) : \lambda > 1, \, \xi = \max \left\{ 1, \, \phi^{-1} \left(\frac{B}{(\lambda - 1)\Phi(0)} \right), \, \phi^{-1} \left(|u^0|_{[-\ell,\ell]} \right) \right\} \right\}.$$
(3.39)

If u is any classical solution to (1.1), (1.2) on Ω_{τ} with $\tau \leq T$ subject to $u(0, \cdot) = u^0$, then $|u|_{\Omega_{\tau}} \leq M$.

Proof. The proof follows along the lines of the proof of [23, Chapt. I, Thm. 2.9] and differs merely where the dynamical boundary conditions come into play. For the reader's ease we give the complete proof here. Put $u = \phi(v)$ and, given $\lambda > 1$, set

$$\hat{v}(t,x) := v(t,x)e^{-\lambda t}, \quad (t,x) \in \overline{\Omega}_{\tau}.$$

Let \hat{v} attain its maximum at some point $(t_0, x_0) \in \overline{\Omega}_{\tau}$. We want to derive an upper bound on \hat{v} and may thus restrict to the case $\hat{v}(t_0, x_0) \ge e^{-\lambda t_0}$, that is, $v(t_0, x_0) \ge 1$ and $u(t_0, x_0) \ge 0$. Let

$$\Gamma:=\{0\}\times [-\ell,\ell]\cup (0,\tau)\times \{\pm\ell\}$$

be the parabolic boundary of Ω_{τ} .

Case 1: Assume $(t_0, x_0) \in \overline{\Omega}_{\tau} \setminus \Gamma$. In this case, (t_0, x_0) does not belong to Γ and at this point we have

$$\hat{v}(t_0, x_0) > 0, \quad \hat{v}_x(t_0, x_0) = 0, \quad \hat{v}_{xx}(t_0, x_0) \le 0, \quad \hat{v}_t(t_0, x_0) \ge 0.$$
 (3.40)

According to (1.1),

$$\hat{v}_t e^{\lambda t} + \lambda \hat{v} e^{\lambda t} - a(t, x, u, u_x) \left(\hat{v}_{xx} e^{\lambda t} + \frac{\phi^{\prime\prime}(v)}{\phi^\prime(v)} e^{2\lambda t} \hat{v}_x^2 \right) = \frac{1}{\phi^\prime(v)} f(t, x, u, u_x)$$

and thus, by (3.40),

$$\lambda \hat{v}(t_0, x_0) e^{\lambda t_0} \le \frac{1}{\phi'(v)} f(t_0, x_0, u, 0).$$

Multiplying the previous inequality by $\phi'(v(t_0, x_0))u(t_0, x_0) \ge 0$ and invoking (3.37) yields

$$\lambda \phi'(v)vu \le f(t_0, x_0, u, 0)u \le B + \Phi(u)u$$
 at $(t, x) = (t_0, x_0)$,

which also reads

$$(\lambda - 1)\Phi(u)u \le B$$
 at $(t, x) = (t_0, x_0)$.

Hence, since Φ is non-decreasing,

$$u(t_0, x_0) \le \frac{B}{(\lambda - 1)\Phi(0)}$$

from which we deduce, since ϕ^{-1} is increasing,

$$v(t,x) \le e^{\lambda t} \hat{v}(t_0,x_0) = \phi^{-1} \left(u(t_0,x_0) \right) e^{\lambda(t-t_0)} \le \phi^{-1} \left(\frac{B}{(\lambda-1)\Phi(0)} \right) e^{\lambda T}, \qquad (t,x) \in \overline{\Omega}_{\tau}.$$
 (3.41)

Case 2: Assume $(t_0, x_0) \in \Gamma$. Suppose first that $t_0 = 0$. Then

$$v(t,x) \le e^{\lambda t} \hat{v}(t_0,x_0) = e^{\lambda t} \phi^{-1} \big(u(0,x_0) \big) \le e^{\lambda T} \phi^{-1} \big(|u^0|_{[-\ell,\ell]} \big), \qquad (t,x) \in \overline{\Omega}_{\tau}.$$
(3.42)

If $t_0 \neq 0$, then $(t_0, x_0) \in (0, \tau) \times \{\pm \ell\}$ in which case we may use the boundary condition (1.2). So, if $(t_0, x_0) \in (0, \tau) \times \{\ell\}$, then $b(\cdot, \ell, \cdot, \cdot) > 0$ and the fact that (t_0, ℓ) is a point of a positive maximum imply

$$v(t_0,\ell) > 0, \quad v_t(t_0,\ell) = 0, \quad b(t_0,\ell,u(t_0,\ell),u_x(t_0,\ell))v_x(t_0,\ell) \ge 0.$$

From (1.2) and (3.37) we deduce, at $(t, x) = (t_0, \ell)$,

$$\lambda \phi'(v) \hat{v} e^{\lambda t} \le e^{\lambda t} \phi'(v) \left(\lambda \hat{v} + \hat{v}_t\right) + b(t, x, u, u_x) \phi'(v) e^{\lambda t} \hat{v}_x = g\left(t, \ell, u, u_x\right) \le B + \Phi(u)u,$$

which again reads

$$(\lambda - 1)\Phi(u)u \le B$$
 at $(t, x) = (t_0, \ell)$

and we may proceed as in Case 1 to derive (3.41). Clearly, the same argument holds when $(t_0, x_0) \in (0, \tau) \times \{-\ell\}$.

Combining the different cases from (3.41) and (3.42) we derive that

$$\max_{\overline{\Omega}_{\tau}} u \leq M$$

with M given in (3.39). Considering -u on $\overline{\Omega}_{\tau}$ yields an estimate from below on u. This proves the assertion.

Corollary 3.8. Let $\alpha \in (0, 1)$ and suppose (2.7). Consider $u^0 \in C^{2+\alpha}([-\ell, \ell])$ satisfying the compatibility condition (2.9). Suppose (3.37) with (3.38) and that (3.2)-(3.5) are satisfied with M given in (3.39). Then there exists a global solution $u \in C^{1+\frac{\alpha}{2},2+\alpha}(\Omega_T)$ to (1.1), (1.3) subject to $u(0, \cdot) = u^0$.

Proof. This now follows from Theorem 2.2, Theorem 3.1, Corollary 3.4, and Proposition 3.7.

Remark 3.9. For simplicity we stated the global existence result of Corollary 3.8 only for the case of the Bernstein-Nagumo condition considered in Theorem 3.1. The result is also true in the case of the weakened Bernstein-Nagumo condition considered in Theorem 3.2.

As a final remark let us point out that, as in [29], we could replace (1.1) by the more general equation

$$u_t = F(t, x, u, u_x, u_{xx})$$

and derive similar results provided that F is differentiable with respect to its last variable.

4. APPENDIX: NOTATION AND PARABOLIC HÖLDER SPACES

We provide in this appendix the definition of parabolic Hölder spaces, for more details we refer e.g. to [23,24]. Recall that $\Omega = (-\ell, \ell)$ and $\Omega_T = (0, T) \times \Omega$. Let Q be an interval in \mathbb{R} and $\alpha \in (0, 1]$. Then $u : Q \to \mathbb{R}$ is uniformly α -Hölder continuous if

$$[u]_Q^{(\alpha)} := \sup_{x,y \in Q} \frac{|u(x) - u(y)|}{|x - y|^{\alpha}} < \infty$$

We say that $u : Q \to \mathbb{R}$ is α -Hölder continuous if every point in Q has a neighborhood $Q' \subset Q$ such that $u|_{Q'}$ is uniformly α -Hölder continuous. Note that α -Hölder continuous functions are uniformly α -Hölder continuous on compact sets. Clearly, 1-Hölder continuous functions are Lipschitz continuous. Given $k \in \mathbb{N}$ and $\alpha \in (0, 1)$ we let $C^{k+\alpha}(Q)$ be the space of all k-times continuously differentiable functions on Q

such that the k-th derivative is α -Hölder continuous. If $u : S \to \mathbb{R}$ is bounded on a set S, we denote its supremum norm by $|u|_S$, i.e.

$$|u|_S := \sup_{x \in S} |u(x)|.$$

If Q is bounded and $k \in \mathbb{N}$, $\alpha \in (0, 1)$ given, we put

$$|u|_Q^{(k+\alpha)} := \sum_{j \le k} |D^j u|_Q + [D^k u]_Q^{(\alpha)}$$

for $u \in C^{k+\alpha}(\overline{Q})$.

Let T > 0 and $Q_T = (0, T) \times Q$. We let $C^{1,2}(Q_T)$ be the space of all continuous real-valued functions defined on Q_T having continuous derivatives u_t, u_{xx} on Q_T and we use a similar notation on \overline{Q}_T .

The parabolic Hölder spaces $\mathcal{C}^{\frac{k+\alpha}{2},k+\alpha}(Q_T)$ for $k \in \mathbb{N}$ and $\alpha \in (0,1)$ are defined as the set of functions $u: Q_T \to \mathbb{R}$ having finite norm

$$|u|_{Q_T}^{(\frac{k+\alpha}{2},k+\alpha)} := \sum_{2j_0+j \le k} |D_t^{j_0} D_x^j u|_{Q_T} + \langle u \rangle_{Q_T}^{(\frac{k+\alpha}{2},k+\alpha)} < \infty,$$

where, if k = 0,

$$\langle u \rangle_{Q_T}^{(\frac{\alpha}{2},\alpha)} := \langle u \rangle_{t,Q_T}^{(\frac{\alpha}{2})} + \langle u \rangle_{x,Q_T}^{(\alpha)}$$

with

$$\langle u \rangle_{t,Q_T}^{(\alpha)} := \sup_{x \in Q} [u(\cdot, x)]_{(0,T)}^{(\alpha)}, \qquad \langle u \rangle_{x,Q_T}^{(\alpha)} := \sup_{t \in (0,T)} [u(t, \cdot)]_{\Omega}^{(\alpha)},$$

and, if $k \geq 1$,

$$\langle u \rangle_{Q_T}^{(\frac{k+\alpha}{2}, k+\alpha)} := \sum_{2j_0+j=k} \langle D_t^{j_0} D_x^j u \rangle_{Q_T}^{(\frac{\alpha}{2}, \alpha)} + \sum_{2j_0+j=k-1} \langle D_t^{j_0} D_x^j u \rangle_{t, Q_T}^{(\frac{1+\alpha}{2})}.$$
(4.1)

Since globally Hölder continuous functions are uniformly continuous, it follows that

$$D_t^{j_0} D_x^j u \in C(\overline{Q}_T), \quad 2j_0 + j \le k,$$

for $u \in \mathcal{C}^{\frac{k+\alpha}{2},k+\alpha}(Q_T)$.

The following interpolation inequalities are quite useful. We refer to [22, Thm.8.8.1] (see also [22, Exercise 8.8.2]) for a proof.

Proposition 4.1. There is a constant c = c(Q) such that for $u \in C^{1+\frac{\alpha}{2},2+\alpha}(Q_T)$:

$$|u_t|_{Q_T} + |u_{xx}|_{Q_T} \le c U_{2+\alpha}^{1/(2+\alpha)} U_0^{1-1/(2+\alpha)},$$

$$\langle u_x \rangle_{Q_T}^{(\frac{\alpha}{2},\alpha)} \le c U_{2+\alpha}^{(1+\alpha)/(2+\alpha)} U_0^{1-(1+\alpha)/(2+\alpha)},$$

$$\langle u \rangle_{Q_T}^{(\frac{\alpha}{2},\alpha)} \le c U_{2+\alpha}^{\alpha/(2+\alpha)} U_0^{1-\alpha/(2+\alpha)},$$
(4.2)

where $U_0 := |u|_{Q_T}$ and $U_{2+\alpha} := \langle u \rangle_{Q_T}^{(1+\frac{\alpha}{2},2+\alpha)}$.

REFERENCES

- B. V. Bazaliy, N. Vasylyeva. On the solvability of a transmission problem for the Laplace operator with a dynamical boundary condition on a nonregular interface. J. Math. Anal. Appl. 393 (2012), no. 2, 651-670.
- [2] B. V. Bazaliy, S. P. Degtyarev. On the classical solvability of a multidimensional Stefan problem with convective movement of a viscous incompressible fluid. Mat. Sb. (N.S.) 132(174) (1987), no. 1, 3-19, 142 (Russian); translation in Math. USSR-Sb. 60 (1988), no. 1, 1-17.
- [3] J. von Below, C. De Coster. A qualitative theory for parabolic problems under dynamical boundary conditions. J. Inequal. Appl. 5 (2000), no. 5, 467-486.
- [4] J. von Below, G.P. Mailly. Blow up for some nonlinear parabolic problems with convection under dynamical boundary conditions. Discrete Contin. Dyn. Syst. 2007, Dynamical Systems and Differential Equations. Proceedings of the 6th AIMS International Conference, suppl., 1031-1041.

- [5] G. I. Bizhanova, V. A. Solonnikov. On the solvability of an initial-boundary value problem for a second-order parabolic equation with a time derivative in the boundary condition in a weighted Hlder space of functions. Algebra i Analiz 5 (1993), no. 1, 109-142 (Russian); translation in: St. Petersburg Math. J. 5 (1994), no. 1, 97-124.
- [6] G. I. Bizhanova, V. A. Solonnikov. Some model problems for second-order parabolic equations with a time derivative in the boundary conditions. Algebra i Analiz 6 (1994), no. 6, 30-50 (Russian); translation in St. Petersburg Math. J. 6 (1995), no. 6, 1151-1166.
- [7] A. Constantin, J. Escher. Global existence for fully parabolic boundary value problems. NoDEA Nonlinear Differential Equations Appl. 13 (2006), no. 1, 91-118.
- [8] R. Denk, J. Prüß, R. Zacher Maximal L_p-regularity of parabolic problems with boundary dynamics of relaxation type. Journ. Funct. Anal. 255 (2008) 3149–3187.
- [9] C. M. Elliott, Y. Giga, S. Goto. Dynamic boundary conditions for Hamilton-Jacobi equations. SIAM J. Math. Anal. 34 (2003), no. 4, 861-881 (electronic).
- [10] J. Escher. Über quasilineare parabolische Probleme. Dissertation. Universität Zürich, Zenstralstelle der Studentenschaft, Zürich, 1991.
- [11] J. Escher. Global existence and nonexistence for semilinear parabolic systems with nonlinear boundary conditions. Math. Ann. 284 (1989), no. 2, 285-305.
- [12] J. Escher. Quasilinear parabolic systems with dynamical boundary conditions. Comm. Partial Differential Equations 18 (1993), no. 7-8, 1309-1364.
- [13] M. Fila, K. Ishige, T. Kawakami. Convergence to the Poisson kernel for the Laplace equation with a nonlinear dynamical boundary condition. Commun. Pure Appl. Anal. 11 (2012), no. 3, 1285-1301.
- [14] M. Fila, P. Quittner. Global solutions of the Laplace equation with a nonlinear dynamical boundary condition. Math. Methods Appl. Sci. 20 (1997), no. 15, 1325-1333.
- [15] M. Fila, P. Quittner. Large time behavior of solutions of a semilinear parabolic equation with a nonlinear dynamical boundary condition. Topics in nonlinear analysis, 251-272, Progr. Nonlinear Differential Equations Appl., 35, Birkhuser, Basel, 1999.
- [16] C. G. Gal. On a class of degenerate parabolic equations with dynamical boundary conditions. J. Differential Equations 253 (2012), no. 1, 126-166.
- [17] G. R. Goldstein. Derivation and physical interpretation of general boundary conditions. Adv. Differential Equations 11 (2006), no. 4, 457-480.
- [18] D. Guidetti. *Classical solutions to quasilinear parabolic problems with dynamic boundary conditions.* arXiv:1504.06112v1, 2015.
- [19] S. Gvelesiani. Globale klassische Lösungen nichtlinearer parabolischer Probleme mit dynamischen Randbedingungen in einer Raumdimension. Dissertation, Leibniz Univeristät Hannover, 2014.
- [20] S. N. Kruzhkov. Quasilinear parabolic equations and systems with two independent variables. Trudy Sem. Petrovsk. 5 (1979), 217272 (Russian). English transl. in: Topics in Modern Math., Consultant Bureau, New York 1985.
- [21] S. N. Kruzhkov. Nonlinear parabolic equations in two independent variables. Transact. Moscow Math. Soc. 16 (1968), 355373.
- [22] N. V. Krylov. *Lectures on elliptic and parabolic equations in Hlder spaces*. Graduate Studies in Mathematics, 12. American Mathematical Society, Providence, RI, 1996.
- [23] O. A. Ladyzhenskaya, V. A. Solonnikov, N. N. Ural'ceva. Linear and quasi-linear equations of parabolic type. Vol. 23 American Mathematical Society, Providence, R.I. 1967.
- [24] G. M. Lieberman. Second order parabolic differential equations. World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [25] M. Meyries. Maximal Regularity in Weighted Spaces, Nonlinear Boundary Conditions, and Global Attractors. Dissertation. Karlsruher Institut fr Technologie, Karlsruhe, 2010.
- [26] J. Petersson. A note on quenching for parabolic equations with dynamical boundary conditions. Nonlinear Anal. 58 (2004), no. 3-4, 417-423.
- [27] P. Quittner, P. Souplet. Superlinear parabolic problems. Blow-up, global existence and steady states. Birkhuser Advanced Texts: Basler Lehrbcher. Birkhuser Verlag, Basel, 2007.
- [28] S. I. Temirbulatov. The problem of heat conduction with a derivative with respect to time in the boundary condition. Differentsial'nye Uravneniya 19 (1983), no. 4, 666-673 (Russian).
- [29] A. Tersenov, A. Tersenov. On the Bernstein-Nagumo's condition in the theory of nonlinear parabolic equations. J. Reine Angew. Math. 572 (2004), 197-217.
- [30] J. L. Vázquez, E. Vitillaro. Heat equation with dynamical boundary conditions of reactive-diffusive type. J. Differential Equations 250 (2011), no. 4, 2143-2161.
- [31] N. V. Vasylyeva. The elliptic and parabolic boundary value problems with the time derivative in the boundary condition in a plane corner and their application to free boundary problems. Dissertation. Institute of Applied Mathematics and Mechanics, National Academy of Science of Ukraine, Donetsk, 2002 (Russian).

Leibniz Universität Hannover, Institut für Angewandte Mathematik, Welfengarten 1, D–30167 Hannover, Germany

E-mail address: gvelesiani@ifam.uni-hannover.de

Leibniz Universität Hannover, Institut für Angewandte Mathematik, Welfengarten 1, D–30167 Hannover, Germany

E-mail address: lippoth@ifam.uni-hannover.de

Leibniz Universität Hannover, Institut für Angewandte Mathematik, Welfengarten 1, D–30167 Hannover, Germany

E-mail address: walker@ifam.uni-hannover.de