# KHOVANOV HOMOTOPY TYPE, BURNSIDE CATEGORY, AND PRODUCTS

#### TYLER LAWSON, ROBERT LIPSHITZ, AND SUCHARIT SARKAR

ABSTRACT. In this paper, we give a new construction of a Khovanov stable homotopy type, or spectrum. We show that this construction gives a space stably homotopy equivalent to the Khovanov spectra constructed in [LS14a] and [HKK16] and, as a corollary, that those two constructions give equivalent spectra. We show that the construction behaves well with respect to disjoint unions, connected sums and mirrors, verifying several conjectures from [LS14a]. Finally, combining these results with computations from [LS14c] and the refined s-invariant from [LS14b] we obtain new results about the slice genera of certain knots.

### Contents

1. Introduction	2
1.1. Basic notation	6
2. Review of Khovanov homology	7
2.1. The cube category	7
2.2. The Khovanov construction	7
3. Cubical flow categories	8
3.1. Manifolds with corners and $\langle n \rangle$ -manifolds	9
3.2. Flow categories	10
3.3. Permutohedra	11
3.4. The cube flow category	13
3.5. Definition of a cubical flow category	16
3.6. Cubical neat embeddings	17
3.7. Cubical realization	19
3.8. Cubical realization agrees with the Cohen-Jones-Segal realization	23
4. Functors from the cube to the Burnside category and their realizations	25
4.1. The Burnside category and 2-functors to it	26
4.2. Homotopy colimits and homotopy coherent diagrams	28
4.3. Cubical flow categories are functors from the cube to the Burnside category	31
4.4. The thickened diagram	33
4.5. The realization	37
4.6. An invariance property of the realization	38
4.7. Products and realization	39
5. Building a smaller cube from little box maps	41
5.1. Box maps	42

Date: September 6, 2019.

 $<sup>2010\</sup> Mathematics\ Subject\ Classification.\ 57M25,\ 55P42.$ 

TL was supported by NSF Grant DMS-1206008.

RL was supported by NSF Grant DMS-1149800.

 $<sup>{\</sup>rm SS}$  was supported by NSF Grant DMS-1350037.

5.2. Refining diagrams via box maps	44
5.3. A coherent cube of box maps	46
5.4. The realizations of the small cube and big cube agree	46
6. A CW complex structure on the realization of the small cube	51
7. The Khovanov homotopy type	57
8. Relationship with Hu-Kriz-Kriz	57
8.1. The functors from the cube to the Burnside category agree	58
8.2. Iterated mapping cones	61
8.3. Another kind of homotopy coherent diagram	61
8.4. The Elmendorff-Mandell machine	62
8.5. Proof that the Khovanov homotopy types agree	65
9. Khovanov homotopy type of a disjoint union and connected sum	65
10. Khovanov homotopy type of a mirror	77
11. Applications	80
Appendix A. Flow chart of sections	85
References	85

#### 1. Introduction

In a sequence of revolutionary papers in the 1980s, Floer introduced a family of invariants in low-dimensional and symplectic topology, now known as Floer homologies [Flo88a, Flo88b, Flo88c]. While the definitions of these invariants appear to be a semi-infinite dimensional version of Morse homology, unlike classical (finite- or infinite-dimensional) Morse homology, Floer homologies were not apparently isomorphic to the singular homologies of any natural space. In the 1990s, under appropriate hypotheses Cohen-Jones-Segal proposed a construction of a stable homotopy type whose singular homology was Floer homology [CJS95]. Their construction builds a CW complex cell-by-cell, using a version of the Pontrjagin-Thom construction for manifolds with corners to define the attaching maps; the input data to this Pontrjagin-Thom construction is what Cohen-Jones-Segal call a framed flow category. Analytic difficulties mean that their proposal has not yet been carried out rigorously, though Manolescu constructed a stable homotopy refinement of the Seiberg-Witten Floer homology of rational homology 3-spheres by other techniques [Man03] (compare [LM18]).

Around the turn of the millennium, in a seminal paper Khovanov gave a bigraded homology theory whose graded Euler characteristic is the Jones polynomial [Kho00]. The definition of this *Khovanov homology* is purely combinatorial, though equivalent or conjecturally equivalent definitions have been given using Floer homology [SS06] (see also [AS]) and algebraic geometry [CK08].

Inspired by Cohen-Jones-Segal's work and the relation with Floer homology, in [LS14a] we introduced a stable homotopy refinement of Khovanov homology. That is, to each link diagram K we associated a spectrum  $\mathcal{X}_{Kh}(K)$ , with the following properties:

- (1) The spectrum  $\mathcal{X}_{Kh}(K)$  is a finite CW spectrum, that is, a formal desuspension of a finite CW complex.
- (2) The spectrum  $\mathcal{X}_{Kh}(K)$  comes with a wedge-sum decomposition  $\mathcal{X}_{Kh}(K) = \bigvee_{j} \mathcal{X}_{Kh}^{j}(K)$ .
- (3) For each j, the cellular cochain complex of  $\mathcal{X}^{j}_{Kh}(K)$  is isomorphic to the Khovanov complex  $\mathcal{C}^{*,j}_{Kh}(K)$  in quantum grading j, via an isomorphism taking the standard generators for  $C^*_{cell}(\mathcal{X}^{j}_{Kh}(K))$  to the standard generators for  $\mathcal{C}^{*,j}_{Kh}(K)$ .

(4) For each j, the stable homotopy type of  $\mathcal{X}_{Kh}^{j}(K)$  is an invariant of the isotopy class of the link represented by the diagram K.

There is also a reduced version  $\widetilde{\mathcal{X}}_{Kh}(K)$  of  $\mathcal{X}_{Kh}(K)$ , which satisfies properties (1)–(4) with  $\mathcal{C}_{Kh}$  replaced by the reduced Khovanov complex  $\widetilde{\mathcal{C}}_{Kh}$ . The constructions of  $\mathcal{X}_{Kh}(K)$  and  $\widetilde{\mathcal{X}}_{Kh}(K)$  involve defining a framed flow category combinatorially, and then applying Cohen-Jones-Segal's Pontrjagin-Thom construction.

Largely independently, inspired by their homotopy-theoretic investigations of topological and conformal field theories, Hu-Kriz-Kriz gave another construction of a Khovanov stable homotopy type with the same basic properties [HKK16]. Roughly, they turn the Khovanov cube into a functor from the cube category to the Burnside category of finite sets and correspondences. They then apply the Elmendorf-Mandell infinite loop space machine to obtain a functor from the cube to symmetric spectra, and then take an iterated mapping cone to obtain a spectrum.

These refinements give extra structure to the Khovanov homology groups. Because the Khovanov homology groups of K are cohomology groups of  $\mathcal{X}_{Kh}(K)$ , the mod-p cohomology groups possess Steenrod power operations such as the squares  $\operatorname{Sq}^n$  at p=2. It also allows one to apply generalized comohology theories, such as topological K-theory, Morava K-theory, and stable homotopy groups. Each of these has its own Atiyah-Hirzebruch spectral sequence determining a potentially new collection of knot invariants, and maps in the stable homotopy category give natural transformations relating these different invariants. In the philosophically related case of Seiberg-Witten Floer theory, for instance, equivariant KO-theory has been used to obtain interesting low-dimensional applications [Man14, Lin15].

One goal of the present paper is to study the behavior of  $\mathcal{X}_{Kh}(K)$  under disjoint unions and connected sums of links. In particular, we will prove:

**Theorem 1.** [LS14a, Conjecture 10.3] Let  $L_1$  and  $L_2$  be links, and  $L_1 \coprod L_2$  their disjoint union. Then

(1.1) 
$$\mathcal{X}_{Kh}^{j}(L_1 \coprod L_2) \simeq \bigvee_{j_1+j_2=j} \mathcal{X}_{Kh}^{j_1}(L_1) \wedge \mathcal{X}_{Kh}^{j_2}(L_2).$$

Moreover, if we fix a basepoint p in  $L_1$ , not at a crossing, and consider the corresponding basepoint for  $L_1 \coprod L_2$ , then

(1.2) 
$$\widetilde{\mathcal{X}}_{Kh}^{j}(L_1 \coprod L_2) \simeq \bigvee_{j_1+j_2=j} \widetilde{\mathcal{X}}_{Kh}^{j_1}(L_1) \wedge \mathcal{X}_{Kh}^{j_2}(L_2).$$

**Theorem 2.** [LS14a, Conjecture 10.4] Let  $L_1$  and  $L_2$  be based links and  $L_1 \# L_2$  the connected sum of  $L_1$  and  $L_2$ , where we take the connected sum near the basepoints. Then

(1.3) 
$$\widetilde{\mathcal{X}}_{Kh}^{j}(L_1 \# L_2) \simeq \bigvee_{j_1 + j_2 = j} \widetilde{\mathcal{X}}_{Kh}^{j_1}(L_1) \wedge \widetilde{\mathcal{X}}_{Kh}^{j_2}(L_2).$$

(We also compute the unreduced Khovanov spectrum of a connected sum [LS14a, Conjecture 10.6], as Theorem 10.)

These theorems, though unsurprising themselves, have some interesting corollaries:

Corollary 1.4. For any n there exists an n-component link  $L_n$  so that the operation

$$\operatorname{Sq}^n \colon Kh^{i,j}(L_n) \to Kh^{i+n,j}(L_n)$$

is non-zero, for some  $i, j \in \mathbb{Z}$ . Similarly, there exists a knot  $K_n$  so that the operation

$$\operatorname{Sq}^n \colon \widetilde{Kh}^{i,j}(K_n) \to \widetilde{Kh}^{i+n,j}(K_n)$$

is non-zero, for some  $i, j \in \mathbb{Z}$ . Further, for this knot, the operation

$$\operatorname{Sq}^n \colon Kh^{i,j}(K_n) \to Kh^{i+n,j}(K_n)$$




FIGURE 1.1. The knots appearing in Corollary 1.5. We have labeled the knot K by the pair  $(K, g_4(K))$ . The value of  $g_4(K)$  is extracted from Knotinfo [CL]; the knot diagrams have been produced using Knotilus [FFFR]. Crossings giving a minimal unknotting for each knot are circled.

is also non-zero for some  $i, j \in \mathbb{Z}$ .

Corollary 1.5. Let K be one of the knots  $9_{42}$ ,  $10_{136}$ ,  $m(11_{19}^n)$ ,  $m(11_{20}^n)$ ,  $11_{70}^n$ , or  $11_{96}^n$ . (Here m denotes the mirror.) Let L be a knot which is the closure of a positive braid. Letting  $g_4$  denote the four-ball genus, we have

$$g_4(K \# L) = g_4(K) + g_4(L).$$

Remark 1.6. There is some confusion for nomenclature of knots regarding mirrors. See Figure 1.1 for our convention, and compare with the discussion around Table 11.1. The value of  $g_4(K)$  can be extracted from Figure 1.1 or Table 11.1; and  $g_4(L)$  equals the genus of the Seifert surface obtained by applying Seifert's algorithm to the positive braid closure knot diagram for L [Ras10, Theorem 4].

Corollary 1.5 is not implied by computations of Rasmussen's s-invariant or the Heegaard Floer  $\tau$ -invariant or the signature; see Table 11.1. At least for  $9_{42}$ , the result is not implied by the Heegaard Floer concordance invariant  $\Upsilon$ ; the Heegaard Floer d invariant of +1 surgery [Krc14]; or Hom-Rasmussen-Wu's  $\nu^+$ -invariant [Ras03, HW16, Hom14]. (These observations are not entirely independent.)

The construction of  $\mathcal{X}_{Kh}(K)$  from [LS14a] uses Cohen-Jones-Segal's notion of flow categories. To give a proof of Theorems 1 and 2 in this language seems tedious at best: it involves understanding the combinatorics of (broken) Morse flows on product manifolds, which turns out to be rather intricate.

Fortunately, as noted above, Hu-Kriz-Kriz gave another construction of a Khovanov stable homotopy type, and from their construction the behavior under connected sums and disjoint unions is clear. In this paper, we describe three additional constructions:

- (1) A framing-free reformulation of the construction from [LS14a] for a special family of flow categories, called cubical flow categories, in Section 3.
- (2) A version similar to the construction from [HKK16] as a homotopy colimit, but using the thickened cube instead of the Elmendorf-Mandell machinery, in Section 4.
- (3) An intermediate object between the two in Section 5, using little k-cubes.

#### We then prove:

**Theorem 3.** The Khovanov stable homotopy types constructed in Section 3, Section 4, and Section 5, the Khovanov stable homotopy type constructed in [HKK16], and the Khovanov stable homotopy type constructed in [LS14a] are all stably homotopy equivalent.

This theorem is proved in parts:

- Theorem 4 asserts that the cubical flow category realization (Section 3) agrees with the Cohen-Jones-Segal construction used in [LS14a].
- Theorem 8 asserts that the cubical flow category realization (Section 3) agrees with the little k-cubes realization (Section 5).
- Theorem 7 asserts that the little k-cubes realization (Section 5) agrees with the thickening construction (Section 4).
- Theorem 9 asserts that the thickening construction (Section 4) agrees with the construction in [HKK16].

Theorems 1 and 2 follow easily. Corollary 1.4 follows immediately from these theorems and computations in [LS14c]. We also obtain, via a TQFT-style argument, that the Khovanov homotopy type of the mirror knot m(K) is the Spanier-Whitehead dual to the Khovanov homotopy type of K (Theorem 11). Finally, Corollary 1.5 follows from these results, the refined s invariant in [LS14b], the computations in [LS14c], and a brief further argument.

This paper is organized as follows. We collect some basic notation in Section 1.1. Section 2 has background on the cube category and the Khovanov construction, the latter of which is not needed again until Section 7 except in some examples. Other background appears at the beginning of the section in which it is first used. In Section 3, after recalling the basics of flow categories, we introduce a special class of them, cubical flow categories, which live over the cube, and give a reformulation of the Cohen-Jones-Segal realization for this class of flow categories ("cubical realization"). (The Khovanov flow category of [LS14a] is a cubical flow category; this is a crucial tool in its construction.) In Section 4 we show that cubical flow categories are equivalent to 2-functors from the cube to the Burnside category, and give a different, choice-free way to realize such a functor. In Section 5 we give a smaller but less canonical way to realize a 2-functor from the cube to the Burnside category, and prove the two ways to realize such a functor are equivalent. Section 6 shows that the realization from Section 5 agrees with the cubical realization from Section 3. Section 7 is a brief interlude to summarize these results and recall the Khovanov homotopy type. Section 8 shows that these realizations agree with the Hu-Kriz-Kriz-construction [HKK16].

In Sections 9–11 we use these reformulated realizations to prove new properties of the Khovanov homotopy type. The realizations of the product (smash product) and disjoint union (wedge sum) of functors from the cube to the Burnside category have properties as one would expect; Section 9 uses these properties to study the Khovanov homotopy type of a disjoint union and connected sum, Theorems 1, 2, and 10,

and verifies Corollary 1.4. Section 10 uses a TQFT-style argument suggested by the referee for [LS14c] to deduce a formula for the Khovanov homotopy type of a mirror, verifying another conjecture from [LS14a]. Finally, Section 11 gives an additivity property for the refined s invariant introduced in [LS14b] and obtains Corollary 1.5.

Appendix A has a flow chart of how the different sections depend on each other, so a reader only interested in a particular result can choose the most efficient path to it.

Remark 1.7. It may be interesting to compare the homotopy colimit definition of the Khovanov homotopy type with [ET14]; but see also [ELST16].

Acknowledgments. We thank Brent Everitt, Jennifer Hom, David Krcatovich, Peter Ozsváth and Paul Turner for interesting conversations. We also thank the referee for [LS14c] for suggesting the argument in Section 10. RL thanks Princeton University for its hospitality; most of this research was completed while he visited there. We thank Michael Willis for corrections to an early version of this manuscript. We also thank the contributors to nLab: while we have cited published references for the relevant background, nLab has often been helpful in finding those references. Finally, we thank the referees for their helpful comments.

- 1.1. **Basic notation.** The "cube"  $\{0,1\}^n$  will appear in a number of contexts in this paper, as will some auxiliary notions related to it:
  - There is a partial order on  $\{0,1\}^n$  defined by  $v \ge w$  if v is obtained from w by replacing some 0's by 1's. Define v > w if  $v \ge w$  and  $v \ne w$ , and define  $\le$  and < in the corresponding ways. The maximum and minimum elements under this partial order are denoted  $\vec{1}$  and  $\vec{0}$ , respectively.
  - We denote the Manhattan (or  $\ell^1$ ) norm on  $\{0,1\}^n$  by  $|v| = \sum_{i=1}^n v_i$ .
  - A sign assignment s on the cube is the following: for every u > v with |u| |v| = 1, we associate an element  $s_{u,v} \in \mathbb{F}_2$  such that for any u > w with |u| - |w| = 2, we have

$$\sum_{\substack{v \\ v > v}} (s_{u,v} + s_{v,w}) = 1.$$

A number of categories will appear in this paper:

- The category **Sets** of finite sets and set maps.
- The Burnside (2-) category  $\mathscr{B}$  of sets, correspondences, and isomorphisms of correspondences (see Section 4.1).
- The cube category  $\underline{2}^n = \{1 \longrightarrow 0\}^n$  (see Section 2.1).
- The category Top. of (well) based topological spaces.
- The subcategory CW• of Top• generated by based CW complexes and cellular maps.
- $\bullet$  The category  $\mathscr S$  of spectra. For concreteness, we can take the category of symmetric spectra in topological spaces [MMSS01].
- The category *R*-Mod of *R*-modules.
- The category Permu of permutative categories (see Section 8.4).
- The category  $\mathsf{Cob}_{\mathrm{emb}}^{1+1}$  of oriented 1-manifolds embedded in  $S^2$  and oriented cobordisms embedded in  $[0,1] \times S^2$  (see Section 8.1).

Some other notation:

- Let  $\mathbb{R}_+ = [0, \infty)$ .
- Let  $\mathbb{N} = \{0, 1, 2, \dots\}$  denote the set of non-negative integers.




FIGURE 2.1. Some resolutions of a knot diagram. Left: A knot diagram for the figure-eight knot with crossings  $c_1, \ldots, c_4$ . Right: The complete resolutions at the vertices v = (1, 0, 0, 0) and u = (1, 1, 1, 0), along with the embedded 1-handle cores representing the embedded cobordism between  $\mathcal{P}(u)$  and  $\mathcal{P}(v)$ .

### 2. Review of Khovanov homology

2.1. The cube category. Let  $\underline{2}^1$  denote the category with two objects, denoted 0 and 1, and a single non-identity morphism, from 1 to 0:

$$\underline{2}^1 = \{1 \longrightarrow 0\}.$$

For  $n \in \mathbb{Z}$ , n > 1 let  $2^n = 2^1 \times 2^{n-1}$ . That is,  $2^n$  is the small category with object set  $\{0,1\}^n$ . Given objects  $v, w \in \{0,1\}^n$  the morphism set  $\operatorname{Hom}(v,w)$  is empty unless  $v \geq w$  in the partial order induced by the relation 1 > 0 on  $2^1$ . If  $v \geq w$  the set  $\operatorname{Hom}(v,w)$  has a single element, which we will denote  $\varphi_{v,w}$ .

The reader is warned that many authors refer to the opposite category  $(\underline{2}^n)^{op}$  as the cube category. (Our choice is made to agree with the grading conventions on Khovanov homology [Kho00] and the Khovanov stable homotopy type [LS14a].)

Often, we will view  $\underline{2}^n$  as a 2-category with no non-identity 2-morphisms. That is, for  $f, g \in \text{Hom}(v, w)$ , we define Hom(f, g) to be empty unless f = g and to have a single element if f = g.

2.2. **The Khovanov construction.** The material in the rest of this section is used in Examples 3.24 and 4.21 and Remark 4.22, and then is not used again until Section 7.

Khovanov homology, and several of its generalizations, are all constructed from the cube of resolutions of a link diagram. Let K be a link diagram in  $S^2$  with n crossings, numbered  $c_1, \ldots, c_n$ . Each of these crossings can be resolved locally in two different ways, called the 0-resolution and the 1-resolution (see, for instance, [Kho00, Figure 14]). Therefore, to each  $v \in \{0,1\}^n$  there is an associated complete resolution  $\mathcal{P}(v)$  obtained by replacing the crossing  $c_i$  by its 0-resolution if  $v_i = 0$  and its 1-resolution if  $v_i = 1$ . The complete resolution  $\mathcal{P}(v)$  consists of a collection of disjoint circles in  $S^2$ .

For any  $u \geq v \in \{0,1\}^n$ , there is a cobordism embedded in  $[0,1] \times S^2$  that connects the resolutions  $\mathcal{P}(u)$  and  $\mathcal{P}(v)$ : to obtain  $\mathcal{P}(u)$ , one attaches embedded 1-handles to  $\mathcal{P}(v)$  with cores certain arcs near the crossings  $c_i$  for all i with  $u_i > v_i$ . These arcs are illustrated in Figure 2.1. (See also [Kho00, Figure 18]. This is also the first step in Bar-Natan's "picture world" approach to Khovanov homology in [BN05].) For the special case when |u| - |v| = 1, the cobordism either merges two circles into one, or splits a single circle into two.

To construct the Khovanov complex, we apply a (1+1)-dimensional TQFT to this cube of cobordisms to obtain a commutative cube of abelian groups. Specifically, consider the rank-2 Frobenius algebras over

 $\mathbb{Z}[h,t]$  with basis  $\{x_+,x_-\}$  and multiplication and comultiplication given by

$$x_{+} \otimes x_{+} \stackrel{m}{\mapsto} x_{+} \qquad x_{+} \otimes x_{-} \stackrel{m}{\mapsto} x_{-} \qquad x_{-} \otimes x_{+} \stackrel{m}{\mapsto} x_{-} \qquad x_{-} \otimes x_{-} \stackrel{m}{\mapsto} hx_{-} + tx_{+}$$

$$x_{+} \stackrel{S}{\mapsto} x_{+} \otimes x_{-} + x_{-} \otimes x_{+} - hx_{+} \otimes x_{+} \qquad x_{-} \stackrel{S}{\mapsto} x_{-} \otimes x_{-} + tx_{+} \otimes x_{+},$$

and the corresponding (1+1)-dimensional TQFT. Applying this TQFT to the cube of resolutions of K gives a commutative cube  $A: (\underline{2}^n)^{\mathrm{op}} \to \mathbb{Z}[h,t]$ -Mod. Explicitly, given a vertex v, a Khovanov generator over v is a labeling of the circles in  $\mathcal{P}(v)$  by elements of the set  $\{x_+,x_-\}$ . The module A(v) is freely generated by the set of Khovanov generators F(v) over v. For an edge of the cube  $\varphi_{u,v}$  (where u > v and |u| - |v| = 1),  $A(\varphi_{u,v}): A(v) \to A(u)$  is the multiplication or comultiplication above, depending on whether the cobordism from  $\mathcal{P}(v)$  to  $\mathcal{P}(u)$  is a merge or split, respectively.

**Definition 2.1.** Fix a sign assignment s on the cube (in the sense of Section 1.1). The chain complex  $\mathcal{C}(K)$  is the totalization of A with respect to s. That is, the chain group is defined to be  $\mathcal{C}(K) = \bigoplus_{v \in \{0,1\}^n} A(v)$  and the differential  $\delta \colon \mathcal{C}(K) \to \mathcal{C}(K)$  is defined by stipulating the component  $\delta_{u,v}$  of  $\delta$  that maps from A(v) to A(u) to be

$$\delta_{u,v} = \begin{cases} (-1)^{s_{u,v}} A(\varphi_{u,v}^{\text{op}}) & \text{if } u > v \text{ and } |u| - |v| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

The Khovanov chain complex  $(\mathcal{C}_{Kh}(K), \delta_{Kh})$  is the specialization h = t = 0. The homology of  $\mathcal{C}_{Kh}(K)$  is the Khovanov homology Kh(K).

(The Khovanov complex was introduced by Khovanov [Kho00]. The specializations (h, t) = (0, 1) and (h, t) = (1, 0) were studied by Lee [Lee05] and Bar-Natan [BN05], respectively; the case of general (h, t) was studied by Khovanov [Kho06b], Naot [Nao06], and others.)

The homological grading of the summand  $A(v) \subseteq \mathcal{C}(K)$  is  $|v| - n_-$ , where  $n_-$  denotes the number of negative crossings in the link diagram. There is additionally an internal grading, called the *quantum grading*, that persists throughout; the quantum grading of any Khovanov generator in F(v) is

$$n-3n_- + |v| + \#\{\text{circles in } \mathcal{P}(v) \text{ labeled } x_+\} - \#\{\text{circles in } \mathcal{P}(v) \text{ labeled } x_-\}.$$

The quantum gradings of the formal variables (h, t) are (-2, -4).

In the presence of a basepoint on the link diagram, and after setting t = 0, there is also a reduced theory. In the reduced theory, for any  $v \in \{0,1\}^n$ , we only consider the Khovanov generators in F(v) that label the pointed circle in  $\mathcal{P}(v)$  as  $x_-$ , and shift quantum gradings up by 1. The reduced Khovanov chain complex and the reduced Khovanov homology are denoted  $\widetilde{C}_{Kh}$  and  $\widetilde{Kh}$ , respectively. (Alternatively, we could have defined a reduced theory by setting t = 0 and only considering the Khovanov generators that label the pointed circle as  $x_+$ , and shifting quantum gradings down by 1. When h = t = 0, these two reduced theories are canonically isomorphic; see also [Wig16].)

The Khovanov homology is an invariant of the link, not just of the link diagram, and the reduced Khovanov homology is an invariant of the pointed link. More generally, the chain homotopy type of the chain complex  $\mathcal{C}(K)$  over  $\mathbb{Z}[h,t]$  is a link invariant as well.

### 3. Cubical flow categories

The Khovanov stable homotopy type is defined using an auxiliary object, the Khovanov flow category. We review flow categories in Section 3.2, after first recalling some notions related to manifolds with corners, on which flow categories are based. The Khovanov flow category from [LS14a] is defined as a kind of cover of another flow category, the cube flow category (Definition 3.16), which in turn is based on permutohedra, a family of polytopes reviewed in Section 3.3. After reviewing the cube flow category and some of its basic

properties in Section 3.4, new material begins in Section 3.5, where we abstract the notion of a cover of the cube flow category, into a cubical flow category (Definition 3.21). In Section 3.6 we give a slightly different notion of neat embeddings for cubical flow categories. Using this notion, in Section 3.7 we give a realization procedure for a cubical flow category, the cubical realization. In Section 3.8 we show that the cubical realization and the Cohen-Jones-Segal realization (reviewed in Section 3.2) give homotopy equivalent spaces.

3.1. Manifolds with corners and  $\langle n \rangle$ -manifolds. The original construction of the Khovanov stable homotopy type from [LS14a] relies on Cohen-Jones-Segal's notion of flow categories [CJS95], which in turn uses a particular notion of manifolds with corners, called  $\langle n \rangle$ -manifolds [Jän68, Lau00]. For the reader's convenience, we review the relevant definitions here.

**Definition 3.1.** A k-dimensional manifold with corners is a topological space X together with an atlas  $\{(U_{\alpha}, \phi_{\alpha} \colon U_{\alpha} \to (\mathbb{R}_{+})^{k})\}$  modeled on open subsets of  $(\mathbb{R}_{+})^{k}$ , so that the transition functions are smooth. Given a point x in a chart  $(U, \phi)$  let c(x) be the number of coordinates in  $\phi(x)$  which are 0; c(x) is independent of the choice of chart. The codimension-i boundary of X is  $\{x \in X \mid c(x) = i\}$ . A k-dimensional manifold with corners X has a well-defined tangent space TX, which is an  $\mathbb{R}^{k}$ -plane bundle; a Riemannian metric on X means a Riemannian metric on TX.

**Definition 3.2.** A facet of X is the closure of a connected component of the codimension-1 boundary of X. (If X is a polytope then this agrees with the usual definition of facets.) A multifacet of X is a (possibly empty) union of disjoint facets of X. A manifold with corners X is a multifaceted manifold if every  $x \in X$  belongs to exactly c(x) facets of X. An  $\langle n \rangle$ -manifold is a multifaceted manifold X along with an ordered n-tuple  $(\partial_1 X, \ldots, \partial_n X)$  of multifacets of X such that:  $\bigcup_i \partial_i X = \partial X$ ; and for all distinct  $i, j, \partial_i X \cap \partial_j X$  is a multifacet of both  $\partial_i X$  and  $\partial_j X$ . (The number n need not be the dimension of X.) See Laures [Lau00] for more details. (Laures uses the terms 'connected face', 'face', and 'manifold with faces' for 'facet', 'multifacet', and 'multifaceted manifold', respectively. We have changed the terminology since 'face' means something different for polytopes in Section 3.3.) Given a  $\langle n \rangle$ -manifold X and a vector  $v \in \{0,1\}^n$  let  $X(v) = \bigcap_{v_i=0} \partial_i X$ , with the convention that  $X(\vec{1}) = X$ .

Example 3.3. An n-gon (polygon with n sides) is a multifaceted manifold if n > 1, while a 1-gon (disk with one corner on the boundary) is a manifold with corners but not a multifaceted manifold. Only the 2n gons can be made into  $\langle 2 \rangle$ -manifolds, though (2n+1)-gons can be viewed as 2-dimensional  $\langle 3 \rangle$ -manifolds. Of the Platonic solids only the tetrahedron, cube, and dodecahedron are manifolds with corners, and all three are multifaceted manifolds. The cube can be made into a  $\langle 3 \rangle$ -manifold by defining  $\partial_1 X$  to be the front and back facets,  $\partial_2 X$  to be the top and bottom facets, and  $\partial_3 X$  to be the left and right facets. The tetrahedron and dodecahedron cannot be given the structure of  $\langle 3 \rangle$ -manifolds, although both can be made into  $\langle 4 \rangle$ -manifolds. An even more fundamental example is  $(\mathbb{R}_+)^n$  itself, which is a  $\langle n \rangle$ -manifold by setting  $\partial_i (\mathbb{R}_+)^n = \{v \in \mathbb{R}_+^n \mid v_i = 0\}$ . Similarly,  $\mathbb{R}^N \times (\mathbb{R}_+^n)$  is an (n + N)-dimensional  $\langle n \rangle$ -manifold.

**Construction 3.4.** Given an  $\langle n \rangle$ -manifold X and an  $\langle m \rangle$ -manifold Y, the product  $X \times Y$  inherits the structure of a  $\langle n+m \rangle$ -manifold, by declaring

$$\partial_i(X \times Y) = \begin{cases} (\partial_i X) \times Y & 1 \le i \le n \\ X \times (\partial_{i-n} Y) & n+1 \le i \le n+m. \end{cases}$$

We end this subsection with the definition of neat embeddings.

**Definition 3.5.** Consider  $\langle n \rangle$ -manifolds X and Y and fix a Riemannian metric on Y. A neat embedding of X into Y is a smooth map  $f: X \to Y$  so that:

- $f^{-1}(Y(v)) = X(v)$  for all  $v \in \{0, 1\}^n$ .
- $f|_{X(v)}: X(v) \to Y(v)$  is an embedding for each  $v \in \{0,1\}^n$ .
- For all  $w < v \in \{0,1\}^n$ , f(X(v)) is perpendicular to Y(w) with respect to the Riemannian metric on Y, and in particular is transverse to Y(w).
- 3.2. Flow categories. Next we recall some notions about flow categories, from Cohen-Jones-Segal [CJS95] (see also LS14a, Section 3), partly to fix terminology for this paper.

**Definition 3.6.** A flow category  $\mathscr{C}$  is a topological category whose objects  $Ob(\mathscr{C})$  form a discrete space, equipped with a grading function gr:  $Ob(\mathscr{C}) \to \mathbb{Z}$ , and whose morphism spaces satisfy the following conditions:

- (FC-1) For any  $x \in Ob(\mathscr{C})$ ,  $Hom(x, x) = \{Id\}$ .
- (FC-2) For distinct  $x, y \in Ob(\mathscr{C})$  with gr(x) gr(y) = k, Hom(x, y) is a (possibly empty) compact (k-1)dimensional  $\langle k-1 \rangle$ -manifold; and
- (FC-3) The composition maps combine to produce a diffeomorphism of  $\langle k-2 \rangle$ -manifolds

aps combine to produce a diffeomorphism of 
$$\langle k-2 \rangle$$
-
$$\coprod_{\substack{z \in \mathrm{Ob}(\mathscr{C}) \setminus \{x,y\} \\ \mathrm{gr}(z) - \mathrm{gr}(y) = i}} \mathrm{Hom}(z,y) \times \mathrm{Hom}(x,z) \cong \partial_i \, \mathrm{Hom}(x,y).$$

(In [CJS95], it is not required that the space of objects be discrete. In Morse theory, this corresponds to allowing Morse-Bott functions.)

The identity morphisms in a flow category are somewhat special, and it is often convenient to ignore them. So, for objects x, y in  $\mathscr{C}$ , the moduli space from x to y,  $\mathcal{M}(x, y)$ , is defined to be  $\mathrm{Hom}(x, y)$  if  $x \neq y$ , and empty if x = y. (In Morse theory, this corresponds to the moduli space of non-constant downwards gradient flows from x to y.)

For any flow category  $\mathscr{C}$ , let  $\Sigma^k\mathscr{C}$  denote the flow category obtained by increasing the gradings of each object by k.

**Definition 3.7.** For each integer i, fix an integer  $D_i \ge 0$ , and let **D** denote this sequence. A neat embedding of a flow category  $\mathscr{C}$  relative to **D** is a collection  $j_{x,y}$  of neat embeddings (with the standard Riemannian metric on the target space)

$$j_{x,y} \colon \mathcal{M}(x,y) \hookrightarrow \mathbb{R}^{D_{\mathrm{gr}(y)}} \times \mathbb{R}_+ \times \mathbb{R}^{D_{\mathrm{gr}(y)+1}} \times \mathbb{R}_+ \times \cdots \times \mathbb{R}_+ \times \mathbb{R}^{D_{\mathrm{gr}(x)-1}}$$

of  $\langle \operatorname{gr}(x) - \operatorname{gr}(y) - 1 \rangle$ -manifolds for all  $x, y \in \operatorname{Ob}(\mathscr{C})$ , subject to the following:

(1) For all integers i, j,

$$\coprod_{\substack{x,y\\ \operatorname{gr}(x)=i,\operatorname{gr}(y)=j}} \jmath_{x,y} : \coprod_{\substack{x,y\\ \operatorname{gr}(x)=i,\operatorname{gr}(y)=j}} \mathcal{M}(x,y) \to \mathbb{R}^{D_j} \times \mathbb{R}_+ \times \cdots \times \mathbb{R}_+ \times \mathbb{R}^{D_{i-1}}$$

is a neat embedding of (i - j - 1)-manifolds.

(2) For all  $x, y, z \in \text{Ob}(\mathscr{C})$ , and all points  $(q, p) \in \mathcal{M}(y, z) \times \mathcal{M}(x, y)$ 

$$j_{x,z}(q \circ p) = (j_{y,z}(q), 0, j_{x,y}(p)).$$

A coherent framing for a neat embedding j is a collection of framings of the normal bundles  $\nu_{j_{x,y}}$  of  $j_{x,y}$  for all  $x, y \in \text{Ob}(\mathscr{C})$ , such that for all  $x, y, z \in \text{Ob}(\mathscr{C})$ , the product framing of  $\nu_{\jmath_{y,z}} \times \nu_{\jmath_{x,y}}$  equals the pullback framing of  $\circ^* \nu_{j_{x,z}}$ , where  $\circ$  denotes composition.

**Definition 3.8** ([LS14a, Definition 3.21]). A framed flow category is a flow category  $\mathscr{C}$ , along with a coherent framing for some neat embedding of  $\mathscr{C}$  (relative to some  $\mathbf{D}$ ).

For a framed flow category, there is an associated cochain complex  $C^*(\mathscr{C})$ , defined as follows. The  $n^{\text{th}}$  chain group  $C^n$  is the  $\mathbb{Z}$ -module freely generated by the objects of  $\mathscr{C}$  of grading n. The differential  $\delta$  is of degree one. For  $x, y \in \text{Ob}(\mathscr{C})$  with gr(x) - gr(y) = 1, the coefficient  $\langle \delta y, x \rangle$  of x in  $\delta(y)$  is the number of points in  $\mathcal{M}(x, y)$ , counted with sign. We say a framed flow category refines its associated chain complex.

Remark 3.9. In order to define the associated chain complex, one only needs the framing of the 0-dimensional moduli spaces; and in order to check that  $\delta^2 = 0$ , one only needs to ensure that the framing extends to the 1-dimensional moduli spaces.

To a framed flow category, Cohen-Jones-Segal associate a based CW complex  $|\mathcal{C}|$  whose cells (except the basepoint) correspond to the objects of the flow category [CJS95]. The following formulation of the Cohen-Jones-Segal construction is described in more detail in [LS14a, Definition 3.24].

**Definition 3.10.** Let  $\mathscr{C}$  be a framed flow category with a neat embedding  $\jmath$  relative to some  $\mathbf{D}$ , and assume all objects of  $\mathscr{C}$  have grading in [B,A] for some fixed  $A,B\in\mathbb{Z}$ . Using the framing of  $\nu_{\jmath_{x,y}}$ , extend  $\jmath_{x,y}$  to a map

$$\bar{\jmath}_{x,y} \colon \mathcal{M}(x,y) \times [-\delta, \delta]^{D_{\operatorname{gr}(y)} + \dots + D_{\operatorname{gr}(x) - 1}} \to \mathbb{R}^{D_{\operatorname{gr}(y)}} \times \mathbb{R}_+ \times \dots \times \mathbb{R}_+ \times \mathbb{R}^{D_{\operatorname{gr}(x) - 1}}.$$

Choose  $\delta$  small enough and T large enough so that the map  $\coprod_{x,y|gr(x)=i,gr(y)=j} \overline{\jmath}_{x,y}$  is an embedding into  $(-T,T)^{D_j} \times [0,T) \times \cdots \times [0,T) \times (-T,T)^{D_{i-1}}$  for all integers i,j. In the based CW complex  $|\mathscr{C}|$ , the cell associated to an object x of grading m is

$$C(x) = \prod_{i=B}^{m-1} ([0,T] \times [-T,T]^{D_i}) \times [-\delta, \delta]^{D_m + \dots + D_{A-1}}.$$

For any other object y with gr(y) = n < m, the embedding  $\bar{\jmath}_{x,y}$  identifies  $C(y) \times \mathcal{M}(x,y)$  with the following subset of  $\partial C(x)$ ,

$$C_y(x) = \prod_{i=B}^{n-1} \left( [0,T] \times [-T,T]^{D_i} \right) \times \{0\} \times \operatorname{im}(\overline{\jmath}_{x,y}) \times [-\delta,\delta]^{D_m + \dots + D_{A-1}}.$$

The attaching map for C(x) sends  $C_y(x) \cong C(y) \times \mathcal{M}(x,y)$  via the projection map to C(y), and sends  $\partial C(x) \setminus \bigcup_y C_y(x)$  to the basepoint.

**Lemma 3.11.** [LS14a, Lemma 3.25] Definition 3.10 defines a CW complex, whose reduced cellular cochain complex is isomorphic (after shifting the gradings down by  $D_B + \cdots + D_{A-1} - B$ ) to the chain complex associated to the framed flow category  $\mathscr C$  from Definition 3.8. The isomorphism sends cells of  $|\mathscr C|$  to the corresponding objects of  $\mathscr C$ .

**Definition 3.12.** The Cohen-Jones-Segal realization of  $\mathscr C$  is the formal (de)suspension  $\Sigma^{B-D_B-\cdots-D_{A-1}}|\mathscr C|$ .

3.3. **Permutohedra.** We will be interested in a particular family of flow categories, in which the moduli spaces are unions of permutohedra. So, we recall some basic facts about permutohedra.

Before starting, let us fix some notations about polytopes, mostly following Ziegler [Zie95]. Let  $P \subset \mathbb{R}^n$  be a polytope or an  $\mathcal{H}$ -polyhedron (as described in [Zie95, Definition 0.1]). If there is an affine half-space of  $\mathbb{R}^n$  which contains P, then the intersection of its boundary with P is called a face of P; and if  $\dim(P) = d$ , then we declare the entire polytope P to be its unique d-dimensional face. This gives a CW complex structure on P, with the cells being the nonempty faces. The faces of dimension 0, 1, and (d-1) are called vertices, edges, and facets, respectively. A d-dimensional polytope is called simple if every vertex is contained in exactly d facets; simple polytopes are multifaceted manifolds.

For  $\sigma \in \mathcal{S}_n$  a permutation, let  $v_{\sigma} = (\sigma^{-1}(1), \dots, \sigma^{-1}(n)) \in \mathbb{R}^n$ . The (n-1)-dimensional permutohedron  $\Pi_{n-1}$  is the convex hull in  $\mathbb{R}^n$  of the n! points  $v_{\sigma}$  [Zie95, Example 0.10]. The permutohedron  $\Pi_{n-1}$  lies in

the affine subspace  $\mathbb{A}^{n-1} := \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid \sum_i x_i = n(n+1)/2\}$  of  $\mathbb{R}^n$ . As its name suggests,  $\Pi_{n-1}$  is (n-1)-dimensional, and the  $v_{\sigma}$  are its vertices.

For each non-empty, proper subset S of  $\{1,\ldots,n\}$  of cardinality say k, let  $H_S\subset\mathbb{A}^{n-1}\subset\mathbb{R}^n$  be the half-space  $\{(x_1,\ldots,x_n)\in\mathbb{A}^{n-1}\mid\sum_{i\in S}x_i\geq k(k+1)/2\}$ . The permutohedron  $\Pi_{n-1}$  can also be defined as the intersection of the  $2^n-2$  half-spaces  $H_S$ . In fact, the facets of  $\Pi_{n-1}$  are exactly the  $F_S:=\Pi_{n-1}\cap\partial H_S$ .

The facets  $F_S$  are identified with products of lower-dimensional permutohedra:

**Lemma 3.13.** Let  $a_1 < a_2 < \cdots < a_k$  be the elements in S, and let  $b_1 < b_2 < \cdots < b_{n-k}$  be the elements in  $\{1, 2, \ldots, n\} \setminus S$ . Then the map  $f_S : \mathbb{R}^n \to \mathbb{R}^k \times \mathbb{R}^{n-k}$ 

$$f_S(x_1,\ldots,x_n)=((x_{a_1},\ldots,x_{a_k}),(x_{b_1}-k,\ldots,x_{b_{n-k}}-k))$$

identifies the facet  $F_S \subset \mathbb{R}^n$  with  $\Pi_{k-1} \times \Pi_{n-1-k} \subset \mathbb{R}^k \times \mathbb{R}^{n-k}$ .

*Proof.* It suffices to show that  $f_S$  takes the vertices of  $F_S \subset \Pi_{n-1}$  to the vertices of  $\Pi_{k-1} \times \Pi_{n-1-k}$ . The vertices of  $F_S$  are the points  $(x_1, \ldots, x_n)$  so that  $\{x_{a_1}, \ldots, x_{a_k}\} = \{1, \ldots, k\}$  and  $\{x_{b_1}, \ldots, x_{b_{n-k}}\} = \{k+1, \ldots, n\}$ . It is immediate that  $f_S$  takes these vertices bijectively to the vertices of  $\Pi_{k-1} \times \Pi_{n-1-k}$ .  $\square$ 

The permutohedron  $\Pi_{n-1}$  is simple, i.e., each vertex lies in exactly n-1 facets:  $v_{\sigma}$  lies in the facet  $F_{\{\sigma(1),\ldots,\sigma(k)\}}$  for each  $1 \leq k < n$ , and no others. Therefore, every d-dimensional face belongs to exactly n-1-d facets; and the subsets corresponding to those facets are nested. Hence, d-dimensional faces correspond to sequences of n-1-d nested proper, non-empty subsets of  $\{1,\ldots,n\}$ . Further:

**Lemma 3.14.** The space  $\Pi_{n-1}$  can be treated as an (n-1)-manifold by declaring

$$\partial_i \Pi_{n-1} = \bigcup_{\substack{S \\ |S|=i}} F_S$$

for  $1 \le i \le n - 1$ .

*Proof.* We must check:

- (1) Every point x belongs to c(x) facets.
- (2) Each  $\partial_i \Pi_{n-1}$  is a multifacet, that is, a disjoint union of facets.
- (3)  $\bigcup_{i} \partial_{i} \Pi_{n-1} = \partial \Pi_{n-1}$ .
- (4) For each  $i \neq j$ ,  $\partial_i \Pi_{n-1} \cap \partial_j \Pi_{n-1}$  is a multifacet of  $\partial_i \Pi_{n-1}$  (and  $\partial_j \Pi_{n-1}$ ).

Point (1) follows from the fact that  $\Pi_{n-1}$  is a simple polyhedron.

For point (2), we claim that if |S| = |T| = i and  $S \neq T$  then  $F_S \cap F_T = \emptyset$ ; it follows that  $\partial_i \Pi_{n-1}$  is the disjoint union of the facets  $F_S$  (with |S| = i). But if  $v_\sigma$  is a vertex in  $F_S \cap F_T$ , then

$$\sum_{j \in S} \sigma^{-1}(j) = \sum_{j \in T} \sigma^{-1}(j) = i(i+1)/2$$

$$\iff \{\sigma^{-1}(j) \mid j \in S\} = \{\sigma^{-1}(j) \mid j \in T\} = \{1, \dots, i\}$$

$$\iff S = T = \{\sigma(1), \dots, \sigma(i)\}.$$

Point (3) is immediate from the definitions.

For point (4), suppose that |S| = i. After identifying  $F_S$  with  $\Pi_{i-1} \times \Pi_{n-i-1}$  using Lemma 3.13, we get

$$F_S \cap \partial_j \Pi_{n-1} = \begin{cases} \Pi_{i-1} \times (\partial_{j-i} \Pi_{n-i-1}) & i < j \\ (\partial_j \Pi_{i-1}) \times \Pi_{n-i-1} & i > j. \end{cases}$$

Therefore,  $F_S \cap \partial_j \Pi_{n-1}$  is a disjoint union of facets of  $F_S \cong \Pi_{i-1} \times \Pi_{n-i-1}$ . Since the  $F_S$  for |S| = i are disjoint,  $\partial_i \Pi_{n-1} \cap \partial_j \Pi_{n-1}$  is a disjoint union of facets of  $\partial_i \Pi_{n-1}$  as well.

We will also use the following well-known cubical complex structure on  $\Pi_{n-1}$ . For any permutation  $\sigma \in \mathcal{S}_n$ , let  $C_{\sigma}$  be the convex hull of the barycenters of all the faces that contain the vertex  $v_{\sigma}$ .

**Lemma 3.15.** Each  $C_{\sigma}$  is combinatorially equivalent to an (n-1)-dimensional cube, and these cubes form a cubical complex subdivision of  $\Pi_{n-1}$ .

*Proof.* We present the proof from Ovchinnikov [Ovc08, Section 3], for which he cites Ziegler. Consider the following intersection of half-spaces

$$FC_{\sigma} = \bigcap_{S|v_{\sigma} \in F_S} H_S.$$

The space  $FC_{\sigma}$  is an (n-1)-dimensional cone with cone point  $v_{\sigma}$ , with n-1 facets (corresponding to the facets of  $\Pi_{n-1}$  that contain  $v_{\sigma}$ ), and therefore is a simplicial cone. (For comparison with [Ovc08, Section 3],  $FC_{\sigma}$  is the dual of the facet cone of the dual of  $v_{\sigma}$  (in the face fan of the dual of  $\Pi_{n-1}$ ).)

Next, consider the vertex cone of  $v_{\sigma}$  (in the normal fan of  $\Pi_{n-1}$ ),  $VC_{\sigma}$ . By definition, the cone point of  $VC_{\sigma}$  is the barycenter of  $\Pi_{n-1}$ , and the edges of  $VC_{\sigma}$  are obtained by dropping perpendiculars from the barycenter to the facets of  $\Pi_{n-1}$  that contain  $v_{\sigma}$ . The cone  $VC_{\sigma}$  is an (n-1)-dimensional cone with (n-1) edges, and therefore,  $VC_{\sigma}$  is a simplicial cone. The d-dimensional faces of  $VC_{\sigma}$  correspond to the (n-1-d)-dimensional faces of  $\Pi_{n-1}$  that contain  $v_{\sigma}$ . Given corresponding faces  $f_{VC}$  of  $VC_{\sigma}$  and f of  $\Pi_{n-1}$ ,  $f_{VC}$  is perpendicular to f and passes through the barycenter of f.

Therefore,  $C_{\sigma}$  is the intersection of the two simplicial cones  $FC_{\sigma}$  and  $VC_{\sigma}$ . Since the edges of  $VC_{\sigma}$  pass through the interiors of the facets of  $FC_{\sigma}$ ,  $VC_{\sigma}$  and  $FC_{\sigma}$  intersect transversely, and therefore,  $VC_{\sigma} \cap FC_{\sigma}$  is combinatorially equivalent to a cube.

The facets of  $C_{\sigma}$  are of two types: the ones contained in  $FC_{\sigma}$ , which are not identified with the facets of any other cube and lie in the boundary of  $\Pi_{n-1}$ ; and the ones contained in  $VC_{\sigma}$ , which are identified with facets of other cubes and lie in the interior of  $\Pi_{n-1}$ . Indeed the facets of the latter type correspond to the edges e of  $\Pi_{n-1}$  that contain  $v_{\sigma}$ : the facet corresponding to e is formed by taking the convex hull of the barycenters of all the faces of  $\Pi_{n-1}$  that contain e. With these identifications, it is clear that these cubes  $C_{\sigma}$  come together to form a cubical subdivision of the permutohedron.

3.4. The cube flow category. We start by recalling the cube flow category from [LS14a, Definition 4.1]. There we gave a definition in terms of Morse flows on  $[0,1]^n$ . Here, we give a more directly combinatorial definition in terms of permutohedra (see also Remark 3.18):

**Definition 3.16.** The objects of the cube flow category  $\mathscr{C}_C(n)$  are the same as the objects of the cube category  $\underline{2}^n$ , i.e., tuples  $u = (u_1, \ldots, u_n) \in \{0, 1\}^n$ . The grading on the objects is defined by  $gr(u) = |u| = \sum_i u_i$ .

The space  $\mathcal{M}(u,v)$  is defined to be empty unless u > v. If u > v and gr(u) - gr(v) = k > 0 then we define  $\mathcal{M}(u,v) = \Pi_{k-1}$ , the (k-1)-dimensional permutohedron. Note that, by Lemma 3.14,  $\mathcal{M}(u,v)$  is a  $\langle k-1 \rangle$ -manifold.

The composition map  $\mathcal{M}(v,w) \times \mathcal{M}(u,v) \to \mathcal{M}(u,w)$  is defined as follows. Assume that u > v > w,  $\operatorname{gr}(u) - \operatorname{gr}(v) = k$  and  $\operatorname{gr}(v) - \operatorname{gr}(w) = l$ . Suppose that  $u_{a_1} = \cdots = u_{a_{k+l}} = 1$  and  $w_{a_1} = \cdots = w_{a_{k+l}} = 0$  (i.e., the  $a_i$  are the coordinates in which u and w differ), where  $a_1 < a_2 < \cdots < a_{k+l}$ . Let S be the subset of  $\{1, \ldots, k+l\}$  such that  $v_{a_s} = 1$  for  $s \in S$ . (The set S has cardinality l.) By Lemma 3.13, there is a corresponding facet  $F_S \subset \Pi_{k+l-1}$ , and  $F_S$  is identified with  $\Pi_{l-1} \times \Pi_{k-1} = \mathcal{M}(v,w) \times \mathcal{M}(u,v)$ . The composition map is the corresponding inclusion map  $\mathcal{M}(v,w) \times \mathcal{M}(u,v) = F_S \hookrightarrow \mathcal{M}(u,w)$ .

**Lemma 3.17.** Definition 3.16 defines a flow category.

*Proof.* Conditions (FC-1) and (FC-2) of Definition 3.6 are immediate from the definitions and Lemma 3.14. For Condition (FC-3), it is enough to recall from Lemma 3.14 that  $\partial_l \Pi_{k+l-1} = \bigcup_{|S|=l} F_S$ .

Finally, we need to check that this defines a category, or in other words, that composition is associative. Towards this end, for any u>v with  $\operatorname{gr}(u)-\operatorname{gr}(v)=k>0$ , it is convenient to treat  $\mathcal{M}(u,v)=\Pi_{k-1}$  as a subset of  $\prod_{i|u_i>v_i}\mathbb{R}$  instead of  $\mathbb{R}^k$ , where the two ambient spaces are identified by linearly ordering the coordinates  $a_1< a_2<\cdots< a_k$  in which u and v differ. With this viewpoint, for u>v>w, with  $\operatorname{gr}(u)-\operatorname{gr}(v)=k$  and  $\operatorname{gr}(v)-\operatorname{gr}(w)=l$ , the composition map  $\Pi_{l-1}\times\Pi_{k-1}\to\Pi_{k+l-1}$  is induced from the map

$$\prod_{i \mid v_i > w_i} \mathbb{R} \times \prod_{i \mid u_i > v_i} \mathbb{R} \ \xrightarrow{+(\vec{0}, \vec{l})} \prod_{i \mid v_i > w_i} \mathbb{R} \times \prod_{i \mid u_i > v_i} \mathbb{R} \ \xrightarrow{\cong} \prod_{i \mid u_i > w_i} \mathbb{R},$$

where the first map adds l to each of the coordinates of  $\prod_{i|u_i>v_i}\mathbb{R}$ , and the second map is coordinate-wise identification. (See also Lemma 3.13.) Now, given u>v>w>x, with gr(u)-gr(v)=k, gr(v)-gr(w)=l and gr(w)-gr(x)=m, the corresponding double compositions are:

$$\begin{split} \prod_{i|w_{i}>x_{i}} \mathbb{R} \times \prod_{i|v_{i}>w_{i}} \mathbb{R} \times \prod_{i|u_{i}>v_{i}} \mathbb{R} \xrightarrow{+(\vec{0},\vec{0},\vec{l})} & \prod_{i|w_{i}>x_{i}} \mathbb{R} \times \prod_{i|u_{i}>w_{i}} \mathbb{R} \\ & \downarrow +(\vec{0},\vec{m},\vec{0}) & \downarrow +(\vec{0},\vec{m}) \\ \prod_{i|v_{i}>x_{i}} \mathbb{R} \times \prod_{i|u_{i}>v_{i}} \mathbb{R} \xrightarrow{+(\vec{0},\vec{l}+\vec{m})} & \prod_{i|u_{i}>x_{i}} \mathbb{R}, \end{split}$$

where we have suppressed the reshuffling of factors from the notation. So, both compositions are given by  $+(\vec{0}, \vec{m}, \vec{l} + \vec{m})$  (and the same reshuffling of factors).

Remark 3.18. We have not shown that the definition of the cube flow category from Definition 3.16 agrees with the Morse-theoretic definition from [LS14a, Definition 4.1], as doing so would seem to require a nontrivial digression about the smooth structures on moduli spaces of Morse flows. For the purposes of this paper (and future work), note that the combinatorial definition used here works just as well for the construction of the Khovanov homotopy type in [LS14a], and all the results stated in [LS14a, LS14c, LS14b] remains true with this combinatorial definition. To be more precise, the only statements that involve the moduli spaces on the cube flow category coming the Morse flows are [LS14a, Lemmas 4.2–4.3], which are immediate for the combinatorial definition. Therefore, when we talk about the Khovanov homotopy type in this paper (and future work), we mean the homotopy type defined using the cube flow category from Definition 3.16.

Before moving on to the definition of cubical flow categories in Section 3.5, we digress a little to study the cubical complex structure from Lemma 3.15 on the permutohedron  $\mathcal{M}(u, v)$ .

**Definition 3.19.** For u > v in  $\{0,1\}^n$ , define the space

$$M_{u,v} = \left( \coprod_{\substack{m \in \{1, \dots, |u| - |v|\} \\ \dots, 0 > \dots, m-1}} [0, 1]^{m-1} \right) / \sim$$

where, for each chain  $u=u^0>\cdots>u^m=v$ , each  $1\leq i\leq m-1$ , and each point  $(t_1,\ldots,t_{m-2})$  in the cube  $[0,1]^{m-2}$  corresponding to the chain  $u^0>\cdots>u^{i-1}>u^{i+1}>\cdots>u^m$  the equivalence relation  $\sim$  identifies

$$(t_1,\ldots,t_{m-2})_{u=u^0>\cdots>u^{i-1}>u^{i+1}>\cdots>u^m=v} \sim (t_1,\ldots,t_{i-1},1,t_i,\ldots,t_{m-2})_{u=u^0>\cdots>u^m=v}.$$




FIGURE 3.1. A cubical decomposition of the permutohedron. Left: The 2-dimensional permutohedron  $\mathcal{M}(111,000)$  with vertices labeled by the corresponding permutations of (1,2,3). Right: The corresponding cubical complex  $M_{111,000}$ , where we have labeled the cube corresponding to a sequence  $111 = u^0 > \cdots > u^m = 000$  by  $(u^1, \ldots, u^{m-1})$ .

For u > v > w in  $\{0,1\}^n$ , define a map  $M_{v,w} \times M_{u,v} \to M_{u,w}$  by:

$$((t_1, \dots, t_{\ell-1})_{v=v^0 > \dots > v^\ell = w}, (s_1, \dots, s_{m-1})_{u=u^0 > \dots > u^m = v})$$

$$\mapsto (s_1, \dots, s_{m-1}, 0, t_1, \dots, t_{\ell-1})_{u=u^0 > \dots > u^m = v = v^0 > \dots, v^\ell = w}.$$

**Lemma 3.20.** There are homeomorphisms  $h_{u,v}: M_{u,v} \xrightarrow{\cong} \mathcal{M}(u,v)$  from the spaces of Definition 3.19 to the permutohedra  $\mathcal{M}(u,v)$  so that:

- (1) The  $h_{u,v}$  identify the cubes in the definition of  $M_{u,v}$  with the cubes in the cubical complex structure on  $\mathcal{M}(u,v)$  from Lemma 3.15 which are not entirely contained in  $\partial \mathcal{M}(u,v)$ .
- (2) The  $h_{u,v}$  identify the points in the cubes for  $M_{u,v}$  where some coordinate is 0 with the points in  $\partial \mathcal{M}(u,v)$ .
- (3) For any u > v > w, the following diagram commutes:

$$M_{v,w} \times M_{u,v} \xrightarrow{\cong} \mathcal{M}(v,w) \times \mathcal{M}(u,v)$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$M_{u,w} \xrightarrow{\cong} \mathcal{M}(u,w).$$

Here, the left vertical arrow is the map from Definition 3.19, and the right vertical arrow is the composition in  $\mathcal{C}_{C}(n)$ .

(See Figure 3.1.)

Proof. The chain  $c = \{u = u^0 > \cdots > u^m = v\}$  in  $\{0,1\}^n$  corresponds to the (|u| - |v| - m)-dimensional face  $F_c = \mathcal{M}(u^{m-1}, u^m) \times \cdots \times \mathcal{M}(u^0, u^1)$  of the permutohedron  $\mathcal{M}(u, v)$ . If we take the convex hull of the barycenters of all the faces of the permutohedron that contain  $F_c$ , we get an (m-1)-dimensional cube  $C_c$  which appears in the cubical complex structure from Lemma 3.15. (If  $F_c$  is a vertex of the permutohedron,

or equivalently if c is a maximal chain, then the cube  $C_c$  is one of the  $C_\sigma$  from Lemma 3.15.) We will identify  $C_c$  with the cube  $[0,1]^{m-1}$  corresponding to c in  $M_{u,v}$ .

Let  $t_1, \ldots, t_{m-1}$  be the coordinates of  $[0,1]^{m-1}$ . As a first step, we identify the vertices of  $C_c$  and  $[0,1]^{m-1}$ . A vertex of  $C_c$  corresponds to a barycenter of some face containing  $F_c$ , which in turn corresponds to some subchain c' of c; the corresponding vertex in  $[0,1]^{m-1}$  has  $t_i=0$  if  $u^i \in c'$ , and has  $t_i=1$  otherwise. The identification on the vertices leads to our desired identification as follows. Construct the simplicial complex subdivision of  $C_c$  (respectively,  $[0,1]^{m-1}$ ) by joining every face to the barycenter of  $\mathcal{M}(u,v)$  (respectively, the vertex  $\vec{1} \in [0,1]^{m-1}$ ), and extend the identification linearly over each simplex.

It is fairly straightforward to check that such identifications induce a cubical complex homeomorphism between the cubical complex  $M_{u,v}$  and the cubical complex structure on  $\mathcal{M}(u,v)$ , and these homeomorphisms satisfy the conditions of the lemma. Further details are left to the reader.

### 3.5. Definition of a cubical flow category.

**Definition 3.21.** A cubical flow category is a flow category  $\mathscr{C}$  equipped with a grading-preserving functor  $\mathfrak{f}: \Sigma^k \mathscr{C} \to \mathscr{C}_C(n)$  for some  $k \in \mathbb{Z}, n \in \mathbb{N}$  so that for each  $x, y \in \mathrm{Ob}(\mathscr{C})$ ,  $\mathfrak{f}: \mathcal{M}(x,y) \to \mathcal{M}(\mathfrak{f}(x),\mathfrak{f}(y))$  is a (trivial) covering map.

Thus, if  $\mathfrak{f}: \mathscr{C} \to \mathscr{C}_C(n)$  is a cubical flow category and  $x, y \in \mathrm{Ob}(\mathscr{C})$  then  $\mathrm{Hom}(x, y)$  is empty unless  $\mathfrak{f}(x) \geq \mathfrak{f}(y)$ . If x = y, then  $\mathrm{Hom}(x, y) = \{\mathrm{Id}\}$ , and if  $\mathfrak{f}(x) = \mathfrak{f}(y)$  but  $x \neq y$  then  $\mathrm{Hom}(x, y)$  is empty. If  $\mathfrak{f}(x) > \mathfrak{f}(y)$ , then the moduli space  $\mathcal{M}_{\mathscr{C}}(x, y) = \mathrm{Hom}(x, y)$  is a (possibly empty) disjoint union of permutohedra.

Convention 3.22. Sometimes we suppress the grading information if it is inessential to the discussion, and drop the grading shift  $\Sigma^k$  from the notation.

A framing of the cube flow category  $\mathscr{C}_C(n)$  in the sense of Definition 3.8 induces a sign assignment s on the cube (see Section 1.1) as follows: for u > v with |u| - |v| = 1,  $s_{u,v} = 0$  if the point  $\mathcal{M}(u,v)$  is framed positively, and  $s_{u,v} = 1$  otherwise. Furthermore, every sign assignment on the cube is induced from an essentially unique framing of  $\mathscr{C}_C(n)$ ; see [LS14a, Section 4.2]. The chain complex associated to  $\mathscr{C}_C(n)$ , framed according to some sign assignment s, is defined as follows. The chain group is generated by the vertices of the cube  $\{0,1\}^n$ , and differential is given by

$$\delta(v) = \sum_{\substack{u > v \\ |u| - |v| = 1}} (-1)^{s_{u,v}} u.$$

This is an acyclic chain complex, and is often referred to as the *cube chain complex*.

Furthermore, if  $(\mathscr{C}, \mathfrak{f}: \mathscr{C} \to \mathscr{C}_C(n))$  is a cubical flow category, then any sign assignment s on the cube induces a framing of the 0-dimensional moduli spaces in  $\mathscr{C}$  as well: all the points in  $\mathcal{M}_{\mathscr{C}}(x,y)$  are framed positively if  $s_{\mathfrak{f}(x),\mathfrak{f}(y)}=0$ , and otherwise, all the points in  $\mathcal{M}_{\mathscr{C}}(x,y)$  are framed negatively. The pullback of the (essentially unique) framing on  $\mathscr{C}_C(n)$  inducing s produces an essentially canonical extension of this framing to the entire cubical flow category  $\mathscr{C}$ . The chain complex associated to  $\mathscr{C}$  for this framing has the following differential: for  $x,y\in \mathrm{Ob}(\mathscr{C})$  with  $\mathrm{gr}(x)-\mathrm{gr}(y)=1$ , the coefficient of x in  $\delta y$  is

$$\langle \delta(y), x \rangle = \begin{cases} (-1)^{s_{\mathfrak{f}(x), \mathfrak{f}(y)}} & \mathfrak{f}(x) > \mathfrak{f}(y) \\ 0 & \text{otherwise.} \end{cases}$$

This chain complex only depends on the sign assignment s and not the entire framing of  $\mathscr{C}_{C}(n)$ .

Even though the definition of cubical flow categories seems fairly restrictive, there are many examples:

Example 3.23. Given any (finite) simplicial complex  $S_{\bullet}$  with vertices  $v_1, \ldots, v_n$ , there is a corresponding cubical flow category  $(\mathscr{C}, \mathfrak{f}: \Sigma\mathscr{C} \to \mathscr{C}_C(n))$ , defined as follows. The objects of  $\mathscr{C}$  correspond to the simplices

of  $S_{\bullet}$ , which in turn can be viewed as non-empty subsets of  $\{v_1, \ldots, v_n\}$ . Given an object in  $\mathscr{C}$ , corresponding to a subset  $T \subseteq \{v_1, \ldots, v_n\}$ , define  $\mathfrak{f}(T)$  to be the vector in  $\{0,1\}^n$  whose  $i^{\text{th}}$  coordinate is 1 if and only if  $v_i \in T$ . Note that the map  $\mathfrak{f}$  is injective on objects. Let  $\mathscr{C}$  be the full subcategory of  $\mathscr{C}_C(n)$  generated by the objects in the image of  $\mathfrak{f}$ . The chain complex associated to  $\mathscr{C}$  is isomorphic to the simplicial chain complex for  $S_{\bullet}$ .

(One can think of the category  $\mathscr{C}$  as coming from choosing a Morse function on each simplex in  $S_{\bullet}$  with a unique interior maximum and no other interior critical points, and so that these Morse functions are compatible under restriction. The moduli spaces in  $\mathscr{C}$  are then the corresponding Morse moduli spaces.)

Example 3.24. The Khovanov flow category [LS14a, Definition 5.3]  $\mathscr{C}_K(K)$  associated to a link diagram K (with n crossings  $c_1, \ldots, c_n$ ) is, by construction, a cubical flow category. For any  $v \in \{0, 1\}^n$ , the subset of  $\mathrm{Ob}(\mathscr{C}_K(K))$  that maps to v are precisely the Khovanov generators over v (as defined in Section 2.2):

$$\mathfrak{f}^{-1}(v) = F(v).$$

For any  $u > v \in \{0,1\}^n$  with |u| - |v| = 1, and any  $x \in \mathfrak{f}^{-1}(u) = F(u)$ ,  $y \in \mathfrak{f}^{-1}(v) = F(v)$ , the moduli space is

$$\mathcal{M}_{\mathscr{C}_K(K)}(x,y) = \begin{cases} \{ \mathrm{pt} \} & \text{if } x \text{ appears in } \delta_{Kh}(y), \\ \varnothing & \text{otherwise.} \end{cases}$$

Therefore, the chain complex associated to  $\mathscr{C}_K(K)$  is isomorphic to the Khovanov chain complex  $\mathscr{C}_{Kh}(K)$  (from Definition 2.1).

3.6. Cubical neat embeddings. Consider the cube flow category  $\mathscr{C}_C(n)$ , and fix a tuple  $\mathbf{d} = (d_0, \dots, d_{n-1}) \in \mathbb{N}^n$  and a real number R > 0. For any u > v in  $\mathrm{Ob}(\mathscr{C}_C(n)) = \{0, 1\}^n$ , let

$$E_{u,v} = \left[\prod_{i=|v|}^{|u|-1} (-R,R)^{d_i}\right] \times \mathcal{M}_{\mathscr{C}_C(n)}(u,v).$$

Equip  $E_{u,v}$  with the Riemannian metric induced from the standard metric on the Euclidean space after viewing the permutohedron  $\mathcal{M}_{\mathscr{C}_C(n)}(u,v)$  as a polyhedron in  $\mathbb{R}^{|u|-|v|}$ . For any u>v>w in  $\mathrm{Ob}(\mathscr{C}_C(n))$ , there is a map  $E_{v,w}\times E_{u,v}\to E_{u,w}$  given by:

$$E_{v,w} \times E_{u,v} = \prod_{i=|w|}^{|v|-1} (-R,R)^{d_i} \times \mathcal{M}_{\mathscr{C}_C(n)}(v,w) \times \prod_{i=|v|}^{|u|-1} (-R,R)^{d_i} \times \mathcal{M}_{\mathscr{C}_C(n)}(u,v)$$

$$\cong \prod_{i=|w|}^{|u|-1} (-R,R)^{d_i} \times \mathcal{M}_{\mathscr{C}_C(n)}(v,w) \times \mathcal{M}_{\mathscr{C}_C(n)}(u,v)$$

$$\stackrel{\mathrm{Id} \times \circ}{\longleftrightarrow} \prod_{i=|w|}^{|u|-1} (-R,R)^{d_i} \times \mathcal{M}_{\mathscr{C}_C(n)}(u,w).$$

**Definition 3.25.** Fix a cubical flow category  $(\mathscr{C}, \mathfrak{f}: \Sigma^k\mathscr{C} \to \mathscr{C}_C(n))$ . A cubical neat embedding  $\iota$  of  $(\mathscr{C}, \mathfrak{f})$  (or, more succinctly, of  $\mathscr{C}$ ) relative to a tuple  $\mathbf{d} = (d_0, \ldots, d_{n-1}) \in \mathbb{N}^n$  consists of neat embeddings

$$\iota_{x,y} \colon \mathcal{M}_{\mathscr{C}}(x,y) \hookrightarrow E_{\mathfrak{f}(x),\mathfrak{f}(y)},$$

such that:

(1) For each  $x, y \in \mathrm{Ob}(\mathscr{C}),$  the following diagram commutes:

$$\mathcal{M}_{\mathscr{C}}(x,y) \xrightarrow{\iota_{x,y}} E_{\mathfrak{f}(x),\mathfrak{f}(y)}$$

$$\downarrow^{\text{projection}}$$
 $\mathcal{M}_{\mathscr{C}_{C}(n)}(\mathfrak{f}(x),\mathfrak{f}(y)).$ 

(2) For each  $u, v \in \mathrm{Ob}(\mathscr{C}_C(n))$ , the induced map

$$\coprod_{\substack{x,y\\\mathfrak{f}(x)=u,\mathfrak{f}(y)=v}}\iota_{x,y}\colon \coprod_{\substack{x,y\\\mathfrak{f}(x)=u,\mathfrak{f}(y)=v}}\mathcal{M}_{\mathscr{C}}(x,y)\to E_{u,v}$$

is a neat embedding.

(3) For each  $x, y, z \in Ob(\mathscr{C})$ , the following diagram commutes:

$$\mathcal{M}_{\mathscr{C}}(y,z) \times \mathcal{M}_{\mathscr{C}}(x,y) \longrightarrow \mathcal{M}_{\mathscr{C}}(x,z)$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$E_{\mathfrak{f}(y),\mathfrak{f}(z)} \times E_{\mathfrak{f}(x),\mathfrak{f}(y)} \longrightarrow E_{\mathfrak{f}(x),\mathfrak{f}(z)}.$$

In order to construct the cubical realization, we need to extend these embeddings  $\iota_{x,y}$  to maps

$$\bar{\iota}_{x,y}: \left[\prod_{i=|\mathfrak{f}(x)|}^{|\mathfrak{f}(x)|-1} [-\epsilon,\epsilon]^{d_i}\right] \times \mathcal{M}_{\mathscr{C}}(x,y) \to E_{\mathfrak{f}(x),\mathfrak{f}(y)},$$

for some  $\epsilon > 0$ , so that that the analogue of the diagram from Condition (3) still commutes, and the extension of the map from Condition (2) is still an embedding. One way to choose such a family of extensions would be to coherently frame the normal bundles of the embeddings  $\iota_{x,y}$  (in a similar sense to Definition 3.7) and then use the construction from Definition 3.10. Instead, we will use the following explicit extension.

For any  $x, y \in \text{Ob}(\mathscr{C})$ , let u and v denote  $\mathfrak{f}(x)$  and  $\mathfrak{f}(y)$ , respectively, and let  $\pi_{u,v}^R$  and  $\pi_{u,v}^M$  denote the

For any  $x, y \in \mathrm{Ob}(\mathscr{C})$ , let u and v denote  $\mathfrak{f}(x)$  and  $\mathfrak{f}(y)$ , respectively, and let  $\pi^{M}_{u,v}$  and  $\pi^{M}_{u,v}$  denote the projections of  $\left[\prod_{i=|v|}^{|u|-1}(-R,R)^{d_i}\right] \times \mathcal{M}_{\mathscr{C}_{C}(n)}(u,v)$  onto the two factors. Given sufficiently small  $\epsilon > 0$ , extend the embedding  $\iota_{x,y}$  to a map

(3.26) 
$$\bar{\iota}_{x,y} \colon \left[ \prod_{i=|v|}^{|u|-1} [-\epsilon, \epsilon]^{d_i} \right] \times \mathcal{M}_{\mathscr{C}}(x,y) \to E_{u,v} = \left[ \prod_{i=|v|}^{|u|-1} (-R, R)^{d_i} \right] \times \mathcal{M}_{\mathscr{C}_{C}(n)}(u,v)$$

$$(a, \gamma) \stackrel{\bar{\iota}_{x,y}}{\mapsto} (a + \pi_{u,v}^R \iota_{x,y}(\gamma), \pi_{u,v}^M \iota_{x,y}(\gamma)).$$

The definition of  $\bar{\iota}$  ensures that the analogue of the diagram from Condition (3) still commutes. By condition 1,  $\iota(\mathcal{M}_{\mathscr{C}}(x,y))$  is transverse to the fibers  $\left[\prod_{i=|v|}^{|u|-1}(-R,R)^{d_i}\right]\times\{\gamma\}$ . So, for  $\epsilon$  sufficiently small, the extension of the map from Condition (2) is still an embedding. We make this a requirement on  $\epsilon$ :

Convention 3.27. When talking about extensions  $\bar{\iota}_{x,y}$  of cubical neat embeddings, we will always assume that  $\epsilon$  is chosen to be small in the sense that the induced map

$$\coprod_{\substack{x,y\\\mathfrak{f}(x)=u,\mathfrak{f}(y)=v\\}} \bar{\iota}_{x,y} \colon \coprod_{\substack{x,y\\\mathfrak{f}(x)=u,\mathfrak{f}(y)=v\\}} \left[ \prod_{i=|v|}^{|u|-1} [-\epsilon,\epsilon]^{d_i} \right] \times \mathcal{M}_{\mathscr{C}}(x,y) \to E_{u,v}$$

is an embedding.

Remark 3.28. In [LS14a], we used a framing of the normal bundles, rather than the kind of explicit extension above, to trivialize tubular neighborhoods. When identifying the cubical realization with the Cohen-Jones-Segal realization (see Section 3.8), we will need an isotopy between these two trivializations.

Let  $V_0 = \prod_{i=|v|}^{|u|-1} \mathbb{R}^{d_i} \times \{0\} \subset (TE_{u,v})|_{\iota_{x,y}(\mathcal{M}_{\mathscr{C}}(x,y))}$  and let  $V_1$  be the normal bundle to  $\iota_{x,y}\mathcal{M}_{\mathscr{C}}(x,y)$ . Since  $\pi^M_{u,v} \circ \iota_{x,y}$  is a covering map, the projection  $d\pi^R_{u,v} \colon V_1 \to V_0$  is an isomorphism. For  $t \in [0,1]$  let  $\pi_t \colon V_1 \to (TE_{u,v})|_{\iota_{x,y}(\mathcal{M}_{\mathscr{C}}(x,y))}$  be  $\pi_t = t \operatorname{Id} + (1-t)d\pi^R_{u,v}$  and let  $V_t = \pi_t(V_1)$ . The  $V_t$  are a 1-parameter family of subbundles connecting  $V_0$  to  $V_1$ , and each  $V_t$  is a complement to  $T(\iota_{x,y}\mathcal{M}_{\mathscr{C}}(x,y))$ .

The bundle  $V_0$  is a trivial bundle, and in particular framed. Pushing forward this framing by  $\pi_t$  gives a framing of each  $V_t$ . Exponentiating these framings gives a 1-parameter family of extensions  $\bar{\iota}_{x,y}^t$  of  $\iota_{x,y}$ , each satisfying the analogue of Condition (3). The framing  $\bar{\iota}_{x,y}^0$  is our explicit extension  $\bar{\iota}_{x,y}$  from Equation (3.26), and  $\bar{\iota}_{x,y}^1$  is an extension coming from coherently framing the normal bundles of  $\iota_{x,y}$ . Since each  $V_t$  is a complement to  $T(\iota_{x,y}\mathcal{M}_{\mathscr{C}}(x,y))$ , each  $\bar{\iota}_{x,y}^t$  satisfies the analog of Condition (2) for some  $\epsilon_t$ ; compactness allows us to choose a uniform  $\epsilon$  for which each  $\bar{\iota}_{x,y}^t$  satisfies the analog of Condition (2). This produces the required isotopy between the extension from Equation (3.26) and an extension coming from some framing of the normal bundle.

### 3.7. Cubical realization.

**Definition 3.29.** Fix a cubical neat embedding  $\iota$  of a cubical flow category  $(\mathscr{C}, \mathfrak{f}: \Sigma^k \mathscr{C} \to \mathscr{C}_C(n))$ , relative to a tuple  $\mathbf{d} = (d_0, \ldots, d_{n-1}) \in \mathbb{N}^d$ , and fix  $\epsilon > 0$  satisfying Convention 3.27. We build a CW complex  $\|\mathscr{C}\| = \|\mathscr{C}\|_{\mathfrak{f}, \iota}$  from this data as follows:

• The CW complex has one cell for each  $x \in \text{Ob}(\mathscr{C})$ . Letting u denote  $\mathfrak{f}(x)$ , this cell is given by

$$\mathcal{C}(x) = \prod_{i=0}^{|u|-1} [-R, R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon, \epsilon]^{d_i} \times \widetilde{\mathcal{M}}_{\mathscr{C}_C(n)}(u, \vec{0}),$$

where  $\widetilde{\mathcal{M}}_{\mathscr{C}_{G}(n)}(u,\vec{0})$  is defined to be  $[0,1] \times \mathcal{M}_{\mathscr{C}_{G}(n)}(u,\vec{0})$  if  $u \neq \vec{0}$ , or the point  $\{0\}$  if  $u = \vec{0}$ .

• For any  $x, y \in \text{Ob}(\mathscr{C})$  with  $\mathfrak{f}(x) = u > \mathfrak{f}(y) = v$ , the cubical neat embedding  $\iota$  furnishes an embedding

$$\begin{split} &\mathcal{C}(y)\times\mathcal{M}_{\mathscr{C}}(x,y)\\ &=\prod_{i=0}^{|v|-1}[-R,R]^{d_{i}}\times\prod_{i=|v|}^{n-1}[-\epsilon,\epsilon]^{d_{i}}\times\widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(v,\vec{0})\times\mathcal{M}_{\mathscr{C}}(x,y)\\ &\cong\prod_{i=0}^{|v|-1}[-R,R]^{d_{i}}\times\prod_{i=|u|}^{n-1}[-\epsilon,\epsilon]^{d_{i}}\times\widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(v,\vec{0})\times\left(\prod_{i=|v|}^{|u|-1}[-\epsilon,\epsilon]^{d_{i}}\times\mathcal{M}_{\mathscr{C}}(x,y)\right)\\ &\stackrel{\mathrm{Id}}{\longleftrightarrow}\prod_{i=0}^{\bar{\iota}_{0}}[-R,R]^{d_{i}}\times\prod_{i=|u|}^{n-1}[-\epsilon,\epsilon]^{d_{i}}\times\widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(v,\vec{0})\times\left(\prod_{i=|v|}^{|u|-1}[-R,R]^{d_{i}}\times\mathcal{M}_{\mathscr{C}_{C}(n)}(u,v)\right)\\ &\cong\prod_{i=0}^{|u|-1}[-R,R]^{d_{i}}\times\prod_{i=|u|}^{n-1}[-\epsilon,\epsilon]^{d_{i}}\times\widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(v,\vec{0})\times\mathcal{M}_{\mathscr{C}_{C}(n)}(u,v)\\ &\hookrightarrow\prod_{i=0}^{|u|-1}[-R,R]^{d_{i}}\times\prod_{i=|u|}^{n-1}[-\epsilon,\epsilon]^{d_{i}}\times\partial(\widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(u,\vec{0}))\\ &\subset\partial\mathcal{C}(x). \end{split}$$


Here, the second inclusion comes from the composition map if  $v \neq \vec{0}$ , or the inclusion  $\{0\} \hookrightarrow [0,1]$  if  $v = \vec{0}$ . Let  $C_y(x) \subset \partial C(x)$  denote the image of this embedding.

• The attaching map for C(x) sends  $C_y(x) \cong C(y) \times \mathcal{M}_{\mathscr{C}}(x,y)$  by the projection map to C(y) and sends the complement of  $\bigcup_y C_y(x)$  in  $\partial C(x)$  to the basepoint.

The cubical realization of  $(\mathscr{C}, \mathfrak{f})$  is defined to be the formal desuspension  $\mathcal{X}(\mathscr{C}) = \Sigma^{-(k+d_0+\cdots+d_{n-1})} \|\mathscr{C}\|$ . (The desuspension ensures that the gradings of the objects in  $\mathscr{C}$  agree with the dimensions of the corresponding cells in  $\mathcal{X}(\mathscr{C})$ .)

**Lemma 3.30.** The attaching maps in the cubical realization are well-defined.

*Proof.* As in the proof of [LS14a, Lemma 3.25], we must show that for any  $x, y, z \in \text{Ob}(\mathscr{C})$  with gr(x) > gr(y) > gr(z), the dashed arrow in the following diagram exists such that the diagram commutes.



Let u, v, w denote  $\mathfrak{f}(x), \mathfrak{f}(y), \mathfrak{f}(z)$ , respectively. We may assume u > v > w; otherwise, it is not hard to verify that  $\mathcal{C}_z(x)$  and  $\mathcal{C}_y(x)$  are disjoint. In a similar vein to Definition 3.29, let  $\mathcal{C}_{y,z}(x) \subset \partial \mathcal{C}(x)$  be the image

of the following embedding:

$$\begin{split} &\mathcal{C}(z) \times \mathcal{M}_{\mathscr{C}}(y,z) \times \mathcal{M}_{\mathscr{C}}(x,y) \\ &= \prod_{i=0}^{|w|-1} [-R,R]^{d_i} \times \prod_{i=|w|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times \widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(w,\vec{0}) \times \mathcal{M}_{\mathscr{C}}(y,z) \times \mathcal{M}_{\mathscr{C}}(x,y) \\ &\cong \prod_{i=0}^{|w|-1} [-R,R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times \widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(w,\vec{0}) \times \left( \prod_{i=|w|}^{|v|-1} [-\epsilon,\epsilon]^{d_i} \times \mathcal{M}_{\mathscr{C}}(y,z) \right) \\ &\times \left( \prod_{i=|v|}^{|u|-1} [-\epsilon,\epsilon]^{d_i} \times \mathcal{M}_{\mathscr{C}}(x,y) \right) \\ &\stackrel{\mathrm{Id}}{\times} \overline{\iota}_{y,z} \times \overline{\iota}_{x,y} \stackrel{|w|-1}{\longrightarrow} [-R,R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times \widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(w,\vec{0}) \\ &\times \left( \prod_{i=|w|}^{|u|-1} [-R,R]^{d_i} \times \mathcal{M}_{\mathscr{C}_{C}(n)}(v,w) \right) \times \left( \prod_{i=|v|}^{|u|-1} [-R,R]^{d_i} \times \mathcal{M}_{\mathscr{C}_{C}(n)}(u,v) \right) \\ &\cong \prod_{i=0}^{|u|-1} [-R,R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times \widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(w,\vec{0}) \times \mathcal{M}_{\mathscr{C}_{C}(n)}(v,w) \times \mathcal{M}_{\mathscr{C}_{C}(n)}(u,v) \\ &\hookrightarrow \prod_{i=0}^{|u|-1} [-R,R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times \widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(u,\vec{0}) \right) \\ &\subset \partial \mathcal{C}(x). \end{split}$$

(As in Definition 3.29, the second inclusion usually comes from the composition map; the only special case is if  $w = \vec{0}$ , when it comes partly from the inclusion  $\{0\} \hookrightarrow [0,1]$ .)

We claim that  $C_{y,z}(x) = C_z(x) \cap C_y(x)$ . The direction  $C_{y,z}(x) \subseteq C_z(x) \cap C_y(x)$  is immediate since the inclusion

$$\mathcal{C}(z) \times \mathcal{M}_{\mathscr{C}}(y,z) \times \mathcal{M}_{\mathscr{C}}(x,y) \xrightarrow{\cong} \mathcal{C}_{y,z}(x) \subset \partial \mathcal{C}(x)$$

factors in both of the following ways (using Condition (3) of Definition 3.25):

$$\mathcal{C}(z) \times \mathcal{M}_{\mathscr{C}}(y, z) \times \mathcal{M}_{\mathscr{C}}(x, y) \xrightarrow{\operatorname{Id} \times \circ} \mathcal{C}(z) \times \mathcal{M}_{\mathscr{C}}(x, z) \xrightarrow{\cong} \mathcal{C}_{z}(x) \subset \partial \mathcal{C}(x)$$

$$\mathcal{C}(z) \times \mathcal{M}_{\mathscr{C}}(y, z) \times \mathcal{M}_{\mathscr{C}}(x, y) \xrightarrow{\cong} \mathcal{C}_{z}(y) \times \mathcal{M}_{\mathscr{C}}(x, y) \hookrightarrow \mathcal{C}(y) \times \mathcal{M}_{\mathscr{C}}(x, y) \xrightarrow{\cong} \mathcal{C}_{y}(x) \subset \partial \mathcal{C}(x).$$

The other direction requires more work. We will abuse notation slightly and identify points with their images under the composition map in  $\mathscr{C}_C(n)$ . View any point  $p \in \mathscr{C}_z(x) \cap \mathscr{C}_y(x)$  as a point  $(p_1, p_2, p_3, p_4)$  in the following subspace of  $\partial \mathscr{C}(x)$ :

$$\bigg(\prod_{i=0}^{|w|-1} [-R,R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i}\bigg) \times \prod_{i=|w|}^{|v|-1} [-R,R]^{d_i} \times \prod_{i=|v|}^{|u|-1} [-R,R]^{d_i} \times \partial \big(\widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(u,\vec{0})\big).$$

For p to lie in  $C_z(x)$ ,  $p_4$  must lie in the subspace

$$\widetilde{\mathcal{M}}_{\mathscr{C}_C(n)}(w,\vec{0}) \times \mathcal{M}_{\mathscr{C}_C(n)}(u,w);$$

similarly, for p to lie in  $C_u(x)$ ,  $p_4$  must lie in the subspace

$$\widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(v,\vec{0}) \times \mathcal{M}_{\mathscr{C}_{C}(n)}(u,v).$$

Therefore,  $p_4$  must lie in the subspace  $\widetilde{\mathcal{M}}_{\mathscr{C}_C(n)}(w,\vec{0}) \times \mathcal{M}_{\mathscr{C}_C(n)}(v,w) \times \mathcal{M}_{\mathscr{C}_C(n)}(u,v)$ . Write  $p_4$  also in component form as  $(p_{4,1},p_{4,2},p_{4,3})$ . Since p lies in  $\mathcal{C}_y(x)$ , we know  $(p_3,p_{4,3}) \in \operatorname{im}(\bar{\iota}_{x,y})$ , and since p lies in  $\mathcal{C}_z(x)$ , we know  $(p_2,p_3,p_{4,2},p_{4,3}) \in \operatorname{im}(\bar{\iota}_{x,z})$ . Moreover, since  $\iota$  is a cubical neat embedding (Definition 3.25),

$$\operatorname{im}(\bar{\iota}_{x,z}) \cap \Big( \prod_{i=|w|}^{|v|-1} [-R,R]^{d_i} \times \mathcal{M}_{\mathscr{C}_C(n)}(v,w) \times \prod_{i=|v|}^{|u|-1} [-R,R]^{d_i} \times \mathcal{M}_{\mathscr{C}_C(n)}(u,v) \Big) = \operatorname{im} \Big( \prod_{y' \mid \mathfrak{f}(y') = v} \bar{\iota}_{y',z} \times \bar{\iota}_{x,y'} \Big),$$

and therefore, there exists y' with  $\mathfrak{f}(y') = v$  such that  $(p_3, p_{4,3}) \in \operatorname{im}(\overline{\iota}_{x,y'})$  and  $(p_2, p_{4,2}) \in \operatorname{im}(\overline{\iota}_{y',z})$ . Condition (2) from Definition 3.25 (but with  $\overline{\iota}$  instead of  $\iota$ ) ensures that y' = y, and then it is straightforward to see that p lies in  $C_{y,z}(x)$ . This completes the proof that  $C_{y,z}(x) = C_z(x) \cap C_y(x)$ .

Define the dashed arrow from  $C_z(x) \cap C_y(x) = C_{y,z}(x) \cong C(z) \times \mathcal{M}_{\mathscr{C}}(y,z) \times \mathcal{M}_{\mathscr{C}}(x,y)$  to  $C_z(y) \cong C(z) \times \mathcal{M}_{\mathscr{C}}(y,z)$  to be the projection map. From the definition of  $C_{y,z}(x)$ , it is easy to verify that the resulting diagram commutes.

**Proposition 3.31.** Up to stable homotopy equivalence, the cubical realization is independent of the choice of tuple  $\mathbf{d}$ . More precisely, let  $\iota$  be a cubical neat embedding relative to  $\mathbf{d} = (d_0, \ldots, d_{n-1})$  and let  $\|\mathscr{C}\|$  be the cubical realization corresponding to  $\iota$ . Fix a tuple  $\mathbf{d}' = (d'_0, \ldots, d'_{n-1})$  with  $d'_i \geq d_i$  for all i. There is an induced cubical neat embedding of  $\mathscr{C}$  relative to  $\mathbf{d}'$ , gotten by identifying the space  $E_{u,v}$  for  $\iota$  with the subspace

$$\prod_{i=|v|}^{|u|-1} (-R,R)^{d_i} \times \{0\}^{d'_i - d_i} \times \mathcal{M}_{\mathscr{C}_C(n)}(u,v)$$

of the space  $E'_{u,v}$  for  $\iota'$ . Let  $\|\mathscr{C}\|'$  be the cubical realization corresponding to  $\iota'$  and let  $N = |\mathbf{d}'| - |\mathbf{d}| = \sum_{i=0}^{n-1} d'_i - \sum_{i=0}^{n-1} d_i$ . Then there is a homotopy equivalence

$$\Sigma^N \|\mathscr{C}\| \simeq \|\mathscr{C}\|',$$

taking cells to the corresponding cells.

*Proof.* The proof is the same as Case (3) in the proof of [LS14a, Lemma 3.27].

One can also show, by following the proofs of [LS14a, Lemmas 3.25–3.27], that the stable homotopy type of  $\|\mathcal{C}\|_{\iota}$  is independent of the cubical neat embedding  $\iota$  and the parameters R and  $\epsilon$ . Since this result also follows from Theorem 4, we do not give further details.

We conclude this section by returning to our simplicial complex example:

**Proposition 3.32.** Let  $S_{\bullet}$  be a simplicial complex with n vertices and let  $(\mathcal{C}, \mathfrak{f}: \Sigma \mathcal{C} \to \mathcal{C}_{C}(n))$  be the corresponding cubical flow category, as in Example 3.23. Let  $|S_{\bullet}|_{+}$  denote the disjoint union of the geometric realization of  $S_{\bullet}$  and a basepoint. Then there is a stable homotopy equivalence

$$\mathcal{X}(\mathscr{C}) \simeq |S_{\bullet}|_{\perp}.$$

Moreover, the stable homotopy equivalence may be chosen so that it induces an isomorphism between the simplicial cochain complex of  $S_{\bullet}$  and the reduced cellular cochain complex of  $\mathcal{X}(\mathscr{C})$  (with the CW complex structure from Definition 3.29).

We leave the proof as a (somewhat involved) exercise to the reader.

3.8. Cubical realization agrees with the Cohen-Jones-Segal realization. The main aim of this subsection is to prove the following.

**Theorem 4.** Let  $(\mathscr{C}, \mathfrak{f}: \mathscr{C} \to \mathscr{C}_C(n))$  be a cubical flow category. For any cubical neat embedding  $\iota$  relative to any tuple  $\mathbf{d} = (d_0, \dots, d_{n-1})$ , and parameters  $R, \epsilon$ , the cubical realization  $\mathcal{X}(\mathscr{C})$  (from Definition 3.29) is stably homotopy equivalent to the Cohen-Jones-Segal realization (from Section 3.2) of C with framing induced from some framing of  $\mathscr{C}_{C}(n)$ . Furthermore, the stable homotopy equivalence sends cells to corresponding cells by degree  $\pm 1$  maps.

*Proof.* Using Proposition 3.31, we may assume that the  $d_i$ 's are sufficiently large so that there is a neat embedding j of the cube flow category  $\mathscr{C}_C(n)$  relative to  $\mathbf{D} = (\dots, 0, \dots, 0, d_0, \dots, d_{n-1}, 0, \dots, 0, \dots)$  in the sense of Definition 3.7. Fix some framing of the cube flow category  $\mathscr{C}_{C}(n)$ , and construct the extension  $\bar{\jmath}$  as in Definition 3.10. Let  $\delta$  and T be the corresponding parameters. After scaling if necessary, we may assume  $\delta = R$ , and after increasing T if necessary, we may assume T > 1.

Note that  $\bar{\jmath} \circ \iota$  is a neat embedding (in the sense of Definition 3.7) of our flow category  $\mathscr{C}$  relative to  $\mathbf{D}$ ; it has an extension  $\bar{j} \circ \bar{\iota}$  (again, as in Definition 3.10) with respect to the parameters  $\epsilon$  and T, and  $\bar{j} \circ \bar{\iota}$  is isotopic to the extension coming from the framing of  $\mathscr{C}$  induced from the framing of  $\mathscr{C}_{C}(n)$ ; see also Remark 3.28.

Let  $\|\mathscr{C}\|$  be the CW complex constructed from the cubical neat embedding  $\iota$  and its extension  $\bar{\iota}$  with respect to the parameters  $\epsilon$  and R; and let  $|\mathscr{C}|$  be the CW complex constructed from the neat embedding  $\bar{\jmath} \circ \iota$  and its extension  $\bar{\jmath} \circ \bar{\iota}$  with respect to the parameters  $\epsilon$  and T. Cells of both  $\|\mathscr{C}\|$  and  $|\mathscr{C}|$  correspond to objects of  $\mathscr{C}$ , and hence to each other. We will produce a quotient map from  $|\mathscr{C}|$  to  $|\mathscr{C}|$  which will send each cell via a degree  $\pm 1$  map to the corresponding cell. It follows that the quotient map is a stable homotopy equivalence.

For any  $x \in \mathrm{Ob}(\mathscr{C})$  with  $\mathfrak{f}(x) = u \in \mathrm{Ob}(\mathscr{C}_C(n))$ , the cell associated to x in  $\|\mathscr{C}\|$  is

$$C(x)' = \begin{cases} \prod_{i=0}^{|u|-1} [-R, R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon, \epsilon]^{d_i} \times [0, 1] \times \mathcal{M}_{\mathscr{C}_C(n)}(u, \vec{0}) & \text{if } u \neq \vec{0}, \\ \prod_{i=0}^{n-1} [-\epsilon, \epsilon]^{d_i} \times \{0\} & \text{if } u = \vec{0}. \end{cases}$$

while the cell associated to x in  $|\mathscr{C}|$  is

$$C(x) = \prod_{i=0}^{|u|-1} [-T, T]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon, \epsilon]^{d_i} \times \prod_{i=0}^{|u|-1} [0, T].$$

Let  $[\mathcal{C}(x)]$  (respectively  $[\mathcal{C}(x)']$ ) denote the image of  $\mathcal{C}(x)$  (respectively  $\mathcal{C}(x)'$ ) in  $|\mathcal{C}|$  (respectively  $|\mathcal{C}(x)'|$ ). We will define a map  $\mathcal{C}(x) \to [\mathcal{C}(x)']$  and check that this induces a well-defined map  $[\mathcal{C}(x)] \to [\mathcal{C}(x)']$ . If  $u = \vec{0}$ , identify  $\mathcal{C}(x)' \cong \prod_{i=0}^{n-1} [-\epsilon, \epsilon]^{d_i} \cong \mathcal{C}(x)$ . If  $u \neq \vec{0}$ , the embedding

$$\overline{\jmath}_{u,\vec{0}} \colon \mathcal{M}_{\mathscr{C}_{C}(n)}(u,\vec{0}) \times \prod_{i=0}^{|u|-1} [-R,R]^{d_{i}} \hookrightarrow (-T,T)^{d_{0}} \times [0,T) \times \dots \times [0,T) \times (-T,T)^{d_{|u|-1}}$$

$$\cong \prod_{i=1}^{|u|-1} [0,T) \times \prod_{i=0}^{|u|-1} (-T,T)^{d_{i}}$$

induces the following codimension-zero embedding of C(x)' into C(x):

$$\begin{split} \mathcal{C}(x)' &= \prod_{i=0}^{|u|-1} [-R,R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times [0,1] \times \mathcal{M}_{\mathscr{C}_{C}(n)}(u,\vec{0}) \\ &\cong \mathcal{M}_{\mathscr{C}_{C}(n)}(u,\vec{0}) \times \prod_{i=0}^{|u|-1} [-R,R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times [0,1] \\ &\stackrel{\bar{\jmath}_{u,\vec{0}} \times \operatorname{Id}}{\longleftrightarrow} \prod_{i=1}^{|u|-1} [0,T] \times \prod_{i=0}^{|u|-1} [-T,T]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times [0,1] \\ &\cong \prod_{i=0}^{|u|-1} [-T,T]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times [0,1] \times \prod_{i=1}^{|u|-1} [0,T] \\ &\hookrightarrow \prod_{i=0}^{|u|-1} [-T,T]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i} \times \prod_{i=0}^{|u|-1} [0,T] \\ &= \mathcal{C}(x). \end{split}$$

(Here, the  $\cong$  arrows correspond to the obvious reshuffling of the factors, and the second inclusion is induced from the inclusion  $[0,1] \hookrightarrow [0,T]$ .) In either case, map  $\mathcal{C}(x)$  to  $[\mathcal{C}(x)']$  by identifying the image of this embedding with  $\mathcal{C}(x)'$ , and quotienting everything else to the basepoint. To see that this gives a well-defined, continuous map from the CW complex  $|\mathscr{C}|$  to the CW complex  $|\mathscr{C}|$ , we need to check that for any other  $y \in \mathrm{Ob}(\mathscr{C})$  with  $\mathfrak{f}(y) = v < u$ , the following commutes:

$$\mathcal{C}(y)' \times \mathcal{M}_{\mathscr{C}}(x,y) \xrightarrow{} \mathcal{C}(y) \times \mathcal{M}_{\mathscr{C}}(x,y) 
\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow 
\mathcal{C}(x)' \xrightarrow{} \mathcal{C}(x).$$

(The horizontal arrows are induced from the inclusions defined above. The right vertical arrow is the inclusion defined in Definition 3.10 for  $\bar{\jmath} \circ \bar{\iota}$ , while the left vertical arrow is the inclusion defined in Definition 3.29 for  $\bar{\iota}$ .)

When  $v = \vec{0}$ , after removing the constant factor of  $\prod_{i=|u|}^{n-1} [-\epsilon, \epsilon]^{d_i}$  and doing some consistent reshuffling, the diagram is

$$\{0\} \times \prod_{i=0}^{|u|-1} [-\epsilon, \epsilon]^{d_i} \times \mathcal{M}_{\mathscr{C}}(x, y) \stackrel{\cong}{\longrightarrow} \prod_{i=0}^{|u|-1} [-\epsilon, \epsilon]^{d_i} \times \mathcal{M}_{\mathscr{C}}(x, y)$$

$$\downarrow^{(\kappa_1, \bar{\iota}_{x,y})} \qquad \qquad \downarrow^{(\kappa_3^0, \operatorname{Id}) \circ \bar{\jmath}_{u, \vec{0}} \circ \bar{\iota}_{x,y}}$$

$$[0, 1] \times \prod_{i=0}^{|u|-1} [-R, R]^{d_i} \times \mathcal{M}_{\mathscr{C}_C(n)}(u, \vec{0}) \stackrel{(\kappa_2, \bar{\jmath}_{u, \vec{0}})}{\longrightarrow} \prod_{i=0}^{|u|-1} [0, T] \times \prod_{i=0}^{|u|-1} [-T, T]^{d_i},$$

where  $\kappa_1$  is the inclusions  $\{0\} \hookrightarrow [0,1]$ ,  $\kappa_2$  is the inclusion  $[0,1] \hookrightarrow [0,T]$ , and  $\kappa_3^{\ell}$  (for  $0 \leq \ell < |u|$ ) is the inclusion

$$\prod_{i=\ell+1}^{|u|-1}[0,T]\cong\{0\}\times\prod_{i=\ell+1}^{|u|-1}[0,T]\stackrel{(\kappa_2\circ\kappa_1,\operatorname{Id})}{\longleftrightarrow}\prod_{i=\ell}^{|u|-1}[0,T].$$

Therefore, the diagram commutes.

When  $v \neq \vec{0}$ , once again after removing the constant factor of  $\prod_{i=|u|}^{n-1} [-\epsilon, \epsilon]^{d_i}$ , and some consistent reshuffling, the diagram factors as

$$\begin{bmatrix}
[0,1] \times \mathcal{M}_{\mathscr{C}_{C}(n)}(v,\vec{0}) \times \prod_{i=0}^{|v|-1} [-R,R]^{d_{i}} \\
\times \prod_{i=|v|}^{|u|-1} [-\epsilon,\epsilon]^{d_{i}} \times \mathcal{M}_{\mathscr{C}}(x,y)
\end{bmatrix}
\xrightarrow{(\kappa_{2},\bar{\jmath}_{v},\vec{0},\mathrm{Id})} \begin{bmatrix}
\prod_{i=0}^{|v|-1} [0,T] \times \prod_{i=0}^{|v|-1} [-T,T]^{d_{i}} \\
\times \prod_{i=|v|}^{|u|-1} [-\epsilon,\epsilon]^{d_{i}} \times \mathcal{M}_{\mathscr{C}}(x,y)
\end{bmatrix}$$

$$\begin{bmatrix}
(\mathrm{Id},\bar{\iota}_{x,y}) \\
\downarrow^{(\mathrm{Id},\bar{\jmath}_{u,0}\circ\bar{\iota}_{x,y})}
\end{bmatrix}
\xrightarrow{(\kappa_{2},\bar{\jmath}_{v},\vec{0},\bar{\jmath}_{u,v})} \begin{bmatrix}
\prod_{i=0}^{|v|-1} [0,T] \times \prod_{i=0}^{|u|-1} [-T,T]^{d_{i}} \times \prod_{i=|v|+1}^{|u|-1} [0,T]
\end{bmatrix}$$

$$\times \mathcal{M}_{\mathscr{C}_{C}(n)}(u,v)$$

$$\downarrow^{(\mathrm{Id},o)\circ\rho}$$

$$\begin{bmatrix}
(\mathrm{Id},o)\circ\rho
\end{bmatrix}
\xrightarrow{(\kappa_{2},\bar{\jmath}_{u},\vec{0})}
\end{bmatrix}
\xrightarrow{(\kappa_{2},\bar{\jmath}_{u},\vec{0})}$$

$$\begin{bmatrix}
\prod_{i=0}^{|v|-1} [0,T] \times \prod_{i=0}^{|u|-1} [-T,T]^{d_{i}} \times \prod_{i=|v|+1}^{|u|-1} [0,T]
\end{bmatrix}$$

$$\begin{bmatrix}
\rho \circ (\mathrm{Id},\kappa_{3}^{|v|})
\end{bmatrix}$$

$$\begin{bmatrix}
\rho \circ (\mathrm{Id},\kappa_{3}^{|v|})
\end{bmatrix}$$

$$\begin{bmatrix}
[0,1] \times \prod_{i=0}^{|u|-1} [-R,R]^{d_{i}} \times \mathcal{M}_{\mathscr{C}_{C}(n)}(u,\vec{0})
\end{bmatrix}
\xrightarrow{(\kappa_{2},\bar{\jmath}_{u},\vec{0})}$$

$$\begin{bmatrix}
[0,1] \times \prod_{i=0}^{|u|-1} [-R,R]^{d_{i}} \times \mathcal{M}_{\mathscr{C}_{C}(n)}(u,\vec{0})
\end{bmatrix}
\xrightarrow{(\kappa_{2},\bar{\jmath}_{u},\vec{0})}$$

where  $\rho$  throughout denotes some (further) consistent reshuffling of factors, and  $\kappa_2$ ,  $\kappa_3^{\ell}$  from before. The top square commutes automatically; the bottom square commutes since Condition (2) of Definition 3.7 holds for  $\bar{\jmath}$  as well (because the extension was defined via coherent framings of the normal bundles of  $\jmath$ ).

Therefore, we get a well-defined map from  $|\mathcal{C}|$  to  $||\mathcal{C}||$  which sends the cell  $[\mathcal{C}(x)]$  to the corresponding cell  $[\mathcal{C}(x)']$  by a degree  $\pm 1$  map. It is easy to check that the grading shifts match up, and therefore, we get the required stable homotopy equivalence.

## 4. Functors from the cube to the Burnside category and their realizations

In this section we give a reformulation of cubical flow categories, as 2-functors from the cube category to the Burnside category. We then give a choice-free way of realizing such a functor, in terms of a thickening construction. (A smaller but choice-dependent way to realize such a functor is given in Section 5.) The proof that this realization agrees with the cubical realization is deferred until Section 6. The main results of this section can be summarized in two theorems:

**Theorem 5.** The data of a cubical flow category is equivalent to the data of a strictly unitary, lax 2-functor from the cube category to the Burnside category.

This is stated more precisely and proved as Lemmas 4.18 and 4.20.

**Theorem 6.** Given a functor F from the cube category to the Burnside category there is a canonically associated CW spectrum |F|, the realization of F.

This is stated precisely as Construction 4.39 and Lemma 4.42. In Section 6, we show that this construction agrees with the cubical realization (a combination of Theorems 7 and 8). A feature of the realization procedure introduced in this section is that it behaves well with respect to products (Proposition 4.46).

We start by reviewing the Burnside (2-)category and the data required to define a functor from the cube to it, in Section 4.1. We then recall some properties of homotopy colimits, in Section 4.2, before turning to new material in Section 4.3, where we formulate precisely and prove Theorem 5. The thickening and realization procedure are given in Sections 4.4 and 4.5, while invariance under natural isomorphisms of 2-functors is proved in Section 4.6. Section 4.7 discusses the behavior of this realization procedure under certain kinds of products.

4.1. The Burnside category and 2-functors to it. Given sets X and Y, a correspondence (or span) from X to Y is a set A and maps  $s: A \to X$  and  $t: A \to Y$  (for source and target). Given a correspondence  $(A, s_A, t_A)$  from X to Y and  $(B, s_B, t_B)$  from Y to Z the composition  $(B, s_B, t_B) \circ (A, s_A, t_A)$  is the correspondence (C, s, t) from X to Z given by

$$C = B \times_Y A = \{(b, a) \in B \times A \mid t(a) = s(b)\}$$
  $s(b, a) = s_A(a)$   $t(b, a) = t_B(b)$ .

Given correspondences  $(A, s_A, t_A)$  and  $(B, s_B, t_B)$  from X to Y, a morphism of correspondences from  $(A, s_A, t_A)$  to  $(B, s_B, t_B)$  is a bijection of sets  $f: A \to B$  which commutes with the source and target maps, i.e., so that  $s_A = s_B \circ f$  and  $t_A = t_B \circ f$ . Composition (of set maps) makes the set of correspondences from X to Y into a category. Further, composition of correspondences makes (Sets, Correspondences, Morphisms of correspondences) into a weak 2-category (bicategory in the language of [Bén67]). By the Burnside category we mean the sub-2-category of finite sets and finite correspondences. We denote the Burnside category by  $\mathcal{B}$ . (More typically, one defines the Burnside category of a group G in terms of G-sets and G-equivariant correspondences; for us, G is the trivial group.)

Remark 4.1. Some authors refer to our Burnside category as the Burnside 2-category, and refer to the 1-category with objects sets and morphisms isomorphism classes of correspondences as the Burnside category.

We will typically drop the maps s and t from the notation, referring simply to a correspondence A from X to Y.

As mentioned above, the Burnside category is a weak 2-category: the identity and associativity axioms only hold up to natural isomorphism. That is, given a set X, the *identity correspondence* of X is simply the set X itself, with the identity map as source and target maps. Given another correspondence A from W to X there is a natural isomorphism

$$\lambda \colon X \times_X A \stackrel{\cong}{\longrightarrow} A$$

defined by  $\lambda(x,a)=a$ . Similarly, given a correspondence B from X to Y there is a natural isomorphism

$$\rho \colon B \times_X X \stackrel{\cong}{\longrightarrow} B$$

defined by  $\rho(b,x)=b$ . Finally, given correspondences A from W to X, B from X to Y and C from Y to Z there is a natural isomorphism

$$\alpha \colon (C \times_Y B) \times_X A \to C \times_Y (B \times_X A)$$

defined by  $\alpha((c,b),a)=(c,(b,a)).$ 

This distinction between weak and strict 2-categories may seem superficial here, but the distinction between weak and strict 2-functors, to which we turn next, will be crucial.

**Definition 4.2.** Given (weak) 2-categories  $\mathscr C$  and  $\mathscr D$ , a lax 2-functor  $F \colon \mathscr C \to \mathscr D$  consists of the following data:

- For each object  $x \in \text{Ob}(\mathscr{C})$  an object  $F(x) \in \text{Ob}(\mathscr{D})$ .
- For each pair of objects  $x, y \in \text{Ob}(\mathscr{C})$  a functor  $F_{x,y} \colon \text{Hom}(x,y) \to \text{Hom}(F(x),F(y))$ .
- For each object  $x \in \text{Ob}(\mathscr{C})$  a 2-morphism  $F_{\text{Id}_x} : \text{Id}_{F(x)} \to F_{x,x}(\text{Id}_x)$ .
- For any three objects  $x, y, z \in Ob(\mathscr{C})$  a natural transformation

$$F_{x,y,z}(\cdot,\cdot)\colon F_{y,z}(\cdot)\circ_1 F_{x,y}(\cdot)\to F_{x,z}(\cdot\circ_1\cdot).$$

(Here, both  $F_{y,z}(\cdot) \circ_1 F_{x,y}(\cdot)$  and  $F_{x,z}(\cdot \circ_1 \cdot)$  are functors  $\operatorname{Hom}(y,z) \times \operatorname{Hom}(x,y) \to \operatorname{Hom}(F(x),F(z))$ ;  $\circ_1$  denotes the 1-composition in  $\mathscr C$  or  $\mathscr D$ , not the composition of functors.)

These data must satisfy the following compatibility conditions:

(Fn-1) For each pair of objects  $x, y \in \text{Ob}(\mathscr{C})$ , the following diagrams commute:

(Here,  $\lambda$  is the natural isomorphism from  $\mathrm{Id}_y \circ_1 \cdot$  to the identity functor of  $\mathscr{C}$  or  $\mathscr{D}$ , as appropriate, and  $\rho$  is the natural isomorphism from  $\cdot \circ_1 \mathrm{Id}_x$  to the identity functor of  $\mathscr{C}$  or  $\mathscr{D}$ .)

(Fn-2) For each quadruple of objects  $x, y, z, w \in Ob(\mathscr{C})$ , the following diagram commutes:

(Here,  $\alpha$  is the natural isomorphism from  $((\cdot \circ_1 \cdot) \circ_1 \cdot)$  to  $(\cdot \circ_1 (\cdot \circ_1 \cdot))$ , the two different orders of triple-compositions, associated to  $\mathscr{C}$  or  $\mathscr{D}$ , as appropriate.)

(See, for instance, [Bén67, Definition 4.1 and Remark 4.2], where lax 2-functors are called homomorphisms.)

We will often drop the subscript from F: given an element  $f \in \operatorname{Hom}_{\mathscr{C}}(x,y)$  and a lax 2-functor  $F \colon \mathscr{C} \to \mathscr{D}$  we will often write F(f) for  $F_{x,y}(f)$ .

**Definition 4.3.** [Bén67, Remark 4.2] We call a lax 2-functor  $F: \mathscr{C} \to \mathscr{D}$  strictly unitary if for all objects  $x \in \mathrm{Ob}(\mathscr{C}), F_{x,x}(\mathrm{Id}_x) = \mathrm{Id}_{F(x)}$  and  $F_{\mathrm{Id}_x}$  is the identity 2-morphism.

As mentioned above, we will be mainly interested in 2-functors from  $\underline{2}^n$  to  $\mathcal{B}$ , which moreover will be strictly unitary. In this case, Definition 4.2 simplifies substantially:

**Lemma 4.4.** A strictly unitary, lax 2-functor  $F: \underline{2}^n \to \mathcal{B}$  is determined by the following data:

- For each object  $v \in \text{Ob}(\underline{2}^n) = \{0,1\}^n$  the set  $X_v = F(v)$ .
- For each pair of objects  $v, w \in \text{Ob}(\underline{2}^n)$  such that v > w, a correspondence  $A_{v,w} = F(\varphi_{v,w})$  from  $X_v$  to  $X_w$ .
- For each triple of objects  $u, v, w \in \text{Ob}(\underline{2}^n)$  such that u > v > w, a bijection  $F_{u,v,w} \colon A_{v,w} \times_{X_v} A_{u,v} \to A_{u,w}$ .

These data satisfy the compatibility condition:

(CF-1) For any  $u, v, w, x \in Ob(\underline{2}^n)$  with u > v > w > x, the following diagram commutes:

(We have suppressed some non-confusing parentheses.)

Moreover, any collection of data satisfying this compatibility condition determines a strictly unitary, lax 2-functor  $F: 2^n \to \mathcal{B}$ .

Proof. Since F is strictly unitary, we have  $F(\varphi_{v,v}) = \operatorname{Id}_{X_v}$  (which is  $X_v$ , viewed as a correspondence from itself to itself). If  $v \not\geq w$  then  $\operatorname{Hom}(v,w) = \varnothing$ , so  $F(\varphi_{v,w})$  is the unique functor from the empty category. Thus, the  $F(\varphi_{v,w})$  are entirely specified by the correspondences  $A_{v,w} = F_{v,w}(\varphi_{v,w})$  with v > w. Next, the source  $\operatorname{Hom}(v,w) \times \operatorname{Hom}(u,v)$  of  $F_{u,v,w}$  is nonempty if and only if  $u \geq v \geq w$ , in which case  $\operatorname{Hom}(v,w) \times \operatorname{Hom}(u,v)$  consists of the single element  $(\varphi_{v,w},\varphi_{u,v})$ . Since  $2^n$  is a strict 2-category and F is strictly unitary, Condition (Fn-1) is equivalent to the statement that  $F_{v,v,w} \colon A_{v,w} \times X_v \times A_{v,w}$  is the canonical isomorphism  $\rho$ , and  $F_{v,w,w} \colon X_w \times_{X_w} A_{v,w} \to A_{v,w}$  is the canonical isomorphism  $\lambda$ . So, F is determined by the specified data. Condition (Fn-2) is equivalent to Condition (CF-1); in Condition (CF-1) we have abused notation to identify the two sides of the top row of Condition (Fn-2), and the bottom arrow in Condition (Fn-2) is an equality because  $2^n$  is a strict 2-category and F is strictly unitary. The result follows.

**Lemma 4.5.** Up to natural isomorphism, a strictly unitary, lax 2-functor  $F: \underline{2}^n \to \mathcal{B}$  is determined by the sets F(v), the correspondences  $F(\varphi_{v,w})$  with v > w and |v| - |w| = 1, and the maps  $F_{u,v',w}^{-1} \circ F_{u,v,w} : F(\varphi_{v,w}) \circ F(\varphi_{u,v}) \to F(\varphi_{v,v}) \circ F(\varphi_{v,v})$  with u > v, v' > w and |u| - |w| = 2.

*Proof.* This follows from the observations that:

- Any morphism in  $\underline{2}^n$  is a composition of maps associated to edges, so,  $F(\varphi_{v,w})$  is determined for all  $v \geq w$ .
- Any two (directed) edge paths from v to w in  $\underline{2}^n$  are related by a sequence of swaps across 2-dimensional faces, so  $F_{u,v,w}$  is determined for all  $u \geq v \geq w$ .

Further details are left to the reader.

4.2. Homotopy colimits and homotopy coherent diagrams. The realization of a 2-functor  $\underline{2}^n \to \mathcal{B}$  in this section and Section 5 will involve an iterated mapping cone, which can be described as a homotopy colimit. So, we review briefly the notion of homotopy colimits here.

Given a diagram  $F: \mathcal{D} \to \mathsf{Top}_{\bullet}$  of based topological spaces, the *homotopy colimit* of F, hocolim  $F = \mathsf{hocolim}_{\mathcal{D}} F$ , is another based topological space. Similarly, if  $F: \mathcal{D} \to \mathcal{S}$  is a diagram of spectra then we can again form the homotopy colimit of F, hocolim F, which is a spectrum. We will give a construction in a slightly more general setting presently, but first we note some key properties of the homotopy colimit, all of which hold for both diagrams of spaces and diagrams of spectra:

(ho-1) Suppose that  $F,G:\mathscr{C}\to \mathsf{Top}_{\bullet}$  are diagrams and  $\eta\colon F\to G$  is a natural transformation. Then  $\eta$  induces a map hocolim  $\eta\colon \mathsf{hocolim}\, F\to \mathsf{hocolim}\, G$ . If  $\eta(c)$  is a stable homotopy equivalence for each  $c\in \mathsf{Ob}(\mathscr{C})$  then hocolim  $\eta$  is a stable homotopy equivalence as well.

- (ho-2) Suppose that  $F, G: \mathscr{C} \to \mathsf{Top}_{\bullet}$  are diagrams and  $F \vee G: \mathscr{C} \to \mathsf{Top}_{\bullet}$  is the diagram obtained by taking their wedge sum,  $(F \vee G)(v) = F(v) \vee G(v)$ . Then the natural map hocolim  $F \vee G \to \mathsf{hocolim}\, F \vee \mathsf{hocolim}\, G$  is an equivalence.
- (ho-3) Suppose that  $F: \mathscr{C} \to \mathsf{Top}_{\bullet}$  and  $G: \mathscr{D} \to \mathsf{Top}_{\bullet}$ . Then there is an induced functor  $F \wedge G: \mathscr{C} \times \mathscr{D} \to \mathsf{Top}_{\bullet}$ , with  $(F \wedge G)(v \times w) = F(v) \wedge G(w)$ , and a natural weak equivalence (hocolim F)  $\wedge$  (hocolim G)  $\to$  hocolim F ho
- (ho-4) Let  $G: \mathscr{C} \to \mathscr{D}$  be a map of diagrams (i.e., a functor between small categories). Given  $d \in \mathrm{Ob}(\mathscr{D})$ , the undercategory of d has objects  $\{(c,f) \mid c \in \mathscr{C}, f \colon d \to G(c)\}$ , and  $\mathrm{Hom}((c,f),(c',f')) = \{g \colon c \to c' \mid f' = G(g) \circ f\}$ . Let  $d \downarrow G$  denote the undercategory of d. The functor G is called homotopy cofinal if for each  $d \in \mathrm{Ob}(\mathscr{D})$ ,  $d \downarrow G$  has contractible nerve.

Now, let  $F: \mathcal{D} \to \mathsf{Top}_{\bullet}$  be a diagram. Then there is an induced functor  $F \circ G: \mathscr{C} \to \mathsf{Top}_{\bullet}$ . Suppose that G is homotopy cofinal. Then

$$\operatorname{hocolim} F \circ G \simeq \operatorname{hocolim} F$$
.

(In the case of homotopy limits, this is [BK72, Cofinality Theorem XI,9.2]. The homotopy colimit can be characterized by knowing that the mapping space Map(hocolim F, Z) is equivalent to the homotopy limit of the diagram of spaces Map(F, Z).)

In Section 5, we will need to talk about homotopy colimits of diagrams which are only homotopy-commutative, but where the homotopies are part of the data, and are coherent up to higher homotopies (also part of the data). The rest of this subsection focuses on one formulation of this generalization.

We start with an appropriate notion of diagrams:

**Definition 4.6.** [Vog73, Definition 2.3] A homotopy coherent diagram in Top<sub>●</sub> consists of:

- A small category  $\mathscr{C}$ .
- For each  $x \in Ob(\mathscr{C})$  a space  $F(x) \in \mathsf{Top}_{\bullet}$ .
- For each  $n \ge 1$  and each sequence

$$x_0 \xrightarrow{f_1} x_1 \xrightarrow{f_2} \cdots \xrightarrow{f_n} x_n$$

of composable morphisms in  $\mathscr{C}$  a continuous map

$$F(f_n, \dots, f_1) \colon [0, 1]^{n-1} \times F(x_0) \to F(x_n)$$

with  $F(f_n, ..., f_1)([0, 1]^{n-1} \times \{*\}) = *$ , the basepoint in  $F(x_n)$ .

Letting  $(t_1, \ldots, t_{n-1})$  denote points in  $[0,1]^{n-1}$ , these maps F are required to satisfy the conditions:

$$(4.7)$$
  $F(f_n,\ldots,f_1)(t_1,\ldots,t_{n-1})$ 

$$= \begin{cases} F(f_n,\ldots,f_2)(t_2,\ldots,t_{n-1}) & f_1 = \operatorname{Id} \\ F(f_n,\ldots,f_{i+1},f_{i-1},\ldots,f_1)(t_1,\ldots,t_{i-1}\cdot t_i,\ldots,t_{n-1}) & f_i = \operatorname{Id}, \ 1 < i < n \\ F(f_{n-1},\ldots,f_1)(t_1,\ldots,t_{n-2}) & f_n = \operatorname{Id} \\ [F(f_n,\ldots,f_{i+1})(t_{i+1},\ldots,t_{n-1})] \circ [F(f_i,\ldots,f_1)(t_1,\ldots,t_{i-1})] & t_i = 0 \\ F(f_n,\ldots,f_{i+1} \circ f_i,\ldots,f_1)(t_1,\ldots,t_{i-1},t_{i+1},\ldots,t_{n-1}) & t_i = 1. \end{cases}$$

(A homotopy coherent diagram is what Vogt [Vog73] calls an  $h\mathscr{C}$ -diagram. We are restricting to the case that his topological category  $\mathscr{C}$  is discrete.)

We will abuse notation and denote a homotopy coherent diagram as above by  $F \colon \mathscr{C} \to \mathsf{Top}_{\bullet}$ . It will be clear from context when we mean a commutative diagram or a homotopy coherent diagram.

Example 4.8. Any commutative diagram  $F: \mathscr{C} \to \mathsf{Top}_{\bullet}$  can be viewed as a homotopy coherent diagram by defining

$$F(f_n,\ldots,f_1)(t_1,\ldots,t_{n-1})=F(f_n\circ\cdots\circ f_1).$$

Remark 4.9. Associated to an ordinary category  $\mathscr{C}$ , there is a simplicially enriched category  $\mathfrak{C}[\mathscr{C}]$ , introduced by Leitch [Lei75] and further developed by many authors (e.g. [Cor82, DK80, Lur09]) such that homotopy coherent diagrams  $\mathscr{C} \to \mathsf{Top}_{\bullet}$  are the same (up to the replacement of the continuous function  $t_{i-1} \cdot t_i$  by an equivalent piecewise linear one) as simplicial functors from  $\mathfrak{C}[\mathscr{C}]$  to spaces. Lemma 3.20 essentially proves that the cube flow category  $\mathscr{C}_{C}(n)$  is the topological category obtained from  $\mathfrak{C}[\underline{2}^{n}]$  by taking the realization of each morphism set. So, homotopy coherent  $\underline{2}^{n}$ -diagrams give functors out of  $\mathscr{C}_{C}(n)$ .

**Definition 4.10.** [Vog73, Paragraph (5.10)] Given a homotopy coherent diagram  $F: \mathscr{C} \to \mathsf{Top}_{\bullet}$ , the homotopy colimit of F is defined by

(4.11) 
$$\operatorname{hocolim} F = \{*\} \coprod \coprod_{n \geq 0} \coprod_{x_0 \xrightarrow{f_1} \dots \xrightarrow{f_n} x_n} [0, 1]^n \times F(x_0) / \sim,$$

where the second coproduct is over n-tuples of composable morphisms in  $\mathscr{C}$  and the case n=0 corresponds to the objects  $x_0 \in \text{Ob}(\mathscr{C})$ . Letting  $(t_1, \ldots, t_n)$  denote points in  $[0,1]^n$  and p a point in  $F(x_0)$ , the equivalence relation  $\sim$  is given by

$$(f_n, \dots, f_1; t_1, \dots, t_n; p) \sim \begin{cases} (f_n, \dots, f_2; t_2, \dots, t_n; p) & f_1 = \text{Id} \\ (f_n, \dots, f_{i+1}, f_{i-1}, \dots, f_1; t_1, \dots, t_{i-1}t_i, \dots, t_n; p) & f_i = \text{Id}, i > 1 \\ (f_n, \dots, f_{i+1}; t_{i+1}, \dots, t_n; F(f_i, \dots, f_1)(t_1, \dots, t_{i-1}, p)) & t_i = 0 \\ (f_n, \dots, f_{i+1} \circ f_i, \dots, f_1; t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_n; p) & t_i = 1, i < n \\ (f_{n-1}, \dots, f_1; t_1, \dots, t_{n-1}; p) & t_n = 1 \\ * & p = *, \end{cases}$$

where \* denotes the basepoint.

Observation 4.12. The first three cases in the compatibility condition (4.7) for a homotopy coherent diagram imply that F is determined by its values on sequences of non-identity morphisms (and on objects). If one restricts to only non-identity morphisms, however, the compatibility condition for F becomes more complicated. In the special case that  $\mathscr C$  has no isomorphisms except for identity maps, however, the compatibility condition for sequences  $(f_n, \ldots, f_1)$  of non-identity morphisms is simply the last two cases of Formula (4.7).

Similarly, the first two relations in the definition of  $\sim$  mean that we can write

$$\operatorname{hocolim} F = \{*\} \coprod \coprod_{\substack{n \geq 0}} \coprod_{\substack{x_0 \xrightarrow{f_1} \dots \xrightarrow{f_n} x_n \\ \forall i \in \{1, \dots, n\}, \ f_i \neq \operatorname{Id}}} [0, 1]^n \times F(x_0) / \sim',$$

for some equivalence relation  $\sim'$ , the difference being that we consider only non-identity morphisms when n > 0. The equivalence relation  $\sim'$  is more complicated than  $\sim$ . In the special case that  $\mathscr{C}$  has no isomorphisms except for identity maps,  $\sim'$  is simply given by the last four cases of the definition of  $\sim$ .

In the homotopy coherent diagrams and homotopy colimits considered in this paper, the categories  $\mathscr{C}$  will have no non-identity isomorphisms, and so we will work with these smaller formulations.

There is a notion of a morphism between homotopy coherent diagrams (h-morphisms [Vog73, Definition 2.7])  $F, G: \mathscr{C} \to \mathsf{Top}_{\bullet}$ , which relaxes the notion of a morphism (natural transformation) between diagrams. In particular, a morphism  $F \to G$  of homotopy coherent diagrams includes the data of maps  $F(v) \to G(v)$ 

for each  $v \in \mathrm{Ob}(\mathscr{C})$ ; we call these maps the underlying maps of the morphism. There is also the notion of a (simplicial) homotopy of morphisms [Vog73, Definition 2.7], and hence the notion of a homotopy equivalence of homotopy coherent diagrams. A special case is that any morphism of homotopy coherent diagrams whose underlying maps are homotopy equivalences is a homotopy equivalence of diagrams [Vog73, Proposition 4.6]. Further:

**Proposition 4.13.** [Vog73, Theorem 5.12] If  $F,G:\mathscr{C}\to \mathsf{Top}_{\bullet}$  are homotopy equivalent diagrams then hocolim  $F\simeq \mathrm{hocolim}\, G$ .

There is also a rectification result, that any homotopy coherent diagram can be made coherent:

**Proposition 4.14.** [Vog73, Theorem 5.6] Given any homotopy coherent diagram  $F: \mathcal{C} \to \mathsf{Top}_{\bullet}$  there is an honest diagram  $G: \mathcal{C} \to \mathsf{Top}_{\bullet}$  which is homotopy equivalent to F.

Finally:

**Proposition 4.15.** [Vog73, Section 8] If  $F: \mathscr{C} \to \mathsf{Top}_{\bullet}$  is an honest diagram then the homotopy colimits of F, viewed as an honest diagram and as a homotopy coherent diagram, are homotopy equivalent.

**Corollary 4.16.** Properties (ho-1)-(ho-4), or their obvious analogues, hold for homotopy colimits of homotopy coherent diagrams.

### 4.3. Cubical flow categories are functors from the cube to the Burnside category.

Construction 4.17. Fix a cubical flow category  $\mathfrak{f}\colon \mathscr{C} \to \mathscr{C}_C(n)$ . We will construct a strictly unitary, lax 2-functor  $F\colon \underline{2}^n \to \mathscr{B}$ , as follows. By Lemma 4.4, it suffices to define the sets  $X_v$   $(v\in\{0,1\}^n)$ , correspondences  $A_{v,w}$   $(v>w\in\{0,1\}^n)$  and isomorphisms of correspondences  $F_{u,v,w}$   $(u>v>w\in\{0,1\}^n)$ . We do so as follows:

- Given  $v \in \{0,1\}^n$  define  $F(v) = \mathfrak{f}^{-1}(v)$ .
- Given v > w, define  $A_{v,w}$  to be the set of path components of

$$\coprod_{\substack{x \in \mathfrak{f}^{-1}(v) \\ y \in \mathfrak{f}^{-1}(w)}} \operatorname{Hom}(x, y).$$

We will write  $\pi_0(X)$  for the set of path components of X. Then the source (respectively target) map  $s: A_{v,w} \to X_v$  (respectively  $t: A_{v,w} \to X_w$ ) is defined by  $s(\pi_0(\operatorname{Hom}(x,y))) = x$  (respectively  $t(\pi_0(\operatorname{Hom}(x,y))) = y$ ).

• Given u > v > w the composition map in  $\mathscr C$  induces a continuous map

$$\circ \colon \left(\coprod_{\substack{y \in \mathfrak{f}^{-1}(v) \\ z \in \mathfrak{f}^{-1}(w)}} \mathrm{Hom}(y,z)\right) \times_{\mathfrak{f}^{-1}(v)} \left(\coprod_{\substack{x \in \mathfrak{f}^{-1}(u) \\ y \in \mathfrak{f}^{-1}(v)}} \mathrm{Hom}(x,y)\right) \to \coprod_{\substack{x \in \mathfrak{f}^{-1}(u) \\ z \in \mathfrak{f}^{-1}(w)}} \mathrm{Hom}(x,z).$$

Taking path components gives a map  $A_{v,w} \times_{X_v} A_{u,v} \to A_{u,w}$ , which we define to be  $F_{u,v,w}$ .

Lemma 4.18. Construction 4.17 defines a strictly unitary, lax 2-functor.

*Proof.* By Lemma 4.4, we only need to check the compatibility Condition (CF-1), which is immediate from associativity of composition in  $\mathscr{C}$ .

Construction 4.19. Fix a strictly unitary, lax 2-functor  $F: \underline{2}^n \to \mathcal{B}$ . We will construct a cubical flow category  $\mathfrak{f}: \mathcal{C} \to \mathcal{C}_C(n)$ , as follows.

•  $Ob(\mathscr{C}) = \coprod_{v \in \{0,1\}^n} F(v)$ . The functor  $\mathfrak{f}$  sends an object  $x \in F(v)$  to v.




FIGURE 4.1. An example showing that the Lee complex does not come from a functor  $2^n \to \mathcal{B}$  that extends  $F_{Kh}$ . Left: A particular diagram for the two-component unlink, and an ordering of its crossings. Center: The corresponding resolution configuration (the  $\vec{0}$ -resolution with dashed lines recording the crossings) and two labelings of this resolution. Both labelings are in the image of  $F(\varphi_{1100,0000})$  of the labeling  $x_-$  of the circle in the (1,1,0,0)-resolution (right). Further, the two labelings give incompatible restrictions on the map  $F_{(1111,1110,1100)}^{-1} \circ F_{(1111,1101,1100)}$  associated to the subcube (1,1,\*,\*).

- For any object x, Hom(x, x) consists of the identity morphism.
- Given objects x and y, with v = f(x) > f(y) = w, consider the set

$$B_{x,y} = s^{-1}(x) \cap t^{-1}(y) \subset A_{v,w} = F(\varphi_{v,w}).$$

Define  $\operatorname{Hom}(x,y) = B_{x,y} \times \mathcal{M}_{\mathscr{C}_{C}(n)}(v,w)$ . The map  $\mathfrak{f} \colon \operatorname{Hom}(x,y) \to \operatorname{Hom}(\mathfrak{f}(x),\mathfrak{f}(y))$  is projection to the permutohedron  $\mathcal{M}_{\mathscr{C}_{C}(n)}(v,w)$ .

• Given objects x, y, z with f(x) > f(y) > f(z) define the composition map  $\operatorname{Hom}(y, z) \times \operatorname{Hom}(x, y) \to \operatorname{Hom}(x, z)$  as follows. Let u = f(x), v = f(y), w = f(z). The 2-functor includes a map  $F_{u,v,w} \colon A_{v,w} \times_{F(v)} A_{u,v} \to A_{u,w}$ . The composition map in  $\mathscr{C}_C(n)$  gives a map  $\circ \colon \mathcal{M}_{\mathscr{C}_C(n)}(v, w) \times \mathcal{M}_{\mathscr{C}_C(n)}(u, v) \to \mathcal{M}_{\mathscr{C}_C(n)}(u, w)$ . Define the composition map in  $\mathscr{C}$  to be

$$F_{u,v,w} \times \circ \colon (B_{y,z} \times \mathcal{M}_{\mathscr{C}_C(n)}(v,w)) \times (B_{x,y} \times \mathcal{M}_{\mathscr{C}_C(n)}(u,v)) \to (B_{x,z} \times \mathcal{M}_{\mathscr{C}_C(n)}(u,w)).$$

(That is, we apply  $F_{u,v,w}$  to the B factors and  $\circ$  to the M factors.)

**Lemma 4.20.** Construction 4.19 defines a cubical flow category.

*Proof.* The proof is similar to the proof of Lemma 3.17, and is left to the reader.

It is straightforward to verify that Constructions 4.17 and 4.19 are inverse to each other, in a sense which does not seem worth spelling out precisely.

Example 4.21. Applying Construction 4.17 to the Khovanov flow category from Example 3.24 yields a functor  $F_{Kh}: \underline{2}^n \to \mathcal{B}$ . For any  $v \in \{0,1\}^n$ , the set  $F_{Kh}(v)$  consists of the Khovanov generators over v, denoted F(v) in Section 2.2. For  $u > v \in \{0,1\}^n$  with |u| - |v| = 1, and for  $x \in F_{Kh}(u), y \in F_{Kh}(v)$ , the set

$$B_{x,y} = s^{-1}(x) \cap t^{-1}(y) \subseteq A_{u,v} = F_{Kh}(\varphi_{u,v})$$

consists of one element if x appears in  $\delta_{Kh}(y)$  (see Definition 2.1), and is empty otherwise. The maps  $F_{u,v,w}$  when |u| - |w| = 2 are defined using the *ladybug matching* [LS14a, Section 5.4]; see also Section 8.1.

Remark 4.22. In Section 2.2, we discussed a generalization of the Khovanov theory that works over the ring  $\mathbb{Z}[h,t]$ : a functor from  $(2^n)^{\mathrm{op}}$  to the category of (graded)  $\mathbb{Z}[h,t]$ -modules. Setting h=t=0 recovers Khovanov's original functor to  $\mathbb{Z}$ -Mod [Kho00], setting (h,t)=(0,1) gives the theory studied by Lee [Lee05], and setting (h,t)=(1,0) gives a theory introduced by Bar-Natan [BN05].

There is a functor  $(\mathscr{B})^{\mathrm{op}} \to \mathbb{Z}$ -Mod given as follows: to a set X, associate the abelian group  $\mathbb{Z}\langle X\rangle$  freely generated by the elements of X; to a correspondence (A, s, t) from X to Y, associate the following map  $\mathbb{Z}\langle Y\rangle \to \mathbb{Z}\langle X\rangle$ :

$$(4.23) y \mapsto \sum_{x \in X} \# |\{a \in A \mid s(a) = x, t(a) = y\}| x.$$

Example 4.21 lifts Khovanov's functor  $(\underline{2}^n)^{\text{op}} \to \mathbb{Z}$ -Mod to the functor  $F_{Kh} : \underline{2}^n \to \mathscr{B}$ . It is natural to ask whether any other specializations of h and t comes from a strictly unitary, lax 2-functor  $\underline{2}^n \to \mathscr{B}$ . Any candidate must have h = 0 and  $t \in \mathbb{N}$ , since the coefficients in Equation (4.23) need to be positive and integral. The special case (h, t) = (0, 1) (i.e., Lee's theory) does not come from any functor  $\underline{2}^n \to \mathscr{B}$  for arbitrary link diagrams extending the functor  $F_{Kh}$ , as can be seen by considering the diagram in Figure 4.1. The question of whether there is such an extension for h = 0, t > 1 is, as far as we know, open.

4.4. The thickened diagram. Fix a small category  $\mathscr{D}$ , which we regard as a strict 2-category whose only 2-morphisms are identity maps. Fix also a strictly unitary, lax 2-functor  $F: \mathscr{D} \to \mathscr{B}$ , i.e., a  $\mathscr{D}$ -diagram in  $\mathscr{B}$ . In this section, we will associate to  $(\mathscr{D}, F)$  a new (1-)category  $\widehat{\mathscr{D}}$  and, for each  $k \geq 1$ , an (honest) functor  $\widehat{F}_k: \widehat{\mathscr{D}} \to \mathsf{Top}_{\bullet}$ , the category of based topological spaces. There will also be natural transformations  $\Sigma \circ \widehat{F}_k \to \widehat{F}_{k+1}$  (where  $\Sigma$  denotes suspension), so that we get an induced diagram  $\widehat{F}: \widehat{\mathscr{D}} \to \mathscr{S}$ , the category of symmetric spectra. To realize a functor from the cube category to the Burnside category we will apply this construction and then take an iterated mapping cone; see Section 4.5.

We start by defining  $\widehat{\mathscr{D}}$ :

**Definition 4.24.** Let  $\mathscr{D}$  be a small category. The *thickening* of  $\mathscr{D}$  is the small category  $\widehat{\mathscr{D}}$  defined as follows:

- The objects of  $\widehat{\mathcal{D}}$  are composable pairs of morphisms  $u \xrightarrow{f} v \xrightarrow{g} w$  in  $\mathscr{D}$ .
- The morphisms in  $\widehat{\mathscr{D}}$  are commutative diagrams: given composable pairs  $u \xrightarrow{f} v \xrightarrow{g} w$  and  $u' \xrightarrow{f'} v' \xrightarrow{g'} w'$ ,

$$\operatorname{Hom}((f,g),(f',g')) = \Big\{ (\alpha \colon u \to u',\beta \colon v' \to v, \gamma \colon w \to w') \mid \begin{array}{c} u \xrightarrow{f} v \xrightarrow{g} w \\ \downarrow^{\alpha} & \uparrow_{\beta} & \downarrow^{\gamma} \\ u' \xrightarrow{f'} v' \xrightarrow{g'} w' \end{array} \text{ commutes} \Big\}.$$

(Note the direction of the middle vertical arrow.)

• Composition of morphisms is given by stacking diagrams vertically:  $(\alpha', \beta', \gamma') \circ (\alpha, \beta, \gamma) = (\alpha' \circ \alpha, \beta \circ \beta', \gamma' \circ \gamma)$ .

Example 4.25. For  $\mathscr{D}=\underline{2}^1$ ,  $\widehat{\mathscr{D}}$  has four objects:  $1\to 1\to 1,\ 1\to 1\to 0,\ 1\to 0\to 0$  and  $0\to 0\to 0$ . There are unique morphisms

$$(1 \to 1 \to 1) \longrightarrow (1 \to 1 \to 0) \longleftarrow (1 \to 0 \to 0) \longrightarrow (0 \to 0 \to 0).$$

(Again, note the direction of the middle arrow.)

The thickening operation respects products:

**Lemma 4.26.** Given small categories  $\mathscr C$  and  $\mathscr D$ , there is an isomorphism  $q\colon \widehat{\mathscr C}\times\widehat{\mathscr D}\stackrel{\cong}{\longrightarrow} \widehat{\mathscr C}\times \mathscr D$  given on objects by

$$q\big([u_C \overset{f_C}{\to} v_C \overset{g_C}{\to} w_C] \times [u_D \overset{f_D}{\to} v_D \overset{g_D}{\to} w_D]\big) = [u_C \times u_D \overset{f_C \times f_D}{\longrightarrow} v_C \times v_D \overset{g_C \times g_D}{\longrightarrow} w_C \times w_D].$$

*Proof.* This is immediate from the definitions.

Next we define  $\widehat{F}_k$ .

**Definition 4.27.** Let  $\mathscr{D}$  be a small category, which we regard as a 2-category with only identity 2-morphisms and let  $F: \mathscr{D} \to \mathscr{B}$  be a strictly unitary, lax 2-functor. Given an integer  $k \geq 0$  we define a new functor  $\widehat{F}_k: \widehat{\mathscr{D}} \to \mathsf{Top}_{\bullet}$  (of 1-categories) as follows.

On objects, let

$$\widehat{F}_k(u \xrightarrow{f} v \xrightarrow{g} w) = \bigvee_{\substack{a \in F(f) \\ s(b) = t(a)}} \prod_{\substack{b \in F(g) \\ s(b) = t(a)}} S^k.$$

(Here,  $s: F(g) \to F(v)$  and  $t: F(f) \to F(v)$  are the source and target maps of the correspondences F(g) and F(f), respectively.)

To define  $\widehat{F}_k$  on morphisms fix a commutative diagram

$$\begin{array}{ccc}
 u & \xrightarrow{f} v & \xrightarrow{g} w \\
\downarrow^{\alpha} & \uparrow^{\beta} & \downarrow^{\gamma} \\
 u' & \xrightarrow{f'} v' & \xrightarrow{g'} w'.
\end{array}$$

We must construct a map

$$(4.29) \qquad \bigvee_{\substack{a \in F(f) \\ s(b) = t(a)}} S^k \to \bigvee_{\substack{a' \in F(f') \\ s(b') = t(a')}} S^k.$$

It suffices to construct this map one a at a time, so fix  $a \in F(f)$ . The maps  $F_{u,u',v'}$  and  $F_{u,v',v}$  induce a bijection

$$F(f) \cong F(\beta) \times_{F(v')} F(f') \times_{F(u')} F(\alpha);$$

let  $(y, a', x) \in F(\beta) \times_{F(v')} F(f') \times_{F(u')} F(\alpha)$  be the triple corresponding to a. The map  $\widehat{F}_k$  will send  $\prod_{\substack{b \in F(g) \\ s(b) = t(a')}} S^k$ , the summand corresponding to a, to  $\prod_{\substack{b' \in F(g') \\ s(b') = t(a')}} S^k$ , the summand corresponding to a'.

Next, the maps  $F_{v',v,w}$  and  $F_{v',w,w'}$  induce a bijection

$$F(g') \cong F(\gamma) \times_{F(w)} F(g) \times_{F(v)} F(\beta).$$

Consider the map

$$(4.30) \qquad \prod_{\substack{b \in F(g) \\ s(b) = t(a)}} S^k \xrightarrow{\prod_b \Delta_b} \prod_{\substack{b \in F(g) \\ s(b) = t(a)}} \prod_{\substack{b' = (z, \tilde{b}, \tilde{y}) \in F(g') \\ \tilde{y} = y}} S^k \cong \prod_{\substack{b' \in F(g') \\ b' = (z, b, y) \\ s(b) = t(a)}} S^k,$$

where  $\Delta_b$  is the diagonal map  $S^k \to \prod_{b'} S^k$ . Notice that  $\{b' \in F(g') \mid b' = (z, b, y), \ s(b) = t(a)\}$  is a subset of  $\{b' \in F(g') \mid s(b') = t(a')\}$  since s(b') = s(y) = t(a'). We can extend the map (4.30) to a map

$$\prod_{\substack{b \in F(g) \\ s(b) = t(a)}} S^k \longrightarrow \prod_{\substack{b' \in F(g') \\ s(b') = t(a')}} S^k$$

by mapping to the basepoint in the remaining factors. This is the desired map.

(It can be helpful to think of this argument diagrammatically; see Figure 4.2.)

**Lemma 4.31.** Definition 4.27 defines a functor  $\widehat{F}_k$  whose values have natural actions of the symmetric group  $S_k$ .





FIGURE 4.2. Constructing the functor from the thickened diagram. In the first row of arrows, leftwards arrows are source maps, and rightwards arrows are target maps. All squares are fiber products. To define the map (4.30), fix  $a \in F(f)$  and  $b \in F(g)$ . The element a specifies  $a' \in F(f')$  and  $y \in F(\beta)$ . We then consider the elements  $b' \in F(g')$  which map down to g and g. The diagonal map g corresponds to this set. Any such g has source equal to the target of g'.

We omit the proof, which is straightforward, albeit a bit elaborate.

Example 4.32. Consider the functor  $F: \underline{2}^1 \to \mathcal{B}$  given by  $F(1) = \{x,y\}$ ,  $F(0) = \{z,w\}$ , and  $F(\varphi_{1,0}) = \{a,b,c,d\}$  with s(a) = x, s(b) = s(c) = s(d) = y, t(a) = t(b) = t(c) = z, t(d) = w. Graphically, F is given by:



Recall the thickening  $\underline{\widehat{2}}^1$  from Example 4.25. The induced diagram  $\widehat{F}_k : \underline{\widehat{2}}^1 \to \mathsf{Top}_{\bullet}$  is given by

$$S^k_{x,x} \vee S^k_{y,y} \stackrel{\operatorname{Id} \vee \Delta}{\longrightarrow} (S^k_{x,a}) \vee (S^k_{y,b} \times S^k_{y,c} \times S^k_{y,d}) \longleftarrow S^k_{a,z} \vee S^k_{b,z} \vee S^k_{c,z} \vee S^k_{d,w} \longrightarrow S^k_{z,z} \vee S^k_{w,w}.$$

(Here, for instance, the sphere  $S_{x,a}^k$  corresponds to the pair  $(x,a) \in F(\varphi_{1,1}) \times F(\varphi_{1,0})$  over the object  $1 \to 1 \to 0$ .) We claim that the second map is the inclusion of the k-skeleton. To see this, note that it corresponds to the diagram

$$\begin{array}{c}
1 \xrightarrow{\varphi_{1,0}} 0 \xrightarrow{\varphi_{0,0}} 0 \\
\downarrow \varphi_{1,1} & \downarrow \varphi_{1,0} & \downarrow \varphi_{0,0} \\
1 \xrightarrow{\varphi_{1,1}} 1 \xrightarrow{\varphi_{1,0}} 0.
\end{array}$$

The map decomposes along wedge sums. For b, for instance, we get the map  $S_{b,z}^k \to (S_{y,b}^k \times S_{y,c}^k \times S_{y,d}^k)$  as follows:

$$\prod_{\{z\}} S^k = \left[\prod_{\substack{p \in F(\varphi_{0,0}) \\ s(p) = t(b)}} S^k\right] \xrightarrow{\prod \Delta} \left[\prod_{\substack{q' \in F(\varphi_{1,0}) \\ q' = (r,p,b) \in F(\varphi_{0,0}) \times F(\varphi_{0,0}) \times F(\varphi_{1,0}) \\ s(p) = t(b)}} S^k\right] = \prod_{\{b\}} S^k \hookrightarrow \prod_{\{b,c,d\}} S^k.$$

Similarly, the third map sends  $S_{a,z}^k$ ,  $S_{b,z}^k$  and  $S_{c,z}^k$  by the identity map to  $S_{z,z}^k$  and  $S_{d,w}^k$  by the identity map to  $S_{w,w}^k$ .

Finally, we show that the  $\widehat{F}_k$  induce a functor  $\widehat{F}:\widehat{\mathscr{D}}\to\mathscr{S}$ . Given a based space X, identify  $S^n\wedge X$  as  $[0,1]^n\times X/(\partial[0,1]^n\times X\cup[0,1]^n\times \{*\})$ , where \* is the basepoint in X. Given  $n\geq 0$  and based spaces  $X_i$ , there is a canonical map

$$\sigma^n \colon S^n \wedge \prod_i X_i \to \prod_i S^n \wedge X_i,$$
$$(y, x_1, \dots, x_n) \mapsto ((y, x_1), \dots, (y, x_n)).$$

Given  $k, n \geq 0$ , define a map

$$(4.33) \eta(u \xrightarrow{f} v \xrightarrow{g} w) \colon S^n \wedge \widehat{F}_k(u \xrightarrow{f} v \xrightarrow{g} w) \to \widehat{F}_{n+k}(u \xrightarrow{f} v \xrightarrow{g} w)$$

as the composition

$$(4.34) S^{n} \wedge \widehat{F}_{k}(u \xrightarrow{f} v \xrightarrow{g} w) = S^{n} \wedge \left(\bigvee_{a \in F(f)} \prod_{\substack{b \in F(g) \\ s(b) = t(a)}} S^{k}\right) \cong \bigvee_{a \in F(f)} S^{n} \wedge \prod_{\substack{b \in F(g) \\ s(b) = t(a)}} S^{k}$$

$$\bigvee_{a \in F(f)} \bigcap_{\substack{b \in F(g) \\ s(b) = t(a)}} S^{n+k} = \widehat{F}_{k+n}(u \xrightarrow{f} v \xrightarrow{g} w).$$

**Lemma 4.35.** The maps  $\eta$  are  $S_n \times S_k$ -equivariant and form a natural transformation, in the sense that for each  $u \xrightarrow{f} v \xrightarrow{g} w$  the square

$$S^{n} \wedge \widehat{F}_{k}(u \xrightarrow{f} v \xrightarrow{g} w) \xrightarrow{S^{n} \wedge \widehat{F}_{k} \begin{pmatrix} u \to v \to w \\ \downarrow & \uparrow & \downarrow \\ u' \to v' \to w' \end{pmatrix}} S^{n} \wedge \widehat{F}_{k}(u' \xrightarrow{f'} v' \xrightarrow{g'} w')$$

$$\downarrow \eta(u \xrightarrow{f} v \xrightarrow{g} w) \downarrow \qquad \qquad \downarrow \eta(u' \xrightarrow{f'} v' \xrightarrow{g'} w')$$

$$\widehat{F}_{k+n}(u \xrightarrow{f} v \xrightarrow{g} w) \xrightarrow{\widehat{F}_{k+n} \begin{pmatrix} u \to v \to w \\ \downarrow & \uparrow & \downarrow \\ u' \to v' \to w' \end{pmatrix}} \widehat{F}_{k+n}(u' \xrightarrow{f'} v' \xrightarrow{g'} w')$$

commutes. Further, each of the maps  $\eta(u \xrightarrow{f} v \xrightarrow{g} w)$  induces an isomorphism on  $\pi_i$  for  $0 \le i \le 2k-2$ .

Proof. The proof is straightforward, and is left to the reader.

Corollary 4.36. The functors  $\widehat{F}_k$  and natural transformations  $\eta$  assemble to define a diagram of symmetric spectra  $\widehat{F} \colon \widehat{\mathcal{D}} \to \mathscr{S}$ .

Remark 4.37. Lemma 4.35 amounts to a verification that we can more concisely express  $\hat{F}$  within the category of spectra itself by the formula

$$\widehat{F}(u \xrightarrow{f} v \xrightarrow{g} w) = \bigvee_{\substack{a \in F(f) \\ s(b) = t(a)}} \prod_{\substack{b \in F(g) \\ s(b) = t(a)}} \mathbb{S}.$$

The wedge and product play the roles of coproduct and product in this category.

Remark 4.38. The construction of the thickened functor  $\widehat{F}_k \colon \widehat{\mathscr{D}} \to \mathsf{Top}_{\bullet}$  from a functor  $F \colon \mathscr{D} \to \mathscr{B}$  would work with any based space  $X_k$  in place of the sphere  $S^k$ : the construction only uses properties of the wedge sum and product operations. Further, if the spaces  $X_k$  form a sequential spectrum then there is a corresponding thickened diagram in spectra. However, this construction is scarcely more general than the one we gave: in spectra, the diagram coming from  $(X_k)$  is naturally equivalent to the smash product of the spectrum X and the diagram obtained from the sphere spectrum  $(S^k)$ .

### 4.5. The realization.

Construction 4.39. Let  $\widehat{\underline{2}^n}_+$  be the category obtained from  $\widehat{\underline{2}^n}$  by adding a new object \* and a unique morphism  $(u \to v \to w) \to *$  from each vertex  $(u \to v \to w)$  of  $\widehat{\underline{2}^n}$  with  $w \neq \vec{0}$ .

Given a functor  $F: \underline{2}^n \to \mathscr{B}$ , define  $\widehat{F}^+: \widehat{\underline{2}^n}_+ \to \mathscr{S}$  by  $\widehat{F}^+|_{\widehat{\underline{2}^n}} = \widehat{F}$  and  $\widehat{F}^+(*) = \{\text{pt}\}$ , a single point.

Let |F| be the homotopy colimit of  $\widehat{F}^+$ . We call |F| the realization of F.

Example 4.40. Continuing with Example 4.32, we have

$$|F| = \operatorname{hocolim} \left( \begin{array}{c} \mathbb{S}_{x,x} \vee \mathbb{S}_{y,y} & \longrightarrow (\mathbb{S}_{x,a}) \vee (\mathbb{S}_{y,b} \times \mathbb{S}_{y,c} \times \mathbb{S}_{y,d}) \longleftarrow \mathbb{S}_{a,z} \vee \mathbb{S}_{b,z} \vee \mathbb{S}_{c,z} \vee \mathbb{S}_{d,w} & \longrightarrow \mathbb{S}_{z,z} \vee \mathbb{S}_{w,w} \\ \downarrow \\ \{\operatorname{pt}\} \end{array} \right),$$

where  $\mathbb{S}$  denotes the sphere spectrum.

Instead of  $\widehat{\underline{2}^n}_+$ , it will be convenient, sometimes, to work with the larger enlargement  $\widehat{\underline{2}^n}_\dagger = \left(\widehat{\underline{2}^1}_+\right)^n$  of  $\widehat{\underline{2}^n}$ . Given a functor  $F:\underline{2}^n \to \mathscr{B}$  extend  $\widehat{F}$  to a functor  $\widehat{F}^\dagger:\widehat{\underline{2}^n}_\dagger \to \mathscr{S}$  by setting  $\widehat{F}^\dagger|_{\widehat{\underline{2}^n}} = \widehat{F}$  and  $\widehat{F}^\dagger(d) = \{\text{pt}\}$  if  $d \notin \text{Ob}(\widehat{\underline{2}^n})$ .

**Lemma 4.41.** For any functor  $F: \underline{2}^n \to \mathscr{B}$  there is a stable homotopy equivalence hocolim  $\widehat{F}^+ \simeq \operatorname{hocolim} \widehat{F}^{\dagger}$ .

*Proof.* Consider the functor  $G: \widehat{2^n}_\dagger \to \widehat{2^n}_+$  which is the identity on  $\widehat{2^n}$  and sends all objects not in  $\widehat{2^n}$  to \*. We claim that G is homotopy cofinal. To see this, we divide the computation of the undercategories into three cases:

- (a) The undercategory  $*\downarrow G$  of \* is the full subcategory of  $\widehat{\underline{2}^n}_{\dagger}$  spanned by the objects not in  $\widehat{\underline{2}^n}$ . The nerve of this category is homeomorphic to  $\{\vec{x}=(x_1,\ldots,x_n)\in[0,1]^n\mid \exists i \text{ such that } x_i=0\}$ , which is contractible.
- (b) For any object  $d = (u \to v \to \vec{0})$  of  $\widehat{\underline{2}^n}$ , the undercategory  $d \downarrow G$  of d is the full subcategory of  $\widehat{\underline{2}^n}$  of objects d' for which there is a map  $d \to d'$ . The object d is an initial object for this subcategory, so the nerve is contractible.
- (c) Fix an object  $d = (u \to v \to w)$  of  $\widehat{\underline{2}^n}$  with  $w \neq \overline{0}$ . The undercategory  $d \downarrow G$  of d is the full subcategory of  $\widehat{\underline{2}^n}_{\dagger}$  spanned by the objects d' not in  $\widehat{\underline{2}^n}$  and the objects d' in  $\widehat{\underline{2}^n}$  for which there is

a map  $d \to d'$ . Let  $\mathscr{D}$  be the full subcategory of  $\widehat{2^n}_{\dagger}$  spanned by the objects d' not in  $\widehat{2^n}$  and let  $\mathscr{E}$  be the full subcategory of  $\widehat{2^n}_{\dagger}$  consisting of objects d' for which there is a morphism  $d \to d'$ . The categories  $\mathscr{D}$  and  $\mathscr{E}$  are each downwards closed in  $d \downarrow G$ , i.e., there are no morphisms out of  $\mathscr{D}$  or  $\mathscr{E}$ . Thus, the nerve of  $d \downarrow G$  is the union of the nerves of  $\mathscr{D}$  and  $\mathscr{E}$ , glued along the nerve of  $\mathscr{D} \cap \mathscr{E}$ . We already saw in part (a) that the nerve of  $\mathscr{D}$  is contractible. The category  $\mathscr{E}$  has an initial object, d, and hence the nerve of  $\mathscr{E}$  is contractible. Finally, the realization of the category  $\mathscr{D} \cap \mathscr{E}$  is similar to the realization of  $\mathscr{D}$  (i.e., a union of coordinate hyperplanes), and so contractible. It follows that the realization of  $d \downarrow G$  is contractible.

Thus, the functor G is homotopy cofinal. It is immediate from the definitions that  $\hat{F}^{\dagger} = \hat{F}^{+} \circ G$ , so the result follows from property (ho-4) of homotopy colimits (Section 4.2).


## 4.6. An invariance property of the realization.

**Lemma 4.42.** If  $F,G: \underline{2}^n \to \mathcal{B}$  are naturally isomorphic 2-functors then the realizations of F and G are stably homotopy equivalent.

*Proof.* A natural isomorphism T from F to G specifies:

- A bijection  $T_v : F(v) \to G(v)$  for each  $v \in \{0,1\}^n$ , and
- A bijection  $T_{v,w} : F(\varphi_{v,w}) \to G(\varphi_{v,w})$  for each  $v > w \in \{0,1\}^n$

satisfying the conditions that



and

$$F(\varphi_{v,w}) \times_{F(v)} F(\varphi_{u,v}) \xrightarrow{F_{u,v,w}} F(\varphi_{u,w})$$

$$\downarrow^{T_{v,w} \times T_{u,v}} \qquad \downarrow^{T_{u,w}}$$

$$G(\varphi_{v,w}) \times_{G(v)} G(\varphi_{u,v}) \xrightarrow{G_{u,v,w}} G(\varphi_{u,w})$$

commute. (See [Gra74, Section I,2.4] and note that an isomorphism in  $\mathcal{B}$  between two sets induces a bijection between them.) Given a natural transformation T, there is a corresponding map of diagrams defined as follows. Given u > v > w we want a map

$$\bigvee_{\substack{a \in F(\varphi_{u,v}) \\ s(b) = t(a)}} \prod_{\substack{b \in F(\varphi_{v,w}) \\ s(b) = t(a)}} \mathbb{S} \to \bigvee_{\substack{a \in G(\varphi_{u,v}) \\ s(b) = t(a)}} \prod_{\substack{b \in G(\varphi_{v,w}) \\ s(b) = t(a)}} \mathbb{S}.$$

This map sends the wedge summand corresponding to  $a \in F(\varphi_{u,v})$  to the wedge summand corresponding to  $T_{u,v}(a)$ , and sends the factor corresponding to  $b \in F(\varphi_{v,w})$  to the factor corresponding to  $T_{v,w}(b)$ . It is straightforward to verify that this gives a map of diagrams, which is by definition an isomorphism. It follows that the map induces a stable homotopy equivalence of homotopy colimits. (In fact, if we work with diagrams of  $S^k$ 's, the map would be a homeomorphism.)

4.7. **Products and realization.** In the language of flow categories, the product is a rather complicated object. In the language of functors from the cube to the Burnside category, however, the product is quite simple:

**Definition 4.43.** There is a map  $\times : \mathcal{B} \times \mathcal{B} \to \mathcal{B}$  which sends a pair of objects (X,Y) to their direct product  $X \times Y$ , a pair of correspondences  $((A, s_A, t_A), (B, s_B, t_B))$  to  $(A \times B, s_A \times s_B, t_A \times t_B)$ , and a pair of isomorphisms of correspondences  $(F: A \to A', G: B \to B')$  to the isomorphism  $F \times G: A \times B \to A' \times B'$ . Given functors  $F: \underline{2}^m \to \mathcal{B}$  and  $G: \underline{2}^n \to \mathcal{B}$ , we define the *product*  $F \times G$  of F and G, to be the composition

$$\underline{2}^{m+n} \stackrel{\cong}{\longrightarrow} \underline{2}^m \times \underline{2}^n \stackrel{(F,G)}{\longrightarrow} \mathscr{B} \times \mathscr{B} \stackrel{\times}{\longrightarrow} \mathscr{B}.$$

Explicitly,  $(F \times G): \underline{2}^{m+n} \to \mathcal{B}$  is given as follows. Identify  $\{0,1\}^m \times \{0,1\}^n$  with  $\{0,1\}^{m+n}$ . Then:

• For  $(u_1, u_2) \in \{0, 1\}^m \times \{0, 1\}^n = \{0, 1\}^{m+n}$  define

$$(F \times G)(u_1, u_2) = F(u_1) \times G(u_2).$$

• For  $(u_1, u_2) > (v_1, v_2) \in \{0, 1\}^{m+n}$  define  $(F \times G)(\varphi_{(u_1, u_2), (v_1, v_2)})$  to be the correspondence

$$F(\varphi_{u_1,v_1})\times G(\varphi_{u_2,v_2})$$

$$\downarrow^{s_F\times s_G}$$

$$F(u_1)\times G(u_2)$$

$$F(v_1)\times G(v_2).$$

• For  $(u_1, u_2) > (v_1, v_2) > (w_1, w_2) \in \{0, 1\}^{m+n}$  define  $(F \times G)_{(u_1, u_2), (v_1, v_2), (w_1, w_2)}$  by  $(F \times G)_{(u_1, u_2), (v_1, v_2), (w_1, w_2)}(x, y) = (F_{u_1, v_1, w_1}(x), G_{v_1, v_2, v_3}(y)).$ 

**Lemma 4.44.** Definition 4.43 specifies a strictly unitary, lax 2-functor.

*Proof.* This is immediate.

Note that smash products distribute across wedge sums. Moreover, while  $X \wedge (Y \times Z)$  is not homotopy equivalent to  $(X \wedge Y) \times (X \wedge Z)$ , there is a natural map  $X \wedge (Y \times Z) \rightarrow (X \wedge Y) \times (X \wedge Z)$  defined by  $(x, y, z) \mapsto ((x, y), (x, z))$ . This generalizes to a map

$$p: \left(\prod_{a \in A} X_a\right) \wedge \left(\prod_{b \in B} Y_b\right) \to \prod_{(a,b) \in A \times B} X_a \wedge Y_b.$$

The map p is natural in both factors.

Given functors  $F: \mathscr{C} \to \mathsf{Top}_{\bullet}$  and  $G: \mathscr{D} \to \mathsf{Top}_{\bullet}$  we can take the smash product of F and G to obtain a functor  $F \wedge G: \mathscr{C} \times \mathscr{D} \to \mathsf{Top}_{\bullet}$  (cf. (ho-3)). At a vertex (c,d) of  $\mathscr{C} \times \mathscr{D}$ ,  $(F \wedge G)(c,d) = F(c) \wedge G(d)$ .

**Lemma 4.45.** The thickening construction from Section 4.4 respects products, in the following sense. Fix functors  $F: \underline{2}^m \to \mathcal{B}$  and  $G: \underline{2}^n \to \mathcal{B}$ . Let  $F \times G: \underline{2}^m \times \underline{2}^n \to \mathcal{B}$  be as in Definition 4.43. Let  $q: \widehat{\mathcal{C}} \times \widehat{\mathcal{D}} \to \widehat{\mathcal{C}} \times \widehat{\mathcal{D}}$  be the isomorphism from Lemma 4.26. Then, the map p induces a natural transformation from  $\widehat{F}_k \wedge \widehat{G}_l$  to  $(\widehat{F} \times \widehat{G})_{k+l} \circ q$  so that on vertices the natural transformation is a weak homotopy equivalence up to dimension  $k+l+\min(k,l)-1$ . These natural transformations respect the spectrum structure, and so induce maps of diagrams  $\widehat{F} \wedge \widehat{G} \to (\widehat{F} \times \widehat{G}) \circ q$  so that the map at each vertex is a stable homotopy equivalence.

*Proof.* This is straightforward from the definitions. To illustrate, we describe the natural transformation at the level of vertices. At a vertex  $[u_C \stackrel{f_C}{\to} v_C \stackrel{g_C}{\to} w_C] \times [u_D \stackrel{f_D}{\to} v_D \stackrel{g_D}{\to} w_D]$  we have

$$\widehat{F}(u_C \xrightarrow{f_C} v_C \xrightarrow{g_C} w_C) \wedge \widehat{G}(u_D \xrightarrow{f_D} v_D \xrightarrow{g_D} w_D) = \left( \bigvee_{\substack{a_C \in F(f_C) \\ s(b_C) = t(a_C)}} \prod_{\substack{b_C \in F(g_C) \\ s(b_C) = t(a_D)}} S^k \right) \wedge \left( \bigvee_{\substack{a_D \in G(f_D) \\ s(b_D) = t(a_D)}} \prod_{\substack{b_D \in G(g_D) \\ s(b_D) = t(a_D)}} S^l \right)$$

while

$$(\widehat{F \times G})_{k+l} (u_C \times u_D \overset{f_C \times f_D}{\longrightarrow} v_C \times v_D \overset{g_C \times g_D}{\longrightarrow} w_C \times w_D) = \bigvee_{\substack{a_C \in F(f_C) \\ a_D \in G(f_D) \\ s(b_C) = t(a_C) \\ s(b_D) = t(a_D)}} \sum_{\substack{b_C \in F(g_C) \\ b_D \in G(g_D) \\ s(b_D) = t(a_D)}} S^{k+l}.$$

The map p sends the first to the second in an obvious way, and is an equivalence up to dimension  $k + l + \min(k, l) - 1$ . Moreover, this map is  $S_k \times S_l$ -equivariant and respects the structure maps of the spectrum.  $\square$ 

**Proposition 4.46.** Given functors  $F: \underline{2}^n \to \mathcal{B}$  and  $G: \underline{2}^m \to \mathcal{B}$ , we have  $|F \times G| \simeq |F| \wedge |G|$ .

*Proof.* Let  $\simeq$  denote stable homotopy equivalence. By Lemma 4.41,

$$|F| \simeq \operatorname{hocolim} \widehat{F}^{\dagger}$$
  $|G| \simeq \operatorname{hocolim} \widehat{G}^{\dagger}$   $|F \times G| \simeq \operatorname{hocolim} (\widehat{F \times G})^{\dagger}$ .

From the definitions, it follows that the natural transformation from  $\widehat{F} \wedge \widehat{G}$  to  $(\widehat{F} \times \widehat{G})$  of Lemma 4.45 extends to a natural transformation from  $\widehat{F}^{\dagger} \wedge \widehat{G}^{\dagger}$  to  $(\widehat{F} \times \widehat{G})^{\dagger}$ , which is a stable homotopy equivalence on objects. By point (ho-1) in Section 4.2,

$$\operatorname{hocolim}(\widehat{F \times G})^{\dagger} \simeq \operatorname{hocolim}(\widehat{F}^{\dagger} \wedge \widehat{G}^{\dagger}).$$

By point (ho-3) in Section 4.2,

$$\operatorname{hocolim}\left(\widehat{F}^\dagger \wedge \widehat{G}^\dagger\right) \simeq \left(\operatorname{hocolim}\widehat{F}^\dagger\right) \wedge \left(\operatorname{hocolim}\widehat{G}^\dagger\right).$$

The result follows.

An even simpler operation is disjoint union:


**Definition 4.47.** There is a map  $\coprod : \mathscr{B} \times \mathscr{B} \to \mathscr{B}$  which sends a pair of objects (X,Y) to their disjoint union  $X \coprod Y$ , a pair of correspondences  $((A, s_A, t_A), (B, s_B, t_B))$  to  $(A \coprod B, s_A \coprod s_B, t_A \coprod t_B)$ , and a pair of isomorphisms of correspondences  $(F \colon A \to A', G \colon B \to B')$  to the isomorphism  $F \coprod G \colon A \coprod B \to A' \coprod B'$ . Given functors  $F, G \colon \underline{2}^n \to \mathscr{B}$ , we define the *disjoint union*  $F \coprod G$  of F and G, to be the composition

$$\underline{2^n} \xrightarrow{\Delta} \underline{2^n} \times \underline{2^n} \overset{(F,G)}{\longrightarrow} \mathscr{B} \times \mathscr{B} \xrightarrow{\mathrm{II}} \mathscr{B}.$$

where the first arrow is the diagonal map.

Explicitly,  $(F \coprod G): 2^n \to \mathcal{B}$  is given as follows:

- For  $v \in \{0,1\}^n$ ,  $(F \coprod G)(v) = F(v) \coprod G(v)$ .
- For  $v > w \in \{0,1\}^n$ ,  $(F \coprod G)_{v,w}(\varphi_{v,w})$  is defined to be the correspondence



• For  $u > v > w \in \{0,1\}^n$ ,  $(F \coprod G)_{u,v,w}$  is defined by the commutative diagram

where the vertical arrow is the obvious bijection.

**Lemma 4.48.** Definition 4.47 specifies a strictly unitary, lax 2-functor.

*Proof.* This is immediate from the definitions.

Given diagrams  $F,G\colon\mathscr{C}\to\operatorname{\mathsf{Top}}_{ullet}$  there is an induced diagram  $F\vee G\colon\mathscr{C}\to\operatorname{\mathsf{Top}}_{ullet}$  with  $(F\vee G)(v)=F(v)\vee G(v)$  and  $(F\vee G)(f)=F(f)\vee G(f)$ .

**Lemma 4.49.** The thickening construction respects disjoint unions in the sense that given  $F, G: \mathscr{C} \to \mathscr{B}$ ,  $(\widehat{F \coprod G})_k \cong \widehat{F}_k \vee \widehat{G}_k$ .

*Proof.* Again, this is immediate from the definitions.

**Proposition 4.50.** Given functors  $F, G: 2^n \to \mathcal{B}$ ,  $|F \coprod G| \simeq |F| \vee |G|$ .

*Proof.* Lemma 4.49 extends immediately to the statement that  $\widehat{F \coprod G}^+ \cong \widehat{F}^+ \vee \widehat{G}^+$ , and homotopy colimit commutes with wedge sum (point (ho-2) in Section 4.2).

#### 5. Building a smaller cube from little box maps

In this section we show that the realization of a functor  $\underline{2}^n \to \mathcal{B}$  can be understood in terms of a smaller diagram.

**Definition 5.1.** Let  $\underline{2}_{+}^{n}$  be the result of adding a single object, which we will denote \*, to  $\underline{2}^{n}$  and declaring that

$$\operatorname{Hom}(v,*) = \begin{cases} \text{one element } \varphi_{v,*} & v \neq \vec{0} \\ \varnothing & v = \vec{0} \end{cases} \qquad \operatorname{Hom}(*,v) = \begin{cases} \{\operatorname{Id}\} & v = * \\ \varnothing & v \neq *. \end{cases}$$

In other words,  $\underline{2}_{+}^{n}$  is the result of replacing the terminal object in  $\underline{2}^{n}$  by two copies of itself, with no maps between them.

The goal of this section is to prove:

**Theorem 7.** Fix a strictly unitary, lax 2-functor  $F: \underline{2}^n \to \mathcal{B}$ . Then there is a homotopy coherent diagram  $\widetilde{F}^+: \underline{2}^n_+ \to \mathcal{S}$  so that:

- (1) For each  $v \in \text{Ob}(\underline{2}^n)$ ,  $\widetilde{F}^+(v) = \bigvee_{a \in F(v)} \mathbb{S}$ .
- (2)  $\widetilde{F}^+(*)$  is a single point.
- (3) hocolim  $\widetilde{F}^+ \simeq |F|$ .

The diagram  $\widetilde{F}^+$  is a special case of a more general construction, which we give in Section 5.2. We specialize to  $\widetilde{F}^+$  in Section 5.3, and prove that this homotopy colimit is equivalent to |F| in Section 5.4. Before turning to these constructions, we introduce a particular class of maps between spheres in Section 5.1.




FIGURE 5.1. **Spaces of boxes.** Left: A correspondence A from a 2-element set X to another 2-element set Y. Center: A point in  $E(\{B_x\}, s)$ . Right: A point in  $F(\{B_x\}, s, t)$  which is not in  $E(\{B_x\}, s)$ . The boxes  $B_{a_i}$  are shaded.

## 5.1. Box maps.

**Definition 5.2.** By a box in  $\mathbb{R}^k$  we mean a subset of the form  $B = [a_1, b_1] \times \cdots \times [a_k, b_k]$  for some  $a_1, \ldots, b_k$  with  $a_i < b_i$ . Given a box B in  $\mathbb{R}^k$ , a sub-box B' of B is a box B' in  $\mathbb{R}^k$  so that  $B' \subset B$ .

**Definition 5.3.** Given a box B in  $\mathbb{R}^k$ , let  $E(B,\ell)$  be the space of ordered  $\ell$ -tuples of disjoint boxes in B, topologized as a subspace of  $(\mathbb{R}^{2k})^{\ell}$ .

Given a correspondence  $(A, s: A \to X, t: A \to Y)$ , consider a collection  $\{B_x \mid x \in X\}$  of distinct copies of the standard unit box. Let  $F(\{B_x\}, s, t)$  be the space of collections  $\{B_a \subset B_{s(a)} \mid a \in A\}$  of sub-boxes labeled by elements of A, satisfying the condition that  $B_a \cap B_{a'} = \emptyset$  if  $a \neq a'$  and s(a) = s(a') and t(a) = t(a'). See Figure 5.1.

Let  $E(\{B_x\}, s) \subset F(\{B_x\}, s, t)$  be the space of collections  $\{B_a \subset B_{s(a)} \mid a \in A\}$  satisfying the stronger condition that  $B_a \cap B_{a'} = \emptyset$  if  $a \neq a'$  and s(a) = s(a'). Again, see Figure 5.1.

Example 5.4. If A is an  $\ell$ -element correspondence and X has one element x then  $E(\{B_x\}, s) = E(B_x, \ell)$ .

**Definition 5.5.** Fix once and for all an identification  $S^k = [0,1]^k/\partial$ . When we view  $S^k$  as a pointed space, we assume that this identification identifies the basepoint and  $\partial [0,1]^k$ . For any box B in  $\mathbb{R}^k$ , there is a standard homeomorphism  $B \to [0,1]^k$ , given by  $(x_1,\ldots,x_k) \mapsto (\frac{x_1-a_1}{b_1-a_1},\ldots,\frac{x_k-a_k}{b_k-a_k})$ , an hence an induced identification  $S^k \cong B/\partial B$ .

Given  $\{B_1, \ldots, B_\ell\} \in E(B, \ell)$  the associated basic box map is the composition

$$(5.6) S^k = B/\partial B \to B/(B \setminus (\mathring{B}_1 \cup \dots \cup \mathring{B}_\ell)) = \bigvee_{a=1}^\ell B_a/\partial B_a = \bigvee_{a=1}^\ell S^k \to S^k,$$

where the last map is the identity on each summand (so the composition has degree  $\ell$ ). This defines a continuous map  $E(B,\ell) \to Map(S^k,S^k)$ .

Remark 5.7. The map  $E(B, \ell) \to Map(S^k, S^k)$  is the composition of the coaction of the little k-cubes operad with the fold map.

**Definition 5.8.** Given a correspondence A from X to Y and a collection of boxes  $e = \{B_a \subset B_{s(a)} \mid a \in A\} \in E(\{B_x\}, s)$ , there is an induced map

(5.9) 
$$\Phi(e,A) \colon \bigvee_{x \in X} S^k \to \bigvee_{y \in Y} S^k.$$

defined by

$$(5.10) \qquad \Phi_{e,A}|_{S_x^k} \colon S_x^k = B_x/\partial B_x \to B_x/\left(B_x \setminus \left(\bigcup_{\substack{a \in A \\ s(a) = x}} \mathring{B_a}\right)\right) = \bigvee_{\substack{a \in A \\ s(a) = x}} B_a/\partial B_a = \bigvee_{\substack{a \in A \\ s(a) = x}} S_a^k \to \bigvee_{y \in Y} S_y^k$$

where the last map sends  $S_a^k$  by the identity map to  $S_{t(a)}^k$ . This defines a continuous map  $\Phi(\cdot, A) : E(\{B_x\}, s) \to Map(\bigvee_{x \in X} S^k, \bigvee_{y \in Y} S^k)$ . We call a map of the form  $\Phi(e, A)$  for some  $e \in E(\{B_x\}, s)$  a disjoint box map, and say it refines the correspondence (A, s, t).

Example 5.11. The basic box maps are exactly the disjoint box maps where X and Y each consists of a single element.

**Definition 5.12.** Given a correspondence A from X to Y and a collection of boxes  $e = \{B_a \subset B_{s(a)} \mid a \in A\} \in F(\{B_x\}, s, t)$ , there is an induced map

$$(5.13) \qquad \bigvee_{x \in X} S^k \to \prod_{y \in Y} S^k$$

whose component sending  $S_x^k$  to  $S_y^k$  is the basic box map associated to  $(B_x, \{B_a \mid s(a) = x, t(a) = y\})$ . We call a map arising this way an overlapping box map, and say it refines the correspondence (A, s, t).

By a box map we mean either a disjoint box map or an overlapping box map.

Example 5.14. Given  $e \in E(\{B_x\}, s) \subset F(\{B_x\}, s, t)$ , the overlapping box map associated to e is obtained by composing the disjoint box map  $\Phi(e, A)$  with the standard inclusion  $\bigvee_{y \in Y} S^k \hookrightarrow \prod_{y \in Y} S^k$ .

Remark 5.15. We have not defined, and will not need, box maps from products, just box maps to products.

Example 5.16. The diagonal map  $S^k \to \prod_{i=1}^m S^k$  is a box map: take  $X = \{x\}$  to have a single element, A and Y to have m elements each, t to be a bijection, and each box  $B_a$  to be all of  $B_x$ .

**Lemma 5.17.** A composition of box maps is a box map.

*Proof.* Given composable box maps F and G, the preimages under F of the boxes for G are the boxes for  $G \circ F$ .

The value to us of these constructions comes from the following:

**Lemma 5.18.** If B is k-dimensional then the space  $E(B,\ell)$  is (k-2)-connected. More generally,  $E(\{B_x\},s)$  and  $F(\{B_x\},s,t)$  are (k-2)-connected.

Proof. The space  $E(B,\ell)$  is homotopy equivalent to the ordered configuration space of  $\ell$  points in the interior of B, i.e.,  $B^{\ell} \setminus \Delta$ , where  $\Delta$  is the fat diagonal. Since  $\Delta$  is a finite union of smooth submanifolds of codimension k, the result for  $E(B,\ell)$  follows. Next,  $E(\{B_x\},s) \cong \prod_{x\in X} E([0,1]^k,|s^{-1}(x)|)$  is a product of (k-2)-connected spaces, and hence is (k-2)-connected. Finally,  $F(\{B_x\},s,t) \cong \prod_{(x,y)\in X\times Y} E([0,1]^k,|s^{-1}(x)|)$   $t^{-1}(y)$ , and so again is (k-2)-connected.

Informally, Lemma 5.18 says that the space of box maps is highly connected. (For this statement to be correct, we must think of a box map as not just the map but also the labeling of the boxes.) Indeed, we will show next that the space of box maps is also highly connected in relative terms.

**Definition 5.19.** For any set map  $s: A \to X$ , define  $E^{\circ}(\{B_x\}, s)$  to be the subspace of  $E(\{B_x\}, s)$  where we require the box  $B_a$  to lie in the interior of the box  $B_{s(a)}$  for all  $a \in A$ .

If  $A_0 \subset A$  is a subset and  $s_0 \colon A_0 \to X$  is the restriction of s then there is a map  $E^{\circ}(\{B_x\}, s) \to E^{\circ}(\{B_x\}, s_0)$  gotten by forgetting the boxes labeled by  $A \setminus A_0$ .

**Lemma 5.20.** Fix a set map  $s: A \to X$  and a subset  $A_0 \subset A$ . Let  $s_0 = s|_{A_0}$ . If the boxes  $\{B_x\}$  are k-dimensional, then for any  $i \leq k-1$  and any commutative diagram

$$S^{i-1} \longrightarrow E^{\circ}(\{B_x\}, s)$$

$$\downarrow$$

$$D^i \longrightarrow E^{\circ}(\{B_x\}, s_0),$$

there exists a lift  $D^i \to E^{\circ}(\{B_x\}, s)$  making the diagram commute.

*Proof.* By induction, we may assume that  $A \setminus A_0$  consists of a single element, say  $a_1$ , and let  $x_1 = s(a_1)$  and  $\ell = |s_0^{-1}(x_1)|$ . The projection  $\pi \colon E^{\circ}(\{B_x\}, s) \to E^{\circ}(\{B_x\}, s_0)$  is seen to be a fiber bundle by the following argument.

Let  $\pi_{\bullet} : E_{\bullet} \to E^{\circ}(\{B_x\}, s_0)$  be the fiber bundle where the fiber over a point is the complement of the  $\ell$  boxes in the interior of  $B_{x_1}$ . It is easy to see that  $\pi_{\bullet}$  is a fiber bundle as follows. For  $z \in E^{\circ}(\{B_x\}, s_0)$ , fix a minimal triangulation of the complement of the  $\ell$  boxes in z. This triangulation persists in some small neighborhood U of z in  $E^{\circ}(\{B_x\}, s_0)$ , and induces a PL homeomorphism between  $\pi_{\bullet}^{-1}(z)$  and  $\pi_{\bullet}^{-1}(z')$  for any  $z' \in U$ .

Now for any  $z \in E^{\circ}(\{B_x\}, s_0)$ , construct a coordinate chart on  $\pi^{-1}(z)$  by the following variables: the center C of the box  $B_{a_1}$  viewed as a point in  $\pi_{\bullet}^{-1}(z)$ ; the 'aspect ratio' R of the box  $B_{a_1}$ , presented as a (k-1)-tuple of ratios of the k side-lengths of  $B_{a_1}$ ; and a proportion  $P \in (0,1)$  of the volume of the box  $B_{a_1}$  relative to the volume of the largest box with the same center and same aspect ratio that lies in  $B_{x_1}$  in the complement of the interiors of other  $\ell$  boxes. This identifies  $\pi^{-1}(z)$  with  $\pi_{\bullet}^{-1}(z) \times (0,\infty)^{k-1} \times (0,1)$ , and the identification holds for small open sets around z. But  $\pi_{\bullet}$  is a fiber bundle, and therefore, so is  $\pi$ .

The fiber over each point is homeomorphic to the space of boxes in the complement of  $\ell$  disjoint boxes in the interior of  $B_{x_1}$ . The fiber, being homotopy equivalent to the complement of  $\ell$  points in  $\mathbb{R}^k$ , is (k-2)-connected, so the statement follows.

#### 5.2. Refining diagrams via box maps.

**Definition 5.21.** Fix a small category  $\mathscr{D}$  and a strictly unitary, lax 2-functor  $F: \mathscr{D} \to \mathscr{B}$  (i.e., a  $\mathscr{D}$ -diagram in  $\mathscr{B}$ ). A k-dimensional spatial refinement of F is a homotopy coherent diagram  $\widetilde{F}_k: \mathscr{D} \to \mathsf{Top}_{\bullet}$  so that

• For any 
$$u \in \text{Ob}(\mathscr{D})$$
,  $\widetilde{F}_k(u) = \bigvee_{x \in F(u)} S^k = \left(\coprod_{x \in F(u)} B_x\right)/\partial;$ 

- For any  $u, v \in \text{Ob}(\mathcal{D})$  and  $f: u \to v$ ,  $\widetilde{F}_k(f)$  is a disjoint box map which refines the correspondence F(f) from F(u) to F(v) (see Section 5.1); and, more generally,
- For any sequence of morphisms

$$u_0 \xrightarrow{f_1} u_1 \xrightarrow{f_2} \cdots \xrightarrow{f_n} u_n$$

in  $\mathcal{D}$ ,

$$\widetilde{F}_k(f_n,\ldots,f_1)\colon [0,1]^{n-1}\to Map(\bigvee_{x\in F(u_0)}S^k,\bigvee_{x\in F(u_n)}S^k)$$

is a family of box maps induced by a map  $[0,1]^{n-1} \to E(\{B_x\}_{x \in F(u_0)}, s_{F(f_n \circ \cdots \circ f_0)})$  (refining the correspondence

$$F(f_n \circ \cdots \circ f_1) \cong F(f_n) \times_{F(u_{n-1})} \cdots \times_{F(u_1)} F(f_1)$$

from  $F(u_0)$  to  $F(u_n)$ ).

**Proposition 5.22.** Let  $\mathscr{D}$  be a small category in which every sequence of composable non-identity morphisms has length at most n. Fix a  $\mathscr{D}$ -diagram F in  $\mathscr{B}$ .

- (1) If  $k \geq n$  then there is a k-dimensional spatial refinement of F.
- (2) If  $k \ge n+1$  then any two k-dimensional spatial refinements of F are homotopic (as homotopy coherent diagrams).
- (3) If  $\widetilde{F}_k$  is a k-dimensional spatial refinement of F then the result of suspending each  $\widetilde{F}_k(u)$  and  $\widetilde{F}_k(f_n,\ldots,f_1)(\vec{t})$  gives a (k+1)-dimensional spatial refinement of F.

Proof. We start with point (1). Given  $u \in \text{Ob}(\mathcal{D})$ , define  $\widetilde{F}_k(u) = \bigvee_{x \in F(u)} S^k$ ; write the  $S^k$  summand corresponding to x as  $B_x/\partial$ , where  $B_x$  is a box in  $\mathbb{R}^k$  (e.g.,  $B_x = [0,1]^k$ ). Next, by Observation 4.12 it suffices to consider only non-identity morphisms. For each non-identity morphism  $f \colon u \to v$  in  $\mathcal{D}$  choose a disjoint box map which refines the correspondence F(f). Let  $e_f \in E(\{B_x \mid x \in F(u)\}, s_{F(f)})$  be the collection of little boxes corresponding to F(f).

We have now defined  $F_k$  on vertices and arrows. The diagram does not commute, so it remains to define the coherence homotopies associated to sequences of composable morphisms. We will build these inductively. As a warm up, we spell out the first case carefully before proceeding to the general case.

Fix a composable pair of morphisms  $u \xrightarrow{f} v \xrightarrow{g} w$  in  $\mathscr{D}$ . There are two points in  $E(\{B_x \mid x \in F(u)\}, s_{F(g \circ f)})$  associated to (g, f). One is the point  $e_{g \circ f}$ . The other is defined as follows. The point  $e_g$  corresponds to a collection of boxes  $\mathcal{B}_g$  in  $\{B_y \mid y \in F(v)\}$ , labeled by elements of F(g). The inverse image  $\Phi(e_f, F(f))^{-1}(\mathcal{B}_g)$  of these boxes is a collection of boxes in  $\{B_x \mid x \in F(u)\}$ . The boxes  $\Phi(e_f, F(f))^{-1}(\mathcal{B}_g)$  inherit a labeling by elements of  $F(g) \times_{F(v)} F(f) \cong_{F_{u,v,w}} F(g \circ f)$ . This labeling makes  $\Phi(e_f, F(f))^{-1}(\mathcal{B}_g)$  into a second point in  $E(\{B_x \mid x \in F(u)\}, s_{F(g \circ f)})$ , which by abuse of notation we will call  $e_g \circ e_f$  (see also Lemma 5.17).

By Lemma 5.18, since  $k \geq 2$  (or else we would not have a composable pair (g, f)), the space  $E(\{B_x \mid x \in F(u)\}, s_{F(g \circ f)})$  is connected, so we can find a path from  $e_{g \circ f}$  to  $e_g \circ e_f$ . Fix such a path, and call it  $e_{g,f} : [0,1] \to E(\{B_x \mid x \in F(u)\}, s_{F(g \circ f)})$ . Then  $e_{g,f}$  defines a homotopy  $\Phi(e_{g,f}, F(g \circ f))$  from  $\Phi(e_g, F(g)) \circ \Phi(e_f, F(f))$  to  $\Phi(e_{g \circ f}, F(g \circ f))$ .

More generally, suppose that for any sequence  $v_0 \xrightarrow{f_1} \cdots \xrightarrow{f_\ell} v_\ell$  of non-identity morphisms we have chosen a map  $e_{f_\ell,\dots,f_1} : [0,1]^{\ell-1} \to E(\{B_x \mid x \in F(v_0)\}, s_{F(f_\ell \circ \dots \circ f_1)})$ , and these maps are compatible in the following sense. Let  $(t_1,\dots,t_{\ell-1})$  be the coordinates on  $[0,1]^{\ell-1}$ . Then for any  $1 \le i \le \ell-1$  we require that:

(5.23) 
$$e_{f_{\ell},\dots,f_{1}}(t_{1},\dots,t_{i-1},0,t_{i+1},\dots,t_{\ell-1}) = e_{f_{\ell},\dots,f_{i}}(t_{i+1},\dots,t_{\ell-1}) \circ e_{f_{i-1},\dots,f_{1}}(t_{1},\dots,t_{i-1}) \\ e_{f_{\ell},\dots,f_{1}}(t_{1},\dots,t_{i-1},1,t_{i+1},\dots,t_{\ell-1}) = e_{f_{\ell},\dots,f_{i}}\circ f_{i-1},\dots,f_{1}}(t_{1},\dots,t_{i-1},t_{i+1},\dots,t_{\ell-1}).$$

Then given  $v_0 \xrightarrow{f_1} \cdots \xrightarrow{f_{\ell+1}} v_{\ell+1}$  there is a map  $S^{\ell-1} = \partial([0,1]^{\ell}) \to E(\{B_x \mid x \in F(v_0)\}, s_{F(f_{\ell+1} \circ \cdots \circ f_1)})$  defined by

$$(5.24) (t_1, \ldots, t_{i-1}, 0, t_{i+1}, \ldots, t_{\ell}) \mapsto e_{f_{\ell+1}, \ldots, f_{i+1}}(t_{i+1}, \ldots, t_{\ell}) \circ e_{f_i, \ldots, f_1}(t_1, \ldots, t_{i-1}) (t_1, \ldots, t_{i-1}, 1, t_{i+1}, \ldots, t_{\ell}) \mapsto e_{f_{\ell+1}, \ldots, f_{i+1} \circ f_i, \ldots, f_1}(t_1, \ldots, t_{i-1}, t_{i+1}, \ldots, t_{\ell}).$$

The inductive hypothesis implies that this map is continuous. Since  $k \geq \ell + 1$ , by Lemma 5.18, the space  $E(\{B_x \mid x \in F(v_0)\}, s_{F(f_{\ell+1} \circ \cdots \circ f_1)})$  is  $(\ell-1)$ -connected, so the map (5.24) extends to a map  $[0,1]^{\ell} \to E(\{B_x \mid x \in F(v_0)\}, s_{F(f_{\ell+1} \circ \cdots \circ f_1)})$ . Define  $e_{f_{\ell+1}, \dots, f_1}$  to be any such extension.

Now, Equation (5.9) gives a map

$$\Phi(e_{v_0,\dots,v_{\ell+1}}, F(f_{\ell+1} \circ \dots \circ f_1)) \colon [0,1]^{\ell} \times \bigvee_{x \in F(v_0)} S^k \to \bigvee_{x \in F(v_{\ell+1})} S^k.$$

It follows from the compatibility conditions (5.23) that these maps define a homotopy coherent diagram.

Next, for point (2), fix spatial refinements  $\widetilde{F}_k$  and  $\widetilde{F}'_k$  of F. Consider the category  $\underline{2}^1 \times \mathscr{D}$ . It suffices to define a homotopy coherent diagram  $G \colon \underline{2}^1 \times \mathscr{D} \to \mathsf{Top}_{\bullet}$  so that  $G|_{\{0\} \times \mathscr{D}} = \widetilde{F}_k$ ,  $G|_{\{1\} \times \mathscr{D}} = \widetilde{F}'_k$ , and for any  $u \in \mathrm{Ob}(\mathscr{D})$ ,  $G(\varphi_{1,0} \times \mathrm{Id}_u)$  is a homotopy equivalence [Vog73, Proposition 4.6]. To define G, note that  $G|_{\{0\} \times \mathscr{D}}$  and  $G|_{\{1\} \times \mathscr{D}}$  are already specified. Let  $G(\varphi_{1,0} \times \mathrm{Id}_u)$  be the identity map. More generally, define (somewhat arbitrarily)  $G(\varphi_{1,0} \times g) = \widetilde{F}_k(g)$ . It follows from the fact that both  $\widetilde{F}_k$  and  $\widetilde{F}'_k$  refine F that the resulting diagram G is homotopy commutative. Extend G to a homotopy coherent diagram inductively, as in the proof of point (1).

Finally, point (3) is immediate from the definitions.

### 5.3. A coherent cube of box maps.

**Definition 5.25.** Given a strictly unitary, lax 2-functor  $F: \underline{2}^n \to \mathcal{B}$  let  $\widetilde{F}_k: \underline{2}^n \to \mathsf{Top}_{\bullet}$  be a spatial refinement of F. Let  $\widetilde{F}_k^+: \underline{2}_+^n \to \mathsf{Top}_{\bullet}$  be the diagram obtained from  $\widetilde{F}_k$  by defining  $\widetilde{F}_k^+(*)$  to be a single point. Let  $\widetilde{F}^+$  be the diagram obtained from  $\widetilde{F}_k^+$  by replacing each vertex  $\widetilde{F}_k^+(u)$  with  $\Sigma^{-k}(\Sigma^{\infty}\widetilde{F}_k^+(u))$ , the k-fold formal desuspension of its suspension spectrum.

Corollary 5.26. Up to stable homotopy equivalence, the spectrum hocolim  $\widetilde{F}^+$  depends only on the functor F. In fact, for any k > n, the homotopy type of hocolim  $\widetilde{F}_k^+$  is independent of the choices in its construction.

*Proof.* This is immediate from Proposition 5.22, together with the fact that the homotopy colimits of homotopy equivalent homotopy coherent diagrams are homotopy equivalent ([Vog73, Theorem 5.12], quoted as Proposition 4.13).  $\Box$ 

As in Section 4.5, we can also work with a larger enlargement  $\underline{2}_{\dagger}^{n} = (* \leftarrow 1 \to 0)^{\times n}$  of  $\underline{2}^{n}$ . Extend  $\widetilde{F}_{k}$  to a functor  $\widetilde{F}_{k}^{\dagger} : \underline{2}_{\uparrow}^{n} \to \mathsf{Top}_{\bullet}$  by setting  $\widetilde{F}_{k}^{\dagger}|_{\underline{2}^{n}} = \widetilde{F}_{k}$  and  $\widetilde{F}_{k}^{\dagger}(v) = \{\mathsf{pt}\}$  if v is an object which is not in  $\underline{2}^{n}$ , i.e., if some coordinate of v is \*.

**Lemma 5.27.** For any functor  $F: \underline{2}^n \to \mathcal{B}$  and any spatial refinement  $\widetilde{F}_k$  of F there is a stable homotopy equivalence hocolim  $\widetilde{F}_k^+ \simeq \operatorname{hocolim} \widetilde{F}_k^{\dagger}$ .

*Proof.* The proof is similar to but easier than the proof of Lemma 4.41, and is left to the reader.  $\Box$ 

5.4. The realizations of the small cube and big cube agree. Before proving Theorem 7, we introduce an auxiliary category, the arrow category of  $2^n$ , and study its relationship with  $2^n$  and  $\widehat{2^n}$ .

**Definition 5.28.** Given a small category  $\mathscr{C}$ , the arrow category of  $\mathscr{C}$ , which we denote  $\mathsf{Arr}(\mathscr{C})$  has  $\mathsf{Ob}(\mathsf{Arr}(\mathscr{C})) = \bigcup_{u,v \in \mathsf{Ob}(\mathscr{C})} \mathsf{Hom}(u,v)$  the set of morphisms in  $\mathscr{C}$ . Given objects  $f \colon u \to v$  and  $g \colon w \to x$  in the arrow category,  $\mathsf{Hom}(f,g)$  consists of pairs  $(\alpha \colon u \to w, \beta \colon v \to x)$  so that

$$\begin{array}{ccc}
 u & \xrightarrow{f} v \\
 \alpha \downarrow & \downarrow \beta \\
 w & \xrightarrow{q} x
\end{array}$$

commutes. Maps compose in the obvious way:  $(\gamma, \delta) \circ (\alpha, \beta) = (\gamma \circ \alpha, \delta \circ \beta)$ .

There is a functor  $A \colon \mathscr{C} \to \mathsf{Arr}(\mathscr{C})$  defined by

$$A(u) = \mathrm{Id}_{\iota}$$

$$A(f \colon u \to v) = \begin{array}{c} u \xrightarrow{\operatorname{Id}_u} u \\ \downarrow_f & \downarrow_f \\ v \xrightarrow{\operatorname{Id}_v} v. \end{array}$$

There is also a functor  $B \colon \widehat{\mathscr{C}} \to \mathsf{Arr}(\mathscr{C})$  defined by

$$B(u \xrightarrow{f} v \xrightarrow{g} w) = g \circ f$$

$$B\left(\begin{matrix} u \xrightarrow{f} v \xrightarrow{g} w \\ \downarrow^{\alpha} & \uparrow^{\beta} & \downarrow^{\gamma} \\ u' \xrightarrow{f'} v' \xrightarrow{g'} w' \end{matrix}\right) = \begin{matrix} u \xrightarrow{g \circ f} w \\ \downarrow^{\gamma} & \downarrow^{\gamma} \\ u' \xrightarrow{g' \circ f'} w'.$$

In the special case of the cube category,

$$\operatorname{Arr}(\underline{2}^1) = (\varphi_{1,1} \longrightarrow \varphi_{1,0} \longrightarrow \varphi_{0,0}) \qquad \operatorname{Arr}(\underline{2}^n) = \left(\operatorname{Arr}(\underline{2}^1)\right)^n.$$

We will need a version with extra objects added, analogous to  $\underline{2}_{\dagger}^{n}$ :

## Definition 5.30. Let


$$\operatorname{Arr}(\underline{2}^1)_{\dagger} = (* \longleftarrow \varphi_{1,1} \longrightarrow \varphi_{1,0} \longrightarrow \varphi_{0,0})$$

$$\operatorname{Arr}(\underline{2}^n)_{\dagger} = \left(\operatorname{Arr}(\underline{2}^1)_{\dagger}\right)^n.$$

The functors A and B have extensions

$$A_{\dagger} : \underline{2}_{\dagger}^{n} \to \operatorname{Arr}(\underline{2}^{n})_{\dagger}$$
$$B_{\dagger} : \underline{\widehat{2}^{n}}_{\dagger} \to \operatorname{Arr}(\underline{2}^{n})_{\dagger}.$$

These are products of the 1-dimensional case, which is given by:



The dashed arrows denote  $A_{\dagger}$ , and the dotted arrows denote  $B_{\dagger}$ .

To relate various diagrams, we will need to know  $A_{\dagger}$  and  $B_{\dagger}$  are homotopy cofinal:

**Lemma 5.31.** The functor  $A_{\dagger}$  is homotopy cofinal.

*Proof.* Recall from point (ho-4) in Section 4.2 that homotopy cofinality means that each undercategory  $d \downarrow A_{\dagger}$  has contractible nerve. Since taking undercategories commutes with taking products, it suffices to verify the one-dimensional case. This verification is straightforward, and is left to the reader.

**Lemma 5.32.** The functor  $B_{\dagger}$  is homotopy cofinal.

*Proof.* As in the proof of Lemma 5.31, it suffices to verify the 1-dimensional case, which is straightforward.  $\Box$ 

Next we see that for any small category  $\mathscr{C}$ , any functor  $F \colon \mathscr{C} \to \mathscr{B}$  lifts to a functor  $\vec{F} \colon \mathsf{Arr}(\mathscr{C}) \to \mathscr{B}$ .

**Definition 5.33.** Given a functor  $F: \mathscr{C} \to \mathscr{B}$ , define a functor  $\vec{F}: \mathsf{Arr}(\mathscr{C}) \to \mathscr{B}$  as follows. For  $f \in \mathsf{Ob}(\mathsf{Arr}(\mathscr{C}))$ , F(f) is a correspondence, and in particular a set; define  $\vec{F}(f) = F(f)$ . This defines  $\vec{F}$  on  $\mathsf{Ob}(\mathsf{Arr}(\mathscr{C}))$ . Given a diagram as in Formula (5.29), define  $\vec{F}(\alpha,\beta) = F(g \circ \alpha)$  (which is exactly the same as  $F(\beta \circ f)$ , but merely in bijection with  $F(g) \circ F(\alpha)$  and  $F(\beta) \circ F(f)$ ). The source and target maps are given by

$$F(\beta) \times_{F(v)} F(f) \overset{F_{u,v,x}^{-1}}{\longleftarrow} \vec{F}(\alpha,\beta) \xrightarrow{F_{u,w,x}^{-1}} F(g) \times_{F(w)} F(\alpha)$$

$$\downarrow \qquad \qquad \downarrow$$

$$\vec{F}(f) = F(f) \qquad \qquad F(g) = \vec{F}(g).$$

For any pair of composable morphisms

$$\begin{array}{ccc} u & \xrightarrow{f} v \\ \alpha \downarrow & \downarrow \beta \\ w & \xrightarrow{g} x \\ \gamma \downarrow & \downarrow \delta \\ y & \xrightarrow{h} z, \end{array}$$

 $\vec{F}_{f,g,h}$  should specify an isomorphism from  $\vec{F}(\gamma,\delta) \times_{F(g)} \vec{F}(\alpha,\beta) = F(\delta \circ g) \times_{F(g)} F(g \circ \alpha)$  to  $F(\delta \circ g \circ \alpha) = F(\gamma \circ \alpha, \delta \circ \beta)$ . Define this isomorphism to be the composition of the isomorphisms

$$F(\delta \circ g) \times_{F(g)} F(g \circ \alpha) \cong F(\delta) \times_{F(x)} F(g) \times_{F(g)} F(g) \times_{F(w)} F(\alpha)$$
$$\cong F(\delta) \times_{F(x)} F(g) \times_{F(w)} F(\alpha)$$
$$\cong F(\delta \circ g \circ \alpha).$$

**Lemma 5.34.** These maps make  $\vec{F}$  into a strictly unitary, lax 2-functor, and  $\vec{F} \circ A = F$ .

We leave the proof as an exercise to the reader.

**Lemma 5.35.** In  $Arr(\underline{2}^n)_{\dagger}$ , any sequence of composable, non-identity morphisms has length at most 2n.

*Proof.* This is immediate from the definitions.

Corollary 5.36. If  $k \geq 2n$  then any strictly unitary, lax 2-functor  $\vec{F} : Arr(\underline{2}^n) \to \mathcal{B}$  admits a k-dimensional spatial refinement  $\widetilde{F} : Arr(\underline{2}^n) \to Top_{\bullet}$ .

*Proof.* This is immediate from Lemma 5.35 and Proposition 5.22.

**Lemma 5.37.** Fix k > 2n. Given  $F: \underline{2}^n \to \mathcal{B}$ , let  $\widetilde{F}$  be a k-dimensional spatial refinement of F and consider the homotopy coherent diagram  $\widetilde{F} \circ B: \widehat{\underline{2}^n} \to \mathsf{Top}_{\bullet}$ . There is a morphism of homotopy coherent diagrams  $G_k: \widetilde{F} \circ B \to \widehat{F}_k$  so that on each object the underlying map induces an isomorphism on  $H_i$  for i < 2k - 1.

*Proof.* Recall that a morphism from  $\widetilde{F} \circ B$  to  $\widehat{F}_k$  is a diagram over  $\underline{2}^1 \times \widehat{\underline{2}^n}$  whose restriction to  $\{1\} \times \widehat{\underline{2}^n}$  is  $\widetilde{F} \circ B$  and whose restriction to  $\{0\} \times \widehat{\underline{2}^n}$  is  $\widehat{F}_k$ . We will build such a diagram inductively, using box maps from wedges to products.

On  $\{0\} \times \widehat{2^n}$  and  $\{1\} \times \widehat{2^n}$ , G is already specified. Notice that for each object  $(u \xrightarrow{\varphi_{u,v}} v \xrightarrow{\varphi_{v,w}} w) \in Ob(\widehat{2^n})$ , the space  $(\widetilde{F} \circ B)(u \to v \to w) = \bigvee_{a \in F(\varphi_{v,w} \circ \varphi_{u,v})} S^k$  is the k-skeleton of  $\widehat{F}_k(u \to v \to w) = \bigvee_{a \in F(\varphi_{u,v})} \prod_{b \in F(\varphi_{v,w}), \ s(b)=t(a)} S^k$ . For each arrow of the form  $\varphi_{1,0} \times \operatorname{Id}_{u \to v \to w}$ , define  $G(\varphi_{1,0} \times \operatorname{Id}_{u \to v \to w})$  to be the inclusion of the k-skeleton (which is an isomorphism on  $H_i$  for  $i \leq 2k-1$ ). More generally, given a morphism  $(\alpha,\beta,\gamma)$  as in Formula (4.28), define  $G(\varphi_{1,0} \times (\alpha,\beta,\gamma))$  to be the composition  $\widehat{F}_k(\alpha,\beta,\gamma) \circ G(\varphi_{1,0} \times \operatorname{Id}_{u \to v \to w})$ . (Factoring in the other order would work just as well, though it would give a different map.)

The result is a homotopy commutative diagram G, so that the restriction to  $\{0\} \times \widehat{2^n}$  is commutative and the restriction to  $\{1\} \times \widehat{2^n}$  is homotopy coherent. By construction, the maps on the 1-side are disjoint box maps. The maps associated to the edges  $\varphi_{1,0} \times \operatorname{Id}_{u \to v \to w}$  from the 1-side to the 0-side are wedge sums of overlapping box maps: the inclusion

$$\bigvee_{x \in F(\varphi_{v,w} \circ \varphi_{u,v})} S^k = \bigvee_{a \in F(\varphi_{u,v})} \bigvee_{\substack{b \in F(\varphi_{v,w}) \\ s(b) = t(a)}} S^k \hookrightarrow \bigvee_{\substack{a \in F(\varphi_{u,v}) \\ s(b) = t(a)}} \prod_{\substack{b \in F(\varphi_{v,w}) \\ s(b) = t(a)}} S^k$$

restricts to  $\bigvee_{b\in F(\varphi_{v,w}),\ s(b)=t(a)} S^k$  as the overlapping box map associated to the correspondence  $(\{b\in F(\varphi_{v,w})\mid s(b)=t(a)\}, \mathrm{Id}, \mathrm{Id})$  from  $\{b\in F(\varphi_{v,w})\mid s(b)=t(a)\}$  to itself where each sub-box is the whole box; cf. Example 5.14. More succinctly, the map  $S^k_{(a,b)}\to \prod_{b'\in F(\phi_{v,w}),\ s(b')=t(a)} S^k$  is the box map corresponding to the 1-element correspondence from  $\{(a,b)\}$  to  $b\in \{b'\in F(\phi_{v,w}),\ s(b')=t(a)\}$ , where the sub-box  $B_{(a,b)}\subset B_b$  is equal to the whole box  $B_b$ .

To extend the diagram constructed so far to an entire homotopy coherent diagram, we need to define families of maps

$$\bigvee_{x \in F(\varphi_{u,w})} S^k \to \bigvee_{\substack{a \in F(\varphi_{u',v'}) \\ s(b) = t(a)}} \prod_{\substack{b \in F(\varphi_{v',w'}) \\ s(b) = t(a)}} S^k$$

corresponding to sequences of morphism starting at  $(1, u \to v \to w)$  and ending at  $(0, u' \to v' \to w')$ . Observe that for there to be such a sequence we must have  $u \ge u'$  and  $w \ge w'$ . We choose these extensions inductively, maintaining the following restrictions:

(X-1) Fix a morphism  $(u \to v \to w) \to (u' \to v' \to w')$  in  $\widehat{\underline{2}^n}$ . Let  $x \in F(\varphi_{u,w})$ . Decompose x as

$$(x_1, x_2, x_3, x_4) \in F(\varphi_{u,u'}) \times_{F(u')} F(\varphi_{u',v'}) \times_{F(v')} F(\varphi_{v',v}) \times_{F(v)} F(\varphi_{v,w}).$$

Then  $G((u \to v \to w) \to (u' \to v' \to w'))$  sends the sphere associated to x to the wedge summand of

$$\bigvee_{\substack{a \in F(\varphi_{u',v'}) \\ s(b)=t(a)}} \prod_{\substack{b \in F(\varphi_{v',w'}) \\ s(b)=t(a)}} S^k$$

corresponding to  $x_2$ .

(X-2) The map

$$h_x \colon S^k \to \prod_{\substack{b \in F(\varphi_{v',w'})\\ s(b) = t(x_2)}} S^k$$

associated to x in the previous condition is an overlapping box map.

For the inductive step, we need to show that:

- (Y-1) If we post-compose a map satisfying Properties (X-1)-(X-2) with the map in Formula (4.29) we obtain another map satisfying (X-1)-(X-2), and
- (Y-2) If we pre-compose a map satisfying Properties (X-1)-(X-2) with one of the box maps used to define G on the 1-side we obtain another map satisfying (X-1)-(X-2).


Then, Lemma 5.18 implies we can find the desired family of maps to continue the induction.

Statement (Y-2) is clear. For Statement (Y-1), let  $h_x$  be as in Condition (X-2), for  $x \in F(\varphi_{u,w})$ . Given a map  $g \colon (u' \to v' \to w') \to (u'' \to v'' \to w'')$ , the map  $\widehat{F}_k(g)$  sends the wedge summand corresponding to  $x_2 \in F(\varphi_{u',v'})$  to the summand corresponding to  $x_2'$  where  $x_2$  corresponds to  $(x_1', x_2', x_3') \in F(\varphi_{u',u''}) \times_{F(u'')} F(\varphi_{u'',v''}) \times_{F(v'')} F(\varphi_{v'',v'})$ . Since x decomposes as  $(x_1, x_1', x_2', x_3', x_3, x_4)$ , Property (X-1) is satisfied. Further, the map

$$\widehat{F}_k(g) \colon \prod_{\substack{b \in F(\varphi_{v',w'}) \\ s(b) = t(x_2)}} S^k \to \prod_{\substack{b' \in F(\varphi_{v'',w''}) \\ s(b') = t(x_2')}} S^k$$

sends  $S_b^k$  to  $\prod_{b'=(x_3',b,y)} S_{b'}^k = \prod_{y|s(y)=t(b)} S_{(x_3',b,y)}^k$  by the diagonal map. If we replace each box for h labeled by c with  $|\{y \in F(\varphi_{w',w''}) \mid s(y)=t(b)\}|$  boxes, labeled by the corresponding  $b'=(x_3',b,y)$ , then the resulting box map is  $\widehat{F}_k(g) \circ h$ . In particular, we have verified that  $\widehat{F}_k(g) \circ h$  is a box map, as desired.

(The following diagram may be helpful in keeping track of the source and targets of the various elements in the previous paragraph:



The letter labeling each arrow  $\rightarrow$  is the element of  $F(\rightarrow)$  under consideration.) As noted above, Lemma 5.18 and induction now complete the proof.

Corollary 5.38. There is a stable homotopy equivalence  $\operatorname{hocolim}(\widetilde{F} \circ B)^{\dagger} \simeq \operatorname{hocolim} \widehat{F}^{\dagger}$ .

*Proof.* The morphism of diagrams  $G_k$  from Lemma 5.37 extends uniquely to a morphism of thickened diagrams  $G_k^{\dagger} : (\widetilde{F} \circ B)^{\dagger} \to \widehat{F}_k^{\dagger}$ . Further, the diagram  $\widetilde{F}$  and the morphisms  $G_k^{\dagger}$  can be chosen so that  $G_{k+1}^{\dagger}$  is the suspension of  $G_k^{\dagger}$ . It follows that there is an induced map of diagrams of spectra  $G^{\dagger}$ , and the underlying maps of  $G^{\dagger}$  are equivalences.

Proof of Theorem 7. Let  $\vec{F} : \mathsf{Arr}(\underline{2}^n) \to \mathscr{B}$  be the functor from Lemma 5.34. By Corollary 5.36 there is a spatial refinement  $\widetilde{F}$  of  $\vec{F}$ . The composition  $\widetilde{F} = \widetilde{F} \circ A$  is a spatial refinement of F. We will show that the

corresponding diagram  $\widetilde{F}^+: \underline{2}^n_+ \to \mathscr{S}$  satisfies the conditions of the theorem. Indeed, all of the conditions except that the homotopy colimit is |F| are immediate. We compute the homotopy colimit.

By Lemma 5.27,

$$\operatorname{hocolim}_{\underline{2}_{+}^{n}} \widetilde{F}^{+} \simeq \operatorname{hocolim}_{\underline{2}_{+}^{n}} \widetilde{F}^{\dagger}.$$

By Lemma 5.31, Lemma 5.32, and Property (ho-4) of homotopy colimits,

$$\operatorname{hocolim}_{\underline{2_{\dagger}^n}} \widetilde{F}^{\dagger} \simeq \operatorname{hocolim}_{\operatorname{Arr}(\underline{2^n})_{\dagger}} \widetilde{F}^{\dagger} \simeq \operatorname{hocolim}_{\widehat{2^n}_{\bullet}} \widetilde{F}^{\dagger} \circ B^{\dagger}.$$

By Corollary 5.38, there is a homotopy equivalence

$$\operatorname{hocolim}_{\widehat{2^n}_+} \widetilde{F}^{\dagger} \circ B^{\dagger} \simeq \operatorname{hocolim}_{\widehat{2^n}_+} \widehat{F}^{\dagger}.$$

By Lemma 4.41,

$$\operatorname{hocolim}_{\widehat{2^n}_+} \widehat{F}^\dagger \simeq \operatorname{hocolim}_{\widehat{2^n}_+} \widehat{F}^+ = |F|,$$

proving the result.

## 6. A CW complex structure on the realization of the small cube

In this section, we prove that the realization in terms of little cubes (Section 5) is stably homotopy equivalent to the cubical realization (Section 3). We start by studying the cell structure on the little cubes realization:

**Proposition 6.1.** Let  $F: \underline{2}^n \to \mathcal{B}$  be a strictly unitary, lax 2-functor and  $\widetilde{F}_k: \underline{2}^n \to \mathsf{Top}_{\bullet}$  a spatial refinement of F (Definition 5.21). Then the homotopy colimit of  $\widetilde{F}_k^+$  carries a CW complex structure whose cells except the basepoint correspond to the elements of the set  $\coprod_{u \in \{0,1\}^n} F(u)$ . Further, the equivalences  $\Sigma$  hocolim  $\widetilde{F}_k^+$  can be chosen to be cellular, so hocolim  $\widetilde{F}^+$  inherits the structure of a CW spectrum (in the sense of [Ada95, Section III.2]).

*Proof.* Per Observation 4.12, when taking the homotopy colimit we may (and will) consider only chains of non-identity arrows.

For  $u \in \text{Ob}(\underline{2}^n)$  and  $x \in F(u)$ , let  $B_x$  be the box that is associated to x during the construction of  $\widetilde{F}_k^+$ ; that is,  $B_x/\partial B_x$  is the  $S^k$ -summand corresponding to x in  $\widetilde{F}_k^+(u) = \bigvee_{x \in F(u)} S^k$ . Following Definition 4.10 (and with  $\sim$  denoting the same equivalence relation), we can write the homotopy colimit as

$$\operatorname{hocolim} \widetilde{F}_{k}^{+} = \left( \{*\} \coprod \coprod_{u \in \{0,1\}^{n}} \left( \coprod_{m \geq 0} \coprod_{\substack{u = u^{0} \xrightarrow{f_{1}, \dots, f_{m} \\ u^{i} \in \{0,1\}^{n} \cup \{*\}}}} [0,1]^{m} \right) \times \left( \bigvee_{x \in F(u)} B_{x} / \partial B_{x} \right) \right) / \sim$$

$$= \left( \left[ \{*\} \coprod \coprod_{u \in \{0,1\}^{n}} \left( \coprod_{\substack{m \geq 0 \\ u \in \{0,1\}^{n} \cup \{*\} \\ t \neq Id}} \prod_{\substack{t \in \{0,1\}^{n} \\ t \neq Id}} [0,1]^{m} \right) / \sim_{1} \right] \times \left( \coprod_{x \in F(u)} B_{x} \right) \right) / \sim_{2}.$$

where we have broken up the identification  $\sim$  into a two-step identification  $\sim_1$  and  $\sim_2$ , defined as follows:

$$(f_{m}, \dots, f_{1}; t_{1}, \dots, t_{m})$$

$$\sim_{1} \begin{cases} (f_{m}, \dots, f_{i+1} \circ f_{i}, \dots, f_{1}; t_{1}, \dots, t_{i-1}, t_{i+1}, \dots, t_{m}) & t_{i} = 1, \ i < m \\ (f_{m-1}, \dots, f_{1}; t_{1}, \dots, t_{m-1}) & t_{m} = 1 \end{cases}$$

$$(f_{m}, \dots, f_{1}; t_{1}, \dots, t_{m}; y)$$

$$\sim_{2} \begin{cases} (f_{m}, \dots, f_{i+1}; t_{i+1}, \dots, t_{m}; \widetilde{F}_{k}^{+}(f_{i}, \dots, f_{1})(t_{1}, \dots, t_{i-1})(y)) & t_{i} = 0 \\ * & y \in \partial B_{x} \end{cases}$$

Now fix  $u \in \{0,1\}^n$ , and let us study the cubical complex

$$M_{u} := \left( \coprod_{\substack{m \geq 0 \\ u = u^{0}}} \coprod_{\substack{u = u^{0} \xrightarrow{f_{1}, \dots, f_{m} \\ u^{i} \in \{0,1\}^{n} \cup \{*\} \\ f_{i} \neq \operatorname{Id}}}} [0, 1]^{m} \right) / \sim_{1}.$$

If  $u = \vec{0}$ , then  $M_u$  is a single point, which we write as  $\{0\}$  for reasons that will soon be apparent. When  $u \neq \vec{0}$ , we divide the chains  $u = u^0 \xrightarrow{f_1} \cdots \xrightarrow{f_m} u^m$  into two types: the ones ending at  $\vec{0}$  or \*, and the ones ending at neither. In the first case, when  $u^m \in \{\vec{0}, *\}$ , the facet  $[0, 1]^{m-1} \times \{1\}$  is identified with the cube  $[0, 1]^{m-1}$  coming from the sub-chain  $u = u^0 \xrightarrow{f_1} \cdots \xrightarrow{f_{m-1}} u^{m-1}$ . Therefore, we can write

$$M_{u} = \left( \left( \prod_{m \geq 1} \prod_{u = u^{0}} \prod_{\substack{f_{1}, \dots, f_{m} \\ f_{i} \neq \text{Id}}} [0, 1]^{m-1} \times [0, 1] \right) \coprod \left( \prod_{m \geq 1} \prod_{u = u^{0}} \prod_{\substack{f_{1}, \dots, f_{m} \\ u^{i} \in \{0, 1\}^{n} \setminus \{\vec{0}\}}} [0, 1]^{m-1} \right) \right) / \sim_{1}$$

$$\cong \left( \left( \prod_{m \geq 1} \prod_{u = u^{0}} \prod_{\substack{f_{1}, \dots, f_{m} \\ f_{i} \neq \text{Id}}} [0, 1]^{m-1} \times [0, 1] \right) \coprod \left( \prod_{m \geq 1} \prod_{u = u^{0}} \prod_{\substack{f_{1}, \dots, f_{m} \\ f_{i} \neq \text{Id}}} [0, 1]^{m-1} \times [1, 2] \right) \right)$$

$$\coprod \left( \prod_{m \geq 1} \prod_{\substack{u = u^{0} > \dots > u^{m-1} \\ u^{i} \in \{0, 1\}^{n} \setminus \{\vec{0}\}}} [0, 1]^{m-1} \right) / \sim_{1}$$

$$= \left( \prod_{m \geq 1} \prod_{\substack{u = u^{0} > \dots > u^{m-1} \\ u^{i} \in \{0, 1\}^{n} \setminus \{\vec{0}\}}} [0, 1]^{m-1} \right) / \sim_{1} \times [0, 2]$$

where the second identification is via the linear map  $[0,1] \to [1,2]$  that sends 0 to 2 and 1 to 1. This quotient space is just  $M_{u,\vec{0}} \times [0,2]$ , where  $M_{u,\vec{0}} \cong \mathcal{M}_{\mathscr{C}(n)}(u,\vec{0})$  is the cubical complex from Definition 3.19.

Therefore, we can write

$$\operatorname{hocolim} \widetilde{F}_k^+ = \left( \{*\} \coprod \left[ \coprod_{u \in \{0,1\}^n \backslash \{\vec{0}\}} \coprod_{x \in F(u)} \mathcal{M}_{\mathscr{C}_C(n)}(u,\vec{0}) \times [0,2] \times B_x \right] \coprod \left[ \coprod_{x \in F(\vec{0})} \{0\} \times B_x \right] \right) / \sim_2.$$

This gives the required CW complex on the homotopy colimit, where the cell corresponding to x is

$$C(x) = \begin{cases} \mathcal{M}_{\mathcal{C}_C(n)}(u, \vec{0}) \times [0, 2] \times B_x & \text{if } u \neq \vec{0} \\ \{0\} \times B_x & \text{if } u = \vec{0}. \end{cases}$$

The identification  $\sim_2$  glues the boundary  $\partial \mathcal{C}(x)$  to lower-dimensional cells. Specifically, everything over  $\partial B_x$  is identified with the basepoint \*. If  $u \neq \vec{0}$ , everything over  $\{2\} \subset \partial([0,2])$  is identified with \* as well, and everything over  $\{0\} \subset \partial([0,2])$  is identified with cells corresponding to  $u = \vec{0}$ . Finally, points on  $\partial \mathcal{M}_{\mathscr{C}(n)}(u,\vec{0})$  correspond to points on some cube  $[0,1]^l \subset \mathcal{M}_{\mathscr{C}(n)}(u,\vec{0})$  where some coordinate is 0 (Lemma 3.20 (2)), and such points are identified by  $\sim_2$  to points of  $\mathcal{M}_{\mathscr{C}(n)}(u',\vec{0})$  for some  $u > u' > \vec{0}$ .

The fact that the suspension maps are cellular is trivial if we choose compatible spatial refinements. Specifically, take the boxes used to define  $\widetilde{F}_{k+1}^+$  to be [0,1] times the boxes used to define  $\widetilde{F}_k^+$ .

It is not hard to show that if  $\mathscr{C}$  is the cubical flow category corresponding to F (from Construction 4.19) then its associated chain complex (from Definition 3.8) is isomorphic to the reduced cellular cochain complex of the above CW complex, via an isomorphism sending the objects of  $\mathscr{C}$  to the corresponding cells in the CW complex. We will not prove this now, since it follows from Theorem 8.

**Theorem 8.** Let  $(\mathcal{C}, \mathfrak{f}: \mathcal{C} \to \mathcal{C}_{\mathcal{C}}(n))$  be a cubical flow category, and let  $F: \underline{2}^n \to \mathcal{B}$  be the corresponding functor (Construction 4.17). Then the cubical realization of  $\mathcal{C}$  (Definition 3.29) is stably homotopy equivalent to the realization of F as the homotopy colimit of the homotopy coherent diagram  $\widetilde{F}^+: \underline{2}^n_+ \to \mathcal{F}$  from Theorem 7; and the homotopy equivalence sends the cells in the CW complex structure on the homotopy colimit from Proposition 6.1 to the corresponding cells in the cubical realization of  $\mathcal{C}$  via maps of degree  $\pm 1$ .

*Proof.* Fix a cubical neat embedding  $\iota$  of  $\mathscr C$  relative to  $\mathbf d=(d_0,\ldots,d_{n-1})$  and let  $k=\sum_i d_i$ . Let  $\epsilon$  and R be the parameters from Section 3.6, and let

$$\bar{\iota}_{x,y} : \prod_{i=|\mathfrak{f}(y)|}^{|\mathfrak{f}(x)|-1} [-\epsilon, \epsilon]^{d_i} \times \mathcal{M}_{\mathscr{C}}(x, y) \to \left[ \prod_{i=|\mathfrak{f}(y)|}^{|\mathfrak{f}(x)|-1} [-R, R]^{d_i} \right] \times \mathcal{M}_{\mathscr{C}_{C}(n)}(\mathfrak{f}(x), \mathfrak{f}(y))$$

be the extension of  $\iota$  from Formula (3.26). Recall that given  $u \in \text{Ob}(\underline{2}^n)$  and  $x \in F(u)$  the cubical realization  $\|\mathscr{C}\|$  has a corresponding cell

$$C(x) = \begin{cases} \prod_{i=0}^{|u|-1} [-R, R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon, \epsilon]^{d_i} \times [0, 1] \times \mathcal{M}_{\mathscr{C}_C(n)}(u, \vec{0}) & \text{if } u \neq \vec{0}, \\ \prod_{i=0}^{n-1} [-\epsilon, \epsilon]^{d_i} \times \{0\} & \text{if } u = \vec{0}. \end{cases}$$

The strategy of the proof is to use the cubical neat embedding to build a particular spatial refinement  $\tilde{F}_k$  of F and construct a map from hocolim  $\tilde{F}_k^+$ , with the CW structure from Proposition 6.1, to the cubical realization  $\|\mathscr{C}\|$  of the cubical flow category  $\mathscr{C}$  associated to F, that sends cells to cells by degree  $\pm 1$  maps, and hence is a stable homotopy equivalence.

The diagram  $F_k$  is defined as follows. The box associated to  $x \in F(u)$  is

$$B_x = \prod_{i=0}^{|u|-1} [-R, R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon, \epsilon]^{d_i}.$$

Next, consider a sequence of composable non-identity morphisms  $u = u^0 \xrightarrow{f_1} \cdots \xrightarrow{f_m} u^m = v$  in  $\underline{2}^n$ . We will define

$$\widetilde{F}_{k}^{+}(f_{m},\ldots,f_{1}) = \Phi(e_{t_{1},\ldots,t_{m-1}},F(\varphi_{u,v})) : [0,1]^{m-1} \times \widetilde{F}_{k}^{+}(u) \to \widetilde{F}_{k}^{+}(v)$$

for an appropriate family of boxes  $e_{t_1,\dots,t_{m-1}}: [0,1]^{m-1} \to E(\{B_x \mid x \in F(u)\}, s_{F(\varphi_{u,v})})$ . In other words, if for  $\gamma \in F(\varphi_{u,v})$  we write  $B_{\gamma} = \prod_{i=0}^{|v|-1} [-R, R]^{d_i} \times \prod_{i=|v|}^{n-1} [-\epsilon, \epsilon]^{d_i}$  then  $e_{t_1,\dots,t_{m-1}}$  is a  $[0,1]^{m-1}$ -parameter family of embeddings (as disjoint sub-boxes)

$$\coprod_{\gamma|s(\gamma)=x} B_{\gamma} \hookrightarrow B_x, \qquad \forall x \in F(u).$$

To define  $e_{t_1,...,t_{m-1}}$ , fix  $\gamma \in F(\varphi_{u,v})$  with  $s(\gamma) = x$ , and let  $y = t(\gamma)$ . Let  $v_{\gamma}$  be the section of the covering map  $\mathcal{M}_{\mathscr{C}}(x,y) \to \mathcal{M}_{\mathscr{C}_{C}(n)}(u,v)$  whose image is the path component corresponding to  $\gamma$ . Consider the map

$$\mathcal{M}_{\mathscr{C}(n)}(u,v) \times B_{\gamma} = \mathcal{M}_{\mathscr{C}(n)}(u,v) \times \prod_{i=0}^{|v|-1} [-R,R]^{d_i} \times \prod_{i=|v|}^{n-1} [-\epsilon,\epsilon]^{d_i}$$

$$\stackrel{(v_{\gamma},\operatorname{Id})}{\longrightarrow} \mathcal{M}_{\mathscr{C}}(x,y) \times \prod_{i=0}^{|v|-1} [-R,R]^{d_i} \times \prod_{i=|v|}^{n-1} [-\epsilon,\epsilon]^{d_i}$$

$$\cong \prod_{i=0}^{|v|-1} [-R,R]^{d_i} \times \left( \prod_{i=|v|}^{|u|-1} [-\epsilon,\epsilon]^{d_i} \times \mathcal{M}_{\mathscr{C}}(x,y) \right) \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i}$$

$$\stackrel{(\operatorname{Id},\bar{\iota}_{x,y},\operatorname{Id})}{\longrightarrow} \prod_{i=0}^{|v|-1} [-R,R]^{d_i} \times \left( \prod_{i=|v|}^{|u|-1} [-R,R]^{d_i} \times \mathcal{M}_{\mathscr{C}(n)}(u,v) \right) \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i}$$

$$\stackrel{(\operatorname{Id},\pi^R,\operatorname{Id})}{\longrightarrow} \prod_{i=0}^{|v|-1} [-R,R]^{d_i} \times \prod_{i=|v|}^{|u|-1} [-R,R]^{d_i} \times \prod_{i=|u|}^{n-1} [-\epsilon,\epsilon]^{d_i}$$

$$\cong \prod_{i=0}^{|u|-1} [-R,R]^{d_i} \times \prod_{i=|v|}^{n-1} [-\epsilon,\epsilon]^{d_i} = B_x$$

and the induced map

(6.3) 
$$\mathcal{M}_{\mathscr{C}(n)}(u,v) \times \coprod_{\substack{\gamma \in F(\varphi_{u,v})\\s(\gamma)=x}} B_{\gamma} \to B_{x}.$$

It follows from the definition of cubical neat embeddings and the formula for  $\bar{\iota}_{x,y}$  that for any point pt  $\in \mathcal{M}_{\mathscr{C}(n)}(u,v)$ , the restriction  $\{\text{pt}\} \times \coprod_{\gamma \in F(\varphi_{u,v})|s(\gamma)=x} B_{\gamma} \to B_x$  is an inclusion of disjoint sub-boxes. Therefore, we may view the map from Equation (6.3) as a  $\mathcal{M}_{\mathscr{C}(n)}(u,v)$ -parameter family of sub-boxes  $\coprod_{\gamma|s(\gamma)=x} B_{\gamma} \subset B_x$ .

The chain  $u = u^0 > \cdots > u^m = v$  corresponds to some cube  $[0,1]^{m-1}$  in the cubical complex  $M_{u,v}$  from Definition 3.19, which via Lemma 3.20 is identified with some cube  $[0,1]^{m-1} \subset \mathcal{M}_{\mathscr{C}_C(n)}(u,v)$ . Restrict the map from Formula (6.3) to  $[0,1]^{m-1} \times \coprod_{\gamma \mid s(\gamma) = x} B_{\gamma}$  to obtain the required  $[0,1]^{m-1}$ -parameter family of sub-boxes  $\coprod_{\gamma \mid s(\gamma) = x} B_{\gamma} \subset B_x$ .

We check that  $\widetilde{F}_k$  is indeed a homotopy coherent diagram. For any sequence of composable non-identity morphisms  $u = u^0 \xrightarrow{f_1} \cdots \xrightarrow{f_m} u^m = v$  in  $2^n$ , we need to show that

$$\widetilde{F}_{k}(f_{m},\ldots,f_{1})(t_{1},\ldots,t_{i-1},1,t_{i+1},\ldots,t_{m-1}) 
= \widetilde{F}_{k}(f_{m},\ldots,f_{i+1}\circ f_{i},\ldots,f_{1})(t_{1},\ldots,t_{i-1},t_{i+1},\ldots,t_{m-1}) 
\widetilde{F}_{k}(f_{m},\ldots,f_{1})(t_{1},\ldots,t_{i-1},0,t_{i+1},\ldots,t_{m-1}) 
= \widetilde{F}_{k}(f_{m},\ldots,f_{i+1})(t_{i+1},\ldots,t_{m-1})\circ \widetilde{F}_{k}(f_{i},\ldots,f_{1})(t_{1},\ldots,t_{i-1}).$$

The first equation is immediate from Definition 3.19 since the facet of  $[0,1]^{m-1}$  which has  $t_i=1$  is identified with the cube  $[0,1]^{m-2}$  coming from the sequence of composable morphisms  $u=u^0 \xrightarrow{f_1} \cdots \xrightarrow{f_{i-1}} u^{i-1} \xrightarrow{f_{i+1} \circ f_i} u^{i-1} \xrightarrow{f_{i+1} \circ f_i} u^{i+1} \xrightarrow{f_{i+2}} \cdots \xrightarrow{f_m} u^m = v$ . The second equation follows from Lemma 3.20 (3) since the facet of  $[0,1]^{m-1}$  that has  $t_i=0$  lies in the facet  $\mathcal{M}_{\mathscr{C}(n)}(u^i,v) \times \mathcal{M}_{\mathscr{C}(n)}(u,u^i)$  of  $\mathcal{M}_{\mathscr{C}(n)}(u,v)$  and is identified with the product  $[0,1]^{m-i-1} \times [0,1]^{i-1}$ , coming from the sequences  $u^i \xrightarrow{f_{i+1}} \cdots \xrightarrow{f_m} u^m = v$  and  $u^0 \xrightarrow{f_1} \cdots \xrightarrow{f_i} u^i$ , respectively. Since the maps  $\widetilde{F}_k^+$  were defined via cubical neat embeddings that satisfied Definition 3.25 (3), the second equation holds.

Finally, we construct the desired cellular map from hocolim  $\widetilde{F}_k^+$  to  $\|\mathscr{C}\|$ .

For any  $u \in \text{Ob}(\underline{2}^n)$  and any  $x \in F(u)$ , the cell associated to x in hocolim  $\widetilde{F}_k^+$  is

$$C(x)' = \begin{cases} \mathcal{M}_{\mathscr{C}_C(n)}(u, \vec{0}) \times [0, 2] \times B_x & \text{if } u \neq \vec{0} \\ \{0\} \times B_x & \text{if } u = \vec{0}, \end{cases}$$

while the cell associated to x in  $\|\mathscr{C}\|$  is

$$\mathcal{C}(x) = \begin{cases} \mathcal{M}_{\mathscr{C}_C(n)}(u, \vec{0}) \times [0, 1] \times B_x & \text{if } u \neq \vec{0} \\ \{0\} \times B_x & \text{if } u = \vec{0}. \end{cases}$$

Map C(x)' to C(x) by the quotient map  $[0,2] \to [0,2]/[1,2] \cong [0,1]$ , and the identity map on all other factors. This map certainly has degree  $\pm 1$  on each cell. To check that it produces a well-defined map on CW complexes, we must check that it commutes with the attaching maps. Everything over  $\partial B_x$  was quotiented to the basepoint on either side. If  $u = \vec{0}$ , this is the only identification on either side. If  $u \neq \vec{0}$ , everything over  $\{2\} \subset [0,2]$  was quotiented to the basepoint for C(x)', while everything over  $\{1\} \subset [0,1]$  was quotiented to the basepoint for C(x). Therefore, we only need to consider  $u \neq \vec{0}$  and concentrate on the attaching maps on the portion of the boundary of C(x) (respectively, C(x)') that lives over  $\partial \mathcal{M}_{\mathcal{C}(n)}(u,\vec{0})$  or  $\{0\} \subset \partial([0,1])$  (respectively,  $\{0\} \subset \partial([0,2])$ ).

Consider the subcomplex

$$\widetilde{M}_{u} := \left( \prod_{\substack{u=u^{0} > \dots > u^{m} \\ u^{i} \in \{0,1\}^{n}}} [0,1]^{m} \right) / \sim_{1} \cong \mathcal{M}_{\mathscr{C}(n)}(u,\vec{0}) \times [0,1]$$

of the cubical complex  $M_u \cong \mathcal{M}_{\mathscr{C}_C(n)}(u, \vec{0}) \times [0, 2]$  from Equation (6.2) in the proof of Proposition 6.1. We are interested in the part of  $\partial \mathcal{C}(x)'$  that lives over the following subset  $N_u$  of  $\partial \widetilde{M}_u$ :

$$N_{u} := \left(\partial \mathcal{M}_{\mathscr{C}(n)}(u, \vec{0}) \times [0, 1]\right) \cup \left(\mathcal{M}_{\mathscr{C}(n)}(u, \vec{0}) \times \{0\}\right)$$
$$= \left(\prod_{u=u^{0} > \dots > u^{m}} \prod_{i=1}^{m} [0, 1]^{i-1} \times \{0\} \times [0, 1]^{m-i}\right) / \sim_{1}.$$

Let  $p \in \partial \mathcal{C}(x)'$  be some point living over  $N_u$ , and write  $p = (p_1, p_2)$ , where  $p_1 \in N_u$  and  $p_2 \in B_x$ . Assume  $p_1$  lies in the cube  $[0, 1]^m$  corresponding to some chain  $u = u^0 > \cdots > u^m$ . Let  $p_{1,1}, \ldots, p_{1,m}$  be the coordinates of  $p_1$  as a point in the cube, and assume  $p_{1,\ell} = 0$ . Let  $\phi$  denote the restriction of  $\widetilde{F}_k(\varphi_{u^{\ell-1},u_\ell}, \ldots, \varphi_{u^0,u^1})$  to  $[0, 1]^{\ell-1} \times (B_x/\partial B_x)$ . Under the CW complex attaching map (denoted  $\sim_2$  in the proof of Proposition 6.1), p is attached to the point

$$((p_{1,\ell+1},\ldots,p_{1,m}),\phi((p_{1,1},\ldots,p_{1,\ell-1}),p_2)) \in [0,1]^{m-\ell} \times \bigvee_{y \in F(u^{\ell})} (B_y/\partial B_y),$$

where  $[0,1]^{m-\ell}$  is the cube in  $\widetilde{M}_{u^\ell}$  corresponding to the chain  $u^\ell > \cdots > u^m$ . The map  $\phi$  is constructed as a  $[0,1]^{m-\ell}$ -parameter family of box maps. Therefore, the attaching map glues p to the basepoint \* unless  $p_2$  lies in the interior of one of the sub-boxes at the point  $(p_{1,1},\ldots,p_{1,\ell-1})$  in the family; and if  $p_2$  lies in the interior of some box  $B_{\gamma_0}$  then p is glued to the point

$$q := ((p_{1,\ell+1}, \dots, p_{1,m}), q_2) \in [0,1]^{m-\ell} \times B_{y_0} \subset \widetilde{M}_{u^\ell} \times B_{y_0} \subset \mathcal{C}(y_0)'$$

where  $y_0 = t(\gamma_0)$  and  $q_2 := \phi((p_{1,1}, \dots, p_{1,\ell-1}), p_2) \in B_{y_0}$ .

We need to check that p, now viewed as a point in  $\partial \mathcal{C}(x)$ , is also glued to q, now viewed as a point in  $\mathcal{C}(y_0)$ , in the CW complex  $\|\mathscr{C}\|$ . In the construction of  $\|\mathscr{C}\|$  (Definition 3.29), we extended the cubical neat embedding

$$\iota_{x,y_0} \colon \mathcal{M}_{\mathscr{C}}(x,y_0) \hookrightarrow \mathcal{M}_{\mathscr{C}_C(n)}(u^{\ell},\vec{0}) \times \prod_{i=|u^{\ell}|}^{|u|-1} (-R,R)^{d_i}$$

to an embedding

$$\bar{\iota}_{x,y_0} \colon \mathcal{M}_{\mathscr{C}}(x,y_0) \times \prod_{i=|u^\ell|}^{|u|-1} [-\epsilon,\epsilon]^{d_i} \hookrightarrow \mathcal{M}_{\mathscr{C}_C(n)}(u^\ell,\vec{0}) \times \prod_{i=|u^\ell|}^{|u|-1} [-R,R]^{d_i},$$

and used it to define embeddings  $i: \mathcal{M}_{\mathscr{C}}(x, y_0) \times B_{y_0} \hookrightarrow \mathcal{M}_{\mathscr{C}_{C}(n)}(u, u^{\ell}) \times B_x$  and  $j: \mathcal{M}_{\mathscr{C}}(x, y_0) \times \mathcal{C}(y_0) \hookrightarrow \partial \mathcal{C}(x)$ . (Note that  $\widetilde{\mathcal{M}}_{\mathscr{C}_{C}(n)}(u^{\ell}, \vec{0}) = \widetilde{M}_{u^{\ell}}$ .)

For any path component  $\gamma$  of  $\mathcal{M}_{\mathscr{C}}(x,y_0)$ , let  $\iota_{\gamma}$  and  $\jmath_{\gamma}$  denote the restrictions  $\iota|_{\gamma\times B_{y_0}}$  and  $\jmath|_{\gamma\times\mathcal{C}(y_0)}$ ; and as before, let  $\upsilon_{\gamma}$  denote the section of  $\mathcal{M}_{\mathscr{C}}(x,y_0) \to \mathcal{M}_{\mathscr{C}(n)}(u,u^{\ell})$  whose image is  $\gamma$ . Since  $\iota_{x,y_0}$  satisfies Definition 3.25 (1), and its extension  $\bar{\iota}_{x,y_0}$  was defined via Equation (3.26), there exists a map  $\mu_{\gamma} \colon B_{y_0} \to \mathcal{M}_{\mathscr{C}(n)}(u,u^{\ell}) \times B_x$ , so that  $\iota_{\gamma}(\upsilon_{\gamma}(a),b) = (a,\mu_{\gamma}(a,b))$ , for all  $(a,b) \in \mathcal{M}_{\mathscr{C}(n)}(u,u^{\ell}) \times B_{y_0}$ .

 $\mu_{\gamma} \colon B_{y_0} \to \mathcal{M}_{\mathscr{C}(n)}(u, u^{\ell}) \times B_x$ , so that  $\iota_{\gamma}(v_{\gamma}(a), b) = (a, \mu_{\gamma}(a, b))$ , for all  $(a, b) \in \mathcal{M}_{\mathscr{C}(n)}(u, u^{\ell}) \times B_{y_0}$ . Let  $q_1 = (p_{1,\ell+1}, \dots, p_{1,m}) \in [0, 1]^{m-\ell}$  and  $q' = (p_{1,1}, \dots, p_{1,\ell-1}) \in [0, 1]^{\ell-1}$ ; treat q' as a point in  $\mathcal{M}_{\mathscr{C}(n)}(u, u^{\ell})$ , after viewing the cube  $[0, 1]^{\ell-1}$  as the cube corresponding to the chain  $u = u^0 > \dots > u^{\ell}$  in the cubical complex structure on  $\mathcal{M}_{\mathscr{C}(n)}(u, u^{\ell})$  from Lemma 3.20. Let  $\kappa$  be the inclusion map  $\mathcal{M}_{\mathscr{C}(n)}(u, u^{\ell}) \times \widetilde{M}_{u^{\ell}} \times B_x \hookrightarrow \mathcal{C}(x)$ . Since  $\widetilde{F}_k(\varphi_{u^{\ell-1}, u_{\ell}}, \dots, \varphi_{u^0, u^1})$  (used to define  $\phi$ ) is defined via Equation (6.3) using the cubical neat embeddings  $\iota_{x,y_0}$  (which are used to define the maps  $\iota_{\gamma}$ , and consequently,  $\mu_{\gamma}$ ),  $p_2$  is in the interior of the box  $B_{\gamma_0}$ , and  $\phi(q', p_2) = q_2$ , it follows that  $\mu_{\gamma_0}(q', q_2) = p_2$ . Therefore,

$$j_{\gamma_0}(\upsilon_{\gamma}(q'), q) = j_{\gamma_0}(\upsilon_{\gamma}(q'), q_1, q_2) = \kappa(q', q_1, \mu_{\gamma_0}(q', q_2)) 
 = \kappa(q', q_1, p_2) = \kappa((p_{1,1}, \dots, p_{1,\ell-1}), (p_{1,\ell+1}, \dots, p_{1,m}), p_2) 
 = ((p_{1,1}, \dots, p_{1,\ell-1}, 0, p_{1,\ell+1}, \dots, p_{1,m}), p_2) = (p_1, p_2) = p.$$

The last equation is justified by the fact that the cubical complex structures on  $\mathcal{M}_{\mathscr{C}_{C}(n)}(u, u^{\ell})$  and  $\widetilde{M}_{u^{\ell}}$  respect the product structure on facets of  $\widetilde{M}_{u} = \mathcal{M}_{\mathscr{C}_{C}(n)}(u, \vec{0}) \times [0, 1]$  (Lemma 3.20 (3)). Therefore, p is glued to q in the CW complex  $\|\mathscr{C}\|$  as well.

#### 7. The Khovanov homotopy type

We pause briefly to review where we stand. We have introduced a special kind of flow categories, cubical flow categories (Section 3), and shown that the data of a cubical flow category is equivalent to a strictly unitary, lax 2-functor from the cube  $2^n$  to the Burnside 2-category (Section 4). Given a cubical flow category (or functor from the cube to the Burnside category) we have four ways of realizing the functor as a spectrum:

- The original Cohen-Jones-Segal realization (Section 3.2).
- The cubical realization, a modification of the Cohen-Jones-Segal construction taking into account the map to the cube (Section 3.7).
- Thickening the diagram, producing a canonical diagram in spectra, and taking the iterated mapping cone (Section 4).
- Using the "little box" construction to produce a homotopy coherent cube in spectra, and then taking the iterated mapping cone (Section 5.3).

Moreover, Theorems 4, 7, and 8 together imply that, up to stable homotopy equivalence, these realizations all agree.

For the rest of the paper, we turn to a particular cubical flow category: the Khovanov flow category constructed in [LS14a] (see also Section 2.2 and Examples 3.24 and 4.21).

**Definition 7.1.** Given an oriented link diagram K, let  $F_{Kh}(K): \underline{2}^n \to \mathscr{B}$  be the Khovanov functor constructed in Example 4.21. Let  $|F_{Kh}(K)|$  be the result of applying Construction 4.39 to  $F_{Kh}(K)$  and let  $\mathcal{X}_{Kh}(K) = \Sigma^{-n-}|F_{Kh}(K)|$ , where  $n_-$  is the number of negative crossings in K.

Theorems 4, 7, and 8 imply that, up to stable homotopy equivalence,  $\mathcal{X}_{Kh}(K)$  is the same as applying any of the other three realization constructions to  $F_{Kh}(K)$ . A more direct description of  $F_{Kh}(K)$  was given by Hu-Kriz-Kriz [HKK16]; see Section 8.1.

Similar constructions can be carried out for the reduced Khovanov flow category:

**Definition 7.2.** Let  $F_{\widetilde{Kh}}(K): \underline{2}^n \to \mathscr{B}$  be the functor corresponding to the reduced Khovanov flow category from [LS14a, Section 8], via Construction 4.17. Let  $\widetilde{\mathcal{X}}_{Kh}(K)$  be the corresponding spectrum, obtained by applying Construction 4.39 to  $F_{\widetilde{Kh}}(K)$  and desuspending  $n_-$  times.

## 8. Relationship with Hu-Kriz-Kriz

The goal of this section is to prove:

**Theorem 9.** Fix a link diagram K. Let M(K) be the homotopy type associated to K by Hu-Kriz-Kriz [HKK16, Theorem 5.4]. Then  $\mathcal{X}_{Kh}(K)$  is stably homotopy equivalent to M(K).

The Hu-Kriz-Kriz construction has four steps:

- (1) Construct a 2-functor  $\underline{2}^n \to \mathcal{B}$ . (The category  $\underline{2}^n$  is denoted  $I^n$  in [HKK16], and the category  $\mathcal{B}$  is denoted  $\mathcal{S}_2$ .)
- (2) Use the Elmendorf-Mandell machine [EM06] to turn the 2-functor  $\underline{2}^n \to \mathscr{B}$  into an  $A_{\infty}$ -functor  $B_2(2^n)' \to \mathscr{S}$ , where  $B_2(2^n)'$  is an auxiliary category which we review below.
- (3) Use Elmendorf-Mandell's rectification result [EM06, Theorem 1.4] to lift this composition to a strict functor  $2^n \to \mathscr{S}$ .
- (4) Expand  $\underline{2}^n$  to a category  $\mathcal{I}$ , analogous to the expansion of  $\underline{2}^n$  to  $\underline{2}^n_{\dagger}$ , extend the functor  $\underline{2}^n \to \mathscr{S}$  to a functor  $\mathcal{I} \to \mathscr{S}$ , and take the homotopy colimit.

We will prove that the two constructions agree, step-by-step.

8.1. The functors from the cube to the Burnside category agree. Hu-Kriz-Kriz's construction of the Khovanov homotopy type uses the following auxiliary category:

**Definition 8.1.** The (1+1)-dimensional embedded cobordism category  $\mathsf{Cob}^{1+1}_{\mathrm{emb}}$  is the 2-category defined as follows. The objects of  $\mathsf{Cob}^{1+1}_{\mathrm{emb}}$  are oriented 1-manifolds C embedded in  $S^2$  along with a 2-coloring of the components of  $S^2 \setminus C$ , by the colors "white" and "black", so that if  $B(S^2 \setminus C)$  denotes the closure of the black region, then C is the oriented boundary of  $B(S^2 \setminus C)$ . The morphisms from  $C_1$  to  $C_0$  are oriented cobordisms  $\Sigma$  embedded in  $[0,1] \times S^2$  satisfying  $\Sigma \cap (\{i\} \times S^2) = \{i\} \times C_i$  for  $i \in \{0,1\}$ , along with a 2-coloring of  $([0,1] \times S^2) \setminus \Sigma$ , so that if  $B(([0,1] \times S^2) \setminus \Sigma)$  denotes the closure of the black region, then  $\Sigma$  is oriented as the boundary of  $B(([0,1] \times S^2) \setminus \Sigma)$ . The 2-morphisms are isotopy classes of isotopies of cobordisms relative boundary.

The Hu-Kriz-Kriz functor  $\underline{2}^n \to \mathscr{B}$  is constructed in two steps [HKK16, Section 5], which we give as Constructions 8.2 and 8.3.

Construction 8.2. [HKK16, Section 4.3] Given a link diagram L in  $S^2$  with n crossings  $c_1, \ldots, c_n$ , along with a checkerboard coloring of the link diagram, we construct a lax 2-functor from  $\underline{2}^n$  to  $\mathsf{Cob}_{\mathrm{emb}}^{1+1}$ . This functor was partially described in Section 2.2: to  $v \in \{0,1\}^n$ , associate the complete resolution  $\mathcal{P}(v)$  which is a collection of disjoint circles in  $S^2$ . The checkerboard coloring for L induces a 2-coloring of the complement of these circles; orient the circles as the boundary of the black region. To  $u > v \in \{0,1\}^n$ , associate the embedded cobordism  $\Sigma \subset [0,1] \times S^2$ , which is a product cobordism outside a neighborhood of the crossings where u and v differ, and has a saddle for each such crossing. We declare the cobordism to be running from  $\mathcal{P}(u) = \Sigma \cap (\{1\} \times S^2)$  to  $\mathcal{P}(v) = \Sigma \cap (\{0\} \times S^2)$ . Up to isotopy,  $\Sigma$  is independent of the order of the saddles, and in fact the isotopies changing the order of saddles are themselves well-defined up to isotopy. Thus, this construction gives a lax 2-functor  $\underline{2}^n \to \mathsf{Cob}_{\mathrm{emb}}^{1+1}$ .

Construction 8.3. [HKK16, Section 3.4] There is a lax 2-functor  $\mathcal{L} \colon \mathsf{Cob}^{1+1}_{\mathsf{emb}} \to \mathscr{B}$  defined as follows. On objects,  $\mathcal{L}$  sends a 1-manifold  $C \subset S^2$  to the set of all possible labelings of the components of C by elements of  $\{x_+, x_-\} \colon \mathcal{L}(C) = \prod_{C_i \in \pi_0(C)} \{x_+, x_-\}$ . The value of  $\mathcal{L}$  on morphisms is more complicated. For any embedded cobordism  $\Sigma$  and any connected component  $\Sigma_0$  of  $\Sigma$ , consider the 2-coloring of  $([0, 1] \times S^2) \setminus \Sigma_0$  that agrees with the given 2-coloring of  $([0, 1] \times S^2) \setminus \Sigma$  near  $\Sigma_0$ , and let  $\mathcal{B}(([0, 1] \times S^2) \setminus \Sigma_0)$  denote the closure of the black region. Observe that

$$H_1(([0,1] \times S^2) \setminus \Sigma_0) / H_1((\{0,1\} \times S^2) \setminus \partial \Sigma_0) \cong \mathbb{Z}^{2g(\Sigma_0)}$$
  
 $H_1(B(([0,1] \times S^2) \setminus \Sigma_0)) / H_1(B((\{0,1\} \times S^2) \setminus \partial \Sigma_0)) \cong \mathbb{Z}^{g(\Sigma_0)}.$ 

A valid labeling of a cobordism  $\Sigma \subset [0,1] \times S^2$  consists of:

- A labeling of each boundary component of  $\Sigma$  by  $x_+$  or  $x_-$ , and
- A labeling of each genus 1 component  $\Sigma_0$  of  $\Sigma$  by  $\alpha$  or  $-\alpha$ , where  $\{\pm \alpha\}$  are the generators of  $H_1(B(([0,1] \times S^2) \setminus \Sigma_0))/H_1(B((\{0,1\} \times S^2) \setminus \partial \Sigma_0)) \cong \mathbb{Z}$ .

so that:

- Each connected component of  $\Sigma$  has genus 0 or 1.
- For each genus 0 connected component of  $\Sigma$ , the number of boundary components in  $\{0\} \times S^2$  labeled  $x_-$  plus the number of boundary components in  $\{1\} \times S^2$  labeled  $x_+$  is 1.
- For each genus 1 connected component of  $\Sigma$ , all boundary components in  $\{0\} \times S^2$  are labeled  $x_+$  and all boundary components in  $\{1\} \times S^2$  are labeled  $x_-$ .

(See [HKK16, Formula (12)].) Define  $\mathcal{L}(\Sigma)$  to be the set of valid labelings of  $\Sigma$ . The source and target maps of  $\mathcal{L}(\Sigma)$  send a labeling of  $\Sigma$  to the induced labeling of the boundary components.

For  $\Sigma_0$  and  $\Sigma_1$  composable cobordisms in  $[0, \frac{1}{2}] \times S^2$  and  $[\frac{1}{2}, 1] \times S^2$ , respectively, the composition 2-isomorphism  $\mathcal{L}(\Sigma_0) \circ \mathcal{L}(\Sigma_1) \stackrel{\cong}{\longrightarrow} \mathcal{L}(\Sigma_0 \circ \Sigma_1)$  is obvious except for the situation when one gets a cobordism  $\Sigma \subset [0, 1] \times S^2$  with a genus 1 component by stacking two genus 0 cobordisms. The composition map decomposes as a product over connected components of  $\Sigma$ , so for simplicity assume that  $\Sigma$  is connected. Given valid labelings on  $\Sigma_0$  and  $\Sigma_1$  that agree on  $(\partial \Sigma_0) \cap (\partial \Sigma_1) \subset \{\frac{1}{2}\} \times S^2$ , we want to construct a valid labeling on  $\Sigma$ , which amounts to labeling  $\Sigma$  by  $\alpha$  or  $-\alpha$ . It follows from the labeling conditions that there is a unique component C of  $(\partial \Sigma_0) \cap (\partial \Sigma_1)$  that is non-separating in  $\Sigma$  and is labeled  $x_+$ . Orient C as the boundary of the black region, and let  $C_b$  and  $C_w$  be the push-offs of C into the black and the white regions, respectively. One of  $C_b$  and  $C_w$  is a generator of  $H_1(([0,1] \times S^2) \setminus \Sigma)/H_1((\{0,1\} \times S^2) \setminus \partial \Sigma) \cong \mathbb{Z}^2$  and the other one is zero. If  $C_b$  is the generator, label  $\Sigma$  by [C]. If  $C_w$  is the generator, let D be a curve on  $\Sigma$ , oriented so that the algebraic intersection number  $D \cdot C = 1$ , and label  $\Sigma$  by [D]. It is clear that these composition 2-isomorphisms satisfy the coherence condition (Fn-2).

Remark 8.4. It is not hard to see that choosing the other checkerboard coloring on the link diagram yields a naturally isomorphic functor  $2^n \to \mathcal{B}$ . On the other hand, one could have required  $D \cdot C$  to be -1 instead of 1; this would have produced a different functor. This global choice is essentially the choice of ladybug matching from [LS14a, Section 5.4].

**Lemma 8.5.** Hu-Kriz-Kriz's 2-functor  $F_{HKK}: \underline{2}^n \to \mathscr{B}$  [HKK16] is naturally isomorphic to the 2-functor  $F_{Kh}$  constructed in Example 4.21 by applying Construction 4.17 to the Khovanov flow category from [LS14a].

*Proof.* The functors  $F_{HKK}$  and  $F_{Kh}$  are identical on objects. By Lemma 4.5, we only need to show that they agree on the edges and that the composition 2-isomorphisms for the functors agree on the 2-dimensional faces of the cube.

For  $u > v \in \{0,1\}^n$  with |u| - |v| = 1, the corresponding embedded cobordism is a merge or split. Let F denote either  $F_{HKK}$  or  $F_{Kh}$ . It is straightforward from the definitions that for any  $x \in F(u)$  and  $y \in F(v)$ ,  $s^{-1}(x) \cap t^{-1}(y) \subset F(\varphi_{u,v})$  is empty if x does not appear in the Khovanov differential of y, and consists of one point otherwise, so in particular there is a unique bijection  $F_{HKK}(\varphi_{u,v}) \cong F_{Kh}(\varphi_{u,v})$ .

Now consider  $u > w \in \{0,1\}^n$  with |u| - |w| = 2, and let v, v' be the two intermediate vertices. The composition 2-isomorphisms for  $F_{HKK}$  and  $F_{Kh}$  produce bijections between  $F(\varphi_{v,w}) \circ F(\varphi_{u,v'})$  and  $F(\varphi_{v',w}) \circ F(\varphi_{u,v'})$ . We want to show that these two bijections are the same.

Fix  $x \in F(u)$  and  $z \in F(w)$ , and consider  $s^{-1}(x) \cap t^{-1}(z) \subset F(\varphi_{u,w})$ . This set can have 0, 1, or 2 elements. The only nontrivial case to check is when the set has 2 elements, which occurs precisely when x and z are related by a ladybug configuration (Figure 8.1).

Therefore, assume x and z are related by a ladybug configuration, and without loss of generality assume that the corresponding embedded genus 1 cobordism  $\Sigma$  is connected. The composition 2-isomorphisms for F are unchanged under isotopy in  $S^2$ : this is [LS14a, Lemma 5.8] for  $F_{Kh}$ , and is immediate from the definition for  $F_{HKK}$ . Therefore, we may further assume that the ladybug configuration is as shown in Figure 8.1, in the following sense. The circle  $C_0$  is the complete resolution at w, and it is labeled  $x_+$  by z; and the circle  $C_1$ , the complete resolution at u, is obtained by attaching embedded 1-handles along the arcs A and A', and  $C_1$  is labeled  $x_-$  by x. The embedded cobordism  $\Sigma \subset [0,1] \times S^2$  connects  $C_1$  to  $C_0$ , that is,  $\Sigma \cap (\{i\} \times S^2) = \{i\} \times C_i$ . Further, the black region contains the arc A. Consider the labelings of the circles at the complete resolutions at v and v' shown in Figure 8.1; the corresponding generators  $y \in F(v)$  and  $y' \in F(v')$  are matched by the ladybug matching for  $F_{Kh}$  [LS14a, Figure 5.1b].

To show that y and y' are also matched by  $F_{HKK}$ , let C (respectively, C') be the circle that is labeled  $x_+$  by y (respectively, y'); therefore, C (respectively, C') is a homology generator of the white (respectively, black) component of the complement of  $\Sigma$ . Isotope the cobordism  $\Sigma$  inside  $[0,1] \times S^2$  relative boundary so that both the saddles occur at  $\{\frac{1}{2}\} \times S^2$ . Isotope the curves C and C' on  $\Sigma$  so that they intersect transversally




FIGURE 8.1. The ladybug matching. Left: The ladybug configuration. Right: Two intermediate configurations, obtained by attaching embedded 1-handles along the two arcs. Black regions are indicated by dark shading.





FIGURE 8.2. The embedded cobordism  $\Sigma$ . The cobordism connects  $C_0$  at the bottom to  $C_1$  at the top. The portion of the cobordism near the saddle point p, the oriented curves C and C', and the normal vector  $\vec{n}$  are shown.

at one point, C lies in  $[0, \frac{1}{2}] \times S^2$ , and C' lies in  $[\frac{1}{2}, 1] \times S^2$ . Therefore, C and C' intersect at the saddle point p corresponding to the arc A. The portion of the cobordism  $\Sigma$  near p is shown in Figure 8.2. Let  $\vec{n}$  be the normal vector to  $\Sigma$  at p pointing away from the black region, and let  $\vec{t}$  and  $\vec{t}'$  be the tangent vectors to C and C' at p. It is clear from Figure 8.2 that  $[\vec{n}, \vec{t'}, \vec{t}]$  constitute a positive basis; consequently in  $\Sigma$ , oriented as the boundary of the black region, the intersection number  $C' \cdot C = 1$ . Therefore, for both p and p, the surface  $\Sigma$  is labeled by [C'], and thus p and p are matched by p are matched by p and p are matched p and p are match

8.2. Iterated mapping cones. The easiest part of the identification is to see that the Hu-Kriz-Kriz notion of iterated mapping cone agrees with ours. Specifically, to take the iterated mapping cone of their functor  $K(F_{HKK}): \underline{2}^n \to \mathscr{S}$  (see Sections 8.1 and 8.4) they enlarge  $\underline{2}^n$  slightly to a category  $\mathcal{I}$ , and extend  $F_{HKK}$  to a functor  $F_{HKK}: \mathcal{I} \to \mathscr{S}$  by declaring that the new vertices in  $\mathcal{I}$  map to  $\{*\}$ , a one-point space [HKK16, Section 5.2]. We observe that their enlargement is the same as  $\underline{2}_{+}^n$ :

**Lemma 8.6.** There is an isomorphism  $\mathcal{I} \cong 2^n_1$  which commutes with the inclusions of  $2^n$ :



*Proof.* The category  $\mathcal{I}$  has, as objects, pairs  $(J \subseteq \{1,\ldots,n\},\phi\colon J \to \{0,1\})$ , and there is a morphism  $\phi \to \psi$  (which is unique) if and only if  $\phi$  is a restriction of  $\psi$ . The cube  $\underline{2}^n$  sits in  $\mathcal{I}$  as the subcategory  $\{(J,\phi)\mid 0\not\in \operatorname{im}(\phi)\}$ . Recall that  $\underline{2}^n_\dagger=(\underline{2}^1_+)^n$ , and  $\operatorname{Ob}(\underline{2}^1_+)=\{0,1,*\}$ . Given an object  $o=(v_1,\ldots,v_n)\in\underline{2}^n_\dagger$ ,

define  $J = \{i \mid v_i \in \{0, *\}\}$  and  $\phi(i) = \begin{cases} 0 & v_i = * \\ 1 & v_i = 0 \end{cases}$ . With this dictionary, the rest of the verification is straightforward.

8.3. Another kind of homotopy coherent diagram. Hu-Kriz-Kriz use a slightly different notion from Vogt of homotopy coherent diagrams (Section 4.2), which is defined in two steps:

**Definition 8.7.** Given a small category  $\mathscr{C}$ , let  $\mathscr{C}'$  be the 2-category with the same objects as  $\mathscr{C}$ ,

$$\operatorname{Hom}_{\mathscr{C}'}(x,y) = \coprod_{x=x_0,x_1,\dots,x_n=y} \operatorname{Hom}_{\mathscr{C}}(x_{n-1},x_n) \times \operatorname{Hom}_{\mathscr{C}}(x_{n-2},x_{n-1}) \times \dots \times \operatorname{Hom}_{\mathscr{C}}(x_0,x_1)$$

the set of finite sequences of composable morphisms starting at x and ending at y, and a unique 2-morphism from  $(f_n, \ldots, f_1)$  to  $(g_m, \ldots, g_1)$  whenever  $f_n \circ \cdots \circ f_1 = g_m \circ \cdots \circ g_1$  (compare [HKK16, Section 4.1]). Composition is given by concatenation of sequences. There is a projection  $\mathscr{C}' \to \mathscr{C}$ , where we view  $\mathscr{C}$  as a 2-category with only identity 2-morphisms, which sends  $(f_n, \ldots, f_1)$  to  $f_n \circ \cdots \circ f_1$ .

**Definition 8.8.** Given a 2-category  $\mathscr{D}$ , we can form a simplicially enriched category  $B_2(\mathscr{D})$  by replacing each Hom category in  $\mathscr{D}$  by the realization of its nerve.

Combining Definitions 8.7 and 8.8, the analogue of a homotopy coherent diagram in [HKK16] is a (simplicially enriched) functor  $B_2(\mathscr{C}') \to \mathscr{S}$ , or more generally an  $A_{\infty}$  functor  $B_2(\mathscr{C}') \to \mathscr{S}$ . Note that there is a projection  $\Pi \colon B_2(\mathscr{C}') \to \mathscr{C}$  induced by the projection  $\mathscr{C}' \to \mathscr{C}$  (and the triviality that  $B_2(\mathscr{C}) = \mathscr{C}$ ). So, given a functor  $F \colon \mathscr{C} \to \mathscr{S}$  (say) there is an induced (simplicially enriched) functor  $B_2(F') = F \circ \Pi \colon B_2(\mathscr{C}') \to \mathscr{S}$ .

**Lemma 8.9.** For any small category  $\mathscr{C}$ , the projection map  $\Pi: B_2(\mathscr{C}') \to \mathscr{C}$  is a homotopy equivalence on each Hom space.

*Proof.* The category  $\operatorname{Hom}_{\mathscr{C}'}(x,y)$  decomposes as a disjoint union of subcategories

$$\operatorname{Hom}_{\mathscr{C}'}(x,y) = \coprod_{f \in \operatorname{Hom}(x,y)} \{ (f_n, \dots, f_1) \mid f_n \circ \dots \circ f_1 = f \},$$

and for each of these subcategories, every object is initial.

The notion of homotopy colimits extends easily to functors from simplicially enriched categories: in Formula (4.11), say, one replaces the disjoint union over sequences of composable morphisms with the disjoint union of products  $\coprod_{x_0,...,x_n} \operatorname{Hom}(x_{n-1},x_n) \times \cdots \times \operatorname{Hom}(x_0,x_1) \times [0,1]^n \times F(x_0)$ , and quotient by the same equivalence relation from Definition 4.10. The properties of homotopy colimits stated in Section 4.2 extend without change to diagrams from simplicially enriched categories.

**Lemma 8.10.** Let  $F: \mathscr{C} \to \mathscr{S}$  be a diagram from a small category  $\mathscr{C}$ . Then there is a homotopy equivalence  $\operatorname{hocolim}_{\mathscr{C}} F \simeq \operatorname{hocolim}_{B_2(\mathscr{C}')} B_2(F')$ .

*Proof.* By definition  $B_2(F') = F \circ \Pi$ . By Lemma 8.9, the projection  $\Pi$  is a quasi-equivalence, and in particular homotopy cofinal, so the result follows from Property (ho-4) of homotopy colimits.

8.4. **The Elmendorff-Mandell machine.** A permutative category is a strictly unital, strictly associative, symmetric monoidal category (see, e.g., [EM06, Definition 3.1]). Let Permu denote the (strict) 2-category of permutative categories [EM06, Definition 3.2]. The Elmendorff-Mandell machine [EM06] is a functor

$$K \colon B_2(\mathsf{Permu}) \to \mathscr{S}$$

from  $(B_2 \text{ of})$  the category of permutative categories to spectra. (Constructions of this type go back to Segal [Seg74].) Rather than explain how the machine works, we list the properties we will need:

(EM-1) Given a permutative category  $\mathscr C$  and an object x in  $\mathscr C$  there is an induced map  $K(x) \colon \mathbb S \to K(\mathscr C)$ . Further, this is natural in the sense that given a functor of permutative categories  $F \colon \mathscr C \to \mathscr D$  the following diagram commutes:

$$K(x) \downarrow K(F(x)) \downarrow K(\mathscr{C}) \xrightarrow{K(F)} K(\mathscr{D}).$$

- (EM-2) The Cartesian product is both the categorical product and coproduct in the category of permutative categories.
- (EM-3) Given permutative categories  $\mathscr{C}$  and  $\mathscr{D}$ ,  $K(\mathscr{C} \times \mathscr{D}) = K(\mathscr{C}) \times K(\mathscr{D})$ .
- (EM-4) The category Sets of finite sets, with disjoint union, is equivalent to a permutative category [Isb69]; to keep the exposition clear we will continue to use the name Sets for this category. The map  $\mathbb{S} \to K(\mathsf{Sets})$  induced by a 1-element set and property (EM-1) is a stable homotopy equivalence. (This is a version of the Barratt-Priddy-Quillen theorem.)
- (EM-5) Given a 2-category  $\mathscr C$  in which all 2-morphisms are isomorphisms and a strict 2-functor  $F:\mathscr C\to \mathsf{Permu}$ , there is an induced (simplicially enriched) functor  $K(B_2(F)):B_2(\mathscr C)\to\mathscr S$ .

Construction 8.11. Given a set X, we can consider the category  $\prod_{x \in X} \mathsf{Sets}$ . Given a correspondence (C, s, t) from X to Y, there is an induced functor  $\prod_{x \in X} \mathsf{Sets} \to \prod_{y \in Y} \mathsf{Sets}$  which sends

$$(A_x)_{x \in X} \mapsto \left(\bigcup_{x \in X} (s^{-1}(x) \cap t^{-1}(y)) \times A_x\right)_{y \in Y}.$$

(Note that the union operation in the above formula is actually a disjoint union.) An isomorphism between correspondences (C, s, t) and (C', s', t') can be viewed as simply a relabeling of the elements of C; and this relabeling induces a natural isomorphism between the two functors.

**Lemma 8.12.** Construction 8.11 defines a lax 2-functor  $\mathscr{B} \to \mathsf{Permu}$ .

Sketch of proof. Verify that the category  $\prod_{x \in X} \mathsf{Sets}$  is naturally isomorphic to the category  $\mathsf{Sets}/X$ , and that under this identification the functor induced by C is naturally isomorphic to the functor  $A \mapsto C \times_X A$ .  $\square$ 

$$\operatorname{hocolim} K(B_2(F'))^{\dagger} \longleftarrow \underbrace{\hspace{1cm} (1) \\ \operatorname{hocolim} K(B_2((\vec{F} \circ B)'))^{\dagger} \longleftarrow \underbrace{\hspace{1cm} (5) \\ \operatorname{hocolim} K(B_2(G'_p))^{\dagger}}_{}$$

$$\downarrow (4)$$

$$\operatorname{hocolim} \hat{F}^{\dagger} \longleftarrow \underbrace{\hspace{1cm} (2) \\ \operatorname{hocolim} G^{\dagger} \longleftarrow \underbrace{\hspace{1cm} (3) \\ \operatorname{hocolim} K(G_p)^{\dagger}}_{}$$

FIGURE 8.3. The steps in the proof of Proposition 8.15. Each arrow is labeled by the step showing it is an equivalence.

Remark 8.13. Using properties (EM-3) and (EM-4),  $K(\prod_{x \in X} \mathsf{Sets}) \simeq \prod_{x \in X} K(\mathsf{Sets}) \simeq \prod_{x \in X} \mathbb{S}$ . With respect to this decomposition, however, the map  $\prod_{x \in X} \mathbb{S} \to \prod_{y \in Y} \mathbb{S}$  induced by a correspondence  $C \colon X \to Y$  is not geometrically obvious.

**Construction 8.14.** Given a category  $\mathscr{C}$ , viewed as a 2-category with only identity 2-morphisms, and a strictly unitary, lax 2-functor  $F:\mathscr{C}\to\mathscr{B}$  there is an induced strict 2-functor  $F':\mathscr{C}'\to\mathscr{B}$ , where  $\mathscr{C}'$  is as in Definition 8.7. Composing with the  $\prod_{x\in(-)}\mathsf{Sets}$  construction gives a strict 2-functor  $\mathscr{C}'\to\mathsf{Permu}$ , which we will still denote F', with  $F'(u)=\prod_{x\in F(u)}\mathsf{Sets}$ . Finally, by (EM-5), the K-theory functor gives a functor  $K(B_2(F'))\colon B_2(\mathscr{C}')\to\mathscr{S}$ .

Hu-Kriz-Kriz apply this construction to the Khovanov functor  $F_{HKK}: \underline{2}^n \to \mathcal{B}$ , and then take the iterated mapping cone as in Section 8.2 to obtain their Khovanov stable homotopy type.

**Proposition 8.15.** Given a strictly unitary, lax 2-functor  $F: \underline{2}^n \to \mathcal{B}$ , there is a stable homotopy equivalence between the Hu-Kriz-Kriz realization hocolim  $K(B_2(F'))^{\dagger}$  and the realization hocolim  $\widehat{F}^{\dagger}$  from Section 4.5.

*Proof.* To identify the Hu-Kriz-Kriz construction and the thickening construction from Section 4 we use a sequence of intermediate diagrams, somewhat in the spirit of Section 5.4. (See Figure 8.3 for an overview.)

(1) By Lemma 5.34, we can lift the functor  $F: \underline{2}^n \to \mathcal{B}$  to a functor  $\vec{F}: \mathsf{Arr}(\underline{2}^n) \to \mathcal{B}$ , so that  $F = \vec{F} \circ A$ . Composing with the composition map  $B: \underline{\widehat{2}^n} \to \mathsf{Arr}(\underline{2}^n)$  gives a functor  $\vec{F} \circ B: \underline{\widehat{2}^n} \to \mathcal{B}$ .

There is a homotopy equivalence hocolim  $K(B_2(F'))^{\dagger} \simeq \operatorname{hocolim} K(B_2((\vec{F} \circ B)'))^{\dagger}$ ; the argument is similar to the one in Section 5.4. In the commutative diagram

$$B_{2}(\underline{2}^{n'})_{\dagger} \longrightarrow B_{2}(\mathsf{Arr}(\underline{2}^{n})')_{\dagger} \longleftarrow B_{2}((\widehat{\underline{2}^{n}})')_{\dagger}$$

$$\cong \bigcup_{\cong} \bigcup_{\cong$$

the vertical arrows are quasi-equivalences by Lemma 8.9, and  $A_{\dagger}$  and  $B_{\dagger}$  are homotopy cofinal by Lemma 5.31 and Lemma 5.32, so the functors in the top row are also homotopy cofinal. Hence, by Property (ho-4) of homotopy colimits hocolim  $K(B_2(F'))^{\dagger} \simeq \operatorname{hocolim} K(B_2(\vec{F}'))^{\dagger} \simeq \operatorname{hocolim} K(B_2(\vec{F}'))^{\dagger}$ .

(2) Recall that  $\widehat{F}$  sends an object  $(u \xrightarrow{f} v \xrightarrow{g} w)$  to  $\bigvee_{a \in F(f)} \prod_{b \in F(g), \ s(b) = t(a)} \mathbb{S}$ . Further, the definition of  $\widehat{F}$  uses only the universal properties of product and coproduct, so for any spectrum X we could define a functor  $\widehat{F}_X : \widehat{\underline{2}^n} \to \mathscr{S}$  with  $\widehat{F}_X(u \xrightarrow{f} v \xrightarrow{g} w) = \bigvee_{a \in F(f)} \prod_{b \in F(g), \ s(b) = t(a)} X$  (cf. Remark 4.38). Moreover, this thickening procedure is clearly natural in X. Taking the special case  $X = K(\mathsf{Sets})$  and

using Property (EM-4) gives a diagram  $G: \widehat{\underline{2^n}} \to \mathscr{S}$  with

$$G(u \xrightarrow{f} v \xrightarrow{g} w) = \bigvee_{\substack{a \in F(f) \\ s(b) = t(a)}} K(\mathsf{Sets}),$$

and a natural transformation of diagrams  $\widehat{F} \to G$  which is an equivalence on objects. In particular, hocolim  $\widehat{F}^{\dagger} \simeq \text{hocolim } G^{\dagger}$ .

(3) There is also a functor  $G_p: \widehat{\underline{2}^n} \to \mathsf{Permu}$  defined similarly to  $\widehat{F}$ , using the product and coproduct on  $\mathsf{Permu}$  in place of the product and wedge sum of spaces, and using  $\mathsf{Sets}$  in place of  $\mathbb{S}$ . That is,

$$G_p(u \xrightarrow{f} v \xrightarrow{g} w) = \coprod_{\substack{a \in F(f) \ s(b) = t(a)}} \prod_{\substack{b \in F(g) \ s(b) = t(a)}} \mathsf{Sets}.$$

Recall from (EM-2) that finite products and coproducts of permutative categories are given by the Cartesian product. Thus, using Property (EM-3),

$$K(G_p)(u \xrightarrow{f} v \xrightarrow{g} w) \simeq \prod_{\substack{a \in F(f) \\ s(b) = t(a)}} K(\mathsf{Sets}).$$

(Note that since  $\widehat{\underline{2}^n}$  has only identity 2-morphisms,  $B_2(\widehat{\underline{2}^n})$  can be identified with  $\widehat{\underline{2}^n}$ , and therefore, we have written  $K(G_p)$  instead of  $K(B_2(G_p))$ .)

Using the stable homotopy equivalence between wedge sum and product we get a natural transformation of diagrams from G to  $K(G_p)$  which is an equivalence on objects. In particular, hocolim  $G^{\dagger} \simeq \operatorname{hocolim} K(G_p)^{\dagger}$ .

- (4) By Lemma 8.10, there is a homotopy equivalence hocolim  $K(G_p)^{\dagger} \simeq \operatorname{hocolim} B_2(K(G_p)')^{\dagger}$ . However, since  $G_p$  is a strict functor from a 1-category to Permu,  $B_2(K(G_p)') = K(G_p) \circ \Pi = K(B_2(G_p'))$ , where  $\Pi \colon B_2((\widehat{2^n})') \to \widehat{2^n}$  is the projection; therefore,  $\operatorname{hocolim} K(G_p)^{\dagger} \simeq \operatorname{hocolim} K(B_2(G_p'))^{\dagger}$ .
- (5) To finish the identification, there are isomorphisms H from  $G'_p$  to  $(\vec{F} \circ B)'$ . On the objects, the natural transformation will induce the equivalence from

$$G_p'(u \xrightarrow{f} v \xrightarrow{g} w) = G_p(u \xrightarrow{f} v \xrightarrow{g} w) = \coprod_{\substack{a \in F(f) \\ s(b) = t(a)}} \prod_{\substack{b \in F(g) \\ s(b) = t(a)}} \mathsf{Sets}$$

to

$$(\vec{F} \circ B)'(u \xrightarrow{f} v \xrightarrow{g} w) = (\vec{F} \circ B)(u \xrightarrow{f} v \xrightarrow{g} w) = \prod_{c \in F(g \circ f)} \mathsf{Sets}$$

which comes from the natural bijection  $F(g \circ f) \to F(g) \times_{F(v)} F(f)$  between their indexing sets, and the identification of coproducts and products in Permu with Cartesian products. Because Cartesian product of sets is not strictly associative, and because of the 2-categorical nature of the identification between coproducts and products in Permu, this does not strictly define a strict natural isomorphism  $G'_p \to (\vec{F} \circ B)'$ .

Instead, it defines a pseudonatural equivalence: for a map  $\varphi: (u \xrightarrow{f} v \xrightarrow{g} w) \to (u' \xrightarrow{f'} v' \xrightarrow{g'} w')$  in  $\widehat{2^n}$ , there is a natural isomorphism of functors

$$(\vec{F} \circ B)'(\varphi) \circ H(u \xrightarrow{f} v \xrightarrow{g} w) \cong H(u' \xrightarrow{f'} v' \xrightarrow{g'} w') \circ G'_p(\varphi).$$

which respects composition in  $\widehat{\underline{2}^n}$ .

We define  $\mathscr E$  to be the category with objects 1 and 0 and a unique morphism between any pair of objects. Let  $\mathscr D=\mathscr E\times\widehat{\underline{2}^n}_\dagger$ , and let  $\mathscr D_0$  and  $\mathscr D_1$  be the full subcategories of  $\mathscr D$  spanned by the objects of  $\{0\}\times\widehat{\underline{2}^n}_\dagger$  and  $\{1\}\times\widehat{\underline{2}^n}_\dagger$ , respectively. Note that the projection  $\mathscr D\to\widehat{\underline{2}^n}_\dagger$  is an equivalence, and restricts to isomorphisms from both  $\mathscr D_0$  and  $\mathscr D_1$  to  $\widehat{\underline{2}^n}_\dagger$ .

The pseudonatural equivalence above defines a lax 2-functor, also denoted H, from  $\mathscr{D}$  to Permu whose restriction to  $\{1\} \times \widehat{\underline{2}^n}$  is  $G'_p$  and whose restriction to  $\{0\} \times \widehat{\underline{2}^n}$  is  $(\vec{F} \circ B)'$ . Therefore,  $K(B_2(H))$  restricts to  $K(B_2(G'_p))^{\dagger}$  and  $K(B_2(\vec{F} \circ B)')^{\dagger}$ , and since both inclusions  $\mathscr{D}_0 \to \mathscr{D}$  and  $\mathscr{D}_1 \to \mathscr{D}$  are homotopy cofinal.

$$\operatorname{hocolim} K(B_2((\vec{F} \circ B)'))^{\dagger} \simeq \operatorname{hocolim} K(B_2(H)) \simeq \operatorname{hocolim} K(B_2(G'_p))^{\dagger}.$$

Putting these all together (cf. Figure 8.3), in conjunction with Lemma 8.6, the Hu-Kriz-Kriz realization hocolim  $K(B_2(F'))^{\dagger}$  agrees with our thickening construction hocolim  $\widehat{F}^{\dagger}$ .

# 8.5. Proof that the Khovanov homotopy types agree.

Proof of Theorem 9. Lemma 8.5 identifies the 2-functors  $\underline{2}^n \to \mathcal{B}$  used in this paper and [HKK16]. Proposition 8.15 identifies Hu-Kriz-Kriz's realization of this functor with ours.

### 9. Khovanov homotopy type of a disjoint union and connected sum

In this section we prove Theorems 1, 2, and 10. For the first two theorems, we merely need to show that the functor associated to a disjoint union (respectively connect sum) of links is the product of the functors of the individual links:

**Proposition 9.1.** Let  $L_1$  and  $L_2$  be link diagrams, and let  $L_1 \coprod L_2$  be their disjoint union. Order the crossings in  $L_1 \coprod L_2$  so that all of the crossings in  $L_1$  come before all of the crossings in  $L_2$ . Then

$$F_{Kh}^{j}(L_1 \coprod L_2) \cong \coprod_{j_1+j_2=j} F_{Kh}^{j_1}(L_1) \times F_{Kh}^{j_2}(L_2),$$

where  $\times$  denotes the product of functors (Definition 4.43),  $\coprod$  denotes the disjoint union of functors (Definition 4.47), and  $\cong$  denotes natural isomorphism of 2-functors. If we fix a basepoint on  $L_1$  then

$$F^j_{\widetilde{Kh}}(L_1 \coprod L_2) \cong \coprod_{j_1+j_2=j} F^{j_1}_{\widetilde{Kh}}(L_1) \times F^{j_2}_{Kh}(L_2).$$

If we fix basepoints on  $L_1$  and  $L_2$  and let  $L_1 \# L_2$  denote the connected sum (at the basepoints) then

$$F_{\widetilde{Kh}}^{j}(L_1 \# L_2) \cong \coprod_{j_1 + j_2 = j} F_{\widetilde{Kh}}^{j_1}(L_1) \times F_{\widetilde{Kh}}^{j_2}(L_2).$$

*Proof.* We will prove the first statement; the proofs of the other two are similar. Let  $n_i$  be the number of crossings in  $L_i$ . To keep notation simple, write  $F = F_{Kh}^j(L_1 \coprod L_2)$ ,  $X_v = F(v)$ ,  $A_{v,w} = F(\varphi_{v,w})$ ,  $G = \coprod_{j_1+j_2=j} F_{Kh}^{j_1}(L_1) \times F_{Kh}^{j_2}(L_2)$ ,  $Y_v = G(v)$  and  $B_{v,w} = G(\varphi_{v,w})$ .

By Lemma 4.5, it suffices to construct bijections  $\phi_v \colon X_v \xrightarrow{\cong} Y_v$  and  $\psi_{v,w} \colon A_{v,w} \xrightarrow{\cong} B_{v,w}$  for all v > w with |v| - |w| = 1 so that  $\psi_{v,w}$  respects the source and target maps and for any u > w with |u| - |w| = 2,

the following diagram commutes:

$$(9.2) A_{v,w} \times_{X_{v}} A_{u,v} \xrightarrow{\psi_{v,w} \times \psi_{u,v}} B_{v,w} \times_{Y_{v}} B_{u,v}$$

$$\downarrow^{F_{u,v,w}} \qquad \qquad \downarrow^{G_{u,v,w}}$$

$$A_{u,w} \qquad B_{u,w}$$

$$\uparrow^{F_{u,v',w}} \qquad \qquad \uparrow^{G_{u,v',w}}$$

$$A_{v',w} \times_{X_{v'}} A_{u,v'} \xrightarrow{\psi_{v',w} \times \psi_{u,v'}} B_{v',w} \times_{Y_{v'}} B_{u,v'}.$$

Here, v and v' are the two vertices so that u > v, v' > w. Note that all arrows in this diagram are isomorphisms.

The map  $\phi_v$  is the canonical identification between Khovanov generators for  $L_1 \coprod L_2$  and pairs of a Khovanov generator for  $L_1$  and a Khovanov generator for  $L_2$ . There is a unique map  $\psi_{v,w} \colon A_{v,w} \to B_{v,w}$  for v > w with |v| - |w| = 1 which commutes with the source and target maps, because:

- (1) Given  $x_v \in X_v$  and  $x_w \in X_w$ ,  $s^{-1}(x_v) \cap t^{-1}(x_w) \subset A_{v,w}$  is either empty, if  $x_v$  does not occur in the Khovanov differential of  $x_w$ , or consists of a single point, if  $x_v$  does occur in the Khovanov differential of  $x_w$ . Similar statements hold for  $Y_v$  and  $B_{v,w}$ . It follows that if  $\psi_{v,w}$  exists then it is unique.
- (2) The canonical identification of Khovanov generators does, in fact, give a chain map. So, by the observations in the previous point, the map  $\psi_{v,w}$  does exist.

Except in one case, the same argument shows that the diagram (9.2) commutes: typically, for each  $x_u \in X_u$  and  $x_w \in X_w$  (with u > w and |u| - |w| = 2),  $s^{-1}(x_u) \cap t^{-1}(x_w) \subset A_{u,w}$  is either empty or has a single element. The exceptional case is the case of a ladybug configuration, as in [LS14a, Section 5.4] (see also Figure 8.1). In the ladybug case, either both crossings under consideration lie in  $L_1$  or both crossings lie in  $L_2$ , from which it follows easily that the diagram commutes. (This is immediate for the present case when we are considering the disjoint union  $L_1 \coprod L_2$ ; the connect-sum case  $L_1 \# L_2$  is also fairly obvious.)

We are now ready to prove Theorems 1 and 2, which we recall for the reader's convenience:

**Theorem 1.** Let  $L_1$  and  $L_2$  be links, and  $L_1 \coprod L_2$  their disjoint union. Then

(1.1) 
$$\mathcal{X}_{Kh}^{j}(L_1 \coprod L_2) \simeq \bigvee_{j_1+j_2=j} \mathcal{X}_{Kh}^{j_1}(L_1) \wedge \mathcal{X}_{Kh}^{j_2}(L_2).$$

Moreover, if we fix a basepoint p in  $L_1$ , not at a crossing, and consider the corresponding basepoint for  $L_1 \coprod L_2$ , then

(1.2) 
$$\widetilde{\mathcal{X}}_{Kh}^{j}(L_1 \coprod L_2) \simeq \bigvee_{j_1+j_2=j} \widetilde{\mathcal{X}}_{Kh}^{j_1}(L_1) \wedge \mathcal{X}_{Kh}^{j_2}(L_2).$$

**Theorem 2.** Let  $L_1$  and  $L_2$  be based links and  $L_1 \# L_2$  the connected sum of  $L_1$  and  $L_2$ , where we take the connected sum near the basepoints. Then

(1.3) 
$$\widetilde{\mathcal{X}}_{Kh}^{j}(L_1 \# L_2) \simeq \bigvee_{j_1 + j_2 = j} \widetilde{\mathcal{X}}_{Kh}^{j_1}(L_1) \wedge \widetilde{\mathcal{X}}_{Kh}^{j_2}(L_2).$$

Proof of Theorems 1 and 2. We will prove Formula (1.1); the proofs of Formulas (1.2) and (1.3) are essentially the same.

Fix a diagram for  $L_1 \coprod L_2$  so that there are no crossings between  $L_1$  and  $L_2$ . Order the crossings in  $L_1 \coprod L_2$  so that all of the crossings in  $L_1$  come before all of the crossings in  $L_2$ . By Proposition 9.1,

$$F_{Kh}^{j}(L_1 \coprod L_2) \cong \coprod_{j_1+j_2=j} F_{Kh}^{j_1}(L_1) \times F_{Kh}^{j_2}(L_2).$$

By Lemma 4.42, naturally isomorphic functors have stably homotopy equivalent realizations. By Propositions 4.46 and 4.50, the realization of  $\coprod_{j_1+j_2=j} F^{j_1}_{Kh}(L_1) \times F^{j_2}_{Kh}(L_2)$  is  $\bigvee_{j_1+j_2=j} \mathcal{X}^{j_1}_{Kh}(K_1) \wedge \mathcal{X}^{j_2}_{Kh}(L_2)$ .

These results quickly imply Corollary 1.4, which we also recall:

Corollary 1.4. For any n there exists a link  $L_n$  so that the operation

$$\operatorname{Sq}^n \colon Kh^{i,j}(L_n) \to Kh^{i+n,j}(L_n)$$

is non-zero, for some  $i, j \in \mathbb{Z}$ . Similarly, there exists a knot  $K_n$  so that the operation

$$\operatorname{Sq}^n \colon \widetilde{Kh}^{i,j}(K_n) \to \widetilde{Kh}^{i+n,j}(K_n)$$

is non-zero, for some  $i, j \in \mathbb{Z}$ . Further, for this knot, the operation

$$\operatorname{Sq}^n \colon \widetilde{Kh}^{i,j}(K_n) \to \widetilde{Kh}^{i+n,j}(K_n)$$

is also non-zero for some  $i, j \in \mathbb{Z}$ .

*Proof.* For the first statement, consider the disjoint union  $L_n$  of n copies of the left-handed trefoil T. It follows from [LS14a, Proposition 9.2] that

$$\mathcal{X}_{Kh}(T) \simeq \Sigma^{-3} \mathbb{S} \vee \Sigma^{-2} \mathbb{S} \vee \mathbb{S} \vee \mathbb{S} \vee \Sigma^{-4} \mathbb{R} P^2$$

(compare [LS14a, Example 9.4]). So, by Theorem 1,

$$\mathcal{X}_{Kh}(L_n) \simeq \Sigma^{-4n}(\overbrace{\mathbb{R}\mathrm{P}^2 \wedge \cdots \wedge \mathbb{R}\mathrm{P}^2}^{n \text{ copies}}) \vee Y$$

for some space Y. It follows from the Cartan formula that

$$\operatorname{Sq}^n \colon H^n((\mathbb{R}P^2)^{\wedge n}; \mathbb{Z}/2\mathbb{Z}) \to H^{2n}((\mathbb{R}P^2)^{\wedge n}; \mathbb{Z}/2\mathbb{Z})$$

is non-trivial. This proves the first part of the result.

For the second part of the result, let K be the knot  $15^n_{41127}$  and let  $K_n$  be the connect sum of n copies of K. According to the calculation in [Shu11, Figure 6],  $\widetilde{Kh}^{-2,0}(K;\mathbb{Z}) \cong \mathbb{Z}$ ,  $\widetilde{Kh}^{0,0}(K;\mathbb{Z}) = \mathbb{Z}/2\mathbb{Z}$ , and  $\widetilde{Kh}^{i,0}(K;\mathbb{Z}) = 0$  for  $i \neq -2,0$ . In particular, there is a class  $\alpha \in \widetilde{Kh}^{-1,0}(K;\mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$  so that  $\operatorname{Sq}^1(\alpha)$  is non-zero and  $\operatorname{Sq}^i(\alpha) = 0$  for i > 1. So, it follows from Theorem 2 and the Cartan formula that for the class  $\beta = \alpha \wedge \cdots \wedge \alpha \in \widetilde{Kh}^{-n,0}(K_n)$ ,  $\operatorname{Sq}^n(\beta)$  is non-trivial.

Finally, we argue that  $\operatorname{Sq}^n$  is nonvanishing on  $Kh^{*,*}(K_n)$ , as well. To see this, recall from [LS14a, Theorem 3] that there is a cofibration sequence

$$\widetilde{\mathcal{X}}_{Kh}^{j-1}(K_n) \to \mathcal{X}_{Kh}^j(K_n) \to \widetilde{\mathcal{X}}_{Kh}^{j+1}(K_n)$$

inducing the long exact sequence

$$\cdots \to \widetilde{Kh}^{i,j+1}(K_n) \to Kh^{i,j}(K_n) \to \widetilde{Kh}^{i,j-1}(K_n) \to \widetilde{Kh}^{i+1,j+1}(K_n) \to \cdots$$

Further, for coefficients in  $\mathbb{Z}/2\mathbb{Z}$ ,

$$\dim(Kh^{i,j}(K_n;\mathbb{Z}/2\mathbb{Z})) = \dim(\widetilde{Kh}^{i,j-1}(K_n;\mathbb{Z}/2\mathbb{Z})) + \dim(\widetilde{Kh}^{i,j+1}(K_n;\mathbb{Z}/2\mathbb{Z}))$$

[ORS, Proposition 1.7], so the map  $Kh^{i,j}(K_n; \mathbb{Z}/2\mathbb{Z}) \to \widetilde{Kh}^{i,j-1}(K_n; \mathbb{Z}/2\mathbb{Z})$  is surjective. In particular, there is a class  $\gamma \in Kh^{-n,1}(K_n; \mathbb{Z}/2\mathbb{Z})$  which maps to  $\beta \in \widetilde{Kh}^{-n,0}(K_n; \mathbb{Z}/2\mathbb{Z})$ . By naturality of  $\operatorname{Sq}^n$ , it follows that  $\operatorname{Sq}^n(\gamma)$  maps to  $\operatorname{Sq}^n(\beta) \neq 0$ , so  $\operatorname{Sq}^n(\gamma) \neq 0$ , as desired.

Finally, we turn to the unreduced Khovanov homology of a connected sum.

**Definition 9.3.** Consider the Khovanov homotopy type associated to the unknot,  $\mathcal{X}_{Kh}(U) = \mathbb{S} \vee \mathbb{S}$ , which is the suspension spectrum of  $S^0 \vee S^0 = \{*, p_-, p_+\}$ . The spectrum  $\mathbb{H}^1 := \mathcal{X}_{Kh}(U)$  has a product  $\mu \colon \mathbb{H}^1 \wedge \mathbb{H}^1 \to \mathbb{H}^1$  induced by the map of spaces  $(S^0 \vee S^0) \wedge (S^0 \vee S^0) \to (S^0 \vee S^0)$  given by

$$p_- \wedge p_- \mapsto p_- \qquad p_+ \wedge p_- \mapsto p_+ \qquad p_- \wedge p_+ \mapsto p_+ \qquad p_+ \wedge p_+ \mapsto *.$$

**Lemma 9.4.** The operation  $\mu$  makes  $\mathbb{H}^1$  into a ring spectrum.

*Proof.* This is immediate from the definitions.

Remark 9.5. The map on reduced cohomology induced by  $\mu$  is the split map  $Kh(U) \to Kh(U) \otimes Kh(U)$ . (The generators of Kh(U) are  $x_-$  corresponding to  $p_-$  and  $x_+$  corresponding to  $p_+$ .)

Remark 9.6. The notation  $\mathbb{H}^1$  is chosen to be reminiscent of the first of Khovanov's arc algebras  $H^n$ .

Next, fix a link diagram K with n crossings and a basepoint  $p \in K$ . We make  $\mathcal{X}_{Kh}(K)$  into a module spectrum over  $\mathbb{H}^1$ . We will use the box map realization from Section 5, applied to the functor  $F_{Kh}: \underline{2}^n \to \mathcal{B}$ , using a particular choice of spatial refinement, which we need to specify. The resulting CW complex is described in Proposition 6.1, producing a finite CW spectrum  $\mathcal{X}_{Kh}(K)$ . Before specifying the spatial refinement, we recall the definitions of the reduced Khovanov functors  $F_{\pm Kh}: \underline{2}^n \to \mathcal{B}$ , to fix notation:

Notation 9.7. For  $u \in \{0,1\}^n$ , define  $F_{+Kh}(u)$  (respectively,  $F_{-Kh}(u)$ ) to be the subset of  $F_{Kh}(u)$  where the circle in the complete resolution  $\mathcal{P}(u)$  containing the basepoint is labeled  $x_+$  (respectively,  $x_-$ ). For  $u > v \in \{0,1\}^n$ , define the correspondence from  $F_{+Kh}(u)$  to  $F_{+Kh}(v)$  (respectively, from  $F_{-Kh}(u)$  to  $F_{-Kh}(v)$ ) to be the subset  $s^{-1}(F_{+Kh}(u)) \cap t^{-1}(F_{+Kh}(v))$  (respectively,  $s^{-1}(F_{-Kh}(u)) \cap t^{-1}(F_{-Kh}(v))$ ) of the correspondence from  $F_{Kh}(u)$  to  $F_{Kh}(v)$ . It is straightforward from the definition of the Khovanov differential (Section 2.2) that this produces well-defined functors  $F_{\pm Kh} \colon 2^n \to \mathcal{B}$ . Furthermore, the map from  $F_{+Kh}(u)$  to  $F_{-Kh}(u)$  which relabels the pointed circle in  $\mathcal{P}(u)$  from  $x_+$  to  $x_-$  induces a natural isomorphism from  $F_{+Kh}$  to  $F_{-Kh}$ ; we often write  $F_{Kh}$  to denote either functor. Finally, for any  $u \in \{0,1\}^n$ ,  $F_{Kh}(u) = F_{+Kh}(u)$  If  $F_{-Kh}(u)$ ; and for any  $u > v \in \{0,1\}^n$ , the correspondence  $F_{Kh}(\varphi_{u,v})$  from  $F_{Kh}(u)$  to  $F_{-Kh}(v)$  is the disjoint union of the correspondence  $F_{+Kh}(\varphi_{u,v})$  from  $F_{+Kh}(u)$  to  $F_{+Kh}(v)$ , the correspondence  $F_{-Kh}(\varphi_{u,v})$  from  $F_{-Kh}(u)$  to  $F_{-Kh}(v)$ , and some correspondence from  $F_{-Kh}(u)$  to  $F_{+Kh}(v)$ .

**Definition 9.8.** A spatial refinement  $\widetilde{F}_{Kh}$  of  $F_{Kh}$  induces spatial refinements  $\widetilde{F}_{+Kh}$  of  $F_{+Kh}$  and  $\widetilde{F}_{-Kh}$  of  $F_{-Kh}$ . We call  $\widetilde{F}_{Kh}$  a pointed spatial refinement if, with respect to the natural isomorphism between  $F_{+Kh}$  and  $F_{-Kh}$ , the boxes used to define  $\widetilde{F}_{+Kh}$  and  $\widetilde{F}_{-Kh}$  are identical. In this case, there is an induced natural isomorphism between  $\widetilde{F}_{+Kh}$  and  $\widetilde{F}_{-Kh}$ . For pointed spatial refinements, the CW complexes hocolim( $\widetilde{F}_{+Kh}^+$ ) and hocolim( $\widetilde{F}_{-Kh}^+$ ) are canonically isomorphic.

**Lemma 9.9.** Every pointed link diagram (K,p) admits a pointed spatial refinement  $\widetilde{F}_{Kh}$ .

Proof. We construct a spatial refinement  $\widetilde{F}_{Kh}$  of  $F_{Kh}$  in several steps. First construct a spatial refinement  $\widetilde{F}_{-Kh}$  of  $F_{-Kh}$  with the additional restriction that the box maps come from the subspaces  $E^{\circ}(\{B_x\}, s)$  of  $E(\{B_x\}, s)$ , that is, the sub-boxes are contained in the interiors of the bigger boxes. Then use the natural isomorphism between  $F_{+Kh}$  and  $F_{-Kh}$  to get a spatial refinement  $\widetilde{F}_{+Kh}$  of  $F_{+Kh}$ . Finally, extend  $\widetilde{F}_{+Kh}$ 

and  $\widetilde{F}_{-Kh}$  to construct a spatial refinement  $\widetilde{F}_{Kh}$  of  $F_{Kh}$ , following the inductive argument in the proof of Proposition 5.22 (1). For the induction step, fix a length- $\ell$  sequence  $v_0 \to \cdots \to v_\ell$  of non-identity morphisms in  $\underline{2}^n$ . There is a correspondence  $F_{Kh}(\varphi_{v_0,v_\ell})$  and a subset  $F_{+Kh}(\varphi_{v_0,v_\ell}) \coprod F_{-Kh}(\varphi_{v_0,v_\ell})$ ; let s be the source map of the correspondence  $F_{Kh}(\varphi_{v_0,v_\ell})$  and s' the restriction of s to  $F_{+Kh}(\varphi_{v_0,v_\ell}) \coprod F_{-Kh}(\varphi_{v_0,v_\ell})$ . Induction and  $\widetilde{F}_{+Kh}$  and  $\widetilde{F}_{-Kh}$  give a diagram

$$\partial([0,1]^{\ell-1}) \longrightarrow E^{\circ}(\{B_x\}, s)$$

$$\downarrow \qquad \qquad \downarrow$$

$$[0,1]^{\ell-1} \longrightarrow E^{\circ}(\{B_x\}, s'),$$

where the right-hand vertical map forgets the boxes labeled by elements of  $F_{Kh}(\varphi_{v_0,v_\ell}) \setminus (F_{+Kh}(\varphi_{v_0,v_\ell}) \coprod F_{-Kh}(\varphi_{v_0,v_\ell}))$ . The inductive step is to construct a lift  $[0,1]^{\ell-1} \to E^{\circ}(\{B_x\},s)$  making the diagram commute. Lemma 5.20 guarantees the existence of such a lift. Thus, induction implies that  $F_{Kh}$  has a spatial refinement extending  $\widetilde{F}_{+Kh}$  and  $\widetilde{F}_{-Kh}$ .

Construction 9.10. Given a pointed spatial refinement, define a map  $\Psi$ : hocolim( $\widetilde{F}_{Kh}^+$ )  $\rightarrow$  hocolim( $\widetilde{F}_{Kh}^+$ ) as follows. Notice that hocolim( $\widetilde{F}_{+Kh}^+$ ) is a subcomplex of hocolim( $\widetilde{F}_{Kh}^+$ ) and hocolim( $\widetilde{F}_{-Kh}^+$ ) is the corresponding quotient complex. Define  $\Psi$  to be the composition

$$\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \twoheadrightarrow \operatorname{hocolim}(\widetilde{F}_{-Kh}^+) \stackrel{\cong}{\longrightarrow} \operatorname{hocolim}(\widetilde{F}_{+Kh}^+) \hookrightarrow \operatorname{hocolim}(\widetilde{F}_{Kh}^+),$$

where the first map is the quotient map, the second map is the canonical isomorphism, and the third map is the subcomplex inclusion. Note that  $\Psi$  is a cellular map. The induced map on  $\mathcal{C}_{Kh}(K)$ , the reduced cellular cochain complex of hocolim( $\widetilde{F}_{Kh}^+$ ), sends generators that label the pointed circle by  $x_-$  to zero and on the rest of the Khovanov generators relabels the pointed circle from  $x_+$  to  $x_-$ .

Now we are ready to define the  $\mathbb{H}^1$ -module structure on  $\mathcal{X}_{Kh}(K)$ .

**Definition 9.11.** Define a map  $\mathcal{X}_{Kh}(K) \wedge \mathbb{H}^1 \to \mathcal{X}_{Kh}(K)$  induced the following map of spaces:

$$(9.12) \quad \operatorname{hocolim}(\widetilde{F}_{Kh}^+) \wedge \{*, p_-, p_+\} = (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_-\}) \vee (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_+\}) \rightarrow \operatorname{hocolim}(\widetilde{F}_{Kh}^+) \wedge \{*, p_-, p_+\} = (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_-\}) \vee (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_+\}) \rightarrow \operatorname{hocolim}(\widetilde{F}_{Kh}^+) \wedge \{*, p_-, p_+\} = (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_-\}) \vee (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_+\}) \rightarrow \operatorname{hocolim}(\widetilde{F}_{Kh}^+) \wedge \{*, p_-, p_+\} = (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_-\}) \vee (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_+\}) \rightarrow \operatorname{hocolim}(\widetilde{F}_{Kh}^+) \wedge \{*, p_-, p_+\} = (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_-\}) \vee (\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_+\}) \wedge (\operatorname{hocolim}(\widetilde{F}_{Kh}^+)$$

On the first summand, the map  $\operatorname{hocolim}(\widetilde{F}_{Kh}^+) \times \{p_-\} \to \operatorname{hocolim}(\widetilde{F}_{Kh}^+)$  is projection to the first factor. On the second summand, the map is projection to the first factor composed with the map  $\Psi \colon \operatorname{hocolim}(\widetilde{F}_{Kh}^+) \to \operatorname{hocolim}(\widetilde{F}_{Kh}^+)$  defined above.

**Lemma 9.13.** Definition 9.11 makes  $\mathcal{X}_{Kh}(K)$  into a module spectrum over  $\mathbb{H}^1$ .

*Proof.* This follows from the fact that  $\Psi \circ \Psi$  sends all of  $\operatorname{hocolim}(\widetilde{F}_{Kh}^+)$  to the basepoint.

The ring spectrum  $\mathbb{H}^1$  is commutative, so we can view the action of  $\mathbb{H}^1$  on  $\mathcal{X}_{Kh}(K)$  as either a left or a right action.

Note that the induced map on the reduced cellular cochain complexes associated to the map (9.12) is the split map  $\mathcal{C}_{Kh}(K) \to \mathcal{C}_{Kh}(K) \otimes Kh(U)$ .

**Proposition 9.14.** Up to weak equivalence of  $\mathbb{H}^1$ -module spectra,  $\mathcal{X}_{Kh}(K)$  is an invariant of pointed links. That is, if (K,p) and (K',p') are pointed link diagrams representing isotopic pointed links, then there exist  $\mathbb{H}^1$ -module spectra  $\mathcal{X}_{Kh}(K) = X_0, X_1, \ldots, X_{\ell-1}, X_{\ell} = \mathcal{X}_{Kh}(K')$ , and for any adjacent pair  $X_i, X_{i+1}$ , either a map  $X_i \to X_{i+1}$  or a map  $X_{i+1} \to X_i$ , which is both an  $\mathbb{H}^1$ -module map and a weak equivalence.

*Proof.* First, observe that the  $\mathbb{H}^1$ -module structure is independent of the choice of box maps; the proof is essentially the same as Proposition 5.22 (2), but using Lemma 5.20 instead of Lemma 5.18.

Next we show that, up to weak equivalence, this  $\mathbb{H}^1$ -module spectrum is invariant under Reidemeister moves. By a standard argument [Kho03, Section 3], we only need to consider Reidemeister moves that do not cross the marked point p. We follow the framework from [LS14a, Section 6]. Let  $K_0$  and  $K_1$  (with  $n_0$  and  $n_1$  crossings respectively) be pointed link diagrams related by any of the three Reidemeister moves of [LS14a, Figure 6.1], and assume  $n_0 < n_1$ . The usual proof of invariance of Khovanov homology shows that  $\mathcal{C}_{Kh}(K_0)$  can be identified with a subquotient complex of  $\mathcal{C}_{Kh}(K_1)$ , inducing a (two-step) zig-zag of isomorphisms connecting  $Kh(K_0)$  and  $Kh(K_1)$  (see also the proofs of [LS14a, Propositions 6.2–6.4]). Indeed, there is a particular vertex  $w \in \{0,1\}^{n_1-n_0}$ , so that for every  $u \in \{0,1\}^{n_0}$ ,  $F_{Kh}(K_0)(u)$  is identified with a certain subset  $S_u \subseteq F_{Kh}(K_1)((u,w))$ , and for every  $u > v \in \{0,1\}^{n_0}$ , the correspondence  $F_{Kh}(K_0)(\varphi_{u,v})$  is identified with the subset  $s^{-1}(S_u) \cap t^{-1}(S_v) \subseteq F_{Kh}(K_1)(\varphi_{(u,w),(v,w)})$ . Furthermore, these identifications identify  $F_{+Kh}(K_0)(u)$  with  $S_u \cap F_{+Kh}(K_1)((u,w))$  and consequently,  $F_{-Kh}(K_0)(u)$  with  $S_u \cap F_{-Kh}(K_1)((u,w))$ .

Construct the  $\mathbb{H}^1$ -module spectrum  $\mathcal{X}_{Kh}(K_1)$  using some pointed spatial refinement  $\tilde{F}_{Kh}(K_1)$  for  $K_1$ . Restricting to the subsets  $S_u$  and the correspondences between them, we get a pointed spatial refinement  $\tilde{F}_{Kh}(K_0)$  for  $K_0$ , which we use to construct the  $\mathbb{H}^1$ -module spectrum  $\mathcal{X}_{Kh}(K_0)$ . With the CW complex structures from Proposition 6.1, hocolim( $\tilde{F}_{Kh}^+(K_0)$ ) can be identified with a subquotient complex of hocolim( $\tilde{F}_{Kh}^+(K_1)$ ), leading to a two-step zig-zag of maps connecting  $\mathcal{X}_{Kh}(K_1)$  and  $\mathcal{X}_{Kh}(K_2)$ . Since the Reidemeister moves do not cross the basepoint, it is immediate from the definitions of these sub- and quotient complexes that the maps are  $\mathbb{H}^1$ -equivariant. Since they also induce isomorphisms on homology, they are weak equivalences.

For the rest of this section, fix a link diagram for  $K_1 \coprod K_2$ , which is a disjoint union of link diagrams for  $K_1$  and  $K_2$ , with  $n_1$  and  $n_2$  crossings respectively, and fix basepoints  $p_i$  on  $K_i$  so that the two basepoints are next to one another (i.e., on the boundary of the same component of  $S^2 \setminus (K_1 \coprod K_2)$ ).

Recall:

**Definition 9.15.** The *(derived) tensor product* of the module spectra  $\mathcal{X}_{Kh}(K_1)$  and  $\mathcal{X}_{Kh}(K_2)$  is the homotopy colimit of the diagram

$$(9.16) \qquad \mathcal{X}_{Kh}(K_1) \wedge \mathcal{X}_{Kh}(K_2) \rightleftharpoons \mathcal{X}_{Kh}(K_1) \wedge \mathbb{H}^1 \wedge \mathcal{X}_{Kh}(K_2) \rightleftharpoons \mathcal{X}_{Kh}(K_1) \wedge \mathbb{H}^1 \wedge \mathcal{X}_{Kh}(K_2) \rightleftharpoons \cdots$$

where the maps are all possible ways of applying  $\mu$  to a pair of consecutive factors. To be more precise, let  $\Delta_{inj}$  be the category with one object  $\underline{\mathbf{n}} = \{0, \dots, n-1\}$  for each positive integer n and  $\operatorname{Hom}(\underline{\mathbf{m}},\underline{\mathbf{n}})$  the set of order-preserving injections  $\{0,\dots,m-1\}\to\{0,\dots,n-1\}$ ; for n>0 and  $0\leq i\leq n$ , let  $f_{\underline{\mathbf{n}},i}\in\operatorname{Hom}_{\Delta_{inj}}(\underline{\mathbf{n}},\underline{\mathbf{n}}+\underline{\mathbf{1}})$  be the morphism  $\underline{\mathbf{n}}\to\underline{\mathbf{n}}+\underline{\mathbf{1}}$  whose image is  $\underline{\mathbf{n}}+\underline{\mathbf{1}}\setminus\{i\}$ . (The category  $\Delta_{inj}$  is the subcategory of the simplex category generated by the face maps, and the  $f_{\underline{\mathbf{n}},i}$  are the face maps themselves.) Then the diagram (9.16) can be treated as a (strict) functor  $F_{\otimes}$  from  $\Delta_{inj}^{\mathrm{op}}$  to  $\mathscr{CW}$ , the category of CW spectra. On objects,  $F_{\otimes}(\underline{\mathbf{n}}) = \mathcal{X}_{Kh}(K_1) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1) \wedge \mathcal{X}_{Kh}(K_2)$ . On morphisms,  $F_{\otimes}(f_{\underline{\mathbf{n}},i}^{\mathrm{op}})$  is the map  $\mathcal{X}_{Kh}(K_1) \wedge (\bigwedge_{i=1}^{n} \mathbb{H}^1) \wedge \mathcal{X}_{Kh}(K_1) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1) \wedge \mathcal{X}_{Kh}(K_2)$  gotten by applying  $\mu$  to the  $(i+1)^{\mathrm{th}}$  pair of consecutive factors. Let  $\mathcal{X}_{Kh}(K_1) \otimes_{\mathbb{H}^1} \mathcal{X}_{Kh}(K_2) := \operatorname{hocolim}(F_{\otimes})$  denote the derived tensor product of  $\mathcal{X}_{Kh}(K_1)$  and  $\mathcal{X}_{Kh}(K_2)$ .

**Theorem 10.** There is a stable homotopy equivalence  $\mathcal{X}_{Kh}(K_1 \# K_2) \simeq \mathcal{X}_{Kh}(K_1) \otimes_{\mathbb{H}^1} \mathcal{X}_{Kh}(K_2)$ .

The proof of Theorem 10 involves three components. First, we show how the functor  $F_{Kh}(K_1 \# K_2)$  is determined by the functors  $F_{Kh}(K_1)$  and  $F_{Kh}(K_2)$ ; this is Lemmas 9.18 and 9.20, which take a little work but are purely combinatorial. Second, in Lemmas 9.21 and 9.23, we prove that Theorem 10 holds at

the level of cellular cochains, for an appropriate CW complex structure on  $\mathcal{X}_{Kh}(K_1) \otimes_{\mathbb{H}^1} \mathcal{X}_{Kh}(K_2)$ . This is essentially immediate from Segal's construction of homotopy colimits and the connected sum theorem for the Khovanov chain complex. Third, using the description of  $F_{Kh}(K_1 \# K_2)$  in terms of  $F_{Kh}(K_1)$  and  $F_{Kh}(K_2)$  and carefully chosen spatial refinements, we produce a (strict) map from the diagram (9.16) to  $\mathcal{X}_{Kh}(K_1 \# K_2)$ , inducing the desired map of cellular cochains. This argument is Lemmas 9.24 and 9.25. From these three steps, Theorem 10 follows easily.

We begin by reconstructing the functor  $F_{Kh}(K_1 \# K_2)$  from the functor  $F_{Kh}(K_1 \coprod K_2): \underline{2}^{n_1+n_2} \to \mathscr{B}$ .

**Definition 9.17.** For  $a, b \in \{*, +, -\}$ , let  $F_{abKh}(K_1 \coprod K_2) : \underline{2}^{n_1+n_2} \to \mathcal{B}$  denote the functor where we only consider the Khovanov generators that label the circle containing  $p_1$  by  $x_a$  if  $a \in \{+, -\}$  and label the circle containing  $p_2$  by  $x_b$  if  $b \in \{+, -\}$ , and we restrict the correspondences correspondingly. (If a or b is \*, we make no restriction on the label of the corresponding circle.)

**Lemma 9.18.** For  $v \in \{0,1\}^{n_1+n_2}$ , the map from  $F_{+-Kh}(K_1 \coprod K_2)(v)$  to  $F_{-+Kh}(K_1 \coprod K_2)(v)$  that interchanges the labelings of the two pointed circles in  $\mathcal{P}(v)$  induces an isomorphism from  $F_{+-Kh}(K_1 \coprod K_2)$  to  $F_{-+Kh}(K_1 \coprod K_2)$ .

*Proof.* The isomorphism from Proposition 9.1 identifies either functor to  $F_{\widetilde{Kh}}(K_1) \times F_{\widetilde{Kh}}(K_2)$ . The given map is the composition  $F_{+-Kh}(K_1 \coprod K_2) \cong F_{\widetilde{Kh}}(K_1) \times F_{\widetilde{Kh}}(K_2) \cong F_{-+Kh}(K_1 \coprod K_2)$ .

$$F(u) = (F_{--Kh}(u) \coprod F_{+-Kh}(u) \coprod F_{-+Kh}(u))/(F_{+-Kh}(u) = F_{-+Kh}(u));$$

and for all  $u > v \in \{0, 1\}^{n_1 + n_2}$ ,

$$F(\varphi_{u,v}) = \left(F_{--Kh}(\varphi_{u,v}) \coprod F_{+-Kh}(\varphi_{u,v}) \coprod F_{-+Kh}(\varphi_{u,v}) \coprod P_{u,v} \coprod Q_{u,v}\right) / (F_{+-Kh}(\varphi_{u,v}) = F_{-+Kh}(\varphi_{u,v})).$$

**Lemma 9.20.** The functor F constructed above is isomorphic to  $F_{Kh}(K_1 \# K_2)$  via the following map: for all  $u \in \{0,1\}^{n_1+n_2}$ , the isomorphism sends  $x \in F(u)$  to  $y \in F_{Kh}(K_1 \# K_2)(u)$  where y labels the connect-sum circle by  $x_-$  if and only if x labels both the pointed circles by  $x_-$ , and x and y label all the circles that are disjoint from the connect-sum region identically.

Proof. The proof is similar to the proof of Proposition 9.1. To keep the notations similar, let  $X_v = F(v)$ ,  $A_{u,v} = F(\varphi_{u,v})$ ,  $G = F_{Kh}(K_1 \# K_2)$ ,  $Y_v = G(v)$ , and  $B_{u,v} = G(\varphi_{u,v})$ . The bijections  $\phi_v \colon X_v \xrightarrow{\cong} Y_v$  are already provided to us. To construct bijections  $\psi_{u,v} \colon A_{u,v} \xrightarrow{\cong} B_{u,v}$ , for all u > v with |u| - |v| = 1, we need to check the conditions (1) and (2) of the proof of Proposition 9.1.

For any  $u \in \{0,1\}^{n_1+n_2}$  and any  $z_u \in F_{+\!\!\!/+\!\!\!/+\!\!\!/}K_h(K_1 \coprod K_2)(u)$ , let  $\pi(z_u)$  denote the image of  $z_u$  in  $X_u$ ; and for any  $x_u \in X_u$ , let  $\iota^1(x_u)$  (respectively,  $\iota^2(x_u)$ ) denote the preimage of  $x_u$  in  $F_{*-Kh}(K_1 \coprod K_2)(u)$  (respectively,  $F_{-*Kh}(K_1 \coprod K_2)(u)$ ). Then for any u > v with |u| - |v| = 1,  $z_u \in F_{+\!\!\!/+\!\!\!/-\!\!\!/}K_h(K_1 \coprod K_2)(u)$ , and  $x_v \in X_v$ , one of the two subsets  $s^{-1}(z_u) \cap t^{-1}(\iota^1(x_v)) \subseteq F_{*-Kh}(K_1 \coprod K_2)(\varphi_{u,v})$  and  $s^{-1}(z_u) \cap t^{-1}(\iota^2(x_v)) \subseteq F_{-*Kh}(K_1 \coprod K_2)(\varphi_{u,v})$  is empty, and the other one is canonically identified with the subset  $s^{-1}(\pi(z_u)) \cap t^{-1}(x_v) \subseteq A_{u,v}$ . This follows from the fact that the correspondences in  $F_{+\!\!\!/-\!\!\!/}K_h(K_1 \coprod K_2)$  preserve two quantum gradings, the one coming

from  $K_1$  and the one coming from  $K_2$ ; however, the double quantum gradings of  $\iota^1(x_v)$  and  $\iota^2(x_v)$  are different, and therefore, at least one of  $s^{-1}(z_u) \cap t^{-1}(\iota^1(x_v))$  and  $s^{-1}(z_u) \cap t^{-1}(\iota^2(x_v))$  is empty.

From this observation, condition (1) is immediate. Condition (2) follows from additionally noting that the composition  $F_{Kh}(K_1 \coprod K_2) \to X \xrightarrow{\phi} Y$  induces the cobordism map  $\mathcal{C}_{Kh}(K_1 \# K_2) \to \mathcal{C}_{Kh}(K_1 \coprod K_2)$ associated to splitting at the connect-sum region, which is a chain map.

Finally, we need to check that diagram (9.2) commutes for all u > w with |u| - |w| = 2. Using the observation in the previous paragraph, this follows from the same arguments as in the proof of Proposition 9.1.

Next we observe that Theorem 10 holds at the level of cellular cochain complexes.

**Lemma 9.21.** There exists a CW complex structure on  $\mathcal{X}_{Kh}(K_1) \otimes_{\mathbb{H}^1} \mathcal{X}_{Kh}(K_2)$  so that the reduced cellular cochain complex is the following chain complex

 $(9.22) \ \mathcal{C}_{Kh}(K_1) \otimes \mathcal{C}_{Kh}(K_2) \to \mathcal{C}_{Kh}(K_1) \otimes Kh(U) \otimes \mathcal{C}_{Kh}(K_2) \to \mathcal{C}_{Kh}(K_1) \otimes Kh(U) \otimes Kh(U) \otimes \mathcal{C}_{Kh}(K_2) \to \cdots$ with the differential given by

$$d(x_0 \otimes \cdots \otimes x_n) = \delta(x_0) \otimes \cdots \otimes x_i \otimes \cdots \otimes x_n + x_0 \otimes \cdots \otimes x_i \otimes \cdots \otimes \delta(x_n)$$
$$+ \sum_{i=0}^n (-1)^{i+n} x_0 \otimes \cdots \otimes x_{i-1} \otimes S(x_i) \otimes x_{i+1} \otimes \cdots \otimes x_n,$$

where S denotes either the Khovanov Frobenius algebra comultiplication map  $Kh(U) \to Kh(U) \otimes Kh(U)$  or the cobordism map  $\mathcal{C}_{Kh}(K_1) \to \mathcal{C}_{Kh}(K_1) \otimes Kh(U)$  (respectively,  $\mathcal{C}_{Kh}(K_2) \to Kh(U) \otimes \mathcal{C}_{Kh}(K_2)$ ) for splitting off a trivial unknot at  $p_1$  (respectively,  $p_2$ ), and  $\delta$  denotes the Khovanov differential on  $\mathcal{C}_{Kh}(K_1)$  and  $\mathcal{C}_{Kh}(K_2)$ .

*Proof.* If  $F: \Delta_{inj}^{op} \to \mathsf{CW}_{\bullet}$  is a functor, Segal [Seg74, Appendix A] defines its geometric realization as

$$||F|| = \left( \{*\} \coprod \prod_{n=1}^{\infty} \Delta^{n-1} \times F(\underline{\mathbf{n}}) \right) / \sim$$

with  $(f_*(\zeta), a) \sim (\zeta, F(f^{\text{op}})(a))$  for all  $\zeta \in \Delta^{m-1}$ ,  $a \in F(\underline{\mathbf{n}})$ , and  $f \in \text{Hom}_{\Delta_{inj}}(\underline{\mathbf{m}}, \underline{\mathbf{n}})$ , and  $\Delta^{n-1} \times \{*\} \sim *$  for all n. (The map  $f_*: \Delta^{m-1} \to \Delta^{n-1}$  is the face inclusion corresponding to f.) If  $\Delta^{n-1}_+$  denotes the disjoint union of  $\Delta^{n-1}$  and a basepoint, then we can rewrite

$$||F|| = \left(\bigvee_n \Delta^{n-1}_+ \wedge F(\underline{\mathbf{n}})\right)/\sim$$

with just the relation  $(f_*(\zeta), a) \sim (\zeta, F(f^{\text{op}})(a))$  for all  $\zeta \in \Delta^{m-1}$ ,  $a \in F(\underline{\mathbf{n}})$ , and  $f \in \text{Hom}_{\Delta_{inj}}(\underline{\mathbf{m}}, \underline{\mathbf{n}})$ . Let  $F_{\otimes,\ell} \colon \Delta^{\text{op}}_{inj} \to \mathsf{CW}_{\bullet}$  be the functor obtained from  $F_{\otimes} \colon \Delta^{\text{op}}_{inj} \to \mathscr{CW}$  by looking at the  $\ell^{\text{th}}$  spaces in the spectra. Define the geometric realization  $||F_{\otimes}||$  as a CW spectrum whose  $\ell^{\text{th}}$  space is  $||F_{\otimes}||_{\ell} = ||F_{\otimes,\ell}||$ 

$$\Sigma \|F_{\otimes}\|_{\ell} \cong \Sigma \Big(\bigvee_{n} \Delta_{+}^{n-1} \wedge F_{\otimes,\ell}(\underline{\mathbf{n}})\Big) / \sim = \Big(\bigvee_{n} \Delta_{+}^{n-1} \wedge \Sigma F_{\otimes,\ell}(\underline{\mathbf{n}})\Big) / \sim \longrightarrow \Big(\bigvee_{n} \Delta_{+}^{n-1} \wedge F_{\otimes,\ell+1}(\underline{\mathbf{n}})\Big) / \sim = \|F_{\otimes}\|_{\ell+1},$$

where the middle arrow is induced by the structure maps  $\Sigma(F_{\otimes}(n))_{\ell} \to (F_{\otimes}(n))_{\ell+1}$ . Equipped with the natural CW complex structure ([Seg74, Proposition A.1(i)]), the reduced cellular cochain complex of  $|F_{\otimes}|$ is easily seen to be the one from Formula (9.22).

To identify  $||F_{\otimes}||$  with  $hocolim(F_{\otimes})$ , use the construction of homotopy colimits of the strict functor  $F_{\otimes}$  via simplices, instead of cubes; Vogt [Vog73, Corollary 8.5] shows the two definitions agree. Using the simplicial model for the homotopy colimit, it is well known that  $hocolim(F_{\otimes})$  is the barycentric subdivision of  $||F_{\otimes}||$ . (See also [Seg74, Proposition A.3] where  $|\text{simp}(\cdot)|$  plays the role of this space.)

**Lemma 9.23.** The cobordism map  $S: \mathcal{C}_{Kh}(K_1 \# K_2) \to \mathcal{C}_{Kh}(K_1) \otimes \mathcal{C}_{Kh}(K_2)$  associated to splitting at the connected sum region induces a quasi-isomorphsim

$$\mathcal{C}_{Kh}(K_1 \# K_2)$$

$$\downarrow^S$$

$$\mathcal{C}_{Kh}(K_1) \otimes \mathcal{C}_{Kh}(K_2) \longrightarrow \mathcal{C}_{Kh}(K_1) \otimes Kh(U) \otimes \mathcal{C}_{Kh}(K_2) \longrightarrow \mathcal{C}_{Kh}(K_1) \otimes Kh(U) \otimes Kh(U) \otimes \mathcal{C}_{Kh}(K_2) \longrightarrow \cdots$$

*Proof.* The Khovanov complex  $\mathcal{C}_{Kh}(K_1 \# K_2)$  is the cotensor product of  $\mathcal{C}_{Kh}(K_1)$  and  $\mathcal{C}_{Kh}(K_2)$  as comodules over Kh(U) [LS14a, Lemma 10.5], while Formula (9.22) is the derived cotensor product of  $\mathcal{C}_{Kh}(K_1)$  and  $\mathcal{C}_{Kh}(K_2)$ . Thus, the statement presumably follows from the fact that  $\mathcal{C}_{Kh}(K_1)$  and  $\mathcal{C}_{Kh}(K_2)$  are co-flat. Rather than going down this rabbit hole, dualize the complex (9.22) over  $\mathbb{Z}$ , which exchanges the split map S and the merge map m, the (derived) cotensor product and the (derived) tensor product, and  $\mathcal{C}_{Kh}(K)$ and  $\mathcal{C}_{Kh}(m(K))$ . (The last assertion is [Kho00, Proposition 32].) The result then follows from the fact that  $\mathcal{C}_{Kh}(m(K_i))$  is free as a Kh(U)-module and Khovanov's connected sum theorem [Kho03, Proposition

We turn to the third part of the argument, constructing compatible spatial refinements for  $F_{Kh}(K_1 \# K_2)$ and  $F_{Kh}(K_1 \coprod U \coprod \cdots \coprod U \coprod K_2)$ .

**Lemma 9.24.** Consider any spatial refinement  $F_{+/+Kh}(K_1 \coprod K_2)$  of  $F_{+/+Kh}(K_1 \coprod K_2)$  whose induced spatial refinements  $\widetilde{F}_{-+Kh}(K_1 \coprod K_2)$  and  $\widetilde{F}_{+-Kh}(K_1 \coprod K_2)$  of  $F_{-+Kh}(K_1 \coprod K_2) \cong F_{+-Kh}(K_1 \coprod K_2)$  agree. Then, identifying  $\widetilde{F}_{-+Kh}(K_1 \coprod K_2)$  and  $\widetilde{F}_{+-Kh}(K_1 \coprod K_2)$  produces a spatial refinement  $\widetilde{F}_{Kh}(K_1 \# K_2)$  of  $F_{Kh}(K_1\#K_2)$ .

*Proof.* It is immediate from the definitions that identifying  $\widetilde{F}_{-+Kh}(K_1 \coprod K_2)$  and  $\widetilde{F}_{+-Kh}(K_1 \coprod K_2)$  produces a spatial refinement of the functor F above. The isomorphism from Lemma 9.20 then produces the spatial refinement  $F_{Kh}(K_1 \# K_2)$  of  $F_{Kh}(K_1 \# K_2)$ .

Let  $\overline{\Delta}_{inj} = \Delta_{inj} \cup \underline{\mathbf{0}}$  be the category obtained by adding an object  $\underline{\mathbf{0}} = \emptyset$  to  $\Delta_{inj}$  and a unique morphism  $\underline{\mathbf{0}} \to \underline{\mathbf{n}}$  for each n; let  $f_{\underline{\mathbf{0}},0}$  denote the unique morphism from  $\underline{\mathbf{0}}$  to  $\underline{\mathbf{1}}$ . We will extend Diagram (9.16) to construct a functor  $\overline{\Delta}_{inj}^{\text{op}} \to \mathscr{C} \mathscr{W}$ .

**Lemma 9.25.** There exists a functor  $\overline{F}_{\times} : \overline{\Delta}_{inj}^{op} \to \mathscr{CW}$  satisfying the following:

- (1)  $\overline{F}_{\times}(\underline{\mathbf{0}}) = \mathcal{X}_{Kh}(K_1 \# K_2)$  and  $\overline{F}_{\times}(\underline{\mathbf{n}}) = \mathcal{X}_{Kh}(K_1 \coprod K_2) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1)$  for all n > 0. (2) Let  $F_{\times}$  denote the restriction  $\overline{F}_{\times}|_{\Delta_{inj}^{\mathrm{op}}}$ . Then there is a natural transformation  $\eta$  from the functor  $F_{\otimes}$ of diagram (9.16) to  $F_{\times}$ , so that for all n > 0,  $\eta_{\underline{\mathbf{n}}} \colon F_{\otimes}(\underline{\mathbf{n}}) \to F_{\times}(\underline{\mathbf{n}})$  sends each cell in  $\mathcal{X}_{Kh}(K_1) \wedge K_{Kh}(K_1) \to K_{Kh}(K_1)$  $(\bigwedge_{i=1}^{n-1} \mathbb{H}^1) \wedge \mathcal{X}_{Kh}(K_2)$  to the corresponding cell in  $\mathcal{X}_{Kh}(K_1 \coprod K_2) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1)$  by a degree one map.
- (3)  $\overline{F}_{\times}(f_{\mathbf{0},\mathbf{0}}^{\mathrm{op}})$  is a map  $\mathcal{X}_{Kh}(K_1 \coprod K_2) \to \mathcal{X}_{Kh}(K_1 \# K_2)$  so that the induced map on reduced cellular cochains is the cobordism map

$$S: \mathcal{C}_{Kh}(K_1 \# K_2) \to \mathcal{C}_{Kh}(K_1 \coprod K_2)$$

induced by splitting at the connected sum region.

from  $\mathcal{C}_{Kh}(K_1 \# K_2)$  to the chain complex from Formula (9.22).

Proof. During the construction of  $\overline{F}_{\times}$ , we will use spatial refinements for  $K_1 \coprod K_2$  that are pointed spatial refinements with respect to both  $p_1$  and  $p_2$ . We will call such spatial refinements doubly pointed spatial refinements. Doubly pointed spatial refinements are spatial refinements that agree on  $F_{*+Kh}(K_1 \coprod K_2)$  and  $F_{*-Kh}(K_1 \coprod K_2)$ , and also on  $F_{*+Kh}(K_1 \coprod K_2)$  and  $F_{-*Kh}(K_1 \coprod K_2)$  (and, therefore, agree on  $F_{++Kh}(K_1 \coprod K_2)$ ,  $F_{+-Kh}(K_1 \coprod K_2)$ ,  $F_{+-Kh}(K_1 \coprod K_2)$ , and  $F_{--Kh}(K_1 \coprod K_2)$ . The CW spectrum  $\mathcal{X}_{Kh}(K_1 \coprod K_2)$  constructed using any such doubly pointed spatial refinement can be viewed as a strict bimodule over  $\mathbb{H}^1$ , with the two actions coming from the two basepoints  $p_1$  and  $p_2$ . Therefore, we can construct a strict functor  $G: \Delta_{inj}^{\mathrm{op}} \to \mathbb{C}W_{\bullet}$  by declaring  $G(\underline{\mathbf{n}}) = \mathcal{X}_{Kh}(K_1 \coprod K_2) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1)$  and by defining the map  $G(f_{\underline{\mathbf{n}},i}^{\mathrm{op}}) \colon \mathcal{X}_{Kh}(K_1 \coprod K_2) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1) \to \mathcal{X}_{Kh}(K_1 \coprod K_2) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1)$  to be the ring multiplication map applied to the  $i^{\mathrm{th}}$  pair of consecutive  $\mathbb{H}^1$ -factors if 0 < i < n, and the bimodule map coming from  $p_1$  (respectively,  $p_2$ ) using the first (respectively, last)  $\mathbb{H}^1$  factor if i = 0 (respectively, n).

Now we are in a position to construct  $F_{\times}$ . The construction proceeds in several stages.

- (F-1) We start with pointed spatial refinements  $\widetilde{F}_{Kh}(K_1)$  and  $\widetilde{F}_{Kh}(K_2)$  of  $F_{Kh}(K_1)$  and  $F_{Kh}(K_2)$  (using  $k_1$ -dimensional and  $k_2$ -dimensional boxes with  $k_1 + k_2 = k$ ). Since  $F_{Kh}(K_1 \coprod K_2) = F_{Kh}(K_1) \times F_{Kh}(K_2)$  (Proposition 9.1),  $\widetilde{F}_{Kh}(K_1 \coprod K_2) := \widetilde{F}_{Kh}(K_1) \wedge \widetilde{F}_{Kh}(K_2)$  (cf. (ho-3)) is a doubly pointed spatial refinement for  $K_1 \coprod K_2$ .
- (F-2) Define  $F_{\times} : \Delta_{inj}^{\text{op}} \to \mathscr{CW}$  to be the functor associated to this doubly pointed spatial refinement. This automatically satisfies the second part of Lemma 9.25 (1). To relate  $F_{\otimes}$  and  $F_{\times}$ , observe that for all n > 0,  $F_{\times}(\underline{\mathbf{n}}) = F_{\times}(\underline{\mathbf{1}}) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1)$  and  $F_{\otimes}(\underline{\mathbf{n}})$  is canonically isomorphic to  $F_{\otimes}(\underline{\mathbf{1}}) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1)$  (preserving the order of the  $\mathbb{H}^1$ -factors); therefore, it is enough to relate  $F_{\times}(\underline{\mathbf{1}})$ , which is a formal desuspension of the suspension spectrum of hocolim  $((\widetilde{F}_{Kh}(K_1) \wedge \widetilde{F}_{Kh}(K_2))^+)$ , and  $F_{\otimes}(\underline{\mathbf{1}})$ , which is a formal desuspension of the suspension spectrum of hocolim  $((\widetilde{F}_{Kh}(K_1))^+) \wedge \text{hocolim}((\widetilde{F}_{Kh}(K_2))^+)$ . However, the former is easily seen to be a quotient of the latter, with the quotient map sending each cell by a degree one map to the corresponding cell. This proves Lemma 9.25 (2).
- (F-3) The doubly pointed spatial refinement  $\widetilde{F}_{Kh}(K_1 \coprod K_2)$  induces a spatial refinement  $\widetilde{F}_{+/+Kh}(K_1 \coprod K_2)$  of  $F_{+/+Kh}(K_1 \coprod K_2)$ ; and its induced spatial refinements  $\widetilde{F}_{-+Kh}(K_1 \coprod K_2)$  and  $\widetilde{F}_{+-Kh}(K_1 \coprod K_2)$  of  $F_{-+Kh}(K_1 \coprod K_2)$  and  $F_{+-Kh}(K_1 \coprod K_2)$  agree (with  $F_{-+Kh}(K_1 \coprod K_2)$  and  $F_{+-Kh}(K_1 \coprod K_2)$  identified by Lemma 9.18). Therefore, by Lemma 9.24, we get a (pointed) spatial refinement  $\widetilde{F}_{Kh}(K_1 \# K_2)$  for  $K_1 \# K_2$  (with the basepoint chosen on either of the two strands near the connect sum region).
- (F-4) We use  $\widetilde{F}_{Kh}(K_1\#K_2)$  to construct the CW spectrum  $\overline{F}_{\times}(\underline{\mathbf{0}})=\mathcal{X}_{Kh}(K_1\#K_2)$ ; this automatically satisfies the first part of Lemma 9.25 (1).
- (F-5) The space hocolim  $((\widetilde{F}_{+/+Kh}(K_1 \coprod K_2))^+)$  is a quotient complex of hocolim  $((\widetilde{F}_{Kh}(K_1 \coprod K_2))^+)$ ; it has two subcomplexes hocolim  $((\widetilde{F}_{-+Kh}(K_1 \coprod K_2))^+)$  and hocolim  $((\widetilde{F}_{+-Kh}(K_1 \coprod K_2))^+)$  that have a canonical isomorphism between them; and the coequalizer is canonically identified with the space hocolim  $((\widetilde{F}_{Kh}(K_1 \# K_2))^+)$ . That is, we have a diagram

$$\operatorname{hocolim}\left((\widetilde{F}_{-+Kh}(K_1 \amalg K_2))^+\right) \cong \operatorname{hocolim}\left((\widetilde{F}_{+-Kh}(K_1 \amalg K_2))^+\right)$$

$$\operatorname{hocolim}\left((\widetilde{F}_{Kh}(K_1 \amalg K_2))^+\right) \longrightarrow \operatorname{hocolim}\left((\widetilde{F}_{++Kh}(K_1 \amalg K_2))^+\right)$$


$$\operatorname{hocolim}\left((\widetilde{F}_{Kh}(K_1 \# K_2))^+\right).$$

Let  $\overline{F}_{\times}(f_{\underline{\mathbf{0}},0}^{\mathrm{op}}) \colon \overline{F}_{\times}(\underline{\mathbf{1}}) = \mathcal{X}_{Kh}(K_1 \coprod K_2) \to \mathcal{X}_{Kh}(K_1 \# K_2) = \overline{F}_{\times}(\underline{\mathbf{0}})$  be the map induced from the composition hocolim  $((\widetilde{F}_{Kh}(K_1 \coprod K_2))^+) \to \operatorname{hocolim}((\widetilde{F}_{Kh}(K_1 \# K_2))^+)$ . This map is a cellular map sending the cells in  $\mathcal{X}_{Kh}(K_1 \coprod K_2)$  to the corresponding cells in  $\mathcal{X}_{Kh}(K_1 \# K_2)$  by degree one maps (with the correspondence described in Lemma 9.20). Therefore, this map satisfies Lemma 9.25 (3).


(F-6) We have to define the map  $\overline{F}_{\times}(\underline{2}) \to \overline{F}_{\times}(\underline{0})$  to be both  $\overline{F}_{\times}(f_{\underline{0},0}^{\text{op}}) \circ \overline{F}_{\times}(f_{\underline{1},0}^{\text{op}})$  and  $\overline{F}_{\times}(f_{\underline{0},0}^{\text{op}}) \circ \overline{F}_{\times}(f_{\underline{1},1}^{\text{op}})$ ; so we merely need to check that the latter two maps agree. The two maps  $\overline{F}_{\times}(f_{\underline{1},0}^{\text{op}})$  and  $\overline{F}_{\times}(f_{\underline{1},1}^{\text{op}})$  are both induced from maps of spaces

$$\begin{aligned} &\operatorname{hocolim}\left((\widetilde{F}_{Kh}(K_1 \amalg K_2))^+\right) \wedge \{*, p_-, p_+\} \\ &= \left(\operatorname{hocolim}\left((\widetilde{F}_{Kh}(K_1 \amalg K_2))^+\right) \times \{p_-\}\right) \vee \left(\operatorname{hocolim}\left((\widetilde{F}_{Kh}(K_1 \amalg K_2))^+\right) \times \{p_+\}\right) \\ &\to \operatorname{hocolim}\left((\widetilde{F}_{Kh}(K_1 \amalg K_2))^+\right). \end{aligned}$$

The two maps agree on the first summand: both are the projection hocolim  $((\widetilde{F}_{Kh}(K_1 \coprod K_2))^+) \times \{p_-\} \to \text{hocolim} ((\widetilde{F}_{Kh}(K_1 \coprod K_2))^+)$ . On the second summand, the two maps are compositions of the projection map hocolim  $((\widetilde{F}_{Kh}(K_1 \coprod K_2))^+) \times \{p_+\} \to \text{hocolim} ((\widetilde{F}_{Kh}(K_1 \coprod K_2))^+)$  with the following two maps:



Therefore, after composing with the map hocolim  $((\widetilde{F}_{Kh}(K_1 \coprod K_2))^+) \to \text{hocolim}((\widetilde{F}_{Kh}(K_1 \# K_2))^+)$  from step (F-5) that induces  $\overline{F}_{\times}(f_{0.0}^{\text{op}})$ , we get the two maps



(The first vertical map is the quotient map to hocolim  $((\widetilde{F}_{--Kh}(K_1 \coprod K_2))^+)$ , as the composition sends the rest of hocolim  $((\widetilde{F}_{Kh}(K_1 \coprod K_2))^+)$  to the basepoint.) However, since hocolim  $((\widetilde{F}_{Kh}(K_1 \coprod K_2))^+)$  is the coequalizer (see step (F-5)), these maps agree.

(F-7) The *n* different morphisms  $\underline{\mathbf{n}} \to \underline{\mathbf{1}}$  in  $\overline{\Delta}_{inj}^{\text{op}}$  induce the following two maps  $\overline{F}_{\times}(\underline{\mathbf{n}}) \to \overline{F}_{\times}(\underline{\mathbf{1}})$ : first apply the ring multiplication on the  $\mathbb{H}^1$ -factors to get a map

$$\phi \colon \overline{F}_{\times}(\underline{\mathbf{n}}) = \mathcal{X}_{Kh}(K_1 \coprod K_2) \wedge (\bigwedge_{i=1}^{n-1} \mathbb{H}^1) \to \mathcal{X}_{Kh}(K_1 \coprod K_2) \wedge \mathbb{H}^1 = \overline{F}_{\times}(\underline{\mathbf{2}}),$$

and then compose with the two maps  $\overline{F}_{\times}(f_{\underline{\mathbf{1}},0}^{\mathrm{op}})$  and  $\overline{F}_{\times}(f_{\underline{\mathbf{1}},1}^{\mathrm{op}})$ . We have already defined the map  $\overline{F}_{\times}(\underline{\mathbf{2}}) \to \overline{F}_{\times}(\underline{\mathbf{0}})$ ; define the map  $\overline{F}_{\times}(\underline{\mathbf{n}}) \to \overline{F}_{\times}(\underline{\mathbf{0}})$  to be its composition with  $\phi$ .

The functor  $\overline{F}_{\times}$  thus constructed satisfies the conditions of the lemma (see steps (F-2), (F-4), and (F-5)), thereby concluding the proof.

Remark 9.26. Steps (F-5) and (F-6) in the proof of Lemma 9.25 actually show that with the specific choices made in the proof,  $\mathcal{X}_{Kh}(K_1 \# K_2)$  is the ordinary tensor product of  $\mathcal{X}_{Kh}(K_1)$  and  $\mathcal{X}_{Kh}(K_2)$  over  $\mathbb{H}^1$ . However, these spectra and their module structures are only defined up to weak equivalence (see Proposition 9.14); and the ordinary tensor product is not invariant under such equivalences, while the derived tensor product is.

Proof of Theorem 10. Let  $F_{\otimes} : \Delta_{inj}^{\mathrm{op}} \to \mathscr{CW}$  be the functor from diagram (9.16), and let  $\overline{F}_{\times} : \overline{\Delta}_{inj}^{\mathrm{op}} \to \mathscr{CW}$  be the functor constructed in Lemma 9.25. The derived tensor product  $\mathcal{X}_{Kh}(K_1) \otimes_{\mathbb{H}^1} \mathcal{X}_{Kh}(K_2)$  is defined to be  $\mathrm{hocolim}(F_{\otimes})$ . The natural transformation  $\eta : F_{\otimes} \to F_{\times} = \overline{F}_{\times}|_{\Delta_{inj}^{\mathrm{op}}}$  from Lemma 9.25 (2) is a stable homotopy equivalence on each object; therefore, by property (ho-1) in Section 4.2,  $\mathrm{hocolim}(F_{\otimes}) \simeq \mathrm{hocolim}(F_{\times})$ . Since  $\underline{\mathbf{0}}$  is a terminal object in  $\overline{\Delta}_{inj}^{\mathrm{op}}$ , the functor  $\overline{F}_{\times}$  induces a map

$$\mathcal{X}_{Kh}(K_1) \otimes_{\mathbb{H}^1} \mathcal{X}_{Kh}(K_2) = \operatorname{hocolim}(F_{\times}) \longrightarrow \overline{F}_{\times}(\underline{\mathbf{0}}) = \mathcal{X}_{Kh}(K_1 \# K_2).$$

Using Lemma 9.21 and Lemma 9.25 (2), we can view  $\operatorname{hocolim}(F_{\times})$  as a CW spectrum whose reduced cellular cochain complex is the complex from Formula (9.22). Lemma 9.25 (3) implies that the map  $\operatorname{hocolim}(F_{\times}) \to \overline{F}_{\times}(\underline{\mathbf{0}})$  induces the quasi-isomorphism from Lemma 9.23 at the level of reduced cellular cochain complexes. Therefore, the map is a stable homotopy equivalence.

## 10. Khovanov homotopy type of a mirror

Given a spectrum X, let  $X^{\vee}$  denote the Spanier-Whitehead dual of X, which is the internal function object parametrizing maps  $X \to \mathbb{S}$ . If X is a finite spectrum, the spectrum  $X^{\vee}$  is also finite, and is characterized by the existence of map of spectra

$$\mu \colon X \wedge X^{\vee} \to \mathbb{S}$$

such that the slant product map

(10.1) 
$$\mu^*(\gamma)/: \widetilde{H}_*(X) \to \widetilde{H}^*(X^{\vee})$$

is an isomorphism. (Here,  $\gamma \in \widetilde{H}^0(\mathbb{S})$  is the fundamental class.) We will say that  $\mu$  witnesses S-duality between X and  $X^{\vee}$ . See Switzer [Swi75, Proposition 14.37] and Adams [Ada95, Section III.9] for more details

**Theorem 11.** [LS14a, Conjecture 10.1] Let L be a link and let m(L) denote the mirror of L. Then

$$\mathcal{X}_{Kh}^{j}(m(L)) \simeq \mathcal{X}_{Kh}^{-j}(L)^{\vee}.$$

Before proving the theorem we recall a reformulation of the Spanier-Whitehead duality criterion, after some homological algebra.

**Lemma 10.2.** Let  $C_*$  and  $D_*$  be finitely-generated chain complexes of abelian groups and let  $f: C_* \to D_*$  be a chain map. Suppose that for any field k, the induced map  $f \otimes \operatorname{Id}_k: C_* \otimes k \to D_* \otimes k$  is a quasi-isomorphism. Then f is a quasi-isomorphism.

*Proof.* (Compare [Hat02, 3.A.7]) It suffices to prove that the mapping cone  $\operatorname{Cone}(f)$  of f is acyclic. Observe that  $\operatorname{Cone}(f \otimes \operatorname{Id}_k) = \operatorname{Cone}(f) \otimes k$ , and by assumption  $\operatorname{Cone}(f \otimes \operatorname{Id}_k)$  is acyclic for any field k. It follows from the universal coefficient theorem that if  $E_* \otimes \mathbb{F}_p$  is acyclic for all primes p then  $E_*$  is acyclic.

**Lemma 10.3.** Let  $C_*$  and  $D_*$  be finitely-generated chain complexes of free abelian groups and  $F: C_* \otimes D_* \to \mathbb{Z}$  a chain map. Write  $D^* = \text{Hom}(D_*, \mathbb{Z})$  for the dual complex to  $D_*$ . Then the following conditions are equivalent:

- (1) The map  $C_* \to D^*$  induced by F is a quasi-isomorphism.
- (2) For any field k, the map  $H_*((C_* \otimes k)) \otimes H_*((D_* \otimes k)) \to k$  induced by F is a perfect pairing.

*Proof.* (1)  $\Longrightarrow$  (2) Tensoring with k, the map  $f: C_* \otimes k \to D^* \otimes k = \operatorname{Hom}(D_*, k)$  induced by F is a quasi-isomorphism. Suppose  $\alpha$  is a non-trivial element of  $H_*(C_* \otimes k)$ . Then  $f_*(\alpha) \neq 0 \in H_*(D^* \otimes k) = \operatorname{Hom}(H_*(D_* \otimes k), k)$ . Let  $\beta \in H_*(D_* \otimes k)$  be an element on which  $f_*(\alpha)$  evaluates non-trivially. Then  $F_*(\alpha \otimes \beta) = f_*(\alpha)(\beta) \neq 0$ .

Similarly, if  $\beta \neq 0 \in H_*(D_* \otimes k)$  then there is an element  $b \in H_*(D^* \otimes k)$  such that  $b(\beta) \neq 0$ . Since f is a quasi-isomorphism there is an  $\alpha \in H_*(C_* \otimes k)$  so that  $b = f(\alpha)$ . Then  $F_*(\alpha \otimes \beta) = f_*(\alpha)(\beta) = b(\beta) \neq 0$ .

(2)  $\Longrightarrow$  (1) We start by checking that for any field k, the map  $f: C_* \otimes k \to D^* \otimes k$  is a quasi-isomorphism. To this end, suppose  $\alpha \neq 0 \in H_*(C_* \otimes k)$ . Then there exists  $\beta \in H_*(D_* \otimes k)$  so that  $F_*(\alpha \otimes \beta) \neq 0 \in k$ . Then  $f_*(\alpha)(\beta) = F_*(\alpha \otimes \beta) \neq 0$ , so  $f_*(\alpha) \neq 0$ . Thus,  $f_*$  is injective. Next, given any element  $b \in H_*(D^* \otimes k) = \text{Hom}(H_*(D_* \otimes k), k)$ , since  $F_*$  is a perfect pairing there is an element  $\alpha \in H_*(C_* \otimes k)$  so that  $b(\cdot) = F_*(\alpha \otimes \cdot)$ . Since  $F_*(\alpha \otimes \cdot) = f_*(\alpha)(\cdot)$ , we get  $b = f_*(\alpha)$ . So,  $f_*$  is surjective.

Thus, the map  $f \otimes \operatorname{Id}_k : C_* \otimes k \to D^* \otimes k$  is a quasi-isomorphism for any field k. So, by Lemma 10.2, the map  $f : C_* \to D^*$  induced by F is a quasi-isomorphism, as desired.

**Proposition 10.4.** Let X and Y be finite CW complexes and let  $\mu: X \wedge Y \to S^n$  be a continuous map. Then  $\mu$  witnesses S-duality between X and  $\Sigma^{-n}Y$  if and only if for every field k and integer i, the induced map

$$\widetilde{H}_i(X;k) \otimes \widetilde{H}_{n-i}(Y;k) \to \widetilde{H}_n(S^n;k) = k$$

is a perfect pairing.

*Proof.* Let  $C_*^{cell}(X)$  denote the reduced cellular chain complex of X and  $C_{cell}^*(X)$  the reduced cellular cochain complex. The slant product map of Formula (10.1) is the map on homology induced by

$$C^{cell}_*(X) \to C^{n-*}_{cell}(Y) = \operatorname{Hom}(C^{cell}_{n-*}(Y), \mathbb{Z})$$
  
 $x \mapsto (y \mapsto \gamma(\mu_{\#}(x \wedge y))).$ 

In other words, this is the map induced by the pairing

$$C_*^{cell}(X) \otimes C_{n-*}^{cell}(Y) \to C_n^{cell}(S^n) = \mathbb{Z}$$
$$x \otimes y \mapsto \gamma(\mu_\#(x \wedge y)).$$

So, the result is immediate from Lemma 10.3.

The other ingredients in the proof of Theorem 11 are some facts about functoriality of Khovanov homology and the Khovanov stable homotopy type.

Suppose that  $F \subset [0,1] \times \mathbb{R}^3$  is a link cobordism from  $L_0$  to  $L_1$ . Associated to F is a map  $\Phi_F \colon Kh(L_0) \to Kh(L_1)$ , well-defined up to multiplication by  $\pm 1$ ; see [Jac04, Kho06a, BN05]. The following properties of  $\Phi$  are immediate from the definition:

- (1) If F and F' are isotopic link cobordisms (relative boundary) then  $\Phi_F = \pm \Phi_{F'}$ .
- (2) If  $F = [0,1] \times L$  is the identity cobordism from L to L then  $\Phi_F = \pm \operatorname{Id}: Kh(L) \to Kh(L)$ .
- (3) If  $F_1$  is a cobordism from  $L_0$  to  $L_1$  and  $F_2$  is a cobordism from  $L_1$  to  $L_2$ , and  $F_2 \circ F_1$  denotes the composition of  $F_2$  and  $F_1$ , then

$$\Phi_{F_2 \circ F_1} = \pm \Phi_{F_2} \circ \Phi_{F_1}.$$

(4) If F is a cobordism from  $L_0$  to  $L_1$  and F' is a cobordism from  $L'_0$  to  $L'_1$ , and  $F \coprod F'$  denotes the disjoint union of F and F', which is a cobordism from  $L_0 \coprod L'_0$  to  $L_1 \coprod L'_1$ , then for any field k,

$$\Phi_{F \coprod F'} = \pm \Phi_F \otimes \Phi_{F'} \colon Kh(L_0; k) \otimes Kh(L'_0; k) = Kh(L_0 \coprod L'_0; k)$$
$$\to Kh(L_1 \coprod L'_1; k) = Kh(L_1; k) \otimes Kh(L'_1; k).$$

Given a link L there is a canonical, genus 0 cobordism F from  $L \coprod m(L)$  to the empty link.

**Proposition 10.5.** For any field k, the map

$$\Phi_F \colon Kh(L;k) \otimes Kh(m(L);k) = Kh(L \coprod m(L);k) \to Kh(\varnothing;k) = k$$

associated to the canonical cobordism from  $L \coprod m(L)$  to the empty link is a perfect pairing.

*Proof.* This follows from Properties (1), (2), (3) and (4) above via the usual snake-straightening argument in topological field theory (see, for instance, [Qui95, Lecture 7]).

As noted above, the cohomology groups of  $\mathcal{X}_{Kh}(L)$  are the Khovanov homology of L. Since we are viewing Khovanov homology as covariant in the cobordism, it is more convenient to work with the homology groups of  $\mathcal{X}_{Kh}(L)$ . These can be understood as follows:

**Lemma 10.6.** Let L be a link diagram. Then the cellular chain complex for  $\mathcal{X}_{Kh}(L)$  agrees with the Khovanov complex for m(L). In particular,  $\widetilde{H}_i(\mathcal{X}_{Kh}^j(L)) = Kh^{-i,-j}(m(L))$ .

Proof. This is immediate from the definitions. To wit,  $\mathcal{X}_{Kh}(L)$  can be constructed as a CW complex whose reduced cellular cochain complex  $C^*_{cell}(\mathcal{X}_{Kh}(L))$  is isomorphic to the Khovanov complex  $\mathcal{C}_{Kh}(L)$  from Section 2.2. Therefore, the reduced cellular chain complex  $C^{cell}_*(\mathcal{X}_{Kh}(L))$  is isomorphic to the dual complex  $\operatorname{Hom}(\mathcal{C}_{Kh}(L), \mathbb{Z})$ . However, the dual complex is isomorphic to  $\mathcal{C}_{Kh}(m(L))$  [Kho00, Proposition 32]. In the language of Section 2.2, this isomorphism takes a Khovanov generator in  $F_L(v)$  to a Khovanov generator in  $F_{m(L)}(\vec{1}-v)$  by changing the labels on the circles of  $\mathcal{P}_L(v) = \mathcal{P}_{m(L)}(\vec{1}-v)$  from  $x_+$  to  $x_-$  and vice versa. The gradings work out, and we get  $\widetilde{H}_i(\mathcal{X}_{Kh}^j(L)) = Kh^{-i,-j}(m(L))$ .

Functoriality for the Khovanov spectrum has not yet been verified, but in [LS14b] we did associate maps to elementary cobordisms:

**Proposition 10.7.** Let  $L_1$  and  $L_2$  be links in  $\mathbb{R}^3$  and F a cobordism from  $L_1$  to  $L_2$ . Then there is a map of spectra

$$\Psi_{m(F)} \colon \mathcal{X}_{Kh}(m(L_1)) \to \mathcal{X}_{Kh}(m(L_2))$$

so that the induced map  $\Psi_{m(F),*}$ :  $\widetilde{H}_*(\mathcal{X}_{Kh}(m(L_1))) = Kh(L_1) \to Kh(L_2) = \widetilde{H}_*(\mathcal{X}_{Kh}(m(L_2)))$  agrees with the cobordism map  $\Phi_F$  up to sign.

*Proof.* The corresponding statement in cohomology is immediate from [LS14b, Proposition 3.4, Lemma 3.6 and Lemma 3.7]; the homology statement comes from dualizing [LS14b, Proposition 3.4, Lemma 3.6 and Lemma 3.7] (cf. Lemma 10.6).  $\Box$ 

With these ingredients, we are now ready to verify the mirror theorem.

*Proof of Theorem 11.* The following argument was suggested to us by the referee for [LS14c].

By Proposition 10.4, it suffices to construct a map  $\mathcal{X}_{Kh}(m(L)) \wedge \mathcal{X}_{Kh}(L) \to \mathbb{S}$  so that for any field k, the induced map

$$\widetilde{H}_i(\mathcal{X}_{Kh}(m(L));k)\otimes\widetilde{H}_{-i}(\mathcal{X}_{Kh}(L);k)\to\widetilde{H}_0(\mathbb{S})=k$$

is a perfect pairing. By Proposition 10.5, applying Proposition 10.7 to the canonical cobordism from  $L \coprod m(L)$  to the empty link gives such a map.

Remark 10.8. Theorem 11 (perhaps) gives an obstruction to knots being amphichiral: if K is amphichiral then  $\mathcal{X}_{Kh}^{j}(K)$  is Spanier-Whitehead dual to  $\mathcal{X}_{Kh}^{-j}(K)$ . Using KnotKit [Seeb] (and the technique in [LS14c]), it is possible to verify that this obstruction does not give any additional restrictions on amphichirality for prime knots up to 15 crossings, beyond those implied by Khovanov homology itself. Indeed, the only Khovanov homology-symmetric knots with 15 or fewer crossings for which  $\operatorname{Sq}^2$  is non-vanishing are  $14_{8440}^n$ ,  $14_{9732}^n$ ,  $14_{21794}^n$ ,  $14_{22073}^n$ , and  $15_{139717}^n$ . In each of these cases, the non-Moore space summands of  $\mathcal{X}_{Kh}$  are copies of (various suspensions and desuspensions of)  $\mathbb{R}P^5/\mathbb{R}P^2$  and  $\mathbb{R}P^4/\mathbb{R}P^1$ , and these summands (with appropriate grading shifts) are exchanged by  $(i,j) \to (-i,-j)$ .

It remains open whether the stable homotopy type gives nontrivial restrictions for larger prime knots, although we expect that it does. As a proof of concept, look at [Seea, Table 1] and consider the link  $L = 14_{17196}^n$  II  $m(14_{17546}^n)$ . Its Khovanov homology is symmetric since Khovanov homologies of  $14_{11196}^n$  and  $14_{17546}^n$  are isomorphic (as bigraded groups); the homology of either knot is supported on three adjacent diagonals, the only torsion present is 2-torsion, and the lowest and highest quantum gradings where 2-torsion appears are -11 and 5. However,  $14_{17546}^n$  has no  $\mathrm{Sq}^2$ , but  $14_{11196}^n$  has a single  $\mathrm{Sq}^2$  in quantum grading -1. Using Cartan formula and Theorem 1, the lowest and highest quantum gradings where Kh(L) has non-trivial

 $\operatorname{Sq}^3$  are -6 and 10, and therefore, its Khovanov homotopy type is not symmetric. (Of course this is not our recommended proof for showing L is achiral!)

## 11. Applications

In this section we give an application of the Künneth theorem for the Khovanov stable homotopy type to knot concordance. We begin with some background, continuing from Section 2.2. Recall that Rasmussen [Ras10] used the Lee deformation of the Khovanov complex [Lee05] (the specialization (h,t) = (0,1) of Definition 2.1) to define a concordance invariant  $s_K^{\mathbb{Q}} \in 2\mathbb{Z}$ . As the notation suggests, to define  $s^{\mathbb{Q}}$ , Rasmussen used Khovanov homology with coefficients in  $\mathbb{Q}$ , though any field of characteristic different from 2 would work as well. Bar-Natan [BN05] gave an analogue  $\mathcal{C}_{BN}$  of the Lee deformation that also works over  $\mathbb{F}_2$  (see also [Nao06, Tur06]). The Bar-Natan complex  $\mathcal{C}_{BN}$  is the specialization (h,t)=(1,0) of the universal complex  $\mathcal{C}$  of Definition 2.1. Let  $s_K = s_K^{\mathbb{F}_2}$  denote the corresponding Rasmussen-type invariant; see also [LS14b, Section 2.2].

Since the formal variable h has quantum grading -2, the differential in the Bar-Natan chain complex either preserves the quantum grading or increases it. Let  $\mathcal{F}_q\mathcal{C}_{BN}$  denote the subcomplex of  $\mathcal{C}_{BN}$  supported in quantum grading q or higher. This defines a filtration on the Bar-Natan complex, and the associated graded object  $\bigoplus_{q} \mathcal{F}_{q} \mathcal{C}_{BN} / \mathcal{F}_{q+2} \mathcal{C}_{BN}$  is the Khovanov chain complex  $\mathcal{C}_{Kh}$ .

The second Steenrod square  $\operatorname{Sq}^2$  for the Khovanov spectrum  $\mathcal{X}_{Kh}(K)$  gives a map

$$\operatorname{Sq}^2 \colon Kh^{i,j}(K; \mathbb{F}_2) \to Kh^{i+2,j}(K; \mathbb{F}_2)$$

for each i, j. Using this, in [LS14b] we constructed a refined s-invariant  $s_+^{\mathrm{Sq}^2}(K)$ , as follows:

**Definition 11.1.** [LS14b, Definition 1.2 and Lemma 4.2] Consider configurations of the following form:

$$(11.2) \qquad \langle \widetilde{a}, \widetilde{b} \rangle \longrightarrow \langle \widehat{a}, \widehat{b} \rangle \longleftarrow \langle a, b \rangle \longrightarrow \langle \overline{a}, \overline{b} \rangle$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad$$

Define

$$s_{+}^{\mathrm{Sq}^2}(K) = \begin{cases} s_K & \text{if there does not exist such a configuration,} \\ s_K + 2 & \text{otherwise.} \end{cases}$$

**Theorem 12.** [LS14b, Theorem 1] The integer  $s_+^{Sq^2}(K)$  is a concordance invariant, and

$$2g_4(K) \ge |s_+^{\operatorname{Sq}^2}(K)|$$

where  $q_4(K)$  is the 4-ball genus of K.

**Lemma 11.3.** Let K and L be two knots such that the following holds:

- (1)  $s_{\perp}^{\text{Sq}^2}(K) = s_K + 2.$
- (2)  $Kh^{0,s_L+1}(L; \mathbb{F}_2) \xrightarrow{\operatorname{Sq}^2} Kh^{2,s_L+1}(L; \mathbb{F}_2)$  is the zero map.
- (3) Either  $Kh^{0,s_L+1}(L;\mathbb{F}_2) \xrightarrow{\operatorname{Sq}^1} Kh^{1,s_L+1}(L;\mathbb{F}_2)$  is the zero map or  $Kh^{-2,s_K-1}(K;\mathbb{F}_2) \xrightarrow{\operatorname{Sq}^1} Kh^{-1,s_K-1}(K;\mathbb{F}_2)$  is the zero map.

Then  $s_{+}^{\text{Sq}^2}(K \# L) = s_{K \# L} + 2 = s_K + s_L + 2$ .




FIGURE 11.1. **Oriented connected sum.** Pieces of two knots and their connected sum, oriented coherently with the obvious saddle cobodism, are shown.

*Proof.* Consider the saddle cobordism from  $K \coprod L$  to K # L. Choose some orientations  $o_K$ ,  $o_L$ , and  $o_{K \# L}$ , of the knots K, L, and K # L, which are coherent with respect to the saddle cobordism (see Figure 11.1). Let  $\{\pm o_K\}$ ,  $\{\pm o_L\}$ , and  $\{\pm o_{K \# L}\}$  denote the sets of orientations of K, L, and K # L. For any orientation o there is a corresponding generator g(o) of  $H_0(\mathcal{C}_{BN}; \mathbb{F}_2)$  [Lee05, Theorem 4.2], [Tur, Theorem 1]. We will use the following facts.

- (1)  $C_{BN}(K \coprod L)$  is canonically identified with  $C_{BN}(K) \otimes C_{BN}(L)$ . The induced identification on the associated graded objects,  $Kh(K \coprod L) \cong Kh(K) \otimes Kh(L)$ , is the one induced from the equivalence from Theorem 1. (This is immediate from the definitions.)
- (2) The sets  $\{g(o_K), g(-o_K)\}$ ,  $\{g(o_L), g(-o_L)\}$ ,  $\{g(o_{K\#L}), g(-o_{K\#L})\}$ , and  $\{g(o_K) \otimes g(o_L), g(-o_K) \otimes g(o_L), g(o_K) \otimes g(-o_L), g(-o_K) \otimes g(-o_L)\}$  form bases of  $H_0(\mathcal{C}_{BN}(K))$ ,  $H_0(\mathcal{C}_{BN}(L))$ ,  $H_0(\mathcal{C}_{BN}(K\#L))$ , and  $H_0(\mathcal{C}_{BN}(K \coprod L))$ , respectively [Lee05, Section 4.4.3], [Tur, Theorem 1].
  - (3)  $g(o_L) + g(-o_L)$  has a cycle representative in  $\mathcal{F}_{s_L+1}\mathcal{C}_{BN}(L)$  [LS14b, Proposition 2.6].
- (4) The saddle cobordism map  $C_{BN}(K \coprod L) \to C_{BN}(K \# L)$  preserves the homological grading and either increases the quantum grading or decreases it by exactly one. The induced map on Kh commutes with the map  $\operatorname{Sq}^2$  [LS14b, Theorem 4]. The induced map on  $H_0(\mathcal{C}_{BN}; \mathbb{F}_2)$  is the following:

$$g(o_K) \otimes g(o_L) \mapsto g(o_{K\#L}) \qquad \qquad g(-o_K) \otimes g(o_L) \mapsto 0$$
  
$$g(o_K) \otimes g(-o_L) \mapsto 0 \qquad \qquad g(-o_K) \otimes g(-o_L) \mapsto g(-o_{K\#L}).$$

(This follows from the same argument as [Ras10, Proposition 4.1], since Turner's change of basis diagonalizes the Bar-Natan Frobenius algebra.

Since  $s_+^{\text{Sq}^2}(K) = s_K + 2$ , there is a configuration as in Formula (11.2). Since  $\{g(o_K), g(-o_K)\}$  also form a basis for  $H_0(\mathcal{C}_{BN}(K); \mathbb{F}_2)$  (Fact (2)), after performing a change of basis if necessary, we may assume that  $\overline{a} = g(o_K)$  and  $\overline{b} = g(-o_K)$ .

Using Fact (3), choose some configuration of the following form

Combining the two configurations and using the identification from Fact (1), we get the configuration

$$\langle \widetilde{a} \otimes \widehat{c}, \widetilde{b} \otimes \widehat{c} \rangle \longrightarrow \langle \widehat{a} \otimes \widehat{c}, \widehat{b} \otimes \widehat{c} \rangle \longleftarrow \langle a \otimes c, b \otimes c \rangle \longrightarrow \langle g(o_K) \otimes (g(o_L) + g(-o_L)), \\ \downarrow \qquad \qquad \downarrow \qquad$$

We should justify the leftmost horizontal arrow; that is, assuming  $\operatorname{Sq}^2(\widetilde{a}) = \widehat{a}$  and  $\operatorname{Sq}^2(\widetilde{b}) = \widehat{b}$ , we need to show that  $\operatorname{Sq}^2(\widetilde{a} \otimes \widehat{c}) = \widehat{a} \otimes \widehat{c}$  and  $\operatorname{Sq}^2(\widetilde{b} \otimes \widehat{c}) = \widehat{b} \otimes \widehat{c}$ . Since the identification  $Kh(K \coprod L) \cong Kh(K) \otimes Kh(L)$  is induced from the identification  $\mathcal{X}_{Kh}(K \coprod L) \cong \mathcal{X}_{Kh}(K) \wedge \mathcal{X}_{Kh}(L)$  of Theorem 1, cf. Fact (1),

$$Sq^{2}(\widetilde{a} \otimes \widehat{c}) = Sq^{2}(\widetilde{a}) \otimes \widehat{c} + Sq^{1}(\widetilde{a}) \otimes Sq^{1}(\widehat{c}) + \widetilde{a} \otimes Sq^{2}(\widehat{c})$$
$$= \widehat{a} \otimes \widehat{c}.$$

The first equality is the Cartan formula; the second uses the lemma's hypotheses. Similarly,  $\operatorname{Sq}^2(\widetilde{b}\otimes\widehat{c}) = \widehat{b}\otimes\widehat{c}$ . Now consider the image of this configuration under the saddle cobordism map. Using Fact (4), we get a configuration

By Fact (2), 
$$\langle g(o_{K\#L}), g(-o_{K\#L}) \rangle = H_0(\mathcal{C}_{BN}(K\#L); \mathbb{F}_2)$$
; therefore,  $s_+^{\text{Sq}^2}(K\#L) = s_{K\#L} + 2$ .

We are almost ready to prove Corollary 1.5. First, we tabulate some invariants of the knots K that appear in its statement.

K	$\sigma(K)$	$s_K$	$\tau(K)$	$s_+^{Sq^2}(K)$	$g_4(K)$	$s_K^{\mathbb{Q}}$	u(K)
942	-2	0	0	2	1	0	1
$10_{136}$	-2	0	0	2	1	0	1
$m(11_{19}^n)$	-4	2	1	4	2	2	2
$m(11_{20}^n)$	-2	0	0	2	1	0	1
$11_{70}^{n}$	-4	2	1	4	2	2	2
$11_{96}^{n}$	-2	0	0	2	1	0	1
Table 11.1							

There is some confusion in the nomenclature of knots regarding mirrors, stemming primarily from the fact that Gauss codes do not distinguish between a knot and its mirror. See Figure 1.1 for our convention. We have followed the convention from the Knot Atlas [BNM<sup>+</sup>], using their planar diagram presentations (which do detect chirality). The knots in Figure 1.1 are produced in Knotilus [FFFR] using Gauss codes from the Knot Atlas; however, for  $11^n_{70}$ , our knot diagram differs from the one produced by Knotilus (as well as the one in the Knot Atlas) by mirroring. Alternatively, one can deduce our convention from the value of the signature  $\sigma(K)$  in Table 11.1 (which, for us, is negative for positive knots).

The values of  $s_K = s_K^{\mathbb{F}_2}$  and  $s_+^{\operatorname{Sq}^2}(K)$  are imported from [LS14c];  $s_K$  can also be computed independently by Knotkit [Seeb]. The values of the four-ball genus  $g_4(K)$  and the unknotting number u(K) are extracted

from Knotinfo [CL]. The absolute values of the signature  $\sigma(K)$  and  $s_K^{\mathbb{Q}}$  are extracted from [BNM<sup>+</sup>] and the absolute value of Ozsváth-Szabó's invariant  $\tau(K)$  come from [BG12]. We deduce the signs of  $\sigma(K)$ ,  $s_K^{\mathbb{Q}}$ , and  $\tau(K)$  by noticing that all the unknotting crossings in Figure 1.1 are positive, and changing a positive crossing to a negative one decreases  $-\sigma(K)/2$ ,  $s^{\mathbb{Q}}/2$ , and  $\tau(K)$  by 0 or 1. We list  $\tau(K)$ ,  $s_K^{\mathbb{Q}}$ , and u(K) purely for the reader's interest.

Finally, we deduce Corollary 1.5 which, to recall, was the following:

Corollary 1.5. Let K be one of the knots  $9_{42}$ ,  $10_{136}$ ,  $m(11_{19}^n)$ ,  $m(11_{20}^n)$ ,  $11_{70}^n$ , or  $11_{96}^n$ . (Here m denotes the mirror.) Let L be a knot which is the closure of a positive braid. Letting  $g_4$  denote the four-ball genus, we have

$$g_4(K \# L) = g_4(K) + g_4(L).$$

*Proof.* Certainly  $g_4(K \# L) \leq g_4(K) + g_4(L)$ , so we only need to show  $g_4(K \# L) \geq g_4(K) + g_4(L)$ .

First consider the knot  $K \in \{9_{42}, 10_{136}, m(11_{19}^n), m(11_{20}^n), 11_{70}^n, 11_{96}^n\}$ . In all cases  $s_+^{\text{Sq}^2}(K) = s_K + 2 = 2g_4(K)$  (see Table 11.1).

Next consider the knot L. Draw L as the closure of a positive braid with n crossings and m strands. We will classify the generators of the Khovanov chain complex  $C_{Kh}(L)$  in quantum grading n+2-m or less.

Let v be some vertex in the cube of resolutions for L. Assume there are  $c_v$  circles in the resolution at v. The smallest quantum grading over v is achieved by the Khovanov generator which labels all the  $c_v$  circles by  $x_-$ . The value of this smallest quantum grading is  $n + |v| - c_v$ .

Since all the crossings of L are positive, the resolution at the zero vertex  $\vec{0}$  is the oriented resolution; therefore  $c_{\vec{0}} = m$ . If u is some vertex of weight one, i.e., |u| = 1, then the resolution at u is obtained from the oriented resolution by a merge; therefore  $c_u = m - 1$ . The resolution at any other vertex v is obtained from some weight one vertex by |v| - 1 merges or splits; therefore  $c_v \le m - 1 + |v| - 1 = m + |v| - 2$ , with equality holding if and only if the resolution at v can be obtained from some weight one resolution by splits only.

It follows that the minimum quantum grading over any vertex  $v \neq \vec{0}$  is at least n-m+2; and over  $\vec{0}$ , the minimum quantum grading is n-m, which is attained only by the Khovanov generator that labels all the m circles by  $x_-$ . Observe that this is enough to compute the values of  $s_L$ ,  $g_4(L)$  and g(L):  $s_L \geq n-m+1$ , and Seifert's algorithm yields a surface of genus (n-m+1)/2; therefore, the inequality

$$n - m + 1 \le s_L \le 2g_4(L) \le 2g(L) \le n - m + 1$$

leads to the equality

$$s_L = 2g_4(L) = 2g(L) = n - m + 1.$$

Next, we compute the Khovanov homology in these quantum gradings. We have  $Kh^{*,q}(L;\mathbb{Z})=0$  for all q< n-m and  $Kh^{*,n-m}(L;\mathbb{Z})$  is  $\mathbb{Z}$  supported in homological grading zero. However, our interest lies in quantum grading n-m+2; we show that  $Kh^{*,n-m+2}(L;\mathbb{Z})$  is  $\mathbb{Z}$  supported in homological grading zero as well.

Number the m strands in the braid diagram from left to right. For  $1 \leq i < m$ , let  $n_i$  be the number of crossings in the diagram between the  $i^{\text{th}}$  and the  $(i+1)^{\text{st}}$  strands, i.e., the number of times  $\sigma_i$  occurs in the braid word. (Then  $n = \sum_i n_i$ , and since L is a knot,  $n_i \geq 1$  for all i.) Number the  $n_i$  crossings from top to bottom. For  $1 \leq i \leq m$ , let  $\mathbf{x}_i = (\vec{0}, x_i)$  be the Khovanov generator where  $x_i$  labels the  $i^{\text{th}}$  circle in the oriented resolution by  $x_+$ , and the rest by  $x_-$ . For  $1 \leq i < m$  and  $\emptyset \neq J \subseteq \{1, \ldots, n_i\}$ , let  $u_{i,J}$  be the weight |J| vertex in the cube of resolutions where the 1-resolution is taken only at the crossings that appear in J between the  $i^{\text{th}}$  and  $(i+1)^{\text{st}}$  strands; and let  $\mathbf{y}_{i,J}$  be the Khovanov generator living over  $u_{i,J}$ 

where all the circles are labeled by  $x_-$ . From the discussion above it is clear that the Khovanov chain group  $C_{Kh}^{*,n+2-m}(L)$  is generated by these generators  $\mathbf{x}_i$  and  $\mathbf{y}_{i,J}$ . The differential is fairly straightforward:

$$\delta \mathbf{x}_{i} = \begin{cases} \sum_{j=1}^{n_{1}} \mathbf{y}_{1,\{j\}} & i = 1\\ \sum_{j=1}^{n_{i}} \mathbf{y}_{i,\{j\}} + \sum_{j=1}^{n_{i-1}} \mathbf{y}_{i-1,\{j\}} & 1 < i < m\\ \sum_{j=1}^{n_{m-1}} \mathbf{y}_{m-1,\{j\}} & i = m \end{cases}$$

$$\delta \mathbf{y}_{i,J} = \sum_{\substack{J' \supset J\\ |J' \setminus J| = 1}} \pm \mathbf{y}_{i,J'}.$$

Here we are using the standard sign assignment on the cube, and the signs are determined by the ordering of the n crossings.

Using the change of basis replacing  $\mathbf{x}_k$  by  $\sum_{i=1}^k (-1)^i \mathbf{x}_i$  for all  $1 \le k \le m$ , it is easy to see that the chain complex  $\mathcal{C}_{Kh}^{*,n+2-m}(L)$  is isomorphic to the following direct sum of cube complexes:

$$\mathcal{C}_{Kh}^{*,n+2-m}(L;\mathbb{Z}) \cong \mathbb{Z} \oplus (\bigoplus_{i=1}^{m-1} (\bigotimes_{j=1}^{n_i} (\mathbb{Z} \to \mathbb{Z}))).$$

Therefore,  $Kh^{*,n+2-m}(L;\mathbb{Z})\cong\mathbb{Z}$ , supported in homological grading zero (and is generated by the cycle  $\sum_{i=1}^{m}(-1)^{i}\mathbf{x}_{i}$ ).

As an immediate consequence, we have that both the maps

$$\operatorname{Sq}^1: Kh^{0,s_L+1}(L; \mathbb{F}_2) \to Kh^{1,s_L+1}(L; \mathbb{F}_2) = 0$$


and

$$\operatorname{Sq}^2: Kh^{0,s_L+1}(L; \mathbb{F}_2) \to Kh^{2,s_L+1}(L; \mathbb{F}_2) = 0$$

vanish. Therefore, K and L satisfy the conditions of Lemma 11.3, so

$$2g_4(K\#L) \ge s_+^{\text{Sq}^2}(K\#L) = s_K + s_L + 2 = 2g_4(K) + 2g_4(L).$$

## APPENDIX A. FLOW CHART OF SECTIONS



Background material is indicated by a different shade. Superficial dependencies (e.g., for examples or small pieces of notation) have been omitted.

## References

- [Ada95] J. F. Adams, Stable homotopy and generalised homology, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1995, Reprint of the 1974 original. MR 1324104 (96a:55002)
- [AS] Mohammed Abouzaid and Ivan Smith, Khovanov homology from Floer cohomology, arXiv:1504.01230.
- [Bén67] Jean Bénabou, Introduction to bicategories, Reports of the Midwest Category Seminar, Springer, Berlin, 1967, pp. 1–77. MR 0220789 (36 #3841)
- [BG12] John A. Baldwin and William D. Gillam, Computations of Heegaard-Floer knot homology, J. Knot Theory Ramifications 21 (2012), no. 8, 1250075, 65. MR 2925428
- [BK72] A. K. Bousfield and D. M. Kan, Homotopy limits, completions and localizations, Lecture Notes in Mathematics, Vol. 304, Springer-Verlag, Berlin, 1972. MR 0365573 (51 #1825)
- [BN05] Dror Bar-Natan, Khovanov's homology for tangles and cobordisms, Geom. Topol.  $\bf 9$  (2005), 1443–1499. MR 2174270 (2006g:57017)
- [BNM+] Dror Bar-Natan, Scott Morrison, et al., The Knot Atlas, http://katlas.org/.
- [CJS95] R. L. Cohen, J. D. S. Jones, and G. B. Segal, Floer's infinite-dimensional Morse theory and homotopy theory, The Floer memorial volume, Progr. Math., vol. 133, Birkhäuser, Basel, 1995, pp. 297–325. MR 1362832 (96i:55012)
- [CK08] Sabin Cautis and Joel Kamnitzer, Knot homology via derived categories of coherent sheaves. I. The \$1(2)-case, Duke Math. J. 142 (2008), no. 3, 511–588. MR 2411561 (2009i:57025)
- [CL] Jae Choon Cha and Charles Livingston, Knotinfo: Table of knot invariants, http://www.indiana.edu/~knotinfo.
- [Cor82] Jean-Marc Cordier, Sur la notion de diagramme homotopiquement cohérent, Cahiers Topologie Géom. Différentielle
   23 (1982), no. 1, 93-112, Third Colloquium on Categories, Part VI (Amiens, 1980). MR 648798 (83g:18018)
- [DK80] W. G. Dwyer and D. M. Kan, Simplicial localizations of categories, J. Pure Appl. Algebra 17 (1980), no. 3, 267–284. MR 579087 (81h:55018)
- [ELST16] Brent Everitt, Robert Lipshitz, Sucharit Sarkar, and Paul Turner, Khovanov homotopy types and the Dold-Thom functor, Homology Homotopy Appl. 18 (2016), no. 2, 177–181. MR 3547241
- [EM06] A. D. Elmendorf and M. A. Mandell, Rings, modules, and algebras in infinite loop space theory, Adv. Math. 205 (2006), no. 1, 163–228. MR 2254311 (2007g:19001)

- [ET14] Brent Everitt and Paul Turner, The homotopy theory of Khovanov homology, Algebr. Geom. Topol. 14 (2014), no. 5, 2747–2781. MR 3276847
- [FFFR] Ortho Flint, Bruce Fontaine, Ralph Furmaniak, and Stuart Rankin, Knotilus, http://knotilus.math.uwo.ca/.
- [Flo88a] Andreas Floer, An instanton-invariant for 3-manifolds, Comm. Math. Phys. 118 (1988), no. 2, 215–240. MR 956166
- [Flo88b] \_\_\_\_\_, Morse theory for Lagrangian intersections, J. Differential Geom. 28 (1988), no. 3, 513–547. MR 965228
- [Flo88c] \_\_\_\_\_, The unregularized gradient flow of the symplectic action, Comm. Pure Appl. Math. 41 (1988), no. 6, 775–813. MR 948771
- [Gra74] John W. Gray, Formal category theory: adjointness for 2-categories, Lecture Notes in Mathematics, Vol. 391, Springer-Verlag, Berlin-New York, 1974. MR 0371990 (51 #8207)
- [Hat02] Allen Hatcher, Algebraic topology, Cambridge University Press, Cambridge, 2002. MR 1867354 (2002k:55001)
- [HKK16] Po Hu, Daniel Kriz, and Igor Kriz, Field theories, stable homotopy theory and Khovanov homology, Topology Proc. 48 (2016), 327–360.
- [Hom14] Jennifer Hom, personal communication, 2014.
- [HW16] Jennifer Hom and Zhongtao Wu, Four-ball genus bounds and a refinement of the Ozváth-Szabó tau invariant, J. Symplectic Geom. 14 (2016), no. 1, 305–323. MR 3523259
- [Isb69] John R. Isbell, On coherent algebras and strict algebras, J. Algebra 13 (1969), 299–307. MR 0249484 (40 #2729)
- [Jac04] Magnus Jacobsson, An invariant of link cobordisms from Khovanov homology, Algebr. Geom. Topol. 4 (2004), 1211–1251 (electronic). MR 2113903 (2005k:57047)
- [Jän68] Klaus Jänich, On the classification of O(n)-manifolds, Math. Ann. 176 (1968), 53-76. MR 0226674 (37 #2261)
- [Kho00] Mikhail Khovanov, A categorification of the Jones polynomial, Duke Math. J. 101 (2000), no. 3, 359–426. MR 1740682 (2002):57025)
- [Kho03] \_\_\_\_\_\_, Patterns in knot cohomology. I, Experiment. Math. 12 (2003), no. 3, 365–374. MR 2034399 (2004m:57022)
   [Kho06a] \_\_\_\_\_, An invariant of tangle cobordisms, Trans. Amer. Math. Soc. 358 (2006), no. 1, 315–327. MR 2171235 (2006g:57046)
- [Kho06b] \_\_\_\_\_, Link homology and Frobenius extensions, Fund. Math. 190 (2006), 179–190. MR 2232858 (2007i:57010)
- [Krc14] David Krcatovich, personal communication, 2014.
- [Lau00] Gerd Laures, On cobordism of manifolds with corners, Trans. Amer. Math. Soc. 352 (2000), no. 12, 5667–5688 (electronic). MR 1781277 (2001i:55007)
- [Lee05] Eun Soo Lee, An endomorphism of the Khovanov invariant, Adv. Math. 197 (2005), no. 2, 554–586. MR 2173845 (2006g:57024)
- [Lei75] R. D. Leitch, The homotopy commutative cube, J. London Math. Soc. (2) 9 (1974/75), 23–29. MR 0365560 (51 #1812)
- [Lin15] Jianfeng Lin, Pin(2)-equivariant KO-theory and intersection forms of spin 4-manifolds, Algebr. Geom. Topol. 15 (2015), no. 2, 863–902. MR 3342679
- [LM18] Tye Lidman and Ciprian Manolescu, The equivalence of two Seiberg-Witten Floer homologies, Astérisque (2018), no. 399, vii+220. MR 3818611
- [LS14a] Robert Lipshitz and Sucharit Sarkar, A Khovanov stable homotopy type, J. Amer. Math. Soc. 27 (2014), no. 4, 983–1042. MR 3230817
- [LS14b] \_\_\_\_\_, A refinement of Rasmussen's s-invariant, Duke Math. J. 163 (2014), no. 5, 923–952. MR 3189434
- [LS14c] \_\_\_\_\_, A Steenrod square on Khovanov homology, J. Topol. 7 (2014), no. 3, 817–848. MR 3252965
- [Lur09] Jacob Lurie, Higher topos theory, Annals of Mathematics Studies, vol. 170, Princeton University Press, Princeton, NJ, 2009. MR 2522659 (2010j:18001)
- [Man03] Ciprian Manolescu, Seiberg-Witten-Floer stable homotopy type of three-manifolds with  $b_1 = 0$ , Geom. Topol. 7 (2003), 889–932 (electronic). MR 2026550 (2005b:57060)
- [Man14] \_\_\_\_\_, On the intersection forms of spin four-manifolds with boundary, Math. Ann. **359** (2014), no. 3-4, 695–728. MR 3231012
- [MMSS01] M. A. Mandell, J. P. May, S. Schwede, and B. Shipley, Model categories of diagram spectra, Proc. London Math. Soc. (3) 82 (2001), no. 2, 441–512. MR 1806878 (2001k:55025)
- [Nao06] Gad Naot, The universal Khovanov link homology theory, Algebr. Geom. Topol. 6 (2006), 1863–1892 (electronic). MR 2263052 (2007k:57022)
- [ORS] Peter Ozsváth, Jacob Rasmussen, and Zoltán Szabó, Odd Khovanov homology, arXiv:0710.4300.
- [Ovc08] Sergei Ovchinnikov, Geometric representations of weak orders, Uncertainty and Intelligent Information Systems (Bernadette Bouchon-Meunier, Christophe Marsala, Maria Rifqi, and Ronald R. Yager, eds.), World Scientific Publishing Co., Inc., 2008, pp. 447–456.

[Qui95] Frank Quinn, Lectures on axiomatic topological quantum field theory, Geometry and quantum field theory (Park City, UT, 1991), IAS/Park City Math. Ser., vol. 1, Amer. Math. Soc., Providence, RI, 1995, pp. 323–453. MR 1338394 (96e:57021)

[Ras03] Jacob Andrew Rasmussen, Floer homology and knot complements, ProQuest LLC, Ann Arbor, MI, 2003, Thesis (Ph.D.)—Harvard University. MR 2704683

[Ras10] Jacob Rasmussen, Khovanov homology and the slice genus, Invent. Math. 182 (2010), no. 2, 419–447. MR 2729272 (2011k:57020)

[Seea] Cotton Seed, Computations of the Lipshitz-Sarkar Steenrod square on Khovanov homology, arXiv:1210.1882.

[Seeb] Cotton Seed, Knotkit, https://github.com/cseed/knotkit.

 $[Seg74] \qquad \text{Graeme Segal, $Categories and cohomology theories}, \\ \text{Topology 13 (1974), 293-312. MR 0353298 (50 \#5782)}$ 

[Shu11] Alexander N. Shumakovitch, Patterns in odd Khovanov homology, J. Knot Theory Ramifications 20 (2011), no. 1, 203–222. MR 2777025

[SS06] Paul Seidel and Ivan Smith, A link invariant from the symplectic geometry of nilpotent slices, Duke Math. J. 134 (2006), no. 3, 453–514. MR 2254624

[Swi75] Robert M. Switzer, Algebraic topology—homotopy and homology, Springer-Verlag, New York, 1975, Die Grundlehren der mathematischen Wissenschaften, Band 212. MR 0385836 (52 #6695)

[Tur] Paul Turner, Khovanov homology and diagonalisable Frobenius algebras, arXiv:1803.10079.

[Tur06] Paul R. Turner, Calculating Bar-Natan's characteristic two Khovanov homology, J. Knot Theory Ramifications 15 (2006), no. 10, 1335–1356. MR 2286127 (2008b:57014)

[Vog73] Rainer M. Vogt, Homotopy limits and colimits, Math. Z. 134 (1973), 11–52. MR 0331376 (48 #9709)

[Wig16] Yuval Wigderson, The Bar-Natan theory splits, J. Knot Theory Ramifications 25 (2016), no. 4, 1650014, 19. MR 3482492

[Zie95] Günter M. Ziegler, Lectures on polytopes, Graduate Texts in Mathematics, vol. 152, Springer-Verlag, New York, 1995. MR 1311028 (96a:52011)

E-mail address: tlawson@math.umn.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455

 $E\text{-}mail\ address{:}\ \mathtt{lipshitz@uoregon.edu}$ 

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403

 $E ext{-}mail\ address: sucharit@math.ucla.edu}$ 

Department of Mathematics, University of California, Los Angeles, CA 90095