Difference of forbidden pairs containing a claw

Guantao Chen¹[∗] Michitaka Furuya²† Songling Shan¹ Shoichi Tsuchiya^{3‡} Ping Yang¹

¹Dept. of Math $\&$ Stat, Georgia State University, Atlanta

 2 Dept. of Mathematical Information Science, Tokyo University of Science, Tokyo 3 School of Network and Information, Senshu University, Kanagawa

May 7, 2022

Abstract

When we study forbidden subgraph conditions guaranteeing graphs to have some properties, a claw (or $K_{1,3}$) frequently appears as one of forbidden subgraphs. Recently, Furuya and Tsuchiya compared two classes generated by different forbidden pairs containing a claw, and characterized one of such classes. In this paper, we give such characterization for three new classes. Furthermore, we give applications of our characterizations to some forbidden subgraph problems.

Key words and phrases. forbidden subgraph, Hamiltonian cycle, Halin graph. AMS 2010 Mathematics Subject Classification. 05C75.

1 Introduction

Let \mathcal{G}_1 and \mathcal{G}_2 be two families of graphs, and let P be a certain property for graphs. We assume that every member of \mathcal{G}_2 satisfies P, and consider the problem whether members of \mathcal{G}_1 satisfy P or not. If we suppose $\mathcal{G}_1 \subseteq \mathcal{G}_2$, then every member of \mathcal{G}_1 satisfies P. Now, we suppose a weaker condition than $\mathcal{G}_1 \subseteq \mathcal{G}_2$.

[∗]This research partially supported by an NSA grant

[†]Email address: michitaka.furuya@gmail.com, This research was partially supported by JSPS KAKENHI Grant number 26800086

[‡]Email address: s.tsuchiya@isc.senshu-u.ac.jp

We first suppose that the family $\mathcal{G}_1 - \mathcal{G}_2$ is finite. Then every member of \mathcal{G}_1 satisfies P with finite exceptions. Since we can check whether finite members of \mathcal{G}_1 satisfy P or not in finite time, we can regard the desired problem as solved.

We next suppose that the members of $\mathcal{G}_1 - \mathcal{G}_2$ is characterized (not necessary finite). Then each member of \mathcal{G}_1 either satisfies P or is characterized. If the characterization has a simple structure, then we may be able to check whether such graphs satisfy P or not. Thus, in this case, it might be possible to solve the desired problem.

In this paper, we try to apply the above strategy for the forbidden subgraph problems.

1.1 Definition and preliminary

For a family $\mathcal F$ of connected graphs, a graph G is said to be $\mathcal F$ -free if G contains no member of $\mathcal F$ as an induced subgraph. We also say that the members of $\mathcal F$ are forbidden subgraphs. If G is $\{F\}$ -free, then G is simply said to be F-free. A family $\mathcal F$ of forbidden subgraphs is called a *forbidden pair* if $|\mathcal F|=2$.

Let $K_{1,3}$ denote the star with three leaves. Let K_n and P_n denote the complete graph and the path of order n, respectively. For nonnegative integers k, l and m , let $N_{k,l,m}$ be a graph obtained from K_3 and vertex disjoint three paths P_{k+1}, P_{l+1} , P_{m+1} by identifying one end-vertex of the paths and distinct three vertices of the K_3 . Commonly, $N_{k,0,0}$ (resp., $N_{k,l,0}$) is usually denoted by Z_k (resp., $B_{k,l}$), and $N_{1,1,1}$ is usually denoted by N (see Figure [1\)](#page-1-0).

Figure 1: Graphs Z_n , $B_{1,n}$ and N.

As we mentioned, our main aim is to characterize connected \mathcal{F}_1 -free but not \mathcal{F}_2 free graphs for two families \mathcal{F}_1 and \mathcal{F}_2 of forbidden subgraphs. Such study derives from [\[9\]](#page-18-0). In [9], Olariu considered the case where $\mathcal{F}_1 = \{Z_1\}$ and $\mathcal{F}_2 = \{K_3\}$, and showed that every connected Z_1 -free but not K_3 -free graph is a complete multipartite graph with at least three partite sets. The result is useful when we investigate the class of Z_1 -free graphs (for example, the characterization was used for research of perfect Z_1 -free graphs in [\[9\]](#page-18-0)). Recently, Furuya and Tsuchiya [\[7\]](#page-17-0) focused on forbidden

pairs for the existence of a Hamiltonian cycle, and studied characterization similar to Olariu's result. A graph H is a *generalized comb* if H is obtained as follows (see Figure [2\)](#page-2-0): Let $m \geq 3$ be an integer. Let L_i $(1 \leq i \leq m)$ and C be vertex-disjoint non-empty cliques with $|C| \geq m$, and let R_i $(1 \leq i \leq m)$ be disjoint non-empty subcliques of C. We define the graph H on $\left(\bigcup_{1\leq i\leq m} L_i\right)\cup C$ such that every vertex in L_i is joined to all vertices in R_i for each i $(1 \leq i \leq m)$. In this context, L_i is called a leaf-clique and R_i is called the root of L_i . The following theorem was proved in [\[7\]](#page-17-0).

Figure 2: Generalized comb

Theorem A (Furuya and Tsuchiya [\[7\]](#page-17-0)) Let G be a connected $\{K_{1,3}, B_{1,2}\}$ -free *graph. Then* G *is not* N*-free if and only if* G *is a generalized comb.*

In other words, they solved a characterization problem for $\mathcal{F}_1 = \{K_{1,3}, B_{1,2}\}$ and $\mathcal{F}_2 = \{K_{1,3}, N\}.$

Our notation and terminology are standard, and mostly taken from [\[5\]](#page-17-1). In particular, we shall use the following terminology. Let G be a graph. For $v \in V(G)$, we let $N_G(v)$ denote the *neighborhood* of v in G. For a set U, we let $G[U]$ denote the subgraph of G induced by $U \cap V(G)$.

1.2 Main results

In this paper, we investigate graphs generated by different families of forbidden subgraphs, and characterize the following classes:

- (F1) connected ${K_{1,3}, Z_2}$ -free but not $B_{1,1}$ -free graphs,
- $(F2)$ connected ${K_{1,3}, B_{1,1}}$ -free but not P_5 -free graphs, and
- (F3) connected $\{K_{1,3}, B_{1,2}\}$ -free but not P_6 -free graphs.

We first give a characterization of graphs as in (F1). A generalized comb is pointed if all of its leaf-cliques consist of exactly one vertex. Let \mathcal{H}_0 be the family of

Figure 3: Graphs H_i

pointed generalized combs. For each i ($1 \le i \le 8$), let H_i be the graph depicted in Figure [3.](#page-3-0) For each integer i ($1 \le i \le 5$), the vertices of H_i enclosed with a circle are called *expandable vertices*. Also, for an expandable vertex a of H_i , expanding of a to a clique C is the operation replacing a to C and adding additional edges between $u \in V(H_i) - \{a\}$ and C if $au \in E(H_i)$. Let U_i be the set of expandable vertices of H_i . For a family $\mathcal{C} = \{C_a \mid a \in U_i\}$ of vertex-disjoint cliques indexed by a, the graph $H_i(\mathcal{C})$ is obtained from H_i by expanding each vertex $a \in U_i$ to the clique C_a (see Figure [4\)](#page-3-1). Let

Figure 4: Expanding vertices to cliques

 $\mathcal{H}_i = \{H_i(\mathcal{C}) \mid \mathcal{C} = \{C_a \mid a \in U_i\}$ is a family of vertex-disjoint cliques indexed by $a\}.$ Note that $H_i \in \mathcal{H}_i$. For each j $(6 \leq j \leq 8)$, let $\mathcal{H}_j = \{H_j\}$.

Theorem 1.1 Let G be a connected $\{K_{1,3}, Z_2\}$ -free graph. Then G is not $B_{1,1}$ -free *if and only if* $G \in \bigcup_{0 \leq i \leq 8} \mathcal{H}_i$ *.*

We next consider giving a characterization of graphs as depicted in (F2) and (F3). Let $l \geq 5$ be an integer, and let L_0, L_1, \ldots, L_l be vertex-disjoint cliques. The graph $F_p = F_p(L_1, \ldots, L_l)$ is obtained from $L_1 \cup \cdots \cup L_l$ by joining every vertex of L_i to all vertices of L_{i+1} for $1 \leq i \leq l-1$, and we call F_p a fat l-path (or simply a *fat path*). In this context, L_i ($1 \leq i \leq l$) are called *fundamental cliques* of F_p . The graph $F_c = F_c(L_0, \ldots, L_l)$ is obtained from $L_0 \cup \cdots \cup L_l$ by joining every vertex of L_i to all vertices of L_{i+1} for $0 \leq i \leq l$ where the indices are calculated modulo $l+1$, and we call F_c a fat l-cycle (or simply a fat cycle). In this context, L_i ($0 \le i \le l$) are called *fundamental cliques* of F_c . Note that fat *l*-paths have *l* fundamental cliques but fat *l*-cycles have $l + 1$ fundamental cliques. Let $\mathcal{P}(l)$ be the family of fat *i*-paths and fat *i*-cycles for all $i > l$.

We give the following characterization.

Theorem 1.2 Let G be a connected $\{K_{1,3}, B_{1,1}\}$ -free graph. Then G is not P_5 -free *if and only if* $G \in \mathcal{P}(5)$ *.*

Theorem 1.3 Let G be a connected $\{K_{1,3}, B_{1,2}\}$ -free graph. Then G is not P_6 -free *if and only if* $G \in \mathcal{P}(6)$ *.*

We prove the following generalization of Theorems [1.2](#page-4-0) and [1.3.](#page-4-1)

Theorem 1.4 Let $m \geq 1$ be an integer, and let G be a connected $\{K_{1,3}, B_{1,m}\}$ -free *graph.* Then *G* is not $P_{\text{max}\{3m,m+4\}}$ -free if and only if $G \in \mathcal{P}(\max\{3m,m+4\})$.

Remark 1 There are infinitely many connected $\{K_{1,3}, B_{1,m}\}$ -free but not $P_{\text{max}\{3m-1,m+3\}}$ *free graphs which are neither fat paths nor fat cycles: Fix an integer* $m \geq 1$ *.* Let $F = F_p(L_1, \ldots, L_{\text{max}3m-1,m+3})$ be a fat path, and let K be a clique with $V(F) \cap K = \emptyset$. Let F' be the graph obtained from $F \cup K$ by joining each vertex of K *to each vertex of* $L_{\max\{m-1,1\}} \cup L_{\max\{m,2\}} \cup L_{\max\{2m,3\}} \cup L_{\max\{2m+1,4\}}$ *(see Figure [5\)](#page-5-0). Then we see that* F' is a connected $\{K_{1,3}, B_{1,m}\}$ *-free but not* $P_{\text{max}\{3m-1,m+3\}}$ *-free graph. Therefore the order of the path in Theorem [1.4](#page-4-2) is best possible if we require the targets to be graphs with a simple structure.*

We prove Theorems [1.1](#page-3-2) and [1.4](#page-4-2) in Subsections [2.1](#page-7-0) and [2.2,](#page-14-0) respectively.

Figure 5: Graph F'

1.3 Applications

Duffus, Jacobson and Gould [\[6\]](#page-17-2) proved that every 2-connected $\{K_{1,3}, N\}$ -free graph has a Hamiltonian cycle. Since we can verify that every 2-connected generalized comb has a Hamiltonian cycle, this together with Theorem [A](#page-2-1) implies that every 2-connected $\{K_{1,3}, B_{1,2}\}$ -free graph has a Hamiltonian cycle. In other words, Duffus-Jacobson-Gould theorem and Theorem [A](#page-2-1) provide an alternative proof of Theorem [D](#page-5-1) in Subsection [1.3.1.](#page-5-2) Our main results have similar applications.

For our argument, we introduce a notation related to forbidden subgraphs. For two families \mathcal{H}_1 and \mathcal{H}_2 of forbidden subgraphs, we write $\mathcal{H}_1 \leq \mathcal{H}_2$ if for every $H_2 \in \mathcal{H}_2$, there exists $H_1 \in \mathcal{H}_1$ such that H_1 is an induced subgraph of H_2 . Note that if $\mathcal{H}_1 \leq \mathcal{H}_2$, then every \mathcal{H}_1 -free graph is also \mathcal{H}_2 -free.

1.3.1 Hamiltonian cycles

In the study of forbidden subgraphs, it is a fundamental problem to characterize the forbidden pairs assuring some properties P . When we consider such problems, we often assume a trivial necessary condition of P (for example, when we consider the existence of a Hamiltonian cycle, it is natural to assume the 2-connectedness). Bedrossian [\[1\]](#page-17-3) characterized the forbidden pairs for the existence of a Hamiltonian cycle as follows:

Theorem B (Bedrossian [\[1\]](#page-17-3)) *Let* H *be a forbidden pair. Then every* 2*-connected* H -free graph has a Hamiltonian cycle if and only if either $H \leq \{K_{1,3}, N\}$ or $H \leq$ ${K_{1,3}, B_{1,2}}$ *or* $\mathcal{H} \leq {K_{1,3}, P_6}$ *.*

Bedrossian's characterization depends on, for example, the following theorems.

Theorem C (Broersma and Veldman [\[3\]](#page-17-4)) *Every* 2*-connected* {K1,3, P6}*-free graph has a Hamiltonian cycle.*

Theorem D (Bedrossian [\[1\]](#page-17-3)) *Every* 2-connected $\{K_{1,3}, B_{1,2}\}$ -free graph has a Hamil*tonian cycle.*

Since any 2-connected fat i-paths and any 2-connected fat i-cycles have a Hamiltonian cycle for $i \geq 6$, Theorems [1.3](#page-4-1) and [C](#page-5-3) give an alternative proof of Theorem [D.](#page-5-1)

1.3.2 Halin graphs

A graph is planar if it can be embedded in the plane without edge-crossing, and such an embedded graph is called a *plane graph*. A *Halin graph*, defined by Halin $[8]$, is a plane graph consisting of a tree T without vertices of degree 2 and a cycle C induced by the leaves of T (and we often write a Halin graph H as $H = T \cup C$). If a graph G contains a Halin graph as a spanning subgraph, then it is called a spanning Halin subgraph of G. In [\[4\]](#page-17-5), the following conjecture was proposed.

Conjecture 1 (Chen, Han, O, Shan and Tsuchiya [\[4\]](#page-17-5)) *Let* H *be a forbidden pair. Then every* 3*-connected* H*-free graph has a spanning Halin subgraph if and only if either* $\mathcal{H} \leq \{K_{1,3}, Z_3\}$ *or* $\mathcal{H} \leq \{K_{1,3}, B_{1,2}\}.$

The "only if" part of Conjecture [1](#page-6-0) was already proved in [\[4\]](#page-17-5). Also, as a partial answer for "if" part of the conjecture, the following theorem was proved.

Theorem E (Chen, Han, O, Shan and Tsuchiya [\[4\]](#page-17-5)) *Every* 3-connected $\{K_{1,3}, F_5\}$ *free graph has a spanning Halin subgraph.*

As corollaries of Theorems [1.1,](#page-3-2) [1.2](#page-4-0) and [E,](#page-6-1) we obtain other partial answers for "if" part of Conjecture [1](#page-6-0) (and we give those detail in Appendix).

Theorem 1.5 *Every* 3-connected $\{K_{1,3}, B_{1,1}\}$ -free graph has a spanning Halin sub*graph.*

Theorem 1.6 *Every* 3-connected $\{K_{1,3}, K_{2}\}$ -free graph has a spanning Halin sub*graph.*

1.3.3 Independence numbers

The *independence number* of a graph G is the maximum cardinality of an independent set of G. In $[2]$, Brandstädt and Hammer found a polynomial-time algorithm for determining the independence number of ${K_{1,3}, P_5}$ -free graphs.

Let H be a graph belonging to $\mathcal{P}(5)$, and let Q be an induced path of H having three vertices. Then any maximal induced paths and any maximal induced cycles containing Q pass through each fundamental cliques of H exactly once. By using the fact above, for a given graph G, we can decide whether G belongs to $\mathcal{P}(5)$ or not in polynomial-time (and we omit its precise algorithm). This together with Theorem [1.2](#page-4-0) assures that we can determine the independence number of ${K_{1,3}, B_{1,1}}$ -free graphs in polynomial-time.

2 Proof of main results

In this section, we prove Theorems [1.1](#page-3-2) and [1.4.](#page-4-2)

2.1 Proof of Theorem [1.1](#page-3-2)

Lemma 2.1 Let G be a connected $\{K_{1,3}, K_{2}\}$ -free graph which contains an induced *subgraph* N*. Then* G *is a pointed generalized comb.*

Proof. Since G is Z_2 -free and Z_2 is an induced subgraph of $B_{1,2}$, G is also $B_{1,2}$ -free. This, together with Theorem [A,](#page-2-1) implies that G is a generalized comb. We only show that every leaf-clique consists of a single vertex. Let L_i $(1 \leq i \leq m)$ be the leaf-cliques of G , and let R_i be the root of L_i . On the contrary, we may assume that $|L_1| \geq 2$. Let $a_1, a_2 \in L_1$ with $a_1 \neq a_2, a_3 \in R_1, a_4 \in R_2$ and $a_5 \in L_2$. Then $G[\{a_1, a_2, a_3, a_4, a_5\}] \cong Z_2$, giving a contradiction. Hence G is a pointed generalized comb. \square

In the following lemmas (Lemmas [2.2](#page-7-1)[–2.8\)](#page-13-0), we follow the labels given in Figures [3](#page-3-0) and [4.](#page-3-1)

Lemma 2.2 Let G be a connected $\{K_{1,3}, \mathbb{Z}_2, N\}$ -free graph which contains an induced subgraph $H = H_1({C_{s_3}, C_{s_4}, C_{s_5}})$, where C_{s_3}, C_{s_4} and C_{s_5} are vertex-disjoint *cliques. Then for each vertex* $a \in V(G) - V(H)$ *with* $N_G(a) \cap V(H) \neq \emptyset$, one of the *following holds:*

- (i) $G[V(H) \cup \{a\}] \in \mathcal{H}_1$,
- (ii) *for some* $i \in \{1,2\}$, $N_G(a) \cap V(H) = \{s_i\} \cup C_{s_5}$ *and* $|C_{s_{5-i}}| = 1$ (*and so* $G[V(H) \cup \{a\}] \in \mathcal{H}_2$,

(iii) *for some* $i \in \{3, 4\}$, $N_G(a) \cap V(H) = \{s_1, s_2\} \cup C_{s_i}$ and $|C_{s_7-i}| = |C_{s_5}| = 1$ (and *so* $G[V(H) \cup \{a\}] \in \mathcal{H}_3$ *)*, *or*

 (iv) $N_G(a) \cap V(H) = \{s_1, s_2\} \cup C_{s_3} \cup C_{s_4}$ and $|C_{s_5}| = 1$ (and so $G[V(H) \cup \{a\}] \in \mathcal{H}_4$).

Proof. For each $i \in \{3, 4, 5\}$, we take a vertex b_i as follows: If $N_G(a) \cap C_{s_i} \neq \emptyset$, let $b_i \in N_G(a) \cap C_{s_i}$; otherwise (i.e., $N_G(a) \cap C_{s_i} = \emptyset$), let $b_i \in C_{s_i}$.

Case 1: $N_G(a) \cap C_{s_5} \neq \emptyset$.

If $as_1, as_2 \in E(G)$, then $G[{a, s_1, s_2, b_5}] \cong K_{1,3}$, giving a contradiction. Thus $as_1 \notin E(G)$ or $as_2 \notin E(G)$. We may assume that $as_1 \notin E(G)$.

If $N_G(a) \cap C_{s_3} \neq \emptyset$ and $ab \notin E(G)$ for some $b \in C_{s_4}$, then $G[\{b_3, a, b, s_1\}] \cong K_{1,3}$; if $N_G(a) \cap C_{s_4} \neq \emptyset$ and $ab \notin E(G)$ for some $b \in C_{s_3}$, then $as_2 \in E(G)$ because $G[\{b_4, a, b, s_2\}] \not\cong K_{1,3}$, and hence $G[\{a, s_2, b_4, b, s_1\}] \cong Z_2$. In either case, we get a contradiction. This implies that either $C_{s_3} \cup C_{s_4} \subseteq N_G(a)$ or $N_G(a) \cap (C_{s_3} \cup C_{s_4}) = \emptyset$.

Subcase 1.1: $C_{s_3} \cup C_{s_4} \subseteq N_G(a)$.

If $ab \notin E(G)$ for some $b \in C_{s_5}$, then $G[\{b_3, a, s_1, b\}] \cong K_{1,3}$, giving a contradiction. Thus $C_{s_5} \subseteq N_G(a)$. If $as_2 \in E(G)$, let $C'_{s_i} = C_{s_i}$ $(i \in \{3, 5\})$ and $C'_{s_4} = C_{s_4} \cup \{a\}$; if $as_2 \notin E(G)$, let $C'_{s_i} = C_{s_i}$ $(i \in \{3,4\})$ and $C'_{s_5} = C_{s_5} \cup \{a\}$. Then $G[V(H) \cup \{a\}] =$ $H_1(\{C'_{s_3}, C'_{s_4}, C'_{s_5}\}) \in \mathcal{H}_1.$

Subcase 1.2: $N_G(a) \cap (C_{s_3} \cup C_{s_4}) = \emptyset$.

Since $G[\{a, s_1, s_2, b_5, b_3, b_4\}] \not\cong N$, we have $as_2 \in E(G)$. If $ab \notin E(G)$ for some $b \in C_{s_5}$, then $G[\{b, b_3, b_4, s_2, a\}] \cong Z_2$, giving a contradiction. Thus $C_{s_5} \subseteq N_G(a)$, and hence $N_G(a) \cap V(H) = \{s_2\} \cup C_{s_5}$. If $|C_{s_3}| \geq 2$, then $G[\{b_3, b, b_4, s_2, a\}] \cong Z_2$ where $b \in C_{s_3} - \{b_3\}$, giving a contradiction. Thus $|C_{s_3}| = 1$.

Case 2: $N_G(a) \cap C_{s_5} = \emptyset$ (i.e., $ab_5 \notin E(G)$).

Claim 2.1 *For each* $i \in \{3, 4\}$ *, if* $N_G(a) \cap C_{s_i} \neq \emptyset$ *, then* $N_G(a) \supseteq \{s_1, s_2\} \cup C_{s_i}$ *.*

Proof. We may assume $i = 3$. Since $G[{b_3, a, b_5, s_1}] \not\cong K_{1,3}$, we have $as_1 \in$ $E(G)$. By the same argument, if $N_G(a) \cap C_{s_4} \neq \emptyset$, then $as_2 \in E(G)$. Since $G[\{a, s_1, b_3, b_4, s_2\}] \ncong Z_2$, we have $as_2 \in E(G)$ or $ab_4 \in E(G)$. In either case, we have $as_2 \in E(G)$. If $ab \notin E(G)$ for some $b \in C_{ss}$, then $G[\{b_5, b, b_3, a, s_2\}] \cong Z_2$, giving a contradiction. Thus $C_{s_3} \subseteq N_G(a)$. \Box

Suppose $N_G(a) \cap (C_{s_3} \cup C_{s_4}) = \emptyset$. Since $N_G(a) \cap V(H) \neq \emptyset$, we have $as_i \in E(G)$ for some $i \in \{1,2\}$. Hence $G[\{b_5, b_{5-i}, b_{i+2}, s_i, a\}] \cong Z_2$, giving a contradiction. Thus $N_G(a) \cap (C_{s_3} \cup C_{s_4}) \neq \emptyset$. We may assume that $N_G(a) \cap C_{s_3} \neq \emptyset$. This together with Claim [2.1](#page-8-0) forces $\{s_1, s_2\} \cup C_{s_3} \subseteq N_G(a)$. If $|C_{s_5}| \geq 2$, then $G[\{b_5, b, b_3, a, s_2\}] \cong Z_2$ where $b \in C_{s_5} - \{b_5\}$, giving a contradiction. Thus $|C_{s_5}| = 1$.

If $N_G(a) \cap C_{s_4} \neq \emptyset$, then $C_{s_4} \subseteq N_G(a)$ by Claim [2.1,](#page-8-0) and hence (iv) holds. Thus we may assume that $N_G(a) \cap C_{s_i} = \emptyset$ (i.e., $N_G(a) \cap V(H) = \{s_1, s_2\} \cup C_{s_3}$). If $|C_{s_4}| \geq 2$, then $G[\{b_4, b, s_2, a, s_1\}] \cong Z_2$ in G where $b \in C_{s_4} - \{b_4\}$, giving a contradiction. Hence $|C_{s_4}| = 1$, and so (iii) holds. \square

Lemma 2.3 Let G be a connected $\{K_{1,3}, \mathbb{Z}_2, \mathbb{N}\}\$ -free graph which contains an in*duced subgraph* $H = H_2(\lbrace C_{t_3}, C_{t_4} \rbrace)$, where C_{t_3} and C_{t_4} are vertex-disjoint cliques. *Then for each vertex* $a \in V(G) - V(H)$ *with* $N_G(a) \cap V(H) \neq \emptyset$, one of the following *holds:*

- (i) $G[V(H) \cup \{a\}] \in \mathcal{H}_2$,
- (ii) *for some* $i \in \{3, 4\}$, $N_G(a) \cap V(H) = \{t_1, t_{i+2}\} \cup C_{t_i}$ and $|C_{t_{7-i}}| = 1$ (and so $G[V(H) \cup \{a\}] \in \mathcal{H}_5$, or
- (iii) $N_G(a) \cap V(H) = \{t_1, t_2, t_5, t_6\}$ and $|C_{t_3}| = |C_{t_4}| = 1$ (and so $G[V(H) \cup \{a\}] \in$ \mathfrak{R}_6 .

Proof. For each $i \in \{3, 4\}$, let $b_i \in C_{t_i}$. For each $i \in \{5, 6\}$, we note that the graph $B_i := H - t_i$ belongs to \mathcal{H}_1 .

Case 1: $N_G(a) \cap (C_{t_3} \cup C_{t_4}) = \emptyset$.

Since $N_G(a) \cap V(H) \neq \emptyset$, $N_G(a) \cap V(B_i) \neq \emptyset$ for some $i \in \{5, 6\}$. We may assume that $N_G(a) \cap V(B_5) \neq \emptyset$. Since $N_G(a) \cap (C_{t_3} \cup C_{t_4}) = \emptyset$, we have $N_G(a) \cap V(B_5) =$ $\{t_1, t_2, t_6\}$ and $|C_{t_4}| = 1$ by Lemma [2.2.](#page-7-1) In particular, $N_G(a) \cap V(B_6) \neq \emptyset$. Then again by Lemma [2.2,](#page-7-1) $N_G(a) \cap V(B_6) = \{t_1, t_2, t_5\}$ and $|C_{t_3}| = 1$. This implies that $N_G(a) \cap V(H) = \{t_1, t_2, t_5, t_6\}$ and $|C_{t_3}| = |C_{t_4}| = 1$, as desired.

Case 2: $N_G(a) \cap (C_{t_3} \cup C_{t_4}) \neq \emptyset$.

We may assume that $N_G(a) \cap C_{t_3} \neq \emptyset$. If $N_G(a) \cap V(B_5) = \{t_6\} \cup C_{t_3}$, then either $N_G(a) \cap V(B_6) = C_{t_3}$ or $N_G(a) \cap V(B_6) = \{t_5\} \cup C_{t_3}$, which contradicts Lemma [2.2.](#page-7-1) Thus, by Lemma [2.2,](#page-7-1) we have either $G[V(B_5) \cup \{a\}] \in \mathcal{H}_1$, or $N_G(a) \cap V(B_5) =$ $\{t_1\} \cup C_{t_3}$ and $|C_{t_4}| = 1$.

Subcase 2.1: $G[V(B_5) \cup \{a\}] \in \mathcal{H}_1$.

We see that $\{t_2\} \cup C_{t_3} \cup C_{t_4} \subseteq N_G(a)$. Since $G[\{a, t_2, b_4, t_6, t_5\}] \not\cong Z_2$, we have $at_5 \in E(G)$ or $at_6 \in E(G)$. We may assume that $at_5 \in E(G)$. If $at_1 \in E(G)$, then $G[\{a, t_1, b_4, t_5\}] \cong K_{1,3}$, giving a contradiction. Thus $at_1 \notin E(G)$. So, $at_6 \notin E(G)$ because $G[\{t_5, t_6, a, t_2, t_1\}] \not\cong Z_2$. Hence we get $N_G(a) \cap V(H) = \{t_2, t_5\} \cup C_{t_3} \cup C_{t_4}$. Consequently, $G[V(H) \cup \{a\}] = H_2(\{C'_{t_3}, C'_{t_4}\})$ where $C'_{s_3} = C_{s_3} \cup \{a\}$ and $C'_{s_4} = C_{s_4}$, as desired.

Subcase 2.2: $N_G(a) \cap V(B_5) = \{t_1\} \cup C_{t_3}$ and $|C_{t_4}| = 1$.

Since $G[{b_3, a, t_2, t_5}] \not\cong K_{1,3}$, we have $at_5 \in E(G)$. Hence $N_G(a) \cap V(H) =$ $\{t_1, t_5\} \cup C_{t_3}$ and $|C_{t_4}| = 1$. \Box

Lemma 2.4 Let G be a connected $\{K_{1,3}, \mathbb{Z}_2, \mathbb{N}\}\)$ -free graph which contains an induced subgraph $H = H_3({C_{u_6}})$, where C_{u_6} is a clique. Then for each vertex $a \in V(G) - V(H)$ with $N_G(a) \cap V(H) \neq \emptyset$, one of the following holds:

- (i) $G[V(H) \cup \{a\}] \in \mathcal{H}_3$,
- (ii) $N_G(a) \cap V(H) = \{u_1, u_i, u_{7-i}\}$ for some $i \in \{2, 3\}$ and $|C_{u_6}| = 1$ (and so $G[V(H) \cup \{a\}] \in \mathcal{H}_6$, or

(iii) $N_G(a) \cap V(H) = \{u_4, u_5\}$ *(and so G[V(H)* $\cup \{a\}$] $\in \mathcal{H}_5$).

Proof. For each $i \in \{2,3\}$, we note that the graph $B_i := H - u_i$ belongs to \mathcal{H}_1 . Since $N_G(a) \cap V(H) \neq \emptyset$, $N_G(a) \cap V(B_i) \neq \emptyset$ for some $i \in \{2,3\}$. If $au_4, au_5 \notin E(G)$, then $N_G(a) \cap V(B_i) \subseteq \{u_1, u_{u_{5-i}}\} \cup C_{u_6}$ for each $i \in \{2, 3\}$, which contradicts Lemma [2.2.](#page-7-1) Thus, $au_4 \in E(G)$ or $au_5 \in E(G)$. We may assume that $au_4 \in E(G)$. Then by Lemma [2.2,](#page-7-1) we have either $G[V(B_3) \cup \{a\}] \in \mathcal{H}_1$, or $N_G(a) \cap V(B_3) = \{u_1, u_4\}$ and $|C_{u_6}| = 1$, or $N_G(a) \cap V(B_3) = \{u_4, u_5\}.$

Case 1: $G[V(B_3) \cup \{a\}] \in \mathcal{H}_1$.

In this case, we have $\{u_2, u_4\} \cup C_{u_6} \subseteq N_G(a)$. Then again by Lemma [2.2,](#page-7-1) we have either $G[V(B_2) \cup \{a\}] \in \mathcal{H}_1$ or $N_G(a) \cap V(B_2) = \{u_1, u_4\} \cup C_{u_6}$ or $N_G(a) \cap V(B_2)$ $V(B_2) = \{u_1, u_3, u_4\} \cup C_{u_6}$. If $N_G(a) \cap V(B_2) = \{u_1, u_4\} \cup C_{u_6}$ (i.e., $N_G(a) \cap V(H) =$ ${u_1, u_2, u_4\} \cup C_{u_6}$, then $G[{u_2, a, u_1, u_3, u_5}] \cong Z_2$; if $N_G(a) \cap V(B_2) = {u_1, u_3, u_4} \cup$ C_{u_6} (i.e., $N_G(a) \cap V(H) = \{u_1, u_2, u_3, u_4\} \cup C_{u_6}$), then $G[\{u_2, u_4, a, u_3, u_5\}] \cong Z_2$. In either case, we get a contradiction. Thus $G[V(B_2) \cup \{a\}] \in \mathcal{H}_1$. Since $au_4 \in$ $E(G)$, we see that $N_G(a) \cap V(B_2) = \{u_3, u_4, u_5\} \cup C_{u_6}$, and hence $N_G(a) \cap V(H) =$ ${u_2, u_3, u_4, u_5} \cup C_{u_6}$. Consequently, $G[V(H) \cup {a}] = H_3({C'_{u_6}})$ where C'_{u_6} = $C_{u_6} \cup \{a\}.$

Case 2: $N_G(a) \cap V(B_3) = \{u_1, u_4\}$ and $|C_{u_6}| = 1$.

Since $au_2 \notin E(G)$ and $G[\{u_1, u_2, u_3, a\}] \ncong K_{1,3}$, we have $au_3 \in E(G)$. Hence $N_G(a) \cap V(H) = \{u_1, u_3, u_4\}$ and $|C_{u_6}| = 1$.

Case 3: $N_G(a) \cap V(B_3) = \{u_4, u_5\}.$

Since $G[{u_3, u_1, b, a}] \ncong K_{1,3}$ for any $b \in C_{u_6}$, we have $au_3 \notin E(G)$. Hence $N_G(a) \cap V(H) = \{u_4, u_5\}.$

Lemma 2.5 Let G be a connected $\{K_{1,3}, \mathbb{Z}_2, \mathbb{N}\}\$ -free graph which contains an induced subgraph $H = H_4({C_{v_4}, C_{v_5}, C_{v_6}})$, where C_{v_4} , C_{v_5} and C_{v_6} are vertex-disjoint *cliques. Then for each vertex* $a \in V(G) - V(H)$ *with* $N_G(a) \cap V(H) \neq \emptyset$, $G[V(H) \cup$ ${a} \in \mathcal{H}_4$.

Proof. For each $i \in \{4, 5, 6\}$, let $b_i \in C_{v_i}$. For each $i \in \{5, 6\}$, we note that the graph $B_i := H - C_{v_i}$ belongs to \mathcal{H}_1 .

Suppose $N_G(a) \cap \{v_1, v_2, v_3\} = \emptyset$. Since $N_G(a) \cap V(H) \neq \emptyset$, we may assume that $ab_4 \in E(G)$. Then $G[{b_4, a, v_1, v_2}] \cong K_{1,3}$, giving a contradiction. Thus, $N_G(a) \cap \{v_1, v_2, v_3\} \neq \emptyset$. We may assume that $av_1 \in E(G)$. Then, by Lemma [2.2,](#page-7-1) we have $G[V(B_5) \cup \{a\}] \in \mathcal{H}_1$ or $N_G(a) \cap V(B_5) = \{v_1, v_i\}$ for some $i \in \{2, 3\}.$

Suppose that $N_G(a) \cap V(B_5) = \{v_1, v_i\}$ for some $i \in \{2, 3\}$. In this case, we may assume that $N_G(a) \cap V(B_5) = \{v_1, v_2\}$. Then by Lemma [2.2,](#page-7-1) $N_G(a) \cap V(B_6) =$ $\{v_1, v_2\}.$ In particular, $N_G(a) \cap V(H) = \{v_1, v_2\}.$ Then $G[\{b_5, v_3, b_6, v_1, a\}] \cong Z_2$, giving a contradiction. Thus $G[V(H) \cup \{a\}] \in \mathcal{H}_1$.

Hence we have $N_G(a) \cap V(B_5) = \{v_1\} \cup C_{v_4} \cup C_{v_6}$ or $N_G(a) \cap V(B_5) = \{v_1, v_i\} \cup C_{v_4} \cup C_{v_5}$ C_{v_6} for some $i \in \{2,3\}$. If $N_G(a) \cap V(B_5) = \{v_1\} \cup C_{v_4} \cup C_{v_6}$, then $ab_5 \in E(G)$ because $G[\{a, v_1, b_6, b_5, v_2\}] \not\cong Z_2$, and hence $G[\{b_5, a, v_2, v_3\}] \cong K_{1,3}$, giving a contradiction. Thus $N_G(a) \cap V(B_5) = \{v_1, v_i\} \cup C_{v_4} \cup C_{v_6}$ for some $i \in \{2, 3\}$. We may assume that $N_G(a) \cap V(B_5) = \{v_1, v_2\} \cup C_{v_4} \cup C_{v_6}$. Since $\{v_1, v_2\} \cup C_{v_4} \subseteq N_G(a) \cap V(B_6) \subseteq$ $\{v_1, v_2\} \cup C_{v_4} \cup C_{v_5}$, we have $C_{v_5} \subseteq N_G(a)$ by Lemma [2.2.](#page-7-1) In particular, $N_G(a) \cap$ $V(H) = \{v_1, v_2\} \cup C_{v_4} \cup C_{v_5} \cup C_{v_6}$. Therefore $G[V(H) \cup \{a\}] = H_4(\{C'_{v_4}, C'_{v_5}, C'_{v_6}\})$ where $C'_{s_4} = C_{s_4} \cup \{a\}$ and $C'_{s_i} = C_{s_i}$ $(i \in \{5, 6\})$. \Box

Lemma 2.6 Let G be a connected $\{K_{1,3}, \mathbb{Z}_2, N\}$ -free graph which contains an induced subgraph $H = H_5({C_{w_7}})$, where C_{w_7} is a clique. Then for each vertex $a \in V(G) - V(H)$ with $N_G(a) \cap V(H) \neq \emptyset$, one of the following holds:

(i) $G[V(H) \cup \{a\}] \in \mathcal{H}_5$, or

(ii) $N_G(a) \cap V(H) = \{w_1, w_2, w_i, w_{9-i}\}$ *for some* $i \in \{3, 4\}$ *and* $|C_{w_7}| = 1$ (*and so* $G[V(H) \cup \{a\}] \cong H_7$.

Proof. For each $i \in \{1, 2\}$, we note that the graph $B_i := H - w_i$ belongs to \mathcal{H}_3 . Since $N_G(a) \cap V(H) \neq \emptyset$, $N_G(a) \cap V(B_i) \neq \emptyset$ for some $i \in \{1,2\}$. We may assume that $N_G(a) \cap V(B_1) \neq \emptyset$. If $N_G(a) \cap V(B_1) = \{w_3, w_5\}$, then either $N_G(a) \cap V(B_2) =$ $\{w_3, w_5\}$ or $N_G(a) \cap V(B_2) = \{w_1, w_3, w_5\}$, which contradicts Lemma [2.4.](#page-10-0) This, together with Lemma [2.4,](#page-10-0) implies that either $G[V(B_1) \cup \{a\}] \in \mathcal{H}_3$ or $N_G(a) \cap$ $V(B_1) = \{w_2, w_i, w_{9-i}\}\text{ for some }i \in \{3, 4\}\text{ and }|C_{w_7}| = 1.$

Case 1: $G[V(B_1) \cup \{a\}] \in \mathcal{H}_3$.

Note that we have either $N_G(a) \cap V(B_2) = \{w_3, w_4, w_5, w_6\} \cup C_{w_7}$ or $N_G(a) \cap V(B_2)$ $V(B_2) = \{w_1, w_3, w_4, w_5, w_6\} \cup C_{w_7}$. This, together with Lemma [2.4,](#page-10-0) leads to $N_G(a) \cap$ $V(H) = \{w_3, w_4, w_5, w_6\} \cup C_{w_7}$. Hence, $G[V(H) \cup \{a\}] = H_5(\{C'_{w_7}\}) \in \mathcal{H}_5$ where $C'_{w_7} = C_{w_7} \cup \{a\}.$

Case 2: $N_G(a) \cap V(B_1) = \{w_2, w_i, w_{9-i}\}\$ for some $i \in \{3, 4\}$ and $|C_{w_7}| = 1$.

We may assume that $N_G(a) \cap V(B_1) = \{w_2, w_3, w_6\}$. Then $N_G(a) \cap V(B_2) =$ $\{w_3, w_6\}$ or $N_G(a) \cap V(B_2) = \{w_1, w_3, w_6\}$. This, together with Lemma [2.4,](#page-10-0) leads to $N_G(a) \cap V(H) = \{w_1, w_2, w_3, w_6\}.$ \Box

Lemma 2.7 Let G be a connected $\{K_{1,3}, \mathbb{Z}_2, \mathbb{N}\}\$ -free graph which contains an in*duced subgraph* $H = H_6$ *. Then for each vertex* $a \in V(G) - V(H)$ *with* $N_G(a) \cap$ $V(H) \neq \emptyset$, $N_G(a) \cap V(H) = \{x_i, x_{i+1}, x_7\}$ for some $i \in \{1,3\}$. Consequently, $G[V(H) \cup \{a\}] \cong H_7.$

Proof. We note that the graph $B := H - x_1$ belongs to \mathcal{H}_3 , and the graph $B^* :=$ $H - x_5$ belongs to \mathcal{H}_2 .

We first suppose that $ax_i, ax_{i+2} \in E(G)$ for some $i \in \{1,2\}$. We may assume that $ax_1, ax_3 \in E(G)$. Then by Lemma [2.3,](#page-9-0) we have $N_G(a) \cap V(B^*) = \{x_1, x_3, x_6, x_7\},\$ and hence either $N_G(a) \cap V(B) = \{x_3, x_6, x_7\}$ or $N_G(a) \cap V(B) = \{x_3, x_5, x_6, x_7\},$ which contradicts Lemma [2.4.](#page-10-0) Thus,

for each
$$
i \in \{1, 2\}
$$
, either $ax_i \notin E(G)$ or $ax_{i+2} \notin E(G)$. (2.1)

If $N_G(a) \cap V(B^*) \neq \emptyset$, then $|N_G(a) \cap V(B^*)| \geq 2$ by Lemma [2.3.](#page-9-0) In particular, we have $N_G(a) \cap V(B) \neq \emptyset$. If $N_G(a) \cap \{x_1, x_2, x_3, x_4\} = \emptyset$, then $N_G(a) \cap V(B) \subseteq$ ${x_5, x_6, x_7}$, which contradicts Lemma [2.4.](#page-10-0) Thus we may assume that $ax_1 \in E(G)$. By [\(2.1\)](#page-12-0), $ax_3 \notin E(G)$. Since $G[{x_1, a, x_2, x_3}]$ $\ncong K_{1,3}$, we have $ax_2 \in E(G)$. So, $ax_4 \notin E(G)$ by [\(2.1\)](#page-12-0). Then, by Lemma [2.4,](#page-10-0) $N_G(a) \cap V(B) = \{x_2, x_7\}$. Consequently, $N_G(a) \cap V(H) = \{x_1, x_2, x_7\}.$

Lemma 2.8 Let G be a connected $\{K_{1,3}, \mathbb{Z}_2, N\}$ -free graph which contains an in*duced subgraph* $H = H_7$. Then for each vertex $a \in V(G) - V(H)$ with $N_G(a) \cap$ $V(H) \neq \emptyset$, $N_G(a) \cap V(H) = \{y_1, y_2, y_7, y_8\}$. Consequently, $G[V(H) \cup \{a\}] \cong H_8$.

Proof. For each $i \in \{1, 2\}$, we note that the graph $B_i := H - y_i$ is isomorphic to H_6 . Since $N_G(a) \cap V(H) \neq \emptyset$, we have $N_G(a) \cap V(B_i) \neq \emptyset$ for some $i \in \{1,2\}$. We may assume that $N_G(a) \cap V(B_1) \neq \emptyset$. Then, by Lemma [2.7,](#page-12-1) $N_G(a) \cap V(B_1) = \{y_2, y_3, y_5\}$ or $N_G(a) \cap V(B_1) = \{y_2, y_7, y_8\}$. In particular, $\{y_3, y_5\} \subseteq N_G(a) \cap V(B_2)$ or $\{y_7, y_8\} \subseteq$ $N_G(a) \cap V(B_2)$. This, together with Lemma [2.7,](#page-12-1) leads to $N_G(a) \cap V(B_1) = \{y_2, y_7, y_8\}$ and $N_G(a) \cap V(B_2) = \{y_1, y_7, y_8\}.$ So, $N_G(a) = \{y_1, y_2, y_7, y_8\}.$ \Box

Proof of Theorem [1.1.](#page-3-2) By routine but tedious argument, we can verify that every graph in $\bigcup_{0 \leq i \leq 8} \mathcal{H}_i$ is $\{K_{1,3}, Z_2\}$ -free but not $B_{1,1}$ -free (and we omit its detail). Thus it suffices to show that, if a connected $\{K_{1,3}, Z_2\}$ -free graph G is not $B_{1,1}$ -free (i.e., G contains $B_{1,1}$ as an induced subgraph), then G belongs to $\bigcup_{0 \leq i \leq 8} \mathcal{H}_i$.

Assume that G contains $B_{1,1}$ ($\in \mathcal{H}_1$) as an induced subgraph. Then G contains a graph $H \in \bigcup_{0 \leq i \leq 8} \mathcal{H}_i$ as an induced subgraph. Choose H so that $|V(H)|$ is as large as possible. It suffices to show that $G = H$. By way of contradiction, suppose that $G \neq H$ (i.e., $V(G) - V(H) \neq \emptyset$). Then by Lemma [2.1,](#page-7-2) G is N-free. Since G is connected, there exists a vertex $a \in V(G) - V(H)$ which is adjacent to a vertex in $V(H)$. By the maximality of H, $G[V(H) \cup \{a\}] \notin \bigcup_{0 \leq i \leq 8} \mathcal{H}_i$. This, together with Lemmas [2.2](#page-7-1)[–2.8,](#page-13-0) gives $H = H_8$. For each $i \in \{7, 8, 9\}$, we note that the graph $B_i := H - z_i$ is isomorphic to H_7 . Since $N_G(a) \cap V(H) \neq \emptyset$, we have $N_G(a) \cap V(B_i) \neq \emptyset$ for some $i \in \{7, 8\}$. We may assume that $N_G(a) \cap V(B_7) \neq \emptyset$. Then by Lemma [2.7,](#page-12-1) $N_G(a) \cap V(B_7) = \{z_3, z_4, z_8, z_9\}.$ In particular, $az_3 \in E(G)$. On the other hand, since $N_G(a) \cap V(B_9) \neq \emptyset$, $N_G(a) \cap V(B_9) = \{z_1, z_2, z_7, z_8\}$, and so $az_3 \notin E(G)$, giving a contradiction.

This completes the proof of Theorem [1.1.](#page-3-2) \Box

2.2 Proof of Theorem [1.4](#page-4-2)

In order to prove Theorem [1.4,](#page-4-2) we give a further definition. For two integers s and t, we let $[s, t] = \{i \in \mathbb{N} \mid s \le i \le t\}$. Note that if $s > t$, then $[s, t] = \emptyset$.

Here we prove Theorem [1.4.](#page-4-2) We can easily verify that every graph in $\mathcal{P}(\max\{3m, m+1\})$ 4}) is $\{K_{1,3}, B_{1,m}\}$ -free but not $P_{\max\{3m, m+4\}}$ -free. Thus it suffices to show that if a connected ${K_{1,3}, B_{1,m}}$ -free graph G is not $P_{\text{max}{3m,m+4}}$ -free (i.e., G contains $P_{\max\{3m,m+4\}}$ as an induced subgraph), then G belongs to $\mathcal{P}(\max\{3m,m+4\})$.

Assume that G contains $P_{\text{max}{3m,m+4}}$ as an induced subgraph. Then G contains a graph $H \in \mathcal{P}(\max\{3m, m+4\})$ as an induced subgraph. Choose H so that $|V(H)|$ is as large as possible. It suffices to show that $G = H$. Otherwise, there exists a vertex $a \in V(G) - V(H)$ such that $N_G(a) \cap V(H) \neq \emptyset$. Let l be the integer so that H is either a fat l-path or a fat l-cycle. Then we can write either $H =$ $F_p(L_1, \ldots, L_l)$ or $H = F_c(L_0, \ldots, L_l)$ for some vertex-disjoint cliques L_0, \ldots, L_l . Let $I = \{i \mid N_G(a) \cap L_i \neq \emptyset\}.$

Claim 2.2 $|I| < 4$ *.*

Proof. Suppose that there are five fundamental cliques $L^{(1)}, \ldots, L^{(5)}$ of H with $N_G(a) \cap L^{(i)} \neq \emptyset \; (1 \leq i \leq 5)$. For each $i \; (1 \leq i \leq 5)$, let $b^{(i)} \in N_G(a) \cap L^{(i)}$. Since max $\{3m, m+4\} \geq 5$, if H is a fat cycle, then H has at least six fundamental cliques. Thus $G[\{b^{(i)} \mid 1 \leq i \leq 5\}]$ has no cycle, and so is a forest of order five and maximum degree at most two. Then we can easily check that $G[\{b^{(i)} \mid 1 \leq i \leq 5\}]$ has an independent set B with $|B| = 3$, and hence $G[{a} \cup B] \cong K_{1,3}$, giving a contradiction. \Box

If H is a fat cycle, then $N_G(a) \cap L_i = \emptyset$ for some $0 \leq i \leq l$ by Claim [2.2.](#page-14-1) By relabeling L_0, \ldots, L_l if necessary, we may assume that

 $(L1)$ $0 \notin I$, and

(L2) subject to (L1), $|I \cap \{1, l\}|$ is as small as possible.

Thus, if H is a fat cycle and there exists an integer i $(1 \leq i \leq l-2)$ with $i, i+1, i+2 \notin$ I, then $I \cap \{0,1,l\} = \emptyset$.

For each i $(1 \leq i \leq l)$, we take a vertex b_i as follows: If $i \in I$, let $b_i \in N_G(a) \cap L_i$; otherwise (i.e., $i \notin I$), let $b_i \in L_i$. Note that, by our choices of indices, $b_1b_l \notin E(G)$ regardless of H being a fat path or a fat cycle.

Claim 2.3 Assume that there exists an index j ($2 \le j \le l-2$) such that $I \cap [2, l-1] =$ $\{j, j+1\}$. Then either $j = 2$ and $ab_1 \in E(G)$ or $j = l-2$ and $ab_l \in E(G)$.

Proof. Recall that $l \ge \max\{3m, m+4\}$. We first consider the case $l - m - 1 \le j \le k$ m+1. Then $l \leq 2m+2$. Since $l \geq 3m$, we have $m \leq 2$; since $l \geq m+4$, we have $m \geq 2$. Hence $m = 2$, and this forces $l = 6$ and $j = 3$. By the assumption of the claim, $ab_2, ab_5 \notin E(G)$. Since $G[\{b_2, b_3, a, b_4, b_5, b_6\}] \ncong B_{1,2}$ and $G[\{b_5, b_4, a, b_3, b_2, b_1\}] \ncong$ $B_{1,2}$, we have $ab_1, ab_6 \in E(G)$. Then $G[{a, b_1, b_3, b_6}] \cong K_{1,3}$, giving a contradiction. Thus either $j \ge m+2$ or $j \le l-m-2$.

We now consider the case $j \geq m+2$ (i.e., $j-m \geq 2$). Then $ab_i \notin E(G)$ for every $j - m \le i \le j - 1$. Since $G[\{b_{j+2}, b_{j+1}, a, b_j, b_{j-1}, \ldots, b_{j-m}\}] \not\cong B_{1,m}$, this forces $b_{j+2} = b_l$ (i.e., $j = l - 2$) and $ab_l \in E(G)$, as desired. Thus we may assume that $j \leq l-m-2$ (i.e., $j+m+1 \leq l-1$). Then $ab_i \notin E(G)$ for every $j+2 \leq i \leq j+m+1$. Since $G[\{b_{j-1}, b_j, a, b_{j+1}, b_{j+2}, \ldots, b_{j+m+1}\}] \ncong B_{1,m}$, this forces $b_{j-1} = b_1$ (i.e., $j = 2$) and $ab_1 \in E(G)$, as desired. \square

Claim 2.4 For each $j \in I$, there exists an index j' $(j' \neq j)$ such that $|j - j'| = 1$ and $j' \in I$.

Proof. If $2 \le j \le l-1$ and $j-1, j+1 \notin I$, then $G[\{b_j, a, b_{j-1}, b_{j+1}\}] \cong K_{1,3}$, giving a contradiction. Hence if $2 \leq j \leq l-1$, then the desired conclusion holds. Thus we may assume that $j \in \{1, l\}.$

For the moment, we assume that $j = 1$ and $2 \notin I$. We further suppose that there exists an index i ($3 \le i \le l-1$) with $i \in I$. Choose i so that i is as small as possible. Then, $i + 1 \in I$ since $3 \leq i \leq l - 1$. If $i + 1 \leq l - 1$ and $I \cap [3, l - 1] = \{i, i + 1\}$, then $i = l-2$ and $l \in I$ by Claim [2.3.](#page-14-2) This implies that if $i+1 \leq l-1$ (i.e., $i \leq l-2$), then there are three indices i_1, i_2, i_3 $(3 \leq i_1 < i_2 < i_3 \leq l)$ with $i_1, i_2, i_3 \in I$, and hence $G[\{a, b_1, b_{i_1}, b_{i_3}\}] \cong K_{1,3}$, giving a contradiction. Thus $i \geq l-1$, and so $i = l-1$. Note that $I \cap [1, l] = \{1, l-1, l\}$. This, together with the fact $l - m - 1 \geq 3$, implies that $G[\{b_1, a, b_l, b_{l-1}, \ldots, b_{l-m-1}\}] \cong B_{1,m}$, giving a contradiction. Thus $I \cap [2, l-1] = \emptyset$. By the choice of L_i , we see that H is a fat path. If there exists a vertex $u \in L_1$ with au $\notin E(G)$, then, since $l \geq m+4$, $G[{a, b_1, u, b_2, ..., b_{m+2}}] \cong B_{1,m}$, giving a contradiction. Thus $L_1 \subseteq N_G(a)$. By the symmetry and the fact $l-1 \notin I$, if $l \in I$, then $L_l \subseteq N_G(a)$. Hence either $N_G(a) \cap V(H) = L_1$ or $N_G(a) \cap V(H) =$ $L_1 \cup L_l$. If $N_G(a) \cap V(H) = L_1$, then $G[V(H) \cup \{a\}] = F_p(\{a\}, L_1, \ldots, L_l)$; if $N_G(a) \cap V(H) = L_1 \cup L_l$, then $G[V(H) \cup \{a\}] = F_c(\{a\}, L_1, \ldots, L_l)$. In either case, $G[V(H) \cup \{a\}] \in \mathcal{P}(\max\{3m, m+4\})$, which contradicts the maximality of H. Thus if j = 1, then $2 \in I$. By the symmetry, if j = l, then $l-1 \in I$. □

Let $i_1 = \min\{i \mid i \in I\}$ and $i_2 = \max\{i \mid i \in I\}$. By Claim [2.4,](#page-15-0) $i_1 + 1$, $i_2 - 1 \in I$.

Claim 2.5 *If* $i_1 \neq 1$ *, then* $L_{i_1+1} \subseteq N_G(a)$ *. If* $i_2 \neq l$ *, then* $L_{i_2-1} \subseteq N_G(a)$ *.*

Proof. If $i_1 \neq 1$ and $L_{i_1+1} \nsubseteq N_G(a)$, say $u \in L_{i_1+1} - N_G(a)$, then $G[\{b_{i_1}, b_{i_1-1}, a, u\}] \cong$ $K_{1,3}$, giving a contradiction. Thus if $i_1 \neq 1$, then $L_{i_1+1} \subseteq N_G(a)$. By the symmetry, we have $L_{i_2-1} \subseteq N_G(a)$ if $i_2 \neq l$. \Box

Since $|I| \leq 4$, we divide the rest of the proof into three cases according to $|I| \leq 2$, $|I| = 3$, and $|I| = 4$.

Case 1: $|I| < 2$.

By Claim [2.4,](#page-15-0) $I = \{i_1, i_2\} = \{i_1, i_1 + 1\}$. If $|I \cap [2, l - 1]| = 2$, then either $1 \in I$ or $l \in I$ by Claim [2.3,](#page-14-2) and so $|I| \geq 3$, giving a contradiction. Thus $|I \cap [2, l-1]| \leq 1$, which implies either $I = \{1, 2\}$ or $I = \{l - 1, l\}$. We may assume that $I = \{1, 2\}$. By the choice of L_i , H is a fat path. If $L_2 \not\subseteq N_G(a)$, say $u \in L_2 - N_G(a)$, then $G[\{a, b_2, u, b_3, b_4, \ldots, b_{m+3}\}] \cong B_{1,m}$, giving a contradiction. Thus $L_2 \subseteq N_G(a)$. This, together with Claim [2.5,](#page-16-0) leads to $N_G(a) \cap V(H) = L_1 \cup L_2$, and hence $G[V(H) \cup \{a\}] =$ $F_p(L_1 \cup \{a\}, L_2, \ldots, L_l) \in \mathcal{P}(\max\{3m, m+4\}),$ which contradicts the maximality of H.

Case 2: $|I| = 3$.

In this case, $I = \{i_1, i_1 + 1 \ (= i_2 - 1), i_2\} = \{i_1, i_1 + 1, i_1 + 2\}$. By the choice of L_i , H is a fat path. Since either $i_1 \neq 1$ or $i_2 \neq l$, L_{i_1+1} $(= L_{i_2-1}) \subseteq N_G(a)$ by Claim [2.5.](#page-16-0) Suppose that either $L_{i_1} \nsubseteq N_G(a)$ or $L_{i_2} \nsubseteq N_G(a)$. We may assume that $L_{i_1} \nsubseteq N_G(a)$. Let $u \in L_{i_1} - N_G(a)$. If $i_1 \leq l - m - 2$ (i.e., $i_1 + m + 2 \leq l$), then $G[\{u, b_{i_1+1}, a, b_{i_1+2}, \ldots, b_{i_1+m+2}\}] \cong B_{1,m}$; if $i_1 \geq m+2$ (i.e., $i_1 - m - 1 \geq 1$), then $G[\{a, b_{i_1}, u, b_{i_1-1}, \ldots, b_{i_1-m-1}\}] \cong B_{1,m}$. In either case, we get a contradiction. Thus $l - m - 1 \leq i_1 \leq m + 1$. This, together with the assumption $l \geq \max\{3m, m + 4\},$ leads to $m = 2$, $l = 6$ and $i_1 = 3$. Then $G[{b_1, b_2, u, b_3, a, b_5}] \cong B_{1,2}$, giving a contradiction. Thus $L_{i_1} \cup L_{i_2} \subseteq N_G(a)$ (i.e., $N_G(a) \cap V(H) = L_{i_1} \cup L_{i_1+1} \cup L_{i_2}$). Hence $G[V(H) \cup \{a\}] = F_p(L_1, \ldots, L_{i_1}, L_{i_1+1} \cup \{a\}, L_{i_2}, \ldots, L_l)$, which contradicts the maximality of H.

Case 3: $|I| = 4$.

In this case, $i_1 + 1 < i_2 - 1$ and $I = \{i_1, i_1 + 1, i_2 - 1, i_2\}$. Let $J_1 = [1, i_1 - 1]$,

 $J_2 = [i_1 + 2, i_2 - 2]$ and $J_3 = [i_2 + 1, l]$ (where J_i may be empty). If $|J_1| \geq m$, then $i_1 - m \geq 1$, and hence $G[\{b_{i_2}, a, b_{i_1+1}, b_{i_1}, b_{i_1-1}, \ldots, b_{i_1-m}\}] \cong B_{1,m}$; if $|J_2| \geq m$, then $i_1 + m + 1 \leq i_2 - 2$, and hence $G[\{b_{i_2}, a, b_{i_1}, b_{i_1+1}, \ldots, b_{i_1+m+1}\}] \cong B_{1,m}$; if $|J_3| \geq m$, then $i_2 + m \leq l$, and hence $G[\{b_{i_1}, a, b_{i_2-1}, b_{i_2}, b_{i_2+1}, \ldots, b_{i_2+m}\}] \cong B_{1,m}$. In either case, we get a contradiction. Thus $\max\{|J_1|, |J_2|, |J_3|\} \leq m-1$. On the other hand, $|J_1| + |J_2| + |J_3| = |[1, l] - {i_1, i_1 + 1, i_2 - 1, i_2}| = l - 4 \ge \max\{3m - 4, m\}.$ Hence we see that $m \geq 2$ and $|J_i| = |J_{i'}| = m - 1$ for some $i, i' \in \{1, 2, 3\}$ with $i \neq i'$. Without loss of generality, we may assume that $|J_1| = m - 1$ (i.e., $i_1 = m$). If $|J_2| = m - 1$, then $G[\{b_{i_2-2}, b_{i_2-1}, b_{i_2}, a, b_m, b_{m-1}, \ldots, b_1\}] \cong B_{1,m}$; if $|J_3| = m - 1$, then $G[\{b_{i_2+1}, b_{i_2}, b_{i_2-1}, a, b_m, b_{m-1}, \ldots, b_1\}] \cong B_{1,m}$. In either case, we again get a contradiction.

This completes the proof of Theorem [1.4.](#page-4-2)

References

- [1] P. Bedrossian, Forbidden subgraph and minimum degree conditions for hamiltonicity, Ph.D. Thesis, Memphis State University, 1991.
- [2] A. Brandstädt and P.L. Hammer, On the stability number of claw-free P_5 -free and more general graphs, Discrete Appl. Math. 95 (1999), 163–167.
- [3] H. Broersma and H.J. Veldman, Restrictions on induced subgraphs ensuring Hamiltonicity or pancyclicity of $K_{1,3}$ -free graphs, *Contemporary methods in* graph theory, Bibliographisches Inst., Mannheim (1990), 181–194.
- [4] G. Chen, J. Han, S. O, S. Shan and S. Tsuchiya, Finding a spanning Halin subgraph in 3-connected ${K_{1,3}, P_5}$ -free graphs, preprint.
- [5] R. Diestel, "Graph Theory" (4th edition), Graduate Texts in Mathematics 173, Springer (2010).
- [6] D. Duffus, M.S. Jacobson and R.J. Gould, Forbidden subgraphs and the Hamiltonian theme, The theory and applications of graphs, Wiley, New York (1981), 297–316.
- [7] M. Furuya and S. Tsuchiya, Claw-free and $N(2, 1, 0)$ -free graphs are almost net-free, Graphs Combin. to appear.
- [8] R. Halin, Studies on minimally n-connected graphs, Combinatorial Mathematics and its Applications, Academic Press, London (1969), 129–136.
- [9] S. Olariu, Paw-free graph, Inform. Process. Lett. 28 (1988), 53–54.

Appendix

Let G be a graph. A sequence $(C; v; Q_1, \ldots, Q_m; x_1, \ldots, x_m)$ is a fan-cycle system of G if

- (1) C is a cycle of G,
- (2) Q_1, \ldots, Q_m are vertex disjoint paths of order at least two on C,
- (3) v, x_1, \ldots, x_m are distinct vertices with $V(G) V(C) = \{v, x_1, \ldots, x_m\},\$
- (4) $|V(C) \bigcup_{1 \leq i \leq m} V(Q_i)| + m \geq 3,$
- (5) v is adjacent to every vertex in $(V(C) \bigcup_{1 \leq i \leq m} V(Q_i)) \cup \{x_1, \ldots, x_m\}$, and
- (6) for i $(1 \leq i \leq m)$, x_i is adjacent to every vertex of Q_i

(see Figure [6\)](#page-18-2). In [\[4\]](#page-17-5), the following lemma was proved in order to construct a

Figure 6: A fan-cycle system

spanning Halin subgraph.

Lemma 2.9 (Chen, Han, O, Shan and Tsuchiya [\[4\]](#page-17-5)) *If a graph* G *has a fancycle system, then* G *has a spanning Halin subgraph.*

Now we show that all 3-connected graphs in $\bigcup_{0 \leq i \leq 8} \mathcal{H}_i$ (defined in Subsection [1.2\)](#page-2-2) have a spanning Halin subgraph.

Lemma 2.10 *For* $G \in \bigcup_{0 \leq i \leq 8} \mathcal{H}_i$, if G is 3-connected, then G has a spanning Halin *subgraph.*

Proof. Since all graphs in $\bigcup_{i \in \{2,3,5,6\}} \mathcal{H}_i$ are not 3-connected, $G \in \mathcal{H}_i$ for some $i \in \{0, 1, 4, 7, 8\}$. By Lemma [2.9,](#page-18-3) it suffices to show that G has a fan-cycle system.

Case 1: $G \in \mathcal{H}_0$.

Let L_1, \ldots, L_m be the leaf-cliques of G, and let R_i be the root of L_i . For each i $(1 \leq i \leq m)$, let $v_i \in R_i$. Since G is 3-connected, $|R_i - \{v_i\}| \geq 2$ for all i, and hence $G - \{v_i \mid 1 \leq i \leq m\}$ has a Hamiltonian cycle C containing $m - 1$ vertex disjoint paths Q_2, \ldots, Q_m with $V(Q_i) = L_i \cup (R_i - \{v_i\})$ $(2 \leq i \leq m)$. Then $(C; v_1; Q_2, \ldots, Q_m; v_2, \ldots, v_m)$ is a fan-cycle system of G.

Case 2: $G \in \mathcal{H}_1$.

Write $G = H_1({C_{s_3}, C_{s_4}, C_{s_5}})$. For each $i \in \{3, 4\}$, let $a_i \in C_{s_i}$. Since G is 3-connected, $|C_{s_i}| \geq 3$ for $i \in \{3, 4\}$, and hence $G - \{a_3, a_4\}$ has a Hamiltonian cycle C containing a path Q with $V(Q) = (C_{s_4} - \{a_4\}) \cup \{s_2\}$. Then $(C; a_3; Q; a_4)$ is a fan-cycle system of G.

Case 3: $G \in \mathcal{H}_4$.

Write $G = H_1({C_{v_4}, C_{v_5}, C_{v_6}})$. Since G is 3-connected, $|C_{v_i} \cup C_{v_j}| \geq 3$ for $i, j \in \{4, 5, 6\}$ with $i \neq j$. By symmetry, we may assume that $|C_{v_4}| \geq 2$ and $|C_{v_5}| \geq 2$. For each $i \in \{4, 5\}$, let $a_i \in C_{v_i}$. Then $G - \{a_4, a_5\}$ has a Hamiltonian cycle C containing a path Q with $V(Q) = (C_{v_5} - \{a_5\}) \cup \{v_3\}$, and hence G has a fan-cycle system $(C; a_4; Q; a_5)$.

Case 4: $G \in \mathcal{H}_7$.

Let $C = y_2y_4y_7y_8y_6$ be a cycle of G, and $Q_1 = y_4y_7$ and $Q_2 = y_8y_6$ be paths on C. Then $(C; y_1; Q_1, Q_2; y_3, y_5)$ is a fan-cycle system of G.

Case 5: $G \in \mathcal{H}_8$.

Let $C = z_2z_4z_7z_9z_8z_6$ be a cycle of G, and $Q_1 = z_4z_7$ and $Q_2 = z_8z_6$ be paths on C. Then $(C; z_1; Q_1, Q_2; z_3, z_5)$ is a fan-cycle system of G.

This completes the proof of Lemma [2.10.](#page-19-0) \Box

Lemma 2.11 *For* $G \in \mathcal{P}(5)$ *, if* G is 3-connected, then G has a spanning Halin *subgraph.*

Proof. We first suppose that G is a fat path, and write $G = F_p(L_1, \ldots, L_l)$. For each i $(2 \leq i \leq l-1)$, let $a_i \in L_i$. Since G is 3-connected, $|L_i - {a_i}| \geq 2$ for i $(2 \leq$ $i \leq l-1$), and hence $G - \{a_2, \ldots, a_{l-1}\}\$ has a Hamiltonian cycle C such that $C[L_i]$ has exactly two components for every i ($2 \leq i \leq l-1$). We take the spanning tree T of G such that $N_T(a_2) = L_1 \cup (L_2 - \{a_2\}) \cup \{a_3\}, N_T(a_{l-1}) = L_l \cup (L_{l-1} - \{a_{l-1}\}) \cup \{a_{l-2}\}$ and $N_T(a_i) = (L_i - \{a_i\}) \cup \{a_{i-1}, a_{i+1}\}$ $(3 \le i \le l-2)$. Then T has no vertices of degree 2 and $V(G) - \{a_2, \ldots, a_{l-1}\}\$ is the set of leaves of T. Hence $T \cup C$ is a spanning Halin subgraph of G.

We next suppose that G is a fat cycle, and write $G = F_c(L_0, \ldots, L_l)$. Since G is 3-connected, G has at most two fundamental cliques of order one. Furthermore, if G has exactly two fundamental cliques of order one, then such cliques are consecutive. By symmetry, we may assume that $|L_i| \geq 2$ for every $i \ (1 \leq i \leq l-1)$. For each i $(1 \leq i \leq l-1)$, let $a_i \in L_i$. Then $G - \{a_1, \ldots, a_{l-1}\}\$ has a Hamiltonian cycle C such that $C[L_i]$ has exactly one component for every $i \ (0 \leq i \leq l)$. We take a spanning tree T of G such that $N_T(a_1) = L_0 \cup (L_1 - \{a_1\}) \cup \{a_2\}, N_T(a_{l-1}) =$ $L_l \cup (L_{l-1} - \{a_{l-1}\}) \cup \{a_{l-2}\}\$ and $N_T(a_i) = (L_i - \{a_i\}) \cup \{a_{i-1}, a_{i+1}\}\ (2 \leq i \leq l-2).$ Then T has no vertices of degree 2 and $V(G) - \{a_1, \ldots, a_{l-1}\}\$ is the set of leaves of T. Hence $T \cup C$ is a spanning Halin subgraph of G . \Box

Theorems [1.1,](#page-3-2) [E](#page-6-1) and Lemma [2.10](#page-19-0) lead to Theorem [1.5.](#page-6-2) Theorems [1.2,](#page-4-0) [1.5](#page-6-2) and Lemma [2.11](#page-19-1) lead to Theorem [1.6.](#page-6-3)