

# Free energy analysis for a system of interacting particles arranged in Bravais lattice

B.I. Lev, V.B. Tymchyshyn, and A.G. Zagorodny

*Bogolyubov Institute for Theoretical Physics, National Academy of Science, Metrolohichna St. 14-b, Kyiv 03680, Ukraine*

We propose a method of free energy calculation for a system of interacting particles arranged in a Bravais lattice. It will be shown how to treat divergences for infinite unbounded systems with “catastrophic” potentials like Coulomb and compare two of them. Besides, we show that current method may be used not only for essentially classical systems, but for some quantum as well. Two systems are considered: grains in dusty plasma and electrons on the liquid-helium surface. For dusty-plasma parameters of grain’s lattice are calculated by numerical solution of obtained equations. Electrons on the liquid-helium surface are analyzed to get localization distance. Besides, conditions for existence of electron lattice are found.

PACS numbers: 52.27.Lw, 71.10.Pm, 02.30.Mv

## INTRODUCTION

There are many soft-matter systems, such as grains in dusty plasma, colloids in various solvents, surfactant solutions, etc., that exhibit self-organization and rearrangement in crystalline structures under certain conditions. Often their inter-particle potential is long-range and similar to the Coulomb one. Usually, this kind of potentials cause divergences (that’s why they are often called “catastrophic”) and complicates consideration. On the other hand, mentioned systems are very interesting due to their application to the studies of a variety of peculiar phenomena in different fields of science [1–3].

It seems to be the most challenging problem to treat Coulomb-like systems with high concentrations of interacting particles [4]. When concentration increases, one can observe crystallization-like phenomena and transition between different lattice symmetries [5–9].

In current contribution we will treat concrete systems along with general formalism development. They were chosen as manifesting a lot of interesting effects. For example, dusty plasma may serve as perfect media for the experimental investigation of classical fluids and solids along with colloids [9–18]. The second system is “electrons on the liquid-helium surface” [19–22]. It may undergo Wigner crystallization [23–29] and transition between triangular and square lattices [30, 31]. Moreover, homogeneous distribution of electron density is not always stable, and there are critical parameters when spatial structures, especially periodic deformations and multi-electron dimples, are formed [32].

But our original goal is some general description of particles arranged in a Bravais lattice when interparticle potential may be “catastrophic”. It is rather difficult, since traditional methods of statistical mechanics cannot be applied to systems with Coulomb-like interactions because of free-energy divergences. Moreover, partition function can be exactly evaluated only for few model systems of interacting particles in the thermodynamical limit [33–36]. Thus, basing on [31, 37–39], we will use alternative approach that employs Hubbard-Stratonovich representation of the partition function [40]. Namely,

we will expand and significantly supplement results obtained in [31] for two-dimensional systems. New results for three-dimensional case, that is far more complicated than two-dimensional, will be obtained.

To be more rigorous: our goal is to find expression for free energy of particles arranged in a Bravais lattice even if inter-particle potential is “catastrophic”. General equations will be applied to two concrete systems: three-dimensional — grains in dusty plasma, and two-dimensional — electrons on the liquid-helium surface. In first case we’ll minimize this energy to find out which lattice should be observed experimentally. And for electrons localization distance will be found. Besides, this will show that presented method allows easily reduce dimension number from three to two.

Present article is organized as follows. It can be contingently divided into two parts: development of technique and its application. Sections I and II focus on general expression for free energy for any Bravais lattice and any potential. The former shows that not only classical, but lots of quantum systems may be treated this way. Besides, an expression for “entropy part” of free energy is calculated. And in section II interaction energy for particles arranged in a Bravais lattice is calculated.

Both these sections are equipped with examples (IA, IIA–IIC) that show how to use developed formalism. These examples are rather useful, because they are later reused when treating physical systems.

Concrete applications of developed formalism start with section III. Here crystal of dust particles is considered. In IIIB it is shown that hexagonal close packing seems to be the lattice we expect to be seen in experiment. This is in agreement with experiments [5] and computer simulations [6].

System of electrons on the liquid-helium surface is analyzed in IV. In IVA localization distance is calculated and compared to classical result [23]. For low temperatures  $T \rightarrow 0$  they coincide.

For not to be confused with lots and lots of complicated mathematical computations we moved them to appendices VI. Thus article mostly contains some physical explanations and results. But using references to appendices, one may find here rigorous exposition of presented

ideas as well.

## I. STATISTICAL DESCRIPTION OF INTERACTING PARTICLES SYSTEM

First of all let's outline class of systems we will be able to consider. Since everything is clear with classical systems, we will give heed to quantum ones.

In this section we aim some general consideration of inhomogeneous system of interacting particles [31, 38]. Nor system, neither interaction potential will be specified to keep generosity and extendability. As result, we will get free energy in some general form and ready for "applying" translation symmetry (section II). Of course, along this section we will do some assumptions about system under consideration. Together they will form restrictions on quantum systems we can treat with proposed method.

In current model macroscopic states of the system are described by occupation numbers. Besides, we suppose that it's Hamiltonian has form:

$$H = \sum_s \varepsilon_s n_s + \frac{1}{2} \sum_{s,s'} V_{ss'} n_s n_{s'}. \quad (1)$$

Here  $\varepsilon_s$  is the additive part of particle energy (usually it is kinetic energy, but it can be particle's energy in external field as well.),  $s$  indicates particle state,  $V_{ss'}$  is interaction energy between particles in states  $s$  and  $s'$ ,  $n_s$  is the occupation number of state  $s$ . We neglect quantum correlations, so only essentially classical systems are considered.

Partition function for this kind of system will be:

$$Z = \sum_{\{n_s\}} \exp(-\beta H),$$

where summation is performed over all possible states  $\{n_s\}$  of the system.

Now we use some properties of Gaussian integrals over auxiliary fields, i.e. Hubbard-Stratonovich transformation [40, 41]:

$$\begin{aligned} \exp\left(\frac{\nu^2}{2\vartheta} \sum_{s,s'} \omega_{ss'} n_s n_{s'}\right) &= \\ &= \int_{-\infty}^{\infty} D\varphi \exp\left(\nu \sum_s n_s \varphi_s - \frac{\vartheta}{2} \sum_{s,s'} \omega_{ss'}^{-1} \varphi_s \varphi_{s'}\right), \end{aligned}$$

with  $D\varphi = \prod_s d\varphi_s / \sqrt{\det(2\pi\beta\omega_{ss'})}$ . Second-order dependence on occupation numbers is avoided by carrying them to introduced field and partition function can be written:

$$Z = \int_{-\infty}^{\infty} D\varphi \exp\left[\sum_s (i\varphi_s - \beta\varepsilon_s) n_s - \frac{1}{2\beta} \sum_{s,s'} (V_{ss'}^{-1} \varphi_s \varphi_{s'})\right].$$

For canonical ensemble number of particles can be fixed by using Cauchy equation:

$$\frac{1}{2\pi i} \oint \xi^{\sum_s n_s - N - 1} d\xi = 1,$$

that leads to  $N$ -particle partition function:

$$\begin{aligned} Z_N &= \frac{1}{2\pi i} \oint d\xi \int D\varphi \exp\left[-\frac{1}{2\beta} \sum_{s,s'} V_{ss'}^{-1} \varphi_s \varphi_{s'} - \right. \\ &\quad \left. -(N+1) \ln \xi\right] \prod_s \sum_{\{n_s\}} [\xi \exp(i\varphi_s - \beta\varepsilon_s)]^{n_s}. \end{aligned}$$

Performing summation over occupation numbers:

$$\begin{aligned} Z_N &= \frac{1}{2\pi i} \oint d\xi \int D\varphi \exp[-\beta F(\varphi, \xi)], \\ \beta F(\varphi, \xi) &= \frac{1}{2} \sum_{s,s'} V_{ss'}^{-1} \varphi_s \varphi_{s'} + \\ &\quad + \delta \sum_s \ln(1 - \delta \xi e^{-\beta\varepsilon_s + i\varphi_s}) + (N+1) \ln \xi. \end{aligned} \quad (2)$$

Here type of statistics is incorporated into  $\delta$  — it equals to +1 for Bose-Einstein, 0 for Maxwell-Boltzmann and -1 for Fermi-Dirac statistics.

Now we have expression for free energy in auxiliary field representation with  $\xi = \exp(\beta\mu)$  for chemical activity. Equation (2) contains all information about probable states of the system and corresponds to the sequence of equilibrium states with regard to their weights. We are interested in asymptotic value of partition function, but want to avoid using perturbation theory. Thus domain is extended to complex plane and saddle-point method is applied.

Dominant contribution is made by states satisfying extrema condition:

$$\frac{\delta\beta F}{\delta\varphi} = \frac{\delta\beta F}{\delta\xi} = 0.$$

Variation of (2) yields expression for saddle-point states:

$$\frac{1}{\beta} \sum_{s'} V_{ss'}^{-1} \varphi_{s'} - \frac{i\xi e^{-\beta\varepsilon_s + i\varphi_s}}{1 - \delta\xi e^{-\beta\varepsilon_s + i\varphi_s}} = 0; \quad (3a)$$

$$\sum_s \frac{\xi e^{-\beta\varepsilon_s + i\varphi_s}}{1 - \delta\xi e^{-\beta\varepsilon_s + i\varphi_s}} = N + 1. \quad (3b)$$

For a certain state expression

$$f_s = \frac{\xi e^{-\beta\varepsilon_s + i\varphi_s}}{1 - \delta\xi e^{-\beta\varepsilon_s + i\varphi_s}}, \quad (4)$$

from (3b) can be treated as an average occupation number. Now saddle-point states (that may be interpreted as thermodynamically stable distributions) can be obtained.

Equation (3a) contains inverse matrix, that is quite inconvenient for us, because its calculation is a challenging mathematical problem even for simple potentials [38, 42, 43]. This problem can be avoided if we use (4) and perform inverse transformation

$$\begin{aligned}\varphi_s &= i\beta \sum_{s'} V_{ss'} f_{s'}, \\ \frac{1}{2\beta} \sum_{s,s'} V_{ss'}^{-1} \varphi_s \varphi_{s'} &= -\frac{\beta}{2} \sum_{s,s'} V_{ss'} f_s f_{s'}.\end{aligned}$$

Then free energy can be written

$$\begin{aligned}\beta F[f, \xi] &= -\frac{\beta}{2} \sum_{s,s'} V_{ss'} f_s f_{s'} - \delta \sum_s \ln(1 + \delta f_s) \\ &\quad + (N + 1) \ln \xi(f).\end{aligned}\quad (5)$$

In terms of canonical ensemble we can write from (4)

$$\begin{aligned}\ln \xi(f) &= \frac{1}{N} \sum_s f_s [\beta(\varepsilon_s + E_s) + \ln f_s - \ln(1 + \delta f_s)], \\ E_s &= \sum_{s'} V_{ss'} f_{s'},\end{aligned}$$

and substitute last expression into free energy (5) to rewrite its expression without chemical potential.

$$\begin{aligned}F[f] &= \sum_s f_s \varepsilon_s + \frac{1}{2} \sum_{s,s'} V_{ss'} f_s f_{s'} + \\ &\quad + \underbrace{\frac{1}{\beta} \sum_s [f_s \ln f_s - (f_s + \delta) \ln(1 + \delta f_s)]}_{F_{ent}}.\end{aligned}\quad (6)$$

First two terms in (6) are kinetic and potential energy respectively, the last one is a contribution due to entropy  $F_{ent}$  (should be equal to zero if  $T = 0$ ).

Now let's do a quick test of obtained results. If a grand canonical ensemble with fixed chemical potential is considered, we can get from (4)

$$f_s = \frac{1}{e^{\beta(\varepsilon_s - \mu_s)} - \delta},\quad (7)$$

which is generalization of the well-known distribution with a chemical potential

$$\mu_s = \mu - E_s.$$

It is obvious that the ‘‘saddle point’’ and ‘‘mean field’’ approximations are equivalents in this case. If an ideal gas ( $\mu_s \equiv \mu$ ) is considered, we can obtain classical statistical distributions.

Before winding up this section let's review the results. We aimed to find some general expression for free energy in a mean-field approximation. As the one, equation (6) should be pointed out (and in some sense (7) that can be used to obtain particles distribution in grand canonical ensemble). Besides, we have got some restrictions on the class of systems we will be able to consider: special form of Hamiltonian (1) and negligible quantum cross-correlations.

### A. Example: two-dimensional system of localized Fermi particles — electrons on the liquid-helium surface

Before we shift our attention to potential energy of particles arranged in a lattice, let's do some example. With this subsection we aim two things: give some idea on how to use the above formalism and secondly — calculate contribution due to entropy for electrons on the liquid-helium surface (will be used in IV).

Electrons on the liquid-helium surface have two degrees of freedom only [23, 24]. Though we set  $\delta = -1$  (Fermi particles) and  $d = 2$  (system is two-dimensional). Then equation (6) can be written

$$\begin{aligned}\beta F[\mu] &= \int \frac{d^2 \vec{p} d^2 \vec{r}}{(2\pi\hbar)^2} \frac{\beta \vec{p}^2}{2m} \frac{1}{e^{\beta(\vec{p}^2/(2m) - \mu(\vec{r}))} + 1} + \\ &\quad + \frac{\beta}{2} \iint \frac{d^2 \vec{p} d^2 \vec{r} d^2 \vec{p}' d^2 \vec{r}'}{(2\pi\hbar)^2 (2\pi\hbar)^2} \times \\ &\quad \times \frac{V(|\vec{r} - \vec{r}'|)}{(e^{\beta(\vec{p}^2/(2m) - \mu(\vec{r}))} + 1)(e^{\beta(\vec{p}'^2/(2m) - \mu(\vec{r}'))} + 1)} - \\ &\quad - \int \frac{d^2 \vec{p} d^2 \vec{r}}{(2\pi\hbar)^2} \left( \frac{\ln(e^{\beta(\vec{p}^2/(2m) - \mu(\vec{r}))} + 1)}{e^{\beta(\vec{p}^2/(2m) - \mu(\vec{r}))} + 1} + \right. \\ &\quad \left. + \frac{\ln(e^{-\beta(\vec{p}^2/(2m) - \mu(\vec{r}))} + 1)}{e^{-\beta(\vec{p}^2/(2m) - \mu(\vec{r}))} + 1} \right).\end{aligned}\quad (8)$$

Here integration sign  $\int$  means integration over the whole phase space.

Since last expression is quite cumbersome we should simplify it somehow. As first approximation we suppose

$$\varepsilon_s = \varepsilon(p) = p^2/(2m).$$

Actually, in the presence of an external field, dispersion relation should be a bit different. But we know, that system of Fermi particles, we will be interested in, is highly degenerated [23]. So quadratic form for dispersion relation looks quite reasonable.

Using last equation and introducing thermal length

$$\lambda_T = \sqrt{2\pi^2 \hbar^2 \beta / m},\quad (9)$$

we get from (7) following expression

$$\begin{aligned}\rho(\vec{r}) &= \int \frac{d^2 \vec{p}}{(2\pi\hbar)^2} \frac{1}{e^{\beta(\vec{p}^2/(2m) - \mu(\vec{r}))} + 1} \\ &= \frac{\pi}{\lambda_T^2} \ln \left( 1 + e^{\beta\mu(\vec{r})} \right),\end{aligned}\quad (10)$$

which connects chemical potential  $\mu(\vec{r})$  and particles density  $\rho(\vec{r})$ .

Now we return to simplification of (8). Performing integration of (8) over momentum and taking (10) and

(9) into account one may get (see appendix VIA)

$$\begin{aligned}
F[\rho] = & \frac{1}{2} \iint d^2\vec{r} d^2\vec{r}' V(|\vec{r} - \vec{r}'|) \rho(\vec{r}) \rho(\vec{r}') + \\
& + \frac{\pi}{\beta \lambda_T^2} \int d^2\vec{r} \left[ \text{Li}_2 \left( e^{-\pi^{-1} \lambda_T^2 \rho(\vec{r})} \right) - \frac{\pi^2}{6} \right] + \\
& + \frac{\lambda_T^2}{2\pi\beta} \int d^2\vec{r} \rho^2(\vec{r}).
\end{aligned} \quad (11)$$

One may notice that in case of Bose statistics we will lose the Bose-condensation effects due to this integration. But we are dealing with Fermi particles, so this is not the case.

We know that electrons on the liquid-helium surface exist in forms of fluid or Wigner crystal [25–27]. We are interested in case when electrons are strongly localized, i.e. their coordinates are quite determined. But since temperature differs from zero, we expect that particle's position will fluctuate near its equilibrium location. Though we assume following distribution function  $\rho_{\text{sp}}$  for *single particle*

$$\rho_{\text{sp}}(\vec{r}) = \frac{1}{(2\pi s^2)^{d/2}} e^{-r^2/(2s^2)}, \quad (12)$$

where  $s$  is dispersion, or localization distance by physical meaning, and  $d = 2$  because system is two-dimensional<sup>1</sup>.

Equation (12) is normal distribution which seems to be quite reasonable. At least this distribution is valid for ground state of quantum harmonic oscillator which is the simplest approximation for particle fluctuating near its equilibrium location.

Last assumption we do is that ‘‘Gaussian’’ (12) is very ‘‘sharp’’ and every particle lays outside localization radius of any other particle. This means that  $s$  is much smaller than other characteristic distances in this system. So free energy per one particle can be written:

$$\begin{aligned}
F_{\text{sp}} = & \frac{1}{2} \iint d^2\vec{r} d^2\vec{r}' V(|\vec{r} - \vec{r}'|) \rho_{\text{sp}}(\vec{r}) \rho_{\text{sp}}(\vec{r}') + \\
& + \underbrace{\frac{\lambda_T^2}{8\pi^2 s^2 \beta} - \frac{\pi^4 s^2}{6\beta \lambda_T^2}}_{F_{\text{ent}}}.
\end{aligned} \quad (13)$$

Here  $F_{\text{ent}}$  is contribution to free energy by *entropy* and temperature. First summand in  $F_{\text{ent}}$  is obtained from direct integration of  $\rho_{\text{sp}}^2$  (see last term in (11)). To get second summand (penultimate term in (11)) we use ‘‘sharpness’’ of (12) and approximate

$$\text{Li}_2 \left( e^{-\pi^{-1} \lambda_T^2 \rho_{\text{sp}}(\vec{r})} \right) \approx \begin{cases} 0, & r \leq s; \\ \pi^2/6, & r > s. \end{cases} \quad (14)$$

With equations (13) and (11) we finish our consideration. We fulfilled our goal to show an example of using section's I formalism. Moreover, now we have expression for entropy part  $F_{\text{ent}}$  of free energy (13), which we use later in IV.

## II. PARTICLES ARRANGED IN A LATTICE

In section I we found some expressions for free energy of inhomogeneous system of interacting particles (6). In this section we will be interested in second term of (6), i.e. potential energy. More accurately, we aim to find method of potential energy calculation for system of identical (from inter-particle's potential point of view) particles arranged in a Bravais lattice. During this section we will show how to use translation symmetry to get in some cases more convenient expressions for potential energy.

Last remark before we dive in. We aim to find appropriate expressions for both two- and three-dimensional cases. But to avoid reduplication all along the section II only three-dimensional case will be treated as far more complicated. For two-dimensional case we will only provide results and short description where difference comes from.

Let  $\mathbb{V}$  designates one subset of exact cover of  $\mathbb{R}^3$  (figure 1) with similar domains containing strictly one particle (relative position of the particle supposed to be the same in every domain)<sup>2</sup>. Then free energy of the particle localized near Bravais lattice site can be written as (we neglect kinetic energy in (6) since particles are localized and potential energy should be bigger)

$$F_{\text{int}} = \iiint_{\mathbb{V}} \iiint_{\mathbb{R}^3} V(|\vec{r} - \vec{r}'|) \rho(\vec{r}) \rho(\vec{r}') d^3\vec{r}' d^3\vec{r}, \quad (15a)$$

$$F_s = \iiint_{\mathbb{V}} \iiint_{\mathbb{V}} V(|\vec{r} - \vec{r}'|) \rho(\vec{r}) \rho(\vec{r}') d^3\vec{r}' d^3\vec{r}, \quad (15b)$$

$$F_{\text{sp}} = F_{\text{int}} - F_s + F_{\text{ent}}. \quad (15c)$$

Equation (15) needs some explanations. Interaction energy is presented here with  $F_{\text{int}} - F_s$ . This ‘‘splitting’’ is just another way of saying we want to integrate over  $\mathbb{R}^3 \setminus \mathbb{V}$ . If one collects both expressions together and uses  $\iiint_{\mathbb{R}^3 \setminus \mathbb{V}} = \iiint_{\mathbb{R}^3} - \iiint_{\mathbb{V}}$  he will get regular integration over  $\mathbb{R}^3 \setminus \mathbb{V}$  for  $\vec{r}'$ .

In other words, since particle is localized near lattice site, we take some ‘‘part of space’’  $\mathbb{V}$  around it. Then by integration we calculate interaction energy between particle in  $\mathbb{V}$  and all the rest particles in  $\mathbb{R}^3 \setminus \mathbb{V}$ .

One may get a good intuitive idea about integration domain  $\mathbb{R}^3 \setminus \mathbb{V}$  through discrete case. When we calculate

<sup>1</sup> We aim to reuse this and some of subsequent equations for systems with different number of dimensions. In such cases we will write them as functions of  $d$  (dimensionality) even if it is redundant at the point we introduce the equation first time.

<sup>2</sup> For example, Wigner-Seitz cells provide this kind of covering. But they often have too complicated shape and we will use parallelepipeds instead (Figure 1).

Coulomb energy for  $i$ -th particle we write  $\sum_{j \neq i} e_i e_j / r_{ij}$ . Integration over  $\mathbb{R}^3 \setminus \mathbb{V}$  is somewhat analogous to  $j \neq i$  for discrete case.

We will further call  $F_{int}$  (15a) *interaction energy* and  $F_s$  (15b) *self-interaction energy*. This “splitting” has one more convenience. Basing on physical background of considered potential we may want or not want to compensate self-interaction. Obviously, Coulomb potential should have self-interaction compensated — potential energy is zero if there is only one particle in a system (considered later in example IIB). On the other hand, self-interaction for effective potentials caused by influence through medium should be uncompensated, because even in absence of other particles medium will influence this one. Surface distortion is a good example. If particle has distorted a surface it has already done some contribution to system’s free energy independently on other particles (considered later in example IIC). Of course, it interacts with distortions created by other particles. But at the same time it interacts with “its own” surface distortion and this may be treated as “self-interaction”.

Now lets consider (15) more carefully. All 3D Bravais lattices can be treated as constructed from parallelepipeds. We will use this property and use parallelepiped as integration region  $\mathbb{V}$ .




Figure 1: One cell from exact cover  $\mathbb{V}$ . Vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are base vectors for Bravais lattice. Angle between  $\vec{a}$  and  $\vec{b}$  is supposed to be  $\alpha$ . Vector  $\vec{c}$  is projected onto  $XY$  plane. Angle between this projection and  $XY$  plane is  $\beta_c$  and between  $\vec{c}$  projection and  $\vec{a}$  is  $\alpha_c$ . One particle is shown in the center.

Next thing we want to do is expressing probability distribution function  $\rho$  for the whole  $\mathbb{R}^3$  through single-particle probability distribution  $\rho_{sp}$ . Basing on the inverse lattice vectors (appendix VIB) we will provide following decomposition

$$\rho(\vec{r}) = \sum_{\vec{k} \in \mathbb{Z}^3} \rho_{\vec{k}} f_{\vec{k}}(\vec{r}), \quad (16a)$$

$$\rho_{\vec{k}} = \bar{\rho} \iiint_{\mathbb{V}} f_{\vec{k}}(\vec{r}) \rho_{sp}(\vec{r}) d^3 \vec{r}, \quad (16b)$$

$$f_{\vec{k}}(\vec{r}) = e^{2\pi i(\vec{k}^T \hat{G} \vec{r})}, \quad (16c)$$

$$\hat{G} = \begin{pmatrix} \frac{1}{a} & -\frac{\cot(\alpha)}{a} & -\frac{\cot(\beta_c) \sin(\alpha - \alpha_c)}{a \sin(\alpha)} \\ 0 & \frac{\csc(\alpha)}{b} & -\frac{\cot(\beta_c) \sin(\alpha_c)}{b \sin(\alpha)} \\ 0 & 0 & \frac{\csc(\beta_c)}{c} \end{pmatrix}. \quad (16d)$$

It may be noticed that all vectors are treated as column vectors.  $\bar{\rho}$  is mean particle density here. To get better intuitive understanding how to compute (16) one may check example IIA for  $\rho_{sp}$  from (12).

It may be shown that presented series are actually Fourier series for  $\rho_{sp}$  and  $\rho$  is periodic along vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  (appendix VIB). So we may treat  $\rho(\vec{r})$  as function on  $\mathbb{R}^3$  “composed” from single particle distributions “arranged in a lattice”. This principle is demonstrated by Fig. 2. Since  $\rho$  has the same symmetry as the lattice and




Figure 2: Two-dimensional analogy of transition from single-particle density probability function  $\rho_{sp}(\vec{r})$  to many particle  $\rho(\vec{r})$ . Function  $\rho(\vec{r})$  is defined for all  $\mathbb{R}^2$  with two-dimensional analogy of (16).

locally is a good approximation for  $\rho_{sp}$  it will be treated as probability distribution function for the whole lattice.

Matrix  $\hat{G}$  comprises all *geometry* of the lattice under consideration. Performing minimization of  $F_{sp}$  over entries of  $\hat{G}$ , one may find which lattice can be formed under certain conditions. We will use later this type of analysis for dust crystal in section III.

If we use equations (16) we can express (15a) in terms of  $\rho_{sp}$  (appendix VID)

$$F_{int} = \frac{1}{\bar{\rho}^2} \sum_{\vec{k} \in \mathbb{Z}^3} |\rho_{\vec{k}}|^2 V_{\vec{k}}, \quad (17a)$$

$$V_{\vec{k}} = \bar{\rho} \iiint_{\mathbb{R}^3} f_{\vec{k}}(\vec{r}') V(|\vec{r}'|) d^3 \vec{r}', \quad (17b)$$

where  $\bar{\rho}$  is mean particle density.

Before moving further, one more thing should be emphasized. This equations are useful because they allow us to deal with “catastrophic” potentials like Coulomb. One may notice that for any descending potential only

$$\frac{1}{\bar{\rho}^2} |\rho_0|^2 V_0 = 4\pi \bar{\rho} \int_0^{\infty} V(r) r^2 dr,$$

can be diverging and only if potential is “catastrophic”.

But this term does not contain “lattice geometry”, it depends on mean particle density only. This means, we can compare two lattices with equal mean particle density even if inter-particle potential is “catastrophic”.

With equations (17), (16) and (15) we achieve our goal and finish consideration of interaction energy for particles arranged in a lattice. Following examples show how to use obtained equations for screened Coulomb (example IIB) and capillary interaction (example IIC). They both use  $\rho$  obtained with  $\rho_{\text{sp}}$  from (12) (example IIA).

### A. Example: calculating distribution function $\rho$ for Gaussian single-particle distribution $\rho_{\text{sp}}$

We suppose that all particles are arranged in a lattice and their positions are quite determined. But since temperature differs from zero we can expect that particle position fluctuates near its lattice site. Using considerations from IA we suppose it to be (12) with  $d = 3$ .

Introduced constant  $s$  means dispersion of presented distribution. From physical point of view it can be treated as localization distance. Later, in section IV, we’ll find its value and compare it with localization distance obtained in different works.

Using that  $\rho_{\text{sp}}(r)$  is a very “sharp” Gaussian, we expand integration limits to  $\mathbb{R}^3$  in (16b) and perform integration (appendix VIC)

$$\rho_{\vec{k}} = \bar{\rho} e^{-2\pi^2 s^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}}. \quad (18)$$

As mentioned previously, we are interested in two-dimensional case as well. One may check, that there exists some sort of analogy of (16) for two dimensions [31]. Difference will appear in changing summations and integrations  $\sum_{\mathbb{Z}^3} \rightarrow \sum_{\mathbb{Z}^2}$ ,  $\iiint_{\mathbb{R}^3} \rightarrow \iint_{\mathbb{R}^2}$  and usage of different matrix

$$\hat{G} = \begin{pmatrix} 1/a & -\cot(\alpha)/a \\ 0 & \csc(\alpha)/b \end{pmatrix}. \quad (19)$$

Performing calculations as in appendix VIC with (12) and  $d = 2$ , one will get

$$\rho_{\vec{k}} = \bar{\rho} e^{-2\pi^2 s^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}}, \quad (20)$$

where  $\bar{\rho}$  is two-, not three-dimensional density as in (18).

Equations (18) and (20) are aimed result of this example. Later they will be used in calculations of potential energy for Coulomb (IIB) and capillary (IIC) interactions.

### B. Example: calculating $F_{\text{int}} - F_s$ for screened Coulomb potential

In section III grains in dusty plasma will be considered and in section IV we’ll set our eyes on electrons on the

liquid-helium surface. In both cases we have to deal with Coulomb potential—screened for dust particles and regular for electrons. The latter will be treated as screened Coulomb with screening distance approaching infinity. During this example we aim to find  $F_{\text{int}} - F_s$  for screened Coulomb potential using methods of section II. Consideration of physics of this two systems will be postponed until sections III and IV.

Screened Coulomb potential can be written as

$$V(r) = \frac{q^2 e^{-r/\lambda_D}}{r}, \quad (21)$$

where  $\lambda_D$  is Debye screening distance. It may be shown (appendix VIE) that expression for  $V_{\vec{k}}$  should be

$$V_{\vec{k}} = \frac{4\pi\bar{\rho}q^2}{1/\lambda_D^2 + 4\pi^2\vec{k}^T \hat{G} \hat{G}^T \vec{k}}. \quad (22)$$

At this point we already have  $V_{\vec{k}}$  (22) and  $\rho_{\vec{k}}$  (18). So we can write  $F_{\text{int}}$  from equation (17a)

$$F_{\text{int}} = \sum_{\vec{k} \in \mathbb{Z}^3} e^{-4\pi^2 s^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}} \frac{4\pi\bar{\rho}q^2}{1/\lambda_D^2 + 4\pi^2\vec{k}^T \hat{G} \hat{G}^T \vec{k}}. \quad (23)$$

Compared to straight-forward method of computing, this one has better convergence. For big values of  $\lambda_D$  we will see, that first summands in (23) descend as  $1/|\vec{k}|^2$ . But if we write series containing interaction energy between this particle and any other, first summands will descend as  $1/|\vec{k}|$ .

In section II we have mentioned that there is one compensatory term we need to take into account. This term will exclude Coulomb self-interaction. If we omit technical details (see appendix VIF) explicit form of  $F_s$  can be obtained

$$F_s = \frac{1}{2\sqrt{\pi}s^3} \int_0^\infty e^{-r'^2/(4s^2)} V(r') r'^2 dr', \quad (24)$$

or substituting screened Coulomb potential (21) as  $V$  and approximating for the case when  $s$  is much smaller then screening distance  $\lambda_D$

$$F_s = \frac{q^2}{\sqrt{\pi}s} - \frac{q^2}{\lambda_D}. \quad (25)$$

Expressions we’ve got for  $F_{\text{int}}$  and  $F_s$  are a bit inconvenient. So we use that  $s$  is small compared to distances in the lattice (“sharp peak” approximation, section II) and perform some approximation.

First of all we introduce mean distance calculated from particle’s density

$$l = \frac{1}{\sqrt[3]{\bar{\rho}}}, \quad (26)$$

and split sum (23) into two parts. Further we will suppose lattice under consideration as not too much degenerated. In other words, we will assume that it may be described as some “deformation” of cubic lattice (appendix VI G)

$$F_{int} - F_s = \frac{\sqrt{\pi}q^2}{l} \sum_{\vec{k} \neq 0} \frac{e^{-\vec{k}^T \hat{G}^{-1} T \hat{G}^{-1} \vec{k} / l^2}}{\vec{k}^T \hat{G}^{-1} T \hat{G}^{-1} \vec{k} / l^2} + \frac{q^2}{\pi l} \sum_{\vec{k} \neq 0} \frac{e^{-l^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}}}{l^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}} + 4\pi \bar{\rho} q^2 \lambda_D^2. \quad (27)$$

One may see that last term is equal to energy in case of evenly distributed charge. Other terms are corrections due to charge pointiness and comprise all information on lattice geometry.

It is very useful to notice, that this is the only summand that approaches infinity if  $\lambda_D \rightarrow \infty$ . This means that (27) can be used for different lattice comparison even without screening. We will only need to measure energy starting from  $4\pi \bar{\rho} q^2 \lambda_D^2$  as “ground-level”.

One more thing to notice. Coulomb is a long-range interaction and localization distance  $s$  should not play any significant role in approximation. Here we see that it is really absent. Rather different may be a picture for short-range interactions or effective interactions with uncompensated  $F_s$  (example II C).

Further we will also need two-dimensional version of (27). Nevertheless, we talk about two-dimensional case, physics is essentially three-dimensional, so previous result should be correct with little modifications. We change the part with uniformly distributed charge to  $2\pi \bar{\rho} q^2 \lambda_D$  ( $\bar{\rho}$  would be two-dimensional density here), use  $\hat{G}$  (19) instead of  $\hat{G}$  (16d) and perform summation over  $\mathbb{Z}^2$ , not  $\mathbb{Z}^3$ .

### C. Example: calculating $F_{int}$ for capillary interaction

Last example we will consider is a rough estimation of capillary interaction energy. This results will be used in section IV. More discussion on physics of the phenomenon and references we postpone until IV.

It is known, that electrons deform liquid-helium surface when pressed against with electric field and this causes effective interaction with potential [30]

$$V(r) = -\frac{q^2 E^2}{2\pi\sigma} K_0(r/l_0), \quad (28)$$

where  $qE$  is force caused by electric field,  $\sigma$  — surface tension of liquid helium,  $r$  — distance between particles,  $K_0$  — modified Bessel function, and  $l_0 = \sqrt{\sigma/g\rho_{He}}$  [30] — capillary length that depends on the fluid properties only.

Since we know that capillary interaction is essentially two-dimensional (particles should be localized on

some surface, no three-dimensional analogue) we will use (17) with (19) and (20). Performing calculations (appendix VI H) we get

$$V_{\vec{k}} = -\frac{q^2 E^2}{\sigma} \frac{\bar{\rho}}{1/l_0^2 + 4\pi^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}}. \quad (29)$$

So we can immediately write

$$F_{int} = -\sum_{\vec{k} \in \mathbb{Z}^2} \frac{\bar{\rho} q^2 E^2 e^{-4\pi^2 s^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}}}{\sigma (1/l_0^2 + 4\pi^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k})}. \quad (30)$$

For our purposes (see section IV) a very rough approximation would be enough. So we use the same trick as in appendix VI G, but perform integration instead of summation and dealing with theta functions (see appendix VII)

$$F_{int} = \frac{q^2 E^2}{4\pi\sigma} \ln(s/l_0). \quad (31)$$

One should notice, that  $s \ll l_0$ , so logarithm value is less then zero and thus whole expression (31) is less then zero.

## III. GRAINS IN DUSTY PLASMA, DUST CRYSTAL AND ITS LATTICE

It is known that grains in dusty plasma interact and even show self-organization in form of melting and crystallization of dust crystals.

We aim to apply methods developed in section II to this system. As a first approximation we suppose that grains interact only as charged particles (screened Coulomb potential). Obtained equation (27) (example II B) and its two-dimension analogue allow to calculate energy of any lattice just inserting correct matrix  $\hat{G}$  and performing summation. But much more interesting is finding the lattice with minimal energy when particles density  $\bar{\rho}$  is fixed. Obtained result can be verified experimentally.

Both sums in (27) are highly convergent and this is very convenient for numeric calculations. Thus we will limit summation with  $|\vec{k}| \leq 4$ . This limit was found experimentally: taking more summands does not change calculated minimal lattice and free energy. On the other hand, taking less results in wrong lattice.

Before considering results, one more thing should be mentioned. Searching for lattice with minimal energy does not need calculation of full expression for  $F_{int} - F_s$ . We need to minimize only parts with  $\hat{G}$ . Moreover, they both contain  $q/l$ , which means, that results of minimization do not depend of particle charge.

### A. 2D lattice.

Before moving to a real-life 3D lattice we start with a 2D. This will help us to emphasize some features that




Figure 3: Same lattice in different representations. It has translation symmetry with respect to two vectors  $\vec{a}$  and  $\vec{b}$ . Upper figure shows different choice for second translation vector  $\vec{b}$  and  $\vec{b}'$ . Using reflections one may show that lattice on bottom figure is the same as on the upper one. So  $\vec{b}''$  is possible choice as well. When performing numerical calculations additional restrictions are used to overcome this uncertainty.

are difficult to understand in 3D without 2D analogue.

Minimization was performed with standard function FindMinimum of Wolfram Mathematica v.9. As function to minimize, modified (27) was used. Minimization was performed with regard to angle  $\alpha$  and length of one of translation vectors. Length of the other one  $b$  is calculated from 2D particles density

$$\bar{\rho} = \frac{1}{ab \sin(\alpha)}.$$

Besides, we want to avoid redundant parameter  $l$ . We may get rid of  $l$  by considering  $a = l(1 + \delta a)$  (by definition  $\bar{\rho} = 1/l^2$ ).

Before considering results, one more remark should be done. Mapping between set of all possible parameter values (angle  $\alpha$  and parameter  $\delta a$ ) and all possible Bravais lattices is not a bijection. It is rather a surjection — one lattice may be described with different  $\alpha$  and  $\delta a$ . For minimization algorithms to be not confused, we provide some restrictions.

Figure 3 shows uncertainty when choosing parameters for lattice description. One may see that it can be eliminated if we claim that projection of  $\vec{b}$  on  $\vec{a}$  is less than a half of  $\vec{a}$ . In terms of parameters this may be expressed as

$$\cot(\alpha) \leq \frac{1 + \delta a}{2}. \quad (32)$$

Here was taken into account that  $\bar{\rho} = 1/l^2$  and should be a constant.

Now results of calculation may be presented. Following table shows values calculated with minimization algorithm and parameters of most similar lattice we can think of.

Parameter	Calculated value	Triangular lattice
$\alpha$	$\approx 1.0472$	$\pi/3 \approx 1.0472$
$a$	$\approx 1.0746$	$\sqrt{2}/\sqrt[4]{3} \approx 1.07457$

It seems that triangular lattice has minimal potential energy that coincides with our intuitive expectations.

## B. 3D lattice.

After considering possible problems and results with 2D lattice we are ready to analyze 3D case. Minimization is performed in a similar way. We use (27) as function to be minimized. Particles density is fixed in a similar way. We consider

$$\frac{1}{l^3} = \bar{\rho} = \frac{1}{abc \sin(\alpha) \sin(\beta_c)},$$

suppose  $a = l(1 + \delta a)$ ,  $b = l(1 + \delta b)$  and express  $c$  through other parameters to keep constant charge density.

Besides, we will meet the same uncertainty problem we had in 2D case. So we should add some restrictions. For vector  $\vec{b}$  it is obviously to use same restriction as previous (32).

Restrictions for  $\vec{c}$  are more sophisticated. But they may be achieved with same considerations as for  $\vec{b}$ . The only difference is that  $\vec{b}$  was projected onto  $\vec{a}$  and we tried to keep its projection less than  $\vec{a}/2$ . Vector  $\vec{c}$  should be projected onto both  $\vec{a}$  and  $\vec{b}$  and its projections should be kept less than half of this vectors respectively. This leads to following equations

$$0 \leq \frac{\cot(\beta_c)}{(1 + \delta a)} \sin(\alpha_c) \leq \frac{(1 + \delta b)^2 \sin^2(\alpha)}{2};$$

$$0 \leq \frac{\cot(\beta_c)}{(1 + \delta b)} (\cos(\alpha_c) - \cos(\alpha)) \leq \frac{(1 + \delta a)^2 \sin^2(\alpha)}{2}.$$

Reader may check that this equations are  $(\vec{c} \cdot \vec{a}) \leq |\vec{a}|/2$  and  $(\vec{c} \cdot \vec{b}) \leq |\vec{b}|/2$  taking relation for  $\bar{\rho}$  into account.

Now we can present results of numerical minimization in form of figure 4 and following table.

Looking at figure 4, one may see, it is very similar to hexagonal close packing (HCP) lattice. HCP consists of two interleaving planes with triangular 2-D lattice in each as well. So we may check, if this is really HCP lattice by comparing their parameters.






Figure 4: Part of lattice obtained from minimization. As is seen, 3-D lattice consists of interleaving planes with 2-D hexagonal (triangular) lattices in each. The only difference between planes is shift: particles from one plane project onto “empty spaces” in another. Translation vectors are shown as well.

Parameter	Calculated value	Hexagonal close packing (HCP)
$\alpha$	$\approx 1.04717$	$\pi/3 \approx 1.0472$
$\alpha_c$	$\approx 0.523589$	$\pi/6 \approx 0.5236$
$\beta_c$	$\approx 0.955205$	$\arcsin(\sqrt{2/3}) \approx 0.9553$
$a$	$\approx 1.122462$	$\sqrt[6]{2} \approx 1.12246$
$b$	$\approx 1.122462$	$\sqrt[6]{2} \approx 1.12246$

Results show that lattice obtained by numerical calculations is really HCP. This complies with experimental data and computer simulation [5, 6].

#### IV. ELECTRONS ON THE LIQUID-HELIUM SURFACE AND THEIR LOCALIZATION DISTANCE

In this section we consider electrons on the liquid helium surface. In this section we will calculate localization distance and compare it to obtained in other works.

In the presence of an external field electrons can be pressed against the helium surface with significant force. But they cannot go through this surface, because quantum effects push them out [23, 28, 29]. Thus they deform surface and this deformation, in its turn, changes interaction potential between electrons. The problem of finding explicit expression of this potential is solved by adding to Coulomb interaction effective capillary interaction. The lateral for two electrons on the helium surface was calculated in [30].

##### A. Localization distance

Now we aim to find localization distance for electrons on the liquid helium surface. To achieve this goal we will

minimize single-particle’s free energy  $F_{sp}$  over  $s$  (localization distance itself, see example IA). So first of all we construct  $F_{sp}$  and find  $\partial F_{sp}/\partial s$ .

In section I we developed a method for treating some quantum systems. Example IA shows how to use it for two-dimensional system of Fermi-particles. So we consider  $F_{ent}$  from (13) as free energy of one particle caused with entropy in the system.

Potential energy is more complicated. In section II we developed methods of its calculation for particles arranged in a lattice. Since approximate expression for Coulomb interaction in 3D lattice, presented in example IIB (27), does not contain  $s$ , the same behavior is expected for 2D. Since we are interested in derivative  $\partial F_{sp}/\partial s$ , consideration of this part of potential energy may be omitted. More interesting is capillary interaction presented in example IIC, approximation (31). It contains explicit dependence on  $s$  and thus should be taken into account.

Bringing it all together we get

$$\frac{\partial F_{sp}}{\partial s} = -\frac{\lambda_T^2}{4\pi^2\beta s^3} - \frac{\pi^4 s}{3\beta\lambda_T^2} + \frac{q^2 E^2}{4\sigma\pi s},$$

independently on lattice we are considering.

Substituting electron charge  $q \rightarrow \bar{e}$  and solving equation  $\partial F/\partial s = 0$  we get

$$s^2 = \frac{3\hbar^2\beta^2\bar{e}^2 E^2}{4\pi^3\sigma m} \left( 1 - \sqrt{1 - \frac{16\sigma^2\pi^4}{3\bar{e}^4 E^4 \beta^2}} \right) \quad (33)$$

If temperature tends to be small  $T \rightarrow 0$  (e.g.  $\beta \rightarrow \infty$ )

$$s = \sqrt{\frac{2\pi\sigma\hbar^2}{m\bar{e}^2 E^2}}. \quad (34)$$

This result does not differ from the classical one presented in [23].

If we consider (33) more carefully, some interesting result may be obtained. Let’s consider low densities so that any value of  $s$  may be treated as much lower than inter-electron distance  $l$ . Then we may notice that solution for  $s$  exists if

$$\frac{E^2}{T} \geq \frac{4\sigma\pi^2 k}{\sqrt{3}\bar{e}^2} \approx 2100 \frac{dyn}{cm \cdot K}. \quad (35)$$

Last equation means that there is no lattice if temperature  $T$  is too high, or if field  $E$  is too weak. This statement is in agreement with our physical intuition.

#### V. CONCLUSIONS

In conclusion we want to make some overview of results we have obtained.

In **section I** we started with a statistical description of inhomogeneous system of interacting particles. It was shown that variety of quantum systems may be described

with no much difference from classical ones. We found an expression for free energy (6). The main feature of this equation appears in expression for “entropy part” of free energy. Obtained result is applicable to a wide variety of systems with different number of dimensions.

To make a step aside from pure theory we considered an example IA of two-dimensional system of Fermi particles. This result was later reused in section IV.

In **section II** we stated that particles are arranged in a lattice. It seems, that adding translation symmetry to consideration we can get some simplification for potential energy expressions (16) and (17). One may notice that for any “catastrophic” potential only there is only one diverging term. But this term does not contain information on “lattice geometry”, it depends on mean particle density only. This means, we can compare two lattices with equal mean particle density even if inter-particle potential is “catastrophic”.

In this approach all potentials are classified into two groups: ones with self-interaction (often effective interactions through medium) and without. Examples IIB and IIC demonstrate how to use proposed technique in both cases. Later they are reused for treating grains in dusty plasma and electrons on the liquid-helium surface.

In **section III** we considered grains in dusty plasma. We started with potential energy obtained in example IIB and minimized it over all possible lattices. Since there are no good approximations for theta-function it was too complicated to perform this minimization analytically. Thus we used numerical computations. This computations have some features first presented in a simplified case IIIA and then used in a real-life case IIIB. Results obtained in IIIB show that we should expect hexagonal close packing (HCP) lattice for grains in dusty plasma. This coincides with numerical simulations in [6] and experiments [5].

Last **section IV** shows how to use developed methods for electrons on the liquid-helium surface. It is shown that in some cases much more simple (but more rough as well) compared to section III approximation may be obtained. This approximation is used to calculate localization distance and then compared to classical result [23]. When temperature  $T \rightarrow 0$  they fully coincide. Besides, basing on this results some “guess” when lattice can exist is done. Obtained equation can be verified experimentally.

## VI. APPENDICES

During this article we tried to provide rigorous exposition of presented ideas, but without physical sole being lost in lots of equations. Thus, calculations that heavily rely on mathematical transformations only, were moved to appendices. Reader, interested only in “physical part”, may omit them.

### A. Integration over momentum in $\beta F[\mu]$ .

We start with (8). First two summands can be easily integrated if we mention

$$\frac{1}{e^x + 1} = \frac{e^{-x}}{1 + e^{-x}},$$

but last summand should be integrated in terms of special functions, namely dilogarithm [44]

$$\text{Li}_2(z) = - \int_0^z \frac{\ln(1-z)}{z} dz.$$

If we mention that

$$\frac{1}{1 + e^x} = 1 - \frac{1}{1 + e^{-x}}.$$

we would get

$$\begin{aligned} \beta F[\mu] = & \frac{m^2}{8\pi^2 \hbar^4 \beta} \int \int d^2 \vec{r} d^2 \vec{r}' V(|\vec{r} - \vec{r}'|) \times \\ & \times \ln(1 + e^{\beta \mu(\vec{r})}) \ln(1 + e^{\beta \mu(\vec{r}')}) + \\ & + \frac{m}{2\pi \hbar^2 \beta} \int d^2 r \left[ \beta \mu(\vec{r}) \ln(1 + e^{\beta \mu(\vec{r})}) - \right. \\ & \left. - \frac{\beta^2 \mu^2(\vec{r})}{2} - \text{Li}_2(-e^{-\beta \mu(\vec{r})}) - \frac{\pi^2}{6} \right], \end{aligned} \quad (36)$$

which can be simplified further.

First of all we will use Landen’s identity [44]

$$\begin{aligned} \text{Li}_2(1-z) + \text{Li}_2\left(1 - \frac{1}{z}\right) = & -\frac{1}{2} \ln^2(z), \\ z \in \mathbb{C} \setminus ]-\infty; 0] \end{aligned} \quad (37)$$

with  $z = 1 + e^{-\beta \mu(\vec{r})}$  and expand  $\ln^2(1 + e^{-\beta \mu(\vec{r})})$ . This leads to

$$\begin{aligned} \beta F[\mu] = & \frac{m^2}{8\pi^2 \hbar^4 \beta} \int \int d^2 \vec{r} d^2 \vec{r}' V(|\vec{r} - \vec{r}'|) \times \\ & \times \ln(1 + e^{\beta \mu(\vec{r})}) \ln(1 + e^{\beta \mu(\vec{r}')}) + \\ & + \frac{m}{2\pi \hbar^2 \beta} \int d^2 r \left[ \text{Li}_2\left(\frac{1}{1 + e^{\beta \mu(\vec{r})}}\right) - \frac{\pi^2}{6} + \right. \\ & \left. + \frac{1}{2} \ln^2(1 + e^{\beta \mu(\vec{r})}) \right]. \end{aligned}$$

Now we substitute (10) and (9) into the last expression and get (11).

### B. On the properties of $f_{\vec{k}}(\vec{r})$ .

In (16) we present how to calculate probability distribution function for the whole system basing on single-particle probability distribution function. It is based on

the decomposition of  $\rho_{\text{sp}}$  using  $\mathbf{f}_{\vec{k}}(\vec{r})$  (16c) as basis functions in inverse lattice space. Here we add some consideration about connection between Fourier series and presented one.

Suppose we have a function  $g(\vec{r})$  defined on unit cube  $\mathbb{U} \equiv [0; 1] \times [0; 1] \times [0; 1]$ . This function can be expressed in terms of Fourier series

$$\begin{aligned} g(\vec{r}) &= \sum_{\vec{k} \in \mathbb{Z}^3} g_{\vec{k}} e^{2\pi i(\vec{k}^T \vec{r})}, \\ g_{\vec{k}} &= \iiint_{\mathbb{U}} g(\vec{r}) e^{2\pi i(\vec{k}^T \vec{r})} d^3 \vec{r}. \end{aligned} \quad (38)$$

Let's consider a bijection  $\hat{G}$  (16d) from  $\mathbb{V}$  to  $\mathbb{U}$  and its inverse  $\hat{G}^{-1}$  (see Fig. 1 for geometrical reasoning)

$$\hat{G}^{-1} = \begin{pmatrix} a & b \cos(\alpha) & c \cos(\alpha_c) \cos(\beta_c) \\ 0 & b \sin(\alpha) & c \sin(\alpha_c) \cos(\beta_c) \\ 0 & 0 & c \sin(\beta_c) \end{pmatrix} \quad (39)$$

It may be seen that rows of  $\hat{G}$  are components of inverse lattice basis vectors. We are interested in case  $g(\vec{r}) = \rho_{\text{sp}}(\hat{G}^{-1}\vec{r})$ . Since we know that for  $\forall \vec{r} \in \mathbb{V}$ :  $\rho_{\text{sp}}(\hat{G}^{-1}\hat{G}\vec{r}) = \rho_{\text{sp}}(\vec{r})$  one can immediately write

$$\rho_{\text{sp}}(\vec{r}) = \sum_{\vec{k} \in \mathbb{Z}^3} g_{\vec{k}} e^{2\pi i(\vec{k}^T \hat{G}\vec{r})}. \quad (40)$$

And same way we consider second equation from (38)

$$g_{\vec{k}} = \iiint_{\mathbb{V}} \rho_{\text{sp}}(\hat{G}^{-1}\vec{r}) e^{2\pi i\vec{k}^T \hat{G}\hat{G}^{-1}\vec{r}} \frac{d^3(\hat{G}^{-1}\vec{r})}{J[\hat{G}^{-1}\vec{r}]}, \quad (41)$$

where  $J$  is Jacobian.

Jacobian  $J[\hat{G}^{-1}\vec{r}] = abc \sin(\alpha) \sin(\beta_c) = 1/\bar{\rho}$  is actually the volume of  $\mathbb{V}$  or inverse mean particle density. Designating  $g_{\vec{k}}$  as  $\rho_{\vec{k}}$  in (40) and (41), changing variable in (41)  $\hat{G}^{-1}\vec{r} \rightarrow \vec{r}$  and designating exponent in (41) as  $\mathbf{f}_{\vec{k}}$  we immediately get equations (16). Moreover, obtained result means that all properties of Fourier series can be applied to decomposition (16).

Last thing to mention are periodical properties of presented series. We are going to expand domain to  $\mathbb{R}^3$  so that  $\rho(\vec{r})$  will be defined everywhere in the space. So now we need to explore behavior of this function. From (16d) one may see, if  $l \in \mathbb{Z}$ ,  $m \in \mathbb{Z}$  and  $n \in \mathbb{Z}$ :

$$\hat{G}(\vec{r} + l\vec{a} + m\vec{b} + n\vec{c}) = \hat{G}\vec{r} + l\vec{e}_x + m\vec{e}_y + n\vec{e}_z, \quad (42)$$

where  $\vec{e}_x$ ,  $\vec{e}_y$  and  $\vec{e}_z$  are unit vectors along coordinate axis. With regard to (16c)

$$\mathbf{f}_{\vec{k}}(\vec{r} + l\vec{a} + m\vec{b} + n\vec{c}) = \mathbf{f}_{\vec{k}}(\vec{r}) \quad (43)$$

and from (16a)

$$\rho(\vec{r} + l\vec{a} + m\vec{b} + n\vec{c}) = \rho(\vec{r}). \quad (44)$$

Last equation justifies our view of connection between  $\rho$  and  $\rho_{\text{sp}}$  as it is presented on Fig. 2.

### C. On the expression of $\rho_{\vec{k}}$ for Gaussian $\rho_{\text{sp}}$

We suppose that  $\rho_{\text{sp}}(\vec{r})$  is either 0 everywhere in  $\mathbb{R}^3 \setminus \mathbb{V}$ , or at least negligibly small. If so, we can change integration limits in (16b) to infinite

$$\rho_{\vec{k}} = \bar{\rho} \iiint_{\mathbb{R}^3} \rho_{\text{sp}}(r) e^{2\pi i(g_{\vec{k}}z + g'_{\vec{k}}x + g''_{\vec{k}}y)} d^3 \vec{r},$$

where  $g_{\vec{k}}$ ,  $g'_{\vec{k}}$  and  $g''_{\vec{k}}$  from (16c) and (16d)

$$\begin{aligned} g_{\vec{k}} &= \frac{k_x \sin(\alpha_c - \alpha)}{a \tan(\beta_c) \sin(\alpha)} - \frac{k_y \sin(\alpha_c)}{b \tan(\beta_c) \sin(\alpha)} + \frac{k_z}{c \sin(\beta_c)}, \\ g'_{\vec{k}} &= \frac{k_x}{a}, \\ g''_{\vec{k}} &= -k_x \frac{\cot(\alpha)}{a} + \frac{k_y}{b \sin(\alpha)}. \end{aligned}$$

Rewriting (16c) in spherical coordinates

$$\mathbf{f}_{\vec{k}}(\vec{r}) = e^{2\pi i r (g_{\vec{k}} \cos(\theta) + [g'_{\vec{k}} \cos(\varphi) + g''_{\vec{k}} \sin(\varphi)] \sin(\theta))},$$

substituting (12) and performing integration we will get

$$\rho_{\vec{k}} = \bar{\rho} e^{-2\pi^2 s^2 (g_{\vec{k}}^2 + g'_{\vec{k}}{}^2 + g''_{\vec{k}}{}^2)}.$$

By direct comparison of this equation with (18) one may check that they are equal.

### D. On the expression of $F_{\text{int}}$ .

Lets start with (15a). Inner integral has infinite borders so we may rewrite this expression as follows

$$F_{\text{int}} = \iiint_{\mathbb{V}} \iiint_{\mathbb{R}^3} V(|\vec{r}'|) \rho(\vec{r}) \rho(\vec{r} + \vec{r}') d^3 \vec{r}' d^3 \vec{r}. \quad (45)$$

Now we can substitute  $\rho(\vec{r})$  from (16a) and mention that  $\mathbf{f}_{\vec{k}}(\vec{r} + \vec{r}') = \mathbf{f}_{\vec{k}}(\vec{r}) \mathbf{f}_{\vec{k}}(\vec{r}')$  (16c). So if we designate  $V_{\vec{k}}$  as in (17b) it can be written

$$F_{\text{int}} = \frac{1}{\bar{\rho}} \sum_{\vec{k} \in \mathbb{Z}^3} \rho_{\vec{k}} V_{\vec{k}} \sum_{\vec{k}' \in \mathbb{Z}^3} \rho_{\vec{k}'} \iiint_{\mathbb{V}} \mathbf{f}_{\vec{k}}(\vec{r}) \mathbf{f}_{\vec{k}'}(\vec{r}) d^3 \vec{r}.$$

Last equation can be simplified if we perform integration. From appendix VI B we expect orthogonality of

$f_{\vec{k}}$  functions. Let's mention from (16c)  $f_{\vec{k}}(\vec{r})f_{\vec{k}'}^*(\vec{r}) = f_{\vec{k}-\vec{k}'}(\vec{r})$  and we will get

$$\iiint_{\mathbb{V}} f_{\vec{k}}(\vec{r})f_{\vec{k}'}^*(\vec{r})d^3\vec{r} = abc \sin(\alpha) \sin(\beta_c) \delta_{\vec{k},\vec{k}'}. \quad (46)$$

Here  $\delta_{\vec{k},\vec{k}'} = \delta_{k_x,k'_x} \delta_{k_y,k'_y} \delta_{k_z,k'_z}$  — product of three Kronecker's delta.

Mentioning that  $abc \sin(\alpha) \sin(\beta_c) = 1/\bar{\rho}$ , where  $\bar{\rho}$  is mean particle density (one particle per  $\mathbb{V}$ , see fig. 1 for geometrical reasoning) we get (17a).

### E. $V_{\vec{k}}$ for screened Coulomb potential

Let's consider equations (16c) and (16d) in spherical coordinates

$$\begin{aligned} f_{\vec{k}}(\vec{r}) &= e^{2\pi i \mathbf{r} \cdot (g_{\vec{k}} \cos(\theta) + g'_{\vec{k}} \cos(\varphi - \delta\varphi_{\vec{k}}) \sin(\theta))}, \\ g_{\vec{k}} &= \frac{k_x \sin(\alpha_c - \alpha)}{a \tan(\beta_c) \sin(\alpha)} - \frac{k_y \sin(\alpha_c)}{b \tan(\beta_c) \sin(\alpha)} + \frac{k_z}{c \sin(\beta_c)}, \\ g'_{\vec{k}} &= \sqrt{\frac{k_x^2}{a^2} + \left( \frac{k_y}{b \sin(\alpha)} - k_x \frac{\cot(\alpha)}{a} \right)^2}, \end{aligned}$$

and rewrite  $V_{\vec{k}}$  from (17b)

$$\begin{aligned} V_{\vec{k}} &= \bar{\rho} \int_0^\infty dr V(r) r^2 \int_0^\pi d\theta \sin(\theta) \times \\ &\quad \times \underbrace{\int_0^{2\pi} d\varphi e^{2\pi i \mathbf{r} \cdot (g_{\vec{k}} \cos(\theta) + g'_{\vec{k}} \cos(\varphi - \delta\varphi_{\vec{k}}) \sin(\theta))}}_{I(\theta)}. \end{aligned}$$

Integrating over  $\varphi$  we get

$$I(\theta) = 2\pi e^{2\pi i g_{\vec{k}} r \cos(\theta)} J_0 \left( 2\pi g'_{\vec{k}} r \sin(\theta) \right).$$

Then we substitute last expression into equation for  $V_{\vec{k}}$

$$\begin{aligned} V_{\vec{k}} &= 2\pi \bar{\rho} \int_0^\infty V(r) r^2 \int_0^\pi e^{2\pi i g_{\vec{k}} r \cos(\theta)} \times \\ &\quad \times J_0 \left( 2\pi g'_{\vec{k}} r \sin(\theta) \right) \sin(\theta) d\theta dr. \end{aligned}$$

We will consider screened Coulomb potential (21), which means integration over  $r$  can be performed. One may use following relation [45]

$$\begin{aligned} \int_0^\infty e^{-\alpha x} J_\nu(\beta x) x^{\nu+1} dx &= \frac{2\alpha(2\beta)^\nu \Gamma(\nu + \frac{3}{2})}{\sqrt{\pi}(\alpha^2 + \beta^2)^{\nu+3/2}}, \\ \Re(\alpha) &> |\Im(\beta)|, \Re(\nu) > -1 \end{aligned}$$

and combine it with equation for  $V_{\vec{k}}$  to get

$$V_{\vec{k}} = \int_0^\pi \frac{2\pi \bar{\rho} q^2 (1/\lambda_D - 2\pi i g_{\vec{k}} \cos(\theta)) \sin(\theta) d\theta}{\left( (1/\lambda_D - 2\pi i g_{\vec{k}} \cos(\theta))^2 + (2\pi g'_{\vec{k}} \sin(\theta))^2 \right)^{3/2}}.$$

Changing variable  $t = \cos(\theta)$  and performing integration over  $t$  we will get

$$V_{\vec{k}} = \frac{4\pi \bar{\rho} q^2}{1/\lambda_D^2 + 4\pi^2 g_{\vec{k}}^2 + 4\pi^2 g'_{\vec{k}}{}^2}.$$

Last expression is equivalent to (22). One may check this by expanding  $g_{\vec{k}}$  and  $g'_{\vec{k}}$ .

### F. Calculation of $F_s$ for screened Coulomb potential.

We will start with equation (15b) and assume that  $\rho_{sp}$  is a very ‘‘sharp’’ function (section I A). With this assumption we can change integration limits to infinite and this, in turn, will allow us to perform variable exchange as we did in (45)

$$F_s = \iiint_{\mathbb{R}^3} \iiint_{\mathbb{R}^3} V(|\vec{r}'|) \rho_{sp}(\vec{r}) \rho_{sp}(\vec{r} + \vec{r}') d^3 \vec{r}' d^3 \vec{r}. \quad (47)$$

First of all we perform some mathematical transformations with (12) (in Cartesian coordinate system) and write the following

$$\rho_{sp}(\vec{r}) \rho_{sp}(\vec{r} + \vec{r}') = \frac{1}{8\pi^3 s^6} e^{-[\vec{r} + \vec{r}'/2]^2/s^2 - \vec{r}'^2/(4s^2)}.$$

Now we can substitute this expression into (47) and perform integration over  $\vec{r}'$ . Writing result in spherical coordinate system and integrating over angles we immediately get (24).

Since we know explicit expression for  $V$  (21) we can substitute it into (24) and perform integration using definition of the complementary error function [46]

$$\operatorname{erfc}(x) = \frac{2}{\sqrt{\pi}} \int_x^\infty e^{-t^2} dt. \quad (48)$$

As result we will get

$$F_s = \frac{q^2}{\sqrt{\pi} s} \left( 1 - \frac{s\sqrt{\pi}}{\lambda_D} e^{s^2/\lambda_D^2} \operatorname{erfc}(s/\lambda_D) \right). \quad (49)$$

Since we may obviously suppose that  $s$  is very small compared to screening distance, then  $s \ll \lambda_D$ . So we can approximate last equation and as result we will get (25).

### G. Approximation of $F_{int} - F_s$ for screened Coulomb potential

We start with expression (23) for  $F_{int}$ . But one may see that it contains expressions  $s^2 \hat{G} \hat{G}^T$ . Components of  $\hat{G} \hat{G}^T$  matrix in (23) are proportional to different products of inverse distances in a lattice, e.g.  $1/a^2$ ,  $1/(ab)$ , and so on (see (16d)). Since  $s \ll a$ ,  $s \ll b$  and  $s \ll c$  (assumption about ‘‘sharp Gaussian’’) we expect  $|s^2 \hat{G} \hat{G}^T| \ll 1$ . This means, that exponent ‘‘starts acting’’ only for terms with very large  $|\vec{k}|$ . In this section we will try to rewrite equations to get even better convergence then we have.

First of all we rewrite expression for  $F_{int}$  as follows

$$F_{int} = \frac{2\bar{\rho}q^2}{\pi} e^{s^2/\lambda_D^2} \int_{2\pi s}^{\infty} \mathfrak{s} e^{-s^2/(2\pi\lambda_D)^2} \sum_{\vec{k} \in \mathbb{Z}^3} e^{-s^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}} d\mathfrak{s}.$$

One may easily integrate this expression and check that it is equal to (23).

We notice, that variable  $\mathfrak{s}$  takes values much smaller then mean distance (26) as well as much bigger. Thus we split this integral into two. First integration we will provide from  $2\pi s$  to  $l$  and second from  $l$  to infinity.

We assume that lattice is not too degenerated (section II B), which means that angles  $\alpha$  and  $\beta_c$  should not significantly differ from  $\pi/2$ . So we may expect elements of  $\mathfrak{s}^2 \hat{G} \hat{G}^T$  to be less than 1 for first integral and greater than 1 for second. If so, then second integration may be performed and we end up with the sum that converges much better than (23).

$$F_{int} = \frac{2\bar{\rho}q^2}{\pi} e^{s^2/\lambda_D^2} \int_{2\pi s}^l \mathfrak{s} e^{-s^2/(2\pi\lambda_D)^2} \underbrace{\sum_{\vec{k} \in \mathbb{Z}^3} e^{-s^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}}}_{\Theta(0; \mathfrak{s}^2 \hat{G} \hat{G}^T)} d\mathfrak{s} + 4\pi\bar{\rho}q^2 \sum_{\vec{k} \in \mathbb{Z}^3} \frac{e^{s^2/\lambda_D^2 - l^2(1/[2\pi\lambda_D]^2 + \vec{k}^T \hat{G} \hat{G}^T \vec{k})}}{1/\lambda_D^2 + 4\pi^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}}.$$

From (16d) one may check with Sylvester’s criterion [47] that  $\hat{G}$  (16d) is a positive-definite matrix. Obviously, we expect  $\hat{G}^T$  and their product  $\hat{G} \hat{G}^T$  to be positive-definite matrices as well. Since  $\mathfrak{s}$  takes only positive values  $\mathfrak{s}^2 \hat{G} \hat{G}^T$  is positive as well. This means highlighted sum in the last equation is a Riemann theta function [48] at point  $z = 0$ , so we appropriately designate it with  $\Theta(0; \mathfrak{s}^2 \hat{G} \hat{G}^T)$ .

Using modular transformation [48]

$$\Theta(z; \hat{A}) = \frac{\pi^{d/2}}{\sqrt{\det \hat{A}}} \Theta(\hat{A}^{-1}z; \hat{A}^{-1}), \quad (50)$$

where  $d$  is number of dimensions (3 in this case) we get

$$\Theta(0; 2\pi^2 \mathfrak{s}^2 \hat{G} \hat{G}^T) = \frac{\pi^{3/2}}{\mathfrak{s}^3 \det \hat{G}} \Theta(0; \hat{G}^{-1T} \hat{G}^{-1} / \mathfrak{s}^2).$$

We used the fact that  $\det \hat{G} = \det \hat{G}^T$  and that multiplication by  $\mathfrak{s}^2$  is equal to multiplication by diagonal matrix which has all elements equal to  $\mathfrak{s}^2$ . Explicit expression for  $\hat{G}^{-1}$  may be used from (39). Besides, we may notice that

$$\det \hat{G} = \frac{1}{abc \sin(\alpha) \sin(\beta_c)} = \bar{\rho}, \quad (51)$$

and rewrite expression for  $F_{int}$  as follows

$$F_{int} = 2\sqrt{\pi}q^2 e^{s^2/\lambda_D^2} \sum_{\vec{k} \in \mathbb{Z}^3} \underbrace{\int_{2\pi s}^l e^{-s^2/(2\pi\lambda_D)^2 - |\hat{G}^{-1}\vec{k}|^2/s^2} d\mathfrak{s}}_{I(\vec{k})} + 4\pi\bar{\rho}q^2 \sum_{\vec{k} \in \mathbb{Z}^3} \frac{e^{s^2/\lambda_D^2 - l^2(1/[2\pi\lambda_D]^2 + |\hat{G}^T \vec{k}|^2)}}{1/\lambda_D^2 + 4\pi^2 \vec{k}^T \hat{G} \hat{G}^T \vec{k}}.$$

Now we can perform integration and find  $I(\vec{k})$ . Since result is very complicated we first introduce two helper functions

$$\phi(L) = \frac{e^{-L^2/(2\pi\lambda_D)^2}}{L} + \frac{1}{2\sqrt{\pi}\lambda_D} \operatorname{erf}\left(\frac{L}{2\pi\lambda_D}\right),$$

$$f_\alpha(L) = e^{\alpha|\hat{G}^{-1}\vec{k}|/(\pi\lambda_D)} \operatorname{erfc}\left[\frac{|\hat{G}^{-1}\vec{k}|}{L} + \frac{\alpha L}{2\pi\lambda_D}\right],$$

and present result in terms of this functions

$$I(\vec{k} = 0) = \phi(2\pi s) - \phi(l),$$

$$I(\vec{k} \neq 0) = \sqrt{\pi}(f_{-1}(l) - f_{-1}(2\pi s)) / \left|4\hat{G}^{-1}\vec{k}\right| + \sqrt{\pi}(f_{+1}(l) - f_{+1}(2\pi s)) / \left|4\hat{G}^{-1}\vec{k}\right|.$$

Using precise expression for  $F_s$  (49) one may see

$$F_s = 2\sqrt{\pi}q^2 e^{s^2/\lambda_D^2} \phi(2\pi s).$$

So subtracting  $F_s$  from  $F_{int}$  simply means neglectation of this term.

Other terms we will approximate taking into account that  $s \ll l \ll \lambda_D$ . First we neglect all terms containing  $l/\lambda_D$  and  $s/\lambda_D$  in  $\operatorname{erfc}$ . It is known that  $\operatorname{erfc}(x \rightarrow \infty) \sim e^{-x^2}/(x\sqrt{\pi})$  [49] and thus  $f_\alpha(l) \gg f_\alpha(2\pi s)$ . So we neglect all  $f$  with  $2\pi s$  arguments. Besides, we approximate all  $\operatorname{erfc}$  with exponentiation and neglect small terms in exponent. To make this procedure more clear we state that  $\forall \vec{k} \in \mathbb{Z}^3 \setminus \vec{0} : |\hat{G}^{-1}\vec{k}|/l > 1$  and thus

$$\left|\hat{G}^{-1}\vec{k}\right|^2/l^2 > \left|\hat{G}^{-1}\vec{k}\right|/l \gg \left|\hat{G}^{-1}\vec{k}\right|/\lambda_D,$$

that leads us to equations

$$\phi(l) \approx \frac{1}{2\sqrt{\pi}\lambda_D},$$

$$I(\vec{k} \neq 0) \approx \frac{l}{2|\hat{G}^{-1}\vec{k}|^2} e^{-|\hat{G}^{-1}\vec{k}|^2/l^2}.$$

To get the first one, identity  $\text{erf}(x) = 1 - \text{erfc}(x)$  and approximation  $\text{erfc}(x \rightarrow \infty) \sim e^{-x^2}/(x\sqrt{\pi})$  [49] were used.

Now approximate equation can be written (27).

### H. $V_{\vec{k}}$ for capillary interaction

Let's consider two-dimensional version of (16c) and (16d) in polar coordinates

$$\mathfrak{f}_{\vec{k}}(\vec{r}) = e^{2\pi i r g_{\vec{k}} \cos(\varphi - \delta\varphi_{\vec{k}})},$$

$$g_{\vec{k}} = \sqrt{\frac{k_x^2}{a^2} + \left(\frac{k_y}{b \sin(\alpha)} - k_x \frac{\cot(\alpha)}{a}\right)^2},$$

and rewrite  $V_{\vec{k}}$  from (17b)

$$V_{\vec{k}} = \bar{\rho} \int_0^\infty dr V(r) r \underbrace{\int_0^{2\pi} d\varphi e^{2\pi i r g_{\vec{k}} \cos(\varphi - \delta\varphi_{\vec{k}})}}_I.$$

Integrating over  $\varphi$  we get

$$I = 2\pi J_0(2\pi g_{\vec{k}} r).$$

Then we substitute last equality and expression for  $V(r)$  from (28) into equation for  $V_{\vec{k}}$ . It is known [45], if  $a > 0$

and  $b > 0$

$$\int_0^\infty x J_0(ax) K_0(bx) dx = \frac{1}{a^2 + b^2}. \quad (52)$$

With this equation we easily obtain  $V_{n,m}$  for capillary interaction part

$$V_{\vec{k}} = -\frac{q^2 E^2}{\sigma} \frac{\bar{\rho}}{4\pi^2 g_{\vec{k}}^2 + 1/l_0^2}. \quad (53)$$

Comparison with (29) shows that they are equal.

### I. Approximation of $F_{int}$ for capillary interaction

We start with expression (30) for  $F_{int}$ . First of all we rewrite it as follows

$$F_{int} = -\frac{\bar{\rho} q^2 E^2}{2\sigma} e^{s^2/l_0^2} \int_{\mathfrak{s}} e^{-s^2/l_0^2} \sum_{\vec{k} \in \mathbb{Z}^2} e^{-4\pi^2 s^2 \vec{k}^T \hat{G} \vec{k}} d\mathfrak{s}.$$

Then we change summation to integration  $\sum_{\vec{k} \in \mathbb{Z}^2} \rightarrow \iint_{\mathbb{R}^2}$ . There is a method of integration for this kind of functions [47]. As result we get

$$F_{int} = -\frac{\bar{\rho} q^2 E^2}{2\sigma} e^{s^2/l_0^2} \int_{\mathfrak{s}} e^{-s^2/l_0^2} \frac{d\mathfrak{s}}{2\mathfrak{s}^2 \det \hat{G}}.$$

For two-dimension density it is true  $\det \hat{G} = \bar{\rho}$ . Besides, we perform integration over  $\mathfrak{s}$  and get

$$F_{int} = -\frac{q^2 E^2}{8\pi\sigma} e^{s^2/l_0^2} \Gamma_0(s^2/l_0^2),$$

where  $\Gamma_0$  is gamma-function. Since we know that  $s \ll l_0$  last equation can be approximated. We use known approximation [49]  $\Gamma_0(x \rightarrow 0) \approx -\gamma - \ln(x)$ , where  $\gamma$  is Euler-Mascheroni constant. We neglect this constant and exponent. The rest can be written as (31).

- 
- [1] V. E. Fortov et al., Phys. Rep. **421**, 1 (2005).
  - [2] H. Lowen, Phys. Rep. **237**, 249 (1994).
  - [3] G. E. Morfill, H. M. Thomas, U. Konopka, and M. Zuzic, Phys. Plasmas **6**, 1769 (1999).
  - [4] E. Helpand and F. H. Stilingler, J. Chem. Phys. **49**, 1232 (1968).
  - [5] B. Klumov, G. Joyce, C. R ath, P. Huber et al, *Structural properties of 3D complex plasmas under microgravity conditions*, EPL, Vol.92, No.1, p.15003, (2010).
  - [6] B. A. Klumov, G. E. Morfill, *Structural Properties of Complex (Dusty) Plasma upon Crystallization and Melting*, JETP Letters, Vol.90, No.6, pp. 444–448, (2009).
  - [7] B. I. Lev and A. G. Zagorodny, Phys. Lett. A **373**, 158 (2009).
  - [8] B. I. Lev, V. B. Tymchyshyn, and A. G. Zagorodny, Condens. Phys. **12**, 593 (2009).
  - [9] H. Thomas, G. E. Morfill, V. Demmel, J. Goree, B. Feuerbacher, and D. Mohlmann, Phys. Rev. Lett. **73**, 652 (1994).
  - [10] J. H. Chu and Lin I, Phys. Rev. Lett. **72**, 4009 (1994).
  - [11] A. Melzer, T. Trottenberg, and A. Piel, Phys. Lett. A **191**, 301 (1994).
  - [12] S. V. Vladimirov, S. A. Khrapak, M. Chaudhuri, and G. E. Morfill, Phys. Rev. Lett. **100**, 055002 (2008).
  - [13] H. Ikezi, Phys. Fluids. **29**, 1764 (1986).
  - [14] A. Melzer, A. Homann, and A. Piel, Phys. Rev. E **53**, 2757 (1996).
  - [15] A. G. Sitenko, A. G. Zagorodny, and V. N. Tsytovich, AIP

- Conf. Proc. **345**, 311 (1995).
- [16] S. A. Brazovskiy, Sov. Phys. JETP **68**, 715 (1975).
- [17] B. I. Lev and H. Yokoyama, IJMP B (International Journal of Modern Physics) **17**, 4913 (2003).
- [18] H. Totsuji, T. Kishimoto, and C. Totsuji, Phys. Rev. Lett. **78**, 3113 (1997).
- [19] P. Leiderer Two-Dimensional Electron Systems by Ed. E. Y. Andrei, (Springer Netherlands 1997).
- [20] V. B. Shikin and P. Leiderer, Sov. Phys. JETP Lett., **54**, 92, (1981).
- [21] D. C. Tsui, H. L. Stormer, and A. C. Gossard, Phys. Rev. Lett. **48**, 1559 (1982).
- [22] R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).
- [23] V. S. Edelman, Uspekhi Fizicheskikh Nauk **23**, 227 (1980).
- [24] T. Ando, A. Fowler, and F. Stern, Rev. Mod. Phys. **54**, 437 (1982).
- [25] W. Wigner, Phys. Rev. **46**, 1002 (1934).
- [26] C. C. Grimes, and G. Adams, Phys. Rev. Lett. **42**, 795 (1979).
- [27] P. M. Platzman and H. Fukuyama, Phys. Rev. B **10**, 3150 (1974).
- [28] I. Skachko, *Phase diagram of a 2-dimensional electron system on the surface of liquid helium*, Ph.D. thesis, Rutgers, The State University of New Jersey (2006).
- [29] D. K. Lambert, *Electrons on the surface of liquid helium*, Ph.D. thesis, Lawrence Berkeley Laboratory (1979).
- [30] M. Haque, I. Paul, and S. Pankov, Phys. Rev. B **68**, 045427 (2003).
- [31] B. I. Lev, V. P. Ostroukh, V. B. Tymchyshyn, and A. G. Zagorodny, *Statistical description of the system electrons on the liquid helium surface*, Eur. Phys. J. B, **87**:253 (2014).
- [32] L. P. Gor'kov and D. M. Chernikova, JETP Lett. **18**, 119 (1973).
- [33] D. Ruelle, *Statistical Mechanics: Rigorous Results*, W.A. Benjamin, New York, (1969).
- [34] A. Isihara, *Statistical Mechanics*, Academic Press, New York-London, (1971).
- [35] K. Huang, *Statistical Mechanics*, J. Wiley and Sons, New York, (1963).
- [36] R. Baxter, *Exactly Solved Models in Statistical Mechanics*, Academic Press, New York, (1982).
- [37] Y. D. Bilotsky and B. I. Lev, Teor. Math. Fiz. **60**, 120 (1984).
- [38] B. I. Lev and A. Y. Zhugaevych, Phys. Rev. E, **57**, 6460 (1998).
- [39] B. I. Lev and A. G. Zagorodny, Statistical description of Coulomb-like systems, Phys. Rev. E **84**, 061115 (2011).
- [40] R. L. Stratonovich, Soviet Physics Doklady, Vol. 2, (1958).
- [41] J. Hubbard, Phys. Rev. Lett. **3**, 77 (1959).
- [42] S. F. Edwards and A. Lennard, J. Math. Phys. **3**, 778 (1962).
- [43] S. Samuel, Phys. Rev. D **18**, 1916 (1978).
- [44] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher Transcendental Functions*, Vol. 1., McGraw-Hill, New York, (1953).
- [45] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, Amsterdam: Elsevier / Academic Press, (2007).
- [46] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher Transcendental Functions*, Vol. 2. Malabar, Krieger, (1981).
- [47] R. Bellman, *Introduction to Matrix Analysis*, Society for Industrial and Applied Mathematics, (1970).
- [48] R. Bellman, *A Brief Introduction to Theta Functions*, Holt, Rinehart and Winston, (1961).
- [49] M. Abramowitz, I. Stegun, *Handbook of Mathematical Functions With Formulas, Graphs, and Mathematical Tables*, Tenth Printing, (1972).