

Transverse instability for nonlinear Schrödinger equation with a linear potential

YOHEI YAMAZAKI*

Department of Mathematics, Kyoto University,
Kitashirakawa-Oiwakecho, Sakyo, Kyoto 606-8502, Japan

Abstract

In this paper we consider the transverse instability for a nonlinear Schrödinger equation with a linear potential on $\mathbb{R} \times \mathbb{T}_L$, where $2\pi L$ is the period of the torus \mathbb{T}_L . Rose and Weinstein [18] showed the existence of a stable standing wave for a nonlinear Schrödinger equation with a linear potential. We regard the standing wave of nonlinear Schrödinger equation on \mathbb{R} as a line standing wave of nonlinear Schrödinger equation on $\mathbb{R} \times \mathbb{T}_L$. We show the stability of line standing waves for all $L > 0$ by using the argument of the previous paper [26].

1 Introduction

We consider the nonlinear Schrödinger equation with linear potential

$$i\partial_t u = -\Delta u + V(x)u - |u|^{p-1}u, \quad (t, x, y) \in \mathbb{R} \times \mathbb{R} \times \mathbb{T}_L, \quad (1.1)$$

where $p > 1$, a potential $V : \mathbb{R} \rightarrow \mathbb{R}$ and $u = u(t, x, y)$ is an unknown complex-valued function for $t \in \mathbb{R}$, $x \in \mathbb{R}$ and $y \in \mathbb{T}_L$. Here, $\mathbb{T}_L = \mathbb{R}/2\pi L\mathbb{Z}$ and $L > 0$.

We assume the following conditions for V .

- (V1) There exist $C > 0$ and $\alpha > 0$ such that $|V(x)| \leq Ce^{-\alpha|x|}$.
- (V2) $-\partial_x^2 + V$ has the lowest eigenvalue $-\lambda_* < 0$.

The Cauchy problem (1.1) is locally well-posed in $H^1(\mathbb{R} \times \mathbb{T}_L)$ by using the argument in [7] and [24]. The equation (1.1) has the following conservation laws:

$$E(u) = \int_{\mathbb{R} \times \mathbb{T}_L} \left(\frac{1}{2} |\nabla u|^2 + \frac{1}{2} V(x) |u|^2 - \frac{1}{p+1} |u|^{p+1} \right) dx dy,$$

$$Q(u) = \frac{1}{2} \int_{\mathbb{R} \times \mathbb{T}_L} |u|^2 dx dy,$$

*E-mail addresses: y-youhei@math.kyoto-u.ac.jp

AMS 1991 subject classifications. 35B32, 35B35, 35Q55.

where $u \in H^1(\mathbb{R} \times \mathbb{T}_L)$.

We define a standing wave $u(t)$ as a non-trivial solution of (1.1) having the form $u(t) = e^{i\omega t}\varphi$. Then, $e^{i\omega t}\varphi$ is a standing wave if and only if φ is a non-trivial solution of

$$-\Delta\varphi + \omega\varphi + V(x)\varphi - |\varphi|^{p-1}\varphi = 0, \quad (x, y) \in \mathbb{R} \times \mathbb{T}_L. \quad (1.2)$$

Using the bifurcation theory, Rose and Weinstein [18] showed the existence of the stable standing wave $e^{i\omega t}\varphi_\omega$ for the following nonlinear Schrödinger equation

$$i\partial_t u = -\partial_x^2 u + V(x)u - |u|^{p-1}u, \quad (t, x) \in \mathbb{R} \times \mathbb{R}. \quad (1.3)$$

Then, the standing wave $e^{i\omega t}\varphi_\omega$ satisfies the following.

Proposition 1.1. *Let ψ_* be the eigenfunction of $-\partial_x^2 + V(x)$ corresponding to $-\lambda_*$ with $\psi_* > 0$ and $\|\psi_*\|_{L^2} = 1$. Then, there exists $\omega_* > \lambda_*$ such that for $\lambda_* < \omega < \omega_*$, $e^{i\omega t}\varphi_\omega$ is a stable standing wave of (1.3) satisfying*

$$\varphi_\omega = \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}} (\omega - \lambda_*)^{\frac{1}{p-1}} \psi_* + r(\omega),$$

where $\|r(\omega)\|_{H^2(\mathbb{R})} = O((\omega_0 - \lambda_*)^{\frac{1}{p-1}+1})$. Moreover, $L_\omega^+ = -\partial_x^2 + \omega + V - p|\varphi_\omega|^{p-1}$ has the exactly one negative eigenvalue $-\lambda_\omega$ and does not have the zero eigenvalue.

We define the line standing wave $e^{i\omega t}\tilde{\varphi}_\omega$ of (1.1) as

$$\tilde{\varphi}_\omega(x, y) = \varphi_\omega(x), \quad (x, y) \in \mathbb{R} \times \mathbb{T}_L.$$

In this paper, we consider the transverse instability of the line standing wave $e^{i\omega t}\tilde{\varphi}_\omega$. The stability of standing waves is defined as follows.

Definition 1.2. We say the standing wave $e^{i\omega t}\varphi$ is orbitally stable in H^1 if for any $\varepsilon > 0$ there exists $\delta > 0$ such that for all $u_0 \in H^1(\mathbb{R} \times \mathbb{T}_L)$ with $\|u_0 - \varphi\|_{H^1} < \delta$, the solution $u(t)$ of (1.1) with the initial data $u(0) = u_0$ exists globally in time and satisfies

$$\sup_{t \geq 0} \inf_{\theta \in \mathbb{R}, y \in \mathbb{T}_L} \|u(t, \cdot, \cdot - y) - e^{i\theta}\varphi(\cdot, \cdot - y)\|_{H^1} < \varepsilon.$$

Otherwise, we say the standing wave $e^{i\omega t}\varphi$ is orbitally unstable in H^1 .

The transverse instability for KP-I or KP-II equation is treated in [1, 15, 16, 19, 20, 21, 22]. In [1], Alexander-Pego-Sachs studied the linear instability for line solitons of KP-I and KP-II. In [16], Mizumachi-Tzvetkov proved the asymptotic stability for line solitons of KP-II on $\mathbb{R} \times \mathbb{T}$. Modulating the local phase and the local amplitude of line solitons, Mizumachi showed the asymptotic stability for line solitons of KP-II on \mathbb{R}^2 in [15]. Rousset-Tzvetkov proved the transverse instability for line solitons of KP-I on \mathbb{R}^2 in [19] and on $\mathbb{R} \times \mathbb{T}_L$ in [20]. In [22], Rousset-Tzvetkov showed the stability of line solitons for KP-I on $\mathbb{R} \times \mathbb{T}_L$ with small $L > 0$. Moreover, Rousset-Tzvetkov proved the existence of the critical period $4/\sqrt{3}$ for the period L of the transverse direction. Namely, a line soliton for KP-I on $\mathbb{R} \times \mathbb{T}_L$ is stable for $0 < L < 4/\sqrt{3}$ and unstable for $L > 4/\sqrt{3}$.

The transverse instability for a nonlinear Schrödinger equation is studied in [2, 19, 20, 25, 26]. In [2], Deconinck-Pelinovsky-Carter studied the linear stability for line standing waves of a hyperbolic Schrödinger equation. Rousset-Tzvetkov proved the transverse instability for cubic nonlinear Schrödinger equation without linear potential on \mathbb{R}^2 in [19] and on $\mathbb{R} \times \mathbb{T}_L$ in [20]. To prove the instability, Rousset-Tzvetkov applied the argument of Grenier [9]. Rousset-Tzvetkov constructed the high order approximate solution with an unstable eigenmode and showed a precise estimate of the growth of the semi-group generated by the linearized operator. To construct the high order approximate solution, we use the regularity of the nonlinearity $|u|^2u$ in the sense of Fréchet differentiation. In [25], the author studied the transverse instability for line standing waves of a system of nonlinear Schrödinger equations on $\mathbb{R} \times \mathbb{T}_L$ which was treated in [4]. In [25], the existence of the critical period for a period L was also proved, which was suggested by Rousset-Tzvetkov. Constructing the estimate for high frequency parts of solutions and using the existence of local solutions, the author showed the transverse instability for line standing waves of equations with the general power nonlinearity. In [26], the author considers the stability for a line standing wave of (1.1) with $V = 0$. The application of the argument in [25] yields the existence of the critical period for a line standing wave of (1.1) with $V = 0$. For (1.1) with $V = 0$ and the critical period, the linearized operator around the line standing wave is degenerate. Therefore, we can not directly apply the argument in Grillakis-Shatah-Strauss [10, 11]. Since the linearized operator around the line standing wave with the critical period does not have any unstable eigenvalues, we can not show the instability by the argument based on the occurrence of unstable eigenmode in [5, 10, 20, 25]. Moreover, the third order term of the Lyapunov functional around the line standing wave with the critical period does not appear. Thus, we can not apply the argument for the degenerate case of the stability in [14]. The transverse instability comes from the symmetry breaking bifurcation. In [26], applying the bifurcation result for symmetry breaking bifurcation and the stability result for the degenerate case in [13], the author showed the stability for the line standing wave with critical period for some exponents $p \geq 2$ of the nonlinearity.

The followings are our main theorems in this paper. In the first theorem, we show the transverse instability of the line standing wave $e^{i\omega t} \tilde{\varphi}_\omega$ and obtain the critical period between the stability and the instability.

Theorem 1.3. *There exists $\omega_{*,0} > \lambda_*$ such that for $\lambda_* < \omega < \omega_{*,0}$ the followings two assertions hold:*

- (i) *If $0 < L < (\lambda_\omega)^{-\frac{1}{2}}$, then the standing wave $e^{i\omega t} \tilde{\varphi}_\omega$ is stable.*
- (ii) *If $(\lambda_\omega)^{-\frac{1}{2}} < L$, then the standing wave $e^{i\omega t} \tilde{\varphi}_\omega$ is unstable.*

In the Second theorem, we show the stability for the line standing wave $e^{i\omega t} \tilde{\varphi}_\omega$ with the critical period $L = (\lambda_\omega)^{-1/2}$.

Theorem 1.4. *Let $p \geq 2$ and*

$$p_* = \frac{9 + \sqrt{57}}{4}.$$

Then there exists $\lambda_ < \omega_p$ satisfying the following two properties:*

- (i) If $p < p_*$ and $\lambda_* < \omega < \omega_p$, then the standing wave $e^{i\omega t} \tilde{\varphi}_\omega$ of (1.1) with $L = (\lambda_\omega)^{-1/2}$ is stable.
- (ii) If $p_* \leq p$ and $\lambda_* < \omega < \omega_p$, then the standing wave $e^{i\omega t} \tilde{\varphi}_\omega$ of (1.1) with $L = (\lambda_\omega)^{-1/2}$ is unstable.

The proof of Theorem 1.3 follows from the spectrum analysis of the linearized operator and the estimate of high frequency parts of solution by the argument in [25]. To show the growth of the semi-group generated by the linearized operator, we use the assumption of the decay for the linear potential V . For the proof of Theorem 1.4, we apply the bifurcation analysis for the symmetry breaking bifurcation and the argument for the stability in [26]. In [26], to prove the stability for the line standing wave with the critical period, we show the increase of L^2 -norm of the symmetry breaking standing wave with respect to the bifurcation parameter or the decrease of it. To show the increase of L^2 -norm, we need to calculate an integral of a solution of an ordinary differential equation which comes from the linearized equation of one dimensional Schrödinger equation around a standing wave. Since it is difficult to obtain the explicit solution of the ordinary differential equation in the argument in [25], we can not calculate the exact value of the integral and we estimate the value of the integral. Therefore, it is not known whether the line standing wave is stable or unstable for some nonlinear Schrödinger equations with the power nonlinearity $|u|^{p-1}u$ which has some exponent $p \in (2, 3)$. In the proof of Theorem 1.4, we treat the small standing wave which bifurcates from the eigenfunction of $-\partial_x^2 + V$ with respect to the lowest eigenvalue. Since the line standing wave of the nonlinear Schrödinger equation studied in [26] comes from the standing wave of the one dimensional nonlinear Schrödinger equation which has the scale invariant, we need to study the fully nonlinear structure of the Lyapunov functional around the line standing wave. In this paper, using the smallness of the line standing wave of (1.1) and the expansion of the standing wave with respect to the parameter ω , we weaken the nonlinear structure of the Lyapunov functional around the line standing wave of (1.1). Therefore, we can evaluate a value of the integral and make a close investigation into the stability for all exponents $p \geq 2$.

The rest of this paper consists of the following three sections. In Section 2, we show the properties of the spectrum and the coerciveness for the linearized operator around line standing waves. In Section 3, applying the variational argument in [10, 4] and the spectrum argument in [25], we prove Theorem 1.3. In Section 4, combining the bifurcation result and the argument for the degenerate case in [13], we prove Theorem 1.4.

2 Preliminaries

In this section, we investigate properties of the linearized operator of (1.1) around the standing wave $e^{i\omega t} \tilde{\varphi}_\omega$.

Let $H^1(X) = \{u : X \rightarrow \mathbb{C} \mid \int_X (|\nabla u|^2 + |u|^2) dx < \infty\}$ and $H^1(X, \mathbb{R}) = \{u : X \rightarrow \mathbb{R} \mid \int_X (|\nabla u|^2 + |u|^2) dx < \infty\}$. Let ψ_ω be the eigenfunction of L_ω^+ corresponding to $-\lambda_\omega$ with $\|\psi_\omega\|_{L^2(\mathbb{R})} = 1$ and $\psi_\omega > 0$. We define the action

$$S_\omega(u) = E(u) + \omega Q(u).$$

Then, the action S_ω is a conservation law of (1.1) and $S'_\omega(\tilde{\varphi}_\omega) = 0$, where S'_ω is the Fréchet derivation of S_ω . Moreover, we have

$$\langle S''_\omega(\tilde{\varphi}_\omega)u, v \rangle_{H^{-1}, H^1} = \langle \mathbb{L}_\omega^+(\operatorname{Re} u), \operatorname{Re} v \rangle_{H^{-1}, H^1} + \langle \mathbb{L}_\omega^-(\operatorname{Im} u), \operatorname{Im} v \rangle_{H^{-1}, H^1},$$

where

$$\mathbb{L}_\omega^+ = -\Delta + \omega + V - p|\tilde{\varphi}_\omega|^{p-1}, \quad \mathbb{L}_\omega^- = -\Delta + \omega + V - |\tilde{\varphi}_\omega|^{p-1}.$$

Let

$$Ju = iu = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \operatorname{Re} u \\ \operatorname{Im} u \end{pmatrix}.$$

For $u, v \in L^2(\mathbb{R} \times \mathbb{T}_L)$, we define

$$\langle u, v \rangle_{L^2} = \operatorname{Re} \int_{\mathbb{R} \times \mathbb{T}_L} u \bar{v} dx dy.$$

In the following proposition, we show properties of the spectrum of the linearized operator for (1.1) around $\tilde{\varphi}_\omega$. This proposition follows Theorem 1.1 of [21] and Lemma 3.1 of [22] (also see Proposition 2.5 of [25].)

Proposition 2.1. *Let $\lambda_* < \omega < \omega_*$.*

- (i) *If $0 < L \leq (\lambda_\omega)^{-1/2}$, then $-JS''_\omega(\tilde{\varphi}_\omega)$ has no positive eigenvalue.*
- (ii) *If $0 < L < (\lambda_\omega)^{-1/2}$, then*

$$\operatorname{Ker}(S''_\omega(\tilde{\varphi}_\omega)) = \operatorname{Span} \{i\tilde{\varphi}_\omega\}.$$

- (iii) *If $L = (\lambda_\omega)^{-1/2}$, then*

$$\operatorname{Ker}(S''_\omega(\tilde{\varphi}_\omega)) = \operatorname{Span} \left\{ i\tilde{\varphi}_\omega, \psi_\omega \cos \frac{y}{L}, \psi_\omega \sin \frac{y}{L} \right\}.$$

- (iv) *If $L > (\lambda_\omega)^{-1/2}$, then $-JS''_\omega(\tilde{\varphi}_\omega)$ has a positive eigenvalue and the number of eigenvalue of $-JS''_\omega(\tilde{\varphi}_\omega)$ with a positive real part is finite.*

Here, $\operatorname{Span}\{v_1, \dots, v_n\}$ is the real vector space spanned by vectors v_1, \dots, v_n .

Proof. We define

$$S(a)u = \begin{pmatrix} L_\omega^+ + a^2 & 0 \\ 0 & L_\omega^- + a^2 \end{pmatrix} \begin{pmatrix} \operatorname{Re} u \\ \operatorname{Im} u \end{pmatrix},$$

where $u \in H^1(\mathbb{R})$, $L_\omega^+ = -\partial_x^2 + \omega + V - p|\varphi_\omega|^{p-1}$ and $L_\omega^- = -\partial_x^2 + \omega + V - |\varphi_\omega|^{p-1}$. Then, for $u \in H^1(\mathbb{R} \times \mathbb{T}_L)$

$$S''_\omega(\tilde{\varphi}_\omega)u(x, y) = \sum_{n=-\infty}^{\infty} S\left(\frac{n}{L}\right) \vec{u}_n(x) e^{\frac{iny}{L}},$$

where

$$u(x, y) = \begin{pmatrix} \operatorname{Re} u \\ \operatorname{Im} u \end{pmatrix} = \sum_{n=-\infty}^{\infty} \begin{pmatrix} u_{R,n} \\ u_{I,n} \end{pmatrix} e^{\frac{iny}{L}} = \sum_{n=-\infty}^{\infty} \vec{u}_n(x) e^{\frac{iny}{L}}, \quad u \in H^1(\mathbb{R} \times \mathbb{T}_L).$$

Therefore, $-JS''_\omega(\tilde{\varphi}_\omega)$ has an eigenvalue λ if and only if there exists $n \in \mathbb{Z}$ such that $-JS(n/L)$ has the eigenvalue λ .

By Proposition 1.1, $S(a)$ has no negative eigenvalues for $a \geq (\lambda_\omega)^{1/2}$. By Theorem 3.1 in [17], the number of eigenvalues of $-JS(a)$ with the positive real part is less than or equal to the number of negative eigenvalues of $S(a)$. Thus, for $a \geq (\lambda_\omega)^{1/2}$, $JS(a)$ has no eigenvalues with the positive real part. (i) follows this. Moreover, the number of eigenvalues of $-JS''_\omega(\tilde{\varphi}_\omega)$ with the positive real part is less than $1 + 2L/(\lambda_\omega)^{1/2}$. Since the kernel of $S(a)$ is trivial for $a > (\lambda_\omega)^{1/2}$, the kernel of $-JS(a)$ is trivial for $a > (\lambda_\omega)^{1/2}$. Then the kernel of $-JS(0)$ is spanned by $i\tilde{\varphi}_\omega$. Therefore, for $a > (\lambda_\omega)^{1/2}$, the kernel of $-JS''_\omega(\tilde{\varphi})$ is spanned by $i\tilde{\varphi}_\omega$ and (ii) is verified.

The kernel of $L_\omega^+ + \lambda_\omega$ is spanned by ψ_ω and $L_\omega^- + \lambda_\omega$ do not has zero eigenvalue. Hence, the kernel of $S''_\omega(\tilde{\varphi}_\omega)$ with $L = (\lambda_\omega)^{-1/2}$ is spanned by $i\tilde{\varphi}_\omega$, $\psi_\omega \cos \frac{y}{L}$ and $\psi_\omega \sin \frac{y}{L}$. This is (iii).

Let

$$M(v, a, \lambda) = S(a)(\psi_\omega + v) + J^{-1}\lambda(\psi_\omega + v),$$

where $v \in \{w \in H^2(\mathbb{R}) \mid \langle v, \psi_\omega \rangle_{L^2(\mathbb{R})} = 0\} =: (\psi_\omega)^\perp$ and $a, \lambda \in \mathbb{R}$. Then, M is a C^∞ function with

$$M(0, (\lambda_\omega)^{1/2}, 0) = 0.$$

Since

$$\left. \frac{\partial M}{\partial(v, a)} \right|_{(v, a, \lambda) = (0, (\lambda_\omega)^{1/2}, 0)} (w, \mu) = 2(\lambda_\omega)^{1/2} \mu \psi_\omega + S((\lambda_\omega)^{1/2})w,$$

by the implicit function theorem, there exist $a(\lambda) \in \mathbb{R}$ and $v(\lambda) \in H^2(\mathbb{R})$ such that $a(\lambda), v(\lambda)$ are the C^∞ functions, where $a(0) = (\lambda_\omega)^{1/2}$, $v(0) = 0$ and $M(v(\lambda), a(\lambda), \lambda) = 0$. Then, we have

$$-JS(a(\lambda))(\psi_\omega + v(\lambda)) = \lambda(\psi_\omega + v(\lambda))$$

for sufficiently small $|\lambda|$. Differentiating with respect to λ , we obtain

$$\frac{\partial M}{\partial \lambda}(v(\lambda), a(\lambda), \lambda) = 2a(\lambda)a'(\lambda)(\psi_\omega + v(\lambda)) + S(a(\lambda))v'(\lambda) + J^{-1}(\psi_\omega + v(\lambda) + \lambda v(\lambda)) = 0.$$

Since $v(0) = 0$, we have

$$\langle 2a(0)a'(0)\psi_\omega, \psi_\omega \rangle_{L^2(\mathbb{R})} = 0,$$

and $a'(0) = 0$. Therefore, we have $S((\lambda_\omega)^{1/2})v'(0) = -J^{-1}\psi_\omega$. Since $v'(0) \in (\psi_\omega)^\perp$ and

$$\frac{\partial^2 M}{\partial \lambda^2}(v(\lambda), a(\lambda), \lambda)|_{\lambda=0} = 2a(0)a''(0)\psi_\omega + S((\lambda_\omega)^{1/2})v''(0) + J^{-1}v'(0) = 0,$$

$$a''(0) = -\frac{\langle S((\lambda_\omega)^{1/2})v'(0), v'(0) \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})}}{2(\lambda_\omega)^{1/2}} < 0.$$

From the proof of (i), for $a > (\lambda_\omega)^{1/2}$, $-JS(a)$ has no positive eigenvalues. Hence, for sufficiently small $\varepsilon > 0$ the function $a(\lambda)$ on $(0, \varepsilon)$ has the inverse function $\lambda(a)$ on $(a(\varepsilon), (\lambda_\omega)^{1/2})$ and $a(\varepsilon) < (\lambda_\omega)^{1/2}$. Namely, $-JS(a)$ has the simple positive eigenvalue on $(a(\varepsilon), (\lambda_\omega)^{1/2})$.

Let

$$a_0 = \inf\{a > 0 \mid -JS(b) \text{ has a simple positive eigenvalue for } a < b < (\lambda_\omega)^{1/2}\},$$

and for $a \in (a_0, (\lambda_\omega)^{1/2})$ the value $\lambda(a)$ be the positive eigenvalue of $-JS(a)$. We assume $a_0 > 0$. By the perturbation theory, there exists $\{a_n\}_{n=1}^\infty \subset (a_0, (\lambda_\omega)^{1/2})$ such that $a_n \rightarrow a_0$ and

$$\lim_{n \rightarrow \infty} \lambda(a_n) = 0$$

or

$$\lim_{n \rightarrow \infty} \lambda(a_n) = \infty.$$

Since there exists $C > 0$ such that $|\langle -JS(a)u, u \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})}| \leq C\|u\|_{L^2(\mathbb{R})}$ for $a \in \mathbb{R}$, $\lambda(a)$ is bounded. Therefore,

$$\lim_{n \rightarrow \infty} \lambda(a_n) = 0.$$

Then, there exists $\{c_n\}_{n=1}^\infty$ such that $\|v_n\|_{H^1(\mathbb{R})} = 1$ and $-JS(a_n)v_n = \lambda(a_n)v_n$. Here,

$$S(a_0)v_n = (a_0^2 - a_n^2)v_n - J^{-1}\lambda(a_n)v_n.$$

Since $S(a_0)$ is invertible and $(S(a_0))^{-1}$ is bounded,

$$v_n = (S(a_0))^{-1}((a_0^2 - a_n^2)v_n - J^{-1}\lambda(a_n)v_n) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

This is contradiction. Therefore, $a_0 = 0$. □

Next we show the coerciveness of L_ω^+ on a function space which follows the proof of Theorem 3.3 in [10].

Lemma 2.2. *There exist $\omega_{*,0} > \lambda_*$ and $k_0 > 0$ such that for $\lambda_* < \omega < \omega_{*,0}$ and $u \in H^1(\mathbb{R}, \mathbb{R})$ with $\langle \varphi_\omega, u \rangle_{L^2(\mathbb{R})} = 0$,*

$$\langle L_\omega^+ u, u \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} \geq k_0 \|u\|_{H^1(\mathbb{R})}^2,$$

where $\langle u, v \rangle_{L^2(\mathbb{R})} = \text{Re} \int_{\mathbb{R}} u \bar{v} dx$.

Proof. Let $u \in H^1(\mathbb{R}, \mathbb{R})$ with $\langle \varphi_\omega, u \rangle_{L^2(\mathbb{R})} = 0$. We decompose $u = a\psi_* + u_\perp$, where $a = \langle u, \psi_* \rangle_{L^2(\mathbb{R})}$ and $\langle \psi_*, u_\perp \rangle_{L^2(\mathbb{R})} = 0$. From the spectrum of $-\partial_x^2 + V + \lambda_*$, there exists $k > 0$ such that

$$\langle (-\partial_x^2 + V + \lambda_*)u_\perp, u_\perp \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} \geq k \|u_\perp\|_{L^2(\mathbb{R})}^2.$$

By $\|V\|_{L^\infty} < \infty$, we have for $\varepsilon > 0$

$$\langle (-\partial_x^2 + V + \lambda_*)u_\perp, u_\perp \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} \geq (k - \varepsilon(\|V\|_{L^\infty} - \lambda_*)) \|u_\perp\|_{L^2(\mathbb{R})}^2 + \varepsilon \|\partial_x u_\perp\|_{L^2(\mathbb{R})}^2.$$

Therefore, there exists $k' > 0$ such that

$$\langle (-\partial_x^2 + V + \lambda_*)u_\perp, u_\perp \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} \geq k' \|u_\perp\|_{H^1(\mathbb{R})}^2.$$

By the assumption $\langle \varphi_\omega, u \rangle_{L^2(\mathbb{R})} = 0$, we have

$$a = -\frac{\langle \varphi_\omega, u_\perp \rangle_{L^2(\mathbb{R})}}{\langle \varphi_\omega, \psi_* \rangle_{L^2(\mathbb{R})}}.$$

Then

$$\begin{aligned}
\langle L_\omega^+ u, u \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} &= \langle (-\partial_x^2 + V + \lambda_*) u_\perp, u_\perp \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} + \lambda_\omega a^2 \\
&\quad + \langle (\omega - \lambda_* - p|\varphi_\omega|^{p-1}) u_\perp, u_\perp \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} - 2\lambda_\omega \frac{\langle \varphi_\omega, u_\perp \rangle_{L^2(\mathbb{R})}^2}{\langle \varphi_\omega, \psi_* \rangle_{L^2(\mathbb{R})}} + o(\|u_\perp\|_{L^2(\mathbb{R})}^2) \\
&\geq k' \|u_\perp\|_{H^1}^2 + \lambda_\omega a^2 + \langle (\omega - \lambda_* - p|\varphi_\omega|^{p-1}) u_\perp, u_\perp \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} \\
&\quad - 2\lambda_\omega \frac{\|u_\perp\|_{L^2(\mathbb{R})}^2 \|\varphi_\omega\|_{L^2(\mathbb{R})}^2}{\langle \varphi_\omega, \psi_* \rangle_{L^2(\mathbb{R})}^2} + o(\|u_\perp\|_{L^2(\mathbb{R})}^2).
\end{aligned}$$

If $|\omega - \lambda_*|$ is sufficiently small, then we obtain the conclusion. \square

3 Proof of Theorem 1.3

In this section, we prove Theorem 1.3. The proof of Theorem 1.3 is similar to the proof of Theorem 1.5 in [25]. We write the detail of the proof of Theorem 1.3 for readers.

3.1 Proof of (i) of Theorem 1.3

In this subsection, we assume $0 < L < (\lambda_\omega)^{-1/2}$. The proof of (i) of Theorem 1.3 follows Section 3.1 in [25].

The following proposition follows Grillakis-Shatah-Strauss [10] or Colin-Ohta [4](see [3]).

Proposition 3.1. *Let $e^{i\omega t}\varphi$ be a standing wave of (1.1). Assume that there exists a constant $\delta > 0$ such that $\langle S_\omega''(\varphi)u, u \rangle_{H^{-1}, H^1} \geq \delta \|u\|_{H^1}^2$ for all $u \in H^1(\mathbb{R} \times \mathbb{T}_L)$ satisfying $\langle \varphi, u \rangle_{L^2} = \langle J\varphi, u \rangle_{L^2} = 0$. Then, the standing wave $e^{i\omega t}\varphi$ is stable.*

Let $u \in H^1(\mathbb{R} \times \mathbb{T}_L)$ satisfy $\langle \tilde{\varphi}_\omega, u \rangle_{L^2} = \langle J\tilde{\varphi}_\omega, u \rangle_{L^2} = 0$. Then, We have

$$\langle S_\omega''(\tilde{\varphi}_\omega)u, u \rangle_{H^{-1}, H^1} = \sum_{n \in \mathbb{Z}} \langle S(n/L)u_n, u_n \rangle_{H^{-1}, H^1},$$

where

$$u(x, y) = \sum_{n \in \mathbb{Z}} u_n(x) e^{\frac{iny}{L}}.$$

Since L_ω^- and $L_\omega^+ + \lambda_\omega$ are nonnegative, there exists $c > 0$ such that

$$\langle S(n/L)v, v \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} \geq c \|v\|_{H^1(\mathbb{R})}^2$$

for $n \in \mathbb{Z} \setminus \{0\}$ and $v \in H^1(\mathbb{R})$. By Proposition 1.1 and Lemma 2.2, there exists $c' > 0$ such that for $v \in H^1(\mathbb{R})$ with $\langle \tilde{\varphi}_\omega, v \rangle_{L^2} = 2\pi L \int_{\mathbb{R}} \varphi_\omega(\operatorname{Re} v) dx = 0$ and $\langle J\tilde{\varphi}_\omega, v \rangle_{L^2} = 2\pi L \int_{\mathbb{R}} \varphi_\omega(\operatorname{Im} v) dx = 0$

$$\langle \mathbb{L}_\omega^+(\operatorname{Re} v), \operatorname{Re} v \rangle_{H^{-1}, H^1} \geq c' \|\operatorname{Re} v\|_{H^1}^2,$$

$$\langle \mathbb{L}_\omega^-(\operatorname{Im} v), \operatorname{Im} v \rangle_{H^{-1}, H^1} \geq c' \|\operatorname{Im} v\|_{H^1}^2.$$

Therefore, (i) of Theorem 1.1 follows from Proposition 3.1.

3.2 Proof of (ii) of Theorem 1.3

The proof of (ii) of Theorem 1.3 follows Section 3.2 in [25].

In this subsection, we assume $L > (\lambda_\omega)^{1/2}$. We define

$$\mu_* = \max\{\lambda > 0 \mid \lambda \in \sigma(-JS''(\tilde{\varphi}_\omega))\},$$

where $\sigma(-JS''(\tilde{\varphi}_\omega))$ is the spectrum of $-JS''(\tilde{\varphi}_\omega)$. Then, there exist $k_0 \in \mathbb{Z}$ and $\chi \in H^1(\mathbb{R} \times \mathbb{T}_L)$ such that $\|\chi\|_{L^2} = 1$, χ is eigenfunction of $-JS''_\omega(\tilde{\varphi}_\omega)$ corresponding to μ_* and

$$\chi(x, y) = \chi_1(x)e^{\frac{ik_0y}{L}} + \chi_2(x)e^{-\frac{ik_0y}{L}},$$

where $\chi_1, \chi_2 \in H^2(\mathbb{R})$. We define the orthogonal projection $P_{\leq k}$ as

$$P_{\leq k}u(x, y) = \sum_{n=-k}^k u_n(x)e^{\frac{iny}{L}}, \quad (x, y) \in \mathbb{R} \times \mathbb{T}_L,$$

where

$$u(x, y) = \sum_{n=-\infty}^{\infty} u_n(x)e^{\frac{iny}{L}}. \quad (x, y) \in \mathbb{R} \times \mathbb{T}_L.$$

A function $u(t)$ is a solution of (1.1) if and only if $v(t)$ is a solution of the equation

$$\partial_t v = -J(S''_\omega(\tilde{\varphi}_\omega)v + g(v)), \quad (3.1)$$

where $u(t) = e^{i\omega t}(\tilde{\varphi}_\omega + v(t))$,

$$g(v) = \begin{pmatrix} |v + \tilde{\varphi}_\omega|^{p-1}(v_R + \tilde{\varphi}_\omega) - p|\tilde{\varphi}_\omega|^{p-1}v_R - |\tilde{\varphi}_\omega|^{p-1}\tilde{\varphi}_\omega \\ |v + \tilde{\varphi}_\omega|^{p-1}v_I - |\tilde{\varphi}_\omega|^{p-1}v_I \end{pmatrix},$$

and $v_R = \operatorname{Re} v$ and $v_I = \operatorname{Im} v$. We define $u_\delta(t)$ as the solution of (1.1) with the initial data $\tilde{\varphi}_\omega + \delta\chi$ and $v_\delta(t)$ as the solution of (3.1) with the data $\delta\chi$. Then, we have that $u_\delta(t) = e^{i\omega t}(\tilde{\varphi}_\omega + v_\delta(t))$.

We show the estimate of nonlinear term in the following lemma which follows Lemma 2.4 of [8].

Lemma 3.2. *There exists $C > 0$ such that*

$$\|g(v)\|_{L^2} \leq \begin{cases} C\|v\|_{H^1}^p, & 1 < p \leq 2, \\ C(\|v\|_{H^1}^2 + \|v\|_{H^1}^p), & 2 < p. \end{cases}$$

Proof. We have

$$\| |a|^{p-1} - |b|^{p-1} \| \leq \begin{cases} |a - b|^{p-1}, & 1 < p \leq 2, \\ p(|a|^{p-2} + |b|^{p-2})|a - b|, & 2 < p. \end{cases}$$

Since

$$g(v(x, y)) = \int_0^1 (|\theta v(x, y) + \tilde{\varphi}_\omega(x, y)|^{p-1} - |\tilde{\varphi}_\omega(x, y)|^{p-1})v(x, y)d\theta,$$

$$\begin{aligned}
\|g(v)\|_{L^2} &\leq \left\| \int_0^1 (|\theta v + \tilde{\varphi}_\omega|^{p-1} - |\tilde{\varphi}_\omega|^{p-1})v d\theta \right\|_{L^2} \\
&\leq \int_0^1 \left\| |\theta v + \tilde{\varphi}_\omega|^{p-1} - |\tilde{\varphi}_\omega|^{p-1} \right\|_{L^4} \|v\|_{L^4} d\theta \\
&\leq \begin{cases} C\|v\|_{H^1}^p, & 1 < p \leq 2, \\ C(\|v\|_{H^1}^2 + \|v\|_{H^1}^p), & 2 < p. \end{cases}
\end{aligned}$$

□

In the following lemma, we estimate the low frequency part of the semi-group.

Lemma 3.3. *For a positive integer k and $\varepsilon > 0$, there exists $C_{k,\varepsilon} > 0$ such that*

$$\left\| e^{-tJS''_\omega(\tilde{\varphi}_\omega)} P_{\leq k} v \right\|_{L^2} \leq C_{k,\varepsilon} e^{(\mu_* + \varepsilon)t} \|v\|_{L^2}, \quad t > 0, v \in L^2(\mathbb{R} \times \mathbb{T}_L).$$

Proof. By the definition of $S(a)$, we have

$$-JS(a) = \begin{pmatrix} 0 & -\partial_x^2 + \omega + a^2 + V - |\tilde{\varphi}_\omega|^{p-1} \\ \partial_x^2 - \omega - a^2 - V + p|\tilde{\varphi}_\omega|^{p-1} & 0 \end{pmatrix}.$$

Using the exponential decay rates of V and $\tilde{\varphi}_\omega$ and applying the argument for the proof of Proposition [5] and Lemma 6 in [6], we obtain

$$\sigma(e^{-JS(a)}) = e^{\sigma(-JS(a))}.$$

By the definition of μ_* , we have that the spectral radius of $e^{-JS(n/L)}$ is less than or equal to e^{μ_*} for $n \in \mathbb{Z}$. Therefore, by Lemma 2 and Lemma 3 in [23] we have

$$\left\| e^{-tJS(n/L)} v \right\|_{L^2(\mathbb{R})} \leq C_{n,\varepsilon} e^{(\mu_* + \varepsilon)t} \|v\|_{L^2(\mathbb{R})}, \quad t > 0, n \in \mathbb{Z}, v \in L^2(\mathbb{R}).$$

Hence, for $t > 0$ and $v \in L^2(\mathbb{R} \times \mathbb{T}_L)$,

$$\left\| e^{-tJS''_\omega(\tilde{\varphi}_\omega)} P_{\leq k} v \right\|_{L^2} \leq \left\| \sum_{n=-k}^k e^{-tJS(n/L)} v_n e^{\frac{iny}{L}} \right\|_{L^2} \leq C_{k,\varepsilon} e^{(\mu_* + \varepsilon)t} \|v\|_{L^2},$$

where

$$v(x, y) = \sum_{n \in \mathbb{Z}} v_n(x) e^{\frac{iny}{L}}.$$

□

In the following lemma, we estimate the high frequency part of $v_\delta(t)$.

Lemma 3.4. *There exist a positive integer K_0 and $C > 0$ such that for $\delta > 0$ and $t > 0$*

$$\|v_\delta(t)\|_{H^1} \leq C\|P_{\leq K_0} v_\delta(t)\|_{L^2} + o(\delta) + o(\|v_\delta(t)\|_{H^1}).$$

Proof. By the Taylor expansion we have that for $v \in H^1(\mathbb{R} \times \mathbb{T}_L)$

$$S_\omega(\tilde{\varphi}_\omega + v) = S_\omega(\tilde{\varphi}_\omega) + \langle S'_\omega(\tilde{\varphi}_\omega), v \rangle_{H^{-1}, H^1} + \frac{1}{2} \langle S''_\omega(\tilde{\varphi}_\omega)v, v \rangle_{H^{-1}, H^1} + o(\|v\|_{H^1}^2).$$

Since S_ω is conservation law, we have $S_\omega(\tilde{\varphi}_\omega + \delta\chi) = S_\omega(\tilde{\varphi}_\omega + v_\delta(t))$ for $t \geq 0$. Using $S'(\tilde{\varphi}_\omega) = 0$ and

$$\langle S''_\omega(\tilde{\varphi}_\omega)\chi, \chi \rangle_{H^{-1}, H^1} = \langle -JS''_\omega(\tilde{\varphi}_\omega)\chi, J^{-1}\chi \rangle_{H^{-1}, H^1} = \langle \mu_*\chi, J^{-1}\chi \rangle_{L^2} = 0,$$

we have

$$\langle S''_\omega(\tilde{\varphi}_\omega)v_\delta(t), v_\delta(t) \rangle_{H^{-1}, H^1} = o(\|v_\delta(t)\|_{H^1}^2) + o(\delta^2).$$

We define K_0 as the integer part of $1 + L(\lambda_\omega)^{1/2}$. Since $S(a)$ is positive for $a > (\lambda_\omega)^{1/2}$, we obtain $S''_\omega(\tilde{\varphi}_\omega)(I - P_{\leq K_0})$ is positive. By the definition of $S(a)$ there exist $c, C > 0$ such that

$$\langle S(a)v, v \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})} \geq c\|v\|_{H^1(\mathbb{R})}^2 - C\|v\|_{L^2(\mathbb{R})}^2$$

for $v \in H^1(\mathbb{R})$ and $a \in \mathbb{R}$. Thus,

$$\begin{aligned} \|v_\delta(t)\|_{H^1}^2 &= \|P_{\leq K_0}v_\delta(t)\|_{H^1}^2 + \|(I - P_{\leq K_0})v_\delta(t)\|_{H^1}^2 \\ &\leq C'\langle S''_\omega(\tilde{\varphi}_\omega)(I - P_{\leq K_0})v_\delta(t), (I - P_{\leq K_0})v_\delta(t) \rangle_{H^{-1}, H^1} \\ &\quad + C'\langle S''_\omega(\tilde{\varphi}_\omega)P_{\leq K_0}v_\delta(t), P_{\leq K_0}v_\delta(t) \rangle_{H^{-1}, H^1} + C''\|P_{\leq K_0}v_\delta(t)\|_{H^1}^2 \\ &\leq C''\|v_\delta(t)\|_{L^2}^2 + o(\delta^2) + o(\|v_\delta(t)\|_{H^1}^2). \end{aligned}$$

□

Let $\varepsilon_0 = \min\{(p-1)\mu_*/2, \mu_*/2\}$. By Lemma 3.2, Lemma 3.3 and Lemma 3.4, we obtain that

$$\begin{aligned} \|v_\delta(t)\|_{H^1} &\leq C\delta e^{\mu_*t} \|\chi\|_{L^2} + C \int_0^t \|e^{-(t-s)JS_\omega(\tilde{\varphi}_\omega)} P_{\leq K_0}g(v_\delta(s))\|_{L^2} ds + o(\delta) + o(\|v_\delta(t)\|_{H^1}) \\ &\leq C\delta e^{\mu_*t} + C \int_0^t e^{(1+\varepsilon_0)\mu_*(t-s)} (\|v_\delta(s)\|_{H^1}^2 + \|v_\delta(s)\|_{H^1}^p) ds + o(\delta) + o(\|v_\delta(t)\|_{H^1}). \end{aligned}$$

There exists $C_0 > 0$ such that for small $\delta > 0$ and $\varepsilon_1 > 0$

$$\|v_\delta(t)\|_{H^1} \leq C_0 e^{\mu_*t}, \quad \text{for } t \in [0, T_{\varepsilon_1, \delta}],$$

where

$$T_{\varepsilon_1, \delta} = \frac{\log(\varepsilon_1/\delta)}{\mu_*}.$$

Then,

$$\begin{aligned} |\langle \chi, v_\delta(T_{\varepsilon_1, \delta}) \rangle_{L^2}| &= \left| \delta e^{\mu_*T_{\varepsilon_1, \delta}} + \int_0^{T_{\varepsilon_1, \delta}} \langle \chi, -Je^{-(T_{\varepsilon_1, \delta}-s)S''_\omega(\tilde{\varphi}_\omega)}g(v_\delta(s)) \rangle_{L^2} ds \right| \\ &\geq \varepsilon_1 - C \int_0^{T_{\varepsilon_1, \delta}} e^{(T_{\varepsilon_1, \delta}-s)\mu_*} (\|v_\delta(s)\|_{H^1}^2 + \|v_\delta(s)\|_{H^1}^p) ds \\ &\geq \varepsilon_1 - C\varepsilon_1^{\min\{p, 2\}}. \end{aligned}$$

Since $P_{\leq 0}\tilde{\varphi}_\omega = \tilde{\varphi}_\omega$, there exists $\varepsilon_1 > 0$ such that for $\varepsilon_1 > \delta > 0$ and $\theta \in \mathbb{R}$

$$\begin{aligned} \|u_\delta(T_{\varepsilon_1, \delta}) - e^{i\theta}\tilde{\varphi}_\omega\|_{L^2} &\geq \|(I - P_{\leq 0})(u_\delta(T_{\varepsilon_1, \delta}) - e^{i\theta}\tilde{\varphi}_\omega)\|_{L^2} \\ &= \|(I - P_{\leq 0})e^{-i\omega T_{\varepsilon_1, \delta}}u_\delta(T_{\varepsilon_1, \delta})\|_{L^2} \\ &= \|(I - P_{\leq 0})(e^{-i\omega T_{\varepsilon_1, \delta}}u_\delta(T_{\varepsilon_1, \delta}) - \tilde{\varphi}_\omega)\|_{L^2}. \end{aligned}$$

By the definition of χ we have

$$\|(P_{\leq k_0} - P_{\leq k_0-1})v\|_{L^2} \geq |\langle \chi, v \rangle_{L^2}|, \quad \text{for } v \in L^2(\mathbb{R} \times \mathbb{T}_L).$$

Therefore,

$$\|(I - P_{\leq 0})(e^{-i\omega T_{\varepsilon_1, \delta}}u_\delta(T_{\varepsilon_1, \delta}) - \tilde{\varphi}_\omega)\|_{L^2} \geq |\langle \chi, v_\delta(T_{\varepsilon_1, \delta}) \rangle_{L^2}| \geq \frac{\varepsilon_1}{2}.$$

This implies that the standing wave $e^{i\omega t}\tilde{\varphi}_\omega$ is unstable.

4 Proof of Theorem 1.4

In this section, we assume $L = (\lambda_{\omega_0})^{-1/2}$ for $0 < \omega_0 - \lambda_* \ll 1$.

The following lemma follows Proposition 1.1.

Lemma 4.1.

$$\begin{aligned} \varphi_\omega &= \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}}(\omega - \lambda_*)^{\frac{1}{p-1}}\psi_* + \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}}(\omega - \lambda_*)^{\frac{1}{p-1}+1}(P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1}\psi_{*,p} \\ &\quad + o((\omega - \lambda_*)^{\frac{1}{p-1}+1}), \\ \partial_\omega \varphi_\omega &= \frac{\|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}}}{p-1}(\omega - \lambda_*)^{\frac{1}{p-1}-1}\psi_* + \frac{p\|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}}}{p-1}(\omega - \lambda_*)^{\frac{1}{p-1}}(P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1}\psi_{*,p} \\ &\quad + o((\omega - \lambda_*)^{\frac{1}{p-1}}). \end{aligned} \tag{4.1}$$

where $\psi_{*,p} = \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)}\psi_*^p - \psi_*$, $H_a = -\partial_x^2 + a + V$ for $a \in \mathbb{R}$ and P_\perp^1 is the orthogonal projection onto $(\psi_*)^\perp = \{u \in L^2(\mathbb{R}) \mid \langle u, \psi_* \rangle_{L^2(\mathbb{R})} = 0\}$.

Proof. Let

$$\varphi_{\omega,0} = (\omega - \lambda_*)^{-\frac{1}{p-1}}\varphi_\omega = \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}}\psi_* + \tilde{r}.$$

By the bifurcation argument, $\langle \psi_*, \tilde{r}(\omega) \rangle_{L^2(\mathbb{R})} = 0$. Since φ_ω is C^1 with respect to ω and

$$(-\partial_x^2 + \omega + V)\varphi_{\omega,0} - |\varphi_{\omega,0}|^{p-1}\varphi_{\omega,0} = 0,$$

we have

$$\begin{aligned} 0 &= \partial_\omega((-\partial_x^2 + \omega + V)\varphi_{\omega,0} - |\varphi_{\omega,0}|^{p-1}\varphi_{\omega,0}) \\ &= \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}}\psi_* + \tilde{r} + H_\omega \partial_\omega \tilde{r} - \left| \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}}\psi_* + \tilde{r} \right|^{p-1} \left(\|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}}\psi_* + \tilde{r} \right) \\ &\quad - p(\omega - \lambda_*) \left| \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}}\psi_* + \tilde{r} \right|^{p-1} \partial_\omega \tilde{r}. \end{aligned}$$

Transverse instability

From $\langle \psi_*, \tilde{r}(\omega) \rangle_{L^2(\mathbb{R})} = \langle \psi_*, \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)} \psi_*^p - \psi_* \rangle_{L^2(\mathbb{R})} = 0$, we obtain

$$\partial_\omega \tilde{r} = \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{p+1}{p-1}} (P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} (\|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)} \psi_*^p - \psi_*) + o(1).$$

□

In the following lemma, we obtain the derivative of the eigenvalue λ_ω .

Lemma 4.2. *Let $p \geq 2$. Then,*

$$\psi_\omega = \psi_* + p(\omega - \lambda_*) (P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} \psi_{*,p} + O((\omega - \lambda_*)^2),$$

$$\begin{aligned} \lambda_\omega &= (p-1)(\omega - \lambda_*) + p(2p-1)(\omega - \lambda_*)^2 \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)} \int_{\mathbb{R}} \psi_*^p (P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} \psi_{*,p} dx \\ &\quad + o((\omega - \lambda_*)^2). \end{aligned}$$

Proof. There exists $\delta_0 > 0$ such that $\{z \in \mathbb{C} \mid |z + \lambda_*| < 2\delta_0\} \cap \sigma(-\Delta + V) = \{-\lambda_*\}$. Let $\Gamma = \{z \in \mathbb{C} \mid |z| = \delta_0\}$ be a simple closed curve and projections

$$P_\omega = \frac{1}{2\pi i} \int_\Gamma (L_\omega^+ - z)^{-1} dz.$$

Then, for $\omega > \lambda_*$ with $0 < \omega - \lambda_* \ll 1$,

$$P_\omega u = \langle u, \psi_\omega \rangle_{L^2(\mathbb{R})} \psi_\omega.$$

Since $p \geq 2$, L_ω^+ is C^1 with respect to ω . Therefore, the projection P_ω is also C^1 . For $\omega, \omega' > \lambda_*$, $(\langle \psi_{\omega'}, \psi_\omega \rangle_{L^2(\mathbb{R})})^2 - 1 = \langle P_{\omega'} \psi_\omega, \psi_\omega \rangle_{L^2(\mathbb{R})} - 1 = o(1)$ as $|\omega' - \omega| \rightarrow 0$. For $\omega > \lambda_*$,

$$\psi_{\omega'} - \psi_\omega = \frac{P_\omega(\psi_\omega + \psi_{\omega'}) - P_{\omega'}(\psi_\omega + \psi_{\omega'})}{1 + \langle \psi_{\omega'}, \psi_\omega \rangle_{L^2(\mathbb{R})}}.$$

Thus, ψ_ω is C^1 with respect to ω . Let $\varphi_{\omega,0} = (\omega - \lambda_*)^{-\frac{1}{p-1}} \varphi_\omega$. Since $L_\omega^+ \psi_\omega = -\lambda_\omega \psi_\omega$, we have

$$-\lambda_\omega = \langle L_\omega^+ \psi_\omega, \psi_\omega \rangle_{H^{-1}(\mathbb{R}), H^1(\mathbb{R})}.$$

Therefore,

$$\begin{aligned} -\frac{d}{d\omega} \lambda_\omega &= 1 - p \int_{\mathbb{R}} (\varphi_{\omega,0})^{p-1} (\psi_\omega)^2 dx - p \int_{\mathbb{R}} (p-1)(\omega - \lambda_*) (\varphi_{\omega,0})^{p-2} (\partial_\omega \varphi_{\omega,0}) (\psi_\omega)^2 dx \\ &= 1 - p + O(\omega - \lambda_*). \end{aligned} \tag{4.2}$$

Since

$$(-\partial_x^2 + \omega + \lambda_\omega + V - p|\varphi_{\omega,0}|^{p-1})\psi_\omega = 0,$$

we have

$$0 = (1 + \partial_\omega \lambda_\omega - p|\varphi_{\omega,0}|^{p-1} - p(p-1)(\omega - \lambda_*)|\varphi_{\omega,0}|^{p-2} \partial_\omega \varphi_{\omega,0})\psi_\omega + L_\omega^+ \partial_\omega \psi_\omega.$$

Therefore,

$$\partial_\omega \psi_\omega = p(P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} (\|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)} \psi_*^p - \psi_*) + O(\omega - \lambda_*).$$

By (4.2) and lemma 4.1, we obtain

$$\begin{aligned} \frac{d^2}{d\omega^2} \lambda_\omega &= 2p(p-1) \int_{\mathbb{R}} (\varphi_{\omega,0})^{p-2} (\partial_\omega \varphi_{\omega,0}) (\psi_\omega)^2 dx + 2p \int_{\mathbb{R}} (\varphi_{\omega,0})^{p-1} \psi_\omega \partial_\omega \psi_\omega dx \\ &\quad + p(p-1) \int_{\mathbb{R}} (\omega - \lambda_*) (\varphi_{\omega,0})^{p-3} (\partial_\omega \varphi_{\omega,0})^2 (\psi_\omega)^2 dx \\ &\quad + p(p-1) \int_{\mathbb{R}} (\omega - \lambda_*) (\varphi_{\omega,0})^{p-2} (\partial_\omega^2 \varphi_{\omega,0}) (\psi_\omega)^2 dx \\ &\quad + 2p(p-1) \int_{\mathbb{R}} (\omega - \lambda_*) (\varphi_{\omega,0})^{p-2} (\partial_\omega \varphi_{\omega,0}) \psi_\omega \partial_\omega \psi_\omega dx \\ &= 2p(2p-1) \int_{\mathbb{R}} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)} \psi_*^p (P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} (\|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)} \psi_*^p - \psi_*) dx + o(1). \end{aligned}$$

□

The following corollary follows Lemma 4.2.

Corollary 4.3. *There exists $\omega_{*,1} > \lambda_*$ such that for $\lambda_* < \omega < \omega_{*,1}$, $\lambda_\omega > 0$. Moreover, if $\lambda_* < \omega_0 < \omega_{*,1}$, then the followings are hold.*

- (i) *If $\omega_0 < \omega < \omega_{*,1}$, then \mathbb{L}_ω^+ has exactly two negative eigenvalue and no kernel.*
- (ii) *If $\lambda_* < \omega < \omega_0$, then \mathbb{L}_ω^+ has exactly one negative eigenvalue and no kernel.*

Applying Lyapunov-Schmidt decomposition and Crandall-Rabinowitz Transversality in [12], we show $\tilde{\varphi}_{\omega_0}$ is a bifurcation point. In this paper, we only write the sketch of the proof of the following proposition (see the proof of Theorem 4 in [12] or Proposition 1 in [26] for the detail of the proof of the following proposition).

Proposition 4.4. *Let $p \geq 2$ and $\lambda_* < \omega_0 < \omega_{*,1}$. There exist $\delta > 0$ and $\phi_{\omega_0} \in C^2([-\delta, \delta], H^2)$ such that $\phi_{\omega_0}(a) > 0$,*

$$\begin{aligned} \phi_{\omega_0}(a)(x, y) &= \phi_{\omega_0}(a)(-x, y) = \phi_{\omega_0}(a)(x, -y), \quad (x, y) \in \mathbb{R} \times [-\pi L, \pi L], \\ -\Delta \phi_{\omega_0}(a) + \omega_{\omega_0}(a) \phi_{\omega_0}(a) + V \phi_{\omega_0}(a) - |\phi_{\omega_0}(a)|^{p-1} \phi_{\omega_0}(a) &= 0, \\ \phi_{\omega_0}(a) &= \tilde{\varphi}_{\omega_0} + a \psi_{\omega_0} \cos \frac{y}{L} + r_{\omega_0}(a), \\ \omega_{\omega_0}(a) &= \omega_0 + \frac{\omega_{\omega_0}''(0)}{2} a^2 + o(a^2), \end{aligned} \tag{4.3}$$

where $r_{\omega_0}(a) \perp \psi_{\omega_0} \cos \frac{y}{L}$, $\|r_{\omega_0}(a)\|_{H^2} = O(a^2)$,

$$\begin{aligned} \omega_{\omega_0}''(0) &= \frac{-p^2(p-1)^2}{\frac{d\lambda_\omega}{d\omega}|_{\omega=\omega_0} \|\psi_{\omega_0} \cos \frac{y}{L}\|_{L^2}^2} \langle (\tilde{\varphi}_{\omega_0})^{p-2} (\psi_{\omega_0} \cos \frac{y}{L})^2, \mathbb{L}_{\omega_0}^{-1} ((\tilde{\varphi}_{\omega_0})^{p-2} (\psi_{\omega_0} \cos \frac{y}{L})^2) \rangle_{L^2} \\ &\quad - \frac{p(p-1)(p-2)}{3 \frac{d\lambda_\omega}{d\omega}|_{\omega=\omega_0} \|\psi_{\omega_0} \cos \frac{y}{L}\|_{L^2}^2} \langle (\psi_{\omega_0} \cos \frac{y}{L})^2, (\tilde{\varphi}_{\omega_0})^{p-3} (\psi_{\omega_0} \cos \frac{y}{L})^2 \rangle_{L^2}, \end{aligned} \tag{4.4}$$

$$\lambda_2(a) = \frac{d\lambda_\omega}{d\omega}\Big|_{\omega=\omega_0} \omega''_{\omega_0}(0) a^2 + o(a^2), \quad (4.5)$$

and

$$\|\phi_{\omega_0}(a)\|_{L^2}^2 = \|\tilde{\varphi}_{\omega_0}\|_{L^2}^2 + \frac{R_{p,\omega_0}}{2} a^2 + o(a^2). \quad (4.6)$$

Here,

$$R_{p,\omega_0} = -2 \frac{d\lambda_\omega}{d\omega}\Big|_{\omega=\omega_0} \left\| \psi_{\omega_0} \cos \frac{y}{L} \right\|_{L^2}^2 + \omega''_{\omega_0}(0) \frac{d\|\tilde{\varphi}_\omega\|_{L^2}^2}{d\omega}\Big|_{\omega=\omega_0},$$

and $\lambda_2(a)$ is the second eigenvalue of $\mathbb{L}(a, \omega_0) = -\Delta + \omega_{\omega_0}(a) + V - |\phi_{\omega_0}(a)|^{p-1}$.

The sketch of the proof. Let F be the function from $H_{sym}^2(\mathbb{R} \times \mathbb{T}_L, \mathbb{R}) \rightarrow L_{sym}^2(\mathbb{R} \times \mathbb{T}_L, \mathbb{R})$ satisfying

$$F(\varphi, \omega) = -\Delta\varphi + \omega\varphi + V\varphi - |\varphi|^{p-1}\varphi,$$

where $L_{sym}^2(\mathbb{R} \times \mathbb{T}_L, \mathbb{R}) = \{u \in L^2(\mathbb{R} \times \mathbb{T}_L, \mathbb{R}) \mid u(x, y) = u(-x, y) = u(x, -y), (x, y) \in \mathbb{R} \times [-\pi L, \pi L]\}$, $H_{sym}^2(\mathbb{R} \times \mathbb{T}_L, \mathbb{R}) = H^2(\mathbb{R} \times \mathbb{T}_L) \cap L_{sym}^2(\mathbb{R} \times \mathbb{T}_L, \mathbb{R})$ and $L^2(\mathbb{R} \times \mathbb{T}_L, \mathbb{R})$ is the set of real valued L^2 -function on $\mathbb{R} \times \mathbb{T}_L$. Then, $\text{Ker}(\partial_\varphi F(\tilde{\varphi}_{\omega_0}, \omega_0))$ is spanned by $\psi_{\omega_0} \cos \frac{y}{L}$. Applying the Lyapunov-Schmidt decomposition, we obtain that there exists a function $h(\omega, a) \in H_{sym}^2(\mathbb{R} \times \mathbb{T}_L, \mathbb{R})$ such that

$$P_\perp F(\tilde{\varphi}_{\omega_0} + a\psi_{\omega_0} \cos \frac{y}{L} + h(\omega, a), \omega) = 0,$$

where P_\perp is the orthogonal projection onto $\{u \in L^2(\mathbb{R} \times \mathbb{T}_L, \mathbb{R}) \mid \langle u, \psi_{\omega_0} \cos \frac{y}{L} \rangle_{L^2} = 0\}$. Then, the problem $F(\tilde{\varphi}_{\omega_0} + a\psi_{\omega_0} \cos \frac{y}{L} + h(\omega, a), \omega) = 0$ is equivalent to the problem

$$F_{||}(\omega, a) = \langle F(\tilde{\varphi}_{\omega_0} + a\psi_{\omega_0} \cos \frac{y}{L} + h(\omega, a), \omega), \psi_{\omega_0} \cos \frac{y}{L} \rangle_{L^2} = 0.$$

We apply the Crandall-Rabinowitz Transversality and we consider the problem $g(\omega, a) = 0$, where

$$g(\omega, a) = \begin{cases} \frac{F_{||}(\omega, a) - F_{||}(\omega, 0)}{a}, & a \neq 0, \\ \frac{\partial F_{||}}{\partial a}(\omega, 0), & a = 0. \end{cases}$$

Here for $a \neq 0$, $F_{||}(\omega, a) = 0$ if and only if $g(\omega, a) = 0$. If $p > 2$, then $F_{||}$ is a C^2 function and g is a C^1 function. In the case $p = 2$, by the positivity of $\tilde{\varphi}_{\omega_0}$ and the Lebesgue dominant converge theorem, we can prove g is C^1 . Then,

$$\frac{\partial g}{\partial \omega}(\omega_0, 0) = \frac{\partial \lambda_\omega}{\partial \omega}\Big|_{\omega=\omega_0} \left\| \psi_{\omega_0} \cos \frac{y}{L} \right\|_{L^2}^2, \quad \frac{\partial g}{\partial a}(\omega_0, 0) = 0.$$

Therefore, by the implicit function theorem there exists $\omega_{\omega_0}(a)$ such that $g(\omega_{\omega_0}(a), a) = 0$. Hence, $\phi_{\omega_0}(a) := \tilde{\varphi}_{\omega_0} + a\psi_{\omega_0} \cos \frac{y}{L} + h(\omega_{\omega_0}(a), a)$ is a solution of $F(\phi_{\omega_0}(a), \omega_{\omega_0}(a)) = 0$ and

$$\omega'_{\omega_0}(0) = -\frac{\frac{\partial g}{\partial a}}{\frac{\partial g}{\partial \omega}}(\omega_0, 0) = 0.$$

Using certain upper and lower exponential decay rates and positivity of $\phi_{\omega_0}(a)$, we can obtain

$$\omega''_{\omega_0}(0) = \lim_{a \rightarrow 0} \frac{\omega'_{\omega_0}(0)}{a} = \frac{-1}{\frac{\partial \lambda_{\omega}}{\partial \omega} |_{\omega=\omega_0}} \lim_{a \rightarrow 0} \frac{1}{a} \frac{\partial g}{\partial a}(\omega_{\omega_0}(a), a),$$

and (4.4).

Since $\mathbb{L}(a, \omega_0)$ is C^1 , there exists an eigenfunction $\chi_*(a)$ of $\mathbb{L}(a, \omega_0)$ corresponding to $\lambda_2(a)$ such that $\|\chi_*(a)\|_{L^2} = 1$, $\chi_*(0) = \left\| \psi_{\omega_0} \cos \frac{y}{L} \right\|_{L^2}^{-1} \psi_{\omega_0} \cos \frac{y}{L}$ and $\chi_*(a)$ is C^1 with respect to a . In the case $p > 2$, since F_{\parallel} is C^2 , $\phi_{\omega_0}(a)$ is C^2 . In the case $p = 2$, since $\mathbb{L}(a, \omega_0)$ is C^1 and

$$\frac{d\phi_{\omega_0}}{da}(a) = \psi_{\omega_0} \cos \frac{y}{L} - (P_{\perp} \mathbb{L}(a, \omega_0) P_{\perp})^{-1} P_{\perp} (\mathbb{L}(a, \omega_0) \psi_{\omega_0} \cos \frac{y}{L} + \omega_{\omega_0}(a) (\tilde{\varphi}_{\omega_0} + h(\omega_{\omega_0}(a), a))),$$

$\phi_{\omega_0}(a)$ is C^2 . Since

$$\lambda_2(a) = \langle \mathbb{L}(a, \omega_0) \chi_*(a), \chi_*(a) \rangle_{L^2},$$

we obtain

$$\begin{aligned} \frac{d\lambda_2}{da} &= \omega''_{\omega_0} - 2p(p-1) \langle (\phi_{\omega_0})^{p-2} \frac{d\phi_{\omega_0}}{da} \frac{d\chi_*}{da}, \chi_* \rangle_{L^2} \\ &\quad - p(p-1) \left\langle \left((p-2)(\phi_{\omega_0})^{p-3} \left(\frac{d\phi_{\omega_0}}{da} \right)^2 + (\phi_{\omega_0})^{p-2} \frac{d^2\phi_{\omega_0}}{da^2} \right) \chi_*, \chi_* \right\rangle_{L^2}, \end{aligned}$$

and (4.5). Finally, calculating $\frac{d^2}{da^2} \|\phi_{\omega_0}(a)\|_{L^2}^2|_{a=0}$, we get (4.6). □

Lemma 4.5. *Let $p \geq 2$. Then, there exists $\omega_p > \lambda_*$ such that for $\omega_0 \in (\lambda_*, \omega_p)$, $\omega''_{\omega_0}(0) > 0$ and*

$$R_{p, \omega_0} \begin{cases} > 0, & 2 \leq p < \frac{9+\sqrt{57}}{4}, \\ < 0, & \frac{9+\sqrt{57}}{4} \leq p. \end{cases}$$

Remark 4.6. The first term of R_{p, ω_0} with respect to $\omega_0 - \lambda_*$ yields the critical exponent p_* . In Lemma 4.5, we show the following expansion:

$$R_{p, \omega_0} = \frac{(-4p^2 + 18p - 6)\pi}{3(p-1)^{3/2}(\omega - \lambda_*)^{1/2}} + O((\omega - \lambda_*)^{1/2}).$$

Proof. First, we prove the positivity of $\omega''_{\omega_0}(0)$. Let

$$\begin{aligned} I_1 &= \left\langle (\tilde{\varphi}_{\omega_0})^{p-2} \left(\psi_{\omega_0} \cos \frac{y}{L} \right)^2, \mathbb{L}_{\omega_0}^{-1} \left((\tilde{\varphi}_{\omega_0})^{p-2} \left(\psi_{\omega_0} \cos \frac{y}{L} \right)^2 \right) \right\rangle_{L^2} \\ I_2 &= \left\langle \left(\psi_{\omega_0} \cos \frac{y}{L} \right)^2, (\tilde{\varphi}_{\omega_0})^{p-3} \left(\psi_{\omega_0} \cos \frac{y}{L} \right)^2 \right\rangle_{L^2}. \end{aligned}$$

Since $(\varphi_{\omega, 0}(x))^{p-3}$ is differentiable with respect to $x \in \mathbb{R}$ and

$$\left| \frac{1}{\omega - \lambda_*} \left((\varphi_{\omega, 0}(x))^{p-3} - \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{(p-3)(p+1)}{p-1}} (\psi_*(x))^{p-3} \right) \right| \leq C(\varphi_{\theta(\omega), 0}(x))^{p-4} |\partial_{\omega} \varphi_{\theta(\omega), 0}(x)|,$$

by the boundedness of $\|\partial_\omega \varphi_{\omega,0}\|_{H^2(\mathbb{R})}$ with respect to ω and certain upper and lower exponential decay rates for φ_ω and ψ_ω we have

$$\begin{aligned} I_2 &= \frac{3}{8}(\omega_0 - \lambda_*)^{\frac{p-3}{p-1}} \int_{\mathbb{R} \times \mathbb{T}_L} \tilde{\varphi}_{\omega_0,0}^{p-3} \psi_{\omega_0}^4 dx dy \\ &= \frac{3\pi L}{4} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{\frac{2(p+1)}{p-1}} (\omega_0 - \lambda_*)^{\frac{p-3}{p-1}} \\ &\quad + \frac{3(5p-3)\pi L}{4} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{(p-3)(p+1)}{p-1}} (\omega_0 - \lambda_*)^{\frac{p-3}{p-1}+1} \int_{\mathbb{R}} \psi_*^p (P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} \psi_{*,p} dx \\ &\quad + o((\omega_0 - \lambda_*)^{\frac{p-3}{p-1}+1} L), \end{aligned}$$

where $\varphi_{\omega,0} = (\omega - \lambda_*)^{-1/(p-1)} \varphi_\omega$ and $\lambda_* < \theta(\omega) < \omega$. On the other hand,

$$\begin{aligned} I_1 &= \frac{1}{4} \langle (\tilde{\varphi}_{\omega_0})^{p-2} \psi_{\omega_0}^2, (L_{\omega_0}^+)^{-1} (\tilde{\varphi}_{\omega_0})^{p-2} \psi_{\omega_0}^2 \rangle_{L^2} \\ &\quad + \frac{1}{8} \langle (\tilde{\varphi}_{\omega_0})^{p-2} \psi_{\omega_0}^2, (L_{\omega_0}^+ + \frac{4}{L^2})^{-1} ((\tilde{\varphi}_{\omega_0})^{p-2} \psi_{\omega_0}^2) \rangle_{L^2} \\ &= I_1' + I_1''. \end{aligned}$$

By $\left\| (L_{\omega_0}^+)^{-1} |_{(\psi_{\omega_0})^\perp} \right\| \leq C$ and the similar calculation for I_2 , we obtain

$$\begin{aligned} I_1' &= \frac{(\omega_0 - \lambda_*)^{\frac{2(p-2)}{p-1}}}{4} \left\{ \left\langle (\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^2, (L_{\omega_0}^+)^{-1} \left(\int_{\mathbb{R}} (\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^3 dx \right) \psi_{\omega_0} \right\rangle_{L^2} \right. \\ &\quad \left. + \left\langle (\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^2, (L_{\omega_0}^+)^{-1} \left((\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^2 - \int_{\mathbb{R}} (\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^3 dx \psi_{\omega_0} \right) \right\rangle_{L^2} \right\} \\ &= - \frac{\pi L (\omega_0 - \lambda_*)^{\frac{p-3}{p-1}} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{\frac{2(p+1)}{p-1}}}{2(p-1)} \\ &\quad + \frac{(-5p^2 + 9p - 3)\pi L (\omega_0 - \lambda_*)^{\frac{p-3}{p-1}+1} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{(p-3)(p+1)}{p-1}}}{2(p-1)^2} \int_{\mathbb{R}} \psi_*^p (P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} \psi_{*,p} dx \\ &\quad + o((\omega_0 - \lambda_*)^{\frac{p-3}{p-1}+1} L), \end{aligned}$$

where $(\psi_\omega)^\perp = \{u \in L^2(\mathbb{R}) \mid \langle u, \psi_\omega \rangle_{L^2(\mathbb{R})} = 0\}$. By the same calculation of I_1' and the

boundedness of $\left\| (L_{\omega_0}^+ + \frac{4}{L^2})^{-1} |_{(\psi_{\omega_0})^\perp} \right\|$,

$$\begin{aligned}
 I_1'' &= \frac{(\omega_0 - \lambda_*)^{\frac{2(p-2)}{p-1}}}{8} \left\langle (\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^2, (L_{\omega_0}^+ + 4/L^2)^{-1} \left(\int_{\mathbb{R}} (\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^3 dx \right) \psi_{\omega_0} \right\rangle_{L^2} \\
 &\quad + \frac{(\omega_0 - \lambda_*)^{\frac{2(p-2)}{p-1}}}{8} \left\langle (\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^2, \right. \\
 &\quad \left. (L_{\omega_0}^+ + 4/L^2)^{-1} \left((\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^2 - \int_{\mathbb{R}} (\varphi_{\omega_0,0})^{p-2} (\psi_{\omega_0})^3 dx \psi_{\omega_0} \right) \right\rangle_{L^2} \\
 &= \frac{\pi L (\omega_0 - \lambda_*)^{\frac{p-3}{p-1}} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{\frac{2(p+1)}{p-1}}}{12(p-1)} \\
 &\quad + \frac{(9p^2 - 17p + 7)\pi L (\omega_0 - \lambda_*)^{\frac{p-3}{p-1}+1} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{(p-3)(p+1)}{p-1}}}{12(p-1)^2} \int_{\mathbb{R}} \psi_*^p (P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} \psi_{*,p} dx \\
 &\quad + o((\omega_0 - \lambda_*)^{\frac{p-3}{p-1}+1} L).
 \end{aligned}$$

Since

$$\frac{1}{\frac{d}{d\omega} \lambda_{\omega_0} \|\psi_{\omega_0} \cos \frac{y}{L}\|_{L^2}^2} = \frac{1}{(p-1)\pi L} - \frac{C_*(\omega_0 - \lambda_*)}{(p-1)\pi L} + o((\omega_0 - \lambda_*)L^{-1}),$$

we obtain

$$\begin{aligned}
 &\omega_{\omega_0}''(0) \\
 &= \frac{p(p+3)(\omega_0 - \lambda_*)^{\frac{p-3}{p-1}} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{\frac{2(p+1)}{p-1}}}{6} \\
 &\quad - \frac{p(2p^3 + 3p^2 + 34p - 18)(\omega_0 - \lambda_*)^{\frac{p-3}{p-1}+1} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{(p-3)(p+1)}{p-1}}}{12(p-1)} \int_{\mathbb{R}} \psi_*^p (P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} \psi_{*,p} dx \\
 &\quad + o((\omega_0 - \lambda_*)^{\frac{p-3}{p-1}+1})
 \end{aligned}$$

where

$$C_* = \frac{2p(2p-1) \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)} \int_{\mathbb{R}} \psi_*^p (P_\perp^1 H_{\lambda_*} P_\perp^1)^{-1} \psi_{*,p} dx}{p-1}.$$

Therefore, if $0 < \omega_0 - \lambda_* \ll 1$, then $\omega_{\omega_0}''(0) > 0$.

Next, we calculate R_{p,ω_0} . Since

$$-2 \frac{d\lambda_\omega}{d\omega} \Big|_{\omega=\omega_0} \left\| \psi_{\omega_0} \cos \frac{y}{L} \right\|_{L^2}^2 = -2(p-1)\pi L - 2C_*(p-1)\pi L(\omega_0 - \lambda_*) + o((\omega_0 - \lambda_*)L)$$

and

$$\frac{d}{d\omega} \|\tilde{\varphi}_\omega\|_{L^2}^2 \Big|_{\omega=\omega_0} = \frac{4\pi L}{p-1} \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-\frac{2(p+1)}{p-1}} (\omega_0 - \lambda_*)^{-\frac{p-3}{p-1}} + o((\omega_0 - \lambda_*)^{\frac{2}{p-1}} L),$$

we have

$$\begin{aligned}
 R_{p,\omega_0} &= -2(p-1)\pi L - 2C_*(p-1)\pi L(\omega_0 - \lambda_*) + \frac{2p(p+3)\pi L}{3(p-1)} + o((\omega_0 - \lambda_*)L) \\
 &\quad - \frac{p(2p^3 + 3p^2 + 34p - 18)\pi L}{3(p-1)^2} (\omega_0 - \lambda_*) \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)} \int_{\mathbb{R}} \psi_*^p (P_{\perp}^1 H_{\lambda_*} P_{\perp}^1)^{-1} \psi_{*,p} dx \\
 &= \frac{(-4p^2 + 18p - 6)\pi L}{3(p-1)} + o((\omega_0 - \lambda_*)L) \\
 &\quad + \frac{p(-26p^3 + 57p^2 - 82p + 30)\pi L}{3(p-1)^2} (\omega_0 - \lambda_*) \|\psi_*\|_{L^{p+1}(\mathbb{R})}^{-(p+1)} \int_{\mathbb{R}} \psi_*^p (P_{\perp}^1 H_{\lambda_*} P_{\perp}^1)^{-1} \psi_{*,p} dx.
 \end{aligned} \tag{4.7}$$

Let

$$p_* = \frac{9 + \sqrt{57}}{4}.$$

Since p_* is the root of $-4p^2 + 18p - 6 = 0$ with $p > 1$, the conclusion for $p \neq p_*$ follows (4.7). Finally, we consider the case $p = p_*$. By $p_* > 4$, we have

$$-26p_*^3 + 57p_*^2 - 82p_* + 30 < 0.$$

Therefore,

$$\begin{aligned}
 R_{p_*,\omega_0} &= \frac{p_*(-26p_*^3 + 57p_*^2 - 82p_* + 30)\pi L(\omega_0 - \lambda_*) \|\psi_*\|_{L^{p_*+1}(\mathbb{R})}^{-(p_*+1)}}{3(p_*-1)^2} \int_{\mathbb{R}} \psi_*^{p_*} (P_{\perp}^1 H_{\lambda_*} P_{\perp}^1)^{-1} \psi_{*,p_*} dx \\
 &\quad + o((\omega_0 - \lambda_*)L) \\
 &= \frac{p_*(-26p_*^3 + 57p_*^2 - 82p_* + 30)\pi L(\omega_0 - \lambda_*)}{3(p_*-1)^2} \int_{\mathbb{R}} \psi_{*,p_*} (P_{\perp}^1 H_{\lambda_*} P_{\perp}^1)^{-1} \psi_{*,p_*} dx \\
 &\quad + o((\omega_0 - \lambda_*)L)
 \end{aligned}$$

The conclusion for $p = p_*$ follows this. \square

Using Lemma 4.5 and applying the argument in Section 3 of [26], we obtain Theorem 1.4.

For the completeness of the proof of Theorem 1.4, we introduce the argument for the stability of standing with the degenerate linearized operator in [13, 26]. Using the following proposition, we show Theorem 1.4.

Proposition 4.7. *Let $\lambda_* < \omega_0 < \omega_{*,0}$.*

- (i) *If $R_{p,\omega_0} > 0$, then $e^{i\omega_0 t} \tilde{\varphi}_{\omega_0}$ is a stable standing wave of (NLS) on $\mathbb{R} \times \mathbb{T}_L$ with $L = (\lambda_{\omega_0})^{-\frac{1}{2}}$.*
- (ii) *If $R_{p,\omega_0} < 0$, then $e^{i\omega_0 t} \tilde{\varphi}_{\omega_0}$ is an unstable standing wave of (NLS) on $\mathbb{R} \times \mathbb{T}_L$ with $L = (\lambda_{\omega_0})^{-\frac{1}{2}}$.*

To modulate the translation symmetry for $y \in \mathbb{T}_L$, we define the polar coordinate $\vec{a} = (a_1, a_2) = (a \cos \frac{\tilde{a}}{L}, -a \sin \frac{\tilde{a}}{L})$ for $\vec{a} \in \mathbb{R}^2$ and

$$\phi_{\omega_0}(\vec{a})(x, y) = \phi_{\omega_0}(a)(x, y + \tilde{a}), \quad \omega_{\omega_0}(\vec{a}) = \omega_{\omega_0}(a).$$

In the following lemma, we construct a curve which captures the degeneracy of the linearized operator $S''_{\omega_0}(\tilde{\varphi}_{\omega_0})$.

Lemma 4.8. *There exist a neighborhood U of $(0, 0)$ in \mathbb{R}^2 and a C^1 function $\rho : U \rightarrow \mathbb{R}$ such that $\rho(0, 0) = 0$ and for $\vec{a} \in U$*

$$\begin{aligned} Q(\phi_{\omega_0}(\vec{a}) + \rho(\vec{a})\partial_{\omega}\tilde{\varphi}_{\omega_0}) &= Q(\tilde{\varphi}_{\omega_0}), \\ \rho(\vec{a})\langle \tilde{\varphi}_{\omega_0}, \partial_{\omega}\tilde{\varphi}_{\omega_0} \rangle_{L^2} &= Q(\tilde{\varphi}_{\omega_0}) - Q(\phi_{\omega_0}(\vec{a})) + o(\rho(\vec{a})). \end{aligned} \quad (4.8)$$

Proof. Since

$$\partial_{\rho}Q(\phi_{\omega_0}(\vec{a}) + \rho\partial_{\omega}\tilde{\varphi}_{\omega_0})|_{\rho=0, \vec{a}=0} = \langle \tilde{\varphi}_{\omega_0}, \partial_{\omega}\tilde{\varphi}_{\omega_0} \rangle_{L^2} > 0,$$

the conclusion follows the implicit function theorem. \square

Let

$$\Phi(\vec{a}) = \phi_{\omega_0}(\vec{a}) + \rho(\vec{a})\partial_{\omega}\tilde{\varphi}_{\omega_0}.$$

for $\vec{a} \in U$.

In the following lemma, we capture the degeneracy of the action S_{ω} .

Lemma 4.9. *For $\vec{a} \in U$,*

$$S_{\omega_0}(\Phi(\vec{a})) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) = \frac{\frac{d}{d\omega}\lambda_{\omega_0}\|\psi_{\omega_0}\cos(y/L)\|_{L^2}^2 R_{p,\omega_0}}{16\langle \tilde{\varphi}_{\omega_0}, \partial_{\omega}\tilde{\varphi}_{\omega_0} \rangle_{L^2}}|\vec{a}|^4 + o(|\vec{a}|^4).$$

Proof. For $\vec{a} \in U$,

$$\begin{aligned} S_{\omega_0}(\Phi(\vec{a})) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) &= S_{\omega_{\omega_0}(\vec{a})}(\Phi(\vec{a})) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) + (\omega_0 - \omega_{\omega_0}(\vec{a}))Q(\tilde{\varphi}_{\omega_0}) \\ &= S_{\omega_{\omega_0}(\vec{a})}(\phi_{\omega_0}(\vec{a})) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) + (\omega_0 - \omega_{\omega_0}(\vec{a}))Q(\tilde{\varphi}_{\omega_0}) \\ &\quad + \frac{1}{2}(\rho(\vec{a}))^2\langle S''_{\omega_0}(\tilde{\varphi}_{\omega_0})\partial_{\omega}\tilde{\varphi}_{\omega_0}, \partial_{\omega}\tilde{\varphi}_{\omega_0} \rangle_{L^2} + o((\rho(\vec{a}))^2). \end{aligned}$$

From $\omega''_{\omega_0}(0) > 0$ and (4.3), $\omega_{\omega_0}(a)$ is increasing on a small interval $(0, \delta)$. Therefore, there exists the inverse function $a^+(\omega)$ of $\omega_{\omega_0}(a)$ from $[\omega_0, \omega_{\omega_0}(\delta)]$ to $[0, \delta]$. By the differentiability of a^+ for $\omega > \omega_0$, $\phi_{\omega_0}(a^+)$ is differentiable for $\omega > \omega_0$. Thus, for ω, ω_1 with $\omega \neq \omega_1$

$$\begin{aligned} &\frac{S_{\omega}(\phi_{\omega_0}(a^+(\omega))) - S_{\omega_1}(\phi_{\omega_0}(a^+(\omega_1)))}{\omega - \omega_1} \\ &= \frac{\langle S''_{\omega_1}(\phi_{\omega_0}(a^+(\omega_1)))(\phi_{\omega_0}(a^+(\omega)) - \phi_{\omega_0}(a^+(\omega_1))), (\phi_{\omega_0}(a^+(\omega)) - \phi_{\omega_0}(a^+(\omega_1))) \rangle_{L^2}}{2(\omega - \omega_1)} \\ &\quad + Q(\phi_{\omega_0}(a^+(\omega))) + \frac{o((\phi_{\omega_0}(a^+(\omega)) - \phi_{\omega_0}(a^+(\omega_1)))^2)}{\omega - \omega_1} \\ &\rightarrow Q(\phi_{\omega_0}(a^+(\omega_1))) \quad \text{as } \omega \rightarrow \omega_1. \end{aligned}$$

Moreover, since $\partial_a \phi_{\omega_0}(a)|_{a=0} = \psi_{\omega_0} \cos \frac{y}{L}$, for $\omega > \omega_0$

$$\begin{aligned} & \frac{S_\omega(\phi_{\omega_0}(a^+(\omega))) - S_{\omega_0}(\tilde{\varphi}_{\omega_0})}{\omega - \omega_0} \\ &= \frac{\langle S''_{\omega_0}(\tilde{\varphi}_{\omega_0})(\phi_{\omega_0}(a^+) - \tilde{\varphi}_{\omega_0}), (\phi_{\omega_0}(a^+) - \tilde{\varphi}_{\omega_0}) \rangle_{L^2}}{\omega''_{\omega_0}(0)(a^+)^2 + o((a^+)^2)} + Q(\tilde{\varphi}_{\omega_0}) + \frac{o((\phi_{\omega_0}(a^+) - \tilde{\varphi}_{\omega_0})^2)}{\omega''_{\omega_0}(0)(a^+)^2 + o((a^+)^2)} \\ &\rightarrow Q(\tilde{\varphi}_{\omega_0}) \quad \text{as } \omega \downarrow \omega_0. \end{aligned}$$

Hence, $S_\omega(\phi_{\omega_0}(a^+))$ is C^1 and

$$\frac{dS_\omega(\phi_{\omega_0}(a^+))}{d\omega} = Q(\phi_{\omega_0}(a^+)).$$

By the equation (4.6), $Q(\phi_{\omega_0}(a^+))$ is C^1 on $(\omega_0, \omega_{\omega_0}(\delta))$ and

$$\lim_{\omega \downarrow \omega_0} \frac{Q(\phi_{\omega_0}(a^+)) - Q(\tilde{\varphi}_{\omega_0})}{\omega - \omega_0} = \frac{R_{p,\omega_0}}{2\omega''_{\omega_0}(0)}.$$

Therefore, $S_\omega(\phi_{\omega_0}(a^+))$ is C^2 with respect to ω on $(\omega_0, \omega_{\omega_0}(\delta))$ and

$$\begin{aligned} S_\omega(\phi_{\omega_0}(a^+)) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) + (\omega_0 - \omega)Q(\tilde{\varphi}_{\omega_0}) &= \frac{R_{p,\omega_0}}{4\omega''_{\omega_0}(0)}(\omega - \omega_0)^2 + o((\omega - \omega_0)^2) \\ &= \frac{\omega''_{\omega_0}(0)R_{p,\omega_0}}{16}(a^+)^4 + o((a^+)^4). \end{aligned} \tag{4.9}$$

From the equation (4.8), we have the expansion

$$(\rho(\vec{a}))^2 \langle \tilde{\varphi}_{\omega_*}, \partial_\omega \tilde{\varphi}_{\omega_0} \rangle_{L^2} = \frac{(R_{p,\omega_0})^2}{16 \langle \tilde{\varphi}_{\omega_*}, \partial_\omega \tilde{\varphi}_{\omega_0} \rangle_{L^2}} |\vec{a}|^4 + o(|\vec{a}|^4). \tag{4.10}$$

Since

$$S_{\omega_{\omega_0}(|\vec{a}|)}(\phi_{\omega_0}(|\vec{a}|)) + (\omega_0 - \omega_{\omega_0}(|\vec{a}|))Q(\tilde{\varphi}_{\omega_0}) = S_{\omega_{\omega_0}(\vec{a})}(\phi_{\omega_0}(\vec{a})) + (\omega_0 - \omega_{\omega_0}(\vec{a}))Q(\tilde{\varphi}_{\omega_0}),$$

by (4.9) and (4.10) we obtain the conclusion. \square

We introduce the distance and tubular neighborhoods of $\tilde{\varphi}_{\omega_0}$ as follows. Set for $\varepsilon > 0$

$$\text{dist}_{\omega_0}(u) = \inf_{\theta \in \mathbb{R}} \|u - e^{i\theta} \tilde{\varphi}_{\omega_0}\|_{H^1},$$

$$\begin{aligned} N_\varepsilon &= \{u \in H^1(\mathbb{R} \times \mathbb{T}_L) \mid \text{dist}_{\omega_0}(u) < \varepsilon\}, \\ N_\varepsilon^0 &= \{u \in N_\varepsilon \mid Q(u) = Q(\tilde{\varphi}_{\omega_0})\}. \end{aligned}$$

Modulating the symmetry, we eliminate the degeneracy of the linearized operator around $\tilde{\varphi}_{\omega_0}$.

Lemma 4.10. *Let $\varepsilon > 0$ sufficiently small. Then, there exist C^2 function $\theta : N_\varepsilon \rightarrow \mathbb{R}$, $\alpha : N_\varepsilon \rightarrow \mathbb{R}$, $\vec{a} : N_\varepsilon \rightarrow U$ and $w : N_\varepsilon \rightarrow H^1(\mathbb{R} \times \mathbb{T}_L)$ such that for $u \in N_\varepsilon$*

$$e^{i\theta(u)}u = \Phi(\vec{a}(u)) + w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u)),$$

where $\langle w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u)), \psi_{\omega_0} \cos(y/L) \rangle_{L^2} = \langle w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u)), \psi_{\omega_0} \sin(y/L) \rangle_{L^2} = \langle w(u), \phi_{\omega_0}(\vec{a}(u)) \rangle_{L^2} = \langle w(u), i\phi_{\omega_0}(\vec{a}(u)) \rangle_{L^2} = 0$.

Proof. Let $\psi_{\omega_0,1} = \psi_{\omega_0} \cos(y/L)$ and $\psi_{\omega_0,2} = \psi_{\omega_0} \sin(y/L)$. We define

$$G(u, \theta, a_1, a_2) = \begin{pmatrix} \langle e^{i\theta}u - \Phi(\vec{a}), i\phi_{\omega_0}(\vec{a}) \rangle_{L^2} \\ \langle e^{i\theta}u - \Phi(\vec{a}), \psi_{\omega_0,1} \rangle_{L^2} \\ \langle e^{i\theta}u - \Phi(\vec{a}), \psi_{\omega_0,2} \rangle_{L^2} \end{pmatrix},$$

where $\vec{a} = (a_1, a_2)$. Since $G(\tilde{\varphi}_{\omega_0}, 0, 0, 0) = 0$ and

$$\frac{\partial G}{\partial(\theta, a_1, a_2)} \Big|_{(u, \theta, a_1, a_2) = (\tilde{\varphi}_{\omega_0}, 0, 0, 0)} = \begin{pmatrix} \|\tilde{\varphi}_{\omega_0}\|_{L^2}^2 & 0 & 0 \\ 0 & -\|\psi_{\omega_0,1}\|_{L^2}^2 & 0 \\ 0 & 0 & -\|\psi_{\omega_0,2}\|_{L^2}^2 \end{pmatrix},$$

by the implicit theorem for sufficiently small $\varepsilon > 0$ there exist C^2 functions $\theta : N_\varepsilon \rightarrow \mathbb{R}$ and $\vec{a} : N_\varepsilon \rightarrow U$ such that for $u \in N_\varepsilon$

$$G(u, \theta(u), \vec{a}(u)) = 0.$$

We define

$$\alpha(u) = \frac{\langle e^{i\theta(u)}u - \Phi(\vec{a}(u)), \phi_{\omega_0}(\vec{a}(u)) \rangle_{L^2}}{\|\phi_{\omega_0}(\vec{a}(u))\|_{L^2}^2},$$

and

$$w(u) = e^{i\theta(u)}u - \Phi(\vec{a}(u)) - \alpha(u)\phi_{\omega_0}(\vec{a}(u)).$$

Then, the conclusion follows the definition of w . □

In the following lemma, we show the estimate of $\alpha(u)$ for $u \in N_\varepsilon^0$.

Lemma 4.11. *Let $\varepsilon > 0$ sufficiently small. There exists $C > 0$ such that for $u \in N_\varepsilon^0$,*

$$|\alpha(u)| \leq C\|w(u)\|_{L^2}(\rho(\vec{a}(u)) + \|w(u)\|_{L^2}).$$

Proof. By Lemma 4.10, for $u \in N_\varepsilon^0$,

$$\begin{aligned} Q(\tilde{\varphi}_{\omega_0}) &= Q(\Phi(\vec{a}(u)) + w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u))) \\ &= Q(\tilde{\varphi}_{\omega_0}) + \alpha(u)\|\phi_{\omega_0}(\vec{a}(u))\|_{L^2}^2 + \rho(\vec{a}(u))\alpha(u)\langle \partial_\omega \tilde{\varphi}_{\omega_0}, \phi_{\omega_0}(\vec{a}(u)) \rangle_{L^2} \\ &\quad + \rho(\vec{a}(u))\langle \partial_\omega \tilde{\varphi}_{\omega_0}, w(u) \rangle_{L^2} + Q(w(u)) + (\alpha(u))^2 Q(\phi_{\omega_0}(\vec{a}(u))). \end{aligned}$$

Since $\rho(\vec{a}(u)) \rightarrow 0$ as $\varepsilon \rightarrow 0$, we obtain the conclusion. □

Next, we prove the coerciveness of the linearized operator around $\tilde{\varphi}_{\omega_0}$.

Lemma 4.12. *There exist $k_0 > 0$ and $\varepsilon_0 > 0$ such that for $a_1, a_2, \alpha \in (-\varepsilon_0, \varepsilon_0)$, if $w \in H^1(\mathbb{R} \times \mathbb{T}_L)$ with $\langle w, \phi_{\omega_0}(\vec{a}) \rangle_{L^2} = \langle w, i\phi_{\omega_0}(\vec{a}) \rangle_{L^2} = \langle w + \alpha\phi_{\omega_0}(\vec{a}), \psi_{\omega_0} \cos(y/L) \rangle_{L^2} = \langle w + \alpha\phi_{\omega_0}(\vec{a}), \psi_{\omega_0} \sin(y/L) \rangle_{L^2} = 0$, then*

$$\langle S''_{\omega_0}(\Phi(\vec{a}))w, w \rangle_{H^{-1}, H^1} \geq k_0 \|w\|_{H^1}^2,$$

where $\vec{a} = (a_1, a_2)$.

Proof. Let $\psi_{\omega_0,1} = \psi_{\omega_0} \cos(y/L)$ and $\psi_{\omega_0,2} = \psi_{\omega_0} \sin(y/L)$. For $w \in H^1(\mathbb{R} \times \mathbb{T}_L)$ with $\langle w, \phi_{\omega_0}(\vec{a}) \rangle_{L^2} = \langle w, i\phi_{\omega_0}(\vec{a}) \rangle_{L^2} = \langle w + \alpha\phi_{\omega_0}(\vec{a}), \psi_{\omega_0,1} \rangle_{L^2} = \langle w + \alpha\phi_{\omega_0}(\vec{a}), \psi_{\omega_0,2} \rangle_{L^2} = 0$, we decompose $w = b_1\tilde{\varphi}_{\omega_0} + b_2i\tilde{\varphi}_{\omega_0} + b_3\psi_{\omega_0,1} + b_4\psi_{\omega_0,2} + w_{\perp}$, where $\langle w_{\perp}, \tilde{\varphi}_{\omega_0} \rangle_{L^2} = \langle w_{\perp}, i\tilde{\varphi}_{\omega_0} \rangle_{L^2} = \langle w_{\perp}, \psi_{\omega_0,1} \rangle_{L^2} = \langle w_{\perp}, \psi_{\omega_0,2} \rangle_{L^2} = 0$, $b_j \in \mathbb{R}$ for $j \in \{1, 2, 3, 4\}$. By the non-negativeness of $L_{\omega_0}^-$ and $L_{\omega_0}^+ + \lambda_{\omega_0}$, Proposition 2.1 and Lemma 2.2, there exists $c > 0$ such that $\langle S''_{\omega_0}(\tilde{\varphi}_{\omega_0})w_{\perp}, w_{\perp} \rangle_{H^{-1}, H^1} \geq c\|w_{\perp}\|_{H^1}^2$, where c is independent of w_{\perp} . Then, from the orthogonal conditions for w , we have for $j \in \{1, 2, 3, 4\}$, $b_j = O((|\vec{a}| + |\alpha|)\|w_{\perp}\|_{L^2})$ as $|\vec{a}| + |\alpha| \rightarrow 0$. Therefore, there exist $\varepsilon_0, k_0 > 0$ such that for $a_1, a_2, \alpha \in (-\varepsilon_0, \varepsilon_0)$,

$$\begin{aligned} \langle S''_{\omega_0}(\Phi(\vec{a}))w, w \rangle_{H^{-1}, H^1} &= \langle S''_{\omega_0}(\tilde{\varphi}_{\omega_0})w_{\perp}, w_{\perp} \rangle_{H^{-1}, H^1} + \sum_{j=1}^4 b_j^2 + o(\|w_{\perp}\|_{L^2}^2) \\ &\geq k_0 \|w\|_{H^1}^2. \end{aligned}$$

□

In the following lemma, we investigate the variational structure of S_{ω_0} around $\tilde{\varphi}_{\omega_0}$.

Lemma 4.13. *Let $\varepsilon > 0$ sufficiently small. For $u \in N_{\varepsilon}^0$*

$$\begin{aligned} S_{\omega_0}(u) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) &= \frac{1}{2} \langle S''_{\omega_0}(\tilde{\varphi}_{\omega_0})w(u), w(u) \rangle_{H^{-1}, H^1} + C_{**} R_{p, \omega_0} |\vec{a}(u)|^4 \\ &\quad + o(\|w(u)\|_{H^1}^2) + o(|\vec{a}(u)|^4), \end{aligned}$$

where $w(u)$ and $\vec{a}(u)$ are defined by Lemma 4.10 and

$$C_{**} = \frac{\frac{d}{d\omega} \lambda_{\omega_0} \|\psi_{\omega_0} \cos(y/L)\|_{L^2}^2}{16 \langle \tilde{\varphi}_{\omega_0}, \partial_{\omega} \tilde{\varphi}_{\omega_0} \rangle_{L^2}}.$$

Proof. Let $u \in N_{\varepsilon}^0$. By Lemma 4.10 and Lemma 4.11, we have

$$\begin{aligned} &S_{\omega_0}(u) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) \\ &= S_{\omega_0}(\Phi(\vec{a}(u)) + w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u))) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) \\ &= S_{\omega_0}(\Phi(\vec{a}(u))) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) + \langle S'_{\omega_0}(\Phi(\vec{a}(u))), w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u)) \rangle_{H^{-1}, H^1} \\ &\quad + \frac{1}{2} \langle S''_{\omega_0}(\Phi(\vec{a}(u)))w(u), w(u) \rangle_{H^{-1}, H^1} + o(\|w(u)\|_{H^1}^2). \end{aligned}$$

Since $\rho(\vec{a}(u)) = O(|\vec{a}(u)|^2)$ as $\text{dist}_{\omega_0}(u) \rightarrow 0$, $\langle \phi_{\omega_0}(\vec{a}(u)), w(u) \rangle_{L^2} = 0$ and $S''_{\omega_0}(\tilde{\varphi}_{\omega_0})\partial_{\omega}\tilde{\varphi}_{\omega_0} = -\tilde{\varphi}_{\omega_0}$, we have

$$\begin{aligned} & \langle S'_{\omega_0}(\Phi(\vec{a}(u))), w(u) \rangle_{H^{-1}, H^1} \\ &= \langle S'_{\omega_{\omega_0}(\vec{a}(u))}(\Phi(\vec{a}(u))) + (\omega_0 - \omega_{\omega_0}(\vec{a}))\Phi(\vec{a}(u)), w(u) \rangle_{H^{-1}, H^1} \\ &= \langle S''_{\omega_{\omega_0}(\vec{a}(u))}(\phi_{\omega_0}(\vec{a}(u)))\rho(\vec{a}(u))\partial_{\omega}\tilde{\varphi}_{\omega_0}, w(u) \rangle_{H^{-1}, H^1} + o(|\vec{a}|^4) + o(\|w(u)\|_{H^1}^2) \\ &= \langle (S''_{\omega_{\omega_0}(\vec{a}(u))}(\phi_{\omega_0}(\vec{a}(u))) - S''_{\omega_0}(\tilde{\varphi}_{\omega_0}))\rho(\vec{a}(u))\partial_{\omega}\tilde{\varphi}_{\omega_0}, w(u) \rangle_{H^{-1}, H^1} \\ &\quad - \rho(\vec{a}(u))\langle \tilde{\varphi}_{\omega_0} - \phi_{\omega_0}(\vec{a}(u)), w(u) \rangle_{L^2} + o(|\vec{a}(u)|^4) + o(\|w(u)\|_{H^1}^2) \\ &= o(|\vec{a}(u)|^4) + o(\|w(u)\|_{H^1}^2). \end{aligned}$$

By Lemma 4.11 and the continuity of $S'_{\omega_0}(\Phi(\vec{a}))$ and $\phi_{\omega_0}(\vec{a})$ at $\vec{a} = 0$, we have

$$\langle S'_{\omega_0}(\Phi(\vec{a}(u))), \alpha(u)\phi_{\omega_0}(\vec{a}(u)) \rangle_{H^{-1}, H^1} = o(|\vec{a}(u)|^4) + o(\|w(u)\|_{H^1}^2).$$

Therefore, from Lemma 4.9, we have the conclusion. \square

4.1 The proof of (i) of Proposition 4.7

In this subsection, we prove (i) of Proposition 4.7. Let $0 < \varepsilon \ll 1$. By Lemma 4.13 and $R_{p, \omega_0} > 0$, for small ε we have that there exists $c > 0$ such that for $u \in N_{\varepsilon}^0$

$$S_{\omega_0}(u) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) \geq c(\|w(u)\|_{H^1}^2 + |\vec{a}(u)|^4), \quad (4.11)$$

where $w(u), \vec{a}(u)$ are defined by Lemma 4.10. We suppose that there exist $\varepsilon_0 > 0$, a sequence $\{u_n\}_n$ of solutions and a sequence $\{t_n\}_n$ such that $t_n > 0$ and $u_n(0) \rightarrow \tilde{\varphi}_{\omega_0}$ in H^1 and

$$\inf_{\theta \in \mathbb{R}} \|u_n(t_n) - e^{i\theta}\tilde{\varphi}_{\omega_0}\|_{H^1} > \varepsilon_0.$$

Let

$$v_n = \sqrt{\frac{Q(\tilde{\varphi}_{\omega_0})}{Q(u_n)}}u_n(t_n).$$

Since $Q(v_n) = Q(\tilde{\varphi}_{\omega_0})$ and $Q(u_n) \rightarrow Q(\tilde{\varphi}_{\omega_0})$ as $n \rightarrow \infty$, $\|v_n - u_n(t_n)\|_{H^1} \rightarrow 0$ and $S_{\omega_0}(v_n) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) \rightarrow 0$ as $n \rightarrow \infty$. By the equation (4.11), $\vec{a}(u_n(t_n)) \rightarrow 0$, $\alpha(u_n(t_n)) \rightarrow 0$ and $w(u_n(t_n)) \rightarrow 0$ in H^1 as $n \rightarrow \infty$. Therefore,

$$\inf_{\theta \in \mathbb{R}} \|u_n(t_n) - e^{i\theta}\tilde{\varphi}_{\omega_0}\|_{H^1} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This is a contradiction. We complete the proof of (i).

4.2 The proof of (ii) of Proposition 4.7

In this subsection, we prove (ii) of Proposition 4.7. Let $0 < \varepsilon \ll 1$. We define the functions $A(u)$ and $P(u)$ as

$$A(u) = \langle e^{i\theta(u)}u, -i[a_1(u)\partial_{a_1}\Phi(\vec{a}(u)) + a_2(u)\partial_{a_2}\Phi(\vec{a}(u))] \rangle_{L^2},$$

$$P(u) = \langle S'_{\omega_{\omega_0}(\vec{a}(u))}(u), iA'(u) \rangle_{H^{-1}, H^1},$$

for $u \in N_\varepsilon$, where $\theta(u)$ and $\vec{a}(u)$ are defined by Lemma 4.10.

Then

$$\begin{aligned} A'(u) &= -ie^{-i\theta(u)}[a_1(u)\partial_{a_1}\Phi(\vec{a}(u)) + a_2(u)\partial_{a_2}\Phi(\vec{a}(u))] \\ &\quad + \langle ie^{i\theta(u)}u, -i[a_1(u)\partial_{a_1}\Phi(\vec{a}(u)) + a_2(u)\partial_{a_2}\Phi(\vec{a}(u))] \rangle_{L^2}\theta'(u) \\ &\quad + \langle e^{i\theta(u)}u, -i[\partial_{a_1}\Phi(\vec{a}(u)) + a_1(u)\partial_{a_1}\partial_{a_1}\Phi(\vec{a}(u)) + a_2(u)\partial_{a_1}\partial_{a_2}\Phi(\vec{a}(u))] \rangle_{L^2}a'_1(u) \\ &\quad + \langle e^{i\theta(u)}u, -i[a_1(u)\partial_{a_1}\partial_{a_2}\Phi(\vec{a}(u)) + \partial_{a_2}\Phi(\vec{a}(u)) + a_2(u)\partial_{a_2}\partial_{a_2}\Phi(\vec{a}(u))] \rangle_{L^2}a'_2(u), \end{aligned}$$

$$\langle iA'(u), Q'(u) \rangle_{L^2} = -\langle A'(u), iu \rangle_{L^2} = \left. \frac{dA(e^{i\theta}u)}{d\theta} \right|_{\theta=0} = 0.$$

Therefore, for any solution $u(t)$ of (1.1)

$$\begin{aligned} \frac{dA(u(t))}{dt} &= \langle A'(u(t)), -iE'(u(t)) \rangle_{H^{-1}, H^1} = \langle iA'(u(t)), E'(u(t)) + \omega(\vec{a}(u(t)))Q'(u(t)) \rangle_{H^{-1}, H^1} \\ &= P(u(t)). \end{aligned}$$

Next, we investigate the function P .

Lemma 4.14. For $\vec{a} \in U$,

$$P(\Phi(\vec{a})) = -|\vec{a}|^2\rho(\vec{a})\frac{d\lambda_\omega}{d\omega}\Big|_{\omega=\omega_0}\left\|\psi_{\omega_0}\cos\frac{y}{L}\right\|_{L^2}^2 + o(\rho(\vec{a})^2).$$

Proof. Let $\vec{a}_0 = (a_{1,0}, a_{2,0}) \in U$. Then $\|\Phi(\vec{a})\|_{L^2} = \|\tilde{\varphi}_{\omega_0}\|_{L^2}$, $\vec{a}(\Phi(\vec{a}_0)) = \vec{a}_0$ and $\theta(\Phi(\vec{a}_0)) = 0$. Therefore,

$$\begin{aligned} S'_{\omega_{\omega_0}(\vec{a}_0)}(\Phi(\vec{a}_0)) &= S''_{\omega_0}(\tilde{\varphi}_{\omega_0})\rho(\vec{a}_0)\partial_\omega\tilde{\varphi}_{\omega_0} - \rho(\vec{a}_0)p(p-1)(\tilde{\varphi}_{\omega_0})^{p-2}\partial_\omega\tilde{\varphi}_{\omega_0}\psi_{\omega_0}\left(a_{1,0}\cos\frac{y}{L} + a_{2,0}\sin\frac{y}{L}\right) \\ &\quad + o(\rho(\vec{a}_0)|\vec{a}_0|), \end{aligned} \tag{4.12}$$

$$\begin{aligned} &iA'(\Phi(\vec{a}_0)) \\ &= a_{1,0}\partial_{a_1}\Phi(\vec{a}_0) + a_{2,0}\partial_{a_2}\Phi(\vec{a}_0) \\ &= \psi_{\omega_0}\left(a_{1,0}\cos\frac{y}{L} + a_{2,0}\sin\frac{y}{L}\right) + (a_{1,0}\partial_{a_1}\rho(\vec{a}_0) + a_{2,0}\partial_{a_2}\rho(\vec{a}_0))\partial_\omega\tilde{\varphi}_{\omega_0} \\ &\quad + (S''_{\omega_0}(\tilde{\varphi}_{\omega_0}))^{-1}\left[-|\vec{a}_0|^2\omega''_{\omega_0}(0)\tilde{\varphi}_{\omega_0} + p(p-1)(\tilde{\varphi}_{\omega_0})^{p-2}\psi_{\omega_0}^2\left(a_{1,0}\cos\frac{y}{L} + a_{2,0}\sin\frac{y}{L}\right)^2\right] \\ &\quad + o(|\vec{a}_0|^2) \end{aligned} \tag{4.13}$$

Hence, we have

$$\begin{aligned}
 & P(\Phi(\vec{a}_0)) \\
 = & \langle S''_{\omega_0}(\tilde{\varphi}_{\omega_0})\rho(\vec{a}_0)\partial_\omega\tilde{\varphi}_{\omega_0}, \psi_{\omega_0} \left(a_{1,0} \cos \frac{y}{L} + a_{2,0} \sin \frac{y}{L} \right) + (a_{1,0}\partial_{a_1}\rho(\vec{a}_0) + a_{2,0}\partial_{a_2}\rho(\vec{a}_0))\partial_\omega\tilde{\varphi}_{\omega_0} \rangle_{L^2} \\
 & + \langle \rho(\vec{a}_0)\partial_\omega\tilde{\varphi}_{\omega_0}, -|\vec{a}_0|^2\omega''_{\omega_0}(0)\tilde{\varphi}_{\omega_0} + p(p-1)(\tilde{\varphi}_{\omega_0})^{p-2}\psi_{\omega_0}^2 \left(a_{1,0} \cos \frac{y}{L} + a_{2,0} \sin \frac{y}{L} \right)^2 \rangle_{L^2} \\
 & + \langle -\rho(\vec{a}_0)p(p-1)(\tilde{\varphi}_{\omega_0})^{p-2}\partial_\omega\tilde{\varphi}_{\omega_0}\psi_{\omega_0} \left(a_{1,0} \cos \frac{y}{L} + a_{2,0} \sin \frac{y}{L} \right), \psi_{\omega_0} \left(a_{1,0} \cos \frac{y}{L} + a_{2,0} \sin \frac{y}{L} \right) \rangle_{L^2} \\
 & + o(\rho(\vec{a}_0)|\vec{a}_0|^2) \\
 = & -\rho(\vec{a}_0)\langle \tilde{\varphi}_{\omega_0}, \partial_\omega\tilde{\varphi}_{\omega_0} \rangle_{L^2} (a_{1,0}\partial_{a_1}\rho(\vec{a}_0) + a_{2,0}\partial_{a_2}\rho(\vec{a}_0) + |\vec{a}_0|^2\omega''_{\omega_0}(0)) + o(\rho(\vec{a}_0)|\vec{a}_0|^2).
 \end{aligned} \tag{4.14}$$

By (4.10), we have

$$\begin{aligned}
 & \langle \tilde{\varphi}_{\omega_0}, \partial_\omega\tilde{\varphi}_{\omega_0} \rangle_{L^2} (a_{1,0}\partial_{a_1}\rho(\vec{a}_0) + a_{2,0}\partial_{a_2}\rho(\vec{a}_0)) \\
 = & -\frac{|\vec{a}_0|^2 R_{p,\omega_0}}{2} + o(|\vec{a}_0|^2) \\
 = & -|\vec{a}_0|^2 \left(-\frac{d\lambda_\omega}{d\omega} \Big|_{\omega=\omega_0} \left\| \psi_{\omega_0} \cos \frac{y}{L} \right\|_{L^2}^2 + \omega''_{\omega_0}(0) \langle \partial_\omega\tilde{\varphi}_{\omega_0}, \tilde{\varphi}_{\omega_0} \rangle_{L^2} \right) + o(|\vec{a}_0|^2).
 \end{aligned}$$

Hence, the conclusion follows the equation (4.14). \square

Lemma 4.15. *Let $\varepsilon > 0$ be sufficiently small and $u \in N_\varepsilon^0$ with $S_{\omega_0}(u) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) < 0$. Then*

$$P(u) = -|\vec{a}|^2 \rho(\vec{a}) \frac{d\lambda_\omega}{d\omega} \Big|_{\omega=\omega_0} \left\| \psi_{\omega_0} \cos \frac{y}{L} \right\|_{L^2}^2 + o(\rho(\vec{a}(u))^2) + o(\|w(u)\|_{H^1}^2).$$

Proof. By the Taylor expansion, we have

$$\begin{aligned}
 P(u) & = \langle S'_{\omega_{\omega_0}(\vec{a}(u))}(\Phi(\vec{a}(u)) + w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u))), iA'(\Phi(\vec{a}(u)) + w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u))) \rangle_{H^{-1}, H^1} \\
 & = \langle S'_{\omega_{\omega_0}(\vec{a}(u))}(\Phi(\vec{a}(u))) + S''_{\omega_{\omega_0}(\vec{a}(u))}(\Phi(\vec{a}(u)))(w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u))), \\
 & \quad iA'(\Phi(\vec{a}(u))) + iA''(\Phi(\vec{a}(u)))(w(u) + \alpha(u)\phi_{\omega_0}(\vec{a}(u))) \rangle_{H^{-1}, H^1} + o(\rho(\vec{a}(u))^2 + \|w(u)\|_{H^1}^2)
 \end{aligned}$$

By (4.12), (4.13), Lemma 4.11 and Lemma 4.14,

$$\begin{aligned}
 P(u) & = P(\Phi(\vec{a}(u))) + \langle S'_{\omega_{\omega_0}(\vec{a}(u))}(\Phi(\vec{a}(u))), iA''(\Phi(\vec{a}(u)))w(u) \rangle_{L^2} \\
 & \quad + \langle S''_{\omega_{\omega_0}(\vec{a}(u))}(\Phi(\vec{a}(u)))w(u), iA'(\Phi(\vec{a}(u))) \rangle_{H^{-1}, H^1} + o(\rho(\vec{a}(u))^2 + \|w(u)\|_{H^1}^2)
 \end{aligned}$$

By the proof of Lemma 4.10, we obtain that

$$\frac{\partial G}{\partial(\theta, a_1, a_2)} \begin{pmatrix} \theta' \\ a'_1 \\ a'_2 \end{pmatrix} = \begin{pmatrix} -ie^{-i\theta(u)}\phi_{\omega_0}(\vec{a}(u)) \\ -e^{-i\theta(u)}\psi_{\omega_0,1} \\ -e^{-i\theta(u)}\psi_{\omega_0,2} \end{pmatrix}.$$

Thus $\theta'(\Phi(\vec{a}(u)))$, $a'_1(\Phi(\vec{a}(u)))$ and $a'_2(\Phi(\vec{a}(u)))$ are linear combinations of $i\phi_{\omega_0}(\vec{a}(u))$, $\psi_{\omega_0,1}$ and $\psi_{\omega_0,2}$. Since $\langle \theta'(\Phi(\vec{a}(u))), w(u) \rangle_{L^2} = \langle a'_1(\Phi(\vec{a}(u))), w(u) \rangle_{L^2} = \langle a'_2(\Phi(\vec{a}(u))), w(u) \rangle_{L^2} = O(\alpha(u)\|w(u)\|_{H^1})$, we have

$$iA''(\Phi(\vec{a}(u)))w(u) = O(\alpha(u)\|w(u)\|_{H^1}).$$

Therefore, by the orthogonal condition of $w(u)$ and $A'(\Phi(\vec{a}(u))) = O(\vec{a}(u))$ we obtain

$$\begin{aligned}
P(u) &= P(\Phi(\vec{a}(u))) + (\omega_{\omega_0}(\vec{a}(u)) - \omega_0) \langle w(u), iA'(\Phi(\vec{a}(u))) \rangle_{L^2} \\
&\quad + \langle p(|\Phi(\vec{a}(u))|^{p-1} - |\tilde{\varphi}_{\omega_0}|^{p-1})w(u), iA'(\Phi(\vec{a}(u))) \rangle_{L^2} \\
&\quad + \langle S''_{\omega_0}(\tilde{\varphi}_{\omega_0})w(u), \psi_{\omega_0} \left(a_1(u) \cos \frac{y}{L} + a_2(u) \sin \frac{y}{L} \right) \rangle_{H^{-1}, H^1} \\
&\quad + \langle S''_{\omega_0}(\tilde{\varphi}_{\omega_0})w(u), (a_1(u) \partial_{a_1} \rho(\vec{a}(u)) + a_2(u) \partial_{a_2} \rho(\vec{a}(u))) \partial_\omega \tilde{\varphi}_{\omega_0} \rangle_{H^{-1}, H^1} \\
&\quad + \langle w(u), -|\vec{a}(u)|^2 \omega''_{\omega_0}(0) \tilde{\varphi}_{\omega_0} + p(p-1) (\tilde{\varphi}_{\omega_0})^{p-2} \psi_{\omega_0}^2 \left(a_1(u) \cos \frac{y}{L} + a_2(u) \sin \frac{y}{L} \right)^2 \rangle_{H^{-1}, H^1} \\
&\quad + o(\rho(\vec{a}(u))^2 + \|w(u)\|_{H^1}^2) \\
&= P(\Phi(\vec{a}(u))) + o(\rho(\vec{a}(u))^2 + \|w(u)\|_{H^1}^2)
\end{aligned}$$

Hence, we obtain the conclusion. \square

We assume $e^{i\omega_0 t} \tilde{\varphi}_{\omega_0}$ is stable. Let $\{\vec{a}_n\}_n$ be a sequence with $\vec{a}_n \rightarrow 0$ and $\{u_n\}_n$ be the sequence of solutions with $u_n(0) = \Phi(\vec{a}_n)$. Since $R_{p, \omega_0} < 0$ and there exists $C > 0$ such that

$$S_{\omega_0}(\Phi(\vec{a}_n)) - S_{\omega_0}(\tilde{\varphi}_{\omega_0}) = CR_{p, \omega_0} |\vec{a}_n|^4 + o(|\vec{a}_n|^4),$$

we obtain $S_{\omega_0}(\tilde{\varphi}_{\omega_0}) > S_{\omega_0}(\Phi(\vec{a}_n))$ for sufficiently large $n > 1$. From Lemma 4.13 and Lemma 4.15 we have for sufficiently large $n > 1$

$$\begin{aligned}
0 &< S_{\omega_0}(\tilde{\varphi}_{\omega_0}) - S_{\omega_0}(\Phi(\vec{a}_n)) \\
&= S_{\omega_0}(\tilde{\varphi}_{\omega_0}) - S_{\omega_0}(u_n(t)) \\
&\leq -C_{**} R_{p, \omega_0} |\vec{a}(u_n(t))|^4 - \frac{k_0}{2} \|w(u_n(t))\|_{H^1}^2 + o(\|w(u_n(t))\|_{H^1}^2) + o(|\vec{a}(u_n(t))|^4).
\end{aligned}$$

By the stability of $e^{i\omega_0 t} \tilde{\varphi}_{\omega_0}$ and the equation (4.10), we obtain there exists $c > 0$ such that for sufficiently large $n > 1$

$$0 < S_{\omega_0}(\tilde{\varphi}_{\omega_0}) - S_{\omega_0}(\Phi(\vec{a}_n)) \leq cP(u_n(t)).$$

Since $\rho(\vec{a}(u_n(t)))$ is positive and bounded for $t \geq 0$ and sufficiently large $n > 1$, there exists $\delta > 0$ such that for $t \geq 0$

$$\frac{dA(u_n(t))}{dt} = P(u_n(t)) > \delta.$$

This contradicts the boundedness of A on N_ε . Hence, $e^{i\omega_0 t} \tilde{\varphi}_{\omega_0}$ is unstable.

Acknowledgments

The author would like to express his great appreciation to Professor Yoshio Tsutsumi for a lot to helpful advices and encouragements. The author would like to thank Professor Mashahito Ohta for his helpful indication.

References

- [1] J. C. Alexander, R. L. Pego and R. L. Sachs, *On the transverse instability of solitary waves in the Kadomtsev-Petviashvili equation*, Phys. Lett. A (1997), no. 3-4, 187–192.
- [2] B. Deconinck, D. E. Pelinovsky and J. D. Carter, *Transverse instabilities of deep-water solitary waves*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **462** (2006), no.2071, 2671–2694.
- [3] M. Colin, T. Colin and M. Ohta, *Stability of solitary waves for a system of nonlinear Schrödinger equations with three wave interaction*, Ann. I. Poincaré-AN, **26** (2009), 2211–2226.
- [4] M. Colin and M. Ohta, *Bifurcation from semitrivial standing waves and ground states for a system of nonlinear Schrödinger equations*, SIAM J. Math. Anal., **44** (2012), no. 1, 206–223.
- [5] V. Georgiev and M. Ohta, *Nonlinear instability of linearly unstable standing waves for nonlinear Schrödinger equations*, Journal of the Mathematical Society of Japan **64** (2012), no. 2, 533–548.
- [6] F. Gesztesy, C. K. R. T. Jones, Y. Latushkin and M. Stanislavova, *A spectral mapping theorem and invariant manifolds for nonlinear Schrödinger equations*, Indiana Univ. Math. J., **49** (2000), 221–243.
- [7] J. Ginibre and G. Velo, *On a class of nonlinear Schrödinger equations. I. The Cauchy problem, general case*, J. Funct. Anal. **32** (1979), 1-32.
- [8] J. Ginibre and G. Velo, *Scattering theory in the energy space for a class of nonlinear wave equations*, Comm. Math. Phys., **123** (1989), 535–573.
- [9] E. Grenier, *On the nonlinear instability of Euler and Prandtl equations*, Comm. Pure Appl. Math. **53** (2000) 1067–1091.
- [10] M. Grillakis, J. Shatah and W. Strauss, *Stability theory of solitary waves in the presence of symmetry. I*, J. Funct. Anal. A **74** (1987), no. 1, 160–197.
- [11] M. Grillakis, J. Shatah and W. Strauss, *Stability theory of solitary waves in the presence of symmetry II*, J. Funct. Anal. **94** (1990), 308–348.
- [12] E. Kirr, P. G. Kevrekidis and D. E. Pelinovsky, *Symmetry-breaking bifurcation in the nonlinear Schrödinger equation with symmetric potentials*, Comm. Math. Phys. **308** (2011), no. 3, 795–844.
- [13] M. Maeda, *Stability of bound states of Hamiltonian PDEs in the degenerate cases*, J. Funct. Anal. **263** (2012), no. 2, 511–528.
- [14] M. Ohta, *Instability of bound states for abstract nonlinear Schrödinger equations*, J. Funct. Anal. **261** (2011), no. 1, 90-110.

- [15] T. Mizumachi, *Stability of line solitons for the KP-II equation in \mathbb{R}^2* , arXiv:1303.3532, 1–76.
- [16] T. Mizumachi and N. Tzvetkov, *Stability of the line soliton of the KP-II equation under periodic transverse perturbations*, Math. Ann. **352** (2012), no. 3, 659–690.
- [17] R. Pego and M. I. Weinstein, *Eigenvalues, and instabilities of solitary waves*, Phil. Trans. R. Soc. London A, **340** (1992), 47–94.
- [18] H. A. Rose and M. I. Weinstein, *On the bound states of the nonlinear Schrödinger equation with a linear potential*, Physica D **30** (1988), 207–218.
- [19] F. Rousset and N. Tzvetkov, *Transverse nonlinear instability of solitary waves for some Hamiltonian PDE's*, J. Math. Pures. Appl. **90** (2008) 550–590.
- [20] F. Rousset and N. Tzvetkov, *Transverse nonlinear instability for two-dimensional dispersive models*, Ann. I. Poincaré-AN **26** (2009) 477–496.
- [21] F. Rousset and N. Tzvetkov, *A simple criterion of transverse linear instability for solitary waves*, Math. Res. Lett., **17** (2010), no. 1, 157–169.
- [22] F. Rousset and N. Tzvetkov, *Stability and instability of the KdV solitary wave under the KP-I flow*, Comm. Math. Phys., **313** (2012), no. 1, 155–173.
- [23] J. Shatah and W. Strauss, *Spectral condition for instability*, Contemp. Math., **255** (2000), 189–198.
- [24] H. Takaoka and N. Tzvetkov, *On 2D nonlinear Schrödinger equations with Data on $\mathbb{R} \times \mathbb{T}$* , J. Funct. Anal. **182** (2001) 427–442.
- [25] Y. Yamazaki, *Transverse instability for a system of nonlinear Schrödinger equations*, Discrete Contin. Dyn. Syst. Ser. B **19** (2014), no.2, 565–588.
- [26] Y. Yamazaki, *Stability of line standing waves near the bifurcation point for nonlinear Schrodinger equations*, appear to Kodai Math. J.

Yohei Yamazaki
 Department of Mathematics
 Kyoto University
 Kyoto 606-8502
 Japan
 E-mail address: y-youhei@math.kyoto-u.ac.jp