

# The discrete-time quaternionic quantum walk on a graph

Norio Konno

Department of Applied Mathematics, Faculty of Engineering, Yokohama National University

Hodogaya, Yokohama 240-8501, Japan

e-mail: konno@ynu.ac.jp

Hideo Mitsunashi

Faculty of Education, Utsunomiya University

Utsunomiya, Tochigi 321-8505, Japan

e-mail: mitsu@cc.utsunomiya-u.ac.jp

Iwao Sato

Oyama National College of Technology

Oyama, Tochigi 323-0806, Japan

e-mail: isato@oyama-ct.ac.jp

**Abstract.** Recently, the quaternionic quantum walk was formulated by the first author as a generalization of discrete-time quantum walks. We treat the right eigenvalue problem of quaternionic matrices to analysis the spectra of its transition matrix. The way to obtain all the right eigenvalues of a quaternionic matrix is given. From the unitary condition on the transition matrix of the quaternionic quantum walk, we deduce some properties about it. Our main results, Theorem 5.3, determine all the right eigenvalues of a quaternionic quantum walk by use of those of the corresponding weighted matrix. In addition, we give some examples of quaternionic quantum walks and their right eigenvalues.

## 1 Introduction

As a quantum version of the classical random walk, the quantum walk has recently been studied in various fields. The discrete-time quantum walk in one dimension lattice was intensively studied by Ambainis et al. [2]. One of the most striking properties is the spreading property of the walker. The standard deviation of the position grows linearly in time,

---

*Abbr. title:* The discrete-time quaternionic quantum walk on a graph

*AMS 2010 subject classifications:* 60F05, 05C50, 15A15, 11R52

*PACS:* 03.67.Lx, 05.40.Fb, 02.50.Cw

*Keywords:* Quantum walk, Ihara zeta function, quaternion, quaternionic quantum walk

quadratically faster than classical random walk. On the other hand, a walker stays at the starting position, i.e., localization occurs. Konno [18] extended the quantum walk to a *quaternionic quantum walk* determined by a unitary matrix whose component is quaternion, and presented some properties of the walk. The reviews and books on quantum walks are Kempe [15], Konno [17], Venegas-Andraca [30], Cantero et al. [6], Manouchehri and Wang [22], Portugal [23] for examples.

For a general graph, a typical case of discrete-time quantum walks is the Grover walk on a graph. The Grover walk on a graph was formulated in [11]. Emms et al. [8] treated spectra of the transition matrix and its positive support of the Grover walk on a graph, and showed that the third power of the transition matrix outperforms the graph spectra methods in distinguishing strongly regular graphs. Godsil and Guo [10] gave new proofs of the results of Emms et al. [8].

The discrete-time quantum walk on a graph is closely related to the Ihara zeta function of a graph. Zeta functions of graphs originally started from the Ihara zeta function for a regular graph by Ihara [14]. The Ihara zeta function of a graph was studied by many researchers [14], [26], [28], [29], [12], [4], [27], [9], [20]. Bass [4] generalized Ihara's result on the zeta function of a regular graph to an irregular graph, and showed that its reciprocal is again a polynomial. Inspired by these works, Sato [25] defined a new zeta function (the second weighted zeta function) of a graph by using not an infinite product but a determinant. This new zeta function and its determinant formula played an essential role in the determination of eigenvalues of quantum walk on a graph in Konno and Sato [19]. Ren et al. [24] found an interesting relation between the Ihara zeta function and the Grover walk on a graph, and showed that the positive support of the transition matrix of the Grover walk is equal to the Perron-Frobenius operator (the edge matrix) related to the Ihara zeta function. Konno and Sato [19] gave the characteristic polynomials of the transition matrix of the Grover walk and its positive support by using the second weighted zeta function and the Ihara zeta function, and so obtained the other proofs of the results on spectra for them by Emms et al. [8].

Quaternion, which can be considered as an extension of complex number, was discovered by Hamilton in 1843. The eigenvalue problem for matrices whose entries are quaternions has been investigated for nearly a century by a number of researchers. A detailed survey on the quaternionic matrices can be found in Zhang [31]. The notable difference from the eigenvalue problem for complex matrices is that it is necessary to treat left eigenvalues and right eigenvalues separately. Right eigenvalues are well studied by, for example, Brenner [5] and Lee [21]. On the contrary, left eigenvalues are less known and not easy to handle as commented in Huang and So [13]. Recently, an extension of quantum walk to the case of quaternions was established by Konno [18] in which a model of quaternionic quantum walk was suggested and its some properties were presented. In the present paper, we extend a discrete-time quantum walk (Grover walk) on a graph to a walk given by a unitary matrix whose components are quaternions (quaternionic Grover walk). We derive the condition for components of transition matrix of quaternionic Grover walk. Furthermore, we deal with right eigenvalue problem, and derive all the right eigenvalues of quaternionic transition matrix from those of the corresponding weighted matrix. Our results can be regarded as a generalization of [8], [19].

The rest of the paper is organized as follows. Section 2 treats the right eigenvalue problem of quaternionic matrices. The way to obtain all the right eigenvalues of a quaternionic matrix

is given. In Section 3, we provide a summary of the Ihara zeta function and the second weighted zeta function of a graph, and present their determinant expressions. In Section 4, we show a brief account of the discrete-time quantum walk on a graph, and define a generalization of it to the case of quaternions, which we call the quaternionic quantum walk on a graph. From the unitary condition on the transition matrix, we deduce some properties about it. In Section 5, our main results, Theorem 5.3, determine all the right eigenvalues of a quaternionic quantum walk by use of those of the corresponding weighted matrix. In addition, we give some examples of quaternionic quantum walks and their right eigenvalues.

## 2 Right eigenvalues of a quaternionic matrix

In this section, we shall show the way to calculate all the right eigenvalues of a quaternionic matrix. Since quaternions do not mutually commute in general, we must treat left eigenvalue and right eigenvalue separately. Since  $\mathbf{M}^n \mathbf{v} = \mathbf{v} \lambda^n$  ( $n \geq 1$ ) holds for an right eigenvalue  $\lambda$  of  $\mathbf{M}$  and an eigenvector  $\mathbf{v}$  corresponding to  $\lambda$ , right eigenvalue is more suitable to calculate the stationary measure of the quaternionic quantum walk than left eigenvalue. Therefore, we treat only right eigenvalues in this paper. Although we cannot apply the ordinary characteristic polynomial for a quaternionic matrix, we can calculate all the complex right eigenvalues of it by embedding it in a complex matrix twice the size of it. From these eigenvalues, we can obtain all the right eigenvalues of the quaternionic matrix. An exposition of these contents can be found in [3]. [31] gives an overview of quaternionic matrix theory.

Let  $\mathbb{H}$  be the set of quaternions.  $\mathbb{H}$  is a noncommutative associative algebra over  $\mathbb{R}$ , whose underlying real vector space has dimension 4 with a basis  $1, i, j, k$  which satisfy the following relations:

$$i^2 = j^2 = k^2 = -1, \quad ij = -ji = k, \quad jk = -kj = i, \quad ki = -ik = j.$$

For  $x = x_0 + x_1i + x_2j + x_3k \in \mathbb{H}$ , we call  $x_0$  the *real part*,  $x_0 + x_1i$  the *complex part*,  $x_1i + x_2j + x_3k$  the *imaginary part* of  $x$ , and set  $\operatorname{Re} x = x_0$ ,  $\operatorname{Co} x = x_0 + x_1i$ ,  $\operatorname{Im} x = x_1i + x_2j + x_3k$  respectively.  $x^*$  denotes the *conjugate* of  $x$  in  $\mathbb{H}$  which is defined by  $x^* = x_0 - x_1i - x_2j - x_3k$ . We call  $|x| = \sqrt{xx^*} = \sqrt{x^*x} = \sqrt{x_0^2 + x_1^2 + x_2^2 + x_3^2}$  the *norm* of  $x$ . Indeed,  $|\cdot|$  satisfies

- (1)  $|x| \geq 0$ , and moreover  $|x| = 0 \Leftrightarrow x = 0$ ,
- (2)  $|xy| = |x||y|$ ,
- (3)  $|x + y| \leq |x| + |y|$ .

For a nonzero element  $x \in \mathbb{H}$ ,  $x^{-1} = x^*/|x|^2$ . Hence,  $\mathbb{H}$  constitutes a skew field.

We can present arbitrary quaternion  $x$  by two complex numbers  $x = a + jb$  uniquely. Such a presentation is called *symplectic decomposition*. Two complex numbers  $a$  and  $b$  are called *simplex part* and *perplex part* of  $x$  respectively. Symplectic decomposition is also valid for a quaternionic matrix, namely a matrix whose entries are quaternions.  $\operatorname{Mat}(m \times n, \mathbb{H})$  denotes the set of  $m \times n$  quaternionic matrices and  $\operatorname{Mat}(n, \mathbb{H})$  the set of  $n \times n$  quaternionic square matrices. For  $\mathbf{M} \in \operatorname{Mat}(m \times n, \mathbb{H})$ , we can write  $\mathbf{M} = \mathbf{A} + j\mathbf{B}$  uniquely where  $\mathbf{A}, \mathbf{B} \in \operatorname{Mat}(m \times n, \mathbb{C})$ .  $\mathbf{A}$  and  $\mathbf{B}$  are called *simplex part* and *perplex part* of  $\mathbf{M}$  respectively. We define  $\psi$  to be the map from  $\operatorname{Mat}(m \times n, \mathbb{H})$  to  $\operatorname{Mat}(2m \times 2n, \mathbb{C})$  as follows:

$$\psi : \operatorname{Mat}(m \times n, \mathbb{H}) \longrightarrow \operatorname{Mat}(2m \times 2n, \mathbb{C}) \quad \mathbf{M} \mapsto \begin{bmatrix} \mathbf{A} & -\overline{\mathbf{B}} \\ \mathbf{B} & \overline{\mathbf{A}} \end{bmatrix},$$

where  $\overline{\mathbf{A}}$  is the complex conjugate of  $\mathbf{A}$ . Then  $\psi$  is an  $\mathbb{R}$ -linear map. We also have:

LEMMA 2.1 *Let  $\mathbf{M} \in \text{Mat}(m \times n, \mathbb{H})$  and  $\mathbf{N} \in \text{Mat}(n \times m, \mathbb{H})$ . Then*

$$\psi(\mathbf{MN}) = \psi(\mathbf{M})\psi(\mathbf{N}).$$

**Proof.** Let  $\mathbf{M} = \mathbf{A} + j\mathbf{B}$  and  $\mathbf{N} = \mathbf{C} + j\mathbf{D}$  be symplectic decompositions of  $\mathbf{M}$  and  $\mathbf{N}$ . Then,

$$\mathbf{MN} = (\mathbf{A} + j\mathbf{B})(\mathbf{C} + j\mathbf{D}) = \mathbf{AC} + \mathbf{A}j\mathbf{D} + j\mathbf{BC} + j\mathbf{B}j\mathbf{D}.$$

Since  $\mathbf{X}j = j\overline{\mathbf{X}}$  for arbitrary complex matrix  $\mathbf{X}$ , it follows that:

$$\mathbf{MN} = \mathbf{AC} - \overline{\mathbf{B}}\mathbf{D} + j(\overline{\mathbf{A}}\mathbf{D} + \mathbf{BC}),$$

and therefore

$$\psi(\mathbf{MN}) = \begin{bmatrix} \mathbf{AC} - \overline{\mathbf{B}}\mathbf{D} & -\overline{\mathbf{A}}\mathbf{D} - \overline{\mathbf{B}}\mathbf{C} \\ \overline{\mathbf{A}}\mathbf{D} + \mathbf{BC} & \overline{\mathbf{AC}} - \mathbf{BD} \end{bmatrix}.$$

On the other hand,

$$\psi(\mathbf{M})\psi(\mathbf{N}) = \begin{bmatrix} \mathbf{A} & -\overline{\mathbf{B}} \\ \mathbf{B} & \overline{\mathbf{A}} \end{bmatrix} \begin{bmatrix} \mathbf{C} & -\overline{\mathbf{D}} \\ \mathbf{D} & \overline{\mathbf{C}} \end{bmatrix} = \begin{bmatrix} \mathbf{AC} - \overline{\mathbf{B}}\mathbf{D} & -\overline{\mathbf{A}}\mathbf{D} - \overline{\mathbf{B}}\mathbf{C} \\ \mathbf{BC} + \overline{\mathbf{A}}\mathbf{D} & -\mathbf{BD} + \overline{\mathbf{AC}} \end{bmatrix}.$$

Thus  $\psi(\mathbf{MN}) = \psi(\mathbf{M})\psi(\mathbf{N})$  holds.  $\square$

PROPOSITION 2.2 *If  $m = n$ , then  $\psi$  is an injective  $\mathbb{R}$ -algebra homomorphism.*

**Proof.** By Lemma 2.1,  $\psi$  is an  $\mathbb{R}$ -algebra homomorphism. Injectivity of  $\psi$  is clear.  $\square$

For a quaternionic square matrix  $\mathbf{M} = (\mathbf{M}_{ij}) \in \text{Mat}(n, \mathbb{H})$ , the *conjugate*  $\mathbf{M}^* = ((\mathbf{M}^*)_{ij})$  is defined by  $(\mathbf{M}^*)_{ij} = (\mathbf{M}_{ji})^*$ . We notice that  $\psi(\mathbf{M}^*) = \psi(\mathbf{M})^*$ , where the right-hand side denotes the conjugate transpose of the complex matrix. A quaternionic square matrix  $\mathbf{M}$  is said to be *unitary* if  $\mathbf{M}^*\mathbf{M} = \mathbf{M}\mathbf{M}^* = \mathbf{I}$ . Since  $\psi$  is an injective  $\mathbb{R}$ -algebra homomorphism,  $\mathbf{M}^*\mathbf{M} = \mathbf{I}$  implies  $\mathbf{M}\mathbf{M}^* = \mathbf{I}$ .

REMARK 2.3  $\psi({}^T\mathbf{M}) = {}^T\psi(\mathbf{M})$  does not hold in general, where  ${}^T\mathbf{M}$  denotes the transpose of  $\mathbf{M}$ .

Let  $\mathbb{H}^n$  be a right vector space, and  $\mathbf{M} \in \text{Mat}(n, \mathbb{H})$ ,  $\mathbf{v} \in \mathbb{H}^n$ ,  $\lambda \in \mathbb{H}$  satisfy  $\mathbf{M}\mathbf{v} = \mathbf{v}\lambda$ .  $\lambda$  is said to be a right eigenvalue of  $\mathbf{M}$ , and  $\mathbf{v}$  a right eigenvector corresponding to  $\lambda$ . We denote by  $\sigma_r(\mathbf{M})$  the set of all the right eigenvalues of  $\mathbf{M}$ . We have  $\mathbf{M}(\mathbf{v}q) = \mathbf{v}\lambda q = \mathbf{v}q(q^{-1}\lambda q)$  for any  $q \in \mathbb{H}^* = \mathbb{H} - \{0\}$ . Hence, the conjugate class  $\lambda^{\mathbb{H}^*} = \{q^{-1}\lambda q \mid q \in \mathbb{H}^*\}$  is contained in  $\sigma_r(\mathbf{M})$ . We notice that if  $\lambda \in \mathbb{R}$ , then  $\lambda^{\mathbb{H}^*} = \{\lambda\}$ . Since  $|q^{-1}xq| = |x|$ , the group homomorphism:

$$\rho : \mathbb{H}^* \longrightarrow \text{Aut } \mathbb{H}, \quad \rho(q)x = qxq^{-1}$$

gives orthogonal transformations  $\rho(q)$  ( $q \in \mathbb{H}^*$ ) on  $\mathbb{H}$ . Let  $\mathbb{H}_R = \mathbb{R}$  and  $\mathbb{H}_P = \mathbb{R}i \oplus \mathbb{R}j \oplus \mathbb{R}k$ . Then  $\mathbb{H} = \mathbb{H}_R \oplus \mathbb{H}_P$  and  $\mathbb{H}_R \perp \mathbb{H}_P$ . We call the elements of  $\mathbb{H}_P$  *pure* quaternion. One can see that  $\rho(q)x = x$  ( $x \in \mathbb{H}_R$ ) and  $\rho(q)\mathbb{H}_P = \mathbb{H}_P$ , hence  $\rho$  also gives orthogonal transformations  $\rho(q)$  ( $q \in \mathbb{H}^*$ ) on  $\mathbb{H}_P$ . Moreover it is well-known that  $\det(\rho(q)) = 1$  and that the group homomorphism:

$$\rho|_{\mathbb{H}_P} : \mathbb{H}^* \longrightarrow SO(\mathbb{H}_P) \cong SO(3)$$

is surjective (for details, see for example [1],[7]). Therefore,  $\mathbb{H}^*$  gives all the rotations in  $\mathbb{H}_P \cong \mathbb{R}^3$ .

If  $\lambda \in \mathbb{H} - \mathbb{R}$ , then we may write  $\lambda = \lambda_0 + \lambda'$  ( $\lambda_0 \in \mathbb{H}_R$ ,  $\lambda' \in \mathbb{H}_P$ ) with  $\lambda' \neq 0$ . Then  $\rho(q)\lambda = q\lambda q^{-1} = \lambda_0 + q\lambda'q^{-1}$ . It follows that  $\lambda^{\mathbb{H}^*}$  is a 2-sphere centering at the origin in  $\mathbb{H}_P$  since the following equations hold:

$$\lambda^{\mathbb{H}^*} = \{q^{-1}\lambda'q \mid q \in \mathbb{H}^*\} = \rho(\mathbb{H}^*)\lambda' = SO(\mathbb{H}_P)\lambda'. \quad (2.1)$$

Thus the intersection of  $\lambda^{\mathbb{H}^*}$  and the complex axis consists of just two points,  $\lambda^{\mathbb{H}^*} \cap \mathbb{R}i = \{\pm|\lambda'|i\}$ . To summarize our discussion, we find out the following:

**PROPOSITION 2.4** *Let  $\mathbf{M} \in \text{Mat}(n, \mathbb{H})$ . The  $\mathbb{H}^*$ -conjugate class of any right eigenvalue of  $\mathbf{M}$  which is not real is contained in  $\sigma_r(\mathbf{M})$ , and has exactly two complex numbers that are conjugate.*

We shall show that one can calculate all the complex right eigenvalues of a quaternionic matrix by use of ordinary determinant.

**PROPOSITION 2.5** *Let  $\mathbf{M} \in \text{Mat}(n, \mathbb{H})$  and  $p \in \mathbb{C}$ . Then*

$$\mathbf{M}\mathbf{v} = \mathbf{v}p \Leftrightarrow \psi(\mathbf{M}) \begin{bmatrix} \mathbf{u} \\ \mathbf{w} \end{bmatrix} = \begin{bmatrix} \mathbf{u} \\ \mathbf{w} \end{bmatrix} p \quad \text{where } \mathbf{v} = \mathbf{u} + j\mathbf{w} \ (\mathbf{u}, \mathbf{w} \in \mathbb{C}^n).$$

**Proof.** Let  $\mathbf{M} = \mathbf{A} + j\mathbf{B}$  be the symplectic decomposition. Then we have:

$$\mathbf{M}\mathbf{v} = (\mathbf{A} + j\mathbf{B})(\mathbf{u} + j\mathbf{w}) = (\mathbf{A}\mathbf{u} - \overline{\mathbf{B}}\mathbf{w}) + j(\mathbf{B}\mathbf{u} + \overline{\mathbf{A}}\mathbf{w}).$$

Hence  $\mathbf{M}\mathbf{v} = \mathbf{v}p$  is equivalent to the following equation:

$$\begin{bmatrix} \mathbf{A} & -\overline{\mathbf{B}} \\ \mathbf{B} & \overline{\mathbf{A}} \end{bmatrix} \begin{bmatrix} \mathbf{u} \\ \mathbf{w} \end{bmatrix} = \begin{bmatrix} \mathbf{u} \\ \mathbf{w} \end{bmatrix} p.$$

□

In the same way as the ordinary triangularization of a complex matrix, the triangularization of a quaternionic matrix can also be obtained. Moreover, we can triangularize a quaternionic matrix into a triangular matrix whose diagonals are complex as follows. The detail of the proof can be found in [21].

**LEMMA 2.6** *For arbitrary  $\mathbf{M} \in \text{Mat}(n, \mathbb{H})$ , there exist  $\mathbf{V} \in \text{Mat}(n, \mathbb{H})$  with  $\mathbf{V}^*\mathbf{V} = \mathbf{V}\mathbf{V}^* = \mathbf{I}_n$  and  $\lambda_1, \dots, \lambda_n \in \mathbb{C}$  such that*

$$\mathbf{V}^*\mathbf{M}\mathbf{V} = \begin{bmatrix} \lambda_1 & * & \cdots & * \\ 0 & \lambda_2 & & * \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix}. \quad (2.2)$$

As a consequence of Proposition 2.5 and Lemma 2.6, we can deduce the following:

PROPOSITION 2.7 Let  $\mathbf{M} \in \text{Mat}(n, \mathbb{H})$ . Then  $2n$  eigenvalues of  $\psi(\mathbf{M}) \in \text{Mat}(2n, \mathbb{C})$  occur in conjugate pairs:

$$\lambda_1, \overline{\lambda_1}, \lambda_2, \overline{\lambda_2}, \dots, \lambda_n, \overline{\lambda_n},$$

counted with multiplicity.

**Proof.** The proof can be found in [21]. Let  $\mathbf{T}$  be the right hand side of (2.2) and  $\mathbf{T} = \mathbf{T}^S + j\mathbf{T}^P$  its symplectic decomposition. Then,

$$\begin{aligned} \det(\lambda \mathbf{I}_{2n} - \psi(\mathbf{M})) &= \det(\lambda \mathbf{I}_{2n} - \psi(\mathbf{V})^{-1} \psi(\mathbf{M}) \psi(\mathbf{V})) \\ &= \det(\lambda \mathbf{I}_{2n} - \psi(\mathbf{V}^* \mathbf{M} \mathbf{V})) \\ &= \det(\lambda \mathbf{I}_{2n} - \psi(\mathbf{T})) = \det \begin{bmatrix} \lambda \mathbf{I}_{2n} - \mathbf{T}^S & \overline{\mathbf{T}}^P \\ -\mathbf{T}^P & \lambda \mathbf{I}_{2n} - \overline{\mathbf{T}}^S \end{bmatrix} \\ &= \det \begin{bmatrix} \lambda - \lambda_1 & * & \cdots & * & 0 & * & \cdots & * \\ 0 & \lambda - \lambda_2 & & * & 0 & 0 & & * \\ \vdots & & \ddots & \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda - \lambda_n & 0 & 0 & \cdots & 0 \\ 0 & * & \cdots & * & \lambda - \overline{\lambda_1} & * & \cdots & * \\ 0 & 0 & & * & 0 & \lambda - \overline{\lambda_2} & & * \\ \vdots & & \ddots & \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & \lambda - \overline{\lambda_n} \end{bmatrix} \\ &= (\lambda - \lambda_1)(\lambda - \overline{\lambda_1}) \cdots (\lambda - \lambda_n)(\lambda - \overline{\lambda_n}). \end{aligned}$$

□

Hence, diagonal elements in  $\mathbf{T}$  and their complex conjugates are eigenvalues of  $\psi(\mathbf{M})$ . We notice that we may replace  $\lambda_i$  by its complex conjugate in (2.2) because of Proposition 2.4. Combining Proposition 2.4, 2.5, 2.7, we can state about eigenvalues of a quaternionic matrix as follows:

THEOREM 2.8 For arbitrary quaternionic matrix  $\mathbf{M} \in \text{Mat}(n, \mathbb{H})$ ,  $2n$  complex right eigenvalues  $\lambda_1, \dots, \lambda_n, \overline{\lambda_1}, \dots, \overline{\lambda_n}$  of  $\mathbf{M}$  exist counted with multiplicity, which can be calculated by solving  $\det(\lambda \mathbf{I}_{2n} - \psi(\mathbf{M})) = 0$ . The set of right eigenvalues  $\sigma_r(\mathbf{M})$  is given by  $\sigma_r(\mathbf{M}) = \lambda_1^{\mathbb{H}^*} \cup \dots \cup \lambda_n^{\mathbb{H}^*}$ .

EXAMPLE 2.9  $\mathbf{M} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$

$$\det(\lambda \mathbf{I}_4 - \psi(\mathbf{M})) = \det \begin{bmatrix} \lambda & -1 & 0 & 0 \\ 1 & \lambda & 0 & 0 \\ 0 & 0 & \lambda & -1 \\ 0 & 0 & 1 & \lambda \end{bmatrix} = (\lambda - i)^2 (\lambda + i)^2. \quad \sigma_r(\mathbf{M}) = i^{\mathbb{H}^*}.$$

EXAMPLE 2.10  $\mathbf{M} = \begin{bmatrix} 1 & 0 \\ 0 & i \end{bmatrix}$

$$\det(\lambda \mathbf{I}_4 - \psi(\mathbf{M})) = \det \begin{bmatrix} \lambda - 1 & 0 & 0 & 0 \\ 0 & \lambda - i & 0 & 0 \\ 0 & 0 & \lambda - 1 & 0 \\ 0 & 0 & 0 & \lambda + i \end{bmatrix} = (\lambda - 1)^2 (\lambda - i) (\lambda + i).$$

$$\sigma_r(\mathbf{M}) = \{1\} \cup i^{\mathbb{H}^*}.$$

EXAMPLE 2.11  $\mathbf{M} = \begin{bmatrix} 0 & i \\ j & 0 \end{bmatrix} = \begin{bmatrix} 0 & i \\ 0 & 0 \end{bmatrix} + j \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$

$$\det(\lambda \mathbf{I}_4 - \psi(\mathbf{M})) = \det \begin{bmatrix} \lambda & -i & 0 & 0 \\ 0 & \lambda & 1 & 0 \\ 0 & 0 & \lambda & i \\ -1 & 0 & 0 & \lambda \end{bmatrix} = (\lambda^2 - i)(\lambda^2 + i)$$

$$= \left( \lambda - \frac{1+i}{\sqrt{2}} \right) \left( \lambda - \frac{-1-i}{\sqrt{2}} \right) \left( \lambda - \frac{1-i}{\sqrt{2}} \right) \left( \lambda - \frac{-1+i}{\sqrt{2}} \right).$$

$$\sigma_r(\mathbf{M}) = \left( \frac{1+i}{\sqrt{2}} \right)^{\mathbb{H}^*} \cup \left( \frac{-1-i}{\sqrt{2}} \right)^{\mathbb{H}^*}.$$

### 3 The Ihara zeta function of a graph

Let  $G = (V(G), E(G))$  be a finite connected graph with the set  $V(G)$  of vertices and the set  $E(G)$  of unoriented edges  $uv$  joining two vertices  $u$  and  $v$ . We assume that  $G$  has neither loops nor multiple edges throughout. For  $uv \in E(G)$ , an arc  $(u, v)$  is the oriented edge from  $u$  to  $v$ . Let  $D(G) = \{(u, v), (v, u) \mid uv \in E(G)\}$  and  $|V(G)| = n$ ,  $|E(G)| = m$ ,  $|D(G)| = 2m$ . For  $e = (u, v) \in D(G)$ ,  $o(e) = u$  denotes the *origin* and  $t(e) = v$  the *terminal* of  $e$  respectively. Furthermore, let  $e^{-1} = (v, u)$  be the *inverse* of  $e = (u, v)$ . The *degree*  $\deg v = \deg_G v$  of a vertex  $v$  of  $G$  is the number of edges incident to  $v$ . A *path*  $P$  of length  $\ell$  in  $G$  is a sequence  $P = (e_1, \dots, e_\ell)$  of  $\ell$  arcs such that  $e_i \in D(G)$  and  $t(e_i) = o(e_{i+1})$  for  $i \in \{1, \dots, \ell - 1\}$ . We set  $o(P) = o(e_1)$  and  $t(P) = t(e_\ell)$ .  $|P|$  denotes the length of  $P$ . We say that a path  $P = (e_1, \dots, e_\ell)$  has a *backtracking* if  $e_{i+1} = e_i^{-1}$  for some  $i(1 \leq i \leq \ell - 1)$ , and that  $P = (e_1, \dots, e_\ell)$  has a *tail* if  $e_\ell = e_1^{-1}$ . A path  $P$  is said to be a *cycle* if  $t(P) = o(P)$ . The *inverse* of a path  $P = (e_1, \dots, e_\ell)$  is the path  $(e_\ell^{-1}, \dots, e_1^{-1})$  and is denoted by  $P^{-1}$ .

We introduce an equivalence relation between cycles. Two cycles  $C_1 = (e_1, \dots, e_\ell)$  and  $C_2 = (f_1, \dots, f_\ell)$  are said to be *equivalent* if there exists  $k$  such that  $f_j = e_{j+k}$  for all  $j$  where indices are treated modulo  $\ell$ . Let  $[C]$  be the equivalence class which contains the cycle  $C$ . Let  $B^r$  be the cycle obtained by going  $r$  times around a cycle  $B$ . Such a cycle is called a *power* of  $B$ . A cycle  $C$  is said to be *reduced* if  $C$  has no backtracking and no tail. Furthermore, a cycle  $C$  is said to be *prime* if it is not a power of a strictly smaller cycle.

The *Ihara zeta function* of a graph  $G$  is a function of  $t \in \mathbf{C}$  with  $|t|$  sufficiently small, defined by

$$\mathbf{Z}(G, t) = \mathbf{Z}_G(t) = \prod_{[C]} (1 - t^{|C|})^{-1},$$

where  $[C]$  runs over all equivalence classes of prime, reduced cycles of  $G$ .

Let  $G$  be a connected graph with  $n$  vertices and  $m$  unoriented edges. Two  $2m \times 2m$  matrices  $\mathbf{B} = (\mathbf{B}_{ef})_{e, f \in D(G)}$  and  $\mathbf{J}_0 = (\mathbf{J}_{ef})_{e, f \in D(G)}$  are defined as follows:

$$\mathbf{B}_{ef} = \begin{cases} 1 & \text{if } t(e) = o(f), \\ 0 & \text{otherwise,} \end{cases} \quad \mathbf{J}_{ef} = \begin{cases} 1 & \text{if } f = e^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$

Then the matrix  $\mathbf{B} - \mathbf{J}_0$  is called the *edge matrix* of  $G$ .

**THEOREM 3.1** (HASHIMOTO [12]; BASS [4]) *Let  $G$  be a connected graph. Then the reciprocal of the Ihara zeta function of  $G$  is given by*

$$\mathbf{Z}(G, t)^{-1} = \det(\mathbf{I}_{2m} - t(\mathbf{B} - \mathbf{J}_0)) = (1 - t^2)^{r-1} \det(\mathbf{I}_n - t\mathbf{A} + t^2(\mathbf{D} - \mathbf{I}_n)),$$

where  $r$  and  $\mathbf{A}$  are the Betti number and the adjacency matrix of  $G$ , respectively, and  $\mathbf{D} = (\mathbf{D}_{uv})_{u,v \in V(G)}$  is the diagonal matrix with  $\mathbf{D}_{uu} = \deg u$  for arbitrary  $u \in V(G)$ .

Consider an  $n \times n$  complex matrix  $\mathbf{W} = (\mathbf{W}_{uv})_{u,v \in V(G)}$  with  $(u, v)$ -entry equals 0 if  $(u, v) \notin D(G)$ . We call  $\mathbf{W}$  an *weighted matrix* of  $G$ . Furthermore, let  $w(u, v) = \mathbf{W}_{uv}$  for  $u, v \in V(G)$  and  $w(e) = w(u, v)$  if  $e = (u, v) \in D(G)$ .

For an weighted matrix  $\mathbf{W}$  of  $G$ , a  $2m \times 2m$  complex matrix  $\mathbf{B}_w = (\mathbf{B}_{ef}^{(w)})_{e,f \in D(G)}$  is defined as follows:

$$\mathbf{B}_{ef}^{(w)} = \begin{cases} w(f) & \text{if } t(e) = o(f), \\ 0 & \text{otherwise.} \end{cases}$$

Then the *second weighted zeta function* of  $G$  is defined by

$$\mathbf{Z}_1(G, w, t) = \det(\mathbf{I}_{2m} - t(\mathbf{B}_w - \mathbf{J}_0))^{-1}.$$

If  $w(e) = 1$  for any  $e \in D(G)$ , then the second weighted zeta function of  $G$  coincides with the Ihara zeta function of  $G$ .

**THEOREM 3.2** (SATO [25]) *Let  $G$  be a connected graph, and let  $\mathbf{W}$  be a weighted matrix of  $G$ . Then the reciprocal of the second weighted zeta function of  $G$  is given by*

$$\mathbf{Z}_1(G, w, t)^{-1} = (1 - t^2)^{m-n} \det(\mathbf{I}_n - t\mathbf{W} + t^2(\mathbf{D}_w - \mathbf{I}_n)),$$

where  $n = |V(G)|$ ,  $m = |E(G)|$  and  $\mathbf{D}_w = (\mathbf{D}_{uv}^{(w)})_{u,v \in V(G)}$  is the diagonal matrix with  $\mathbf{D}_{uu}^{(w)} = \sum_{e:o(e)=u} w(e)$  for arbitrary  $u \in V(G)$ .

**REMARK 3.3** *For later use, we mention that taking transpose, the following equation also holds:*

$$\det(\mathbf{I}_{2m} - t({}^T\mathbf{B}_w - \mathbf{J}_0)) = (1 - t^2)^{m-n} \det(\mathbf{I}_n - t^T\mathbf{W} + t^2(\mathbf{D}_w - \mathbf{I}_n)).$$

## 4 The transition matrix of a quaternionic quantum walk on a graph

A discrete-time quantum walk is a quantum process on a graph whose state vector is governed by a unitary matrix called the transition matrix. An important example of the quantum walk on a graph is the Grover walk, which was introduced in [11]. Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges. Set  $d_u = \deg u$  for  $u \in V(G)$ . The *transition matrix* of the Grover walk  $\mathbf{U}^{\text{Gro}} = (\mathbf{U}_{ef}^{\text{Gro}})_{e,f \in D(G)}$  of  $G$  is defined by

$$\mathbf{U}_{ef}^{\text{Gro}} = \begin{cases} 2/d_{t(f)} (= 2/d_{o(e)}) & \text{if } t(f) = o(e) \text{ and } f \neq e^{-1}, \\ 2/d_{t(f)} - 1 & \text{if } f = e^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$



$\mathbf{U}$  is called the *Grover matrix*. We denote by  $\text{Spec}(\mathbf{A})$  the multiset of spectra of a complex square matrix  $\mathbf{A}$  counted with multiplicity. Let  $\mathbf{T} = (\mathbf{T}_{uv})_{u,v \in V(G)}$  be the  $n \times n$  matrix defined as follows:

$$\mathbf{T}_{uv} = \begin{cases} 1/d_u & \text{if } (u, v) \in D(G), \\ 0 & \text{otherwise.} \end{cases}$$

In [8], Emms et al. determined the spectra of  $\mathbf{U}^{\text{Gro}}$  by use of those of  $\mathbf{T}$ .

**THEOREM 4.1** (EMMS, HANCOCK, SEVERINI AND WILSON [8]) *Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges. The transition matrix  $\mathbf{U}^{\text{Gro}}$  has  $2n$  eigenvalues of the form:*

$$\lambda = \lambda_{\mathbf{T}} \pm i\sqrt{1 - \lambda_{\mathbf{T}}^2},$$

where  $\lambda_{\mathbf{T}}$  is an eigenvalue of the matrix  $\mathbf{T}$ . The remaining  $2(m - n)$  eigenvalues of  $\mathbf{U}^{\text{Gro}}$  are  $\pm 1$  with equal multiplicities.

Now, we extend the Grover walk to the case of quaternions. A discrete-time quaternionic quantum walk is a quantum process on a graph whose state vector, whose entries are quaternions, is governed by a quaternionic unitary matrix called the quaternionic transition matrix. Let  $G$  be a finite connected graph with  $n$  vertices and  $m$  edges, The *quaternionic transition matrix*  $\mathbf{U} = (\mathbf{U}_{ef})_{e,f \in D(G)}$  of  $G$  is defined by

$$\mathbf{U}_{ef} = \begin{cases} q(e) & \text{if } t(f) = o(e) \text{ and } f \neq e^{-1}, \\ q(e) - 1 & \text{if } f = e^{-1}, \\ 0 & \text{otherwise,} \end{cases}$$

where  $q$  is a map from  $D(G)$  to  $\mathbb{H}$ . The unitary condition on  $\mathbf{U}$  is equivalent to the following equations:

**LEMMA 4.2** *Let  $q(e) = q_0(e) + q_1(e)i + q_2(e)j + q_3(e)k$ . Then  $\mathbf{U}$  is unitary if and only if the following five equations hold:*

$$q_0(e)^2 + q_1(e)^2 + q_2(e)^2 + q_3(e)^2 - \frac{2q_0(e)}{d_{o(e)}} = 0, \quad (4.3)$$

$$(q_0(e)q_0(f) + q_1(e)q_1(f) + q_2(e)q_2(f) + q_3(e)q_3(f))d_{o(e)} - (q_0(e) + q_0(f)) = 0, \quad (4.4)$$

$$(-q_0(e)q_1(f) + q_1(e)q_0(f) - q_2(e)q_3(f) + q_3(e)q_2(f))d_{o(e)} - (q_1(e) - q_1(f)) = 0, \quad (4.5)$$

$$(-q_0(e)q_2(f) + q_2(e)q_0(f) - q_3(e)q_1(f) + q_1(e)q_3(f))d_{o(e)} - (q_2(e) - q_2(f)) = 0, \quad (4.6)$$

$$(-q_0(e)q_3(f) + q_3(e)q_0(f) - q_1(e)q_2(f) + q_2(e)q_1(f))d_{o(e)} - (q_3(e) - q_3(f)) = 0, \quad (4.7)$$

where  $e$  and  $f$  are different arcs and have the same origin.

**Proof.** Observing that

$$(\mathbf{U}\mathbf{U}^*)_{ef} = \sum_{g \in D(G), t(g)=o(e)=o(f)} \mathbf{U}_{eg}(\mathbf{U}_{fg})^*,$$

one can readily check by direct calculations that  $(\mathbf{U}\mathbf{U}^*)_{ee} = 1$  is equivalent to (4.3), and  $(\mathbf{U}\mathbf{U}^*)_{ef} = 0$  to (4.4),(4.5),(4.6) and (4.7). If  $o(e) \neq o(f)$ , then clearly  $(\mathbf{U}\mathbf{U}^*)_{ef} = 0$  holds.  $\square$

From (4.3), it follows that  $q_0(e)$  must satisfy

$$0 \leq q_0(e) \leq \frac{2}{d_{o(e)}}.$$

**THEOREM 4.3** *If  $\mathbf{U}$  is unitary, then  $q(e) = q(f)$  holds for any two arcs  $e, f \in D(G)$  with  $o(e) = o(f)$ .*

**Proof.** We set  $q_0(e) = x/d_u$ ,  $q_0(f) = y/d_u$  ( $0 \leq x, y \leq 2$ ) and  $\mathbf{q}(e) = {}^T(q_1(e), q_2(e), q_3(e))$ , where  $u = o(e) = o(f)$ . Then (4.4) turns into:

$$\frac{(x-1)(y-1)-1}{d_u} + d_u \mathbf{q}(e) \cdot \mathbf{q}(f) = 0 \quad (4.8)$$

On the other hand, (4.5),(4.6),(4.7) can be expressed all at once as follows:

$$(y-1)\mathbf{q}(e) - (x-1)\mathbf{q}(f) - d_u \mathbf{q}(e) \times \mathbf{q}(f) = 0. \quad (4.9)$$

We show the assertion divided into three cases depending upon  $\mathbf{q}(e)$  and  $\mathbf{q}(f)$ .

If  $\mathbf{q}(e) = \mathbf{q}(f) = \mathbf{0}$ , then (4.9) trivially holds and (4.8) turns into  $(x-1)(y-1) - 1 = 0$ . In addition, by (4.3) we have

$$\frac{x(x-2)}{d_u} = 0$$

Thus,  $(x, y)$  must be  $(0, 0)$  or  $(2, 2)$ .

If only either one of  $\mathbf{q}(e)$  or  $\mathbf{q}(f)$  equals  $\mathbf{0}$ , say  $\mathbf{q}(f) = \mathbf{0}$ , then (4.9) becomes  $(y-1)\mathbf{q}(e) = \mathbf{0}$ . Hence  $y = 1$  and  $q(f) = 1/d_u$ . But from (4.8) it follows that  $-1/d_u = 0$ , thereby reaching a contradiction.

If both  $\mathbf{q}(e)$  and  $\mathbf{q}(f)$  do not equal  $\mathbf{0}$ , then by (4.9) it follows that:

$$(y-1)\mathbf{q}(e) - (x-1)\mathbf{q}(f) = \mathbf{0}, \quad (4.10)$$

$$\mathbf{q}(e) \times \mathbf{q}(f) = \mathbf{0}. \quad (4.11)$$

By (4.11),  $\mathbf{q}(e)$  and  $\mathbf{q}(f)$  are parallel, so we may write  $\mathbf{q}(f) = c\mathbf{q}(e)$  for some nonzero real number  $c$ . Then (4.10) implies  $y-1 = c(x-1)$ , and (4.8) yields:

$$c\{(x-1)^2 + d_u^2 |\mathbf{q}(e)|^2\} = 1 \quad (4.12)$$

Finally, substituting  $x = q_0(e)d_u$  for (4.12) and using (4.3), we get  $c = 1$ .  $\square$

Conversely, if  $q(e) = q(f)$  for any two arcs  $e, f \in D(G)$  with  $o(e) = o(f)$ , then (4.4) turns into (4.3), and moreover, (4.5), (4.6), (4.7) hold trivially. Thus we obtain:

**COROLLARY 4.4**  *$\mathbf{U}$  is unitary if and only if*

$$q_0(e)^2 + q_1(e)^2 + q_2(e)^2 + q_3(e)^2 - \frac{2q_0(e)}{d_{o(e)}} = 0,$$

*and  $q(e) = q(f)$  for any two arcs  $e, f \in D(G)$  with  $o(e) = o(f)$ .*

Hereafter we impose the following condition on  $q(e)$ :

$$\sum_{e:o(e)=u} q(e) \text{ does not depend on } u. \quad (4.13)$$

Hence, we may write  $\alpha = \alpha_0 + \alpha_1 i + \alpha_2 j + \alpha_3 k = \sum_{e:o(e)=u} q(e)$  for arbitrary  $u \in V(G)$ . Theorem 4.3 and (4.13) imply  $q(e) = \alpha/d_{o(e)}$ . If the imaginary part of  $\alpha$  equals 0, then  $\mathbf{U}$  must equal either  $-\mathbf{J}_0$  or  $\mathbf{U}^{\text{Gro}}$ . Therefore,  $\mathbf{U}$  can be regarded as a quaternionic extension of the Grover walk.

## 5 Eigenvalues of the transition matrix of a quaternionic quantum walks on a graph

We shall calculate all the right eigenvalues of  $\mathbf{U}$ . By Theorem 2.8, it suffices to solve  $\det(\lambda \mathbf{I}_{4m} - \psi(\mathbf{U})) = 0$ . Consider the set of quaternionic conjugates

$$\mathbf{U}^{\mathbb{H}^*} = \{q^{-1} \mathbf{U} q \mid q \in \mathbb{H}^*\},$$

of  $\mathbf{U}$ . By (2.1) and the subsequent argument,  $\alpha^{\mathbb{H}^*} \cap \mathbb{C} = \alpha_{\pm}$  where  $\alpha_{\pm} = \alpha_0 \pm \sqrt{\alpha_1^2 + \alpha_2^2 + \alpha_3^2} i$ .

Let  $\mathbf{U}_{\pm} = ((\mathbf{U}_{\pm})_{ef})_{e,f \in D(G)}$  be two complex matrices defined as follows:

$$(\mathbf{U}_{\pm})_{ef} = \begin{cases} \frac{\alpha_{\pm}}{d_{o(e)}} & \text{if } t(f) = o(e) \text{ and } f \neq e^{-1}, \\ \frac{\alpha_{\pm}}{d_{o(e)}} - 1 & \text{if } f = e^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$

We notice that  $\mathbf{U}_- = \overline{\mathbf{U}_+}$ . Then,  $\mathbf{U}^{\mathbb{H}^*} \cap \text{Mat}(2m, \mathbb{C}) = \{\mathbf{U}_+, \mathbf{U}_-\}$ . Therefore, for the quaternion  $q$  such that  $q^{-1} \alpha q = \alpha_+$ , we have:

$$\begin{aligned} \det(\lambda \mathbf{I}_{4m} - \psi(\mathbf{U})) &= \det(\psi(q \mathbf{I}_{2m})^{-1} (\lambda \mathbf{I}_{4m} - \psi(\mathbf{U})) \psi(q \mathbf{I}_{2m})) \\ &= \det(\lambda \mathbf{I}_{4m} - \psi(q \mathbf{I}_{2m})^{-1} \psi(\mathbf{U}) \psi(q \mathbf{I}_{2m})) \\ &= \det(\lambda \mathbf{I}_{4m} - \psi(q^{-1} \mathbf{U} q)) \\ &= \det(\lambda \mathbf{I}_{4m} - \psi(\mathbf{U}_+)). \end{aligned}$$

Thus, under the condition (4.13), we can calculate all the right eigenvalues of  $\mathbf{U}$  by calculating eigenvalues of  $\psi(\mathbf{U}_+)$ . However,

$$\begin{aligned} \det(\lambda \mathbf{I}_{4m} - \psi(\mathbf{U}_+)) &= \det \left( \begin{bmatrix} \lambda \mathbf{I}_{2m} & \mathbf{O} \\ \mathbf{O} & \lambda \mathbf{I}_{2m} \end{bmatrix} - \begin{bmatrix} \mathbf{U}_+ & \mathbf{O} \\ \mathbf{O} & \mathbf{U}_- \end{bmatrix} \right) \\ &= \det(\lambda \mathbf{I}_{2m} - \mathbf{U}_+) \det(\lambda \mathbf{I}_{2m} - \mathbf{U}_-). \end{aligned} \quad (5.14)$$

Hence,  $\text{Spec}(\psi(\mathbf{U})) = \text{Spec}(\psi(\mathbf{U}_+)) = \text{Spec}(\mathbf{U}_+) \cup \text{Spec}(\mathbf{U}_-)$ . Since  $\text{Spec}(\mathbf{U}_-)$  is obtained from  $\text{Spec}(\mathbf{U}_+)$  by conjugating their elements,  $\text{Spec}(\psi(\mathbf{U}))$  consists of eigenvalues of  $\mathbf{U}_+$  and their complex conjugates. Thus by Theorem 2.8, we obtain:

**THEOREM 5.1** *Let  $\mathbf{U}$  is a quaternionic transition matrix on a graph which satisfies (4.13). Then  $\sigma_r(\mathbf{U})$  is given by:*

$$\sigma_r(\mathbf{U}) = \lambda_1^{\mathbb{H}^*} \cup \dots \cup \lambda_n^{\mathbb{H}^*},$$

where  $\lambda_1, \dots, \lambda_n$  are eigenvalues of  $\mathbf{U}_+$ .

Now we apply Theorem 3.2 to the eigenvalue problem for  $\mathbf{U}$ . Let us define the  $n \times n$  complex matrix  $\mathbf{W}_\pm = ((\mathbf{W}_\pm)_{uv})_{u,v \in V(G)}$  as follows:

$$(\mathbf{W}_\pm)_{uv} = \begin{cases} \frac{\alpha_\pm}{d_u} & \text{if } (u, v) \in D(G), \\ 0 & \text{otherwise.} \end{cases}$$

**PROPOSITION 5.2** *Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges. Then, for the quaternionic transition matrix  $\mathbf{U}$  of  $G$ ,*

$$\det(\lambda \mathbf{I}_{2m} - \mathbf{U}_\pm) = (\lambda^2 - 1)^{m-n} \det((\lambda^2 + \alpha_\pm - 1)\mathbf{I}_n - \lambda^T \mathbf{W}_\pm).$$

**Proof.** Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges. Corresponding to  $\mathbf{W}_\pm$ , we define  $\mathbf{B}_{w\pm} = ((\mathbf{B}_\pm^{(w)})_{ef})_{e,f \in D(G)} \in \text{Mat}(2m, \mathbb{C})$  to be as follows:

$$(\mathbf{B}_\pm^{(w)})_{ef} = \begin{cases} \frac{\alpha_\pm}{d_{o(f)}} & \text{if } t(e) = o(f), \\ 0 & \text{otherwise.} \end{cases}$$

We readily see that  $\mathbf{U}_\pm = {}^T \mathbf{B}_{w\pm} - \mathbf{J}_0$ . Then, by Theorem 3.2 and the subsequent remark, it follows that:

$$\begin{aligned} \det(\mathbf{I}_{2m} - t\mathbf{U}_\pm) &= \det(\mathbf{I}_{2m} - t({}^T \mathbf{B}_{w\pm} - \mathbf{J}_0)) \\ &= (1 - t^2)^{m-n} \det(\mathbf{I}_n - t^T \mathbf{W}_\pm + t^2(\mathbf{D}_{w\pm} - \mathbf{I}_n)) \\ &= (1 - t^2)^{m-n} \det(\mathbf{I}_n - t^T \mathbf{W}_\pm + (\alpha_\pm - 1)t^2 \mathbf{I}_n) \\ &= (1 - t^2)^{m-n} \det((1 + (\alpha_\pm - 1)t^2)\mathbf{I}_n - t^T \mathbf{W}_\pm). \end{aligned}$$

Now, let  $t = 1/\lambda$ . Then we have

$$\det\left(\mathbf{I}_{2m} - \frac{1}{\lambda} \mathbf{U}_\pm\right) = \left(1 - \frac{1}{\lambda^2}\right)^{m-n} \det\left(\left(1 + (\alpha_\pm - 1)\frac{1}{\lambda^2}\right)\mathbf{I}_n - \frac{1}{\lambda} {}^T \mathbf{W}_\pm\right).$$

This implies the following conclusion:

$$\det(\lambda \mathbf{I}_{2m} - \mathbf{U}_\pm) = (\lambda^2 - 1)^{m-n} \det((\lambda^2 + \alpha_\pm - 1)\mathbf{I}_n - \lambda^T \mathbf{W}_\pm).$$

□

Using Proposition 5.2, we can calculate the spectra of  $\psi(\mathbf{U})$  as follows.

**THEOREM 5.3** *Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges.  $\text{Spec}(\psi(\mathbf{U}))$  has  $4m$  elements. If  $G$  is not a tree, then  $4n$  eigenvalues of them are of the form:*

$$\lambda = \frac{\mu \pm \sqrt{\mu^2 - 4(\alpha_+ - 1)}}{2}, \frac{\nu \pm \sqrt{\nu^2 - 4(\alpha_- - 1)}}{2},$$

where  $\mu \in \text{Spec}({}^T\mathbf{W}_+)$ ,  $\nu \in \text{Spec}({}^T\mathbf{W}_-)$ . The remaining  $4(m-n)$  eigenvalues of  $\psi(\mathbf{U})$  are  $\pm 1$  with equal multiplicities. If  $G$  is a tree, then  $\text{Spec}(\psi(\mathbf{U}))$  is as follows:

$$\begin{aligned} & \text{Spec}(\psi(\mathbf{U})) \\ &= \left\{ \frac{\mu \pm \sqrt{\mu^2 - 4(\alpha_+ - 1)}}{2}, \frac{\nu \pm \sqrt{\nu^2 - 4(\alpha_- - 1)}}{2} \mid \mu \in \text{Spec}({}^T\mathbf{W}_+), \nu \in \text{Spec}({}^T\mathbf{W}_-) \right\} \\ & \quad - \{1, 1, -1, -1\}. \end{aligned}$$

**Proof.** By (5.14) and Proposition 5.2, we have

$$\det(\lambda \mathbf{I}_{4m} - \psi(\mathbf{U})) = (\lambda^2 - 1)^{2m-2n} \prod_{\mu \in \text{Spec}({}^T\mathbf{W}_+)} (\lambda^2 + \alpha_+ - 1 - \mu\lambda) \prod_{\nu \in \text{Spec}({}^T\mathbf{W}_-)} (\lambda^2 + \alpha_- - 1 - \nu\lambda).$$

Solving  $\lambda^2 + \alpha_+ - 1 - \mu\lambda = 0$  and  $\lambda^2 + \alpha_- - 1 - \nu\lambda = 0$ , we obtain

$$\lambda = \frac{\mu \pm \sqrt{\mu^2 - 4(\alpha_+ - 1)}}{2}, \frac{\nu \pm \sqrt{\nu^2 - 4(\alpha_- - 1)}}{2}.$$

If  $G$  is not a tree, then  $m \geq n$  and hence the statement clearly holds. If  $G$  is a tree, then  $m = n - 1$ . Since  $\det(\lambda \mathbf{I}_{4m} - \psi(\mathbf{U}))$  is a polynomial of  $\lambda$ ,  $\prod_{\mu \in \text{Spec}({}^T\mathbf{W}_+)} (\lambda^2 + \alpha_+ - 1 - \mu\lambda) \prod_{\nu \in \text{Spec}({}^T\mathbf{W}_-)} (\lambda^2 + \alpha_- - 1 - \nu\lambda)$  must have the factor  $(\lambda^2 - 1)^2$ . Thus the statement holds for trees.  $\square$

**REMARK 5.4** Since  $\text{Spec}({}^T\mathbf{W}_-)$  is obtained from  $\text{Spec}({}^T\mathbf{W}_+)$  by conjugating their elements, it suffices to calculate only the eigenvalues of  $\text{Spec}({}^T\mathbf{W}_+)$ .

**EXAMPLE 5.5** Let  $G = K_3$ , the complete graph with 3 vertices, and  $\alpha = 1 + 1/2i + \sqrt{2}/2j - 1/2k$ . We give a weighted matrix as follows:

$$\mathbf{W} = \begin{bmatrix} 0 & \frac{\alpha}{2} & \frac{\alpha}{2} \\ \frac{\alpha}{2} & 0 & \frac{\alpha}{2} \\ \frac{\alpha}{2} & \frac{\alpha}{2} & 0 \end{bmatrix}.$$

Then the quaternionic transition matrix  $\mathbf{U}$  is as follows:

$$\mathbf{U} = {}^T\mathbf{B}_w - \mathbf{J}_0 = \begin{bmatrix} 0 & \frac{\alpha}{2} - 1 & 0 & 0 & \frac{\alpha}{2} & 0 \\ \frac{\alpha}{2} - 1 & 0 & 0 & \frac{\alpha}{2} & 0 & 0 \\ \frac{\alpha}{2} & 0 & 0 & \frac{\alpha}{2} - 1 & 0 & 0 \\ 0 & 0 & \frac{\alpha}{2} - 1 & 0 & 0 & \frac{\alpha}{2} \\ 0 & 0 & \frac{\alpha}{2} & 0 & 0 & \frac{\alpha}{2} - 1 \\ 0 & \frac{\alpha}{2} & 0 & 0 & \frac{\alpha}{2} - 1 & 0 \end{bmatrix}.$$

Then  $\alpha_{\pm} = 1 \pm i$  and

$$\text{Spec}({}^T\mathbf{W}_+) = \left\{ 1 + i, -\frac{1}{2} - \frac{1}{2}i, -\frac{1}{2} - \frac{1}{2}i \right\}, \text{Spec}({}^T\mathbf{W}_-) = \left\{ 1 - i, -\frac{1}{2} + \frac{1}{2}i, -\frac{1}{2} + \frac{1}{2}i \right\}.$$

and

$$\begin{aligned} \text{Spec}(\mathbf{U}_+) &= \left\{ 1, i, \frac{-1 \pm \sqrt{7}}{4} + \frac{-1 \mp \sqrt{7}}{4}i, \frac{-1 \pm \sqrt{7}}{4} + \frac{-1 \mp \sqrt{7}}{4}i \right\}, \\ \text{Spec}(\mathbf{U}_-) &= \left\{ 1, -i, \frac{-1 \pm \sqrt{7}}{4} + \frac{1 \pm \sqrt{7}}{4}i, \frac{-1 \pm \sqrt{7}}{4} + \frac{1 \pm \sqrt{7}}{4}i \right\}. \end{aligned}$$

Thus we obtain:

$$\sigma_r(\mathbf{U}) = \{1\} \cup i^{\mathbb{H}^*} \cup \left( \frac{-1 + \sqrt{7}}{4} + \frac{-1 - \sqrt{7}}{4}i \right)^{\mathbb{H}^*} \cup \left( \frac{-1 - \sqrt{7}}{4} + \frac{-1 + \sqrt{7}}{4}i \right)^{\mathbb{H}^*}.$$

Next, we state spectra of the Grover matrix. Let  $\alpha = 2$ . Then  $\mathbf{U}$  is the Grover matrix  $\mathbf{U}^{\text{Gro}}$ :

$$\mathbf{U}^{\text{Gro}} = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Furthermore, the transition matrix  $\mathbf{T}$  of the simple random walk of  $K_3$  is given as follows:

$$\mathbf{T} = \begin{bmatrix} 0 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 0 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & 0 \end{bmatrix}.$$

Then

$$\text{Spec}(\mathbf{T}) = \left\{ 1, -\frac{1}{2}, -\frac{1}{2} \right\} \quad \text{and} \quad \text{Spec}(\mathbf{U}^{\text{Gro}}) = \left\{ 1, 1, \frac{-1 \pm \sqrt{3}i}{4}, \frac{-1 \pm \sqrt{3}i}{4} \right\}.$$

Thus we obtain:

$$\sigma_r(\mathbf{U}^{\text{Gro}}) = \{1\} \cup \left( \frac{-1 + \sqrt{3}i}{2} \right)^{\mathbb{H}^*}.$$

**EXAMPLE 5.6** Let  $G = S_4$ , the star graph with 4 vertices, and  $\alpha = 4/3 + 1/3i + 2/3j + \sqrt{3}/3k$ . We give a weighted matrix as follows:

$$\mathbf{W} = \begin{bmatrix} 0 & 0 & 0 & \alpha \\ 0 & 0 & 0 & \alpha \\ 0 & 0 & 0 & \alpha \\ \frac{\alpha}{3} & \frac{\alpha}{3} & \frac{\alpha}{3} & 0 \end{bmatrix}.$$

Then the quaternionic transition matrix  $\mathbf{U}$  is as follows:

$$\mathbf{U} = {}^T\mathbf{B}_w - \mathbf{J}_0 = \begin{bmatrix} 0 & \alpha - 1 & 0 & 0 & 0 & 0 \\ \frac{\alpha}{3} - 1 & 0 & \frac{\alpha}{3} & 0 & \frac{\alpha}{3} & 0 \\ 0 & 0 & 0 & \alpha - 1 & 0 & 0 \\ \frac{\alpha}{3} & 0 & \frac{\alpha}{3} - 1 & 0 & \frac{\alpha}{3} & 0 \\ 0 & 0 & 0 & 0 & 0 & \alpha - 1 \\ \frac{\alpha}{3} & 0 & \frac{\alpha}{3} & 0 & \frac{\alpha}{3} - 1 & 0 \end{bmatrix}.$$

Then  $\alpha_{\pm} = 4/3 \pm 2\sqrt{2}/3i$  and

$$\text{Spec}({}^T\mathbf{W}_+) = \left\{ 0, 0, \frac{4}{3} + \frac{2\sqrt{2}}{3}i, -\frac{4}{3} - \frac{2\sqrt{2}}{3}i \right\}, \quad \text{Spec}({}^T\mathbf{W}_-) = \left\{ 0, 0, \frac{4}{3} - \frac{2\sqrt{2}}{3}i, -\frac{4}{3} + \frac{2\sqrt{2}}{3}i \right\}.$$

and

$$\begin{aligned} \text{Spec}(\mathbf{U}_+) &= \left\{ \pm \left( \frac{1}{\sqrt{3}} - \frac{\sqrt{2}}{\sqrt{3}}i \right), \pm \left( \frac{1}{\sqrt{3}} - \frac{\sqrt{2}}{\sqrt{3}}i \right), 1, \frac{1}{3} + \frac{2\sqrt{2}}{3}i, -\frac{1}{3} - \frac{2\sqrt{2}}{3}i, -1 \right\} - \{1, -1\}, \\ \text{Spec}(\mathbf{U}_-) &= \left\{ \pm \left( \frac{1}{\sqrt{3}} + \frac{\sqrt{2}}{\sqrt{3}}i \right), \pm \left( \frac{1}{\sqrt{3}} + \frac{\sqrt{2}}{\sqrt{3}}i \right), 1, \frac{1}{3} - \frac{2\sqrt{2}}{3}i, -\frac{1}{3} + \frac{2\sqrt{2}}{3}i, -1 \right\} - \{1, -1\}. \end{aligned}$$

Thus we obtain:

$$\sigma_r(\mathbf{U}) = \left( \frac{1}{\sqrt{3}} - \frac{\sqrt{2}}{\sqrt{3}}i \right)^{\mathbb{H}^*} \cup \left( -\frac{1}{\sqrt{3}} + \frac{\sqrt{2}}{\sqrt{3}}i \right)^{\mathbb{H}^*} \cup \left( \frac{1}{3} + \frac{2\sqrt{2}}{3}i \right)^{\mathbb{H}^*} \cup \left( -\frac{1}{3} - \frac{2\sqrt{2}}{3}i \right)^{\mathbb{H}^*}.$$

Next, we state spectra of the Grover matrix. Let  $\alpha = 2$  and  $\mathbf{U} = \mathbf{U}^{\text{Gro}}$ . Then  $\mathbf{U}^{\text{Gro}}$  is given as follows:

$$\mathbf{U}^{\text{Gro}} = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ \frac{-1}{3} & 0 & \frac{2}{3} & 0 & \frac{2}{3} & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ \frac{2}{3} & 0 & \frac{-1}{3} & 0 & \frac{2}{3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ \frac{2}{3} & 0 & \frac{2}{3} & 0 & \frac{-1}{3} & 0 \end{bmatrix}.$$

Furthermore, the transition matrix  $\mathbf{T}$  of the simple random walk of  $S_4$  is

$$\mathbf{T} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{3} & \frac{1}{3} & 0 \end{bmatrix}.$$

Then

$$\text{Spec}(\mathbf{T}) = \left\{ 0, 0, -1, 1 \right\} \quad \text{and} \quad \text{Spec}(\mathbf{U}^{\text{Gro}}) = \left\{ \pm i, \pm i, 1, -1 \right\}.$$

Thus we obtain:

$$\sigma_r(\mathbf{U}^{\text{Gro}}) = \{1\} \cup \{-1\} \cup i^{\mathbb{H}^*}.$$

**Acknowledgment.** The first author was partially supported by the Grant-in-Aid for Scientific Research (C) of Japan Society for the Promotion of Science (Grant No.21540116). The third author is partially supported by the Grant-in-Aid for Scientific Research (C) of Japan Society for the Promotion of Science (Grant No.19540154). We are grateful to K. Tamano, S. Matsutani and Y. Ide for some valuable comments on this work.

## References

- [1] Altmann, S.L.: Rotations, Quaternions, and Double Groups. Dover Publications (2005)
- [2] Ambainis, A., Bach, E., Nayak, A., Vishwanath, A., Watrous, J.: One-dimensional quantum walks. In: Proceedings of the 33rd Annual ACM Symposium on Theory of Computing, pp.37–49 (2001)
- [3] Aslaksen, H.: Quaternionic Determinants. *Math. intelligencer* **18**, 57–65 (1996)
- [4] Bass, H.: The Ihara-Selberg zeta function of a tree lattice. *Internat. J. Math.* **3**, 717–797 (1992)
- [5] Brenner, J.L.: Matrices of quaternions. *Pac. J. Math.* **1**, 329–335 (1951)
- [6] Cantero, M.J., Grunbaum, F.A., Moral, L., Velazquez, L.: The CGMV method for quantum walks. *Quantum Inf. Process.* **11**, 1149–1192 (2012)
- [7] Conway, J.H., Smith, D.: On Quaternions and Octonions. A K Peters, Ltd. (2003)
- [8] Emms, D., Hancock, E.R., Severini, S., Wilson, R.C.: A matrix representation of graphs and its spectrum as a graph invariant. *Electr. J. Combin.* **13**, R34 (2006)
- [9] Foata, D., Zeilberger, D.: A combinatorial proof of Bass’s evaluations of the Ihara-Selberg zeta function for graphs. *Trans. Amer. Math. Soc.* **351**, 2257–2274 (1999)
- [10] Godsil, C., Guo, K.: Quantum walks on regular graphs and eigenvalues. *Electr. J. Combin.* **18**, P165 (2011)
- [11] Grover, Lov K.: A fast quantum mechanical algorithm for database search. Proceedings of the 28th Annual ACM Symposium on the Theory of Computing (STOC), May 1996, pp.212–219 (1996)
- [12] Hashimoto, K.: Zeta functions of finite graphs and representations of  $p$ -adic groups. in "Adv. Stud. Pure Math". Vol.15, pp.211–280, Academic Press, New York (1989)
- [13] Huang, L., So, W.: On left eigenvalues of a quaternionic matrix. *Linear Algebra Appl.* **323**, 105–116 (2001)
- [14] Ihara, Y.: On discrete subgroups of the two by two projective linear group over  $p$ -adic fields. *J. Math. Soc. Japan* **18**, 219–235 (1966)
- [15] Kempe, J.: Quantum random walks - an introductory overview. *Contemporary Physics* **44**, 307–327 (2003)
- [16] Kendon, V.: Decoherence in quantum walks - a review. *Math. Struct. in Comp. Sci.* **17**, 1169–1220 (2007)
- [17] Konno, N.: Quantum Walks. In: Quantum Potential Theory, Franz, U., and Schürmann, M., Eds., Lecture Notes in Mathematics: Vol.1954, pp.309–452, Springer-Verlag, Heidelberg (2008)
- [18] Konno, N.: Quaternionic quantum walks. *Quantum Stud.: Math. Found.* **2**, 63–76 (2015)



- [19] Konno, N., Sato, I.: On the relation between quantum walks and zeta functions. *Quantum Inf. Process.* **11**, 341–349 (2012)
- [20] Kotani, M., Sunada, T.: Zeta functions of finite graphs. *J. Math. Sci. U. Tokyo* **7**, 7–25 (2000)
- [21] Lee, H. C.: Eigenvalues and canonical forms of matrices with quaternion coefficients. *Proc. Roy. Irish. Acad. Sect. A*, **52**, 253–260 (1949)
- [22] Manouchehri, K., Wang, J.: *Physical Implementation of Quantum Walks*. Springer, Berlin (2013)
- [23] Portugal, R.: *Quantum Walks and Search Algorithms*. Springer, Berlin (2013)
- [24] Ren, P., Aleksic, T., Emms, D., Wilson, R.C., Hancock, E.R.: Quantum walks, Ihara zeta functions and cospectrality in regular graphs. *Quantum Inf. Process.* **10**, 405–417 (2011)
- [25] Sato, I.: A new Bartholdi zeta function of a graph. *Int. J. Algebra* **1**, 269–281 (2007)
- [26] Serre, J. -P.: *Trees*, Springer-Verlag, New York (1980)
- [27] Stark, H.M., Terras, A.A.: Zeta functions of finite graphs and coverings. *Adv. Math.* **121**, 124–165 (1996)
- [28] Sunada, T.: *L-Functions in Geometry and Some Applications*. in "Lecture Notes in Math"., Vol.1201, pp.266–284, Springer-Verlag, New York (1986)
- [29] Sunada, T.: *Fundamental Groups and Laplacians (in Japanese)*. Kinokuniya, Tokyo (1988)
- [30] Venegas-Andraca, S.E.: Quantum walks: a comprehensive review. *Quantum Inf. Process.* **11**, 1015–1106 (2012)
- [31] Zhang, Z.: Quaternions and matrices of quaternions. *Linear Algebra Appl.* **251**, 21–57 (1997)