# Fibered Multiderivators and (co)homological descent

Fritz Hörmann

Mathematisches Institut, Albert-Ludwigs-Universität Freiburg

October 20, 2015

2010 Mathematics Subject Classification: 55U35, 14F05, 18D10, 18D30, 18E30, 18G99

Keywords: Derivators, fibered derivators, multiderivators, fibered multicategories, Grothendieck's six-functor-formalism, cohomological descent, homological descent, fundamental localizers, well-generated triangulated categories, equivariant derived categories

## Abstract

The theory of derivators enhances and simplifies the theory of triangulated categories. In this article a notion of fibered (multi-)derivator is developed, which similarly enhances fibrations of (monoidal) triangulated categories. We present a theory of cohomological as well as homological descent in this language. The main motivation is a descent theory for Grothendieck's six operations.

# Introduction

#### Grothendieck's six functors and descent

Let  $S$  be a category, for instance a suitable category of schemes, topological spaces, analytic manifolds, etc. A Grothendieck six functor formalism on  $S$  consists of a collection of (derived) categories  $\mathcal{D}_S$ , one for each "base space" S in S with the following six types of operations:



The fiber  $\mathcal{D}_S$  is, in general, a *derived* category of "sheaves" over S, for example coherent sheaves, l-adic sheaves, abelian sheaves, D-modules, motives, etc. The functors on the left hand side are left adjoints of the functors on the right hand side. The functor  $f_!$  and its right adjoint  $f_!$  are called "push-forward with proper support", and "exceptional pull-back", respectively. The six functors come along with a bunch of compatibility isomorphisms between them (cf. [A.2.19\)](#page-83-0) and it is not easy to make their axioms really precise. In an appendix to this article, we explain that one quite simple precise definition is the following:

**Definition [A.2.16.](#page-82-0)** Let S be a category with fiber products. A (symmetric) Grothendieck sixfunctor-formalism on  $S$  is a bifibration of (symmetric) 2-multicategories with 1-categorical fibers

$$
p: \mathcal{D} \to \mathcal{S}^\text{cor}
$$

where  $\mathcal{S}^{\text{cor}}$  is the symmetric 2-multicategory of correspondences in  $\mathcal{S}$  (cf. Definition [A.2.15\)](#page-81-0).

From such a bifibration we obtain the operations  $f_*$ ,  $f^*$  (resp.  $f^!$ ,  $f_!$ ) as pull-back and push-forward along the correspondences



respectively. We get  $\mathcal{E} \otimes \mathcal{F}$  for objects  $\mathcal{E}, \mathcal{F}$  above X as the target of a Cartesian 2-ary multimorphism from the pair  $\mathcal{E}, \mathcal{F}$  over the correspondence



Given such a six-functor-formalism and a simplicial resolution  $\pi : U_{\bullet} \to S$  of a space  $S \in \mathcal{S}$  (for example arising from a Čech cover w.r.t. a suitable Grothendieck topology)

$$
\cdots \underbrace{\Longrightarrow} U_2 \underbrace{\Longrightarrow} U_1 \underbrace{\Longrightarrow} U_0,
$$

and given an object  $\mathcal E$  in  $\mathcal D_S$ , one can construct complexes in the category  $\mathcal D_S$ :

$$
\cdots \longrightarrow \pi_{2,1}\pi_2^!\mathcal{E} \longrightarrow \pi_{1,1}\pi_1^!\mathcal{E} \longrightarrow \pi_{0,1}\pi_0^!\mathcal{E} \longrightarrow 0
$$
  

$$
\cdots \longleftarrow \pi_{2,*}\pi_2^*\mathcal{E} \longleftarrow \pi_{1,*}\pi_1^*\mathcal{E} \longleftarrow \pi_{0,*}\pi_0^*\mathcal{E} \longleftarrow 0
$$

The first question of homological (resp. cohomological) descent is whether the hyper(co)homology of these complexes recovers the homology (resp. cohomology) of  $\mathcal{E}$ . Without a suitable enhancement of the situation, this question, however, does not make sense because a double complex, once considered as a complex in the derived category, looses the information of the homology of its total complex. There are several remedies for this problem. Classically, if at least the  $\pi_i^*$  are derived functors and  $\mathcal E$  is acyclic w.r.t. them, one can derive the whole construction to get a coherent double complex. This does not work, however, for the functors  $f_1$ ,  $f'$  which are often only constructed on the derived category. One possibility is to consider enhancements of the triangulated categories in question as DG-categories or  $\infty$ -categories. In this article, we have worked out a different approach based on Grothendieck's idea of derivators which is, perhaps, conceptually even simpler. It is sufficiently powerful to glue the six functors and define them for morphisms between stacks, or even higher stacks. Like the ∞-categorical approach, it is also very general, not being restricted to the stable case.

## Fibered multiderivators

The notion of triangulated category developed by Grothendieck and Verdier in the 1960's, as successful as it has been, is not sufficient for many purposes, for both practical reasons (certain natural constructions cannot be performed) as well as for theoretical reasons (the axioms are rather involved and lack conceptual clarity). Grothendieck much later [\[11\]](#page-86-0) and Heller independently, with the notion of derivator, proposed a marvelously simple remedy to both deficiencies. The basic observation is that all problems mentioned above are based on the following fact: Consider a category C and a class of morphisms  $W$  (quasi-isomorphisms, weak equivalences, etc.) which one would like to become isomorphisms. Then homotopy limits and colimits w.r.t.  $(C, W)$  cannot be reconstructed once passed to the homotopy category  $\mathcal{C}[\mathcal{W}^{-1}]$  (for example a derived category, or the homotopy category of a model category). Examples of homotopy (co)limits are the cone (required to exist in a triangulated category in a brute-force way, but not functorially!) and the total complex of a complex of complexes (totally lost in the derived category). Furthermore, very basic and intuitive properties of homotopy limits and colimits, and more general Kan extensions, not only determine the additional structure (triangles, shift functors) on a triangulated category but also imply all of its rather involved axioms. This idea has been successfully worked out by Cisinski, Groth, Grothendieck, Heller, Maltsiniotis, and others. We refer to the introductory article [\[8\]](#page-86-1) for an overview.

The purpose of this article is to propose a notion of fibered (multi-)derivator which enhances the notion of a fibration of (monoidal) triangulated categories in the same way as the notion of usual derivator enhances the notion of triangulated category. We emphasize that this new context is very well suited to reformulate (and reprove the theorems of) the classical theory of cohomological descent and to establish a completely dual theory of homological descent which should be satisfied by the  $f_!, f^!$ -functors.

#### (Co)homological descent with fibered derivators

Pursuing the idea of derivators, there is a neat conceptual solution to the problem of (co)homological descent: Analogously to a derivator which associates a (derived) category with each diagram shape I, we should consider a (derived) category  $\mathbb{D}(I, F)$  for each diagram of correspondences  $F : I \to \mathcal{S}^{\text{cor}}$ . Then, given a simplicial resolution  $\pi: U_{\bullet} \to S$  as before, considered as a morphism  $p: (\Delta^{op}, U_{\bullet}) \to$  $(., S)$  of diagrams in  $\mathcal{S}^{cor}$ , resp.  $i : (\Delta, (U_{\bullet})^{\text{op}}) \to (., S)$  in a dual diagram category (cf. [1.6.2\)](#page-25-0), the question becomes:

Q1: Does the corresponding pull-back  $i^*$  have a right adjoint  $i_*$ , respectively does  $p^*$  have a left adjoint  $p_{\parallel}$  (a straightforward generalization of the question of existence of homotopy (co)limits in usual derivators!) and is the corresponding unit id  $\rightarrow i_*i^*$  (resp. counit  $p_!p^* \rightarrow id$ ) and isomorphism?

Instead of, however, taking an association  $(I, F) \rightarrow \mathbb{D}(I, F)$  as the fundamental datum, we propose to take a morphism of pre-derivators  $p : \mathbb{D} \to \mathbb{S}$  (or even pre-multiderivators) as the fundamental datum, the  $\mathbb{D}(I, F)$  being reconstructed as its fibers  $\mathbb{D}(I)F$ , if S is the pre-derivator associated with a category. This allows more general situations, where S is not associated with an ordinary category. In many situations, in particular for a six-functor-formalism, it will be necessary to consider S which are pre-2-multiderivators instead, a notion which will be introduced and investigated in a forthcoming article [\[14](#page-86-2)]. There we will define (and give examples of) a derivator version of a (symmetric) Grothendieck six-functor formalism, that is, a (symmetric) fibered multiderivator

$$
p:\mathbb{D}\to \mathbb{S}^{\mathrm{cor}},
$$

where  $\mathbb{S}^{\text{cor}}$  is the symmetric pre-2-multiderivator of correspondences in  $\mathcal{S}$ .

Q2: More generally, we may consider Cartesian (resp. coCartesian) objects in the fiber over a diagram  $(\Delta^{\rm op}, U_{\bullet})$  (resp.  $(\Delta, (U_{\bullet})^{\rm op})$ ), and ask whether these categories depend only on  $U_{\bullet}$ up to taking (finite) hypercovers w.r.t. a fixed Grothendieck topology on  $S$ .

This will allow to define the six operations, for example, if the simplicial objects  $U_{\bullet}$  are presentations of stacks. The categories of coCartesian objects are a generalization of the equivariant derived categories of Bernstein and Lunts (cf. [2.4.3\)](#page-41-0).

## **Overview**

In section [1](#page-5-0) we give the general definition of a left (resp. right) fibered multiderivator  $p : \mathbb{D} \to \mathbb{S}$ . The axioms are basically a straight-forward generalization of those of a left, resp. right derivator. To give a priori some conceptual evidence that these axioms are indeed reasonable, we prove that the notion of fibered multiderivator is transitive [\(1.4\)](#page-21-0), and that it gives rise to a pseudo-functor from 'diagrams in S' to categories, for which a neat base-change formula holds [\(1.6\)](#page-25-1).

In section [2,](#page-29-0) a theory of (co)homological descent for fibered derivators is developed (the monoidal, i.e. multi-, aspect does not play any role here). We propose a definition of localizer (resp. of system of relative localizers) in the category of diagrams in S which is a generalization of Grothendieck's notion of fundamental localizer in categories. The latter gives a nice combinatorial description of weak equivalences of categories in terms of the condition of Quillen's theorem A. In our more general setting the notion of fundamental localizer depends on the choice of a Grothendieck (pre-)topology on S. In section [2.3](#page-34-0) we show purely abstractly that a finite hypercover, considered as a morphism of simplicial diagrams, lies in any localizer or system of relative localizers. Thus this more general notion of localizer has a similar relation to weak equivalences of simplicial pre-sheaves, although we will not yet give any precise statement in this direction.

In sections [2.4](#page-41-1) and [2.5](#page-42-0) these notions of localizers are tied to the theory of fibered derivators. We introduce two notions of (co)homological descent for a fibered derivator  $p : \mathbb{D} \to \mathbb{S}$ , namely weak and strong D-equivalences. The notion of weak D-equivalences (related to Q1 above) is a straightforward generalization of Cisinski's notion of D-equivalence for usual derivators. In our relative context, both notions of D-equivalence come in a cohomological as well as in a homological flavour (a phenomenon which does not occur for usual derivators).

Whenever the fibered derivator is (co)local w.r.t. to the Grothendieck pre-(co)topology — a rather weak and obviously necessary condition (see section  $1.5$ ) — then the Main Theorem [2.5.4](#page-43-0) (resp. [2.5.5\)](#page-43-1) of this article states that weak D-equivalences form a system of relative localizers under very general conditions (the easier case) and that strong D-equivalences (related to Q2 above) form an absolute localizer, at least in the case of fibered derivators with stable, compactly generated fibers. The proof uses results from the theory of triangulated categories due to Neeman and Krause (centering around Brown representability type theorems). The link to our theory of fibered (multi-)derivators is explained in section [3.](#page-55-0)

In section [4](#page-59-0) we introduce the notion of fibration of multi-model-categories. This is the most favorable standard context in which a fibered multi-derivator (whose base is associated with a usual multicategory) can be constructed. We will present more general methods of constructing fibered multiderivators in a forthcoming article, in particular those encoding a full six-functor-formalism.

# Notation

We denote by  $\mathcal{CAT}$  the 2-"category"<sup>[1](#page-4-0)</sup> of categories, by  $(\mathcal{S})\mathcal{MCAT}$  the 2-"category" of (symmetric) multicategories, and by Cat the 2-category of small categories. We consider a partially ordered set (poset) X as a small category by considering the relation  $x \leq y$  to be equivalent to the existence of a unique morphism  $x \to y$ . We denote the positive integers (resp. non-negative integers) by N (resp.  $\mathbb{N}_0$ ). The ordered sets  $\{0, \ldots, n\} \subset \mathbb{N}_0$  considered as a small category are denoted by  $\Delta_n$ . We denote by  $\text{Mor}(\mathcal{D})$  (resp. Iso( $\mathcal{D}$ )) the class of morphisms (resp. isomorphisms) in a category  $\mathcal{D}$ . The final category (which consists of only one object and its identity) is denoted by  $\cdot$  or  $\Delta_0$ . The same notation is also used for the final multi-category, i.e. that with one object and precisely one  $n$ -ary morphism for any  $n$ . Our conventions about multicategories and fibered (multi-)categories are summarized in appendix [A.](#page-75-0)

# Contents



<span id="page-4-0"></span><sup>1</sup>where "category" has classes replaced by 2-classes (or, if the reader prefers, is constructed w.r.t. a larger universe).

## <span id="page-5-1"></span><span id="page-5-0"></span>1 Fibered derivators

## <span id="page-5-3"></span>1.1 Categories of diagrams

**Definition 1.1.1.** A diagram category is a full sub-2-category Dia ⊂ Cat, satisfying the following axioms:

- (Dia1) The empty category  $\varnothing$ , the final category  $\cdot$  (or  $\Delta_0$ ), and  $\Delta_1$  are objects of Dia.
- (Dia2) Dia is stable under taking finite coproducts and fibered products.
- (Dia3) For each functor  $\alpha : I \to J$  in Dia and object  $j \in J$  the slice categories  $I \times_{i,j} j$  and  $j \times_{i,j} I$  are in Dia.

A diagram category Dia is called self-dual, if it satisfies in addition:

(*Dia4*) If  $I \in$  Dia then  $I^{op} \in$  Dia.

A diagram category Dia is called infinite, if it satisfies in addition:

(Dia5) Dia is stable under taking arbitrary coproducts.

In the following we mean by a diagram a small category.

**Example 1.1.2.** We have the following diagram categories:

Cat the category of all diagrams. It is self-dual.

Inv the category of inverse diagrams  $C$ , i.e. small categories  $C$  such that there exists a functor  $C \to \mathbb{N}_0$  with the property that the preimage of an identity consists of identities<sup>[2](#page-5-2)</sup>. An example is the injective simplex category  $\Delta^{\circ}$ :

$$
\cdots \overleftarrow{\underleftarrow{\Longleftarrow}} \cdots \overleftarrow{\underleftarrow{\Longleftarrow}} \cdots \overleftarrow{\ominus}
$$

Dir the category of directed diagrams D, i.e. small categories such that  $D^{op}$  is inverse. An example is the opposite of the injective simplex category  $(\Delta^{\circ})^{\mathrm{op}}$ :

$$
\cdots \stackrel{\longrightarrow}{\Longrightarrow} \stackrel{\longrightarrow}{\Longrightarrow} \stackrel{\longrightarrow}{\Longrightarrow} \cdots
$$

- Catf, Dirf, and Invf are defined as before but consisting of finite diagrams. Those are self-dual and  $\text{Dirf} = \text{Invf}.$
- Catlf, Dirlf, and Invlf are defined as before but consisting of **locally finite diagrams**, *i.e.* those which have the property that a morphism  $\gamma$  factors as  $\gamma = \alpha \circ \beta$  only in a finite number of ways.
	- Pos, Posf, Dirpos, and Invpos: the categories of posets, finite posets, directed posets, and inverse posets.

<span id="page-5-2"></span><sup>&</sup>lt;sup>2</sup>In many sources  $\mathbb{N}_0$  is replaced by any ordinal.

<span id="page-6-0"></span>**Definition 1.2.1.** A pre-derivator of domain Dia is a contravariant (strict) 2-functor

$$
\mathbb{D}: \mathrm{Dia}^{1-\mathrm{op}} \to \mathcal{CAT}.
$$

A pre-multiderivator of domain Dia is a contravariant (strict) 2-functor

$$
\mathbb{D}: \mathrm{Dia}^{1-\mathrm{op}} \to \mathcal{MCAT}
$$

into the 2-"category" of multicategories. A morphism of pre-derivators is a natural transformation. For a morphism  $\alpha : I \to J$  in Dia the corresponding functor

$$
\mathbb{D}(\alpha): \mathbb{D}(J) \to \mathbb{D}(I)
$$

will be denoted by  $\alpha^*$ .

We call a pre-multiderivator symmetric (resp. braided), if its images are symmetric (resp. braided), and the morphisms  $\alpha^*$  are compatible with the actions of the symmetric (resp. braid) groups.

**1.2.2.** The pre-derivator associated with a category: Let S be a category. We associate with it the pre-derivator

$$
\mathbb{S}: I \mapsto \text{Hom}(I, \mathcal{S}).
$$

The pull-back  $\alpha^*$  is defined as composition with  $\alpha$ . A 2-morphism  $\kappa : \alpha \to \beta$  induces a natural 2-morphism  $\mathbb{S}(\kappa): \alpha^* \to \beta^*$ .

**1.2.3.** The pre-derivator associated with a simplicial class (in particular with an  $\infty$ -category): Let  $S$  be a simplicial class, i.e. a functor

$$
\mathcal{S}:\Delta\rightarrow\mathcal{CLASS}
$$

into the "category" of classes. We associate with it the pre-derivator

$$
\mathbb{S}: I \mapsto \text{Ho}(\text{Hom}(N(I), \mathcal{S})),
$$

where  $N(I)$  is the nerve of I and Ho is the left adjoint of N. In detail this means the objects of the category  $\mathcal{S}(I)$  are morphisms  $\alpha : N(I) \to \mathcal{S}$ , the class of morphisms in  $\mathcal{S}(I)$  is freely generated by morphisms  $\mu: N(I \times \Delta_1) \to S$  considered to be a morphism from its restriction to  $N(I \times \{0\})$  to its restriction to  $N(I \times \{1\})$  modulo the relations given by morphisms  $\nu : N(I \times \Delta_2) \to S$ , i.e. if  $\nu_1, \nu_2$ and  $\nu_3$  are the restrictions of  $\nu$  to the 3 faces of  $\Delta_2$  then we have  $\mu_3 = \mu_2 \circ \mu_1$ . The pull-back  $\alpha^*$  is defined as composition with the morphism  $N(\alpha): N(I) \to N(J)$ . A 2-morphism  $\kappa : \alpha \to \beta$  can be given as a functor  $I \times \Delta_1 \rightarrow J$  which yields (applying N and composing) a natural transformation which we call  $\mathbb{S}(\kappa)$ .

**1.2.4.** More generally, consider the full subcategory  $^M\Delta \subset \mathcal{MCAT}$  of all *finite connected* multicategories M that are freely generated by a finite set of multimorphisms  $f_1, \ldots, f_n$  such that each object of M occurs at most once as a source and at most once as the target of one of the  $f_i$ . Similarly consider the full subcategory  $T \subset \mathcal{SMCAT}$  which is obtained from  $^M\Delta$  adding images under the operations of the symmetric groups. This category is usually called the symmetric tree category. With a functor

$$
S: {}^M\Delta \to \mathcal{CLASS}
$$
 resp.  $S: T \to \mathcal{CLASS}$ 

we associate the pre-multiderivator (resp. symmetric pre-multiderivator):

$$
\mathbb{S}: I \mapsto \text{Ho}(\text{Hom}(N(I), \mathcal{S})),
$$

where  $N: \mathcal{MCAT} \to \mathcal{CLASS}^{M\Delta}$  (resp.  $N: \mathcal{SMCAT} \to \mathcal{CLASS}^T$  ) is the nerve, I is considered to be a multicategory without any n-ary morphisms for  $n \geq 2$ , and Ho is the left adjoint of N. Objects in  $\mathcal{SET}^T$  are called dendroidal sets in [\[18\]](#page-87-0).

## <span id="page-7-0"></span>1.3 Fibered (multi-)derivators

**1.3.1.** Let  $p : \mathbb{D} \to \mathbb{S}$  be a strict morphism of pre-derivators with domain Dia, and let  $\alpha : I \to J$  be a functor in Dia. Consider an object  $S \in \mathbb{S}(J)$ . The functor  $\alpha^*$  induces a morphism between fibers (denoted the same way)

$$
\alpha^* : \mathbb{D}(J)_S \to \mathbb{D}(I)_{\alpha^*S}.
$$

We are interested in the case that the latter has a left adjoint  $\alpha_!^S$ , resp. a right adjoint  $\alpha_*^S$ . These will be called relative left/right homotopy Kan extension functors with base  $S$ . For better readability we often omit the base from the notation. Though the base is not determined by the argument of  $\alpha_!,$  it will often be understood from the context, cf. also  $(1.3.28)$ .

1.3.2. We are interested in the case in which all morphisms

$$
p(I): \mathbb{D}(I) \to \mathbb{S}(I)
$$

are Grothendieck fibrations, resp. opfibrations [\(A.1\)](#page-75-1) or, more generally, (op)fibrations of multi-categories [\(A.2\)](#page-75-2). We always assume that the functors  $\alpha^* := \mathbb{D}(\alpha)$  map coCartesian morphisms to coCartesian morphisms but map Cartesian morphisms to Cartesian morphisms (for arity  $n \geq 2$ ) only if  $\alpha$  itself is a Grothendieck opfibration.

Then we will choose an associated pseudo-functor, i.e. for each  $f : S \to T$  in  $\mathbb{S}(I)$  a pair of adjoints functors

$$
f_{\bullet} : \mathbb{D}(I)_{S} \to \mathbb{D}(I)_{T},
$$

resp.

$$
f^{\bullet} : \mathbb{D}(I)_T \to \mathbb{D}(I)_S,
$$

characterized by functorial isomorphisms:

$$
\mathrm{Hom}_f\big(\mathcal{E},\mathcal{F}\big)\cong \mathrm{Hom}_{\mathrm{id}_T}\big(\mathcal{E},f^\bullet\mathcal{F}\big)\cong \mathrm{Hom}_{\mathrm{id}_S}\big(f_\bullet\mathcal{E},\mathcal{F}\big).
$$

More generally, in the multicategorical setting, if f is a multimorphism  $f \in Hom(S_1, \ldots, S_n; T)$  for some  $n \geq 1$ , we get an *adjunction* of *n* variables

$$
f_{\bullet}: \mathbb{D}(I)_{S_1} \times \cdots \times \mathbb{D}(I)_{S_n} \to \mathbb{D}(I)_T,
$$

and

$$
f^{i, \bullet}: \mathbb{D}(I)_{S_1}^{\text{op}} \times \frac{\widehat{i}}{\cdots} \times \mathbb{D}(I)_{S_n}^{\text{op}} \times \mathbb{D}(I)_T \to \mathbb{D}(I)_{S_i}.
$$

1.3.3. For a diagram of categories

$$
K \xrightarrow[\beta]{I} \sqrt{\alpha}
$$

the slice category  $K \times_{i} I$  is the category of tripels  $(k, i, \mu)$ , where  $k \in K$ ,  $i \in I$  and  $\mu : \alpha(i) \to \beta(k)$ . It sits in a corresponding 2-commutative square:

$$
K \times_{/J} I \xrightarrow{B} I
$$
  
A  

$$
A \downarrow \qquad \nearrow^{\mu} \qquad \downarrow \alpha
$$
  

$$
K \xrightarrow{\beta} J
$$

which is universal w.r.t. such squares. This construction is associative, but of course not commutative unless J is a groupoid. The projection  $K \times_{\text{IJ}} I \to K$  is a Grothendieck fibration and the projection  $K \times_{I} I \to I$  is a Grothendieck opfibration (see [A.1\)](#page-75-1). There is an adjunction

$$
I\times_{/{\cal J}}{\cal J}\xrightarrow{\hspace{0.5cm}I}{\hspace{0.5cm}I}.
$$

1.3.4. Consider an arbitrary 2-commutative square

<span id="page-8-0"></span>
$$
L \xrightarrow{B} I
$$
  
\n
$$
A \downarrow_{\alpha} \nearrow^{\mu} \downarrow_{\alpha} \qquad (1)
$$
  
\n
$$
K \xrightarrow{\beta} J
$$

and let  $S \in \mathbb{S}(J)$  be an object and  $\mathcal{E}$  a preimage in  $\mathbb{D}(J)$  w.r.t. p. The 2-morphism (natural transformation)  $\mu$  induces a functorial morphism

$$
\mathbb{S}(\mu): A^*\beta^*S \to B^*\alpha^*S
$$

and therefore a functorial morphism

$$
\mathbb{D}(\mu): A^*\beta^*\mathcal{E} \to B^*\alpha^*\mathcal{E}
$$

over  $\mathcal{S}(\mu)$ , or — if we are in the (op)fibered situation — equivalently

$$
A^*\beta^*\mathcal{E} \to (\mathbb{S}(\mu))^{\bullet} B^*\alpha^*\mathcal{E}
$$

respectively

$$
(\mathbb{S}(\mu))_{\bullet} A^* \beta^* \mathcal{E} \rightarrow B^* \alpha^* \mathcal{E}
$$

in the fiber above  $A^*\beta^*S$ , resp.  $B^*\alpha^*S$ ,

Let now  $\mathcal F$  be an object over  $\alpha^* S$ . If relative right homotopy Kan extensions exist, we may form the following composition which will be called the (right) base-change morphism:

<span id="page-8-1"></span>
$$
\beta^* \alpha_* \mathcal{F} \to A_* A^* \beta^* \alpha_* \mathcal{F} \to A_* (\mathbb{S}(\mu))^{\bullet} B^* \alpha^* \alpha_* \mathcal{F} \to A_* (\mathbb{S}(\mu))^{\bullet} B^* \mathcal{F}.
$$
 (2)

(We again omit the base S from the notation for better readability  $-$  it is always determined by the argument.)

Let now  $\mathcal F$  be an object over  $\beta^*S$ . If relative left homotopy Kan extensions exist, we may form the composition, the (left) base-change morphism:

<span id="page-8-2"></span>
$$
B_! (\mathbb{S}(\mu))_{\bullet} A^* \mathcal{F} \to B_! (\mathbb{S}(\mu))_{\bullet} A^* \beta^* \beta_! \mathcal{F} \to B_! B^* \alpha^* \beta_! \mathcal{F} \to \alpha^* \beta_! \mathcal{F}.
$$
 (3)

We will later say that the square  $(1)$  is **homotopy exact** if  $(2)$  is an isomorphism for all right fibered derivators (see Definition [1.3.6](#page-10-0) below) and [\(3\)](#page-8-2) is an isomorphism for all left fibered derivators. It is obvious a priori that for a left and right fibered derivator [\(2\)](#page-8-1) is an isomorphism if and only if [\(3\)](#page-8-2) is, one being the adjoint of the other (see [\[8,](#page-86-1) §1.2] for analogous reasoning in the case of usual derivators).

**Definition 1.[3](#page-9-0).5.** We consider the following axioms<sup>3</sup> on a pre-(multi-)derivator  $\mathbb{D}$ :

- (Der1) For I, J in Dia, the natural functor  $\mathbb{D}(I \coprod J) \to \mathbb{D}(I) \times \mathbb{D}(J)$  is an equivalence of (multi-)categories. Moreover  $\mathbb{D}(\varnothing)$  is not empty.
- (Der2) For I in Dia the 'underlying diagram' functor

$$
dia: \mathbb{D}(I) \to Hom(I, \mathbb{D}(\cdot))
$$

is conservative.

In addition, we consider the following axioms for a strict morphism of pre-(multi-)derivators

$$
p:\mathbb{D}\to\mathbb{S}:
$$

(FDer0 left) For each I in Dia the morphism p specializes to an opfibered (multi-)category and any functor  $\alpha : I \rightarrow J$  in Dia induces a diagram



of opfibered (multi-)categories, i.e. the top horizontal functor maps coCartesian arrows to coCartesian arrows.

(FDer3 left) For each functor  $\alpha : I \to J$  in Dia and  $S \in \mathbb{S}(J)$  the functor  $\alpha^*$  between fibers

$$
\mathbb{D}(J)_S \to \mathbb{D}(I)_{\alpha^*S}
$$

has a left-adjoint  $\alpha_!^S$ .

(FDer4 left) For each functor  $\alpha : I \rightarrow J$  in Dia, and for any object  $j \in J$ , and the 2-cell

$$
I \times_{/J} j \xrightarrow{\iota} I
$$
  
\n
$$
\alpha_j \downarrow \qquad \downarrow \alpha
$$
  
\n
$$
\{j\} \xrightarrow{\iota} J
$$

we get that the induced natural transformation of functors  $\alpha_{j}(\mathbb{S}(\mu))_{\bullet}i^* \to j^*\alpha_!$  is an isomor- $phism<sup>4</sup>$  $phism<sup>4</sup>$  $phism<sup>4</sup>$ .

(FDer5 left) (only needed for the multiderivator case). For any Grothendieck opfibration  $\alpha : I \to J$  in Dia, and for any morphism  $\xi \in \text{Hom}(S_1, \ldots, S_n; T)$  in  $\mathbb{S}(\cdot)$  for some  $n \geq 1$ , the natural transformations of functors

$$
\alpha_! (\alpha^* \xi)_\bullet (\alpha^* -, \cdots, \alpha^* -, \ldots, \alpha^* -, \cdots, \alpha^* - ) \cong \xi_\bullet (-, \cdots, -, \alpha_! - , -, \cdots, -)
$$

are isomorphisms.

<sup>&</sup>lt;sup>3</sup>The numbering is compatible with that of [\[8](#page-86-1)] in the case of non-fibered derivators.

<span id="page-9-1"></span><span id="page-9-0"></span><sup>&</sup>lt;sup>4</sup>This is meant to hold w.r.t. all bases  $S \in \mathcal{S}(J)$ .

and their dual variants:

(FDer0 right) For each I in Dia the morphism p specializes to a fibered (multi-)category and any Grothendieck opfibration  $\alpha : I \rightarrow J$  in Dia induces a diagram



of fibered (multi-)categories, i.e. the top horizontal functor maps Cartesian arrows w.r.t. the i-th slot to Cartesian arrows w.r.t. the i-th slot.

(FDer3 right) For each functor  $\alpha : I \to J$  in Dia and  $S \in \mathbb{S}(J)$  the functor  $\alpha^*$  between fibers

 $\mathbb{D}(J)_S \to \mathbb{D}(I)_{\alpha^*S}$ 

has a right-adjoint  $\alpha_*^S$ .

(FDer4 right) For each morphism  $\alpha : I \rightarrow J$  in Dia, and for any object  $j \in J$ , and the 2-cell



we get that the induced natural transformation of functors  $j^* \alpha_* \to \alpha_{j*} (\mathbb{S}(\mu))^{\bullet} \iota^*$  is an iso- $morphism<sup>5</sup>$  $morphism<sup>5</sup>$  $morphism<sup>5</sup>$ .

(FDer5 right) (only needed for the multiderivator case). For any functor  $\alpha : I \rightarrow J$  in Dia, and for any a morphism  $\xi \in \text{Hom}(S_1, \ldots, S_n; T)$  in  $\mathbb{S}(\cdot)$  for some  $n \geq 1$ , the natural transformations of functors

$$
\alpha_*(\alpha^*\xi)^{\bullet,i}(\alpha^*-,\cdots,\alpha^*-;-)=\xi^{\bullet,i}(-,\cdots,-; \alpha_*-)
$$

are isomorphisms for all  $1 \leq i \leq n$ .

<span id="page-10-0"></span>**Definition 1.3.6.** A strict morphism of pre-(multi-)derivators  $p : \mathbb{D} \to \mathbb{S}$  with domain Dia is called a left fibered (multi-)derivator with domain Dia, if axioms (Der1–2) hold for  $\mathbb D$  and  $\mathbb S$  and  $(FDer0-5 \; left)$  hold for p. Similarly it is called a right fibered (multi)-derivator with domain Dia, if instead the corresponding dual axioms (FDer0–5 right) hold. It is called just fibered if it is both left and right fibered.

The squares in axioms (FDer4 left/right) are in fact homotopy exact and it follows from the axioms (FDer4 left/right) that many more are (see [1.3.23\)](#page-15-0).

There is some reduncancy in the axioms, cf. [1.3.8](#page-10-2) and [1.3.27.](#page-19-1)

Question 1.3.7. It seems natural to allow also (symmetric) multicategories, in particular operads, as domain for a fibered (symmetric) multiderivator. The author however did not succeed in writing down a neat generalization of (FDer3–4) which would encompass (FDer5).

<span id="page-10-2"></span><span id="page-10-1"></span><sup>&</sup>lt;sup>5</sup>This is meant to hold w.r.t. all bases  $S \in \mathcal{S}(J)$ .

**Lemma 1.3.8.** For a strict morphism of pre-derivators  $\mathbb{D} \rightarrow \mathbb{S}$  such that both satisfy (Der1) and (Der2) and such that it induces a bifibration of multi-categories  $\mathbb{D}(I) \to \mathbb{S}(I)$  for all  $I \in$ Dia we have the following implications:

<span id="page-11-0"></span>
$$
(FDer0left) for n-ary morphisms, n \ge 1 \Leftrightarrow (FDer5right) \tag{4}
$$

 $(FDer0 \ right) \Leftrightarrow (FDer5 \ left)$  (5)

Proof. We will only show the implication [\(4\)](#page-11-0), the other being similar. Choosing pull-back functors  $f^{\bullet}$ , the remaining part of (FDer0) says that the natural 2-morphism

$$
\mathbb{D}(J)_{S_1} \times \cdots \times \mathbb{D}(J)_{S_n} \xrightarrow{f_{\bullet}} \mathbb{D}(J)_T
$$
\n
$$
\downarrow \alpha^* \qquad \qquad \downarrow \qquad \qquad \downarrow \alpha^*
$$
\n
$$
\mathbb{D}(I)_{\alpha^* S_1} \times \cdots \times \mathbb{D}(J)_{\alpha^* S_n} \xrightarrow{(\alpha^* f)_{\bullet}} \mathbb{D}(I)_{\alpha^* T}
$$

is an isomorphism. Taking the adjoint of this diagram (of  $f_{\bullet}$  w.r.t. the *i*-th slot) we get the diagram

$$
\mathbb{D}(I)_{\alpha^*S_1}^{\text{op}} \times \frac{\widehat{i}_{\cdot}}{\alpha^* \mathcal{S}_1} \times \mathbb{D}(I)_{\alpha^*S_n}^{\text{op}} \times \mathbb{D}(I)_{\alpha^*T} \xrightarrow{(\alpha^*f)^{\bullet,i}} \mathbb{D}(J)_{\alpha^*S_i}
$$
  
\n
$$
(\alpha^*)^{\text{op}} \downarrow \qquad \qquad \downarrow \qquad \downarrow
$$

That its 2-morphism is an isomorphism is the content of (FDer5 left). Hence (FDer0 left) and (FDer5 right) are equivalent in this situation.

For [\(5\)](#page-11-0) note that for both (FDer0 right) and (FDer5 left), the functor  $\alpha$  in question is restricted to the class of Grothendieck opfibrations.  $\Box$ 

**Remark 1.3.9.** The axioms (FDer0) and (FDer3–5) are similar to the axioms of a six-functorformalism (cf. the introduction or the appendix  $A.2$ ). It is actually possible to make this analogy precise and define a fibered multiderivator as a bifibration of 2-multicategories  $[\mathbb{D}] \to \text{Di}^{\text{cor}}(\mathbb{S})$ where  $\text{Dia}^{\text{cor}}(\mathbb{S})$  is a certain category of multicorrespondences of diagrams in  $\mathbb{S}$ , similar to our definition of a usual six-functor-formalism (cf. Definition [A.2.16\)](#page-82-0). This also clarifies the existence and comparison of the internal and external monoidal structure, resp. duality, in a closed monoidal derivator (i.e. fibered multiderivator over  $\{\cdot\}$ ) or more generally for any fibered multiderivator. We will explain this in detail in a subsequent article [\[14](#page-86-2)].

1.3.10. The pre-derivator associated with an  $\infty$ -category S is actually a left and right derivator (in the usual sense, i.e. fibered over  $\{\cdot\}$ ) if S is complete and co-complete [\[10\]](#page-86-3). This includes the case of pre-derivators associated with categories, which is, of course, classical — axiom (FDer4) expressing nothing else than Kan's formulas.

**1.3.11.** Let  $S \in \mathbb{S}(\cdot)$  be an object and  $p : \mathbb{D} \to \mathbb{S}$  be a (left, resp. right) fibered multiderivator. The association

$$
I \mapsto \mathbb{D}(I)_{p^*S},
$$

where  $p: I \to \cdot$  is the projection, defines a (left, resp. right) derivator in the usual sense which we call its fiber  $\mathbb{D}_S$  over S. The axioms (FDer6–7) stated below involve only these fibers.

**Definition 1.3.12.** More generally, if  $S \in \mathbb{S}(J)$  we may consider the association

$$
I\mapsto \mathbb{D}(I\times J)_{\operatorname{pr}_2^*S},
$$

where  $pr_2: I \times J \rightarrow J$  is the second projection. This defines again a (left, resp. right) derivator in the usual sense which we call its fiber  $\mathbb{D}_S$  over S.

<span id="page-12-0"></span>**Lemma 1.3.13** (left). Let  $\mathbb{D} \to \mathbb{S}$  be a left fibered multiderivator and let I  $\in$  Dia be a diagram and  $f \in \text{Hom}_{S(J)}(S_1, \ldots, S_n; T)$  for some  $n \geq 1$  be a morphism. Then the collection of functors for each  $J \in \mathrm{Dia}$ 

$$
f_{\bullet}: \mathbb{D}(J \times I)_{\mathrm{pr}_2^* S_1} \times \cdots \times \mathbb{D}(J \times I)_{\mathrm{pr}_2^* S_n} \rightarrow \mathbb{D}(J \times I)_{\mathrm{pr}_2^* T}
$$
  

$$
\mathcal{E}_1, \ldots, \mathcal{E}_n \rightarrow (\mathrm{pr}_2^* f)_{\bullet}(\mathcal{E}_1, \ldots, \mathcal{E}_n)
$$

defines a morphism of derivators  $\mathbb{D}_{S_1} \times \cdots \times \mathbb{D}_{S_n} \to \mathbb{D}_T$ . Furthermore, for a collection  $\mathcal{E}_k \in \mathbb{D}(I), k \neq i$ the morphism of derivators:

$$
\mathbb{D}(J \times I)_{\mathrm{pr}_2^* S_i} \rightarrow \mathbb{D}(J \times I)_{\mathrm{pr}_2^* T}
$$
  

$$
\mathcal{E}_i \rightarrow (\mathrm{pr}_2^* f)_\bullet (\mathrm{pr}_2^* \mathcal{E}_1, \dots, \mathcal{E}_i, \dots, \mathrm{pr}_2^* \mathcal{E}_n)
$$

is left continuous (i.e. commutes with left Kan extensions).

Proof. The only point which might not be clear is the left continuity of the bottom morphism of pre-derivators. Consider the following 2-commutative square, where  $I, J, J' \in \text{Dia}, \alpha : J \to J'$  is a functor, and  $j' \in J'$ 

$$
I \times (j' \times_{/J'} J) \xrightarrow{\text{(id, }\iota)} I \times J
$$
  
(id,p)  

$$
I \times j' \xrightarrow{\text{(id, }\iota)} I \times J'
$$

It is homotopy exact by [1.3.23,](#page-15-0) 4. Therefore we have (using FDer3–5 left):

$$
(\mathrm{id}, j')^*(\mathrm{id}, \alpha)_{!}(\mathrm{pr}_1^* f)_{\bullet}(\mathrm{pr}_1^* \mathcal{E}_1, \dots, \mathcal{E}_i, \dots, \mathrm{pr}_1^* \mathcal{E}_n)
$$
\n
$$
\cong (\mathrm{id}, p)_{!}(\mathrm{id}, \iota)^*(\mathrm{pr}_1^* f)_{\bullet}(\mathrm{pr}_1^* \mathcal{E}_1, \dots, \mathcal{E}_i, \dots, \mathrm{pr}_1^* \mathcal{E}_n)
$$
\n
$$
\cong (\mathrm{id}, p)_{!}(\mathrm{pr}_1^* f)_{\bullet}((\mathrm{id}, \iota)^* \mathrm{pr}_1^* \mathcal{E}_1, \dots, (\mathrm{id}, \iota)^* \mathcal{E}_i, \dots, (\mathrm{id}, \iota)^* \mathrm{pr}_1^* \mathcal{E}_n)
$$
\n
$$
\cong (\mathrm{id}, p)_{!}(\mathrm{pr}_1^* f)_{\bullet}((\mathrm{id}, p)^* \mathcal{E}_1, \dots, (\mathrm{id}, \iota)^* \mathcal{E}_i, \dots, (\mathrm{id}, p)^* \mathcal{E}_n)
$$
\n
$$
\cong f_{\bullet}(\mathcal{E}_1, \dots, (\mathrm{id}, j')^*(\mathrm{id}, \alpha)_{!} \mathcal{E}_i, \dots, \mathcal{E}_n)
$$
\n
$$
\cong (\mathrm{id}, j')^*(\mathrm{pr}_1^* f)_{\bullet}(\mathrm{pr}_1^* \mathcal{E}_1, \dots, (\mathrm{id}, \alpha)_{!} \mathcal{E}_i, \dots, \mathrm{pr}_1^* \mathcal{E}_n)
$$

(Note that  $(id, p)$  is trivially a Grothendieck op-fibration). A tedious check shows that the composition of these isomorphisms is  $(id, j')^*$  applied to the exchange morphism

$$
(\mathrm{id}, \alpha)_! (\mathrm{pr}_1^* f)_\bullet (\mathrm{pr}_1^* \mathcal{E}_1, \ldots, \mathcal{E}_i, \ldots, \mathrm{pr}_1^* \mathcal{E}_n) \to (\mathrm{pr}_1^* f)_\bullet (\mathrm{pr}_1^* \mathcal{E}_1, \ldots, (\mathrm{id}, \alpha)_! \mathcal{E}_i, \ldots, \mathrm{pr}_1^* \mathcal{E}_n)
$$

Since the above holds for any  $j' \in J'$  the exchange morphism is therefore an isomorphism by (Der2).  $\Box$ 

In the right fibered situation the analogously defined morphisms  $f^{i, \bullet}$  are not expected to be made into a morphism of fibers this way. For a discussion of how this is solved, we refer the reader to the article [\[14](#page-86-2)] in preparation, where a fibered multiderivator is redefined as a certain type of six-functor-formalism. This will let appear the discussion and results of this section in a much more clear fashion. However, we have:

<span id="page-13-0"></span>**Lemma 1.3.14** (right). Let  $\mathbb{D} \to \mathbb{S}$  be a right fibered multiderivator and let I  $\in$  Dia be a diagram and  $f \in \text{Hom}_{S(J)}(S_1, \ldots, S_n; T)$ , for some  $n \geq 1$ , be a morphism. For each  $J \in$  Dia and for each collection  $\mathcal{E}_k \in \mathbb{D}(I)$ ,  $k \neq i$ , the association

$$
\mathbb{D}(J \times I)_{\mathrm{pr}_2^* T} \rightarrow \mathbb{D}(J \times I)_{\mathrm{pr}_2^* S_i}
$$
  

$$
\mathcal{F} \rightarrow (\mathrm{pr}_2^* f)^{\bullet, i} (\mathrm{pr}_2^* \mathcal{E}_1, \dots, \mathrm{pr}_2^* \mathcal{E}_n; \mathcal{F})
$$

defines a morphism of derivators which is right continuous (i.e. commutes with right Kan extensions). This is the right adjoint in the pre-derivator sense to the morphism of pre-derivators in the previous lemma, as soon as  $\mathbb{D} \to \mathbb{S}$  is left and right fibered.

*Proof.* Consider the following 2-commutative square where  $I, J, J' \in \text{Dia}, \alpha : J \to J'$  is a functor, and  $j' \in J'$ 

$$
I \times (J \times_{/J'} j') \xrightarrow{\text{(id, }\iota)} I \times J
$$
  
(id,p)  

$$
I \times j' \xrightarrow{\text{(id, }\iota)} I \times J'
$$

It is homotopy exact by [1.3.23,](#page-15-0) 4.

Therefore we have (using FDer3–5 right):

$$
(\mathrm{id}, j')^*(\mathrm{id}, \alpha)_{!}(\mathrm{pr}_1^* f)^{i, \bullet}(\mathrm{pr}_1^* \mathcal{E}_1, \widehat{\cdot}, \mathrm{pr}_1^* \mathcal{E}_n; \mathcal{F})
$$
  
\n
$$
\cong (\mathrm{id}, p)_*(\mathrm{id}, \iota)^*(\mathrm{pr}_1^* f)^{i, \bullet}(\mathrm{pr}_1^* \mathcal{E}_1, \widehat{\cdot}, \mathrm{pr}_1^* \mathcal{E}_n, \mathcal{F})
$$
  
\n
$$
\cong (\mathrm{id}, p)_*(\mathrm{pr}_1^* f)^{i, \bullet}((\mathrm{id}, \iota)^* \mathrm{pr}_1^* \mathcal{E}_1, \widehat{\cdot}, \widehat{\cdot}, (\mathrm{id}, \iota)^* \mathrm{pr}_1^* \mathcal{E}_n; (\mathrm{id}, \iota)^* \mathcal{F})
$$
  
\n
$$
\cong (\mathrm{id}, p)_*(\mathrm{pr}_1^* f)^{i, \bullet}((\mathrm{id}, p)^* \mathcal{E}_1, \widehat{\cdot}, \cdot, (\mathrm{id}, p)^* \mathcal{E}_n; (\mathrm{id}, \iota)^* \mathcal{F})
$$
  
\n
$$
\cong f^{i, \bullet}(\mathcal{E}_1, \widehat{\cdot}, \mathcal{E}_n; (\mathrm{id}, p)_*(\mathrm{id}, \iota)^* \mathcal{F})
$$
  
\n
$$
\cong f^{i, \bullet}(\mathcal{E}_1, \widehat{\cdot}, \mathcal{E}_n; (\mathrm{id}, j')^*(\mathrm{id}, \alpha)_* \mathcal{F})
$$
  
\n
$$
\cong (\mathrm{id}, j')^*(\mathrm{pr}_1^* f)^{i, \bullet}(\mathrm{pr}_1^* \mathcal{E}_1, \widehat{\cdot}, \cdot, \mathrm{pr}_1^* \mathcal{E}_n; (\mathrm{id}, \alpha)_! \mathcal{F})
$$

Note that  $(id, \iota)$  is a Grothendieck op-fibration, but  $(id, j')$  is not. Hence the last step has to be justified further. Consider the 2-commutative diagram:

$$
I \times (J \times_{/J'} j') \xrightarrow{\text{(id,i')}} I \times J'
$$
  
(id,p)  

$$
I \times j' \xrightarrow{\text{(id,i')}} I \times J'
$$

It is again homotopy exact by [1.3.23,](#page-15-0) 4. Therefore we have

$$
\begin{array}{lll}\n\cong&& f^{i,\bullet}(\mathcal{E}_1,\hat{i}_-,\mathcal{E}_n;(\mathrm{id},j')^*(\mathrm{id},\alpha)_*\mathcal{F})\\
\cong&& f^{i,\bullet}(\mathcal{E}_1,\hat{i}_-,\mathcal{E}_n;(\mathrm{id},p)_*(\mathrm{id},\iota')^*(\mathrm{id},\alpha)_*\mathcal{F})\\
\cong&(\mathrm{id},p)_*(\mathrm{pr}_1^*f)^{i,\bullet}((\mathrm{id},p)^*\mathcal{E}_1,\hat{i}_-,( \mathrm{id},p)^*\mathcal{E}_n;(\mathrm{id},\iota')^*(\mathrm{id},\alpha)_*\mathcal{F})\\
\cong&(\mathrm{id},p)_*(\mathrm{pr}_1^*f)^{i,\bullet}((\mathrm{id},\iota')^*\mathrm{pr}_1^*\mathcal{E}_1,\hat{i}_-,( \mathrm{id},\iota')^*\mathrm{pr}_1^*\mathcal{E}_n;(\mathrm{id},\iota')^*(\mathrm{id},\alpha)_*\mathcal{F})\\
\cong&(\mathrm{id},p)_*(\mathrm{id},\iota')^*(\mathrm{pr}_1^*f)^{i,\bullet}(\mathrm{pr}_1^*\mathcal{E}_1,\hat{i}_-,\hat{i}_-,\mathcal{E}_n;(\mathrm{id},\iota')^*(\mathrm{id},\alpha)_*\mathcal{F})\\
\cong&(\mathrm{id},j')^*(\mathrm{pr}_1^*f)^{i,\bullet}(\mathrm{pr}_1^*\mathcal{E}_1,\hat{i}_-,\mathrm{pr}_1^*\mathcal{E}_n;(\mathrm{id},\alpha)_*\mathcal{F})\n\end{array}
$$

Note that  $(id, \iota')$  is a Grothendieck opfibration, too. In other words: the reason why  $f^{\bullet, i}$  also commutes with  $(id, j')^*$  in this particular case is that the other argument are constant in the J direction.

A tedious check shows the composition of the isomorphisms of the previous computations yield  $(id, j')^*$  applied to the exchange morphism

$$
(\mathrm{id}, \alpha)_* \left(\mathrm{pr}_1^* f\right)^{i, \bullet} \left(\mathrm{pr}_1^* \mathcal{E}_1, \widehat{\cdot}^i, \mathrm{pr}_1^* \mathcal{E}_n; \mathcal{F}\right) \to \left(\mathrm{pr}_1^* f\right)^{i, \bullet} \left(\mathcal{E}_1, \widehat{\cdot}^i, \mathcal{E}_n; (\mathrm{id}, \alpha)_* \mathcal{F}\right).
$$

Since the above holds for any  $j' \in J'$  it is therefore an isomorphism by (Der2).

**1.3.15.** Let  $p : \mathbb{D} \to \mathbb{S}$  be a (left, resp. right) fibered multi-derivator and  $S : \{\cdot\} \to \mathbb{S}(\cdot)$  a functor of multicategories. This is equivalent to the choice of an object  $S \in \mathbb{S}(\cdot)$  and a collection of morphisms  $\alpha_n \in \text{Hom}_{\mathbb{S}(\cdot)}(S,\ldots,S;S)$  for all  $n \geq 2$ , compatible with composition. Then the fiber

 $\overbrace{ }$  *n* times

$$
I \mapsto \mathbb{D}(I)_{p^*S}
$$

defines even a (left, resp. right) multiderivator (i.e. a fibered multiderivator over  $\{\cdot\}$ ). The same holds analogously for a functor of multicategories  $S: \{\cdot\} \to \mathbb{S}(I)$ .

Axiom (FDer5 left) and Lemma [A.2.6](#page-78-0) imply the following:

**Proposition 1.3.16.** The definition of a left fibered multiderivator  $\mathbb{D} \to \{\cdot\}$  is equivalent to the definition of a monoidal left derivator in the sense of Groth [\[7\]](#page-86-4). It is also, in addition, right fibered if and only if it is a right derivator and closed monoidal in the sense of [loc. cit.].

**Definition 1.3.17.** We call a pre-derivator  $\mathbb{D}$  strong, if the following axiom holds:

(Der8) For any diagram K in Dia the 'partial underlying diagram' functor

$$
dia: \mathbb{D}(K \times \Delta_1) \to \text{Hom}(\Delta_1, \mathbb{D}(K))
$$

is full and essentially surjective.

**Definition 1.3.18.** Let  $p : \mathbb{D} \to \mathbb{S}$  be a fibered (left and right) derivator. We call  $\mathbb{D}$  pointed (relative to  $p$ ) if the following axiom holds:

(FDer6) For any  $S \in \mathbb{S}(\cdot)$ , the category  $\mathbb{D}(\cdot)_S$  has a zero object.

**Definition 1.3.19.** Let  $p : \mathbb{D} \to \mathbb{S}$  be a fibered (left and right) derivator. We call  $\mathbb{D}$  stable (relative to p) if its fibers are strong and the following axiom holds:

 $\Box$ 

(FDer7) For any  $S \in \mathbb{S}(\cdot)$ , in the category  $\mathbb{D}(\square)_{p^*S}$  an object is homotopy Cartesian if and only if it is homotopy coCartesian.

This condition can be weakened (cf. [\[9,](#page-86-5) Corollary 8.13]).

1.3.20. Recall from [\[8\]](#page-86-1) that axiom (FDer7) implies that the fibers of a stable fibered derivator are triangulated categories in a natural way. Actually the proof shows that it suffices to have a derivator of domain Posf (finite posets).

Since, by Lemma [1.3.13](#page-12-0) and Lemma [1.3.14](#page-13-0) push-forward, resp. pull-back w.r.t. any slot commute with homotopy colimits, resp. homotopy limits, they induce triangulated functors between the fibers.

**1.3.21** (left). The following is a consequence of (FDer0): For a functor  $\alpha : I \rightarrow J$  and a morphism in  $f$  :  $S$  →  $T$  ∈  $\mathbb{S}(J)$ , we get a natural isomorphism

$$
\mathbb{S}(\alpha^* f) \bullet \alpha^* \to \alpha^* \mathbb{S}(f) \bullet.
$$

<span id="page-15-1"></span>W.r.t. this natural isomorphism we have the following:

**Lemma 1.3.22** (left). Given a "pasting" diagram

$$
N \xrightarrow{G} L \xrightarrow{B} I
$$
  
\n
$$
\downarrow{A \mathscr{L}} \qquad \downarrow{a \mathscr{L}} \qquad \downarrow{\alpha}
$$
  
\n
$$
M \xrightarrow{\gamma} K \xrightarrow{\beta} J
$$

we get for the pasted natural transformation  $\nu \odot \mu := (\beta * \nu) \circ (\mu * G)$  that the following diagram is commutative:



Here the morphisms going to the right are (induced by) the various base-change morphisms. In particular, the pasted square is homotopy exact if the individual two squares are.

<span id="page-15-0"></span>Proof. This is an analogue of [\[8](#page-86-1), Lemma 1.17] and proven similarly.

Proposition 1.3.23. 1. Any square of the form

$$
I \times_{/J} K \xrightarrow{B} I
$$
  
\n
$$
A \downarrow \qquad \varphi^{\mu} \downarrow \alpha
$$
  
\n
$$
K \xrightarrow{\beta} J
$$

(where  $I \times_{i,j} K$  is the slice category) is homotopy exact (in particular the ones from axiom FDer4 left and FDer4 right are).

 $\Box$ 

#### 2. A Cartesian square

$$
I \times_J K \xrightarrow{B} I
$$
  
A  

$$
K \xrightarrow{\beta} \sqrt{\alpha}
$$
  

$$
K \xrightarrow{\beta} J
$$

(where  $I \times J K$  is the fiber product) is homotopy exact, if  $\alpha$  is a Grothendieck opfibration or if  $\beta$  is a Grothendieck fibration.

3. If  $\alpha : I \to J$  is a morphism of Grothendieck opfibrations over a diagram E, then



is homotopy exact for all objects  $e \in E$ .

4. If a square

$$
L \xrightarrow{\text{B}} I
$$
  
A  

$$
\downarrow^{\text{B}} \downarrow^{\text{A}}
$$
  

$$
K \xrightarrow{\text{B}} J
$$

is homotopy exact then so is the square

$$
L \times X \xrightarrow{\beta} I \times X
$$
  
A  

$$
A \downarrow \qquad \qquad \downarrow \alpha
$$
  

$$
K \times X \xrightarrow{\beta} J \times X
$$

for any diagram X.

Proof. This proof is completely analogous to the non-fibered case. We sketch the arguments here (for the left-case only, the other case follows by logical duality): 3. Let  $j$  be an object in  $J_e$  and consider the cube:



where w is given by the inclusions  $\iota_{I,e}$  resp.  $\iota_{J,e}$ . By standard arguments on homotopy exact squares it suffices to show that the left square is homotopy exact on constant diagrams, i.e. that

$$
p_{e,!}w^* \cong p_!
$$

holds true for all usual derivators. By  $[8,$  Proposition 1.23 it suffices to show that w has a left adjoint.

Denote  $\pi_I : I \to E$  and  $\pi_J : J \to E$  the given opfibrations. Consider the two functors

$$
I_e \times_{J,e} j \xrightarrow{w} I \times_{J} j
$$

where c is given by mapping  $(i, \mu : \alpha(i) \to j)$  to  $(i', \mu' : \alpha(i') \to j)$  where we chose, for any i, a coCartesian morphism  $\xi_{i,\mu} : i \to i'$  over  $\pi_I(\mu) : \pi_I(i) \to e$ . Since  $\alpha$  maps coCartesian morphisms to coCartesian morphisms by assumption,  $\alpha(\xi_{i,\mu}) : \alpha(i) \to \alpha(i')$  is coCartesian, and therefore there is a unique factorization

$$
\alpha(i) \xrightarrow{\alpha(\xi_i)} \alpha(i') \xrightarrow{\mu'} j
$$

of  $\mu$ . A morphism  $\alpha$  :  $(i_1, \mu_1 : \alpha(i_1) \to j) \to (i_2, \mu_2 : \alpha(i_2) \to j)$ , by definition of coCartesian, gives rise to a unique morphism  $\alpha': i'_1 \to i'_2$  over  $\pi_I(i_1) \to \pi_I(i_2)$  such that  $\alpha' \xi_{i_1,\mu_1} = \xi_{i_2,\mu_2} \alpha'$  holds, and we set  $c(\alpha) \coloneqq \alpha'$ . We have  $c \circ w = id$ , and a morphism  $id_{I \times_{/} j} \to w \circ c$  given by  $(i, \mu) \mapsto \xi_{i, \mu}$ . This makes  $w$  right adjoint to  $c$ .

2. By axiom (Der2) it suffices to show that for any object  $k$  of  $K$ , the induced morphism

$$
k^*A_!B^*\to k^*\beta^*\alpha_!
$$

is an isomorphism. Consider the following pasting diagram

$$
I \times_J k \xrightarrow{j} I \times_J K \times_{/K} k \xrightarrow{\iota} I \times_J K \xrightarrow{B} I
$$
  
\n
$$
\downarrow \pi \qquad \qquad \downarrow \qquad
$$

Lemma [1.3.22](#page-15-1) shows that the following composition

$$
\pi_!\mathbb{S}(\beta*\mu * j) \bullet j^* \iota^* B^* \to \pi_!\jmath^* \mathbb{S}(\beta*\mu) \bullet \iota^* B^* \to p_!\mathbb{S}(\beta*\mu) \bullet \iota^* B^* \to k^* A_!\B^* \to k^* \beta^* \alpha_!\
$$

is the base-change associated with the pasting of the 3 squares in the diagram. All morphisms in this sequence are isomorphisms except possibly for the rightmost one. The second from the left is an isomorphism because j is a right adjoint [\[8,](#page-86-1) Proposition 1.23]. The base-change morphism of the pasting is an isomorphism because of 3.

1. By axiom (Der2) it suffices to show that for any object k of K the induced morphism

$$
k^*A_!\mathbb{S}(\mu)_\bullet B^* \to k^*\beta^*\alpha_!
$$

is an isomorphism. Consider the following pasting diagram

$$
I \times_{/J} k \xrightarrow{k} I \times_{/J} K \xrightarrow{B} I
$$
  
\n
$$
\downarrow p
$$
  
\n
$$
k \xrightarrow{k} K \xrightarrow{\beta} J
$$

Lemma [1.3.22](#page-15-1) shows that the following diagram is commutative

$$
p_!\mathbb{S}(\mu\iota)_\bullet \iota^* B^* \xrightarrow{\sim} k^* \beta^* \alpha_!
$$
  
can. 
$$
\downarrow \sim
$$
  

$$
p_!\iota^* \mathbb{S}(\mu)_\bullet B^* \xrightarrow{\sim} k^* A_!\mathbb{S}(\mu)_\bullet B^*
$$

where the bottom horizontal morphism is an isomorphism by 2., and the top horizontal morphism is an isomorphism by (FDer4 left). Therefore the right vertical morphism is also an isomorphism. 4. (cf. also [\[8,](#page-86-1) Theorem 1.30]). For any  $x \in X$  consider the cube

$$
\begin{array}{ccc}\n & L & \xrightarrow{B} & I \\
\hline\n & & & \downarrow \\
 & & & \downarrow \\
A & & & \downarrow \\
 & & & \downarrow \\
A & & & \downarrow \\
 & & & \downarrow \\
A & & & \downarrow \\
 & & & \down
$$

The left and right hand side squares are homotopy exact because of 3., whereas the rear one is homotopy exact by assumption. Therefore the pasting

$$
L \longrightarrow I \times X
$$
  
A  

$$
A \downarrow \qquad \qquad \downarrow \alpha
$$
  

$$
K \longrightarrow J \times X
$$

is homotopy exact. Therefore we have an isomorphism

$$
(id,x)^*A_!\mathbb{S}(\mu)_\bullet B^* \to (id,x)^*\beta^*\alpha_!
$$

where the morphism is induced by the base change of the given 2-commutative square. We may then conclude by axiom (Der2).  $\Box$ 

<span id="page-18-0"></span>1.3.24 (left). If S is strong the pull-backs and push-forwards along a morphism in  $\mathcal{S}(\cdot)$ , or more generally along a morphism in  $\mathcal{S}(I)$ , can be expressed using only the relative Kan-extension functors: Let  $p : \mathbb{D} \to \mathbb{S}$  be a left fibered derivator such that S is strong. Consider the 2-commutative square



and consider a morphism  $f : S \to T$  in  $\mathcal{S}(I)$ . By the strongness of S, the morphism f may be lifted to an object  $F \in \mathcal{S}(I \times \Delta_1)$ , and this means that the morphism

$$
\mathbb{S}(\mu)_\bullet : p^* F \to \iota^* F
$$

is isomorphic to f. Since the square is homotopy exact by Proposition [1.3.23](#page-15-0) 1., we get that the natural transformation

$$
f_{\bullet} \to \iota^* p_!
$$

is an isomorphism.

**1.3.25** (left). Let  $\alpha : I \to J$  a functor in Dia and let  $f : S \to T$  be a morphism in  $\mathbb{S}(J)$ . Axiom (FDer0) of a left fibered derivator implies that we have a canonical isomorphism

$$
(\alpha^*(f))_\bullet\alpha^*=\alpha^*f_\bullet
$$

which is determined by the choice of the push-forward functors. We get an associated exchange morphism

<span id="page-19-2"></span>
$$
\alpha_!(\alpha^*(f))_{\bullet} \to f_{\bullet}\alpha_!.
$$
 (8)

<span id="page-19-4"></span>**Proposition 1.3.26.** If  $p : \mathbb{D} \to \mathbb{S}$  is a left fibered derivator then the natural transformation [\(8\)](#page-19-2) is an isomorphism. The corresponding dual statement holds for a right fibered derivator.

Proof. Consider the following 2-commutative squares (the third and fourth are even commutative on the nose):

<span id="page-19-3"></span>
$$
I = I \t J = J \t J \t (9)
$$
  
\n
$$
I \t N \Delta_1
$$
  
\n
$$
\downarrow \t J \t N \Delta_1
$$
  
\n
$$
I \t N \Delta_1
$$
  
\n
$$
\downarrow \t J \t N \Delta_1
$$
  
\n
$$
I \t N \Delta_1
$$
  
\n
$$
\downarrow \t J \t N \Delta_1
$$
  
\n
$$
I \t N \Delta_1
$$
  
\n
$$
\downarrow \t J \t N \Delta_1
$$
  
\n
$$
I \t N \Delta_1
$$
  
\n
$$
\downarrow \t J \t N \Delta_1
$$
  
\n
$$
I \t N \Delta_1
$$
  
\n
$$
\downarrow \t J \t N \Delta_1
$$
  
\n
$$
\downarrow \t J \t N \Delta_1
$$
  
\n
$$
\downarrow \t J \t N \Delta_1
$$
  
\n
$$
\downarrow \t J \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$
  
\n
$$
\downarrow \t N \t N \Delta_1
$$

They are all homotopy exact. Consider the diagram

$$
\alpha_!(\alpha^*(f)) \bullet \longrightarrow f_{\bullet}\alpha_!
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$
\n
$$
\alpha_!\iota_I^* p_{I,!} \longrightarrow \iota_J^* \alpha_! p_{I,!} \longrightarrow \iota_J^* p_{J,!} \alpha_!
$$

where the vertical morphisms come from  $(1.3.24)$  — these are the base change morphism for the first and second square in  $(9)$  — and the lower horizontal morphisms are respectively the base change for the third diagram in [\(9\)](#page-19-3), and the natural morphism associated with the commutativity of the fourth diagram in [\(9\)](#page-19-3). Repeatedly appying Lemma [1.3.22](#page-15-1) shows that this diagram is commutative. Therefore the upper horizontal morphism is an isomorphism because all the others in the diagram are. П

<span id="page-19-1"></span>1.3.27. The last proposition states that push-forward commutes with homotopy colimits (left case) and pull-back commutes with homotopy limits (right case). This is also the content of (FDer5 left/right) for fibered derivators (not multiderivators), and hence this axiom is implied by the other axioms of left fibered derivators. Even in the multi-case, by Lemma [1.3.8,](#page-10-2) axiom (FDer5 left/right) also follow from both (FDer0 left) and (FDer0 right).

<span id="page-19-0"></span>**1.3.28** (left). Let  $\alpha: I \to J$  be a functor in Dia. Proposition [1.3.26](#page-19-4) allows us to extend the functor  $\alpha_!$  to a functor

$$
\alpha_! : \mathbb{D}(I) \times_{\mathbb{S}(I)} \mathbb{S}(J) \to \mathbb{D}(J)
$$

which is still left adjoint to  $\alpha^*$ , more precisely: to  $(\alpha^*, p(J))$ . Here the fiber product is formed w.r.t.  $p(I)$  and  $\alpha^*$  respectively. We sketch its construction:  $\alpha_!(\mathcal{E},S)$  is given by  $\alpha_!^S \mathcal{E}$ , where  $\alpha_!^S$ is the functor from axiom (FDer3 left) with base S. Let a pair of a morphism  $f : S \to T$  in  $\mathbb{S}(J)$ and  $F: \mathcal{E} \to \mathcal{F}$  in  $\mathbb{D}(I)$  over  $\alpha^*(f)$  be given. We define  $\alpha_!(F,f)$  as follows: F corresponds to a morphism

$$
(\alpha^* f) \bullet \mathcal{E} \bullet \mathcal{F}.
$$

Applying  $\alpha_!^T$  we get a morphism

$$
\alpha_!^T(\alpha^*f)_\bullet \mathcal{E} \to \alpha_!^T \mathcal{F}
$$

and composition with the inverse of the morphism [\(8\)](#page-19-2) yields

$$
f_{\bullet}\alpha_!^S \mathcal{E} \rightarrow \alpha_!^T \mathcal{F}
$$

or, equivalently, a morphism which we define to be  $\alpha_1(F, f)$ 

$$
\alpha_!^S \mathcal{E} \to \alpha_!^T \mathcal{F}
$$

over  $f$ .

For the adjunction, we have to give a functorial isomorphism

$$
\mathrm{Hom}_{\alpha^*f}(\mathcal{E}, \alpha^* \mathcal{F}) \cong \mathrm{Hom}_f(\alpha_! (\mathcal{E}, S), \mathcal{F}),
$$

where  $\mathcal{E} \in \mathbb{D}(I)_{\alpha^*S}$  and  $\mathcal{F} \in \mathbb{D}(J)_T$ . We define it to be the following composition of isomorphisms:

$$
\text{Hom}_{\alpha^* f}(\mathcal{E}, \alpha^* \mathcal{F})
$$
\n
$$
\cong \text{Hom}_{\text{id}_{\alpha^* T}}((\alpha^* f)_\bullet \mathcal{E}, \alpha^* \mathcal{F})
$$
\n
$$
\cong \text{Hom}_{\text{id}_T}(\alpha_! (\alpha^* f)_\bullet \mathcal{E}, \mathcal{F})
$$
\n
$$
\cong \text{Hom}_{\text{id}_T}(f_\bullet \alpha_! \mathcal{E}, \mathcal{F})
$$
\n
$$
\cong \text{Hom}_f(\alpha_! \mathcal{E}, \mathcal{F}).
$$

A dual statement holds for a right fibered derivator and the functor  $\alpha_*$ .

<span id="page-20-0"></span>From Proposition [1.3.26](#page-19-4) we also get a vertical version of Lemma [1.3.22:](#page-15-1)

**Lemma 1.3.29** (left). Given a "pasting" diagram

$$
N \xrightarrow{B} M
$$
  
\n
$$
\Gamma \downarrow \xrightarrow{\varphi \nu} M
$$
  
\n
$$
L \xrightarrow{b} I
$$
  
\n
$$
a \downarrow \xrightarrow{\varphi \mu} a
$$
  
\n
$$
K \xrightarrow{\beta} J
$$

we get for the pasted natural transformation  $\mu \odot \nu = (\mu * \Gamma) \circ (\alpha * \nu)$  that the following diagram is commutative:



Here the morphisms going to the right are (induced by) the various base-change morphisms and the upper horizontal morphism is the isomorphism from Proposition [1.3.26.](#page-19-4) In particular, the pasted square is homotopy exact if the two individual squares are.

#### <span id="page-21-0"></span>1.4 Transitivity

#### Proposition 1.4.1. Let

$$
\mathbb{E} \xrightarrow{p_1} \mathbb{D} \xrightarrow{p_2} \mathbb{S}
$$

be two left (resp. right) fibered multiderivators. Then also the composition  $p_3 = p_2 \circ p_1 : \mathbb{E} \to \mathbb{S}$  is a left (resp. right) fibered multiderivator.

Proof. We will show the statement for left fibered multiderivators. The other statement follows by logical duality.

Axiom (FDer0): For any  $I \in$  Dia, we have a sequence

$$
\mathbb{E}(I) \to \mathbb{D}(I) \to \mathbb{S}(I)
$$

of fibered multicategories. It is well-known that then also the composition  $\mathbb{E}(I) \to \mathbb{S}(I)$  is a fibered multicategory (see [A.2\)](#page-75-2). The other statement of (FDer0) is immediate, too. Let  $\alpha : I \to J$  be a functor as in axioms (FDer3 left) and (FDer4 left). We denote the relative homotopy Kan-extension functors w.r.t. the 2 fibered derivators by  $\alpha_!^1$ , and  $\alpha_!^2$ , respectively. As always, the base will be understood from the context or explicitly given as extra argument as in [\(1.3.28\)](#page-19-0). Axiom (FDer3 left): Let  $S \in \mathbb{S}(J)$  be given. We define a functor

$$
\alpha_!^3:\mathbb{E}(I)_{\alpha^*S}\to\mathbb{E}(J)_S
$$

in the fiber (under  $p_2$ ) of  $\mathcal{E} \in \mathbb{D}(I)_{\alpha^*S}$  as the composition

$$
\mathbb{E}(I)_{\alpha^*S} \xrightarrow{(\nu_{\bullet}, \alpha_!^2 p_1)} \mathbb{E}(I)_{\alpha^*S} \times_{\mathbb{D}(I)_{\alpha^*S}} \mathbb{D}(J)_S \xrightarrow{\alpha_!^1} \mathbb{E}(J)_S
$$

where  $\nu$  is the unit

$$
\nu:\mathcal{E}\rightarrow \alpha^*\alpha^2_!\mathcal{E}
$$

and  $\alpha_!^1$  with two arguments is the extension given in [\(1.3.28\)](#page-19-0). Let  $\mathcal{F}_1 \in \mathbb{E}(I)_{\alpha^*S}$  and  $\mathcal{F}_2 \in \mathbb{E}(J)_S$  be given with images  $\mathcal{E}_1$  and  $\mathcal{E}_2$ , respectively under  $p_1$ . The adjunction is given by the following composition of isomorphisms:

$$
Hom_S(\alpha_1^3 \mathcal{F}_1, \mathcal{F}_2)
$$
\n= Hom<sub>S</sub>(α<sub>1</sub><sup>1</sup>(ν<sub>•</sub>  $\mathcal{F}_1, \alpha_1^2 \mathcal{E}_1), \mathcal{F}_2)$  Definition\n  
\n= {f ∈ Hom<sub>S</sub>(α<sub>1</sub><sup>2</sup>  $\mathcal{E}_1, \mathcal{E}_2); \xi ∈ Homf(α11(ν•  $\mathcal{F}_1, \alpha_1^2 \mathcal{E}_1), \mathcal{F}_2)$ } Definition\n  
\n
$$
f ∈ HomS(α12  $\mathcal{E}_1, \mathcal{E}_2); \xi ∈ Homα* f(ν•  $\mathcal{F}_1, \alpha^2 \mathcal{F}_2$ )}$  Adjunction (1.3.28)\n  
\n
$$
f ∈ Homα* S(\mathcal{E}_1, \alpha^* \mathcal{E}_2); \xi ∈ Homf( $\mathcal{F}_1, \alpha^* \mathcal{F}_2$ )} Note below\n  
\n= Hom<sub>α* S</sub>( $\mathcal{F}_1, \alpha^* \mathcal{F}_2$ )} Denmitian
$$
$$$ 

Note that the composition

$$
\widetilde{f}: \mathcal{E}_1 \longrightarrow \alpha^* \alpha_!^2 \mathcal{E}_1 \xrightarrow{\alpha^* f} \alpha^* \mathcal{E}_2
$$

is determined by f via the adjunction of (FDer3 left) for base S and  $p_2 : \mathbb{D} \to \mathbb{S}$ . Axiom (FDer4 left): Let  $\mathcal E$  be in  $\mathbb E(I)_{\alpha^*S}$  and let  $\mathcal F$  be its image under  $p_1$ . We have to show that the natural morphism

$$
\alpha_j^3 \mathbb{S}(\mu) \partial^3 \iota^* \mathcal{E} \to j^* \alpha_!^3
$$

is an isomorphism. Inserting the definition of the push-forwards, resp. of the Kan extensions for  $p_3$ , we get

$$
\alpha_{j!}^1(\nu_j) \cdot \operatorname{cart}_{\bullet}^1 \iota^* \mathcal{E} \to j^* \alpha_!^1 \nu_{\bullet}^1 \mathcal{E}.
$$

Here  $\nu_j : \mathbb{S}(\mu)\mathbb{I}^2 \iota^* \mathcal{F} \to \alpha_j^* \alpha_j^2 \mathbb{S}(\mu)\mathbb{I}^2 \iota^* \mathcal{F}$  is the unit and  $\nu : \mathcal{F} \to \alpha^* \alpha_j^2 \mathcal{F}$  is the unit. 'cart<sup>1</sup>' is the Cartesian morphism  $\iota^* \mathcal{F} \to \mathbb{S}(\mu)\mathcal{E}^* \mathcal{F}$ . Consider the base-change isomorphism (FDer4 for  $p_2$ )

$$
\mathrm{bc}: \alpha_{j\cdot}^2 \mathbb{S}(\mu)_\bullet^2 \iota^* \mathcal{F} \to j^* \alpha_!^2 \mathcal{F},
$$

and the morphism

$$
\mathbb{D}(\mu): \iota^* \alpha^* \alpha_!^2 \mathcal{F} \to \alpha_j^* j^* \alpha_!^2 \mathcal{F}.
$$

Claim: We have the equality

$$
(\alpha_j^*bc)\circ\nu_j\circ \mathrm{cart}=\mathbb{D}(\mu)\circ\iota^*(\nu).
$$

*Proof of the claim:* Consider the diagram (which affects only the fibered derivator  $p_2 : \mathbb{D} \to \mathbb{S}$ , hence we omit superscripts):



Clearly all squares and triangles in this diagram are commutative. The two given morphisms are the compositions of the extremal paths hence they are equal. We have a natural isomorphism induced by bc:

$$
\alpha_{j!}^1(\cdots,\alpha_{j!}^2\mathbb{S}(\mu)\mathbf{1}_{\bullet}\mathbf{1}^*\mathcal{F})\cong\alpha_{j!}^1((\alpha_j^*\text{bc})_{\bullet}(\cdots),j^*\alpha_!^2\mathcal{F})
$$

(this is true for any isomorphism). We therefore have

$$
\alpha_{j_1}^1(\nu_j)_\bullet^1 \text{cart}_\bullet^1 \iota^* \mathcal{E}
$$
\n
$$
\cong \alpha_{j_1}^1(\alpha_j^* \text{bc})_\bullet^1(\nu_j)_\bullet^1 \text{cart}_\bullet^1 \iota^* \mathcal{E}
$$
\n
$$
\cong \alpha_{j_1}^1 \mathbb{D}(\mu)_\bullet^1 (\iota^* \nu)_\bullet^1 \iota^* \mathcal{E}
$$
\n
$$
\cong \alpha_{j_1}^1 \mathbb{D}(\mu)_\bullet^1 \iota^* \nu_\bullet^1 \mathcal{E}
$$

Thus we are left to show that

$$
\alpha_{j!}^1 \mathbb{D}(\mu)_\bullet^1 \iota^* \nu_\bullet^1 \mathcal{E} \to j^* \alpha_!^1 \nu_\bullet^1 \mathcal{E}
$$

is an isomorphism. A tedious check shows that this is the base change morphism associated with  $p_1$ . It is an isomorphism by (FDer4 left) for  $p_1$ .  $\Box$ 

## <span id="page-23-0"></span>1.5 (Co)Local morphisms

**1.5.1.** Let Dia be a diagram category and let S be a *strong* right derivator with domain Dia. Strongness implies that for each diagram



in S(.) there exists a homotopy pull-back " $U \times_T S$ " which is well-defined up to (non-unique!) isomorphism. A Grothendieck pre-topology on S is basically a Grothendieck pre-topology in the usual sense on  $\mathbb{S}(\cdot)$  except that pull-backs are replaced by homotopy pull-backs. We state the precise definition:

<span id="page-23-2"></span>**Definition 1.5.2.** A Grothendieck pre-topology on  $S$  is the datum consisting of, for any  $S \in \mathbb{S}(\cdot)$ , a collection of families  $\{U_i \to S\}_{i \in S}$  of morphisms in  $\mathbb{S}(\cdot)$  called **covers**, such that

- 1. Every family consisting of isomorphisms is a cover,
- 2. If  $\{U_i \to S\}_{i \in S}$  is a cover and  $T \to S$  is any morphism then the family  $\{U_i \times_S T'' \to T\}_{i \in I}$  is a cover for any choice of particular members of the family  $\{``U_i \times_S T"\}$ .
- 3. If  $\{U_i \to S\}_{i \in I}$  is a cover and for each i, the family  $\{U_{i,j} \to U_i\}_{j \in J_i}$  is a cover then the family of compositions  $\{U_{i,j} \to U_i \to S\}_{i \in I, j \in J_i}$  is a cover.

<span id="page-23-3"></span>**Definition 1.5.3** (left). Let  $p : \mathbb{D} \to \mathbb{S}$  be a left fibered derivator satisfying also (FDer0 right). Assume that pull-backs exist in S. We call a morphism  $f: U \to X$  in  $\mathbb{S}(\cdot)$  D-local if

(Dloc1 left) The morphism f satisfies **base change**: for any diagram  $Q \in \mathbb{D}(\square)$  with underlying diagram



such that  $p(Q)$  in  $\mathbb{S}(\square)$  is a pull-back-diagram, i.e. is (homotopy) Cartesian, the following holds true: If  $\widetilde{F}$  and  $\widetilde{f}$  are Cartesian, and  $\widetilde{g}$  is coCartesian then also  $\widetilde{G}$  is coCartesian.<sup>[6](#page-23-1)</sup>

(Dloc2 left) The morphism of derivators (cf. Lemma [1.3.14\)](#page-13-0)

$$
f^{\bullet}: \mathbb{D}_X \to \mathbb{D}_U
$$

commutes with homotopy colimits.

<span id="page-23-1"></span>6 In other words, if

$$
U \times_X Y'' \xrightarrow{F} Y
$$
  

$$
G \downarrow \qquad \qquad \downarrow g
$$
  

$$
U \xrightarrow{f} X
$$

is the underlying diagram of  $p(Q)$  then the exchange morphism

$$
G_{\bullet}F^{\bullet} \to f^{\bullet}g_{\bullet}
$$

is an isomorphism.

A morphism  $f: U \to X$  in  $\mathbb{S}(\cdot)$  is called **universally D-local** if any homotopy pull-back of f is D-local.

Definition 1.5.4 (left). Assume that S is equipped with a Grothendieck pre-topology (cf. [1.5.2\)](#page-23-2). A left fibered derivator  $p : \mathbb{D} \to \mathbb{S}$  as in Definition [1.5.3](#page-23-3) is called **local** w.r.t. the pre-topology on  $\mathbb{S}$ , if the following conditions hold:

- 1. Every morphism  $U_i \rightarrow S$  which is part of a cover is  $\mathbb{D}$ -local.
- 2. For a cover  $\{f_i: U_i \to S\}$  the family

$$
(f_i)^\bullet : \mathbb{D}(S) \to \mathbb{D}(U_i)
$$

is jointly conservative.

<span id="page-24-0"></span>**Definition 1.5.5** (right). Let  $p : \mathbb{D} \to \mathbb{S}$  be a right fibered derivator satisfying also (FDer0 left). Assume that push-outs exist in S. We call a morphism  $f: X \to U$  in  $\mathbb{S}(\cdot)$  D-colocal if

(Dloc1 right) The morphism f satisfies **base change**: for any diagram  $Q \in \mathbb{D}(\square)$  with underlying diagram:



such that  $p(Q)$  in  $\mathbb{S}(\Box)$  is a pushout-diagram, i.e. is (homotopy) coCartesian, if  $\widetilde{F}$  and  $\widetilde{f}$  are coCartesian, and  $\tilde{g}$  is Cartesian then also  $\tilde{G}$  is Cartesian.

(Dloc2 right) The morphism of derivators (cf. Lemma [1.3.13\)](#page-12-0)

$$
f_{\bullet}: \mathbb{D}_X \to \mathbb{D}_U
$$

commutes with homotopy limits.

A morphism  $f: X \to U$  in  $\mathbb{S}(\cdot)$  is called **universally D-colocal** if any homotopy push-out of f is D-colocal.

Definition 1.5.6 (right). Assume that S is equipped with a Grothendieck pre-cotopology, i.e. that  $\mathbb{S}^{\text{op}}$  is equipped with a Grothendieck pre-topology (cf. [1.5.2\)](#page-23-2). A right fibered derivator  $p : \mathbb{D} \to \mathbb{S}$  as in Definition [1.5.5](#page-24-0) is called **colocal** w.r.t. the pre-cotopology on  $\mathbb{S}$ , if

- 1. Every morphism  $S \to U_i$  which is part of a cocover is  $\mathbb{D}\text{-}colocal$ .
- 2. For a cocover  $\{f_i : S \to U_i\}$  the family

$$
(f_i)_\bullet : \mathbb{D}(\cdot)_S \to \mathbb{D}(\cdot)_{U_i}
$$

is jointly conservative.

#### <span id="page-25-1"></span>1.6 The associated pseudo-functor

Let  $p : \mathbb{D} \to \mathbb{S}$  be a morphism of pre-derivators with domain Dia.

**1.6.1** (left). Let Dia(S) be the 2-category of diagrams over S, where the objects are pairs  $(I, F)$ such that  $I \in \text{Dia}$  and  $F \in \mathcal{S}(I)$ , the morphisms  $(I, F) \to (J, G)$  are pairs  $(\alpha, f)$  such that  $\alpha : I \to I$  $J, f: F \to \alpha^* G$  and the 2-morphisms  $(\alpha, f) \to (\beta, g)$  are the natural transformations  $\mu: \alpha \to \beta$ satisfying  $\mathbb{S}(\mu)(G) \circ f = g$ .

We call a morphism  $(\alpha, f)$  of fixed shape if  $\alpha$  = id, and of diagram type if f consists of identities. Every morphism is obviously a composition of one of diagram type by one of fixed shape.

<span id="page-25-0"></span>**1.6.2** (right). There is a dual notion of a 2-category  $Dia^{op}(\mathbb{S})$ . Explicitly, the objects are pairs  $(I, F)$  such that  $I \in$  Dia and  $F \in \mathcal{S}(I)$ , the morphisms  $(I, F) \to (J, G)$  are pairs  $(\alpha, f)$  such that  $\alpha: I \to J, f: \alpha^* G \to F$  and the 2-morphisms  $(\alpha, f) \to (\beta, g)$  are the natural transformations  $\mu : \alpha \Rightarrow \beta$  satisfying  $f \circ \mathbb{S}(\mu)(G) = g$ .

The association  $(I, F) \mapsto (I^{\text{op}}, F^{\text{op}})$  induces an isomorphism  $\text{Dia}^{\text{op}}(\mathbb{S}) \to \text{Dia}(\mathbb{S}^{\text{op}})^{2-\text{op}}$ .

<span id="page-25-2"></span>We are interested in associating to a fibered derivator a pseudo-functor like for classical fibered categories.

**1.6.3** (left). We associate to a morphism of pre-derivators  $p : \mathbb{D} \to \mathbb{S}$  which satisfies (FDer0 right) a (contravariant) 2-pseudo-functor

$$
\mathbb{D}: \mathrm{Dia}(\mathbb{S})^{1-\mathrm{op}} \to \mathcal{CAT}
$$

mapping a pair  $(I, F)$  to  $\mathbb{D}(I)_F$ , and a morphism  $(\alpha, f) : (I, F) \to (J, G)$  to  $f^{\bullet} \circ \alpha^* : \mathbb{D}(J)_G \to \mathbb{D}(I)_F$ . A natural transformation  $\mu : \alpha \Rightarrow \beta$  is mapped to the natural transformation pasted from the following two 2-commutative triangles:



*Proof of the pseudo-functor property.* For a composition  $(\beta, g) \circ (\alpha, f) = (\beta \circ \alpha, \alpha^*(g) \circ f)$  we have:  $f^{\bullet} \circ \alpha^* \circ g^{\bullet} \circ \beta^* \cong f^{\bullet} \circ (\alpha^* g)^{\bullet} \circ \alpha^* \circ \beta^*$ . This follows from the isomorphism  $\alpha^* \circ g^{\bullet} \cong (\alpha^* g)^{\bullet} \circ \alpha^*$ (FDer0). One checks that this indeed yields a pseudo-functor.  $\Box$ 

**1.6.4** (right). We associate to a morphism of pre-derivators  $p : \mathbb{D} \to \mathbb{S}$  which satisfies (FDer0 left) a (contravariant) 2-pseudo-functor

$$
\mathbb{D}:\mathrm{Dia}^{\mathrm{op}}(\mathbb{S})^{1-\mathrm{op}}\to\mathcal{CAT}
$$

<span id="page-25-3"></span>mapping a pair  $(I, F)$  to  $\mathbb{D}(I)_{F(I)}$ , and a morphism  $(\alpha, f) : (I, F) \to (J, G)$  to  $f_{\bullet} \circ \alpha^*$  from  $\mathbb{D}(J)_G \to \mathbb{D}(I)_F$ . This defines a functor by the same reason as in [1.6.3.](#page-25-2)

1.6.5 (left). We assume that S is a strong right derivator. There is a notion of "comma object" in Dia(S) which we describe here for the case that S is the pre-derivator associated with a category  $S$ and leave it to the reader to formulate the derivator version. In that case the corresponding object will be determined up to (non-unique!) isomorphism only.

Given diagrams  $D_1 = (I_1, F_1), D_2 = (I_2, F_2), D_3 = (I_3, F_3)$  in Dia(S) and morphisms  $\beta_1 : D_1 \rightarrow D_3$ ,  $\beta_2: D_2 \to D_3$ , we form the comma diagram  $D_1 \times_{/D_3} D_2$  as follows: the underlying diagram  $I_1 \times_{/I_3} I_2$ has objects being tripels  $(i_1, i_2, \mu)$  such that  $i_1 \in I_1$ ,  $i_2 \in I_2$ , and  $\mu : \alpha_1(i_1) \to \alpha_2(i_2)$  in  $I_3$ . A morphism is a pair  $\beta_j : i_j \to i'_j$  for  $j = 1, 2$  such that

$$
\alpha_1(i_1) \xrightarrow{\alpha_1(\beta_1)} \alpha_1(i'_1)
$$
  
\n
$$
\downarrow^{\mu} \qquad \qquad \downarrow^{\mu'}
$$
  
\n
$$
\alpha_2(i_2) \xrightarrow{\alpha_2(\beta_2)} \alpha_2(i'_2)
$$

commutes in  $I_3$ . The corresponding functor  $\widetilde{F} \in \mathbb{S}(I_1 \times_{I_3} I_2)$  maps a tripel  $(i_1, i_2, \mu)$  to

$$
F_1(i_1) \times_{F_3(\alpha_2(i_2))} F_2(i_2).
$$

We define  $P_j$  to be  $(\iota_j, p_j)$  for  $j = 1, 2$ , where  $\iota_j$  maps a tripel  $(i_1, i_2, \mu)$  to  $i_j$ , and  $p_j$  is the corresponding projection of the fiber product. We then get a 2-commutative diagram

$$
D_1 \times_{/D_3} D_2 \xrightarrow{P_1} D_1
$$
  
\n
$$
\downarrow P_2 \qquad \downarrow^{\mu} \qquad \downarrow \downarrow^{\beta_1}
$$
  
\n
$$
D_2 \xrightarrow{\beta_2} D_3
$$

If we are given  $I_2, I_3$  only and two maps  $I_1 \rightarrow I_3$  and  $I_2 \rightarrow I_3$  we also form  $D_1 \times_{I_3} I_2$  by the same underlying category, with functor  $F_1 \circ \iota_1$ .

1.6.6 (right). We assume that S is a strong left derivator. There is a dual notion of "comma object" in Dia<sup>op</sup>(S) which we describe here again for the case that S is the pre-derivator associated with a category  $\mathcal S$  and leave it to the reader to formulate the derivator version. In that case the corresponding object will be determined up to (non-unique!) isomorphism only.

Given three diagrams  $D_1^o = (I_1, F_1), D_2^o = (I_2, F_2)$  in  $\text{Dia}^{\text{op}}(\mathbb{S})$  mapping to  $D_3^o = (I_3, F_3)$ , we form the comma diagram  $D_1^o \times_{D_3^o} D_2^o$  as follows: the underlying diagram is  $I_1 \times_{D_3} I_2$  which has object being tripels  $(i_1, i_2, \mu)$  such that  $i_1 \in I_1, i_2 \in I_2$  and  $\mu : \alpha_1(i_1) \to \alpha_2(i_2)$  in  $I_3$ . A morphism is a pair  $\beta_j : i_j \to i'_j$  for  $j = 1, 2$  such that

$$
\alpha_1(i_1) \xrightarrow{\alpha_1(\beta_1)} \alpha_1(i'_1)
$$
  
\n
$$
\downarrow^{\mu}
$$
  
\n
$$
\alpha_2(i_2) \xrightarrow{\alpha_2(\beta_2)} \alpha_2(i'_2)
$$

commutes in  $I_3$ . The corresponding functor  $\widetilde{F}$  maps a tripel  $(i_1, i_2, \mu)$  to

$$
F_1(i_1) \sqcup_{F_3(\alpha_1(i_1))} F_2(i_2).
$$

We then get a 2-commutative diagram

$$
D_2^o \times_{/D_3^o} D_1^o \longrightarrow D_1^o
$$
  
\n
$$
D_2^o \longrightarrow D_3^o
$$

<span id="page-27-0"></span>This language allows us to restate Lemma [1.3.22](#page-15-1) and Lemma [1.3.29](#page-20-0) in a more convenient way: **Lemma 1.6.7** (left). 1. Given a "pasting" diagram in  $Diag(S)$ 

$$
D_1 \xrightarrow{\Gamma} D_3 \xrightarrow{B} D_5
$$
  
\n
$$
A \downarrow^{\nu} \downarrow^{\alpha} \downarrow^{\mu} \downarrow^{\alpha}
$$
  
\n
$$
D_2 \xrightarrow{\gamma} D_4 \xrightarrow{\beta} D_6
$$

the pasted natural transformation  $\nu \odot \mu = \beta \nu \circ \mu \Gamma$  satsisfies

$$
\nu_!\odot\mu_!=(\nu\odot\mu)_!.
$$

2. Given a "pasting" diagram in Dia(S)

$$
D_1 \xrightarrow{B} D_2
$$
  
\n
$$
\Gamma \downarrow \qquad \qquad \nearrow
$$
  
\n
$$
D_3 \xrightarrow{b} D_4
$$
  
\n
$$
a \downarrow \qquad \nearrow
$$
  
\n
$$
B_5 \xrightarrow{\beta} D_6
$$

the pasted natural transformation  $\nu \odot \mu = \alpha \nu \circ \mu \Gamma$  satisfies

$$
\mu_!\odot\nu_! = (\mu\odot\nu)_!.
$$

**Definition 1.6.8.** If S is equipped with a Grothendieck pre-topology (cf. [1.5.2\)](#page-23-2) then we call  $(\alpha, f)$ :  $(I, F) \rightarrow (J, G)$  D-local if  $f_i : F(i) \rightarrow G \circ \alpha(i)$  is D-local (cf. [1.5.3\)](#page-23-3) for all  $i \in I$ . Likewise for the notions of universally  $\mathbb D$ -local,  $\mathbb D$ -colocal, and universally  $\mathbb D$ -colocal.

**Proposition 1.6.9** (left). Let  $\mathbb{D} \to \mathbb{S}$  be a left fibered derivator satisfying also (FDer0 right) and such that  $S$  is a strong right derivator. Then the associated pseudo-functor satisfies the following properties:

1. For a morphism of diagrams  $(\alpha, f): D_1 \rightarrow D_2$  the corresponding pull-back

$$
(\alpha, f)^* : \mathbb{D}(D_2) \to \mathbb{D}(D_1)
$$

has a left-adjoint  $(\alpha, f)$ .

2. For a diagram like in [1.6.5](#page-25-3)

$$
D_1 \times_{/D_3} D_2 \xrightarrow{P_1} D_1
$$
  
\n
$$
P_2 \downarrow \qquad \varphi^{\alpha} \downarrow \beta_1
$$
  
\n
$$
D_2 \xrightarrow{\beta_2} D_3
$$

the corresponding exchange morphism

$$
P_{2!}P_1^*\to \beta_2^*\beta_{1!}
$$

is an isomorphism in  $\mathbb{D}(D_2)$  provided that  $\beta_2$  is  $\mathbb{D}\text{-}local$ .

*Proof.* 1. By (FDer0 left) and (FDer3 left) we can form  $(\alpha, f)$ :  $= \alpha_1 \circ f$  which is clearly left adjoint to  $(\alpha, f)^*$ .

2. We first reduce to the case where  $I_2$  is the trivial category. Indeed consider the diagram

$$
D_1 \times_{/D_3} (\{j_2\}, F_2(j_2)) \xrightarrow{can.} D_1 \times_{/D_3} D_2 \times_{/D_2} (\{j_2\}, F_2(j_2)) \longrightarrow D_1 \times_{/D_3} D_2 \xrightarrow{P_1} D_1
$$
  
\n
$$
\downarrow \qquad \qquad \downarrow \qquad \qquad P_2 \downarrow \qquad \downarrow \qquad \qquad \downarrow
$$
  
\n
$$
(\{j_2\}, F_2(j_2)) \xrightarrow{can.} D_2 \xrightarrow{P_1} D_3
$$

The exchange morphism of the middle square and outmost rectangle are isomorphisms by the reduced case. The morphism can. of the left hand square is of diagram type and its underlying diagram functor has an adjoint. The exchange morphism is therefore an isomorphism by [\[8,](#page-86-1) 1.23]. Using Lemma [1.6.7](#page-27-0) therefore, applying this for all  $j_2 \in I_2$ , also the exchange morphism of the right square has to be an isomorphism (this uses axiom Der2).

Now we may assume  $D_2 = (\{j_2\}, F_2(j_2))$ . Consider the following diagram, in which we denote  $\beta_1 = (\alpha_1, f_1), \ \beta_2 = (\alpha_2, f_2).$ 

$$
(I_{1} \times_{/I_{3}} \{i_{2}\}, \widetilde{F}) \xrightarrow{p_{1}} (I_{1} \times_{/I_{3}} \{i_{2}\}, F_{1} \circ \iota_{1}) \xrightarrow{\iota_{1}} (I_{1}, F_{1})
$$
\n
$$
\downarrow^{p_{2}} \qquad \varnothing \qquad \qquad \downarrow^{i} f_{1} \qquad \varnothing \qquad \qquad \downarrow^{i} f_{1}
$$
\n
$$
(I_{1} \times_{/I_{3}} \{i_{2}\}, \widetilde{F}') \xrightarrow{p_{1}'} (I_{1} \times_{/I_{3}} \{i_{2}\}, F_{3} \circ \alpha_{1} \circ \iota_{1}) \xrightarrow{\iota_{1}} (I_{1}, F_{3} \circ \alpha_{1})
$$
\n
$$
\downarrow^{p_{2}'} \qquad \qquad \varnothing \qquad \qquad \downarrow^{r} f_{3}(\mu) \qquad \qquad \varnothing
$$
\n
$$
(I_{1} \times_{/I_{3}} \{i_{2}\}, F_{2}(i_{2})) \xrightarrow{\iota_{2}^{*} f_{2}} (I_{1} \times_{/I_{3}} \{i_{2}\}, F_{3}(\alpha_{2}(i_{2})))
$$
\n
$$
\downarrow^{i} \qquad \qquad \varnothing \qquad \qquad \downarrow^{i} g
$$
\n
$$
(\{i_{2}\}, F_{2}(i_{2})) \xrightarrow{f_{2}} (\{i_{2}\}, F_{3}(\alpha_{2}(i_{2}))) \xrightarrow{\alpha_{2}} (I_{3}, F_{3})
$$

where  $\widetilde{F}$  is the functor defined in [1.6.5](#page-25-3) mapping a tripel  $(i_1, i_2, \mu : \alpha_1(i_1) \to \alpha_2(i_2))$  to

$$
F_1(i_1) \times_{F_3(\alpha_2(i_2))} F_2(i_2)
$$

and  $\widetilde{F}'$  is the functor mapping a tripel  $(i_1, i_2, \mu : \alpha_1(i_1) \to \alpha_2(i_2))$  to

$$
F_3(\alpha_1(i_1)) \times_{F_3(\alpha_2(i_2))} F_2(i_2).
$$

We have to show that the exchange morphism for the outer square is an isomorphism. Using Lemma [1.6.7](#page-27-0) it suffices to show this for the squares 1–5. That the exchange morphism for the squares 1 and 2, where the morphisms are of fixed shape, is an isomorphism can be checked pointwise by (Der2). Then it boils down to the base change condition (Dloc1 left). Note that the squares are pull-back squares in S by construction of  $\widetilde{F}'$  resp  $\widetilde{F}$ . The exchange morphism for 4 is an isomorphism by (FDer0). The exchange morphism for 3 is an isomorphism because of (Dloc2 left). The exchange morphism for 5 is an isomorphism because of (FDer4 left). □

Dualizing, there is a right-variant of the theorem, which uses  $Dia^{op}(\mathcal{S})$  instead. We leave its formulation to the reader.

# <span id="page-29-1"></span><span id="page-29-0"></span>2 (Co)homological descent

## <span id="page-29-3"></span>2.1 Categories of S-diagrams

**Definition 2.1.1.** Let S be a strong right derivator with Grothendieck pre-topology. A category of S-diagrams in Cat(S) is a full sub-2-category  $DIA \subset Cat(S)$ , satisfying the following axioms:

- (SDia1) The empty diagram  $(\emptyset, -)$ , the diagrams  $(\cdot, S)$  for any  $S \in \mathbb{S}(\cdot)$ , and  $(\Delta_1, f)$  for any  $f \in \mathbb{S}(\Delta_1)$ are objects of DIA.
- (SDia2) DIA is stable under taking finite coproducts and such fibered products, where one of the morphisms is of pure diagram type.
- (SDia3) For each morphism  $\alpha : D_1 \to D_2$  with  $D_i = (I_i, F_i)$  in  $\mathcal{DIA}$  and for each object  $i \in I_2$  and morphism  $U \rightarrow F_2(i)$  being part of a cover in the chosen pre-topology, the slice diagram  $D_1 \times_{/D_2} (i, U)$  is in  $\mathcal{DIA}$ , and if  $\alpha$  is of pure diagram type then also  $(i, F_2(i)) \times_{/D_2} D_1$  is in DIA.

A category of S-diagrams  $\mathcal{DIA}$  is called infinite, if it satisfies in addition:

(SDia5) DIA is stable under taking arbitrary coproducts.

There is an obvious dual notion of a category of  $\mathcal S$ -diagrams in Cat<sup>op</sup>( $\mathcal S$ ). If  $\mathcal S$  is the trivial derivator both definitions boil down to the previous definition of a diagram category [1.1.1.](#page-5-3)

## <span id="page-29-2"></span>2.2 Fundamental (co)localizers

**Definition 2.2.1.** A class of morphisms  $W$  in a category is called weakly saturated, if it satisfies the following properties:

- (WS1) Identities are in W.
- (WS2) W has the 2-out-of-3 property.
- (WS3) If  $p: Y \to X$  and  $s: X \to Y$  are morphisms such that  $p \circ s = id_X$  and  $s \circ p \in W$  then  $p \in W$ (and hence  $s \in W$  by (WS2)).

Definition 2.2.2. Let S be a strong right derivator with Grothendieck pre-topology [\(1.5.2\)](#page-23-2). Let  $\mathcal{DIA} \subset \text{Cat}(\mathbb{S})$  be a category of S-diagrams (cf. [2.1.1\)](#page-29-3).

Consider a family of subclasses W<sub>S</sub> of morphisms in  $\mathcal{DIA} \times_{\mathcal{DIA}} (\cdot, S)$  parametrized by all objects  $S \in \mathbb{S}(\cdot)$ . Such a family  $\{ \mathcal{W}_S \}_S$  is called a **system of relative localizers** if the following properties are satisfied:

- (L0) For any morphism  $S_1 \rightarrow S_2$  the induced functor  $\mathcal{DIA} \times_{\mathcal{DIA}} (\cdot, S_1) \rightarrow \mathcal{DIA} \times_{\mathcal{DIA}} (\cdot, S_2)$  maps  $\mathcal{W}_{S_1}$  to  $\mathcal{W}_{S_2}$ .
- $(L1)$  Each  $W<sub>S</sub>$  is weakly saturated.
- (L2 left) If  $D = (I, F) \in \mathcal{DIA}$ , and I has a final object e, then the projection  $D \to (e, F(e))$  is in  $\mathcal{W}_{F(e)}$ .
- (L3 left) For any commutative diagram in  $\mathcal{DIA}$  over  $(\cdot, S)$



and for any chosen covers  ${U_{e,i} \rightarrow F(e)}$  for all  $e \in E$ , the following implication holds true:

$$
\forall e \in E \ \forall i \quad w \times_{/D_3} (e, U_{e,i}) \in \mathcal{W}_{U_{e,i}} \quad \Rightarrow \quad w \in \mathcal{W}_S.
$$

(L4 left) For any morphism  $w : D_1 \rightarrow D_2 = (E, F)$  of pure diagram type over  $(·, S)$  the following implication holds true:

$$
\forall e \in E \quad (e, F(e)) \times_{/D_2} D_1 \to (e, F(e)) \in \mathcal{W}_{(e, F(e))} \Rightarrow w \in \mathcal{W}_S.
$$

<span id="page-30-0"></span>There is an obvious dual notion of a **system of colocalizers** in  $DIA \subset \text{Cat}^{op}(\mathbb{S})$  where S is supposed to be a strong left derivator with Grothendieck pre-cotopology.

**Definition 2.2.3.** Let S be a strong right derivator. Assume we are given a Grothendieck pre-topology on S (cf. [1.5.2\)](#page-23-2). Let  $\mathcal{DIA} \subset \text{Cat}(\mathbb{S})$  be a category of S-diagrams (cf. [2.1.1\)](#page-29-3). A subclass W of morphisms in  $DIA$  is called an absolute localizer (or just localizer) if the following properties are satisfied:

 $(L1)$  W is weakly saturated.

(L2 left) If  $D = (I, F) \in \mathcal{DIA}$ , and I has a final object e, then the projection  $D \to (e, F(e))$  is in W.

(L3 left) For any commutative diagram in DIA



and chosen covering  ${U_{i,e} \to F_3(e)}$  for all  $e \in E$ , the following implication holds true:

 $\forall e \in E \ \forall i \quad w \times_{D_3} (e, U_i) \in W \Rightarrow w \in W.$ 

(L4 left) For any morphism  $w: D_1 \to D_2 = (E, F)$  of pure diagram type, the following implication holds true:

$$
\forall e \in E \quad (e, F(e)) \times_{/D_2} D_1 \to (e, F(e)) \in W \quad \Rightarrow \quad w \in W.
$$

There is an obvious dual notion of absolute **colocalizer** in  $DIA \subset \text{Cat}^{op}(\mathbb{S})$  where S is supposed to be a strong left derivator with Grothendieck pre-cotopology. Recall the identification

Cat(S) 
$$
\rightarrow
$$
 Cat<sup>op</sup>(S<sup>op</sup>)<sup>2-op</sup>  
(*I, F*)  $\rightarrow$  (*I*<sup>op</sup>, *F*<sup>op</sup>)

By abuse of notation, we denote the image of  $DIA$  under this identification by  $DIA^{op}$ . Note that if S is a strong right derivator with Grothendieck pre-topology, then S<sup>op</sup> is a strong left derivator with Grothendieck pre-cotopology.

- **Remark 2.2.4.** 1. If W is a localizer in DIA, then  $W^{op}$  is a colocalizer in DIA<sup>op</sup> and vice versa. The same holds true for systems of relative localizers.
	- 2. If  $S$  is the trivial derivator, then a system of relative localizers or a localizer are the same notion, and (L1–L3 left) are precisely the definition of fundamental localizer of Grothendieck.

**Proposition 2.2.5** (Grothendieck). If  $\mathbb{S} = \{\cdot\}$  is the trivial derivator, then Cat( $\cdot$ ) = Cat<sup>op</sup>( $\cdot$ ) as 2-categories. If  $\mathcal{DIA}$  is self-dual, i.e. if  $\mathcal{DIA}^{op} = \mathcal{DIA}$  under this identification, then the notions of localizer, localizer without (L4 left), colocalizer, and colocalizer without (L4 right) are all equivalent.

Proof. [\[5,](#page-86-6) Proposition 1.2.6]

Remark 2.2.6. The class of localizers is obviously closed under intersection, hence there is a smallest localizer  $W_{\mathcal{DIA}}^{\min}$ . Furthermore the smallest localizer in  $\mathcal{DIA}$  and the smallest colocalizer in  $\mathcal{DIA}^{\mathrm{op}}$  correspond. If S is the trivial derivator and  $\mathcal{DIA}$  = Cat, Cisinski [\[5](#page-86-6), Théorème 2.2.11] has shown that  $\mathcal{W}_{\text{Cat}}^{\text{min}}$  is precisely the class  $\mathcal{W}_{\infty}$  of functors  $\alpha : I \to J$  such that  $N(\alpha)$  is a weak equivalence in the classical sense (of simplicial sets, resp. topological spaces). For a localizer in the sense of Definition [2.2.3](#page-30-0) this implies the following:

**Theorem 2.2.7.** If  $DIA = \text{Cat}(\mathbb{S})$  and W is an absolute localizer in  $DIA$  and  $\alpha \in W_{\infty}$ , i.e.  $\alpha : I \to J$  is a functor such that  $N(\alpha)$  is a weak equivalence of topological spaces, the morphism  $(\alpha, id) : (I, p_I^*S) \to (J, p_J^*S)$  is in W for all  $S \in \mathbb{S}(\cdot)$ . The same holds analogously for a system of relative localizers.

*Proof.* The class of functors  $\alpha : I \to J$  in Cat such that  $(\alpha, id) : (I, p_I^*S) \to (J, p_J^*S)$  is in W obviously form a fundamental localizer in the classical sense.  $\Box$ 

<span id="page-31-0"></span>2.2.8. We will for (notational) simplicity assume that the following properties hold:

- 1. S has all relative finite coproducts (i.e. for each Grothendieck opfibration with finite discrete fibers  $p: O \to I$  the functor  $p^*$  has a left adjoint  $p_!$  and Kan's formula holds true w.r.t. it).
- 2. For all finite families  $(S_i)_{i\in I}$  of objects in S( $\cdot$ ) the collection  $\{S_i \to \coprod_{j\in I} S_j\}_{i\in I}$  is a cover.

 $\Box$ 

Let  $\varnothing$  be the initial object of  $\mathcal S$  (which exists by 1.). Then the map

$$
\varnothing \to (\cdot, \varnothing),
$$

where  $\emptyset$  on the left denotes the empty diagram, is in W (resp. in  $\mathcal{W}_{\emptyset}$ , and hence in all  $\mathcal{W}_{S}$ ) by (L3 left) applied to the empty cover.

From this and (L3 left) again it follows that for a finite collection  $(S_i)_{i\in I}$  of objects of  $\mathbb{S}(\cdot)$  the map

$$
(I,(S_i)_{i\in I})\to (\cdot,\coprod_{i\in I}S_i)
$$

is in  $W$  (resp. in  $\mathcal{W}_{\coprod_{i\in I}S_i}$ ). More generally, if we have a Grothendieck opfibration with finite *discrete* fibers  $p: O \to I$  and a diagram  $F \in \mathbb{S}(O)$  (over  $S \in \mathbb{S}(\cdot)$ ), then the morphism

$$
(O, F) \rightarrow (I, p_!F)
$$

is in  $W$  (resp. in  $W_S$ ).

Example 2.2.9 (Mayer-Vietoris). For the simplest non-trivial example of a non-constant map in W consider a cover  $\{U_1 \rightarrow S, U_2 \rightarrow S\}$  in  $\mathbb{S}(\cdot)$  consisting of two monomorphisms<sup>[7](#page-32-0)</sup>. Then the projection

$$
p: \left( \begin{array}{c} "U_1 \times_S U_2" \longrightarrow U_1 \\ \downarrow \\ U_2 \end{array} \right) \rightarrow S
$$

is in W (resp. in  $W_S$ ) as is easily derived from the axioms (L1–L4). See [2.5.14](#page-49-0) for how the Mayer-Vietoris long exact sequence is related to this.

<span id="page-32-1"></span>**2.2.10.** Let  $\alpha, \beta : D_1 \rightarrow D_2$  be two morphisms in  $DIA$ . Recall that it is the same to give a 2-morphism  $\alpha \Rightarrow \beta$  or a morphism  $D_1 \times \Delta_1 \rightarrow D_2$  such that for  $i = 1, 2$  the compositions  $D_1 \xrightarrow{e_i} D_1 \times \Delta_1 \longrightarrow D_2$  are  $\alpha$  and  $\beta$  respectively. We call  $\alpha$  and  $\beta$  **homotopic** if they are equivalent for the smallest equivalence relation containing by the following relation:  $\alpha \sim \beta$ , if there exists a 2-morphism  $\alpha \Rightarrow \beta$ . In other words  $\alpha$  and  $\beta$  are homotopic if there is a finite set of 1-morphisms  $\gamma_0, \gamma_1, \ldots, \gamma_n : D_1 \to D_2$  such that  $\gamma_0 = \alpha$  and  $\gamma_n = \beta$  and a chain of 2-morphisms:

$$
\gamma_0 \Leftarrow \gamma_1 \Rightarrow \gamma_2 \Leftarrow \dots \Rightarrow \gamma_n.
$$

<span id="page-32-2"></span>**Proposition 2.2.11.** Let  $DIA$  be a category of S-diagrams (cf. [2.1.1\)](#page-29-3) and let W be localizer in DIA (resp. let  $\{W_S\}_S$  be a system of relative localizers). Then W (resp.  $\{W_S\}_S$ ) satisfies the following properties:

- 1. The localizer W (resp. each  $W<sub>S</sub>$ ) is closed under coproducts.
- 2. Let  $\widetilde{s} = (s, id) : D_2 = (I_2, s^*F) \rightarrow D_1 = (I_1, F)$  be a morphism in  $\mathcal{DIA}$  (resp. over  $(\cdot, S)$ ) of pure diagram type such that s has a left adjoint  $p: I_1 \rightarrow I_2$ . Then the obvious morphisms  $\widetilde{p}: D_1 \to D_2$  and  $\widetilde{s}$  are in W (resp. in W<sub>S</sub>).

<span id="page-32-0"></span><sup>&</sup>lt;sup>7</sup>For an arbitrary S this means that the projections " $U_i \times_S U_i$ "  $\rightarrow U_i$  are isomorphisms.

3. Given a commutative diagram in  $\mathcal{DIA}$  (resp. one over  $(\cdot, S)$ )



where the underlying functors of the morphisms to  $D_3$  are Grothendieck opfibrations and the underlying functor of w is a morphism of opfibrations, and coverings  $\{U_{e,i} \rightarrow F_3(e)\}\$  for all  $e \in I_3$ , then (in the relative case)

$$
\forall e \in I_3 \ \forall i \quad w \times_{D_3} (e, U_{e,i}) \in \mathcal{W}_{U_{e,i}} \Rightarrow w \in \mathcal{W}_S
$$

or (in the absolute case)

$$
\forall e \in I_3 \ \forall i \quad w \times_{D_3} (e, U_{e,i}) \in \mathcal{W} \quad \Rightarrow \quad w \in \mathcal{W}
$$

- 4. If  $f: D_1 \to D_2$  is in W (resp. in W<sub>S</sub>) then also  $f \times E: D_1 \times E \to D_2 \times E$  is in W (resp. in  $W_S$ ) for any E  $\in$  Cat such that the morphism  $f \times E$  is a morphism in DIA.
- 5. Any morphism which is homotopic (in the sense of [2.2.10\)](#page-32-1) to a morphism in W (resp. in  $W_S$ ) is in W (resp. in  $W_S$ ).

Proof. 1. This property follows immediately from (L3 left) applied to a diagram



where  $I$  is considered to be a discrete category. (In the absolute case let  $S$  be the final object of  $\mathbb{S}(\cdot).$ 

2. We first show that  $\widetilde{p} \in \mathcal{W}$ . Using (L3 left), it suffices to show that  $\widetilde{p}_i : D_1 \times_{/I_2} i \to D_2 \times_{/I_2} i$  is in  $\mathcal{W}$ (resp. in  $W_S$ ) for all  $i \in I_2$ , however by the adjunction we have  $I_1 \times_{I_2} i = I_1 \times_{I_1} s(i)$  and therefore  $I_1 \times_{I_2} i$  has a final object. In the diagram

$$
D_1 \times_{/I_2} i \xrightarrow{\widetilde{p}_i} D_2 \times_{/I_2} i
$$
  
\n
$$
\downarrow \qquad \qquad \downarrow
$$
  
\n
$$
(\cdot, s(i)^* F) \longrightarrow (\cdot, s(i)^* F)
$$

the vertical morphisms are thus in W (resp.  $W_S$ ) and so is the upper horizontal morphism. That  $\tilde{s}$ is in W (resp. in  $W_S$ ) will follow from 4. because this implies that  $\tilde{s} \circ \tilde{p}$  and  $\tilde{p} \circ \tilde{s}$  are in W (resp. in  $W_S$ ) therefore by (L1) also  $\tilde{s}$  is in W (resp. in  $W_S$ ). For note that the unit and the counit extend to 2-morphisms of diagrams.

3. Using (L3 left), we have to show that  $D_1 \times_{/D_3}(e, U_{e,i}) \to D_2 \times_{/D_3}(e, U_{e,i})$  is in W (resp. in  $W_{U_{e,i}}$ ). Since the underlying functors of  $D_1 \rightarrow D_3$  and  $D_2 \rightarrow D_3$  are Grothendieck opfibrations, we have a diagram over  $(e, U_{e,i})$ :

$$
D_1 \times_{D_3} (e, U_{e,i}) \longrightarrow D_2 \times_{D_3} (e, U_{e,i})
$$
  
\n
$$
s_e \wedge \cdots \wedge s_e \wedge \cdots \wedge s_e \wedge \cdots \wedge s_e
$$
  
\n
$$
D_1 \times_{/D_3} (e, U_{e,i}) \longrightarrow D_2 \times_{/D_3} (e, U_{e,i})
$$

where the underlying functor of  $\iota_e$  is of diagram type and is right adjoint to  $s_e$ . Therefore  $s_e$  is in W (resp. in  $W_{U_{e,i}}$ ) by 2. and hence the same holds for  $\iota_e$  because  $s_e\iota_e = id$  (using L1). Note: we are not using the not yet proven part of 2. Since the top arrow is in W (resp. in  $W_{U_{e,i}}$ ) the same holds for the bottom arrow.

4. This is a special case of 2.

5. A natural transformation  $\mu : f \Rightarrow g$  for  $f, g : D_1 \rightarrow D_2$  can be seen as a morphism of diagrams  $\mu : \Delta_1 \times D_1 \to D_2$  such that  $\mu \circ e_0 = f$  and  $\mu \circ e_1 = g$ . Since the projection  $p : \Delta_1 \times D_1 \to D_1$  is in W by 3. also the morphisms  $e_{0,1}: D_1 \to \Delta_1 \times D_1$  are in W. Since  $\mu \circ e_0 = f$  and  $\mu \circ e_1 = g$ , the morphism f is in W if and only if  $g \in W$ . □

**Proposition 2.2.12.** Axiom (L4 left) is, in the presence of (L1–L3 left), equivalent to the following, apparently weaker axiom:

 $(L_4'$  left) Let  $w: D_1 \to D_2$  be a morphism (resp. a morphism over  $(\cdot, S)$ ) of pure diagram type such that the underlying functor is a Grothendieck fibration. Then (in the relative case)

 $\forall e \in I_2 \quad (e, F_2(e)) \times_{D_2} D_1 \rightarrow (e, F_2(e)) \in \mathcal{W}_{F_2(e)} \Rightarrow w \in \mathcal{W}_S$ 

or (in the absolute case)

$$
\forall e \in I_2 \quad (e, F_2(e)) \times_{D_2} D_1 \to (e, F_2(e)) \in \mathcal{W} \Rightarrow w \in \mathcal{W}.
$$

*Proof.* (L4' left) implies (L4 left): Consider the following 2-commutative diagram

$$
(E, F) \times_{/(E, F)} (I, p^* F) \longrightarrow (I, p^* F) = D_1
$$
\n
$$
(E, F) \longrightarrow (E, F) = D_2
$$

The underlying diagram functor of the top horizontal map (which is not purely of diagram type) is a Grothendieck opfibration and hence by Proposition [2.2.11,](#page-32-2) 3. it is in  $W$  (resp.  $W<sub>S</sub>$ ), provided that the morphisms of the fibers  $(E \times_{/E} e, pr_1^* F) \to (\cdot, F(e))$  are in W (resp. in  $W_{F(e)}$ ). However  $E \times_{/E} e$  has the final object id<sub>e</sub> whose value under pr<sup>\*</sup><sub>1</sub> F is  $F(e)$ . The morphisms of the fibers are therefore in W (resp. in  $\mathcal{W}_{F(e)}$ ) by (L2 left). The underlying diagram functor of the left vertical map is a Grothendieck fibration and  $\text{pr}_1^* F$  is constant along the fibers. Therefore the fact that all  $(e, F(e)) \times_{D_2} D_1 \to (e, F(e))$  are in W (resp. in  $W_{F(e)}$ ) implies that the left vertical map is in W (resp. in  $W_S$ ) by (L4' left). Thus also the right vertical map is in W (resp. in  $W_S$ ). (This uses Proposition [2.2.11,](#page-32-2) 5. and the fact that the two compositions in the diagram are homotopic). (L4 left) implies (L4' left): If  $D_1 \rightarrow D_2 = (E, F)$  is a morphism whose underlying functor is a

Grothendieck fibration as in Axiom (L4' left), the morphism of *constant* diagrams  $(e, F(e)) \times_{D_2} D_1 \rightarrow$  $(e, F(e)) \times_{D_2} D_1$  is in W (resp.  $W_{F(e)}$ ) (their underlying functors being part of an adjunction), therefore (L4 left) applies.  $\Box$ 

#### <span id="page-34-0"></span>2.3 Simplicial objects in a localizer

2.3.1. In this section, we fix a strong right derivator S equipped with a Grothendieck pre-topology and satisfying the assumptions of [2.2.8](#page-31-0) and a category of S-diagrams  $\mathcal{DIA}$  (cf. [2.1.1\)](#page-29-3). Assume that for all  $S_{\bullet} \in \mathbb{S}(\Delta^{\mathrm{op}})$  the diagrams  $(\Delta^{\mathrm{op}}, S_{\bullet})$  and also all truncations  $((\Delta^{\leq n})^{\mathrm{op}}, S_{\bullet})$  are in  $\mathcal{DIA}$ . Later we will assume that also  $((\Delta^{\circ})^{\text{op}}, S_{\bullet})$  for all  $S_{\bullet} \in \mathbb{S}((\Delta^{\circ})^{\text{op}})$  (injective simplex diagram) and

all truncations  $((\Delta^{\circ, \leq n})^{\text{op}}, S_{\bullet})$  are in  $\mathcal{DIA}$ . The reasoning in this section uses little of the explicit definition of  $\Delta^{op}$ . For comparison with classical texts on cohomological descent we stick to the particular diagram  $\Delta^{op}$ .

Consider the category  $\mathcal{S}(\Delta^{\mathcal{O}p})$ . Since S has all (relative) finite coproducts,  $\mathcal{S}(\cdot)$  is actually tensored over  $\mathcal{SETF}$ , hence  $\mathcal{S}(\Delta^{\rm op})$  will be tensored over  $\mathcal{SETF}^{\Delta^{\rm op}}$ . We sketch this construction. A finite simplicial set, i.e. a functor  $\xi : \Delta^{op} \to \mathcal{SETF}$ , can be seen as a functor with values in finite discrete categories. The corresponding Grothendieck construction yields a Grothendieck op-fibration  $\pi_{\epsilon}$ :  $\int \xi \to \Delta^{\rm op}$ . We define for  $X_{\bullet} \in \mathcal{S}(\Delta^{\rm op})$ :

$$
\xi \otimes X_{\bullet} := (\pi_{\xi})_{!} (\pi_{\xi})^{*} X_{\bullet}.
$$

Recall that the notion 'S has *relative finite coproducts*' means that all functors  $(\pi_{\epsilon})$  arising this way exist and can be computed fiber-wise.

**2.3.2.** Consider the full subcategory  $\Delta^{\leq n}$  of  $\Delta$  consisting of  $\Delta_0, \ldots, \Delta_n$ . Since S is assumed to be a right derivator, the restriction functor

$$
\iota^* : \mathbb{S}(\Delta^{\mathrm{op}}) \to \mathbb{S}((\Delta^{\leq n})^{\mathrm{op}})
$$

has a right adjoint  $\iota_*$ , which is usually called the **coskelet** and denoted  $\cosh^n$ .

Let some simplicial object  $Y_{\bullet} \in \mathbb{S}(\Delta^{\mathrm{op}})$  and a morphism  $\alpha : X_{\leq n} \to \iota^* Y_{\bullet}$  be given. Consider the full subcategory  $(\Delta^{op} \times \Delta_1)^{0-\leq n}$  of all objects  $\Delta_i \times [1]$  for all  $i \in \mathbb{N}_0$ , and  $\Delta_i \times [0]$  for  $i \leq n$ . The restriction

$$
\iota^* : \mathbb{S}(\Delta^{\mathrm{op}} \times \Delta_1) \to \mathbb{S}((\Delta^{\mathrm{op}} \times \Delta_1)^{0-\leq n})
$$

has again an adjoint  $\iota_*$ . Since S is assumed to be strong we can consider  $\alpha$  as an object over  $(\Delta \times \Delta_1)^{0-\leq n}$ . The first row of  $\iota_*\alpha$  is called the **relative coskelet** cosk<sup>n</sup> $(X_{\leq n}|Y_{\bullet})$  of  $X_{\leq n}$ . For  $n = -1$  we understand  $\cosh^{-1}(-|Y_{\bullet}) = Y_{\bullet}$ .

These constructions work the same way with  $\Delta$  replaced by  $\Delta^{\circ}$ . The functor 'coskelet' and 'relative coskelet' is in both cases even the same functor, i.e. these functors commute with the restriction of a simplicial to a semi-simplicial object<sup>[8](#page-35-0)</sup>. This would not at all be true for the corresponding left adjoint, the functor 'skelet'.

We call a diagram I in a diagram category Dia **contractible**, if  $I \rightarrow \cdot$  lies in every fundamental localizer on Dia.

**Lemma 2.3.3.** 1. Let  $\Delta$  be the simplex category, and I a category admitting a final object i. Let  $N(I)$  be the nerve of I. Then the category

$$
\int_{(\Delta^\circ)^{\rm op}} N(I)
$$

is contractible.

2. Let  $\Delta$  be the simplex category, and I a category admitting a final object i. Let  $N(I)$  be the nerve of I. Then the category

$$
\int_{\Delta^{\rm op}} N(I)
$$

is contractible.

<span id="page-35-0"></span> ${}^{8}$ To see this, e.g., for the case of the 'coskelet', observe that there is an adjunction:

$$
\Delta_m \times_{/(\Delta^{\circ})^{\text{op}}} (\Delta_{\leq n}^{\circ})^{\text{op}} \xrightarrow{\text{def}} \Delta_m \times_{/\Delta^{\text{op}}} \Delta_{\leq n}^{\text{op}}.
$$
3. Let  $\Delta^{\circ}$  be the injective simplex category and I a directed category admitting a final object i. Let  $N^{\circ}(I)$  be the semi-simplicial nerve of I, defined by letting  $N^{\circ}(I)_m$  be the set of functors  $[n] \rightarrow I$  such that no non-identity morphism is mapped to an identity. Then the category

$$
\int_{(\Delta^\circ)^{\rm op}} N^\circ(I)
$$

is contractible.

*Proof.* 1. is shown in [\[5](#page-86-0), Proposition 2.2.3]. 2. is the same but considering  $N(I)$  as a functor from  $(\Delta^{\circ})^{\text{op}}$  to  $\mathcal{S}\mathcal{E}\mathcal{T}$ . The same proof works when  $(\Delta^{\circ})^{\text{op}}$  is replaced by  $\Delta^{\text{op}}$ . 3. is also just a small modification of [loc. cit.]. Define a functor  $\xi : \int N^{\circ}(I) \to \int N^{\circ}(I)$  as follows: an object  $(n, x)$ , where  $x \in N^{\circ}(I)_n$  is mapped to  $(n, x)$  if  $x(n) = i$  and to  $(n + 1, x')$  with

$$
x'(k) = \begin{cases} x(k) & k \le n \\ i & k = n+1. \end{cases}
$$

otherwise. There are natural transformations

$$
\mathrm{id}_{\int N^{\circ}(I)} \Rightarrow \xi \qquad i \Rightarrow \xi
$$

where i denotes here the constant functor with value  $(0, i)$ , showing that  $\int N^{\circ}(I)$  is contractible.

Corollary 2.3.4. The diagrams  $\Delta$ ,  $\Delta^{\circ}$ ,  $\int_{\Delta^{op}} \Delta_n$ ,  $\int_{(\Delta^{\circ})^{op}} \Delta_n^{\circ}$ ,  $\int_{\Delta^{op}} \Delta_n \times \Delta_m$  and  $\Delta_m \times_{/\Delta^{op}} (\Delta^{\circ})^{op}$  =  $\int_{(\Delta^{\circ})^{\rm op}} \Delta_m$  are contractible.

*Proof.* The simplicial set  $\Delta_n$  is just the nerve N of [n]. Likewise the semi-simplicial set  $\Delta_n^{\circ}$  is the semi-simplicial nerve  $N^{\circ}$  of  $[n]$ . □

<span id="page-36-0"></span>Note that the diagram  $\int \Delta_n^{\circ}$  is even *finite*.

**Lemma 2.3.5.** Let W be a localizer (resp. let  $\{W_S\}_S$  be a system of relative localizers) in DIA. Let  $((\Delta^{\text{op}})^2, F_{\bullet,\bullet}) \in \mathcal{DIA}$  be a bisimplicial diagram (resp. a bisimplicial diagram over  $(\cdot, S)$ ) and let  $\delta : \Delta^{\mathrm{op}} \to (\Delta^{\mathrm{op}})^2$  be the diagonal. Then the morphism

$$
(\Delta^{\mathrm{op}}, \delta^* F_{\bullet, \bullet}) \to ((\Delta^{\mathrm{op}})^2, F_{\bullet, \bullet})
$$

is in W (resp.  $W_S$ ).

**Remark 2.3.6.** The statement of the Lemma is false when  $\Delta$  is replaced by  $\Delta^{\circ}$ .

Proof of Lemma [2.3.5.](#page-36-0) We focus on the absolute case. For the relative case the proof is identical. Since the morphism in the statement is of pure diagram type, we may check the condition of (L4 left): we have to show that the category

$$
(\Delta_m \times \Delta_n) \times ((\Delta^{\rm op})^2 \Delta^{\rm op})
$$

is contractible, say, on the diagram category of diagrams I such that  $(I, F_{m,n}) \in \mathcal{DIA}$ . Equivalently we may prove this for the dual category. Objects of that category are diagrams of the form:



This is the category  $\Delta/(\Delta_m \times \Delta_n)$  which is contractible by Lemma [2.3.3,](#page-35-0) 2. Note that this is the only feature of  $\Delta^{op}$  used in the proof of this Lemma.  $\Box$  Remark 2.3.7. The previous lemma should be seen in the following context: the Grothendieck construction gives a way of embedding the category of simplicial sets into the category of small categories. This construction maps weak equivalences to weak equivalences and induces an equivalence between the corresponding homotopy categories. A bisimplicial set can be seen as a simplicial object in the category of simplicial sets. Its homotopy colimit is given by the diagonal simplicial set. On the other hand the homotopy colimit in the category of small categories is just given by the Grothendieck construction. From this perspective, the lemma is clear if S is the derivator associated with the category of sets (equipped with the discrete topology).

<span id="page-37-0"></span>**Lemma 2.3.8.** Let W be a localizer (resp. let  $\{W_S\}_S$  be a system of relative localizers) in DIA. Consider a simplicial diagram  $(\Delta^{op}, F_{\bullet}) \in \mathcal{DIA}$  (resp. a simplicial diagram over  $(\cdot, S)$ ). The morphism

$$
(\Delta^{\mathrm{op}}, F_{\bullet} \otimes \Delta_n) \to (\Delta^{\mathrm{op}}, F_{\bullet})
$$

is in W (resp.  $W_S$ ).

Proof. We focus on the absolute case. For the relative case the proof is identical. The diagram  $(\Delta^{\rm op}, F_\bullet \otimes \Delta_n)$  is equivalent to  $(\int \Delta_n, \pi^* F_\bullet)$  by definition (see [2.3.1\)](#page-34-0). We apply the criterion of (L4 left) to the resulting map

$$
(\int \Delta_n, \pi^* F_{\bullet}) \to (\Delta^{\mathrm{op}}, F_{\bullet})
$$

and have to show that

$$
\Delta_m\times_{/\Delta^{\rm op}}\int\Delta_n
$$

is contractible. This category is again (dual to) the category of objects



<span id="page-37-1"></span>and we have already seen in the proof of Lemma [2.3.5](#page-36-0) that it is contractible.

**Corollary 2.3.9.** Let W be a localizer (resp. let  $\{W_S\}_S$  be a system of relative localizers) in DIA. Let  $f, g : (\Delta^{\mathrm{op}}, F_{\bullet}) \to (\Delta^{\mathrm{op}}, G_{\bullet})$  be two homotopic morphisms of simplicial objects (resp. morphisms over  $(\cdot, S)$ ). Then  $f \in \mathcal{W}$  (resp. in  $\mathcal{W}_S$ ) if and only if  $g \in \mathcal{W}$  (resp. in  $\mathcal{W}_S$ ).

*Proof.* The statement follows by the standard argument because the projection  $(\Delta^{\rm op}, F_{\bullet} \otimes \Delta_1) \rightarrow$  $(\Delta^{\rm op}, F_{\bullet})$  is in W (resp. in  $W_S$ ) by Lemma [2.3.8.](#page-37-0)  $\Box$ 

<span id="page-37-2"></span>**Proposition 2.3.10** (Cech resolutions are in W). Let W be a localizer (resp. let  $\{W_S\}_S$  be a system of relative localizers) in DIA.

Let  $U \rightarrow S$  be a local epimorphism in  $\mathbb{S}(\cdot)$ . Then the morphism

$$
p : (\Delta^{\rm op}, \cos \mathrm{k}^0(U|S)) \to (\cdot, S)
$$

is in W (resp.  $W_S$ ).

*Proof.* To simplify the exposition we focus on the case in which  $\mathbb S$  is associated with a category  $\mathcal S$ . The reader may check however that everything goes through in the general case because the only

 $\Box$ 

constructions involved can be expressed as right Kan extensions. The assumption means that there is a cover  $\mathcal{U} = \{U_i \rightarrow S\}$  in the given pre-topology, such that for all indices i, the induced map

$$
p_i: U \times_S U_i \to U_i
$$

has a section  $s_i$ . By axiom (L3 left) it suffices to show that for all i the map

$$
\widetilde{p}_i : (\Delta^{\mathrm{op}}, \mathrm{cosk}^0(U \times_S U_i | U_i)) \to (\cdot, U_i)
$$

is in W (resp. in  $W_{U_i}$ ). Explicitly the simplicial object  $\cos k^0(U \times_S U_i | U_i)$  is given by

$$
\cdots \stackrel{\overrightarrow{\longrightarrow}}{\longrightarrow} U \times_S U \times_S U \times_S U_i \stackrel{\longrightarrow}{\longrightarrow} U \times_S U \times_S U_i \stackrel{\longrightarrow}{\longrightarrow} U \times_S U_i
$$

Since  $\Delta^{op}$  is contractible (in particular the morphism  $(\Delta^{op}, p^*T) \to (\cdot, T)$  is in W, resp. in  $W_T$ , for any  $T \in \mathbb{S}(\cdot)$ , it suffices to show that the map

$$
\widetilde{p}_i : (\Delta^{\mathrm{op}}, \mathrm{cosk}^0(U \times_S U_i | U_i)) \to (\Delta^{\mathrm{op}}, p^* U_i)
$$

is in  $W$  (resp. in  $W_{U_i}$ ). There is a section

$$
\widetilde{s}_i : (\Delta^{\mathrm{op}}, p^*U_i) \to (\Delta^{\mathrm{op}}, \mathrm{cosk}^0(U \times_S U_i | U_i))
$$

induced by  $s_i$  such that  $\widetilde{p}_i \circ \widetilde{s}_i = id$ . By (L1) it then suffices to check that  $\widetilde{s}_i \circ \widetilde{p}_i \in \mathcal{W}$  (resp. in  $\mathcal{W}_{U_i}$ ). We will construct a homotopy between id and  $\widetilde{s}_i \circ \widetilde{p}_i$ 

$$
(\Delta^{\rm op}, \Delta_1 \times \cosh^0(U \times_S U_i | U_i)) \to (\Delta^{\rm op}, \cosh^0(U \times_S U_i | U_i))
$$

in the sense of simplicial objects. This will suffice by Corollary [2.3.9.](#page-37-1) Since  $\Delta_1$  is 1-coskeletal, and the  $\cosh^{0}$ 's anyway, it suffices to construct the homotopy in degrees 0 and 1:

$$
\text{Hom}(\Delta_1, \Delta_1) \times U \times_S U \times_S U_i \longrightarrow \text{Hom}(\Delta_0, \Delta_1) \times U \times_S U_i
$$
\n
$$
\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow
$$
\n
$$
U \times_S U \times_S U_i \longrightarrow U \times_S U_i
$$

This can be achieved by mapping  $\mathrm{id}_{\Delta_1} \times (U \times_S U_i) \times_{U_i} (U \times_S U_i)$  to  $(U \times_S U_i) \times_{U_i} (U \times_S U_i)$  via  $(s_i \circ pr_2) \times id.$  $\Gamma$ 

<span id="page-38-0"></span>**Definition 2.3.11.** A morphism  $X_{\bullet} \rightarrow Y_{\bullet}$  of simplicial objects is called a hypercover if the following two equivalent conditions hold:

1. In any diagram of simplicial objects



there is a cover  $\mathcal{U} = \{U_i \to U\}$  such that for all i there is a lift (indicated by a dotted arrow) in the diagram

$$
\partial \Delta_n \otimes U_i \longrightarrow \partial \Delta_n \otimes U \longrightarrow X_{\bullet}
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$
\n
$$
\Delta_n \otimes U_i \longrightarrow \Delta_n \otimes U \longrightarrow Y_{\bullet}
$$

2. For any  $n \geq 0$  the morphism

$$
X_n \to \operatorname{cosk}^{n-1}(\iota_{\leq n-1}^* X_{\bullet} | Y_{\bullet})_n
$$

admits local sections in the pre-topology on  $\mathcal S$  (i.e. it is a local epimorphism).

- **Remark 2.3.12.** 1. In particular the notion of hypercover depends only on the Grothendieck topology generated by the pre-topology because a morphism is a local epimorphism precisely if the sieve generated by it is a covering sieve.
	- 2. The equivalent condition 1. of the definition of hypercover shows that, if  $\mathbb{S}$  is the derivator associated with the category  $\mathcal{S}\mathcal{E}\mathcal{T}$  equipped with the discrete topology, then a hypercover is precisely a trivial Kan fibration.

**Definition 2.3.13.** If in condition 2. of Definition [2.3.11](#page-38-0) the morphism is even an isomorphism for all sufficiently large n, then  $\alpha$  is called a finite (or bounded) hypercover. Equivalently we have  $X_{\bullet} \cong \cosh^{n}(\iota_{\leq n}^{*} X_{\bullet} | Y_{\bullet})$  for some n.

<span id="page-39-0"></span>**Lemma 2.3.14.** Let W be a localizer (resp.  $\{W_S\}_S$  be a system of relative localizers) in DIA. For a finite hypercover  $X_{\bullet} \to Y_{\bullet}$  (resp. one over  $(\cdot, S)$ ) such that  $X_{\bullet} \cong \cosh^{i+1}(X_{\bullet} | Y_{\bullet})$  and  $\iota_{\leq i-1}^{*} X_{\bullet} \cong Y_{\bullet}$  $\iota_{\leq i-1}^* Y_\bullet$  the morphism  $(\Delta^{\mathrm{op}}, X_\bullet) \to (\Delta^{\mathrm{op}}, Y_\bullet)$  is in W (resp. in W<sub>S</sub>).

Proof. Again, to simplify the exposition we focus on the case in which S is associated with a category S. We may assume  $i \geq 1$  because otherwise we are in the situation of Lemma [2.3.10.](#page-37-2) The assumptions imply that the map  $X_i \to Y_i$  is a local epimorphism. Indeed, this is the map  $X_i \to Y_i = \cosh^{i-1}(\iota_{\leq i-1}^* X_{\bullet} | Y_{\bullet})_i$  in this case. Therefore the morphism  $X_j \to Y_j$  is actually a local epimorphism for all  $j$ .

Consider the following diagram in  $DIA$ :

$$
(\Delta^{\rm op} \times \Delta^{\rm op}, (X_{\bullet} \times_{Y_{\bullet}} X_{\bullet} | X_{\bullet})) \longrightarrow (\Delta^{\rm op} \times \Delta^{\rm op}, (X_{\bullet} | Y_{\bullet}))
$$
  

$$
(\Delta^{\rm op}, X_{\bullet}) \longrightarrow (\Delta^{\rm op}, Y_{\bullet})
$$

where

$$
(X_{\bullet} | Y_{\bullet})_{m,n} \coloneqq \operatorname{cosk}^{0} (X_{n} | Y_{n})_{m} = \underbrace{X_{n} \times_{Y_{n}} \cdots \times_{Y_{n}} X_{n}}_{m+1 \text{ factors}}.
$$

$$
(X_{\bullet} \times_{Y_{\bullet}} X_{\bullet} | X_{\bullet})_{m,n} \coloneqq \operatorname{cosk}^{0} (X_{n} \times_{Y_{n}} X_{n} | X_{n})_{m} = \underbrace{X_{n} \times_{Y_{n}} \cdots \times_{Y_{n}} X_{n}}_{m+2 \text{ factors}}.
$$

The vertical morphisms are in W by Proposition [2.2.11,](#page-32-0) 3. because its columns are in W by Lemma [2.3.10.](#page-37-2) Again by Proposition 2.2.[11,](#page-32-0) 3. it then suffices to show that the rows

$$
p: (\Delta^{\mathrm{op}}, (X_{\bullet} \times Y_{\bullet} | X_{\bullet})_{m, \bullet}) \longrightarrow (\Delta^{\mathrm{op}}, (X_{\bullet} | Y_{\bullet})_{m, \bullet})
$$

of the top horizontal morphism are in  $W$ . These are again hypercovers of the form considered in this Lemma, in particular *i*-coskeletal, where the *i*-truncation is given by

$$
\underbrace{X_i \times_{Y_i} \cdots \times_{Y_i} X_i}_{m+2} \xrightarrow{...} X_{i-1} = Y_{i-1} \cdots \qquad \cdots \Longrightarrow X_0 = Y_0
$$
\n
$$
\underbrace{X_i \times_{Y_i} \cdots \times_{Y_i} X_i}_{m+1} \xrightarrow{...} Y_{i-1} \cdots \qquad \cdots \Longrightarrow Y_0
$$

where the left-most vertical arrow is induced by the map  $\Delta_{m+1} \to \Delta_{m+2}$ ,  $i \to i$ . There is a section s, with  $s_i$  induced by the map

$$
\Delta_{m+2}\to\Delta_{m+1},\quad i\mapsto\begin{cases} i&i
$$

We will construct a homotopy  $\mu : id \Rightarrow s \circ p$  of truncated simplicial objects:

$$
\text{Hom}(\Delta_i, \Delta_1) \times \underbrace{X_i \times_{Y_i} \cdots \times_{Y_i} X_i}_{m+2} \xrightarrow{...} \text{Hom}(\Delta_{i-1}, \Delta_1) \times Y_{i-1} \cdots \qquad \cdots \xrightarrow{...} \text{Hom}(\Delta_0, \Delta_1) \times Y_0
$$
\n
$$
\downarrow^{\mu_i} \qquad \qquad \downarrow^{\mu_i} \qquad \qquad \downarrow^{\mu_i} \qquad \qquad \downarrow^{\mu_i} \qquad \qquad \downarrow^{\mu_0} \qquad \qquad \downarrow^{\mu_0} \qquad \qquad \downarrow^{\mu_0} \qquad \qquad \downarrow^{\mu_1} \qquad \qquad \downarrow^{\mu_0} \qquad \qquad \downarrow^{\mu_1} \qquad \qquad \downarrow^{\mu_2} \qquad \qquad \downarrow^{\mu_2} \qquad \qquad \downarrow^{\mu_3} \qquad \qquad \downarrow^{\mu_4} \qquad \qquad \downarrow^{\mu_5} \qquad \qquad \downarrow^{\mu_6} \qquad \qquad \downarrow^{\mu_7} \qquad \qquad \downarrow^{\mu_8} \qquad \qquad \downarrow^{\mu_9} \qquad \qquad \downarrow^{\mu_1} \qquad \qquad \downarrow^{\mu_1} \qquad \qquad \downarrow^{\mu_2} \qquad \qquad \downarrow^{\mu_6} \qquad \qquad \downarrow^{\mu_7} \qquad \qquad \downarrow^{\mu_8} \qquad \qquad \downarrow^{\mu_9} \qquad \qquad \downarrow^{\mu_9} \qquad \qquad \downarrow^{\mu_9} \qquad \qquad \downarrow^{\mu_1} \qquad \qquad \downarrow^{\mu_1} \qquad \qquad \downarrow^{\mu_2} \qquad \qquad \downarrow^{\mu_6} \qquad \qquad \downarrow^{\mu_7} \qquad \qquad \downarrow^{\mu_8} \qquad \qquad \downarrow^{\mu_9} \qquad \qquad
$$

The morphism  $\mu_i$  at the constant morphism  $0 : \Delta_i \to \Delta_1$  is given by the identity, at the constant morphism  $1: \Delta_i \to \Delta_1$  given by  $s_i \circ p_i$ , and at the other morphisms  $\Delta_i \to \Delta_1$  arbitrarily. The existence of this homotopy allows by Lemma [2.3.9](#page-37-1) and by (L1) to conclude.  $\Box$ 

**Theorem 2.3.15.** Let W be a localizer (resp.  $\{W_S\}_S$  be a system of relative localizers) in DIA. Any finite hypercover (resp. one over S) considered as a morphism of diagrams in  $DIA$ 

<span id="page-40-0"></span>
$$
(\Delta^{\text{op}}, X_{\bullet}) \to (\Delta^{\text{op}}, Y_{\bullet})
$$
\n(10)

is in W (resp. in  $W_S$ ). Let  $\iota : (\Delta^{\circ})^{\text{op}} \to \Delta^{\text{op}}$  be the inclusion. If the morphism [\(10\)](#page-40-0) exists in  $\mathcal{DIA}$  then also

$$
((\Delta^{\circ})^{\mathrm{op}}, \iota^* X_{\bullet}) \to ((\Delta^{\circ})^{\mathrm{op}}, \iota^* Y_{\bullet})
$$

is in W (resp. in  $W_S$ ).

Proof. Any finite hypercover is a finite succession of hypercovers of the form considered in Lemma [2.3.14.](#page-39-0) The additional statement is a consequence of the following Lemma.  $\Box$ 

**Lemma 2.3.16.** Let W be a localizer (resp. let  $\{W_S\}_S$  be a system of relative localizers) in DIA. Let  $\iota : (\Delta^{\circ})^{\text{op}} \to \Delta^{\text{op}}$  be the inclusion and let  $(\Delta^{\text{op}}, X_{\bullet})$  be a simplicial diagram in  $\mathcal{DIA}$  (resp. a simplicial diagram over  $(·, S)$ ). Then the morphism

$$
((\Delta^{\circ})^{\mathrm{op}}, \iota^* X_{\bullet}) \to (\Delta^{\mathrm{op}}, X_{\bullet})
$$

(if in  $\mathcal{DIA}$ ) is in W (resp. in  $W_S$ ).

Proof. We focus on the absolute case. For the relative case the proof is identical. Since the morphism in the statement is of pure diagram type, we may check the condition of (L4 left): we have to show that the category

$$
\Delta_m \times_{/\Delta^{\mathrm{op}}} (\Delta^{\circ})^{\mathrm{op}}
$$

is contractible, say, on the diagram category of diagrams I such that  $(I, X_m) \in \mathcal{DIA}$ . This is true by Lemma [2.3.3,](#page-35-0) 1.  $\Box$ 

#### 2.4 Cartesian and coCartesian objects

**Definition 2.4.1.** Let  $\mathbb{D} \to \mathbb{S}$  be a fibered derivator of domain Dia. Let  $I, E \in$  Dia be diagrams and let  $\alpha : I \to E$  be a functor in Dia. We say that an object

$$
X\in\mathbb{D}(I)
$$

is E-(co-)Cartesian, if for any morphism  $\mu : i \rightarrow j$  in I mapping to an identity in E, the corresponding morphism  $\mathbb{D}(\mu) : i^*X \to j^*X$  is (co-)Cartesian.

If E is the trivial category, we omit it from the notation, and talk about  $(co-)Cartesian$  objects.

These notions define full subcategories  $\mathbb{D}(I)^{E-\text{cart}}$  (resp.  $\mathbb{D}(I)^{E-\text{cocart}}$ ) of  $\mathbb{D}(I)$ , and  $\mathbb{D}(I)^{E-\text{cart}}$ (resp.  $\mathbb{D}(I)_F^{E\text{-cocart}}$ ) of  $\mathbb{D}(I)_F$  for any  $F \in \mathbb{S}(I)$ .

**Lemma 2.4.2.** The functor  $\alpha^*$  w.r.t. a morphism  $\alpha : D_1 \rightarrow D_2$  in Dia(S) maps Cartesian objects to Cartesian objects. The functor  $\alpha^*$  for a morphism  $\alpha : D_1 \to D_2$  in Dia<sup>op</sup>(S) maps coCartesian objects to coCartesian objects.

Remark 2.4.3. The categories of coCartesian objects are a generalization of the equivariant derived categories of Bernstein and Lunts [\[3](#page-86-1)]. For this let  $\mathbb{D} \to \mathbb{S}^{op}$  be the stable fibered derivator of sheaves of abelian groups on (nice) topological spaces, where S is the pre-derivator associated with the category of (nice) topological spaces. Let  $G$  be a topological group acting on a space  $X$ . Then we may form the following simplicial space which is an object of  $S(\Delta^{\mathrm{op}})$ :

$$
[G\backslash X]_{\bullet}: \cdots \xrightarrow{\bullet} G \times G \times X \xrightarrow{\bullet} G \times X \xrightarrow{\bullet} X,
$$

 $cf.$  [\[3](#page-86-1), B1]. Then the category

$$
\mathbb{D}(\Delta)^{\mathrm{cocart}}_{[G\setminus X]_\bullet}
$$

is equivalent to the (unbounded) equivariant derived category, cf.  $[3,$  Proposition  $B_4$ ]. Note that all pull-back functors are exact in this context.

<span id="page-41-0"></span>**Definition 2.4.4.** Let  $\mathbb{D} \to \mathbb{S}$  be a fibered derivator of domain Dia. We say that  $\mathbb{D} \to \mathbb{S}$  admits left Cartesian projections if for all functors  $\alpha : I \to E$  in Dia and  $S \in \mathbb{S}(\cdot)$ , the fully-faithful inclusion

$$
\mathbb{D}(I)_F^{E\text{-cart}} \to \mathbb{D}(I)_F
$$

has a left adjoint  $\Box^E_!$ . More generally we have four notions with the following notations:



We will, in general, only use left Cartesian and right coCartesian projection, the others being somewhat unnatural. In [3.3.1](#page-58-0) we will show (using Brown representability) that for an infinite fibered derivator whose fibers are stable and well-generated a right coCartesian projection exists. Similarly if, in addition, Brown representability for the dual holds, e.g. if the fibers are compactly generated, then a left Cartesian projection exists (see [3.3.2\)](#page-58-1) in many cases. Note that for a usual (non fibered) derivator, the notions 'Cartesian' and 'coCartesian' are equivalent. If for a fibered derivator with stable fibers both left and right Cartesian projections exist, then there is actually a recollement [\[16,](#page-86-2) Proposition 4.13.1]:

$$
\mathbb{D}(I)_{F}^{E-\text{cart}} \xrightarrow{\text{incl.}} \mathbb{D}(I)_{F} \xrightarrow{\text{incl.}} \mathbb{D}(I)_{F}/\mathbb{D}(I)_{F}^{E-\text{cart}}
$$

Example 2.4.5. The projections are difficult to describe explicitly, except in very special situations. Here a rather trivial example where this is possible. Let  $\mathbb D$  be a stable derivator and consider  $I = \Delta_1$ , the projection  $p : \Delta_1 \to \cdot$  and the inclusions  $e_0, e_1 : \cdot \to \Delta_1$ . Then a left and a right Cartesian projection exist and the recollement above is explicitly given by:

$$
\mathbb{D}(\Delta_1)^{\text{cart}} \cong \mathbb{D}(\cdot) \xrightarrow{p^*} \mathbb{D}(\Delta_1) \xrightarrow{c_1} \mathbb{D}(\cdot)
$$

Note that the functor C (Cone) may be described as either  $[1] \circ e_0^!$  or  $e_1^?$  (cf. [\[8,](#page-86-3) §3]) and that the essential image of  $p^*$  is precisely the kernel of C, which also coincides with the full subcategory of Cartesian=coCartesian objects.

#### 2.5 Weak and strong D-equivalences

**Definition 2.5.1** (left). Let Dia be a diagram category and let S be a strong right derivator with domain Dia equipped with a Grothendieck pre-topology. Let  $\mathbb{D} \to \mathbb{S}$  be a left fibered derivator satisfying (FDer0 right) and let  $S \in \mathbb{S}(\cdot)$ . A morphism  $f: D_1 \to D_2$  in  $\text{Dia}(\mathbb{S})/(\cdot, S)$  is called a weak **D-equivalence** relative to  $S$  if the natural transformation

$$
p_{1!}p_1^*\rightarrow p_{2!}p_2^*
$$

is an isomorphism of functors. A morphism  $f \in \text{Di}(S)$  is called a **strong** D-equivalence if the functor  $f^*$  induces an equivalence of categories

$$
f^* : \mathbb{D}(D_2)^{\text{cart}} \to \mathbb{D}(D_1)^{\text{cart}}.
$$

Note that weak is a relative notion whereas strong is absolute.

**Definition 2.5.2** (right). Let Dia be a diagram category, and let  $S$  be a strong left derivator with domain Dia equipped with a Grothendieck pre-cotopology. Let  $\mathbb{D} \to \mathbb{S}$  be a right fibered derivator satisfying (FDer0 left) and  $S \in \mathbb{S}(\cdot)$ . A morphism  $f: D_1 \to D_2$  in  $\text{Dia}^{\text{op}}(\mathbb{S})/(\cdot, S)$  is called a weak  $\mathbb{D}\text{-equivalence relative to } S, \text{ if the natural transformation}$ 

$$
p_{2*}p_2^* \rightarrow p_{1*}p_1^*
$$

is an isomorphism of functors. A morphism  $f \in \text{Dia}^{\text{op}}(\mathbb{S})$  is called a strong  $\mathbb{D}\text{-equivalence}$  if the functor  $f^*$  induces an equivalence of categories

$$
f^* : \mathbb{D}(D_2)^{\text{cocart}} \to \mathbb{D}(D_1)^{\text{cocart}}.
$$

For a (left and right) derivator, i.e. for  $\mathbb{S} = \cdot$ , there is no difference between  $\text{Dia}(\mathbb{S})$  and  $\text{Dia}^{\text{op}}(\mathbb{S})$  and then also the two different definitions of weak, resp. strong D-equivalence coincide (for the case of weak D-equivalences, note that the two conditions become adjoint to each other). These notions of  $\mathbb{D}\text{-}equivalence$  (right version) should be compared to the classical notions of *cohomological descent*, see  $[1, E$ xposé  $V^{\text{bis}}]$ .

**Lemma 2.5.3** (left). Let  $f : D_1 \rightarrow D_2$  be a morphism in Dia(S)/( $\cdot$ , S). Then the following implication holds:

f strong D-equivalence  $\Rightarrow$  f weak D-equivalence relative to S.

*Proof.* The morphism in the definition of weak  $\mathbb{D}$ -equivalence is induced by the counit w.r.t. the adjunction  $f^*, f_!$ :

$$
p_{1!}p_1^* \cong p_{2!}f_!f^*p_2^* \to p_{2!}p_2^*
$$

Now let  $f_{\text{in}}$  be an inverse to  $f^*$ , as required by the definition of strong D-equivalence. From  $f^*p_2^*\cong p_1^*$  follows  $p_{2!}f_{! \square} \cong p_{1!}$  and moreover the diagram



is commutative. Since the diagonal morphism is a natural isomorphism the statement follows.  $\Box$ 

<span id="page-43-1"></span>Of course there is an analogous right version of this lemma. The goal of this section is to prove the following two theorems:

**Main Theorem 2.5.4** (right). Let Dia be a diagram category and let  $\mathbb{S}$  be a strong left derivator with domain Dia equipped with a Grothendieck pre-cotopology.

- 1. Let  $\mathbb{D} \to \mathbb{S}$  be a fibered derivator with domain Dia which is colocal in the sense of Defini-tion [1.5.6](#page-24-0) for the Grothendieck pre-cotopology on S. Then the collection of classes  $\{W_{D,S}\}_{S}$ , where  $\mathcal{W}_{\mathbb{D},S}$  consists of those morphisms  $f : D_1 \to D_2$  in  $\text{Dia}^{\text{op}}(\mathbb{S})$  which are weak  $\mathbb{D}$ equivalences relative to  $S \in \mathbb{S}(\cdot)$ , forms a system of relative colocalizers.
- 2. Let  $\mathbb{D} \to \mathbb{S}$  be an infinite fibered derivator with domain Dia which is colocal in the sense of Definition [1.5.6](#page-24-0) for the Grothendieck pre-cotopology on S, with stable, compactly generated fibers. The class  $W_{\mathbb{D}}$  consisting of those morphisms  $f : D_1 \to D_2$  in Dia<sup>op</sup>(S) which are strong D-equivalences forms an absolute colocalizer.

<span id="page-43-0"></span>**Main Theorem 2.5.5** (left). Let Dia be a diagram category and let  $\mathbb{S}$  be a strong right derivator with domain Dia equipped with a Grothendieck pre-topology.

- 1. Let  $\mathbb{D} \to \mathbb{S}$  be a fibered derivator with domain Dia, which is local in the sense of Definition [1.5.4](#page-24-1) for the Grothendieck pre-topology on S. Then the collection of classes  $\{W_{\mathbb{D},S}\}_S$ , where  $W_{\mathbb{D},S}$ consists of those morphisms  $f: D_1 \to D_2$  in Dia(S) which are weak D-equivalences relative to  $S \in \mathbb{S}(\cdot)$ , forms a system of relative localizers.
- 2. Let  $\text{Dia}'(\mathbb{S})$   $\subset$   $\text{Dia}(\mathbb{S})$  be the full subcategory of the diagrams which consist of universally D-local morphisms.

Let  $\mathbb{D} \to \mathbb{S}$  be an infinite fibered derivator with domain Dia, which is local in the sense of Definition [1.5.4](#page-24-1) for the Grothendieck pre-topology on S, with stable, compactly generated fibers. The class  $W_{\mathbb{D}}$  consisting of those morphisms  $f: D_1 \rightarrow D_2$  in Dia'(S) which are strong  $\mathbb{D}$ equivalences *forms an absolute localizer in* Dia<sup>'</sup>(S).

**Remark 2.5.6.** The restriction onto  $\text{Dia}'(\mathbb{S})$  in the left-variant of the theorem is needed because otherwise we do not know whether left Cartesian projections exist (cf. Theorem [3.3.2\)](#page-58-1).

The weak D-equivalences for the case of usual derivators (i.e. for  $\mathbb{S} = \{\cdot\}$ ) were called just 'Dequivalences' by Cisinski [\[6\]](#page-86-5) and it is rather straight-forward to see from the definition of derivator that they from a fundamental localizer in the classical sense (= absolute localizer for  $\mathbb{S} = \{\cdot\},\$  $=$  system of relative localizers for  $\mathbb{S} = \{\cdot\}$ .

We will only prove the left-variant of the theorem. The other follows by logical duality and the restriction to Dia'(S) is not necessary because Lemma [2.5.11](#page-45-0) is used instead of Lemma [2.5.10.](#page-44-0) Before proving the theorem we need a couple of lemmas. We assume for the rest of this section that Dia is a diagram category and that S is a strong right derivator with domain Dia equipped with a Grothendieck pre-topology.

<span id="page-44-1"></span>**Definition 2.5.7.** Two morphisms (in Dia(S) or in Dia<sup>op</sup>(S))

$$
D_1 \xrightarrow[s]{p} D_2
$$

such that chains of 2-morphisms

 $p \circ s \Rightarrow \dots \Leftarrow \dots \Rightarrow \text{id}_{D_1} \qquad s \circ p \Rightarrow \dots \Leftarrow \dots \Rightarrow \text{id}_{D_2}$ 

<span id="page-44-2"></span>exist are called a **homotopy equivalence** (or p is called as such if an s with this property exists).

**Lemma 2.5.8** (left). Let  $\mathbb D$  be a left fibered derivator satisfying (FDer0 right) and let  $D_1, D_2 \in$ Dia(S). Given any homotopy equivalence  $(p, s)$ , then the functors  $p^*$  and  $s^*$  induce an equivalence

$$
\mathbb{D}(D_2)^{\text{cart}} \stackrel{p^*}{\underset{s^*}{\longrightarrow}} \mathbb{D}(D_1)^{\text{cart}}
$$

*Proof.* The 2-morphisms  $\mu : (\alpha, f) \Rightarrow (\beta, g)$  in Definition [2.5.7](#page-44-1) induce morphisms between the pull-back functors

$$
(\alpha, f)^{*}\mathcal{E} \to (\beta, g)^{*}\mathcal{E}
$$

which are isomorphisms on Cartesian objects.

**Example 2.5.9** (cf. also Proposition [2.2.11,](#page-32-0) 2.). Let  $I_1, I_2$  be diagrams in Dia. If

$$
I_1 \xrightarrow[s]{p} I_2
$$

is an adjunction where p is left adjoint to s, and if  $F \in \mathbb{S}(I_1)$  then we get an equivalence

$$
\mathbb{D}(D_2)^{\text{cart}} \stackrel{p^*}{\underset{s^*}{\rightleftharpoons}} \mathbb{D}(D_1)^{\text{cart}}
$$

<span id="page-44-0"></span>where  $D_1 = (I_1, F)$  and  $D_2 = (I_2, F \circ s)$ .

**Lemma 2.5.10** (left). Let Dia be an infinite diagram category and let  $\mathbb{D} \rightarrow \mathbb{S}$  be an infinite fibered derivator with domain Dia with stable, compactly generated fibers. Consider a morphism  $D = (I, F) \rightarrow (\cdot, S)$  such that F is a diagram of universally  $\mathbb{D}$ -local morphisms. Let  $U \rightarrow S$  be a universally D-local morphism. Write  $D_U = D \times_{(:,S)} (·, U)$  in Dia(S). Then the following diagram is 2-commutative (via the exchange natural isomorphism):



 $\Box$ 

Note that left Cartesian projectors exist for  $D$  and  $D_U$  by Theorem [3.3.2.](#page-58-1)

*Proof.* The functor  $pr_1^*$  has a right adjoint  $pr_{1*}$  by (Dloc2 left) and then Brown representability theorem. (Dloc1 left) says that  $\text{pr}_1^*$  preserves coCartesian morphisms, hence  $\text{pr}_{1*}$  preserves Cartesian morphisms. Therefore the right adjoint of the given diagram is the following commutative diagram:



Consequently the exchange morphism of the diagram in the statement is also a natural isomorphism. □

<span id="page-45-0"></span>**Lemma 2.5.11** (right). Let Dia be an infinite diagram category and let  $\mathbb{D} \to \mathbb{S}$  be an infinite fibered derivator with domain Dia with stable, compactly generated fibers. Consider a morphism  $D = (I, F) \rightarrow (\cdot, S)$ . Let  $S \rightarrow U$  be a universally D-colocal morphism. Write  $D_U = D \times_{(\cdot, S)} (\cdot, U)$  in Dia<sup>op</sup>(S). Then the following diagram is 2-commutative (via the exchange natural isomorphism):



Note that right Cartesian projectors exist for D and  $D_U$  by Theorem [3.3.1.](#page-58-0)

*Proof.* The functor  $pr_1^*$  has a left adjoint  $pr_{1!}$  by (Dloc2 right) and by the Brown representability theorem for the dual. (Dloc1 right) says that  $pr_1^*$  preserves Cartesian morphisms, hence  $pr_1$ ! preserves coCartesian morphisms. Therefore the right adjoint of the given diagram is the following commutative diagram:

$$
\mathbb{D}(D) \longleftarrow \mathbb{D}(D)^{\text{cocart}}
$$
\n
$$
\mathbb{P}_{1!}
$$
\n
$$
\mathbb{D}(D_U) \longleftarrow \mathbb{D}(D_U)^{\text{cocart}}
$$

Consequently the exchange morphism of the diagram in the statement is also a natural isomorphism.  $\Box$ 

<span id="page-45-1"></span>**Lemma 2.5.12** (left). Let  $\mathbb{D} \to \mathbb{S}$  be a fibered derivator with domain Dia admitting a left Cartesian projection (cf. [2.4.4\)](#page-41-0). For any Grothendieck opfibration

$$
\bigg\downarrow_{\pi}^{\pi}
$$

I

in Dia, for any diagram in  $F \in \mathcal{S}(I)$ , and for each element  $e \in E$ , the following diagram is 2commutative:



where  $\iota : I_e \to I$  is the inclusion of the fiber.

**Lemma 2.5.13** (right). Let  $\mathbb{D} \to \mathbb{S}$  be a fibered derivator with domain Dia admitting a right  $coCartesian$  projection (cf. [2.4.4\)](#page-41-0). For a Grothendieck fibration

> I π  $\overline{a}$ E

in Dia, for any diagram in  $F \in \mathbb{S}(I)$ , and for each element  $e \in E$ , the following diagram is 2commutative:

$$
\mathbb{D}(I)_{F} \xrightarrow{\Box_{\ast}^{E}} \mathbb{D}(I)_{F}^{E\text{-cocart}}
$$
\n
$$
\downarrow^* \qquad \qquad \downarrow^* \qquad \qquad \downarrow^* \qquad \qquad \mathbb{D}(I_e)_{F_e}^{\text{cocart}}
$$

where  $\iota: I_e \to I$  is the inclusion of the fiber.

Proof. We restrict to the right-variant, the other being dual. We will show that the functor  $\iota_!$ maps coCartesian objects to E-coCartesian ones. Then the left adjoint of the given diagram is the diagram

$$
\mathbb{D}(I)_F \leftarrow \mathbb{D}(I)_F^{E\text{-cocart}}
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$

which is commutative. Consequently also the diagram of the statement is 2-commutative via the natural exchange morphism.

Let f in E be an object and let  $\nu : i_1 \to i_2$  be a morphism in I mapping to id<sub>f</sub>. Let  $\alpha_k$  be the inclusions of  $\cdot$  into I with image  $i_k$ . The morphism  $\nu$  yields a natural transformation

$$
\nu:\alpha_1\Rightarrow \alpha_2.
$$

Consider the diagram

$$
e \times_{/E} f \xrightarrow{\phantom{+}c_k \phantom{+}} I_e \times_{/I} i_k \xrightarrow{\phantom{+}A_k \phantom{+}} I_e
$$
  
\n
$$
\downarrow p' \phantom{+} p_k \phantom{+} \swarrow p_k \phantom{+} \swarrow p_k
$$
  
\n
$$
\underbrace{\phantom{+}p' \phantom{+} p_k \phantom{+} p_k \phantom{+} \swarrow p_k}_{\phantom{+} \phantom{+} \phantom{+} \phantom{+} \phantom{+} p_k \phantom{+} \swarrow p_k}_{\phantom{+} \phantom{+} \phantom{+} \phantom{+} \phantom{+} \swarrow p_k}
$$

where  $c_k$  is given on a morphism  $\beta: e \to f$  in E by the choice of a Cartesian arrow  $i'_k \to i_k$ . It is right adjoint to  $\pi$  by the definition of Cartesian arrow.

There is a functor (composition with  $\nu$ ):

$$
\widetilde{\nu}: I_e \times_{/I} i_1 \to I_e \times_{/I} i_2
$$

such that  $A_2\tilde{\nu} = A_1$  and  $p_2\tilde{\nu} = p_1$ . We have therefore a natural (point-wise) coCartesian morphism  $\mathbb{S}(\mu_1)_\bullet \widetilde{\nu}^* \to \widetilde{\nu}^* \mathbb{S}(\mu_2)_\bullet$  of functors  $\mathbb{D}(I_e \times_{/I} i_2)_{A_2^* F_e} \to \mathbb{D}(I_e \times_{/I} i_1)$ .

We have also a natural transformation  $\rho : \tilde{\nu}c_1 \to c_2$  defined for a morphism  $\beta : e \to f$  in E as the unique arrow  $\rho(\beta)$  over id<sub>e</sub> making the following diagram commutative:

$$
\rho(\beta) \begin{cases} i'_1 \xrightarrow{c_1(\beta)} i_1 \\ \downarrow \downarrow \\ i'_2 \xrightarrow{c_2(\beta)} i_2 \end{cases}
$$

The resulting morphism  $\mathbb{D}(\rho) : c_1^* \widetilde{\nu}^* \to c_2^*$  is point-wise coCartesian on coCartesian objects. We get a commutative diagram of natural transformations

$$
\mathbb{S}(\mu_1)_\bullet A_1^* \longrightarrow \mathbb{S}(\mu_1)_\bullet A_1^* \iota^* \iota_! \xrightarrow{\mathbb{D}(\mu_1)'} p_1^* \alpha_1^* \iota_!
$$
\n
$$
\downarrow \sim \qquad \qquad \downarrow \sim \qquad \qquad \downarrow \sim
$$
\n
$$
\mathbb{S}(\mu_1)_\bullet \widetilde{\nu}^* A_2^* \longrightarrow \mathbb{S}(\mu_1)_\bullet \widetilde{\nu}^* A_2^* \iota^* \iota_! \xrightarrow{\mathcal{V}^* p_2^* \alpha_1^* \iota_!}
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$

where the first two top vertical morphisms are the natural isomorphisms induced by  $A_2\tilde{\nu} = A_1$ , the third top vertical morphism is the natural isomorphism induced by  $p_2\tilde{\nu} = p_1$ , and the first two lower vertical morphisms are point-wise coCartesian. Here we use the notation  $\mathbb{D}(\mu_1)'$  for the morphism  $\mathbb{S}(\mu_1)_\bullet X \to Y$  induced by a morphism  $\mathbb{D}(\mu_1) : X \to Y$ . Now we apply  $p_{1!}$  to the outer square:

$$
p_{1!} \mathbb{S}(\mu_1)_\bullet A_1^* \longrightarrow p_{1!} p_1^* \alpha_1^* \iota_1
$$
  
\n
$$
\downarrow \qquad \qquad \downarrow
$$
  
\n
$$
p_{1!} \widetilde{\nu}^* \mathbb{S}(\mu_2)_\bullet A_2^* \longrightarrow p_{1!} \widetilde{\nu}^* p_2^* \alpha_2^* \iota_1
$$

The left vertical map is still coCartesian (homotopy colimits preserve coCartesian morphisms). There is a canonical isomorphism  $p'_i c_i^* \rightarrow p_i$  [\[8](#page-86-3), Prop. 1.23] and the natural transformation  $\mathbb{D}(\rho): c_1^*\tilde{\nu}^* \to c_2^*$  is an isomorphism on coCartesian objects over constant diagrams. Consider the commutative diagram:

$$
p_1'c_1^*\widetilde{\nu}^* \longrightarrow p_{2!}\widetilde{\nu}_1c_{1!}c_1^*\widetilde{\nu}^* \longrightarrow p_{2!}
$$
\n
$$
\parallel \qquad \qquad \downarrow \mathbb{D}(\rho)^{ad}
$$
\n
$$
p_1'c_1^*\widetilde{\nu}^* \longrightarrow p_{2!}c_{2!}c_1^*\widetilde{\nu}^* \qquad \qquad \downarrow \mathbb{D}(\rho)
$$
\n
$$
p_{2!}c_{2!}c_2^* \longrightarrow p_{2!}
$$

where the rightmost horizontal morphisms are the respective counits. Since  $\mathbb{D}(\rho)$  is an isomorphism on coCartesian objects over constant diagrams, so is the morphism  $p'_!c_1^*\tilde{\nu}^* \to p_{2!}$ . Now we have the commutative diagram



which shows that also the natural map  $p_{1!}\tilde{\nu}^* \to p_{2!}$  is an isomorphism on coCartesian objects over constant diagrams.

We get a commutative diagram

$$
p_{1!} \mathbb{S}(\mu_1) \bullet A_1^* \longrightarrow p_{1!} p_1^* \alpha_1^* \iota_! \longrightarrow \alpha_1^* \iota_!
$$
  
\n
$$
p_{1!} \widetilde{\nu}^* \mathbb{S}(\mu_2) \bullet A_2^* \longrightarrow p_{1!} \widetilde{\nu}^* p_2^* \alpha_2^* \iota_!
$$
  
\n
$$
p_{2!} \mathbb{S}(\mu_2) \bullet A_2^* \longrightarrow p_{2!} p_2^* \alpha_2^* \iota_! \longrightarrow \alpha_2^* \iota_!
$$

where the composition of the left vertical arrows is coCartesian on coCartesian objects because the functor  $\mathbb{S}(\mu_2)_\bullet A_2^*$  maps coCartesian objects to coCartesian objects over constant diagrams. The composition of the horizontal morphisms in the top and bottom rows are isomorphisms by (FDer4 left). Hence the rightmost vertical map is coCartesian, too.  $\Box$ 

Proof of Main Theorem [2.5.5,](#page-43-0) 1. This is the case of weak D-equivalences.

(L0) and (L1) are clear.

For (L2 left), let  $D_1 = (I, F)$  and  $D_2 = (\{e\}, F(e))$ . The projection p and the inclusion i of the final object induce morphisms:

$$
D_1 \xrightarrow[i]{p} D_2
$$

We have  $p \circ i = id$  and there is a 2-morphism  $\beta : id \Rightarrow i \circ p$ . Therefore the statement is clear for weak D-equivalences over any base S. (L3 left): Let



be a morphism as in (L3 left) over a base  $S \in \mathbb{S}(\cdot)$ . We have to show that

$$
p_{1!} \, p_1^* \to p_{2!} \, p_2^*
$$

is an isomorphism and it suffices to show that the morphism

$$
p'_{1!}(p_1)^* \to p'_{2!}(p_2)^*
$$

is an isomorphism. This may be checked point-wise by (Der2) and after pull-back to an open cover by condition 2. of 'local' for a fibered derivator (see Definition [1.5.4\)](#page-24-1), so fix  $e \in E$  and consider the 2-commutative diagrams

$$
D_i \times_{/D_3} (e, U_j) \xrightarrow{\iota_{i,e,j}} D_i
$$
  
\n
$$
p'_{i,e,j} \downarrow \qquad p'_{i}
$$
  
\n
$$
(e, U_j) \xrightarrow{\epsilon_{e,j}} D_3
$$

and let  $p_{i,e}: D_i \times_{/D_3} (e, U_i) \to D_i$  be the projection. Applying the functor  $\epsilon_{e,j}^*$ , we get

 $\epsilon_{e,j}^* p'_{1!}(p_1)^* \to \epsilon_{e,j}^* p'_{2!}(p_2)^*$ 

which is, using Proposition [1.6.9](#page-27-0) (note that  $\iota_{e,j}$  is D-local by assumption), the same as

$$
(p'_{1,e,j})_! (i_{i,e,j})^*(p_1)^* \to (p'_{2,e,j})_! (i_{i,e,j})^*(p_2)^*.
$$

Now  $p_i \circ \iota_{i,e,j} = \pi_j \circ p'_{i,e,j}$ , where  $\pi : (\cdot, U_j) \to (\cdot, S)$  is the structural morphism. Therefore we get:

$$
(p'_{1,e,j})_! (p'_{1,e,j})^* \pi_j^* \to (p'_{2,e,j})_! (p'_{2,e,j})^* \pi_j^*.
$$

By Lemma [1.6.7](#page-27-1) this is induced by the canonical natural transformation which is an isomorphism by assumption.

(L4 left): By Lemma [2.2.12](#page-34-1) we may prove axiom (L4' left) instead. Consider a morphism  $p : D_1 \rightarrow$  $(E, F) = D_2$  in Dia(S) of pure diagram type, where the underlying functor of p is a Grothendieck fibration. It suffices to show that the counit

$$
p_! \, p^* \to \mathrm{id}
$$

is an isomorphism. This is the same as showing that the unit

$$
\mathrm{id} \to p_* \, p^*
$$

is an isomorphism. Note that  $p_*$  exists because this is a morphism of diagram type and  $\mathbb{D} \to \mathbb{S}$  is assumed to be a right fibered derivator, too (this is the only place, where this assumption is used for the case of weak D-equivalences). Now, since p is a Grothendieck fibration,  $p_*$  can be computed fiber-wise. So we have to show that

$$
id \to p_{e,*} p_e^*
$$

is an isomorphism or, equivalently, that

$$
p_{e,!} p_e^* \rightarrow \text{id}
$$

is an isomorphism. This holds true because by assumption the map of fibers  $I_e \to e$  is in  $\mathcal{W}_{F(e)}$ .

We proceed to state some consequences of the fact that *weak* D-equivalences form a fundamental localizer.

**Example 2.5.14** (Mayer-Vietoris). Let S be a strong right derivator (e.g. associated with a category with limits) with a Grothendieck pre-topology. We saw in Example [2.2.9](#page-32-1) that for a cover  $\{U_1 \rightarrow$  $S, U_2 \rightarrow S$  consisting of 2 monomorphisms, the projection

$$
p: \left( \begin{array}{c} \n\text{``}U_1 \times_S U_2 \n\text{''} \longrightarrow U_1 \\
\downarrow \\
U_2\n\end{array} \right) \rightarrow S
$$

belongs to any fundamental localizer. If  $\mathbb{D} \to \mathbb{S}$  is a fibered derivator which is local w.r.t. the pre-topology on S, Theorem [2.5.5](#page-43-0) implies therefore that p is a weak  $\mathbb{D}\text{-}equivalence$  in  $\text{Dia}(\mathbb{S})/(\cdot, S)$ , i.e. for  $A \in \mathbb{D}(\cdot)_{S}$  we have

$$
p_!p^*A\cong A,
$$

i.e. the homotopy colimit of

$$
i_{1,2,\bullet} i_{1,2}^{\bullet} A \longrightarrow i_{1,\bullet} i_1^{\bullet} A
$$
  
\n
$$
\downarrow
$$
  
\n
$$
i_{2,\bullet} i_2^{\bullet} A
$$

is isomorphic to A. If  $\mathbb D$  has stable fibers, this translates to the usual distinguished triangle

$$
i_{1,2,\bullet} \, i_{1,2}^\bullet A \to i_{1,\bullet} \, i_1^\bullet A \oplus i_{2,\bullet} \, i_2^\bullet A \to A \to i_{1,2,\bullet} \, i_{1,2}^\bullet A[1]
$$

in the language of triangulated categories.

Dually, if  $\mathbb{D} \to \mathbb{S}^{op}$  is a fibered derivator which is colocal w.r.t. the pre-cotopology on  $\mathbb{S}^{op}$ , The-orem [2.5.4](#page-43-1) implies that  $p^{op}$  is a weak D-equivalence in  $\text{Dia}^{op}(\mathbb{S}^{op})/(\cdot, S)$ , i.e. for  $A \in \mathbb{D}(\cdot)_S$  we have

$$
A \cong p_* p^* A.
$$

This means that the homotopy limit of

$$
i_1^{\bullet} i_{1,\bullet} A
$$
\n
$$
i_2^{\bullet} i_{2,\bullet} A \longrightarrow i_{1,2}^{\bullet} i_{1,2,\bullet} A
$$

is isomorphic to A. If  $\mathbb D$  has stable fibers, this translates to the usual distinguished triangle

$$
A \to i_1^{\bullet} i_1, A \oplus i_2^{\bullet} i_2, A \to i_{1,2}^{\bullet} i_{1,2, \bullet} A \to A[1]
$$

in the language of triangulated categories. Note that  $i_{\bullet}$  denotes a left adjoint push-forward along a morphism in  $\mathbb{S}^{op}$ , i.e. a left adjoint pull-back along a morphism in  $\mathbb{S}$ .

**Example 2.5.15** ((Co)homological descent). Let  $\mathbb S$  be a strong right derivator with a Grothendieck pre-topology and let  $X_{\bullet} \in \mathbb{S}(\Delta^{\mathrm{op}})$  be a simplicial diagram over  $S \in \mathbb{S}(\cdot)$  with underlying diagram

$$
\cdots \underbrace{\Longrightarrow} X_2 \underbrace{\Longrightarrow} X_1 \underbrace{\Longrightarrow} X_0
$$

such that  $(id, p) : (\Delta^{op}, X_{\bullet}) \to (\Delta^{op}, \pi^*S)$  is a finite hypercover. Here  $\pi : \Delta^{op} \to \cdot$  denotes the projection. If  $\mathbb{D} \to \mathbb{S}$  is a fibered derivator which is local w.r.t. the pre-topology on  $\mathbb{S}$ , Theorem [2.5.5](#page-43-0) implies that  $(\pi, p)$  is a weak D-equivalence in  $\mathbb{S}(\cdot)/(\cdot, S)$ , i.e. for  $A \in \mathbb{D}(\cdot)_S$  we have

$$
A\cong \pi_! \, p_\bullet \, p^\bullet \, \pi^* A.
$$

This means that the homotopy colimit of  $p_{\bullet} p^{\bullet} \pi^* A$  is equal to A. If the fibers of  $\mathbb{D} \to \mathbb{S}$  are in fact derived categories, this yields a spectral sequence of homological descent because the homotopy colimit over a simplicial complex is the total complex of the associated double complex (a well-known fact). This double complex looks like

$$
\cdots \longrightarrow p_{2,\bullet} p_2^{\bullet} A \longrightarrow p_{1,\bullet} p_1^{\bullet} A \longrightarrow p_{0,\bullet} p_0^{\bullet} A.
$$

The point is that we get a coherent double complex. Knowing the individual morphisms  $p_{i,\bullet} p_i^{\bullet} A \to p_i^{\bullet} A$  $p_{i-1,\bullet} p_{i-1}^{\bullet} A$  as morphisms in the derived category  $\mathbb{D}(\cdot)_S$  would not be sufficient! Dually (applying everything to a fibered derivator  $\mathbb{D} \to \mathbb{S}^{op}$ , and working in  $\text{Dia}^{op}(\mathbb{S}^{op})$ ), one obtains the more classical spectral sequence of cohomological descent.

Proof of Main Theorem [2.5.5,](#page-43-0) 2. This is the case of strong D-equivalences. (L1) is clear.

For (L2 left), let  $D_1 = (I, F)$  and  $D_2 = (e, F(e))$ . The projection p and the inclusion i of the final object induce morphisms:

$$
D_1 \xrightarrow[i]{p} D_2
$$

We have  $p \circ i = \text{id}$  and there is a 2-morphism  $\beta : \text{id} \Rightarrow i \circ p$ . Therefore the statement follows from Lemma [2.5.8.](#page-44-2) (Actually  $(i \circ p)^*$  is left adjoint to the inclusion  $\mathbb{D}(D_1)^{\text{cart}} \to \mathbb{D}(D_1)$ .) (L3 left): It suffices to prove the following two statements:

1. Consider a morphism of diagrams  $w = (\alpha, f) : D_1 = (I_1, F_1) \rightarrow D_2 = (I_2, F_2)$  such that we have a commutative diagram



and such that  $w \times_{E} e$  is a strong D-equivalence for all objects e in E. Then w is a strong D-equivalence.

2. Consider a morphism of diagrams  $w : D_1 = (I_1, F_1) \rightarrow D_2 = (I_2, F_2)$  over  $(·, S)$  and let  $\{U_i \rightarrow S\}$  be a covering. If  $w \times_{(\cdot,S)} (\cdot, U_i)$  is a strong D-equivalence for all i then w is a strong D-equivalence.

We proceed by showing statement 1. Consider the following diagram over  $E$ 

$$
D_1 \xrightarrow{w} D_2
$$
  
\n
$$
D_1 \times_{/E} E \xrightarrow{w'} D_2 \times_{/E} E
$$

where the vertical morphisms are of pure diagram type. We have an adjunction

$$
I_i \xrightarrow[\iota_i]{\kappa_i} I_i \times_{/E} E
$$

where  $\kappa_i$  maps an object i to  $(i, id_{p(i)})$ . We have a natural transformation  $\kappa_i \circ \iota_i \Rightarrow id_{I_i \times_{I_i} E}$ and moreover  $\iota_i \circ \kappa_i = id_E$  holds. Actually this defines an adjunction with  $\kappa_i$  left-adjoint to  $\iota_i$ . Furthermore, we get lifts to diagrams

$$
D_i \xleftarrow{\widetilde{\kappa}_i} (I_i \times_{/E} E, \iota_i \circ F) = D_1 \times_{/E} E,
$$

and a 2-morphism  $\widetilde{\kappa}_i \circ \widetilde{\iota}_i \Rightarrow \mathrm{id}_{D_1 \times_{/E} E}$ , and we have  $\widetilde{\iota}_i \circ \widetilde{\kappa}_i = \mathrm{id}_{D_1}$ .

Hence, by Lemma [2.5.8,](#page-44-2) the pull-backs along  $\tilde{t}_1$  and  $\tilde{t}_2$  induce equivalences on Cartesian objects, so we are reduced to showing that the pull-back along  $w'$  induces an equivalence on Cartesian objects.

The underlying diagrams  $I_k \times_{/E} E$  are Grothendieck opfibratons over E and the functor underlying w' is a map of Grothendieck opfibrations (the push-forward along a map  $\mu : e \to f$  in E being given by mapping  $(i, \nu : p(i) \to e)$  to  $(i, \nu \circ \mu)$ . Hence w.l.o.g. we may assume that  $I_1 \to E$  is a Grothendieck opfibration and the morphism  $I_1 \rightarrow I_2$  underlying f is a morphism of Grothendieck opfibrations.

We keep the notation  $w: D_1 \to D_2$ , however, and the assumption translates to the statement that the composition

$$
\mathbb{D}(D_{2,e})^{\text{cart}} \xrightarrow{w_e^*} \mathbb{D}(D_{1,e})^{\text{cart}}
$$

for the fibers is an equivalence with inverse  $\square_{!}w_{e,}$ . Consider the two functors:

$$
\mathbb{D}(D_2)^{E\text{-cart}} \xrightarrow{incl.} \mathbb{D}(D_2) \xrightarrow{w^*} \mathbb{D}(D_1).
$$

We first show that the counit

$$
\Box^E_! w_! w^* \mathcal{E} \to \mathcal{E}
$$

is an isomorphism for every E-Cartesian  $\mathcal{E}$ .

This can be checked after pulling back to the fibers. Let  $\iota_k : I_{k,e} \to I_k$  be the inclusion of the fiber over some  $e \in E$ .

We have the isomorphisms

$$
\iota_2^*\sqcup^E_! w_!w^*\mathcal{E}\cong \square_!w_{e,!}\iota_1^*w^*\mathcal{E}\cong \square_!w_{e,!}w^{e,*}\iota_2^*\mathcal{E}\cong \iota_2^*\mathcal{E},
$$

where we used the isomorphism  $\iota_2^* \Box^E \cong \Box_! \iota_2^*$  (Lemma [2.5.12\)](#page-45-1) and the isomorphism  $\iota_2^* w_! \cong w_{e,!} \iota_1^*$  (exists for morphisms of pure diagram type because we have a morphism of Grothendieck cofibrations, see Proposition [1.3.23,](#page-15-0) 3. and for morphisms of fixed shape by axiom (FDer0 left)). The morphism  $\Box_! w_{e,} w^{e, *}$   $\mathcal{E} \to \mathcal{E}$  is an isomorphism for Cartesian  $\mathcal{E}$  by assumption.

We now show that the unit

$$
\mathcal{E} \to w^* \sqcup^E_! w_! \mathcal{E}
$$

is an isomorphism for every E-Cartesian  $\mathcal E$ . This can be checked again on the fibers:

$$
\iota_1^* w^* \square^E_! w_! \mathcal{E} \cong w_e^* \iota_2^* \square^E_! w_! \mathcal{E} \cong w_e^* \square_! w_{e,!} \iota_1^* \mathcal{E} \cong \iota_1^* \mathcal{E}.
$$

Therefore we have already proven that the functors

$$
\mathbb{D}(D_2)^{E-\text{cart}} \xrightarrow{\quad w^* \quad \text{D}(D_1)^{E-\text{cart}}}
$$

form an equivalence.

We conclude by showing that  $\Box^E_\cdot w_\cdot$  maps Cartesian objects to Cartesian objects: Let  $\nu : e \to f$  be a morphisms of E. It induces a morphism (choice of push-forward for  $I_k \rightarrow E$ )

$$
\widetilde{\nu}_k: D_{k,e} \to D_{k,f}
$$

(not of diagram type!) and a 2-morphism:  $\iota_{k,e} \to \iota_{k,f} \circ \widetilde{\nu}_{k}$ .

• *Claim:* It suffices to show that for all  $\nu : e \rightarrow f$  the induced morphism

$$
\iota_{2,e}^* \square_{!}^E w_! \mathcal{E} \rightarrow \widetilde{\nu}_2^* \iota_{2,f}^* \square_{!}^E w_! \mathcal{E}
$$

is an isomorphism for every Cartesian  $\mathcal{E}$ .

Proof of the claim: Every morphism  $\mu : i \to i''$  in I with  $p(\mu) = \nu$ , say, is the composition of a coCartesian  $\mu'$  and some morphism  $\mu''$  with  $p(\mu'') = id_f$ . Since  $\mathcal E$  is E-Cartesian, the morphism  $\mathcal{E}(\mu'')$  is Cartesian. Hence to show that  $\mathcal{E}(\mu)$  is Cartesian it suffices to see that  $\mathcal{E}(\mu')$  is Cartesian. A reformulation is, however, that the morphism of the claim be an isomorphism. 口 Using the same argument as in the first part of the proof, we have to show that

$$
\Box_! w_{e,!} \iota_{1,e}^* \mathcal{E} \rightarrow \widetilde{\nu}_2^* \Box_! w_{f,!} \iota_{1,f}^* \mathcal{E}
$$

is an isomorphism for every Cartesian  $\mathcal{E}$ . Since both sides are Cartesian objects, this can be checked after applying  $w_e^*$  which is an equivalence on Cartesian objects:

$$
w_e^* \boxdot_! w_{e,!} \iota_{1,e}^* \mathcal{E} \to w_e^* \widetilde{\nu}_2^* \boxdot_! w_{f,!} \iota_{1,f}^* \mathcal{E}.
$$

We have  $w_e^* \widetilde{\nu}_2^* = \widetilde{\nu}_1^* w_f^*$  because the map of diagrams underlying w is a morphism of Grothendieck opfibrations. Hence, after applying  $w_e^*$ , we get

$$
w_{e^*} \boxdot_{!} w_{e,!} \iota_{1,e}^* \mathcal{E} \to \widetilde{\nu}_1^* w_{f^*} \boxdot_{!} w_{f,!} \iota_{1,f}^* \mathcal{E}.
$$

Since  $w_{e^*} \Box_! w_{e,!}$  and  $w_{f^*} \Box_! w_{f,!}$  are equivalences on Cartesian objects, we get

$$
\iota_{1,e}^* \mathcal{E} \to \widetilde{\nu}_1^* \iota_{1,f}^* \mathcal{E}.
$$

A slightly tedious check shows that this is again the morphism induced by the 2-morphism  $\iota_{1,e} \rightarrow$  $\iota_{1,f} \circ \tilde{\nu}_1$ . It is an isomorphism because  $\mathcal E$  is Cartesian.

We will now show statement 2. Consider a diagram



For any i (index of the cover in L4 left) we have the following commutative diagrams of objects in  $\mathrm{Dia}(\mathbb{S})$ :

$$
D_1 \times_S U_i \xrightarrow{w_i} D_2 \times_S U_i
$$
  
\n
$$
\begin{array}{c}\n\text{pr}_1^{(i)} \\
D_1 \xrightarrow{w} D_2\n\end{array}
$$

The morphisms  $pr_1^{(i)}$  are of fixed shape. We first show that the unit is an isomorphism

$$
\mathcal{E} \to w^* \sqcap_{!} w_! \, \mathcal{E}
$$

for any Cartesian  $\mathcal{E}$ . Note that by the stability axiom of a Grothendieck pre-topology also the collections  $(D_1 \times_S U_i)_j \to D_{1,j}$  are covers for any  $j \in I_1$ , where  $I_1$  is the underlying diagram of  $D_1$ . Since  $\mathbb D$  is local w.r.t. the Grothendieck pre-topology (and by axiom Der2), the family  $(\text{pr}_1^{(i)})$  $i^{(i)}_1$ <sup>\*</sup> is jointly conservative. Therefore it suffices to show that the unit is an isomorphism after applying  $(\mathrm{pr}_1^{(i)}$  $\binom{i}{1}^*$ . We get

$$
(\text{pr}_1^{(i)})^* \mathcal{E} \to (\text{pr}_1^{(i)})^* w^* \square_! w_! \mathcal{E}
$$
  

$$
(\text{pr}_1^{(i)})^* \mathcal{E} \to w_i^* (\text{pr}_1^{(i)})^* \square_! w_! \mathcal{E}.
$$

which is the same as

$$
f_{\rm{max}}
$$

Since  $(\text{pr}_1^{(i)}$  $\binom{i}{1}^*$  commutes with  $\Box_!$  (Lemma [2.5.10\)](#page-44-0) and with  $w_!$  (Proposition [1.6.9,](#page-27-0) 2.), we get

$$
(\text{pr}_1^{(i)})^* \mathcal{E} \to w_i^* \square_! w_{i,!} (\text{pr}_1^{(i)})^* \mathcal{E}.
$$

This morphism is an isomorphism by assumption. In the same way one shows that the counit is an isomorphism.

(L4 left): By Lemma [2.2.12](#page-34-1) we may prove axiom (L4' left) instead. We have shown during the proof for (L4' left) for the case of weak D-equivalences that

$$
p_! \, p^* \to \text{id}
$$

is an isomorphism, hence on Cartesian objects the same holds for the natural transformation

$$
\Box_! \, p_! \, p^* \to \mathrm{id} \, .
$$

We have to show that also the counit

<span id="page-54-0"></span>
$$
\text{id} \to p^* \square! p_! \tag{11}
$$

is an isomorphism on Cartesian objects. First note that  $p_*$  also is a right adjoint of  $p^*$  when restricted to the full subcategories of Cartesian objects because p<sup>∗</sup> preserves Cartesian objects. Indeed,  $p_{\star}$  can be computed fiber-wise because p is a Grothendieck fibration. The fibers being contractible in the sense of any localizer on Dia implies that the functors  $p_e^*, p_{e,*}$  induce an equivalence  $\mathbb{D}(D_e)^{\text{cart}} \cong \mathbb{D}(\cdot)_{F(e)}$ . Note: This uses that (L1–L3 left) hold for the class of strong D-equivalences on the fiber  $\mathbb{D}_{F(e)}$ , a fact which has been proven already. Therefore we pass to the right adjoints of the functors in [\(11\)](#page-54-0) and have to show that the counit

$$
p^* p_* \to \text{id}
$$

is an isomorphism on Cartesian objects. Again this can be checked fiber-wise, i.e. we have to show that the counit

$$
p_e^* p_{e,*} \to \text{id}
$$

is an isomorphism on Cartesian objects. But the pair of functors is an equivalence as we have seen, and the claim follows.  $\Box$ 

We proceed to state some consequences of the fact that *strong* D-equivalences form a fundamental localizer.

**Corollary 2.5.16** (left). Let S be a strong right derivator. If  $\mathbb{D} \rightarrow \mathbb{S}$  is an infinite fibered derivator which is local w.r.t. the pre-topology on  $S$  (cf. [1.5.2\)](#page-23-0) with stable, compactly generated fibers then for any finite hypercover  $f: X_{\bullet} \to Y_{\bullet}$  considered in Dia(S)' the functor  $f^*$  induces an equivalence

$$
\mathbb{D}(Y_{\bullet})^{\mathrm{cart}} \to \mathbb{D}(X_{\bullet})^{\mathrm{cart}}
$$

.

Here  $\text{Dia}(\mathbb{S})'$  is the full subcategory of diagrams with universally  $\mathbb{D}\text{-}local$  morphisms.

**Corollary 2.5.17** (right). Let S be a strong right derivator. If  $\mathbb{D} \to \mathbb{S}^{op}$  is an infinite fibered derivator which is colocal w.r.t. the pre-cotopology on  $\mathbb{S}^{op}$  (cf. [1.5.2\)](#page-23-0) with stable, compactly generated fibers then for any finite hypercover  $f: X_{\bullet} \to Y_{\bullet}$  considered in  $\text{Dia}^{\text{op}}(\mathbb{S}^{\text{op}})$  the functor  $f^*$  induces an equivalence

$$
\mathbb{D}(Y_{\bullet})^{\text{cocart}} \to \mathbb{D}(X_{\bullet})^{\text{cocart}}.
$$

Corollary 2.5.18. If  $D$  is an infinite derivator (not fibered) with domain Cat which is stable and well-generated, then for each homotopy type I, we get a category  $\mathbb{D}(I)^{\text{cart}}$  well-defined up to equivalence of categories. Moreover each morphism  $I \rightarrow J$  of homotopy types gives rise to a corresponding functor  $\alpha^*: \mathbb{D}(J)^{\text{cart}} \to \mathbb{D}(I)^{\text{cart}}$ . It is, however, not possible to arrange those as a pseudo functor  $HOT \to \mathcal{CAT}$ , but it is possible to arrange them as a pseudo-functor  $HOT^{(2)} \to \mathcal{CAT}$ where  $HOT^{(2)}$  is the homotopy 2-category (2-truncation) of any model for the homotopy theory of spaces (cf. also [A.2\)](#page-75-0).

# 3 Representability

In this section we exploit the consequences that Brown representability type results have for fibered derivators. In particular these results are useful to see that under certain circumstances a left fibered (multi-)derivator is already a right fibered (multi-)derivator, provided that its fibers are nice (i.e. stable and well-generated derivators). Furthermore they provide us with (co)Cartesian projectors that are needed for the strong form of (co)homological descent. In contrast to the rest of the article the results are stated in a rather unsymmetric form. This is due to the fact that in applications the stable derivators will rather be well-generated or compactly generated whereas their duals will rather not satisfy this condition. All the auxiliary results are taken from [\[16](#page-86-2)] and [\[19\]](#page-87-0).

#### <span id="page-55-0"></span>3.1 Well-generated triangulated categories and Brown representability

**Definition 3.1.1** (cf. [\[16,](#page-86-2) 5.1, 6.3]). Let  $\mathcal{D}$  be a category with zero object and small coproducts. We call D perfectly generated if there is a set of objects  $\mathcal T$  in D such that the following conditions hold:

- 1. An object  $X \in \mathbb{D}(\cdot)$  is zero if and only if  $\text{Hom}(T, X) = 0$  for all  $T \in \mathcal{T}$ .
- 2. If  $\{X_i \to Y_i\}$  is any set of maps, and  $\text{Hom}(T, X_i) \to \text{Hom}(T, Y_i)$  is surjective for all i, then  $Hom(T, \Pi_i, X_i) \rightarrow Hom(T, \Pi_i, Y_i)$  is also surjective.

The category  $\mathcal D$  is called well-generated if there is a set of objects  $\mathcal T$  in  $\mathcal D$  such that in addition to 1., 2. there is a regular cardinal  $\alpha$  such that the following condition holds:

3. All objects  $T \in \mathcal{T}$  are  $\alpha$ -small, cf. [\[16](#page-86-2), 6.3].

The category  $D$  is called **compactly generated** if there is a set of objects  $T$  in  $D$  such that in addition to 1., 2. the following two equivalent conditions hold:

- 4. All  $T \in \mathcal{T}$  are  $\aleph_0$ -small.
- 4'. All  $T \in \mathcal{T}$  are compact, i.e. for each morphism  $\gamma : T \to \coprod_{i \in I} X_i$  there is a finite subset  $J \subseteq I$ such that  $\gamma$  factors through  $\coprod_{i\in J} X_i$ .

**Definition 3.1.2.** A pre-derivator  $\mathbb{D}$  whose domain Dia is infinite (i.e. closed under infinite disjoint unions) is called **infinite** if the restriction-to- $I_j$  functors induce an equivalence

$$
\mathbb{D}(\coprod_{j\in J}I_j)\cong \prod_{j\in J}\mathbb{D}(I_j)
$$

for all sets J.

Recall (cf. [\[16,](#page-86-2) 4.4]) that a functor from a triangulated category  $\mathcal D$  to an abelian category is called cohomological if it sends distinguished triangles to exact sequences. We recall the following theorem:

<span id="page-56-0"></span>**Theorem 3.1.3** (right Brown representability). Let  $\mathcal{D}$  be a perfectly generated triangulated category with small coproducts. Then a functor  $F : \mathcal{D}^{op} \to \mathcal{AB}$  is cohomological and sends coproducts to products if and only if it is representable. An exact functor  $\mathcal{D} \to \mathcal{E}$  between triangulated categories commutes with coproducts if and only if it has a right adjoint.

Proof. See [\[16](#page-86-2), Theorem 5.1.1].

It can be shown that for a compactly generated triangulated category  $D$  with small coproducts,  $\mathcal{D}^{\text{op}}$  is perfectly generated and has small coproducts. Therefore the dual version of the previous theorem holds in this case:

<span id="page-56-1"></span>**Theorem 3.1.4** (left Brown representability). Let  $\mathcal{D}$  be a compactly generated triangulated category with small coproducts. Then a functor  $F : \mathcal{D} \to \mathcal{AB}$  is homological and sends products to products if and only if F is representable. An exact functor  $\mathcal{D} \to \mathcal{E}$  between triangulated categories commutes with products if and only if it has a left adjoint.

**Theorem 3.1.5.** Let  $D$  be a well-generated triangulated category with small coproducts. Consider a functor  $F: \mathcal{D} \to \mathcal{AB}$  which is cohomological and commutes with coproducts. Then there exists a right adjoint to the inclusion of the full subcategory of objects X such that  $F(X[n]) = 0$  for all  $n \in \mathbb{Z}$  (i.e. this subcategory is reflective).

<span id="page-56-2"></span>Proof. See [\[16](#page-86-2), Theorem 7.1.1].

**Lemma 3.1.6.** Let Dia be an infinite diagram category [\(1.1.1\)](#page-5-0). Let  $\mathbb{D} \to \mathbb{S}$  be an infinite left fibered derivator with domain Dia. If  $\mathbb{D}(\cdot)_{S}$  for all  $S \in \mathbb{S}(\cdot)$  is perfectly generated (resp. well-generated, resp. compactly generated), then the same holds for  $\mathbb{D}(I)_{S'}$  for all  $I \in$  Dia and for all  $S' \in \mathbb{S}(I)$ . Furthermore the categories  $\mathbb{D}(I)_{S'}$  all have small coproducts.

*Proof.* A set of generators as requested is given by the set  $\mathcal{T}_I := \{i_1T\}_{i\in I, T\in\mathcal{T}}$ . Indeed, an object  $X \in \mathbb{D}(I)$  is zero if  $i^*X$  is zero for all  $i \in X$  by (Der2). Therefore X is zero if  $\text{Hom}(i_!T, X)$  = Hom $(T, i^*X) = 0$  for all  $i \in I$  and for every  $T \in \mathcal{T}$ . We have to show that  $\text{Hom}(i_1T, \underline{\Pi}_i X_i) \to$ Hom $(i_!T, \underline{U}_i, Y_i)$  is an isomorphism for a family  $\{X_i \to Y_i\}_{i \in O}$  of morphisms as in Definition [3.1.1,](#page-55-0) 2. We have  $\text{Hom}(i_!T, \coprod_i X_i) = \text{Hom}(T, i^* \coprod_i X_i) = \text{Hom}(T, \coprod_i i^* X_i)$ , where we used that  $i^*$  commutes with coproducts. This follows because the Cartesian diagram



is homotopy exact. Note that, since D is infinite, coproducts exist and are equal to the corresponding homotopy coproducts. The map  $\text{Hom}(T, \coprod_i i^* X_i) \to \text{Hom}(T, \coprod_i i^* Y_i)$  is surjective by assumption. We have to show that a morphism

$$
i_!T \to \coprod_{i \in I} Y_i
$$

in  $\mathbb{D}(I)_{S'}$  factors trough  $\coprod_{i\in J} Y_i$  for some subset  $J \subset I$  of cardinality less than  $\alpha$ . By the same reasoning as above, we get a morphism

$$
T \to \coprod_{i \in I} i^* Y_i
$$

口

 $\Box$ 

Hence, there is some subset  $J \subset I$ , as required, such that this morphism factors through it. The same then holds for the original morphism. Since there is no need to enlarge  $J$ , the same statement holds for finite subsets.

 $\Box$ 

The categories  $\mathbb{D}(I)_{S'}$  have small coproducts because  $\mathbb{D} \to \mathbb{S}$  is infinite and left fibered.

**Definition 3.1.7.** Let  $\mathbb{D} \to \mathbb{S}$  be an infinite left fibered derivator with domain Dia. We will say that  $\mathbb{D} \to \mathbb{S}$  has perfectly-generated (resp. well-generated, resp. compactly-generated) fibers, if all categories  $\mathbb{D}(\cdot)_{S}$  are perfectly-generated (resp. well-generated, resp. compactly-generated) for all  $S \in \mathbb{S}(\cdot)$ . It follows from the previous Lemma that, in this case, for all  $I \in$  Dia and for all  $S' \in \mathbb{S}(I)$ the category  $\mathbb{D}(I)_{S'}$  is also perfectly-generated (resp. well-generated, resp. compactly-generated).

### <span id="page-57-0"></span>3.2 Left and Right

**Theorem 3.2.1** (left). Let Dia be an infinite diagram category (cf. [1.1.1\)](#page-5-0). Let D and E be infinite left derivators with domain Dia such that for all  $I \in$  Dia the pre-derivators  $\mathbb{D}_I$  and  $\mathbb{E}_I$  are stable (left and right) derivators with domain Post. Assume that  $\mathbb D$  is perfectly generated. Then a morphism of derivators  $F: \mathbb{D} \to \mathbb{E}$  commutes with all homotopy colimits w.r.t. Dia if and only if it has a right adjoint.

*Proof.* Let I be in Dia. Since  $\mathbb{D}_I$  and  $\mathbb{E}_I$  are stable,  $\mathbb{D}(I)$  is canonically triangulated, and we may use Theorem [3.1.3](#page-56-0) of right Brown representability. It follows that the functor  $F(I) : \mathbb{D}(I) \to \mathbb{E}(I)$ has a right adjoint  $G(I)$ , because it is triangulated, commutes with small coproducts and  $\mathbb{D}(I)$  is perfectly generated. To construct a morphism of derivators out of this collection, for any  $\alpha : I \to J$ , we have to give an isomorphism:  $G(J)\alpha^* \to \alpha^* G(I)$ . We may take the adjoint of the isomorphism  $\alpha_1 F(J) \to F(I) \alpha_1$  expressing that F commutes with all homotopy colimits (see [\[8,](#page-86-3) Lemma 2.1] for details). □

Analogously, using Theorem [3.1.4](#page-56-1) of left Brown representability, we obtain:

**Theorem 3.2.2** (right). Let Dia be an infinite diagram category (cf. [1.1.1\)](#page-5-0). Let  $\mathbb{D}$  and  $\mathbb{E}$  be infinite right derivators with domain Dia such that for all  $I \in$  Dia, the pre-derivators  $\mathbb{D}_I$  and  $\mathbb{E}_I$  are stable (left and right) derivators with domain Post. Assume that  $\mathbb D$  is compactly generated. Then a morphism of derivators  $F : \mathbb{D} \to \mathbb{E}$  commutes with all homotopy limits w.r.t. Dia if and only if it has a left adjoint.

**Theorem 3.2.3** (left). Let Dia be an infinite diagram category (cf. [1.1.1\)](#page-5-0). Let  $\mathbb{D} \to \mathbb{S}$  be an infinite left fibered (multi-)derivator with domain Dia whose fibers  $\mathbb{D}_S$  for every I  $\in$  Dia and all  $S \in \mathbb{S}(I)$  are stable (left and right) derivators with domain Posf. Assume that D has perfectly generated fibers. Then  $\mathbb D$  is a right fibered (multi-)derivator, too.

*Proof.* Let  $I \in$  Dia and let  $f \in$  Hom<sub>S(I)</sub>( $S_1, \ldots, S_n; T$ ) be a multimorphism. By Lemma [1.3.13,](#page-12-0) fixing  $\mathcal{E}_1, \hat{i}$ ,  $\mathcal{E}_n$ , the association

$$
\mathbb{D}(I \times J)_{p^*S_i} \rightarrow \mathbb{D}(I)_{p^*T}
$$
  

$$
\mathcal{E}_i \rightarrow (p^*f)_{\bullet}(p^*\mathcal{E}_1,\ldots,\mathcal{E}_i,\ldots,p^*\mathcal{E}_n)
$$

defines a morphism of derivators

$$
\mathbb{D}_{S_i} \to \mathbb{D}_T
$$

which is left continuous. Hence by Theorem [3.2.1](#page-57-0) it has a right adjoint. This shows the first part of (FDer0 right), i.e. the functor  $\mathbb{D}(I) \to \mathbb{S}(I)$  is an opfibration, too, for every  $I \in$  Dia. Then axiom (FDer5 left) implies the remaining assertion of (FDer0 right) while (FDer0 left) implies (FDer5 right), see Lemma [1.3.8.](#page-10-0)

Similarly a morphism  $\alpha : I \to J$  in Dia induces a morphism of derivators

$$
\alpha^*: \mathbb{D}_S \to \mathbb{D}_{\alpha^*S}.
$$

It commutes with homotopy colimits by Proposition [1.3.23,](#page-15-0) 2. Therefore  $\alpha^*$  has a right adjoint  $\alpha_{*}$  by the previous theorem, i.e. (FDer3 right) holds. (FDer4 right) is then a consequence of Lemma [1.3.23,](#page-15-0) 1.  $\Box$ 

Analogously, using Theorem [3.1.4](#page-56-1) of left Brown representability, we obtain:

**Theorem 3.2.4** (right). Let Dia be an infinite diagram category (cf. [1.1.1\)](#page-5-0). Let  $\mathbb{D} \to \mathbb{S}$  be an infinite right fibered (multi-)derivator with domain Dia, whose fibers  $\mathbb{D}_S$  for every I  $\epsilon$  Dia and for all  $S \in \mathcal{S}(I)$  are stable (left and right) derivators with domain Post. Assume that  $\mathbb{D}$  has compactly generated fibers. Then  $\mathbb D$  is a left fibered (multi-)derivator, too.

## <span id="page-58-0"></span>3.3 (Co)Cartesian projectors

**Theorem 3.3.1** (right). Let  $\mathbb{D} \to \mathbb{S}$  be an infinite fibered left derivator (w.r.t. Dia) whose fibers are stable derivators w.r.t. Posf. Assume that  $\mathbb{D}(\cdot)_S$  is well-generated for every  $S \in \mathbb{S}(\cdot)$ . Then for all I  $\epsilon$  Dia, for all  $F \in \mathcal{S}(I)$ , and for all functors  $I \to E$  in Dia the fully-faithful inclusion

$$
\mathbb{D}(I)_F^{E\text{-cocart}} \to \mathbb{D}(I)_F
$$

has a right adjoint  $\Box_*^E$ .

If  $\mathbb{D} \to \mathbb{S}$  also satisfies (FDer0 right) and if F is such that  $F(\mu)$  satisfies (Dloc2 left) for every  $\mu$ mapping to an identity in  $E$ , then the fully-faithful inclusion

$$
\mathbb{D}(I)_F^{E-\text{cart}} \to \mathbb{D}(I)_F
$$

has a right adjoint  $\blacksquare^E$ .

*Proof.* Consider the set O of morphisms  $\mu : i \rightarrow j$  which map to an identity in E, and for each morphism  $\mu \in O$  the composition  $D_{\mu}$ :

$$
\mathbb{D}(I)_F \xrightarrow{\mu^*} \mathbb{D}(\rightarrow)_{\mu^*F} \xrightarrow{F(\mu)_{\bullet}} \mathbb{D}(\rightarrow)_{i^*F} \xrightarrow{\text{Cone}} \mathbb{D}(\cdot)_{i^*F}
$$

We define a functor  $D$  by

$$
\prod_{\mu\in O}D_\mu:\mathbb{D}(I)_F\to\prod_{\mu\in O}\mathbb{D}(\cdot)_{i^*F}=\mathbb{D}(O)_{\iota^*F},
$$

where  $\iota: O \to I$  is the map "source". D commutes with coproducts, as all functors in the succession do, and it is exact. Therefore by [\[16](#page-86-2), Theorem 7.4.1] the triangulated subcategory  $\mathbb{D}(I)_F^{E\text{-cocart}}$  =  $ker D$  is well-generated and hence the inclusion in the statement has a right adjoint. In the Cartesian case,  $F(\mu)$ <sup>•</sup> commutes with coproducts only if  $F(\mu)$  satisfies (Dloc2 left).  $\Box$ 

<span id="page-58-1"></span>**Theorem 3.3.2** (left). Let  $\mathbb{D} \to \mathbb{S}$  be an infinite fibered derivator (with domain Dia) whose fibers are stable. Assume that all  $\mathbb{D}(\cdot)_{S}$  for  $S \in \mathbb{S}(\cdot)$  are compactly generated. Let  $I \to E$  be a functor in Dia and let  $F \in \mathcal{S}(I)$ . Suppose that  $F(\mu)$  satisfies (Dloc2 left) for every morphism  $\mu$  in I that maps to an identity in E. Then the fully-faithful inclusion

$$
\mathbb{D}(I)_F^{E\text{-cart}} \to \mathbb{D}(I)_F
$$

has a left adjoint  $\square^E_!$ .

Proof. As in the proof of the previous theorem we have an exact functor

$$
F^{\text{cart}}: \mathbb{D}(I)_F \to \mathcal{T}
$$

into another triangulated category which commutes with coproducts and such that the subcate-gory of E-Cartesian objects is precisely its kernel. Lemma [3.1.6](#page-56-2) implies that  $\mathbb{D}(I)$ <sub>F</sub> is compactly generated, and hence  $\mathbb{D}(I)_F^{\text{op}}$  $F_F^{\text{op}}$  is perfectly generated. Furthermore, Theorem [3.3.1](#page-58-0) implies that the categories  $\mathbb{D}(I)_F/\mathbb{D}(I)_F^{E\text{-carit}}$  are locally small. Note that

$$
\mathbb{D}(I)_F^{\mathrm{op}}/(\mathbb{D}(I)_F^{E\mathrm{-cart}})^{\mathrm{op}} = (\mathbb{D}(I)_F/\mathbb{D}(I)_F^{E\mathrm{-cart}})^{\mathrm{op}}.
$$

Therefore [\[16,](#page-86-2) Proposition 5.2.1] implies that a right adjoint to the inclusion  $(\mathbb{D}(I))^{\text{E-cart}}_F$ <sup>op</sup> →  $(\mathbb{D}(I)_F)^{\text{op}}$  exists. So a left adjoint to the inclusion

$$
\mathbb{D}(I)_F^{E\text{-cart}} \to \mathbb{D}(I)_F
$$

exists.

In the compactly generated case,  $\Box_!^E$  should exist unconditionally, but we were not able to prove this.

# 4 Constructions

## 4.1 The fibered multiderivator associated with a fibered multicategory

4.1.1. The most basic situation in which a (non-trivial) fibered (multi-)derivator can be constructed arises from a bifibration of (locally small) multicategories

$$
p:\mathcal{D}\to\mathcal{S}
$$

where we are given a set of weak equivalences  $W_S \subset \text{Mor}(\mathcal{D}_S)$  for each object S of S. In the examples we have in mind, the objects of  $S$  are spaces (or schemes), the objects of  $D$  are chain complexes of sheaves (coherent, etale Abelian, etc.) on them, and the morphisms in  $\mathcal{W}_S$  are the quasi-isomorphisms. In these examples the multicategory-structure arises from the tensor product and it is even, in most cases, the more natural structure because no particular tensor-product is chosen a priori.

<span id="page-59-1"></span>**Definition 4.1.2.** In the situation above, let  $S$  be the pre-multiderivator associated with the multicategory S. We define a pre-multiderivator  $\mathbb D$  as follows (cf. [A.3.1](#page-84-0) for localizations of multicategories):

$$
\mathbb{D}(I) = \mathrm{Hom}(I,\mathcal{D})[\mathcal{W}_I^{-1}]
$$

where  $W_I$  is the set of natural transformations which are element-wise in the union  $\bigcup_S W_S$ . The functor p obviously induces a morphism of pre-multiderivators

 $\widetilde{p} : \mathbb{D} \to \mathbb{S}$ 

Observe that morphisms in  $W_I$ , by definition, necessarily map to identities in Hom $(I, S)$ .

<span id="page-59-0"></span>In this section we prove that the above morphism of pre-(multi-)derivators is a left (resp. right) fibered (multi-)derivator on directed (resp. inverse) diagrams, provided that the fibers are model categories whose structure is compatible with the structure of bifibration. We use the definition of a model category from [\[15](#page-86-6)]. We denote the cofibrant replacement functor by Q and the fibrant replacement functor by R.

 $\Box$ 

Definition 4.1.3. A bifibration of (multi-)model-categories is a bifibration of (multi-)categories  $p: \mathcal{D} \to \mathcal{S}$  together with the collection of a closed model structure on the fiber

$$
(\mathcal{D}_S,\mathrm{Cof}_S,\mathrm{Fib}_S,\mathcal{W}_S)
$$

for any object  $S$  in  $S$  such that the following two properties hold:

1. For any  $n \geq 1$  and for every multimorphism



the push-forward  $f_{\bullet}$  and the various pull-backs  $f^{\bullet,j}$  define a Quillen adjunction in n-variables

$$
\Pi_i(\mathcal{D}_{S_i},\mathrm{Cof}_{S_i},\mathrm{Fib}_{S_i},\mathcal{W}_{S_i})\longrightarrow (\mathcal{D}_T,\mathrm{Cof}_T,\mathrm{Fib}_T,\mathcal{W}_T)
$$

$$
(\mathcal{D}_T, \mathrm{Cof}_T, \mathrm{Fib}_T, \mathcal{W}_T) \times \prod_{i \neq j} (\mathcal{D}_{S_i}, \mathrm{Cof}_{S_i}, \mathrm{Fib}_{S_i}, \mathcal{W}_{S_i}) \xrightarrow{f^{\bullet, j}} (\mathcal{D}_{S_j}, \mathrm{Cof}_{S_j}, \mathrm{Fib}_{S_j}, \mathcal{W}_{S_j})
$$

2. For any 0-ary morphism f in S, let  $f_{\bullet}$  () be the corresponding unit object (i.e. the object representing the 0-ary morphism functor  $\text{Hom}_{f}(\cdot,-)$  and consider the cofibrant replacement  $Qf_{\bullet}() \rightarrow f_{\bullet}()$ . Then the natural morphism

$$
F_{\bullet}(X_1,\ldots,X_{i-1},Qf_{\bullet}(),X_i,\ldots,X_n)\to F_{\bullet}(X_1,\ldots,X_{i-1},f_{\bullet}(),X_i,\ldots,X_n)\cong (F\circ_i f)_{\bullet}(X_1,\ldots,X_n)
$$

is a weak equivalence if all  $X_i$  are cofibrant. Here F is any morphism which is composable with f.

**Remark 4.1.4.** If  $S = \{\cdot\}$ , the above notion coincides with the notion of monoidal model-category in the sense of [\[15,](#page-86-6) Definition 4.2.6]. In this case it is enough to claim property 1. for  $n = 1, 2$ .

**Theorem 4.1.5.** Under the conditions of Definition [4.1.3](#page-59-0) the morphism of pre-derivators

 $\widetilde{p} : \mathbb{D} \to \mathbb{S}$ 

(defined in [4.1.2\)](#page-59-1) is a left fibered multiderivator (satisfying also FDer0 right) with domain Dir and a right fibered multiderivator (satisfying also FDer0 left) with domain Inv. Furthermore for all  $S \in \mathbb{S}(\cdot)$  its fiber  $\mathbb{D}_S$  (c.f. [1.3.11\)](#page-11-0) is just the pre-derivator associated with the pair  $(\mathcal{D}_S, \mathcal{W}_S)$ .

There are techniques by Cisinski [\[4](#page-86-7)] which allow to extend a derivator to more general diagram categories. We will explain in a forthcoming article how these can be applied to fibered (multi-)derivators.

<span id="page-60-0"></span>The proof of the theorem will occupy the rest of this section. First we have:

**Proposition 4.1.6.** Let  $D \rightarrow S$  be a bifibration of multicategories with complete fibers. For any  $diagram$  category  $I$ , the functors

$$
p_I: \mathrm{Hom}(I,\mathcal{D}) \to \mathrm{Hom}(I,\mathcal{S}) = \mathbb{S}(I)
$$

are bifibrations of multicategories.

Morphisms in  $\text{Hom}(I,\mathcal{D})$  are Cartesian, if and only if they are point-wise Cartesian. The 1-ary morphisms in  $\text{Hom}(I,\mathcal{D})$  are coCartesian, if and only if they are point-wise coCartesian.

*Proof (Sketch)*. We choose push-forward functors  $f_{\bullet}$  and pull-back functors  $f^{i,\bullet}$  for  $\mathcal{D} \to \mathcal{S}$  as usual. Let  $f \in \text{Hom}(S_1, \ldots, S_n; T)$  be a morphism in  $\text{Hom}(I, S)$ . We define a functor

$$
f_{\bullet}: \text{Hom}(I, \mathcal{D})_{S_1} \times \cdots \times \text{Hom}(I, \mathcal{D})_{S_n} \to \text{Hom}(I, \mathcal{D})_T
$$

by

$$
\mathcal{E}_1,\ldots,\mathcal{E}_n\mapsto\{i\mapsto (f_i)_\bullet(\mathcal{E}_1(i),\ldots,\mathcal{E}_n(i))\}.
$$

Note that a morphism  $\alpha : i \to i'$  in I induces a well-defined morphism

$$
(f_i) \bullet (\mathcal{E}_1(i), \ldots, \mathcal{E}_n(i)) \to (f_{i'}) \bullet (\mathcal{E}_1(i'), \ldots, \mathcal{E}_n(i'))
$$

lying over  $T(\alpha)$ . The functor  $f_{\bullet}$  comes equipped with a morphism in

$$
\operatorname{Hom}(\mathcal{E}_1,\ldots,\mathcal{E}_n;f_\bullet(\mathcal{E}_1,\ldots,\mathcal{E}_n))
$$

which is checked to be Cartesian in the strong form of Definition [A.2.5.](#page-77-0)

For 1-ary morphisms we can perform the same construction to produce coCartesian morphisms. For  $n \geq 2$  the construction is more complicated. Let  $f \in Hom(S_1, \ldots, S_n; T)$  be a morphism with  $n \geq 2$ . To ease notation, we construct a pull-back functor w.r.t. first slot. The other constructions are completely symmetric.

For any  $i_1 \in I$  consider the category (a variant of the twisted arrow category)

$$
X_{i_1}(I) \coloneqq \{ (i_2, \ldots, i_n, j, \{\alpha_k\}_{k=1..n}) \mid \alpha_k : i_k \to j \}
$$

which is covariant in j and contravariant in  $i_2, \ldots, i_n$ . For any  $\beta : i_1 \to i'_1$  we have an induced functor  $\tilde{\beta}: X_{i_1}(I) \to X_{i'_1}(I)$ . Any object in  $X_{i_1}(I)$  defines by pre-composition with  $S_k(\alpha_k)$  for all  $1 \leq k \leq n$  a morphism  $f_{\alpha} \in \text{Hom}(S_1(i_1), \ldots, S_n(i_n); T(j)).$ We define a functor

$$
f^{1,\bullet}: (\text{Hom}(I,\mathcal{D})_{S_2})^{\text{op}} \times \cdots \times (\text{Hom}(I,\mathcal{D})_{S_n})^{\text{op}} \times \text{Hom}(I,\mathcal{D})_T \to \text{Hom}(I,\mathcal{D})_{S_1}
$$

assigning to  $\mathcal{E}_2, \ldots, \mathcal{E}_n; \mathcal{F}$  the following functor  $X_{i_1}(I) \to \mathcal{D}_{S_1(i_1)}$ :

$$
\alpha \mapsto (f_{\alpha})^{1,\bullet}(\mathcal{E}_2(i_2),\ldots,\mathcal{E}_n(i_n);\mathcal{F}(j))
$$

and then taking  $\lim_{X_{i_1}(I)}$  which exists because the fibers are required to be complete. For the functoriality note that for  $\beta : i_1 \to i'_1$  we have a natural morphism

$$
\lim_{X_{i_1}(I)} \cdots \to \lim_{X_{i'_1}(I)} \cdots
$$

induced by  $\widetilde{\beta}$ .

We define a morphism

$$
\Xi \in \mathrm{Hom}_f(f^{1,\bullet}(\mathcal{E}_2,\ldots,\mathcal{E}_n;\mathcal{F}),\mathcal{E}_2,\ldots,\mathcal{E}_n;\mathcal{F})
$$

and we will show that it is coCartesian w.r.t. the first slot in a weak sense. At some object  $i \in I$ , the morphism  $\Xi$  is given by composing the projections from  $\lim_{X_{i=i_1}(I)} f_{\alpha}^{1,\bullet}(\mathcal{E}_2(i_2),\ldots,\mathcal{E}_n(i_n); \mathcal{F}(j))$ to  $f_i^{1,\bullet}$  $i^{1,\bullet}(\mathcal{E}_2(i),\ldots,\mathcal{E}_n(i); \mathcal{F}(i))$  (note that  $f_i = f_\alpha$  for  $\alpha = {\text{id}_i}_k$ ) and then composing with the  $coCartesian$  morphism (in  $\mathcal{D}$ ) in

Hom
$$
(f_i^{1,\bullet}(\mathcal{E}_2(i),\ldots,\mathcal{E}_n(i); \mathcal{F}(i)), \mathcal{E}_2(i),\ldots,\mathcal{E}_n(i); \mathcal{F}(i)).
$$

One checks that the so defined  $\Xi$  is functorial in i. It remains to be shown that the composition with Ξ induces an isomorphism

<span id="page-62-0"></span>
$$
\operatorname{Hom}_{\operatorname{id}_{S_1}}(\mathcal{E}_1; f^{1,\bullet}(\mathcal{E}_2,\ldots,\mathcal{E}_n;\mathcal{F})) \to \operatorname{Hom}_f(\mathcal{E}_1,\ldots,\mathcal{E}_n;\mathcal{F}).\tag{12}
$$

We will give a map in the other direction which is inverse to composition with Ξ. Let

$$
a \in \text{Hom}_f(\mathcal{E}_1,\ldots,\mathcal{E}_n;\mathcal{F})
$$

be a morphism. To give a morphism on the left hand side of [\(12\)](#page-62-0), for any  $i_1$  and  $\alpha \in X_{i_1}(I)$  we have to give a morphism (functorial in  $i_1$ )

$$
\mathcal{E}_1(i_1) \to f_\alpha^{1,\bullet}(\mathcal{E}_2(i_2);\ldots,\mathcal{E}_n(i_n); \mathcal{F}(j))
$$

or, which is the same, a morphism

$$
\operatorname{Hom}_{f_\alpha}(\mathcal{E}_1(i_1), \mathcal{E}_2(i_2), \ldots, \mathcal{E}_n(i_n); \mathcal{F}(j)).
$$

But we have such a morphism, namely the pre-composition of  $a_i$  with the n-tuple  $\{\mathcal{E}_k(\alpha_k)\}_k$ . (Because we know already that  $Hom(I, \mathcal{D}) \to Hom(I, \mathcal{S})$  is an opfibration of multicategories, it suffices to establish that Ξ is coCartesian in this weak form.)  $\Box$ 

**Remark 4.1.7.** The construction in the proof of the above Proposition will become much clearer, when we define a fibered multiderivator itself as a six-functor-formalism, similar to the definition mentioned in the introduction. For example, for  $S = \{\cdot\}$  we will get an external and internal monoidal product, resp. right adjoints which a clear relation. We have in that case

$$
\mathbb{Z}: \text{Hom}(I,\mathcal{D}) \times \text{Hom}(J,\mathcal{D}) \to \text{Hom}(I \times J,\mathcal{D})
$$

by applying ⊗ point-wise and

$$
\mathcal{HOM}_{l/r} : \text{Hom}(I,\mathcal{D}) \times \text{Hom}(J,\mathcal{D}) \to \text{Hom}(I^{\text{op}} \times J,\mathcal{D})
$$

by applying  $\text{Hom}_{l/r}$  point-wise. The formula for the internal hom obtained in the proof of the proposition boils down to the formula

$$
\mathrm{Hom}_{l/r}(\mathcal{E},\mathcal{F})(i_1)=\int_i\mathcal{H}\mathcal{OM}_{l/r}(\mathcal{E}(i),\mathcal{F}(i))^{\mathrm{Hom}(i_1,i)}
$$

where  $\int_i$  is the categorical end. We refer to a subsequent article [\[14\]](#page-86-8) for an explanation of this in the language of six-functor-formalisms.

We will need later the following

**Lemma 4.1.8.** Let  $f \in Hom(S_1, \ldots, S_n; T)$  be a morphism in  $Hom(I, S)$  for some  $n \geq 2$ . Consider the pull-back functor  $f^{j,\bullet}$  constructed in the proof of Proposition [4.1.6.](#page-60-0) Let  $p: I \times J \to I$  be the projection and fix objects  $\mathcal{E}_1, \tilde{j}$ ,  $\mathcal{E}_n, \mathcal{F}$  in D lying over  $S_1, \tilde{j}$ ,  $S_n, T$ . Then the natural morphism

$$
p^* f^{j, \bullet}(\mathcal{E}_1, \widehat{\cdot}, \widehat{\cdot}, \mathcal{E}_n; \mathcal{F}) \to (p^* f)^{j, \bullet}(p^* \mathcal{E}_1, \widehat{\cdot}, \widehat{\cdot}, p^* \mathcal{E}_n; p^* \mathcal{F})
$$

is an isomorphism, or, in other words, the functor  $p^* : Hom(I, \mathcal{D}) \to Hom(I \times J, \mathcal{S})$  maps Cartesian arrows to Cartesian arrows.

*Proof.* Again, we assume  $j = 1$  to ease the notation. The statement concerning the other pull-backs is completely symmetric. We have by definition

$$
(f^{1,\bullet}(\mathcal{E}_2,\ldots,\mathcal{E}_n;\mathcal{F}))(i')=\lim_{\alpha\in X_{i_1}(I)}f^{1,\bullet}_{\alpha}(\mathcal{E}_2(i_1),\ldots,\mathcal{E}_n(i_n);\mathcal{F}(i'))
$$

and

$$
((p^*f)^{1,\bullet}(\mathcal{E}_2,\ldots,\mathcal{E}_n;\mathcal{F}))(i',j')=\lim_{\alpha\in X_{i_1,j_1}(I\times J)}(p^*f)^{1,\bullet}_{\alpha}((p^*\mathcal{E}_2)(i_1,j_1),\ldots,(p^*\mathcal{E}_n)(i_n,j_n);(p^*\mathcal{F})(i',j'))
$$

The natural map in question is induced by the functor  $\tilde{p}$  :  $X_{i_1,j_1}(I \times J) \to X_{i_1}(I)$  which forgets all data involving the J direction. Now there is also a functor  $\tilde{s}: X_{i_1}(I) \to X_{i_1,j_1}(I \times J)$  which is constant on the J-component with value  $\{\mathrm{id}_{j_1}\}_{k=1..n}$ . We have  $\widetilde{p} \circ \widetilde{s} = \mathrm{id}$  and a chain of natural transformations  $\widetilde{s} \circ \widetilde{p} \Leftarrow \cdots \Rightarrow$  id involving only data in the *J*-direction. However, all the natural transformations are mapped to identities by the functor

$$
\alpha \mapsto \lim_{\alpha \in X_{i_1,j_1}(I\times J)} (p^*f)_{\alpha}^{1,\bullet}((p^*\mathcal{E}_2)(i_1,j_1),\ldots,(p^*\mathcal{E}_n)(i_n,j_n);(p^*\mathcal{F})(i',j'))
$$

because everything is constant along the J-direction. This shows that the natural morphism in the statement is an isomorphism.  $\Box$ 

If I is directed or inverse we want to show that also  $p<sub>I</sub>$  is a bifibration of multi-model-categories in the sense of Definition [4.1.3.](#page-59-0)

<span id="page-63-0"></span>Afterwards we will apply the following variant and generalization to multicategories of the results in  $[2, Ex$ posé XVII,  $\S 2.4$ .

**Proposition 4.1.9.** Let  $p : \mathcal{D} \to \mathcal{S}$  be a bifibration of (multi-)model-categories in the sense of [4.1.3.](#page-59-0) Let W be the union of the W<sub>S</sub> over all objects  $S \in \mathcal{S}$ . Then the fibers of  $\widetilde{p}: \mathcal{D}[\mathcal{W}^{-1}] \to \mathcal{S}$  (as ordinary categories) are the homotopy categories  $\mathcal{D}_S[\mathcal{W}^{-1}_S]$  and  $\widetilde{p}$  is again a bifibration of multicategories such that the push-forward  $F_{\bullet}$  along any  $F \in \text{Hom}_{\mathcal{S}}(X_1,\ldots,X_n;Y)$  (for  $n \geq 1$ ) is the left derived functor of the corresponding push-forward w.r.t. p. Similarly the pull-back w.r.t. some slot ist the right derived functor of the corresponding pull-back w.r.t. p.

4.1.10. The above proposition and its proof have several well-known consequences which we mention, despite being all elementary, because the proof below gives a uniform treatment of all the cases.

1. The homotopy category of a model category is locally small and can be described as the category of cofibrant/fibrant objects modulo homotopy of morphisms. Apply the proof of the proposition to the (trivial) bifibration of ordinary categories  $\mathcal{D} \rightarrow \{\cdot\}.$ 

- 2. Quillen adjunctions lead to an adjunction of the derived functors on the homotopy categories. Apply the proposition to a bifibration of ordinary categories  $\mathcal{D} \to \Delta_1$ .
- 3. The homotopy category of a closed monoidal model category is a closed monoidal category. Apply the proposition to a bifibration of multicategories  $\mathcal{D} \rightarrow \{\cdot\}.$
- 4. Quillen adjunctions in n variables lead to an adjunction in n variables on the homotopy categories. Apply the proposition to a bifibration of multicategories  $\mathcal{D} \to \Delta_{1,n}$ , where the multicategory  $\Delta_{1,n}$  consists of  $n+1$  objects and one n-ary morphism connecting them.

Before proving Proposition [4.1.9,](#page-63-0) we define homotopy relations on  $\text{Hom}_F(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  where  $F \in$  $Hom(X_1, \ldots, X_n; Y)$  is a multimorphism in S.

**Definition 4.1.11.** 1. Two morphisms f and g in  $\text{Hom}_F(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  are called right ho**motopic** if there is a path object of  $\mathcal F$ 

$$
\mathcal{F}\longrightarrow \mathcal{F}'\smash{\xrightarrow[\text{pr}_2]{\text{pr}_1}} \mathcal{F}
$$

and a morphism Hom $(\mathcal{E}_1,\ldots,\mathcal{E}_n;\mathcal{F}')$  over the same multimorphism F such that the compositions with  $pr_1$  and  $pr_2$  are f and g, respectively.

2. For  $n \geq 1$ , two morphisms f and g in  $\text{Hom}_F(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  are called *i*-left homotopic if there is a cylinder object  $\mathcal{E}'_i$  of  $\mathcal{E}_i$ 

$$
\mathcal{E}_i \xrightarrow[\iota_2]{\iota_1} \mathcal{E}'_i \longrightarrow \mathcal{E}_i
$$

and a morphism  $Hom(\mathcal{E}_1,\ldots,\mathcal{E}'_i,\ldots,\mathcal{E}_n;\mathcal{F})$  over F such that the compositions with  $\iota_1$  and  $\iota_2$ are f and g, respectively.

- **Lemma 4.1.12.** 1. The condition 'right homotopic' is preserved under pre-composition, while the condition 'i-left homotopic' is preserved under post-composition.
	- 2. Let  $n \geq 1$ . If  $f, g \in Hom(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  are *i*-left homotopic and all  $\mathcal{E}_i$  are cofibrant then f and g are right homotopic. If  $f, g \in \text{Hom}(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  are right homotopic,  $\mathcal F$  is fibrant, and all  $\mathcal{E}_j$  for  $j \neq i$  are cofibrant then f and g are *i*-left homotopic.
	- 3. Let  $n \geq 1$ . In Hom $(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  right homotopy is an equivalence relation if all  $\mathcal{E}_i$  are cofibrant. In Hom $(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  i-left homotopy is an equivalence relation if  $\mathcal F$  is fibrant, and all  $\mathcal{E}_i$ ,  $j \neq i$  are cofibrant

In particular on the category  $\mathcal{D}^{\text{Cof,Fib}}$  of fibrant/cofibrant objects, *i*-left homotopy=right homotopy is an equivalence relation, which is compatible with composition.

Proof. 1. is obvious.

2. If all  $\mathcal{E}_i$  are cofibrant then also  $F_{\bullet}(\mathcal{E}_1,\ldots,\mathcal{E}_n)$  is cofibrant and f and g correspond uniquely to morphisms  $f', g' : F_{\bullet}(\mathcal{E}_1, \ldots, \mathcal{E}_n) \to \mathcal{F}$ . Since f and g are *i*-left homotopic, there is a cylinder object

$$
\mathcal{E}_i \Longrightarrow \mathcal{E}'_i \longrightarrow \mathcal{E}_i
$$

realizing the *i*-left homotopy. Since  $\mathcal{E}_i$  is cofibrant so is  $\mathcal{E}'_i$ . Hence also

$$
F_{\bullet}(\mathcal{E}_1,\ldots,\mathcal{E}_n) \longrightarrow F_{\bullet}(\mathcal{E}_1,\ldots,\mathcal{E}'_i,\ldots,\mathcal{E}_n) \longrightarrow F_{\bullet}(\mathcal{E}_1,\ldots,\mathcal{E}_n)
$$

is a cylinder object because all  $\mathcal{E}_j$  are cofibrant, and hence also  $f'$  and  $g'$  are left homotopic. These are therefore also right homotopic and hence so are  $f$  and  $g$ . Dually we obtain the second statement. 3. follows from [\[15,](#page-86-6) Proposition 1.2.5, (iii)]. □

<span id="page-65-0"></span>**Lemma 4.1.13.** Two i-left homotopic morphisms become equal in  $\mathcal{D}^{\text{Cof}}[(\mathcal{W}^{\text{Cof}})^{-1}]$ .

Proof. This follows from the fact that a cylinder object

$$
\mathcal{E}_i \xrightarrow[1]{\iota_1} \mathcal{E}'_i \xrightarrow{p} \mathcal{E}_i
$$

automatically lies in  $\mathcal{D}^{\text{Cof}}$  if  $\mathcal{E}_i$  does, and the two maps  $\iota_1$  and  $\iota_2$  become equal because p becomes invertible.  $\Box$ 

We have to distinguish the easier case, in which all objects  $F_{\bullet}$  () for 0-ary morphisms F are cofibrant. Otherwise we define a category  $\mathcal{D}^{\text{Cof}}[\widetilde{(\mathcal{W}^{\text{Cof}})}^{-1}]$  in which we set  $\text{Hom}_F(\mathcal{F}) \coloneqq \text{Hom}_{\mathcal{D}_S[\mathcal{W}^{-1}_S]}(QF_{\bullet}(\mathcal{F};\mathcal{F})$ for all F, where F is a 0-ary morphism with domain S. Composition is given as follows: For a morphism  $f \in Hom_G(\mathcal{E}_1,\ldots,\mathcal{E}_n;\mathcal{F})$  with cofibrant  $\mathcal{E}_i$  and  $\mathcal{F}$  and  $\xi: QF_{\bullet}() \to \mathcal{E}_i$ , we define the composition  $\xi \circ f$  as the following composition

$$
\varepsilon_1
$$
\n
$$
\overline{\varepsilon}_n
$$
\n
$$
\varepsilon_n
$$
\n
$$
\varepsilon_n
$$
\n
$$
\leftarrow G_{\bullet}(\mathcal{E}_1, \ldots, QF_{\bullet}(), \ldots, \mathcal{E}_n) \longrightarrow G_{\bullet}(\mathcal{E}_1, \ldots, \mathcal{E}_n) \longrightarrow \mathcal{F}.
$$

<span id="page-65-1"></span>One checks that the so-defined composition is associative and independent of the choice of the push-forwards.

**Lemma 4.1.14.** If the object  $F_{\bullet}$  () is cofibrant for every 0-ary morphism F then the natural functor

$$
\mathcal{D}^{\mathrm{Cof}}[(\mathcal{W}^{\mathrm{Cof}})^{-1}] \to \mathcal{D}[\mathcal{W}^{-1}]
$$

is an equivalence of categories. Otherwise it is, if we replace  $\mathcal{D}^{\text{Cof}}[(\mathcal{W}^{\text{Cof}})^{-1}]$  by  $\mathcal{D}^{\text{Cof}}[\widetilde{(\mathcal{W}^{\text{Cof}})^{-1}}]$ .

*Proof.* The inclusion  $\mathcal{D}^{Cof} \to \mathcal{D}$  induces a functor  $\Xi : \mathcal{D}^{Cof}[(\mathcal{W}^{Cof})^{-1}] \to \mathcal{D}[\mathcal{W}^{-1}]$ . If the objects  $F_{\bullet}$ () are not cofibrant then  $\Xi$  may be modified to a functor

$$
\mathcal{D}^{\mathrm{Cof}}\big[\widetilde{(\mathcal{W}^{\mathrm{Cof}})^{-1}}\big]\rightarrow \mathcal{D}[\mathcal{W}^{-1}]
$$

as follows: a 0-ary morphism  $QF_{\bullet}$   $\rightarrow$   $\mathcal F$  is mapped to the composition

$$
\underbrace{\circ \text{ccart}} \to F_{\bullet}() \longleftarrow QF_{\bullet}() \longrightarrow \mathcal{F}
$$

in  $\mathcal{D}[\mathcal{W}^{-1}].$ 

We now specify a functor  $\Phi$  in the other direction.  $\Phi$  maps an object X to a cofibrant replacement QX. For  $n \geq 1$ , a morphism  $f \in Hom(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  over F is mapped to the following morphism. Composing with the morphisms  $Q\mathcal{E}_i \to \mathcal{E}_i$ , we get a morphism  $f' \in Hom(Q\mathcal{E}_1,\ldots,Q\mathcal{E}_n;\mathcal{F})$  or equivalently a morphism  $X_i \to F^{\bullet,i}(Q\mathcal{E}_1,\tilde{X}_i,Q\mathcal{E}_n;\mathcal{F})$ . Now choose a lift (dotted arrow in the diagram)



which exists because the vertical map is again a trivial fibration (because all the  $Q\mathcal{E}_i$  are cofibrant). The resulting map in  $Hom(Q\mathcal{E}_1,\ldots,Q\mathcal{E}_n;P\mathcal{F})$  is actually well-defined in  $\mathcal{D}^{\text{Cof}}[(\mathcal{W}^{\text{Cof}})^{-1}]$ . Indeed, two different lifts are left homotopic because  $Q\mathcal{E}_i$  is cofibrant [\[15](#page-86-6), Proposition 1.2.5. (iv)], and therefore also the two morphisms in  $Hom(Q\mathcal{E}_1,\ldots,Q\mathcal{E}_n;Q\mathcal{F})$  become equal in  $\mathcal{D}^{\text{Cof}}[(\mathcal{W}^{\text{Cof}})^{-1}]$  by Lemma [4.1.13.](#page-65-0) From this it follows that  $\Phi$  is indeed a functor on *n*-ary morphisms for  $n \geq 1$ . For  $n = 0$ , a morphism  $f \in Hom(\mathcal{F})$  over F corresponds to a morphism  $F_{\bullet}() \rightarrow \mathcal{F}$ . If  $F_{\bullet}()$ is cofibrant, this morphism lifts (again uniquely up to right homotopy) to a morphism  $F_{\bullet}() \rightarrow$  $Q\mathcal{F}$ , i.e. to a morphism in Hom  $_F$  (;  $Q\mathcal{F}$ ). If  $F_{\bullet}$  () is not cofibrant then the composition lifts to a morphism:  $QF_{\bullet}(\rightharpoonup P\mathcal{F}$  which is defined to be the image of  $\Phi$ . Furthermore  $\Phi$  is inverse to  $\Xi$  up to isomorphism. 口

**Lemma 4.1.15.** Right homotopic morphisms become equal in  $\mathcal{D}^{\text{Cof},\text{Fib}}[(\mathcal{W}^{\text{Cof},\text{Fib}})^{-1}]$ .

Proof. The assertion follows from the fact that there exists a path object

$$
\mathcal{F} \xleftarrow{\mathrm{pr}_1} \mathcal{F}' \xleftarrow{i} \mathcal{F}
$$

where  $\mathcal{F}'$  is cofibrant and fibrant which realizes the right homotopy [\[15](#page-86-6), Proposition 1.2.6.]. This uses that all sources are cofibrant and the domain is fibrant. The two morphisms  $pr_1$  and  $pr_2$ become equal because  $i$  becomes invertible.  $\Box$ 

Lemma 4.1.16. The functor  ${\cal D}^{\rm Fib, Cof}[(W^{\rm Fib, Cof})^{-1}]\to {\cal D}^{\rm Cof}[(W^{\rm Cof})^{-1}]$ **Lemma 4.1.16.** The functor  $\mathcal{D}^{\text{Fib},\text{Cof}}[(\mathcal{W}^{\text{Fib},\text{Cof}})^{-1}] \to \mathcal{D}^{\text{Cof}}[(\mathcal{W}^{\text{Cof}})^{-1}]$ , respectively, are equivalences of multicategories.

Proof. The proof is analogous to that of Lemma [4.1.14](#page-65-1) but with some minor changes which require, in particular, the chosen order of restriction to cofibrant and fibrant objects. We specify again a functor  $\Phi$  in the other direction. On objects,  $\Phi$  maps  $\mathcal E$  to a fibrant replacement  $R\mathcal E$ . Note that RE is still cofibrant. A morphism  $f \in Hom(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  over F corresponds to a morphism  $F_{\bullet}(\mathcal{E}_1,\ldots,\mathcal{E}_n) \to \mathcal{F}$ . Now choose a lift (dotted arrow in the diagram)



It exists because the vertical maps are again trivial cofibrations (because all the  $\mathcal{E}_i$  and  $R\mathcal{E}_i$  are cofibrant). The lift is well-defined in  $\mathcal{D}^{\text{Cof,Fib}}[(\mathcal{W}^{\text{Cof,Fib}})^{-1}]$ , because two lifts in the triangle above become right homotopic (because  $R\mathcal{F}$  is fibrant by [\[15](#page-86-6), Proposition 1.2.5. (iv)]). Therefore also the corresponding morphisms in  $Hom(R\mathcal{E}_1,\ldots,R\mathcal{E}_n;R\mathcal{F})$  become equal in  $\mathcal{D}^{Cof,Fib}[(\mathcal{W}^{Cof,Fib})^{-1}]$ by the previous lemma. It follows that  $\Phi$  is indeed a functor which is inverse to the inclusion up to isomorphism. □

**Lemma 4.1.17.** If the objects  $F_{\bullet}$ () for all 0-ary morphisms in S are cofibrant then the natural functor

$$
\mathcal{D}^{\mathrm{Fib},\mathrm{Cof}}\big[(\mathcal{W}^{\mathrm{Fib},\mathrm{Cof}})^{-1}\big]\rightarrow\mathcal{D}^{\mathrm{Fib},\mathrm{Cof}}/\sim
$$

is an isomorphism of categories. Otherwise it is, if we modify the 0-ary morphisms as before.

*Proof.* The natural functor  $\mathcal{D}^{\text{Fib},\text{Cof}} \to \mathcal{D}^{\text{Fib},\text{Cof}}/ \sim$  takes weak equivalences to isomorphisms [\[15](#page-86-6), Proposition 1.2.8] and has the universal property of  $\mathcal{D}^{\text{Fib,Cof}}[(\mathcal{W}^{\text{Fib,Cof}})^{-1}]$  by the same argument as in [\[15](#page-86-6), Proposition 1.2.9].  $\Box$ 

*Proof of Proposition [4.1.9.](#page-63-0)* The previous lemmas showed that  $\mathcal{D}[\mathcal{W}^{-1}]$  is equivalent to  $\mathcal{D}^{\text{Fib},\text{Cof}}/\sim$  if all objects of the form  $F_{\bullet}$ () are cofibrant, or if we replace the second multicategory by  $\mathcal{D}^{\widehat{\text{Fib},\text{Cof}}}/\sim$ , where we set  $\text{Hom}_{F,\mathcal{D}^{\text{Fib},\text{Cof}}/\sim}((X)) \coloneqq \text{Hom}_{\mathcal{D}_S[\mathcal{W}_S^{-1}]}(F_{\bullet}(),\mathcal{F})$  for all 0-ary morphism F in S with domain S and for every  $\mathcal{F} \in \mathcal{D}_S$ .

It remains to show that the functor

$$
p/\sim: \mathcal{D}^{\text{Fib},\text{Cof}}/\sim \rightarrow \mathcal{S}
$$

is bifibered if all  $F_{\bullet}$  () are cofibrant or otherwise bifibered for  $n \geq 1$  (i.e. (co)Cartesian morphisms exist for  $n \geq 1$ ). (The modification  $\mathcal{D}^{\widehat{\text{Fib},\text{Cof}}}$   $\sim$  has been constructed in such a way that it has  $\text{coCartesian morphisms}$  for  $n = 0.$ )

We show that  $p / \sim$  is opfibered, the other case being similar. Let F be a multimorphism in S with codomain S. The set  $\text{Hom}_F(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  modulo right homotopy is in bijection with the set  $\text{Hom}_{\mathcal{D}_Y}(F_\bullet(\mathcal{E}_1,\ldots,\mathcal{E}_n),\mathcal{F})$  modulo right homotopy. Since  $\mathcal F$  is fibrant, the latter set is the same as  $\text{Hom}_{\mathcal{D}_S}(R(F_\bullet(\mathcal{E}_1,\ldots,\mathcal{E}_n)),\mathcal{F})$  modulo right homotopy. Hence morphisms in  $\text{Hom}_F(\mathcal{E}_1,\ldots,\mathcal{E}_n;\mathcal{F})$ 

uniquely decompose as the composition



followed by a morphism in  $\text{Hom}_{\mathcal{D}_S}(R(F_\bullet(\mathcal{E}_1,\ldots,\mathcal{E}_n)),\mathcal{F})$  modulo right homotopy. More generally, by the same argument, a morphism in some  $\text{Hom}_{GF}(\mathcal{F}_1,\ldots,\mathcal{E}_1,\ldots,\mathcal{E}_n,\ldots,\mathcal{F}_m;\mathcal{G})$ , where G is another multimorphism in  $S$ , modulo right homotopy factorizes uniquely into the above composition followed by a morphism in

$$
\mathrm{Hom}_G(\mathcal{F}_1,\ldots,R(F_{\bullet}(\mathcal{E}_1,\ldots,\mathcal{E}_n)),\ldots,\mathcal{F}_m;\mathcal{G})
$$

modulo right homotopy.

It remains to see that the push-forward in  $\mathcal{D}[\mathcal{W}^{-1}]$  corresponds to the left derived functor of  $F_{\bullet}$ . For any objects  $\mathcal{E}_1, \ldots, \mathcal{E}_n$  the composition



is a coCartesian morphism lying over F, with domains isomorphic to the  $\mathcal{E}_i$ . However, the object  $R(F_{\bullet}(RQ\mathcal{E}_1,\ldots,RQ\mathcal{E}_n))$  is isomorphic to the value of the left derived functor of  $F_{\bullet}$  at  $\mathcal{E}_1, \ldots, \mathcal{E}_n$ .  $\Box$ 

<span id="page-68-0"></span>4.1.18. We now focus on the left case. If I is a directed diagram, we proceed to construct a model structure on the fibers of the bifibration of multicategories (cf. [4.1.6\)](#page-60-0):

$$
\mathrm{Hom}(I,\mathcal{D})\to \mathrm{Hom}(I,\mathcal{S})=\mathbb{S}(I).
$$

This model structure is an analogue of the classical Reedy model structure and it has the property that pull-backs w.r.t. diagrams and the corresponding relative left Kan extension functors form a Quillen adjunction.

Let I ∈ Dir and let  $F: I \to S$  be a functor. We will define a model-category structure

$$
({\mathcal D}_F,{\rm Cof}_F,{\rm Fib}_F,\mathcal{W}_F)
$$

where  $\mathcal{D}_F$  is the fiber of  $\text{Hom}(I,\mathcal{D})$  over F and where  $\mathcal{W}_F$  is the class of morphisms which are element-wise in the corresponding  $\mathcal{W}_{F(i)}$ .

For any  $G \in \mathcal{D}_F$ , and for any  $i \in I$ , we define a **latching object** 

$$
L_iG \coloneqq \mathrm{colim}_{I_i}\{F(\alpha)_\bullet G(j)\}_{\alpha:j\to i},
$$

Here  $I_i$  is the full subcategory of  $I \times_{I} i$  consisting of all objects except id<sub>i</sub>. We have a canonical morphism

$$
L_iG \to G(i)
$$

in  $\mathcal{D}_{F(i)}$ . We define Fib<sub>F</sub> to be the class of morphisms which are element-wise in the corresponding Fib<sub>F(i)</sub>. We define Cof<sub>F</sub> to be the class of morphisms  $G \to H$  such that for any  $i \in I$  the induced morphism  $\delta$  in the diagram

$$
L_i G \longrightarrow L_i H
$$
  
\n
$$
G(i) \longrightarrow \text{push-out } \cdots \stackrel{\delta}{\longrightarrow} H(i)
$$

belongs to  $\mathrm{Cof}_{F(i)}$ . We call a morphism  $G \to H$  in  $\mathrm{Cof}_F$  temporarily an **acyclic cofibration** if  $\delta$  is, in addition, a weak equivalence. The proof that this yields a model-category structure is completely analogous to the classical case [\[15](#page-86-6),  $\S 5.1$ ] (if S is trivial). We need a couple of lemmas:

**Lemma 4.1.19.** The class of cofibrations (resp. acyclic cofibrations) in  $\mathcal{D}_F$  consists precisely of the morphisms which have the left lifting property w.r.t. trivial fibrations (resp. fibrations). These are stable under retracts.

Proof. This is shown as in the classical case: we first prove that acyclic cofibrations have the lifting property w.r.t. fibrations. Consider a diagram



where  $\alpha$  is an acyclic cofibration and  $\beta$  is a fibration. We proceed by induction on n and assume that for all  $i \in I$  with  $\nu(i) < n$  a map  $G_2(i) \to H_1(i)$  has been constructed such that it is a lift in the above diagram, evaluated at i. For each i of degree n consider the following diagram (where the morphism  $L_iG_2 \rightarrow L_iH_1 \rightarrow H_1(i)$  is formed using the already constructed lifts):

$$
G_1(i) \amalg_{L_iG_1} L_iG_2 \longrightarrow H_1(i)
$$
  
\n
$$
\downarrow^{\alpha'(i)} \qquad \qquad \downarrow^{\beta(i)}
$$
  
\n
$$
G_1(i) \longrightarrow H_2(i)
$$

Here  $\alpha'(i)$  is a trivial Cof<sub>F(i)</sub>-cofibration by definition, and  $\beta(i)$  is a Fib<sub>F(i)</sub>-fibration by definition. Hence a lift exists. In the same way the statement for cofibrations and for trivial fibrations is shown. Closure under retracts is left as an exercise for the reader. The assertion that the class of acyclic cofibrations (resp. cofibrations) is precisely the class of morphisms that have the left lifting property w.r.t. fibrations (resp. trivial fibrations) follows from the retract argument as for model categories.  $\Box$ 

**Lemma 4.1.20.** There exists a functorial factorization of morphisms in  $\mathcal{D}_F$  into a fibration followed by an acyclic cofibration and into a trivial fibration followed by a cofibration.

*Proof.* We show this again by induction on  $n$ . We do the first case, the other being similar. Let  $G \rightarrow K$  a morphism in  $\mathcal{D}_F$ . We have the following diagram:



Here the top row is constructed using the already defined factorizations. The object  $H(i)$  and the dotted maps are constructed as the factorization in the model category  $\mathcal{D}_{F(i)}$  into a trivial  $\mathrm{Cof}_{F(i)}$ -cofibration followed by  $\mathrm{Fib}_{F(i)}$ -fibration.  $\Box$ 

Lemma 4.1.21. The classes of cofibrations, acyclic cofibrations, fibrations and weak equivalences are stable under composition.

Proof. This follows from the characterization by a lifting property (resp. by definition for the case of the weak equivalences).  $\Box$ 

Lemma 4.1.22. Acyclic cofibrations are precisely the trivial cofibrations.

Proof. We begin by showing that an acyclic cofibration is a weak equivalence. It suffices to show that in the diagram



the top horizontal morphism is a trivial cofibration. Then the lower horizontal morphism is a composition of two trivial cofibrations and hence is a weak equivalence. The top morphism is indeed a trivial cofibration because the morphism of  $I_i$ -diagrams (cf. [4.1.18\)](#page-68-0)

$$
\{F(\alpha)_\bullet G(j)\}_{\alpha:j\to i}\to \{F(\alpha)_\bullet G(j)\}_{\alpha:j\to i}
$$

is a trivial cofibration in the classical sense (i.e. over the constant diagram over  $I_i$  with value  $F(i)$ ) because of Lemma[s4.1.23](#page-70-0) and [4.1.24.](#page-71-0)

In the other direction, let f be a trivial cofibration and factor it as  $f = pq$ , where q is an acyclic cofibration and  $p$  is a fibration. It follows that  $p$  is a weak-equivalence. Now construct a lift in the diagram



<span id="page-70-0"></span>This shows that  $f$  is a retract of  $g$ , and hence is an acyclic cofibration, too.

**Lemma 4.1.23.** For each (1-ary) morphism of diagrams  $f \in Hom_S(X_1;Y)$  there is an associated push-forward and an associated pull-back, defined by taking the point-wise push-forward  $f_{\bullet}$ , and point-wise pull-back  $f^{\bullet}$  (cf. [4.1.6\)](#page-60-0), respectively. The push-forward  $f_{\bullet}$  respects the classes of cofibrations and acyclic cofibrations. The pull-back  $f^{\bullet}$  respects the classes of fibrations and trivial fibrations.

 $\Box$ 

*Proof.* It suffices (by the lifting property) to show that  $f^{\bullet}$  respects fibrations and trivial fibrations. This is clear because they are defined point-wise.  $\Box$ 

<span id="page-71-0"></span>A posteriori this will say that the pair of functors  $f^{\bullet}, f_{\bullet}$  form a Quillen adjunction between the corresponding model categories (cf. [4.1.28\)](#page-72-0).

**Lemma 4.1.24.** Let  $i \in I$  be an object, let  $\iota : I_i \to I$  be the corresponding latching category with its natural functor to I, and let  $F_i := \iota^* F : I_i \to S$  be the restriction of F to I<sub>i</sub>. The pull-back  $\iota^* : \mathcal{D}_F \to \mathcal{D}_{F_i}$  respects cofibrations and acyclic cofibrations.

Proof. It is easy to see that the pull-back induces an isomorphism of the corresponding latching  $\Box$ objects as in the classical case.

Corollary 4.1.25. The structure constructed in [4.1.18](#page-68-0) defines a model category.

Proof. This follows from the previous Lemmas.

**Proposition 4.1.26.** For any morphism of directed diagrams  $\alpha : I \rightarrow J$ , and for any functor  $F: J \rightarrow S$ , the functor

$$
\alpha^*:\mathcal{D}_F\to\mathcal{D}_{\alpha^*F}
$$

has a left adjoint  $\alpha_1^F$ . The pair  $\alpha^*$ ,  $\alpha_1^F$  define a Quillen adjunction.

*Proof.* That the two functors define a Quillen adjunction is clear once we have shown that  $\alpha_!$  exists because  $\alpha^*$  preserves fibrations and weak equivalences. Let G be an object of  $\mathcal{D}_F$ . We define

$$
(\alpha_!G)(j)\coloneqq \mathrm{colim}_{I\times_{J}J} \mathbb{S}(\mu) \mathbf{.}\iota_j^*G.
$$

For each morphism  $\mu : j \to j'$  we get a functor

$$
\widetilde{\mu}: I \times_{/{\cal J}} j \to I \times_{/{\cal J}} j'
$$

and hence an induced morphism

$$
F(\mu)_{\bullet} \mathbb{S}(\mu)_{\bullet} \iota_j^* G \to \widetilde{\mu}^* \mathbb{S}(\mu')_{\bullet} \iota_{j'}^*.
$$

Sincd  $F(\mu)$  commutes with colimits we get a morphism

$$
F(\mu) \bullet \operatorname{colim}_{I \times_{J} J} \mathbb{S}(\mu) \bullet \iota_j^* G \to \operatorname{colim}_{I \times_{J} J'} \mathbb{S}(\mu') \bullet \iota_{J'}^*
$$

which we define to be  $(\alpha_0 G)(\mu)$ . We now proceed to show that the functor we have constructed is indeed adjoint to  $\alpha^*$ . A morphism  $\mu: G \to \alpha^* H$  is given by a collection of maps  $a(i): G(i) \to$  $H(\alpha(i))$  for all objects  $i \in I$ , subject to the condition that the diagram

$$
F(\alpha(\lambda)) \bullet G(i) \xrightarrow{F(\alpha(\lambda)) \bullet a(i)} F(\alpha(\lambda)) \bullet H(\alpha(i))
$$
  
\n
$$
\downarrow \overline{G(\lambda)} \qquad \qquad \downarrow \overline{H(\alpha(\lambda))}
$$
  
\n
$$
G(i') \xrightarrow{a(i')} H(\alpha(i'))
$$

commutes for each morphism  $\lambda : i \to i'$  in *I*. For each  $j \in J$  and morphism  $\mu : \alpha(i) \to j$  we get a morphism

$$
\overline{H(\mu)} \circ (F(\mu) \bullet a(i)) : F(\mu) \bullet G(i) \to H(j)
$$

 $\Box$
and therefore for fixed  $j$  a morphism

$$
\operatorname{colim}_{I \times_{jJ} j} \mathbb{S}(\mu) \bullet \iota_j^* G \to H(j).
$$

One checks that this yields a morphism  $\alpha_! G \to H$ . On the other hand, let  $b : \alpha_! G \to H$  be a morphism given by

$$
b(j): \operatorname{colim}_{I \times_{jJ} j} \mathbb{S}(\mu) \bullet \iota_j^* G \to H(j)
$$

or equivalently for all  $\mu : \alpha(i) \to j$  by morphisms

$$
F(\mu) \bullet G(i) \to H(j).
$$

In particular, if  $\mu$  is the identity of  $\alpha(i)$ , we get morphisms

$$
G(i) \to H(\alpha(i))
$$

which constitute a morphism of diagrams  $G \to \alpha^* H$ . One checks that these associations are inverse to each other.  $\Box$ 

**Lemma 4.1.27.** Let  $\alpha: I \rightarrow J$  be a morphism of directed diagrams and let j be an object of J. The functor  $\iota_j^* : \mathcal{D}_I \to \mathcal{D}_{I \times_{I} J}$  respects cofibrations and trivial cofibrations.

*Proof.* This follows easily from the fact that  $\iota_j$  induces a canonical identification

$$
I_i = (I \times_{j} j)_\mu
$$

for any  $\mu = (i, \alpha(i) \to j)$ . For this implies that we have a canonical isomorphism  $L_i G \cong L_{\mu} \iota_j^* G$ .

<span id="page-72-0"></span>**Lemma 4.1.28.** The bifibration of multicategories, defined in  $4.1.6$ 

$$
\operatorname{Hom}(I,\mathcal{D})\to \operatorname{Hom}(I,\mathcal{S})=\mathbb{S}(I)
$$

equipped with the model-category structures constructed in [4.1.18](#page-68-0) is a bifibration of multi-modelcategories in the sense of [4.1.3.](#page-59-0)

*Proof.* First for each multi-morphism of diagrams  $f \in Hom_{\mathcal{S}}(X_1, \ldots, X_n; Y)$  we have to see that the push-forward and the various pull-backs form a Quillen adjunction in n variables. The case  $n = 1$ has been treated above. We only work out the case  $n = 2$ , the proof for higher n being similar. It suffices to check the following: for any cofibration  $\mathcal{E}_1 \to \mathcal{E}'_1$  and for any fibration  $\mathcal{F} \to \mathcal{F}'$  the dotted induced morphism in the following diagram

$$
f^{\bullet,2}(\mathcal{E}'_1;\mathcal{F}) \longrightarrow \text{pull-back} \longrightarrow f^{\bullet,2}(\mathcal{E}'_1;\mathcal{F}')
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$
\n
$$
f^{\bullet,2}(\mathcal{E}_1;\mathcal{F}) \longrightarrow f^{\bullet,2}(\mathcal{E}_1;\mathcal{F}')
$$

<span id="page-72-1"></span>is a fibration. Since fibrations are defined point-wise and fibered products are computed point-wise, we have only to see that the assertion holds point-wise. Now  $\mathcal{F} \to \mathcal{F}'$  is a point-wise fibration and  $\mathcal{E}_1 \to \mathcal{E}'_1$  is a Reedy cofibration, so by the reasoning in the proof of Lemma [4.1.22](#page-70-0) it is in particular a point-wise cofibration. Hence the assertion holds because of the assumption that  $\mathcal{D} \to \mathcal{S}$  is a bifibration of multi-model-categories [\(4.1.3\)](#page-59-0). The requested property for the 0-ary push-forward is easier and is left to the reader.  $\Box$  **Proposition [4.1.2](#page-59-1)9.** The functor  $\mathbb{D}(I) \to \mathbb{S}(I)$  defined in 4.1.2 is a bifibration of multicategories whose fibers are equivalent to  $\mathcal{D}_F[\mathcal{W}_F^{-1}]$ . The pull-back and push-forward functors are given by the left derived functors of  $f_{\bullet}$ , and by the right derived functors of  $f^{\bullet,j}$ , respectively.

*Proof.* We have seen in [4.1.28](#page-72-0) that the fibers of  $Hom(I, \mathcal{D}) \rightarrow S(I)$  are a bifibration of multi-model-categories in the sense of [4.1.3.](#page-59-0) Therefore by Proposition [4.1.9](#page-63-0) we get that  $\mathbb{D}(I) \to \mathbb{S}(I)$  are bifibered multicategories with the requested properties.  $\Box$ 

*Proof of Theorem [4.1.5.](#page-60-1)* (Der1) and (Der2) for  $\mathbb D$  and  $\mathbb S$  are obvious.

(FDer0 left) and the first part of (FDer0 right) follow from Theorem [4.1.29.](#page-72-1)

(FDer3 left) follows from [4.1.26.](#page-71-0)

(FDer4 left): By construction of  $\alpha_1$  the natural base-change

<span id="page-73-0"></span>
$$
\operatorname{colim} \mathbb{S}(\mu) \bullet \iota_j^* G \to j^* \alpha_! G \tag{13}
$$

is an isomorphism for the non-derived functors. For the derived functors the same follows because all functors in the equation respect cofibrations and trivial cofibrations and all functors which have to be derived in [\(13\)](#page-73-0) are left Quillen functors and hence can be derived by composing them with cofibrant replacement.

(FDer3 right) and (FDer4 right) are shown precisely the same way.

(FDer5 left): Fixing a morphism  $f \in Hom(S_1, ..., S_n; T)$  in S and objects  $\mathcal{E}_2, ..., \mathcal{E}_n$  over  $S_2, ..., S_n$ we have by Theorem [4.1.29](#page-72-1) a push-forward functor

$$
\mathbb{D}(I \times J)_{p^*S_1} \rightarrow \mathbb{D}(I \times J)_{p^*T}
$$
  

$$
\mathcal{E}_1 \rightarrow (p^*f)_{\bullet}(\mathcal{E}_1, p^*\mathcal{E}_2, \dots, p^*\mathcal{E}_n)
$$

(we denote it with the same letter as the underived version) which, by (FDer0 left), defines a morphism of pre-derivators

$$
\mathbb{D}_{S_1}\to \mathbb{D}_T.
$$

We first show that it preserves colimits, i.e. that for  $p: J \to \cdot$  we have that for all  $\mathcal{E}_1 \in \mathcal{D}_{p^*S_1}(I \times J)$ the natural morphism

$$
f_{\bullet}(p_{*}\mathcal{E}_{1},\mathcal{E}_{2},\cdots,\mathcal{E}_{n})\rightarrow p_{*}(p^{*}f)_{\bullet}(\mathcal{E}_{1},p^{*}\mathcal{E}_{2},\cdots,p^{*}\mathcal{E}_{n})
$$

(where we wrote p also for the projection  $p: I \times J \to I$ ) is an isomorphism. This is the same as showing that

$$
p^* f^{1,\bullet}(\mathcal{E}_1,\ldots,\mathcal{E}_n) \to (p^* f)^{1,\bullet}(p^* \mathcal{E}_1,\ldots,p^* \mathcal{E}_n)
$$

is an isomorphism. This follows from Lemma [4.1.8](#page-62-0) because it suffices to check this for the underived functors. Now let  $\alpha : I \to J$  be a Grothendieck opfibration. To show that

$$
f_{\bullet}(\alpha_{*}\mathcal{E}_{1},\mathcal{E}_{2},\ldots,\mathcal{E}_{n}) \to \alpha_{*}(\alpha^{*}f)_{\bullet}(\mathcal{E}_{1},\alpha^{*}\mathcal{E}_{2},\ldots,\alpha^{*}\mathcal{E}_{n})
$$

is an isomorphism we may show this point-wise. Indeed, after applying  $j^*$  we get

$$
(j^*f) \bullet (j^*\alpha_* \mathcal{E}_1, j^*\mathcal{E}_2, \dots, j^*\mathcal{E}_n) \to j^*\alpha_* (\alpha^* f) \bullet (\mathcal{E}_1, \alpha^* \mathcal{E}_2, \dots, \alpha^* \mathcal{E}_n)
$$
  

$$
(j^*f) \bullet (p_* \iota_j^* \mathcal{E}_1, j^* \mathcal{E}_2, \dots, j^* \mathcal{E}_n) \to p_* \iota_j^* (\alpha^* f) \bullet (\mathcal{E}_1, \alpha^* \mathcal{E}_2, \dots, \alpha^* \mathcal{E}_n)
$$

where  $\iota_j : I_j \to I$  is the inclusion of the fiber. Note that the commutative diagram



is homotopy exact by Lemma [1.3.23,](#page-15-0) 2. because  $\alpha$  is a Grothendieck opfibration. Finally we get the morphism

$$
(j^*f) \bullet (p_* \iota_j^* \mathcal{E}_1, j^* \mathcal{E}_2, \dots, j^* \mathcal{E}_n) \to p_* (j^*f) \bullet (\iota_j^* \mathcal{E}_1, p^* j^* \mathcal{E}_2, \dots, p^* j^* \mathcal{E}_n)
$$

which is an isomorphism by the above reasoning.

By Lemma [1.3.8](#page-10-0) the full content of (FDer0 right) follows from (FDer5 left) while (FDer5 right) follows from (FDer0 left).  $\Box$ 

## <span id="page-75-0"></span>A Fibrations of categories

### A.1 Grothendieck (op-)fibrations

**A.1.1** (right). Let  $p : \mathcal{D} \to \mathcal{S}$  be a functor, and let  $f : S \to T$  be a morphism in S. A morphism  $\xi : \mathcal{E}' \to \mathcal{E}$  over f is called **Cartesian** if the composition with  $\xi$  induces an isomorphism

$$
\operatorname{Hom}_{g}(\mathcal{F},\mathcal{E}') \cong \operatorname{Hom}_{f \circ g}(\mathcal{F},\mathcal{E})
$$

for any morphism  $q: R \to S$  in S and for every  $\mathcal{F} \in \mathcal{D}_R$ .

The functor p is called a Grothendieck fibration if for any  $f : S \to T$  and for every object E in  $\mathcal{D}_T$  (i.e. such that  $p(\mathcal{E}) = T$ ) there exists a Cartesian morphism  $\mathcal{E}' \to \mathcal{E}$ .

**A.1.2** (left). Let  $p : \mathcal{D} \to \mathcal{S}$  be a functor, and let  $f : S \to T$  be a morphism in S. A morphism  $\xi : \mathcal{E} \to \mathcal{E}'$  over f is called **coCartesian** if the composition with  $\xi$  induces an isomorphism

$$
\operatorname{Hom}_{g}(\mathcal{E}',\mathcal{F})\cong \operatorname{Hom}_{g\circ f}(\mathcal{E},\mathcal{F})
$$

for any morphism  $q: T \to U$  in S and for every  $\mathcal{F} \in \mathcal{D}_U$ .

The functor p is called a Grothendieck opfibration if for any  $f : S \to T$  and for every object  $\mathcal E$ in  $\mathcal{D}_S$  there exists a coCartesian morphism  $\mathcal{E} \to \mathcal{E}'$ .

**A.1.3.** The functor p is a Grothendieck opfibration if and only if  $p^{op} : \mathcal{D}^{op} \to \mathcal{S}^{op}$  is a Grothendieck fibration. We say that  $p$  is a **bifibration** if is a fibration and an opfibration at the same time. If  $p: \mathcal{D} \to \mathcal{S}$  is a Grothendieck fibration we may choose an associated pseudo-functor, i.e. to each  $S \in \mathcal{S}$  we associate the category  $\mathcal{D}_S$ , and to each  $f : S \to T$  we associate a push-forward functor

$$
f_{\bullet}: \mathcal{D}_S \to \mathcal{D}_T
$$

such that for each  $\mathcal E$  in  $\mathcal D_S$  there is a Cartesian morphism  $\mathcal E \to f_*\mathcal E$ . The same holds similarly for an opfibration with the pull-back  $f^{\bullet}$  instead of the push-forward. If the functor p is a bifibration,  $f_{\bullet}$  is left adjoint to  $f^{\bullet}$ . Situations where this is the opposite can be modeled by considering bifibrations  $\mathcal{D} \rightarrow \mathcal{S}^{\text{op}}$ .

#### A.2 Fibered multicategories and the six functors

A.2.1. We give a definition of a (op-)fibered multicategory. This is a straightforward generalization of the notion of (op-)fibered category given in the section [A.1.](#page-75-0) It is very useful to encode the formalism of the Grothendieck six functors. Details about (op-)fibered multicategories can be found, for instance, in [\[12](#page-86-0), [13](#page-86-1)].

The reader should keep in mind that a multicategory abstracts the properties of multilinear maps, and indeed every monoidal category gives rise to a multicategory setting

<span id="page-75-1"></span>
$$
\text{Hom}(A_1, \dots, A_n; B) \coloneqq \text{Hom}((A_1 \otimes (A_2 \otimes (\dots))), B). \tag{14}
$$

Definition A.2.2. A multicategory  $D$  consists of

- a class of objects  $Ob(\mathcal{D})$ ;
- for every  $n \in \mathbb{Z}_{\geq 0}$ , and objects  $X_1, \ldots, X_n, Y$  a class

 $\text{Hom}(X_1,\ldots,X_n;Y);$ 

• a composition law, i.e. for objects  $X_1, \ldots, X_n, Y_1, \ldots, Y_m, Z$  and for each integer  $1 \leq i \leq m$  a map:

 $Hom(X_1, ..., X_n; Y_i) \times Hom(Y_1, ..., Y_m; Z) \rightarrow Hom(Y_1, ..., Y_{i-1}, X_1, ..., X_n, Y_{i+1}, ..., Y_m; Z);$ 

• for each object  $X \in Ob(\mathcal{D})$  an identity  $id_X \in Hom(X; X)$ ;

satisfying associativity and identity laws. A symmetric (braided) multicategory is given by an action of the symmetric (braid) groups, i.e. isomorphisms

 $\alpha: \text{Hom}(X_1,\ldots,X_n;Y) \to \text{Hom}(X_{\alpha(1)},\ldots,X_{\alpha(n)};Y)$ 

for  $\alpha \in S_n$  (resp.  $\alpha \in B_n$ ) forming an action which is compatible with composition in the obvious way (substitution of strings in the braid group).

In some references the composition is defined in a seemingly more general way; in the presence of identities these descriptions are, however, equivalent. We denote a multimorphism in  $f \in \text{Hom}(X_1, \ldots, X_n; Y)$  also by



for  $n \geq 1$ , or by



for  $n = 0$ .

We will also need the definition of a strict 2-multicategory which is a multicategory enriched in (usual) categories:

**Definition A.2.3.** A (strict) 2-multicategory  $D$  consists of

- a class of objects  $Ob(\mathcal{D})$ ;
- for every  $n \in \mathbb{Z}_{\geq 0}$ , and objects  $X_1, \ldots, X_n, Y$  a category

 $\text{Hom}(X_1, \ldots, X_n; Y);$ 

• a composition, i.e. for objects  $X_1, \ldots, X_n$ ,  $Y_1, \ldots, Y_m$ , Z and for each integer  $1 \leq i \leq m$  a functor:

Hom
$$
(X_1,...,X_n; Y_i) \times \text{Hom}(Y_1,...,Y_m; Z) \to \text{Hom}(Y_1,...,Y_{i-1},X_1,...,X_n,Y_{i+1},...,Y_m; Z);
$$

• for each object  $X \in Ob(\mathcal{D})$  an identity object  $\mathrm{id}_X$  in the category  $\mathrm{Hom}(X;X)$ ;

satisfying strict associativity and identity laws. A symmetric (braided) 2-multicategory is given by an action of the symmetric (braid) groups, i.e. isomorphisms of categories

$$
\alpha: \text{Hom}(X_1, \ldots, X_n; Y) \to \text{Hom}(X_{\alpha(1)}, \ldots, X_{\alpha(n)}; Y)
$$

for  $\alpha \in S_n$  (resp.  $\alpha \in B_n$ ) forming an action which is strictly compatible with composition in the obvious way (substitution of strings in the braid group).

The 1-composition of 2-morphisms is (as for usual 2-categories) determined by the following whiskering operations: Let  $f, g \in \text{Hom}(X_1, \ldots, X_n; Y_i)$  be 1-morphisms and let  $h \in \text{Hom}(Y_1, \ldots, Y_m; Z)$  be a 1-morphism and let  $\mu : f \Rightarrow g$  be a 2-morphism in Mor(Hom( $X_1, \ldots, X_n; Y_i$ )). Then we define

$$
h\ast\mu\coloneqq\mathrm{id}_h\!\cdot\!\mu
$$

where the right hand side is the image of the morphism  $id_h \times \mu$  under the composition functor. Similarly we define  $\mu * h$  for  $\mu : f \Rightarrow g$  with  $f, g \in Hom(Y_1, \ldots, Y_m; Z)$  and  $h \in Hom(X_1, \ldots, X_n; Y_i)$ .

A.2.4. We leave it to the reader to state the obvious definition of a functor between multicategories. Similarly there is a definition of a opmulticategory, in which we have classes

$$
\mathrm{Hom}(X;Y_1,\ldots,Y_n)
$$

and similar data. For a multicategory  $\mathcal D$  we get a natural opmulticategory  $\mathcal D^{op}$  by reversing the arrows.

The trivial category  $\{\cdot\}$  is considered as a multicategory setting all Hom $(\cdot, \ldots, \cdot; \cdot)$  to the 1-element set. It is the final object in the "category" of multicategories.

To clarify the precise relation between multicategories and monoidal categories we have to define Cartesian and coCartesian morphisms. It turns out that we can actually give a definition which is a common generalization of coCartesian morphisms in opfibered categories and the morphisms expressing the existence of a tensor product:

<span id="page-77-0"></span>**Definition A.2.5.** Consider a functor of multicategories  $p : D \rightarrow S$ . We call a morphism

$$
\xi \in \text{Hom}(X_1,\ldots,X_n;Y)
$$

in D coCartesian w.r.t. p, if for all  $Y_1, \ldots, Y_m, Z$  with  $Y_i = Y$ , and for all

$$
f \in \mathrm{Hom}(p(Y_1), \ldots, p(Y_m); p(Z))
$$

the map

$$
Hom_f(Y_1, \ldots, Y_m; Z) \rightarrow Hom_{f \circ p(\xi)}(Y_1, \ldots, Y_{i-1}, X_1, \ldots, X_n, Y_{i+1}, \ldots, Y_m; Z)
$$
  

$$
\alpha \mapsto \alpha \circ \xi
$$

is bijective. We call a morphism

$$
\xi \in \mathrm{Hom}(X_1,\ldots,X_n;Y)
$$

in D Cartesian w.r.t. p at the i-th slot, if for all  $Y_1, \ldots, Y_m$ , and for all  $f \in \text{Hom}(p(Y_1), \ldots, p(Y_m); p(X_i))$ the map

$$
\operatorname{Hom}_f(Y_1,\ldots,Y_m;X_i)\to \operatorname{Hom}_{p(\xi)\circ f}(X_1,\ldots,X_{i-1},Y_1,\ldots,Y_m,X_{i+1},\ldots,X_n;Z).
$$
  

$$
\alpha\mapsto \xi\circ\alpha
$$

is bijective.

The functor  $p : \mathcal{D} \to \mathcal{S}$  is called an **opfibered multicategory** if for every  $g \in \text{Hom}(S_1, \ldots, S_n; T)$ in S, and for every collection of objects  $X_i$  with  $p(X_i) = S_i$  there is some object Y over T and some coCartesian morphism  $\xi \in \text{Hom}(X_1, \ldots, X_n; Y)$  such that  $p(\xi) = g$ .

The functor  $p : \mathcal{D} \to \mathcal{S}$  is called a fibered multicategory if for every  $1 \leq j \leq n$ , for each g  $\in$  $Hom(S_1, \ldots, S_n; T)$  in S, for every collection of objects  $X_i$  for  $i \neq j$  with  $p(X_i) = S_i$ , and for every Y over T, there is some object  $X_j$  and some Cartesian morphism w.r.t. the j-th slot  $\xi \in$  $Hom(X_1, \ldots, X_n; Y)$  such that  $p(\xi) = g$ .

The functor  $p: \mathcal{D} \to \mathcal{S}$  is called a **bifibered multicategory** if it is both fibered and opfibered. A morphism of (op)fibered multicategories is a commutative diagram of functors



such that F maps (co-)Cartesian morphisms to (co-)Cartesian morphisms.

<span id="page-78-2"></span>It turns out that the composition of Cartesian morphisms is Cartesian (and similarly for coCartesian morphisms if they are composed w.r.t. the right  $\text{slot}$ <sup>[9](#page-78-0)</sup>.

**Lemma A.2.6.** 1. An opfibered multicategory  $p : \mathcal{D} \to \{\cdot\}$  is a monoidal category defining  $X \otimes Y$ to be the target of a coCartesian arrow from the pair X, Y over the unique map in Hom $(\cdot, \cdot; \cdot)$ of the final multicategory  $\{\cdot\}.$ 

Conversely any monoidal category gives rise to an opfibered multicategory  $p : \mathcal{D} \to \{\cdot\}$  via [\(14\)](#page-75-1). A multicategory  $\mathcal D$  is a closed category if and only if it is fibered over  $\{\cdot\}$ . In particular, the fibers of an (op)fibered multicategory  $p : \mathcal{D} \to \mathcal{S}$  are always closed/monoidal in the following sense: given any functor of multicategories<sup>[10](#page-78-1)</sup>  $x: {\{\cdot\}} \to S$ , the category  $\mathcal{D}_x$  of objects over x is closed/monoidal.

2. Given (op)fibered multicategories  $p: C \to D$  and  $q: D \to \mathcal{E}$  also the composition  $q \circ p$  is an (op)fibered multicategory. In particular, if we have an opfibered multicategory  $p: \mathcal{C} \to \mathcal{S}$  and if  $S \rightarrow \{\cdot\}$  is opfibered (i.e. S is monoidal) then also  $C \rightarrow \{\cdot\}$  is opfibered (i.e. C is monoidal). The same holds dually. A morphism  $\alpha$  is (co)Cartesian for  $q \circ p$  if and only if  $\alpha$  is (co)Cartesian for p and  $p(\alpha)$  is (co)Cartesian for q.

Similarly, the unit 1 is just the target of a coCartesian morphism in  $Hom(1)$  which exists by definition (the existence is also required for the empty set of objects).

The second part of the lemma encapsulates the distinction between internal and external tensor product in a four (or six) functor context, see [A.2.17.](#page-82-0)

**A.2.7.** Let  $\mathcal{D}, \mathcal{S}$  be (usual) multicategories. More generally any opfibered multicategory  $\mathcal{D} \rightarrow \mathcal{S}$ gives rise to a pseudo-functor of 2-multi-categories

$$
S \to \mathcal{MCAT}^{\text{2-op}}
$$

where  $MCAT$  is the 2-multicategory of categories, whose objects are categories and the morphism categories are defined to be:

$$
\mathrm{Hom}_{\mathcal{MCAT}}(\mathcal{C}_1,\ldots,\mathcal{C}_n;\mathcal{D})\coloneqq \mathrm{Fun}(\mathcal{C}_1\times\cdots\times\mathcal{C}_n,\mathcal{D})
$$

<span id="page-78-1"></span><sup>10</sup>This specifies also morphisms in  $Hom(X, \ldots, X; X)$ , for all *n*, compatible with composition. ´¹¹¹¹¹¹¹¹¹¸¹¹¹¹¹¹¹¹¹¶

n

<span id="page-78-0"></span><sup>9</sup>As with fibered categories there are weaker notions of Cartesian which still uniquely determine a Cartesian morphism (up to isomorphism) from given objects over a given multimorphism, however, do not imply stability under composition. Similarly for coCartesian morphisms.

Here by a pseudo-functor  $\Psi : \mathcal{S} \to \mathcal{T}$ , where  $\mathcal{T}$  is a 2-multicategory, we understand the obvious generalization of the usual concept of a pseudo-functor. This means that for each  $f \in Hom_{\mathcal{S}}(S_1, \ldots, S_n; T)$ we are given a functor  $\Psi(f) \in \text{Hom}(\Psi(S_1), \ldots, \Psi(S_n); T)$  and for each composition  $g \cdot f$  a natural isomorphism

<span id="page-79-0"></span>
$$
\Psi_{f,g} : \Psi(g)\Psi(f) \Rightarrow \Psi(g \cdot f) \tag{15}
$$

satisfying the usual relation for composable morphisms  $f, g$  and  $h$ :

$$
(\Psi(h) * \Psi_{f,g}) \Psi_{gf,h} = (\Psi_{g,h} * \Psi(f)) \Psi_{f,hg}.
$$

This definition generalizes readily to the case in which also  $S$  is a 2-multicategory, the only modification being that, on morphisms, we are given functors

$$
\text{Hom}_{\mathcal{S}}(S_1,\ldots,S_n;T)\to\text{Hom}_{\mathcal{T}}(\Psi(S_1),\ldots,\Psi(S_n); \Psi(T))
$$

and the 2-morphisms  $(15)$  have to be functorial in f and g.

A.2.8. Translated back to the language of fibrations we arrive at the following definition: see [A.2.9.](#page-79-1) First note that the definition of coCartesian morphism (cf. [A.2.5\)](#page-77-0) may be stated in the following way: A morphism

$$
\xi \in \text{Hom}(X_1,\ldots,X_n;Y)
$$

in D coCartesian w.r.t. p, if for all  $Y_1, \ldots, Y_m, Z$  with  $Y_i = Y$  the diagram of sets

$$
\text{Hom}(Y_1,\ldots,Y_m;Z) \longrightarrow \text{Hom}(Y_1,\ldots,Y_{i-1},X_1,\ldots,X_n,Y_{i+1},\ldots,Y_m;Z)
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$
\n
$$
\text{Hom}(p(Y_1),\ldots,p(Y_m);p(Z)) \xrightarrow{\circ p(\xi)} \text{Hom}(p(Y_1),\ldots,p(Y_{i-1}),p(X_1),\ldots,p(X_n),p(Y_{i+1}),\ldots,p(Y_m);p(Z))
$$

<span id="page-79-1"></span>is Cartesian.

**Definition A.2.9.** Let  $p : \mathcal{D} \to \mathcal{S}$  be a strict functor of 2-multicategories. A 1-morphism

$$
\xi \in \mathrm{Hom}_f(\mathcal{E}_1,\ldots,\mathcal{E}_n;\mathcal{F})
$$

in D over  $f \in \text{Hom}(S_1, \ldots, S_n; T)$  is called coCartesian w.r.t. p, if for all  $\mathcal{F}_1, \ldots, \mathcal{F}_m, \mathcal{G}$  with  $\mathcal{F}_i = \mathcal{F}$  the diagram of categories

$$
\text{Hom}(\mathcal{F}_1, \dots, \mathcal{F}_m; \mathcal{G}) \xrightarrow{\circ \xi} \text{Hom}(\mathcal{F}_1, \dots, \mathcal{F}_{i-1}, \mathcal{E}_1, \dots, \mathcal{E}_n, \mathcal{F}_{i+1}, \dots, \mathcal{F}_m; \mathcal{G})
$$
\n
$$
\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow
$$
\n
$$
\text{Hom}(T_1, \dots, T_m; U) \xrightarrow{\circ p(\xi)} \text{Hom}(T_1, \dots, T_{i-1}, S_1, \dots, S_n, T_{i+1}, \dots, T_m; U)
$$

is Cartesian (where we set  $T_k := p(\mathcal{F}_k)$  and  $U := p(\mathcal{G})$ ).

The strict functor p is called a 2-opfibered 1-opfibered multicategory (with 1-categorical **fibers)** if for all  $f \in \text{Hom}(S_1, \ldots, S_n; T)$  and objects  $\mathcal{E}_1, \ldots, \mathcal{E}_n$  with  $p(\mathcal{E}_i) = S_i$  there is a coCartesian 1-morphism with domains  $\mathcal{E}_1, \ldots, \mathcal{E}_n$ . Furthermore the functors

$$
\operatorname{Hom}(\mathcal{E}_1,\ldots,\mathcal{E}_n;\mathcal{F})\to \operatorname{Hom}(p(\mathcal{E}_1),\ldots,p(\mathcal{E}_n);p(\mathcal{F}))
$$

have to be Grothendieck opfibrations (with discrete fibers) and composition has to be a morphism of Grothendieck opfibrations.

The functor  $p: \mathcal{D} \to \mathcal{S}$  is called a 2-fibered 1-opfibered multicategory (with 1-categorical fibers) if for every  $1 \leq j \leq n$  and for each  $g \in \text{Hom}(S_1, \ldots, S_n; T)$  in S, and for each collection of objects  $\mathcal{E}_i$  for  $i \neq j$  with  $p(\mathcal{E}_i) = S_i$ , and for each  $\mathcal F$  over  $T$ , there is some object  $\mathcal{E}_j$  and some Cartesian 1-morphism w.r.t. the j-th slot  $\xi \in \text{Hom}(\mathcal{E}_1, \ldots, \mathcal{E}_n; \mathcal{F})$  with  $p(\xi) = g$ . Furthermore the functors

 $\text{Hom}(\mathcal{E}_1,\ldots,\mathcal{E}_n;\mathcal{F}) \to \text{Hom}(p(\mathcal{E}_1),\ldots,p(\mathcal{E}_n);p(\mathcal{F}))$ 

have to be Grothendieck opfibrations (with discrete fibers) and composition has to be a morphism of Grothendieck opfibrations.

There are several other, partly more general, definitions of an (op-)fibration with 2-categorical fibers which we will not need in this section. We will discuss them in a subsequent article [\[14\]](#page-86-2). Note that for (op-)fibrations with 1-categorical fibers the composition is automatically a morphism of Grothendieck opfibrations.

**A.2.10.** An opfibration  $p : \mathcal{D} \to \mathcal{S}$  of 2-multicategories with 1-categorical fibers is in particular (forgetting 2-morphisms) a usual opfibration. The additional datum, which makes it into a 2 opfibration is the following: For each 2-morphism  $\mu : f \Rightarrow g$  in S a map of sets (the 2-push-forward):

$$
p^*(\mu): \operatorname{Hom}_f(X_1, \ldots, X_n; Y) \to \operatorname{Hom}_g(X_1, \ldots, X_n; Y)
$$

such that

$$
p^*(id_f)(\beta) = \beta
$$
  

$$
p^*(\mu) \circ p^*(\nu) = p^*(\mu \circ \nu)
$$

(composition of 2-coCartesian morphisms are 2-coCartesian) and

$$
p^*(p(\alpha) * \mu)(\alpha \circ \xi) = \alpha \circ (p^*(\mu)(\xi))
$$
  

$$
p^*(\mu * p(\alpha))(\xi \circ \alpha) = (p^*(\mu)(\xi)) \circ \alpha
$$

(1-composition maps coCartesian morphisms to coCartesian morphisms).

The 2-morphisms between  $\alpha$  and  $\beta$  in  $\mathcal D$  lying over f, resp. g in S can be reconstructed from the datum  $p^*$  as

$$
\mathrm{Hom}(\alpha,\beta)=\{\mu\in\mathrm{Hom}(f,g)\mid p^*(\mu)(\alpha)=\beta\}.
$$

A.2.11. With a pseudo-functor

$$
\Psi: \mathcal{S} \rightarrow \mathcal{MCAT}^{\mathrm{2-op}}
$$

where  $S$  is any strict 2-multicategory, we associate the opfibration

$$
\mathcal{D}_{\Psi} \to \mathcal{S}.
$$

The objects of  $\mathcal{D}_{\Psi}$  are pairs

$$
(S,X\in\Psi(S))
$$

in which  $S$  is an object of  $S$ . The 1-morphisms



are pairs of (multi-)morphisms

$$
f \in \text{Hom}(S_1, \ldots, S_n; T) \qquad \alpha : \Psi(f)(X_1, \ldots, X_n) \to Y.
$$

The 2-morphisms

$$
(f,\alpha) \Rightarrow (f',\alpha')
$$

are given by 2-morphisms  $\mu : f \Rightarrow f'$  such that  $\alpha \circ (\Psi(\mu)(X)) = \alpha'$ . The fiber<sup>[11](#page-81-0)</sup> of  $\mathcal{D}_{\Psi} \to \mathcal{S}$  over S is actually a 1-category, namely precisely the category  $\Psi(S)$ .

<span id="page-81-3"></span>We have the following generalization of Lemma [A.2.6,](#page-78-2) 2.:

**Lemma A.2.12.** Given (op)fibered 2-multicategories  $p : C \to D$  and  $q : D \to \mathcal{E}$  then the composition  $q \circ p$  is an (op)fibered 2-multicategory, too. A 1-morphism  $\alpha$  is (co)Cartesian for  $q \circ p$  if and only if  $\alpha$  is (co)Cartesian for p and  $p(\alpha)$  is (co)Cartesian for q.

<span id="page-81-1"></span>**Example A.2.13.** Let S be a usual category. Then S may be turned into a symmetric multicategory by setting

$$
\operatorname{Hom}(X_1,\ldots,X_n;Y)\coloneqq \operatorname{Hom}(X_1;Y)\times\cdots\times\operatorname{Hom}(X_n;Y).
$$

If S has coproducts, then S (with this multi-category structure) is opfibered over  $\{\cdot\}$ . Let  $p : \mathcal{D} \to \mathcal{S}$ be an opfibered (usual) category. Any object  $X$  induces a canonical functor of multicategories  $x: \{\cdot\} \to S$  with image X, hence the fibers of an opfibered multicategory  $p: \mathcal{D} \to S$  are monoidal and the datum p is equivalent to giving a monoidal structure on the fibers such that the push-forwards  $f_{\bullet}$  are monoidal functors and such that the compatibility morphisms between them are morphisms of monoidal functors. This is called a covariant monoidal pseudo-functor in [\[17,](#page-86-3) (3.6.7)].

<span id="page-81-2"></span>**Example A.2.14.** Let S be a usual category. Then  $S^{op}$  may be turned into a symmetric multicategory (or equivalently S into a symmetric opmulticategory) by setting

$$
\operatorname{Hom}(X_1,\ldots,X_n;Y)\coloneqq \operatorname{Hom}(Y;X_1)\times\cdots\times\operatorname{Hom}(Y;X_n).
$$

If S has products then  $S^{op}$  (with this multi-category structure) is opfibered over  $\{\cdot\}$ . Let  $p : \mathcal{D} \to$  $\mathcal{S}^{\mathrm{op}}$  be an opfibered (usual) category. Then an opfibered multicategory structure on p, w.r.t. this multicategory structure on  $S^{op}$ , is equivalent to a monoidal structure on the fibers such that pullbacks  $f^*$  (along morphisms in S) are monoidal functors and such that the compatibility morphisms between them are morphisms of monoidal functors. This is called a contravariant monoidal pseudo-functor in [\[17,](#page-86-3) (3.6.7)].

Definition A.2.15. The point is that the notion of (op)fibered multicategory is not restricted to the situation of Examples [A.2.13](#page-81-1) and [A.2.14.](#page-81-2) Let  $S$  be a category with fiber products and define  $\mathcal{S}^{\text{cor}}$ , denoted the symmetric 2-multicategory of correspondences in S to be the symmetric 2-multicategory having the same objects as  $S$ , and where the category of morphisms  $Hom(S_1, \ldots, S_n; T)$  is the category of objects



<span id="page-81-0"></span><sup>&</sup>lt;sup>11</sup>i.e. the 2-category of those objects, morphisms, and 2-morphisms which  $\Psi$  maps to S, id<sub>S</sub>, and id<sub>ids</sub>, respectively

and where the 2-morphisms  $(A, f, g_1, \ldots, g_n) \Rightarrow (A', f', g'_1, \ldots, g'_n)$  are isomorphisms  $A \rightarrow A'$  compatible with  $f, f'$  and  $g_1, g'_1, \ldots, g_n, g'_n$ .

Composition is given by:



where strictly associative fiber products have been chosen in  $S$ . This 2-multicategory is representable (i.e. opfibered over {⋅}), closed (i.e. fibered over {⋅}) and selfdual, with tensor product and internal hom both given by  $\times$  and having as unit the final object of S.

**Definition A.2.16.** Let S be a category with fiber products. A (symmetric) Grothendieck six-functor-formalism on S is a bifibered (symmetric) 2-multicategory with 1-categorical fibers

$$
p:\mathcal{D}\to \mathcal{S}^\mathrm{cor}.
$$

<span id="page-82-0"></span>**A.2.17.** We have a morphism of opfibered (over  $\{\cdot\}$ ) symmetric multicategories  $S^{\text{op}} \to S^{\text{cor}}$  where  $S^{op}$  is equipped with the symmetric multicategory structure as in [A.2.14.](#page-81-2) However there is no reasonable morphism of opfibered multicategories  $S \rightarrow S^{\text{cor}}$ . (There is no compatibility involving only '⊗' and '!'.) From a Grothendieck six-functor-formalism we get operations  $g_*$ ,  $g^*$  as the pull-back and the push-forward along the correspondence



We get  $f'$  and  $f_!$  as the pull-back and the push-forward along the correspondence



We get the monoidal product  $A \otimes B$  for objects  $A, B$  above X as the target of any Cartesian morphism ⊗ over the correspondence



Alternatively, we have

$$
A\otimes B=\Delta^*(A\boxtimes B)
$$

where  $\Delta^*$  is the push-forward along the correspondence



induced by the canonical  $\xi_X \in \text{Hom}(X, X; X)$ , and where  $\boxtimes$  is the absolute monoidal product which exists because by Lemma [A.2.12](#page-81-3) the composition  $\mathcal{D} \to {\{\cdot\}}$  is opfibered, too, i.e.  $\mathcal D$  is monoidal.

**A.2.18.** It is easy to derive from the definition of bifibered multicategory over  $\mathcal{S}^{\text{cor}}$  that the absolute monoidal product  $A \boxtimes B$  can be reconstructed from the fiber-wise product as  $\text{pr}_1^* A \otimes \text{pr}_2^* B$ on  $X\times Y$  , whereas the absolute  $\mathbf{HOM}(A,B)$  is given by  $\mathcal{HOM}(\mathrm{pr}_1^*A,\mathrm{pr}_2^!B)$  on  $X\times Y.$  In particular  $DA = HOM(A, 1)$  is given by  $HOM(A, \pi^1)$  for  $\pi : X \to \cdot$  being the final morphism.

**Lemma A.2.19.** Given a Grothendieck six-functor-formalism on  $S$ 

$$
p: \mathcal{D} \to \mathcal{S}^\mathrm{cor}
$$

for the six operations as extracted in  $A.2.17$  there exist naturally the following compatibility isomophisms:



Here f, g, F, G are morphisms in S which, in the  $(!,*)$ -row, are related by the Cartesian diagram



**Remark A.2.20.** In the right column the corresponding adjoint natural transformations are listed. In each case the left hand side natural isomorphism determines the right hand side one and conversely. (In the  $(\otimes,!)$ -case there are 2 versions of the commutation between the right adjoints; in this case any of the three isomorphisms determines the other two.) The  $(!,*)$ -isomorphism (between left adjoints) is called **base change**, the  $(\otimes,!)$ -isomorphism is called the **projection for**mula, and the  $(*, \otimes)$ -isomorphism is usually part of the definition of a **monoidal functor**. The (⊗, ⊗)-isomorphism is the associativity of the tensor product and part of the definition of a monoidal category. The  $(*,*)$ -isomorphism, and the  $(!,!)$ -isomorphism express that the corresponding functors arrange as a pseudo-functor with values in categories.

Proof. The existence of all isomorphisms is a consequences of the fact that the composition of coCartesian morphisms is coCartesian. For example, the projection formula  $(\otimes,!)$  is derived from the following composition in  $\mathcal{S}^{\text{cor}}$ :



where  $\circ_1$  means that we compose w.r.t. the first slot. The "monoidality of  $f^{**}$  (\*, ⊗) is derived from the following composition in  $\mathcal{S}^{\text{cor}}$ :



Base change  $(l, *)$  is derived from:



 $\Box$ 

All compatibilities between these isomorphisms can be derived, too. Each of these compatibilities corresponds to an associativity relation in the fibered multicategory. One can also axiomatize the properties of the morphism  $f_! \rightarrow f_*$  that often accompanies a six-functor-formalism. Can one give a finite list of compatibility diagrams from which all the others would follow?

A.2.21. The goal and motivation for this research is, as said in the introduction, to define (and to construct in reasonable contexts) a derivator version of a Grothendieck six-functor-formalism, i.e. a fibered multiderivator

 $\mathbb{D} \to \mathbb{S}^{\mathrm{cor}}$ 

where  $\mathbb{S}^{\text{cor}}$  is the pre-2-multiderivator associated with the 2-category  $\mathcal{S}^{\text{cor}}$ . We will give the definition of a pre-2-multiderivator and of a fibered derivator over such in a subsequent article [\[14](#page-86-2)].

#### A.3 Localization of multicategories

**Proposition A.3.1.** Let  $D$  be a (symmetric, braided) multicategory and let  $W$  be a subclass of 1-ary morphisms. Then there exists a (symmetric, braided) multicategory  $\mathcal{D}[\mathcal{W}^{-1}]$ , which is not necessarily locally small, together with a functor  $\iota : \mathcal{D} \to \mathcal{D}[\mathcal{W}^{-1}]$  of (symmetric, braided) multicategories with the property that  $\iota(w)$  is an isomorphism for all  $w \in \mathcal{W}$  and which is universal w.r.t. this property.

Proof. This construction is completely analogous to the construction for usual categories. Morphisms  $Hom(X_1, \ldots, X_n; Y)$  are formal compositions of *i*-ary morphisms in D and formal inverses of morphisms in  $W$ , for example:



More precisely: Morphisms are defined to be the class of lists of  $n_i$ -ary morphisms  $f_i \in Hom(X_{i,1}, \ldots, X_{i,n_i}; Y_i)$ , morphisms  $w_i: Y'_i \to Y_i$  in W and integers  $k_i$  as follows

$$
(f_1, w_1), k_1, (f_2, w_2), k_2, \ldots, k_{n-1}, (f_n, w_n)
$$

such that  $Y_i' = X_{i+1,k_i}$ , modulo relations coming from commutative squares and forcing the  $(id, w_i)$  $\Box$ to become the left and right inverse of  $(w_i, id)$ .

# References

- $[1]$  Théorie des topos et cohomologie étale des schémas. Tome 2. Lecture Notes in Mathematics, Vol. 270. Springer-Verlag, Berlin-New York, 1972. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat.
- [2] Théorie des topos et cohomologie étale des schémas. Tome 3. Lecture Notes in Mathematics, Vol. 305. Springer-Verlag, Berlin-New York, 1973. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de P. Deligne et B. Saint-Donat.
- [3] J. Bernstein and V. Lunts. Equivariant sheaves and functors, volume 1578 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1994.
- [4] D. C. Cisinski. Images directes cohomologiques dans les catégories de modèles. Ann. Math. Blaise Pascal, 10(2):195–244, 2003.
- [5] D. C. Cisinski. Le localisateur fondamental minimal. *Cah. Topol. Géom. Différ. Catég.*, 45 (2):109–140, 2004.
- [6] D. C. Cisinski. Propriétés universelles et extensions de Kan dérivées. *Theory Appl. Categ.*, 20: No. 17, 605–649, 2008.
- [7] M. Groth. Monoidal derivators and additive derivators. arXiv:1203.5071, 2012.
- [8] M. Groth. Derivators, pointed derivators and stable derivators. Algebr. Geom. Topol., 13(1): 313–374, 2013.
- [9] M. Groth and J. Stovicek. Tilting theory via stable homotopy theory. arXiv:1401.6451, 2012.
- [10] M. Groth, K. Ponto, and M. Shulman. Mayer-Vietoris sequences in stable derivators. arXiv:1306.2072, 2013.
- [11] A. Grothendieck. Les dérivateurs. Edited by M. Künzer, J. Malgoire and G. Maltsiniotis, <https://www.imj-prg.fr/~georges.maltsiniotis/groth/Derivateurs.html>, 1991.
- <span id="page-86-0"></span>[12] C. Hermida. Representable multicategories. Adv. Math., 151(2):164–225, 2000.
- <span id="page-86-1"></span>[13] C. Hermida. Fibrations for abstract multicategories. In Galois theory, Hopf algebras, and semiabelian categories, volume 43 of Fields Inst. Commun., pages 281–293. 2004.
- <span id="page-86-2"></span>[14] F. Hörmann. Fibered multiderivators and Grothendieck's six functors. In preparation, 2015.
- [15] M. Hovey. Model categories, volume 63 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1999.
- [16] H. Krause. Localization theory for triangulated categories. In Triangulated categories, volume 375 of *London Math. Soc. Lecture Note Ser.*, pages 161–235. Cambridge Univ. Press, Cambridge, 2010.
- <span id="page-86-3"></span>[17] J. Lipman and M. Hashimoto. Foundations of Grothendieck duality for diagrams of schemes, volume 1960 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2009.
- [18] I. Moerdijk and I. Weiss. Dendroidal sets. Algebr. Geom. Topol., 7:1441–1470, 2007.
- [19] A. Neeman. Triangulated categories, volume 148 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2001.