

SN 2011A: AN LOW-LUMINOSITY INTERACTING TRANSIENT, WITH A DOUBLE PLATEAU AND STRONG SODIUM ABSORPTION¹

T. de Jaeger^{2,3*}, J. P. Anderson⁴, G. Pignata^{5,2}, M. Hamuy^{3,2}, E. Kankare⁶, M. D. Stritzinger⁷, S. Benetti⁸, F. Bufano^{2,5}, N. Elias-Rosa^{8,9}, G. Folatelli^{10,11}, F. Förster², S. González-Gaitán^{2,3}, C.P. Gutiérrez^{2,3,4}, C. Inserra⁶, R. Kotak⁶, P. Lira³, N. Morrell¹², F. Taddia¹³, L. Tomasella⁸

(2) Millennium Institute of Astrophysics, Casilla 36-D, Santiago, Chile

(3) Departamento de Astronomía – Universidad de Chile, Camino el Observatorio 1515, Santiago, Chile.

(4) European Southern Observatory, Alonso de Cordova 3107, Vitacura, Santiago, Chile

(5) Departamento de Ciencias Físicas – Universidad Andres Bello, Avda. República 252, Santiago, Chile.

(6) Astrophysics Research Center, School of Mathematics and Physics, Queens University Belfast, Belfast, BT7 1NN, UK.

(7) Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, DK-8000 Aarhus C, Denmark.

(8) INAF-Osservatorio Astronomico di Padova, vicolo dell'Osservatorio 5, I-35122, Padova, Italia.

(9) Institut de Ciències de l'Espai (IEEC-CSIC), Facultat de Ciències, Campus UAB, 08193 Bellaterra, Spain.

(10) Instituto de Astrofísica de La Plata, CONICET, Paseo del Bosque S/N, B1900FWA, La Plata, Argentina.

(11) Kavli Institute for the Physics and Mathematics of the Universe (WPI), Todai Institutes for Advanced Study, the University of Tokyo, Kashiwa, Japan 277-8583 (Kavli IPMU, WPI).

(12) Las Campanas Observatory, Carnegie Observatories, Casilla 601, La Serena, Chile.

(13) The Oskar Klein Centre, Department of Astronomy, Stockholm University, AlbaNova, 10691 Stockholm, Sweden.

Accepted for publication in ApJ

ABSTRACT

We present optical photometry and spectroscopy of the optical transient SN 2011A. Our data spans 140 days after discovery including $BVRiug'r'i'z'$ photometry and 11 epochs of optical spectroscopy. Originally classified as a type II_n supernova (SN II_n) due to the presence of narrow H α emission, this object shows exceptional characteristics. Firstly, the light curve shows a double plateau; a property only observed before in the impostor SN 1997bs. Secondly, SN 2011A has a very low luminosity ($M_V = -15.72$), placing it between normal luminous SNe II_n and SN impostors. Thirdly, SN 2011A shows low velocity and high equivalent width absorption close to the sodium doublet, which increases with time and is most likely of circumstellar origin. This evolution is also accompanied by a change of line profile; when the absorption becomes stronger, a P-Cygni profile appears. We discuss SN 2011A in the context of interacting SNe II_n and SN impostors, which appears to confirm the uniqueness of this transient. While we favour an impostor origin for SN 2011A, we highlight the difficulty in differentiating between terminal and non-terminal interacting transients.

Subject headings: supernovae: general; supernovae: individual: 2011A; stars: mass loss, circumstellar matter

1. INTRODUCTION

It is generally accepted that the majority of massive stars with zero-age main-sequence mass $\geq 8 M_\odot$ (Smartt

2009) end their lives as core-collapse supernovae (CC SNe), with the possibility that some directly form black holes with no visible supernovae (Smartt 2015). CC SNe are classified in two groups according to the absence (SNe Ib/c) or presence (SNe II) of H I lines (Filippenko 1997). The ejected material from a SN explosion is sometimes observed to interact with surrounding circumstellar material (CSM), related to progenitor mass-loss episodes prior to explosion (Chevalier 1981; Fransson 1982).

When the CSM is sufficiently dense, strong CSM-ejecta interaction can begin shortly after explosion; this is often observed as a type II_n supernova (SN II_n henceforth) (Schlegel 1990; Chugai & Danziger 1994). These objects are characterized by prevalent blue continua and narrow emission lines in their spectra superimposed on broader emission profiles. The H α emission typically dominates the spectrum and often has multiple components consisting of (e.g., Salamanca et al. 1998):

1. narrow emission ($v_{FWHM} \sim$ few hundred km s⁻¹) formed by the photoionization of the high density pre-existing CSM from the prompt emission of the SN. The high density has furthermore been attributed to a high mass-loss rate (Chugai et al. 2004).

*dthomas@das.uchile.cl

¹This paper includes data obtained with the 6.5-m Magellan Telescopes and du Pont telescope; the Gemini-North Telescope, Mauna Kea, USA (Gemini Program GN-2010B-Q67, PI: Stritzinger); the PROMPT telescopes at Cerro Tololo Inter-American Observatory in Chile; with the Liverpool Telescope operated on the island of La Palma by Liverpool John Moores University in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias with financial support from the UK Science and Technology Facilities Council; based on observations made with the Nordic Optical Telescope, operated by the Nordic Optical Telescope Scientific Association at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofísica de Canarias; the NTT from ESO Science Archive Facility under allocations 184.D-1151 and 184.D-1140 (PI: Benetti. S), at the Centro Astronomico Hispano Alemn (CAHA) at Calar Alto, operated jointly by the Max-Planck Institut für Astronomie and the Instituto de Astrofísica de Andalucía (CSIC), on observations collected at Asiago Observatory and the Southern Astrophysical Research (SOAR) telescope, which is a joint project of the Ministério da Ciência, Tecnologia, e Inovação (MCTI) da República Federativa do Brasil, the U.S. National Optical Astronomy Observatory (NOAO), the University of North Carolina at Chapel Hill (UNC), and Michigan State University (MSU).

2. intermediate emission ($v_{FWHM} \sim$ few thousand km s^{-1}) possibly produced from the shock interaction of the SN blast within a dense shell of clumpy CSM or with a dense equatorial wind (Chugai & Danziger 1994).
3. broad emission ($v_{FWHM} \geq 10^4 \text{ km s}^{-1}$) most likely due to SN ejecta as a result of multiple scattering by thermal electrons in the opaque CS gas (Chugai 2001). The presence of this broad H α emission without broad P–Cygni absorption is usually considered to be caused by a dense wind (Chugai 1990).

Measured parameters of these components can provide valuable information on progenitor properties and its late time evolution (such as the mass–loss rate or wind velocity), through the observed properties of their CSM.

Observations of SNe IIn published to date (see e.g., Kiewe et al. 2012; Taddia et al. 2013; Zhang et al. 2012) have allowed researchers to probe the immediate environment of these explosions and point towards significant diversity among various SNe IIn. The high mass–loss rates calculated from the observations of various SNe IIn (e.g., Kiewe et al. 2012; Taddia et al. 2013; Moriya et al. 2014) have prompted several authors to assign Luminous Blue Variable stars (LBVs) in several cases as immediate progenitors of SNe IIn. LBVs, as defined by Humphreys & Davidson (1994), are some of the most luminous stars, with episodic and violent mass–loss events with mass–loss rates $\geq 10^{-3} M_{\odot} \text{ yr}^{-1}$. Classic examples of LBVs are Eta Carinae and P–Cygni. The direct detection of the extremely luminous progenitor of SN 2005gl (Gal–Yam & Leonard 2009) also supports this possibility. However, in stellar evolutionary models, the LBV phase follows a blue giant star state (Crowther 2007; Gräfener & Hamann 2008), most likely with mass loss driven by the bi–stability jump (Vink 2011), and it is then followed by a H–poor WN star (nitrogen rich Wolf–Rayet). In this scenario the LBV star loses its H envelope, becomes a Wolf–Rayet star and then explodes as a SN. Thus, LBVs have been placed as a post–main sequence, but not a final pre–SN phase (Schaller et al. 1992; Langer 1993; Stothers & Chin 1996; Maeder et al. 2005; Maeder & Meynet 2008; Dwarkadas et al. 2010), although see Groh et al. (2013) for recent modeling suggesting that some stars could indeed end their lives during this phase.

There have been a number of cases where transients have originally been defined as SNe IIn, but a link to a definitive SN event (i.e. the terminal stage of a star’s life) is questionable. Such events have been defined as SN “impostors” (Van Dyk et al. 2000; Maund et al. 2006) which are believed to be luminous non–terminal eruptions of massive stars instead of SNe. There are multiple cases of such events e.g., the on–going transient SN 2009ip (Pastorello et al. 2012a; Prieto et al. 2013; Mauerhan et al. 2013; Fraser et al. 2013; Margutti et al. 2014; Smith et al. 2014). SN impostors are generally claimed to be LBV transients with peak luminosities lower than SNe ($M \geq -14$), such as SN 1997bs ($M_V \sim -13.8$; Van Dyk et al. 2000), SN 2000ch

($M_R \sim -12.8$; Wagner et al. 2004; Pastorello et al. 2010), SN 2002kg ($M_V \sim -9.6$; Van Dyk et al. 2006; Maund et al. 2006; Weis & Bomans 2005), SN 2007sv ($M_R \sim -14.2$; Tartaglia et al. 2014), SN 2008S ($M_R \sim -13.9$; Kochanek 2011; Smith et al. 2009, and 2008 NGC 300–OT ($M_V \sim -12$ to -13 ; aBond et al. 2009 (see also the compilation paper by Smith et al. 2011). However, unambiguous evidence that all transients classified as “impostors” are actually non–terminal events is lacking. Indeed for the first transient identified as a SN impostor, SN 1997bs, it has been claimed that no surviving star is present (Li et al. 2002; Adams & Kochanek 2015). Adams & Kochanek (2015) have argued that the scenario where the surviving star is obscured by dust created in the eruption is excluded.

Given that SN impostors and SNe IIn show similar spectroscopic features, such as the presence of narrow emission lines (in particular H I lines), the differentiation based solely on spectral analysis is rendered difficult, and constraining the nature of their progenitors (e.g., ZAMS mass) has been problematic. Indeed, the detection of narrow emission lines is merely evidence for high density material close to the SN and it appears that this high density CSM can be produced by a number of different progenitor scenarios. Indeed, it has even been observed in SNe Ia (e.g., SN 2002ic Hamuy et al. 2003, SN 2011kx Dilday et al. 2012).

While observations of individual SNe IIn have been used to argue for LBV progenitors or LBV transients, statistical investigations of their environments within host galaxies have pointed to lower mass progenitors. Anderson et al. (2012) and Haberman et al. (2014) found a low correlation between SNe IIn and host–galaxy H II regions. It was found that SNe IIn show a similar degree of association with star formation (SF) as SNe IIP, which are thought to arise from stars at the lower end of the CC SNe scale; with masses between 8–16 M_{\odot} (Smartt 2009). Therefore, these results suggest that a significant fraction of SNe IIn arise from relatively low mass progenitors. This is somewhat surprising given that the LBV phase is thought to be associated only with very massive stars (~ 30 – $80 M_{\odot}$). Contrary to the above, we note that recent results suggest that a significant fraction of SNe IIn could arise from electron capture SNe (Smith 2013) with AGB progenitor stars (Prieto et al. 2009; Botticella et al. 2009). Also, Smith & Tombleson (2015) argue that LBV stars are the product of binary kicks, due to their isolated environments.

In this paper we present optical photometry and spectroscopy of the transient SN 2011A, originally classified as a SN IIn (Stritzinger et al. 2011). Given the properties of the transient we discuss its characteristics in comparison to other SN IIn and SN impostors with the aim of defining the nature of the transient and constraining its properties. Due to the fact that it is unclear whether SN 2011A was a true SN, we refer to it as “transient” throughout the paper.

The paper is organised as follows. Section 2 contains a description of the observations and in Section 3 we dis-




FIG. 1.— The field of SN 2011A showing the location of sequence stars (designated numerically) listed in Table 1. The location of SN 2011A is marked by a grey circle. North is up and East is left. The image is $10' \times 10'$ and was taken with PROMPT-5 on 2011 January 23 in r' band.

cuss our results, first analysing light- and color-curves and then using spectroscopy to derive physical parameters for the transient. In Section 4 we discuss our results and we conclude with a summary in Section 5.

2. OBSERVATIONS

2.1. Discovery

SN 2011A was discovered on an unfiltered image (apparent magnitude ~ 16.9) on 2011 January 2.30 UT with the 0.41-m ‘PROMPT 4’ telescope located at Cerro Tololo (Pignata et al. 2011) on behalf of the CHASE project (Pignata et al. 2009). The transient was located at $\alpha = 13^{\text{h}}01^{\text{m}}01^{\text{s}}19 \pm 0^{\text{s}}.2$, $\delta = -14^{\circ}31'34''.8 \pm 0''.2$ (J2000.0), $21''.3$ east and $46''.5$ south of center of the galaxy NGC 4902 at $z=0.008916$ (Theureau et al. 2007, see NED²). NGC 4902 is a luminous Sb galaxy ($M_B = -21.40$). Stritzinger et al. (2011) classified the object to be a SNe IIn due to the presence of prevalent narrow Balmer lines in the spectrum. The most recent pre-explosion non detection on archival images was dated 2010 July 13.05 (limiting magnitude 18.5) providing extremely weak constraints on any explosion time or previous variability. All observations are presented with respect to the discovery date: 2011 January 2.30 UT. In Fig. 1 the field of the transient is displayed.

2.2. Photometry

Optical photometry was obtained with the Panchromatic Robotic Optical Monitoring and Polarimetry Telescopes (PROMPT; Reichart et al. 2005) at the Cerro Tololo Inter-American Observatory, RATCam mounted on the 2.0m Liverpool Telescope (Steele et al. 2004) (LT) at the Observatorio del Roque de los Muchachos on the Canary Island of La Palma, with EFOSC2

(Buzzoni et al. 1984) mounted on the 3.6m New Technology Telescope (NTT) at the La Silla Observatory, the 2.2m telescope at the Calar Alto (CA) Observatory with the optical imager/spectrograph CAFOS, and from the imager AFOSC mounted at the 1.82m reflector telescope at the Asiago Observatory. All images were analysed after automated reductions (bias-, flatfield-correction and astrometric solution) had been applied and all the science images were host-galaxy subtracted.

Photometry of the transient was computed relative to a sequence of 19 stars in the field of NGC 4902, which we calibrate using catalogs from Landolt (1992, 2007) for B, V, R, I bands and Smith et al. (2002) for u', g', r', i', z' bands. Instrument-specific colours terms were derived using several epochs of standard field observations. The resulting $B, V, R, I, u', g', r', i', z'$ magnitudes of the local sequence stars are reported in Table 1, which correspond to the averages from n photometric nights ($n=2$ for the z' filter and $n \geq 3$ for the remaining filters), and the uncertainties are the standard deviations from the mean.

Differential photometry of the transient was done relative to the local sequence. To obtain instrumental magnitudes, we used the IRAF³ package SNOOPY⁴. Photometry for SN 2011A is presented in Table 2 and the resulting light curve is displayed in Fig. 2. Errors come from the point spread function (PSF) fit and the local sequence magnitudes.

2.3. Visual-wavelength spectroscopy

Optical spectroscopy of SN 2011A was acquired using the GOODMAN spectrograph at the Southern Astrophysical Research (SOAR) 4.1 meter Telescope, the WFCCD at the du Pont telescope located at Las Campanas Observatory (LCO), the GMOS spectrograph mounted on Gemini-North on the summit of Mauna Kea, EFOSC2 on the 3.6m NTT at La Silla, the ALFOSC imager/spectrograph on the 2.5m Nordic Optical Telescope (NOT) at La Palma and IMACS mounted on the Baade telescope at LCO. In total we obtained 11 optical spectra mostly covering all the optical range, from ~ 3500 – 9000 Å except for three spectra, one covering only the $H\alpha$ emission and the others ~ 4200 – 7000 Å. A log of the optical observations is given in Table 3.

Spectroscopic reductions were performed using standard IRAF routines. All data were debiased, flat-fielded and cleaned of cosmic rays. Extracted 1D spectra were wavelength calibrated using He-Ne-Ar/Hg-Ar/Th-Ar lamps (depending on the instrument) and the calibrations were corrected using bright night-sky emission lines. However for the Gemini spectrum the wavelength calibration was computed before 1D spectrum extraction. Flux calibration was determined

³ IRAF (Image Reduction and Analysis Facility) is distributed by the National Optical Astronomy Observatories, which are operated by AURA, Inc., under cooperative agreement with the National Science Foundation.

⁴ SNOOPY, originally presented in Patat (1996), has been implemented in IRAF by E. Cappellaro.

² <http://ned.ipac.caltech.edu/>

TABLE 1
 $B, V, R, I, u', g', r', i', z'$ LOCAL SEQUENCE STAR MAGNITUDES.

STAR	R.A	Decl	B	V	R	I	u'	g'	r'	i'	z'
1	13:01:01.523	-14:28:21.35	15.78±0.02	15.20±0.04	14.83±0.03	14.46±0.05	16.65±0.02	15.45±0.02	15.05±0.02	14.90±0.04	14.91±0.01
2	13:01:09.706	-14:28:07.32	17.69±0.05	17.06±0.04	16.69±0.03	16.32±0.05	18.57±0.01	17.31±0.03	16.91±0.01	16.74±0.03	16.76±0.02
3	13:01:12.549	-14:29:48.59	17.20±0.02	16.41±0.02	15.95±0.03	15.51±0.04	18.35±0.09	16.75±0.02	16.19±0.02	15.96±0.03	15.94±0.01
4	13:01:13.685	-14:29:56.09	15.68±0.02	14.70±0.03	14.12±0.03	13.65±0.04	17.66±0.05	15.16±0.04	14.38±0.01	14.11±0.03	14.06±0.01
5	13:01:09.906	-14:32:40.66	15.96±0.01	15.10±0.03	14.61±0.02	14.21±0.41	17.60±0.03	15.50±0.04	14.86±0.01	14.65±0.03	14.65±0.01
6	13:01:06.445	-14:34:08.13	13.42±0.01	12.43±0.03	11.87±0.03	11.36±0.03	15.19±0.02	12.88±0.04	12.14±0.01	11.83±0.03	11.74±0.01
7	13:00:54.730	-14:33:11.10	17.33±0.02	16.17±0.02	15.44±0.02	14.84±0.04	19.52±0.10	16.74±0.04	15.72±0.01	15.32±0.03	15.19±0.01
8	13:00:47.126	-14:31:26.73	16.46±0.01	15.55±0.02	14.98±0.02	14.48±0.04	18.27±0.04	15.99±0.03	15.23±0.02	14.94±0.03	14.87±0.02
9	13:00:45.583	-14:31:31.08	17.99±0.03	17.36±0.01	17.00±0.02	16.66±0.04	18.72±0.02	17.62±0.02	17.24±0.02	17.08±0.05	17.08±0.01
10	13:00:42.987	-14:32:14.39	16.06±0.01	15.42±0.03	15.03±0.02	14.64±0.04	16.95±0.02	15.69±0.02	15.26±0.02	15.08±0.03	15.07±0.01
11	13:00:44.202	-14:28:44.77	13.21±0.03	12.73±0.03	12.44±0.03	12.17±0.04	14.21±0.02	12.92±0.03	12.65±0.02	12.58±0.03	12.64±0.01
12	13:00:44.752	-14:27:31.79	12.69±0.04	12.09±0.03	11.72±0.02	11.38±0.03	13.69±0.03	12.36±0.01	11.94±0.02	11.81±0.04	11.83±0.01
13	13:00:53.129	-14:28:40.67	13.90±0.02	13.11±0.03	12.65±0.03	12.20±0.04	15.15±0.02	13.46±0.02	12.90±0.02	12.66±0.03	12.61±0.01
14	13:01:16.297	-14:32:11.47	14.07±0.01	13.48±0.03	13.15±0.03	12.83±0.04	14.99±0.02	13.72±0.02	13.37±0.01	13.25±0.03	13.30±0.01
15	13:01:00.164	-14:35:22.03	15.98±0.01	15.33±0.02	14.93±0.02	14.83±0.04	16.74±0.01	15.60±0.02	15.16±0.02	14.97±0.03	14.96±0.01
16	13:00:52.187	-14:32:20.12	13.95±0.01	13.16±0.02	12.70±0.03	12.31±0.04	15.35±0.02	13.52±0.02	12.94±0.02	12.75±0.03	12.72±0.01
17	13:00:54.651	-14:29:18.73	11.16±0.03	10.38±0.02	10.65±0.03	10.57±0.04	10.39±0.03	9.74±0.06
18	13:00:56.730	-14:29:37.00	19.93±0.05	18.61±0.03	17.74±0.01	17.10±0.06	20.60±1.56	19.19±0.10	18.09±0.01	17.57±0.03	17.35±0.02
19	13:00:51.075	-14:31:37.80	10.84±0.02	10.33±0.03	10.48±0.06	10.58±0.05	10.41±0.01	9.66±0.05

Note. – Errors correspond to the standard deviation from the mean.

TABLE 2
 OPTICAL PHOTOMETRY OF SN 2011A.

JD-2450000+	Epoch ^a	B	V	R	I	u'	g'	r'	i'	z'	Telescope/Instrument
5566.3	2.5	18.11±0.02	17.86±0.02	17.51±0.02	17.38±0.02	PROMPT/Apoge
5567.3	3.5	18.19±0.04	17.86±0.02	17.53±0.01	17.26±0.05	PROMPT/Apoge
5569.3	5.5	18.18±0.03	17.85±0.02	17.49±0.04	17.40±0.02	PROMPT/Apoge
5570.3	6.5	...	17.87±0.02	17.50±0.04	17.45±0.02	PROMPT/Apoge
5573.3	9.5	18.23±0.03	17.87±0.02	17.54±0.04	17.41±0.02	...	18.05±0.02	17.82±0.02	17.89±0.03	17.78±0.09	PROMPT/Apoge
5573.6	9.8	18.23±0.04	17.90±0.03	LT/RATCam
5575.3	11.5	...	17.89±0.02	17.56±0.06	17.33±0.04	PROMPT/Apoge
5575.7	11.9	18.22±0.06	19.04±0.08	LT/RATCam
5576.2	12.4	18.26±0.02	17.98±0.03	17.65±0.04	17.41±0.05	CA-2.2/CAFOS
5577.3	13.5	...	17.89±0.02	17.56±0.06	17.30±0.04	PROMPT/Apoge
5579.6	15.8	18.22±0.03	17.96±0.02	18.77±0.08	18.13±0.03	17.92±0.02	17.91±0.03	18.00±0.04	LT/RATCam
5581.3	17.5	18.35±0.03	17.95±0.02	17.57±0.04	17.55±0.03	PROMPT/Apoge
5583.3	19.5	18.50±0.04	18.02±0.03	17.62±0.06	17.45±0.05	17.97±0.02	17.68±0.06	...	PROMPT/Apoge
5584.3	20.5	...	18.10±0.03	17.70±0.04	17.51±0.03	18.02±0.02	18.05±0.03	18.02±0.05	PROMPT/Apoge
5585.7	21.9	18.41±0.03	18.20±0.12	17.91±0.12	NTT/EFOSC2
5588.3	24.5	18.70±0.04	18.24±0.03	17.90±0.03	17.67±0.13	...	18.43±0.04	18.04±0.04	18.1±0.080	18.42±0.10	PROMPT/Apoge
5591.3	27.5	17.97±0.05	17.65±0.04	PROMPT/Apoge
5600.3	36.5	...	18.27±0.02	17.98±0.02	17.62±0.03	PROMPT/Apoge
5600.7	36.9	18.81±0.02	18.29±0.02	19.65±0.06	18.51±0.02	18.20±0.02	18.20±0.01	18.14±0.04	LT/RATCam
5602.3	38.5	18.20±0.02	18.17±0.02	18.15±0.03	PROMPT/Apoge
5606.3	42.5	18.99±0.04	18.29±0.02	18.24±0.02	18.11±0.02	18.05±0.04	PROMPT/Apoge
5610.2	46.4	...	18.40±0.04	18.06±0.03	17.58±0.05	17.98±0.07	18.10±0.03	18.11±0.01	PROMPT/Apoge
5610.6	46.8	20.45±0.20	18.80±0.06	18.30±0.04	18.22±0.03	18.22±0.05	LT/RATCam
5613.2	49.4	18.24±0.06	PROMPT/Apoge
5615.3	51.5	19.39±0.05	18.55±0.03	18.02±0.06	17.63±0.03	PROMPT/Apoge
5615.6	51.8	18.96±0.03	18.37±0.02	18.29±0.03	18.14±0.03	LT/RATCam
5617.3	53.5	19.36±0.04	18.55±0.01	18.06±0.02	17.80±0.03	PROMPT/Apoge
5621.6	57.8	19.16±0.02	18.50±0.01	18.28±0.02	18.26±0.02	LT/RATCam
5623.2	59.4	19.53±0.01	18.71±0.05	18.08±0.04	17.61±0.05	18.25±0.05	...	18.22±0.05	PROMPT/Apoge
5633.2	69.4	...	18.94±0.03	18.07±0.06	17.83±0.02	PROMPT/Apoge
5646.2	82.4	19.15±0.02	18.69±0.02	18.63±0.04	PROMPT/Apoge
5647.1	83.3	20.40±0.05	18.91±0.04	18.80±0.03	...	PROMPT/Apoge
5651.2	87.4	...	19.54±0.06	...	18.44±0.05	PROMPT/Apoge
5651.5	87.7	20.43±0.09	19.31±0.14	18.87±0.33	18.85±0.88	LT/RATCam
5653.1	89.3	...	19.59±0.06	19.07±0.02	PROMPT/Apoge
5661.4	97.6	...	20.20±0.22	19.30±0.11	18.79±0.20	1.82m Reflector/AFOSC
5666.2	102.4	19.51±0.12	19.34±0.10	...	PROMPT/Apoge
5677.1	113.3	19.54±0.06	19.14±0.05	PROMPT/Apoge

Note. – Errors correspond to one sigma uncertainties and come from the PSF fit and the local sequence magnitudes.

^aDays after discovery 2011 January 2.30 UT.

TABLE 3
 LOG OF OPTICAL SPECTROSCOPY OF SN 2011A.

JD-2450000+	UT date	Epoch ^a	Telescope/Instrument	Exposure(s)	Grating	Resolution FWHM (Å)	Resolution (km s ⁻¹)	Range (Å)
5565.8	Jan 4.3	2.0	du Pont/WFCDD WF4K-1	600	Blue Grism	8.0	365	3630-9100
5566.2	Jan 4.7	2.4	Gemini-North/GMOS	2 × 600	B600+G5307	5.2	240	3580-9100
5578.7	Jan 17.2	14.9	NOT/ALFOSC	3 × 1200	Grism 17	1.5	70	6330-8900
5620.6	Feb 28.1	56.8	SOAR/GOODMAN	2 × 1200	RALC 300	14	640	4200-8800
5627.8	Mar 7.3	64.0	du Pont/WFCDD	1000	Blue Grism	8.0	365	3700-9180
5633.8	Mar 13.3	70.0	SOAR/GOODMAN	2 × 2700	KOSI 600	5.1	235	4380-7015
5644.7	Mar 24.2	80.9	NTT/EFOSC2	2 × 3600	Grism 11/16	14.0	640	3740-9000
5649.7	Mar 29.2	85.9	Baade/IMACS	4 × 2000	300-4.3 Grating	2.7	120	4190-9000
5662.3	Apr 10.8	98.5	NTT/EFOSC2	2 × 3600	Grism 11/16	14.0	640	3610-9050
5675.6	Apr 24.1	111.8	SOAR/GOODMAN	2 × 2700	KOSI 600	7.0	320	4350-7015
5708.5	May 27.0	144.7	SOAR/GOODMAN	2 × 2400	RALC 300	10.2	470	3650-8840

^aDays after discovery 2011 January 2.30 UT.




FIG. 2.— SN 2011A $u'Bg'VRr'Ii'z'$ band light curves. The top figure represents the light curves for Landolt filters and the bottom figure for Sloan filters. Epochs are with respect to the discovery date (2011 January 2.30 UT= 0 days). Vertical red lines represent epochs of optical spectroscopy. The apparent magnitude is shown on the left hand side y-axis, while the absolute magnitude (calculated using only the distance modulus) is represented on the right hand side.

with spectrophotometric standard stars (Hamuy et al. 1992, 1994). Atmospheric absorption features were not removed except for spectra 2.0, 64.0, and 85.9 using a telluric standard (Hamuy et al. 2006). The spectra are not corrected for Milky Way or host-galaxy extinction. The spectral sequence is shown in Fig. 3.

3. RESULTS

3.1. Photometry

3.1.1. A rare double plateau

One of the defining photometric features of SN 2011A, displayed in Fig. 2 is the presence of a double plateau. The V -band light curve exhibits an initial plateau which lasts ~ 15 days with a slope $\sim 0.37 \text{ mag} \times 100 \text{ days}^{-1}$. Subsequently, the light curve declines $\sim 0.39 \text{ mag}$ in ~ 10 days. Then there is a second plateau phase which lasts ~ 15 days with a slope of $0.29 \text{ mag} \times 100 \text{ days}^{-1}$, which is flatter and longer lasting at redder wavelengths. For the remaining observed epochs the light curve declines with a slope of $2.8 \text{ mag} \times 100 \text{ days}^{-1}$. If we look at the entire V -band light curve, SN 2011A declines $\sim 2.1 \text{ mag} \times 100 \text{ days}^{-1}$. To our knowledge this double plateau has only been seen previously in the transient SN 1997bs (see Section 4.1.2). Some other SNe such as SN 1993J (Richmond et al. 1994), SN 2006aj (Campana et al. 2006), or SN 2011dh (Arcavi et al. 2011) present interesting characteristics (double-humped profile) which could be similar to our double plateau, but we think that the physics is most likely distinct. Indeed the double plateau seen in SN 2011A is powered by interaction whereas the double peak (in the above events) is first due to the cooling of the shock surface material and then to radioactive decay (Nakar & Piro 2014). We note that the double plateau is not so evident in the Sloan light curves, due to less well sampled photometry and noisier data. In the Landolt filters the double plateau is clearly visible in V and R , but less evident in B and I . We are confident that this is a real feature of the transient.

3.1.2. Colours

The $(B - V)$, $(V - R)$, and $(R - I)$ colour curves of SN 2011A corrected for Milky Way extinction (Schlafly & Finkbeiner 2011⁵, $A_V = 0.137 \text{ mag}$ and assuming $R_V = 3.1$ Cardelli et al. 1989) are presented in Fig. 4. In $(B - V)$ the object shows a nearly constant colour $\sim 0.20 \text{ mag}$ during the first 20 days, after which it grows steadily redder until 60 post discovery. A similar behavior is observed in $(V - R)$ except that the constant colour phase $\sim 0.30 \text{ mag}$ extends through day ~ 40 , before growing redder. For $(R - I)$ we can see an increase from 0 to 0.4 mag in 100 days, but the noise prevents us from ruling out an initial phase of constant color as observed in $(B - V)$ and $(V - R)$.

3.1.3. Extinction and Absolute Magnitude

We attempt to obtain constraints on host-galaxy extinction by analysing the early colour. If we assume that the initial $(B - V) = 0.2 \text{ mag}$ corresponds to

maximum light and that the intrinsic colour was close to $(B - V) = 0.0 \text{ mag}$, (which is what one might expect for SNe II cf Faran et al. 2014), then $E(B - V) \sim 0.20 \pm 0.03 \text{ mag}$ (aware of the possibility that SNe II_n may have different colours at maximum than normal SNe II). Using $R_V = 3.1$ this implies a host-galaxy extinction $A_V \sim 0.62 \pm 0.10 \text{ mag}$.

In addition we also examined the interstellar Na I D lines $\lambda\lambda 5889, 5895$ to estimate host-galaxy extinction using the spectrum with highest resolution taken close to discovery, i.e., that one obtained at Gemini-North on 2011 January 4.7 UT. The measurement of the Na I D absorption feature equivalent width (EW) yields $EW(D_{1,2}) = 2.55 \pm 0.45 \text{ \AA}$. Unfortunately, the relation between A_V and EW for unresolved Na I D lines breaks down at EWs higher than 1 \AA (Phillips et al. 2013), preventing us from deriving an independent host-galaxy extinction for SN 2011A. If this detection is due to interstellar material (ISM) then this would imply significant host-galaxy extinction. However, later we argue that the absorption is due to CSM material, and hence any connection with host galaxy-extinction is unclear.

Galactic extinction for SN 2011A in the V band is $\sim 0.137 \text{ mag}$ (Schlafly & Finkbeiner 2011⁶) assuming $R_V = 3.1$. Using a host-galaxy redshift $z = 0.008916$ (Theureau et al. 2007) and assuming a Λ CDM cosmology with $\Omega_\Lambda = 0.73$, $\Omega_M = 0.27$, and $H_0 = 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$, we obtain a distance of 36.7 Mpc , which implies an absolute magnitude uncorrected for host extinction of $M_V = -15.10 \text{ mag}$ close to discovery. Aware of the uncertainties in the A_V from the host-galaxy ($0.62 \pm 0.10 \text{ mag}$), we obtain an absolute V -band magnitude of -15.72 mag . This is low for the SN II_n class ($-16 \geq M_V \geq -20$, Richardson et al. 2002), and is high compared to the SN impostors (e.g., SN 1997bs has $M_V > -14$ Van Dyk et al. 2000 or see compilation in Smith et al. 2011). It is therefore difficult to differentiate between a SN II_n or SN impostor origin for this transient using solely its M_V .

3.2. Spectroscopy

As expected for an interacting transient, the spectral sequence presented in Fig. 3 is dominated by relatively narrow $H\alpha$ emission. However, there is significant evolution with time, as will now be discussed.

3.2.1. Spectral evolution

The spectrum with the highest resolution taken close to discovery (+2.4 days), is dominated by Balmer lines, most prominently $H\alpha$ and $H\beta$ in emission, with a characteristic width of $\sim 2400 \text{ km s}^{-1}$ measured using the Full Width at Half Maximum (FWHM). In what follows we will refer to these lines as “broad”, although the reader should keep in mind that these velocities are much lower than what is observed in most SNe II ($v \sim 10^4 \text{ km s}^{-1}$). We observe also many narrow ($\sim 500\text{--}1000 \text{ km s}^{-1}$) P-Cygni profiles attributed to Ca II $\lambda\lambda 3933, 3968$, the Ca II near-infrared triplet $\lambda\lambda 8498, 8542, 8662$, and many Fe II

⁵ via NED.

⁶ via NED.




FIG. 3.— Optical spectra of SN 2011A taken with SOAR, NTT, Gemini–North, NOT, Baade and du Pont telescopes. The epochs post discovery (2011 January 2.30 UT) are listed. Atmospheric absorption features were not removed except for spectra at 2.0, 64.0, and 85.9 days, and are indicated with a \oplus symbol. Spectra were smoothed using a boxcar of 5 pixels size. Flux calibration of spectra was revised using the multi-color photometry. The spectra have not been corrected for Milky Way or host-galaxy extinction.

lines. After \sim day 50, $H\alpha$ emission is still present but $H\beta$ is significantly weaker. We see a similar evolution of the Fe II lines which are present during the first few days, but disappear from day 56.8 onwards. The Ca II triplet is not clearly seen in spectra between days 56.8 and 85.9 due to noisy data but is still visible on the +98.5 day spectrum, and possibly 144.7 days after discovery.

To understand SNe IIn, it is common practice to analyse the $H\alpha$ line profile and its evolution. This is shown in Fig. 5 where we see considerable temporal variations. Indeed, from discovery to 56.8 days later, our spectra are characterised by prominent “broad” emission ($v \sim 2000$ km s $^{-1}$). Then between 56.8 days and 85.9 days post discovery, a narrow low-velocity (600–1100 km s $^{-1}$) P-Cygni absorption appears. After day 85.9 we see again a “broad” emission with no signs of the P-Cygni profile. Again, we emphasize the absence in the spectra at all epochs of any kind of high velocity components ($\sim 10^4$ km s $^{-1}$) characteristic of SN ejecta.

In Fig. 6 we show the $H\alpha$ line evolution splitting it into epochs with and without P-Cygni profiles. The P-Cygni component is seen in spectra obtained on days +56.8, +64.0, +70.0, and +80.9. The blueshifted absorption velocity has values between 575 km s $^{-1}$ to 1060 km s $^{-1}$. It is unlikely that these low-velocity P-Cygni profiles can be attributed to the ejecta, which is expected to have velocities of several thousand km s $^{-1}$. Therefore, we argue that this P-Cygni component is related to a pre-SN wind. The wind velocity is consistent with an LBV progenitor (100 – 1000 km s $^{-1}$ Smith & Townsend 2007); but too high for red supergiants (20 – 40 km s $^{-1}$; Smith et al. 2007) and possibly too low for Wolf-Rayet stars (1000 – 5000 km s $^{-1}$; Abbott & Conti 1987).

Our first spectra, taken 2.0 days and 2.4 days after discovery, show many strong Fe II features with P-Cygni profiles. The velocity measured with respect to the emission from the Fe II multiplet $\lambda\lambda 4923, 5018, 5169$ lines in the +2.4 day spectrum is between 430–460 km s $^{-1}$, arguing for a CSM rather than ejecta origin. We also note




FIG. 4.— Observed $(B-V)$, $(V-R)$, $(R-I)$ colour evolution of SN 2011A corrected for Milky Way extinction.




FIG. 5.— $H\alpha$ line profile evolution of SN 2011A. The epochs are indicated in black and are with respect to the discovery date (2011 Jan 2.30 UT = 0 days). The dotted red vertical line represents the rest-frame $H\alpha$ wavelength, at 6562.81 Å.

that these velocities have the same order of magnitude as those derived from the $H\alpha$ P-Cygni component. The same is seen in the red part with the calcium triplet $\lambda\lambda 8498, 8542, 8662$. Indeed in the +2.4 day spectrum we measured low velocity with respect to the emission ($\sim 440 \text{ km s}^{-1}$), again supporting the hypothesis of CSM origin and consistent with the velocities derived from the Fe II lines. Looking at the continuum we can also





FIG. 6.— $H\alpha$ line profile evolution. The top figure shows the $H\alpha$ line profile showing a broad component without absorption. We also present a zoom on $H\alpha$ for late spectra. The bottom figure shows the $H\alpha$ line profile with P-Cygni absorption. We normalized all spectra with respect to the Gemini spectrum continuum. The scale has been adjusted to overlap the broad wings of $H\alpha$.

estimate the temperature through a blackbody fit. The temperature evolution shows a rapid decline from $\approx 9000 \text{ K}$ in the +2.0 days spectrum to $\approx 5000 \text{ K}$ in the +56.8 days spectrum and to $\approx 3300 \text{ K}$ at the later epochs.

3.2.2. Variable low velocity absorption close to Na I D

As seen in the Type IIIn SN 1994W (Chugai et al. 2004), we observe a strong absorption near the Na I D doublet wavelength ($\lambda\lambda 5889, 5895$) which becomes stronger with time and has low velocity. This is shown




FIG. 7.— Zoom on the absorption close Na I D of our SN 2011A optical spectra. The epochs are listed with respect to the discovery date (2011 January 2.30 UT= 0 days). The red line represents the center of the Na I D corrected to the redshift of the host-galaxy. We note that this appears offset by $\sim 200 \text{ km s}^{-1}$ from the emission component of Na I D.

in Fig. 7, where the evolution of this feature is presented. The equivalent width and velocity of the line profile were calculated for each epoch. Because the majority of our spectra were taken in low resolution, the sodium doublet is unresolved, hence, we will refer to this feature as absorption at the wavelength of Na I D, due to the fact that the Na I D can be blended with the He I $\lambda 5876$ line (we also note the possible presence of He I at the wavelength $\lambda 6678$ and $\lambda 7065$ in the 2.4 days spectrum). Therefore the total blended doublet is measured as a single line. In Fig. 8 we present both the equivalent width and velocity evolution and see that the absorption becomes stronger with time. Indeed the equivalent width is initially equal to $\sim 3 \text{ \AA}$ and increases to 10 \AA after the first 70.0 days. At the same time, the feature profile evolves. In the first two spectra, only an absorption line is observed while, between days 56.8 and 110.3 a P-Cygni profile appears. The velocities are measured from the Na I D center, 5892 \AA with respect to the emission are shown in Table 4. Fig. 8 also shows the average Na I D equivalent widths from ISM for all the *Carnegie Supernova Project* (CSP; Hamuy et al. 2006) SNe II (as used in Anderson et al. 2014b). We measure the equivalent width for each SN regardless of the epoch and we averaged these values. This is again further evidence of a CSM origin for this line: the strength is much higher than seen in any other SNe II from the CSP database. Even if the low resolution spectra prevent us to be sure that the measured absorption close to Na I D position is only due to Na I D and not blending with other lines, the absorption strength is unprecedented.

3.2.3. Wind velocity

TABLE 4
EVOLUTION OF THE ABSORPTION CLOSE Na I D WITH TIME.

Epoch ^a	Equivalent width (\AA) ^b	Velocity (km s^{-1}) ^b	Line profile
2.0	3.2(0.5)	530(25)	absorption
2.4	2.6(0.7)	400(30)	absorption
56.8	6.9(0.35)	500(75)	P-Cygni
64.0	11.5(0.7)	510(55)	P-Cygni
70.0	9.5(0.3)	490(70)	P-Cygni
80.9	8.0(1.1)	490(50)	P-Cygni
85.9	10.8(0.5)	600(75)	P-Cygni
98.5	7.7(0.8)	830(125)	P-Cygni
111.8	10.6(1.6)	400(130)	P-Cygni
144.7

^aDays after discovery 2011 January 2.30 UT.

^bWe note in parenthesis the associated error. The errors come from the uncertainties of defining the continuum for measurement of the absorption and emission components. This was achieved using multiple measurements changing the continuum position each time. The velocities are measured from the Na I D center, 5892 \AA with respect to the emission.




FIG. 8.— Top figure: the absorption close Na I D equivalent width evolution with time for SN 2011A. The solid blue line shows the average Na I D EWs (averaged over time) for all CSP SNe II (including SNe IIn, but not SNe IIb) where the EW is most likely of ISM origin. The lower dashed blue line shows the 1 sigma error on the EW estimation, and the upper shows 3 sigma error. The lower panel shows the absorption close Na I D velocity evolution measured with respect to the emission.

In order to measure the wind velocity we estimate the width of the narrow component of the $H\alpha$ emission line. For this estimation we decompose the $H\alpha$ feature using a least-squares PYTHON script, which solves the best-fit multi Gaussian decomposition. The best fit was using three components, so nine parameters are used as input: the FWHM, the amplitude and the center of each Gaussian corresponding to the three components. Using these values our script provides the best-fit Gaussian parameters, the best FWHM, the amplitude and the center of each component. The errors on these parameters are derived assuming a reduced chi squared equal to one. In Fig. 9 we show the $H\alpha$ emission line decomposition for spectra taken with ALFOSC (highest resolu-

tion spectrum available) on 2011 January 17.2 UT. Our three components fit the $H\alpha$ emission line yields velocities of $107 \pm 2 \text{ km s}^{-1}$, $467 \pm 14 \text{ km s}^{-1}$, and $1971 \pm 35 \text{ km s}^{-1}$. Each velocity component is low compared to other SNe IIn. For comparison, Kiewe et al. (2012) found typical velocities of 2000 km s^{-1} for the intermediate component. We also note that Pastorello et al. (2004) found that for low-luminosity SNe II the expansion velocities range from $3000\text{--}5000 \text{ km s}^{-1}$ during early epochs. Therefore, SN 2011A appears to have ejecta velocities lower than the majority of normal SNe events. This could be evidence for a SN impostor origin, as will be discussed below. In addition, recently Tartaglia et al. (2014) have reported a case of an interesting transient, SN 2007sv. From this object the authors, also analysing the $H\alpha$ emission line, derived very similar velocity values. Indeed the broad component velocity was estimated to 2000 km s^{-1} (1971 km s^{-1} for SN 2011A), an intermediate component velocity of $600\text{--}800 \text{ km s}^{-1}$ (467 km s^{-1} for SN 2011A) and a narrow component velocity of $120\text{--}150 \text{ km s}^{-1}$ (107 km s^{-1} for SN 2011A). In their paper, they conclude that the transient SN 2007sv is most likely a SN impostor, based mainly on its absolute magnitude. Note that the absolute magnitude derived for this transient is ~ 2 mag fainter than SN 2011A.

Note that from the Gaussian decomposition of the highest resolution spectrum we can calculate the progenitor mass-loss rate using the following formula (Chugai & Danziger 1994; Kiewe et al. 2012):

$$L_{H\alpha} = \frac{1}{2} \dot{M} \psi \frac{v_{shock}^3}{v_{wind}} \quad (1)$$

where ψ is the efficiency of the conversion of mechanical energy into optical energy in the shock wave (we adopt as Kiewe et al. 2012, $\psi = 0.1$). \dot{M} is the mass loss rate, v_{wind} is the unshocked wind velocity derived as described above and v_{shock} is the shock velocity. The shock velocity was taken as the FWHM of the intermediate width component. The absolute $L_{H\alpha}$ was then obtained by integrating over the intermediate width feature and taking account of the distance to the SN. We used the same luminosity distance used above, i.e., 36.70 Mpc. We found $\dot{M} = 0.038 M_{\odot} \text{ yr}^{-1}$ with a wind velocity of 110 km s^{-1} and a shock velocity about 470 km s^{-1} . This mass loss rate is consistent with values found by Kiewe et al. (2012) and is too high for any class of massive stars other than LBVs in the eruptive phase (Humphreys & Davidson 1994).

4. DISCUSSION

SN 2011A shows a number of interesting properties. In order to understand the nature of the transient, we compare in this section our object with other interacting transients. We discuss the transient in comparison to other events which possibly show similarities. However, in any case where similarities exist, we show that there are other features which confirm the uniqueness of SN 2011A.

4.1. Photometry




FIG. 9.— The decomposition of the $H\alpha$ emission line for the spectrum taken with ALFOSC at NOT on 2011 January 17.2 UT.

4.1.1. Photometric comparison to other SNe IIn

As stated in the introduction, the SN IIn group is very heterogeneous, in terms of luminosity, light curve shape and spectral evolution. Based on Kiewe et al. (2012) who proposed to differentiate SNe IIn according to their photometric and spectroscopic characteristics, Kankare et al. (2012) identify a subclass of SNe IIn in which they included SN 2009kn (Kankare et al. 2012), SN 1994W (Chugai et al. 2004) and SN 2005cl (Kiewe et al. 2012). To see if SN 2011A shares common properties with these SNe and belongs to this subclass, we compare the light curves and spectra (see Section 4.2.2). We also include comparison with SN 1998S (Fassia et al. 2001) a well studied SN IIn, SN 2005kj (Taddia et al. 2013) which, similarly to SN 2011A shows evidence of low velocity absorption at Na I D wavelengths, SN 1997bs (Van Dyk et al. 2000) classified as SN impostor (note that we compare in more detail SN 2011A and SN 1997bs in the Section 4.1.2.), and SN 2007sv, also recently classified as an SN impostor (Tartaglia et al. 2014). These comparisons are shown in Fig. 10, where the V -band light curves are plotted. Note that we do not present the SN 1994W light curve due to the lack of data and because it is very similar to SN 2009kn (Kankare et al. 2012). As we can see in Fig. 10 SN 2009kn is brighter than SN 2011A with $M_V \simeq -17.5/-18$ and the SN 2009kn light curve does not show a double plateau like SN 2011A. Indeed, after an initially declining plateau phase with a slope equal to $0.018 \text{ mag day}^{-1}$, we see a quick fall around $0.08 \text{ mag day}^{-1}$ followed by a slow decline phase with a slope of $\sim 1 \text{ mag } 100 \text{ days}^{-1}$. However, the photometry of SN 2009kn does not cover the first few tens of days from the assumed explosion date. As can be seen in Fig. 10, SN 2011A is photometrically very different that the other SNe IIn. None of the other selected SNe IIn show a double plateau and they are all ≥ 2 mag brighter than SN 2011A. Also we remark that SN 2011A is ~ 2 mag brighter than the SN impostors. Fig. 10 allows us to conclude that SN 2011A is a very rare event which does not share any similarities in the light curve with other SN 1994W-like SNe IIn. An




FIG. 10.— Comparison of the absolute V -band light curves of SN 2011A, SN 1998S (Fassia et al. 2001), SN 2009kn (Kankare et al. 2012), SN 2005cl (Kiewe et al. 2012), SN 2005kj (Taddia et al. 2013), SN 1997bs (Van Dyk et al. 2000), and SN 2007sv (Tartaglia et al. 2014). Each SN was corrected for Milky Way extinction. Full red triangles represent SN 2011A corrected for host-galaxy extinction ($A_V = 0.62$ mag). For SN 2009kn we use the explosion date estimation from Kankare et al. (2012). For SN 1998S, SN 2005kj, SN 2005cl, SN 1997bs, and SN 2007sv we use the discovery date as the explosion date. For each SN we applied the host-galaxy extinction estimated by the authors.

important additional point to note is the origin of the two SNe II_n; SN 1994W and SN 2009kn used to compare SN 2011A to other SNe II_n. Indeed the origin is not so clear, for example Dessart et al. (2009) argued that maybe SN 1994W was not a SN, but rather the result of two interacting shells of material related to LBV-like eruptions. Also, in the case of SN 2009kn (Kankare et al. 2012) a non-SN origin could not be completely excluded, but also found the observations to be consistent with an electron capture SN. We note that later an argument will be presented that constrains the epoch of the photometry of SN 2011A to be ~ 50 days later than shown in Fig. 10. However, the main conclusions remain valid: SN 2011A has a luminosity between the two classes of interacting transients, SNe II_n and SN impostors; this event shows a rare double plateau.

In Fig. 11 we present the optical colours of SN 2011A together with the colour evolution of SN 2009kn, one SN 1994W-like, and the well-studied SN 1998S. We do not present the SN 1994W and SN 2005cl colours due to the lack of data. All magnitudes were corrected for Milky Way extinction. Taking the discovery date as the explosion date, the colour evolution of SN 2011A looks fairly similar to SN 1998S. Indeed, the $(B - V)$ colour becomes redder in SN 1998S between days 0 and 60. However, in $(V - R)$, SN 1998S does not show an initial plateau as SN 2011A.

The colour evolution of SN 2009kn looks similar to




FIG. 11.— $(B - V)$, $(V - R)$, $(R - I)$ colour evolution of SN 2011A (red triangles) compared to other SNe, SN 2009kn (Kankare et al. 2012) shown in black squares and SN 1998S (Fassia et al. 2000) represented by blue circles. All colours were corrected for Milky Way extinction and by host-galaxy extinction. For SN 2011A and SN 1998S the epochs are with respect to the discovery date, whereas for SN 2009kn they are with respect to the explosion date estimated by the authors.

SN 2011A in $(B - V)$ and $(V - R)$, but shifted by ~ 50 days. This could be explained by the lack of precision in the explosion date for SN 2011A. If we shift our epochs by 50 days (motivated by the spectral analysis in section 4.2.1), the colour evolution in $(B - V)$ and $(V - R)$ is very similar to SN 2009kn. Indeed for $(V - R)$, SN 2009kn shows a small plateau during ~ 50 days and then an increase from 0.3 to 0.7 mag. It is more difficult to compare the behavior of the $(R - I)$ colour, due to their larger errors, however we can see the same trend: for SN 2011A, $(R - I)$ increases from ~ 0 to ~ 0.4 mag in 100 days, and for SN 2009kn, $(R - I)$ increases from ~ 0.2 to ~ 0.55 mag in the same temporal windows. The colours between SN 2011A and SN 2009kn are very similar, but shifted by 50 days which could be consistent with the hypothesis proposed in section 4.2.1, namely, that the explosion date is not close to the discovery date but more consistent with 50 days earlier.

4.1.2. SN 2011A versus SN 1997bs

The light curve comparison between SN 2011A and SN 1997bs (which was classified as a SN impostor by Van Dyk et al. 2000) shows similarities. The light curves of both objects in B , V , R , and I bands are shown in Fig. 12. For the SN 1997bs explosion date, we choose 1997 April 15th (UT discovery date), which should be valid because an image taken on 1997 April 10th does not show anything at the location of the SN. During the first 15 days of evolution in R and I bands both objects exhibit a plateau. After this first plateau the two transients show a second plateau but the drop between the two is much larger for SN 1997bs: ~ 0.5 mag for

SN 2011A and ~ 1 mag for SN 1997bs in R . The B and V bands have also some similarities. Note, while the light curve evolution displayed in Fig. 12 appears very similar, this may be coincidental given the uncertainty of the SN 2011A explosion date. However, the important observation is the similar double plateau features in both light curves, which may be evidence for the same underlying mechanism; namely interaction with two separate CSM shells. The optical colours of SN 2011A compared with SN 1997bs are presented in Fig. 13. From the $(B - V)$ colour we can compare the colours near maximum. For SN 2011A we find ~ 0.25 mag whereas for SN 1997bs Van Dyk et al. (2000) found 0.67 mag, adding a host extinction for 1997bs. Additionally we can compare the colour evolution. In the $(V - R)$ colour, the evolution looks similar during the first 30 days. For SN 1997bs, a plateau at 0.3 mag during 30 days is seen and then a reddening as for SN 2011A. But in SN 2011A the plateau lasts 50 days. Despite the noise, $(B - V)$ colour exhibits the same trend with a plateau during 20 days and then a reddening. Again, in $(R - I)$, SN 2011A and the SN impostor show the same evolution with a reddening during the first 40 days. Although the data are noisy the shape of the reddening is pretty similar. Finally, note that the SN impostor nature of SN 1997bs is not entirely clear as there is some evidence that the progenitor did not survive to the explosion (Li et al. 2002; Adams & Kochanek 2015). Hence, while SN 2011A shows similarities to SN 1997bs, this is not concluding proof that the former was an impostor event.

4.1.3. ^{56}Ni mass estimate

One diagnostic to discriminate between SNe and SN impostors is the amount of synthesized ^{56}Ni . For low mass core-collapse SNe (SNe IIP) one expects to find a ^{56}Ni mass around $0.0016 - 0.26 M_{\odot}$ (Hamuy 2003) whereas for non-terminal explosions we do not expect any ^{56}Ni production because only the outer layers of the star are ejected. For very massive CCSNe produced by progenitors with initial masses between $30 - 100 M_{\odot}$, ^{56}Ni masses between $2.3 - 6.6 M_{\odot}$ have been predicted (Umeda & Nomoto 2008), and masses between $0.07 - 0.6 M_{\odot}$ have been estimated for progenitor masses between $16 - 46 M_{\odot}$ (Mazzali et al. 2009). We can obtain a ^{56}Ni mass estimate using the formula from Hamuy (2003) and the last V -band magnitude (epoch 97.6 days). This value should be taken as an upper limit of the ^{56}Ni mass because the spectrum at this epoch does not indicate that the emission is being solely powered by radioactivity. Indeed in this spectrum narrow lines are still present due to interaction.

As stated previously the explosion date is not well constrained and could be 50 days before discovery. Therefore the ^{56}Ni mass is calculated assuming the explosion date as the discovery date and also using an explosion date 50 days before discovery. We find $0.01 - 0.015 M_{\odot}$ without extinction and $0.03 - 0.05 M_{\odot}$ with extinction. In both cases the upper value is using an explosion date 50 days before discovery. These values are consistent with Hamuy (2003) and lower than Mazzali et al. (2009). Indeed Hamuy (2003) found for SNe II values between $0.0016 - 0.26 M_{\odot}$ and Mazzali et al. values between 0.07

$- 0.6 M_{\odot}$. If we apply this same approach to SN 1997bs, we find a ^{56}Ni of $0.003 M_{\odot}$ based on the V -band magnitude measured 63.5 days after discovery. In conclusion it is very difficult to set significant constraints on the nature of SN 2011A based on these uncertain ^{56}Ni mass estimates.

4.2. Spectroscopy

4.2.1. Explosion date

Unfortunately, the explosion date of SN 2011A is not well determined by pre-explosion images. A fit to the early time spectrum with a blackbody function yields a high temperature (~ 10000 K), thus suggesting that the observations possibly began soon after explosion. However, given the interacting nature of SN 2011A, it is unclear how valid such a constraint is. To better estimate this parameter, we compare SN 2011A with two other SNe with well-constrained explosion times.

Firstly, SN 1994W (Cortini et al. 1994), for which Sollerman et al. (1998) estimated from its R -band data a precise explosion date of 1994 July 14 $^{+2}_{-4}$. In Fig. 14 we compare our spectrum taken 2.0 days after discovery with a SN 1994W spectrum taken 57 days after explosion. As we can see, the spectra are remarkably similar. A blue continuum characterizes both spectra and the slope/shape are nearly identical. Also, both SNe show strong Fe II multiplet $\lambda\lambda 4923, 5018, 5169$ lines with low velocity (\sim hundreds km s^{-1}) P-Cygni profiles (Fig. 14, top right). We also distinguish in the blue part of the spectrum a strong absorption attributed to CaII $\lambda\lambda 3933, 3968$. SN 2011A shows two noticeable differences from SN 1994W: (1) the Ca II near-infrared triplet is more prominent than in SN 1994W and (2) SN 1994W shows a narrow P-Cygni absorption on top of the broad of $H\alpha$ component whereas in SN 2011A this feature is not seen and only appears in the spectrum taken +56.8 days after discovery.

Secondly, in Fig. 15 we compare our +2.0 day spectrum with the SN 1997bs spectrum taken on 1997 April 16th (day +1 after discovery) to search for similarities as seen in their light curves. Given that the non-detection was very close to discovery, the explosion date is well constrained for SN 1997bs. The continua look very similar when we apply a total reddening of $E(B - V) = 0.24$ mag to SN 1997bs (Van Dyk et al. 2000 estimated this total color excess at 0.21 mag) and null for SN 2011A. We choose this colour excess value to match the two continua. The spectra show some differences, most notably the lack of narrow P-Cygni Fe II lines in SN 1997bs. In contrast, their $H\alpha$ emission profiles are virtually identical. Note also that if these spectra were taken at similar epochs then this would imply that the host-galaxy extinction for SN 2011A is less than that of SN 1997bs. For SN 1997bs Van Dyk et al. (2000) derived an total extinction $A_V \sim 0.65$ mag. This value is close to that found for SN 2011A in section 3.1.3 but not significantly lower.

Thanks to these comparisons and the fact that our spectrum taken 2.0 days after discovery has a good match with the SN 1994W spectrum taken 57 days past explo-




FIG. 12.— Comparison of the *BVRI* light curves of SN 2011A and the impostor SN 1997bs. Photometric data of the SN impostor were taken from Van Dyk et al. (2000). For each band empty symbols represent SN 2011A and full symbols SN 1997bs. Squares represent *I*-band light curve, triangles *R* band, hexagons *V* band and circles *B* band.

sion, it is possible that SN 2011A maybe exploded $\sim 2 - 15$ km s^{-1} (Fassia et al. 2001). If the component comes from the CSM, the velocity will be higher (few hundred km s^{-1} as for SN 2011A) and depends on the progenitor. Finally, the component from a SN ejecta would have the highest values of a few thousand km s^{-1} . Though the majority of spectra are low resolution and prevent us to resolve the Na I D absorption line, we believe that the absorption close to the Na I D wavelength position is mostly due to Na I D. Even if some lines like Ba II $\lambda 5854$ can be blended with the Na I D, the absorption strength and velocity are very rare. Indeed others SNe have shown similar remarkable increase of Na I D absorption, such as SN 1998A and SN 2009E (Pastorello et al. 2005, 2012b), but Pastorello et al. (2012a) measured for SN 2009E the Na I D velocity and found a value of $\sim 3000 - 5000$ km s^{-1} . This implies that the absorption line is due the SN ejecta, whereas the velocities measured for Na I D absorption in SN 2011A never gets higher than 1100 km s^{-1} , inconsistent with an ejecta origin, but consistent with a CSM origin interpretation.

4.2.2. Na I D evolution

Na I D absorption can have three origins. From the interstellar medium (ISM) in the line of sight, the circumstellar material ejected by the progenitor ionized by the radiation from the SN and recombined over the following several days to weeks, or from the SN ejecta. The main distinguishing feature between these three components is the velocity. If the absorption is due to the

ISM, the velocity will be very low, $\sim 2 - 15$ km s^{-1} (Fassia et al. 2001). If the component comes from the CSM, the velocity will be higher (few hundred km s^{-1} as for SN 2011A) and depends on the progenitor. Finally, the component from a SN ejecta would have the highest values of a few thousand km s^{-1} . Though the majority of spectra are low resolution and prevent us to resolve the Na I D absorption line, we believe that the absorption close to the Na I D wavelength position is mostly due to Na I D. Even if some lines like Ba II $\lambda 5854$ can be blended with the Na I D, the absorption strength and velocity are very rare. Indeed others SNe have shown similar remarkable increase of Na I D absorption, such as SN 1998A and SN 2009E (Pastorello et al. 2005, 2012b), but Pastorello et al. (2012a) measured for SN 2009E the Na I D velocity and found a value of $\sim 3000 - 5000$ km s^{-1} . This implies that the absorption line is due the SN ejecta, whereas the velocities measured for Na I D absorption in SN 2011A never gets higher than 1100 km s^{-1} , inconsistent with an ejecta origin, but consistent with a CSM origin interpretation.

In the literature we found a small number of cases




FIG. 13.— $(B-V)$, $(V-R)$, $(R-I)$ colour evolution of SN 2011A (red triangles) compared to impostor SN 1997bs (Van Dyk et al. 2000) shown in black circles. All colours were corrected by Milky Way extinction and by host-galaxy extinction.




FIG. 14.— Comparison between the SN 2011A spectrum taken 2.0 days after discovery, in black, and that of SN 1994W taken +57 days after explosion, in red. In the top right inset we zoom on of the Fe II lines, $\lambda\lambda 4923, 5018, 5169$. On the bottom left inset we zoom on H α emission line. For both spectra we did not apply any MW or host-galaxy extinction correction.

of CSM Na I D in SNe IIn. Indeed for SN 2009kn (Kankare et al. 2012), SN 1994W (Chugai et al. 2004), SN 1997bs (Van Dyk et al. 2000), and SN 2005kj (Taddia et al. 2013), the velocity is low as for SN 2011A, less than 1100 km s^{-1} . The increase of Na I D absorp-




FIG. 15.— Comparison between the SN 2011A spectrum taken 2.0 days after discovery, in black, and that of SN 1997bs taken on 16th April 1997, one day after discovery, in red (no detection on 10th Apr 1997). In the top right inset we zoom on the Fe II lines, $\lambda\lambda 4923, 5018, 5169$. On the bottom left inset we zoom on the H α emission line. We applied a total reddening of $E(B-V)=0.24$ mag to SN 1997bs in order to match the continua.

tion, according to Chugai & Danziger (1994), is a result of the increasing Na I ionization fraction.

4.2.3. Dust formation

Most SN impostors candidates in the literature have no surviving star at the location of the transient, which could contradict the nature of the transient, i.e., a non-terminal eruptions of massive stars. However, a simple reason could explain this. The dust formed in a dense shell ejected by the star will obscure the SN impostor. The presence of dust can be deduced by different ways. First, by the presence of an infrared excess due to the thermal emission from dust condensed in the ejecta or from heated pre-existing dust. Unfortunately we do not have useful infrared data for this transient. Secondly, looking at the emission lines. If the spectral feature profiles show a blueshift, this can be attributed to the fact that the red side of the ejecta is blocked by the new dust (Smith et al. 2012, but see Anderson et al. 2014a and Dessart et al. 2015 for an alternative explanation). This effect is maybe seen in our transient. Indeed in Fig. 5 where the H α emission line evolution is shown, we can see for the spectrum +111.8 days that the red side is weaker compared to the blue part. This characteristic is not clearly observed after and before this epoch especially due to the appearance of a P-Cygni profile.

The estimated black body temperature at late times (3300K) is also possibly indicative of dust formation. This is because this temperature appears to be too low for the spectra to be dominated by strong Balmer emission lines. This low temperature could be due to dust

formation which reddens the object. Multi-band photometry fitting seems to favor an extincted black body but it is difficult to put strong constraints on the black body temperature and extinction due to the small wavelength baseline and because our transient is mainly powered by interaction which implies that using a black body fitting could be not realistic. Any increase in reddening could also related to the observed increase in Na I D. However we note that the increase in the latter is more likely due to CSM effects.

4.2.4. CSM properties

Analysing the H α emission line evolution with time together with the light curve allows us to constrain CSM properties. The SN 2011A light curve shows a double plateau. The presence of this rare double plateau leads us to speculate that the CSM is composed of two shells ejected by the pre SN wind. During ~ 15 days there is an initial plateau which corresponds to the interaction between the SN blast wave and the first shell. When the ejecta reaches the edge of the first shell the light curve drops. If we assume that the blast wave speed is constant and taking the value found in section 3.2.4, $\sim 2000 \text{ km s}^{-1}$, we find a lower limit width of the first shell of $\geq 2.6 \times 10^{14} \text{ cm}$. It is a lower limit due to the fact the beginning of this shell is not well constrained (because of the uncertainty in the explosion date). Then, there is another weaker plateau which lasts ~ 15 days in the V band. Again, assuming that the ejecta velocity is constant with time, we find the width for the second shell, $\sim 2.6 \times 10^{14} \text{ cm}$. A cartoon showing the CSM and the velocities of the different components is presented in Fig. 16. This scenario would be similar to that proposed by Dessart et al. (2009) for SN 1994W.

This photometric analysis seems be also consistent with the spectral evolution. In Fig. 5 we see that the H α profile changes with time, from the very broad component at early epochs during the first plateau phase (spectra +2.0, +2.4, and +14.9 days), to a narrower broad component, accompanied with low velocity H α absorption (+56.8, +64, +70, +80.9, +85.9 days) during and after the second plateau.

5. CONCLUSIONS

We have presented a detailed study of SN 2011A, initially classified as a SN IIn based on the presence of a narrow H α emission in the spectrum. While the exact nature of the transient remains unclear, SN 2011A shows a number of interesting and unusual properties:

1. Double plateau in light curve likely due to CSM composed of two shells.
2. Low luminosity, $-15.10 \geq M_V \geq -15.72$ (depending on assumed host-galaxy extinction, 0 or 0.62

mag in V -band).

3. Low P-Cygni H α velocity $\leq 1200 \text{ km s}^{-1}$.
4. Low velocity absorption close to Na I D $\leq 1100 \text{ km s}^{-1}$.
5. Variable absorption close to Na I D increasing in strength, from 3 to 10 \AA in equivalent width.

Given the photometric and spectroscopic properties we have derived for SN 2011A; most importantly its low luminosity and low ejecta velocity together with their comparison to other interacting transients, we believe that SN 2011A is most likely an SN impostor event. However, we also note that our analysis has demonstrated the difficulty in determining whether interacting transients are true terminal events or indeed SN impostors. The reader also must keep in mind that the characteristics used to separate the two events are in general arbitrary and hence detailed studies of future events are needed to further elucidate this issue.

6. ACKNOWLEDGEMENTS

The referee is thanked for their through reading of the manuscript, which helped clarify and improve the paper. Support for TdJ, GP, MH, FF, CG, FB, and SG is provided by the Ministry of Economy, Development, and Tourism's Millennium Science Initiative through grant IC120009, awarded to The Millennium Institute of Astrophysics, MAS. S.G., and F.B. acknowledges support by CONICYT through FONDECYT grants 3130680 and 3120227. SB and LT are partially supported by the PRIN-INAF 2011 with the project "Transient Universe: from ESO Large to PESSTO". N.E.R. acknowledges financial support by the MICINN grant AYA2011-24704/ESP, and by the ESF EUROCORES Program EuroGENESIS (MINECO grants EUI2009-04170). M. S. gratefully acknowledge generous support provided by the Danish Agency for Science and Technology and Innovation realized through a Sapere Aude Level 2 grant. E.K. acknowledges financial support from the Jenny and Whhuri Foundation. M. D. S. gratefully acknowledges generous support provided by the Danish Agency for Science and Technology and Innovation realized through a Sapere Aude Level 2 grant. The authors thank J. Mauerhan for providing us with the SN 1994W spectrum (taken from the A. Filippenko's database in Berkeley) and also S. D. Van Dyk for the SN 1997bs spectrum. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration and of data provided by the Central Bureau for Astronomical Telegrams.

REFERENCES

- Abbott, D. C., & Conti, P. S. 1987, *ARA&A*, 25, 113
 Adams, S. M., & Kochanek, C. S. 2015, ArXiv e-prints
 Anderson, J. P., Haberman, S. M., James, P. A., & Hamuy, M. 2012, *MNRAS*, 424, 1372
 Anderson, J. P., et al. 2014a, *MNRAS*, 441, 671
 —. 2014b, *ApJ*, 786, 67
 Arcavi, I., et al. 2011, *ApJ*, 742, L18
 Bond, H. E., et al. 2009, *ApJ*, 695, L154
 Botticella, M. T., et al. 2009, *MNRAS*, 398, 1041
 Buzzoni, B., et al. 1984, *The Messenger*, 38, 9




FIG. 16.— Cartoon illustration of the CSM composition with the velocity of the different components. The black star represents the transient. Not to scale.

- Campana, S., et al. 2006, *Nature*, 442, 1008
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
 Chevalier, R. A. 1981, *ApJ*, 251, 259
 Chugai, N. N. 1990, *Soviet Astronomy Letters*, 16, 457
 —. 2001, *MNRAS*, 326, 1448
 Chugai, N. N., & Danziger, I. J. 1994, *MNRAS*, 268, 173
 Chugai, N. N., et al. 2004, *MNRAS*, 352, 1213
 Cortini, G., et al. 1994, *IAU Circ.*, 6042, 1
 Crowther, P. A. 2007, *ARA&A*, 45, 177
 Dessart, L., Audit, E., & Hillier, D. J. 2015, *ArXiv e-prints*
 Dessart, L., Hillier, D. J., Gezari, S., Basa, S., & Matheson, T. 2009, *MNRAS*, 394, 21
 Dilday, et al. 2012, *Science*, 337, 942
 Dwarkadas, V. V., Dewey, D., & Bauer, F. 2010, *MNRAS*, 407, 812
 Faran, T., et al. 2014, *MNRAS*, 442, 844
 Fassia, et al. 2001, *MNRAS*, 325, 907
 Filippenko, A. V. 1997, *ARA&A*, 35, 309
 Fransson, C. 1982, *A&A*, 111, 140
 Fraser, M., et al. 2013, *MNRAS*, 433, 1312
 Gal-Yam, A., & Leonard, D. C. 2009, *Nature*, 458, 865
 Gräfener, G., & Hamann, W.-R. 2008, *A&A*, 482, 945
 Groh, J. H., Meynet, G., & Ekström, S. 2013, *A&A*, 550, L7
 Habergham, S. M., et al. 2014, *MNRAS*, 441, 2230
 Hamuy, M. 2003, *ApJ*, 582, 905
 Hamuy, M., Suntzeff, N. B., Heathcote, S. R., Walker, A. R., Gigoux, P., & Phillips, M. M. 1994, *PASP*, 106, 566
 Hamuy, M., Walker, A. R., Suntzeff, N. B., Gigoux, P., Heathcote, S. R., & Phillips, M. M. 1992, *PASP*, 104, 533
 Hamuy, M., et al. 2003, *Nature*, 424, 651
 —. 2006, *PASP*, 118, 2
 Humphreys, R. M., & Davidson, K. 1994, *PASP*, 106, 1025
 Kankare, et al. 2012, *MNRAS*, 424, 855
 Kiewe, M., et al. 2012, *ApJ*, 744, 10
 Kochanek, C. S. 2011, *ApJ*, 741, 37
 Landolt, A. U. 1992, *AJ*, 104, 340
 —. 2007, *AJ*, 133, 2502
 Langer, N. 1993, *Space Sci. Rev.*, 66, 365
 Li, W., Filippenko, A. V., Van Dyk, S. D., Hu, J., Qiu, Y., Modjaz, M., & Leonard, D. C. 2002, *PASP*, 114, 403
 Maeder, A., & Meynet, G. 2008, in *Astronomical Society of the Pacific Conference Series*, Vol. 388, *Mass Loss from Stars and the Evolution of Stellar Clusters*, ed. A. de Koter, L. J. Smith, & L. B. F. M. Waters, 3
 Maeder, A., Meynet, G., & Hirschi, R. 2005, in *Astronomical Society of the Pacific Conference Series*, Vol. 336, *Cosmic Abundances as Records of Stellar Evolution and Nucleosynthesis*, ed. T. G. Barnes, III & F. N. Bash, 79
 Margutti, R., et al. 2014, *ApJ*, 780, 21
 Mauerhan, J., Smith, N., Filippenko, A. V., Silverman, J., Cenko, B., & Clubb, K. 2013, in *American Astronomical Society Meeting Abstracts*, Vol. 221, *American Astronomical Society Meeting Abstracts*, 233.03
 Maund, J. R., et al. 2006, *MNRAS*, 369, 390
 Mazzali, P. A., Deng, J., Hamuy, M., & Nomoto, K. 2009, *ApJ*, 703, 1624
 Moriya, T. J., Maeda, K., Taddia, F., Sollerman, J., Blinnikov, S. I., & Sorokina, E. I. 2014, *MNRAS*, 439, 2917
 Nakar, E., & Piro, A. L. 2014, *ApJ*, 788, 193
 Pastorello, A., et al. 2004, *MNRAS*, 347, 74
 —. 2005, *MNRAS*, 360, 950
 —. 2010, *MNRAS*, 408, 181
 —. 2012a, *ArXiv e-prints*
 —. 2012b, *A&A*, 537, A141
 Patat, F. 1996, *PhD thesis*, Univ. Padova
 Phillips, M. M., et al. 2013, *ApJ*, 779, 38
 Pignatta, G., et al. 2009, in *American Institute of Physics Conference Series*, Vol. 1111, *American Institute of Physics Conference Series*, ed. G. Giobbi, A. Tornambe, G. Raimondo, M. Limongi, L. A. Antonelli, N. Menci, & E. Brocato, 551–554

- Pignata, G., et al. 2011, Central Bureau Electronic Telegrams, 2623, 1
- Prieto, J. L., Brinacombe, J., Drake, A. J., & Howerton, S. 2013, ApJ, 763, L27
- Prieto, J. L., et al. 2009, ApJ, 705, 1425
- Reichert, D., et al. 2005, Nuovo Cimento C Geophysics Space Physics C, 28, 767
- Richardson, D., et al. 2002, AJ, 123, 745
- Richmond, M. W., et al. 1994, AJ, 107, 1022
- Salamanca, I., Cid-Fernandes, R., Tenorio-Tagle, G., Telles, E., Terlevich, R. J., & Munoz-Tunon, C. 1998, MNRAS, 300, L17
- Schaller, G., Schaerer, D., Meynet, G., & Maeder, A. 1992, A&AS, 96, 269
- Schlafly, E. F., & Finkbeiner, D. P. 2011, ApJ, 737, 103
- Schlegel, E. M. 1990, MNRAS, 244, 269
- Smartt, S. J. 2009, ARA&A, 47, 63
- . 2015, ArXiv e-prints
- Smith, J. A., et al. 2002, AJ, 123, 2121
- Smith, N. 2013, MNRAS, 434, 102
- Smith, N., Mauerhan, J. C., & Prieto, J. L. 2014, MNRAS, 438, 1191
- Smith, N., & Tombleson, R. 2015, MNRAS, 447, 598
- Smith, N., & Townsend, R. H. D. 2007, ApJ, 666, 967
- Smith, N., et al. 2007, ApJ, 666, 1116
- . 2009, ApJ, 697, L49
- . 2011, MNRAS, 415, 773
- . 2012, AJ, 143, 17
- Sollerman, J., Cumming, R. J., & Lundqvist, P. 1998, ApJ, 493, 933
- Steele, I. A., et al. 2004, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 5489, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, ed. J. M. Oschmann, Jr., 679–692
- Stothers, R. B., & Chin, C.-W. 1996, ApJ, 468, 842
- Stritzinger, M., Prieto, J. L., Morrell, N., & Pignata, G. 2011, Central Bureau Electronic Telegrams, 2623, 2
- Taddia, F., et al. 2013, A&A, 555, A10
- Tartaglia, L., et al. 2014, ArXiv e-prints
- Theureau, G., Hanski, M. O., Coudreau, N., Hallet, N., & Martin, J.-M. 2007, A&A, 465, 71
- Umeda, H., & Nomoto, K. 2008, ApJ, 673, 1014
- Van Dyk, S. D., Peng, C. Y., King, J. Y., Filippenko, A. V., Treffers, R. R., Li, W., & Richmond, M. W. 2000, PASP, 112, 1532
- Van Dyk, S. D., et al. 2006, ArXiv Astrophysics e-prints
- Vink, J. S. 2011, Ap&SS, 336, 163
- Wagner, R. M., et al. 2004, PASP, 116, 326
- Weis, K., & Bomans, D. J. 2005, A&A, 429, L13
- Zhang, T., et al. 2012, AJ, 144, 131