

A mathematical framework for exact milestoning

David Aristoff*, Juan M. Bello-Rivas†, Ron Elber‡

Abstract

We give a mathematical framework for Exact Milestoning, a recently introduced algorithm for mapping a continuous time stochastic process into a Markov chain or semi-Markov process that can be efficiently simulated and analyzed. We generalize the setting of Exact Milestoning and give explicit error bounds for the error in the Milestoning equation for mean first passage times.

1 Introduction

Molecular Dynamics (MD) simulations, in which classical equations of motions are solved for molecular systems of significant complexity, have proven useful for interpreting and understanding many chemical and biological phenomena (for textbooks see [47, 24, 3]). However, a significant limitation of MD is of time scales. Many molecular processes of interest occur on time scales significantly longer than the temporal scales accessible to straightforward simulations. For example, permeation of molecules through membranes can take hours [9] while MD is usually restricted to the microseconds time scale. One approach to extend simulation times is to use faster hardware [48, 49, 45]. Other approaches focus on developing theories and algorithms for long time phenomena. Most of the emphasis has been on methodologies for activated processes with a single dominant barrier, as in Transition Path Sampling [16, 7, 15]. Approaches for dynamics on rough energy landscapes, and for more general and/or diffusive dynamics, have also been developed [41, 46, 14, 50, 43]. The techniques of Exact Milestoning [5] and Milestoning [22] belong to the last category. They are theories and algorithms to accelerate trajectory calculations of kinetics and thermodynamics in complex molecular systems. The acceleration is based on the use of a large number of short trajectories instead of complete trajectories between reactants and products (Figure 1). The simulation of short trajectories is trivial to implement in parallel, making the formulation efficient to use on modern computing resources. Moreover, the use of short trajectories makes it

*Department of Mathematics, Colorado State University, Fort Collins, CO

†Institute for Computational Engineering and Sciences, University of Texas at Austin, Austin, TX

‡Institute for Computational Engineering and Sciences, Department of Chemistry, University of Texas at Austin, Austin, TX

possible to enhance sampling of improbable but important events by initiating the short trajectories near bottlenecks of reactions. A challenge is how to start the short trajectories, and how to analyze the result to obtain correct long time behavior.

While Milestoning is an approximate procedure, it shares the same philosophy and core algorithm as the Exact Milestoning approach. In both algorithms the phase space Ω is partitioned by hypersurfaces, which we call milestones $M \subset \Omega$, into cells. The short trajectories are initiated on milestones and are terminated the first time they reach a neighboring milestone (Figure 1). The short trajectories can be simulated in parallel.

Milestoning uses an approximate distribution for the initial conditions of the trajectories at the hypersurfaces. The results are then analyzed within the Milestoning theory. The approximation is typically the (normalized) canonical distribution restricted to the milestone interface M . In Exact Milestoning the distribution of hitting points at the interface is estimated numerically by iteratively computing trajectory fragments between milestones. In a straightforward implementation of the iterations (see also [5]) the final phase points of trajectories that were terminated on one milestone are continued until they hit another milestone. This type of trajectory continuation procedure is also used in Non-Equilibrium Umbrella Sampling [55] and Trajectory Tilting [53]. The continuation does not mean that full trajectories from reactants to products are computed. The calculations stop when the stationary distribution at the interface, or observables of interest, converge. In practice, and depending of course on the initial guess, the calculation ends significantly earlier than complete trajectories from R to P are computed. The fast convergence of the iterations leads to significant computational savings.

A number of other algorithms build on the use of short trajectories to estimate long time kinetics by “patching” these short trajectories at milestones or interfaces. These technologies include the Weighted Ensemble (WE) [57, 32], Transition Interface Sampling (TIS) [52], Partial Path Transition Interface Sampling (PPTIS) [43], Forward Flux Sampling (FFS) [2], Non-Equilibrium Umbrella Sampling (NEUS) [55], Trajectory Tilting [53], and Boxed Molecular Dynamics (BMD) [25]. Some of these techniques are similar; however, many subtle differences remain. Some of the differences are as follows. WE is the only method that makes it necessary to use stochastic dynamics. The trajectory sampling in NEUS, Trajectory Tilting and Exact Milestoning is similar, even though the theories are quite different. Exact Milestoning allows for the calculations of all the moments of the first passage time [5], a result which is not available for other technologies. Boxed Molecular Dynamics, Milestoning and PPTIS are approximate methods leading to greater efficiency. TIS, PPTIS, FFS, and BMD are focused on one-dimensional reaction coordinates. Other technologies (e.g. WE, Milestoning, NEUS, and Trajectory Tilting) focus on a space of one or several coarse variables.

Hence, the overall scopes of these techniques differ significantly, which make direct comparison between them less obvious. We have compared in the past the accuracy and efficiency of the methods of Milestoning and Exact Milestoning

with Forward Flux [5, 10]. Forward Flux is one of the closest algorithms (in one dimension) to Milestoning and Exact Milestoning. Numerous examples of kinetics of molecular systems studied with Milestoning were published [9, 11, 36, 34, 38, 21, 20, 56, 19]. We have also discussed extensively the features of alternate technologies that exploit trajectory fragments [56, 40].

The Milestoning theory has not yet been subject to rigorous mathematical analysis, which is the goal of the present manuscript. In this manuscript we show that the Exact Milestoning method can be derived and analyzed in the framework of probability theory. The result is a useful link between physical intuition and a more formal approach. Readers that are interested in the efficiency of the algorithm on concrete examples, and comparison to other technologies, are referred to the sources mentioned the above paragraph.




Figure 1: Representation of the state space Ω and the milestones. Each milestone is one of the line segments traced by dashed grey lines. The reactant state R is highlighted as a square dot in the left-bottom corner while the product state P is comprised of the two line segments shown in blue at the upper-right corner. A particular realization of a long trajectory appears as a continuous black line and the corresponding values of (J_n) are marked with dots.

This article is organized as follows. In Section 2, we describe the setting for Exact Milestoning and introduce notation used throughout. In Section 3, we show existence of and convergence to a stationary flux under very general conditions. In Section 4 we state precisely the Exact Milestoning algorithm [5]. In Section 5, we establish conditions under which convergence to the stationary flux is consistent in the presence of numerical error (Lemma 7 and Theorem 8), and we give a natural upper bound for the numerical error arising in Exact Milestoning (Theorem 9). Finally, in Section 6 we consider some instructive

examples.

2 Setup and notation

2.1 The dynamics and MFPT

In Milestoning we spatially coarse-grain a dynamics (X_t) . The basic idea is to stop and start trajectories on certain interfaces, called milestones, and then reconstruct functions of (X_t) using these short trajectories, which can be efficiently simulated in parallel. We assume here the dynamics is stochastic, and focus on using Milestoning for the efficient computation of *mean first passage times* (MFPTs) of (X_t) , although similar ideas can be used to compute other non-equilibrium quantities.

To make our arguments we need some assumptions on (X_t) . We let (X_t) be a time homogeneous strong Markov process with càdlàg paths taking values in a standard Borel space Ω . These assumptions allow us to stop and restart (X_t) on the milestones without knowing its history. In applications, usually (X_t) is Langevin or overdamped Langevin dynamics, and Ω is a subset of Euclidean space.

We write \mathbb{P} , \mathbb{E} for all probability measures and expectations, with superscripts \mathbb{P}^x (resp. \mathbb{P}^ξ) to indicate a starting point x (resp. distribution ξ). The symbol \sim will indicate equality in probability law. We will use the words *distribution* and *probability measure* interchangeably. Total variation norm will be denoted by $\|\cdot\|_{TV}$. Our analysis below will mostly take place in an idealized setting where we assume infinite sampling on the milestones. In this setting, distributions are smooth (if state space is continuous) and the total variation norm is the appropriate one.

Recall we are interested in computing the MFPT of (X_t) from a reactant set R to a product set P . Throughout we consider fixed disjoint product and reactant sets $P, R \subset \Omega$. When R is not a single point, we will start (X_t) from a fixed probability measure ρ on R . If R is a single point, $\rho = \delta_R$, the delta distribution at R . As discussed above, Milestoning allows for an efficient computation of the MFPT of (X_t) to P , starting at ρ . It is useful to think of P as a sink, and R as a source for (X_t) . More precisely, *we assume that when (X_t) reaches P , it immediately restarts on R according to ρ* . Obviously, this assumption has no effect on the MFPT to P . It will be useful, however, for computational and theoretical considerations.

Many of the results below follow from well-known theorems in probability theory. However, because of the special source-sink structure of (X_t) , simpler proofs are often available, and we include them for clarity and completeness.

2.2 The milestones and semi-Markov viewpoint

We write $M \subset \Omega$ for the space of milestones used for parallelizing the computation of the MFPT. Each point $x \in M$ belongs to a *milestone* $M_x \subset M$.

Thus, M is the union of all the milestones. We assume there are finitely many milestones, each of which is a closed set. Moreover, we demand that (X_t) passes through the intersection of two milestones with probability 0 – thus, (X_t) can only cross one milestone at a time. This can be accomplished for Langevin or overdamped Langevin dynamics by taking the milestones to be codimension 1 with pairwise intersections of codimension 2 or larger; see Figure 1. The sets P and R will be two of the milestones. We always start (X_t) on M .

By following the sequence of milestones crossed by (X_t) , we obtain a sequence of points (J_n) in M . See Figure 1. We now describe (J_n) more precisely. Let θ_n be the n th milestone crossing time for (X_t) , defined recursively by $\theta_0 = 0$ and

$$\text{if } X_{\theta_n} = x, \text{ then } \theta_{n+1} := \inf\{t > \theta_n : X_t \in M_y \text{ for some } M_y \neq M_x\}.$$

Note that by a milestone crossing, we mean a crossing of a milestone different from the previous one. The sequence of milestone crossing points is $J_n = X_{\theta_n}$.

We show now that (X_t) can be partially reconstructed from (J_n) and (θ_n) . Let (Y_t) be defined¹ by setting $Y_t = J_n$ whenever $\theta_n \leq t < \theta_{n+1}$. Then (X_t) and (Y_t) agree at each milestone crossing time $t = \theta_n$, $n = 0, 1, 2, \dots$ and (Y_t) is obtained from (X_t) by throwing away the path of (X_t) between milestone crossings, keeping only the endpoints. It follows that (X_t) and (Y_t) have the same MFPT to P . Thus, for our purposes it is enough to study (Y_t) . We note that (Y_t) , like (X_t) , immediately restarts at ρ upon reaching P .

By our assumptions above, (J_n) is a Markov chain on M , and (Y_t) is a semi-Markov process on M , meaning it has the Markov property at crossing times. We write $K(x, dy)$ for the transition kernel of (J_n) . Thus, if the initial distribution of (J_n) is $J_0 \sim \xi$, then the distribution at time n is $\mathbb{P}^\xi(J_n \in \cdot) = \xi K^n$. We also write $\xi K^n f := \mathbb{E}^\xi[f(J_n)]$ and $\xi f := \int_M f(x) \xi(dx)$ for suitable functions f .

The following notation will be needed. For $x \in M$, define local first passage times

$$\tau_M^x = \inf\{t > 0 : Y_t \in M_y \text{ for some } M_y \neq M_x\}.$$

Thus, τ_M^x is the first time for (Y_t) to cross some milestone other than M_x , starting at $Y_0 = x$. In particular, if $X_{\theta_{n-1}} = x$, then $\theta_n \sim \theta_{n-1} + \tau_M^x$. We also define τ_P to be the first time to cross P and σ_P the number of crossings before reaching P ,

$$\tau_P = \inf\{t > 0 : Y_t \in P\}, \quad \sigma_P = \min\{n \geq 0 : J_n \in P\}.$$

We are interested in $\mathbb{E}^\rho[\tau_P]$, the MFPT from ρ to P .

¹When (Y_t) has a probability density, it corresponds to the density $p(x, t)$ from [5] for the last milestone point passed.

3 Invariant measure and MFPT

3.1 Stationary distribution on the milestones

The MFPT will be estimated via short trajectories between milestones. An important ingredient is the correct starting distribution for these trajectories. Exact Milestoning makes use of a *stationary flux* of (X_t) on the milestones, which corresponds² to the stationary distribution μ of (J_n) . It is worth noting that Milestoning can also be made exact by choosing milestones as isocommittor surfaces [54]. The advantage of the formulation here is that the milestones can be arbitrary.

Some assumption is required to guarantee the existence of a stationary flux. We adopt the following sufficient condition, which we assume holds throughout:

$$\mathbb{E}^\xi[\tau_P] \text{ and } \mathbb{E}^\xi[\sigma_P] \text{ are finite for all probability measures } \xi \text{ on } M.$$

This ensures that (Y_t) reaches P in finite expected time and does not have infinitely many milestone crossings in finite time. The condition can be readily verified in the standard settings for milestoning discussed above. Using this assumption and the source-sink structure of the dynamics – namely, that (Y_t) immediately restarts at ρ upon reaching P – we show in Theorem 1 below that μ exists.

Theorem 1. (J_n) has an invariant probability measure μ defined by

$$\mu(\cdot) := \mathbb{E}^\rho \left[\sum_{n=0}^{\sigma_P} \mathbb{1}_{\{J_n \in \cdot\}} \right] \mathbb{E}^\rho[\sigma_P + 1]^{-1}.$$

where $\mathbb{1}_{\{J_n \in C\}} = 1$ if $J_n \in C$ and otherwise $\mathbb{1}_{\{J_n \in C\}} = 0$.

Proof. Define $\nu(\cdot) = \mathbb{E}^\rho \left[\sum_{n=0}^{\sigma_P} \mathbb{1}_{\{J_n \in \cdot\}} \right]$ and observe that

$$\nu(\cdot) = \sum_{n=0}^{\infty} \sum_{m=0}^n \mathbb{P}^\rho(J_m \in \cdot \mid \sigma_P = n) \mathbb{P}^\rho(\sigma_P = n) = \sum_{n=0}^{\infty} \mathbb{P}^\rho(J_n \in \cdot, \sigma_P \geq n).$$

If $C \cap R = \emptyset$, by bounded convergence,

$$\int_M \nu(dx) K(x, C) = \sum_{n=0}^{\infty} \mathbb{P}^\rho(J_{n+1} \in C, \sigma_P \geq n) = \sum_{n=0}^{\infty} \mathbb{P}^\rho(J_n \in C, \sigma_P \geq n) = \nu(C),$$

where the second equality uses $\mathbb{P}^\rho(J_0 \in C) = 0$ and $J_{n+1} \notin R \Rightarrow \sigma_P \neq n$. If $C \subset R$,

$$\begin{aligned} \int_M \nu(dx) K(x, C) &= \sum_{n=0}^{\infty} \mathbb{P}^\rho(J_{n+1} \in C, \sigma_P = n) + \sum_{n=0}^{\infty} \mathbb{P}^\rho(J_{n+1} \in C, \sigma_P \geq n+1) \\ &= \rho(C) - \mathbb{P}^\rho(J_0 \in C, \sigma_P \geq 0) + \sum_{n=0}^{\infty} \mathbb{P}^\rho(J_n \in C, \sigma_P \geq n) = \nu(C). \end{aligned}$$

²Our μ is the same as the appropriately normalized stationary flux q in other Milestoning papers. We use μ instead of q to emphasize that here it is a probability measure, not a density.

□

We will show below that (J_n) converges to μ under appropriate conditions. In that case μ is unique and we will call μ the *stationary distribution* of (J_n) . A successful application of Exact Milestoning will require some technique for sampling μ . The algorithm we present (Algorithm 1 below) is based on convergence of the distribution of (J_n) to μ in total variation. We demonstrate this convergence in Theorem 4 under an additional assumption on (J_n) .

It is worth noting that the proof of Theorem 1 leads to the following representation of μ as a Neumann series. The representation is given in Corollary 2 below. This representation can be used, in principle, to sample μ without additional assumptions on (J_n) . The Neumann series is written in terms of the *transient kernel*

$$(1) \quad \bar{K}(x, dy) = \begin{cases} K(x, dy), & x \notin P \\ 0, & x \in P \end{cases}.$$

$\bar{K}(x, dy)$ corresponds to a modified version of (J_n) that is absorbed (killed) on P .

Corollary 2. *We have*

$$(2) \quad \lim_{n \rightarrow \infty} \|\nu(M)^{-1} \sum_{i=0}^{n-1} \rho \bar{K}^i - \mu\|_{TV} = 0.$$

Proof. Recall that $\mu = \nu/\nu(M)$ where

$$\nu(\cdot) = \sum_{n=0}^{\infty} \mathbb{P}^\rho(J_n \in \cdot, \sigma_P \geq n) = \sum_{n=0}^{\infty} \rho \bar{K}^n.$$

Moreover,

$$(3) \quad \sup_{|f| \leq 1} \left| \nu(M)^{-1} \sum_{i=0}^{n-1} \rho \bar{K}^i f - \mu f \right| \leq \nu(M)^{-1} \sum_{i=n}^{\infty} \mathbb{P}^\rho(\sigma_P \geq i),$$

and the right hand side of (3) is summable since by assumption $\mathbb{E}^\rho[\sigma_P] < \infty$. □

3.2 Milestoning equation for the MFPT

Equipped with an invariant measure μ , we are now able to state the Milestoning equation (4) for the MFPT. In Exact Milestoning, this equation is used to efficiently compute the MFPT. The algorithm is based on two principles: first, many trajectories can be simulated in parallel to estimate τ_M^x for various x ; and second, the stationary distribution μ can be efficiently estimated through a technique based on power iteration. See the right hand side of equation (4) below.

The gain in efficiency comes from the fact that the trajectories used to estimate τ_M^x are much shorter than trajectories from R to P . Whether we can efficiently sample μ may depend somewhat on whether we have a good initial guess. When (X_t) is Langevin dynamics, we have found in some cases the canonical Gibbs distribution is a sufficiently good guess. See [5] and [6] for details and discussion.

Theorem 3. *Let μ be defined as above. Then $\mu(P) > 0$ and*

$$(4) \quad \mu(P)\mathbb{E}^\rho[\tau_P] = \int_M \mu(dx)\mathbb{E}^x[\tau_M^x] := \mathbb{E}^\mu[\tau_M].$$

Proof. The assumption $\mathbb{E}^\rho[\sigma_P] < \infty$ shows that $\mu(P) > 0$. For any $x \in M$,

$$\begin{aligned} \mathbb{E}^x[\tau_P] &= \int_M \mathbb{E}^x[\tau_P | Y_{\tau_M^x} = y] K(x, dy) \\ &= \int_M \mathbb{E}^x[\tau_M^x | Y_{\tau_M^x} = y] K(x, dy) + \int_{M \setminus P} \mathbb{E}^x[\tau_P - \tau_M^x | Y_{\tau_M^x} = y] K(x, dy) \\ &= \mathbb{E}^x[\tau_M^x] + \int_{M \setminus P} \mathbb{E}^y[\tau_P] K(x, dy). \end{aligned}$$

Thus,

$$\mathbb{E}^\mu[\tau_P] = \mathbb{E}^\mu[\tau_M] + \int_{M \setminus P} \int_M \mu(dx)\mathbb{E}^y[\tau_P] K(x, dy) = \mathbb{E}^\mu[\tau_M] + \int_{M \setminus P} \mu(dy)\mathbb{E}^y[\tau_P],$$

and so

$$\mathbb{E}^\mu[\tau_M] = \int_P \mu(dy)\mathbb{E}^y[\tau_P] = \mu(P)\mathbb{E}^\rho[\tau_P].$$

□

In Section 4 below we present the Exact Milestoning algorithm (Algorithm 1) recently used in [5] and [6]. The algorithm uses a technique which combines coarse-graining and power iteration to sample μ . Consistency of power iteration algorithms are justified via Theorem 4 below, where we show $\xi K^n \rightarrow \mu$ as $n \rightarrow \infty$. Though we emphasize that there are a range of possibilities for sampling μ (for example, algorithms based on (2) or (7) below) we note that Algorithm 1 was shown to be efficient for computing the MFPT in the entropic barrier example of [5] and the random energy landscapes example of [6].

3.3 Convergence to stationarity

In this section we justify the consistency of power iteration-based methods for sampling μ by showing that ξK^n converges to μ in total variation norm as $n \rightarrow \infty$. The theorem requires an extra assumption – aperiodicity of the jump chain (J_n) .

Theorem 4. *Suppose that (J_n) is aperiodic in the following sense:*

$$(5) \quad g.c.d. \{n \geq 1 : \mathbb{P}^\rho(\sigma_P = n - 1) > 0\} = 1.$$

Then for all probability measures ξ on M ,

$$(6) \quad \lim_{n \rightarrow \infty} \|\mathbb{P}^\xi(J_n \in \cdot) - \mu\|_{TV} \equiv \lim_{n \rightarrow \infty} \|\xi K^n - \mu\|_{TV} = 0.$$

In particular, μ is unique.

Proof. We use a simple coupling argument. Let (H_n) be an independent copy of (J_n) and let $J_0 \sim \xi$ and $H_0 \sim \mu$. For $n \geq 0$, let S_n (resp. T_n) be the times at which (J_n) (resp. (H_n)) hit P for the $(n + 1)$ st time. Then $S_{n+1} - S_n$, $n \geq 0$, are iid random variables with finite expected value and nonlattice distribution, and $(S_{n+1} - S_n)_{n \geq 0} \sim (T_{n+1} - T_n)_{n \geq 0}$. It follows that $(S_n - T_n)_{n \geq 0}$ is a mean zero random walk with nonlattice step distribution. Thus, its first time to hit 0 is finite almost surely. So

$$\zeta := \inf\{n \geq 0 : J_n \in P, H_n \in P\}$$

obeys $\mathbb{P}(\zeta \geq n) \rightarrow 0$ as $n \rightarrow \infty$. Note that $J_n \sim H_n$ whenever $\zeta < n$. Thus

$$|\mathbb{P}^\xi(J_n \in C) - \mathbb{P}^\mu(H_n \in C)| \leq 2\mathbb{P}(\zeta \geq n).$$

Since μ is stationary for (H_n) we have $\mathbb{P}^\mu(H_n \in C) = \mu(C)$. Now

$$\|\mathbb{P}^\xi(J_n \in \cdot) - \mu\|_{TV} = \sup_{C \subset M} |\mathbb{P}^\xi(J_n \in C) - \mu(C)| \leq 2\mathbb{P}(\zeta \geq n),$$

which establishes the convergence result. To see uniqueness, suppose ξ is another invariant probability measure for (J_n) ; then the last display becomes $\|\xi - \mu\|_{TV} \leq 2\mathbb{P}(\zeta \geq n)$. Letting $n \rightarrow \infty$ shows that $\xi \sim \mu$. \square

We now consider a class of problems where there is a smooth one-dimensional *reaction coordinate* $\psi : \Omega \rightarrow [0, 1]$ tracking progress of (X_t) from R to P . In this case $\psi|_R \equiv 0$, $\psi|_P \equiv 1$, the milestones M_1, \dots, M_m are disjoint level sets of ψ , and $R = M_1$, $P = M_m$. The jump chain (J_n) can only hop between neighboring milestones, unless it is at P . That is, if $J_n \in M_i$ for $i \notin \{1, m\}$, then $J_{n+1} \in M_{i-1}$ or $J_{n+1} \in M_{i+1}$; if $J_n \in M_1$ then $J_{n+1} \in M_2$; and if $J_n \in M_m$, then $J_{n+1} \in M_1$. Suppose that if $J_n \in M_i$ for $i \notin \{1, m\}$, then $J_{n+1} \in M_{i-1}$ with probability in $(0, 1)$. Then the aperiodicity assumption (5) is satisfied if and only if m is odd. This is due to the fact that, if $J_0 \in M_1$, then $J_{m-1} \in M_m$ and $J_{m+1} \in M_m$ with positive probability, and m and $m + 2$ are coprime when m is odd. On the other hand, if m is even then the conclusion of Theorem 4 cannot hold. To see this, let m be even and suppose J_0 is supported in an odd-indexed milestone. Then J_{2n} is always supported on an odd-indexed milestone, while J_{2n+1} is always supported on an even-indexed milestone.

Theorem 4 establishes convergence the distribution of J_n to μ in total variation norm. Even when (J_n) is not aperiodic, it converges in a time-averaged sense. Thus, problems in sampling μ arising from aperiodicity can be managed by averaging over time. More precisely, we have the following version of the Birkhoff ergodic theorem:

Theorem 5. Let $J_0 \sim \xi$, with ξ a probability measure on M . For bounded measurable $f : M \rightarrow \mathbb{R}$,

$$(7) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} f(J_i) \stackrel{\text{a.s.}}{=} \int_M f d\mu \equiv \mu f.$$

Proof. Let S_n be the times at which (J_n) hits P for the $(n+1)$ st time, and define

$$f_n = \sum_{i=S_n+1}^{S_{n+1}} f(J_i).$$

Note that f_n , $n \geq 0$, are iid. Let $k(n) = \max\{k : S_k \leq n\}$ and write

$$\frac{1}{n} \sum_{i=0}^{n-1} f(J_i) = \frac{1}{n} \sum_{i=0}^{S_0} f(J_i) + \frac{1}{n} \sum_{i=0}^{k(n)-1} f_i + \frac{1}{n} \sum_{i=S_{k(n)}+1}^n f(J_i)$$

Since (J_n) hits P in finite time a.s., $n - S_{k(n)}$ and S_0 are finite a.s. Thus,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} f(J_i) = \lim_{n \rightarrow \infty} \frac{k(n)-1}{n} \frac{1}{k(n)-1} \sum_{i=0}^{k(n)-1} f_i.$$

Notice $R_n := S_{n+1} - S_n$, $n \geq 0$ are iid with finite expectation and

$$\frac{R_0 + \dots + R_{k(n)-1}}{k(n)} \leq \frac{n - S_0}{k(n)} \leq \frac{R_0 + \dots + R_{k(n)}}{k(n)}.$$

By the previous two displays and the law of large numbers,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} f(J_i) = \frac{\mathbb{E}[f_0]}{\mathbb{E}[R_0]} = \mathbb{E}^\rho[\sigma_P + 1]^{-1} \sum_{i=0}^{\sigma_P} \mathbb{E}^\rho[f(J_i)] = \frac{\nu f}{\nu(M)} = \mu f,$$

with ν defined as in Theorem 4. □

Markov chains for which the conclusion of Theorem 4 hold are called *Harris ergodic*. It is worth noting that a slightly stronger aperiodicity condition leads to a limit for the distribution of Y_t . More precisely, suppose (5) holds and for each $x \in M \setminus P$ and $y \in M$, $\mathbb{P}^x(\tau_1 \in \cdot | J_1 = y)$ is nonlattice. Then for any $C \subset M$ and μ -a.e. x ,

$$(8) \quad \lim_{t \rightarrow \infty} \mathbb{P}^x(Y_t \in C) = \frac{\int_C \mu(dy) \mathbb{E}^y[\tau_M^y]}{\int_M \mu(dy) \mathbb{E}^y[\tau_M^y]}.$$

See [4] for details³ and proof.

³When the right hand side of (8) has a density, it is the same as the stationary probability density $p(x)$ in [5] for the last milestone point passed.

4 Exact Milestoning algorithm

We now describe in detail an algorithm for sampling μ and the MFPT $\mathbb{E}^\rho[\tau_P]$, used successfully in [5] and [6]. We assume throughout this section that the conclusion of Theorem 4 holds. Let ξ be an initial guess for μ . (If (X_t) is Brownian or Langevin dynamics, we usually take ξ to be the canonical Gibbs distribution.) We write M_i for the distinct milestones, so that $M = \cup_i M_i$. The algorithm will produce approximations

$$\xi \equiv \mu^{(0)}, \mu^{(1)}, \mu^{(2)}, \dots$$

of μ . Let $\mu_i^{(n)}$ be the *non-normalized* restriction of $\mu^{(n)}$ to M_i , and define

$$\mathbb{E}^{\mu_i^{(n)}}[\tau_M] := \mu^{(n)}(M_i)^{-1} \int_{M_i} \mu_i^{(n)}(dx) \mathbb{E}^x[\tau_M^x],$$

For $C \subset M_j$ we will also use the notation

$$a_{ij}^{(n)}(C) = \mu^{(n-1)}(M_i)^{-1} \int_{M_i} \mu_i^{(n-1)}(dx) K(x, C).$$

Below we think of $a_{ij}^{(n)}$ and $\mu_i^{(n)}$ as either distributions or densities. The $a_{ij}^{(n)}$ are obtained from trajectory fragments between milestone crossings. A simple Monte Carlo scheme for estimating these distributions is as follows. Let x_1, \dots, x_L be iid samples from the distribution $\mu_i^{(n-1)}/\mu^{(n-1)}(M_i)$. Starting at each $x_\ell \in M_i$, simulate (X_t) until it crosses the next milestone, say at the point $y_\ell \in M_j$. If we idealize by assuming the simulation of (X_t) is done exactly, then by Chebyshev's inequality,

$$\mathbb{P} \left(\left| a_{ij}^{(n)}(C) - \int_C \frac{1}{L} \sum_{\ell=1}^L \delta_{y_\ell}(dy) \right| > \epsilon \right) \leq \frac{a_{ij}^{(n)}(C) - a_{ij}^{(n)}(C)^2}{L\epsilon^2},$$

where δ_y is the Dirac delta distribution at y . We therefore write, for $y \in M$,

$$(9) \quad a_{ij}^{(n)}(y) \approx \frac{1}{L} \sum_{\ell=1}^L \tilde{\delta}_{y_\ell}(y),$$

where $\tilde{\delta}_{y_\ell}$ is either some suitable approximation to the identity at y_ℓ , or simply a delta function at y_ℓ . Thus, in Algorithm 1 we think of $a_{ij}^{(n)}$ and $\mu_i^{(n-1)}$ as either densities in the former case, or as distributions in the latter. The local mean first passage times (i.e., the times between successive milestone crossings) are approximated by the sample means

$$\mathbb{E}^{\mu_i^{(n-1)}}[\tau_M] \approx \frac{1}{L} \sum_{\ell=1}^L \tau_M^{x_\ell}.$$

Algorithm 1 Exact Milestoning algorithm.

Input: Milestones $M = \cup_{j=1}^m M_j$, initial guess ξ , and tolerance $\varepsilon > 0$ for the absolute error in the MFPT.

Output: Estimates for μ , local MFPTs $\mathbb{E}^\mu[\tau_M]$, and overall MFPT $\mathbb{E}^\rho[\tau_P]$.

```

 $\mu^{(0)} \leftarrow \xi$ 
 $T^{(0)} \leftarrow +\infty$ 
for all  $n = 1, 2, \dots$  do
  for  $i = 1$  to  $m$  do
    Estimate  $a_{ij}^{(n)}$  and  $\mathbb{E}^{\mu^{(n-1)}}[\tau_M]$ 
     $\mathbf{A}_{ij}^{(n)} \leftarrow a_{ij}^{(n)}(M_j)$ 
  end for
  Solve  $\mathbf{w}^T \mathbf{A} = \mathbf{w}^T$  (with  $\mathbf{A} = (\mathbf{A}_{ij}^{(n)}) \in \mathbb{R}_{\geq 0}^{m \times m}$  and  $\mathbf{w} = (w_1, \dots, w_m) \in \mathbb{R}_{\geq 0}^m$ )

  for  $j = 1$  to  $m$  do
     $\mu_j^{(n)} \leftarrow \sum_{i=1}^m w_i a_{ij}^{(n)}$ 
  end for
  Normalize  $\mu^{(n)}$ 
   $T^{(n)} \leftarrow \mu^{(n)}(P)^{-1} \mathbb{E}^{\mu^{(n-1)}}[\tau_M]$ 
  if  $|T^{(n)} - T^{(n-1)}| < \varepsilon$  then
    break
  end if
end for
return  $(\mu^{(n)}, \mathbb{E}^{\mu^{(n-1)}}[\tau_M], T^{(n)})$ 

```

It is important to realize that we do not need to store the full coordinates of each y_ℓ in memory. Instead, it suffices to use a data-structure that keeps track of the pairs (y_ℓ, M_j) . The actual coordinates of each point can be written to disk and read from it as needed.

The eigenvalue problem in Algorithm 1 involves a stochastic matrix $\mathbf{A} \in \mathbb{R}_{\geq 0}^{m \times m}$ that is sparse. Indeed, the i -th row corresponds to milestone M_i and may have only as many non-zero entries as the number of neighboring milestones M_j . In practice, to solve the eigenvalue problem we can use efficient and accurate Krylov subspace solvers [28] such as Arnoldi iteration [39] to obtain \mathbf{w} without computing all the other eigenvectors.

In Algorithm 1, if $\mathbf{w}_i := \mu_i^{(n-1)}(M_i)$ is used instead of the solution \mathbf{w} to $\mathbf{w}^T \mathbf{A} = \mathbf{w}^T$, then the algorithm approximates μ by simple power iteration, $\mu^{(n)} = \xi K^n$. The reason for defining the weights as the solution to $\mathbf{w}^T \mathbf{A} = \mathbf{w}^T$ is practical: we have found that it gives faster convergence of the iterations, at no apparent cost to accuracy. It can be seen as a version of power iteration that uses coarse-graining. See [5, 6] for applications of the algorithm in Exact Milestoning and [28, 39] for related discussions.

Finally, we mention the fact that pseudo-random number generators (PRNGs) can only produce a finite amount of pseudo-random numbers. Once the maximum amount is reached, the generators may silently reuse the previous random

numbers in the same order. It has been noted [13] that this phenomenon leads to unphysical artifacts in simulations. The simplest approach to properly use PRNGs (and avoid the aforementioned artifacts altogether) consists of reseeding the generator from time to time, obtaining the new seeds from high-quality entropy sources such as those available in modern computer hardware (see [17, 33] for more details).

5 Error analysis

5.1 Stationary distribution error

In practice, due to time discretization error, we cannot generate trajectories exactly according to the transition kernel K . Instead, we can generate trajectories according to a numerical approximation K_ϵ . We investigate here whether such schemes are consistent, that is, whether powers of K_ϵ or K converge to a distribution $\mu_\epsilon \approx \mu$. We emphasize that, even though we account for time discretization here, we still assume infinite sampling, and thus for a given $x \in M$, $K_\epsilon(x, dy)$ may be a continuous distribution. See Section 5.2 below for related remarks and a discussion of how time discretization errors affect the Exact Milestoning estimate of the MFPT.

The following theorem, restated from [23], establishes consistency of iteration schemes based on Theorem 4 when K_ϵ is sufficiently close to K and (J_n) is geometrically ergodic. After the theorem, in Lemma 7 and Theorem 8 we give natural conditions for geometric ergodicity of (J_n) .

Theorem 6. *Suppose (J_n) is geometrically ergodic: there exists $\kappa \in (0, 1)$ such that*

$$\sup_{x \in M} \|\delta_x K^n - \mu\|_{TV} = O(\kappa^n).$$

Let $\{K_\epsilon\}$ be a family of stochastic kernels with $K_0 = K$, assumed to act continuously on \mathcal{B} , such that

$$(10) \quad \lim_{\epsilon \rightarrow 0} \sup_{|f| \leq 1} \|K_\epsilon f - Kf\|_\infty = 0.$$

Then for each $\hat{\kappa} \in (\kappa, 1)$, there is $\delta > 0$ such that for each $\epsilon \in [0, \delta)$, K_ϵ has a unique invariant probability measure μ_ϵ , and

$$\begin{aligned} \sup_{\epsilon < \delta} \sup_{x \in M} \|\delta_x K_\epsilon^n - \mu_\epsilon\|_{TV} &= O(\hat{\kappa}^n), \\ \lim_{\epsilon \rightarrow 0} \|\mu_\epsilon - \mu\|_{TV} &= 0. \end{aligned}$$

Geometric ergodicity is inconvenient to check directly. We give two sufficient conditions for geometric ergodicity of (J_n) . The first condition is a uniform lower bound on the probability to reach P in N steps; see Lemma 7. We use this to obtain a strong Feller condition in Theorem 8. The latter is a very natural condition and is easy to verify in some cases, for instance when (X_t) is a nondegenerate diffusion and the milestones are sufficiently regular.

Lemma 7. *Suppose that there exists $\lambda \in (0, 1)$ and $N \in \mathbb{N}$ such that for all $x \in M$, $\mathbb{P}^x(J_{N-1} \in P) \geq \lambda > 0$. Then (J_n) is geometrically ergodic:*

$$\sup_{x \in M} \|\delta_x K^n - \mu\|_{TV} \leq \lambda^{-1}(1 - \lambda)^{\lfloor n/N \rfloor}.$$

Proof. Let $\xi_1, \xi_2 \in \mathcal{P}$, consider the signed measure $\xi = \xi_1 - \xi_2$ and compute

$$\begin{aligned} \|\xi_1 K^N - \xi_2 K^N\|_{TV} &= \sup_{|f| \leq 1} \left| \int_M \int_M \xi(dy) K^N(y, dz) f(z) \right| \\ &= \sup_{|f| \leq 1} \left| \int_M \int_M \xi(dy) (K^N(y, dz) - \lambda \rho(dz)) f(z) \right| \\ &= \sup_{|f| \leq 1} \left| \int_M \xi(dy) \int_M (K^N(y, dz) - \lambda \rho(dz)) f(z) \right| \\ &\leq (1 - \lambda) \sup_{|f| \leq 1} \left| \int_M \xi(dy) f(y) \right| = (1 - \lambda) \|\xi_1 - \xi_2\|_{TV}. \end{aligned}$$

The last line uses the fact that $K^N(y, dz) - \lambda \rho(dz)$ is a positive measure. This shows that K^N is a contraction mapping on \mathcal{P} with contraction constant $(1 - \lambda)$. Observe also that $\|\xi_1 K - \xi_2 K\|_{TV} \leq \|\xi_1 - \xi_2\|_{TV}$. The result now follows from the contraction mapping theorem. See for instance Theorem 6.40 of [18]. \square

Note that the λ in Lemma (7) is a quantity that can be estimated, at least in principle, by running trajectories of (X_t) starting at x which cross $N - 1$ milestones before reaching P . However, this is likely impractical for the same reason direct estimation of the MFPT is impractical – the trajectories would be too long. One alternative would be to compute the probability $P^i(J_{N-1}^A \in P)$ for the Markov chain (J_n^A) on $\{1, \dots, m\}$ with transition matrix A , and use the minimum over $i \in \{1, \dots, m\}$ as a proxy for λ . Even without a practical way to estimate λ , we believe the characterization of Lemma 7 is useful for understanding the convergence rate.

Lemma 7 leads to the following condition for geometric ergodicity of (J_n) .

Theorem 8. *Suppose that M is compact and (J_n) is a strong Feller chain which is aperiodic in the sense of (5). Then (J_n) is geometrically ergodic.*

Proof. Let $\epsilon \in (0, \mu(P))$. By Theorem 4, for each $x \in M$ there is $N_x \in \mathbb{N}$ such that $\mathbb{P}^x(J_n \in P) \geq \epsilon$ for all $n \geq N_x$. Because (J_n) is strong Feller, the map $x \rightarrow \mathbb{P}^x(J_n \in P)$ is continuous. By compactness of M , it follows that for any $\lambda \in (0, \epsilon)$ there is $N \in \mathbb{N}$ such that $\mathbb{P}^y(J_n \in P) \geq \lambda$ for all $y \in M$ and $n \geq N - 1$. Theorem 7 now yields the result. \square

5.2 MFPT error

As discussed above, Equation 4 can be used to estimate the MFPT $\mathbb{E}^\rho[\tau_P]$ by sampling μ and local MFPTs τ_M^x . The error in this estimate has two sources. First, in general we only have an approximation $\tilde{\mu} \equiv \mu_\epsilon$ of μ . The second source

of error is in the sampling of τ_M^x , due to the fact that we can only simulate a time discrete version ($\tilde{X}_{n\delta t}$) of (X_t) . In Theorem 9 below we give an explicit formula for the numerical error of the MFPT in terms of these two sources. We first need the following notation. Let $\tilde{\tau}_M^x$ be the minimum of all $n\delta t > 0$ such that the line segment between $\tilde{X}_{n\delta t}$ and $\tilde{X}_{(n+1)\delta t}$ intersects $M \setminus M_x$, and define

$$\mathbb{E}^{\tilde{\mu}}[\tilde{\tau}_M] := \int_M \tilde{\mu}(dx) \mathbb{E}^x[\tilde{\tau}_M^x].$$

Theorem 9 below gives an expression for the error in the *original* Milestoning as well as in Exact Milestoning.

Theorem 9. *There exists a nonnegative function ϕ such that*

$$(11) \quad \begin{aligned} |\mathbb{E}^\rho[\tau_P] - \tilde{\mu}(P)^{-1} \mathbb{E}^{\tilde{\mu}}[\tilde{\tau}_M]| &\leq c_1 |\mu(P)^{-1} - \tilde{\mu}(P)^{-1}| \\ &\quad + \tilde{\mu}(P)^{-1} (c_2 \|\mu - \tilde{\mu}\|_{TV} + \phi(\delta t)), \end{aligned}$$

where

$$c_1 := \mathbb{E}^\mu[\tau_M], \quad c_2 := \sup_{x \in M} \mathbb{E}^x[\tau_M^x].$$

Proof. Note that

$$\begin{aligned} |\mathbb{E}^\rho[\tau_P] - \tilde{\mu}(P)^{-1} \mathbb{E}^{\tilde{\mu}}[\tilde{\tau}_M]| &= |\mu(P)^{-1} \mathbb{E}^\mu[\tau_M] - \tilde{\mu}(P)^{-1} \mathbb{E}^{\tilde{\mu}}[\tilde{\tau}_M]| \\ &\leq |\mu(P)^{-1} \mathbb{E}^\mu[\tau_M] - \tilde{\mu}(P)^{-1} \mathbb{E}^\mu[\tau_M]| \\ &\quad + |\tilde{\mu}(P)^{-1} \mathbb{E}^\mu[\tau_M] - \tilde{\mu}(P)^{-1} \mathbb{E}^{\tilde{\mu}}[\tau_M]| \\ &\quad + |\tilde{\mu}(P)^{-1} \mathbb{E}^{\tilde{\mu}}[\tau_M] - \tilde{\mu}(P)^{-1} \mathbb{E}^{\tilde{\mu}}[\tilde{\tau}_M]| \end{aligned}$$

where we have written $\mathbb{E}^{\tilde{\mu}}[\tau_M] := \int_M \tilde{\mu}(dx) \mathbb{E}^x[\tau_M^x]$. We may write

$$\phi(\delta t) = |\mathbb{E}^{\tilde{\mu}}[\tau_M] - \mathbb{E}^{\tilde{\mu}}[\tilde{\tau}_M]|$$

for the term depending only on time stepping error. Note that

$$\begin{aligned} |\mathbb{E}^\mu[\tau_M] - \mathbb{E}^{\tilde{\mu}}[\tau_M]| &= \left| \int_M \mu(dx) \mathbb{E}^x[\tau_M^x] - \int_M \tilde{\mu}(dx) \mathbb{E}^x[\tau_M^x] \right| \\ &\leq \left(\sup_{x \in M} \mathbb{E}^x[\tau_M^x] \right) \|\mu - \tilde{\mu}\|_{TV}. \end{aligned}$$

Combining the last three expressions yields the result. \square

Recall that in the above we have ignored errors from finite sampling. We now discuss the implications of those errors. In the original Milestoning, $\tilde{\mu}$ is the canonical Gibbs distribution on the milestones. In that setting, we can typically sample independently from $\tilde{\mu}$ on the milestones. Thus, the central limit theorem implies that the *true* error in the Milestoning approximation of $\mathbb{E}^\rho[\tau_P]$ is bounded above with high probability by the right hand side of (11) plus a constant times $1/\sqrt{N}$, where N is the number of samples. An analogous

argument applies to Exact Milestoning if $\tilde{\mu}$ is sampled by simple power iteration. For our coarse-grained version of power iteration in Algorithm 1, however, we obtain samples of $\tilde{\mu}$ which are not independent, and thus a more detailed analysis would be required to determine the additional error from finite sampling.

We do not analyze the time discretization error $\phi(\delta t)$ and instead refer the reader to [27] and references therein. Here we simply remark that, if (X_t) is a diffusion process, then under certain smoothness assumptions on the drift and diffusion coefficients of (X_t) and on M , we have $\phi(\delta t) = \theta(\sqrt{\delta t})$ when $(X_{n\delta t})$ is the standard Euler time discretization with time step δt . See [26] for details and proof. See also [8, 31] for numerical schemes that mitigate time discretization error in the MFPTs.

6 Illustrative examples

In this section we discuss two examples of Milestoning to illustrate the method.

We consider the solution, (X_t) , of the Brownian dynamics equation,

$$(12) \quad \begin{cases} dX_t = -\nabla U(X_t) dt + \sqrt{2\beta^{-1}} dB_t, \\ X_0 \sim \rho \end{cases}$$

where $U: \Omega \rightarrow \mathbb{R}$ is a smooth potential energy function, $\beta > 0$ is the inverse temperature, and (B_t) is a standard Brownian motion.

6.1 Müller-Brown potential

We begin with a system characterized by the Müller-Brown potential [44]. The energy function $U: \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ is given by the formula (see also the corresponding energy landscape in Figure 2)

$$\begin{aligned} U(x_1, x_2) = & -200 e^{-(x_1-1)^2-10x_2^2} - 100 e^{-x_1^2-10(x_2-\frac{1}{2})^2} \\ & - 170 e^{-\frac{13}{2}(x_1+\frac{1}{2})^2+11(x_1+\frac{1}{2})(x_2-\frac{3}{2})-\frac{13}{2}(x_2-\frac{3}{2})^2} \\ & + 15 e^{\frac{7}{10}(x_1+1)^2+\frac{3}{5}(x_1+1)(x_2-1)+\frac{7}{10}(x_2-1)^2}. \end{aligned}$$

This system is a commonly used benchmark for numerical methods for obtaining reaction rates.

We chose to partition Ω using a Voronoi tessellation (displayed in Figures 2 and 3) generated from a set of points gathered by the method of locally updated planes [51]. However, any other set of points could have been chosen (as we shall discuss in the next example). Figure 2 also shows our choice of reactant and product milestones.

For the numerical experiments to be detailed below, we solve the stochastic differential equation in (12) using the Euler-Maruyama scheme [42] with a time step length $\Delta t = 10^{-5}$ at a temperature determined by $\beta^{-1} = 5$. We use the




Figure 2: Graph of the Müller-Brown potential energy function. The milestones are shown as the overlaid line segments.

number of force evaluations as a measure of the computational cost of our methods and we note that the Euler-Maruyama method requires one force evaluation per time step.

We compared two types of experiments that we now describe. The first experiment consists of running Brownian dynamics trajectories started at the reactant milestone until they reach the product milestone. As soon as a trajectory reaches the product, we initiate a new trajectory from the reactant milestone and so on. We refer to these as *long* trajectories. In the second experiment we run Exact Milestoning starting the first iteration with exactly one phase space point at each of the milestones along the reaction path. Next, we run ten *short* trajectories per milestone per iteration. These short trajectories start at each milestone and stop whenever they reach any neighboring milestone, as described in Section 4.

Despite allowing the long trajectories to go on for approximately 2.5×10^9 force evaluations, only seven reach the product milestone. This leads to a poor approximation of the mean first passage time. By contrast, running the Exact Milestoning method for approximately 2×10^9 force evaluations, we obtain good estimates of the stationary distribution μ and the local mean first passage times.

The values of $\mu(M_i)$ are displayed in Figures 4 and 5. The empirical distributions corresponding to μ on some of the milestones are shown in Figure 6.

Figures 4 and 5 illustrate the non-equilibrium nature of Exact Milestoning. The stationary distribution that we compute differs noticeably from the equilibrium (canonical) distribution. Recall from Figures 2 and 3 that the reactant




Figure 3: Milestones represented as line segments. Some of the milestones are labeled and the reactant is colored in green while the product is shown in blue.




Figure 4: Phase space points in the empirical distributions of the stationary flux μ for two types of simulations: long trajectories using straight-forward Brownian dynamics (left) and Exact Milestoning (right). Despite the fact that the two types of simulations involved comparable amounts of computational effort, we see that the sampling in Exact Milestoning is much more exhaustive.

milestone, M_7 , is located at the lower-right minimum while the product milestone, M_0 , is at the global minimum in the upper-left side of the graph. With this in mind, we see that trajectories initiated at M_7 arrive at the intermediate minimum located close to the center of the graph and many of those trajectories return to the lower minimum, crossing M_7 again. This results in high values of $\mu(M_6)$ at the transition state, while the density at M_0 (the global minimum)




Figure 5: Values of $\mu(M_i)$ at some of the milestones in the Müller-Brown potential. The values correspond to the long trajectories (in red) and to Exact Milestoning (in blue), as discussed in Section 6.1. Not shown are the milestones other than M_i for $i = 0, \dots, 7$, where the sampling obtained from the long trajectories is insufficient for comparison.




Figure 6: Empirical distributions of the stationary flux obtained by long trajectories (in red) and Exact Milestoning (in blue) corresponding to the system in Section 6.1. Notice that the sampling of the long trajectories is very sparse at the milestones close to the product.

is significantly lower. Equilibrium considerations, which are inappropriate here, would suggest that most of the stationary density (and the stationary probability) is concentrated at the global minimum and that the weight at the transition state would be small.

6.2 Rough energy landscape

In this case, we present an example of Milestoning on the torus $\Omega = \mathbb{R}^2/\mathbb{Z}^2$. For our computations, we consider a uniformly spaced mesh of milestones with fixed product and reactant sets P, R ; see Figure 8. We model a rough energy

landscape by a potential energy function of the form:

$$(13) \quad U(x_1, x_2) = \operatorname{Re} \sum_{k_1=-N}^N \sum_{k_2=-N}^N z_{k_1, k_2} e^{2\pi i(k_1 x_1 + k_2 x_2)}$$

where Re denotes the real part of a complex number and $N \in \mathbb{N}$ is a constant that tunes the ruggedness of the potential. Each coefficient $z_{k_1, k_2} = a_{k_1, k_2} + i b_{k_1, k_2} \in \mathbb{C}$ is determined by the random variables a_{k_1, k_2} and b_{k_1, k_2} , which are distributed according to

$$\begin{cases} c, & \text{with probability } \frac{1}{2}, \\ 0, & \text{with probability } \frac{1}{2}, \end{cases}$$

with c itself being a uniform random variable in the interval $(-1, 1)$. Since N is fixed, a particular realization of the coefficients specified above completely determines the potential energy function U . The graph of the canonical density of a potential energy of the form discussed above is shown in Figure 7. Notice




Figure 7: Graph of the density of the canonical distribution corresponding to a rough energy landscape with $N = 7$ at temperature $\beta^{-1} = 1$.

that this class of energy functions generalizes the model for rough landscapes introduced by [58] and that similar potential energy functions have been used to model Wigner glasses [1].

We carry out Exact Milestoning in this example by solving boundary value problems, as described in [6]. The resulting stationary density obtained after convergence is shown in Figure 9.




Figure 8: Diagram showing the reactant state (orange square) and the product state (blue square) within the set of all milestones for the example in Section 6.2. Each milestone is an edge of one of the small squares in the diagram. (The total number of milestones has been decreased to enhance visibility.)

It is interesting to note that it has been argued [54] that an optimal choice of milestones would consist of using the level sets (also called *isocommittors*) of the committor function. These surfaces (see Figure 10) are typically hard to compute in practice, which makes the use of Exact Milestoning more appealing, as its results are independent of how the milestones are set up.

7 Conclusions

The main goal of this manuscript is to present a rigorous mathematical derivation, based on probability theory, of Exact Milestoning. While the theory of Exact Milestoning and accompanying numerical examples were discussed elsewhere [5, 6], the mathematical formulation in the earlier paper was not as rigorous as in this manuscript. Once this formulation is established, it opens the way for further communication between chemical physicists and mathematicians, and it bridges the gap between the communities for further development of an important tool for computer simulation.

Exact Milestoning belongs to a class of enhanced sampling methods for the calculation of kinetics. Most closely related approaches to Milestoning are the




Figure 9: Stationary density μ on the rough energy landscape of Section 6.2. The contour lines are the level sets of U . There are $2 \times 40 \times 40$ total milestones (shown as the segments in the overlaid grid).

Non-Equilibrium Umbrella sampling [55] and Trajectory Tilting [53]. The way in which trajectories are sampled is similar in all of these methods; however, the theoretical frameworks are different. For example, Milestoning allows the calculation of all the moments of the first passage time (FPT) distribution [5], and hence better estimates of the FPT can be constructed, a result that was not reported for other methods.

It is not necessary in Milestoning to establish or rely on metastability to estimate the average transition time. From this perspective the method is different from another exact approach —Transition Path Sampling [16]— that exploits the short duration of rare trajectories between metastable states. Exact Milestoning makes the sampled trajectories short by sampling trajectory fragments between boundaries of phase space cells or milestones. The statistics of short trajectories between milestones make it possible to investigate wide ranges of types of energy landscapes, which may be corrugated or not, as illustrated in the two examples in this manuscript. We have shown in the present manuscript that Exact Milestoning is both accurate and highly efficient.




Figure 10: Isocommittor surfaces for the rough energy landscape discussed in Section 6.2.

It is important to emphasize that the choice of the milestones in Exact Milestoning is arbitrary from a formal viewpoint. Efficiency considerations suggest that it is beneficial to select them following two criteria: (i) the milestones should be sufficiently close in the kinetic sense to make the trajectories short, and (ii) milestones should be chosen to make the number of iterations as small as possible. For example, the number of iterations can be small if the system is close to equilibrium and the milestones are expressed in a space of slow variables. Then an initial choice of the canonical distribution is quite accurate.

We comment that the method of Milestoning that was broadly used in the past (e.g., [35]) is approximate and assumes local equilibrium within the milestones. While corrections and further refinements were proposed [40, 30, 29], these approximations cannot be made exact and are similar in spirit to the local equilibrium and lag time approximations of Markov State Models [14]. Nevertheless, these approximations can be accurate with a proper choice of coarse variables. These types of approximations are very useful as the system grows in complexity and size and exact calculations become prohibitively expensive. Milestoning made it possible to investigate kinetics of enzymes [37] and transport through membranes [12] in agreement with experimental observations. These are systems of tens to hundreds of thousands of particles and time scales of milliseconds. It will be of considerable interest to re-evaluate these approximations for large systems with the method of Exact Milestoning. As the efficiency of Exact Milestoning increases with faster hardware, we are breaking scale barriers that were not accessible before to atomically detailed simulations.

8 Acknowledgments

This research was supported in part by a grant from the NIH GM59796 and from the Welch Foundation Grant No. F-1783. D. Aristoff gratefully acknowledges support from the National Science Foundation via the award NSF-DMS-1522398.

References

- [1] S. AKHANJEE AND J. RUDNICK, *Disorder induced transition into a one-dimensional Wigner glass*, Physical Review Letters, 99 (2007), pp. 236403, 0706.4462.
- [2] R.J. ALLEN, D. FRENKEL, AND P.R. TEN WOLDE, *em* Forward flux sampling-type schemes for simulating rare events: Efficiency analysis, The Journal of Chemical Physics, 124(19) (2006), pp. 463102.
- [3] P. ALLEN AND D. J. TILDESLEY, *Computer simulation of liquids*, Oxford Science Publications, Clarendon Press, 1989.
- [4] G. ALSMEYER, *On the Markov renewal theorem*, Stochastic Processes and their Applications, 50 (1994), pp. 37–56.
- [5] J. M. BELLO-RIVAS AND R. ELBER, *Exact milestoning*, The Journal of Chemical Physics, 142 (2015), pp. 094102.
- [6] J. M. BELLO-RIVAS AND R. ELBER, *Simulations of thermodynamics and kinetics on rough energy landscapes with milestoning*, Journal of Computational Chemistry, (2015).
- [7] P.G. BOLHUIS ET AL., *Transition path sampling: Throwing ropes over roughmountain passes, in the dark*. Annual Review of Physical Chemistry, 53 (2002) pp. 291-318.
- [8] B. BOUCHARD, S. GEISS, AND E. GOBET, *First time to exit of a continuous Itô process: general moment estimates and L_1 -convergence rate for discrete time approximations*, (2013), pp. 1–31, 1307.4247.
- [9] A.E. CARDENAS ET AL., *Unassisted Transport of N-Acetyl-L-tryptophanamide through Membrane: Experiment and Simulation of Kinetics*, Journal of Physical Chemistry B, 116(9) (2012), pp. 2739–2750.
- [10] A.E. CARDENAS AND R. ELBER, *Enhancing the capacity of molecular dynamics simulations with trajectory fragments*, in *Innovation in Biomolecular Modeling and Simulation*, T. Schlick, Editor. 2012, Royal Society of Chemistry: London.
- [11] A.E. CARDENAS AND R. ELBER, *Computational study of peptide permeation through membrane: searching for hidden slow variables*, Molecular Physics, 111 (2013), pp. 356–3578.

- [12] A. E. CARDENAS, R. SHRESTHA, L. J. WEBB, AND R. ELBER, *Membrane Permeation of a Peptide: It Is Better to be Positive*, The Journal of Physical Chemistry B, 119 (2015), pp. 6412–6420.
- [13] D. S. CERUTTI, R. DUKE, P. L. FREDDOLINO, H. FAN, AND T. P. LYBRAND, *A vulnerability in popular Molecular Dynamics packages concerning Langevin and Andersen dynamics*, Journal of Chemical Theory and Computation, 4 (2008), pp. 1669–1680.
- [14] J. D. CHODERA, N. SINGHAL, V. S. PANDE, K. A. DILL, AND W. C. SWOPE, *Automatic discovery of metastable states for the construction of Markov models of macromolecular conformational dynamics*, The Journal of Chemical Physics, 126 (2007), pp. 155101.
- [15] C. DELLAGO ET AL., *Transition path sampling and the calculation of rate constants*, The Journal of Chemical Physics, 108(5) (1998), pp. 1964–1977.
- [16] C. DELLAGO, P. G. BOLHUIS, AND P. L. GEISSLER, *Transition Path Sampling*, vol. 123, 2002.
- [17] Y. DODIS, A. SHAMIR, N. STEPHENS-DAVIDOWITZ, AND D. WICHS, *How to eat your entropy and have it too — Optimal recovery strategies for compromised RNGs*, in Advances in Cryptology CRYPTO 2014 SE - 3, J. Garay and R. Gennaro, eds., vol. 8617 of Lecture Notes in Computer Science, Springer Berlin Heidelberg, 2014, pp. 37–54.
- [18] R. DOUC, E. MOULINES, AND D. STOFFER, *Nonlinear time series: theory, methods and applications with R examples*, Chapman & Hall/CRC Texts in Statistical Science, Taylor & Francis, 2014.
- [19] ELBER, R., *A milestoning study of the kinetics of an allosteric transition: Atomically detailed simulations of deoxy Scapharca hemoglobin*, Biophysical Journal, 92(9) (2007), pp. L85–L87.
- [20] R. ELBER, K. KUCZERA, AND G.S. JAS, *The kinetics of helix unfolding: Molecular dynamics simulations with Milestoning*, Journal of Physical Chemistry A, 113 (2009), pp. 7461–7473.
- [21] R. ELBER AND A. WEST, *Atomically detailed simulation of the recovery stroke in myosin by Milestoning*, Proceedings of the National Academy of Sciences USA 107, (2010), pp. 5001–5005.
- [22] A. K. FARADJIAN AND R. ELBER, *Computing time scales from reaction coordinates by Milestoning*, Journal of Chemical Physics, 120 (2004), pp. 10880–10889.
- [23] D. FERRÉ, L. HERVÉ, AND J. LEDOUX, *Regular perturbation of V-geometrically ergodic Markov chains*, Journal of Applied Probability, 50 (2013), pp. 184–194.

- [24] D. FRENKEL AND B. SMIT, *Understanding molecular simulation: from algorithms to applications*, 2002.
- [25] D.R. GLOWACKI, E. PACI, AND D.V. SHALASHILIN, *Boxed Molecular Dynamics: Decorrelation Time Scales and the Kinetic Master Equation*, *Journal of Chemical Theory and Computation*, 7(5) (2011), pp. 1244-1252.
- [26] E. GOBET AND S. MENOZZI, *Exact approximation rate of killed hypoelliptic diffusions using the discrete Euler scheme*, *Stochastic Processes and their Applications*, 112 (2004), pp. 201–223.
- [27] ———, *Stopped diffusion processes: Boundary corrections and overshoot*, *Stochastic Processes and their Applications*, 120 (2010), pp. 130–162, 0706.4042.
- [28] G. H. GOLUB AND C. F. VAN LOAN, *Matrix computations*, Johns Hopkins Studies in the Mathematical Sciences, Johns Hopkins University Press, Baltimore, MD, fourth ed., 2013.
- [29] A. T. HAWK, *Milestoning with coarse memory*, *The Journal of chemical physics*, 138 (2013), pp. 154105.
- [30] A. T. HAWK AND D. E. MAKAROV, *Milestoning with transition memory*, *The Journal of Chemical Physics*, 135 (2011), pp. 224109.
- [31] D. J. HIGHAM, X. MAO, M. ROJ, Q. SONG, AND G. YIN, *Mean exit times and the Multilevel Monte Carlo method*, *SIAM/ASA Journal on Uncertainty Quantification*, 1 (2013), pp. 2–18.
- [32] G.A. HUBER AND S. KIM, *Weighted-ensemble Brownian dynamics simulations for protein association reactions*, *Biophysical Journal*, 70(1) (1996), pp. 97–110.
- [33] INTEL CORPORATION, *Intel[®] Digital Random Number Generator (DRNG) Software Implementation Guide*, 2014.
- [34] G.S. JAS ET AL., *Experiments and Comprehensive Simulations of the Formation of a Helical Turn*, *Journal of Physical Chemistry B*, 116(23) (2012), pp. 6598-6610.
- [35] S. KIRMIZIALTIN AND R. ELBER, *Revisiting and computing reaction coordinates with Directional Milestoning*, *The journal of physical chemistry. A*, 115 (2011), pp. 6137–48.
- [36] S. KIRMIZIALTIN ET AL., *How Conformational Dynamics of DNA Polymerase Select Correct Substrates: Experiments and Simulations*, *Structure*, 20(4) (2012), pp. 618-627.

- [37] S. KIRMIZIALTIN, K. A. JOHNSON, AND R. ELBER, *Enzyme Selectivity of HIV Reverse Transcriptase: Conformations, Ligands, and Free Energy Partition*, The Journal of Physical Chemistry B, 119 (2015), pp. 11513–11526.
- [38] S.M. KREUZER, R. ELBER, AND T.J. MOON, *Early Events in Helix Unfolding under External Forces: A Milestoning Analysis*, The Journal of Physical Chemistry B, 116(29) (2012), pp. 8662–8691.
- [39] R. B. LEHOUCQ, D. C. SORENSEN, AND C. YANG, *ARPACK users' guide*, vol. 6 of Software, Environments, and Tools, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1998.
- [40] P. MÁJEK AND R. ELBER, *Milestoning without a reaction coordinate*, Journal of chemical theory and computation, 6 (2010), pp. 1805–1817.
- [41] P. METZNER, C. SCHÜTTE, AND E. VANDEN-EIJNDEN, *Transition Path Theory for Markov jump processes*, 2009.
- [42] G. N. MILSTEIN AND M. V. TRETYAKOV, *Stochastic numerics for Mathematical Physics*, Scientific Computation, Springer Berlin Heidelberg, Berlin, Heidelberg, 2004.
- [43] D. MORONI, P. G. BOLHUIS, AND T. S. VAN ERP, *Rate constants for diffusive processes by partial path sampling*, Journal of Chemical Physics, 120 (2004), pp. 4055–4065, 0310466.
- [44] K. MÜLLER AND L. D. BROWN, *Location of saddle points and minimum energy paths by a constrained simplex optimization procedure*, Theoretica Chimica Acta, 53 (1979), pp. 75–93.
- [45] A. P. RUYMGAART, A. E. CARDENAS, AND R. ELBER, *MOIL-opt: Energy-conserving molecular dynamics on a GPU/CPU system*, Journal of chemical theory and computation, 7 (2011), pp. 3072–3082.
- [46] M. SARICH, F. NOÉ, AND C. SCHÜTTE, *On the approximation quality of Markov State Models*, 2010.
- [47] T. SCHLICK, *Molecular modeling and simulation: an interdisciplinary guide*, 2010.
- [48] D. E. SHAW, J. C. CHAO, M. P. EASTWOOD, J. GAGLIARDO, J. P. GROSSMAN, C. R. HO, D. J. LERARDI, I. KOLOSSVÁRY, J. L. KLEPEIS, T. LAYMAN, C. MCLEAVEY, M. M. DENEROFF, M. A. MORAES, R. MUELLER, E. C. PRIEST, Y. SHAN, J. SPENGLER, M. THEOBALD, B. TOWLES, S. C. WANG, R. O. DROR, J. S. KUSKIN, R. H. LARSON, J. K. SALMON, C. YOUNG, B. BATSON, AND K. J. BOWERS, *Anton, a special-purpose machine for molecular dynamics simulation*, 2008.

- [49] J. E. STONE, J. C. PHILLIPS, P. L. FREDDOLINO, D. J. HARDY, L. G. TRABUCO, AND K. SCHULTEN, *Accelerating molecular modeling applications with graphics processors*, Journal of Computational Chemistry, 28 (2007), pp. 2618–2640.
- [50] D. W. H. SWENSON AND P. G. BOLHUIS, *A replica exchange transition interface sampling method with multiple interface sets for investigating networks of rare events*, The Journal of Chemical Physics, 141 (2014).
- [51] A. ULITSKY AND R. ELBER, *A new technique to calculate steepest descent paths in flexible polyatomic systems*, The Journal of Chemical Physics, 92 (1990), pp. 1510.
- [52] AN ERP, T.S., D. MORONI, AND P.G. BOLHUIS, *A novel path sampling method for the calculation of rate constants*, The Journal of Chemical Physics, 118(17) (2003), pp. 7762–7774.
- [53] E. VANDEN-EIJNDEN AND M. VENTUROLI, *Exact rate calculations by trajectory parallelization and tilting*, The Journal of chemical physics, 131 (2009), pp. 1–7, 0904.3763.
- [54] E. VANDEN-EIJNDEN, M. VENTUROLI, G. CICCOTTI, AND R. ELBER, *On the assumptions underlying milestoning*, The Journal of chemical physics, 129 (2008), pp. 174102.
- [55] A. WARMFLASH, P. BHIMALAPURAM, AND A. R. DINNER, *Umbrella sampling for nonequilibrium processes*, The Journal of chemical physics, 127 (2007), pp. 154112.
- [56] A.M.A WEST, R. ELBER, AND D. SHALLOWAY, *Extending molecular dynamics time scales with milestoning: Example of complex kinetics in a solvated peptide*, Journal of Chemical Physics, 126(14) (2007).
- [57] B.W. ZHANG, D. JASNOW, AND D.M. ZUCKERMAN, *The “weighted ensemble” path sampling method is statistically exact for a broad class of stochastic processes and binning procedures*, Journal of Chemical Physics, 132(5) (2010).
- [58] R. ZWANZIG, *Diffusion in a rough potential*, Proceedings of the National Academy of Sciences of the United States of America, 85 (1988), pp. 2029–2030.