On the algebraic K-theory of the Hilbert modular group

MAURICIO BUSTAMANTE LUIS JORGE SÁNCHEZ SALDAÑA

We give formulas for the Whitehead groups and the rational *K*-theory groups of the (integer group ring of the) Hilbert modular group in terms of its maximal finite subgroups.

1 Introduction

The algebraic K-theory of the integral group ring of a discrete group G is known to encode topological invariants of (high dimensional) manifolds X or spaces whose fundamental group $\pi_1 X$ is isomorphic to G. Examples come in many guises: the obstructions to a finitely dominated space having the homotopy type of finite CW-complex and to an open smooth manifold being the interior of a compact smooth manifold with boundary are elements of $K_0(\mathbb{Z}G)$ (or $Wh_0(G)$ in our notation below). The obstruction to an h-cobordism admitting a product structure $X \times I$ is an element in the Whitehead group Wh(G)(certain quotient of $K_1(\mathbb{Z}G)$ denoted in this article by $Wh_1(G)$) and the uniqueness up to isotopy of these product structures on a compact h-cobordism has to do with a quotient of $K_2(\mathbb{Z}G)$ named $Wh_2(G)$. In fact an entire sequence of groups $Wh_n(G)$ arising from the higher algebraic K-theory of a group ring can be defined to contain the type of invariants needed to deal with parametrized phenomena in topology, e.g. pseudoisotopy theory. These groups $Wh_n(G)$ associated to a group G are called Whithead groups of G and appeared defined (for larger n) by Waldhausen in [Wal78]. Thus, explicit calculations of algebraic K-theory groups and their associated Whitehead groups Wh_n become relevant in geometric topology. Unfortunately such calculation is generally a difficult task to complete, for instance the algebraic Ktheory of \mathbb{Z} is not known yet in full generality. However, sometimes it is possible to approach the lower K-theory and the higher K-theory modulo torsion via certain generalized homology theory. One instance where this can be done is the integral group ring of the Hilbert modular group $PSL_2(\mathcal{O}_k)$. In fact, we prove:

Theorem 1.1 Let k be a totally real number field of finite degree and let \mathcal{O}_k be its ring of algebraic integers. Then for all q,

$$Wh_q(PSL_2(\mathcal{O}_k)) \cong \bigoplus_{(M)} Wh_q(M)$$

where the sum runs over the conjugacy classes of maximal finite subgroups of $PSL_2(\mathcal{O}_k)$.

As for the non-projective Hilbert modular group $SL_2(\mathcal{O}_k)$, we find that its Whitehead group is determined by $K_1(\mathbb{Z}[PSL_2(\mathcal{O}_k)])$. In fact, we have:

Theorem 1.2 Let k be a totally real number field of finite degree and let O_k be its ring of algebraic integers. Then

$$Wh_q(SL_2(\mathcal{O}_k)) \cong K_q(\mathbb{Z}[PSL_2(\mathcal{O}_k)])$$

for $q \leq 1$.

Although the higher K-theory groups of $PSL_2(\mathcal{O}_k)$ seem to be harder to compute, a lot more can be said about its higher rational K-theory. For a finite group F, let

- r(F) denote the number distinct real irreducible representations of F,
- c(F) be the number of those real representations that are of complex type,
- q(F) be the number of distinct rational irreducible representations of F,
- $k_p(F)$ be the number of irreducible representations of F over the p-adic numbers \mathbb{Q}_p and
- $r_p(F)$ be the number of irreducible representations of F over the field with p elements \mathbb{F}_p .

Denote the rank of an abelian group by rk. Then we have the following

Theorem 1.3 Let *k* be a totally real number field of finite degree and let \mathcal{O}_k be its ring of algebraic integers. If $G = PSL_2(\mathcal{O}_k)$, then

$$rk\left(K_{q}(\mathbb{Z}[G])\right) - rk\left(H_{q}(BG;\mathcal{K}(\mathbb{Z}))\right) = \begin{cases} \sum_{(M)}^{(M)} r(M) - m & \text{if } q > 2 \text{ and } q \equiv 1 \mod 4, \\ \sum_{(M)}^{(M)} c(M) & \text{if } q > 2 \text{ and } q \equiv 3 \mod 4, \\ \sum_{(M)}^{(M)} (r(M) - q(M)) & \text{if } q = 1, \\ m - \sum_{(M)} \left[\sum_{p \mid |M|} q(M) - (k_{p}(M) - r_{p}(M))\right] \text{if } q = -1, \\ 0 & \text{otherwise} \end{cases}$$

where *m* is the number of conjugacy classes of maximal finite subgroups in *G*, the sums range over the conjugacy classes of maximal finite subgroups of $G = PSL_2(\mathcal{O}_k)$ and $H_q(BG; \mathcal{K}(\mathbb{Z}))$ denotes the *q*-th homology of *BG* with coefficients in the *K*-theory spectrum $\mathcal{K}(\mathbb{Z})$.

The proof of Theorem 1.3, together with Theorem 1.2 lead us to expressions for the the classical Whitehead group of the non-projective Hilbert modular group and the lower reduced *K*-theory of $SL_2(\mathcal{O}_k)$.

Corollary 1.4

$$Wh_1(SL_2(\mathcal{O}_k)) \simeq Wh_1(PSL_2(\mathcal{O}_k)) \oplus PSL_2(\mathcal{O}_k)^{ab} \oplus \mathbb{Z}_2,$$

where $PSL_2(\mathcal{O}_k)^{ab}$ is the abelianization of $PSL_2(\mathcal{O}_k)$. Also

$$Wh_0(SL_2(\mathcal{O}_k)) \simeq Wh_0(PSL_2(\mathcal{O}_k)) \oplus \mathbb{Z};$$

and

$$Wh_{-1}(SL_2(\mathcal{O}_k)) \simeq \bigoplus_{(M)} K_{-1}(\mathbb{Z}[M]).$$

where the sum are taken over the conjugacy classes of maximal finite subgroups of $PSL_2(\mathcal{O}_k)$.

The main tool we use to prove our main results is the *K*-theoretic *Farrell-Jones Isomorphism conjecture* for a group. When this conjecture is verified for a group Γ , then one can potentially compute the algebraic *K*-theory of the group ring $\mathbb{Z}[\Gamma]$ by first determining the algebraic *K*-theory of its virtually cyclic subgroups and the structure of the restricted orbit category $Or(\Gamma, V_{CYC})$. This will be explained in more detail in Section 2. Then a spectral sequence argument can give rise to the calculation of some of the *K*-theory groups. This method has proved to be effective in several cases, for example 2- and 3-dimensional crystallographic groups [Pea98], [AO06], cocompact Fuchsian groups [BJPP01], [BJPP02], [LS00], Bianchi groups [BFJPP00], braid groups [JPML06], [JPML10], hyperbolic reflexion groups [LO09], virtually free groups [JPLMVP11], [JPSS14], etc.

The spectral sequence we use in our calculation is the p-chain spectral sequence of Davis and Lück which we review in Section 3. This spectral sequence turned out to be convenient due to the structure of the finite subgroups of the Hilbert modular group. We go over this in Section 4. Section 5 is devoted to the proof of Theorems 1.1, 1.2 and 1.3.

Acknowledgements

We are grateful to Tom Farrell for suggesting this problem to us and D. Juan-Pineda for helpful comments on a preliminary version of this paper. The first author acknowledges the second author for teaching him how to navigate through the somewhat dense literature that surrounds this topic. The second author wishes to thank the Binghamton University for its hospitality during the time where part of this paper was developed. The first author was supported by a Binghamton University Dissertation Fellowship. The second author was supported by PAPIIT-UNAM grant 105614, CONACYT Research grant 151338 and CONACYT doctoral scholarship.

2 Preliminaries

The essential tool in our calculation is the Farrell-Jones Isomorphism conjecture. To state this conjecture in a convenient way, we recall some definitions from [DL98].

From now on we only consider discrete groups. A family of subgroups \mathcal{F} of a group G is always assumed to be closed under conjugation and under taking subgroups. We are specially interested in the following families of subgroups:

- A_{LL} of all subgroups of G;
- V_{CYC} of all virtually cyclic subgroups of G, i.e. subgroups which have a possibly finite cyclic subgroup of finite index;
- *FbC* of finite subgroups and all virtually cyclic subgroups of the form $F \rtimes \mathbb{Z}$ with F finite;
- *F*_{*IN*} of all finite subgroups;
- *Tr* consisting of the trivial subgroup.

Note that each family is contained in the previous one.

Definition 2.1 Let G be a group and \mathcal{F} be a family of subgroups. The restricted orbit category $Or(G, \mathcal{F})$ is the category whose objects are homogenous spaces, also called orbits, G/H, $H \in \mathcal{F}$ and whose morphisms are G-maps.

Let Or(G) denote $Or(G, A_{LL})$, and the set of G maps between the orbits G/H and G/K is denoted by $mor_G(G/H, G/K)$.

Note that every element in $mor_G(G/H, G/K)$ is of the form $R_a : G/H \to G/K$ $gH \mapsto ga^{-1}K$, provided $aHa^{-1} \subseteq K$. And that $R_a = R_b$ if and only if $ab^{-1} \in K$.

Definition 2.2 Let G be a group and \mathcal{F} a family of subgroups of G.

- An Or(G, F)-space (resp. Or(G, F)-spectrum) is a functor Or(G, F) → SPACES (resp. Or(G, F) → SPECTRA), where SPACES (resp. SPECTRA) is the cateogry of compactly generated topological spaces (resp. spectra). See [DL98, p. 207].
- We denote by $*_{\mathcal{F}}$ the Or(G)-space defined as

$$*_{\mathcal{F}}(G/H) = \begin{cases} point & if H \in \mathcal{F} \\ \emptyset & otherwise. \end{cases}$$

Given a $G - \mathcal{F}$ -space Z, i.e. a G-space Z all of whose isotropy groups belong to \mathcal{F} , we define the fixed point (contravariant) $Or(G, \mathcal{F})$ -space $map_G(_, Z) : Or(G, \mathcal{F}) \to SPACES$ by $G/H \mapsto map_G(G/H, Z) = Z^H$. In particular, if Z is a $G - \mathcal{F}$ -CW complex we say that $map_G(_, Z)$ is a free $Or(G, \mathcal{F})$ -CW complex.

Remark 2.3 An unreduced homology theory for contravariant $Or(G, \mathcal{F})$ -spaces with coefficientes in a covariant $Or(G, \mathcal{F})$ -spectrum can be defined. This homology theory, denoted by $H^{Or(G,\mathcal{F})}_*(X, Y; \mathbf{E})$, is constructed in [DL98] in order to establish the Farrell-Jones conjecture. It satisfies the Weak Homotopy Equivalence (WHE) axiom, i.e., given a weak homotopy equivalence of pairs of contravariant $Or(G, \mathcal{F})$ -spaces $(f, g) : (X, A) \to (Y, B)$ then the homology groups $H^{Or(G,\mathcal{F})}_p(X, A; \mathbf{E})$ and $H^{Or(G,\mathcal{F})}_p(Y, B; \mathbf{E})$ are isomorphic for all $p \in \mathbb{Z}$.

It can be shown that $H_n^{Or(G)}(*_{\mathcal{F}}; \mathbb{K}) \cong \pi_n(hocolim_{Or(G,\mathcal{F})} \mathbb{K})$, where \mathbb{K} denotes the *K*-theory spectrum defined in [DL98, Section 2]. Also

$$H_n^{Or(G)}(*_{ALL}; \mathbb{K}) \cong \pi_n(hocolim_{Or(G)} \mathbb{K}) \cong K_n(\mathbb{Z}[G])$$

since Or(G) has a final object G/G and $\pi_n(\mathbb{K}(G/G)) \cong K_n(\mathbb{Z}[G])$. It is worth mentioning that in [DL98, Section 2] an $Or(G, \mathcal{F})$ -spectrum \mathbb{K} is defined for every group G, in such a way that $\pi_i(\mathbb{K}(G/H)) = K_n(\mathbb{Z}H)$ for all $n \in \mathbb{Z}$.

In their seminal paper ([FJ93]) Farrell and Jones established their famous isomorphism conjecture for the *K*-theory, *L*-theory and Pseudoisotopy functors. Here we consider the *K*-theoretic version of the conjecture as stated by Davis and Lück in [DL98].

Conjecture 2.4 (The Farrell-Jones isomorphism conjecture) For any group G the following assembly map, induced by inclusion of Or(G)-spaces, is an isomorphism

(*)
$$A_{VCYC,ALL}: H_n^{Or(G)}(*_{VCYC}; \mathbb{K}) \to H_n^{Or(G)}(*_{ALL}; \mathbb{K}) \cong K_n(\mathbb{Z}G).$$

Once the Farrell-Jones conjecture has been verified for a group G, one can hope to compute $K_n(\mathbb{Z}[G])$ by computing the left hand side of (*). The later is a generalized homology theory that can be approached, for example, via Mayer-Vietoris sequences, Atiyah-Hirzebruch-type spectral sequences or, as in our case, the *p*-chain spectral sequence.

The Whitehead groups $Wh_n(G)$ of G appear in this context as follows

Proposition 2.5 [Wal78, Prop. 15.7] Let G be a group. Then $Wh_n(G) \cong H_n^{Or(G)}(*_{ALL}, *_{Tr}; \mathbb{K})$ for all $n \in \mathbb{Z}$. In fact they fit in a long exact sequence

$$\cdots \to H_n(BG; \mathbb{K}(G/1)) \to K_n(\mathbb{Z}G) \to Wh_n(G) \to H_{n-1}(BG; \mathbb{K}(G/1)) \to \cdots$$

where $H_n(BG; \mathbb{K}(G/1))$ is the classical generalized homology theory with coefficients in the spectrum $\mathbb{K}(G/1)$ which has as homotopy groups the algebraic *K*-theory of the group ring $\mathbb{Z}G$.

When the Farrell-Jones conjecture holds for a group G we obtain:

Corollary 2.6 Let *G* be a group. Suppose that the Farrell-Jones conjecture holds for *G*, then $Wh_n(G)$ is isomorphic to $H_n^{Or(G)}(*_{VCYC}, *_{Tr}; \mathbb{K})$ for all $n \in \mathbb{Z}$.

One usually wonders whether a smaller family of subgroups than V_{crc} would suffice to succeed in the calculation of the Whitehead groups of a group or the *K*-theory of a group ring. The next two theorems tell us something about it.

Theorem 2.7 [FJ93, A.10] Let G be a group. Suppose that for any finite by cyclic subgroup V of G, the assembly map

$$A_{F_{IN},ALL}: H_n^{Or(V)}(*_{F_{IN}}; \mathbb{K}) \to H_n^{Or(V)}(*_{ALL}; \mathbb{K}) \cong K_n(\mathbb{Z}V)$$

is an isomorphism. Then $H_n^{Or(G)}(*_{FIN}; \mathbb{K}) \cong H_n^{Or(G)}(*_{FbC}; \mathbb{K})$.

Remark 2.8 The obstructions to $A_{FIN,ALL}$ being an isomorphism is the so-called Nil groups. The vanishing of these groups allows one to consider only the family of finite subgroups. Thus under the hypothesis of Theorem 2.7 and using results of [DKR11], [DQR11], one can actually compute the Whitehead groups of a group G by considering only its family of finite groups. In other words, if G satisfies the Farrell-Jones conjecture then

$$Wh_n(G) \cong H_n^{Or(G)}(*_{ALL}, *_{Tr}; \mathbb{K})$$
$$\cong H_n^{Or(G)}(*_{VCYC}, *_{Tr}; \mathbb{K})$$
$$\cong H_n^{Or(G)}(*_{FbC}, *_{Tr}; \mathbb{K})$$
$$\cong H_n^{Or(G)}(*_{FIN}, *_{Tr}; \mathbb{K}).$$

Here is one more way in which one can replace the family of subgroups of a group by a smaller family.

Theorem 2.9 [DL03, Corollary 3.9] Let \mathcal{F} be a family of subgroups of G. Denote by $M\mathcal{F} \subset \mathcal{F}$ the subfamily consisting of: all maximal elements in \mathcal{F} , the groups in \mathcal{F} which are contained in more than one maximal element and the groups in \mathcal{F} which are contained in no maximal element of \mathcal{F} . Then for all n

$$H_n^{Or(G)}(*_{\mathcal{F}};\mathbb{K})\cong H_n^{Or(G)}(*_{M\mathcal{F}};\mathbb{K}).$$

3 The *p*-chain spectral sequence

In this section we establish a special case of the *p*-chain spectral sequence of Davis and Lück [DL03]. Besides the Farrell-Jones conjecture, this will be the main ingredient in our calculation. The *p*-chain spectral sequence converges to $H^{Or(G,\mathcal{F})}(X; \mathbf{E})$. By reasons that will become evident in the next section, it will be enough to restrict ourselves to the restricted orbit category $Or(G, F_{IN})$ and X the one point $Or(G, F_{IN})$ -space. This category has the property that every endomorphism is an isomorphism and for each object G/K, Aut(G/K) acts freely in $mor_G(G/H, G/K)$. Let $\overline{G/H}$ denote the isomorphism class of G/H. Note that there is a partial order on the set of isomorphism classes defined by $\overline{G/H} \leq \overline{G/K}$ if $mor_G(G/H, G/K)$ is nonempty. We also write $\overline{G/H} < \overline{G/K}$ if $\overline{G/H} \leq \overline{G/K}$ and $\overline{G/K} \notin \overline{G/H}$. **Definition 3.1** Let *G* be a group and \mathcal{F} a family of subgroups. A sequence of isomorphism classes of objects $c := \{\overline{G/H_0}, \overline{G/H_1}, \cdots, \overline{G/H_p}\}$ in $Or(G, \mathcal{F})$ is called a *p*-chain if

$$\overline{G/H_0} < \overline{G/H_1} < \cdots < \overline{G/H_p}.$$

Associated to a *p*-chain $c = \{\overline{G/H_0}, \overline{G/H_1}, \cdots, \overline{G/H_p}\}$ there are $Aut(G/H_p)$ - $Aut(G/H_0)$ -spaces S(c) defined by

$$S(c) = \begin{cases} Aut(G/H_0) & \text{if } p = 0\\ mor_G(G/H_0, G/H_1) & \text{if } p = 1\\ mor_G(G/H_{p-1}, G/H_p) \times_{Aut(G/H_{p-1})} \cdots \times_{Aut(G/H_1)} mor_G(G/H_0, G/H_1) & \text{if } p \ge 2 \end{cases}$$

The next result is proven (in more generality) in [DL98, Lemma 2.11 and Remark 2.13]

Theorem 3.2 (The *p*-chain spectral sequence) Let *G* be a group and let $\mathcal{J} \subset F_{\mathbb{N}}$ be any subfamily of the family of finite subgroups of *G*. Then there is a spectral sequence whose first page is given by

$$E_{p,q}^{1} = \bigoplus_{\substack{p-\text{ chains } c\\\text{ in } Or(G,\mathcal{J})}} H_{q}^{Aut(G/H_{0})}(pt \times_{Aut(G/H_{p})} S(c); \mathbb{K})$$

which converges to $H_{p+q}^{Or(G)}(*_{\mathcal{J}}; \mathbb{K})$.

Remark 3.3 This spectral sequence is very manageable when one can control the length of the *p*-chains. In our case, when *G* becomes the Hilbert modular group, we will see that we can work in a family of subgroups in which no *p*-chains for $p \ge 2$ have to be considered in the restricted orbit category. This will make the spectral sequence collapse quickly and explicit calculations can be done.

4 The Hilbert Modular Group

In this section we review the definition and some basic properties of the Hilbert modular group. The results we state without proof in this section can be found for example in [Fre90]. For additional information about the Hilbert modular group we refer the reader to [Hir73], [vdG88].

A totally real number field k is an algebraic extension of \mathbb{Q} such that all its embeddings $\sigma_i : k \to \mathbb{C}$ have image contained in \mathbb{R} . Let k denote a totally real number field of degree n and \mathcal{O}_k its ring of integers. The *Hilbert modular group* $PSL_2(\mathcal{O}_k)$ is defined to be the quotient of the special linear group of 2 by 2 matrices $SL_2(\mathcal{O}_k)$ with entries in \mathcal{O}_k by the subgroup consisting of $\{I, -I\}$, where I denotes the identity matrix; in other words

$$PSL_2(\mathcal{O}_k) = SL_2(\mathcal{O}_k) / \{I, -I\}.$$

Note that if $k = \mathbb{Q}$, then $PSL_2(\mathcal{O}_k) = PSL_2(\mathbb{Z})$ is nothing but the classical modular group. However $PSL_2(\mathcal{O}_k)$ is not a discrete subgroup of $PSL_2(\mathbb{R})$ if $n \ge 2$. Yet it does act properly and discontinuously on the *n*-fold product $\mathbb{H}^n = \mathbb{H} \times \cdots \times \mathbb{H}$ of upper half planes, by fractional linear transformations in each of the *n* factors, via the *n* different embeddings of *k* into \mathbb{R} . Thus the Hilbert modular group is a discrete subgroup of $PSL_2(\mathbb{R})^n = PSL_2(\mathbb{R}) \times \cdots \times PSL_2(\mathbb{R})$. For example, if we let *d* be a square-free positive integer and $k = \mathbb{Q}(\sqrt{d})$, then there are two embeddings of *k* into \mathbb{R} , namely

$$\sigma_1: s + t\sqrt{d} \mapsto s + t\sqrt{d}$$
 and $\sigma_2: s + t\sqrt{d} \mapsto s - t\sqrt{d}$,

K-theory of Hilbert modular group

for $s, t \in \mathbb{Q}$. In this case, a matrix $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in PSL_2(O_k)$ acts on $(z_1, z_2) \in \mathbb{H} \times \mathbb{H}$ by $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} (z_1, z_2) = \begin{pmatrix} \frac{\sigma_1(\alpha)z_1 + \sigma_1(\beta)}{\sigma_1(\gamma)z_1 + \sigma_1(\delta)}, \frac{\sigma_2(\alpha)z_2 + \sigma_2(\beta)}{\sigma_2(\gamma)z_2 + \sigma_2(\delta)} \end{pmatrix}.$

The action of $PSL_2(\mathcal{O}_k)$ on \mathbb{H}^n is not free. One can detect points with non-trivial isotropy by the following lemma (Compare with the analogous situation of $PSL_2(\mathbb{Z})$ acting on \mathbb{H} by Möbius transformations):

Lemma 4.1 Let $h \in PSL_2(\mathcal{O}_k)$ and denote by $\sigma_i : PSL_2(\mathcal{O}_k) \hookrightarrow PSL_2(\mathbb{R})$, i = 1, ..., n the canonical embeddings of $PSL_2(\mathcal{O}_k)$ into $PSL_2(\mathbb{R})$. Then the following conditions are equivalent:

(1) $\sigma_1(h), \ldots, \sigma_n(h) \in PSL_2(\mathbb{R})$ are elliptic matrices, i.e. their traces satisfy

$$[Tr \sigma_i(h)]^2 - 4 < 0.$$

- (2) h has finite order.
- (3) h has a unique fixed point.

Moreover the stabilizer Γ_z of any point $z \in \mathbb{H}^n$ in $PSL_2(\mathcal{O}_k)$ is a finite cyclic group.

For example the matrix $\begin{pmatrix} \sqrt{2} & -1 \\ 1 & 0 \end{pmatrix}$ is an elliptic element of $PSL_2\left(\mathcal{O}_{\mathbb{Q}(\sqrt{2})}\right)$ of order 4 that fixes the point $\frac{\sqrt{2}}{2}((1+i),(-1+i)) \in \mathbb{H} \times \mathbb{H}$.

Remark 4.2 \mathbb{H}^n turns out to be a classifying space for proper actions of the Hilbert modular group $PSL_2(\mathcal{O}_k)$, where the fixed point sets by finite subgroups are not only contractible but consist of a single point. This property will simplify the calculation of the *K*-theory groups enormously.

Perhaps more importantly for our purposes, $PSL_2(\mathcal{O}_k)$ satisfies the conditions M, NM and FJ specified in [DL03, Section 4]:

Lemma 4.3 The Hilbert modular group $PSL_2(\mathcal{O}_k)$ satisfies the following three properties:

- (M) Every finite subgroup of $PSL_2(\mathcal{O}_k)$ is contained in a unique maximal finite subgroup.
- (NM) If M is a maximal finite subgroup of $PSL_2(\mathcal{O}_k)$ then N(M) = M, where N(M) denotes the normalizer of M in $PSL_2(\mathcal{O}_k)$.
- (FJ) The Isomorphism Conjecture of Farrell and Jones for algebraic K-theory is true for $PSL_2(\mathcal{O}_k)$.

Proof Note that by the lemma above if *H* is a finite subgroup of $PSL_2(\mathcal{O}_k)$ then *H* is contained in the finite stabilizer Γ_z of a point $z \in \mathbb{H}^n$. Γ_z is a finite maximal subgroup of $PSL_2(\mathcal{O}_k)$. For if *F* is a finite subgroup of $PSL_2(\mathcal{O}_k)$ containing Γ_z , then there must exist $y \in \mathbb{H}^n$ such that $f \cdot y = y$ for all $f \in F$. In particular, every element of Γ_z would fix *y* and by uniqueness of the fixed points, y = z, that is to say $f \in \Gamma_z$. This proves that $PSL_2(\mathcal{O}_k)$ satisfies property (M).

Let *M* be a maximal finite subgroup of $PSL_2(\mathcal{O}_k)$. Then by the previous paragraph, $M = \Gamma_z$, where Γ_z is the stabilizer of some point $z \in \mathbb{H}^n$. Let now $g \in N(M)$, then $gfg^{-1} \in M$ for some all $f \in M = \Gamma_z$. Hence $gfg^{-1}z = z$ which implies that $fg^{-1}z = g^{-1}z$, and by the uniqueness of the fixed points $g^{-1}z = z$, i.e. $g \in M$. This proves that $PSL_2(\mathcal{O}_k)$ satisfies the property (NM).

As for property (FJ) we have to note that $PSL_2(\mathcal{O}_k)$ is a lattice in the Lie group $PSL_2(\mathbb{R})^n$. The Farrell-Jones conjecture has been proven for this type of groups in [KLR].

Remark 4.4 Conditions (M) and (NM) can be interpreted in a somewhat geometric way as follows: Let \mathcal{J} be the family of groups G satisfying (FJ) and for which there is a model for $\underline{E}G$ with the property that every fixed point set by a finite subgroup of G consists of a single point. Then it is clear that if $G \in \mathcal{J}$, then G satisfies (M) and (NM) and (FJ). It is worth noticing that such family \mathcal{J} is closed under free products.

If a group *G* has properties (M) and (NM) its only possible virtually cyclic subgroups are very limited (Compare [DL03, p.100]).

Lemma 4.5 Let *G* be a group having properties (M) and (NM), then every infinite virtually cyclic subgroup of *G* is isomorphic to either \mathbb{Z} or $\mathbb{Z}_2 * \mathbb{Z}_2$.

Proof Every infinite virtually cyclic subgroup V of G fits into an extension

$$1 \to F \to V \to \Gamma \to 1$$

where Γ is either \mathbb{Z} or $\mathbb{Z}_2 * \mathbb{Z}_2$ and *F* is finite. Since *F* is normal in *V* then $V \subset N(F)$ where N(F) is the normalizer of *F* in *G*. Suppose that *F* is non-trivial and let *M* be the unique maximal finite subgroup of *G* containing *F*. It is clear that $F \subset gMg^{-1}$ if $g \in N(F)$. Hence, by uniqueness, $g \in M$. This shows that $N(F) \subset M$ which is finite, contradicting that $V \subset N(F)$. Therefore *F* is trivial and the conclusion follows.

5 Whitehead groups and rational *K*-theory of the Hilbert modular group

In this section we prove Theorems 1.1, 1.2 and 1.3.

Proof of Theorem 1.1 Let $G = PSL_2(\mathcal{O}_k)$. Recall that $Wh_q(G) \simeq H_q^{Or(G)}(*_{ALL}, *_{Tr}; \mathbb{K})$. The first step in our calculation is to verify the Farrell-Jones conjecture for *G*. As mentioned in Lemma 4.3, this follows from [KLR]. The next step is trying to reduce the family of subgroups as much as possible. In fact, by Lemma 4.5 and the fact that the Nil groups of the integer group rings of \mathbb{Z} and $\mathbb{Z}_2 * \mathbb{Z}_2$ vanish [LS00, Lemma 2.5], we have that the assembly map

$$A_{FIN,ALL}: H_q^{Or(V)}(*_{FIN}; \mathbb{K}) \to H_q^{Or(V)}(*_{ALL}; \mathbb{K})$$

is an isomorphism. Hence $Wh_q(G)$ can be computed using only the family of finite subgroups of G. That is to say

$$Wh_q(G) \simeq H_q^{Or(G)}(*_{FIN}, *_{Tr}; \mathbb{K})$$

Furthermore, by combining Lemma 4.3 (property M) and Theorem 2.9, we see that the family of subgroups considered can be reduced further to the subfamily MF_{IN} , where MF_{IN} denotes the family of maximal finite subgroups of G union the trivial subgroup, as defined in Theorem 2.9. Hence

$$Wh_q(G) \cong H_q^{Or(G)}(*_{MFIN}, *_{Tr}; \mathbb{K}).$$

Now that we have replaced V_{CYC} by MF_{IN} , we analyze the structure of the *p*-chains in $Or(G, MF_{IN})$.

0-chains. These are chains of the form $\{G/H\}$, where $H \in MF_{IN}$.

1-chains. Here we only have chains of the form $\{\overline{G/1}, \overline{G/H}\}$. Note that there are no 1-chains of the form $\{\overline{G/H}, \overline{G/K}\}$, for $H, K \in MF_{IN}, H \neq 1$, because every morphism $G/H \to G/K$ has to be an isomorphism.

p-chains, $p \ge 2$. There are none because every morphism $G/H \to G/K$, $H \ne 1$, is an isomorphism.

Also notice that by considering the pair $(*_{MFIN}, *_{Tr})$ we are neglecting terms coming from the 0-chain $\{G/1\}$ in the E^1 page of the *p*-chain spectral sequence for pairs. Particularly, no 1-chains have to be considered in the calculation of the Whitehead groups. Thus the spectral sequence will only have contributions from isomorphism classes of orbits of the form G/H with $H \in MF_{IN}$, $H \neq 1$. Now recall that $Aut(G/H) = N(H)/H = \{e\}$, where N(H) is the normalizer of H in G and the last equality follows from Lemma 4.3 (property NM). Also there is a correspondence between 0-chains in $Or(G, MF_{IN})$ and conjugacy classes of maximal finite subgroups in G. Thus we end up with the following E^1 -term. Therefore the spectral sequence collapses at E^1 and the result follows.

2	$\bigoplus_{(M)} Wh_2(M)$	0	0
1	$\bigoplus_{(M)} Wh_1(M)$	0	0
0	$\bigoplus_{(M)} Wh_0(M)$	0	0
-1	$\bigoplus_{(M)} Wh_{-1}(M)$	0	0
	0	1	2

Proof of Theorem 1.2 We follow the same strategy as in the proof of Theorem 1.1. Let $G = SL_2(\mathcal{O}_k)$. It satisfies the Farrell-Jones conjecture for the same reasons that $PSL_2(\mathcal{O}_k)$ does. Thus we can guarantee that

$$H_q^{Or(G)}(*_{V_{Cyc}}; \mathbb{K}) \cong K_q(\mathbb{Z}[G])$$

for all $q \in \mathbb{Z}$. The idea now is trying to reduce the family of subgroups in order to compute the left side of this equation. Recall that by [DKR11], [DQR11] it is always possible to reduce the family from V_{CPC} to FbC (see also Section 2). Hence if we want to replace the family V_{CPC} with F_{IN} it is enough to prove that for any subgroup of G of the form $F \rtimes \mathbb{Z}$ with F finite, the assembly map $f : H_q^{OP(F \rtimes \mathbb{Z})}(*_{FIN}; \mathbb{K}) \to K_q(\mathbb{Z}[F \rtimes \mathbb{Z}])$ is an isomorphism. To see this, note that $F \times \mathbb{Z} < F \rtimes \mathbb{Z}$ and the only non-trivial finite-order element that commutes with an element of infinite order is $-I = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$, then every group in the family FbC is isomorphic to either \mathbb{Z} or $\mathbb{Z}_2 \times \mathbb{Z}$. By [LS00, Lemma 2.5] f is an isomorphism for $q \leq 1$ when F is the trivial group or $F \cong \mathbb{Z}_2$. Therefore

$$H_q^{Or(G)}(*_{VCYC}; \mathbb{K}) \cong H_q^{Or(G)}(*_{FbC}; \mathbb{K}) \cong H_q^{Or(G)}(*_{FIN}; \mathbb{K}),$$

whenever $q \leq 1$.

We can actually work with a smaller family of subgroups. First note that by the same reasoning as in the proof of Lemma 4.3, every finite maximal subgroup of *G* is of the form G_z , $z \in \underline{E}G = \mathbb{H} \times \cdots \times \mathbb{H}$, the *n*-fold product of upper half-spaces. Hence every finite subgroup different from $\{\pm I\}$ is contained in a unique finite maximal subgroup. Let MF_{IN} as defined in Theorem 2.9, note that every element in MF_{IN} is a maximal finite subgroup, or the trivial subgroup or the group $\{I, -I\}$. By Theorem 2.9 we have

$$H_q^{Or(G)}(*_{FIN};\mathbb{K})\cong H_q^{Or(G)}(*_{MFIN};\mathbb{K}).$$

Now, using the Five Lemma and the previous reductions of the family of subgroups, it is straightforward to show that, for $q \leq 1$

$$Wh_q(G) \cong H_q^{Or(G)}(*_{All}, *_{Tr}; \mathbb{K}) \cong H_q^{Or(G)}(*_{MFIN}, *_{Tr}; \mathbb{K}).$$

We now analyze the *p*-chains that appear in $Or(G, MF_{IN})$

- 0-chains. These are chains of the form $\{\overline{G/1}\}, \{\overline{G/\{\pm I\}}\}\$ and $\{\overline{G/H}\}$, where *H* is a finite maximal subgroup.
- 1-chains. We have chains of the form $\{\overline{G/1}, \overline{G/\{\pm I\}}\}, \{\overline{G/1}, \overline{G/H}\}$ and $\{\overline{G/\{\pm I\}}, \overline{G/H}\}$, with *H* a finite maximal subgroup.
- 2-chains. The only 2-chains that we can form are of the type $\{\overline{G/1}, \overline{G/\{\pm I\}}, \overline{G/H}\}$, with *H* a finite maximal subgroup.
- *p*-chains, $p \ge 3$. There are none because for *H*, *K* maximal every morphism $G/H \to G/K$ has to be an isomorphism

Also notice that by considering the pair $(*_{MFIN}, *_{Tr})$ we are neglecting terms coming from *p*-chains whose least element is the 0-chain $\{\overline{G/1}\}$. Particularly, no 2-chains have to be considered. Thus the E^1 term of *p*-chain spectral sequence for pairs will only have contributions from orbits in $Or(G, MF_{IN}^0)$, where $MF_{IN}^0 := MF_{IN} - Tr$.

It is clear that $Or(G, MF_{IN}^0)$ is equivalent to $Or(PSL_2(\mathcal{O}_k), MF_{IN})$. Thus the E^1 -term of the *p*-chain spectral sequence (for pairs) in $Or(G, MF_{IN})$ is isomorphic to the E^1 -term of the *p*-chain spectral sequence in $Or(PSL_2(\mathcal{O}_k), MF_{IN})$. Since the former converges to $Wh_*(G)$ and the latter converges to $K_*(\mathbb{Z}[PSL_2(\mathcal{O}_k)])$ the result follows.

Proof of theorem 1.3 Let $G = PSL_2(\mathcal{O}_k)$. Recall that MF_{IN} from Theorem 2.9 is the family of subgroups of G consisting of all maximal finite subgroups together with the trivial subgroup. In [Gru08, Theorem 5.6], it is proven that for any group Γ that satisfies the Farrell-Jones conjecture

$$H^{Or(\Gamma)}_*(*_{FIN};\mathbb{K})\otimes\mathbb{Q}\cong H^{Or(\Gamma)}_*(*_{VCYC};\mathbb{K})\otimes\mathbb{Q}\cong K_*(\mathbb{Z}[\Gamma])\otimes\mathbb{Q}$$

Now we use Theorem 2.9 to reduce the family from F_{IN} to MF_{IN} . We start analyzing the *p*-chains that appear in $Or(G, MF_{IN})$. This has been done already in the proof of Theorem 1.1. Note, though, that this time we are not computing the homology of a pair, so we do need to consider 0-chains and 1-chains (there are no *p*-chains in this category for $p \ge 2$.) The first page of the *p*-chain spectral sequence is given (after some simplifications, see [BJPP01, Proposition 12]) by

$$E_{0q}^{1} = H_{q}(BG; \mathcal{K}(\mathbb{Z})) \oplus \bigoplus_{(M)} K_{q}(\mathbb{Z}[M]))$$
$$E_{1q}^{1} = \bigoplus_{(M)} H_{q}(BM; \mathcal{K}(\mathbb{Z}))$$
$$E_{pq}^{1} = 0 \text{ for } p \neq 0, 1,$$

where $H_q(_; \mathcal{K}(\mathbb{Z}))$ refers to the generalized homology theory with coefficients in the Pedersen-Weibel *K*-theory spectrum and the sum runs over conjugacy classes of maximal finite subgroups. Thus the *p*-chain spectral sequence in this case looks like this:

The differentials $d^1 : E_{1q}^1 \to E_{0q}^1$ are all induced by two types of maps between *p*-chains: Those arising from $\{\overline{G/1}, \overline{G/M}\} \mapsto \{\overline{G/1}\}$ and those that come from $\{\overline{G/1}, \overline{G/M}\} \mapsto \{\overline{G/M}\}$. It is shown in [Pea98, Lemma 3.10] that in the first case the induced map in homology is inclusion whereas the second map induces the classical assembly map $H_q(BM; \mathcal{K}(\mathbb{Z})) \to K_q(\mathbb{Z}[M])$ which in our notation is

$$A_{Tr,All}: H_q^{Or(M)}(*_{Tr}; \mathbb{K}) \to H_q^{Or(M)}(*_{All}; \mathbb{K}).$$

In our case this assembly map is known to be rationally injective for all q (see [BHM93]). Hence, at least rationally the differentials d^1 are injective and the spectral sequence (rationally) collapses at E^2 . Then

$$\operatorname{rk}\left(K_q(\mathbb{Z}[G])\right) = \operatorname{rk}\left(E_{0q}^{\infty}\right) = \operatorname{rk}\left(E_{0q}^{2}\right)$$

Note that E_{0q}^2 fits in an exact sequence:

$$0 \to E_{1q}^1 \otimes \mathbb{Q} \xrightarrow{d^1 \otimes id_{\mathbb{Q}}} E_{0q}^1 \otimes \mathbb{Q} \to E_{0q}^2 \otimes \mathbb{Q} \to 0.$$

Finally, by taking into account that every finite subgroup of the Hilbert modular group G is cyclic, an Atiyah-Hirzebruch spectral sequence calculation shows that

$$\operatorname{rk} (H_q(BM; \mathcal{K}(\mathbb{Z})) = \begin{cases} 1 & \text{if } q = 0 \text{ or } q \equiv 1 \mod 4, q > 2\\ 0 & \text{otherwise} \end{cases}$$

Hence,

$$\operatorname{rk} (K_q(\mathbb{Z}[G])) = \operatorname{rk} (E_{0q}^2)$$

= $\operatorname{rk} (E_{0q}^1) - \operatorname{rk} (E_{1q}^1)$
= $\sum_{(M)} \operatorname{rk} (K_q(\mathbb{Z}[M])) + \operatorname{rk} (H_q(BG; \mathcal{K}(\mathbb{Z})) - \sum_{(M)} \operatorname{rk} (H_q(BM; \mathcal{K}(\mathbb{Z}))).$

Thus we have

(**) rk
$$(K_q(\mathbb{Z}[G]))$$
 - rk $(H_q(BG; \mathcal{K}(\mathbb{Z})) = \begin{cases} \sum_{(M)} \operatorname{rk} (K_q(\mathbb{Z}[M])) - m & \text{if } q = 0 \text{ or } q \equiv 1 \mod 4, q > 2 \\ \sum_{(M)} \operatorname{rk} (K_q(\mathbb{Z}[M])) & \text{otherwise,} \end{cases}$

where *m* denotes the number of conjugacy classes of maximal finite subgroups of *G*. By results of Bass [Bas65], Carter [Car80] and Jahren [Jah09] on the *K*-theory of finite groups, we have the following equalities:

$$\operatorname{rk} (K_q(\mathbb{Z}[M])) = \begin{cases} r(M) & \text{if } q > 2 \text{ and } q \equiv 1 \mod 4 \\ c(M) & \text{if } q > 2 \text{ and } q \equiv 3 \mod 4 \\ r(M) - q(M) & \text{if } q = 1 \\ 1 & \text{if } q = 0 \\ 1 - q(M) + \sum_{p||M|} (k_p(M) - r_p(M)) & \text{if } q = -1 \\ 0 & \text{otherwise.} \end{cases}$$

The result then follows by substituting in the equation (**).

Proof of Corollary 1.4 Recall that by Theorem 1.2 $Wh_1(SL_2(\mathcal{O}_k)) \simeq K_1(PSL_2(\mathcal{O}_k))$. Now notice that the differential of the spectral sequence that appears in the proof of Theorem 1.3, is injective for $q \le 1$, see [BJPP01, Proof of Proposition 14]. This implies that $K_1(PSL_2(\mathcal{O}_k)) = E_{01}^{\infty} = E_{01}^2$ and then we obtain an exact sequence

$$0 \to E_{11}^1 \xrightarrow{d^1} E_{01}^1 \to K_1(PSL_2(\mathcal{O}_k)) \to 0.$$

Following [BJPP01, Proof of Proposition 14] is straightforward to check that

$$E_{01}^{1} = PSL_{2}(\mathcal{O}_{k})^{ab} \oplus \mathbb{Z}_{2} \oplus \bigoplus_{(M)} K_{1}(\mathbb{Z}[M])$$

and also

$$E_{11}^1 = \bigoplus_{(M)} (M \oplus \mathbb{Z}_2)$$

Since *M* is abelian, $K_1(\mathbb{Z}[M]) \simeq M \oplus \mathbb{Z}_2 \oplus Wh_1(M)$ and the image of the assembly map $H_1(BM; \mathcal{K}(\mathbb{Z})) \simeq M \oplus \mathbb{Z}_2 \to K_1(\mathbb{Z}[M])$ splits for each *M*. This gives us the desired formula for the classical Whitehead group:

$$Wh_1(SL_2(\mathcal{O}_k)) \simeq Wh_1(PSL_2(\mathcal{O}_k)) \oplus PSL_2(\mathcal{O}_k)^{ab} \oplus \mathbb{Z}_2.$$

Similarly we obtain an exact sequence

$$0 \to E_{10}^1 \xrightarrow{d^1} E_{00}^1 \to K_0(PSL_2(\mathcal{O}_k)) \to 0,$$

where

$$E_{10}^{1} = \bigoplus_{(M)} H_{0}(BM; \mathcal{K}(\mathbb{Z})) \simeq \bigoplus_{(M)} \mathbb{Z}$$

and

$$E_{00}^1\simeq \mathbb{Z}\oplus igoplus_{(M)} K_0(\mathbb{Z}[M]).$$

Note that $K_0(\mathbb{Z}[M]) \simeq \mathbb{Z} \oplus Wh_0(M)$ and $Wh_0(M)$ is finite because M is finite. The result then follows by noticing that the assembly map $H_0(BM; \mathcal{K}(\mathbb{Z})) \simeq \mathbb{Z} \to K_0(\mathbb{Z}[M]) \simeq \mathbb{Z} \oplus Wh_0(M)$ is split injective for each M.

The same analysis at the -1-st level yields $Wh_{-1}(SL_2(\mathcal{O}_k)) \simeq \bigoplus K_{-1}(\mathbb{Z}[M])$. This completes the proof of the corollary.

We conclude this section working out a concrete example. Let $k = \mathbb{Q}(\sqrt{5})$ the quadratic extension of the rational numbers obtained by adjoining $\sqrt{5}$ to \mathbb{Q} . In this case, $PSL_2(\mathcal{O}_k)$ acts on $\mathbb{H} \times \mathbb{H}$ via the embeddings induced by $\sigma_1 : \sqrt{5} \mapsto \sqrt{5}$ and $\sigma_2 : \sqrt{5} \mapsto -\sqrt{5}$. By Theorem 1.1, the Whitehead groups of $PSL_2(\mathcal{O}_k)$ are determined by its maximal finite subgroups. Hence our problem reduces to finding all conjugacy classes of maximal finite subgroups. By Lemma 4.3, every maximal finite subgroup of $PSL_2(\mathcal{O}_k)$ appears as a stabilizer of some point in $\mathbb{H} \times \mathbb{H}$. We will say that two fixed points in $\mathbb{H} \times \mathbb{H}$ are inequivalent if their isotropy groups are not conjugate. Thus we have a bijection between conjugacy classes of maximal finite subgroups of $PSL_2(\mathcal{O}_k)$ and the number of inequivalent fixed points in $\mathbb{H} \times \mathbb{H}$. The problem of finding the number of fixed points of the action of the Hilbert modular group of a quadratic number field has been addressed in [Pre68]. We sketch here the way to proceed: since only elliptic elements of $PSL_2(\mathcal{O}_k)$ can have fix points, then the trace of such an element must satisfy

$$[Tr \sigma_i(h)]^2 - 4 < 0, \quad i = 1, 2.$$

Also, every elliptic matrix is conjugate (in $PSL_2(\mathbb{C})$) to one of the form $\begin{pmatrix} \omega & 0 \\ 0 & \overline{\omega} \end{pmatrix}$ where ω is a primitive

root of unity. Thus, in our case, each of these traces should be an algebraic integer of $\mathbb{Q}(\sqrt{5})$, i.e. an element of $\mathbb{Z}[\frac{1+\sqrt{5}}{2}]$. The only possibilities are: $0, \pm 1$ and $\pm \frac{1\pm\sqrt{5}}{2}$, corresponding to a 4th, 6th and 10th root of unity respectively. This says that in $PSL_2(\mathcal{O}_k)$ we will only have elliptic elements of order 2, 3 and 5, consequently the only finite subgroups that can appear will be isomorphic to \mathbb{Z}_2 , \mathbb{Z}_3 and \mathbb{Z}_5 . The number of conjugacy classes of these maximal finite subgroups is calculated in [Pre68]: each of these groups has exactly two conjugacy classes in $PSL_2(\mathcal{O}_k)$. Therefore, by Theorem 1.1 we have

$$Wh_q(PSL_2(\mathcal{O}_{\mathbb{Q}(\sqrt{5})})) = Wh_q(\mathbb{Z}_2)^2 \oplus Wh_q(\mathbb{Z}_3)^2 \oplus Wh_q(\mathbb{Z}_5)^2$$

In the special case of the classical Whithead group, i.e. q = 1, it is known (See [Oli88]) that $Wh(\mathbb{Z}_2) \simeq Wh(\mathbb{Z}_3) \simeq 0$ and $Wh(\mathbb{Z}_5) \simeq \mathbb{Z}$. Therefore

$$Wh(PSL_2(\mathcal{O}_{\mathbb{Q}(\sqrt{5})})) \simeq \mathbb{Z} \oplus \mathbb{Z}.$$

Also, by Corollary 1.4 we have

$$Wh(SL_2(\mathcal{O}_{\mathbb{Q}(\sqrt{5})})) \simeq \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}_2$$

since $PSL_2(\mathcal{O}_{\mathbb{Q}(\sqrt{5})})$ is a perfect group.

Finally for the higher *K*-theory we obtain (here $G = PSL_2(\mathcal{O}_{\mathbb{Q}(\sqrt{5})}))$:

$$\operatorname{rk} K_{q}(\mathbb{Z}[G]) - \operatorname{rk} H_{q}(BG; \mathcal{K}(\mathbb{Z})) = \begin{cases} 2r(\mathbb{Z}_{2}) + 2r(\mathbb{Z}_{3}) + 2r(\mathbb{Z}_{5}) & \text{if } q > 2 \text{ and } q \equiv 1 \mod 4 \\ 2c(\mathbb{Z}_{2}) + 2c(\mathbb{Z}_{3}) + 2c(\mathbb{Z}_{5}) & \text{if } q > 2 \text{ and } q \equiv 3 \mod 4 \\ 2(r(\mathbb{Z}_{2}) - q(\mathbb{Z}_{2})) + 2(r(\mathbb{Z}_{3}) - q(\mathbb{Z}_{3})) \\ + 2(r(\mathbb{Z}_{5}) - q(\mathbb{Z}_{5})) & \text{if } q = 1 \end{cases}$$
$$= \begin{cases} 4 + 4 + 6 & \text{if } q > 2 \text{ and } q \equiv 1 \mod 4 \\ 0 + 2 + 4 & \text{if } q > 2 \text{ and } q \equiv 3 \mod 4 \\ 0 + 0 + 2 & \text{if } q = 1. \end{cases}$$
$$= \begin{cases} 14 & \text{if } q > 2 \text{ and } q \equiv 1 \mod 4 \\ 6 & \text{if } q > 2 \text{ and } q \equiv 3 \mod 4 \\ 2 & \text{if } q = 1. \end{cases}$$

Remark 5.1 It is worth noticing that the formula of Corollary 1.4 also provides a calculation for the Whitehead group of $SL_2(\mathbb{Z})$:

$$Wh_1(SL_2(\mathbb{Z})) \simeq Wh_1(PSL_2(\mathbb{Z})) \oplus PSL_2(\mathbb{Z})^{ab} \oplus \mathbb{Z}_2$$
$$\simeq Wh_1(\mathbb{Z}_2 * \mathbb{Z}_3) \oplus \mathbb{Z}_6 \oplus \mathbb{Z}_2$$
$$\simeq \mathbb{Z}_6 \oplus \mathbb{Z}_2,$$

where we have used the fact that $Wh_1(\mathbb{Z}_2 * \mathbb{Z}_3) \simeq Wh_1(\mathbb{Z}_2) \oplus Wh_1(\mathbb{Z}_3) \simeq 0$.

References

[AO06] A. Alves and P. Ontaneda. A formula for the Whitehead group of a three-dimensional crystallographic group. *Topology*, 45(1):1–25, 2006.

- [Bas65] Hyman Bass. The Dirichlet unit theorem, induced characters, and Whitehead groups of finite groups. *Topology*, 4:391–410, 1965.
- [BFJPP00] E. Berkove, F. T. Farrell, D. Juan-Pineda, and K. Pearson. The Farrell-Jones isomorphism conjecture for finite covolume hyperbolic actions and the algebraic *K*-theory of Bianchi groups. *Trans. Amer. Math. Soc.*, 352(12):5689–5702, 2000.
- [BHM93] M. Bökstedt, W. C. Hsiang, and I. Madsen. The cyclotomic trace and algebraic *K*-theory of spaces. *Invent. Math.*, 111(3):465–539, 1993.
- [BJPP01] Ethan Berkove, Daniel Juan-Pineda, and Kimberly Pearson. The lower algebraic *K*-theory of Fuchsian groups. *Comment. Math. Helv.*, 76(2):339–352, 2001.
- [BJPP02] E. Berkove, D. Juan-Pineda, and K. Pearson. A geometric approach to the lower algebraic *K*-theory of Fuchsian groups. *Topology Appl.*, 119(3):269–277, 2002.
- [Car80] David W. Carter. Lower K-theory of finite groups. Comm. Algebra, 8(20):1927–1937, 1980.
- [DKR11] James F. Davis, Qayum Khan, and Andrew Ranicki. Algebraic *K*-theory over the infinite dihedral group: an algebraic approach. *Algebr. Geom. Topol.*, 11(4):2391–2436, 2011.
- [DL98] James F. Davis and Wolfgang Lück. Spaces over a category and assembly maps in isomorphism conjectures in *K* and *L*-theory. *K*-Theory, 15(3):201–252, 1998.
- [DL03] James F. Davis and Wolfgang Lück. The *p*-chain spectral sequence. *K*-*Theory*, 30(1):71–104, 2003. Special issue in honor of Hyman Bass on his seventieth birthday. Part I.
- [DQR11] James F. Davis, Frank Quinn, and Holger Reich. Algebraic *K*-theory over the infinite dihedral group: a controlled topology approach. *J. Topol.*, 4(3):505–528, 2011.
- [FJ93] F. T. Farrell and L. E. Jones. Isomorphism conjectures in algebraic K-theory. J. Amer. Math. Soc., 6(2):249–297, 1993.
- [Fre90] Eberhard Freitag. Hilbert modular forms. Springer-Verlag, Berlin, 1990.
- [Gru08] Joachim Grunewald. The behavior of Nil-groups under localization and the relative assembly map. *Topology*, 47(3):160–202, 2008.
- [Hir73] Friedrich E. P. Hirzebruch. Hilbert modular surfaces. *Enseignement Math.* (2), 19:183–281, 1973.
- [Jah09] Bjørn Jahren. Involutions on the rational *K*-theory of group rings of finite groups. In *Alpine perspectives on algebraic topology*, volume 504 of *Contemp. Math.*, pages 189–202. Amer. Math. Soc., Providence, RI, 2009.
- [JPLMVP11] D. Juan-Pineda, J.-F. Lafont, S. Millan-Vossler, and S. Pallekonda. Algebraic K-theory of virtually free groups. Proc. Roy. Soc. Edinburgh Sect. A, 141(6):1295–1316, 2011.
- [JPML06] Daniel Juan-Pineda and Silvia Millan-López. Invariants associated to the pure braid group of the sphere. *Bol. Soc. Mat. Mexicana* (3), 12(1):27–32, 2006.
- [JPML10] Daniel Juan-Pineda and Silvia Millan-López. The Whitehead group and the lower algebraic *K*-theory of braid groups on \mathbb{S}^2 and \mathbb{RP}^2 . *Algebr. Geom. Topol.*, 10(4):1887–1903, 2010.
- [JPSS14] Daniel Juan-Pineda and Luis Jorge Sánchez Saldaña. On the ranks of the algebraic *K*-theory of hyperbolic groups. *Bol. Soc. Mat. Mex.* (3), 20(2):277–285, 2014.
- [KLR] Holger Kammeyer, Wolfgang Lück, and Henrik Rüping. The Farrell-Jones conjecture for artbitrary lattices in virtually connected lie groups. *arXiv:1401.0876v1*.
- [LO09] Jean-François Lafont and Ivonne J. Ortiz. Lower algebraic K-theory of hyperbolic 3-simplex reflection groups. Comment. Math. Helv., 84(2):297–337, 2009.
- [LS00] Wolfgang Lück and Roland Stamm. Computations of *K* and *L*-theory of cocompact planar groups. *K*-*Theory*, 21(3):249–292, 2000.
- [Oli88] Robert Oliver. *Whitehead groups of finite groups*, volume 132 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1988.
- [Pea98] Kimberly Pearson. Algebraic K-theory of two-dimensional crystallographic groups. K-Theory, 14(3):265–280, 1998.
- [Pre68] Alexander Prestel. Die elliptischen Fixpunkte der Hilbertschen Modulgruppen. Math. Ann., 177:181–209, 1968.

- [vdG88] Gerard van der Geer. Hilbert modular surfaces, volume 16 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1988.
- [Wal78] Friedhelm Waldhausen. Algebraic K-theory of generalized free products. I, II. Ann. of Math. (2), 108(1):135–204, 1978.

Department of Mathematical Sciences, Binghamton University 4400 Vestal Pkwy E, Binghamton, NY 13902, USA Centro de Ciencias Matemáticas, UNAM, Campus Morelia Morelia, Michoacán C.P. 58190, México

bustamante@math.binghamton.edu, luisjorge@matmor.unam.mx