

# Tiling with arbitrary tiles

Vytautas Gruslys\*      Imre Leader\*      Ta Sheng Tan†

August 23, 2016

## Abstract

Let  $T$  be a tile in  $\mathbb{Z}^n$ , meaning a finite subset of  $\mathbb{Z}^n$ . It may or may not tile  $\mathbb{Z}^n$ , in the sense of  $\mathbb{Z}^n$  having a partition into copies of  $T$ . However, we prove that  $T$  does tile  $\mathbb{Z}^d$  for some  $d$ . This resolves a conjecture of Chalcraft.

## 1 Introduction

Let  $T$  be a *tile*, by which we mean a finite non-empty subset of  $\mathbb{Z}^n$  for some  $n$ . It is natural to ask if  $\mathbb{Z}^n$  can be partitioned into copies of  $T$ , that is, into subsets each of which is isometric to  $T$ . If such a partition exists, we say that  $T$  *tiles*  $\mathbb{Z}^n$ .

For instance, consider the following tiling of  $\mathbb{Z}^2$  by copies of the  $C$ -shaped pentomino.




Figure 1: The  $C$ -shaped pentomino tiles  $\mathbb{Z}^2$ .

As another example, the one-dimensional tile  $X.X$  (to be understood as  $\{1, 3\}$ ) tiles  $\mathbb{Z}$ , and so does  $XX.X$ . On the other hand,  $XX.XX$  is a one-dimensional tile that does not tile  $\mathbb{Z}$ . Does it tile some space of higher dimension? The following diagram shows that  $XX.XX$  does tile  $\mathbb{Z}^2$ .

---

\*Department of Pure Mathematics and Mathematical Statistics, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, CB3 0WB Cambridge, United Kingdom; e-mail: {v.gruslys,i.leader}@dpmmms.cam.ac.uk.

†Institute of Mathematical Sciences, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia; e-mail: tstan@um.edu.my. This author acknowledges support received from the Ministry of Education of Malaysia via FRGS grant FP048-2014B.




Figure 2: This pattern is formed from disjoint copies of  $XX.XX$ ; copies of the pattern may be stacked vertically to tile  $\mathbb{Z}^2$ .

A similar pattern works for  $XXX.XX$  in  $\mathbb{Z}^2$ . However, one can check by hand that  $XXX.XXX$  does not tile  $\mathbb{Z}^2$ . Does it tile  $\mathbb{Z}^3$ , or  $\mathbb{Z}^d$  for some  $d$ ? What about more complicated one-dimensional tiles?

Let us now consider a couple of two-dimensional examples. Let  $T$  denote the  $3 \times 3$  square with the central point removed. Clearly  $T$  does not tile  $\mathbb{Z}^2$ , since the hole in a copy of  $T$  cannot be filled. However, in  $\mathbb{Z}^3$  there is enough space for one copy of  $T$  to fill the hole of another. (Of course, this in no way implies that  $T$  does tile  $\mathbb{Z}^3$ .)

For a ‘worse’ example, consider the  $5 \times 5$  square with the central point removed. Two copies of such tile cannot be interlinked in  $\mathbb{Z}^3$ . However, there is, of course, enough space in  $\mathbb{Z}^4$  to fill the hole, as demonstrated in the following diagram.




Figure 3: The diagram on the right is four-dimensional and shows a  $5 \times 5 \times 5 \times 5$  region of  $\mathbb{Z}^4$ . Let  $x_1, x_2, x_3, x_4$  be the directions of  $\mathbb{Z}^4$ . Each of the five  $5 \times 5 \times 5$  cubes corresponds to a fixed value of  $x_1$ . Increasing the value of  $x_1$  by 1 means jumping from a cube to the cube on its right. This four-dimensional diagram contains two copies of the two-dimensional tile depicted on the left side. One copy is horizontal and can be found in the top left part of the diagram. The second copy is formed by the vertical columns.

Chalcraft [8, 9] made the remarkable conjecture that every tile  $T \subset \mathbb{Z}$ , or even  $T \subset \mathbb{Z}^n$ , does tile  $\mathbb{Z}^d$  for some  $d$ .

**Conjecture 1** (Chalcraft). *Let  $T \subset \mathbb{Z}^n$  be a tile. Then  $T$  tiles  $\mathbb{Z}^d$  for some  $d$ .*

It is not important if reflections are allowed when forming copies of a tile. Indeed, any reflection of an  $n$ -dimensional tile can be obtained by rotating it in  $n + 1$  dimensions. It is

also not important if only connected tiles are considered, as it is an easy exercise to show that any disconnected tile in  $\mathbb{Z}^n$  tiles a connected tile in  $\mathbb{Z}^{2n}$ .

In this paper we prove Chalcraft's conjecture.

**Theorem 2.** *Let  $T \subset \mathbb{Z}^n$  be a tile. Then  $T$  tiles  $\mathbb{Z}^d$  for some  $d$ .*

Interestingly, the problem is not any easier for tiles  $T \subset \mathbb{Z}$ . Indeed, the proof for one-dimensional tiles seems to us to be as hard as the general problem.

The plan of the paper is as follows. In Section 2 we prove a special case of the theorem, namely when  $T$  is an interval in  $\mathbb{Z}$  with one point removed. The aim of this section is to demonstrate some of the key ideas in a simple setting. The proof of the general case builds on these ideas and on several additional ingredients. We give a proof of Theorem 2 in Section 3. Finally, in Section 4 we give some open problems.

We end the section with some general background. A lot of work has been done about tiling  $\mathbb{Z}^2$  by polyominoes (a polyomino being a connected tile in  $\mathbb{Z}^2$ ). Golomb [3] proved that every polyomino of size at most 6 tiles  $\mathbb{Z}^2$ . In [4] he also proved that there is no algorithm which decides, given a finite set of polyominoes, if  $\mathbb{Z}^2$  can be tiled with their copies – this is based on the work of Berger [1], who showed a similar undecidability result for Wang tiles (which are certain coloured squares). However, it is not known if such an algorithm exists for single polyominoes. A related unsolved problem is to determine whether there is a polyomino which tiles  $\mathbb{Z}^2$  but such that every tiling is non-periodic. On the other hand, Wijshoff and van Leeuwen [10] found an algorithm which determines if disjoint translates (rather than translates, rotations and reflections) of a single given polyomino tile  $\mathbb{Z}^2$ . A vast number of results and questions regarding tilings of  $\mathbb{Z}^2$  by polyominoes and other shapes are compiled in Grünbaum and Shephard [6].

One may also wish to know if a given polyomino tiles some finite region of  $\mathbb{Z}^2$ , say a rectangle. This class of questions has also received significant attention, producing many beautiful techniques and invariants – see, for example, [2, 5, 7]. In the context of this paper, we observe that there are tiles which cannot tile any (finite) cuboid of any dimension. For example, consider the plus-shaped tile of size 5 in  $\mathbb{Z}^2$ : this tile cannot cover the corners of any cuboid. In fact, there are one-dimensional such tiles. For example, let  $T \subset \mathbb{N}$ , where  $\mathbb{N} = \{1, 2, \dots\}$ , be a symmetric tile (meaning that  $-T$  is a translate of  $T$ ) whose associated polynomial  $p(x) = \sum_{t \in T} x^t$  does not have all of its non-zero roots on the unit circle – it turns out that such  $T$  cannot tile a cuboid (see [9]).

## 2 Tiling $\mathbb{Z}^d$ by an interval minus a single point

### 2.1 Overview

Before starting the proof of Theorem 2, we demonstrate some of the key ideas in a simple setting, where the tile is a one-dimensional interval with one point removed. We give a self-contained proof of the general case in Section 3, but it will build on the ideas in this section.

We write  $[k] = \{1, \dots, k\}$ .

**Theorem 3.** *Fix integers  $k \geq 3$  and  $i \in \{2, \dots, k-1\}$  and let  $T$  be the tile  $[k] \setminus \{i\}$ . Then  $T$  tiles  $\mathbb{Z}^d$  for some  $d$ .*

The tile  $T = [k] \setminus \{i\}$  will remain fixed throughout this section.

The proof is driven by two key ideas. A first natural idea is to use *strings*, where a string is a one-dimensional infinite line in  $\mathbb{Z}^d$  with every  $k$ -th point removed. Note that any string is a union of disjoint copies of  $T$ . An obvious way to use strings would be to partition  $\mathbb{Z}^d$  into them. Although this is an attractive idea, it is not possible, for the following simple reason: if we consider just the fixed cuboid  $[k]^d \subset \mathbb{Z}^d$ , then every string intersects it in exactly 0 or  $k - 1$  points, but the order of  $[k]^d$  is not divisible by  $k - 1$ .

This suggests a refinement of the idea. We will try to use strings parallel to  $d - 1$  of the  $d$  directions, while the remaining direction will be special and copies of  $T$  parallel to it will be used even without forming strings. In other words, we will view  $\mathbb{Z}^d$  as  $\mathbb{Z} \times \mathbb{Z}^{d-1}$ , that is, as being partitioned into  $(d - 1)$ -dimensional *slices* according to the value of the first coordinate. We will first put down some tiles parallel to the first direction (each such tile intersects multiple slices), and then complete the tilings in each slice separately by strings.

To do this we need another idea. What subsets of  $\mathbb{Z}^{d-1}$  can be tiled by strings? Note that a partial tiling of  $\mathbb{Z}^{d-1}$  by strings can be identified with a partial tiling of the discrete torus  $\mathbb{Z}_k^{d-1}$  (where  $\mathbb{Z}_k$  denotes the integers modulo  $k$ ), where a tile in  $\mathbb{Z}_k^{d-1}$  means any line with one point removed. The size of  $\mathbb{Z}_k^{d-1}$  is  $k^{d-1} \equiv 1 \pmod{k-1}$ , so any such partial tiling of  $\mathbb{Z}_k^{d-1}$  must leave out  $1 \pmod{k-1}$  points. Of course, it is far from true that any subset of  $\mathbb{Z}_k^{d-1}$  of size a multiple of  $k - 1$  may be partitioned into tiles. However, our plan is to find a large supply of sets that do have this property. In particular, it turns out that a key idea will be to find a large set  $C \subset \mathbb{Z}_k^{d-1}$  such that for any choice of distinct elements  $x_1, \dots, x_m \in C$  with  $m \equiv 1 \pmod{k-1}$ ,  $T$  does tile  $\mathbb{Z}_k^{d-1} \setminus \{x_1, \dots, x_m\}$ .




Figure 4: A partial tiling of  $\mathbb{Z}_k^2$  (here  $k = 5$ ) corresponds to a partial tiling of  $\mathbb{Z}^2$  by strings.

These ideas work together as follows (see Figure 5). First, in  $\mathbb{Z} \times \mathbb{Z}_k^{d-1}$  (for large  $d$ ) we find a subset  $X$  which is a disjoint union of translates of  $T \times \{0\}^{d-1}$  and has the property that for any  $n \in \mathbb{Z}$  the set  $\{x \in \mathbb{Z}_k^{d-1} : (n, x) \in X\}$  is a subset of  $C$  of size congruent to 1 modulo  $k - 1$ . Then  $T$  tiles  $(\{n\} \times \mathbb{Z}_k^{d-1}) \setminus X$ . This holds for all  $n \in \mathbb{Z}$ , so in fact  $T$  tiles  $\mathbb{Z} \times \mathbb{Z}_k^{d-1}$ , and hence it tiles  $\mathbb{Z}^d$ , establishing Theorem 3.

The rest of this section is organised as follows. In Section 2.2 we consider partial tilings by strings. In Section 2.3 we consider the special direction. Both ideas are combined in Section 2.4, where a full proof of Theorem 3 is given.




Figure 5: The aim is to put down tiles parallel to one of the directions so that the remainder of each slice could be tiled by strings. This diagram only symbolically visualises this principle. In particular, the slices here are two-dimensional, while in the proof they can have much higher dimension.

## 2.2 Tiling $\mathbb{Z}_k^d$ with some elements removed

For any  $1 \leq j \leq d$ , define the  $j$ -th corner of  $\mathbb{Z}_k^d$  to be  $c_{j,d}$  where

$$c_{j,d} = \underbrace{(0, \dots, 0, k-1, 0, \dots, 0)}_{d \text{ coordinates}} \in \mathbb{Z}_k^d.$$

$\downarrow$   
 $j$ -th coordinate

Write  $C_d = \{c_{j,d} : j = 1, \dots, d\}$  for the set of corners.




Figure 6: The set of corners  $C_4$  when  $k = 6$ . In this diagram the space  $\mathbb{Z}_6^4 = \{(x_1, x_2, x_3, x_4) : x_i \in \{0, \dots, 5\}\}$  is split from left to right, according to the value of  $x_4$ , into 6 three-dimensional slices.

Looking ahead, our aim later will be to provide some copies of  $T$  in the  $x_1$ -direction in  $\mathbb{Z} \times \mathbb{Z}_k^d$ , at heights corresponding to points of  $C_d$ , and in such a way that what remains in each  $\mathbb{Z}_k^d$  can be partitioned into lines with one point removed. But first we need to create a useful supply of such subsets of  $\mathbb{Z}_k^d$ .

Recall that  $|\mathbb{Z}_k^d| \equiv 1 \pmod{k-1}$ , so if  $T$  tiles some set  $X \subset \mathbb{Z}_k^d$  (here and in the remainder of this section  $T$  is identified with its image under the projection  $\mathbb{Z} \rightarrow \mathbb{Z}_k$ , so its copies in  $\mathbb{Z}_k^d$  are lines with one point removed), then  $|\mathbb{Z}_k^d \setminus X| \equiv 1 \pmod{k-1}$ . In this section we will prove Lemma 4, which is an approximate converse of this statement.

**Lemma 4.** *Let  $d \geq 1$  and suppose that  $S \subset C_d$  is such that  $|S| \equiv 1 \pmod{k-1}$  and  $|S| \leq d - \log_k d$ . Then  $T$  tiles  $\mathbb{Z}_k^d \setminus S$ .*

In fact, this lemma holds even without the assumption that  $|S| \leq d - \log_k d$ , but we keep it for the sake of simpler presentation.

We will prove Lemma 4 at the end of this section. Meanwhile, we collect the tools needed for the proof. In fact, there are several ways to prove Lemma 4. The method outlined here is quite general, and we will build on it in Section 3.

We start with a simple proposition.

**Proposition 5.** *Let  $d \geq 1$  and  $x \in \mathbb{Z}_k^d$ . Then  $T$  tiles  $\mathbb{Z}_k^d \setminus \{x\}$ .*

*Proof (see Figure 7).* Use induction on  $d$ . If  $d = 1$ , then  $\mathbb{Z}_k \setminus \{x\}$  is itself a translate of  $T$ . Now suppose that  $d \geq 2$  and write  $x = (x_1, \dots, x_d)$ ,  $\hat{x} = (x_1, \dots, x_{d-1})$ . By the induction hypothesis, for each  $j \in \mathbb{Z}_k$ ,  $(\mathbb{Z}_k^{d-1} \setminus \{\hat{x}\}) \times \{j\}$  can be tiled with copies of  $T$ . It remains to tile  $\{\hat{x}\} \times (\mathbb{Z}_k \setminus \{x_d\})$ , but this is itself a copy of  $T$ .  $\square$




Figure 7: The induction step in the proof of Proposition 5. The grey cube represents  $x$ . The vertical column in which  $x$  lies, without  $x$  itself, is a copy of  $T$ . Each horizontal slice minus the point in this column can be tiled by the induction hypothesis.

Let  $X \subset \mathbb{Z}_k^d$  (for any  $d \geq 1$ ) be such that  $T$  tiles  $\mathbb{Z}_k^d \setminus X$ . We will say that such  $X$  is a *hole* in  $\mathbb{Z}_k^d$ . The intuition for  $X$  is that it is a set that remains uncovered after an attempt to tile  $\mathbb{Z}_k^d$  by copies of  $T$ .

We can identify  $X$  with a higher-dimensional set  $X' = X \times \{0\} \subset \mathbb{Z}_k^{d+1}$ . One can easily verify that  $X'$  is a hole in  $\mathbb{Z}_k^{d+1}$ . More importantly, we will show in the following proposition that a single additional point of  $X'$  can be covered in exchange for leaving the  $(d+1)$ -st corner of  $\mathbb{Z}_k^{d+1}$  uncovered (see Figure 9). This is why, for any  $S \subset \mathbb{Z}_k^d$ , we define

$$S^\dagger = (S \times \{0\}) \cup \{c_{d+1, d+1}\} \subset \mathbb{Z}_k^{d+1}.$$

Note that the definition of  $S^\dagger$  and the definition of  $S$  being a hole depend not only on  $S$ , but also on the dimension of the underlying discrete torus  $\mathbb{Z}_k^d$ . For  $m \geq 1$ , we will use the shorthand  $S^{\dagger(m)}$  to denote the result of  $m$  consecutive applications of the  $\dagger$  operation to  $S$ , that is,

$$S^{\dagger(m)} = \underbrace{S^{\dagger \dots \dagger}}_m \subset \mathbb{Z}_k^{d+m}.$$




Figure 8: Suppose  $S$  is the subset of  $\mathbb{Z}_k^2$  given in the diagram on the left (here  $k = 6$ ). The diagram in the middle depicts  $S^\dagger$ , and the diagram on the right depicts  $S^{\dagger(2)}$ . Observe that  $S$  is a hole in  $\mathbb{Z}_k^2$ , but  $S^\dagger$  and  $S^{\dagger(2)}$  are not holes in  $\mathbb{Z}_k^3$  and  $\mathbb{Z}_k^4$ , respectively.

**Proposition 6.** *Let  $d \geq 1$  and let  $X \subset \mathbb{Z}_k^d$  be a hole. Then for each  $x \in X$  the set  $(X \setminus \{x\})^\dagger$  is a hole in  $\mathbb{Z}_k^{d+1}$ .*




Figure 9: An illustration of the statement of Proposition 6. The aim is to show that  $T$  tiles  $\mathbb{Z}_k^{d+1} \setminus (X \setminus \{x\})^\dagger$ .

*Proof (see Figure 10).* Use (i) a tiling of  $\mathbb{Z}_k^d \setminus X$  for  $(\mathbb{Z}_k^d \setminus X) \times \{0\}$ , and (ii) one copy of  $T$  to cover  $\{x\} \times (\mathbb{Z}_k \setminus \{k-1\})$ . By Proposition 5, (iii)  $(\mathbb{Z}_k^d \setminus \{(0, \dots, 0)\}) \times \{k-1\}$  and (iv)  $(\mathbb{Z}_k^d \setminus \{x\}) \times \{i\}$ ,  $i \in \{1, \dots, k-2\}$ , can each be tiled by copies of  $T$ .  $\square$




Figure 10: The bottom horizontal piece (i) is tilable because  $X$  is a hole, and the other horizontal pieces (iii) and (iv) are tilable by Proposition 5. The remaining vertical column (ii) is a copy of  $T$ .

We will apply Proposition 6 inductively, that is, in the form of the following corollary.

**Corollary 7.** *Let  $d \geq 1$  and let  $X \subset \mathbb{Z}_k^d$  be a hole. Then for any distinct elements  $x_1, \dots, x_m \in X$ , the set  $(X \setminus \{x_1, \dots, x_m\})^{\dagger(m)}$  is a hole in  $\mathbb{Z}_k^{d+m}$ .  $\square$*

We are now ready to prove Lemma 4.

*Proof of Lemma 4.* Write  $|S| = m$  and  $r = d - m$ . By symmetry, we can assume that

$$S = \{c_{j,d} : j = r + 1, \dots, d\}.$$

Our aim is to prove that  $S$  is a hole in  $\mathbb{Z}_k^d$ . Note that  $S = \emptyset^{\dagger(m)}$ , where the empty set  $\emptyset$  is considered as a subset of  $\mathbb{Z}_k^r$ . Therefore by Corollary 7 it suffices to find a hole  $X \subset \mathbb{Z}_k^r$  with  $|X| = m$ .

This can be done by partitioning  $\mathbb{Z}_k^r$  into a singleton  $\{x\}$  and copies of  $T$  (this can be done by Proposition 5), and letting  $X$  be the union of  $\{x\}$  and the appropriate number of copies of  $T$ . By assumption,  $m \equiv 1 \pmod{k-1}$  so the only potential problem with this construction of  $X$  is if  $|\mathbb{Z}_k^r| < m$ . However, this is ruled out by the assumption that  $m \leq d - \log_k d$ .  $\square$

### 2.3 Using one special direction to get $T$ -tilable slices

The purpose of this section is to demonstrate that tiles in the first direction in  $\mathbb{Z} \times \mathbb{Z}_k^{d-1}$  (that is, translates of  $T \times \{0\}^{d-1}$ ) can be combined in such a way that the uncovered part of each slice can be tiled by copies of  $T$  using Lemma 4. The exact claim is as follows.

**Lemma 8.** *There exists a number  $\ell \geq 1$  such that for any  $d \geq 1$  and any set  $C \subset \mathbb{Z}_k^{d-1}$  of order  $|C| \geq \ell$  there is a set  $X \subset \mathbb{Z} \times C$ , satisfying:*

- (a)  $X$  is a union of disjoint sets of the form  $(T + n) \times \{c\}$  with  $n \in \mathbb{Z}$  and  $c \in C$ ;
- (b)  $|(\{n\} \times C) \cap X| \equiv 1 \pmod{k-1}$  for every  $n \in \mathbb{Z}$ ;
- (c)  $|(\{n\} \times C) \cap X| \leq \ell$  for every  $n \in \mathbb{Z}$ .




Figure 11: A possible construction of  $X$ . In this example the aim is to have 1 modulo 6 elements covered in each column.



We start with the following trivial proposition.

**Proposition 9.** *There is a function  $f : \mathbb{Z} \rightarrow \{0, \dots, k-2\}$  such that for each  $x \in \mathbb{Z}$*

$$\sum_{y \in T} f(x-y) \equiv 1 \pmod{k-1}.$$

*Proof.* Start by defining  $f(n) = 0$  for  $-k+1 \leq n \leq -1$ . Now define  $f(n)$  for  $n \geq 0$  as follows. Suppose that for some  $n \geq 0$  the values of  $f(j)$  are already defined for all  $j$  such that  $-k+1 \leq j \leq n-1$ . Then the value of  $f(n)$  is uniquely defined by

$$f(n) \equiv 1 - \sum_{y \in T \setminus \{1\}} f(n+1-y) \pmod{k-1}.$$

Define  $f(n)$  for all  $n \leq -k$  in a similar way. □

Now Lemma 8 can be proved quickly.

*Proof of Lemma 8.* Write  $\ell = 2k(k-2)$  and suppose that  $|C| = \ell$ . Let  $f : \mathbb{Z} \rightarrow \{0, \dots, k-2\}$  be as given by Proposition 9. The aim is to choose subsets  $S_n \subset C$  for every  $n \in \mathbb{Z}$ , with orders satisfying  $|S_n| = f(n)$ , and such that  $S_m \cap S_n = \emptyset$  whenever  $m \neq n$  and  $(T+m) \cap (T+n) \neq \emptyset$ . Then  $X$  can be taken to be  $\bigcup_{n \in \mathbb{Z}} (T+n) \times S_n$ .

Fix any enumeration of  $\mathbb{Z}$ , and define the sets  $S_n$  one by one in that order. When defining  $S_n$ , there can be at most  $2k-1$  choices of  $m$  with  $S_m$  already defined and  $m-n \in T-T$ . Moreover,  $|S_m| \leq k-2$  for each  $m$ . Therefore to be able to find  $S_n$  it is enough to have  $|C| - (2k-1)(k-2) \geq f(n)$ . Finally, this condition is ensured by the choice of  $\ell$ , completing the proof in the case when  $|C| = \ell$ . If  $|C| > \ell$ , we are done by restricting to a subset of  $C$  of size exactly  $\ell$ . □

## 2.4 Completing the proof of Theorem 3

It was noted in Section 2.1 that Lemmas 4 and 8 together imply that for some  $d \geq 1$

$$T \text{ tiles } \mathbb{Z} \times \mathbb{Z}_k^{d-1}, \tag{1}$$

and therefore

$$T \text{ tiles } \mathbb{Z}^d, \tag{2}$$

implying Theorem 3. However, some abuse of notation is already present in the statement of (1). In this section we will carefully explain what is meant by (1), why it follows from the two lemmas and how it implies (2). In doing so, we will complete the proof of Theorem 3.

To avoid confusion, within this section we use quite precise language. Although this might seem pedantic here, for later it will be very important to have precise notation available. We denote the elements of  $\mathbb{Z}_k$  by  $\bar{x}$  for  $x \in \mathbb{Z}$  (instead of identifying them with  $x$ , which was our preferred notation in the rest of the section), and we will denote the image of  $T$  under the natural projection  $\pi : \mathbb{Z} \rightarrow \mathbb{Z}_k$  by  $\pi(T)$  rather than simply by  $T$ .

**Theorem 3.** *Fix integers  $k \geq 3$  and  $i \in \{2, \dots, k-1\}$  and let  $T$  be the tile  $[k] \setminus \{i\}$ . Then  $T$  tiles  $\mathbb{Z}^d$  for some  $d$ .*

*Proof.* Fix a large  $d$  (more precisely, first let  $\ell$  be as given by Lemma 8 and then fix  $d$  such that  $d - 1 - \log_k(d - 1) \geq \ell$ ).

Denote the projection map  $\mathbb{Z} \rightarrow \mathbb{Z}_k$  by  $\pi$ , and consider the following subsets of  $\mathbb{Z} \times \mathbb{Z}_k^{d-1}$ :

$$\begin{aligned} \mathbb{T}_1 &= T \times \{\bar{0}\} \times \{\bar{0}\} \times \cdots \times \{\bar{0}\}, \\ \mathbb{T}_2 &= \{0\} \times \pi(T) \times \{\bar{0}\} \times \cdots \times \{\bar{0}\}, \\ &\vdots \\ \mathbb{T}_d &= \{0\} \times \{\bar{0}\} \times \{\bar{0}\} \times \cdots \times \pi(T). \end{aligned}$$

Recall from Section 2.2 the definition of

$$C_{d-1} = \left\{ \underbrace{(\bar{0}, \dots, \bar{0}, \overbrace{k-1}^{j\text{-th coordinate}}, \bar{0}, \dots, \bar{0})}_{d-1 \text{ coordinates}} : j = 1, \dots, d-1 \right\} \subset \mathbb{Z}_k^{d-1}.$$

By Lemma 8, there is a set  $X \subset \mathbb{Z} \times C_{d-1}$ , which is a union of disjoint translates of  $\mathbb{T}_1$  and for each  $n \in \mathbb{Z}$  satisfies  $|(\{n\} \times C_{d-1}) \cap X| \leq d - 1 - \log_k(d - 1)$  and  $|(\{n\} \times C_{d-1}) \cap X| \equiv 1 \pmod{k-1}$ . Hence, by Lemma 4,  $(\{n\} \times \mathbb{Z}_k^{d-1}) \setminus X$  is a union of disjoint translates of  $\mathbb{T}_2, \dots, \mathbb{T}_d$  for each  $n \in \mathbb{Z}$ . Therefore  $\mathbb{Z} \times \mathbb{Z}_k^{d-1}$  is a union of disjoint translates of  $\mathbb{T}_1, \dots, \mathbb{T}_d$  (this is exactly what is meant by (1)).

More explicitly, there are integers  $1 \leq t(\alpha) \leq d$  and  $x_1(\alpha), \dots, x_d(\alpha) \in \mathbb{Z}$ , indexed by  $\alpha \in A$ , such that  $\mathbb{Z} \times \mathbb{Z}_k^{d-1}$  is the disjoint union

$$\mathbb{Z} \times \mathbb{Z}_k^{d-1} = \bigsqcup_{\alpha \in A} \left[ \mathbb{T}_{t(\alpha)} + (x_1(\alpha), \overline{x_2(\alpha)}, \dots, \overline{x_d(\alpha)}) \right].$$

From this it follows that, in fact,  $\mathbb{Z}^d$  is  $T$ -tilable. Indeed, consider the following subsets of  $\mathbb{Z}^d$ :

$$\begin{aligned} \mathbb{T}'_1 &= T \times \{0\} \times \{0\} \times \cdots \times \{0\}, \\ \mathbb{T}'_2 &= \{0\} \times T \times \{0\} \times \cdots \times \{0\}, \\ &\vdots \\ \mathbb{T}'_d &= \{0\} \times \{0\} \times \{0\} \times \cdots \times T. \end{aligned}$$

Then we can express  $\mathbb{Z}^d$  as the disjoint union

$$\mathbb{Z}^d = \bigsqcup_{\substack{\alpha \in A \\ c_2, \dots, c_d \in \mathbb{Z}}} \left[ \mathbb{T}'_{t(\alpha)} + (x_1(\alpha), x_2(\alpha) + kc_2, \dots, x_d(\alpha) + kc_d) \right]. \quad \square$$

### 3 The general case

Recall the statement of the main theorem.

**Theorem 2.** *Let  $T \subset \mathbb{Z}^n$  be a tile. Then  $T$  tiles  $\mathbb{Z}^d$  for some  $d$ .*

In this section we prove the main theorem by generalising the approach demonstrated in Section 2. We have to account for two ways in which Theorem 3 is a special case: firstly, the tile can be multidimensional; secondly, even in the one-dimensional case the tile can have more complicated structure than in Section 2.

It turns out that dealing with the first issue does not add significant extra difficulty to the proof, provided that the right setting is chosen. Namely, most of the intermediate results will be stated in terms of abelian groups rather than integer lattices. This way a multidimensional tile  $T \subset \mathbb{Z}^b$  can be considered as being one-dimensional, if  $\mathbb{Z}^b$  (rather than  $\mathbb{Z}$ ) is chosen as the underlying abelian group. Moreover, this point of view is vital for comparing periodic tilings of an integer lattice with tilings of a discrete torus, already an important idea in the proof of the special case.

On the other hand, dealing with the second issue requires significant effort. It involves finding the right way to generalise the two key ideas from Section 2, as well as introducing a new ingredient that allows the argument to be applied iteratively.

We now introduce some definitions. Given an abelian group  $G$ , we call any non-empty subset  $T \subset G$  a *tile* in  $G$ . Given abelian groups  $G_1, \dots, G_d$  and corresponding tiles  $T_i \subset G_i$ , consider the following subsets of  $G_1 \times \dots \times G_d$ :

$$\begin{aligned} \mathbb{T}_1 &= T_1 \times \{0\} \times \dots \times \{0\}, \\ \mathbb{T}_2 &= \{0\} \times T_2 \times \dots \times \{0\}, \\ &\vdots \\ \mathbb{T}_d &= \{0\} \times \{0\} \times \dots \times T_d. \end{aligned}$$

Any translate of such  $\mathbb{T}_i$  (that is, a set of the form  $\mathbb{T}_i + x$  for  $x \in G_1 \times \dots \times G_d$ ) is called a *copy* of  $T_i$ . We say that a subset  $X \subset G_1 \times \dots \times G_d$  is  $(T_1, \dots, T_d)$ -tilable if  $X$  is a disjoint union of copies of  $T_1, \dots, T_d$ .

It will often be the case that  $(G_1, T_1) = \dots = (G_d, T_d) = (G, T)$ . Then we will use the term  $T$ -tilable as a shorthand for  $(T, \dots, T)$ -tilable.

More generally, we may consider subsets of  $G_1^{d_1} \times \dots \times G_m^{d_m}$  where  $G_1, \dots, G_m$  are abelian groups with tiles  $T_i \subset G_i$ . In this setting we would say that a subset is  $(d_1 \cdot T_1, \dots, d_m \cdot T_m)$ -tilable. In other words, each  $d_i \cdot T_i$  replaces

$$\underbrace{T_i, \dots, T_i}_{d_i}.$$

However, we suppress “1.” in the notation. So, for example, we could say that a subset of  $G_1^7 \times G_2 \times G_3^{10}$  is  $(7 \cdot T_1, T_2, 10 \cdot T_3)$ -tilable.

### 3.1 A summary of the proof

Let  $T$  be a fixed finite tile in  $\mathbb{Z}^b$ . Without loss of generality assume that  $T \subset [k]^b$  for some  $k \geq 1$ . Then, writing  $\pi : \mathbb{Z}^b \rightarrow \mathbb{Z}_k^b$  for the projection map,  $\pi(T)$  is a tile in  $G = \mathbb{Z}_k^b$ .

In the light of the argument from Section 2, one might hope to find a positive integer  $d$  and a large family  $\mathcal{F}$  of disjoint subsets of  $G^d$  with the property that whenever a subfamily  $\mathcal{S} \subset \mathcal{F}$  with  $|\mathcal{S}| \equiv 1 \pmod{|T|}$  is chosen, the set  $G^d \setminus (\cup_{S \in \mathcal{S}} S)$  is  $\pi(T)$ -tilable. However, this seems to be achievable only in the case when  $\pi(T)$  is in a certain sense a ‘dense’ subset of  $G$ .

If  $\pi(T)$  is sparse, we achieve a weaker aim. Namely, we find a certain set  $X \subset G^d$  which has sufficiently nice structure and is a denser subset of  $G^d$  than  $\pi(T)$  is of  $G$ . Also, we find a large family  $\mathcal{F}$  of disjoint subsets of  $X$  such that for any  $\mathcal{S} \subset \mathcal{F}$  of appropriate size  $X \setminus (\bigcup_{S \in \mathcal{S}} S)$  is  $\pi(T)$ -tilable. Taking copies of  $T$  in the special direction, we can now tile  $\mathbb{Z}^b \times X$ .

Repeating this process, we can use copies of  $T$  and  $\mathbb{Z}^b \times X$  to tile  $\mathbb{Z}^p \times Y$  for an even denser subset  $Y \subset G^l$ . After finitely many iterations of this procedure we tile the whole of  $\mathbb{Z}^q \times G^m$  for some possibly large  $q$  and  $m$ . From this it follows that  $\mathbb{Z}^{q+bm}$  is  $T$ -tilable.

The rest of this section is organised as follows. In Section 3.2 we show how any tile in a (finite) abelian group  $H$  can be used to *almost* tile a sufficiently nice denser subset of  $H^d$  for some  $d$ . This is the most complicated part of the proof, but it shares a similar structure with the simpler argument in Section 2.2.

In Section 3.3 we show how one special dimension can be used to cover the gaps in every slice. The argument is almost identical to the one in Section 2.3.

In Section 3.4 we observe some simple transitivity properties of tilings. They enable the iterative application of the process. The ideas in this section are fairly straightforward.

Finally, in Section 3.5 we compile the tools together and complete the proof of Theorem 2.

### 3.2 Almost tiling denser multidimensional sets

Our goal is to prove the following lemma.

**Lemma 10.** *Let  $T \not\subset G$  be a tile in a finite abelian group  $G$ . Then there is a set  $A \subset G$ , with  $T \not\subset A$ , having the following property. Given any  $d_0 \geq 1$ , there is some  $d \geq d_0$  and a family  $\mathcal{F}$  consisting of at least  $d_0$  pairwise disjoint subsets of  $A^d$  such that*

$$G \times \left( A^d \setminus \bigcup_{S \in \mathcal{S}} S \right) \subset G^{d+1}$$

*is  $T$ -tilable whenever  $\mathcal{S} \subset \mathcal{F}$  satisfies  $|\mathcal{S}| \equiv 1 \pmod{|T|}$ .*

Before presenting the proof, we make a few definitions that will hold throughout this section. First, let  $G$  and  $T$  be fixed as in the statement of Lemma 10. Since  $T \neq G$ , we can fix an  $x \in G$  such that  $T + x \neq T$ . Define

$$\begin{aligned} T^{\text{up}} &= T + x, \\ C^{\text{up}} &= T^{\text{up}} \setminus T, \\ C_{\text{down}} &= T \setminus T^{\text{up}}, \\ A &= T \cup T^{\text{up}} \end{aligned}$$

(see Figures 12 and 13).

We will use  $A$  from this definition in the proof of Lemma 10. For the family  $\mathcal{F}$  we will take all sets of the following form. For any integers  $1 \leq i \leq d$ , write

$$C_{i,d} = \underbrace{C_{\text{down}} \times \cdots \times C_{\text{down}} \times \overset{i\text{-th component}}{\downarrow} C^{\text{up}} \times C_{\text{down}} \times \cdots \times C_{\text{down}}}_{d \text{ components}} \subset A^d$$




Figure 12: An illustration of the definitions.




Figure 13: A four-dimensional diagram of  $A^4$ . The sets  $C_{\text{down}}$  and  $C^{\text{up}}$  are marked on two of the axes. In this example  $|A| = 5$  and  $|C_{\text{down}}| = |C^{\text{up}}| = 2$ . This and the following four-dimensional diagrams in this section should be understood more generally as depicting  $A^d$  for any  $d$ , the three-dimensional slices representing copies of  $A^{d-1}$ .

(see Figure 14). Also write  $C_{0,d} = (C_{\text{down}})^d$ . Note that if  $i \neq j$ , then  $C_{i,d} \cap C_{j,d} = \emptyset$ .

Finally, as  $T$  is fixed, we can simply say *tilable* instead of  $T$ -tilable.

One of the reasons why these definitions are useful is that they allow the following analogue of Proposition 5.

**Proposition 11.** *For any integers  $d \geq 1$  and  $0 \leq i \leq d$ , the set  $A^d \setminus C_{i,d}$  is tilable.*

*Proof (see Figure 15).* Use induction on  $d$ . If  $d = 1$ , observe that  $A = C^{\text{up}} \sqcup T = C_{\text{down}} \sqcup T^{\text{up}}$ , and so  $A \setminus C_{i,1}$  ( $= A \setminus C^{\text{up}}$  or  $A \setminus C_{\text{down}}$ ) is a translate of  $T$ .

Now suppose that  $d \geq 2$  and without loss of generality assume that  $i \neq d$ . By the induction hypothesis, for each  $g \in A$ , the slice  $(A^{d-1} \setminus C_{i,d-1}) \times \{g\}$  can be  $T$ -tiled. It remains to tile the set  $C_{i,d-1} \times (A \setminus C_{\text{down}}) = C_{i,d-1} \times T^{\text{up}}$ , but this is obviously a union of disjoint copies of  $T$ .  $\square$

We now make a series of definitions that are useful for lifting subsets of lower-dimensional spaces to higher-dimensional spaces.

A *basic set* is a set of the form  $A^d$ ,  $G \times A^d$  or  $\{g\} \times A^d$  for some  $g \in G$ , with  $d$  any positive integer. Let  $X$  be a subset of a basic set  $\Omega$  and write  $\Omega = W \times A^d$  (so  $W = A^0$ ,  $G$  or  $\{g\}$ ) for




Figure 14: A four-dimensional diagram, which extends the previous diagram. Note that  $C_{3,4}$  and  $C_{4,4}$  both intersect two three-dimensional slices, because in this example  $|C_{\text{down}}| = |C^{\text{up}}| = 2$ .




Figure 15: The induction step in the proof of Proposition 11. The set  $C_{i,d-1} \times T^{\text{up}}$  is a union of copies of  $T^{\text{up}}$ . In each slice it remains to tile a copy of  $A^{d-1} \setminus C_{i,d-1}$ . This can be done by the induction hypothesis.

some  $g \in G$ ). We define

$$X^\dagger = (X \times C_{\text{down}}) \cup (W \times C_{d+1,d+1}) \subset W \times A^{d+1}$$

(see Figure 16).

Moreover, for any  $m \geq 1$  we use the shorthand  $X^{\dagger(m)}$  to denote the result of  $m$  consecutive applications of the  $\dagger$  operation to  $X$ , that is,

$$\begin{aligned} X^{\dagger(m)} &= X \underbrace{\dagger \dots \dagger}_m \\ &= (X \times C_{0,m}) \cup (W \times C_{d+1,d+m}) \cup \dots \cup (W \times C_{d+m,d+m}) \\ &\subset W \times A^{d+m}. \end{aligned}$$

For the final definition, we say that  $X$  is a *hole* in  $\Omega$  if  $\Omega \setminus X$  is tilable. Note that these definitions depend not only on  $X$ , but also on the underlying basic set  $\Omega$ . Therefore we will only use them when the underlying set is explicitly stated or clear from the context.

**Proposition 12.** *Let  $d \geq 1$  and let  $X$  be a hole in  $A^d$ . Suppose that  $C_{i,d} \subset X$  for some  $0 \leq i \leq d$ . Then  $(X \setminus C_{i,d})^\dagger$  is a hole in  $A^{d+1}$ .*

*Proof (see Figure 17).* Partition  $A^{d+1} \setminus (X \setminus C_{i,d})^\dagger$  into four sets




Figure 16: An illustration of the definition of  $X^\dagger$ , building on Figure 13. The diagram on the left is four-dimensional and represents a generic set  $X \subset W \times A^d$ . The diagram on the right is five-dimensional and represents the corresponding  $X^\dagger$ . We stress that this is an abstract illustration. In particular, here  $|A| = 5$  and  $|W| = 3$ , while in fact we always have either  $|W| = 1$  or  $|W| = |G| \geq |A|$ .

- (i)  $C_{i,d} \times (A \setminus C^{\text{up}})$  — tilable, because  $A \setminus C^{\text{up}} = T$ ;
- (ii)  $(A^d \setminus X) \times C_{\text{down}}$  — tilable, because  $A^d \setminus X$  is tilable;
- (iii)  $(A^d \setminus C_{i,d}) \times (A \setminus (C^{\text{up}} \cup C_{\text{down}}))$  — tilable by Proposition 11;
- (iv)  $(A^d \setminus C_{0,d}) \times C^{\text{up}}$  — tilable by Proposition 11. □

This proposition is the most useful for us in the form of the following corollary.

**Corollary 13.** *Let  $d \geq 1$  and suppose that  $0 \leq i_1, \dots, i_m \leq d$  are distinct integers. Then*

$$(A^d \setminus (C_{i_1,d} \cup \dots \cup C_{i_m,d}))^{\dagger(m)}$$

*is a hole in  $A^{d+m}$ .*

*Proof.* Use induction on  $m$ . The base case  $m = 1$  is a special case of Proposition 12, so suppose that  $m \geq 2$ . Note that

$$\begin{aligned} & (A^d \setminus (C_{i_1,d} \cup \dots \cup C_{i_m,d}))^{\dagger(m)} \\ &= \left( (A^d \setminus (C_{i_1,d} \cup \dots \cup C_{i_{m-1},d}))^{\dagger(m-1)} \setminus C_{i_m,d+m-1} \right)^{\dagger} \end{aligned}$$

so it is a hole in  $A^{d+m-1}$  by the induction hypothesis and Proposition 12. □

Now we have the tools needed for the proof of Lemma 10.

*Proof of Lemma 10.* Fix any  $d \geq (1 + |G|/|T|)d_0$  and write  $\mathcal{F} = \{C_{i,d} : i = 1, \dots, d\}$ . By symmetry, it is enough to find a tiling for the set

$$M_m = G \times (A^d \setminus (C_{d-m+1,d} \cup \dots \cup C_{d,d}))$$

for every choice of  $m \leq d_0$  with  $m \equiv 1 \pmod{|T|}$ . Fix one such value of  $m$ , and let  $M = M_m$  be the corresponding set that we have to tile.

Define  $r = d - m$  and  $\Omega = G \times A^r$ . We will construct a partition  $\mathcal{B}$  of the set  $\Omega$ , satisfying:




Figure 17: An illustration of the proof of Proposition 12. The three-dimensional diagram on the left represents a hole  $X \subset A^d$  which contains  $C_{i,d}$ . The four-dimensional diagram on the right represents  $(X \setminus C_{i,d})^\dagger$  and demonstrates why it is a hole in  $A^{d+1}$ .

- $\mathcal{B}$  consists of the set  $Y_0 = G \times C_{0,r}$  and copies of the tile  $T$ ;
- for each  $1 \leq i \leq r$ , there is some  $y_i \in G$  such that the set  $Y_i = (T + y_i) \times C_{i,r}$  is exactly the union of some copies of  $T$  in  $\mathcal{B}$ ;
- each  $y \in G$  appears at least  $t = (m - 1)/|T|$  times in the list  $y_1, \dots, y_r$ .




Figure 18: By constructing the partition  $\mathcal{B}$  we show that the set  $\bigcup_{i=0}^r Y_i$  (grey in this diagram) is a hole in  $\Omega = G \times A^r$ . In fact,  $\bigcup_{i \in I \cup \{0\}} Y_i$  is a hole for any  $I \subset [r]$ .

We start the construction by fixing any list  $y_1, \dots, y_r$  such that each member of  $G$  appears exactly  $t$  times in  $y_1, \dots, y_{t|G|}$  (in particular, this list satisfies the final condition displayed above). Note that such a list exists since  $r \geq t|G|$ .



Now we use induction to construct, for each  $0 \leq j \leq r$ , a partition  $\mathcal{B}_j$  of  $G \times A^j$  such that the first two conditions are satisfied when  $\mathcal{B}$  and  $r$  are replaced by  $\mathcal{B}_j$  and  $j$ .

Let  $\mathcal{B}_0 = \{G\}$ . Having defined  $\mathcal{B}_{j-1}$ , let  $\mathcal{B}_j$  consist of the following sets (see Figure 19):

- (i)  $G \times C_{0,j}$ ,
- (ii)  $X \times \{b\}$  for each  $X \in \mathcal{B}_{j-1}$  that is a copy of  $T$  and each  $b \in C_{\text{down}}$ ,
- (iii)  $\{g\} \times \{a\} \times T^{\text{up}}$  for each  $g \in G \setminus (T + y_j)$  and each  $a \in A^{j-1}$ ,
- (iv)  $(T + y_j) \times \{a\} \times \{b\}$  for each  $a \in A^{j-1}$  and each  $b \in T^{\text{up}} = A \setminus C_{\text{down}}$ .

One can easily check that  $\mathcal{B}_j$  is a partition of  $G \times A^j$  with the required properties. In particular, the sets of the first two types cover  $G \times A^{j-1} \times C_{\text{down}}$ , and the remaining sets cover  $G \times A^{j-1} \times (A \setminus C_{\text{down}})$ .

This concludes the construction of  $\mathcal{B}$ .




Figure 19: The induction step in the construction of the partition  $\mathcal{B}$ .

Define (recalling that  $Y_0 = G \times C_{0,r}$  and  $Y_i = (T + y_i) \times C_{i,r}$  for  $1 \leq i \leq t|G|$ )

$$S = \Omega \setminus \left( \bigcup_{i=0}^{t|G|} Y_i \right).$$

The point is that  $S$  is tilable by the restriction of  $\mathcal{B}$ , and hence  $S \times C_{0,m}$  is also tilable. Therefore it only remains to prove that  $M \setminus (S \times C_{0,m})$  is tilable, because this would imply that  $M$  is tilable. Observe that  $M \setminus (S \times C_{0,m}) = (G \times A^d) \setminus S^{\dagger(m)}$ , so it remains to prove that  $S^{\dagger(m)}$  is a hole.

To prove this, fix any  $g \in G$  and write  $\Omega_g = \{g\} \times A^r$ . Then  $\Omega_g$  intersects  $Y_0$  and exactly  $t|T| = m - 1$  of the  $Y_1, \dots, Y_{t|G|}$ . In other words,

$$\Omega_g \cap S = \{g\} \times \left( A^r \setminus \bigcup_{k=1}^m C_{j_k, r} \right)$$

for some  $0 = j_1 < j_2 < \dots < j_m \leq r$ . By Corollary 13,  $(\Omega_g \cap S)^{\dagger(m)}$  is a hole in  $\{g\} \times A^d$ . This holds for any  $g \in G$ , so in fact  $S^{\dagger(m)} = \bigcup_{g \in G} (\Omega_g \cap S)^{\dagger(m)}$  is a hole in  $G \times A^d$ , completing the proof.  $\square$

### 3.3 Using one special dimension to cover certain subsets in slices

In this section we show how one special dimension can be used to lay foundations for a tiling so that the tiling can be completed in each slice separately using Lemma 10.

Here is the main result of this section. Its statement and proof are very similar to Lemma 8 from Section 2.

**Lemma 14.** *Let  $t, b \geq 1$  be integers and  $T$  a finite tile in  $\mathbb{Z}^b$ . Further, let  $S$  be a set and let  $\mathcal{F}$  be a family consisting of at least  $(t-1)|T|^2$  pairwise disjoint subsets of  $S$ . Then there is a set  $X \subset \mathbb{Z}^b \times S$ , satisfying:*

- $X$  is a union of disjoint sets of the form  $(T+x) \times A$  with  $x \in \mathbb{Z}^b$  and  $A \in \mathcal{F}$ , and
- for each  $x \in \mathbb{Z}^b$  there is some  $m \equiv 1 \pmod{t}$  such that  $\{y \in S : (x, y) \in X\}$  is a union of  $m$  distinct members of  $\mathcal{F}$ .

We will deduce Lemma 14 from the following simple deconvolution type statement.

**Proposition 15.** *Let  $t \geq 1$  and  $b \geq 0$  be integers,  $T$  a finite tile in  $\mathbb{Z}^b$ , and  $f : \mathbb{Z}^b \rightarrow \mathbb{Z}$  a function. Then there is a function  $g : \mathbb{Z}^b \rightarrow \{0, \dots, t-1\}$  such that for each  $x \in \mathbb{Z}^b$*

$$\sum_{y \in T} g(x-y) \equiv f(x) \pmod{t}.$$

*Proof.* Use induction on  $b$ . The base case  $b = 0$  is trivial ( $\mathbb{Z}^0$  being the trivial group), so suppose that  $b \geq 1$ . For any  $n \in \mathbb{Z}$ , write

$$T_n = \{x \in \mathbb{Z}^{b-1} : (x, n) \in T\}.$$

Without loss of generality, assume that  $T_0 \neq \emptyset$  and  $T_n = \emptyset$  for all  $n < 0$ . Write  $k$  for the greatest integer such that  $T_k \neq \emptyset$ . In other words,  $[0, k]$  is the minimal interval containing the projection of  $T$  in the last coordinate.

Set  $g(x, n) = 0$  for all  $x \in \mathbb{Z}^{b-1}$  and all  $n \in \mathbb{Z}$  such that  $-k \leq n \leq -1$ . The next step is to define  $g(x, n)$  for all  $n \geq 0$  and  $x \in \mathbb{Z}^{b-1}$ . Consider  $N = 0, 1, \dots$  in turn, at each step having defined  $g(x, n)$  whenever  $-k \leq n \leq N-1$  and  $x \in \mathbb{Z}^{b-1}$ . By the induction hypothesis, we can define  $g(x, N)$  so that for all  $x \in \mathbb{Z}^{b-1}$

$$\sum_{y \in T_0} g(x-y, N) \equiv f(x, N) - \sum_{\substack{1 \leq j \leq k \\ z \in T_j}} g(x-z, N-j) \pmod{t}.$$

The final step is to define  $g(x, n)$  when  $n \leq -k-1$ . The argument is similar. Consider  $N = -k-1, -k-2, \dots$  in turn, at each step having defined  $f(x, n)$  whenever  $n \geq N+1$ . By the induction hypothesis, we can define  $g(x, N)$  so that for each  $x \in \mathbb{Z}^{b-1}$

$$\sum_{y \in T_k} g(x-y, N) \equiv f(x, N+k) - \sum_{\substack{0 \leq j \leq k-1 \\ z \in T_j}} g(x-z, N+k-j) \pmod{t}.$$

These three steps together define  $g$  completely, and it is easy to see that  $g$  satisfies the required condition.  $\square$

We get Lemma 14 as a quick corollary. In its proof we write  $\mathbb{N}$  for  $\{1, 2, \dots\}$ .

*Proof of Lemma 14.* Let  $g: \mathbb{Z}^b \rightarrow \{0, \dots, t-1\}$  be such that for each  $x \in \mathbb{Z}^b$

$$\sum_{y \in T} g(x-y) \equiv 1 \pmod{t}.$$

Let  $z_1, z_2, \dots$  be any enumeration of the elements of  $\mathbb{Z}^b$ . We will define sets  $F_1, F_2, \dots \subset \mathcal{F}$  such that  $|F_n| = g(z_n)$  for any  $n \in \mathbb{N}$ , and  $F_m \cap F_n = \emptyset$  for any distinct  $m, n \in \mathbb{N}$  with  $(T + z_m) \cap (T + z_n) \neq \emptyset$ . Then we will be done by taking

$$X = \bigcup_{\substack{n \in \mathbb{N} \\ A \in F_n}} [(T + z_n) \times A].$$

The  $F_n$  can be defined inductively. Indeed, suppose that for some  $N \in \mathbb{N}$  the sets  $F_1, \dots, F_{N-1}$  are already defined. Then we can define  $F_N$  to consist of exactly  $g(z_N)$  elements of  $\mathcal{F}$  that are not contained in  $F_n$  for any  $n \leq N-1$  with  $(T + z_n) \cap (T + z_N) \neq \emptyset$ . This is possible, because we have at most  $|T|^2 - 1$  choices for such  $n$ , and

$$g(z_N) + (|T|^2 - 1) \max_{n \leq N-1} |F_n| \leq (t-1)|T|^2 \leq |\mathcal{F}|. \quad \square$$

### 3.4 General properties of tilings

In this section we prove some transitivity results for tilings. The underlying theme is, expressed very roughly, ‘if  $B$  is  $A$ -tilable with the help of  $k$  extra dimensions, and  $C$  is  $B$ -tilable with the help of  $\ell$  extra dimensions, then  $C$  is  $A$ -tilable with the help of  $k+\ell$  extra dimensions’.

To avoid making the notation, which is already somewhat cumbersome, even more complicated we allow ourselves to abuse it in places where this is unlikely to create ambiguity. For example, given a tiling  $X = \bigsqcup X_\alpha$  we may refer to the sets  $X_\alpha$  as tiles (technically, they are not tiles, but copies of tiles). Otherwise, the proofs in this section are fairly straightforward.

**Proposition 16.** *Let  $G, G_1, \dots, G_m$  and  $H_1, \dots, H_n$  be abelian groups with tiles  $A \subset B \subset C \subset G$ ,  $T_i \subset B_i \subset G_i$  and  $U_i \subset C_i \subset H_i$ . Suppose that*

$$B_1 \times \dots \times B_m \times B \text{ is } (T_1, \dots, T_m, A)\text{-tilable} \quad (1)$$

and that

$$C_1 \times \dots \times C_n \times C^d \text{ is } (U_1, \dots, U_n, d \cdot B)\text{-tilable}. \quad (2)$$

Then

$$B_1 \times \dots \times B_m \times C_1 \times \dots \times C_n \times C^d \text{ is } (T_1, \dots, T_m, U_1, \dots, U_n, d \cdot A)\text{-tilable}.$$

Let us unravel the statement of this proposition. Intuitively, condition (1) asserts that ‘ $B$  is almost  $A$ -tilable’ – the extra dimensions  $B_1, \dots, B_m$  are used to fill the gaps. Similarly, condition (2) asserts that ‘ $C^d$  is almost  $B$ -tilable’ – here we use the extra dimensions  $C_1, \dots, C_n$ . Finally, the conclusion states that ‘ $C^d$  is almost  $A$ -tilable’ – we use all the extra dimensions,  $B_1, \dots, B_m$  and  $C_1, \dots, C_n$ , to complete this tiling.

*Proof.* For each tile  $X$  in the  $(U_1, \dots, U_n, d \cdot B)$ -tiling of  $C_1 \times \dots \times C_n \times C^d$ , partition the set  $B_1 \times \dots \times B_m \times X$  in one of the two following ways:

- if  $X$  is a copy of  $B$ , then partition  $B_1 \times \dots \times B_m \times X$  into its  $(T_1, \dots, T_m, A)$ -tiling;
- otherwise (that is, if  $X$  is a copy of one of the  $U_1, \dots, U_n$ ), partition the set into copies of  $X$ , namely  $\{b\} \times X$  for each  $b \in B_1 \times \dots \times B_m$ .

This produces a  $(T_1, \dots, T_m, U_1, \dots, U_n, d \cdot A)$ -tiling of  $B_1 \times \dots \times B_m \times C_1 \times \dots \times C_n \times C^d$ .  $\square$

In the proof of the main theorem, we will apply this result in the following more compact form.

**Corollary 17.** *Let  $G$  and  $H$  be abelian groups with tiles  $T \subset G$  and  $A \subset B \subset H$ . Suppose that*

$$G^k \times H^\ell \times B^d \text{ is } (k \cdot T, (\ell + d) \cdot A)\text{-tilable} \quad (1)$$

and that

$$G^u \times H^v \text{ is } (u \cdot T, v \cdot B)\text{-tilable.} \quad (2)$$

Then

$$G^{du+k} \times H^{dv+\ell} \text{ is } ((du+k) \cdot T, (dv+l) \cdot A)\text{-tilable.}$$

*Proof.* Use induction on  $d$ . The base case  $d = 0$  is trivial, so suppose that  $d \geq 1$ .

Rewrite (1) to state that

$$G^k \times H^\ell \times B^{d-1} \times B \text{ is } (k \cdot T, (\ell + d - 1) \cdot A, A)\text{-tilable.}$$

Now Proposition 16 applied to this and (2) implies that

$$G^k \times H^\ell \times B^{d-1} \times G^u \times H^v \text{ is } (k \cdot T, (\ell + d - 1) \cdot A, u \cdot T, v \cdot A)\text{-tilable,}$$

which after reordering and combining terms becomes the statement that

$$G^{u+k} \times H^{v+\ell} \times B^{d-1} \text{ is } ((u+k) \cdot T, (v+l+d-1) \cdot A)\text{-tilable.}$$

Finally, apply the induction hypothesis to this and (2) to conclude the proof.  $\square$

The following straightforward proposition allows tilings to be lifted via surjective homomorphisms.

**Proposition 18.** *Let  $G, H$  and  $G_1, \dots, G_n$  be abelian groups with tiles  $T \subset G$  and  $U_i \subset G_i$ , and let  $\rho: G \rightarrow H$  be a surjective homomorphism that is injective on  $T$ . If  $G_1 \times \dots \times G_n \times H$  is  $(U_1, \dots, U_n, \rho(T))$ -tilable, then  $G_1 \times \dots \times G_n \times G$  is  $(U_1, \dots, U_n, T)$ -tilable.*

*Proof.* For any tile  $X$  in the  $(U_1, \dots, U_n, \rho(T))$ -tiling of  $G_1 \times \dots \times G_n \times H$ , let  $\hat{X}$  denote the set

$$\hat{X} = \{(x_1, \dots, x_n, x) \in G_1 \times \dots \times G_n \times G : (x_1, \dots, x_n, \rho(x)) \in X\}.$$

For every  $X$ , partition  $\hat{X}$  in one of the two following ways:

- if  $X$  is a copy of  $U_i$  for some  $1 \leq i \leq n$ , then partition  $\hat{X}$  into copies of  $U_i$  in the obvious way;

- if  $X$  is a copy of  $\rho(T)$ , then  $X = \{(x_1, \dots, x_n)\} \times \rho(T+x)$  for some  $x_i \in G_i$  and  $x \in G$ . Hence  $\hat{X} = \{(x_1, \dots, x_n)\} \times (T+x+\ker(\rho))$ , and as  $\rho$  is injective on  $T$ , this can be partitioned into copies of  $T$ .

Since the sets  $\hat{X}$  partition  $G_1 \times \dots \times G_n \times G$ , this produces a  $(U_1, \dots, U_n, T)$ -tiling for it.  $\square$

An inductive application of this proposition gives the following result, which we will use in the proof of the main theorem.

**Corollary 19.** *Let  $G$  and  $H$  be abelian groups, and let  $T \subset G$  be a tile. Moreover, suppose that a surjective homomorphism  $\rho: G \rightarrow H$  is injective on  $T$ . If  $G^k \times H^\ell$  is  $(k \cdot T, \ell \cdot \rho(T))$ -tilable, then  $G^{k+\ell}$  is  $T$ -tilable.  $\square$*

### 3.5 Proof of the main theorem

The tools needed for the proof Theorem 2 are now available.

**Theorem 2.** *Let  $T \subset \mathbb{Z}^n$  be a tile. Then  $T$  tiles  $\mathbb{Z}^d$  for some  $d$ .*

*Proof.* Without loss of generality assume that  $T \subset [k]^b$ , where  $k \in \mathbb{N}$ . Write  $G = \mathbb{Z}_k^b$  and let  $\pi: \mathbb{Z}^b \rightarrow G$  be the projection map. In particular,  $\pi(T)$  is a tile in  $G$ .

**Claim.** *Suppose that  $A \subset G$  is a tile. Then there exist integers  $p \geq 0$  and  $q \geq 1$  such that  $(\mathbb{Z}^b)^p \times G^q$  is  $(p \cdot T, q \cdot A)$ -tilable.*

*Proof of claim.* Use reverse induction on  $|A|$ . If  $|A| = |G|$  then in fact  $A = G$ , and the claim holds with  $p = 0, q = 1$ . So suppose that  $|A| \leq |G| - 1$ .

Applying Lemma 10 to the tile  $A$  with fixed large  $d_0$  produces a number  $d_1 \geq d_0$ , a set  $B$  such that  $A \not\subset B \subset G$  and a family  $\mathcal{F}$  ( $|\mathcal{F}| \geq d_0$ ) of pairwise disjoint subsets of  $B^{d_1}$  with the property that for any subfamily  $\mathcal{S} \subset \mathcal{F}$  of size satisfying  $|\mathcal{S}| \equiv 1 \pmod{|A|}$ , the set

$$G \times \left( B^{d_1} \setminus \bigcup_{S \in \mathcal{S}} S \right)$$

is  $A$ -tilable.

Since  $d_0$  is large, Lemma 14 gives a set  $X \subset \mathbb{Z}^b \times G \times B^{d_1}$  that is a disjoint union of copies of  $T$ , and such that for every  $x \in \mathbb{Z}^b$  the slice  $\{y \in G \times B^{d_1} : (x, y) \in X\}$  is a hole in  $G \times B^{d_1}$ . Therefore  $\mathbb{Z}^b \times G \times B^{d_1}$  is  $(T, (d_1 + 1) \cdot A)$ -tilable.

By the induction hypothesis, there exist  $u \geq 0$  and  $v \geq 1$  such that  $(\mathbb{Z}^b)^u \times G^v$  is  $(u \cdot T, v \cdot B)$ -tilable. Now apply Corollary 17 to conclude that the claim holds with  $p = d_1 u + 1$  and  $q = d_1 v + 1$ . This proves the claim.  $\square$

To finish the proof of the theorem, apply the claim to the tile  $\pi(T)$ . This gives  $p \geq 0$  and  $q \geq 1$  such that  $(\mathbb{Z}^b)^p \times G^q$  is  $(p \cdot T, q \cdot \pi(T))$ -tilable. Hence, by Corollary 19,  $(\mathbb{Z}^b)^{p+q}$  is  $T$ -tilable.  $\square$

## 4 Concluding remarks and open problems

We mention in passing that all our tilings are (or can be made to be) periodic. Also, our copies of  $T$  arise only from translations and permutations of the coordinates – in particular, ‘positive directions stay positive’.

We have made no attempt to optimise the dimension  $d$  in Theorem 2. What can be read out of the proof is the following.

**Theorem 2’.** *Let  $T \subset \mathbb{Z}^n$  be a tile and suppose that  $T \subset [k]^n$ . Then  $T$  tiles  $\mathbb{Z}^d$ , where  $d = \lceil \exp(100(n \log k)^2) \rceil$ .*

Thus our upper bound on  $d$  is superpolynomial in the variable  $k^n$ . We believe that there should be an upper bound on  $d$  in terms only of the size and dimension of  $T$ . Even in the case  $n = 1$  this seems to be a highly non-trivial question.

**Conjecture 20.** *For any positive integer  $t$  there is a number  $d$  such that any tile  $T \subset \mathbb{Z}^d$  with  $|T| \leq t$  tiles  $\mathbb{Z}^d$ .*

On the other hand, it is easy to see that there cannot be a bound just in terms of the dimension of the tile. Indeed, given any  $d$  it is possible to find a one-dimensional tile that does not tile  $\mathbb{Z}^d$ . Such a tile  $T$  can be constructed by fixing an integer  $k$  and taking two intervals of length  $k$ , distance  $k^2 - 1$  apart, where in between the intervals only every  $k$ -th point is present in the tile. For example, if  $k = 4$  then the resulting tile would be

XXXX...X...X...X...XXXX

Suppose that  $T$  tiles  $\mathbb{Z}^d$ . Choose a large integer  $N$  and consider the cuboid  $[N]^d$ . Fix one of the  $d$  directions and only consider the copies of  $T$  in this direction that intersect  $[N]^d$ . There can be at most  $N^{d-1}(N/(k^2 + 2k - 1) + 2) = O(N^d/k^2)$  such tiles and they cover at most  $O(N^d/k)$  elements of  $[N]^d$ . Since there are  $d$  directions, at most  $O(dN^d/k)$  elements of the cuboid can be covered with tiles, but this number is less than  $N^d$  for large  $k$ . Therefore, if  $k$  is large enough, then  $T$  does not tile  $\mathbb{Z}^d$ .

Finally, we do not know how to find reasonable lower bounds, even for seemingly simple tiles. In particular, we do not know the smallest dimension that can be tiled by a given interval with the central point removed. The largest tile of this shape for which we know the answer is XXX.XXX: it does not tile  $\mathbb{Z}^2$  (by case analysis), but it tiles  $\mathbb{Z}^3$ . (One way to achieve this is to adapt the argument of Section 2, to make it work in the case of this tile for  $d = 3$ .)

**Question 21.** *Let  $T$  be the tile  $\underbrace{\text{XXXXX}}_k . \underbrace{\text{XXXXX}}_k$ . What is the least  $d$  for which  $T$  tiles  $\mathbb{Z}^d$ ?*

Just getting rough bounds on  $d$  would be very interesting. We do not even know if  $d \rightarrow \infty$  as  $k \rightarrow \infty$ .

## References

- [1] R. Berger, *The undecidability of the domino problem*, *Memoirs of the American Mathematical Society* **66** (1966), 1–72.

- [2] J.H. Conway and J.C. Lagarias, *Tiling with polyominoes and combinatorial group theory*, Journal of Combinatorial Theory, Series A **53** (1990), no. 2, 183–208.
- [3] S.W. Golomb, *Tiling with polyominoes*, Journal of Combinatorial Theory **1** (1966), no. 2, 280–296.
- [4] S.W. Golomb, *Tiling with sets of polyominoes*, Journal of Combinatorial Theory **9** (1970), no. 1, 60–71.
- [5] S.W. Golomb, *Polyominoes which tile rectangles*, Journal of Combinatorial Theory, Series A **51** (1989), no. 1, 117–124.
- [6] B. Grünbaum and G.C. Shephard, *Tilings and patterns*, Dover Books on Mathematics Series, Dover Publications, Incorporated, 2013.
- [7] D.A. Klarner, *Packing a rectangle with congruent  $n$ -ominoes*, Journal of Combinatorial Theory **7** (1969), no. 2, 107–115.
- [8] The Math Forum, *Two tiling problems*, <http://mathforum.org/kb/message.jspa?messageID=6223965> (retrieved on 15 April 2015).
- [9] MathOverflow, *Does every polyomino tile  $\mathbb{R}^n$  for some  $n$ ?*, <http://mathoverflow.net/questions/49915/does-every-polyomino-tile-rn-for-some-n> (retrieved on 15 April 2015).
- [10] H.A.G. Wijshoff and J. van Leeuwen, *Arbitrary versus periodic storage schemes and tessellations of the plane using one type of polyomino*, Information and Control **62** (1984), no. 1, 1–25.