FUNDAMENTAL LIMITS ON THE ACCURACY OF DEMOGRAPHIC INFERENCE BASED ON THE SAMPLE FREQUENCY SPECTRUM

By Jonathan Ternorst^{*} and Yun S. Song[†]

University of California, Berkeley

The sample frequency spectrum (SFS) of DNA sequences from a collection of individuals is a summary statistic which is commonly used for parametric inference in population genetics. Despite the popularity of SFS-based inference methods, currently little is known about the information-theoretic limit on the estimation accuracy as a function of sample size. Here, we show that using the SFS to estimate the size history of a population has a minimax error of at least $O(1/\log s)$, where s is the number of independent segregating sites used in the analysis. This rate is exponentially worse than known convergence rates for many classical estimation problems in statistics. Another surprising aspect of our theoretical bound is that it does not depend on the dimension of the SFS, which is related to the number of sampled individuals. This means that, for a fixed number sof segregating sites considered, using more individuals does not help to reduce the minimax error bound. Our result pertains to populations that have experienced a bottleneck, and we argue that it can be expected to apply to many populations in nature.

1. Introduction. The past decade has seen a revolution in our ability to interrogate the genome at the molecular level. Fueled by technological advances in DNA sequencing, studies now routinely query thousands or tens of thousands of individuals [1, 19, 28, 31, 32] in order to better understand disease susceptibility, heritability, population history, and other phenomena. In most cases, the conclusions of these studies come in the form of statistical estimates obtained from models that relate the effect of interest to mutation patterns arising in sampled DNA sequences. As genetic sample sizes explode, it is natural to wonder how additional data improve the quality of these estimates. While this general question has received intense focus in theoretical statistics, certain aspects of the genetics setting (for example, non-Gaussianity and lack of independence among samples) complicate efforts to study such models using classical techniques. New methods are needed to theoretically characterize some common models in statistical genetics.

^{*}Supported in part by a Citadel Fellowship.

[†]Supported in part by an NIH grant R01-GM109454, a Packard Fellowship for Science and Engineering, and a Miller Research Professorship.

Here, we address this need for a specific estimation problem in population genetics known as *demographic inference*. As we explain in further detail below, the aim of this problem is to reconstruct the sequence of historical events—including population size changes, migration, and admixture—that gave rise to present-day populations, using DNA samples obtained from those populations. We focus on the simplest problem of estimating the size history of a single population backwards in time.

A summary statistic known as the sample frequency spectrum (SFS; defined below) is often employed in empirical studies [3, 5–8, 10, 19, 20], but there have been fewer attempts to understand SFS-based estimation from a theoretical perspective. The main result of this paper is to show that, for a common class of estimators which analyze the SFS, there is a fundamental limit on their accuracy as a function of the sample size. More precisely, we show that, under a standard statistical error metric known as minimax error, the rate at which these estimators converge to the truth for certain populations is at best inversely logarithmic in the number of independent segregating sites analyzed, and does not depend at all on the number of individuals sampled. Compared to other types of statistical estimation problems (for example, linear regression), this is an extremely slow rate of convergence. Our proof is information-theoretic in nature and applies to any estimator that operates solely on the SFS. This is the first result we are aware of that characterizes the convergence rate of demographic history estimates as a function of sample size.

The remainder of this paper is organized as follows. In Section 2 we formally define our notation and model. In Section 3 we state our main theoretical results, followed by a discussion of their practical implications in Section 4. To streamline our exposition, all mathematical proofs are deferred until Section 5.

2. Preliminaries. The stochastic process underlying the inference procedure we consider is Kingman's coalescent [13–15], which evolves backward in time and describes the genealogy of a collection of chromosomes randomly sampled from a population. The population size is assumed to change deterministically over time and is described by a function $\eta : [0, \infty) \to (0, \infty)$, with $\eta(t)$ being the population size at time t in the past. The instantaneous rate of coalescence between any pair of lineages at time t is $1/\eta(t)$.

As in the standard infinite-sites model of mutation [12], we assume that every dimorphic site (i.e., a site with exactly two observed allelic types) has experienced mutation exactly once in the evolutionary history relating the sample. Further, for each such site, we assume that it is known which allele is the ancestral type versus the mutant type. In what follows, we use the terms dimorphic and segregating interchangeably.

A population size function $\eta(t)$ induces a probability distribution on the number of derived alleles found at a particular segregating site. Specifically, for a sample of $n \ge 2$ randomly sampled individuals, let $\xi_{n,b}^{(\eta)}$, for $1 \le b \le n-1$, denote the probability that a segregating site contains b mutant alleles in a sample of n individuals under model η . The vector $\boldsymbol{\xi}_n^{(\eta)} \stackrel{\text{def}}{=} (\xi_{n,1}^{(\eta)}, \dots, \xi_{n,n-1}^{(\eta)})$ is called the expected SFS. In the coalescent setting, a general expression for $\boldsymbol{\xi}_{n,b}^{(\eta)}$ is given by [9]

$$\xi_{n,b}^{(\eta)} \propto \sum_{k=2}^{n-b+1} \frac{\binom{n-b-1}{k-2}}{\binom{n-1}{k-1}} \cdot k \cdot \mathbb{E}T_{n,k}^{(\eta)},$$

where $\mathbb{E}T_{n,k}^{(\eta)}$ denotes the amount of time (in coalescent units) during which the genealogy of the sample contained k lineages under model η . The expected waiting time $\mathbb{E}T_{m,m}^{(\eta)}$ to the first coalescence in a sample of m individuals is given by

(1)
$$c_m^{(\eta)} \stackrel{\text{def}}{=} \mathbb{E}T_{m,m}^{(\eta)} = \int_0^\infty t \frac{a_m}{\eta(t)} \exp\left\{-a_m R_\eta(t)\right\} \, \mathrm{d}t,$$

where $a_m \stackrel{\text{def}}{=} \binom{m}{2}$ and $R_\eta(t) \stackrel{\text{def}}{=} \int_0^t \frac{1}{\eta(s)} \mathrm{d}s$ is the cumulative rate of coalescence up to time t. It turns out [22] that there is an invertible linear transformation which relates $(\mathbb{E}T_{n,2}^{(\eta)}, \mathbb{E}T_{n,3}^{(\eta)}, \dots, \mathbb{E}T_{n,n}^{(\eta)})$ to $\mathbf{c}^{(\eta)} \stackrel{\text{def}}{=} (c_2^{(\eta)}, c_3^{(\eta)}, \dots, c_n^{(\eta)})$. Using this relation, the quantity $\xi_{n,b}^{(\eta)}$ can be written as [23]

(2)
$$\xi_{n,b}^{(\eta)} = \frac{\left\langle \mathbf{c}^{(\eta)}, \mathbf{W}_{n,b} \right\rangle}{\left\langle \mathbf{c}^{(\eta)}, \mathbf{V}_{n} \right\rangle},$$

where $\mathbf{W}_{n,b} = (W_{n,b,2}, \ldots, W_{n,b,n})$ and $\mathbf{V}_n = (V_{n,2}, \ldots, V_{n,n})$ are vectors of universal constants that do not depend on the population size function η , and $\langle \cdot, \cdot \rangle$ denotes the l_2 -inner product. Under model η , the quantity $\langle \mathbf{c}^{(\eta)}, \mathbf{W}_{n,b} \rangle$ is the total expected length of edges subtending b out of n individuals sampled at time 0, while the quantity $\langle \mathbf{c}^{(\eta)}, \mathbf{V}_n \rangle$ is the total expected tree length for a sample of size n. Both quantities are positive for all population size functions η . For an arbitrary population size function η , we have $\sum_{b=1}^{n-1} W_{n,b,m} = V_{n,m}$ for all $2 \leq m \leq n$, which implies

(3)
$$\sum_{b=1}^{n-1} \left\langle \mathbf{c}^{(\eta)}, \mathbf{W}_{n,b} \right\rangle = \left\langle \mathbf{c}^{(\eta)}, \mathbf{V}_n \right\rangle.$$

For a constant function $\eta(t) \equiv N$,

$$c_m^{(\eta)}=rac{N}{a_m}, \ \langle {f c}^{(\eta)}, {f W}_{n,b}
angle=rac{2}{b}N,$$

(5)
$$\langle \mathbf{c}^{(\eta)}, \mathbf{V}_n \rangle = 2NH_{n-1},$$

where $H_{n-1} \stackrel{\text{def}}{=} \sum_{b=1}^{n-1} \frac{1}{b}$.

To formulate the problem, we employ the following notation. We suppose that a sample of $n \geq 2$ randomly sampled individuals has been typed at *s* independent segregating sites. These data are used to form the *empirical* sample frequency spectrum, which is an (n-1)-tuple $(\hat{\xi}_{n,1}, \ldots, \hat{\xi}_{n,n-1})$, where $\hat{\xi}_{n,b}$ denotes the proportion of segregating sites with *b* copies of the mutant allele and n-b copies of the ancestral allele. A *frequency-based estimator* is any statistic $\hat{\eta}$ which maps an empirical SFS to a population size history.

3. Main Results. Here, we establish a minimax lower bound on the ability of *any* estimator $\hat{\eta}$ to accurately reconstruct population size functions.

3.1. A general bound on the KL divergence between two SFS distributions. Abusing notation, we use $D(\eta || \eta')$ to denote the Kullback-Leibler (KL) divergence between the probability distributions $\boldsymbol{\xi}_n^{(\eta)}$ and $\boldsymbol{\xi}_n^{(\eta')}$. In Section 5, we prove the following general upper bound on the KL divergence between two SFS distributions:

Theorem 1. Let \mathcal{M} denote a general space of population size functions and suppose $\eta, \eta' \in \mathcal{M}$ satisfy $\eta(t) = \eta'(t)$ for all $0 \leq t \leq t_c$ and $\max_{t>t_c} \eta(t) \leq \min_{t>t_c} \eta'(t)$. Then,

(6)
$$D(\eta \| \eta') \leq \frac{\left\langle \mathbf{c}^{(\eta')} - \mathbf{c}^{(\eta)}, \mathbf{V}_n \right\rangle}{\left\langle \mathbf{c}^{(\eta)}, \mathbf{V}_n \right\rangle}.$$

3.2. Bounds for a family of piecewise-constant models. We now focus on a particular class of population size functions which are easier to analyze and popular in the literature [2,3,16]. For a fixed positive integer K > 1, let $\mathcal{M}_K \subset \mathcal{M}$ denote the space of piecewise-constant size functions with exactly

(4)

K pieces. A population size function η is a member of \mathcal{M}_K if and only if there exist positive real numbers $t_1 < \cdots < t_{K-1}$ and N_1, N_2, \ldots, N_K such that

(7)
$$\eta(t) = \sum_{k=1}^{K} N_k \mathbf{1}\{t_{k-1} \le t < t_k\},$$

where by convention we define $t_0 = 0$ and $t_K = \infty$. For such an η , define

(8)
$$S_k^{(\eta)} \stackrel{\text{def}}{=} \sum_{j=1}^k \frac{t_j - t_{j-1}}{N_j}$$

For $\eta \in \mathcal{M}_K$, the expected waiting time $c_m^{(\eta)}$ defined in (1) is given by

(9)
$$c_m^{(\eta)} = \frac{1}{a_m} \sum_{k=1}^K N_k (e^{-a_m S_{k-1}^{(\eta)}} - e^{-a_m S_k^{(\eta)}}).$$

Note that since $t_K = \infty$,

(10)
$$e^{-a_m S_K^{(\eta)}} \equiv 0$$
, for all $\eta \in \mathcal{M}_K$.

To formulate our result, we let I, J denote positive integers that satisfy I + J = K, and introduce a subfamily $\mathcal{F}_{I,J} \subset \mathcal{M}_K$ of piecewise-constant functions defined as follows. See Figure 1 for illustration. We assume that all change points $t_1 < \cdots < t_{I+J-1}$ are fixed and that the sizes N_1, \ldots, N_I of the first I epochs are also fixed, with N_I being the smallest size. So, all functions in $\mathcal{F}_{I,J}$ are identical to each other for the first I epochs, and there is a population bottleneck in the last epoch. Then, for $t \geq t_I$, every function $\eta \in \mathcal{F}_{I,J}$ undergoes jumps according to the following rules:

- 1. For the interval $t_I \leq t < t_{I+1}$, $\eta(t)$ takes a constant value of either h or $h + \delta$, where $h > N_I$ and $\delta > 0$.
- 2. At later change points $\{t_{I+1}, \ldots, t_{I+J-1}\}$, η either stays the same or jumps upward by δ .

Hence, $\mathcal{F}_{I,J}$ consists of 2^J distinct piecewise-constant functions that are nondecreasing functions of t for $t \ge t_I$. Note that $\min_t \eta(t) = N_I$ for all $\eta \in \mathcal{F}_{I,J}$. For ease of notation, we use $\varepsilon \stackrel{\text{def}}{=} N_I$ to denote the bottleneck size and $\tau_B \stackrel{\text{def}}{=} t_I - t_{I-1}$ to denote the bottleneck duration. To facilitate analysis later, we fix $t_{I+j} - t_{I+j-1}$ to some positive constant τ_A for all $j = 1, \ldots, J - 1$.

For any two models in $\mathcal{F}_{I,J}$, we obtain the following bound on the difference of their waiting times to the first coalescence:



FIG 1. A family $\mathcal{F}_{I,J}$ of piecewise-constant population size models with K = I + J epochs.

Lemma 2. For all $\eta, \eta' \in \mathcal{F}_{I,J}$,

(11)
$$\left|c_m^{(\eta)} - c_m^{(\eta')}\right| \le J \frac{\delta}{a_m} e^{-a_m \tau_B/\varepsilon}.$$

Then, together with Theorem 1, this lemma can be used to show

Theorem 3. Let $\eta, \eta' \in \mathcal{F}_{I,J}$ that satisfy $\max_{t \geq t_I} \eta(t) \leq \min_{t \geq t_I} \eta'(t)$. Then,

(12)
$$D(\eta \| \eta') \le J \frac{\delta}{\varepsilon} e^{-\tau_B/\varepsilon}$$

Our proofs of the above results are deferred to Section 5. It is interesting that the above bound does not depend on the number n of sampled individuals.

3.3. Minimax lower bounds. Before using the above results to obtain a minimax lower bound, we first note a subtle fact. Given any population size function η , consider a function ζ which satisfies $\zeta(t) = \kappa \cdot \eta(t/\kappa)$ for all $t \in [0, \infty)$, where κ is some positive constant. Such functions are equivalent, as it turns out that $\xi_{n,b}^{(\zeta)} = \xi_{n,b}^{(\eta)}$ for all $n \geq 2$ and $1 \leq b \leq n-1$. To mod out by this equivalence, we assume that every $\eta \in \mathcal{M}$ satisfies $\eta(0) = N_{\text{fix}}$, where N_{fix} is some fixed positive constant.

Let $\|\cdot\|_*$ denote a generic norm (specific examples will be given later) and let $\mathbb{E}_{\eta}(\cdot)$ denote expectation with respect to the SFS distribution $\boldsymbol{\xi}_n^{(\eta)} =$

$$(\xi_{n,1}^{(\eta)},\ldots,\xi_{n,n-1}^{(\eta)})$$
 induced by population size function η . Then, note that

$$\inf_{\hat{\eta}} \sup_{\eta \in \mathcal{M}} \mathbb{E}_{\eta} \| \hat{\eta} - \eta \|_{*} \geq \inf_{\hat{\eta}} \sup_{\eta \in \mathcal{M}_{K}} \mathbb{E}_{\eta} \| \hat{\eta} - \eta \|_{*} \geq \inf_{\hat{\eta}} \sup_{\eta \in \mathcal{F}_{I,J}} \mathbb{E}_{\eta} \| \hat{\eta} - \eta \|_{*}$$

In what follows, we will put a lower bound on the last quantity. We first fix a sensible distance metric on \mathcal{M} . An intuitive way to measure distance between two population size functions is their L_1 distance, $\|\eta_a - \eta_b\|_1 = \int_0^\infty |\eta_a(t) - \eta_b(t)| dt$, but this is unreasonably stringent in that $\|\eta_a - \eta_b\|_1 = \infty$ if η_a and η_b do not agree infinitely far back into the past. Instead we will focus on the following truncated L_1 distance: $\|\eta_a - \eta_b\|_{1,T} \stackrel{\text{def}}{=} \int_0^T |\eta_a(t) - \eta_b(t)| dt$, which measures the discrepancy between η_a and η_b back to some fixed time T in the past.

Henceforth, let $\hat{\eta}$ be any estimator of the population size function which operates on a sample of *s* independent segregating sites obtained from a sample of *n* randomly sampled individuals. In Section 5, we prove the following main results of our paper:

Theorem 4. Consider the subfamily $\mathcal{F}_{I,J}$ of models described above, and suppose J > 8 and $T \ge t_{I+J-1} + \tau_A$. Then,

(13)
$$\inf_{\hat{\eta}} \sup_{\eta \in \mathcal{F}_{I,J}} \mathbb{E}_{\eta} \|\hat{\eta} - \eta\|_{1,T} \ge C\tau_A \frac{(J-8)^2}{J} \frac{\varepsilon}{s} e^{\tau_B/\varepsilon},$$

where C is a positive constant.

The above theorem applies to all models in $\mathcal{F}_{I,J}$. We now consider the subset $\mathcal{F}_{I,J}^M = \{\eta \in \mathcal{F}_{I,J} : ||\eta||_{\infty} < M\}$, which is the set of all models in $\mathcal{F}_{I,J}$ that are bounded by some constant M. For this family of bounded population size functions, a sharper asymptotic lower bound can be obtained as follows.

Theorem 5. Suppose J > 8 and $T \ge t_{I+J-1} + \tau_A$. Then,

(14)
$$\inf_{\hat{\eta}} \sup_{\eta \in \mathcal{F}_{I,J}^{M}} \mathbb{E}_{\eta} \| \hat{\eta} - \eta \|_{1,T} \ge C' \frac{(J-8)^{2}}{J} \frac{\tau_{B} \tau_{A}}{\log s},$$

where C' is a positive constant.

By specializing $\mathcal{F}_{I,I}^{M}$, a simplified version of Theorem 5 can be obtained:

Corollary 6. Suppose $T \ge t_{I+J-1} + \tau_A$ and let $\mathcal{F}_{I,\star}^M = \bigcup_{J\ge 1} \mathcal{F}_{I,J}^M$. Then, (15) $\inf_{\hat{\eta}} \sup_{\eta\in\mathcal{F}_{I,\star}^M} \mathbb{E}_{\eta} \|\hat{\eta}-\eta\|_{1,T} \ge C''(T-t_I)\frac{\tau_B}{\log s},$

where C'' is a positive constant.

Note that the above lower bounds do not depend on the dimension of the SFS (which is equal to n - 1). Hence, for a fixed number s of segregating sites considered, using more individuals does not help to reduce the error bounds.

4. Discussion. In this paper, we have theoretically characterized fundamental limits on the accuracy of demographic inference from data. The paper that most closely relates to the present work is by Kim *et al.* [11], who obtain lower bounds on the amount of *exact* coalescence time data necessary to distinguish between size histories in a hypothesis testing framework. Since coalescence times are never observed and must be estimated from data, these bounds place a limit on the accuracy with which a population size function can be inferred. The authors also describe an estimator which uses coalescence times (again observed without noise) to accurately recover the underlying population size function with high probability, at a rate that roughly matches the lower bound. The results presented here are complementary to those of Kim *et al.* in the following sense: access to coalescence times is equivalent to knowing the site frequency spectrum, so that their results apply to SFS-based estimators as well as other types of procedures which also incorporate e.g. linkage information. By focusing specifically on SFS-based estimators, we are able to more precisely quantify the effects of sample size and underlying demography on the ability to accurately reconstruct a population size function. Ultimately, we derive minimax lower bounds on the error of estimating a size history using the SFS.

Another line of work centers around the *identifiability* of the parameter $\eta(t)$ using the SFS. Roughly speaking, a family of statistical models $\{P_{\theta}\}_{\theta \in \Theta}$ defined over a parameter space Θ is identifiable if for any $\theta_1, \theta_2 \in \Theta$ with $\theta_1 \neq \theta_2$, the sampling distributions induced by P_{θ_1} and P_{θ_2} are different. In our context this simply says that, for all $n, \boldsymbol{\xi}_n^{(\eta_1)} \neq \boldsymbol{\xi}_n^{(\eta_2)}$ unless $\eta_1 = \eta_2$ almost everywhere. Standard desiderata for statistical estimators (e.g., consistency or unbiasedness) are impossible without identifiability, so it is the weakest possible regularity condition one can impose on a useful family of models.

Perhaps surprisingly, it turns out that in general a population size function is not identifiable from the SFS [18]. Indeed, for any given $\eta(t)$, it has been shown that an infinite number of smooth functions F(t) exist such that $\boldsymbol{\xi}_n^{(\eta)} = \boldsymbol{\xi}_n^{(\eta+F)}$. Moreover, explicit examples can be constructed which demonstrate this phenomenon [18]. On the other hand, these counter-examples consist of functions which exhibit an unbounded frequency of oscillatory behavior near the present time, which is perhaps unrealistic when modeling naturally occurring populations. More recently, it has been shown [2] that identifiability holds for many classes of population size functions employed by practitioners (including piecewise-constant, -exponential and -generalized-exponential). Furthermore, the number n of sampled individuals sufficient for identifiability can be explicitly given and is a function of the complexity of the underlying class of models being studied [2].

Identifiability asserts that, given an *infinite* amount of data (specifically, taking the number of segregating sites $s \to \infty$), the model parameter $\eta(t)$ can be uniquely recovered. In practice, s is finite and only a perturbed version of the expected frequency spectrum, say $\hat{\boldsymbol{\xi}}_n^{(\eta)}$, is observed. From a practical standpoint, it is important to understand how these perturbations ultimately affect the parameter estimate $\hat{\eta}(t)$. It is this question that forms the starting point for the present work.

We have shown that the minimax error rate for estimating the piecewiseconstant demography of a single population is at least $O(1/\log s)$, where s is the number of independent segregating sites analyzed. In contrast, the minimax error for many classical estimation problems in statistics (for example, non-parametric regression or density estimation) decays inverse-polynomially in the sample size [29]. Compared with these problems, exponentially more samples would be required to estimate a population size history function to within a similar magnitude of error.

A single population evolving under a piecewise-constant demography is a special case of many richer classes of demographic models. For example, it is a (limiting) member of the family of exponential growth models, seen by taking each exponential growth parameter to zero. In the multi-species coalescent setting [4,6], multiple population size histories must be estimated, and the error of that estimate must necessarily be lower bounded by that of estimating a single such history. Thus, our result can be expected to apply to a broader class of models than the one we have studied here.

As detailed in Section 5, the result in Theorem 5 follows from setting $\varepsilon = \tau_B / \log s$ and $\delta \propto \frac{\varepsilon}{s} \exp(\tau_B/\varepsilon)$ in the subfamily $\mathcal{F}_{I,J}^M$. The size $\tau_B / \log s$ is in coalescent units. In terms of the number of individuals, it is proportional to $g_B / \log s$, where g_B is the number of generations corresponding to duration τ_B in the coalescent limit. Intuitively, as the severity of the bottleneck increases, the population is increasingly likely to find its most recent common ancestor (MRCA) during that time; further back in time than the MRCA, no information is conveyed concerning the demographic events experienced by the population.

One might object to considering models with a bottleneck size that scales inversely with the number s of segregating sites in the data, and it is indeed possible that a better convergence rate may be achievable for populations

which are known not to contain a bottleneck. On the other hand, we note that $1/\log s$ decreases sufficiently slowly with s that our result can be expected to apply to many real-world examples. For example, for $s \approx 10^8$, which is a conservative upper bound for most organisms, $g_B/\log s \approx 0.054g_B$. This implies that for populations which have experienced roughly an order-of-magnitude increase in effective population size during their history, accurate estimation of demographic events which occurred before this expansion is difficult using SFS-based methods. Additionally, an interesting aspect of our work is that our minimax lower bounds do not depend on the number n of sampled individuals; increasing n is not enough to overcome the information barrier imposed by the presence of a bottleneck. This is intuitively plausible since, as n increases, the (n + 1)th sampled lineage becomes more likely to coalesce early on.

An interesting question which we have not attempted to analyze is whether the $O(1/\log s)$ rate is optimal, i.e. whether there exists some estimator $\hat{\eta}(t)$ which achieves the minimax lower bound established here. In practice, from equations (2), (8), and (9), it can be seen that naively maximizing the likelihood of the observed SFS with respect to $\eta(t)$ requires solving a non-convex optimization problem, so that convergence to the global maximum is not even guaranteed. Computational issues aside, finding such an estimator remains an open theoretical challenge.

In closing, we stress that our result is specific to SFS-based estimators, which analyze only independent sites. The main allure of these estimators is their mathematical tractability, rather than their realism. In fact, a rich source of additional information exists in the correlation structure found among linked sites in the genome. Methods which seek to exploit this structure by modeling the action of recombination pose greater mathematical and computational difficulties, but there has been recent progress in this area [16,21,24–27]. Our result serves to underscore the importance of pursuing ever more realistic models of genomic evolution, challenging though they may be.

5. Proofs.

PROOF OF THEOREM 1. To simplify the notation, we write $\mathbf{c} = \mathbf{c}^{(\eta)}$ and $\mathbf{c}' = \mathbf{c}^{(\eta')}$. Then, using (2) we can write

$$D(\eta \| \eta') = \sum_{b=1}^{n-1} \xi_{n,b}^{(\eta)} \log \frac{\xi_{n,b}^{(\eta)}}{\xi_{n,b}^{(\eta')}} = \sum_{b=1}^{n-1} \xi_{n,b}^{(\eta)} \left[\log \left(\frac{\langle \mathbf{c}, \mathbf{W}_{n,b} \rangle}{\langle \mathbf{c}', \mathbf{W}_{n,b} \rangle} \right) + \log \left(\frac{\langle \mathbf{c}', \mathbf{V}_{n} \rangle}{\langle \mathbf{c}, \mathbf{V}_{n} \rangle} \right) \right]$$

The assumption $\min_{t>t_c} \eta'(t) \ge \max_{t>t_c} \eta(t)$ implies that, for all times $t, t' > t_c$

 t_c , the instantaneous rate of coalescence at time t in model η is \geq the instantaneous rate of coalescence at time t' in model η' . Hence, this assumption together with $\eta(t) = \eta'(t)$ for all $0 \leq t \leq t_c$ implies $\langle \mathbf{c} - \mathbf{c}', \mathbf{W}_{n,b} \rangle \leq 0$ for all $1 \leq b \leq n-1$; equivalently, $\log \left(\frac{\langle \mathbf{c}, \mathbf{W}_{n,b} \rangle}{\langle \mathbf{c}', \mathbf{W}_{n,b} \rangle}\right) < 0$. Additionally, $\frac{\langle \mathbf{c}' - \mathbf{c}, \mathbf{V}_n \rangle}{\langle \mathbf{c}, \mathbf{V}_n \rangle} > -1$ and $\log(1 + x) \leq x$ for all $x \geq -1$. Combining these facts, we obtain

$$D(\eta \| \eta') \le \sum_{b=1}^{n-1} \xi_{n,b}^{(\eta)} \log \left(\frac{\langle \mathbf{c}', \mathbf{V}_n \rangle}{\langle \mathbf{c}, \mathbf{V}_n \rangle} \right) \le \sum_{b=1}^{n-1} \xi_{n,b}^{(\eta)} \frac{\langle \mathbf{c}' - \mathbf{c}, \mathbf{V}_n \rangle}{\langle \mathbf{c}, \mathbf{V}_n \rangle} = \frac{\langle \mathbf{c}' - \mathbf{c}, \mathbf{V}_n \rangle}{\langle \mathbf{c}, \mathbf{V}_n \rangle},$$

where we have used $\sum_{b=1}^{n-1} \xi_{n,b}^{(\eta)} = 1$ in the final equality.

PROOF OF LEMMA 2. We distinguish two particular models $\eta_{\ell}, \eta_u \in \mathcal{F}_{I,J}$ which are the *lower* and the *upper* envelopes of $\mathcal{F}_{I,J}$. The function η_{ℓ} stays constant at h for all $t \geq t_I$, while η_u jumps upward by δ at every change point t_I, \ldots, t_{I+J-1} . Hence, $\eta_{\ell} \leq \eta \leq \eta_u$ pointwise for all $\eta \in \mathcal{F}_{I,J}$. The two enveloping functions will form the basis of subsequent analysis.

Fix $\eta, \eta' \in \mathcal{F}_{I,J}$ and note that, by the definition of $\mathcal{F}_{I,J}$, one of these functions must pointwise-dominate the other. So, assume without loss of generality that $\eta(t) \leq \eta'(t)$ for all t. Then for all t,

$$\eta_{\ell}(t) \le \eta(t) \le \eta'(t) \le \eta_u(t),$$

which implies

$$c_m^{(\eta_\ell)} \le c_m^{(\eta)} \le c_m^{(\eta')} \le c_m^{(\eta_u)}$$

for all $m = 2, \ldots, n$. Using these inequalities, we conclude

$$c_m^{(\eta')} - c_m^{(\eta)} \le c_m^{(\eta_u)} - c_m^{(\eta_\ell)},$$

so it suffices to demonstrate (11) for $c_m^{(\eta_u)} - c_m^{(\eta_\ell)}$. Now, by equation (9) and the definition of η_ℓ ,

$$a_m c_m^{(\eta_\ell)} = \sum_{i=1}^I N_i \left[e^{-a_m S_{i-1}^{(\eta_\ell)}} - e^{-a_m S_i^{(\eta_\ell)}} \right] + \sum_{j=1}^J h \left[e^{-a_m S_{I+j-1}^{(\eta_\ell)}} - e^{-a_m S_{I+j}^{(\eta_\ell)}} \right]$$
$$= \sum_{i=1}^I N_i \left[e^{-a_m S_{i-1}^{(\eta_\ell)}} - e^{-a_m S_i^{(\eta_\ell)}} \right] + h e^{-a_m S_I^{(\eta_\ell)}},$$

where we have used equation (10). Similarly,

$$a_{m}c_{m}^{(\eta_{u})} = \sum_{i=1}^{I} N_{i} \left[e^{-a_{m}S_{i-1}^{(\eta_{u})}} - e^{-a_{m}S_{i}^{(\eta_{u})}} \right] + \sum_{j=1}^{J} (h+j\delta) \left[e^{-a_{m}S_{I+j-1}^{(\eta_{u})}} - e^{-a_{m}S_{I+j}^{(\eta_{u})}} \right] = \sum_{i=1}^{I} N_{i} \left[e^{-a_{m}S_{i-1}^{(\eta_{u})}} - e^{-a_{m}S_{i}^{(\eta_{u})}} \right] + he^{-a_{m}S_{I}^{(\eta_{u})}} + \sum_{j=1}^{J} j\delta \left[e^{-a_{m}S_{I+j-1}^{(\eta_{u})}} - e^{-a_{m}S_{I+j}^{(\eta_{u})}} \right].$$

Now, using the fact that η_{ℓ} and η_u agree on the first I epochs, we obtain

$$a_m [c_m^{(\eta_u)} - c_m^{(\eta_\ell)}] = \sum_{j=1}^J j\delta \left[e^{-a_m S_{I+j-1}^{(\eta_u)}} - e^{-a_m S_{I+j}^{(\eta_u)}} \right]$$
$$= \delta \sum_{j=1}^J e^{-a_m S_{I+j-1}^{(\eta_u)}}$$
$$\leq J\delta e^{-a_m \tau_B/\varepsilon},$$

where the second line follows from telescoping and the fact $S_{I+J}^{(\eta_u)} = \infty$, while the last line follows from the fact that $\frac{\tau_B}{\varepsilon} \leq S_{I+j-1}^{(\eta_u)}$ for all $j = 1, \ldots, J$. \Box

PROOF OF THEOREM 3. For ease of notation, define $\mathbf{c} = \mathbf{c}^{(\eta)}$ and $\mathbf{c}' = \mathbf{c}^{(\eta')}$. By Lemma 2,

$$\begin{split} \langle \mathbf{c}' - \mathbf{c}, \mathbf{V}_n \rangle &= \sum_{m=2}^n (c'_m - c_m) V_{n,m} \le J \delta \sum_{m=2}^n \frac{V_{n,m}}{a_m} e^{-a_m \tau_B/\varepsilon} \\ &\le J \delta e^{-\tau_B/\varepsilon} \sum_{m=2}^n \frac{V_{n,m}}{a_m}, \end{split}$$

where the second inequality follows from $e^{-a_m\tau_B/\varepsilon} \leq e^{-\tau_B/\varepsilon}$ for all $m = 2, \ldots, n$. Now, noting that $\sum_{m=2}^{n} \frac{V_{n,m}}{a_m}$ corresponds to the total tree length for the constant population size function $\eta \equiv 1$ and using (5), we obtain

(16) $\langle \mathbf{c}' - \mathbf{c}, \mathbf{V}_n \rangle \leq J \delta e^{-\tau_B/\varepsilon} 2H_{n-1}.$

To finish the proof, recall that $\langle \mathbf{c}, \mathbf{V}_n \rangle$ is the total expected branch length of the coalescent tree under model η . Since $\min_t \eta(t) = \varepsilon$, we have that $\langle \mathbf{c}, \mathbf{V}_n \rangle$ is at least as large as the corresponding quantity under a model with constant population size ε . By (5), the total expected tree length under the latter model equals $2\varepsilon H_{n-1}$. Thus, $\langle \mathbf{c}, \mathbf{V}_n \rangle \geq 2\varepsilon H_{n-1}$ and combining this result with (16) gives

$$\frac{\langle \mathbf{c}' - \mathbf{c}, \mathbf{V}_n \rangle}{\langle \mathbf{c}, \mathbf{V}_n \rangle} \le J \frac{\delta}{\varepsilon} e^{-\tau_B/\varepsilon}.$$

Finally, (12) follows from this inequality and Theorem 1.

PROOF OF THEOREM 4. Our proof uses a generalized form of Fano's inequality [30]. Adapted to our setting and notation, the method reads as follows.

Theorem 7 (Fano's method). Consider a space \mathcal{M} of population size models. Let $r \geq 2$ be an integer, and let $S_n^r = \{\eta_1, \eta_2, \ldots, \eta_r\} \subset \mathcal{M}$ contain r population size functions such that for all $a \neq b$, $\|\eta_a - \eta_b\|_* \geq \alpha_r$ and $D(\xi_n^{(\eta_a)}\|\xi_n^{(\eta_b)}) \leq \beta_r$. Let $\hat{\eta}^{(n,s)} = \hat{\eta}^{(n,s)}(X_1, \ldots, X_s)$ be an estimator of η based on the SFS data X_1, \ldots, X_s sampled independently from $\xi_n^{(\eta)}$; i.e., X_1, \ldots, X_s are SFS data for n individuals at s independent segregating sites. Then,

(17)
$$\inf_{\hat{\eta}} \sup_{\eta \in \mathcal{M}} \mathbb{E}_{\eta} \| \hat{\eta}^{(n,s)} - \eta \|_{*} \ge \frac{\alpha_{r}}{2} \left(1 - \frac{s \cdot \beta_{r} + \log 2}{\log r} \right).$$

This theorem places a lower bound on the minimax rate of convergence of a population size history estimator based on the SFS.

For $\eta \in \mathcal{F}_{I,J}$, let w_j denote the variable $\in \{0,1\}$ indicating whether η jumps by δ at change point t_{I+j} . Let $\mathcal{Y} = \{\mathbf{w} = (w_0, \ldots, w_{J-1}) \mid w_i \in \{0,1\}\}$, where $J \geq 8$. By the Varshamov-Gilbert lemma (see [17, Lemma 4.7]), there exist $\mathcal{X} = \{\mathbf{w}^0, \ldots, \mathbf{w}^M\} \subset \mathcal{Y}$ such (i) $\mathbf{w}^0 = (0, \ldots, 0)$, (ii) $M \geq 2^{J/8}$, and (iii) $H(\mathbf{w}^i, \mathbf{w}^j) \geq J/8$, where $H(\cdot, \cdot)$ denotes the Hamming distance.

Let $\mathcal{F}_{I,J}^{\mathcal{X}}$ denote the subset of $2^{J/8} + 1$ functions in $\mathcal{F}_{I,J}$ with the indicator variable for δ -jumps at t_I, \ldots, t_{I+J-1} given by $\mathbf{w} \in \mathcal{X}$. Then, for any two $\eta_a \neq \eta_b \in \mathcal{F}_{I,J}^{\mathcal{X}}$, we have

(18)
$$\|\eta_a - \eta_b\|_{1,T} \ge \frac{J}{8} \cdot \tau_A \cdot \delta.$$

Using Theorem 7 via (18) and Theorem 3, we obtain

(19)
$$\inf_{\hat{\eta}} \sup_{\eta \in \mathcal{F}_{I,J}} \mathbb{E}_{\eta} \| \hat{\eta}^{(n,s)} - \eta \|_{1,T} \geq \frac{J \cdot \tau_A \cdot \delta}{16} \left[1 - \frac{sJ\frac{\delta}{\varepsilon}e^{-\tau_B/\varepsilon} + \log 2}{\log(2^{J/8} + 1)} \right] \\ \geq \frac{J \cdot \tau_A \cdot \delta}{16} \left[1 - \frac{sJ\frac{\delta}{\varepsilon}e^{-\tau_B/\varepsilon} + \log 2}{\frac{J}{8}\log 2} \right].$$

We now optimize the bound with respect to δ . A straightforward calculation shows that the maximum is attained at

(20)
$$\delta^* = \frac{(J-8)\log 2}{16J} \left(\frac{\varepsilon}{s}\right) e^{\tau_B/\varepsilon}$$

and setting $\delta = \delta^*$ in (19) yields the result.

PROOF OF THEOREM 5. The result is obtained by scaling ε with the number of segregating sites s. Denote this scaling by $\varepsilon(s)$; we will determine $\varepsilon(s)$ which produces the largest possible lower bound. Starting from Equation (20) in the proof of Theorem 4, note that δ^* scales as $\frac{\varepsilon}{s}e^{\tau_B/\varepsilon} =: f(\varepsilon)$. In order to satisfy the constraint that $\|\eta\|_{\infty} < M$ for all $\eta \in \mathcal{F}_{I,J}^M$ and s, the condition

(21)
$$\limsup_{s \to \infty} \max\left\{\frac{\varepsilon(s)}{s} e^{\tau_B/\varepsilon(s)}, \varepsilon(s)\right\} < \infty$$

must therefore hold. This implies that $\varepsilon(s)s^p \to \infty$ as $s \to \infty$ for all p > 0. Suppose that $q \stackrel{\text{def}}{=} \liminf_{s \to \infty} \frac{\varepsilon(s)\log s}{\tau_B} < 1$; note that $\varepsilon(s) > 0$ implies q > 0. Then there exists a diverging sequence $s_1, s_2, \dots \to \infty$ with $\log(s_i) < \frac{1+q}{2} \frac{\tau_B}{\varepsilon(s_i)}$ for all i, whence

$$\limsup_{s \to \infty} \frac{\varepsilon(s)}{s} e^{\tau_B/\varepsilon(s)} \ge \limsup_{i \to \infty} \frac{\varepsilon(s_i)}{s_i} e^{\frac{2}{1+q}\log(s_i)} = \limsup_{i \to \infty} \varepsilon(s_i) s_i^{\frac{1-q}{1+q}} = \infty.$$

From this it follows that $\varepsilon(s) \geq \tau_B/\log s$ for sufficiently large s. Now, on the interval $(0,\infty)$, the function $f(\varepsilon)$ is convex with a unique minimum at $\varepsilon = \tau_B$. Let ε' be a point where $f(\varepsilon') > f(\tau_B/\log s) = \tau_B/\log s$. Then $\varepsilon' \notin [\frac{\tau_B}{\log s}, \tau_B]$. If $\varepsilon' > \tau_B$, then $f(\varepsilon') < \frac{\varepsilon'}{s}e^1$. Since $\frac{\tau_B}{\log s} < f(\varepsilon')$, we then conclude $\varepsilon' > \frac{s\tau_B}{e^1\log s}$, which is not bounded as $s \to \infty$.

In summary, we see that the largest possible lower bound which obeys (21) must have $f(\varepsilon)$ asymptotically $\leq \tau_B / \log s$, and that this bound is achieved by setting $\varepsilon(s) = \tau_B / \log s$. Plugging this in to equation (17) yields the claim.

PROOF OF COROLLARY 6. For $c \in (0, 1)$, choose J large enough so that (J-8)/J > c, and fix τ_A so that $T = t_I + J\tau_A$. Then $(J-8)\tau_A \ge cJ\tau_A = c(T-t_I)$. Substituting the above inequalities into equation (14) and letting $C'' = C'c^2$ yields the desired result.

Acknowledgments. We thank Anand Bhaskar for helpful comments on a draft of this paper and for suggesting Corollary 6 to simplify the presentation of the main result. We also thank Jack Kamm and Jeff Spence for useful feedback.

REFERENCES

- 1000 GENOMES PROJECT CONSORTIUM, (2010). A map of human genome variation from population-scale sequencing. *Nature* 467 1061-1073.
- [2] BHASKAR, A. and SONG, Y. S. (2014). Descartes' rule of signs and the identifiability of population demographic models from genomic variation data. *Annals of Statistics* 42 2469–2493.
- [3] BHASKAR, A., WANG, Y. X. R. and SONG, Y. S. (2015). Efficient inference of population size histories and locus-specific mutation rates from large-sample genomic variation data. *Genome Research* 25 268-279.
- [4] CHEN, H. (2012). The joint allele frequency spectrum of multiple populations: A coalescent theory approach. *Theoretical Population Biology* 81 179–195.
- [5] COVENTRY, A., BULL-OTTERSON, L. M., LIU, X., CLARK, A. G., MAXWELL, T. J., CROSBY, J., HIXSON, J. E., REA, T. J., MUZNY, D. M., LEWIS, L. R. and et al. (2010). Deep resequencing reveals excess rare recent variants consistent with explosive population growth. *Nature Communications* 1 131.
- [6] EXCOFFIER, L., DUPANLOUP, I., HUERTA-SÁNCHEZ, E., SOUSA, V. C. and FOLL, M. (2013). Robust Demographic Inference from Genomic and SNP Data. *PLoS Genetics* 9 e1003905.
- [7] GAZAVE, E., MA, L., CHANG, D., COVENTRY, A., GAO, F., MUZNY, D., BOER-WINKLE, E., GIBBS, R. A., SING, C. F., CLARK, A. G. and et al. (2014). Neutral genomic regions refine models of recent rapid human population growth. *Proceedings* of the National Academy of Sciences 111 757–762.
- [8] GRAVEL, S., HENN, B. M., GUTENKUNST, R. N., INDAP, A. R., MARTH, G. T., CLARK, A. G., YU, F., GIBBS, R. A., BUSTAMANTE, C. D., ALTSHULER, D. L. and et al. (2011). Demographic history and rare allele sharing among human populations. *Proceedings of the National Academy of Sciences* 108 11983–11988.
- [9] GRIFFITHS, R. and TAVARÉ, S. (1998). The age of a mutation in a general coalescent tree. Communications in Statistics. Stochastic Models 14 273–295.
- [10] GUTENKUNST, R. N., HERNANDEZ, R. D., WILLIAMSON, S. H. and BUSTA-MANTE, C. D. (2009). Inferring the Joint Demographic History of Multiple Populations from Multidimensional SNP Frequency Data. *PLoS Genetics* 5 e1000695.
- [11] KIM, J., MOSSEL, E., RÁCZ, M. Z. and ROSS, N. (2015). Can one hear the shape of a population history? *Theoretical Population Biology* **100** 26–38.
- [12] KIMURA, M. (1969). The number of heterozygous nucleotide sites maintained in a finite population due to steady flux of mutations. *Genetics* 61 893.
- [13] KINGMAN, J. F. C. (1982). The coalescent. Stoch. Process. Appl. 13 235-248.
- [14] KINGMAN, J. F. C. (1982). Exchangeability and the evolution of large populations. In *Exchangeability in Probability and Statistics* (G. Koch and F. Spizzichino, eds.) 97–112. North-Holland Publishing Company.
- [15] KINGMAN, J. F. C. (1982). On the genealogy of large populations. J. Appl. Prob. 19A 27-43.
- [16] LI, H. and DURBIN, R. (2011). Inference of human population history from individual whole-genome sequences. *Nature* 475 493–496.
- [17] MASSART, P. (2007). Concentration inequalities and model selection **1896**. Berlin: Springer.
- [18] MYERS, S., FEFFERMAN, C. and PATTERSON, N. (2008). Can one learn history from the allelic spectrum? *Theor. Popul. Biol.* **73** 342–348.

- [19] NELSON, M. R., WEGMANN, D., EHM, M. G., KESSNER, D., JEAN, P. S., VERZILLI, C., SHEN, J., TANG, Z., BACANU, S.-A., FRASER, D. and et al. (2012). An Abundance of Rare Functional Variants in 202 Drug Target Genes Sequenced in 14,002 People. *Science* **337** 100–104.
- [20] NIELSEN, R. (2000). Estimation of population parameters and recombination rates from single nucleotide polymorphisms. *Genetics* 154 931–942.
- [21] PAUL, J. S., STEINRÜCKEN, M. and SONG, Y. S. (2011). An accurate sequentially Markov conditional sampling distribution for the coalescent with recombination. *Genetics* 187 1115–1128. (PMC3070520).
- [22] POLANSKI, A., BOBROWSKI, A. and KIMMEL, M. (2003). A note on distributions of times to coalescence, under time-dependent population size. *Theoretical Population Biology* **63** 33–40.
- [23] POLANSKI, A. and KIMMEL, M. (2003). New Explicit Expressions for Relative Frequencies of Single-Nucleotide Polymorphisms With Application to Statistical Inference on Population Growth. *Genetics* 165 427–436.
- [24] RASMUSSEN, M. D., HUBISZ, M. J., GRONAU, I. and SIEPEL, A. (2014). Genomewide inference of ancestral recombination graphs. *PLoS Genetics* 10 e1004342.
- [25] SCHIFFELS, S. and DURBIN, R. (2014). Inferring human population size and separation history from multiple genome sequences. *Nature Genetics* 46 919-925.
- [26] SHEEHAN, S., HARRIS, K. and SONG, Y. S. (2013). Estimating variable effective population sizes from multiple genomes: A sequentially Markov conditional sampling distribution approach. *Genetics* **194** 647–662. (PMC3697970).
- [27] STEINRÜCKEN, M., PAUL, J. S. and SONG, Y. S. (2013). A sequentially Markov conditional sampling distribution for structured populations with migration and recombination. *Theor. Popul. Biol.* 87 51–61. (PMC3532580).
- [28] TENNESSEN, J. A., BIGHAM, A. W., O'CONNOR, T. D., FU, W., KENNY, E. E., GRAVEL, S., MCGEE, S., DO, R., LIU, X., JUN, G. and et al. (2012). Evolution and Functional Impact of Rare Coding Variation from Deep Sequencing of Human Exomes. *Science* 337 64–69.
- [29] TSYBAKOV, A. B. (2009). Introduction to Nonparametric Estimation. Springer Science+Business Media, New York.
- [30] YU, B. (1997). Assouad, Fano, and Le Cam. In *Festschrift for Lucien Le Cam* (G. L. Y. David Pollard Erik Torgersen, ed.) 423–435. Springer.
- [31] The Exome Aggregation Consortium. http://exac.broadinstitute.org/.
- [32] The UK10K Project: Rare Genetic Variants in Health and Disease. http://www. uk10k.org/.

J. TERHORST DEPARTMENT OF STATISTICS UNIVERSITY OF CALIFORNIA, BERKELEY BERKELEY, CA 94720 USA E-MAIL: terhorst@stat.berkeley.edu Y. S. Song Department of Statistics and Computer Science Division University of California, Berkeley Berkeley, CA 94720 USA E-mail: yss@stat.berkeley.edu