HETEROGENEOUS CHANGE POINT INFERENCE

FLORIAN PEIN¹, HANNES SIELING¹ AND AXEL MUNK¹,²

ABSTRACT. We propose H-SMUCE (heterogeneous simultaneous multiscale changepoint estimator) for the detection of multiple change-points of the signal in a heterogeneous gaussian regression model. A piecewise constant function is estimated by minimizing the number of change-points over the acceptance region of a multiscale test which locally adapts to changes in the variance. The multiscale test is a combination of local likelihood ratio tests which are properly calibrated by scale dependent critical values in order to keep a global nominal level α , even for finite samples.

We show that H-SMUCE controls the error of over- and underestimation of the number of change-points. To this end, new deviation bounds for F-type statistics are derived. Moreover, we obtain confidence sets for the whole signal. All results are non-asymptotic and uniform over a large class of heterogeneous change-point models. H-SMUCE is fast to compute, achieves the optimal detection rate and estimates the number of changepoints at almost optimal accuracy for vanishing signals, while still being robust.

We compare H-SMUCE with several state of the art methods in simulations and analyse current recordings of a transmembrane protein in the bacterial outer membrane with pronounced heterogeneity for its states. An R-package is available online.

1. INTRODUCTION

1.1. Change-point regression. Multiple change-point detection is a long standing task in statistical research and related areas. One of the most fundamental models in this context is (homogeneous) gaussian change-point regression

(1.1) $Y_i = \mu(i/n) + \sigma \epsilon_i, \ i = 1, \dots, n.$

Here, $Y = (Y_1, \ldots, Y_n)$ denotes the observations, μ is an unknown piecewise constant mean function, σ^2 a constant (homogeneous) variance and $\epsilon_1, \ldots, \epsilon_n$ are independent standard gaussian distributed errors. For simplicity, we restrict ourself in this paper to an equidistant sampling scheme $x_{i,n} = i/n$, but extensions to other designs are straightforward.

¹Institute for Mathematical Stochastics, Georg-August-University of Göttingen, Goldschmidtstrasse 7, 37077 Göttingen

²MAX PLANCK INSTITUTE FOR BIOPHYSICAL CHEMISTRY, AM FASSBERG 11, 37077 GÖTTINGEN *E-mail address*: {fpein, hsieling, munk}@math.uni-goettingen.de.

Date: September 24, 2018.

²⁰¹⁰ Mathematics Subject Classification. 62G08,62G15,90C39.

Key words and phrases. change-point regression, deviation bounds, dynamic programming, heterogeneous noise, honest confidence sets, ion channel recordings, multiscale methods, robustness, scale dependent critical values.

Methods for estimating the change-points in (1.1) and in related models are vast, see for instance (Yao, 1988; Donoho and Johnstone, 1994; Csörgo and Horváth, 1997; Bai and Perron, 1998; Braun et al., 2000; Birgé and Massart, 2001; Kolaczyk and Nowak, 2005; Boysen et al., 2009; Harchaoui and Lévy-Leduc, 2010; Jeng et al., 2010; Killick et al., 2012; Rigollet and Tsybakov, 2012; Zhang and Siegmund, 2012; Fryzlewicz, 2014) and the references in these papers.

A crucial condition in most of the afore-mentioned papers is the assumption of homogeneous noise, i.e. a constant variance σ^2 in (1.1). In many applications, however, this assumption is violated and the variance σ^2 varies over time, $\sigma^2(i/n)$, say. This problem arises for instance in the analysis of array CGH data, see (Muggeo and Adelfio, 2011; Arlot and Celisse, 2011). Further examples include economic applications, e.g. the real interest rate is modelled by Bai and Perron (2003) as piecewise linear regression with covariates and heterogeneous noise. In this paper we will discuss an example from membrane biophysics, the recordings of ion channels, see Section 5. It is well known that the noise of the open state can be much larger than the background noise, see (Sakmann and Neher, 1995, Section 3.4.4) and the references therein, rendering the different states as a potential source for variance heterogeneity.

To illustrate the effects of missing heterogeneity we show in Figure 1 a reconstruction by SMUCE¹ (Frick et al., 2014), a method that has been designed for homogeneous noise. The constant variance assumption of SMUCE leads to an overestimation of the standard deviation (which is pre-estimated by a global IQR type estimator) in the first half and an underestimation in the second half. Therefore, in Figure 1 SMUCE misses the first change-point and includes artificial change-points in the second half to compensate for the too small variance it is forced to use, see also (Zhou, 2014). Note, that this flaw is not a particular feature of SMUCE, it will occur for any sensible segmentation method which relies on a constant variance assumption. Hence, from Figure 1 the fundamental difficulty of the heterogeneous (multiscale) change-point regression problem becomes apparent: How to decide whether a change of fluctuations of the data result from high frequent changes in the mean μ or merely from an increase of the noise level? Apparently, if changes can occur on any scale (i.e. the length of an interval of neighbouring observations) this is a notoriously difficult issue and proper separation of signal and noise cannot be performed without extra information.

Indeed, the basis of the presented theory is that often a reasonable assumption is to exclude changes of the variance in constant segments of μ (see Section 1.2). Under this relatively weak assumption, we show in this paper that estimation of μ for heterogeneous data in a multiscale fashion becomes indeed feasible. In addition, we also aim for a

¹http://cran.r-project.org/web/packages/stepR, v. 1.0-3, 2015-06-18



(b) Simulated observations (black dots) together with the true signal (black line), the confidence band (grey), the confidence intervals for the change-point locations (brackets and thick lines), estimated change-points locations (red dashes) as well as the estimates by H-SMUCE (red dotted line) and by SMUCE (blue dashed line), both with $\alpha = 0.1$.

FIGURE 1. Illustration of missing heterogeneity.

method which is robust when changes in the variance occur at locations where the signal is constant, as we believe that this cannot be excluded in many practical cases. To this end, we introduce a new estimator H-SMUCE (heterogeneous simultaneous multiscale change-point estimator) which recovers the signal under heterogeneous noise over a broad range of scales, controls the family-wise error rate to overestimate the number of change-points, allows for confidence statements for all unknown quantities, obeys certain statistical optimality properties, and can be efficiently computed. At the same hand it is robust against heterogeneous noise on constant signal segments, which as a byproduct reveals it also as robust against more heavily tailed errors.

1.2. Heterogeneous change-point model. To be more specific, from now on we consider the *heterogeneous* gaussian change-point model

(1.2)
$$Y_i = \mu(i/n) + \sigma(i/n)\epsilon_i, \ i = 1, \dots, n_i$$

where now the variance σ^2 is also given by an unknown piecewise constant function. For the following theoretical results we assume that it only can have possible change-points at the same locations as the mean function μ . In other words, (μ, σ^2) is a pair of unknown piecewise constant functions in

(1.3)
$$\mathcal{S} := \left\{ (\mu, \sigma^2) : [0, 1] \mapsto \mathbb{R}^2, \ \mu = \sum_{k=0}^K m_k \mathbb{1}_{[\tau_k, \tau_{k+1})}, \ \sigma^2 = \sum_{k=0}^K s_k^2 \mathbb{1}_{[\tau_k, \tau_{k+1})}, \ K \in \mathbb{N} \right\},$$

with unknown change-point locations $\tau_0 = 0 < \tau_1 < \cdots < \tau_K < 1 = \tau_{K+1}$ for some unknown number of change-points $K \in \mathbb{N}$ and also unknown function values $m_k \in \mathbb{R}$ and $s_k^2 \in \mathbb{R}_+$ of μ and σ^2 . By technical reasons, we define $\mu(1)$ and $\sigma^2(1)$ by continuous extension of μ and σ^2 , respectively. For identifiability of μ we assume $m_k \neq m_{k+1} \forall k =$ $0,\ldots,K$ and exclude isolated changes in the signal by assuming that $\mu:[0,1]\to\mathbb{R}$ is a right continuous function. It is important to stress that in (1.3) we allow the variance to potentially have changes at the locations of the changes of the signal, but the variance σ^2 need not necessarily change when μ changes, as we do not assume $s_k^2 \neq s_{k+1}^2$. In particular, homogeneous observations are still part of the model. The other way around, we assume that within a constant segment of μ it may not happen that the variance changes, i.e. the local signal to noise ratio is assumed to be constant on $[\tau_k, \tau_{k+1})$ for all $k = 0, \ldots, K$. We argue that this is a reasonable assumption in many applications (recall the examples given above and see our data example in Section 5), since a changepoint represents typically a change of the condition of the underlying state. Moreover, for example, in many engineering applications locally a constant signal to noise ratio is assumed (Guillaume et al., 1990), which motivates our modelling as well. However, we stress that the restriction to model (1.3) is only required for our theory. For the practical application we will show in simulations in Section 4.2 that H-SMUCE is in addition robust against a violation of this assumption (i.e. when a variance change may occur without a signal change) and hence works still well in the general heterogeneous change-point model (1.2) with arbitrary variance changes.

1.3. Heterogeneous change-point regression. Up to our best knowledge there are only few methods which explicitly take into account the heterogeneity of the noise in change-point regression, either in the model considered here or in related models. Thereby, we have to distinguish two settings.

First, that also changes in the variance are considered as relevant structural changes of the underlying data (even when the mean does not change) and to seek for changes in the mean and in the variance, respectively. In this spirit are local search methods, such as binary segmentation (BS) (Scott and Knott, 1974; Vostrikova, 1981) (if the corresponding single change-point detection method takes the heterogeneous variance into account), but also global methods can achieve this goal, e.g. PELT (Killick et al., 2012). For a Bayesian approach in this context see (Du et al., 2015) and the references therein. In addition, methods which search for more general structural changes in the distribution potentially apply to this setup as well, see e.g. (Csörgo and Horváth, 1997; Arlot et al., 2012; Matteson and James, 2014).

This is in contrast to the setting we address in this paper: The variance is considered as a nuisance parameter and we primarily seek for changes in the signal μ . Hence, we aim for statistically efficient estimation of the mean function, but still being robust against heterogeneous noise. Obviously, this cannot be achieved by methods addressing the first setting. Although of great practical relevance, this situation has only rarely been considered and in particular no theory exists, to our knowledge. The cross-validation method LOOVF (Arlot and Celisse, 2011) and cumSeg (Muggeo and Adelfio, 2011) have been designed specifically to be robust against heterogeneous noise. Moreover, also circular binary segmentation (CBS), see (Venkatraman and Olshen, 2007), applies to this.

For a better understanding of the problem considered here it is illustrative to distinguish our setting further, namely from the case when it is known before hand that changes in the variance will necessarily occur with changes in the signal. This will potentially increase the detection power as under this assumption variance changes can be used for finding signal changes, as well. The information gain due to the variance changes for this case has been recently quantified by Enikeeva et al. (2015) in terms of the minimax detection boundary for single vanishing signal bumps of size $\delta_n \searrow 0$. More precisely, if the base line variance is σ_0^2 and the variance at the bump is $\sigma_0^2 + \sigma_n^2$ then the constant in the minimax detection boundary is $b = \sqrt{2}\sigma_0\sqrt{2/(2+c^2)}$ for $c = \sigma_0^{-1} \lim_{n\to\infty} \sigma_n/\delta_n$, see (Enikeeva et al., 2015, Theorems 3.1-3.3). For the particular case of homogeneous variance, i.e. $\sigma_n^2 = 0$, we obtain $b = \sqrt{2}\sigma_0$ and the factor $\sqrt{2/(2+c^2)} = 1$ becomes maximal, see also (Dümbgen and Walther, 2008; Frick et al., 2014). This reflects that no additional information on the location of a change can be gained from the variance in the homogeneous case. Comparing this to the inhomogeneous case we see that when the variance change is known to be large enough, i.e. $\sigma_0^{-1} \lim_{n\to\infty} \sigma_n/\delta_n > 0$, additional information for the signal change can be gained from the variance change, as then $b < \sqrt{2}\sigma_0$, provided it is known that signal and variance change *simultaneously*.

In contrast, in the present setting the variance need not necessarily change when the signal changes, hence the "worst case" of no variance change from above is contained in our model, which lower bounds the detection boundary. The situation is further complicated due to the fact that missing knowledge of a variance change can potentially even have an adverse effect because in model (1.3) detection power will be potentially decreased further as the nuisance parameter $\sigma^2(\cdot)$ hinders estimation of change-points of μ . For this situation the optimal minimax *constants* are unknown to us, but from the fact that the model with a constant variance is a submodel of our model (1.3) it immediately follows that the minimax constant for a single bump has to be at least $\sqrt{2}\sigma_0$. This will allow us to show that H-SMUCE attains the same optimal minimax detection *rate* as for the homogeneous case and $4\sigma_0$ instead of $\sqrt{2}\sigma_0$ as the constant appearing in the minimax detection boundary. Remarkably, the only extra assumption we have to suppose is that signal and variance have to be constant on segments at least of order $\log(n)/n$, see Theorem 3.10. This reflects the additional difficulty to separate "locally" signal and noise levels in a multiscale fashion. In other words, when we assume that the number of i.i.d. neighbouring observations (no change in signal and variance) in each segment is at least of order $\log(n)$, separation of signal and noise will be done by H-SMUCE in an optimal way (possibly up to a constant).

1.4. Heterogeneous change-point inference. We define H-SMUCE as the multiscale constrained maximum likelihood estimator restricted to all solutions of the following optimisation problem

(1.4)
$$\operatorname{argmin}_{\mu \in \mathcal{M}} |\mathcal{I}(\mu)| \quad \text{s.t.} \quad \max_{[\frac{i}{n}, \frac{j}{n}] \in \mathcal{D}(\mu)} \left[T_i^j(Y, \mu([i/n, j/n])) - q_{ij} \right] \le 0,$$

see also (Boysen et al., 2009; Davies et al., 2012; Frick et al., 2014) for related approaches. Here, \mathcal{M} (as a subset of \mathcal{S}) is the set of all piecewise constant mean functions, $|\mathcal{I}(\mu)|$ the cardinality of the set of change-points of μ and the right hand side of (1.4) a multiscale constraint to be explained now. Given a candidate function μ this tests *simultaneously* over the system of all intervals $\mathcal{D}(\mu)$ on which μ is constant, whether its function value $\mu([i/n, j/n])$ is the mean value of the observations on the respective interval [i/n, j/n]. In order to perform each test, i.e. to decide whether the observations Y_i, \ldots, Y_j have constant mean $\mu([i/n, j/n])$, the local log-likelihood-ratio statistic

(1.5)
$$T_i^j(Y,\mu([i/n,j/n])) := (j-i+1) \frac{\left(\overline{Y}_{ij} - \mu([i/n,j/n])\right)^2}{\hat{s}_{ij}^2},$$

with $\overline{Y}_{ij} := (j-i+1)^{-1} \sum_{l=i}^{j} Y_l$ and local variance estimate $\hat{s}_{ij}^2 := (j-i)^{-1} \sum_{l=i}^{j} (Y_l - \overline{Y}_{ij})^2$, is compared with a local threshold q_{ij} in a multiscale fashion, to be discussed now. In what follows, we restrict the multiscale test to intervals in the *dyadic partition*

(1.6)
$$\mathcal{D} := \bigcup_{k=1}^{d_n} \mathcal{D}_k,$$

where $d_n := \lfloor \log_2(n) \rfloor$ is the number of different scales and

(1.7)
$$\mathcal{D}_k := \bigcup_{l=1}^{\left\lfloor \frac{n}{2^k} \right\rfloor} \left[\frac{1 + (l-1)2^k}{n}, \frac{l2^k}{n} \right]$$

the set of intervals from the dyadic partition with length $n^{-1}2^k$. This allows fast computation and simplifies the asymptotic analysis. Nevertheless, our methodology can be adapted to other intervals systems, see Remark 2.2.

It remains to determine thresholds q_{ij} for \mathcal{D} in (1.6) that combine the local tests appropriately. To this end, note that logarithmic (or related) scale penalisation as in the

er, 2008; F

homogeneous case (Dümbgen and Spokoiny, 2001; Dümbgen and Walther, 2008; Frick et al., 2014) does not balance scales anymore appropriately in the heterogeneous case. In particular, this will give a multiscale statistic which diverges, since due to the local variance estimation the test statistic fails to have subgaussian (but still has subexponential) tails. To overcome this burden we introduce in Section 2 scale dependent critical values such that the multiscale test has global significance level α , see (2.3). To this end, the different scales are balanced appropriately by weights $\beta_1, \ldots, \beta_{d_n}$, with $d_n := \lfloor \log_2(n) \rfloor$, see (2.4) and (2.5). More precisely, these weights determine the ratios between the rejection probabilities of the multiscale test on a corresponding scale. Existence and uniqueness of the so defined scale dependent critical values is shown in Lemma 2.1 and explicit bounds are given in Lemma 3.1. The weights also allow to incorporate prior scale information, see Section 3.4.

Using the so obtained thresholds q_{ij} allows to obtain several confidence statements which are a main feature of H-SMUCE. First of all, we show in Section 3 that the probability to overestimate the number of change-points is bounded by the significance level α uniformly over S in (1.3), $\mathbb{P}(\hat{K} > K) \leq \alpha$, see Theorem 3.3. More specifically, we show the *overestimation* bound

(1.8)
$$\sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} > K + 2k\right) \le \alpha^{k+1}, \ \forall \ k \in \mathbb{N}_0,$$

see Theorem 3.4. In Theorem 3.5 we provide an exponential bound for the underestimation of the number of change-points by H-SMUCE, $\mathbb{P}(\hat{K} < K)$. To this end, we show new exponential deviation bounds for *F*-statistics (Section C.3), which might be of interest by its own. Combining the over- and the underestimation bound provides upper bounds for the errors $\mathbb{P}(\hat{K} \neq K)$ and $\mathbb{E}[|\hat{K} - K|]$. For a fixed signal both bounds vanish super polynomially in *n* if $\alpha = \alpha_n \searrow 0$ when the weights are chosen appropriately, see Remark 3.6. Consequently, the estimated number of change-points converges almost surely to the true number, see Theorem 3.7. Further, these exponential bounds enable us to obtain a confidence band for the signal μ as well as confidence intervals for the locations of the change-points, for an illustration see Figures 1 and 2. We show that the diameters of these confidence intervals decrease asymptotically as fast as the (optimal) sampling rate up to a log factor. All confidence statements hold uniformly over $S_{\Delta,\lambda} \subset S$, all functions with minimal signal to noise ratio $\geq \Delta$ and minimal scale $\geq \lambda := \min_{k=0,\dots,K} (\tau_{k+1} - \tau_k)$, with Δ and λ arbitrarily, but fixed, see Theorems 3.8 and 3.9.

1.5. **H-SMUCE in action.** Figure 2 illustrates the performance of H-SMUCE in an example with $n = 1\,000$ observations and K = 10 change-points. We found that H-SMUCE misses for $\alpha = 0.1$ one change-point (as the choice $\alpha = 0.1$ tunes H-SMUCE to provide



(a) True standard deviation (black) and estimates resulting from H-SMUCE at $\alpha = 0.1$ (red line) and $\alpha = 0.3, 0.5$ (blue line).



FIGURE 2. b-d: Observations (black dots), true signal (black line), confidence band (grey), confidence intervals for the change-point locations (brackets and thick lines), estimated change-points locations (red dashes) and estimate (red line) H-SMUCE at given α and with equal weights $\beta_1 = \cdots = \beta_{d_n} = 1/d_n$, see (2.4) and (2.5).

the strong guarantee not to overestimate the number of change-points K with probability 0.9, see (1.8)), whereas for α between 0.15 and 0.99 (only displayed for $\alpha = 0.3$ and $\alpha = 0.5$) the correct number of change-points is detected always (while providing a weaker guarantee for not overestimating K). In addition, for α between 0.15 and 0.99 each true change-point is covered by the associated confidence interval at level $1 - \alpha$. This illustrates the influence of the significance level α . Notably, we find that the reconstructions are remarkably stable in α . In fact, combining Lemma 3.1 and (A.1) shows that the width of the confidence band is proportional to $\sqrt{\log(1/\alpha)}$ which decreases only logarithmically for increasing α .

We compare the performance of H-SMUCE with CBS (Venkatraman and Olshen, 2007), cumSeg (Muggeo and Adelfio, 2011) and LOOVF (Arlot and Celisse, 2011) in several simulation studies in Section 4 (see also Figure 9 in Supplement B for their performance on the data in Figure 2), where we also examine robustness issues, see Section 4.2. In all of these simulations and in the subsequent application H-SMUCE performs very robust and includes too many change-points only rarely in accordance with (1.8).

In Section 5 we apply H-SMUCE to current recordings of a transmembrane protein with pronounced heterogeneity for its states. In contrast to segmentation methods which rely on homogeneous noise, we found that H-SMUCE provides a reasonable reconstruction, where all visible gating events are detected.

Finally, we stress that the confidence band and confidence intervals for the change-point locations provided by H-SMUCE can be used to accompany any segmentation method to assess significance of its estimated change-points. This is illustrated in Section 5 as well. Computation of the estimator by a pruned dynamic program and of the critical values based on Monte-Carlo simulation is explained carefully in Supplement A. There we also study the theoretical and empirical computation time of H-SMUCE. Due to the underlying dyadic partition the computation of H-SMUCE is very fast, in some scenarios even linear in the number of observations. Additional simulations results are collected in Supplement B and all proofs are given together with some auxiliary statements in Supplement C. An R-package is available online².

2. Scale dependent critical values

For the definition of H-SMUCE it remains to determine the local thresholds q_{ij} in (1.4). First of all, the multiscale test on the r.h.s. in (1.4) should be a level α test, i.e.

(2.1)
$$\sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{P}_{(\mu,\sigma^2)} \left(\max_{\left[\frac{i}{n},\frac{j}{n}\right]\in\mathcal{D}(\mu)} \left[T_i^j \left(Y, \mu(\left[i/n, j/n\right])\right) - q_{ij} \right] > 0 \right) \le \alpha.$$

²http://www.stochastik.math.uni-goettingen.de/hsmuce

Here, we use the same threshold for all intervals of the same length as no a-priori information on the change-point locations is assumed. More precisely, as we have restricted the multiscale test to the dyadic partition in (1.6) we aim to find a vector of critical values $\mathbf{q} := (q_1, \ldots, q_{d_n})$, where now $q_{ij} := q_k$ if and only if $j - i + 1 = 2^k$. To this end, w.l.o.g., we may consider standard gaussian observations Z_1, \ldots, Z_n instead of Y_1, \ldots, Y_n , since the supremum in (2.1) is attained at $\mu \equiv 0$ and $\sigma^2 \equiv 1$, see the proof of Theorem 3.3. We then define the statistics T_1, \ldots, T_{d_n} with \mathcal{D}_k in (1.7) as

(2.2)
$$T_k := \max_{[i/n, j/n] \in \mathcal{D}_k} T_i^j(Z, 0) \text{ for } k = 1, \dots, d_n$$

Then, the critical values q_1, \ldots, q_{d_n} fulfil (2.1) if

(2.3)
$$\mathbb{P}\left(\max_{k=1,\dots,d_n} \left[T_k - q_k\right] > 0\right) = 1 - F\left(q_1,\dots,q_{d_n}\right) = \alpha,$$

with F the cumulative distribution function of (T_1, \ldots, T_{d_n}) .

As the critical values q_1, \ldots, q_{d_n} are not uniquely determined by (2.3) they can be chosen to render the multiscale test particularly powerful for certain scales. To this end, we introduce weights

(2.4)
$$\beta_1, \dots, \beta_{d_n} \ge 0, \text{ with } \sum_{k=1}^{d_n} \beta_k = 1,$$

where $\beta_k = 0$ means to omit the k-th scale, i.e. $q_k = \infty$. Finally, we define q_1, \ldots, q_{d_n} implicitly through

(2.5)
$$\frac{1 - F_1(q_1)}{\beta_1} = \dots = \frac{1 - F_{d_n}(q_{d_n})}{\beta_{d_n}}$$

with F_k the cumulative distribution function of T_k . If $\beta_k = 0$ this will not enter the systems of equations in (2.5). The weights determine the fractions between the probabilities that a test on a certain scale rejects, and hence regulate the allocation of the level α among the single scales. In summary, the choice of the local thresholds q_{ij} boils down to choosing the significance level α and the weights $\beta_1, \ldots, \beta_{d_n}$, we discuss these choices in Section 3.4 more carefully. If no prior information on scales is available a default option is always to set all weights equal, i.e. $\beta_1 = \ldots = \beta_{d_n} = 1/d_n$.

The next result shows that the vector of critical values satisfying (2.3)-(2.5) is always well-defined.

Lemma 2.1 (Existence and uniqueness). For any $\alpha \in (0, 1)$ and for any weights $\beta_1, \ldots, \beta_{d_n}$, s.t. (2.4) holds, there exits a unique vector of critical values $\boldsymbol{q} = (q_1, \ldots, q_{d_n}) \in \mathbb{R}^{d_n}_+$ which fulfils the equations (2.3) and (2.5). An explicit computation of the vector \mathbf{q} (or F) appears to be very hard, since the statistics T_1, \ldots, T_{d_n} are dependent, although the dependence structure is explicitly known. Alternatively, it would be helpful to have an approximation for the distribution (and hence its quantiles) of the maximum in (2.1), which, however, appears to be rather difficult, as well. For the case when $q_{ij} \equiv q$ (which does not apply to H-SMUCE), see Davies (1987, 2002). Therefore, we determine in Section A.2 the vector \mathbf{q} by Monte-Carlo simulations. Note that the distribution does not depend on the specific element $(\mu, \sigma^2) \in S$ and hence the critical values can be computed in a universal manner. We stress that the determination of the scale dependent critical values is not restricted to our setting and can also be applied to multiscale testing in other contexts. Different to scale penalisation and like the block criterion in (Rufibach and Walther, 2010) no model dependent derivations are required and the critical values are adapted to the exact finite sample distribution of the local test statistics. However, our approach allows additionally a flexible scale calibration by the choice of the weights (see Section 3.4) and arbitrary interval sets can be used as the following remark points out.

Remark 2.2 (Other interval sets). H-SMUCE can be easily adjusted to other interval sets as follows. Let \mathcal{I} be an arbitrary set of intervals. Then, we replace in the definition of H-SMUCE in (3.3) and (3.7) the set \mathcal{D} by the set \mathcal{I} and the vector (T_1, \ldots, T_{d_n}) by the vector $(\tilde{T}_2, \ldots, \tilde{T}_n)$ (empty scales should be omitted) in Section 2, with

(2.6)
$$\widetilde{T}_k := \max_{\substack{[i/n,j/n] \in \mathcal{I}, \\ j-i+1=k}} T_i^j(Z,0).$$

Again it remains to choose the significance level $\alpha \in (0, 1)$ and the weights β_2, \ldots, β_n to determine the critical values required for H-SMUCE. Note, however, that the critical values and its bounds in Lemma 3.1 and therefore the results in Section 3 (besides of Theorems 3.3 and 3.4) will depend on the specific system \mathcal{I} and have to be computed for each \mathcal{I} separately.

Employing a larger interval set than \mathcal{D} may lead to a better detection power, but at the price of a larger computation time. Hence, in practice, a trade-off between computational and statistical efficiency may guide this choice as well. Our R-package includes beside of the dyadic partition also the system of all intervals (of order $\mathcal{O}(n^2)$, statistically most efficient, but computationally expensive) and the system of all intervals of dyadic length ($\mathcal{O}(n \log(n))$, intermediate efficiency and computational time). Interesting choices might be also approximating sets like \mathcal{J}_{app} introduced in (Walther, 2010; Rivera and Walther, 2013) which are larger than the dyadic partition, but achieve the minimax boundary in the context of density estimation.

3. Theory

In this section we collect our theoretical results. We start with finite bounds for the critical values. These will allow to bound $\mathbb{P}(\hat{K} \neq K)$. With these bounds we obtain confidence statements for the signal μ and its main characteristics. Finally, we investigate asymptotic detection rates of H-SMUCE for vanishing signals.

3.1. Finite bounds for over- and underestimation. In the following we require upper bounds for the critical values, since the definition of the critical values by the equations (2.3)-(2.5) is implicit.

Lemma 3.1 (Bound on critical values). Let $\boldsymbol{q} = (q_1, \ldots, q_{d_n})$ be the vector of critical values defined by (2.3)-(2.5), then for every $k \in \{2, \ldots, d_n\}$ such that

(3.1)
$$2^{-k} \log\left(\frac{n}{2^k \alpha \beta_k}\right) \le \frac{1}{2}$$

we have

(3.2)
$$q_k \le 8 \log\left(\frac{n}{2^k \alpha \beta_k}\right).$$

Remark 3.2. The log term of the bound (3.2) can be split into a scale dependent penalty term $\log(n2^{-k})$ which is of the same order as the penalties in the homogeneous case in (Dümbgen and Spokoiny, 2001; Frick et al., 2014), and into the term $\log((\alpha\beta_k)^{-1})$ which incorporates the significance level α and the weight β_k .

The following theorem shows that the significance level α controls the probability to overestimate the number of change-points.

Theorem 3.3 (Overestimation control I). Assume the heterogeneous gaussian changepoint model (1.2). Let $K := |\mathcal{I}(\mu)|$ be the number of change-points of a signal $\mu \in \mathcal{M}$. Let further \hat{K} be the estimated number of change-points by H-SMUCE, *i.e.*

(3.3)
$$\hat{K} := \min\left\{ |\mathcal{I}(\mu)| : \mu \in \mathcal{M} \text{ with } \max_{[\frac{i}{n}, \frac{j}{n}] \in \mathcal{D}(\mu)} \left[T_i^j \left(Y, \mu([i/n, j/n]) \right) - q_{ij} \right] \le 0 \right\}.$$

Then, for any vector of critical values \mathbf{q} with significance level $\alpha \in (0,1)$ and weights $\beta_1, \ldots, \beta_{d_n}$ in (2.3)-(2.5), uniformly over \mathcal{S} in (1.3) it holds

$$\sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} > K\right) \le \alpha.$$

The theorem gives us a direct interpretation of the parameter α as the probability to overestimate the number of change-points. This even holds locally, i.e. on every union of adjoining segments of the estimator H-SMUCE with probability $1 - \alpha$ there are at least as many change-points as detected. Moreover, we strengthen the result by showing that

the probability to estimate additional changes decays exponentially fast and hence the expected overestimation is small.

Theorem 3.4 (Overestimation control II). Under the assumptions of Theorem 3.3, we have

$$\sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} > K + 2k\right) \le \alpha^{k+1}, \ \forall \ k \in \mathbb{N}_0.$$

Moreover,

$$\sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{E}_{(\mu,\sigma^2)} \left[(\hat{K} - K)_+ \right] \le \frac{2\alpha}{1-\alpha}$$

To control the probability $\mathbb{P}(\hat{K} \neq K)$ we need additionally an upper bound for the probability to *underestimate* K. Unlike to the *overestimation* bounds in the Theorems 3.3 and 3.4 the probability to underestimate cannot be bounded uniformly over S, since size and scale of changes could be arbitrarily small. This is made more precise in Theorem 3.10 which gives the detection boundary in terms of the smallest (standardized) jump size Δ and the smallest scale λ . The next theorem provides an exponential bound uniformly over the subset

$$(3.4) \quad \mathcal{S}_{\Delta,\lambda} := \left\{ (\mu, \sigma^2) \in \mathcal{S} : \Delta \leq \inf_{1 \leq k \leq K} \frac{|\mu_k - \mu_{k-1}|}{\max(s_{k-1}, s_k)} \text{ and } \lambda \leq \inf_{0 \leq k \leq K} (\tau_{k+1} - \tau_k) \right\},$$

with $\Delta, \lambda > 0$ arbitrary, but fixed.

Theorem 3.5 (Underestimation control). Let $S_{\Delta,\lambda}$ be as in (3.4) with $\Delta, \lambda > 0$ arbitrary, but fixed, and $k_n := \lfloor \log_2(n\lambda/4) \rfloor$. We define

$$\eta := \left[1 - 3\exp\left(-\frac{1}{48}\left(\sqrt{\frac{n\lambda\Delta^2}{32}} - \sqrt{16\log\left(\frac{8}{\lambda\alpha\beta_{k_n}}\right)}\right)_+^2\right)\right]_+^2.$$

Under the assumptions of Theorem 3.3 and if $n\lambda \geq 32$ and

$$(n\lambda)^{-1}\log\left(\frac{8}{\lambda\alpha\beta_{k_n}}\right) \le \frac{1}{512}$$

are satisfied, then uniformly in $\mathcal{S}_{\Delta,\lambda}$

(3.5)
$$\mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} < K\right) \le 1 - \eta^K \text{ and } \mathbb{E}_{(\mu,\sigma^2)}\left[\left(K - \hat{K}\right)_+\right] \le K \left(1 - \eta\right).$$

Roughly speaking, H-SMUCE detects any change-point of the signal μ under assumptions of Theorem 3.5 at least with probability η . A sharper version with different probabilities η_1, \ldots, η_K is given in Theorem C.5 in the supplement. Such a result clarifies the dependence on the different weights, but is technically way more difficult. Combining Theorems 3.3, 3.4 and 3.5 gives upper bounds for the probability $\mathbb{P}(\hat{K} \neq K)$ and the expectation $\mathbb{E}[|\hat{K} - K|]$ that H-SMUCE missspecifies the number of change-points. Remark 3.6 (Vanishing errors). For a fixed signal (fixed Δ and λ are sufficient) both errors vanish asymptotically if $\alpha = \alpha_n \to 0$ is chosen such that $\log(\alpha_n \beta_{k_n,n})/n \to 0$, with triangular scheme $\beta_{1,n}, \ldots, \beta_{d_n,n}$ for the weights in (2.5). We can achieve a rate arbitrary close to the exponential rate by the choice $\alpha_n = \exp(-n/r_n)$, with $r_n \to \infty$ arbitrarily slow. The condition on the sequence $\beta_{k_n,n}$ allows a variety of possible choices of the weights, too. For instance, the choice $\beta_{1,n} = \cdots = \beta_{d_n,n} = 1/d_n$, which weights all scales equally, fulfils this condition.

A direct consequence is the strong model consistency of H-SMUCE.

Theorem 3.7 (Strong model consistency). Assume the setting of Theorem 3.3 and let $(\hat{K}_n)_n$ be the sequence of estimated numbers of change-points by H-SMUCE, where \hat{K}_n is as \hat{K} with significance level α_n and corresponding weights $\beta_{1,n}, \ldots, \beta_{d_n,n}$. Moreover, let $S_{\Delta,\lambda}$ be as in (3.4) with $\Delta, \lambda > 0$ arbitrary, but fixed, and $k_n := \lfloor \log_2(n\lambda/4) \rfloor$. Let $\rho > 0$ be arbitrary, but fixed. If

(3.6)
$$\lim_{n \to \infty} \frac{n^{1+\rho}}{\alpha_n} = 0 \text{ and } \lim_{n \to \infty} \frac{\log(\alpha_n \beta_{k_n,n})}{n} = 0$$

holds, then $\hat{K}_n \to K$, almost surely and uniformly in $\mathcal{S}_{\Delta,\lambda}$.

Again, there is a wide range of sequences α_n and $\beta_{k_n,n}$ to satisfy (3.6). Moreover, we still have (weak) model consistency, if $\alpha_n \to 0$ and the second condition of (3.6) holds.

3.2. Confidence sets. In this section we obtain confidence sets for the signal μ and for the locations of the change-points. First, we show that the set of all solutions of (1.4)

(3.7)
$$C(\mathbf{q}) := \left\{ \mu \in \mathcal{M} : |\mathcal{I}(\mu)| = \hat{K} \text{ and } \max_{\left[\frac{i}{n}, \frac{j}{n}\right] \in \mathcal{D}(\mu)} \left[T_i^j \left(Y, \mu(\left[i/n, j/n\right]) \right) - q_{ij} \right] \le 0 \right\}$$

is a confidence set for the unknown signal μ .

Theorem 3.8 (Confidence set). Assume the setting of Theorem 3.3 and let $S_{\Delta,\lambda}$ be as in (3.4) with $\Delta, \lambda > 0$ arbitrary, but fixed, and $k_n := \lfloor \log_2(n\lambda/4) \rfloor$. Let $C(\cdot)$ be as in (3.7) and \mathbf{q}_n be a vector of critical values determined by significance level α and weights $\beta_{1,n}, \ldots, \beta_{d_n,n}$, with $\lim_{n\to\infty} n^{-1} \log(\beta_{k_n,n}) = 0$. Then,

(3.8)
$$\lim_{n \to \infty} \inf_{(\mu, \sigma^2) \in \mathcal{S}_{\Delta, \lambda}} \mathbb{P}_{(\mu, \sigma^2)} \left(\mu \in C \left(\boldsymbol{q}_n \right) \right) \ge 1 - \alpha.$$

This shows that the asymptotic coverage of $C(\mathbf{q}_n)$ is at least $1 - \alpha$. Lemma C.6 gives an exponential inequality similar to (3.5) which shows that $C(\mathbf{q}_n)$ is also a non-asymptotic confidence set. We further derive from this set confidence intervals for the change-point locations.

Theorem 3.9 (Change-point locations). Assume the setting of Theorem 3.8, where α is replaced by a sequence $\alpha_n \to 0$. Let $c_n := r_n/n \leq \lambda/2$ and $k_n := \lfloor \log_2(nc_n/2) \rfloor$ s.t.

(3.9)
$$\liminf_{n \to \infty} \frac{r_n}{\log(n)} > \frac{216}{\min(\Delta^2, 1)} \text{ and } \lim_{n \to \infty} \frac{\log(\alpha_n \beta_{k_n, n})}{r_n} = 0.$$

Then,

(3.10)
$$\lim_{n \to \infty} \sup_{(\mu, \sigma^2) \in \mathcal{S}_{\Delta, \lambda}} \mathbb{P}_{(\mu, \sigma^2)} \left(\sup_{\hat{\mu} \in C(q_n)} \max_{k=1, \dots, K} c_n^{-1} \left| \tau_k - \hat{\tau}_k \right| > 1 \right) = 0.$$

Here, the rate c_n is equal to the sampling rate 1/n up to the (logarithmic) rate r_n depending on the tuning parameters $\alpha_n\beta_{k_n,n}$. For example, if $\alpha_n\beta_{k_n,n} \simeq n^{-\gamma}, \gamma \ge 0$, $r_n/\log(n) \to \infty$ is sufficient to satisfy (3.9). A non-asymptotic statement is given in Lemma C.7 in the supplement. For visualization of the confidence statements it is useful to further derive a confidence band $B(\mathbf{q}_n)$ for the signal as in (Frick et al., 2014, Corollary 3 and the explanation around). It can be shown that also the collection $I(\mathbf{q}_n) = {\hat{K}_n, B(\mathbf{q}_n), [L_k, R_k]_{k=1,...,\hat{K}_n}}$, with $[L_k, R_k]$ confidence intervals for the changepoint locations according to Theorem 3.9, satisfies (3.8). Recall Figures 1 and 2 for an illustration. It is also possible to strengthen the statements of this section to sequences of vanishing signals with $\Delta_n \to 0$ and $\lambda_n \to 0$ slow enough, but we omit such results.

3.3. Asymptotic detection rates for vanishing signals. For the detection of a single vanishing bump against a noisy background see Theorem C.8 in the supplement. The following theorem deals with the detection of a signal with several vanishing change-points.

Theorem 3.10 (Multiple vanishing change-points). Assume the heterogeneous gaussian change-point model (1.2). Let $(K_n)_n := (|\mathcal{I}(\mu_n)|)_n$ be the sequence of true number of change-points. Let further $(\hat{K}_n)_n$ be the sequence of the estimated numbers of changepoints by H-SMUCE (3.3), with significance levels α_n and weights $\beta_{1,n}, \ldots, \beta_{d_n,n}$. Let $\mathcal{S}_{\Delta_n,\lambda_n} \subset \mathcal{S}$ be a sequence of submodels as in (3.4) and $k_n := \lfloor \log_2(n\lambda_n/4) \rfloor$. We further assume

(3.11)
$$\liminf_{n \to \infty} \frac{n\lambda_n}{\log(n)} > 512 \text{ and } \lim_{n \to \infty} \frac{\log(\alpha_n \beta_{k_n, n})}{n\lambda_n} = 0$$

as well as

(1) for large scales, i.e. $\liminf_{n>0} \lambda_n > 0$, the limit $n\lambda_n \Delta_n^2 \log(1/(\alpha_n \beta_{k_n,n}))^{-1} \to \infty$,

(2) for small scales, i.e. $\lambda_n \to 0$, the inequality

(3.12)
$$\sqrt{n\lambda_n}\Delta_n \ge \left(\sqrt{512} + C + \epsilon_n\right)\sqrt{-\log(\lambda_n)}$$

with possibly $\epsilon_n \to 0$, but such that $\epsilon_n \sqrt{-\log(\lambda_n)} \to \infty$ and $\limsup_{n \to \infty} \frac{\sqrt{\log(8/(\alpha_n \beta_{k_n,n}))}}{\epsilon_n \sqrt{-\log(\lambda_n)}} < \frac{1}{\sqrt{512}},$

with C = 0 for K_n bounded and $C = 16\sqrt{6}$ for K_n unbounded.

Then,

$$\lim_{n \to \infty} \sup_{(\mu_n, \sigma_n^2) \in \mathcal{S}_{\Delta_n, \lambda_n}} \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(\hat{K}_n < K_n \right) = 0.$$

Theorems C.8 and 3.10 state conditions on the tuning parameters α_n and $\beta_{k_n,n}$ as well as on the length of the minimal scale $|I_n| =: \lambda_n$ (to simplify notations we only write λ_n in the following) and the standardized jump size Δ_n to detect the vanishing signals uniformly over $\mathcal{S}_{\Delta_n,\lambda_n}$. If, in addition, $\lim_{n\to\infty} \alpha_n = 0$ holds, then we control also the probability to overestimate the number of change-points and therefore the estimation of the number of change-points is still consistent in the case of a vanishing signal. The main condition in both theorems is that $\sqrt{n\lambda_n}\Delta_n$ has to be at least of order $\sqrt{-\log(\lambda_n)}$, see (C.9) and (3.12). This is optimal in the sense that no signal with a smaller rate can be detected asymptotically with probability one, see (Dümbgen and Spokoiny, 2001; Chan and Walther, 2013; Frick et al., 2014) for the case of homogeneous observations, and note that this is a sub-model of our model. But different to the homogeneous case we need, in addition, that λ_n is at least of order $\log(n)/n$, see (C.10) and (3.11). Such a restriction appears reasonable, since for the additional variance estimation only the number of observation on the segment is relevant and not the size of the change. Finally, we observe that the constants encountered in the lower detection bound for H-SMUCE in (C.9) and (3.12) increase with the difficulty of the estimation problem, where the difficulty is represented by the number of vanishing segments. All of these constants are a little bit larger as the analogue constants for SMUCE in (Frick et al., 2014, Theorem 5 and 6) reflecting the additional difficulty encountered by the heterogeneous noise. More precisely, we have 4 instead of the optimal $\sqrt{2}$ for one vanishing segment, $\sqrt{512}$ instead of 4 for a bounded number of vanishing segments and $\sqrt{512} + 16\sqrt{6}$ instead of 12 for an unbounded number of vanishing segments. Note again, that the optimal constants for the heterogeneous case are unknown to us.

3.4. Choice of the tuning parameters. In this section we discuss the choice of the tuning parameters α and $\beta_1, \ldots, \beta_{d_n}$.

Choice of α . As illustrated in Figure 2 the choice depends on the application. If a strict overestimation control of the number of change-points K is desirable α should be chosen small, e.g. 0.05 or 0.1, recall Theorems 3.3 and 3.4. This might come at the expense of missing change-points but with large probability not detecting too many (recall Figure 2 and see also the simulations in Section 4). If change-point screening is the primarily

goal, i.e. we aim to avoid missing of change-points, α should be increased, e.g. $\alpha = 0.5$ or even higher, since Theorem 3.5 shows that the error probability to underestimate the number of change-points decreases with increasing α . If model selection, i.e. $\hat{K} = K$, is the major aim, an intermediate level that balances the over- and underestimation error should be chosen, e.g. α between 0.1 and 0.5. Both errors vanish super polynomially for the asymptotic choice $\alpha = \alpha_n \in \exp(-o(n))$, see Remark 3.6. A finite sample approach is to weight these error probabilities $\gamma \mathbb{P}(\hat{K} > K) + (1 - \gamma)\mathbb{P}(\hat{K} < K)$, with $\gamma \in (0, 1)$, and to choose α such that its upper bound

$$\gamma \alpha + (1 - \gamma) \left(1 - \left[1 - 3 \exp\left(-\frac{1}{48} \left(\sqrt{\frac{n\lambda\Delta^2}{32}} - \sqrt{16\log\left(\frac{8}{\lambda\alpha\beta_k}\right)} \right)_+^2 \right) \right]_+^{2K} \right)$$

is minimized. This also allows to incorporate prior information on (λ, Δ) . Alternatively, the bound on the expectation $\mathbb{E}[|\hat{K} - K|]$ by combining Theorems 3.4 and 3.5 can be minimized to take the size of the missestimation into account. Despite of all possibilities to choose the 'best' α for a given application, comparing estimates at different α can be helpful to trace the "stability of evidence" of the estimated change-points at different significance levels. Of course, the interpretation of such a "significance screening" does not allow for a frequentist interpretation of a significance level anymore as α has to be fixed in advance, see e.g. (Schervish, 1996). Nevertheless, it might give for instance some indication whether to perform further experiments. Despite of this, for a fixed α the confidence statements of H-SMUCE can also be used to support findings by other estimators. This is illustrated in Section 5 for the ion channel application.

Choice of $\beta_1, \ldots, \beta_{d_n}$. As a default choice we recommend equal weights $\beta_1 = \cdots = \beta_{d_n} = 1/d_n$. This choice fulfils (together with many other choices) the conditions of the Theorems 3.7 and 3.8. Unlike as for the significance level α only the bound for the underestimation of the number of change-points depends on these weights. Note, that this gives the user the possibility to incorporate prior information on the scales without violating the overestimation control in Theorems 3.3 and 3.4: If for instance changes are expected to occur only on small segments then the detection power on these scales can be increased if the first weights are chosen large and the other ones small (or even zero). In contrast, if the general signal to noise ratio is expected to be very small then it is nearly impossible to detect changes on small scales and larger scales should be weighted more to detect at least the changes on these scales. A quantitative influence of the weights on the detection power can be seen in the underestimation bound in Theorem C.5 in Supplement C which is a refinement of Theorem 3.5. We also investigate such choices quantitatively in simulations in Section 4.

4. Simulations

In this section we compare H-SMUCE³ in simulations with CBS (Venkatraman and Olshen, 2007), cumSeg (Muggeo and Adelfio, 2011) and LOOVF (Arlot and Celisse, 2011) as they are also designed to be robust against heterogeneous noise. Moreover, we include SMUCE (Frick et al., 2014) in simulations with a constant variance as a benchmark to examine how much the detection power of H-SMUCE decreases in this case, which may be regarded as the price for adaptation to heterogeneous noise. We fix the weights $\beta_1, \ldots, \beta_{d_n} = 1/d_n$ and vary the significance level α . A simulation with tuned weights can be found in Section B.2 in the Supplement. For circular binary segmentation (CBS) we call the function segmentByCBS⁴ with the standard parameters. For the cross-validation method LOOVF we use the Matlab function proc_LOOVF⁵ with the parameter choice of the demo file. For cumSeg we call the method jumpoints⁶ with the parameter k large enough such that the estimation is not influenced by this choice. For SMUCE we call the function smuceR⁷ with the standard parameters, in particular the interval set of all intervals is used if $n \leq 1000$.

To avoid specific interactions between the signal and the dyadic partition we generate in each repetition a random pair $(\mu_R, \sigma_R^2) \in \mathcal{S}$ (all random variables are independent from each other).

- (a) We fix the number of observations n, the number of change-points K, a constant C and a minimum value for the smallest scale λ_{\min} .
- (b) We draw the locations of the change-points $\tau_0 := 0 < \tau_1 < \cdots < \tau_K < 1 =: \tau_{K+1}$ uniformly distributed with the restriction that $\lambda := \min_{k=0,\dots,K} |\tau_{k+1} - \tau_k| \ge \lambda_{\min}$.
- (c) We choose the function values s_0, \ldots, s_K of the standard deviation function σ_R by $s_k := 2^{U_k}$, where U_0, \ldots, U_K are uniform distributed on [-2, 2].
- (d) We determine the function values m_0, \ldots, m_K of the signal μ_R such that

(4.1)
$$|m_k - m_{k-1}| = \sqrt{\frac{C}{n} \min\left(\frac{\tau_{k+1} - \tau_k}{s_k^2}, \frac{\tau_k - \tau_{k-1}}{s_{k-1}^2}\right)^{-1}} \quad \forall \ k = 1, \dots, K$$

Thereby, we start with $m_0 = 0$ and choose randomly with probability 1/2 whether the expectation increases or decreases.

By (4.1) we provide a situation where all change-points are similarly hard to find, recall the minimax detection boundary from Section 3.3. An example has been displayed in Figure 2 in the introduction, where H-SMUCE misses at $\alpha = 0.1$ one change-point and

³http://www.stochastik.math.uni-goettingen.de/hsmuce, v. 0.0.0.9000, 2015-04-15

⁴http://cran.r-project.org/web/packages/PSCBS/, v. 0.40.4, 2014-02-04

⁵http://www.di.ens.fr/~arlot/code/CHPTCV.htm, v. 1.0, 2010-10-27

⁶http://cran.r-project.org/web/packages/cumSeg/, v. 1.1, 2011-10-14

⁷http://cran.r-project.org/web/packages/stepR, v. 1.0-3, 2015-06-18

detects for α between 0.15 and 0.99 (only displayed for $\alpha = 0.3$ and $\alpha = 0.5$) the correct number of change-points. In Figure 9 (Supplement B) we see that CBS (Venkatraman and Olshen, 2007) finds also all change-points, but detects further changes. Less good is the performance of cumSeg (Muggeo and Adelfio, 2011) and LOOVF (Arlot and Celisse, 2011) which both miss several changes and LOOVF adds also a false positive. We examine these methods now more extensively. All simulations are repeated 10 000 times.

In the following we report the difference between the estimated \hat{K} and the true number K of change-points as well as the mean of the absolute value of this difference. Additionally, we use the false positive sensitive location error

$$\text{FPSLE} = \frac{n}{2\hat{K}} \sum_{k=1}^{\hat{K}+1} |\tau_{l_k-1} - \hat{\tau}_{k-1}| + |\tau_{l_k} - \hat{\tau}_k|$$

with $l_k \in \{1, \ldots, K+1\}$ such that $(\hat{\tau}_{k-1} + \hat{\tau}_k)/2 \in (\tau_{l_k-1}, \tau_{l_k}]$, i.e. the left and right neighbouring change-points to the middle point of $(\hat{\tau}_{k-1}, \hat{\tau}_k]$, and the false negative sensitive location error

FNSLE =
$$\frac{n}{2K} \sum_{k=1}^{K+1} |\tau_{k-1} - \hat{\tau}_{l_k-1}| + |\tau_k - \hat{\tau}_{l_k}|,$$

with $l_k \in \{1, \ldots, \hat{K} + 1\}$ such that $(\tau_{k-1} + \tau_k)/2 \in (\hat{\tau}_{l_k-1}, \hat{\tau}_{l_k}]$, see (Futschik et al., 2014, Section 3.1), to rate the estimation of the locations of the change-points. We also show the mean integrated squared (absolute) error MISE (MIAE) for all methods.

4.1. Simulation results. In this section we discuss the results of the simulations for model (1.2) and (1.3). We start in Table 1 (Supplement B) with the simple setting of a single change at the midpoint, where we vary the variances on the adjoining segments. In Table 2 (Supplement B) we display results for a constant variance and in Table 3 (Supplement B) for heterogeneous errors. We excluded LOOVF from simulations for larger n due to its large computation time, confer the run time simulations in Section A.3 in the supplement.

All simulations confirm the overestimation control α for H-SMUCE from Theorem 3.3 and the exponential decay of the overestimation in Theorem 3.4. The simulations with a single change-point confirm that the size of the variance change has no influence, rather the size of the variances matters. We found that H-SMUCE performs well compared to all other methods. A small α avoids overestimation, but risks to miss changes that are harder to detect. Thus, the comparison of the estimates of H-SMUCE for different α shows in accordance with our theory that it is reasonable to relax α if changes are expected to be harder to detect (recall the discussion in Section 3.4). From the other methods cumSeg performs best in the easier and LOOVF in the difficult scenarios, whereby CBS and in particular LOOVF shows a tendency to overestimate the number of change-points.

For a constant signal (corresponding to K = 0 in Table 2) H-SMUCE overestimates the

number of change-points even slightly less than SMUCE, whereas CBS and cumSeg overestimate hardly ever. In the case of a constant variance we found that the detection power of H-SMUCE is only slightly worse than SMUCE for K = 2, although SMUCE used instead of the dyadic partition \mathcal{D} the system of all intervals. The difference is larger for K = 10 and in this case also CBS and cumSeg performs better than H-SMUCE, since the detection power of H-SMUCE depends strongly on the lengths of the constant segments. Moreover, λ_{\min} plays a similar role as the number of change-points K, since the average constant segments length decreases if λ_{\min} decreases or K increases. Worse results for smaller lengths are due to the familywise error control α of H-SMUCE as it guarantees a strict control of overestimating the number of change-points.

Similar results can be observed for n = 100 with heterogeneous errors. CBS performs better than cumSeg and LOOVF, and in particular better than in the single change-point setting. CBS outperforms H-SMUCE for K = 5, although H-SMUCE has a much smaller tendency to overestimate the number of change-points, whereas in particular CBS and LOOVF tend to overestimation. This can also be seen for the MISE and MIAE as these measures are much more affected by underestimation than by overestimation. These findings are also supported by the FPSLE and the FNSLE, the FPSLE is heavily affected by overestimation, whereas the FNSLE is larger in case of underestimation.

In all simulations with heterogeneous errors and 1000 observations H-SMUCE outperforms the other methods, for 10000 observations this becomes even more pronounced. In comparison to the simulation with 100 observations the tendency of CBS to overestimate the number of change-points becomes then also more prominent. Finally, in further simulations (not displayed) we found that the detection power of all methods decreases for smaller C in (d), but all results remain qualitatively the same. All in all, we found that H-SMUCE performs well as sample size becomes larger, in particular if the constant segments are not too short as indicated by assumption (3.11) in Theorem 3.10.

A comparison of Table 3 and 4 (Supplement B) shows that tuned weights increase the detection power of H-SMUCE for all significance levels, so we encourage the user to adapt the weights if prior information on the scales where changes occur is available. Details how the weights are chosen can be found in Section B.2 in the supplement.

4.2. Robustness against model violations. We begin by investigating how robust the methods are against a violation of the assumption that the standard deviation changes only at the same locations as the mean changes. We consider continuous changes as well as abrupt changes. The exact functions for the standard deviation can be seen in Figure 10 (Supplement B). In Table 5 (Supplement B) we see that H-SMUCE and CBS perform very robust against heterogeneous noise on the constant segments, whereas, remarkably, the detection power of cumSeg is even improved. Moreover, in additional simulations (not displayed) with less observations we found that LOOVF is very robust, too.

Moreover, we examine robustness against small periodic trends in the mean in simulations similar to those in (Venkatraman et al., 2004), also adapted to the inhomogeneous variance. The exact simulation setting can be found in Section B.3 (Supplement B). We obtain from Table 6 that H-SMUCE shows similar results for small trends compared to the simulation without trend for small trends, but is affected by larger trends, in particular if these are not scaled by the standard deviation. CBS overestimates heavily in all cases, whereas cumSeg (although not affected by the trend) shows over- and underestimation. Furthermore, we investigate robustness against heavy tails of the error distribution. In Table 7 (Supplement B) we consider t_3 -distributed errors which are scaled such that the expectation and the standard deviation are the same as in Section 4.1. As expected SMUCE is not robust against heavy tails (as it misinterprets extreme values as a change in the signal, whereas H-SMUCE provides reasonable results. In comparison to gaussian errors H-SMUCE is not influenced for K = 0, underestimation is more distinct in the constant variance scenario and detection power is even increased in the scenario with heterogeneous errors. In comparison, CBS is not influenced for K = 0, too, underestimates and overestimates in the constant variance scenario and is slightly worse with a tendency to underestimation in the scenario with heterogeneous errors, whereas cumSeg overestimates rarely, but heavily for K = 0, underestimates and overestimates in the constant variance scenario and is robust in the last scenario.

In summary, H-SMUCE seems to be robust against a wide range of variance changes on constant segments and seems to be only slightly affected by larger tails than gaussian, in particular no tendency to overestimation was visible in our simulations. This may be explained by the fact that the local likelihood tests of H-SMUCE are quite robust against heterogeneous noise, see for instance (Bakirov and Szekely, 2006; Ibragimov and Müller, 2010), and against non-normal errors, see (Lehmann and Romano, 2005) and the references therein. Unlike the number of change-points, the locations are sometimes missestimated, since the restricted maximum likelihood estimator is influenced by changes of the variance. Instead, more robust estimators, for instance local median and MAD estimators, could be used.

5. Application to ion channel recordings

In this section we apply H-SMUCE to current recordings of a porin in planar lipid bilayers performed in the Steinem lab (Institute of Organic and Biomolecular Chemistry, University of Göttingen). Porins are β -barrel proteins present in the outer membrane of bacteria and in the outer mitochondrial membrane of eukaryotes (Benz, 1994; Schirmer, 1998). Due to their large pore diameter they enable passive diffusion of small solutes like ions or sugars. The partial blockade of the pore by an internal loop results in gating that can be detected using the voltage clamp technique (Sakmann and Neher, 1995). We aim to detect the gating automatically, since in many ion channel applications hundred or more datasets each with several hundredthousands data points have to be analysed. For noise reduction the data was automatically preprocessed in the amplifier with an analogue four-pole Bessel low-pass filter of 1 kHz. Hence, the noise is coloured, but the correlation is less than 10^{-3} if the trace is subsampled by eleven or more observations, see (Hotz et al., 2013, (6)), which has been done in the following. Finally, we apply to 882 subsampled observations H-SMUCE, CBS, cumSeg and LOOVF.

In Figure 3a we see that the signal fluctuates around two or more levels, the so called open (higher conductivity, larger current measurements) and closed (lower conductivity, smaller current measurements) states. Moreover, the variance in the open states is larger than in the closed state, a well known phenomenon denoted as open channel noise (Sakmann and Neher (1995, Section 3.4.4) and the references therein) which arises for larger ion channels such as porins from conformational fluctuations in the channel protein (Sigworth, 1985). Due to the pronounced heterogeneity in the variance, methods which assume a constant variance fail to reconstruct the gating, see Figure 1 in the introduction for an illustration. In contrast, H-SMUCE at $\alpha = 0.05$ provides a reasonable fit that covers the main features of the data. Additional smaller changes are found by CBS, cumSeg and LOOVF, see Figure 3c. These changes might be explained by some uncontrollable base line fluctuations caused for instance by small holes in the membrane due to movements of the lipids. On the other hand, we found in the simulations, see Table 3 and the example in Figure 9 (both Supplement B), that CBS and LOOVF tend to include small artificial changes, whereas we saw in Table 6 (Supplement B) that H-SMUCE is quite robust against small periodic trends in the signal. For illustrative purposes, in order to examine these changes further we increase in Figure 3b the significance level $\alpha = 0.5$ and detect for instance changes around 33.0s and 33.2s, too. Taking also the confidence regions of H-SMUCE into account confirms several changes with high "significance" (e.g. the reconstruction of CBS between 33.5 and 33.8) and further changes with less "significance" (e.g. the changes around 33.0s and 33.2s). Other changes could not be confirmed by H-SMUCE at any reasonable significance level (e.g. the peaks of CBS and LOOVF at 33.85). In this spirit H-SMUCE can always be used to accompany any segmentation method to help to identify its significant changes. Recall, that of course a frequentist statistical error control is only given when α is fixed in advance.

Acknowledgement

Support of DFG CRC 803 - Z2 and FOR 916 - B3 is gratefully acknowledged. We are grateful to various comments by two referees and an associate editor which lead to a significant improvement of the paper. We thank A. Bartsch, O. Schütte and C. Steinem (Institute of Organic and Biomolecular Chemistry, University of Göttingen) for providing



(c) Estimates by CBS (green), cumSeg (purple) and LOOVF (blue).

FIGURE 3. a, b: Subsampled observations (black points) together with the confidence band (grey), the confidence intervals for the change-point locations (brackets and thick lines), the estimated change-points locations (red dashes) and the estimate (red line) by H-SMUCE at different α . c: other estimates.

HETEROGENEOUS CHANGE POINT INFERENCE

24

the ion channel data and helpful discussions. We also thank T. Aspelmeier, M. Behr, A. Hartmann, H. Li and I. Tecuapetla for helpful comments to the paper and the R-package.

Supplement to Heterogeneous Change Point Inference by Florian Pein, Hannes Sieling and Axel Munk

APPENDIX A. COMPUTATION

In this section we detail the computation of the estimator H-SMUCE (Section A.1) and of the critical values q_1, \ldots, q_{d_n} (Section A.2). We also examine the computation time (Section A.3) theoretically and empirically. An R-package is available online⁸.

A.1. Computation of the estimator. First of all, we obtain from the multiscale test the bounds

(A.1)
$$[\underline{b}_{i,j}, \overline{b}_{i,j}] := \left[\overline{Y}_{ij} - \sqrt{\frac{q_{ij}\hat{s}_{ij}^2}{j-i+1}}, \overline{Y}_{ij} + \sqrt{\frac{q_{ij}\hat{s}_{ij}^2}{j-i+1}}\right]$$

for μ on the interval $[i/n, j/n] \in \mathcal{D}$. Therefore, H-SMUCE can be computed as in (Frick et al., 2014, Section 3) for SMUCE described. However, in what follows we give a modification of the algorithm which reduces the computation time remarkably due to the small number of intervals $\mathcal{O}(n)$ in the dyadic partition \mathcal{D} . Here, we compute first left and right limits for the location of the change-points and then start the dynamic program restricted to these intervals. A notable difference to (Killick et al., 2012; Frick et al., 2014) is that this approach leads also to pruning in the forward step of the dynamic program. More precisely, we define the intersected bounds as

$$\underline{B}_{i,j} := \max_{\substack{i \le s < t \le j \\ [s/n,t/n] \in \mathcal{D}}} \underline{b}_{s,t} \text{ and } \overline{B}_{i,j} := \min_{\substack{i \le s < t \le j \\ [s/n,t/n] \in \mathcal{D}}} \overline{b}_{s,t}$$

and set recursively

$$L_k := \min \left\{ 1 < r \le L_{k+1} - 1 : \underline{B}_{r, L_{k+1} - 1} \le \overline{B}_{r, L_{k+1} - 1} \right\},\$$

for $k = \hat{K}, \ldots, 1$, with $L_{\hat{K}+1} := n + 1$. The right limits are defined as

$$R_k := \min \left\{ R_{k-1} < r \le n : \underline{B}_{R_{k-1},r} > \overline{B}_{R_{k-1},r} \right\},$$

for $k = 1, \ldots, \hat{K}$, with $R_0 := 1$. In other words, the left limit for the k-th change-point L_k is the smallest number $1 < r \leq n$ such that between Y_r and Y_n a piecewise constant solution with $\hat{K} - k$ change-points exists which respects the bounds (A.1). Analogously, the right limit R_k is the smallest number $1 < r \leq n$ such that between Y_1 and Y_r no piecewise constant solution with k-1 change-points exists which fulfils the bounds (A.1). Note, that we do not have to compute the right limits separately, since we can just start the dynamic program at L_k and stop if another change-point has to be included. It

⁸http://www.stochastik.math.uni-goettingen.de/hsmuce

follows that the k-th change-point $\hat{\tau}_k$ has to be in the confidence interval $[L_k/n, R_k/n]$, since otherwise an additional change-point would be necessary to fulfil the multiscale constraints.

A.2. Computation of the critical values. In this section we show how the critical values can be computed by Monte-Carlo simulations. Note first that the following method uses only the continuity and the monotonicity of the cumulative distribution functions of the statistics T_1, \ldots, T_{d_n} and therefore the methodology can also be used for other multiscale tests, see for instance the extension to other interval sets in Remark 2.2.

Let M be the number of simulations and $(T_{1,1}, \ldots, T_{d_n,1}), \ldots, (T_{1,M}, \ldots, T_{d_n,M})$ be i.i.d. copies of the vector (T_1, \ldots, T_{d_n}) . Moreover, we denote by $F_M(\cdot)$ the empirical distribution function of (T_1, \ldots, T_{d_n}) and by $F_{M,k}(\cdot)$ the empirical distribution function of the random variable T_k . Then, we aim to find a vector of critical values $\widehat{\mathbf{q}}_M = (\widehat{q}_{M,1}, \ldots, \widehat{q}_{M,d_n})$ which satisfies with

(A.2)
$$\alpha - \frac{1}{M} < 1 - F_M(\widehat{\mathbf{q}}_M) \le \alpha,$$

an empirical version of condition (2.3), and with

(A.3)
$$\frac{1 - F_{M,j_1}(\widehat{q}_{M,j_1})}{\beta_{j_1}} \le \frac{1 - F_{M,j_2}(\widehat{q}_{M,j_2}) + \frac{1}{M}}{\beta_{j_2}} \quad \text{for all} \quad j_1, j_2 \in \{1, \dots, d_n\}$$

an empirical version of condition (2.5). In the following we propose an iterative method to determine such a vector and show afterwards that this vector converges almost surely to the vector of critical values defined by (2.3) and (2.5). As the k-th entry of the starting vector we choose the empirical $(1 - \alpha \beta_k)$ -quantile of the statistic T_k , since the vector with these values satisfies condition (A.3) and the inequality

$$1 - F_M(\cdot) \le \alpha.$$

Afterwards, we reduce the entries until the lower bound from condition (A.2) is satisfied, too. To ensure condition (A.3) in every iteration, we always reduce the entry which has the smallest ratio

$$\frac{1-F_{M,k}(\widehat{q}_{M,k})}{\beta_k}.$$

In Algorithm 1 the determination of the critical values is summarized in pseudocode. The method has the advantage that we do not need specific assumptions on the distribution of the vector (T_1, \ldots, T_{d_n}) and still get critical values which are adapted to the exact finite sample distribution of (T_1, \ldots, T_{d_n}) and ensure therefore even for a finite number of observations the significance level α .

The following theorem shows the convergence of this algorithm to $\mathbf{q} = (q_1, \ldots, q_{d_n})$.

Algorithm 1 Determination of the critical values.

Input: The statistics T_1, \ldots, T_{d_n} as well as the significance level $\alpha \in (0, 1)$, the weights $\beta_1, \ldots, \beta_{d_n} > 0$, with $\sum_{k=1}^{d_n} \beta_k = 1$, and the number of simulations $M \in \mathbb{N}$.

Output: The vector of critical values $\widehat{\mathbf{q}}_M = (\widehat{q}_{M,1}, \dots, \widehat{q}_{M,d_n})$ which fulfils the conditions (A.2) and (A.3).

1: for i = 1, ..., M do $(T_{1,i},\ldots,T_{d_n,i}) \leftarrow \text{ realisation of } (T_1,\ldots,T_{d_n})$ 2: 3: end for 4: for $k = 1, ..., d_n$ do $(S_{k,1},\ldots,S_{k,M}) \leftarrow \operatorname{sort}((T_{k,1},\ldots,T_{k,M}))$ 5: $w_k \leftarrow M - |\alpha \beta_k M|$ 6: 7: end for 8: repeat $\hat{k} \leftarrow \operatorname{argmin}_{k=1,\dots,d_n} \beta_k^{-1} \left(1 - F_{M,k}(S_{k,w_k}) \right)$ $w_{\hat{k}} \leftarrow w_{\hat{k}} - 1$ 9: 10:11: **until** $1 - F_M(S_{1,w_1}, \dots, S_{m,w_{d_n}}) > \alpha$ 12: $w_{\hat{k}} \leftarrow w_{\hat{k}} + 1$ 13: return $S_{1,w_1}, \ldots, S_{m,w_{d_n}}$

Theorem A.1 (Considency of Monte-Carlo critical values). The empirical vector of critical values $\widehat{\boldsymbol{q}}_M = (\widehat{q}_{M,1}, \ldots, \widehat{q}_{M,d_n})$ converges almost surely in the number of simulations M to the vector of critical values $\boldsymbol{q} = (q_1, \ldots, q_{d_n})$ defined by (2.3) and (2.5).

The computation time is dominated by the generation of the M i.i.d. copies of the vector (T_1, \ldots, T_{d_n}) . Therefore, we store the generated realizations and recycle them. To avoid memory problems we only store the realizations for every dyadic number, because the significance level α is still satisfied if we determine the critical values based on realizations with a larger number of observations, since then the maxima in (T_1, \ldots, T_{d_n}) are taken over more intervals. To this end, the choice $M = 10\,000$ seems to be a good trade-off between computation time and approximation accuracy.

A.3. Computation time. In this section we discuss the theoretical computation time of H-SMUCE and compare it later in simulations with CBS, cumSeg and LOOVF. We stress that the computation time for the bounds, for the limits $L_1, \ldots, L_{\hat{K}}$ (and so for \hat{K}) and for the optimization problem (1.4), and therefore of all confidence sets, is always $\mathcal{O}(n)$. Hence, the computation time is dominated by the determination of the restricted maximum likelihood estimator by dynamic programming.

Lemma A.2 (Computation time). The algorithm has data depended computation time

(A.4)
$$\mathcal{O}\left(n + \sum_{k=1}^{\hat{K}-1} (R_k - L_k + 1)(R_{k+1} - L_{k+1} + 1)\right).$$

This can be bounded by $\mathcal{O}(n^2)$ in the worst case, but the computation time is in many cases much smaller. In particular, if the signal to noise ratios are large enough such that the change-points are easy to detect, i.e. $R_k - L_k$ is small. This is for instance the case for a fixed signal, where $R_k - L_k$ stays more or less constant. More precisely, by combining (A.4) with equation (3.10) we see that with probability tending to one the computation time of H-SMUCE is even linear, if $\alpha_n \to 0$, but $n^{-\frac{1}{2}} \log((\alpha_n \beta_{k_n,n})^{-1}) \to 0$. In comparison to the computation time of SMUCE, see (Sieling, 2013, (4.3)), which is dominated by the term

$$\mathcal{O}\left(\sum_{k=1}^{\hat{K}-1} (R_k - R_{k-1})(R_{k+1} - R_k)\right),\$$

we see that the computation time is further reduced. In particular, if no change-point is present the computation time is $\mathcal{O}(n)$ instead of $\mathcal{O}(n^2)$. The computation time is also $\mathcal{O}(n)$ if the number of change-points increases linear in the number of observations and the change-points are evenly enough distributed.

In the following we examine the computation time empirically in a similar simulation study as in (Maidstone and Pickering, 2014). More precisely, we generate data with varying number of observations n and equidistant change-points. Thereby, we consider $K = 10, K = \sqrt{n}$ and K = n/100. In all scenarios we choose the values of the mean and the standard deviation function randomly like in Section 4, once again with C = 200. All simulations are repeated 100 times and terminated after ten seconds. The simulations were performed on a single core system with 1.8 GHz and 8 GB RAM in a 64-bit OS.

We fix the significance level $\alpha = 0.1$ as well as the weights $\beta_1 = \cdots = \beta_{d_n} = 1/d_n$ and compare H-SMUCE with CBS, LOOVF and cumSeg. Note, that we restore the Monte-Carlo simulations at the first use to reduce further loading times, here we only take the already restored simulations into account. Furthermore, we set for cumSeg the maximal number of change-points $k = \max(2K, 10)$, since for the default parameter $k = \min(30, n/10)$ the program requires manual increase of k for many simulations runs. Note, that the choice above already incorporates prior knowledge about the true signal. We stress (not displayed) that the computation time (and the required memory space) increases severely in the parameter k.

From Figure 8 we draw that H-SMUCE is much faster than the other methods, in particular if the number of change-points increases. For K = n/100 the computation time increases almost linearly in the number of observations. For example, when $n = 10^7$ it is still less than a minute. The second shortest computation time has CBS for larger numbers of observations, whereas cumSeg is superior for smaller numbers of observations. The computation time of CBS for $n = 10^5$ observations is still less than a minute in all scenarios, whereas cumSeg has a similar computation time for K = 10, but lasts several



FIGURE 8. Mean computation time of H-SMUCE (red crosses), CBS (green triangles), cumSeg (purple circles) and LOOVF (blue squares) for different number of observations n and different number of change-points K. Note that for purposes of visualization the x-axis is displayed non-equidistantly.

minutes in the other cases. Lastly, LOOVF exceeds ten seconds already for n = 400 observations and is always found to be the slowest method.

APPENDIX B. ADDITIONAL FIGURES AND TABLES

In this section we collect additional figures and tables.

B.1. Simulations. We start with estimates by CBS, cumSeg and LOOVF for the data from Figure 2.

The following three tables collect the results of the simulations in Section 4.1. Recall the random pair $(\mu_R, \sigma_R^2) \in \mathcal{S}$ (all random variables are independent from each other):

- (a) We fix the number of observations n, the number of change-points K, a constant C and a minimum value for the smallest scale λ_{\min} .
- (b) We draw the locations of the change-points $\tau_0 := 0 < \tau_1 < \cdots < \tau_K < 1 =: \tau_{K+1}$ uniformly distributed with the restriction that $\lambda := \min_{k=0,\dots,K} |\tau_{k+1} - \tau_k| \ge \lambda_{\min}$.
- (c) We choose the function values s_0, \ldots, s_K of the standard deviation function σ_R by $s_k := 2^{U_k}$, where U_0, \ldots, U_K are uniform distributed on [-2, 2].
- (d) We determine the function values m_0, \ldots, m_K of the signal μ_R such that

$$|m_k - m_{k-1}| = \sqrt{\frac{C}{n} \min\left(\frac{\tau_{k+1} - \tau_k}{s_k^2}, \frac{\tau_k - \tau_{k-1}}{s_{k-1}^2}\right)^{-1}} \ \forall \ k = 1, \dots, K$$

Thereby, we start with $m_0 = 0$ and choose randomly with probability 1/2 whether the expectation increases or decreases.

All simulations are repeated 10000 times.

B.2. Prior information on scales. To demonstrate the effect of incorporating prior knowledge about those scales where change-points are likely to happen we consider again the observations from Table 3 with $n = 10\,000$, K = 10 and $\lambda_{\min} = 50$. To this end, we use the adapted weights, where we eliminate the smallest three scales k = 1, 2, 3, since all constant segments contain at least 50 observations and therefore these small scales are not needed for detection. Moreover, we choose $\tilde{\beta}_4 = 1/4$, $\tilde{\beta}_5 = 1/4$, $\tilde{\beta}_6 = 1/6$, $\tilde{\beta}_7 = 1/6$, $\tilde{\beta}_8 = 1/12$, $\tilde{\beta}_9 = 1/12$ in decreasing order, since change-points on smaller scales are more likely and harder to detect. For the same reasons we eliminate the four largest scales k = 10, 11, 12, 13, too.

A comparison of Table 3 and 4 shows that the modified weights increase the detection power of H-SMUCE for all significance levels, so we encourage the user to adapt the weights if prior information on the scales where changes occur is available.



FIGURE 9. Observations (black points) and true signal (black line) together with estimates by CBS, cumSeg and LOOVF for the data from Figure 2. All parameters are chosen as described in Section 4.

HETEROGENEOUS CHANGE POINT INFERENCE

Setting	Method	-1	0	+1	$\geq +2$	$ \hat{K} - K $	FPSLE	FNSLE	MISE	MIAE
$\sigma_0 = 0.5,$	HS(0.1)	0.000	0.995	0.004	0.000	0.005	0.82	0.74	0.0119	0.0644
$\sigma_1 = 0.5,$	HS(0.3)	0.000	0.975	0.025	0.000	0.025	1.38	0.95	0.0129	0.0672
	HS(0.5)	0.000	0.929	0.070	0.001	0.072	2.67	1.40	0.0144	0.0706
	CBS	0.000	0.949	0.036	0.015	0.066	2.31	0.94	0.0128	0.0660
	cumSeg	0.000	0.995	0.005	0.000	0.005	1.37	1.28	0.0172	0.0707
	LOOVF	0.000	0.774	0.142	0.084	0.378	10.10	2.42	0.1402	0.2897
$\sigma_0 = 0.5,$	HS(0.1)	0.112	0.886	0.002	0.000	0.114	3.99	6.77	0.0543	0.1405
$\sigma_1 = 1,$	HS(0.3)	0.020	0.961	0.019	0.000	0.039	2.38	2.56	0.0321	0.1086
	HS(0.5)	0.005	0.940	0.054	0.001	0.061	3.12	2.30	0.0314	0.1090
	CBS	0.042	0.873	0.068	0.017	0.147	5.87	4.93	0.0496	0.1315
	cumSeg	0.008	0.969	0.021	0.003	0.034	3.09	2.78	0.0375	0.1126
	LOOVF	0.006	0.791	0.112	0.091	0.373	11.30	3.90	0.1720	0.3004
$\sigma_0 = 0.5,$	HS(0.1)	0.484	0.515	0.001	0.000	0.485	12.77	24.89	0.1736	0.3110
$\sigma_1 = 1.5,$	HS(0.3)	0.209	0.778	0.012	0.000	0.222	6.81	11.92	0.1025	0.2075
	HS(0.5)	0.089	0.872	0.039	0.000	0.129	4.92	6.54	0.0725	0.1690
	CBS	0.417	0.454	0.105	0.024	0.577	17.63	25.40	0.1845	0.3385
	cumSeg	0.231	0.731	0.032	0.006	0.276	9.60	14.62	0.1149	0.2317
	LOOVF	0.135	0.683	0.098	0.085	0.490	15.78	11.92	0.2307	0.3322
$\sigma_0 = 1,$	HS(0.1)	0.453	0.547	0.001	0.000	0.453	13.49	24.97	0.1514	0.3140
$\sigma_1 = 1,$	HS(0.3)	0.171	0.818	0.011	0.000	0.182	8.11	12.53	0.0942	0.2170
	HS(0.5)	0.062	0.900	0.038	0.000	0.101	6.75	7.99	0.0745	0.1847
	CBS	0.156	0.744	0.091	0.008	0.265	9.51	11.41	0.0943	0.2127
	cumSeg	0.120	0.876	0.004	0.000	0.124	5.93	8.88	0.0748	0.1839
	LOOVF	0.039	0.749	0.132	0.081	0.405	13.29	6.87	0.1947	0.3472
$\sigma_0 = 1,$	HS(0.1)	0.727	0.272	0.000	0.000	0.728	19.44	37.71	0.2237	0.4244
$\sigma_1 = 1.5,$	HS(0.3)	0.410	0.584	0.006	0.000	0.416	13.35	23.77	0.1644	0.3256
	HS(0.5)	0.218	0.753	0.028	0.000	0.247	10.25	15.64	0.1283	0.2669
	CBS	0.491	0.406	0.096	0.008	0.604	18.00	28.42	0.2013	0.3741
	cumSeg	0.409	0.580	0.010	0.000	0.420	12.91	22.99	0.1571	0.3155
	LOOVF	0.184	0.638	0.105	0.072	0.501	16.55	14.92	0.2410	0.3626
$\sigma_0 = 1.5,$	HS(0.1)	0.844	0.156	0.000	0.000	0.844	22.41	43.65	0.2581	0.4713
$\sigma_1 = 1.5,$	HS(0.3)	0.574	0.423	0.003	0.000	0.577	18.21	33.12	0.2219	0.4101
	HS(0.5)	0.352	0.629	0.018	0.000	0.371	15.47	25.01	0.1915	0.3582
	CBS	0.659	0.258	0.079	0.003	0.746	20.73	35.81	0.2449	0.4379
	cumSeg	0.629	0.369	0.002	0.000	0.631	17.56	33.32	0.2147	0.4067
	LOOVF	0.297	0.534	0.104	0.066	0.589	19.34	21.26	0.2715	0.4046

TABLE 1. Simulations with a single change (fixed signal and variances): n = 100 observations and a single change at 0.5, from 0 to 1 for different standard deviations changing from σ_0 to σ_1 at 0.5, too. Columns from left to right: setting, method, proportions of $\hat{K} - K$ and averages of the corresponding error criteria. $\text{HS}(\alpha)$ denotes H-SMUCE at significance level α .

B.3. Robustness. Figure 10 shows the standard deviation functions in Table 5 to examine robustness against variance changes on constant segments. We consider the sinusshaped standard deviation σ_1 (continuous changes), the piecewise linear standard deviation σ_2 (continuous and abrupt changes at the same time) and the piecewise constant standard deviation σ_3 (abrupt changes). Moreover, we analyse in Table 6 robustness against small periodic trends in simulations similar to those in (Venkatraman et al., 2004). More precisely, we generate the random pairs (μ_R, σ_R^2) $\in S$ as in (a)-(d) described, but

Setting	Method	≤ -2	-1	0	+1	$\geq +2$	$ \hat{K} - K $	FPSLE	FNSLE	MISE	MIAE
n = 1000,	HS(0.1)	-	-	0.965	0.035	0.000	0.035	17.75	4.73	0.0035	0.0365
$\mathbf{K}=0,$	HS(0.3)	-	-	0.867	0.128	0.005	0.138	68.95	18.42	0.0045	0.0401
$\mu = \mu_R \equiv 0,$	HS(0.5)	-	-	0.719	0.256	0.025	0.307	153.45	41.25	0.0061	0.0454
$\sigma = \sigma_R$	S(0.1)	-	-	0.965	0.034	0.001	0.036	17.90	5.03	0.0039	0.0371
$\equiv \text{const}$	S(0.3)	-	-	0.832	0.160	0.008	0.177	88.45	24.80	0.0059	0.0435
	S(0.5)	-	-	0.667	0.298	0.035	0.370	184.90	50.94	0.0082	0.0499
	CBS	-	-	0.991	0.000	0.009	0.018	8.90	1.26	0.0037	0.0351
	cumSeg	-	-	0.999	0.001	0.000	0.001	0.30	0.06	0.0029	0.0345
n = 1000,	HS(0.1)	0.010	0.174	0.802	0.014	0.000	0.208	26.32	72.66	0.0132	0.0613
K = 2,	HS(0.3)	0.004	0.108	0.819	0.067	0.002	0.187	38.10	52.90	0.0114	0.0571
$\lambda_{\min} = 30,$	HS(0.5)	0.002	0.070	0.768	0.150	0.010	0.244	64.14	48.50	0.0111	0.0573
$\mu = \mu_R,$	S(0.1)	0.003	0.074	0.912	0.011	0.000	0.092	16.96	34.03	0.0092	0.0513
$\sigma \equiv 1$	S(0.3)	0.001	0.040	0.892	0.065	0.002	0.112	32.24	27.39	0.0090	0.0513
	S(0.5)	0.001	0.025	0.806	0.155	0.013	0.209	63.30	32.33	0.0095	0.0536
	CBS	0.005	0.060	0.821	0.082	0.033	0.221	37.55	37.57	0.0111	0.0527
	cumSeg	0.025	0.116	0.749	0.099	0.011	0.289	65.32	82.63	0.0364	0.0738
n = 1000,	HS(0.1)	0.009	0.160	0.815	0.015	0.000	0.194	27.14	68.91	0.0127	0.0611
K = 2,	HS(0.3)	0.004	0.098	0.829	0.067	0.001	0.176	37.77	49.63	0.0111	0.0572
$\lambda_{\min} = 50,$	HS(0.5)	0.002	0.063	0.774	0.152	0.009	0.237	63.46	46.06	0.0109	0.0573
$\mu = \mu_R,$	S(0.1)	0.003	0.068	0.919	0.009	0.000	0.084	16.82	31.94	0.0091	0.0515
$\sigma \equiv 1$	S(0.3)	0.001	0.035	0.899	0.063	0.002	0.104	31.19	25.81	0.0090	0.0515
	S(0.5)	0.001	0.020	0.819	0.147	0.013	0.195	59.86	30.23	0.0095	0.0537
	CBS	0.005	0.058	0.824	0.083	0.031	0.215	37.50	36.27	0.0112	0.0532
	cumSeg	0.023	0.110	0.769	0.090	0.008	0.262	59.74	79.25	0.0336	0.0741
n = 1000,	HS(0.1)	0.508	0.330	0.161	0.001	0.000	1.634	54.37	172.66	0.1112	0.1842
K = 10,	HS(0.3)	0.354	0.377	0.263	0.006	0.000	1.233	44.53	127.81	0.0817	0.1561
$\lambda_{\min} = 30,$	HS(0.5)	0.253	0.384	0.346	0.017	0.000	0.987	40.88	102.88	0.0679	0.1419
$\mu = \mu_R,$	S(0.1)	0.163	0.352	0.485	0.001	0.000	0.721	29.14	77.49	0.0424	0.1193
$\sigma \equiv 1$	S(0.3)	0.093	0.301	0.598	0.007	0.000	0.513	24.23	56.17	0.0366	0.1099
	S(0.5)	0.062	0.258	0.657	0.022	0.001	0.415	23.34	46.37	0.0342	0.1060
	CBS	0.033	0.129	0.531	0.204	0.102	0.644	42.69	45.08	0.0417	0.1078
	cumSeg	0.163	0.216	0.403	0.165	0.053	0.904	65.16	105.59	0.1107	0.1492
n = 1000,	HS(0.1)	0.445	0.356	0.198	0.001	0.000	1.474	59.32	162.03	0.0913	0.1801
K = 10,	HS(0.3)	0.303	0.384	0.307	0.005	0.000	1.104	47.34	120.10	0.0682	0.1532
$\lambda_{\min} = 50,$	HS(0.5) $ $	0.213	0.379	0.390	0.018	0.001	0.881	41.98	96.70	0.0577	0.1398
$\mu = \mu_R,$	S(0.1)	0.155	0.351	0.494	0.000	0.000	0.697	32.51	77.29	0.0426	0.1235
$\sigma \equiv 1$	S(0.3)	0.085	0.299	0.612	0.004	0.000	0.485	26.14	55.78	0.0368	0.1131
	S(0.5)	0.054	0.252	0.680	0.014	0.000	0.381	23.81	45.39	0.0344	0.1086
	CBS	0.027	0.135	0.524	0.203	0.111	0.653	45.64	44.88	0.0425	0.1116
	cumSeg	0.165	0.217	0.389	0.179	0.050	0.904	63.73	104.37	0.1037	0.1522

TABLE 2. Simulations with constant variance and C = 200. Columns from left to right: setting, method, proportions of $\hat{K} - K$ and averages of the corresponding error criteria. $HS(\alpha)$ and $S(\alpha)$ denote H-SMUCE and SMUCE at significance level α , respectively.

replace the signal μ_R by

$$\mu_T(i/n) = \mu_R + b\sin(a\pi i)$$

and $\mu_{T_\sigma}(i/n) = \mu_R + b\sigma_R(i/n)\sin(a\pi i) + b(\sigma_R(i/n) - \sigma_R((i-1)/n)\sin(a\pi i),$
 $i = 1, \dots, n,$

respectively. The signal μ_T reflects the situation of a fixed periodic trend, whereas in $\mu_{T_{\sigma}}$ the trend is scaled by the local standard deviation. The last term corrects the size of changes such that still σ_R determines the changes. We consider as in (Venkatraman et al.,

Setting	Method	< -2	-1	0	+1	> +2	$ \hat{K} - K $	FPSLE	FNSLE	MISE	MIAE
n = 100,	HS(0.1)	0.000	0.125	0.873	0.002	0.000	0.128	1.51	4.07	0.8182	0.3308
K = 2,	HS(0.3)	0.000	0.042	0.945	0.013	0.000	0.055	1.04	1.70	0.4217	0.2482
$\lambda_{\min} = 15,$	HS(0.5)	0.000	0.016	0.940	0.043	0.000	0.060	1.63	1.26	0.2776	0.2291
$\mu = \mu_R,$	CBS	0.000	0.001	0.925	0.058	0.016	0.092	2.03	0.79	0.2220	0.2143
$\sigma = \sigma_R$	cumSeg	0.000	0.066	0.720	0.167	0.047	0.343	6.50	4.39	0.4898	0.3053
	LOOVF	0.000	0.031	0.700	0.163	0.106	0.683	12.83	3.36	0.3167	0.2639
n = 100,	HS(0.1)	0.608	0.364	0.028	0.000	0.000	1.610	13.51	32.33	9.5104	1.8626
K = 5,	HS(0.3)	0.212	0.577	0.211	0.000	0.000	1.003	8.63	19.80	6.5362	1.3263
$\lambda_{\min} = 15,$	HS(0.5) $ $	0.061	0.466	0.473	0.001	0.000	0.588	5.27	11.65	3.9992	0.9047
$\mu = \mu_R,$	CBS	0.001	0.008	0.884	0.089	0.018	0.137	1.65	1.02	0.4539	0.3130
$\sigma = \sigma_R$	cumSeg	0.098	0.230	0.544	0.117	0.012	0.588	6.93	12.13	1.2454	0.5441
	LOOVF	0.031	0.112	0.520	0.152	0.184	1.648	14.61	6.92	0.5887	0.4042
n = 1000,	HS(0.1) $ $	0.000	0.007	0.974	0.018	0.000	0.026	8.42	5.83	0.0195	0.0617
$\mathbf{K}=2,$	HS(0.3) $ $	0.000	0.001	0.921	0.075	0.002	0.080	24.72	9.57	0.0193	0.0636
$\lambda_{\min} = 30,$	HS(0.5)	0.000	0.000	0.827	0.162	0.012	0.185	53.23	17.09	0.0204	0.0668
$\mu = \mu_R,$	CBS	0.005	0.019	0.774	0.146	0.056	0.298	52.95	21.17	0.0347	0.0711
$\sigma = \sigma_R$	cumSeg	0.022	0.161	0.683	0.103	0.030	0.387	64.04	92.66	0.0765	0.1112
n = 1000,	HS(0.1) $ $	0.000	0.002	0.982	0.017	0.000	0.018	7.25	4.35	0.0182	0.0630
K = 2,	HS(0.3)	0.000	0.000	0.926	0.071	0.002	0.076	22.64	8.49	0.0196	0.0657
$\lambda_{\min} = 50,$	HS(0.5)	0.000	0.000	0.830	0.160	0.010	0.181	50.02	16.22	0.0214	0.0692
$\mu = \mu_R,$	CBS	0.003	0.011	0.776	0.153	0.057	0.296	53.69	15.85	0.0355	0.0730
$\sigma = \sigma_R$	cumSeg	0.016	0.155	0.699	0.098	0.031	0.370	60.63	84.69	0.0739	0.1132
n = 1000,	HS(0.1)	0.123	0.429	0.446	0.002	0.000	0.080	22.83	55.00	0.4045	0.2402
K = 10,	HS(0.3)	0.010	0.199	0.770	0.015	0.000	0.245	11.98	21.12 10.71	0.1803	0.1018
$\lambda_{\min} \equiv 30,$	HS(0.5)	0.002	0.088	0.803	0.045	0.001	0.140	11.84	12.71	0.1220	0.1404 0.1425
$\mu \equiv \mu_R,$ $\sigma = \sigma$	CD5	0.002	0.008	0.405 0.187	0.510	0.211	0.845	47.20	10.20	0.1274	0.1455
$\frac{0 = 0_R}{n = 1000}$	$\frac{\text{Cumbeg}}{\text{US}(0,1)}$	0.439	0.243	0.107	0.000	0.040	0.215	94.91 16.04	220.44	0.3120	0.2800
II = 1000, K = 10	HS(0.1)	0.025	0.202	0.711	0.002	0.000	0.010	10.94 9.46	32.39 10.59	0.2102	0.1000
K = 10,	HS(0.5)	0.002	0.058 0.017	0.925	0.013	0.000	0.070	0.40	10.36 7 79	0.1009	0.1372 0.1307
$\lambda_{\min} = 50,$	CBS	0.000	0.017 0.007	0.940 0.451	0.045	0.001	0.001	9.03	15.10	0.0800	0.1307
$\mu = \mu_R,$ $\sigma = \sigma_R$	cumSeg	0.001	0.001 0.254	0.451 0.197	0.019	0.222	1 601	97.00	223.47	0.1233 0.2771	0.1403 0.2794
$\frac{0 - 0_R}{n - 10000}$	HS(0 1)	0.100	0.004	0.101	0.002	0.000	0.017	50.65	30.94	0.0016	0.0183
K = 2.	HS(0.3)	0.000	0.001	0.936	0.061	0.001	0.065	188.73	63.72	0.0016	0.0188
$\lambda_{\min} = 30.$	HS(0.5)	0.000	0.001	0.865	0.128	0.006	0.142	407.41	125.46	0.0016	0.0197
$\mu = \mu_B$,	CBS	0.012	0.036	0.532	0.200	0.220	0.886	1548.96	373.22	0.0057	0.0235
$\sigma = \sigma_R$	cumSeg	0.054	0.245	0.600	0.084	0.017	0.477	682.64	1457.08	0.0090	0.0379
n = 10000,	HS(0.1)	0.000	0.001	0.984	0.015	0.000	0.016	53.23	24.89	0.0014	0.0182
K = 2,	HS(0.3)	0.000	0.000	0.941	0.057	0.002	0.060	181.06	59.83	0.0014	0.0188
$\lambda_{\min} = 50,$	HS(0.5)	0.000	0.000	0.870	0.124	0.007	0.137	394.16	115.62	0.0016	0.0197
$\mu = \mu_R,$	CBS	0.012	0.035	0.521	0.208	0.225	0.917	1601.54	366.42	0.0058	0.0238
$\sigma = \sigma_R$	cumSeg	0.052	0.241	0.603	0.087	0.016	0.473	673.81	1430.47	0.0084	0.0377
n = 10000,	HS(0.1)	0.023	0.231	0.741	0.005	0.000	0.282	58.42	165.72	0.0178	0.0431
K = 10,	HS(0.3) $ $	0.006	0.123	0.844	0.027	0.000	0.162	68.27	98.25	0.0122	0.0385
$\lambda_{\min} = 30,$	HS(0.5) $ $	0.003	0.079	0.854	0.064	0.002	0.151	108.19	87.63	0.0103	0.0377
$\mu = \mu_R,$	CBS	0.024	0.043	0.180	0.222	0.531	2.088	1286.59	525.95	0.0198	0.0475
$\sigma = \sigma_R$	cumSeg	0.619	0.169	0.130	0.059	0.024	2.345	1000.55	3122.28	0.0433	0.0917
n = 10000,	HS(0.1)	0.009	0.165	0.819	0.007	0.000	0.190	59.11	124.05	0.0132	0.0418
K = 10,	HS(0.3) $ $	0.001	0.064	0.905	0.029	0.001	0.097	67.32	65.54	0.0089	0.0375
$\lambda_{\min} = 50,$	HS(0.5)	0.000	0.029	0.900	0.067	0.003	0.102	103.42	60.04	0.0078	0.0368
$\mu = \mu_R,$	CBS	0.019	0.034	0.162	0.228	0.557	2.203	1317.31	467.47	0.0198	0.0475
$\sigma = \sigma_R$	cumSeg	0.607	0.188	0.131	0.051	0.023	2.277	997.64	3105.88	0.0405	0.0925
n = 10000,	HS(0.1) $ $	0.609	0.284	0.107	0.001	0.000	1.908	155.65	504.02	0.1016	0.1031
n = 25,	HS(0.3)	0.278	0.399	0.318	0.000	0.000	1.044	94.53	203.3U 199.54	0.0640	0.0702
$\lambda_{\min} = 30,$	DS(0.5) CPC		0.371	0.470	0.019	0.000	0.090	04.07	102.04	0.0483	0.0703
$\mu = \mu_R, \ \sigma = \sigma$		0.015	0.024	0.009	0.128	0.700	0.048 6.099	921.91 1042.99	409.98 3488 49	0.0411	0.0723
$\frac{0 - 0_R}{n - 10000}$		0.934	0.000	0.010	0.009	0.000	1 224	1/16 7/	387 66	0.1109	0.1040
M = 10000, K = 25	HS(0.1)	0.390	0.303	0.220	0.001	0.000	0 501	140.74	175 02	0.0099	0.0940
$\lambda = 20,$ $\lambda = -50$	HS(0.5)	0.103	0.559	0.020	0.010	0.000	0.391	78 74	11/ 01	0.0390	0.0715
$n_{\min} = 00,$	CBS	0.000	0.241 0.017	0.055	0.120	0 700	3 520	934 20	346.33	0.0291	0.0047
$ \begin{array}{c} \mu = \mu_R, \\ \sigma = \sigma_R \end{array} $	cumSeg	0.934	0.036	0.019	0.008	0.003	5.849	1053.35	3462.62	0.1022	0.0120 0.1547
											··-· + •

TABLE 3. Simulations with heterogeneous errors and C = 200. Columns from left to right: setting, method, proportions of $\hat{K} - K$ and averages of the corresponding error criteria. $HS(\alpha)$ denotes H-SMUCE at significance level α

Method	≤ -2	-1	0	+1	$\geq +2$	$ \hat{K} - K $	FPSLE	FNSLE	MISE	MIAE
HS(0.1)	0.005	0.117	0.876	0.002	0.000	0.130	50.82	113.50	0.0107	0.0406
HS(0.3)	0.000	0.032	0.952	0.016	0.000	0.049	48.39	49.84	0.0075	0.0368
HS(0.5)	0.000	0.013	0.940	0.045	0.001	0.061	78.86	48.19	0.0072	0.0368

TABLE 4. $n = 10\,000$ observations, K = 10 change-points, C = 200 and $\lambda_{\min} = 50$ from Table 3. Columns from left to right: setting, method, proportions of $\hat{K} - K$ and averages of the corresponding error criteria. $\text{HS}(\alpha)$ denotes H-SMUCE at significance level α , but with weights $\tilde{\beta}_4, \ldots, \tilde{\beta}_9$.

2004) long (a = 0.01) and short (a = 0.025) trends. Finally, Table 7 reports result of t_3 distributed errors.



FIGURE 10. a: Continuous sinus-shaped standard deviation $\sigma_1(t)$:= $1 + 0.5 \sin(20\pi t).$ \mathbf{b} : Piecewise linear standard deviation $\sigma_2(t)$:=0.5 + $\sum_{i=0}^{9} (10t-i) \mathbb{1}_{(0.1i,0.1(i+1)]}(t).$ c: Piecewise constant standard deviation $\sigma_3(t)$:= $\sum_{i=1}^{n/200} 0.5\mathbb{1}_{(200(i-1)/n, 200(i-1)/n+100/n]}(t) + \mathbb{1}_{(200(i-1)/n+100/n, 200i/n]}(t),$ exemplary for $n = 1\,000.$

Appendix C. Proofs

In this section we collect the proofs together with some auxiliary statements.

C.1. Proof of Lemma 2.1.

Proof of Lemma 2.1. A single statistic $T_i^j(Z,0)$ has the c.d.f. $F_{1,j-i}(\cdot)$ of an F-distribution with (1, j - i) degrees of freedom. Thus, $F_k(\cdot) = F_{1,2^{k}-1}(\cdot)^{|\mathcal{D}_k|}$ is continuous and strictly monotonically increasing for positive arguments. Now, it follows from equation (2.5) that

(C.1)
$$q_k = F_k^{-1} \left(1 - \frac{\beta_k}{\beta_1} \left(1 - F_1(q_1) \right) \right) \text{ for } k = 2, \dots, d_n$$

This together with equation (2.3) yields

$$G(q_1) := F\left(q_1, F_2^{-1}\left(1 - \frac{\beta_2}{\beta_1}\left(1 - F_1(q_1)\right)\right), \dots, F_{d_n}^{-1}\left(1 - \frac{\beta_{d_n}}{\beta_1}\left(1 - F_1(q_1)\right)\right)\right) = 1 - \alpha.$$

Note, that F is continuous and $\lim_{q_k\to 0} F(q_1,\ldots,q_{d_n}) = 0$ for all $k = 1,\ldots,d_n$ as well as $\lim_{q_1,\ldots,q_{d_n}\to\infty} F(q_1,\ldots,q_{d_n}) = 1$. Thus, the function G is continuous, strictly monotonically increasing on $[0,\infty)$ and attains all values in [0,1). Therefore, the existence of the

Setting	Method	≤ -2	-1	0	+1	$\geq +2$	$ \hat{K} - K $	FPSLE	FNSLE	MISE	MIAE
n = 1000,	HS(0.1)	-	-	0.968	0.032	0.000	0.033	16.30	4.12	0.0013	0.0277
$\mathbf{K}=0,$	HS(0.3) $ $	-	-	0.876	0.118	0.005	0.129	64.60	15.91	0.0018	0.0306
$\mu = \mu_R \equiv 0,$	HS(0.5) $ $	-	-	0.734	0.239	0.027	0.293	146.75	36.45	0.0023	0.0338
$\sigma = \sigma_1$	CBS	-	-	0.916	0.001	0.083	0.186	93.25	11.21	0.0045	0.0288
	cumSeg	-	-	1.000	0.000	0.000	0.000	0.20	0.04	0.0011	0.0264
n = 1000,	HS(0.1)	-	-	0.968	0.031	0.001	0.032	16.10	4.12	0.0013	0.0278
$\mathbf{K}=0,$	HS(0.3)	-	-	0.876	0.118	0.005	0.129	64.55	15.73	0.0017	0.0306
$\mu = \mu_R \equiv 0,$	HS(0.5)	-	-	0.734	0.241	0.024	0.292	145.80	35.28	0.0022	0.0340
$\sigma = \sigma_2$	CBS	-	-	0.937	0.004	0.060	0.135	67.70	8.96	0.0034	0.0281
	cumSeg	-	-	0.999	0.001	0.000	0.001	0.40	0.12	0.0011	0.0264
n = 1000,	HS(0.1)	-	-	0.969	0.030	0.001	0.032	15.75	3.91	0.0007	0.0210
$\mathbf{K}=0,$	HS(0.3)	-	-	0.875	0.119	0.006	0.130	65.10	16.31	0.0009	0.0227
$\mu = \mu_R \equiv 0,$	HS(0.5)	-	-	0.737	0.236	0.026	0.290	145.15	36.09	0.0012	0.0250
$\sigma = \sigma_3$	CBS	-	-	0.937	0.002	0.061	0.134	67.10	8.64	0.0019	0.0213
	cumSeg	-	-	0.999	0.001	0.000	0.001	0.35	0.10	0.0006	0.0199
n = 10000,	HS(0.1)	0.013	0.185	0.796	0.005	0.000	0.218	661.16	755.83	0.0212	0.0684
K = 10,	HS(0.3)	0.003	0.076	0.890	0.031	0.001	0.113	543.91	548.21	0.0167	0.0585
$\lambda_{\min} = 50,$	HS(0.5) $ $	0.001	0.041	0.886	0.069	0.003	0.117	513.55	468.37	0.0147	0.0542
$\mu = \mu_R,$	CBS	0.000	0.001	0.191	0.155	0.653	2.636	1590.35	276.51	0.0092	0.0358
$\sigma = \sigma_1$	cumSeg	0.206	0.118	0.413	0.193	0.070	0.984	790.10	1054.73	0.0146	0.0502
n = 10000,	HS(0.1)	0.014	0.205	0.776	0.006	0.000	0.238	421.19	513.32	0.0156	0.0556
K = 10,	HS(0.3)	0.001	0.077	0.894	0.027	0.001	0.108	348.50	358.14	0.0119	0.0475
$\lambda_{\min} = 50,$	HS(0.5) $ $	0.000	0.038	0.897	0.062	0.002	0.105	344.93	311.35	0.0106	0.0446
$\mu = \mu_R,$	CBS	0.000	0.000	0.215	0.174	0.611	2.362	1454.85	247.26	0.0085	0.0346
$\sigma = \sigma_2$	cumSeg	0.114	0.102	0.467	0.236	0.082	0.795	756.12	720.95	0.0136	0.0478
n = 10000,	HS(0.1)	0.019	0.233	0.744	0.004	0.000	0.276	161.27	251.06	0.0053	0.0301
K = 10,	HS(0.3)	0.002	0.069	0.904	0.025	0.000	0.099	137.86	136.79	0.0036	0.0254
$\lambda_{\min} = 50,$	HS(0.5) $ $	0.000	0.029	0.906	0.062	0.003	0.096	170.29	128.56	0.0033	0.0248
$\mu = \mu_R,$	CBS	0.000	0.000	0.246	0.173	0.582	2.189	1134.85	214.71	0.0047	0.0263
$\sigma = \sigma_3$	cumSeg	0.054	0.051	0.516	0.279	0.101	0.669	749.33	499.10	0.0070	0.0346

TABLE 5. Simulations with standard deviations $\sigma_1(\cdot)$ - $\sigma_3(\cdot)$ from Figure 10 and C = 200. Columns from left to right: setting, method, proportions of $\hat{K} - K$ and averages of the corresponding error criteria. HS(α) denotes H-SMUCE at significance level α .

vector of critical values follows from the intermediate value theorem and the vector is also unique. $\hfill \Box$

C.2. **Proof of Lemma 3.1.** First of all, recall from the proof of Lemma 2.1 that the statistic T_k has c.d.f. $F_{1,2^k-1}(\cdot)^{|\mathcal{D}_k|}$. For every $k = 1, \ldots, d_n$ we use the transformation

$$U_k := F_{1,2^k-1} \left(T_k \right)^{|\mathcal{D}_k|}$$

and the identity

$$T_k = F_{1,2^k-1}^{-1} \left(U_k^{|\mathcal{D}_k|^{-1}} \right).$$

Here, $F_{1,2^{k}-1}^{-1}(\cdot)$ denotes the quantile function of an F-distribution with $(1, 2^{k} - 1)$ degrees of freedom. Analogously, we define

$$q_{k,U} := F_{1,2^k-1} \left(q_k \right)^{|\mathcal{D}_k|}$$

Setting	Method	≤ -2	-1	0	+1	$\geq +2$	$ \hat{K} - K $	FPSLE	FNSLE	MISE	MIAE
$\mu = \mu_T,$	HS(0.1)	0.137	0.421	0.439	0.003	0.000	0.709	24.74	58.06	0.3976	0.2449
a = 0.01,	HS(0.3)	0.019	0.209	0.741	0.032	0.000	0.279	15.65	24.13	0.1814	0.1681
b = 0.1,	HS(0.5)	0.003	0.091	0.822	0.081	0.003	0.184	17.32	15.63	0.1206	0.1474
$\sigma = \sigma_R$	CBS	0.001	0.011	0.383	0.290	0.315	1.141	72.93	22.14	0.1321	0.1535
	cumSeg	0.443	0.237	0.192	0.085	0.043	1.663	95.08	225.57	0.3080	0.2823
$\mu = \mu_T,$	HS(0.1)	0.149	0.360	0.370	0.104	0.016	0.821	73.52	94.89	0.4410	0.3070
a = 0.01,	HS(0.3)	0.029	0.181	0.496	0.226	0.067	0.611	78.00	62.38	0.2304	0.2376
b = 0.3,	HS(0.5)	0.007	0.092	0.466	0.306	0.129	0.697	88.73	53.42	0.1595	0.2169
$\sigma = \sigma_R$	CBS	0.001	0.006	0.082	0.135	0.776	3.243	249.67	74.97	0.1646	0.2325
	cumSeg	0.439	0.233	0.200	0.086	0.043	1.652	107.57	237.10	0.3394	0.3222
$\mu = \mu_T,$	HS(0.1)	0.140	0.287	0.323	0.176	0.075	0.936	134.92	135.90	0.5279	0.4052
a = 0.01,	HS(0.3)	0.032	0.146	0.327	0.298	0.197	0.970	146.40	103.03	0.3106	0.3393
b = 0.5,	HS(0.5)	0.009	0.076	0.258	0.329	0.328	1.223	162.18	92.24	0.2410	0.3207
$\sigma = \sigma_R$	CBS	0.002	0.004	0.020	0.043	0.932	5.623	435.32	124.66	0.2462	0.3440
	cumSeg	0.420	0.244	0.190	0.093	0.054	1.641	127.88	248.73	0.3921	0.3823
$\mu = \mu_T,$	HS(0.1)	0.128	0.424	0.446	0.002	0.000	0.693	23.50	56.36	0.4066	0.2415
a = 0.025,	HS(0.3)	0.017	0.201	0.759	0.023	0.001	0.259	13.43	22.21	0.1854	0.1628
b = 0.1,	HS(0.5)	0.003	0.086	0.843	0.066	0.002	0.162	14.51	13.77	0.1218	0.1416
$\sigma = \sigma_R$	CBS	0.002	0.008	0.395	0.287	0.308	1.135	64.07	19.55	0.1304	0.1471
	cumSeg	0.440	0.240	0.188	0.086	0.046	1.672	95.36	229.25	0.3058	0.2796
$\mu = \mu_T$,	HS(0.1)	0.108	0.344	0.411	0.111	0.027	0.738	58.57	73.26	0.4223	0.2606
a = 0.025,	HS(0.3)	0.016	0.138	0.468	0.252	0.126	0.715	77.74	50.19	0.2143	0.1929
b = 0.3,	HS(0.5)	0.003	0.058	0.382	0.315	0.243	0.989	101.66	48.03	0.1503	0.1749
$\sigma = \sigma_R$	CBS	0.002	0.003	0.050	0.065	0.880	5.370	356.54	85.59	0.1585	0.2027
	cumSeg	0.438	0.241	0.184	0.091	0.046	1.678	101.67	234.23	0.3243	0.2958
$\mu = \mu_T$	HS(0.1)	0.054	0.180	0.276	0.226	0.264	1.247	164.83	114.26	0.4732	0.3127
a = 0.025,	HS(0.3)	0.007	0.060	0.195	0.229	0.509	1.945	214.89	100.21	0.2748	0.2586
b = 0.5,	HS(0.5)	0.001	0.027	0.127	0.191	0.654	2.591	256.78	101.30	0.2115	0.2465
$\sigma = \sigma_R$	CBS	0.000	0.001	0.006	0.011	0.982	10.383	709.91	149.02	0.2261	0.2993
	cumSeg	0.439	0.238	0.184	0.088	0.050	1.698	113.20	245.71	0.3520	0.3206
$\mu = \mu_{T_{\sigma}},$	HS(0.1)	0.151	0.435	0.411	0.002	0.000	0.755	26.56	64.22	0.4469	0.3000
a = 0.01,	HS(0.3)	0.021	0.230	0.725	0.023	0.000	0.297	15.80	27.08	0.2289	0.2245
b = 0.2,	HS(0.5)	0.004	0.103	0.819	0.071	0.003	0.188	17.53	17.34	0.1622	0.2019
$\sigma = \sigma_R$	CBS	0.004	0.012	0.342	0.305	0.338	1.225	80.55	26.89	0.1689	0.2056
	cumSeg	0.422	0.233	0.193	0.095	0.056	1.653	107.36	234.27	0.3431	0.3292
$\mu = \mu_{T_{\sigma}},$	HS(0.1)	0.254	0.410	0.298	0.036	0.001	1.012	68.27	115.70	0.6346	0.4582
a = 0.01.	HS(0.3)	0.055	0.261	0.488	0.176	0.019	0.591	81.26	80.78	0.4483	0.3953
b = 0.5,	HS(0.5)	0.015	0.129	0.466	0.321	0.070	0.624	100.70	72.34	0.3826	0.3694
$\sigma = \sigma_R$	CBS	0.002	0.004	0.022	0.059	0.914	4.231	357.42	117.42	0.2907	0.3140
	cumSeg	0.332	0.211	0.198	0.139	0.121	1.575	179.79	280.65	0.4522	0.4317
$\mu = \mu_{T_{\sigma}},$	HS(0.1)	0.136	0.440	0.422	0.002	0.000	0.726	24.82	59.14	0.4567	0.3165
a = 0.025,	HS(0.3)	0.017	0.219	0.746	0.018	0.000	0.273	14.12	24.13	0.2432	0.2435
b = 0.2,	HS(0.5)	0.003	0.096	0.843	0.056	0.002	0.162	14.44	14.77	0.1756	0.2222
$\sigma = \sigma_R$	CBS	0.003	0.011	0.353	0.295	0.338	1.231	71.51	23.32	0.1892	0.2287
-	cumSeg	0.432	0.238	0.182	0.094	0.054	1.688	103.19	236.34	0.3604	0.3501
$\mu = \mu_{T_{\sigma}},$	HS(0.1)	0.181	0.433	0.370	0.016	0.000	0.831	37.89	75.31	0.7365	0.5244
a = 0.025,	HS(0.3)	0.033	0.240	0.594	0.125	0.009	0.450	42.28	44.37	0.5518	0.4736
b = 0.5,	HS(0.5)	0.007	0.110	0.541	0.281	0.061	0.534	64.52	41.39	0.4981	0.4582
$\sigma = \sigma_R$	CBS	0.002	0.002	0.023	0.043	0.929	5.589	365.42	103.32	0.4594	0.4362
	cumSeg	0.316	0.184	0.179	0.139	0.182	1.735	158.60	254.38	0.6153	0.5317

TABLE 6. Simulations with small periodic trends in the mean and n = 1000, K = 10, $\lambda_{\min} = 30$ and C = 200. Columns from left to right: setting, method, proportions of $\hat{K} - K$ and averages of the corresponding error criteria. HS(α) denotes H-SMUCE at significance level α .

Setting	Method	≤ -2	-1	0	+1	$\geq +2$	$ \hat{K} - K $	FPSLE	FNSLE	MISE	MIAE
n = 1000,	HS(0.1)	-	-	0.982	0.018	0.000	0.018	9.05	2.53	0.0031	0.0347
$\mathbf{K}=0,$	HS(0.3)	-	-	0.927	0.071	0.001	0.074	37.05	10.31	0.0034	0.0361
$\mu = \mu_R \equiv 0,$	HS(0.5)	-	-	0.824	0.167	0.009	0.185	92.40	26.35	0.0043	0.0392
$\sigma = \sigma_R,$	S(0.1)	-	-	0.001	0.001	0.999	11.859	5929.70	369.55	0.8710	0.1491
	S(0.3)	-	-	0.000	0.000	1.000	14.803	7401.65	397.77	0.9338	0.1674
	S(0.5)	-	-	0.000	0.000	1.000	16.862	8431.00	411.30	0.9730	0.1787
	CBS	-	-	0.991	0.000	0.009	0.018	9.05	1.13	0.0058	0.0340
	cumSeg	-	-	0.955	0.001	0.044	0.188	93.90	11.98	0.0682	0.0375
n = 1000,	HS(0.1)	0.008	0.136	0.848	0.007	0.000	0.160	25.70	62.95	0.0120	0.0578
K = 2,	HS(0.3)	0.003	0.086	0.876	0.035	0.000	0.127	29.74	44.61	0.0103	0.0537
$\lambda_{\min} = 30,$	HS(0.5)	0.001	0.055	0.851	0.090	0.003	0.152	44.21	38.62	0.0097	0.0524
$\mu = \mu_R,$	S(0.1)	0.000	0.000	0.001	0.001	0.998	11.104	2683.40	250.21	0.3046	0.1232
$\sigma \equiv 1,$	S(0.3)	0.000	0.000	0.000	0.000	1.000	13.984	3361.80	283.17	0.3264	0.1340
	S(0.5)	0.000	0.000	0.000	0.000	1.000	15.991	3836.28	302.43	0.3400	0.1419
	CBS	0.053	0.161	0.726	0.043	0.018	0.346	46.69	119.74	0.0241	0.0712
	cumSeg	0.025	0.097	0.722	0.093	0.063	0.456	108.11	81.86	0.0557	0.0707
n = 10000,	HS(0.1)	0.002	0.079	0.916	0.004	0.000	0.086	93.09	119.69	0.0130	0.0425
K = 10,	HS(0.3)	0.000	0.025	0.957	0.017	0.000	0.043	86.32	81.78	0.0105	0.0397
$\lambda_{\min} = 50,$	HS(0.5)	0.000	0.012	0.950	0.038	0.000	0.050	99.93	76.24	0.0097	0.0389
$\mu = \mu_R,$	CBS	0.467	0.148	0.167	0.107	0.111	2.516	1356.25	6254.20	0.0877	0.1308
$\sigma = \sigma_R$	cumSeg	0.586	0.192	0.136	0.055	0.032	2.242	997.13	3005.71	0.0433	0.0906

TABLE 7. Simulations with t_3 distributed errors and C = 200. Columns from left to right: setting, method, proportions of $\hat{K} - K$ and averages of the corresponding error criteria. $HS(\alpha)$ and $S(\alpha)$ denote H-SMUCE and SMUCE at significance level α , respectively.

and have the identity

(C.2)
$$q_k = F_{1,2^{k-1}}^{-1} \left(q_{k,U}^{|\mathcal{D}_k|^{-1}} \right).$$

Then, the events $U_k > q_{k,U}$ and $T_k > q_k$ are equivalent and therefore the vector $\mathbf{q}_U = (q_{1,U}, \ldots, q_{d_n,U})$ satisfies similar conditions to the equations (2.3) and (2.5), i.e.

(C.3)
$$1 - \mathbb{P}\left(U_1 \le q_{1,U}, \dots, U_{d_n} \le q_{d_n,U}\right) = \alpha$$

and

(C.4)
$$\frac{1 - \mathbb{P}\left(U_1 \le q_{1,U}\right)}{\beta_1} = \dots = \frac{1 - \mathbb{P}\left(U_{d_n} \le q_{d_n,U}\right)}{\beta_{d_n}}$$

The following bounds can be interpreted as a weighted version of the Bonferroni-inequality.

Lemma C.1. $q_{k,U} \leq 1 - \alpha \beta_k$ for $k = 1, \ldots, d_n$.

Proof. We have $\mathbb{P}(U_j \leq q_{j,U}) = q_{j,U}$ for $j = 1, \ldots, d_n$, since U_j is uniformly distributed. Moreover, it follows from condition (C.4) that $1 - q_{j,U} = (1 - q_{k,U})\beta_j/\beta_k$. Combining this with equation (C.3) and $\sum_{j=1}^{d_n} \beta_j = 1$ yields

$$\alpha = 1 - \mathbb{P}\left(U_1 \le q_{1,U}, \dots, U_{d_n} \le q_{d_n,U}\right)$$
$$\leq \sum_{j=1}^{d_n} \mathbb{P}\left(U_j > q_{j,U}\right) = \sum_{j=1}^{d_n} (1 - q_{j,U}) = \sum_{j=1}^{d_n} (1 - q_{k,U}) \frac{\beta_j}{\beta_k} = \frac{1 - q_{k,U}}{\beta_k}$$

which proves the assertion.

Lemma C.2 bounds the quantile function of an F-distribution with (1, c) degrees of freedom.

Lemma C.2 (Bounds on the F-quantiles). Let $F_{1,c}^{-1}(y)$ be the quantile function of an *F*-distribution with (1, c) degrees of freedom, then

$$c\left[\left(1-y^2\right)^{-\frac{1}{c}}-1\right] \le F_{1,c}^{-1}(y) \le c\left[\left(1-y^2\right)^{-\frac{2}{c-\frac{1}{2}}}-1\right].$$

Proof. We have from (Fujikoshi and Mukaihata, 1993, Theorem 4.2) that

$$c\left[\exp\left(\frac{(\chi_1^2)^{-1}(y)}{c}\right) - 1\right] \le F_{1,c}^{-1}(y) \le c\left[\exp\left(\frac{(\chi_1^2)^{-1}(y)}{c - \frac{1}{2}}\right) - 1\right],$$

with $(\chi_1^2)^{-1}(y)$ the quantile function of the chi-squared distribution with one degree of freedom. Moreover, we obtain for all $y \ge 0$

$$\mathbb{P}\left(\chi_1^2 \le y\right) = \mathbb{P}\left(-\sqrt{y} \le Z \le \sqrt{y}\right) = 2\Phi\left(\sqrt{y}\right) - 1$$
$$\iff \left(\chi_1^2\right)^{-1}(y) = \Phi^{-1}\left(\frac{y+1}{2}\right)^2,$$

where $\Phi^{-1}(y)$ is the quantile of the standard gaussian distribution. Furthermore, we have from (Johnson et al., 1994, (13.48), p. 115) that

$$\frac{1}{2} \left[1 + \left(1 - \exp\left(-\frac{x^2}{2}\right) \right)^{\frac{1}{2}} \right] \le \Phi(x) \le \frac{1}{2} \left[1 + \left(1 - \exp\left(-x^2\right) \right)^{\frac{1}{2}} \right]$$

and so for the quantile function one finds

$$\sqrt{-\log\left(1 - (2y - 1)^2\right)} \le \Phi^{-1}\left(y\right) \le \sqrt{-2\log\left(1 - (2y - 1)^2\right)}.$$

Combining the formulas proves the assertion.

Proof of Lemma 3.1. First of all, (C.2) and the equation $|\mathcal{D}_k| = \lfloor n2^{-k} \rfloor$ yields

$$q_{k} = F_{1,2^{k}-1}^{-1} \left(q_{k,U}^{|\mathcal{D}_{k}|^{-1}} \right) = F_{1,2^{k}-1}^{-1} \left(q_{k,U}^{\lfloor n2^{-k} \rfloor^{-1}} \right) \le F_{1,2^{k}-1}^{-1} \left(q_{k,U}^{2^{k}/n} \right).$$

Moreover, it follows from the Lemmas C.1 and C.2 that

$$q_{k} \leq F_{1,2^{k}-1}^{-1} \left(q_{k,U}^{2^{k}/n} \right) \leq F_{1,2^{k}-1}^{-1} \left((1 - \alpha \beta_{k})^{2^{k}/n} \right)$$
$$\leq \left(2^{k} - 1 \right) \left[\left(1 - \left((1 - \alpha \beta_{k})^{2^{k}/n} \right)^{2} \right)^{-\frac{2}{(2^{k}-1)-\frac{1}{2}}} - 1 \right]$$
$$\leq 2^{k} \left[\left(1 - (1 - \alpha \beta_{k})^{2^{k}/n} \right)^{-\frac{4}{2^{k+1}-3}} - 1 \right].$$

Applying Bernoulli's inequality $(1-x)^c \leq 1-cx$ gives

$$q_k \le 2^k \left[\left(1 - (1 - \alpha \beta_k)^{2^k/n} \right)^{-\frac{4}{2^{k+1}-3}} - 1 \right] \le 2^k \left[\left(\frac{2^k \alpha \beta_k}{n} \right)^{-\frac{4}{2^{k+1}-3}} - 1 \right].$$

Moreover, for x, c > 0 the inequality $c^x \le 1 + 2x \log(c)$ holds whenever $x \log(c) \le 1$. Together with the assumption $k \ge 2$ we finally obtain

$$q_{k} \leq 2^{k} \left[\left(\frac{2^{k} \alpha \beta_{k}}{n} \right)^{-\frac{4}{2^{k+1}-3}} - 1 \right] \leq 4 \frac{2^{k+1}}{2^{k+1}-3} \log \left(\frac{n}{2^{k} \alpha \beta_{k}} \right) \leq 8 \log \left(\frac{n}{2^{k} \alpha \beta_{k}} \right)$$
$$2^{-k} \log \left(\frac{n}{2^{k} \alpha \beta_{k}} \right) \leq \frac{1}{2} \frac{2^{k+1}-3}{2^{k+1}} \leq \frac{1}{2}.$$

C.3. Exponential deviation bounds. For the subsequent proofs we need a bound for the distribution function of a single test statistic T_i^j (1.5) which is in our setting a bound for the c.d.f. of a non-central F-distribution.

Lemma C.3. Let Y_1, \ldots, Y_n be *i.i.d.* gaussian random variables with expectation $m \in \mathbb{R}$ and variance $s^2 > 0$. Let $x_+ := \max(x, 0)$. Then, for any $\delta \neq 0$, q > 0

$$\mathbb{P}\left(T_1^n(Y, m+\delta) \le q\right)$$
(C.5)
$$\leq \min_{z\ge 0} \left\{ \exp\left(-\frac{1}{2}\left(\frac{\Delta\sqrt{n}}{2} - \frac{q(1+z)}{\Delta\sqrt{n}}\right)_+^2\right) + \exp\left(-(n-1)\frac{z-\log(1+z)}{2}\right) \right\},$$

where $\Delta := |\delta|/s$.

Proof. Let $\widetilde{T}_i^j(Y,m) := (j-i+1) \left(\overline{Y}_{ij} - m\right)^2 / s^2$. Then, $\widetilde{T}^n(Y,m+\delta)$

$$T_1^n(Y, m+\delta) = \frac{T_1^n(Y, m+\delta)}{\hat{s}_{1n}^2/s^2}.$$

if

The statistics \hat{s}_{1n}^2/s^2 and $\widetilde{T}_1^n(Y, m + \delta)$ are independent, since $\widetilde{T}_1^n(Y, m + \delta)$ depends only on the mean \overline{Y}_{1n} . Hence, for all $z \ge 0$

$$\begin{split} \mathbb{P}\left(T_1^n(Y,m+\delta) \le q\right) &= \mathbb{P}\left(\widetilde{T}_1^n(Y,m+\delta) \le q\frac{\hat{s}_{1n}^2}{s^2}\right) \\ &= \mathbb{P}\left(\widetilde{T}_1^n(Y,m+\delta) \le q\frac{\hat{s}_{1n}^2}{s^2} \middle| \frac{\hat{s}_{1n}^2}{s^2} \le 1+z\right) \mathbb{P}\left(\frac{\hat{s}_{1n}^2}{s^2} \le 1+z\right) \\ &+ \mathbb{P}\left(\widetilde{T}_1^n(Y,m+\delta) \le q\frac{\hat{s}_{1n}^2}{s^2} \middle| \frac{\hat{s}_{1n}^2}{s^2} > 1+z\right) \mathbb{P}\left(\frac{\hat{s}_{1n}^2}{s^2} > 1+z\right) \\ &\le \mathbb{P}\left(\widetilde{T}_1^n(Y,m+\delta) \le q(1+z)\right) + \mathbb{P}\left(\frac{\hat{s}_{1n}^2}{s^2} > 1+z\right) \\ &\le \exp\left(-\frac{1}{2}\left(\frac{\Delta\sqrt{n}}{2} - \frac{q(1+z)}{\Delta\sqrt{n}}\right)_+^2\right) + \exp\left(-(n-1)\frac{z-\log(1+z)}{2}\right). \end{split}$$

The first term of the last inequality follows from (Frick et al., 2014, Lemma 7.3 and the proof) and the second from (Spokoiny and Zhilova, 2013, Theorem 2.1), since $(n - 1)\hat{s}_{1n}^2/s^2 \sim \chi_{n-1}^2$.

It remains to show that the minimum in (C.5) is attained for some $z \ge 0$. The function $(\Delta\sqrt{n}/2 - q(1+z)/(\Delta\sqrt{n}))^2_+$ is strictly monotonically decreasing for z > 0 until the function value zero is attained for some finite z. The function $(n-1)(z - \log(1+z))$ is zero for z = 0 and strictly monotonically increasing on $[0, \infty)$. Therefore, the two continuous functions intersect and the minimum is attained for some $z \ge 0$.

The minimum in the last lemma cannot be determined analytically, but it can be computed numerically. In Lemma C.4 we estimate the right hand side further to obtain an explicit exponential bound.

Lemma C.4. Let Y_1, \ldots, Y_n , $n \ge 4$, be i.i.d. gaussian random variables with expectation $m \in \mathbb{R}$ and variance $s^2 > 0$, then we have for all q > 0 with

(C.6)
$$\frac{q}{n} \le \frac{1}{8}$$

as well as for all $\delta \neq 0$ and $\Delta := |\delta|/s$ the bound

(C.7)
$$\mathbb{P}\left(T_1^n(Y, m+\delta) \le q\right) \le 2\exp\left(-\frac{1}{48}\left(\sqrt{n\Delta} - \sqrt{2q}\right)_+^2\right).$$

Proof. Let z > 0 be arbitrary, but fixed. Then, it follows from Lemma C.3 that

$$\mathbb{P}\left(T_1^n(Y, m+\delta) \le q\right) \le \exp\left(-\frac{1}{2}\left(\frac{\Delta\sqrt{n}}{2} - \frac{q(1+z)}{\Delta\sqrt{n}}\right)_+^2\right) + \exp\left(-(n-1)\frac{z - \log(1+z)}{2}\right)$$
$$\le 2\exp\left(-\min\left[\frac{1}{2}\left(\frac{\Delta\sqrt{n}}{2} - \frac{q(1+z)}{\Delta\sqrt{n}}\right)_+^2, (n-1)\frac{z - \log(1+z)}{2}\right]\right).$$

The inequality

$$z - \log(1+z) \ge \frac{1}{2} \frac{z^2}{1+z} \ge \frac{1}{4} \min(z^2, z)$$

yields

$$\mathbb{P}\left(T_1^n(Y, m+\delta) \le q\right)$$

$$\le 2\exp\left(-\min\left[\frac{1}{8}n\left(\Delta - \frac{2q(1+z)}{\Delta n}\right)_+^2, \frac{1}{8}(n-1)\min\left(z^2, z\right)\right]\right)$$

$$\le 2\exp\left(-\frac{1}{8}(n-1)\min\left[\min\left[\left(\Delta - \frac{2q(1+z)}{\Delta n}\right)_+^2, z^2\right], \min\left[\left(\Delta - \frac{2q(1+z)}{\Delta n}\right)_+^2, z\right]\right]\right)$$

Now, we minimize the r.h.s in $z \ge 0$. The functions z and z^2 are both increasing, the function $(\Delta - 2q(1+z)/(\Delta n))^2_+$ in contrast is decreasing in z. Therefore, both inner minima are attained and by solving the corresponding quadratic equations (note that we have to take the solution with $\Delta - 2q(1+z)/(\Delta n) \ge 0$) we get

$$\mathbb{P}\left(T_{1}^{n}(Y, m+\delta) \leq q\right)$$

$$\leq 2\exp\left(-\frac{1}{8}(n-1)\min\left[\left(\frac{\Delta-\frac{2q}{\Delta n}}{1+\frac{2q}{\Delta n}}\right)_{+}^{2}, \frac{1+2\frac{2q}{\Delta n}\left(\Delta-\frac{2q}{\Delta n}\right)_{+}-\sqrt{1+4\frac{2q}{\Delta n}\left(\Delta-\frac{2q}{\Delta n}\right)_{+}}}{2\left(\frac{2q}{\Delta n}\right)^{2}}\right]\right).$$

Using the inequality $\sqrt{1+4x} \le 1+2x-2x^2+4x^3$ for all x > -1/4 with $x = 2q/(\Delta n)(\Delta - 2q/(\Delta n))_+$ we find

$$\mathbb{P}\left(T_{1}^{n}(Y, m+\delta) \leq q\right)$$

$$\leq 2\exp\left(-\frac{1}{8}\frac{n-1}{n}n\min\left[\left(\frac{\Delta-\frac{2q}{\Delta n}}{1+\frac{2q}{\Delta n}}\right)_{+}^{2}, \left(\Delta-\frac{2q}{\Delta n}\right)_{+}^{2}\left[1-2\frac{2q}{\Delta n}\left(\Delta-\frac{2q}{\Delta n}\right)_{+}\right]\right]\right).$$

Next, we consider the two terms in the minimum separately. We assume w.l.o.g. that $\sqrt{2q}/(\Delta\sqrt{n}) \leq 1$, since otherwise the r.h.s. in (C.7) is two. For the first term we distinguish the cases $2q > \Delta n$ and $2q \leq \Delta n$. If $2q \leq \Delta n$ is satisfied, then

$$n\left(\frac{\Delta-\frac{2q}{\Delta n}}{1+\frac{2q}{\Delta n}}\right)_{+}^{2} \ge \frac{1}{4}n\left(\Delta-\frac{2q}{\Delta n}\right)_{+}^{2} = \frac{1}{4}\left(\sqrt{n}\Delta-\frac{2q}{\Delta\sqrt{n}}\right)_{+}^{2} \ge \frac{1}{4}\left(\sqrt{n}\Delta-\sqrt{2q}\right)_{+}^{2}.$$

For the other case, when $2q > \Delta n$ holds, we obtain with $q/n \le 1/8$

$$n\left(\frac{\Delta - \frac{2q}{\Delta n}}{1 + \frac{2q}{\Delta n}}\right)_{+}^{2} \ge \frac{1}{4}n\left(\frac{\Delta - \frac{2q}{\Delta n}}{\frac{2q}{\Delta n}}\right)_{+}^{2} = \frac{1}{4}n\left(\frac{n\Delta^{2}}{2q} - 1\right)_{+}^{2}$$
$$= \frac{1}{4}\frac{n}{2q}\left(\frac{(\sqrt{n}\Delta)^{2}}{\sqrt{2q}} - \sqrt{2q}\right)_{+}^{2} \ge \left(\sqrt{n}\Delta - \sqrt{2q}\right)_{+}^{2}.$$

For the second term it follows with $q/n \leq 1/8$ that

$$n\left(\Delta - \frac{2q}{\Delta n}\right)_{+}^{2} \left[1 - 2\frac{2q}{\Delta n}\left(\Delta - \frac{2q}{\Delta n}\right)_{+}\right] = \left(\sqrt{n}\Delta - \frac{2q}{\Delta\sqrt{n}}\right)_{+}^{2} \left[1 - 4\frac{q}{n}\left(1 - \frac{2q}{\Delta^{2}n}\right)_{+}\right]$$
$$\geq \left(\sqrt{n}\Delta - \sqrt{2q}\right)_{+}^{2} \frac{1}{2}.$$

This yields

$$\mathbb{P}\left(T_1^n(Y, m+\delta) \le q\right) \le 2 \exp\left(-\frac{1}{32}\frac{n-1}{n}\left(\sqrt{n\Delta} - \sqrt{2q}\right)_+^2\right)$$
$$\le 2 \exp\left(-\frac{1}{48}\left(\sqrt{n\Delta} - \sqrt{2q}\right)_+^2\right).$$

C.4. Proofs of Section 3.

Proof of Theorem 3.3. The estimated number of change-points \hat{K} is by its definition in (3.3) equal to the minimal number of change-points of all feasible functions. Therefore, all functions with the true number of change-points (or less change-points) have to be infeasible, if the number of change-points is overestimated. Hence, by (2.3)

$$\sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} > K\right) \leq \sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{P}_{(\mu,\sigma^2)}\left(\max_{\left[\frac{i}{n},\frac{j}{n}\right]\in\mathcal{D}(\mu)} \left[T_i^j\left(Y,\mu(\left[i/n,j/n\right]\right)) - q_{ij}\right] > 0\right)$$
$$\leq \mathbb{P}_{(0,1)}\left(\max_{\left[\frac{i}{n},\frac{j}{n}\right]\in\mathcal{D}} \left[T_i^j\left(Y,0\right)\right) - q_{ij}\right] > 0\right) = \alpha,$$

where the last inequality follows from $\mathcal{D}(\mu) \subset \mathcal{D}$ and the fact that the distribution of $T_i^j(Y, \mu([i/n, j/n]))$ does not depend on $\mu(\cdot)$ and $\sigma(\cdot)$, as these are constant on intervals in $\mathcal{D}(\mu)$.

Proof of Theorem 3.4. First of all, we show that it is enough to prove the result for $\mu \equiv 0$ and $\sigma^2 \equiv 1$ and hence K = 0. We have

$$\begin{split} \sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{P}_{(\mu,\sigma^2)} \left(\hat{K} > K + 2k \right) \\ &= \sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{P}_{(\mu,\sigma^2)} \left(\max_{\left[\frac{i}{n},\frac{j}{n}\right]\in\mathcal{D}(\tilde{\mu})} \left[T_i^j \left(Y, \tilde{\mu}([i/n, j/n])\right) - q_{ij} \right] > 0 \ \forall \ \tilde{\mu} \in \mathcal{M} \ s.t. \ |\mathcal{I}(\tilde{\mu})| \le K + 2k \right) \\ &\leq \sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{P}_{(\mu,\sigma^2)} \left(\max_{\left[\frac{i}{n},\frac{j}{n}\right]\in\mathcal{D}(\tilde{\mu})} \left[T_i^j \left(Y, \tilde{\mu}([i/n, j/n])\right) - q_{ij} \right] > 0 \\ &\quad \forall \ \tilde{\mu} \in \mathcal{M} \ s.t. \ \mathcal{I}(\mu) \subset \mathcal{I}(\tilde{\mu}), |\mathcal{I}(\tilde{\mu})| \le K + 2k \right) \\ &\leq \mathbb{P}_{(0,1)} \left(\max_{\left[\frac{i}{n},\frac{j}{n}\right]\in\mathcal{D}(\tilde{\mu})} \left[T_i^j \left(Y, \tilde{\mu}([i/n, j/n])\right) - q_{ij} \right] > 0 \ \forall \ \tilde{\mu} \in \mathcal{M} \ s.t. \ |\mathcal{I}(\tilde{\mu})| \le 2k \right) \\ &= \mathbb{P}_{(0,1)} \left(\hat{K} > 2k \right), \end{split}$$

where the last inequality follows from the same argument as in the proof of Theorem 3.3. Now, we define $R_0 := 0$ and iteratively

$$R_{k+1} := \min\{t > R_k : \exists s \ s.t. \ R_k < s < t \ \text{and} \ [s/n, t/n] \in \mathcal{D}, T_s^t(Y, 0) > q_{\log_2(t-s+1)}\},$$

with the convention $\min \emptyset = \infty$. Then,

$$\mathbb{P}_{0,1}(R_{k+1} \le n | R_1 = t) \le \mathbb{P}_{0,1}(R_k \le n)$$
 for all $t \in \{1, \dots, n\}$

since for the l.h.s. the remaining k rejections R_2, \ldots, R_{k+1} have to be in $\{t+1, \ldots, n\}$ instead of $\{1, \ldots, n\}$. It follows

$$\mathbb{P}_{0,1}(\hat{K} > 2k) \le \mathbb{P}_{0,1}(R_{k+1} \le n) = \sum_{t=1}^{n} \mathbb{P}_{0,1}(R_{k+1} \le n | R_1 = t) \mathbb{P}_{0,1}(R_1 = t)$$
$$\le \mathbb{P}_{0,1}(R_1 \le n) \mathbb{P}_{0,1}(R_k \le n) \le \dots \le \mathbb{P}_{0,1}(R_1 \le n)^{k+1} \le \alpha^{k+1},$$

where the last inequality is given by Theorem 3.3. It follows

$$\sup_{(\mu,\sigma^2)\in\mathcal{S}} \mathbb{E}_{(\mu,\sigma^2)} \left[(\hat{K} - K)_+ \right] = \sup_{(\mu,\sigma^2)\in\mathcal{S}} \sum_{k=0}^{\infty} \mathbb{P}_{(\mu,\sigma^2)} \left(\hat{K} - K > k \right)$$
$$\leq \sup_{(\mu,\sigma^2)\in\mathcal{S}} 2 \sum_{k=0}^{\infty} \mathbb{P}_{(\mu,\sigma^2)} \left(\hat{K} - K > 2k \right) \leq 2 \sum_{k=0}^{\infty} \alpha^{k+1} = \frac{2\alpha}{1-\alpha}.$$

The following theorem is sharper version of 3.5 that shows different probabilities for the detection of the change-points.

Theorem C.5 (Underestimation control II). Let $\lambda_j := \tau_{j+1} - \tau_j$ and $k_{n,j} := \lfloor \log_2(n\lambda_j/4) \rfloor$, $j = 0, \ldots, K$, as well as $\delta_j := |m_j - m_{j-1}|$ and

$$\eta_j := \left[1 - 3 \exp\left(-\frac{1}{48} \left(\sqrt{\frac{n\lambda_{j-1}\delta_j^2}{32\sigma_{j-1}^2}} - \sqrt{16\log\left(\frac{8}{\lambda_j\alpha\beta_{k_{n,j-1}}}\right)} \right)_+^2 \right) \right]_+ \times \left[1 - 3 \exp\left(-\frac{1}{48} \left(\sqrt{\frac{n\lambda_j\delta_j^2}{32\sigma_j^2}} - \sqrt{16\log\left(\frac{8}{\lambda_j\alpha\beta_{k_{n,j}}}\right)} \right)_+^2 \right) \right]_+,$$

 $j = 1, \ldots, K$. Under the assumptions of Theorem 3.3 and if $n\lambda_j \geq 32$ and

$$(n\lambda_j)^{-1}\log\left(\frac{8}{\lambda_j\alpha\beta_{k_{n,j}}}\right) \le \frac{1}{512}$$

are satisfied for all $j = 1, \ldots, K$, then

$$\mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} < K\right) \le 1 - \prod_{j=1}^K \eta_j \text{ and } \mathbb{E}_{(\mu,\sigma^2)}\left[\left(K - \hat{K}\right)_+\right] \le \sum_{j=1}^K (1 - \eta_j) + \sum_{j=1}^K (1 - \eta_$$

Proof. For each j = 1, ..., K we consider the disjoint intervals $I_j := [\tau_j - \lambda_{j-1}/2, \tau_j + \lambda_j/2)$ and split them into disjoint intervals $I_j^+ \cup I_j^- = I_j$ such that $\mu(t) = \mu^+ \forall t \in I_j^+$ and $\mu(t) = \mu^- \forall t \in I_j^-$, with $\mu^+ := \max(m_{j-1}, m_j)$ and $\mu^- := \min(m_{j-1}, m_j)$. Without loss of generality we assume $\mu^+ = m_{j-1}$ and $\mu^- = m_j$ in the following. Then, there exists subintervals $J_j^+ \subset I_j^+$ and $J_j^- \subset I_j^-$ with $J_j^+, J_j^- \in \mathcal{D}$ that have length $\lambda_{j-1}^* := n^{-1} 2^{\lfloor \log_2(n\lambda_{j-1}/4) \rfloor} = n^{-1} 2^{k_{n,j-1}} \geq \lambda_{j-1}/8$, since $n|I_j^+| = n\lambda_{j-1}/2 \geq 3$, and $\lambda_j^* := n^{-1} 2^{\lfloor \log_2(n\lambda_j/4) \rfloor} = n^{-1} 2^{k_{n,j}} \geq \lambda_j/8$, since $n|I_j^-| = n\lambda_j/2 \geq 3$, respectively. It follows

$$\begin{split} \mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} < K\right) &= 1 - \mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} \ge K\right) \\ &\leq 1 - \mathbb{P}_{(\mu,\sigma^2)}\left(\nexists \ \hat{\mu} \in C(\mathbf{q}), \ j \in \{1, \dots, K\} : \ \hat{\mu} \text{ is constant on } I_j\right) \\ &\leq 1 - \mathbb{P}_{(\mu,\sigma^2)}\left(\forall \ j \in \{1, \dots, K\} : \ \nexists \ \hat{m} \le (m_{j-1} + m_j)/2 : T_{J_j^+}(Y, \hat{m}) \le q_{k_{n,j-1}} \text{ and} \\ & \nexists \ \hat{m} \ge (m_{j-1} + m_j)/2 : T_{J_j^-}(Y, \hat{m}) \le q_{k_{n,j}}\right) \\ &\leq 1 - \prod_{j=1}^K \mathbb{P}_{(\mu,\sigma^2)}\left(\nexists \ \hat{m} \le (m_{j-1} + m_j)/2 : T_{J_j^+}(Y, \hat{m}) \le q_{k_{n,j-1}} \text{ and} \\ & \nexists \ \hat{m} \ge (m_{j-1} + m_j)/2 : T_{J_j^-}(Y, \hat{m}) \le q_{k_{n,j-1}} \text{ and} \\ & \nexists \ \hat{m} \ge (m_{j-1} + m_j)/2 : T_{J_j^-}(Y, \hat{m}) \le q_{k_{n,j}}\right), \end{split}$$

where we used in the last inequality that the events are independent, since all intervals are disjoint. We denote by Z_1, \ldots, Z_n i.i.d. standard normally distributed random variables. It follows from once again from the independence due to disjoint intervals and from the

Lemmas 7.1 in (Frick et al., 2014), C.4 and 3.1 that

$$\mathbb{P}_{(\mu,\sigma^2)} \Big(\nexists \ \hat{m} \le (m_{j-1} + m_j)/2 : T_{J_j^+}(Y, \hat{m}) \le q_{k_{n,j-1}} \text{ and} \nexists \ \hat{m} \ge (m_{j-1} + m_j)/2 : T_{J_j^-}(Y, \hat{m}) \le q_{k_{n,j}} \Big) \ge \Big[1 - \mathbb{P}_{(\mu,\sigma^2)} \Big(\exists \ \hat{m} \le (m_{j-1} + m_j)/2 : T_{J_j^+}(Y, \hat{m}) \le q_{k_{n,j-1}} \Big) \Big] \times \Big[1 - \mathbb{P}_{(\mu,\sigma^2)} \Big(\exists \ \hat{m} \ge (m_{j-1} + m_j)/2 : T_{J_j^-}(Y, \hat{m}) \le q_{k_{n,j}} \Big) \Big] \ge \eta_j,$$

since

$$\begin{split} & \mathbb{P}_{(\mu,\sigma^2)} \left(\exists \ \hat{m} \le (m_{j-1} + m_j)/2 : T_{J_j^+}(Y, \hat{m}) \le q_{k_{n,j-1}} \right) \\ & \le \mathbb{P}_{(\mu,\sigma^2)} \left(\overline{Y}_{J_j^+} \le (m_{j-1} + m_j)/2 \text{ or } T_{J_j^+}(Y, (m_{j-1} + m_j)/2) \le q_{k_{n,j-1}} \right) \\ & \le \mathbb{P}_{(\mu,\sigma^2)} \left(\overline{Y}_{J_j^+} \le (m_{j-1} + m_j)/2 \right) + \mathbb{P}_{(\mu,\sigma^2)} \left(T_{J_j^+}(Y, (m_{j-1} + m_j)/2) \le q_{k_{n,j-1}} \right) \\ & \le \mathbb{P} \left(\overline{Z}_{[0,\lambda_{j-1}^*]} \ge \frac{\delta_j}{2\sigma_{j-1}} \right) + \mathbb{P} \left(T_{[0,\lambda_{j-1}^*]} \left(Z, \frac{\delta_j}{2\sigma_{j-1}} \right) \le q_{k_{n,j-1}} \right) \\ & \le \exp \left(-\frac{1}{64} \frac{n\lambda_{j-1}\delta_j^2}{\sigma_{j-1}^2} \right) + 2 \exp \left(-\frac{1}{48} \left(\sqrt{\frac{n\lambda_{j-1}\delta_j^2}{32\sigma_{j-1}^2}} - \sqrt{2q_{k_{n,j-1}}} \right)_+^2 \right) \\ & \le 3 \exp \left(-\frac{1}{48} \left(\sqrt{\frac{n\lambda_{j-1}\delta_j^2}{32\sigma_{j-1}}} - \sqrt{16 \log \left(\frac{8}{\lambda_{j-1}\alpha\beta_{k_{n,j-1}}} \right)} \right)_+^2 \right) \end{split}$$

and the second term by symmetry arguments. Moreover, it follows

$$\begin{split} & \mathbb{E}_{(\mu,\sigma^2)} \left[\left(K - \hat{K} \right)_+ \right] \\ \leq & \mathbb{E}_{(\mu,\sigma^2)} \left[\sum_{j=1}^K \mathbb{1}_{\exists \, \hat{m} \le (m_{j-1} + m_j)/2: \, T_{J_j^+}(Y, \hat{m}) \le q_{k_{n,j-1}} \text{ or } \exists \, \hat{m} \ge (m_{j-1} + m_j)/2: \, T_{J_j^-}(Y, \hat{m}) \le q_{k_{n,j}} \right] \\ \leq & \sum_{j=1}^K \left(1 - \eta_j \right). \end{split}$$

Proof of Theorem 3.5. The proof is analogue to the proof of Theorem C.5, but with $I_j = [\tau_j - \lambda/2, \tau_j + \lambda/2)$.

Proof of Theorem 3.7. We prove the theorem with the Borel-Cantelli lemma. It follows from Theorems 3.3 and 3.5 that

$$\sup_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}}\mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K}_n\neq K\right)$$

$$=\sup_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}}\mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K}_n>K\right) + \sup_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}}\mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K}_n< K\right)$$

$$\leq \alpha_n + 1 - \left[1 - 3\exp\left(-\frac{1}{48}\left(\sqrt{\frac{n\lambda\Delta^2}{32}} - \sqrt{16\log\left(\frac{8}{\lambda\alpha_n\beta_{k_n,n}}\right)}\right)_+^2\right)\right]_+^{2K}$$

$$\leq \alpha_n + 6K\exp\left(-\frac{1}{48}\left(\sqrt{\frac{n\lambda\Delta^2}{32}} - \sqrt{16\log\left(\frac{8}{\lambda\alpha_n\beta_{k_n,n}}\right)}\right)_+^2\right),$$

since under the given assumptions the conditions of Theorem 3.5 are satisfied. The upper bounds for the error probabilities are summable if (3.6) is satisfied.

Lemma C.6 (Confidence set). Assume the setting and assumptions of Theorem 3.5 and let C(q) be as in (3.7) with significance level α and weights $\beta_1, \ldots, \beta_{d_n}$. Let $S_{\Delta,\lambda}$ be as in (3.4) with $\Delta, \lambda > 0$ arbitrary, but fixed, and $k_n := \lfloor \log_2(n\lambda/4) \rfloor$. If $n\lambda \geq 32$ and

$$\frac{\log\left(\frac{8}{\lambda\alpha_n\beta_{k_n}}\right)}{n\lambda} \le \frac{1}{512}$$

hold, then uniformly in $\mathcal{S}_{\Delta,\lambda}$

$$\mathbb{P}_{(\mu,\sigma^2)}\left(\mu \in C\left(\boldsymbol{q}\right)\right) \ge 1 - \alpha - (1 - \eta^K),$$

with η like in Theorem 3.5.

Proof. It follows from the definition of $C(\mathbf{q})$ in (3.7) as well as from Theorems 3.3 and 3.5 that

$$\inf_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}} \mathbb{P}_{(\mu,\sigma^2)} \left(\mu \in C\left(\mathbf{q}\right)\right) \\
= \inf_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}} \mathbb{P}_{(\mu,\sigma^2)} \left(\max_{\left[\frac{i}{n},\frac{j}{n}\right]\in\mathcal{D}(\mu)} \left[T_i^j\left(Y,\mu(\left[i/n,j/n\right]\right)\right) - q_{ij}\right] \le 0, \hat{K} = K \right) \\
= \inf_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}} \mathbb{P}_{(\mu,\sigma^2)} \left(\max_{\left[\frac{i}{n},\frac{j}{n}\right]\in\mathcal{D}(\mu)} \left[T_i^j\left(Y,\mu(\left[i/n,j/n\right]\right)\right) - q_{ij}\right] \le 0, \hat{K} \ge K \right) \\
\ge \inf_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}} \mathbb{P}_{(\mu,\sigma^2)} \left(\max_{\left[\frac{i}{n},\frac{j}{n}\right]\in\mathcal{D}(\mu)} \left[T_i^j\left(Y,\mu(\left[i/n,j/n\right]\right)\right) - q_{ij}\right] \le 0 \right) - \sup_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}} \mathbb{P}_{(\mu,\sigma^2)} \left(\hat{K} < K\right) \\
\ge 1 - \alpha - (1 - \eta^K).$$

Proof of Theorem 3.8. The statement is a direct consequence of Lemma C.6.

Lemma C.7 (Change-point locations). Assume the setting of Lemma C.6. If c_n is a sequence with $0 < c_n \le \lambda/2$ and $k_n := \lfloor \log_2(nc_n/2) \rfloor$ such that $nc_n \ge 16$ and

(C.8)
$$\frac{\log\left(\frac{4}{c_n\alpha\beta_{k_n}}\right)}{nc_n} \le \frac{1}{256}$$

hold, then uniformly in $\mathcal{S}_{\Delta,\lambda}$

$$\mathbb{P}_{(\mu,\sigma^2)}\left(\sup_{\hat{\mu}\in C(q_n)} \max_{\tau\in\mathcal{I}(\mu)} \min_{\hat{\tau}\in\mathcal{I}(\hat{\mu})} |\hat{\tau}-\tau| > c_n\right)$$

$$\leq 1 - \left[1 - 3\exp\left(-\frac{1}{48}\left(\sqrt{\frac{nc_n\Delta^2}{16}} - \sqrt{16\log\left(\frac{4}{c_n\alpha\beta_{k_n}}\right)}\right)_+^2\right)\right]_+^{2K}.$$

Proof. Analogously to the proof of Theorem C.5 we have

$$\sup_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}} \mathbb{P}_{(\mu,\sigma^2)} \left(\sup_{\hat{\mu}\in C(\mathbf{q}_n)} \max_{\tau\in\mathcal{I}(\mu)} \min_{\hat{\tau}\in\mathcal{I}(\hat{\mu})} |\hat{\tau}-\tau| > c_n \right)$$

$$\leq \sup_{(\mu,\sigma^2)\in\mathcal{S}_{\Delta,\lambda}} \mathbb{P}_{(\mu,\sigma^2)} \left(\exists \ j \in \{1,\ldots,K\} \text{ and } \hat{\mu}\in C(\mathbf{q}_n) : \hat{\mu} \text{ is constant on } [\tau_j - c_n, \tau_j + c_n) \right)$$

$$\leq 1 - \left[1 - 3 \exp\left(-\frac{1}{48} \left(\sqrt{\frac{nc_n\Delta^2}{16}} - \sqrt{16\log\left(\frac{4}{c_n\alpha\beta_{k_n}}\right)} \right)_+^2 \right) \right]_+^{2K} .$$

Proof of Theorem 3.9. For n large enough such that (3.9) guarantees the assumption of Lemma C.7 it follows

$$\mathbb{P}_{(\mu,\sigma^2)}\left(\sup_{\hat{\mu}\in C(\mathbf{q}_n)} \max_{j=1,\dots,K} c_n^{-1} |\tau_j - \hat{\tau}_j| > 1\right) \\
\leq \mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} > K \text{ or } \exists \ \hat{\mu}\in C(\mathbf{q}_n), \ j\in\{1,\dots,K\}: \hat{\mu} \text{ is constant on } [\tau_j - c_n, \tau_j + c_n]\right) \\
\leq \mathbb{P}_{(\mu,\sigma^2)}\left(\hat{K} > K\right) + \mathbb{P}_{(\mu,\sigma^2)}\left(\exists \ \hat{\mu}\in C(\mathbf{q}_n), \ j\in\{1,\dots,K\}: \hat{\mu} \text{ is constant on } [\tau_j - c_n, \tau_j + c_n]\right) \\
\leq \alpha_n + \left(1 - \left[1 - 3\exp\left(-\frac{1}{48}\left(\sqrt{\frac{nc_n\Delta^2}{16}} - \sqrt{16\log\left(\frac{4}{c_n\alpha_n\beta_{k_n,n}}\right)}\right)_+^2\right)\right]_+^{2K}\right).$$

The assertion follows from $\alpha_n \to 0$ and

$$\lim_{n \to \infty} \sqrt{\frac{nc_n \Delta^2}{16}} - \sqrt{16 \log\left(\frac{4}{c_n \alpha_n \beta_{k_n, n}}\right)} = \infty,$$

whereby latter one is direct consequence of (3.9).

The following theorem deals with the detection of single vanishing bump against a noisy background.

Theorem C.8 (Single vanishing bump). Assume the heterogeneous gaussian changepoint model (1.2) with sequences of bump signals $\mu_n(t) := m_0 + \delta_n \mathbb{1}_{I_n}(t)$ and $\sigma_n(t) := \mathbb{1}_{I_n^C}(t) + s_n \mathbb{1}_{I_n}(t)$, where $\delta_n \neq 0$ is a sequence of change-point sizes, $s_n > 0$ a sequence of standard deviations on $I_n \in \mathcal{D}$, which is a sequence of intervals with $|I_n| \to 0$. Let $k_n := \lfloor \log_2(n|I_n|) \rfloor$ and $\Delta_n := |\delta_n|/s_n$ be the sequence of the signal to noise ratios. Let $(\hat{K}_n)_n, \alpha_n$ and $\beta_{1,n}, \ldots, \beta_{d_n,n}$ be as in Theorem 3.10. We further assume

(C.9)
$$\sqrt{n|I_n|}\Delta_n \ge (4+\epsilon_n)\sqrt{-\log(|I_n|)},$$

with possibly $\epsilon_n \to 0$, but such that $\epsilon_n \sqrt{-\log(|I_n|)} \to \infty$ and

$$\limsup_{n \to \infty} \frac{\sqrt{-\log\left(\alpha_n \beta_{k_n,n}\right)}}{\epsilon_n \sqrt{-\log(|I_n|)}} < \frac{1}{4},$$

(C.10)
$$\liminf_{n \to \infty} \frac{n|I_n|}{\log(n)} > 64 \text{ and } \lim_{n \to \infty} \frac{\log(\alpha_n \beta_{k_n, n})}{n|I_n|} = 0.$$

(C.11)
$$\lim_{n \to \infty} s_n \frac{\sqrt{|I_n|}}{\sqrt{|I_n|}} = \infty \ and$$

(C.12)
$$\liminf_{n \to \infty} \frac{\log(\beta_{k_n,n})}{\log(\beta_{\min,n})} > 0, \text{ with } \beta_{\min,n} := \min\{\beta_{1,n}, \dots, \beta_{d_n,n}\}.$$

Then,

(C.13)
$$\lim_{n \to \infty} \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(\hat{K}_n > 0 \right) = 1$$

Conditions (C.9) and (C.10) are the main assumptions of the theorem to detect the vanishing signal on I_n . We discussed them together with the conditions of Theorem 3.10 in Section 3.3. We also need the weak technical conditions (C.11) and (C.12) on the size of $|I_n^C|$ and the minimal weight $\beta_{\min,n}$ to ensure that the detection power on the complement I_n^C is large enough, too. Condition (C.12) is for instance fulfilled by uniform weights $\beta_{1,n} = \cdots = \beta_{d_n,n} = 1/d_n$, but many other choices are possible, too. We further assumed $I_n \in \mathcal{D}$, otherwise we have to replace I_n by the largest subinterval which is an element of the dyadic partition. Such an interval exists always and has at least length $n^{-1}2^{\lfloor \log_2(n|I_n|/2) \rfloor} > |I_n|/4$. Therefore, omitting the condition $I_n \in \mathcal{D}$ would not change the rate. It is possible to strengthen (C.13) further to $\lim_{n\to\infty} \mathbb{P}_{(\mu_n,\sigma_n^2)}(\hat{K}_n \geq K) = 1$ if we increase all constants a little bit.

Proof of Theorem C.8. We denote by J_n the longest subinterval $J_n \subset I_n^C$ which is part of the dyadic partition. Such an interval exists (at least for n large enough) always, since $|I_n| \to 0$, and has at least length $|I_n^C|/8$. Moreover, let $k_n := \log_2(n|I_n|)$ and $l_n := \log_2(n|J_n|)$. Then, the Lemmas 7.1 in (Frick et al., 2014) and C.4 yield for any $\theta_n > 0$

$$\begin{split} \lim_{n \to \infty} \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(\hat{K}_n > 0 \right) \\ &= \lim_{n \to \infty} 1 - \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(\hat{\mu} \text{ is constant} \right) \\ &\geq \lim_{n \to \infty} 1 - \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(\exists \ \hat{m} \le m_0 + \theta_n : T_{I_n}(Y, \hat{m}) \le q_{k_n} \text{ or } \exists \ \hat{m} \ge m_0 + \theta_n : T_{J_n}(Y, \hat{m}) \le q_{l_n} \right) \\ &\geq \lim_{n \to \infty} 1 - \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(\exists \ \hat{m} \le m_0 + \theta_n : T_{I_n}(Y, \hat{m}) \le q_{k_n} \right) \\ &- \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(\exists \ \hat{m} \ge m_0 + \theta_n : T_{J_n}(Y, \hat{m}) \le q_{l_n} \right) \\ &\geq \lim_{n \to \infty} 1 - \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(\overline{Y}_{I_n} \le m_0 + \theta_n \right) - \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(T_{I_n}(Y, m_0 + \theta_n) \le q_{k_n} \right) \\ &- \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(\overline{Y}_{J_n} \ge m_0 + \theta_n \right) - \mathbb{P}_{(\mu_n, \sigma_n^2)} \left(T_{J_n}(Y, m_0 + \theta_n) \le q_{l_n} \right) \\ &\geq \lim_{n \to \infty} 1 - 2\mathbb{P}_{(\mu_n, \sigma_n^2)} \left(T_{I_n}(Y, m_0 + \theta_n) \le q_{k_n} \right) - 2\mathbb{P}_{(\mu_n, \sigma_n^2)} \left(T_{J_n}(Y, m_0 + \theta_n) \le q_{l_n} \right) \\ &\geq \lim_{n \to \infty} 1 - 4 \exp\left(-\frac{1}{48} \left(\Gamma_{I_n} \right)_+^2 \right) - 4 \exp\left(-\frac{1}{48} \left(\Gamma_{J_n} \right)_+^2 \right) = 1, \end{split}$$

i

$$\Gamma_{I_n} := \sqrt{n|I_n|} \frac{\delta_n - \theta_n}{s_n} - \sqrt{2q_{k_n}} \to \infty \text{ and } \Gamma_{J_n} := \sqrt{n|J_n|} \theta_n - \sqrt{2q_{l_n}} \to \infty,$$

and if the conditions of Lemma C.4 are satisfied. This is the case, since $n|I_n| \to \infty$ and $n|J_n| \to \infty$, because of (C.10) and $|I_n| \to 0$, as well as $q_{k_n}/(n|I_n|) \le 1/8$ and $q_{l_n}/(n|J_n|) \leq 1/8$ hold at least for n large enough: The first one is a direct consequence of Lemma 3.1 and (C.10)

$$\frac{q_{k_n}}{n|I_n|} \le \frac{8\log\left(\frac{1}{|I_n|\alpha_n\beta_{k_n,n}}\right)}{n|I_n|} \le \frac{1}{8},$$

since then the assumptions of Lemma 3.1 are also fulfilled. The second inequality follows from Lemma 3.1, (C.10) and (C.12) as well as the fact that $|I_n|/|J_n| \to 0$

$$\lim_{n \to \infty} \frac{q_{l_n}}{n|J_n|} \le \lim_{n \to \infty} \frac{8 \log\left(\frac{1}{|J_n|\alpha_n\beta_{l_n,n}}\right)}{n|J_n|} \le \lim_{n \to \infty} \frac{8 \log\left(\frac{1}{|J_n|\alpha_n\beta_{l_n,n}}\right)}{8 \log\left(\frac{1}{|I_n|\alpha_n\beta_{k_n,n}}\right)} \frac{|I_n|}{|J_n|} \frac{8 \log\left(\frac{1}{|I_n|\alpha_n\beta_{k_n,n}}\right)}{n|I_n|} \to 0,$$

since then the assumptions of Lemma 3.1 are also fulfilled. We define now $\theta_n = \sqrt{\gamma_n/n}$ via the equation

$$\sqrt{\frac{\gamma_n |I_n|}{s_n^2}} = c\epsilon_n \sqrt{\log\left(\frac{1}{|I_n|}\right)}$$

for 0 < c < 1. Then, it follows from Lemma 3.1 and from $\sqrt{x+y} \le \sqrt{x} + \sqrt{y}$ for x, y > 0 together with the assumptions of the theorem that

$$\begin{split} \Gamma_{I_n} &= \sqrt{n|I_n|} \frac{\delta_n - \theta_n}{s_n} - \sqrt{2q_{k_n}} \\ &= \sqrt{n|I_n|\delta_n^2/s_n^2} - \sqrt{\gamma_n|I_n|/s_n^2} - \sqrt{2q_{k_n}} \\ &\geq \sqrt{n|I_n|\Delta_n^2} - \sqrt{\gamma_n|I_n|/s_n^2} - \sqrt{16\log\left(\frac{1}{|I_n|}\right)} \\ &\geq (4 + \epsilon_n)\sqrt{\log\left(\frac{1}{|I_n|}\right)} - c\epsilon_n\sqrt{\log\left(\frac{1}{|I_n|}\right)} - 4\sqrt{\log\left(\frac{1}{|I_n|}\right)} - 4\sqrt{\log\left(\frac{1}{\alpha_n\beta_{k_n,n}}\right)} \\ &\geq (1 - c)\epsilon_n\sqrt{\log\left(\frac{1}{|I_n|}\right)} - 4\sqrt{\log\left(\frac{1}{\alpha_n\beta_{k_n,n}}\right)} \to \infty, \end{split}$$

since the conditions of Lemma 3.1 are satisfied, as shown above. Moreover, we have $\Gamma_{J_n} := \sqrt{n|J_n|}\theta_n - \sqrt{2q_{l_n}} = \sqrt{|J_n|\gamma_n} - \sqrt{2q_{l_n}} \to \infty$ if

$$\sqrt{\frac{|J_n|\gamma_n}{2q_{l_n}}} \ge \sqrt{\frac{|J_n|\gamma_n}{16\log\left(\frac{1}{|J_n|\alpha_n\beta_{l_n,n}}\right)}} \to \infty,$$

where we used Lemma 3.1 again. Finally, it follows from the assumptions of the theorem that $\liminf_{n\to\infty} |J_n| \ge \liminf_{n\to\infty} |I_n^C|/8 > 0$ and thus

$$\sqrt{\frac{|J_n|\gamma_n}{\log\left(\frac{1}{\alpha_n\beta_{l_n,n}}\right)}} = \frac{\sqrt{|I_n|\gamma_n}}{s_n\sqrt{\log\left(\frac{1}{\alpha_n\beta_{k_n,n}}\right)}} \frac{s_n\sqrt{|J_n|}}{\sqrt{|I_n|}} \frac{\sqrt{\log\left(\frac{1}{\alpha_n\beta_{l_n,n}}\right)}}{\sqrt{\log\left(\frac{1}{\alpha_n\beta_{l_n,n}}\right)}} \\
\geq \frac{c\epsilon_n\sqrt{\log\left(\frac{1}{|I_n|\right)}}}{\sqrt{\log\left(\frac{1}{\alpha_n\beta_{k_n,n}}\right)}} \frac{s_n\sqrt{|I_n^C|/8}}{\sqrt{|I_n|}} \frac{\sqrt{\log\left(\frac{1}{\alpha_n\beta_{k_n,n}}\right)}}{\sqrt{\log\left(\frac{1}{\alpha_n\beta_{min,n}}\right)}} \to \infty.$$

Proof of Theorem 3.10. It follows from Theorem 3.5 that

$$\mathbb{P}_{(\mu_n,\sigma_n^2)}\left(\hat{K}_n < K_n\right) \le 1 - \left[1 - 3\exp\left(-\frac{1}{48}\left(\Gamma_n\right)_+^2\right)\right]_+^{2K_n} \le 6K_n \exp\left(-\frac{1}{48}\left(\Gamma_n\right)_+^2\right),$$

with

$$\Gamma_n := \sqrt{\frac{n\lambda_n\Delta_n^2}{32}} - \sqrt{16\log\left(\frac{8}{\lambda_n\alpha_n\beta_{k_n,n}}\right)},$$

since the assumptions of Theorem 3.5 are satisfied by (3.11). In case (1) it is enough to show $\Gamma_n \to \infty$, because K_n is bounded. Finally, $\Gamma_n \to \infty$ follows from

$$\frac{n\lambda_n\Delta_n^2}{\log\left(\frac{8}{\lambda_n\alpha_n\beta_{k_n,n}}\right)}\to\infty.$$

In case (2) for bounded K_n , $\Gamma_n \to \infty$ follows from

$$\Gamma_{n} = \sqrt{\frac{n\lambda_{n}\Delta_{n}^{2}}{32}} - \sqrt{16\log\left(\frac{8}{\lambda_{n}\alpha_{n}\beta_{k_{n},n}}\right)}$$

$$\geq \left(\frac{\sqrt{512}}{\sqrt{32}} + \frac{\epsilon_{n}}{\sqrt{32}}\right)\sqrt{\log\left(\frac{1}{\lambda_{n}}\right)} - \sqrt{16}\sqrt{\log\left(\frac{1}{\lambda_{n}}\right)} - \sqrt{16}\sqrt{\log\left(\frac{8}{\alpha_{n}\beta_{k_{n},n}}\right)}$$

$$= \frac{1}{\sqrt{32}}\left(\epsilon_{n}\sqrt{\log\left(\frac{1}{\lambda_{n}}\right)} - \sqrt{512}\sqrt{\log\left(\frac{8}{\alpha_{n}\beta_{k_{n},n}}\right)}\right) \to \infty.$$

For unbounded K_n we have $K_n \leq 1/\lambda_n$. It follows

$$K_{n} \exp\left(-\frac{1}{48}\left(\Gamma_{n}\right)_{+}^{2}\right)$$

$$\leq \exp\left(\log\left(\frac{1}{\lambda_{n}}\right) - \frac{1}{48}\left(\frac{C}{\sqrt{32}}\sqrt{\log\left(\frac{1}{\lambda_{n}}\right)} + \frac{1}{\sqrt{32}}\epsilon_{n}\sqrt{\log\left(\frac{1}{\lambda_{n}}\right)} - \sqrt{16}\sqrt{\log\left(\frac{8}{\alpha_{n}\beta_{k_{n},n}}\right)}\right)_{+}^{2}\right)$$

$$\leq \exp\left(-\frac{1}{48}\left(\frac{1}{\sqrt{32}}\epsilon_{n}\sqrt{\log\left(\frac{1}{\lambda_{n}}\right)} - \sqrt{16}\sqrt{\log\left(\frac{8}{\alpha_{n}\beta_{k_{n},n}}\right)}\right)^{2}\right) \rightarrow 0.$$

C.5. Proofs of Section A.

Proof of Theorem A.1. We prove the assertion with (van der Vaart, 2007, Theorem 5.9) which states three conditions for the convergence of a Z-estimator. Note that the convergence in probability can be replaced by almost sure convergence, if the assumptions hold almost surely. We define

$$\Psi(\theta) := |F(\theta) - (1 - \alpha)| + \sum_{k=2}^{d_n} \left| \frac{1 - F_1(\theta_1)}{\beta_1} - \frac{1 - F_k(\theta_k)}{\beta_k} \right|$$

and

$$\Psi_{M}(\theta) := |F_{M}(\theta) - (1 - \alpha)| + \sum_{k=2}^{d_{n}} \left| \frac{1 - F_{M,1}(\theta_{1})}{\beta_{1}} - \frac{1 - F_{M,k}(\theta_{k})}{\beta_{k}} \right|$$

as well as $\Theta := [0, \infty)^{d_n}$, $\theta_0 := \mathbf{q}$ and $\hat{\theta}_M := \widehat{\mathbf{q}}_M$. Now, (A.2) and (A.3) yield

$$\Psi_M\left(\widehat{\mathbf{q}}_M\right) \le \frac{1}{M} \left(1 + \frac{d_n - 1}{\min\{\beta_1, \dots, \beta_{d_n}\}}\right) = \mathbf{o}(1)$$

almost surely. In addition, Lemma 2.1 shows that the vector of critical values \mathbf{q} is unique. Moreover, $\sup_{\theta \in [0,\infty)^{d_n}} \|F_M(\theta) - F(\theta)\|$ and $\sup_{\theta_k \ge 0} \|F_{M,k}(\theta_k) - F_k(\theta_k)\|$ for all

 $k \in \{1, \ldots, d_n\}$ converge to zero almost surely. Thus, all assumptions of (van der Vaart, 2007, Theorem 5.9) are satisfied and the assertion follows.

Proof of Lemma A.2. The computation time for the bounds $\underline{b}_{i,j}$ and $\overline{b}_{i,j}$ is $\mathcal{O}(1)$ for every fixed interval $[i/n, j/n] \in \mathcal{D}$, since they depend only on the sums $\sum_{l=i}^{j} Y_{l}$ as well as $\sum_{l=i}^{j} Y_{l}^{2}$ and these can be obtained from (precomputed) cumulative sums. The computation time for the intersected bounds $\underline{B}_{i,j}$ and $\overline{B}_{i,j}$ are also $\mathcal{O}(1)$ for a fixed interval [i/n, j/n], since they can be computed iteratively. Therefore, the total time to compute the bounds is $\mathcal{O}(n)$, since the dyadic partition contains less than n intervals.

It follows from its iterative definition that the left limits $L_1, \ldots, L_{\hat{K}}$ can be computed in $\mathcal{O}(n)$. Therefore, the dynamic programming algorithm has cost $\mathcal{O}(\sum_{k=1}^{\hat{K}-1}(R_k - L_k + 1) (R_{k+1} - L_{k+1} + 1))$ besides some linear costs, since for each point in the interval $[L_{k+1}, R_{k+1}]$ the optimal change-point in the interval $[L_k, R_k]$ has to be determined by computing the cost functional for each of these points. But, for a single interval the computation time for the restricted maximum likelihood estimator and for the cost functional is $\mathcal{O}(1)$ if the constraints $\underline{B}_{i,j}$ and $\overline{B}_{i,j}$ are given, since the restricted maximum likelihood estimator and the cost functional depend besides these constraints again only on the sums $\sum_{l=i}^{j} Y_l$ and $\sum_{l=i}^{j} Y_l^2$. This proves the assertion.

References

- S. Arlot and A. Celisse. Segmentation of the mean of heteroscedastic data via cross-validation. *Statist. Comput.*, 21(4):613–632, 2011.
- S. Arlot, A. Celisse, and Z. Harchaoui. Kernel change-point detection. arXiv:1202.3878, 2012.
- J. Bai and P. Perron. Estimating and testing linear models with multiple structural changes. *Econometrica*, 66(1):47–78, 1998.
- J. Bai and P. Perron. Computation and analysis of multiple structural change models. J. Appl. Econometrics, 18:1–22, 2003.
- N. K. Bakirov and G. J. Szekely. Students t-test for gaussian scale mixtures. J. Math. Sci., 139(3):6497–6505, 2006.
- R. Benz. Permeation of hydrophilic solutes through mitochondrial outer membranes: review on mitochondrial porins. *BBA-Rev. Biomembranes*, 1197(2):167–196, 1994.
- L. Birgé and P. Massart. Gaussian model selection. J. Eur. Math. Soc., 3(3):203–268, 2001.
- L. Boysen, A. Kempe, V. Liebscher, A. Munk, and O. Wittich. Consistencies and rates of convergence of jump-penalized least squares estimators. Ann. Statist., 37(1):157–183, 2009.

- J. V. Braun, R. K. Braun, and H.-G. Müller. Multiple changepoint fitting via quasilikelihood, with application to DNA sequence segmentation. *Biometrika*, 87(2):301–314, 2000.
- H. P. Chan and G. Walther. Detection with the scan and the average likelihood ratio. *Statist. Sinica*, 23:409–428, 2013.
- M. Csörgo and L. Horváth. Limit Theorems in Change-Point Analysis. Wiley, 1997.
- L. Davies, C. Höhenrieder, and W. Krämer. Recursive computation of piecewise constant volatilities. *Comput. Statist. Data Anal.*, 56:3623–3631, 2012.
- R. B. Davies. Hypothesis testing when a nuisance parameter is present only under the alternatives. *Biometrika*, 74(1):33–43, 1987.
- R. B. Davies. Hypothesis testing when a nuisance parameter is present only under the alternative: Linear model case. *Biometrika*, 89(2):484–489, 2002.
- D. L. Donoho and I. M. Johnstone. Ideal spatial adaptation by wavelet shrinkage. *Biometrika*, 81(3):425–455, 1994.
- C. Du, C.-L. M. Kao, and S. C. Kou. Stepwise signal extraction via marginal likelihood. J. Amer. Statist. Assoc., 2015. to appear.
- L. Dümbgen and V. G. Spokoiny. Multiscale testing of qualitative hypotheses. Ann. Statist., 29(1):124–152, 2001.
- L. Dümbgen and G. Walther. Multiscale inference about a density. Ann. Statist., 36(4): 1758–1785, 2008.
- F. Enikeeva, A. Munk, and F. Werner. Bumb detection in a heterogeneous Gaussian model. arXiv:1504.07390, 2015.
- K. Frick, A. Munk, and H. Sieling. Multiscale change-point inference. With discussion and rejoinder by the authors. J. Roy. Statist. Soc. Ser. B, 76(3):495–580, 2014.
- P. Fryzlewicz. Wild binary segmentation for multiple change-point detection. Ann. Statist., 42(6):2243–2281, 2014.
- Y. Fujikoshi and S. Mukaihata. Approximations for the quantiles of Student's t and F distributions and their error bounds. *Hiroshima Math. J.*, 23(3):557–564, 1993.
- A. Futschik, T. Hotz, A. Munk, and H. Sieling. Multiscale DNA partitioning: Statistical evidence for segments. *Bioinformatics*, 30(16):2255–2262, 2014.
- P. Guillaume, R. Pintelon, and J. Schoukens. On the use of signals with a constant signalto-noise ration in the frequency domain. *IEEE Trans. Instrum. Meas.*, 39(6):835–842, 1990.
- Z. Harchaoui and C. Lévy-Leduc. Multiple change-point estimation with a total variation penalty. J. Amer. Statist. Assoc., 105:1480–1493, 2010.
- T. Hotz, O. Schütte, H. Sieling, T. Polupanow, U. Diederichsen, C. Steinem, and A. Munk. Idealizing ion channel recordings by jump segmentation and statistical multiresolution analysis. *IEEE Trans. Nanobiosci.*, 12:376–386, 2013.

- R. Ibragimov and U. K. Müller. t-test based correlation and heterogeneity robust inference. J. Bus. Econom. Statist., 28(4):453–468, 2010.
- X. J. Jeng, T. T. Cai, and H. Li. Optimal sparse segment identification with application in copy number variation analysis. J. Amer. Statist. Assoc., 105(491):1156–1166, 2010.
- N. L. Johnson, S. Kotz, and N. Balakrishnan. Continuous Univariate Distributions, volume 1. Wiley, 2. edition, 1994.
- R. Killick, P. Fearnhead, and I. A. Eckley. Optimal detection of changepoints with a linear computational cost. J. Amer. Statist. Assoc., 107(500):1590–1598, 2012.
- E. D. Kolaczyk and R. D. Nowak. Multiscale generalised linear models for nonparametric function estimation. *Biometrika*, 92(1):119–133, 2005.
- E. L. Lehmann and J. P. Romano. *Testing Statistical Hypotheses*. Springer Texts in Statistics, 3. edition, 2005.
- R. Maidstone and B. Pickering. Discussion of "Multiscale change-point inference" by K. Frick, A. Munk and H. Sieling. J. Roy. Statist. Soc. Ser. B, 76(3):560–561, 2014.
- D. S. Matteson and N. A. James. A nonparametric approach for multiple change point analysis of multivariate data. J. Amer. Statist. Assoc., 109(505):334–345, 2014.
- V. M. R. Muggeo and G. Adelfio. Efficient change point detection for genomic sequences of continuous measurements. *Bioinformatics*, 27(2):161–166, 2011.
- P. Rigollet and A. Tsybakov. Sparse estimation by exponential weighting. *Statist. Sci.*, 27(4):558–575, 2012.
- C. Rivera and G. Walther. Optimal detection of a jump in the intensity of a Poisson process or in a density with likelihood ratio statistics. *Scand. J. Stat.*, 40:752–769, 2013.
- K. Rufibach and G. Walther. The block criterion for multiscale inference about a density, with applications to other multiscale problems. J. Comput. Graph. Statist., 19(1): 175–190, 2010.
- B. Sakmann and E. Neher, editors. Single-Channel Recording. Plenum Press, 2nd edition, 1995.
- M. J. Schervish. P values: what they are and what they are not. Am. Stat., 50(3):203–206, 1996.
- T. Schirmer. General and specific porins from bacterial outer membranes. J. Struct. Biol., 121(2):101–109, 1998.
- A. J. Scott and M. Knott. A cluster analysis method for grouping means in the analysis of variance. *Biometrics*, 30(3):507–512, 1974.
- H. Sieling. Statistical Multiscale Segmentation: Inference, Algorithms and Applications. PhD thesis, Georg-August University Göttingen, 2013.
- F. J. Sigworth. Open channel noise. I. Noise in acetylcholine receptor currents suggests conformational fluctuations. *Biophys. J.*, 47(5):709–720, 1985.

- V. Spokoiny and M. Zhilova. Sharp deviation bounds for quadratic forms. *Math. Methods Statist.*, 22(2):100–113, 2013.
- A. W. van der Vaart. *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics, 8th printing edition, 2007.
- E. S. Venkatraman and A. B. Olshen. A faster circular binary segmentation algorithm for the analysis of array CGH data. *Bioinformatics*, 23(6):657–663, 2007.
- E. S. Venkatraman, A. B. Olshen, R. Lucito, and M. Wigler. Circular binary segmentation for the analysis of array-based DNA copy number data. *Biostatistics*, 5(4):557–572, 2004.
- L. J. Vostrikova. Detecting "disorder" in multidimensional random processes. Soviet Math. Dokl., 24(1):55–59, 1981.
- G. Walther. Optimal and fast detection of spatial clusters with scan statistics. Ann. Statist., 38(2):1010–1033, 2010.
- Y. Yao. Estimating the number of change-points via Schwarz criterion. Statist. Probab. Lett., 6:181–189, 1988.
- N. R. Zhang and D. O. Siegmund. Model selection for high-dimensional, multi-sequence change-point problems. *Statist. Sinica*, 22:1507–1538, 2012.
- Z. Zhou. Discussion of "Multiscale change-point inference" by K. Frick, A. Munk and H. Sieling. J. Roy. Statist. Soc. Ser. B, 76(3):566–567, 2014.