REGULARITY OF POWERS OF EDGE IDEAL OF WHISKERED CYCLES

M. MOGHIMIAN, S. A. SEYED FAKHARI, AND S. YASSEMI

ABSTRACT. Let $G = W(C_n)$ be a whiskered cycle graph with edge ideal I = I(G). We prove that for every $s \ge 1$, the equality $\operatorname{reg}(I^s) = 2s + \lceil \frac{n-1}{2} \rceil - 1$ holds.

1. INTRODUCTION AND PRELIMINARIES

Let I be a homogeneous ideal in the polynomial ring $R = \mathbb{K}[x_1, \ldots, x_n]$. Suppose that the minimal free resolution of I is given by

$$0 \longrightarrow \cdots \longrightarrow \bigoplus_{j} R(-j)^{\beta_{1,j}(I)} \longrightarrow \bigoplus_{j} R(-j)^{\beta_{0,j}(I)} \longrightarrow I \longrightarrow 0.$$

The Castelnuovo-Mumford regularity (or simply, regularity) of I, denote by reg(I), is defined as follows:

 $\operatorname{reg}(I) = \max\{j - i | \beta_{i,j}(I) \neq 0\}.$

The regularity of I is an important invariant in commutative algebra.

There is a natural correspondence between quadratic squarefree monomial ideals of R and finite simple graphs with n vertices. To every simple graph G with vertex set $V(G) = \{x_1, \ldots, x_n\}$ and edge set E(G), we associate an ideal I = I(G) defined by

$$I(G) = (x_i x_j : \{x_i, x_j\} \in E(G)) \subseteq R.$$

Computing and finding bounds for the regularity of edge ideals and their powers have been studied by a number of researchers (see for example[1], [2], [3], [5], [7], [11] and [12]). It is well-known that $\operatorname{reg}(I^s)$ is asymptotically a linear function for $s \gg 0$. When I = I(G) is an edge ideal, then there exist integers b and s_0 such that for all $s \geq s_0$, $\operatorname{reg}(I^s) = 2s + b$. The simplest case is when b = 0, i.e $\operatorname{reg}(I^s) = 2s$, in this case, I^s has linear minimal free resolution. For example, Banerjee [2] proves that if G is a gap-free and cricket-free gragh then $\operatorname{reg}(I(G)^s) = 2s$. Using [10, Theorem 3.2] together with Fröberg's result [8], we know that if the complement graph of G is chordal, then $I(G)^s$ has linear resolution for all $s \geq 1$.

Recently, Beyarslam, Hà and Trung [3] proved that for every graph G and every integer $s \geq 1$, the inequality

 $\operatorname{reg}(I(G)^s) \ge 2s + \operatorname{indmatch}(G) - 1$

²⁰⁰⁰ Mathematics Subject Classification. Primary: 13D02, 05E40 Secondary: 05C38.

Key words and phrases. Edge ideal, Castelnuovo-Mumford regularity, Even-connected path, Whiskered cycle graph.

holds, where indmatch(G) denotes the induced matching number of G, and it is the maximum cardinality of the induced matching of G (see [3, Theorem 4.5]). In the same paper, the authors proved the equality for every $s \ge 1$, if G is a forest and for every $s \ge 2$, if G is a cycle (see [3, Theorems 4.7 and 5.2]). In this paper, we determine a new class of graphs for which the equality

$$\operatorname{reg}(I(G)^s) = 2s + \operatorname{indmatch}(G) - 1$$

holds, for every $s \ge 1$.

Let G be a graph. Adding a whisker to G at a vertex v means adding a new vertex u and the edge $\{u, v\}$ to G. The graph which is obtained from G by adding a whisker to each vertex of G is denoted by W(G). For every integer $n \ge 3$, let C_n be the n-cycle graph and set $G = W(C_n)$. It immediately follows from [4, Theorem 13] that $\operatorname{reg}(I(G)) = \lceil \frac{n-1}{2} \rceil + 1$ and one can easily check that $\operatorname{indmath}(G) = \lceil \frac{n-1}{2} \rceil$. Thus, we have the following result.

Proposition 1.1. For every integer $n \geq 3$, let C_n be the n-cycle graph and set $G = W(C_n)$. Then $\operatorname{reg}(I(G)) = \lceil \frac{n-1}{2} \rceil + 1 = \operatorname{indmath}(G) + 1$

As the main result of this paper, we extend Proposition 1.1 by proving than for every $s \ge 1$, $\operatorname{reg}(I(W(C_n))^s) = 2s + \lceil \frac{n-1}{2} \rceil - 1$ (see Theorem 2.5).

We first need to recall some basic definitions from graph theory.

If two vertices of a graph G are joined by an edge then these vertices are called neighbors. The set of all neighbors of a vertex v is called the neighborhood set of vand is denoted by N(v). Moreover, we set $N[v] = N(v) \cup \{v\}$. The degree of a vertex v of G is the number of its neighbors and is denoted by $\deg_G(v)$.

A subgraph H of G is called induced provided that two vertices of H are adjacent if and only if they are adjacent in G. For an edge e in G, let $G \setminus e$ be the subgraph of G obtained by removing the edge e from G. For a subset $S \subseteq V(G)$ of vertices, let $G \setminus S$ be the induced subgraph on $V(G) \setminus S$ and we write $G \setminus v$ instead of $G \setminus \{v\}$. For an edge $e = \{u, v\}$ of G, we set $N_G[e] = N_G[u] \cup N_G[v]$ and define G_e to be the induced subgraph $G \setminus N_G[e]$ of G.

Convention: Throughout this paper, for simplicity, we use reg(G) instead of reg(I(G)).

We need the following result due to Hà [9] on the regularity of edge ideals.

Theorem 1.2. Let G = (V(G), E(G)) be a graph.

- (i) [9, Lemma 3.1] For any induced subgraph H of G we have $reg(H) \leq reg(G)$.
- (ii) [9, Theorem 3.4] Let $x \in V(G)$. Then

 $\operatorname{reg}(G) \le \max\{\operatorname{reg}(G \setminus x), \operatorname{reg}(G \setminus N[x]) + 1\}.$

(iii) [9, Theorem 3.5] Let $e \in E(G)$. Then

 $\operatorname{reg}(G) \leq \max\{2, \operatorname{reg}(G \setminus e), \operatorname{reg}(G_e) + 1\}.$

Our method for proving the main result is based on the recent work of Banerjee [2]. We recall the following definition and theorem from [2].

Definition 1.3. Let G be a graph. Two vertices u and v (u may be equal to v) are said to be even-connected with respect to an s-fold product $e_1 \ldots e_s$ of edges of G, if there is a path $p_0, p_1, \ldots, p_{2l+1}, l \ge 1$ in G such that the following conditions hold:

- (i) $p_0 = u$ and $p_{2l+1} = v$;
- (ii) for all $0 \le k \le l 1$, $\{p_{2k+1}, p_{2k+2}\} = e_i$ for some *i*; and
- (iii) for all i, $|\{k \mid \{p_{2k+1}, p_{2k+2}\} = e_i\} |\leq |\{j \mid e_i = e_j\}|$.

Theorem 1.4. [2, Theorems 6.1 and 6.7]

Assume that $s \ge 1$ is an integer, G is a graph and I = I(G) is its edge ideal. Let M be a minimal generator of I^s . Then the ideal $(I^{s+1}: M)$ is generated by monomials of degree two and for every generator uv (u may be equal to v) of this ideal, either $\{u, v\}$ is an edge of G or u and v are even-connected with respect to M.

2. Main results

The aim of this section is to prove that for every $s \ge 1$, we have

$$\operatorname{reg}(I^s) = 2s + \lceil \frac{n-1}{2} \rceil - 1,$$

where I is the edge ideal of a whiskered cycle $W(C_n)$ (see Theorem 2.5). We first need the following technical lemma. The proof of this lemma is long and we postpone it to the next section.

Lemma 2.1. Let G be a graph with 2n vertices, say $\{x_1, x_2, \ldots, x_{2n}\}$. Assume that

- (i) $\{x_i, x_{i+1}\} \in E(G)$ for every $1 \le i \le n-1$.
- (ii) $\{x_i, x_{n+i}\} \in E(G)$ for every $1 \le i \le n$.
- (iii) For $2 \le i, j \le n-1$, if x_{n+i} is adjacent to x_{n+j} in G, then $\{x_{n+i}, x_{j-1}\}$, $\{x_{n+i}, x_{j+1}\}$, $\{x_{n+j}, x_{i-1}\}$ and $\{x_{n+j}, x_{i+1}\}$ are edges of G.
- (iv) For $2 \le i \le n-1$, if x_{n+i} is adjacent to x_{n+1} in G, then $\{x_{n+i}, x_2\}$, $\{x_{n+1}, x_{i-1}\}$ and $\{x_{n+1}, x_{i+1}\}$ are edges of G.
- (v) For $2 \le i \le n-1$, if x_{n+i} is adjacent to x_{2n} in G, then $\{x_{n+i}, x_{n-1}\}$, $\{x_{i-1}, x_{2n}\}$ and $\{x_{i+1}, x_{2n}\}$ are edges of G; and
- (vi) If x_{n+1} and x_{2n} are adjacent in G, then $\{x_{n+1}, x_{n-1}\}$ and $\{x_2, x_{2n}\}$ are edges of G.

Then

$$\operatorname{reg}(G) \le \lceil \frac{n+2}{2} \rceil.$$

The following lemma has a crucial role in the proof of our main result.

Lemma 2.2. Let $n \ge 3$ be an integer and G be a graph with 2n vertices, say $\{x_1, x_2, \ldots, x_{2n}\}$. Assume that

- (i) $\{\{x_1, x_2\}, \{x_2, x_3\}, \dots, \{x_{n-1}, x_n\}, \{x_n, x_1\}\} \subseteq E(G).$
- (ii) $\{x_i, x_{n+i}\} \in E(G)$, for every $1 \le i \le n$.

(iii) For $1 \le i, j \le n$, if x_{n+i} is adjacent to x_{n+j} in G, then $\{x_{n+i}, x_{j-1}\}$, $\{x_{n+i}, x_{j+1}\}$, $\{x_{n+j}, x_{i-1}\}$ and $\{x_{n+j}, x_{i+1}\}$ are edges of G, where we consider the indices i-1, i+1, j-1 and j+1 modulo n.

Then

$$\operatorname{reg}(G) \le \lceil \frac{n+1}{2} \rceil.$$

Proof. We use induction on the number of vertices of G.

If n = 3, then one can check that G^c (the complement graph of G) is a chordal graph and thus reg(G) = 2. If n = 4, then G contains a whiskered path of length 4, which satisfies the assumption of lemma 2.1 and thus its regularity is at most 3. Now assume that $n \ge 5$. By assumption $|E(G)| \ge 2n$. If |E(G)| = 2n, then G is a whiskered cycle graph and it follows from Proposition 1.1 that

$$\operatorname{reg}(G) = \lceil \frac{n-1}{2} \rceil + 1 = \lceil \frac{n+1}{2} \rceil.$$

Suppose now that $|E(G)| \ge 2n + 1$. By induction on the number of edges of G, we prove that $\operatorname{reg}(G) \le \lfloor \frac{n+1}{2} \rfloor$. We consider the following cases.

Case 1. Assume that there exist $1 \le i < j \le n$ such that j - i is not congruent to one modulo n and $e = \{x_i, x_j\} \in E(G)$. Using induction hypothesis on the number of edges, it follows that

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+1}{2} \rceil.$$

By Theorem 1.2, it is enough to show that

$$\operatorname{reg}(G_e) \leq \lceil \frac{n-1}{2} \rceil.$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_i, x_{n+i}, x_j, x_{n+j}\}$ and let H' be the graph with the same vertex set as H and the edge set

$$E(H') = E(H) \cup \{\{x_{i-1}, x_{i+1}\}, \{x_{j-1}, x_{j+1}\}\} \cup \{\{x_{i-1}, x_{n+k}\} \mid 1 \le k \le n, k \ne i, j\} \cup \{\{x_{i+1}, x_{n+k}\} \mid 1 \le k \le n, k \ne i, j\} \cup \{\{x_{j-1}, x_{n+l}\} \mid 1 \le l \le n, l \ne i, j\} \cup \{\{x_{j+1}, x_{n+l}\} \mid 1 \le l \le n, l \ne i, j\},\$$

where we consider the indices i - 1, i + 1, j - 1 and j + 1 modulo n.

One can easily check that H' satisfies the assumptions of the lemma. Thus, by induction on the number of vertices,

$$\operatorname{reg}(H') \leq \left\lceil \frac{(n-2)+1}{2} \right\rceil = \left\lceil \frac{n-1}{2} \right\rceil.$$

Moreover, G_e is an induced subgraph of H' which implies that

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H') \le \lceil \frac{n-1}{2} \rceil.$$

4

Case 2. Assume that there exist $1 \le i < j \le n$ such that j - i is not congruent to one modulo n and $e = \{x_{n+i}, x_{n+j}\} \in E(G)$. By induction hypothesis on the number of edges, we have

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+1}{2} \rceil.$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_i, x_{n+i}, x_j, x_{n+j}\}$ and let H' be the graph with the same vertex set as H and the edge set

$$E(H') = E(H) \cup \{\{x_{i-1}, x_{i+1}\}, \{x_{j-1}, x_{j+1}\}\} \cup \{\{x_{i-1}, x_{n+k}\} \mid 1 \le k \le n, k \ne i, j\} \cup \{\{x_{i+1}, x_{n+k}\} \mid 1 \le k \le n, k \ne i, j\} \cup \{\{x_{j-1}, x_{n+l}\} \mid 1 \le l \le n, l \ne i, j\} \cup \{\{x_{j+1}, x_{n+l}\} \mid 1 \le l \le n, l \ne i, j\},\$$

where we consider the indices i - 1, i + 1, j - 1 and j + 1 modulo n.

One can easily check that H' satisfies the assumptions of the lemma. On the other hand, it follows from the assumptions $\{x_{n+i}, x_{j-1}\}$, $\{x_{n+i}, x_{j+1}\}$, $\{x_{n+j}, x_{i-1}\}$ and $\{x_{n+j}, x_{i+1}\}$ are edges of G. Thus, G_e is an induced subgraph of H'. Hence, the induction hypothesis on n and Theorem 1.2 show that

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H') \le \left\lceil \frac{(n-2)+1}{2} \right\rceil = \left\lceil \frac{n-1}{2} \right\rceil.$$

The conclusion that $\operatorname{reg}(G) \leq \lfloor \frac{n+1}{2} \rfloor$, follows from Theorem 1.2.

Case 3. Assume that there exists $1 \le i \le n$ such that $e = \{x_{n+i}, x_{n+i+1}\} \in E(G)$, (where by x_{2n+1} , we mean x_{n+1}). Then, by induction hypothesis on the number of edges,

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+1}{2} \rceil.$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_i, x_{n+i}, x_{i+1}, x_{n+i+1}\}$ and let H' be the graph with the same vertex set as H and the edge set

$$E(H') = E(H) \cup \{\{x_{i-1}, x_{i+2}\}\} \cup \{\{x_{i-1}, x_{n+k}\} \mid 1 \le k \le n, k \ne i, i+1\} \cup \{\{x_{i+2}, x_{n+k}\} \mid 1 \le k \le n, k \ne i, i+1\},\$$

where we consider the indices i - 1 and i + 2 modulo n.

One can easily check that H' satisfies the assumptions of the lemma. Thus, by induction hypothesis on n,

$$\operatorname{reg}(H') \leq \left\lceil \frac{(n-2)+1}{2} \right\rceil = \left\lceil \frac{n-1}{2} \right\rceil.$$

By assumptions, $\{x_{n+i}, x_{i+2}\}$ and $\{x_{n+i+1}, x_{i-1}\}$ are edges of G. Hence, G_e is an induced subgraph of H'. Therefore, by Theorem 1.2,

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H') \le \lceil \frac{n-1}{2} \rceil,$$

and then again by Theorem 1.2, we conclude that $\operatorname{reg}(G) \leq \lceil \frac{n+1}{2} \rceil$.

Case 4. Assume that there exist $1 \leq i < j \leq n$ such that j - i is not congruent to one modulo n and $e = \{x_{n+i}, x_j\} \in E(G)$. By cases 2 and 3, we may assume that for every $1 \leq k, l \leq n$ the vertices x_{n+k} and x_{n+l} are not adjacent in G. Then, by induction hypothesis on the number of edges,

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+1}{2} \rceil.$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_i, x_{n+i}, x_j, x_{n+j}\}$ and let H' be the graph with the same vertex set as H and the edge set

$$E(H') = E(H) \cup \{\{x_{i-1}, x_{j-1}\}, \{x_{i+1}, x_{j+1}\}\},\$$

where we consider the indices i - 1, i + 1, j - 1 and j + 1 modulo n. By considering the cycle

$$x_1, x_2, \ldots, x_{i-1}, x_{j-1}, x_{j-2}, \ldots, x_{i+1}, x_{j+1}, x_{j+2}, \ldots, x_n, x_1$$

one easily check that H' satisfies the assumptions of the lemma. On the other hand, G_e is an induced subgraph of H'. Thus, by induction hypothesis on the number of vertices, we have

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H') \le \left\lceil \frac{(n-2)+1}{2} \right\rceil = \left\lceil \frac{n-1}{2} \right\rceil.$$

It follows from Theorem 1.2 that $\operatorname{reg}(G) \leq \left\lceil \frac{n+1}{2} \right\rceil$.

Case 5. Assume that there exists $1 \le i \le n$ such that $e = \{x_{n+i}, x_{i+1}\} \in E(G)$, (where we consider the index i + 1 modulo n). By cases 2 and 3, we may assume that for every $1 \le k, l \le n$,

$$\{x_{n+k}, x_{n+l}\} \notin E(G).$$

Then, by induction hypothesis on the number of edges,

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+1}{2} \rceil$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_i, x_{n+i}, x_{i+1}, x_{n+i+1}\}$ and let H' be the graph with the same vertex set as H and the edge set

$$E(H') = E(H) \cup \{\{x_{i-1}, x_{i+2}\}\},\$$

where we consider the indices i - 1 and i + 2 modulo n. One easily check that H' satisfies the assumptions of the lemma. On the other hand, G_e is an induced subgraph of H' and by induction hypothesis on n,

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H') \le \lceil \frac{(n-2)+1}{2} \rceil = \lceil \frac{n-1}{2} \rceil$$

Therefore, by Theorem 1.2, $\operatorname{reg}(G) \leq \left\lceil \frac{n+1}{2} \right\rceil$.

To prove the main result, we also need the following two lemmas.

Lemma 2.3. Let $G = W(C_n)$ be a whiskered cycle graph with edge ideal I = I(G)and assume that the vertices of the n-cycle (in order) are x_1, \ldots, x_n and the whiskers are the edges $\{x_1, x_{n+1}\}, \ldots, \{x_n, x_{2n}\}$. Then for a minimal generator M of I^s , we have $x_n^2 \in (I^{s+1}: M)$ if and only if n is odd, say n = 2m + 1 for some $1 \le m \le s$ and

$$M = (x_1 x_2)(x_3 x_4) \dots (x_{2m-1} x_{2m})N,$$

for some $N \in I^{s-m}$.

Moreover, in this case $x_n x_j \in (I^{s+1} : M)$ for all j = 1, ..., 2n.

Proof. First assume that n = 2m + 1 is odd and $M = (x_1 x_2)(x_3 x_4) \dots (x_{2m-1} x_{2m})N$ for some $N \in I^{s-m}$. Then $x_n^2 M = (x_n x_1)(x_2 x_3) \dots (x_{2m} x_n)N \in I^{s+1}$.

Now, assume that $x_n^2 \in (I^{s+1}: M)$. Using an argument similar to the proof of [3, Lemma 5.1], one concludes that n is odd, say n = 2m + 1 for some $1 \le m \le s$ and

$$M = (x_1 x_2)(x_3 x_4) \dots (x_{2m-1} x_{2m}) N,$$

for some $N \in I^{s-m}$.

By Theorem 1.4, to prove the last statement of the lemma, it is enough to show that for every j = 1, ..., 2n, the vertices x_n and x_j are even-connected with respect to M. By assumptions, there is nothing to prove if j = n or 2n. Hence assume that $j \neq n, 2n$. We consider the following cases.

Case 1. If $1 \leq j < n$ is odd, then x_n, x_1, \ldots, x_j is an even-connected path between x_n and x_j with respect to M (remember that $M = (x_1x_2)(x_3x_4) \ldots (x_{2m-1}x_{2m})N$, for some $N \in I^{s-m}$).

Case 2. If $1 \leq j < n$ is even, then $x_j, x_{j+1}, \ldots, x_n$ is an even-connected path between x_j and x_n with respect to M (remember that n is odd).

Case 3. Assume that $n + 1 \leq j < 2n$. If j is even, then $x_n, x_{n-1}, \ldots, x_{j-n}, x_j$ is an even-connected path between x_n and x_j with respect to M. If j is odd, then $x_n, x_1, \ldots, x_{j-n}, x_j$ is an even-connected path between x_n and x_j with respect to M.

Thus, $x_n x_j \in (I^{s+1} : M)$ for all j = 1, ..., 2n and the lemma is proved.

Lemma 2.4. With the same assumptions as Lemma 2.3, $x_{n+1}^2 \in (I^{s+1} : M)$ if and only if n is odd, say n = 2m - 1 for some $2 \le m \le s$ and

$$M = (x_1 x_2)(x_3 x_4) \dots (x_{2m-3} x_{2m-2})(x_{2m-1} x_1)N,$$

for some $N \in I^{s-m}$.

Moreover, in this case $x_{n+1}x_j \in (I^{s+1}:M)$ for all $j = 1, \ldots, 2n$.

Proof. First suppose that n = 2m - 1 is odd and

$$M = (x_1 x_2)(x_3 x_4) \dots (x_{2m-3} x_{2m-2})(x_{2m-1} x_1) N$$

for some $N \in I^{s-m}$. Then

$$x_{n+1}^2 M = (x_{n+1}x_1)(x_2x_3)\dots(x_{2m-2}x_{2m-1})(x_1x_{n+1})N \in I^{s+1}.$$

Now, assume $x_{n+1}^2 \in (I^{s+1} : M)$. Then Theorem 1.4 implies that x_{n+1} is evenconnected to itself with respect to M. Let $x_{n+1} = p_0, p_1, \ldots, p_{2m+1} = x_{n+1}$ be a shortest even-connected path between x_{n+1} and itself. Assume that there exists $1 \leq j \leq 2k$ such that $p_j = x_{n+1}$. If j is odd then $x_{n+1} = p_0, \ldots, p_j = x_{n+1}$ is a shorter even-connected path between x_{n+1} and itself, a contradiction. If j is even then $x_{n+1} = p_j, \ldots, p_{2m+1} = x_{n+1}$ is again a shorter even-connected path between x_{n+1} and itself, a contradiction. Thus, we assume that x_{n+1} does not appear in the path p_0, \ldots, p_{2m+1} except at its endpoints.

We note that since $\deg_G(x_{n+1}) = 1$, then the even-connected path p_0, \ldots, p_{2m+1} can not be simple. Hence, there exist indices i and j with $1 \le i < j \le 2k$ such that $p_i = p_j$ and we choose i and j such that j - i is minimal. Then p_i, \ldots, p_j is a simple closed path in G. This can only occur if this simple path is C_n . If the path p_0, \ldots, p_{2m+1} contains at least two copies of C_n , then by removing the edges of these two copies, we obtain a shorter even-connected path. Thus, the path p_0, \ldots, p_{2m+1} contains exactly one copy of C_n . This shows that the even-connected path p_0, \ldots, p_{2m+1} is of the form $x_{n+1}, x_1, x_2, \ldots, x_n, x_1, x_{n+1}$. Therefore, 2m + 2 = n + 3 and hence n = 2m - 1 is odd. By re-indexing if necessary, we may assume that $p_i = x_i$ for $i = 1, \ldots, 2m - 1$ and $p_{2m} = x_1$. Moreover, by the definition of even-connected path, we have

$$M = (p_1 p_2) \dots (p_{2m-1} p_{2m}) N = (x_1 x_2) \dots (x_{2m-1} x_1) N$$

where $N \in I^{s-m}$.

Using Theorem 1.4, in order to prove the last statement of the lemma, we must show that for every j = 1, ..., 2n, the vertices x_{n+1} and x_j are even-connected with respect to M. By assumptions, there is nothing to prove if j = 1 or n + 1. Hence assume that $j \neq 1, n + 1$. We consider the following cases.

Case 1. If $1 < j \leq n$ is odd, then $x_{n+1}, x_1, \ldots, x_j$ is an even-connected path between x_{n+1} and x_j .

Case 2. If $1 < j \le n$ is even, then $x_j, x_{j+1} \ldots, x_1, x_{n+1}$ is an even-connected path between x_j and x_n .

Case 3. Assume that $n + 1 < j \leq 2n$.

If j is odd, then $x_{n+1}, x_1, x_2, x_3, \ldots, x_{j-n}, x_j$ is an even-connected path between x_{n+1} and x_j .

If j is even, then $x_{n+1}, x_1, x_n, x_{n-1}, x_{n-2}, \ldots, x_{j-n}, x_j$ is an even-connected path between x_{n+1} and x_j .

Therefore, $x_{n+1}x_j \in (I^{s+1}:M)$ for all $j = 1, \dots, 2n$.

We are now ready to prove the main result of this paper.

Theorem 2.5. Let $G = W(C_n)$ be a whiskered cycle graph and I = I(G) be its edge ideal. Then for all $s \ge 1$, we have

$$\operatorname{reg}(I^s) = 2s + \lceil \frac{n-1}{2} \rceil - 1 = 2s + \operatorname{indmatch}(G) - 1.$$

Proof. One can easily check that $\operatorname{indmatch}(G) = \lceil \frac{n-1}{2} \rceil$. Thus, the second inequality is obvious. To prove the first inequality, note that by [3, Theorem 4.5], the inequality

$$\operatorname{reg}(I^s) \ge 2s + \operatorname{indmatch}(G) - 1$$

is known. Therefore, we must prove that for every $s \ge 1$ the inequality

$$\operatorname{reg}(I^s) \le 2s + \lceil \frac{n-1}{2} \rceil - 1$$

holds. For s = 1 the above inequality is known (see Proposition 1.1). By applying [2, Theorem 5.2] and using induction on s, it is enough to prove that for every $s \ge 1$ and every minimal generator M of I^s ,

$$\operatorname{reg}(I^{s+1}:M) \le \lceil \frac{n-1}{2} \rceil + 1.$$

By Theorem 1.4, the ideal $(I^{s+1}: M)$ is generated by the quadratics uv, where it is either an edge ideal of G or u and v are even-connected with respect to M. Let J denote the polarization of the ideal $(I^{s+1}: M)$. Assume that $x_{i_1}^2, \ldots, x_{i_t}^2$ are the non-squarefree minimal generators of $(I^{s+1}: M)$. Then

$$J = I(G') + (x_{i_1}y_{i_1}, \dots, x_{i_t}y_{i_t}),$$

where G' is a graph over the vertices x_1, \ldots, x_{2n} and y_{i_1}, \ldots, y_{i_t} are new variables. Since polarization does not change the regularity, we have $\operatorname{reg}(J) = \operatorname{reg}(I^{s+1}:M)$.

On the other hand, since $(I^{s+1}: M)$ has all edges of G as minimal generators, G is a subgraph of G'. For every $j = 0, \ldots, t$, let H_j be the graph whose edge ideal is

$$I(G') + (x_{i_1}y_{i_1}, \dots, x_{i_j}y_{i_j}).$$

Then, $H_0 = G'$ and $J = I(H_t)$. By Lemmas 2.3 and 2.4, we observe that $\{x_{i_j}, x_l\} \in E(G')$ for every $j = 1, \ldots, t$ and every $l = 1, \ldots, 2n$ with $i_j \neq l$. This implies that for every $1 \leq j \leq t$, the graph $H_j \setminus N_{H_j}[x_{i_j}]$ consists of isolated vertices $\{y_{i_1}, \ldots, y_{i_{j-1}}\}$. Hence,

$$\operatorname{reg}(H_j \setminus N_{H_j}[x_{i_j}]) = 0.$$

Now, by [6, Lemma 2.10] we have

$$\operatorname{reg}(H_j) = \operatorname{reg}(H_j \setminus x_{i_j}).$$

On the other hand, y_{i_j} is an isolated vertex in $H_j \setminus x_{i_j}$ and $H_j \setminus \{x_{i_j}, y_{i_j}\}$ is an induced subgraph of $H_j \setminus y_{i_j} = H_{j-1}$. Hence,

$$\operatorname{reg}(H_j) = \operatorname{reg}(H_j \setminus x_{i_j}) = \operatorname{reg}(H_j \setminus \{x_{i_j}, y_{i_j}\}) \le \operatorname{reg}(H_j \setminus y_{i_j}) = \operatorname{reg}(H_{j-1}).$$

Note that H_{j-1} is an induced subgraph of H_j and this implies that $\operatorname{reg}(H_{j-1}) \leq \operatorname{reg}(H_j)$. Therefore, we obtain that $\operatorname{reg}(H_j) = \operatorname{reg}(H_{j-1})$ for all $j = 1, \ldots, t$.

This, in particular, implies that

$$\operatorname{reg}(J) = \operatorname{reg}(H_t) = \operatorname{reg}(H_0) = \operatorname{reg}(G')$$

To complete the proof, it is enough to show that $\operatorname{reg}(G') \leq \lfloor \frac{n-1}{2} \rfloor + 1$.

If $\{x_{n+i}, x_{n+j}\} \in E(G')$, for some $1 \leq i, j \leq n$, then $x_{n+i}x_{n+j} \in (I^{s+1}: M)$. This means that there exists an even-connected path $x_{n+i} = p_0, p_1, \ldots, p_{2m+1} = x_{n+j}$ in G, between x_{n+i} and x_{n+j} . Since $N_G(x_{n+i}) = \{x_i\}$, we conclude that $p_1 = x_i$. Thus, $x_{i+1}, p_1, \ldots, p_{2m+1} = x_{n+j}$ is an even-connected path between x_{i+1} and x_{n+j} . Also, $x_{i-1}, p_1, \ldots, p_{2m+1} = x_{n+j}$ is an even-connected path between x_{i-1} and x_{n+j} . Similarly, one can show that x_{n+i} is even-connected to both x_{j-1} and x_{j+1} (where we consider the indices i - 1, i + 1, j - 1 and j + 1 modulo n). This shows that

$$\{x_{i-1}, x_{n+j}\}, \{x_{i+1}, x_{n+j}\}, \{x_{j-1}, x_{n+i}\}, \{x_{j+1}, x_{n+i}\} \in E(G')$$

It follows from Lemma 2.2 that $\operatorname{reg}(G') \leq \lceil \frac{n+1}{2} \rceil$ and this completes the proof. \Box

3. Proof of Lemma 2.1

In this section, we prove Lemma 2.1.

Proof. We use induction on n. Since every connected graph with at most four vertices has chordal complement, the result is true for $n \leq 2$. Set

$$d = \deg_G(x_{n+1}) + \deg_G(x_{n+2})$$

We prove the assertion by induction on d. Note that $d \ge 2$.

We first consider the case d = 2. In this case, let $H = G \setminus \{x_1, x_{n+1}\}$ and note that in the graph $G \setminus x_1$, the vertex x_{n+1} is an isolated vertex. Thus, $\operatorname{reg}(G \setminus x_1) = \operatorname{reg}(H)$. Hence, by the induction hypothesis on n, we get

$$\operatorname{reg}(G \setminus x_1) = \operatorname{reg}(H) \le \left\lceil \frac{(n-1)+2}{2} \right\rceil \le \left\lceil \frac{n+2}{2} \right\rceil.$$
(3.1)

Let K be the induced subgraph of G over the vertices $V(G) \setminus \{x_1, x_{n+1}, x_2, x_{n+2}\}$. Clearly, $G \setminus (N[x_1] \cup \{x_{n+2}\})$ is an induced subgraph of K. So, by Theorem 1.2,

$$\operatorname{reg}(G \setminus (N[x_1] \cup \{x_{n+2}\})) \le \operatorname{reg}(K).$$

On the other hand, $\deg_G(x_{n+2}) = 1$ and x_{n+2} is an isolated vertex in $G \setminus N[x_1]$. Thus,

$$\operatorname{reg}(G \setminus N[x_1]) = \operatorname{reg}(G \setminus (N[x_1] \cup \{x_{n+2}\}))$$

and hence, $\operatorname{reg}(G \setminus N[x_1]) \leq \operatorname{reg}(K)$. Moreover, by induction on n, we have

$$\operatorname{reg}(K) \le \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$
(3.2)

It then follows from Theorem 1.2, together with (3.1) and (3.2) that

$$\operatorname{reg}(G) \le \max\{\operatorname{reg}(G \setminus x_1), \operatorname{reg}(G \setminus N[x_1]) + 1\} \le \lceil \frac{n+2}{2} \rceil.$$

Now assume that d > 2. In this case, there is a vertex $x_t \in \{x_1, \ldots, x_n, x_{n+1}, \ldots, x_{2n}\}$ such that either

- (i) $t \neq 1$ and $\{x_{n+1}, x_t\} \in E(G)$ or
- (ii) $t \neq 2$ and $\{x_{n+2}, x_t\} \in E(G)$.

Case 1. Assume that $e = \{x_{n+1}, x_{n+2}\} \in E(G)$. Then

 $\deg_{G\setminus e}(x_{n+1}) + \deg_{G\setminus e}(x_{n+2}) < \deg_G(x_{n+1}) + \deg_G(x_{n+2}).$

So, by induction hypothesis on d, we have

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+2}{2} \rceil.$$

Let *H* be the induced subgraph of *G* over the vertices $V(G) \setminus \{x_1, x_{n+1}, x_2, x_{n+2}\}$. Then, by induction on *n*,

$$\operatorname{reg}(H) \leq \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$

Moreover, G_e is an induced subgraph of H and it follows from Theorem 1.2 that $\operatorname{reg}(G_e) \leq \operatorname{reg}(H) \leq \lceil \frac{n}{2} \rceil$. Hence, by Theorem 1.2, we conclude that

$$\operatorname{reg}(G) \le \lceil \frac{n+2}{2} \rceil.$$

Case 2. Assume that $e = \{x_{n+1}, x_2\} \in E(G)$ and $\{x_{n+1}, x_{n+3}\} \notin E(G)$. Then

 $\deg_{G\setminus e}(x_{n+1}) + \deg_{G\setminus e}(x_{n+2}) < \deg_G(x_{n+1}) + \deg_G(x_{n+2}),$

and since $\{x_{n+1}, x_{n+3}\} \notin E(G)$, it follows that $G \setminus e$ satisfies the assumptions of the lemma. So, by induction hypothesis on d, we have

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+2}{2} \rceil.$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_1, x_{n+1}, x_2, x_{n+2}\}$. Then, by induction on n,

$$\operatorname{reg}(H) \leq \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$

Moreover, G_e is an induced subgraph of H and it follows from Theorem 1.2 that $\operatorname{reg}(G_e) \leq \operatorname{reg}(H) \leq \lceil \frac{n}{2} \rceil$. Hence, by Theorem 1.2, we conclude that

$$\operatorname{reg}(G) \le \lceil \frac{n+2}{2} \rceil.$$

Case 3. Assume that $e = \{x_{n+1}, x_2\}$ and $e' = \{x_{n+1}, x_{n+3}\}$ are edges of G. Since in $G \setminus e'$ we have

$$\deg_{G\setminus e'}(x_{n+1}) + \deg_{G\setminus e'}(x_{n+2}) < \deg_G(x_{n+1}) + \deg_G(x_{n+2})$$

by induction on d, it follows that

$$\operatorname{reg}(G \setminus e') \le \lceil \frac{n+2}{2} \rceil.$$

Using Theorem 1.2, to show that $\operatorname{reg}(G) \leq \lceil \frac{n+2}{2} \rceil$, it suffices to prove that $\operatorname{reg}(G_{e'}) \leq \lceil \frac{n}{2} \rceil$.

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_1, x_{n+1}, x_3, x_{n+3}\}$ and let H' be the graph with the same vertex set as H and the edge set

 $E(H') = E(H) \cup \{\{x_2, x_4\}\} \cup \{\{x_2, x_{n+i}\} \mid 4 \le i \le n\} \cup \{\{x_4, x_{n+i}\} \mid 2 \le i \le n, i \ne 3\}.$ Then, H' satisfies the assumptions of the lemma. Hence, by induction hypothesis on n, we see that

$$\operatorname{reg}(H') \leq \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$

Moreover, $G_{e'}$ is an induced subgraph of H' (since x_2 and x_4 are not vertices of $G_{e'}$), which implies that

$$\operatorname{reg}(G_{e'}) \le \operatorname{reg}(H') \le \lceil \frac{n}{2} \rceil.$$

Case 4. Assume that $e = \{x_{n+2}, x_1\}$ is an edge of G. Then

$$\deg_{G\setminus e}(x_{n+1}) + \deg_{G\setminus e}(x_{n+2}) < \deg_G(x_{n+1}) + \deg_G(x_{n+2})$$

Thus, by induction on d,

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+2}{2} \rceil.$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_1, x_{n+1}, x_2, x_{n+2}\}$. Then, G_e is an induced subgraph of H and the induction hypothesis implies that

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H) \le \lceil \frac{n}{2} \rceil.$$

It follows from Theorem 1.2 that

$$\operatorname{reg}(G) \leq \lceil \frac{n+2}{2} \rceil.$$

Case 5. Assume that $3 \le t \le n$ and $e = \{x_{n+1}, x_t\} \in E(G)$ and $\{x_{n+1}, x_{n+t+1}\} \notin E(G)$.

If t = 3, then $e = \{x_{n+1}, x_3\} \in E(G)$ and $\{x_{n+1}, x_{n+4}\} \notin E(G)$. If $\{x_{n+1}, x_{n+2}\}$ is an edge of G, then the assertion follows from case 1.

If $t \ge 4$ and $\{x_{n+1}, x_{n+t-1}\}$ is an edge of G, then by assumption, $\{x_{n+1}, x_{t-2}\}$ is an edge of G, which is a contradiction by the choice of t.

Thus, we assume that $\{x_{n+1}, x_{n+t-1}\} \notin E(G)$. Since $\{x_{n+1}, x_{n+t-1}\}$ and $\{x_{n+1}, x_{n+t+1}\}$ are not edges of G, it follows that $G \setminus e$ satisfies the assumptions of the lemma. Now,

$$\deg_{G\setminus e}(x_{n+1}) + \deg_{G\setminus e}(x_{n+2}) < \deg_G(x_{n+1}) + \deg_G(x_{n+2}).$$

Thus, by induction on d,

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+2}{2} \rceil.$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_1, x_{n+1}, x_3, x_{n+3}\}$ and let H' be the graph with the same vertex set as H and the edge set

$$E(H') = E(H) \cup \{\{x_2, x_4\}\} \cup \{\{x_2, x_{n+i}\} \mid 4 \le i \le n\} \cup \{\{x_4, x_{n+i}\} \mid 2 \le i \le n, i \ne 3\}$$

Then G_e is an induced subgraph of H' (since x_2 and x_4 are not vertices of G_e). Therefore, by induction on n and Theorem 1.2, we have

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H') \le \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$

The conclusion that $\operatorname{reg}(G) \leq \left\lceil \frac{n+2}{2} \right\rceil$ now follows from Theorem 1.2.

Case 6. Assume that $3 \leq t \leq n$ and $e = \{x_{n+1}, x_t\}$ and suppose that $e' = \{x_{n+1}, x_{n+t+1}\}$ are edges of G. Then

$$\deg_{G\setminus e'}(x_{n+1}) + \deg_{G\setminus e'}(x_{n+2}) < \deg_G(x_{n+1}) + \deg_G(x_{n+2})$$

and by induction on d,

$$\operatorname{reg}(G \setminus e') \le \lceil \frac{n+2}{2} \rceil.$$

According to Theorem 1.2, to prove that $\operatorname{reg}(G) \leq \lceil \frac{n+2}{2} \rceil$, it remains to show that

$$\operatorname{reg}(G_{e'}) \leq \lceil \frac{n}{2} \rceil.$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_1, x_{n+1}, x_{t+1}, x_{n+t+1}\}$ and let H' be the graph with the same vertex set as H and the edge set

$$E(H') = E(H) \cup \{\{x_t, x_{t+2}\}\} \cup \{\{x_t, x_{n+i}\} \mid 2 \le i \le n, i \ne t+1\} \\ \cup \{\{x_{t+2}, x_{n+i}\} \mid 2 \le i \le n, i \ne t+1\}.$$

One can easily check that H' satisfies the assumptions of the lemma. Hence by induction on n, we conclude that

$$\operatorname{reg}(H') \leq \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$

Moreover, $G_{e'}$ is an induced subgraph of H' (since x_t and x_{t+2} are not vertices of $G_{e'}$). This implies that $\operatorname{reg}(G_{e'}) \leq \operatorname{reg}(H') \leq \lceil \frac{n}{2} \rceil$ and the result follows.

Case 7. Assume that $n + 3 \le t \le 2n$ and $\{x_{n+1}, x_t\}$ is an edge of G. Then, by assumption, $\{x_{n+1}, x_{t-n-1}\}$ is an edge of G, which is contradiction by the choice of t.

Case 8. Assume that $3 \le t \le n$ and suppose that $e = \{x_{n+2}, x_t\}$ and $e' = \{x_{n+2}, x_{n+t+1}\}$ are edges of G. In the graph $G \setminus e'$, we have

$$\deg_{G\setminus e'}(x_{n+1}) + \deg_{G\setminus e'}(x_{n+2}) < \deg_G(x_{n+1}) + \deg_G(x_{n+2})$$

So, by induction on d,

$$\operatorname{reg}(G \setminus e') \le \lceil \frac{n+2}{2} \rceil.$$

It follows from Theorem 1.2 that

$$\operatorname{reg}(G) \le \max\{\operatorname{reg}(G \setminus e'), \operatorname{reg}(G_{e'}) + 1\}.$$

Thus, in order to prove that $\operatorname{reg}(G) \leq \lceil \frac{n+2}{2} \rceil$, it is enough to show that

$$\operatorname{reg}(G_{e'}) \le \lceil \frac{n}{2} \rceil. \tag{3.3}$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_2, x_{n+2}, x_{t+1}, x_{n+t+1}\}$ and let H' be the graph with the same vertex set as H and the edge set

$$E(H') = E(H) \cup \{\{x_1, x_3\}, \{x_t, x_{t+2}\}\} \cup \{\{x_1, x_{n+i}\} \mid 3 \le i \le n, i \ne t+1\} \cup \{\{x_3, x_{n+i}\} \mid 1 \le i \le n, i \ne 2, t+1\} \cup \{\{x_t, x_{n+i}\} \mid 1 \le i \le n, i \ne 2, t+1\} \cup \{\{x_{t+2}, x_{n+i}\} \mid 1 \le i \le n, i \ne 2, t+1\}.$$

One can easily check that H' satisfies the assumptions of the lemma. Hence by induction on n,

$$\operatorname{reg}(H') \le \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$

Moreover, $G_{e'}$ is an induced subgraph of H' (since x_1, x_3, x_t and x_{t+2} are not vertices of $G_{e'}$). So, by Theorem 1.2,

$$\operatorname{reg}(G_{e'}) \le \operatorname{reg}(H') \le \lceil \frac{n}{2} \rceil$$

and (3.3) holds.

Case 9. Assume that $e = \{x_{n+2}, x_{n+3}\}$ is an edge of G. By induction on d,

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+2}{2} \rceil.$$

Now, using Theorem 1.2

$$\operatorname{reg}(G) \le \max\{\operatorname{reg}(G \setminus e), \operatorname{reg}(G_e) + 1\}.$$

Thus, it is enough to show that

$$\operatorname{reg}(G_e) \leq \lceil \frac{n}{2} \rceil.$$

Let H be the induced subgraph of G over the vertices $V(G) \setminus \{x_2, x_{n+2}, x_3, x_{n+3}\}$ and let H' be the graph with the same vertex set as H and the edge set

$$E(H') = E(H) \cup \{\{x_1, x_4\}\} \cup \{\{x_1, x_{n+i}\} \mid 4 \le i \le n\} \cup \{\{x_4, x_{n+i}\} \mid 1 \le i \le n, i \ne 2.3\}.$$

One can easily check that H' satisfies the assumptions of the lemma. Therefore, by induction on n, we have

$$\operatorname{reg}(H') \leq \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$

Moreover, G_e is an induced subgraph of H' (since x_1 and x_4 are not vertices of G_e) which implies that

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H') \le \lceil \frac{n}{2} \rceil.$$

Case 10. Assume that $e = \{x_{n+2}, x_3\}$ is an edge of G and $\{x_{n+2}, x_{n+4}\}$ is not an edge of G. Note that

$$\deg_{G\setminus e}(x_{n+1}) + \deg_{G\setminus e}(x_{n+2}) < \deg_G(x_{n+1}) + \deg_G(x_{n+2})$$

Since $\{x_{n+2}, x_{n+4}\}$ is not an edge of G, we conclude that $G \setminus e$ satisfies the assumptions of the lemma. So, by induction on d,

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+2}{2} \rceil.$$

Let *H* be the induced subgraph of *G* over the vertices $V(G) \setminus \{x_2, x_{n+2}, x_3, x_{n+3}\}$ and let *H'* be the graph with the same vertex set as *H* and the edge set

$$E(H') = E(H) \cup \{\{x_1, x_4\}\} \cup \{\{x_4, x_{n+i}\} \mid 1 \le i \le n, i \ne 2, 3\}$$
$$\cup \{\{x_1, x_{n+i}\} \mid \{x_{n+4}, x_{n+i}\} \in E(G), i \ne 2, 3\}.$$

One can easily check that H' satisfies the assumptions of the lemma. Note that x_4 is not a vertex of G_e . On the other hand, if $\{x_{n+4}, x_{n+i}\} \in E(G)$, for some $i \neq 2, 3$, then the assumptions of the lemma implies that $\{x_3, x_{n+i}\} \in E(G)$. Consequently, x_{n+i} is not a vertex of G_e . Therefore, G_e is an induced subgraph of H'. Hence, by Theorem 1.2 and by induction on n,

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H') \le \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$

Finally, using Theorem 1.2, we have,

$$\operatorname{reg}(G) \le \lceil \frac{n+2}{2} \rceil.$$

Case 11. Assume that $4 \le t \le n$ and suppose that $e = \{x_{n+2}, x_t\}$ is an edge of G and $\{x_{n+2}, x_{n+t+1}\}$ is not an edge of G.

If t = 4 and $\{x_{n+2}, x_{n+3}\} \in E(G)$, then the assertion follows from case 9.

If $t \geq 5$ and $\{x_{n+2}, x_{n+t-1}\} \in E(G)$, then by assumption, $\{x_{n+2}, x_{t-2}\} \in E(G)$, which is a contradiction by the choice of t.

Thus, we assume that $\{x_{n+2}, x_{n+t-1}\} \notin E(G)$. In this case, we observe that

$$\deg_{G\setminus e}(x_{n+1}) + \deg_{G\setminus e}(x_{n+2}) < \deg_G(x_{n+1}) + \deg_G(x_{n+2}).$$

Since $\{x_{n+2}, x_{n+t-1}\}$ and $\{x_{n+2}, x_{n+t+1}\}$ are not edges of G, we conclude that $G \setminus e$ satisfies the assumptions of the lemma. So, by induction on d,

$$\operatorname{reg}(G \setminus e) \le \lceil \frac{n+2}{2} \rceil$$

Let *H* be the induced subgraph of *G* over the vertices $V(G) \setminus \{x_2, x_{n+2}, x_t, x_{n+t}\}$ and let *H'* be the graph with the same vertex set as *H* and the edge set

$$E(H') = E(H) \cup \{\{x_1, x_{t-1}\}, \{x_3, x_{t+1}\}\} \cup \{\{x_{t-1}, x_{n+i}\} \mid 1 \le i \le n, i \ne 2, t\} \cup \{\{x_{t+1}, x_{n+i}\} \mid 1 \le i \le n, i \ne 2, t\} \cup \{\{x_1, x_{n+i}\} \mid \{x_{n+t-1}, x_{n+i}\} \in E(G), i \ne 2, t\} \cup \{\{x_3, x_{n+i}\} \mid \{x_{n+t+1}, x_{n+i}\} \in E(G), i \ne 2, t\}.$$

By considering the path

$$x_1, x_{t-1}, x_{t-2}, \ldots, x_4, x_3, x_{t+1}, x_{t+2}, \ldots, x_n,$$

one can easily check that H' satisfies the assumptions of the lemma. Note that x_{t-1} and x_{t+1} are not vertices of G_e . On the other hand, if $\{x_{n+t-1}, x_{n+i}\} \in E(G)$, for some $i \neq 2, t$, then the assumptions of the lemma implies that $\{x_t, x_{n+i}\} \in E(G)$. Consequently, x_{n+i} is not a vertex of G_e . Similarly, if $\{x_{n+t+1}, x_{n+i}\} \in E(G)$, for some $i \neq 2, t$, then x_{n+i} is not a vertex of G_e . Therefore, G_e is an induced subgraph of H'. Hence, by Theorem 1.2 and by induction on n,

$$\operatorname{reg}(G_e) \le \operatorname{reg}(H') \le \lceil \frac{(n-2)+2}{2} \rceil = \lceil \frac{n}{2} \rceil.$$

Finally, using Theorem 1.2, we have,

$$\operatorname{reg}(G) \le \lceil \frac{n+2}{2} \rceil.$$

Case 12. Assume that $n + 4 \le t \le 2n$ and $e = \{x_{n+2}, x_t\}$ is an edge of G. Then by assumption, $\{x_{n+2}, x_{t-n-1}\} \in E(G)$, which is a contradiction by the choice of t. \Box

References

- [1] A. Aliooee, A. Banerjee, Powers of edge ideals of regularity three bipartite graphs, preprint.
- [2] A. Banerjee, The regularity of powers of edge ideals, preprint.
- [3] S. Beyarslan, H. T. Hà, T. N. Trung, Regularity of powers of forests and cycles, preprint.
- [4] J. Biermann, A. Van Tuyl, Balanced vertex decomposable simplicial complexes and their hvectors, Electron. J. Combin. 20 (2013), no. 3, Paper 15.
- [5] A. Conca, Regularity jumps for powers of ideals, Commutative algebra,2132, Lect. Notes pure Appl., Queens Papers in Pure and Appl. *Math.*, 244, 1–40. Chapman Hall/CRC, Boca Raton, FL, 2006. Math. 20, Kingston, Ontario: Queens University, 1969.
- [6] H. Dao, C. Huneke, J. Schweig, Bounda on the regularity and projective dimension of ideals associated to graphs, J. Algebraic Combin. 38 (2013), 37–55.
- [7] C. Ferro, M. Murgia, O. Olteanu, Powers of edge ideals, Matematiche (Catania), 67 (2012), no. 1, 129–144.
- [8] R. Fröberg, On Stanley-Reisner rings, Topics in algebra, Part 2(Warsaw, 1998), 5770, Banach Center Publ., 26, Part 2, PWN, Warsaw, 1990.
- [9] H. T. Hà, Regularity of squarefree monomial ideals, In S.M. Copper and S. Sather-Wagstaff(Ed.) Connections Between Algebra, Combinatorics, and Geometry. Springer Proceedings in Mathematics Statistics 76 (2014), 251–276.
- [10] J. Herzog, T. Hibi, X. Zheng, Monomial ideals whose powers have a linear resolution, *Math. Scand.* 95 (2004), no. 1, 23–32.
- [11] S. Jacques, Betti numbers of graph ideals, ph.D. Thesis, University of sheffield, 2004.
- [12] S. Morey, R. H. Villarreal, Edge ideals: algebraic and combinatorial properties, Progress in commutative algebra 1, 85–126, de Gruyter, Berlin, 2012.

M. Moghimian, Science and Research Branch, Islamic Azad University (IAU), Tehran, Iran.

E-mail address: math_moghimi@yahoo.com

S. A. Seyed Fakhari, School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box 19395-5746, Tehran, Iran.

E-mail address: fakhari@ipm.ir *URL*: http://math.ipm.ac.ir/~fakhari/

S. YASSEMI, SCHOOL OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE, COLLEGE OF SCIENCE, UNIVERSITY OF TEHRAN, TEHRAN, IRAN, AND SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), P.O. BOX 19395-5746, TEHRAN, IRAN.

E-mail address: yassemi@ipm.ir

URL: http://math.ipm.ac.ir/~yassemi/