

Adaptive importance sampling via minimization of estimators of cross-entropy, mean square, and inefficiency constant

Tomasz Badowski *

Freie Universität Berlin

July 26, 2022

Abstract

Inefficiency constant, equal to product of variance of an estimator and its mean computation cost, can be used for quantifying the inefficiency of using unbiased estimators in Monte Carlo procedures. We develop single- and multi-stage minimization methods for obtaining parameters of importance sampling (IS) change of measure via minimization of well-known estimators of cross-entropy and mean square (of the IS estimator), as well as of new estimators of mean square and inefficiency constant. We prove convergence and asymptotic properties of minimization results in our methods, in the proofs for the multi-stage methods using a new uniform strong law of large numbers. We show that if the zero-variance IS parameter exists, then, under appropriate assumptions, minimization results of the new estimators converge to such parameter at a faster rate than such results of the well-known estimators, and a positive definite asymptotic covariance matrix of minimization results of the cross-entropy estimator is four times such matrix for the well-known mean square estimator. We provide an analytical example in which all relations between the asymptotic variances of minimization results of different estimators converging to the same point are possible, except that such variance for the well-known mean square estimator is always not higher than for the cross-entropy one. We introduce criteria for comparing asymptotic efficiency of stochastic optimization methods, applicable to methods minimizing estimators of functions considered in this work. In our numerical experiments for computing expectations of functionals of Euler scheme, minimization of the new estimators led to the lowest inefficiency constants and variances, followed by the well-known mean square estimator, and the cross-entropy one.

*Supported by the Berlin Mathematical School.

Contents

1	Introduction	4
1.1	Problem statement and literature overview	4
1.2	Overview of own work	7
1.3	Outline of the paper	12
1.4	Basic notations	12
2	Interpretations of inefficiency constants	13
3	Importance sampling	17
3.1	Background on densities	17
3.2	IS and zero- and optimal-variance IS distributions	17
3.3	Mean costs and inefficiency constants in IS	19
3.4	IS using parametric families of distributions	20
4	Examples of parametric families of IS distributions	22
4.1	Exponential change of measure	22
4.2	IS for independently parametrized product distributions	24
4.3	IS for stopped sequences	25
4.3.1	Change of measure for stopped sequences using a tilting process	25
4.3.2	Parametrizations of IS for stopped sequences	27
4.3.3	Change of measure for Gaussian stopped sequences	27
4.3.4	Linearly parametrized exponential tilting for stopped sequences	29
4.4	IS for functionals of Brownian motion up to a stopping time	30
4.5	IS for diffusions and Euler scheme	31
4.6	Some special functionals of diffusions and Euler scheme, and zero-variance IS for diffusions	33
4.7	Some special functionals of diffusions and Euler scheme	34
4.8	Diffusion in a potential	35
4.9	The special cases considered in our numerical experiments	36
5	The minimized functions and their estimators	38
5.1	The minimized functions	38
5.2	Estimators of the minimized functions	40
5.3	Cross-entropy and its estimator for ECM	42
5.4	Some conditions in the LETS setting	43
5.5	Some conditions in the LETGS setting	44
5.6	Discussion of Condition 62 in the Euler scheme case	47
5.7	Cross-entropy and its estimator in the LETGS setting	49
5.8	Background theory for mean square, inefficiency constant, and their estimators	50
5.9	Mean square and its estimators in the ECM setting	53
5.10	Strongly convex functions and ϵ -minimizers	56
5.11	Mean square and its first estimator in the LETGS setting	57
5.12	Smoothness of functions	58

6	Minimization methods of estimators and their convergence properties	60
6.1	A simple adaptive IS procedure used in our numerical experiments	61
6.2	Helper strong laws of large numbers	62
6.3	Locally uniform convergence of estimators	68
6.4	Exact minimization of estimators	72
6.5	Helper theorems for proving convergence properties of minimization methods with gradient-based stopping criteria	75
6.6	Minimization of estimators with gradient-based stopping criteria	78
6.7	Helper theorems for proving convergence properties of multi-phase minimization methods	81
6.8	Two-phase minimization of estimators with gradient-based stopping criteria and constraints or function modifications	82
6.9	Three-phase minimization of estimators with gradient-based stopping criteria and function modifications	86
6.10	Comparing the first-order asymptotic efficiency of minimization methods	92
6.11	Finding exactly the zero- or optimal-variance IS parameter	94
7	Asymptotic properties of minimization methods	96
7.1	Helper theory for proving asymptotic properties of minimization results	97
7.2	Asymptotic properties of functions of minimization results and comparing the second-order asymptotic efficiency of minimization methods	99
7.3	Some sufficient conditions for asymptotic properties in our methods	101
7.4	Asymptotic properties of minimization results of the new estimators when the zero- or optimal-variance IS parameter exists	102
7.5	Asymptotic properties of SSM results	103
7.6	A helper CLT	106
7.7	Asymptotic properties of MSM results	109
7.8	Properties of matrices characterizing the asymptotic distributions when the optimal- or zero-variance IS parameter exists	111
7.9	Analytical example of a symmetric three-point distribution	115
8	Two-stage estimation of the quantity of interest	117
9	Numerical experiments	119
9.1	Testing statistical hypotheses	120
9.2	Estimation experiments	121
9.3	Experiments comparing the spread of IS drifts.	126
9.4	Intuitions behind some results of our numerical experiments	130
10	Conclusions and further ideas	131

1 Introduction

1.1 Problem statement and literature overview

Consider a probability \mathbb{Q}_1 on a measurable space $\mathcal{S}_1 = (\Omega_1, \mathcal{F}_1)$ and let Z be an \mathbb{R} -valued random variable on \mathcal{S}_1 such that $\mathbb{E}_{\mathbb{Q}_1}(|Z|) < \infty$. We are interested in estimation of $\alpha := \mathbb{E}_{\mathbb{Q}_1}(Z)$. Monte Carlo (MC) method relies on approximating α using a MC average $\hat{\alpha}_n := \frac{1}{n} \sum_{i=1}^n Z_i$ of independent random variables Z_i , $i = 1, \dots, n$, each having the same distribution as Z under \mathbb{Q}_1 (shortly called n independent replicates of Z under \mathbb{Q}_1). Variance of $\hat{\alpha}_n$ measures its mean square error of approximation of α , and for $\text{var} := \text{Var}_{\mathbb{Q}_1}(Z)$ we have $\text{Var}(\hat{\alpha}_n) = \frac{\text{var}}{n}$. When performing a MC procedure on a computer it is often the case that there exists a nonnegative random variable \hat{C} on \mathcal{S}_1 such that for generated independent replicates (Z_i, \hat{C}_i) , $i = 1, \dots, n$, of (Z, \hat{C}) under \mathbb{Q}_1 , \hat{C}_i are typically approximately equal to some practical simulation costs, like computation times, needed to generate Z_i . We call such \hat{C} an approximate practical cost. In a number of situations $\hat{C} = p_C C$ for some $p_C \in (0, \infty)$, which may be different for different computers and implementations (shortly, for different practical realizations) considered and a random variable C on \mathcal{S}_1 , called theoretical cost, which is common for these practical realizations. In case when practical costs of generating Z_i are approximately constant, one can take $C = 1$. A random C can be e.g. an internal duration time of a stochastic process from which Z is computed. For instance, such C can be the number of steps of an acceptance-rejection algorithm. In molecular dynamics applications, when computing the probability that a discretization of a diffusion exits some set before a given time T (see e.g. [53, 15] and our numerical examples), as such C one can typically take the minimum of T and the exit time of the set. In computational finance, when pricing a knock-out barrier option, as such C one can take the minimum of the hitting time of the asset of the barrier and the expiry date of the option (see [18, 31, 29]). Assuming that such \hat{C} and C are given, let us define mean costs $\hat{c} = \mathbb{E}_{\mathbb{Q}_1}(\hat{C})$ and $c = \mathbb{E}_{\mathbb{Q}_1}(C)$, practical inefficiency constant $\hat{\text{ic}} = \hat{c} \text{var}$, and theoretical inefficiency constant $\text{ic} = c \text{var}$. For \hat{c} and var finite, values of practical inefficiency constants are reasonable measures of inefficiency of MC procedures as above, i.e. higher such values imply lower efficiency. The name inefficiency constant was coined and some its interpretations were given in [5, 6], but the idea of using a reciprocal of such constant to quantify the efficiency of MC methods was conceived much earlier, see [22] for a historical review. Glynn and Whitt [22] propose to quantify the asymptotic efficiency of simulation estimators using asymptotic efficiency rates and values and the above inefficiency constant is equal to the reciprocal of their efficiency value in the special case of a MC method, in which the efficiency rate equals $\frac{1}{2}$. See [19], Section 10 of Chapter 3 in [4], and Section 1.1.3 in [18] for accessible descriptions of their approach in the special case of MC method, and Section 2 of this work for a review of interpretations of inefficiency constants in MC methods as well as for some new interpretations. If for different Z , \mathbb{Q}_1 , and C as above (but in each case $\mathbb{E}_{\mathbb{Q}_1}(Z) = \alpha$ for the same α), for some practical realizations of their corresponding MC procedures the proportionality constants p_C as above are the same, then comparing practical inefficiency constants in these realizations is equivalent to comparing the theoretical inefficiency constants. For instance in [5, 6] this idea was used for theoretical costs C being the numbers of the most

costly function evaluations involved in computation of estimators of variance-based sensitivity indices and orthogonal projection coefficients, and inequalities between the resulting theoretical inefficiency constants (called there inefficiency constants of schemes) were derived. If the mean practical costs of two MC methods are the same, then comparing their practical inefficiency constants reduces to comparing their variances.

Efficiency improvement techniques (EITs), the term having been proposed in [22], try to improve the efficiency of estimation of α , e.g. when using MC methods by reducing the values of their practical inefficiency constants. One way to improve the efficiency is to enhance the practical realization of a method, e.g. by increasing the computational power available or improving the implementation details. In case of MC method this can reduce the practical inefficiency constant by reducing the mean practical cost but without changing the variance. An alternative is to use statistical EITs, like control variates, importance sampling (IS), antithetic variables, or stratified sampling (see e.g. [4, 19]). Control variates method relies on generating in a MC procedure i.i.d. replicates under \mathbb{Q}_1 of a sum of Z and a \mathbb{Q}_1 -zero-mean random variable, called control variate [4, 21]. In IS, for some probability \mathbb{Q}_2 on \mathcal{S}_1 , called IS distribution, and a random variable L , called IS density, such that $\mathbb{E}_{\mathbb{Q}_2}(ZL) = \alpha$, one performs a MC procedure generating i.i.d. replicates of the IS estimator ZL under \mathbb{Q}_2 . IS has found numerous applications among others in queueing theory, computational finance, and molecular modelling, and can be a useful tool for rare-event simulation [4, 20, 53, 15, 9]. Using IS and control variates typically changes both mean practical cost and variance as compared to crude MC (CMC). Implementing these methods often introduces additional computational overhead over crude MC (CMC). In particular, if one can use the same theoretical cost variable C in definitions of theoretical inefficiency constants in these methods as for CMC, then their corresponding proportionality constants p_C as above are often higher. In such case one receives higher mean practical cost for control variates than for CMC, while the mean practical cost for IS, equal to $p_C \mathbb{E}_{\mathbb{Q}_2}(C)$, can change in various ways from that in CMC.

Adaptive EITs use information available from the random drawings to make the estimation method more efficient, e.g. by tuning some parameter of the method from some set $A \in \mathcal{B}(\mathbb{R}^l)$. For instance, in adaptive control variates one typically tunes the parameter in some parametrization of control variates, while in IS — in parametrizations $b \in A \rightarrow \mathbb{Q}(b)$ of IS distributions and $b \in A \rightarrow L(b)$ of IS densities. We call $b \in A$ for these cases the control variates and IS parameters respectively. Typically in the literature the adaptive EITs for control variates and IS have attempted to find a parameter minimizing or maximizing some function $b \in A \rightarrow v(b) \in \mathbb{R}$. Frequently, such v was variance or equivalently mean square of the estimator and it was minimized. We say that two functions $f_i, i = 1, 2$ are positively (negatively) linearly equivalent, if $f_1 = af_2 + b$ for some $a \in (0, \infty)$ ($a \in (-\infty, 0)$) and $b \in \mathbb{R}$. As discussed below, in a number of adaptive IS approaches it has been proposed to maximize certain functions v negatively linearly equivalent to some f -divergences of an IS distribution from the zero variance one (if it exists), like cross-entropy distance or Hellinger divergence [42, 2, 38].

Two classes of stochastic optimization methods have been typically used for optimizing (i.e. minimizing or maximizing) some v as above. One of them are stochastic approximation algorithms. These are multi-stage stochastic opti-

mization methods using stochastic gradient descent, in which estimates of values of gradients of v are computed in each stage. When applying such methods in EITs one typically estimates the quantity α using an average of its estimates from different stages. See e.g. [31] for an application of such methods to variance minimization in adaptive control variates and [3, 37, 35] in IS. A problem with such methods is that their practical performance heavily depends on the choice of step sizes, and some heuristic tuning of them may be needed to achieve a reasonable performance [31].

Consider a probability \mathbb{V} on a measurable space \mathcal{D} , and let $X \sim \mathbb{V}$. Let $\hat{v} : (A, \mathcal{B}(A)) \otimes \mathcal{D} \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ be an estimator of v , i.e. the values of $\hat{v}(b, X)$ are thought of as approximations of $v(b)$, $b \in A$. Another popular class of stochastic optimization methods of v is based on optimization of the function $b \rightarrow \hat{v}(b, X)$. Methods from this class have been called stochastic counterpart methods, alternative names including sample path and sample average approximation methods [27, 31, 34, 51]. See Chapter 6 of Section 9 in [51] for a historical review of such methods, related to M-estimation and in particular maximum likelihood estimation in statistics [52]. The most well-known example of application of the stochastic counterpart method to efficiency improvement are linearly parametrized control variates [4, 21, 40], in which to obtain the control variates parameter one minimizes sample variance of the control variates estimator by solving a certain system of linear equations. In [31] the authors considered minimization of sample variance estimator of a nonlinearly parametrized control variate estimator.

Consider some functions $\hat{w}_k : (A, \mathcal{B}(A))^2 \otimes \mathcal{S}_1^k \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$, $k \in \mathbb{N}_+$. In some works using stochastic counterpart method for adaptive IS, for some $n \in \mathbb{N}_+ := \{1, 2, \dots\}$ and $b' \in A$, one uses $\mathbb{V} = \mathbb{Q}(b')^n$ and $\hat{v}(b, X) = \hat{w}_n(b', b, X)$. We call such case single-stage optimization of estimators. In a number of works, multi-stage optimization of estimators was used in which for some $b_0 \in A$ and $m \in \mathbb{N}_+$, for each $k \in \{1, \dots, m\}$, for some $n_k \in \mathbb{N}_+$ and $Y_k \sim \mathbb{Q}(b_{k-1})^{n_k}$, the k -th stage result b_k is an optimization result of $b \rightarrow \hat{w}_{n_k}(b_{k-1}, b, Y_k)$, and b_m is used as the final estimate of an optimum point of v . To our knowledge, in the literature such n_k were always taken equal to the same $N \in \mathbb{N}_+$ for different k , but we consider above more general n_k with a view to discussing our new methods in the next section.

In [45] (see Algorithm 2.1 in that work), multi-stage minimization of estimators of mean square (of the IS estimator) was proposed, see also Algorithm 3.2.1 in [46]. In [29], single-stage minimization of mean square estimator was considered for computing means of functions of a fixed number of i.i.d. normal random variables using IS via exponential change of measure. Under an idealistic assumption that one can find the minimum points of mean square estimators exactly, the authors proved convergence and asymptotic properties of minimization results in their method as the number n of independent normal random vectors used goes to infinity. In [44] it was proposed to maximize an estimator of a certain function v negatively linearly equivalent to cross-entropy distance of an IS distribution from the zero-variance one (if it exists). A multi-stage maximization algorithm of such estimator is presented as Algorithm 3.3.1 in [46]. The advantage of maximizing such estimator over minimizing mean square estimator is that for a number of important parametric families of IS distributions its maximum point can be found explicitly and one does not need to use iterative optimization algorithms. Explicit formulas for such maximum

points often exist when using for IS products of distributions from natural exponential families (see Example 3.5 in [46]) or when using Girsanov transformation with linear parametrization of drifts for diffusions [53]. If the zero-variance IS distribution does not belong to the parametric family of IS distributions used, then the minimum point of cross-entropy distance can be different from such point of variance (or equivalently mean square). However, in a number of rare-event problems distance of these points can be proved to converge to zero in the limit in some limiting rare-event regime [10]. In [42] the author considered maximization of an estimator of a function negatively linearly equivalent to Hellinger divergence of an IS distribution from the zero-variance one, whose maximizers, unfortunately, typically cannot be found analytically.

For a real-valued random variable X on \mathcal{S}_1 and some level $\lambda \in (0, \infty)$, a so-called multi-level approach was proposed for adaptive IS for computing expectation of a rare event indicator variable $Z = 1(X > \lambda)$, in which λ is increased in the consecutive stages of multi-stage optimization of estimators to make the event less rare in the initial and more rare in the later stages [44, 46]. In [53], when computing expectations of some functionals of certain diffusions, it was proposed to increase the temperature in the initial stages to speed up the sampling and make certain rare events less rare in these stages.

A popular approach to estimation of α using stochastic counterpart method is a two-stage approach, in the first stage of which one obtains a value b of the adaptive parameter via optimization of some estimators as above, and in the second stage one performs a separate MC procedure using such b and generating new independent random variables [44, 46, 31]. An alternative approach is to reuse in the second stage random variables generated in the first stage [29, 4].

1.2 Overview of own work

In this work we investigate various methods for obtaining IS parameters via single- or multi-stage minimization of estimators of different functions, abbreviated further as SSM and MSM respectively. Such received IS parameter can be later used in a separate IS MC procedure to estimate the quantity of interest, as in a two-stage estimation approach in the previous section.

We define cross-entropy to be a certain function of the IS parameter, positively linearly equivalent to cross-entropy distance of the corresponding IS distribution from the zero-variance IS distribution (if the latter exists), even though this name is typically used in the literature as a synonym of the cross-entropy distance. We consider minimization of an estimator of such cross-entropy, which is equivalent to maximization of a linearly negatively equivalent to it estimator typically considered in the literature, as discussed in the previous section. Therefore, we call such estimator of cross-entropy well-known. We also investigate minimization of an estimator of mean square (of the IS estimator), already minimized in some works mentioned in the previous section in various special cases of the IS settings considered in this work, and thus also called well-known. Furthermore, we consider minimization of new estimators of mean square and theoretical inefficiency constant (in which we further often drop the word theoretical). As discussed in the previous section, our methods can be viewed as stochastic counterpart approaches to minimization of the corresponding functions. To our knowledge it is the first time when inefficiency constant is being adaptively minimized. In numerous current and potential applications

of IS, mean approximate practical simulation cost \hat{c} significantly depends on the IS parameter used, so that practical inefficiency constant is not proportional to variance and their minimization is not equivalent. This often happens when computation of replicates of Z involves simulating a stochastic process until some random time, as discussed in the previous section. For a number of parametrizations of IS distributions and densities, for some theoretical cost variable C , and for typical joint practical realizations of IS MC methods corresponding to different IS parameters, the approximate practical costs \hat{C} are equal to $p_C C$ for the same $p_C \in (0, \infty)$ for different IS parameters. This is for instance typically the case when applying IS using discrete Girsanov transformation to certain molecular dynamics problems as in our numerical experiments or to the problem of pricing knock-out barrier options e.g. using discretizations of Black-Scholes model, as discussed in the previous section. In such cases practical inefficiency constants are equal to the theoretical ones multiplied by p_C and their minimization is equivalent.

We pay special attention to two classes of families of IS distributions. One of them arises from exponential tilting with fixed tilting parameter, also called exponential change of measure (ECM). The second class is obtained through exponential tilting for Gaussian stopped sequences with a linear parametrization of tilting processes (LETGS) and it can be also viewed as a result of applying discrete version of Girsanov transformation up to a stopping time and using such tilting processes. IS in the LETGS setting can be used for computing expectations of functionals of sequences of i.i.d. Gaussian random variables up to a stopping time, which can be functionals of various discretizations of SDEs [32], like Euler scheme. Using IS in some special cases of the ECM and LETGS settings was demonstrated to lead to significant variance reductions e.g. in rare event simulation [9, 4] and when pricing options in computational finance [29, 37].

We provide necessary and sufficient assumptions under which there exist unique minimum points of cross-entropy and mean square (and thus also of variance) of the IS estimator in the ECM and LETGS settings, and we give some sufficient conditions for these assumptions to hold in the Euler scheme case. In the LETGS setting and in some important special cases of the ECM setting we have analytical formulas for the unique minimum points of the cross-entropy estimators. In both ECM and LETGS settings, the well-known mean square estimator is convex. We prove that in the ECM setting the new mean square estimator is also convex and a necessary and sufficient condition for it to have a unique minimum point is for a certain system of linear inequalities to have only a trivial solution.

We define various single- and multi-stage minimization methods of different estimators for which we prove appropriate convergence properties of minimization results in the ECM and LETGS settings as the budget of SSM or the number of stages of MSM (jointly, the budget) goes to infinity. We define various single- and multi-stage minimization methods of different estimators for which we prove appropriate convergence properties of minimization results in the ECM and LETGS settings as the budget of SSM or the number of stages of MSM (jointly, the budget) goes to infinity. To ensure such convergence properties in the MSM methods we use the numbers n_k of samples in the subsequent stages going to infinity and as IS parameters b_k needed for constructing estimators $b \rightarrow \hat{w}_{n_k}(b_{k-1}, b, Y_k)$ minimized in the subsequent stages as discussed in the

previous section, we use e.g. minimization results of these estimators projected onto some bounded sets. Furthermore, in the proofs we apply a newly developed uniform multi-stage strong law of large numbers. For the cross-entropy estimators in both settings we consider their exact minimization utilising expressions for their minimum points, and we prove convergence of their minimization results to the unique minimum point of cross-entropy. For the well-known mean square estimators in both settings and the new mean square estimators in the ECM setting we consider their minimization with gradient-based stopping criteria and prove a.s. convergence of their minimization results to the unique minimum point of mean square for gradients in these results decreasing to zero (as the budget goes to infinity). Under appropriate assumptions, the new mean square estimators in the LETGS setting and the inefficiency constant ones in the ECM setting for the theoretical cost variable $C = 1$ can be shown to a.s. become strongly convex on each bounded convex set for sufficiently large budget of SSM or MSM. We prove a.s. convergence of their minimization results to the unique minimum point of mean square when using a following two-phase minimization procedure. In its first phase some convex estimator of mean square as above is minimized, and then, using its minimization result as a starting point, a constrained minimization of the considered estimator or an unconstrained minimization but of an appropriately modified such estimator is carried out. For inefficiency constant estimators in the LETGS setting with cost variable equal to the stopping time, we propose three-phase minimization. Its first phase is the same as above and in the second phase, starting from the first-phase minimization result, a single gradient descent is performed on a modified inefficiency constant estimator with estimated modifications until Wolfe condition is satisfied. In the third phase one performs further unconstrained minimization of such modified estimator with a gradient stopping criterion with gradient decreasing to zero. This ensures convergence of distance of the minimization results of the inefficiency constant estimators to the set of first-order critical points of inefficiency constant which have not higher values of inefficiency constant than in the minimum point of variance, or even by at least some positive constant lower such values in case when gradient of inefficiency constant in the minimum point of variance does not vanish.

If $\mathbb{Q}_1(Z \neq 0) > 0$, then there exists a unique IS distribution leading to the lowest variance (of the IS estimator) and called the optimal-variance one. If additionally $Z \geq 0$, \mathbb{Q}_1 a.s., then the optimal-variance IS distribution leads to a zero-variance IS estimator. The IS parameters leading to such distributions are called the optimal-variance or zero-variance ones respectively. If b^* is the optimal-variance IS parameter for the new mean square estimator or the zero-variance IS parameter for the inefficiency constant estimator, then under appropriate assumptions a.s. the minimization results d_t of exact SSM and MSM of these estimators are equal to b^* for sufficiently large budget t used. Furthermore, for such results of SSM and MSM of these estimators with gradient-based stopping criteria, for arbitrary $\lim_{t \rightarrow \infty} r_t = \infty$ and for gradients of estimators in d_t going to zero sufficiently quickly, $\sqrt{r_t}(d_t - b^*)$ converges to zero in probability.

Let in SSM, for $b' \in \mathbb{R}^l$ denoting an IS parameter used to construct estimators, $u(b')$ denote the mean cost of generating replicates under $\mathbb{Q}(b')$ and doing some additional computations per replicate needed for later estimator minimization. We prove that under appropriate assumptions, including existence of a local minimum point b^* of the minimized function, for all estimators

considered, for some corresponding covariance matrices $V(b) \in \mathbb{R}^{l \times l}$, we have following asymptotic results. The SSM results d_t corresponding to budget t fulfill $\sqrt{t}(d_t - b^*) \Rightarrow \mathcal{N}(0, u(b')V(b'))$. Let $n_k \in \mathbb{N}_+$, $k \in \mathbb{N}_+$, be such that $\lim_{k \rightarrow \infty} n_k = \infty$. In MSM methods in which n_k samples are generated in the k th stage, and in which the IS parameters b_k from subsequent stages converge in probability to some $d^* \in A$ (like e.g. $d^* = b^*$), their MSM results d_k from the k th stage fulfill $\sqrt{n_k}(d_k - b^*) \Rightarrow \mathcal{N}(0, V(d^*))$. If b^* is the optimal-variance IS parameter for the new mean square estimator or the zero-variance one for the inefficiency constant estimator, then from the above discussion, for r_t equal to t or n_t respectively we have in such case $V(b) = 0$ in the above formulas for $b = b'$ or $b = d^*$ respectively, and for r_t growing faster than t or n_t we can have faster convergence rate of d_t to b^* than in the above formulas. We also show that if b^* is the zero-variance IS parameter, then, under appropriate assumptions, for the cross-entropy estimator the covariance matrix $V(b)$ as above is positive definite and is four times such matrix for the well-known mean square estimator. We provide an analytical example of ECM for a symmetric three-point distribution with cost variable $C = 1$, in which $b^* = 0$ is the unique minimum point of both cross-entropy and mean square and in which all possible relations between the values $V(b) \in \mathbb{R}$ for each b (i.e. equalities and both strict inequalities) for different pairs of estimators are achieved for different parameters of the example, except that using cross-entropy estimator always leads to not lower $V(b)$ than using the well-known mean square estimator.

For an unbounded set $T \subset \mathbb{R}_+$, we propose first- and second-order methods for comparing the asymptotic efficiency of processes $d_t \in A$, $t \in T$, for minimization of some functions $f : A \rightarrow \mathbb{R}$, like inefficiency constant or variance. For each $t \in T$, such d_t can be e.g. a result of some stochastic optimization algorithm minimizing f and using simulation budget t . A process more efficient in the first-order sense leads to lower values of the minimized function by at least a fixed positive constant with probability going to one as $t \rightarrow \infty$. For instance, the above discussed process of results of three-phase minimization of inefficiency constant estimators in the LETGS setting can be more efficient in this sense than a one from minimization of some mean square estimators if gradient of the inefficiency constant in the minimum point of mean square does not vanish. For processes d_t fulfilling $r_t(f(d_t) - f(b^*)) \Rightarrow \mu$ for the same f , $b^* \in A$, and $r_t \in \mathbb{R}_+$, $t \in T$, such that $r_t \rightarrow \infty$, as well as for possibly different probability distributions μ , these processes are equally efficient in the first-order sense. Their second-order asymptotic efficiency for minimization of f can be compared using different properties of probabilities μ , like their means. In case of mean, the process with its lower value can be considered more efficient. For instance, if for some covariance matrix M , $\sqrt{r_t}(d_t - b^*) \Rightarrow \mathcal{N}(0, M)$, then for appropriate f , $r_t(f(d_t) - f(b^*))$ converges in distribution to a generalized χ^2 distribution with mean $\frac{1}{2} \text{Tr}(M \nabla^2 f(b^*))$. In particular, for processes from SSM for $r_t = t$ and MSM for $r_k = n_k$, enjoying appropriate asymptotic properties as above, such mean is $\frac{u(b')}{2} \text{Tr}(V(b') \nabla^2 f(b^*))$ and $\frac{1}{2} \text{Tr}(V(d^*) \nabla^2 f(b^*))$ respectively. As we discuss, such means can be potentially estimated and adaptively minimized. We show that for b^* minimizing f , $b \rightarrow \frac{1}{2} \text{Tr}(V(b) \nabla^2 f(b^*))$ may in general assume a minimum in a different point than b^* , so that the usual choice of parameters b_k in MSM converging to $d^* = b^*$ does not need to lead to the lowest mean as above.

In our numerical experiments we consider Euler scheme discretization of a diffusion in a one-dimensional three-well potential. We address the problem of estimating moment-generating function (MGF) of the exit time of such Euler scheme of a domain, probability to exit it before a fixed time, and probabilities to leave it through given parts of the boundary, called committors. Such quantities for some diffusions in potentials are of interest among others in molecular dynamics and have been estimated in a number of works [25, 53, 16, 24]. We consider IS in the LETGS setting, for which under the IS distribution we receive again an Euler scheme but this time with an additional drift depending on the IS parameter, called IS drift. For estimation of the above quantities we use a two-stage method as discussed in the previous section, in the first stage of which to obtain the IS parameter we use simple MSM of estimators without various enhancements like projections or function modifications needed to ensure appropriate convergence properties as discussed above. In our numerical experiments, MSM of the new estimators of inefficiency constant and mean square led to the lowest variances and inefficiency constants as well as to visually fastest convergence of IS drifts, followed by the well-known mean square and cross-entropy estimators. In one case, minimization of the inefficiency constant estimator outperformed minimization of the new mean square estimator by arriving at a lower mean stopping time and higher variance but so that their product, equal to inefficiency constant, was lower. When computing expectations of nonincreasing functions of the exit time, like MGF and probability to leave the domain before a fixed time, the mean exit times under the IS parameters obtained from MSM were significantly smaller than for CMC. An intuition behind this result is provided by a theorem, which we prove, that for Z being nonincreasing functions of the (theoretical) cost variable C , under the zero-variance IS distribution we should have not higher mean cost than for CMC, and under some additional assumptions this inequality is sharp. For a committor we also performed experiments comparing the spread of IS drifts obtained from SSM, both visually and in terms of histograms and interquartile ranges (IQRs) of IS drifts at zero. Minimization results of the new estimators led to the lowest spreads, followed by such results of mean square and cross-entropy estimators, and the received IQRs are lower when using in SSM an IS parameter b' from some preliminary MSM than $b' = 0$. Furthermore, the received ratios of IQRs from minimization of the cross-entropy estimator to the ones from minimization of the well-known mean square estimator are close to 2. We show that if the zero-variance IS parameter existed, then from the above discussed properties of asymptotic distributions of minimization results in such case, we would have zero asymptotic variance and thus also zero IQR of the asymptotic distribution of IS drifts from SSM and MSM in all points for the new estimators, and ratio of such IQRs of IS drifts for the cross-entropy estimator to the well-known mean square estimator would be exactly equal 2. Thus, a possible intuition behind the results of our numerical experiments concerning the relations of spreads and convergence speeds of IS drifts from SSM and MSM of different estimators is that the IS distributions and densities corresponding to minimum points of the minimized functions can be close to the zero-variance ones and we have approximately the same asymptotic distributions of IS drifts as if the zero-variance IS parameters existed. We also found out that using a two-phase approach in which minimization of inefficiency constant estimator is preceded by minimization of mean square estimator can help avoid receiving outlying IS drifts, which sometimes occur when only the

inefficiency constant estimator is minimized. Furthermore, the variances and inefficiency constants of adaptive IS estimators in our experiments strongly depended on the parametrization of the IS drifts used and could be reduced by adding appropriate positive constants to the variables Z .

1.3 Outline of the paper

The rest of this work is organized as follows. After introducing basic notations in Section 1.4 and describing different interpretations of inefficiency constants in Section 2, we discuss IS and related concepts in Section 3. Then, in Section 4, we present various examples of parametric families of IS distributions, including the ones later used in our numerical experiments. In Section 5 we provide definitions of the minimized functions and their estimators, and we discuss their important properties. In Section 6 we provide definitions of single- and multi-stage minimization methods of estimators of different functions and prove strong convergence properties of their results. In its Subsection 6.1 we describe the two-stage estimation methods used in our numerical experiments. In Section 7 we investigate asymptotic properties of such minimization results and their functions. In Section 8 we discuss a two-stage method for estimation of α . In Section 9 we present results of our numerical experiments. Finally, in Section 10 we make some additional comments on our results and discuss some further ideas, including possible alternative approaches.

1.4 Basic notations

In the further sections we frequently use objects defined in Section 1.1, like Z , \mathbb{Q}_1 , and α . We assume that $x^0 = 1$, $x \in \mathbb{R}$. We denote $\mathbb{N}_+ = \{1, 2, \dots\}$, $\mathbb{N} = \mathbb{N}_+ \cup \{0\}$, for $p \in \mathbb{N}_+$, $\mathbb{N}_p = \{p, p+1, \dots\}$, $\mathbb{R}_+ = (0, \infty)$, and $\overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, \infty\}$. For $a, b \in \overline{\mathbb{R}}$, $a \wedge b$ denotes their minimum, and $a \vee b$ their maximum, but these symbols are also used to denote logical conjunction and disjunction. By default, all vectors are considered to be column vectors. In particular, gradient of a function is considered to be a column vector. Expectation of a random variable Z with respect to a probability \mathbb{Q} is denoted as $\mathbb{E}_{\mathbb{Q}}(Z)$, and variance as $\text{Var}_{\mathbb{Q}}(Z)$. Subscript \mathbb{Q} in the above notations is omitted when \mathbb{Q} is the default probability \mathbb{P} on the default measurable space (Ω, \mathcal{F}) . For a real-valued random variable Y we denote $Y_+ = Y\mathbb{1}(Y > 0)$ and $Y_- = -Y\mathbb{1}(Y < 0)$. We say that q is a random condition on some measurable space $\mathcal{S} = (B, \mathcal{B})$ if $\{x \in B : q(x)\} \in \mathcal{B}$. Often the event $\{x \in B : q(x)\}$ will be denoted simply as $\{q\}$ and we frequently write q in place of $\{q\}$ in some notations or statements. For some probability \mathbb{Q} on \mathcal{S} we say that an event $A \in \mathcal{B}$ holds \mathbb{Q} a.s. on some event $B \in \mathcal{B}$, if $\mathbb{Q}(A \cap B) = \mathbb{Q}(B)$. Unless stated otherwise by minimum (point) of a function we mean its global minimum (point). By $|\cdot|$ we denote the standard Euclidean norm and by $\|\cdot\|_{\infty}$ the supremum norm induced by $|\cdot|$. For a set $A \in \mathcal{B}(\mathbb{R}^l)$ or $A \in \mathcal{B}(\overline{\mathbb{R}})$ (where $\mathcal{B}(B)$ is the borel σ -field on B), the default measurable space considered on it is $(A, \mathcal{B}(A))$, which shall be denoted simply as $\mathcal{S}(A)$. For $a \in \mathbb{R}^l$ and $\epsilon \in \mathbb{R}_+$, we define a sphere $S_l(a, \epsilon) = \{x \in \mathbb{R}^l : |x - a| = \epsilon\}$, a ball $B_l(a, \epsilon) = \{x \in \mathbb{R}^l : |x - a| < \epsilon\}$, and a closed ball $\overline{B}_l(a, \epsilon) = \overline{B_l(a, \epsilon)} = \{x \in \mathbb{R}^l : |x - a| \leq \epsilon\}$. By $\xrightarrow{\mathbb{P}}$ we denote convergence in probability, by \Rightarrow convergence in distribution or right implication depending on the context, and by \rightrightarrows uniform convergence. We denote $\mathbb{1}(B)$ or $\mathbb{1}_B$ the indicator function of a subset B of some set X (which

can be inferred from the context). If \mathbb{R} -valued random variables Y_n , $n \in \mathbb{N}_+$, are well-defined only on some subsets $A_n \in \mathcal{F}$ (e.g. for \mathbb{R} -valued random variables V_n and W_n we may have $Y_n = \frac{V_n}{W_n}$ and $A_n = \{W_n \neq 0\}$), then by $Y_n \xrightarrow{p} Y$ or $Y_n \Rightarrow Y$ we mean that $\lim_{n \rightarrow \infty} \mathbb{P}(A_n) = 1$ and these statements hold for Y_n replaced by $Y_n \mathbb{1}(A_n)$ which is defined to be equal to zero on A_n and to Y_n otherwise.

2 Interpretations of inefficiency constants

In this section we provide some interpretations of inefficiency constants, both from the literature and new ones, justifying their utility for quantifying the inefficiency of MC procedures. The theorems introduced in the process will be frequently used further on. We focus on theoretical inefficiency constants and costs (dropping further the word theoretical), but analogous interpretations hold for the practical ones. We use notations as in Section 1.1, including var , c , and $\text{ic} = c \text{var}$, characterizing some sequence of MC procedures for estimating α indexed by different numbers of replicates $n \in \mathbb{N}_+$ used in these procedures. The following interpretation of inefficiency constants was given in Section 2.6 of [6]. The ratio of positive finite inefficiency constants ic of different sequences of MC procedures as above for estimating α is equal to the limit of ratios of their mean costs nc corresponding to the minimum numbers of replicates $n = \lceil \frac{\text{var}}{\epsilon} \rceil$ needed to reduce the variances $\frac{\text{var}}{n}$ of the MC averages $\hat{\alpha}_n$ below a given threshold $\epsilon > 0$ for $\epsilon \rightarrow 0$. Consider a function $f : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ such that for $x, y \in \mathbb{R}_+$, $f(x, y) = f(y, x)$ and for $a \in \mathbb{R}_+$, $af(x, y) = f(ax, ay)$. Let $g : \mathbb{R}_+^2 \rightarrow [0, \infty)$ be such that $g(x, y) = \frac{|x-y|}{f(x, y)}$, so that $g(x, y) = g(y, x)$ and $g(ax, ay) = g(x, y)$, $a \in \mathbb{R}_+$. For instance, $f(x, y)$ can be equal to $\max(x, y)$, $\min(x, y)$, or $\frac{x+y}{2}$, in which case $g(x, y)$ can be interpreted as the relative difference of x and y . For some $\delta > 0$, we say that $x, y \in \mathbb{R}_+$ are δ -approximately equal, which we denote $x \approx_\delta y$, if $g(x, y) \leq \delta$. Note that $x \approx_0 y$ implies that $x = y$. The below simple interpretations of inefficiency constants were given by Badowski in sections 1.9 and 2.6 of [6] in the special case of $f = \min$ as above. For two MC procedures for estimating α , one like above using n replicates and an analogous primed one, assuming that $\text{ic}, \text{ic}' \in \mathbb{R}_+$, from an easy calculation we have

$$g\left(\frac{\frac{\text{var}}{n}}{\frac{\text{var}'}{n'}}, \frac{\text{ic}}{\text{ic}'}\right) = g(nc, n'c') \quad (1)$$

and

$$g\left(\frac{nc}{n'c'}, \frac{\text{ic}}{\text{ic}'}\right) = g\left(\frac{\text{var}}{n}, \frac{\text{var}'}{n'}\right). \quad (2)$$

In particular, the ratio of positive finite inefficiency constants of these procedures is δ -approximately equal to the ratio of variances of their respective MC averages for δ -approximately equal respective mean total costs and it is also δ -approximately equal to the ratio of their average costs for δ -approximately equal variances of their MC averages.

Let (Z_i, C_i) , $i \in \mathbb{N}_+$, be independent replicates of (Z, C) under \mathbb{Q}_1 . Before providing further interpretations of inefficiency constants, let us recall some basic facts about MC procedures as above. From strong law of large numbers (SLLN), for $\hat{\alpha}_n = \frac{1}{n} \sum_{i=1}^n Z_i$ it holds a.s. $\lim_{n \rightarrow \infty} \hat{\alpha}_n = \alpha$, and if $\text{var} < \infty$, then

from central limit theorem (CLT), $\sqrt{n}(\hat{\alpha}_n - \alpha) \Rightarrow \mathcal{N}(0, \text{var})$. Consider following sample variance estimators

$$\widehat{\text{var}}_n = \frac{n}{n-1} \left(\frac{1}{n} \sum_{i=1}^k Z_i^2 - \hat{\alpha}_n^2 \right), \quad n \in \mathbb{N}_2. \quad (3)$$

If $\text{var} < \infty$, then from SLLN a.s. $\lim_{n \rightarrow \infty} \widehat{\text{var}}_n = \text{var}$ and if further $\text{var} > 0$, then from Slutsky's lemma (see e.g. 2.14 in [52])

$$\sqrt{\frac{n}{\widehat{\text{var}}_n}} (\hat{\alpha}_n - \alpha) \Rightarrow \mathcal{N}(0, 1), \quad (4)$$

which can be used to construct confidence intervals for $\hat{\alpha}_n$, as discussed e.g. in Chapter 3, Section 1 in [4]).

For $n \in \mathbb{N}_+$, let $\hat{c}_n = \frac{1}{n} \sum_{i=1}^n C_i$ and for $n \in \mathbb{N}_2$, let $\hat{\text{ic}}_n = \hat{c}_n \widehat{\text{var}}_n$. Assuming that $c, \text{var} < \infty$, from SLLN a.s. $\lim_{n \rightarrow \infty} \hat{c}_n \rightarrow c$ and $\lim_{n \rightarrow \infty} \hat{\text{ic}}_n \rightarrow \text{ic}$. Let $S_n = \sum_{i=1}^n C_i$, $n \in \mathbb{N}$ (in particular $S_0 = 0$), so that S_n is the cost of generating first n replicates of Z . For $t \in \mathbb{R}_+$ consider

$$N_t = \sup\{n \in \mathbb{N} : S_n \leq t\}, \quad (5)$$

or

$$N_t = \inf\{n \in \mathbb{N} : S_n \geq t\}. \quad (6)$$

The above defined N_t are reasonable choices of numbers of simulations to perform if we want to spend approximate total budget t (like e.g. some internal simulation time) on the whole MC procedure. Definition (5) ensures that we do not exceed the budget t . Under definition (6) we let ourselves finish the last computation started before the budget t is exceeded and thus we do not waste the computational effort already invested in it. Note that if $t > 0$, then under (6) we have $N_t > 0$, i.e. at least one simulation is carried out, which does not need to be the case under (5). If $C < \infty$, \mathbb{Q}_1 a.s., then a.s. $C_i < \infty$, $i \in \mathbb{N}_+$, and thus under both definitions a.s.

$$\lim_{t \rightarrow \infty} N_t = \infty. \quad (7)$$

We have a following well-known slight generalization of the ordinary SLLN (see the corollary on page 292 in [7]).

Theorem 1. *If an $\overline{\mathbb{R}}$ -valued random variable Y is such that $\mathbb{E}(Y_-) < \infty$, then for Y_1, Y_2, \dots , i.i.d. $\sim Y$ a.s.*

$$\frac{1}{n} \sum_{i=1}^n Y_i \rightarrow \mathbb{E}(Y) \in \mathbb{R} \cup \{\infty\}. \quad (8)$$

Let $c > 0$ (in particular we can have $c = \infty$). Then, from the above lemma a.s. $\lim_{n \rightarrow \infty} \frac{S_n}{n} \rightarrow c$ and thus

$$\lim_{n \rightarrow \infty} S_n = \infty, \quad (9)$$

so that under both definitions a.s.

$$N_t < \infty, \quad t \geq 0. \quad (10)$$

From renewal theory (see Theorem 5.5.2 in [11]), under definition (5) we have a.s.

$$\lim_{t \rightarrow \infty} \frac{N_t}{t} = \frac{1}{c}. \quad (11)$$

Since, marking N_t given by (6) with a prim, $N_t \leq N'_t \leq N_t + 1$, (11) holds also when using definition (6).

Let us further consider general $\mathbb{N} \cup \{\infty\}$ -valued random variables N_t , $t \in \mathbb{R}_+$. Let Y be an \mathbb{R}^m -valued random vector such that $\mathbb{E}(Y_i^2) < \infty$, $i = 1, \dots, m$, with mean $\mu = \mathbb{E}(Y)$ and covariance matrix $W = \mathbb{E}((Y - \mu)(Y - \mu)^T)$. Let X_1, X_2, \dots be i.i.d. $\sim Y$. Let $\hat{\mu}_n = \frac{1}{n} \sum_{i=1}^n X_i$, $n \in \mathbb{N}_+$. For the string λ in the below expressions substituted by each of strings α , var, ic, c , or μ , consider estimator $\tilde{\lambda}_t$ corresponding to the total budget $t \in \mathbb{R}_+$ and with initial value λ_0 (where $\lambda_0 \in \mathbb{R}^m$ for λ substituted by μ and $\lambda_0 \in \mathbb{R}$ otherwise), defined as follows

$$\tilde{\lambda}_t = \hat{\lambda}_{N_t} \mathbb{1}(N_t \in \mathbb{N}_+) + \lambda_0 \mathbb{1}(N_t = 0). \quad (12)$$

Note that under definition (6) we have $\tilde{\lambda}_t = \hat{\lambda}_{N_t} \mathbb{1}(N_t < \infty)$, $t > 0$. We shall need a following trivial remark.

Remark 2. For $k \in \mathbb{N}_+$, let τ_k be a.s. \mathbb{N} -valued random variables (i.e. $\mathbb{P}(\tau_k \in \mathbb{N}) = 1$) such that a.s. $\lim_{k \rightarrow \infty} \tau_k = \infty$. If for $k \in \mathbb{N}_+$, a_k are random variables such that a.s. $\lim_{k \rightarrow \infty} a_k = a$ then a.s. $\lim_{k \rightarrow \infty} \mathbb{1}(\tau_k \in \mathbb{N}) a_{\tau_k} = a$.

When we have a.s. (7), (10), and $\lim_{n \rightarrow \infty} \hat{\lambda}_n = \lambda$, then from Remark 2, a.s.

$$\lim_{t \rightarrow \infty} \tilde{\lambda}_t = \lambda. \quad (13)$$

Lemma 3. Let a_n , $n \in \mathbb{N}_+$, be \mathbb{N}_+ -valued random variables such that for some $t_n \in \mathbb{R}_+$, $n \in \mathbb{N}_+$, such that $\lim_{n \rightarrow \infty} t_n = \infty$, for some $b \in \mathbb{R}_+$ we have $\frac{a_n}{t_n} \xrightarrow{P} b$. Then,

$$\frac{\sum_{i=1}^{a_n} (X_i - \mu)}{\sqrt{a_n}} \Rightarrow \mathcal{N}(0, W). \quad (14)$$

Proof. Using Cramér-Wold device (see page 16 in [52]) it is sufficient to consider the case of $m = 1$, which let us assume. For $W = 0$ we have a.s. $X_i = \mu$, $i \in \mathbb{N}_+$, so that the thesis is obvious. The general case with $W > 0$ can be easily inferred from the special case in which $\mu = 0$ and $W = 1$, which can be proved analogously as Theorem 7.3.2 in [11]. \square

For general N_t as above, consider a following condition (which for $c \in \mathbb{R}_+$ follows e.g. from (11)).

Condition 4. It holds $c \in \mathbb{R}_+$ and

$$\frac{N_t}{t} \xrightarrow{P} \frac{1}{c} \quad (15)$$

(as $t \rightarrow \infty$).

Theorem 5. Under Condition 4,

$$\sqrt{t}(\tilde{\mu}_t - \mu) \Rightarrow \mathcal{N}(0, cW). \quad (16)$$

Proof. Let $M_t = (\mathbb{1}(N_t \neq \infty)N_t) \vee 1$, which is an \mathbb{N}_+ -valued random variable. From Condition 4,

$$\frac{M_t}{t} \xrightarrow{P} \frac{1}{c} \in \mathbb{R}_+. \quad (17)$$

Thus, from Lemma 3

$$R_t := \sqrt{M_t}(\hat{\mu}_{M_t} - \mu) \Rightarrow \mathcal{N}(0, W), \quad t \rightarrow \infty. \quad (18)$$

Let $\tilde{R}_t = \mathbb{1}(N_t \in \mathbb{N}_+)R_t = \mathbb{1}(N_t \in \mathbb{N}_+)\sqrt{N_t}(\tilde{\mu}_t - \mu)$. From Condition 4, $\lim_{t \rightarrow \infty} \mathbb{P}(N_t \in \mathbb{N}_+) = 1$, and thus $R_t - \tilde{R}_t \xrightarrow{P} 0$. Therefore, from (18) and Slutsky's lemma, $\tilde{R}_t \Rightarrow \mathcal{N}(0, W)$, and thus

$$\sqrt{c}\tilde{R}_t \Rightarrow \mathcal{N}(0, cW). \quad (19)$$

Let $G_t = \sqrt{t}(\tilde{\mu}_t - \mu)$ and $\tilde{G}_t = \mathbb{1}(N_t \in \mathbb{N}_+)G_t$. Then, $G_t - \tilde{G}_t \xrightarrow{P} 0$, so that to prove (16) it is sufficient to prove that

$$\tilde{G}_t \Rightarrow \mathcal{N}(0, cW). \quad (20)$$

From Condition 4, the continuous mapping theorem, and Slutsky's lemma, $S_t := \mathbb{1}(N_t \in \mathbb{N}_+)\sqrt{\frac{t}{cN_t}} \xrightarrow{P} 1$. Thus, (20) follows from (19) and the fact that from Slutsky's lemma

$$\sqrt{c}\tilde{R}_t - \tilde{G}_t = \sqrt{c}\tilde{R}_t(1 - S_t) \xrightarrow{P} 0. \quad (21)$$

□

In the below theorem and remark we extend the interpretations of inefficiency constants provided at the beginning of Section 10, Chapter 3 in [4] (see also [19] and Example 1 in [22]).

Theorem 6. *If $\text{var} < \infty$ and Condition 4 holds, then*

$$\sqrt{t}(\tilde{\alpha}_t - \alpha) \Rightarrow \mathcal{N}(0, \text{ic}), \quad t \rightarrow \infty. \quad (22)$$

If we further have $\text{var} > 0$ and a.s. (7) and (10), then

$$\sqrt{\frac{t}{\tilde{\text{ic}}_t}}(\tilde{\alpha}_t - \alpha) \Rightarrow \mathcal{N}(0, 1), \quad t \rightarrow \infty. \quad (23)$$

Proof. (22) follows immediately from Theorem 4 and (23) follows from (22), (13) for $\lambda = \text{ic}$, and Slutsky's lemma. □

Remark 7. *Let $X \sim \mathcal{N}(0, 1)$ and let for $\beta \in (0, 1)$, z_β be the β -quantile of normal distribution, i.e. $\mathbb{P}(X \leq z_\beta) = \beta$. Then, for $\gamma \in (0, 1)$ and $p_\gamma = z_{1-\frac{\gamma}{2}}$, we have $\mathbb{P}(|X| \leq p_\gamma) = 1 - \gamma$. Assuming (23), for the random interval $I_{\gamma,t} = (\tilde{\alpha}_t - p_\gamma\sqrt{\frac{\tilde{\text{ic}}_t}{t}}, \tilde{\alpha}_t + p_\gamma\sqrt{\frac{\tilde{\text{ic}}_t}{t}})$ we have*

$$\lim_{t \rightarrow \infty} \mathbb{P}(\alpha \in I_{\gamma,t}) = \mathbb{P}(|X| \leq p_\gamma) = 1 - \gamma, \quad (24)$$

i.e. $I_{\gamma,t}$ is an asymptotic $1 - \gamma$ confidence interval for α . It follows that inefficiency constant and its estimator play the same role when constructing asymptotic confidence intervals for α for $t \rightarrow \infty$ using $\tilde{\alpha}_t$, as variance and its estimator do for $n \rightarrow \infty$ using $\hat{\alpha}_n$ as discussed below (4). For $C = 1$, both approaches to constructing asymptotic confidence intervals become equivalent.

3 Importance sampling

In this section we provide a background on IS.

3.1 Background on densities

For probability distributions \mathbb{P}_1 and \mathbb{P}_2 on the same measurable space $\mathcal{S} = (D, \mathcal{D})$, we say that \mathbb{P}_1 has density L (also called Radon-Nikodym derivative) with respect to \mathbb{P}_2 on some $A \in \mathcal{D}$, or simply that $L = (\frac{d\mathbb{P}_1}{d\mathbb{P}_2})_A$, if L is a random variable such that for each $B \in \mathcal{D}$, $\mathbb{P}_1(\mathbb{1}(A \cap B)) = \mathbb{E}_{\mathbb{P}_2}(L\mathbb{1}(A \cap B))$, or equivalently for each random variable Y on \mathcal{D} such that $\mathbb{1}_A Y$ is nonnegative or \mathbb{P}_1 -integrable,

$$\mathbb{E}_{\mathbb{P}_1}(\mathbb{1}(A)Y) = \mathbb{E}_{\mathbb{P}_2}(L\mathbb{1}(A)Y). \quad (25)$$

Such L is uniquely defined only \mathbb{P}_2 a.s. on A , i.e. for some L' we also have $L' = (\frac{d\mathbb{P}_1}{d\mathbb{P}_2})_A$ only if $L' = L$, \mathbb{P}_2 a.s. on A . From Radon-Nikodym theorem, L exists only if \mathbb{P}_1 is absolutely continuous with respect to \mathbb{P}_2 on A , which we denote $\mathbb{P}_1 \ll_A \mathbb{P}_2$, and which means that for each $B \in \mathcal{D}$, from $\mathbb{P}_2(A \cap B) = 0$ it follows that $\mathbb{P}_1(A \cap B) = 0$. We say that \mathbb{P}_1 and \mathbb{P}_2 are mutually absolutely continuous on A if $\mathbb{P}_1 \ll_A \mathbb{P}_2$ and $\mathbb{P}_2 \ll_A \mathbb{P}_1$, which we also denote $\mathbb{P}_1 \sim_A \mathbb{P}_2$.

Lemma 8. *If $L = (\frac{d\mathbb{P}_1}{d\mathbb{P}_2})_A$, then $\mathbb{P}_1 \sim_A \mathbb{P}_2$ only if $\mathbb{P}_2(\{L = 0\} \cap A) = 0$, in which case $\frac{\mathbb{1}(L \neq 0)}{L} = (\frac{d\mathbb{P}_2}{d\mathbb{P}_1})_A$.*

Proof. If $\mathbb{P}_2(\{L = 0\} \cap A) = 0$, then for $B \in \mathcal{D}$, from (25),

$$\mathbb{E}_{\mathbb{P}_2}(\mathbb{1}(A \cap B)) = \mathbb{E}_{\mathbb{P}_2}(L(\frac{\mathbb{1}(L \neq 0)}{L}\mathbb{1}(A \cap B))) = \mathbb{E}_{\mathbb{P}_1}(\frac{\mathbb{1}(L \neq 0)}{L}\mathbb{1}(A \cap B)). \quad (26)$$

On the other hand, since $\mathbb{P}_1(\{L = 0\} \cap A) = \mathbb{E}_{\mathbb{P}_2}(L\mathbb{1}(\{L = 0\} \cap A)) = 0$, if $\mathbb{P}_2(\{L = 0\} \cap A) > 0$ then we cannot have $\mathbb{P}_2 \ll_A \mathbb{P}_1$. \square

For $A = D$ we omit A in the above notations, e.g. we write $\mathbb{P}_1 \ll \mathbb{P}_2$, $\mathbb{P}_1 \sim \mathbb{P}_2$, and $L = \frac{d\mathbb{P}_1}{d\mathbb{P}_2}$.

3.2 IS and zero- and optimal-variance IS distributions

If $\mathbb{Q}_1 \ll_{Z \neq 0} \mathbb{Q}_2$, then for $L = (\frac{d\mathbb{Q}_1}{d\mathbb{Q}_2})_{Z \neq 0}$ we have $\alpha = \mathbb{E}_{\mathbb{Q}_2}(ZL)$. Importance sampling (IS) relies on estimating α by using in a MC method independent replicates of such IS estimator ZL under \mathbb{Q}_2 . Variance of the IS estimator fulfills

$$\text{Var}_{\mathbb{Q}_2}(ZL) = \mathbb{E}_{\mathbb{Q}_2}((ZL)^2) - \alpha^2 = \mathbb{E}_{\mathbb{Q}_1}(Z^2L) - \alpha^2. \quad (27)$$

Condition 9. *It holds $\mathbb{Q}_1 \ll_{Z \neq 0} \mathbb{Q}_2$ and for some $L = (\frac{d\mathbb{Q}_1}{d\mathbb{Q}_2})_{Z \neq 0}$ it holds $\text{Var}_{\mathbb{Q}_2}(ZL) = 0$ or equivalently \mathbb{Q}_2 a.s. $ZL = \alpha$.*

Theorem 10. *Condition 9 holds only if it holds with 'for some' replaced by 'for each'.*

Proof. Let L be as in Condition 9 and $L' = (\frac{d\mathbb{Q}_1}{d\mathbb{Q}_2})_{Z \neq 0}$. Then, $L = L'$, \mathbb{Q}_2 a.s. on $Z \neq 0$ and $0 = ZL = ZL'$ on $Z = 0$. Thus, from \mathbb{Q}_2 a.s. $ZL = \alpha$ it also holds \mathbb{Q}_2 a.s. $ZL' = \alpha$. \square

Condition 11. It holds $\mathbb{Q}_1(Z \neq 0) > 0$.

Condition 12. We have $\alpha \neq 0$ and \mathbb{Q}_1 a.s. $Z \operatorname{sgn}(\alpha) \geq 0$. Equivalently, for $S = Z$ or $S = -Z$ we have $\mathbb{Q}_1(S > 0) > 0$ and \mathbb{Q}_1 a.s. $S \geq 0$.

Note that Condition 12 implies Condition 11.

Theorem 13. If Condition 12 holds, then for probability \mathbb{Q}^* given by

$$\frac{d\mathbb{Q}^*}{d\mathbb{Q}_1} = \frac{Z}{\alpha}, \quad (28)$$

Condition 9 holds for $\mathbb{Q}_2 = \mathbb{Q}^*$. Furthermore, $\mathbb{Q}^*(Z \neq 0) = 1$ and $\mathbb{Q}^* \sim_{Z \neq 0} \mathbb{Q}_1$ with

$$L^* := \mathbb{1}(Z \neq 0) \frac{\alpha}{Z} = \left(\frac{d\mathbb{Q}_1}{d\mathbb{Q}^*} \right)_{Z \neq 0}. \quad (29)$$

Proof. Condition 12 implies that \mathbb{Q}^* is well-defined. Furthermore, $\mathbb{Q}^*(Z \neq 0) = \mathbb{E}_{\mathbb{Q}_1}(\frac{Z}{\alpha}) = 1$ and from Lemma 8 we have (29). In particular, $ZL^* = \alpha$, \mathbb{Q}^* a.s., that is Condition 9 holds. \square

Lemma 14. Assuming Condition 11, if there exists \mathbb{Q}_2 fulfilling Condition 9, then Condition 12 holds and such \mathbb{Q}_2 is equal to \mathbb{Q}^* as in Theorem 13.

Proof. Condition 9 implies that \mathbb{Q}_2 a.s.

$$\mathbb{1}(Z \neq 0) \frac{\alpha}{Z} = \mathbb{1}(Z \neq 0) L = \left(\frac{d\mathbb{Q}_1}{d\mathbb{Q}_2} \right)_{Z \neq 0}. \quad (30)$$

Thus, from Condition 11, $\mathbb{E}_{\mathbb{Q}_2}(\mathbb{1}(Z \neq 0) \frac{\alpha}{Z}) = \mathbb{Q}_1(Z \neq 0) > 0$, which implies that $\alpha \neq 0$. From (30) and Lemma 8 we have $\mathbb{Q}_1 \sim_{Z \neq 0} \mathbb{Q}_2$ and $\frac{Z}{\alpha} = \left(\frac{d\mathbb{Q}_2}{d\mathbb{Q}_1} \right)_{Z \neq 0}$. Thus, $\mathbb{Q}_2(Z \neq 0) = \mathbb{E}_{\mathbb{Q}_1}(\mathbb{1}(Z \neq 0) \frac{Z}{\alpha}) = 1$ and $\frac{Z}{\alpha} = \frac{d\mathbb{Q}_2}{d\mathbb{Q}_1}$. In particular, \mathbb{Q}_1 a.s. $Z \operatorname{sgn}(\alpha) \geq 0$ and thus Condition 12 holds and $\mathbb{Q}_2 = \mathbb{Q}^*$. \square

Theorem 15. If Condition 12 holds, then \mathbb{Q}^* as in Theorem 13 is the unique distribution \mathbb{Q}_2 for which Condition 9 holds.

Proof. Since Condition 12 implies Condition 11 and Theorem 13 implies existence of \mathbb{Q}_2 fulfilling Condition 9, from Lemma 14, $\mathbb{Q}_2 = \mathbb{Q}^*$. \square

We shall call \mathbb{Q}^* as in Theorem 13 the zero-variance IS distribution. Assuming that $L = \left(\frac{d\mathbb{Q}_1}{d\mathbb{Q}_2} \right)_{Z \neq 0}$, from (27),

$$\operatorname{Var}_{\mathbb{Q}_2}(|ZL|) = \mathbb{E}_{\mathbb{Q}_1}(Z^2 L) - (\mathbb{E}_{\mathbb{Q}_1}(|Z|))^2, \quad (31)$$

and thus

$$\operatorname{Var}_{\mathbb{Q}_2}(ZL) = \operatorname{Var}_{\mathbb{Q}_2}(|ZL|) + (\mathbb{E}_{\mathbb{Q}_1}(|Z|))^2 - \alpha^2 \geq (\mathbb{E}_{\mathbb{Q}_1}(|Z|))^2 - \alpha^2, \quad (32)$$

with equality holding only if Condition 9 holds for $|Z|$ (i.e. for Z replaced by $|Z|$ and in particular for α replaced by $\mathbb{E}_{\mathbb{Q}_1}(|Z|)$). Let Condition 11 hold. Then, Condition 12 holds for $|Z|$ and from Theorem 15, \mathbb{Q}^* as in Theorem 13 but for Z replaced by $|Z|$, i.e. such that

$$\frac{d\mathbb{Q}^*}{d\mathbb{Q}_1} = \frac{|Z|}{\mathbb{E}_{\mathbb{Q}_1}(|Z|)}, \quad (33)$$

is the unique distribution \mathbb{Q}_2 for which Condition 9 holds for $|Z|$. Furthermore,

$$L^* := \mathbb{1}(Z \neq 0) \frac{\mathbb{E}_{\mathbb{Q}_1}(|Z|)}{|Z|} = \left(\frac{d\mathbb{Q}_1}{d\mathbb{Q}^*}\right)_{Z \neq 0}. \quad (34)$$

Note that from Condition 9 holding for $|Z|$ and (31), \mathbb{Q}^* a.s. (or equivalently \mathbb{Q}_1 a.s. on $Z \neq 0$)

$$|Z|L^* = \mathbb{E}_{\mathbb{Q}_1}(|Z|) = \sqrt{\mathbb{E}_{\mathbb{Q}_1}(Z^2 L^*)}. \quad (35)$$

We call such \mathbb{Q}^* the optimal-variance IS distribution. Under Condition 12 the optimal-variance IS distribution is also the zero-variance one. In some places in the literature our optimal-variance IS distribution is called simply the optimal IS distribution (see e.g. page 127 in [4]). However, since as argued in Section 2 it may be more optimal to minimize inefficiency constant than variance and optimal-variance IS distribution does not need to lead to the lowest inefficiency constant achievable via IS, calling it optimal may be misleading.

3.3 Mean costs and inefficiency constants in IS

Let $L = \left(\frac{d\mathbb{Q}_1}{d\mathbb{Q}_2}\right)_{Z \neq 0}$ and let C be a nonnegative (theoretical) cost variable on \mathcal{S}_1 for computing replicates of ZL under \mathbb{Q}_2 . We shall consider C to be the same for different \mathbb{Q}_2 under consideration. Mean cost under \mathbb{Q}_2 is $\mathbb{E}_{\mathbb{Q}_2}(C)$ and such inefficiency constant is

$$\text{Var}_{\mathbb{Q}_2}(ZL)\mathbb{E}_{\mathbb{Q}_2}(C), \quad (36)$$

assuming that it is well-defined (i.e. we do not multiply zero by infinity). Note that if the zero-variance IS distribution \mathbb{Q}^* exists and mean cost $\mathbb{E}_{\mathbb{Q}^*}(C)$ is finite, then inefficiency constant under \mathbb{Q}^* is zero.

The below theorem provides an intuition why in our numerical experiments in Section 9, for some $a, s \in \mathbb{R}_+$, for a nonincreasing function $f(x) = \mathbb{1}(x < s) + a$ and a strictly decreasing one $f(x) = \exp(-sx)$, for $Z = f(C)$, we observed mean cost reduction after changing the initial distribution to a one in a sense closer to the respective zero-variance IS distribution \mathbb{Q}^* .

Theorem 16. *Let $f : \mathcal{S}(\mathbb{R}) \rightarrow \mathcal{S}([0, \infty))$, $Z = f(C)$, $\mathbb{E}_{\mathbb{Q}_1}(Z) \in \mathbb{R}_+$, and $\mathbb{E}_{\mathbb{Q}_1}(C) < \infty$. Let \mathbb{Q}^* be the zero-variance IS distribution.*

1. *If f is nonincreasing, then*

$$\mathbb{E}_{\mathbb{Q}^*}(C) \leq \mathbb{E}_{\mathbb{Q}_1}(C), \quad (37)$$

and if further for some $0 \leq x_1 < x_2 < \infty$ we have $f(x_1) > f(x_2)$, $\mathbb{Q}_1(C \in [0, x_1]) > 0$, and $\mathbb{Q}_1(C \in [x_2, \infty)) > 0$ (which is the case e.g. if f is strictly decreasing and C is not \mathbb{Q}_1 a.s. constant), then the inequality in (37) is sharp.

2. *If f is nondecreasing, then*

$$\mathbb{E}_{\mathbb{Q}^*}(C) \geq \mathbb{E}_{\mathbb{Q}_1}(C), \quad (38)$$

and if further for some $0 \leq x_1 < x_2 < \infty$ we have $f(x_1) < f(x_2)$, $\mathbb{Q}_1(C \in [0, x_1]) > 0$, and $\mathbb{Q}_1(C \in [x_2, \infty)) > 0$, then the inequality in (38) is sharp.

Proof. We have

$$\mathbb{E}_{\mathbb{Q}^*}(C) = \frac{\mathbb{E}_{\mathbb{Q}_1}(f(C)C)}{\mathbb{E}_{\mathbb{Q}_1}(f(C))}. \quad (39)$$

For C_1 and C_2 being independent replicates of C under \mathbb{Q}_1 , we have

$$\mathbb{E}_{\mathbb{Q}_1}(f(C)C) - \mathbb{E}_{\mathbb{Q}_1}(C)\mathbb{E}_{\mathbb{Q}_1}(f(C)) = \frac{1}{2}\mathbb{E}_{\mathbb{Q}_1}((f(C_1) - f(C_2))(C_1 - C_2)), \quad (40)$$

which is nonpositive if f is nonincreasing and negative under the additional assumptions of point one, or nonnegative if f is nondecreasing and positive under the additional assumptions of point two. From this and (39), the thesis easily follows. \square

3.4 IS using parametric families of distributions

For some nonempty set A , let us consider a family $\mathcal{W} = \{\mathbb{Q}(b) : b \in A\}$ of probability distributions on \mathcal{S}_1 . Typically, we shall assume that

$$A \in \mathcal{B}(\mathbb{R}^l). \quad (41)$$

Consider a function $L : A \times \Omega_1 \rightarrow \mathbb{R}$, for which we denote $L(b) = L(b, \cdot)$, $b \in A$. If the following condition is fulfilled, then one can perform IS with respect to distributions from \mathcal{W} as in Section 3.2.

Condition 17. *It holds $\mathbb{Q}_1 \ll_{Z \neq 0} \mathbb{Q}(b)$ and $L(b) = (\frac{\mathbb{Q}_1}{\mathbb{Q}(b)})_{Z \neq 0}$, $b \in A$.*

The following conditions will be useful further on.

Condition 18. *We have (41) and L is measurable from $\mathcal{S}(A) \times \mathcal{S}_1$ to $\mathcal{S}(\mathbb{R})$.*

Condition 19. *We have (41) and probability measure \mathbb{P}_1 on a measurable space \mathcal{C}_1 and $\xi : \mathcal{C}_1 \otimes \mathcal{S}(A) \rightarrow \mathcal{S}_1$ are such that for each $b \in A$*

$$\mathbb{Q}(b)(B) = \mathbb{P}_1(\xi(\cdot, b)^{-1}[B]), \quad B \in \mathcal{F}_1, \quad (42)$$

or equivalently, for each random variable $X \sim \mathbb{P}_1$, $\xi(X, b) \sim \mathbb{Q}(b)$.

Remark 20. *Let conditions 17, 18, and 19 hold and let b be some A -valued random variable, which can be e.g. some adaptive IS parameter. Let $\beta_i \sim \mathbb{P}_1$, $i \in \mathbb{N}_+$, be i.i.d. and independent of b . Then, from Fubini's theorem it follows that random variables $\hat{\alpha}_n = \frac{1}{N} \sum_{i=1}^N (ZL(b))(\xi(\beta_i, b))$, $n \in \mathbb{N}_+$, are unbiased and strongly consistent estimators of α , i.e. $\mathbb{E}(\hat{\alpha}_n) = \alpha$, $n \in \mathbb{N}_+$, and a.s. $\lim_{n \rightarrow \infty} \hat{\alpha}_n = \alpha$.*

In the further sections we shall often deal with families of distributions and densities satisfying the following condition.

Condition 21. *For some $B_1 \in \mathcal{F}_1$, we have $\mathbb{Q}(b) \sim_{B_1} \mathbb{Q}_1$ and $L(b) = (\frac{d\mathbb{Q}_1}{d\mathbb{Q}(b)})_{B_1}$, $b \in A$.*

Let us formulate separately the special important case of the above condition.

Condition 22. *Condition 21 holds for $B_1 = \Omega_1$, or equivalently $\mathbb{Q}(b) \sim \mathbb{Q}_1$ and $L(b) = \frac{d\mathbb{Q}_1}{d\mathbb{Q}(b)}$, $b \in A$.*

The following condition will be useful to avoid different technical problems like when dividing by L or taking its logarithm.

Condition 23. It holds $L(b)(\omega) > 0$, $b \in A$, $\omega \in \Omega_1$.

Condition 24. Condition 21 holds and $\mathbb{Q}(b)(\{Z \neq 0\} \setminus B_1) = 0$, $b \in A$.

Remark 25. Note that for Z such that Condition 24 holds, we have $\mathbb{Q}(b) \sim_{Z \neq 0 \cup B_1} \mathbb{Q}_1$, $b \in A$, and

$$L(b) = \left(\frac{d\mathbb{Q}_1}{d\mathbb{Q}(b)} \right)_{B_1 \cup \{Z \neq 0\}}, \quad b \in A, \quad (43)$$

so that Condition 17 holds.

Definition 26. We say that $b^* \in A$ is the zero-variance (optimal-variance) IS parameter if Condition 12 (Condition 11) holds and $\mathbb{Q}(b^*)$ is the zero-variance (optimal-variance) IS distribution.

Note that in the literature the name optimal IS parameter is sometimes used for the parameter minimizing variance $b \in A \rightarrow \text{Var}_{\mathbb{Q}(b)}(ZL(b))$ of the IS estimator (see e.g. [35]), which may be not equal to the optimal-variance IS parameter in the sense of the above definition.

The below theorem characterizes random variables Z as above for which there exists zero-variance IS parameter, under some of the above conditions.

Theorem 27. Let us assume Condition 21. Then, Condition 24 holds and there exists zero-variance IS parameter b_1 (for which we denote $\mathbb{Q}^* = \mathbb{Q}(b_1)$), only if for some $b_2 \in A$, $\mathbb{Q}(b_2)(B_1) = 1$ and for some $\beta \in \mathbb{R} \setminus 0$,

$$Z = \mathbb{1}_{B_1} \mathbb{1}(L(b_2) \neq 0) \frac{\beta}{L(b_2)}, \quad \mathbb{Q}_1 \text{ a.s.} \quad (44)$$

Furthermore, in the latter case we have $\beta = \alpha$ and $\mathbb{Q}(b_2) = \mathbb{Q}^*$.

Proof. Let us first show the right implication. From Condition 24 and $\mathbb{Q}^* = \mathbb{Q}(b_1)$ it follows for $b_2 = b_1$ that $\mathbb{Q}(b_2)(B_1) = \mathbb{Q}^*(B_1) = \mathbb{Q}^*(Z \neq 0 \cap B_1) = \mathbb{Q}^*(Z \neq 0) - \mathbb{Q}(b_2)(\{Z \neq 0\} \setminus B_1) = 1$. From (29), \mathbb{Q}^* a.s. $L(b_2)Z = \mathbb{1}(Z \neq 0)\alpha$, which from $\mathbb{Q}^* \sim_{Z \neq 0} \mathbb{Q}_1$ holds also \mathbb{Q}_1 a.s. Thus, since from Condition 12 we have $\alpha \neq 0$, it holds \mathbb{Q}_1 a.s. that if $Z \neq 0$ then also $L(b_2) \neq 0$. Therefore, we have \mathbb{Q}_1 a.s. $Z = \mathbb{1}(Z \neq 0 \wedge L(b_2) \neq 0) \frac{1}{L(b_2)} \alpha$. Thus, from Condition 24,

$$Z = \mathbb{1}_{B_1} \frac{1}{L(b_2)} \mathbb{1}(Z \neq 0 \wedge L(b_2) \neq 0) \alpha, \quad \mathbb{Q}_1 \text{ a.s.} \quad (45)$$

From Condition 21, $\mathbb{Q}^* \sim_{B_1} \mathbb{Q}_1$, and thus from Lemma 8 and (28), $0 = \mathbb{Q}_1(\{L(b_2) = 0\} \cap B_1) = \mathbb{Q}_1(\{Z = 0\} \cap B_1)$, so that from (45) and Condition 12 we have (44) only for $\beta = \alpha$. For the left implication note that for Z as in (44) Condition 24 holds and since from (44)

$$\alpha = \mathbb{E}_{\mathbb{Q}(b_2)}(ZL(b_2)) = \beta \mathbb{Q}(b_2)(B_1) = \beta, \quad (46)$$

Condition 12 holds as well. Thus

$$\frac{d\mathbb{Q}(b_2)}{d\mathbb{Q}_1} = \mathbb{1}(B_1) \mathbb{1}(L(b_2) \neq 0) \frac{1}{L(b_2)} = \frac{Z}{\beta} = \frac{d\mathbb{Q}^*}{d\mathbb{Q}_1}, \quad (47)$$

where in the first equality we used Condition 21, $\mathbb{Q}(b_2)(B_1) = 1$, and Lemma 8, in the second (44), and in the last (46). \square

Remark 28. From discussion in Section 3.2, the optimal-variance IS distribution for Z is the zero-variance one for $|Z|$. Thus, from the above theorem for Z replaced by $|Z|$ we receive a characterization of variables Z for which there exists optimal-variance IS parameter under certain assumptions.

4 Examples of parametric families of IS distributions

In this section we present a number of parametric families of distributions, most of which shall be used in theoretical reasonings or numerical experiments in this work.

4.1 Exponential change of measure

Exponential change of measure (ECM), also known as exponential tilting, is a popular method for obtaining a family of IS distributions from a given one. It has found numerous applications among others in rare event simulation, as discussed in Section 1.2. Consider an \mathbb{R}^l -valued random vector X on \mathcal{S}_1 . Let us define moment generating function $b \in \mathbb{R}^l \rightarrow \Phi(b) = \mathbb{E}_{\mathbb{Q}_1}(\exp(b^T X))$. Let A be the set of all $b \in \mathbb{R}^l$ for which $\Phi(b) < \infty$. Note that $0 \in A$ and from convexity of exponential function A is convex. Cumulant generating function is defined as $\Psi(b) = \ln(\Phi(b))$, $b \in A$.

Condition 29. For each $b_1, b_2 \in A$, $b_1 \neq b_2$, $(b_1 - b_2)^T X$ is not \mathbb{Q}_1 a.s. constant.

Lemma 30. Ψ is convex on A and it is strictly convex on A only if Condition 29 holds.

Proof. Let $b_1, b_2 \in A$ and $q_1, q_2 \in \mathbb{R}_+$ be such that $q_1 + q_2 = 1$. From Hölder inequality

$$\Phi\left(\sum_{i=1}^2 q_i b_i\right) \leq \prod_{i=1}^2 \Phi(b_i)^{q_i} \quad (48)$$

and taking the logarithms of both sides we receive

$$\Psi\left(\sum_{i=1}^2 q_i b_i\right) \leq \sum_{i=1}^2 q_i \Psi(b_i), \quad (49)$$

and thus Ψ is convex. Equality in (48) or equivalently in (49) holds only if for some $a \in \mathbb{R}_+$, \mathbb{Q}_1 a.s. $\exp(b_1^T X) = a \exp(b_2^T X)$ (see page 63 in [48]) or equivalently if for some $c \in \mathbb{R}$, \mathbb{Q}_1 a.s. $(b_1 - b_2)^T X = c$. Ψ is strictly convex only if there do not exist $b_1, b_2 \in A$, $b_1 \neq b_2$, such that equality in (49) holds, and thus only if Condition 29 holds. \square

Condition 31. For each $t \in \mathbb{R}^l \setminus \{0\}$, $t^T X$ is not \mathbb{Q}_1 a.s. constant.

Note that Condition 31 implies Condition 29 and if A has nonempty interior then these conditions are equivalent. If A contains some neighbourhood of zero, then X has finite all mixed moments, i.e. $\mathbb{E}(\prod_{i=1}^l |X_i|^{v_i}) < \infty$, $v \in \mathbb{N}^l$. Let us denote for $v \in \mathbb{N}^l$, $\partial_v = \frac{\partial^{v_1 + \dots + v_l}}{\partial_{b_1}^{v_1} \dots \partial_{b_l}^{v_l}}$.

Condition 32. A is open, Φ is smooth (i.e. infinitely continuously differentiable) on A , and for each $v \in \mathbb{N}^l$ we have

$$\partial_v \Phi(b) = \mathbb{E}_{\mathbb{Q}_1}(\partial_v \exp(b^T X)) = \mathbb{E}_{\mathbb{Q}_1}\left(\prod_{i=1}^l X_i^{v_i} \exp(b^T X)\right). \quad (50)$$

Remark 33. It is easy to show using inductively mean value theorem and Lebesgue dominated convergence theorem that Condition 32 holds when $A = \mathbb{R}^l$ or when $\mathbb{Q}_1([0, \infty)^l) = 1$ and for some $\lambda > 0$, $A = (-\infty, \lambda)^l$.

We define an exponentially tilted family of probability distributions $\mathcal{W} = \{\mathbb{Q}(b), b \in A\}$ corresponding to the above \mathbb{Q}_1 and X by formula

$$\frac{d\mathbb{Q}(b)}{d\mathbb{Q}_1} = \exp(b^T X - \Psi(b)), \quad b \in A. \quad (51)$$

Note that $\mathbb{Q}(0) = \mathbb{Q}_1$ and

$$L(b) := \exp(-b^T X + \Psi(b)) = \frac{d\mathbb{Q}_1}{d\mathbb{Q}(b)}, \quad b \in A. \quad (52)$$

In particular, conditions 18, 22, and 23 hold (for the above L). From Lemma 30, for each $\omega \in \Omega_1$, $b \in A \rightarrow L(b)(\omega)$ is log-convex (and thus convex) and if Condition 29 holds, then it is strictly log-convex (and thus strictly convex). Let us define means $\mu(b) = \mathbb{E}_{\mathbb{Q}(b)}(X)$ and covariance matrices $\Sigma(b) = \mathbb{E}_{\mathbb{Q}(b)}(X - \mu(b))(X - \mu(b))^T$, for $b \in A$ for which they exist. Note that functions Φ , Ψ , Σ , and μ depend only on the law of X under \mathbb{Q}_1 . Assuming that $\Sigma(b) \in \mathbb{R}^{l \times l}$, we have $t^T \Sigma(b) t = \mathbb{E}_{\mathbb{Q}(b)}((t^T(X - \mu(b)))^2)$, $t \in \mathbb{R}^l$, and thus $\Sigma(b)$ is positive definite only if Condition 31 holds. When Condition 32 holds, then we receive by direct calculation that $\nabla \Psi(b) = \mu(b)$ and $\nabla^2 \Psi(b) = \Sigma(b)$, $b \in A$. Let U be an open subset of \mathbb{R}^l . The following well-known lemma is an easy consequence of the inverse function theorem.

Lemma 34. If $f : U \rightarrow \mathbb{R}^l$ is injective and differentiable with invertible derivative Df on U , then f is a diffeomorphism of the open sets U and $f(U)$.

Lemma 35. If U is convex and a function $g : U \rightarrow \mathbb{R}$ is strictly convex and differentiable, then function $b \in U \rightarrow \nabla g(b)$ is injective.

Proof. If for some $b_1, b_2 \in U$, $b_1 \neq b_2$, we had $\nabla g(b_1) = \nabla g(b_2)$, then for $v = \frac{b_2 - b_1}{|b_2 - b_1|}$ it would hold

$$\left(\frac{dg(b_1 + tv)}{dt}\right)_{t=0} = v^T \nabla g(b_1) = v^T \nabla g(b_2) = \left(\frac{dg(b_1 + tv)}{dt}\right)_{t=|b_2 - b_1|} \quad (53)$$

which is impossible since $t \in [0, |b_2 - b_1|] \rightarrow g(b_1 + tv)$ is strictly convex. \square

Theorem 36. If conditions 31 and 32 hold, then $b \in A \rightarrow \mu(b) = \nabla \Psi(b)$ is a diffeomorphism of the open sets A and $\mu[A]$.

Proof. From Condition 31 and Lemma 30, Ψ is strictly convex. From Condition 32, $D\mu = \nabla^2 \Psi$ is positive definite. Thus, for $U = A$ the thesis follows from Lemma 35 for $g = \Psi$ and Lemma 34 for $f = \mu$. \square

Some important special cases of ECM for $l = 1$ are when X has binomial, Poisson, or gamma distribution under $\mathbb{Q}(b)$, $b \in A$, while for general $l \in \mathbb{N}_+$ — when X has such multivariate normal distribution (see page 130 in [4]). In all these cases, from Remark 33, Condition 32 holds. Furthermore, for the first three cases and non-degenerate multivariate normal distributions, Condition 31 is satisfied and we have analytical formulas for μ^{-1} . In the gamma case, for some $\alpha, \lambda \in \mathbb{R}_+$, and $A = (-\infty, \lambda)$, for each $b \in A$, for $\lambda_b = \lambda - b$, under $\mathbb{Q}(b)$, X has distribution with density

$$\frac{1}{\Gamma(\alpha)} \lambda_b^\alpha x^{\alpha-1} \exp(-\lambda_b x) \quad (54)$$

with respect to the Lebesgue measure on $(0, \infty)$. Furthermore, it holds $\Psi(b) = \alpha \ln(\frac{\lambda}{\lambda-b})$, $\mu(b) = \frac{\alpha}{\lambda-b}$, and $\mu^{-1}(x) = \lambda - \frac{\alpha}{x}$, $x \in \mu[A] = \mathbb{R}_+$. In the Poisson case we have $A = \mathbb{R}$, and for some initial mean $\mu_0 \in \mathbb{R}_+$, for each $b \in A$ we have $\mu(b) = \mu_0 \exp(b)$ and

$$\mathbb{Q}(b)(X = k) = \frac{\mu(b)^k}{k!} \exp(-\mu(b)), \quad k \in \mathbb{N}, \quad (55)$$

i.e. $X \sim \text{Pois}(\mu(b))$ under $\mathbb{Q}(b)$. Furthermore, it holds $\Psi(b) = \mu_0(\exp(b) - 1)$, $\mu^{-1}(x) = \ln(\frac{x}{\mu_0})$, $x \in \mu[A] = (0, \infty)$, and $\Sigma(b) = \mu(b)$. In the multivariate normal case we have $A = \mathbb{R}^l$ and for $M \in \mathbb{R}^{l \times l}$ being some positive semidefinite covariance matrix and $\mu_0 \in \mathbb{R}^l$ some initial mean, the mean function has form $\mu(b) = \mu_0 + Mb$ and under $\mathbb{Q}(b)$, $X \sim \mathcal{N}(\mu(b), M)$. Moreover, it holds $\Psi(b) = b^T \mu_0 + \frac{1}{2} b^T M b$ and $\Sigma(b) = M$. An important special case are non-degenerate normal distributions in which M is positive definite, $\mu[A] = A$, and $\mu^{-1}(x) = M^{-1}(x - \mu_0)$. In the standard multivariate normal case we have $M = I_l$ and $\mu_0 = 0$, so that $\mathbb{Q}(b) = \mathbb{N}(b, I_l)$, $b \in A$.

For exponentially tilted distributions with $A = \mathbb{R}^l$ we shall further need the following function defined for $a \in [0, \infty)$

$$F(a) = \sup\{\Psi(b) : b \in \mathbb{R}^l, |b| \leq a\}. \quad (56)$$

For instance, for $X \sim \mathcal{N}(0, I_l)$ under \mathbb{Q}_1 as above we have $F(a) = \frac{a^2}{2}$, while for the Poisson case it holds $F(a) = \mu_0(\exp(a) - 1)$.

Remark 37. *In some practical realizations of ECM, the computation times on a computer needed to generate i.i.d. replicates of the IS estimator $ZL(b)$ under $\mathbb{Q}(b)$ for different $b \in A$ are approximately equal to the same constant. This is typically the case e.g. when $X \sim \mathcal{N}(0, I_l)$ under \mathbb{Q}_1 . In such case one can take the theoretical cost $C = 1$.*

4.2 IS for independently parametrized product distributions

For $n \in \mathbb{N}_+$ and each $i \in \{1, \dots, n\}$, consider some probability distribution $\tilde{\mathbb{Q}}_{1,i}$ on a measurable space $\mathcal{S}_{1,i} = (\Omega_{1,i}, \mathcal{F}_{1,i})$, a nonempty set A_i , and a parametric family of distributions $\mathcal{W}_i = \{\tilde{\mathbb{Q}}_i(b_i) : b_i \in A_i\}$ and densities $\tilde{L}_i(b_i) = \frac{d\tilde{\mathbb{Q}}_{1,i}}{d\tilde{\mathbb{Q}}_i(b_i)}$, $b_i \in A_i$. Let us define a product measure $\mathbb{Q}_1 = \prod_{i=1}^n \tilde{\mathbb{Q}}_{1,i}$, a product parameter set $A = \prod_{i=1}^n A_i$, a family of independently parametrized product measures

$\mathcal{W} = \{\mathbb{Q}(b) = \prod_{i=1}^n \tilde{\mathbb{Q}}(b_i) : b \in A\}$, and densities $L(b) = \prod_{i=1}^n \tilde{L}_i(b_i)$, $b = (b_i)_{i=1}^n \in A$. Note that $\mathbb{Q}(b) \sim \mathbb{Q}_1$ and $L(b) = \frac{d\mathbb{Q}_1}{d\mathbb{Q}(b)}$, $b \in A$.

Let us consider the special case of \mathcal{W}_i being exponentially tilted families of distributions given by probabilities $\tilde{\mathbb{Q}}_{1,i}$ and random variables \tilde{X}_i , having moment-generating functions Φ_i , and cumulant generating functions Ψ_i , $i = 1, \dots, n$. Then, \mathcal{W} is an exponentially tilted family of distributions corresponding to probability \mathbb{Q}_1 and random variable $X(\omega) = (\tilde{X}_i(\omega_i))_{i=1}^n$, $\omega \in \prod_{i=1}^n \Omega_{1,i}$, with moment-generating function $\Phi(b) = \prod_{i=1}^n \Phi_i(b_i)$, and cumulant generating function $\Psi(b) = \sum_{i=1}^n \Psi_i(b_i)$. If Condition 31 or 32 holds for \mathcal{W}_i for each $i \in \{1, \dots, n\}$, then such condition holds also for \mathcal{W} . If for $b = (b_i)_{i=1}^n \in A$, $\mu_i(b_i)$ are mean vectors for \mathcal{W}_i , $i = 1, \dots, n$, then $\mu(b) = (\mu_i(b_i))_{i=1}^n$ is such mean vector for \mathcal{W} and if all μ_i^{-1} exist, then for $x = (x_i)_{i=1}^n \in \mu[A] = \prod_{i=1}^n \mu_i[A_i]$, $\mu^{-1}(x) = (\mu_i^{-1}(x_i))_{i=1}^n$.

4.3 IS for stopped sequences

4.3.1 Change of measure for stopped sequences using a tilting process

Let $\tilde{\mathbb{U}}_1$ be a probability measure on a measurable space $\tilde{\mathcal{C}} = (\tilde{E}, \tilde{\mathcal{E}})$, let $\mathcal{C} = (E, \mathcal{E}) := \tilde{\mathcal{C}}^{\mathbb{N}_+}$, let $\eta = (\eta_i)_{i \in \mathbb{N}_+} = \text{id}_E$ be the coordinate process on E , and let $\tilde{\eta}_k = (\eta_i)_{i=1}^k$, $k \in \mathbb{N}_+$. Let \mathbb{U} be the unique probability measure on \mathcal{C} such that η_1, η_2, \dots , are i.i.d. $\sim \tilde{\mathbb{U}}_1$ under \mathbb{U} (see Theorem 16, Chapter 9 in [17]). Let $\mathcal{F}_k = \sigma(\tilde{\eta}_k)$, $k \in \mathbb{N}_+$, i.e. it is the natural filtration of η , and let $\mathcal{F}_0 = \{\emptyset, E\}$, i.e. it is a trivial σ -field. For some $d \in \mathbb{N}_+$ and nonempty $B \in \mathcal{B}(\mathbb{R}^d)$, let conditions 18, 22, and 23 hold for $A = B$, $\mathbb{Q}_1 = \tilde{\mathbb{U}}_1$, and some probabilities $\mathbb{Q}(b)$ and densities $L(b)$ denoted further as $\tilde{\mathbb{U}}(b)$ and $\tilde{L}(b)$, $b \in B$. Let $\kappa(b) = \tilde{L}(b)^{-1} = \frac{d\tilde{\mathbb{U}}(b)}{d\tilde{\mathbb{U}}_1}$, $b \in B$.

Definition 38. We define \mathcal{J} to be the set of all B -valued stochastic processes $\lambda = (\lambda_k)_{k \in \mathbb{N}}$ on \mathcal{C} such that there exist $h_0 \in B$ and measurable functions $h_k : \tilde{\mathcal{C}}^k \rightarrow \mathcal{S}(B)$, $k \in \mathbb{N}_+$, such that $\lambda_0 = h_0$ and $\lambda_k = h_k(\tilde{\eta}_k)$, $k \in \mathbb{N}_+$.

Processes λ as in the above definition shall be called tilting processes. Note that such λ are adapted to $(\mathcal{F}_k)_{k \in \mathbb{N}}$. If $\tilde{\mathcal{C}}$ is a Borel space like $\mathcal{S}(\mathbb{R}^d)$, then \mathcal{J} is simply the set of all $(\mathcal{F}_k)_{k \in \mathbb{N}}$ -adapted B -valued stochastic processes on \mathcal{C} (see Chapter 21 in [17]). Let further in this section λ and h_i be as in Definition 38. For a probability \mathbb{S} on \mathcal{C} and a stopping time τ for $(\mathcal{F}_k)_{k \in \mathbb{N}}$, we shall denote $\mathbb{S}_{|\tau} = \mathbb{S}_{|\mathcal{F}_\tau}$. Identifying each $n \in \mathbb{N}_+$ with a constant random variable we thus have $\mathbb{S}_{|n} = \mathbb{S}_{|\mathcal{F}_n}$. The following theorem is an easy consequence of Theorem 3, Chapter 22 in [17].

Theorem 39. There exists a unique probability \mathbb{V} on \mathcal{C} satisfying one of the following equivalent conditions.

1. Under \mathbb{V} , η_1 has density $\kappa(h_0)$ with respect to $\tilde{\mathbb{U}}_1$ and for each $k \in \mathbb{N}_+$, η_{k+1} has conditional density $\kappa(\lambda_k)$ with respect to $\tilde{\mathbb{U}}_1$ given \mathcal{F}_k (see Definition 14, Chapter 21 in [17]).

2. For each $n \in \mathbb{N}_+$,

$$\frac{d\mathbb{V}|_n}{d\mathbb{U}|_n} = \gamma_n := \prod_{k=0}^{n-1} \kappa(\lambda_k)(\eta_{k+1}). \quad (57)$$

Let \mathbb{V} and γ_n , $n \in \mathbb{N}_+$, be as in the above theorem, and let τ be a stopping time for $(\mathcal{F}_n)_{n \in \mathbb{N}}$.

Lemma 40. *It holds*

$$\mathbb{V}|_\tau \sim_{\tau < \infty} \mathbb{U}|_\tau, \quad (58)$$

with

$$\mathbb{1}(\tau < \infty)\gamma_\tau = \left(\frac{d\mathbb{V}|_\tau}{d\mathbb{U}|_\tau} \right)_{\tau < \infty}. \quad (59)$$

Proof. To prove (59) we notice that for each $B \in \mathcal{F}_\tau$ we have

$$\begin{aligned} \mathbb{E}_\mathbb{U}(\mathbb{1}(B \cap \{\tau < \infty\})\gamma_\tau) &= \sum_{n=0}^{\infty} \mathbb{E}_\mathbb{U}(\mathbb{1}(\{\tau = n\} \cap B)\gamma_n) \\ &= \sum_{n=0}^{\infty} \mathbb{V}(\{\tau = n\} \cap B) = \mathbb{V}(B \cap \{\tau < \infty\}), \end{aligned} \quad (60)$$

where in the second equality we used (57) and the fact that $\{\tau = n\} \cap B \in \mathcal{F}_n$, $n \in \mathbb{N}_+$. Now, (58) follows from (59) and Lemma 8. \square

From the above lemma, if $\tau < \infty$ both \mathbb{V} and \mathbb{U} a.s., then $\mathbb{V}|_\tau \sim \mathbb{U}|_\tau$. For some $\epsilon \in \mathbb{R}_+$, considered to avoid some technical problems as discussed above Condition 23, let us define

$$L = \mathbb{1}(\tau < \infty) \frac{1}{\gamma_\tau} + \epsilon \mathbb{1}(\tau = \infty) = \mathbb{1}(\tau < \infty) \prod_{k=0}^{\tau-1} \tilde{L}(\lambda_k(\eta_{k+1})) + \epsilon \mathbb{1}(\tau = \infty). \quad (61)$$

Let Z be an $\bar{\mathbb{R}}$ -valued random variable measurable from (E, \mathcal{F}_τ) and such that $\mathbb{E}_\mathbb{U}(|Z|) < \infty$ (shortly, $Z \in L^1(\mathbb{U}|_\tau)$). Let us assume that $\mathbb{U}(Z \neq 0, \tau = \infty) = \mathbb{V}(Z \neq 0, \tau = \infty) = 0$, so that from (58), $\mathbb{U}|_\tau \sim_{Z \neq 0 \vee \tau < \infty} \mathbb{V}|_\tau$ and from Lemma 40,

$$L = \left(\frac{d\mathbb{U}|_\tau}{d\mathbb{V}|_\tau} \right)_{Z \neq 0 \vee \tau < \infty}. \quad (62)$$

Then, one can perform IS as in Section 3.2 for $\mathbb{Q}_1 = \mathbb{U}|_\tau$, $\mathbb{Q}_2 = \mathbb{V}|_\tau$, and L as above. Note that such \mathbb{Q}_1 is defined on $\mathcal{S}_1 = (\Omega_1, \mathcal{F}_1) = (E, \mathcal{F}_\tau)$.

Remark 41. *Consider two stopping times τ_1, τ_2 for $(\mathcal{F}_n)_{n \in \mathbb{N}}$, such that $\tau_1 \leq \tau_2$, $Z \in L^1(\mathbb{U}_{\tau_1})$, and $\mathbb{U}(Z \neq 0, \tau_2 = \infty) = \mathbb{V}(Z \neq 0, \tau_2 = \infty) = 0$. Then, we also have $Z \in L^1(\mathbb{U}_{\tau_2})$ and $\mathbb{U}(Z \neq 0, \tau_1 = \infty) = \mathbb{V}(Z \neq 0, \tau_1 = \infty) = 0$. Furthermore, denoting L as in (61) for $\tau = \tau_i$ as L_{τ_i} , we have*

$$\mathbb{E}_\mathbb{V}(ZL_{\tau_2} | \mathcal{F}_{\tau_\infty}) = ZL_{\tau_1}. \quad (63)$$

Indeed, for each $D \in \mathcal{F}_{\tau_1}$ and $i = 1, 2$ from (62) it holds

$$\mathbb{E}_\mathbb{V}(ZL_{\tau_i} \mathbb{1}(D)) = \mathbb{E}_\mathbb{V}(ZL_{\tau_i} \mathbb{1}(D \cap \{Z \neq 0\})) = \mathbb{E}_\mathbb{U}(Z \mathbb{1}(D \cap \{Z \neq 0\})). \quad (64)$$

From (63) and conditional Jensen's inequality we have $\text{Var}_{\mathbb{V}}(ZL_{\tau_2}) \geq \text{Var}_{\mathbb{V}}(ZL_{\tau_1})$, i.e. using τ_1 for IS leads to not higher variance than using τ_2 . Furthermore, $\mathbb{E}_{\mathbb{V}}(\tau_1) \leq \mathbb{E}_{\mathbb{V}}(\tau_2)$, so that for theoretical costs C equal to the stopping times, using τ_1 also leads to not higher mean cost and inefficiency constant than τ_2 (assuming that such constants are well-defined).

4.3.2 Parametrizations of IS for stopped sequences

For some nonempty $A \in \mathcal{B}(\mathbb{R}^l)$, let us consider a parametrization $\lambda : A \rightarrow \mathcal{J}$ of tilting processes such that the following condition is satisfied.

Condition 42. For each $n \in \mathbb{N}$, $(b, x) \rightarrow \lambda_n(b)(x)$ is measurable from $\mathcal{S}(A) \otimes (E, \mathcal{F}_n)$ to $\mathcal{S}(B)$.

For each $b \in A$, let $\mathbb{V}(b)$ and $\mathbb{V}_{|\tau}(b)$ be given by $\lambda(b)$ similarly as \mathbb{V} and $\mathbb{V}_{|\tau}$ are given by λ in the unparametrized case above. Let $\mathbb{Q}(b) = \mathbb{V}_{|\tau}(b)$, $b \in A$, let $\mathbb{Q}_1 = \mathbb{U}_{|\tau}$ be as above, and let $L(b)$ be defined by formula (61) but using $\lambda(b)$ in place of λ . Note that such L satisfies Condition 23. From Condition 42 and assumptions on \tilde{L} above, L also satisfies Condition 18. From (58), Condition 21 is satisfied for $B_1 = \{\tau < \infty\}$. Thus, for such B_1 , Condition 24 is equivalent to a following one.

Condition 43. For each $b \in A$, $\mathbb{Q}(b)(Z \neq 0, \tau = \infty) = 0$.

Condition 22 is implied by a following one.

Condition 44. For each $b \in A$, $\tau < \infty$, $\mathbb{V}(b)$ a.s. (or equivalently $\mathbb{Q}(b)$ a.s.).

Consider a following simple linear parametrization of tilting processes, for which we assume that $B = \mathbb{R}^d$. Let an $\mathbb{R}^{d \times l}$ -valued process $\Lambda = (\Lambda_k)_{k \geq 0}$ on \mathcal{C} be such that for each $j \in \{1, \dots, l\}$, $((\Lambda_k)_{i,j})_{i=1}^d)_{k \in \mathbb{N}} \in \mathcal{J}$, let $A = \mathbb{R}^l$, and for each $b \in A$ let $\lambda(b)$ be such that

$$\lambda_k(b) = \Lambda_k b, \quad k \in \mathbb{N}. \quad (65)$$

Note that Condition 42 holds for such λ and we have $\mathbb{Q}(0) = \mathbb{Q}_1$.

4.3.3 Change of measure for Gaussian stopped sequences

Let $\tilde{\mathbb{U}}_1 = \mathcal{N}(0, I_d)$, $X = \text{id}_{\mathbb{R}^d}$, and let $\tilde{\mathbb{U}}(b)$ and $\tilde{L}(b)$, $b \in B := \mathbb{R}^d$, be exponentially tilted distributions and densities corresponding to such X , $\mathbb{Q}_1 = \tilde{\mathbb{U}}_1$, and $A = B$, as in Section 4.1. For such distributions and densities, let us consider IS for stopped sequences for some tilting process $\lambda \in \mathcal{J}$ and h_k , $k \in \mathbb{N}$, as in Section 4.3.1. In particular, $\kappa(b)(x) = \exp(-\frac{1}{2}|b|^2 + b^T x)$. Let $\dot{\eta}_k = \eta_k - \lambda_{k-1}$, $k \in \mathbb{N}_+$. The following theorem is a discrete version of Girsanov theorem.

Theorem 45. Under \mathbb{V} , random variables $\dot{\eta}_k$, $k \in \mathbb{N}$, are i.i.d. $\sim \mathcal{N}(0, I_d)$.

Proof. Writing h_k in place of $h_k(x_1, \dots, x_k)$, $k \in \mathbb{N}_+$, for each $n \in \mathbb{N}_+$ and

$$\Gamma \in (\mathcal{B}(\mathbb{R}^d))^n$$

$$\begin{aligned} \mathbb{V}((\dot{\eta}_i)_{i=1}^n \in \Gamma) &= \mathbb{E}_{\mathbb{U}}(\mathbb{1}((\dot{\eta}_i)_{i=1}^n \in \Gamma) \gamma_n) \\ &= \int_{(\mathbb{R}^d)^n} \mathbb{1}((x_k - h_{k-1})_{k=1}^n \in \Gamma) \frac{1}{(2\pi)^{nd/2}} \exp\left(-\frac{1}{2} \sum_{k=1}^n (x_k - h_{k-1})^2\right) dx_n \dots dx_1 \\ &= \int_{(\mathbb{R}^d)^n} \mathbb{1}(y \in \Gamma) \frac{1}{(2\pi)^{nd/2}} \exp\left(-\frac{1}{2} \sum_{k=1}^n y_k^2\right) dy_n \dots dy_1 \\ &= \mathbb{U}((\eta_i)_{i=1}^n \in \Gamma), \end{aligned} \tag{66}$$

where we used Fubini's theorem and a sequence of changes of variables $y_k(x_k) = x_k - h_{k-1}$, $k = 1, \dots, n$, each of which is a diffeomorphism with Jacobian 1. \square

Let us consider a function $\pi : E \rightarrow E$ such that $\pi = (\dot{\eta}_i)_{i \in \mathbb{N}_+}$. Its inverse function π^{-1} is given by formula

$$\pi^{-1} = (\eta_k + \lambda_{k-1}(\pi^{-1}))_{k \in \mathbb{N}_+}, \tag{67}$$

or in more detail we have $\pi^{-1} = (\ddot{\eta}_i)_{i=1}^\infty$ for $\ddot{\eta}_i = \eta_i + \ddot{\lambda}_{i-1}$, $i \in \mathbb{N}_+$, where $\ddot{\lambda}_0 = h_0$ and $\ddot{\lambda}_k = h_k((\ddot{\eta}_i)_{i=1}^k)$, $k \in \mathbb{N}_+$. Note that both π and π^{-1} are measurable from $\mathcal{U}_n := (E, \mathcal{F}_n)$ to \mathcal{U}_n , $n \in \mathbb{N}$, i.e. π is an isomorphism of \mathcal{U}_n , $n \in \mathbb{N}$, and thus also of $\mathcal{U}_\infty := (E, \mathcal{F}_\infty) = \mathcal{C}$. From Theorem 45 we have $\mathbb{U}(B) = \mathbb{V}(\pi^{-1}[B])$, $B \in \mathcal{E}$, so that

$$\mathbb{U}(\pi[B]) = \mathbb{V}(B), \quad B \in \mathcal{E}. \tag{68}$$

In particular, for each random variable Y on \mathcal{C} the distribution of $Y\pi^{-1} := Y(\pi^{-1})$ under \mathbb{U} is the same as of Y under \mathbb{V} . For $\overline{\pi}$ denoting the image function of π we have

$$\begin{aligned} \overline{\pi}[\mathcal{F}_\tau] &= \{\pi[B] : B \in \mathcal{F}_\tau\} \\ &= \{\pi[B] : B \in \mathcal{F}_\infty, B \cap \{\tau = k\} \in \mathcal{F}_k, k \in \mathbb{N} \cup \{\infty\}\} \\ &= \{C \in \mathcal{F}_\infty : \pi^{-1}[C] \cap \{\tau = k\} \in \mathcal{F}_k, k \in \mathbb{N} \cup \{\infty\}\} \\ &= \{C \in \mathcal{F}_\infty : C \cap \{\tau\pi^{-1} = k\} \in \mathcal{F}_k, k \in \mathbb{N} \cup \{\infty\}\} \\ &= \mathcal{F}_{\tau\pi^{-1}}, \end{aligned} \tag{69}$$

where in the fourth equality we used the fact that π is an isomorphism of \mathcal{U}_n , $n \in \mathbb{N} \cup \{\infty\}$. In particular, if a random variable Y on \mathcal{C} is \mathcal{F}_τ -measurable, then $Y\pi^{-1}$ is $\mathcal{F}_{\tau\pi^{-1}}$ -measurable, i.e. it depends only on the information available until time $\tau\pi^{-1}$.

For some parametrization $\lambda(b)$, $b \in A$, of tilting processes as in Section 4.3.2, let π_b be given by $\lambda(b)$ in a way that π is given by λ above. Let $\mathbb{Q}(b)$ and $L(b)$, $b \in A$, correspond to such parametrization as in Section 4.3.2. Then, for $\mathcal{C}_1 = \mathcal{C}$ and $\mathbb{P}_1 = \mathbb{U}$, we can choose ξ in Condition 19 to be given by formula

$$\xi(\eta, b) = \pi_b^{-1}(\eta), \quad b \in A. \tag{70}$$

Indeed, from (68),

$$\mathbb{U}(\xi(\cdot, b)^{-1}[B]) = \mathbb{V}_{|\tau}(b)(B), \quad B \in \mathcal{F}_\tau, \tag{71}$$

i.e. (42) holds. For each random variable Y on \mathcal{C} and $b \in A$ we shall denote

$$Y^{(b)} = Y\pi_b^{-1} = Y(\xi(\cdot, b)). \quad (72)$$

Note that from (67) we have

$$(\xi(\eta, b))_k = \eta_k^{(b)} = \eta_k + \lambda_{k-1}(b)^{(b)}, \quad k \in \mathbb{N}_+. \quad (73)$$

For each $b \in A$ it holds

$$L(b) = \mathbb{1}(\tau < \infty) \exp\left(\sum_{k=0}^{\tau-1} \left(\frac{1}{2}|\lambda_k(b)|^2 - \lambda_k(b)^T \eta_{k+1}\right)\right) + \mathbb{1}(\tau = \infty)\epsilon. \quad (74)$$

Thus, for each $b', b \in A$ we have

$$L(b)^{(b')} = \mathbb{1}(\tau^{(b')} < \infty) \exp\left(\sum_{k=0}^{\tau^{(b')}-1} \left(\frac{1}{2}|\lambda_k(b)^{(b')}|^2 - (\lambda_k(b)^{(b')})^T (\eta_{k+1} + \lambda_k(b')^{(b')})\right)\right) + \mathbb{1}(\tau^{(b')} = \infty)\epsilon \quad (75)$$

and in particular

$$L(b)^{(b)} = \mathbb{1}(\tau^{(b)} < \infty) \exp\left(-\sum_{k=0}^{\tau^{(b)}-1} \left(\frac{1}{2}|\lambda_k(b)^{(b)}|^2 + (\lambda_k(b)^{(b)})^T \eta_{k+1}\right)\right) + \mathbb{1}(\tau^{(b)} = \infty)\epsilon. \quad (76)$$

4.3.4 Linearly parametrized exponential tilting for stopped sequences

Let $\tilde{U}_1, \tilde{X}, \tilde{U}(b), \tilde{L}(b), \tilde{\Psi}(b)$, $b \in B = \mathbb{R}^d$, and \tilde{F} , be as some $\mathbb{Q}_1, X, \mathbb{Q}(b), L(b), \Psi(b)$, $b \in A = B$, and F , in the ECM setting in Section 4.1. Considering additionally some Λ as in Section 4.3.2, let $\mathbb{Q}(b)$ and $L(b)$, $b \in A$, be the corresponding linearly parametrized IS distributions and densities. Note that we now have from (61), for $U(b) = \mathbb{1}(\tau < \infty) \sum_{k=0}^{\tau-1} \tilde{\Psi}(\lambda_k(b))$ and $H = -\mathbb{1}(\tau < \infty) \sum_{k=0}^{\tau-1} (\tilde{X}(\eta_{k+1}))^T \Lambda_k$, that

$$L(b) = \mathbb{1}(\tau < \infty) \exp(U(b) + Hb) + \epsilon \mathbb{1}(\tau = \infty). \quad (77)$$

We shall call the above IS setting linearly parametrized exponentially tilted stopped sequences (LETS) setting. Its special case in which $\tilde{U}_1 = \mathcal{N}(0, I_d)$ and $\tilde{X} = \text{id}_{\mathbb{R}^d}$ shall be called linearly parametrized exponentially tilted Gaussian stopped sequences (LETGS) setting. Note that the LETGS setting is a special case of parametrized IS for Gaussian stopped sequences from Section 4.3.3. In the LETGS setting $H = -\mathbb{1}(\tau < \infty) \sum_{k=0}^{\tau-1} (\eta_{k+1})^T \Lambda_k$ and for $G := \mathbb{1}(\tau < \infty) \frac{1}{2} \sum_{k=0}^{\tau-1} \Lambda_k^T \Lambda_k$ we have $U(b) = b^T G b$, so that

$$L(b) = \mathbb{1}(\tau < \infty) \exp(b^T G b + Hb) + \mathbb{1}(\tau = \infty)\epsilon. \quad (78)$$

Furthermore, for $G^{(b')} = \mathbb{1}(\tau^{(b')} < \infty) \frac{1}{2} \sum_{k=0}^{\tau^{(b')}-1} (\Lambda_k^{(b')})^T \Lambda_k^{(b')}$ and

$$H^{(b')} = -\mathbb{1}(\tau^{(b')} < \infty) \sum_{k=0}^{\tau^{(b')}-1} (\eta_{k+1} + \Lambda_k^{(b')} b')^T \Lambda_k^{(b')}, \quad (79)$$

formula (75) reduces to

$$L(b)^{(b')} = \mathbb{1}(\tau^{(b')} < \infty) \exp(b'^T G^{(b')} b' + H^{(b')} b') + \mathbb{1}(\tau^{(b')} = \infty)\epsilon. \quad (80)$$

Remark 46. Note that in the LETGS setting, on $\tau < \infty$ we have

$$\inf_{b \in \mathbb{R}^d} \ln(L(b)) \geq \sum_{k=0}^{\tau-1} \inf_{y \in \mathbb{R}^d} \left(\frac{1}{2} |y|^2 - \eta_{k+1}^T y \right) = -\frac{1}{2} \sum_{k=1}^{\tau-1} |\eta_k|^2 \in \mathbb{R}. \quad (81)$$

Remark 47. In our numerical experiments performing IS for computing expectations of functionals of Euler scheme in the LETGS setting, the simulation times were roughly proportional to replicates of τ under $\mathbb{Q}(b)$. Thus, on several occasions in this work when dealing with the LETGS setting, we shall consider theoretical cost $C = \tau$.

Returning to the general LETS setting, consider its special case in which Λ is a sequence of constant matrices and $\tau = n \in \mathbb{N}$ is deterministic. Then, for the above constructed family $\{\mathbb{Q}(b) : b \in A\}$, the family of distributions $\mathcal{W}' = \{\mathbb{Q}'(b) : b \in A\}$ on $\tilde{\mathcal{C}}^n$ such that $\mathbb{Q}'(b)(\tilde{\eta}_n[C]) = \mathbb{Q}(b)(C)$, $C \in \mathcal{F}_n$, $b \in A$, is the exponentially tilted family corresponding to $\mathbb{Q}'_1 := \tilde{\mathbb{U}}_1^n$ and $X'(\omega) := \sum_{i=1}^n \Lambda_{i-1}^T \tilde{X}(\omega_i)$, $\omega = (\omega_i)_{i=1}^n \in \tilde{E}^n$, as in Section 4.1. Note that when $\tau = 1$ and $\Lambda_0 = I_d$, then such \mathcal{W}' coincides with the exponentially tilted family $\{\tilde{\mathbb{U}}(b) : b \in A\}$ used for its construction. Furthermore, if $\tilde{X} \sim \mathcal{N}(0, I_d)$ under $\tilde{\mathbb{U}}_1$, then $X' \sim \mathcal{N}(0, \sum_{i=0}^{n-1} \Lambda_i^T \Lambda_i)$ under \mathbb{Q}'_1 .

4.4 IS for functionals of Brownian motion up to a stopping time

Let us now briefly discuss IS for computing expectations of functionals of Brownian motion. Let $B = (B_t)_{t \geq 0}$ be the coordinate process on the Wiener space $\mathcal{C}([0, \infty), \mathbb{R}^d)$, whose measurable space let us denote \mathcal{W} , and let $(\tilde{\mathcal{F}}_t)_{t \geq 0}$ be the natural filtration of B . Let $\tilde{\mathbb{U}}$ be the unique probability on \mathcal{W} for which B is a d -dimensional Brownian motion (see Chapter 1, Section 3 in [43]). For a probability \mathbb{S} on \mathcal{W} and a stopping time τ for $(\tilde{\mathcal{F}}_t)_{t \geq 0}$, we denote $\mathbb{S}|_t = \mathbb{S}|_{\tilde{\mathcal{F}}_t}$. From Girsanov theorem, if $(\tilde{\lambda}_t)_{t \geq 0}$ is a predictable locally square-integrable \mathbb{R}^d -valued process on \mathcal{W} for which

$$\tilde{\gamma}_t = \exp \left(\int_0^t \tilde{\lambda}_s^T dB_s - \frac{1}{2} \int_0^t |\tilde{\lambda}_s|^2 ds \right), \quad t \geq 0, \quad (82)$$

is a martingale under $\tilde{\mathbb{U}}$ (for which e.g. Novikov's condition suffices), then from Kolmogorov's extension theorem there exists a unique measure $\tilde{\mathbb{V}}$ on \mathcal{W} such that $\frac{d\tilde{\mathbb{U}}|_t}{d\tilde{\mathbb{V}}|_t} = \tilde{\gamma}_t$, $t \geq 0$. Furthermore,

$$\tilde{B}_t = B_t - \int_0^t \tilde{\lambda}_s ds, \quad t \geq 0, \quad (83)$$

is a Brownian motion under $\tilde{\mathbb{V}}$. From Proposition 1.3, Chapter 8 in [43], for a stopping time $\tilde{\tau}$ for $(\tilde{\mathcal{F}}_t)_{t \geq 0}$, we have $\mathbb{1}(\tilde{\tau} < \infty) \tilde{\gamma}_{\tilde{\tau}} = \left(\frac{d\tilde{\mathbb{V}}|_{\tilde{\tau}}}{d\tilde{\mathbb{U}}|_{\tilde{\tau}}} \right)_{\tilde{\tau} < \infty}$ and thus $\tilde{L} = \mathbb{1}(\tilde{\tau} < \infty) \frac{1}{\tilde{\gamma}_{\tilde{\tau}}} = \left(\frac{d\tilde{\mathbb{U}}|_{\tilde{\tau}}}{d\tilde{\mathbb{V}}|_{\tilde{\tau}}} \right)_{\tilde{\tau} < \infty}$, similarly as in the discrete case. Thus, if for some $\tilde{Z} \in L^1(\tilde{\mathbb{U}}|_{\tilde{\tau}})$ we have $\tilde{\mathbb{U}}$ and $\tilde{\mathbb{V}}$ a.s. that $\tilde{\tau} = \infty$ implies $\tilde{Z} = 0$,

then $\tilde{\mathbb{U}}_{|\tilde{\tau}} \sim_{Z \neq 0} \mathbb{V}_{|\tilde{\tau}}$ and we can perform IS for computing $\mathbb{E}_{\tilde{\mathbb{U}}}(\tilde{Z})$ analogously as in the discrete case. For adaptive IS we can use e.g. linear parametrization $\tilde{\lambda}_t(b) = \tilde{\Lambda}_t b$, $b \in A := \mathbb{R}^l$ of tilting processes for some $\mathbb{R}^{d \times l}$ -valued predictable process $(\tilde{\Lambda}_t)_{t \geq 0}$ with locally square integrable coordinates.

Due to the fact that sequence $(B_{k+1} - B_k)_{k \in \mathbb{N}}$ has i.i.d. $\sim \mathcal{N}(0, I_d)$ coordinates under $\tilde{\mathbb{U}}$, under appropriate identifications the LETGS setting can be viewed a special discrete case of IS for Brownian motion with linear parametrization of tilting processes as above. In the further sections we focus mainly on the discrete case, both for simplicity and due to its important applications in numerical methods. However, many of our reasonings can be generalized to the Brownian case.

4.5 IS for diffusions and Euler scheme

Using notations as for IS for Brownian motion in the previous section, let us consider functions $\mu : \mathcal{S}(\mathbb{R}^m) \rightarrow \mathcal{S}(\mathbb{R}^m)$ and $\sigma : \mathcal{S}(\mathbb{R}^m) \rightarrow \mathcal{S}(\mathbb{R}^{m \times d})$ such that there exists a unique strong solution Y of the SDE

$$dY_t = \mu(Y_t)dt + \sigma(Y_t)dB_t, \quad Y_0 = x_0. \quad (84)$$

Such Y is called a diffusion, μ a drift, and σ a diffusion matrix. For $\tilde{\tau}$ being some stopping time for $(\tilde{\mathcal{F}}_t)_{t \geq 0}$ (like e.g. some hitting time of Y an appropriate set) and $\tilde{Z} \in L^1(\tilde{\mathbb{U}}_{|\tilde{\tau}})$, one can be interested in estimating

$$\tilde{\phi}(x_0) = \mathbb{E}_{\tilde{\mathbb{U}}}(\tilde{Z}). \quad (85)$$

A popular way of discretizing Y , especially in many dimensions, is by using an Euler scheme $X = (X_k)_{k \in \mathbb{N}}$ with a time step $h \in \mathbb{R}_+$, which, for some η_1, η_2, \dots , i.i.d. $\sim \mathcal{N}(0, I_m)$ and some starting point $x_0 \in \mathbb{R}^m$, fulfills $X_0 = x_0$ and

$$X_{k+1} = X_k + h\mu(X_k) + \sqrt{h}\sigma(X_k)\eta_{k+1}, \quad k \in \mathbb{N}. \quad (86)$$

Let further η_i , $i \in \mathbb{N}_+$, be as in Section 4.3.1 for $\tilde{\mathbb{U}}_1 = \mathcal{N}(0, I_d)$, so that X as above is an Euler scheme under \mathbb{U} as in that section. As discussed further on, in some cases, for sufficiently small h , for appropriate stopping time τ for $(\mathcal{F}_n)_{n \geq 0}$ and appropriate $Z \in L^1(\mathbb{U}_{|\tau})$, $\tilde{\phi}(x_0)$ can be approximated well using

$$\phi(x_0) = \mathbb{E}_{\mathbb{U}}(Z). \quad (87)$$

For some function $r : \mathcal{S}(\mathbb{R}^m) \rightarrow \mathcal{S}(\mathbb{R}^d)$, called an IS drift, let us consider a tilting process $\lambda_k = \sqrt{h}r(X_k)$, $k \in \mathbb{N}$. Then, for

$$\tilde{\mu} = \mu + \sigma r \quad (88)$$

and $\dot{\eta}_k$, $k \in \mathbb{N}$, as in Section 4.3.3, we have

$$X_{k+1} = X_k + h\tilde{\mu}(X_k) + \sqrt{h}\sigma(X_k)\dot{\eta}_{k+1}, \quad k \in \mathbb{N}, \quad (89)$$

so that from Theorem 45, X is an Euler scheme under \mathbb{V} with drift $\tilde{\mu}$. As discussed in Section 4.3.3, the distribution of X under \mathbb{V} is the same as of

$\widehat{X} := X(\pi^{-1})$ under \mathbb{U} . Since $\dot{\eta}_i = \eta_i \pi$, we have $\dot{\eta}_i \pi^{-1} = \eta_i$, $i \in \mathbb{N}_+$, so that \widehat{X} satisfies $\widehat{X}_0 = x_0$ and

$$\widehat{X}_{k+1} = \widehat{X}_k + h\tilde{\mu}(\widehat{X}_k) + \sqrt{h}\sigma(\widehat{X}_k)\eta_{k+1}, \quad k \in \mathbb{N}, \quad (90)$$

i.e. it is also an Euler scheme with drift $\tilde{\mu}$, but this time under \mathbb{U} .

For some nonempty $A \in \mathcal{B}(\mathbb{R}^l)$, let us consider some parametrization $r : A \rightarrow \{f : \mathbb{R}^m \rightarrow \mathbb{R}^d\}$ of IS drifts, such that $(b, x) \rightarrow r(b)(x)$ is measurable from $\mathcal{S}(A) \otimes \mathcal{S}(\mathbb{R}^m)$ to $\mathcal{S}(\mathbb{R}^d)$, and let $\tilde{\mu}(b) = \mu + \sigma r(b)$, $b \in A$. Consider a parametrization $\lambda : A \rightarrow \mathcal{J}$ of tilting processes such that

$$\lambda(b) = (\lambda_k(b))_{k \in \mathbb{N}} = (\sqrt{h}r(b)(X_k))_{k \in \mathbb{N}}, \quad b \in A. \quad (91)$$

Note that Condition 42 holds for such parametrization. Note also that, using notation (72), from (90) we have

$$X_{k+1}^{(b)} = X_k^{(b)} + h\tilde{\mu}(b)(X_k^{(b)}) + \sqrt{h}\sigma(X_k^{(b)})\eta_{k+1}, \quad k \in \mathbb{N}. \quad (92)$$

Let us now describe the linear case of the above parametrization, leading to IS in the special case of the LETGS setting. We take $A = \mathbb{R}^l$ and for some functions $\tilde{r}_i : \mathcal{S}(\mathbb{R}^m) \rightarrow \mathcal{S}(\mathbb{R}^d)$, $i = 1, \dots, l$, called IS basis functions, we set

$$r(b)(x) = \sum_{i=1}^l b_i \tilde{r}_i(x), \quad b \in \mathbb{R}^l, \quad x \in \mathbb{R}^m. \quad (93)$$

Let $\Theta : \mathbb{R}^m \rightarrow \mathbb{R}^{d \times l}$ be such that for $i = 1, \dots, l$ and $j = 1, \dots, d$

$$\Theta_{j,i}(x) = \sqrt{h}(\tilde{r}_i)_j(x). \quad (94)$$

Then, a process Λ leading to $\lambda(b)$ given by (65) and such that (91) holds, can be defined as

$$\Lambda_k = \Theta(X_k), \quad k \in \mathbb{N}. \quad (95)$$

We have $\Lambda_k^{(b)} = \Theta(X_k^{(b)})$, $b \in A$.

An example of a stopping time τ for $(\mathcal{F}_k)_{k \geq 0}$ is an exit time of X of some $D \in \mathcal{B}(\mathbb{R}^m)$, that is $\tau = \inf\{k \in \mathbb{N} : X_k \notin D\}$, for which we have $\tau^{(b)} = \inf\{k \in \mathbb{N} : X_k^{(b)} \notin D\}$, $b \in A$.

Theorem 48. *Let us consider some linear parametrization of IS drifts as above. Let τ be the exit time of X of $D \in \mathcal{B}(\mathbb{R}^m)$ such that $x_0 \in D$, let $B \subset \mathbb{R}^l$ be nonempty, and let there exist $v \in \mathbb{R}^m$, $v \neq 0$, such that*

$$M_1 := \sup_{x, y \in D} |v^T(x - y)| < \infty \quad (96)$$

and

$$M_2 := \sup_{x \in D, b \in B} |v^T \tilde{\mu}(b)(x)| < \infty. \quad (97)$$

For some $i \in \{1, \dots, d\}$, let there exist $\delta_i \in \mathbb{R}_+$ and $\delta_j \in [0, \infty)$, $j \in \{1, \dots, d\}$, $j \neq i$, such that $|(v^T \sigma(x))_i| \geq \delta_i$ and $|(v^T \sigma(x))_j| \leq \delta_j$, $j \neq i$, $x \in D$. Let $M = M_1 + hM_2$ and consider the following random conditions on \mathcal{C} for $k \in \mathbb{N}_+$

$$q_k(\omega) = (|\eta_{k,i}(\omega)\delta_i| > \frac{M}{\sqrt{h}} + |\sum_{j \in \{1, \dots, d\}, j \neq i} \eta_{k,j}(\omega)\delta_j|), \quad \omega \in E. \quad (98)$$

Then, a random variable $\hat{\tau}$ on \mathcal{C} such that $\hat{\tau}(\omega) = \inf\{k \in \mathbb{N}_+ : q_k(\omega)\}$, $\omega \in E$, fulfills $\tau^{(b)} \leq \hat{\tau}$, $b \in B$. Under \mathbb{U} , the variable $\hat{\tau}$ has geometric distribution with parameter $q = \mathbb{U}(q_1)$, that is $\mathbb{U}(\hat{\tau} = k) = q(1 - q)^{k-1}$, $k \in \mathbb{N}_+$.

Proof. From (92), for each $k \in \mathbb{N}_+$

$$\begin{aligned} (X_k^{(b)} \notin D) \wedge (X_{k-1}^{(b)} \in D) &\equiv (\sqrt{h}\sigma(X_{k-1}^{(b)})\eta_k \notin D - X_{k-1}^{(b)} - h\tilde{\mu}(b)(X_{k-1}^{(b)})) \wedge (X_{k-1}^{(b)} \in D) \\ &\Leftrightarrow (\sqrt{h}v^T\sigma(X_{k-1}^{(b)})\eta_k \notin v^T(D - X_{k-1}^{(b)} - h\tilde{\mu}(b)(X_{k-1}^{(b)}))) \wedge (X_{k-1}^{(b)} \in D) \\ &\Leftrightarrow (\sqrt{h}|v^T\sigma(X_{k-1}^{(b)})\eta_k| > M) \wedge (X_{k-1}^{(b)} \in D) \\ &\Leftrightarrow q_k \wedge (X_{k-1}^{(b)} \in D). \end{aligned} \tag{99}$$

Thus, $q_k \Rightarrow (X_{k-1}^{(b)} \notin D \vee X_k^{(b)} \notin D) \Rightarrow \tau^{(b)} \leq k$. For $\omega \in E$ such that $\hat{\tau}(\omega) < \infty$ it holds $q_{\hat{\tau}(\omega)}$, and thus $\tau^{(b)}(\omega) \leq \hat{\tau}(\omega)$. \square

Remark 49. Note that if for each $b \in A$ the assumptions of Theorem 48 hold for $B = \{b\}$, then from $\tau^{(b)}$ having the same distribution under \mathbb{U} as τ under $\mathbb{V}(b)$, we receive that τ has all finite moments under $\mathbb{V}(b)$, $b \in A$, and in particular Condition 44 holds.

We say that a matrix- or vector-valued function f is uniformly bounded on some set B if for some arbitrary vector or matrix norm $\|\cdot\|$ we have $\sup_{x \in B} \|f(x)\| < \infty$.

Remark 50. Note that (96) holds for D bounded. Furthermore, if for some $v \in \mathbb{R}^m$, $v^T\mu$, $v^T\sigma$, and Θ are uniformly bounded on D (which holds e.g. when they are continuous on \mathbb{R}^l and D is bounded) then (97) holds for each bounded B .

4.6 Some special functionals of diffusions and Euler scheme, and zero-variance IS for diffusions

To provide an intuition when variance of IS estimators of expectations of functionals of Euler scheme as in the previous section can be small, let us briefly describe a situation when their diffusion counterparts have zero variance. See Section 4 in [20] for details. For $\tilde{\tau}$ being the hitting time of Y a boundary of an appropriate set D , as well as for appropriate $g : \mathbb{R}^m \rightarrow [0, \infty)$ and $\beta : \mathbb{R}^m \rightarrow \mathbb{R}$, consider

$$\tilde{Z} = \mathbb{1}(\tilde{\tau} < \infty)g(Y_{\tilde{\tau}})\exp\left(\int_0^{\tilde{\tau}} \beta(Y_s)ds\right). \tag{100}$$

If there exists $u : \mathbb{R}^m \rightarrow \mathbb{R}$, such that for $Lu = \text{tr}(\sigma\sigma^T)\Delta u + \mu^T\nabla u + \beta u$ we have

$$Lu(x) = 0, \quad x \in D, \tag{101}$$

and

$$u(x) = g(x), \quad x \in \partial D, \tag{102}$$

then, from Feynmann-Kac theorem, $\tilde{\phi}(x_0) = u(x_0)$. Under certain assumptions it can be proved (see Theorem 4 in [20]) that for r equal to

$$r^* := \frac{\sigma^T\nabla u}{u} = \sigma^T\nabla(\ln(u)), \tag{103}$$

for IS for Brownian motion as above with $\tilde{\lambda}_t = r^*(Y_t)$, we have $\tilde{Z}\tilde{L} = \tilde{\phi}(x_0)$, $\tilde{\mathbb{V}}$ a.s., i.e. the unbiased IS estimator for the diffusion case has zero variance. Furthermore, from (83)

$$dY_t = (\mu + \sigma r^*)(Y_t)dt + \sigma(Y_t)d\tilde{B}_t, \quad Y_0 = x_0. \quad (104)$$

For τ being the exit time of X of some set E , a possible Euler scheme counterpart of (100) is

$$Z = \mathbb{1}(\tau < \infty)g(X_\tau) \exp\left(\sum_{k=0}^{\tau-1} h\beta(X_k)\right). \quad (105)$$

In [23] it was proved that under appropriate assumptions (holding e.g. in our numerical experiments discussed in Section 4.9) for such Z we have

$$\lim_{h \rightarrow 0} \phi(x_0) = \tilde{\phi}(x_0) \quad (106)$$

for $E = D$, and that the rate of convergence in (106) can be increased by taking as E appropriately shifted set D . Further on for Z as in (105) we assume that $E = D$, but one can easily modify the below reasonings to consider the shifted case instead. It seems intuitive that for some such Z , for r close to r^* , and for small h , we can receive low variance of the Euler scheme IS estimator ZL . This intuition shall be confirmed in our numerical experiments in Section 9. Note that for such Z , $Z^{(b)} = \mathbb{1}(\tau^{(b)} < \infty)g((X^{(b)})_{\tau^{(b)}}) \exp(\sum_{k=0}^{\tau^{(b)}-1} h\beta(X_k^{(b)}))$.

4.7 Some special functionals of diffusions and Euler scheme

We shall now provide several examples of quantities of form $\tilde{\phi}(x_0) = \mathbb{E}_{\tilde{\mathbb{U}}}(\tilde{Z})$ for some \tilde{Z} as in (100). As discussed in Section 1.2, these quantities can be of interest among others in molecular dynamics, and their Euler scheme counterparts $\phi(x_0) = \mathbb{E}_{\mathbb{U}}(Z)$ for Z as in (105) are estimated in our numerical experiments in Section 9.

Taking $\beta(x) = -p$, for some $p \in \mathbb{R}_+$ and $g(x) = 1$, we receive $\tilde{Z} = \exp(-p\tilde{\tau})\mathbb{1}(\tilde{\tau} < \infty)$ and $Z = \exp(-ph\tau)\mathbb{1}(\tau < \infty)$. The quantities $\widetilde{\text{mgf}}(x_0) := \tilde{\phi}(x_0)$ and $\text{mgf}(x_0) := \phi(x_0)$ for this case are called moment-generating functions (MGFs) of $\tilde{\tau}$ and $h\tau$ respectively.

For the next example let us assume that

$$\mathbb{U}(\tau < \infty) = \tilde{\mathbb{U}}(\tilde{\tau} < \infty) = 1 \quad (107)$$

and let $D' = \mathbb{R}^m \setminus D = A \cup B$ for two closed disjoint sets A and B from $\mathcal{S}(\mathbb{R}^m)$. Let us consider $a \in [0, \infty)$, called an added constant, let $\beta(x) = 0$, $g(x) = a + 1$, $x \in B$, and $g(x) = a$, $x \in A$. We receive $\tilde{Z} = \mathbb{1}(\tilde{\tau} < \infty)(a + \mathbb{1}(Y_{\tilde{\tau}} \in B))$ and $\tilde{\phi}(x_0)$ equal to $\tilde{q}_{AB,a}(x_0) := a + \tilde{\mathbb{U}}(Y_{\tilde{\tau}} \in B)$, which we shall call translated committor. For the added constant $a = 0$, we denote $\tilde{q}_{AB,a}(x_0)$ simply as $\tilde{q}_{AB}(x_0)$ and call it committor. For the Euler scheme case we have analogous definitions but with omitted tilde sign.

For some $T \in \mathbb{R}_+$, let us now consider $\tilde{Z} = \mathbb{1}(\tilde{\tau} < T) + a$, $\tilde{p}_{T,a}(x_0) = \mathbb{E}_{\tilde{\mathbb{U}}}(\tilde{Z})$, and $\tilde{p}_T(x_0) = \tilde{p}_{T,0}(x_0)$, while for the Euler scheme case $Z = \mathbb{1}(h\tau < T) + a$, $p_{T,a}(x_0) = \mathbb{E}_{\mathbb{U}}(Z)$, and $p_T(x_0) = p_{T,0}(x_0)$. Note that this time \tilde{Z} and Z do not have forms as in (100) and (105). However, they can be rewritten in such forms for some time-extended processes as discussed in the below remark.

Remark 51. Consider $Y' = (Y_t, t)_{t \in [0, \infty)}$ and $X' = (X_k, kh)_{k \in \mathbb{N}}$, which are a diffusion and a corresponding Euler scheme in the definition of which, in place of m , x_0 , μ and σ , one uses $m' = m + 1$, $x'_0 = (x_0, 0)$, as well as $\mu' : \mathbb{R}^{m'} \rightarrow \mathbb{R}^{m'}$ and $\sigma' : \mathbb{R}^{m'} \rightarrow \mathbb{R}^{m' \times d}$ such that for each $x \in \mathbb{R}^m$ and $t \in \mathbb{R}$ we have $\mu'(x, t) = (\mu(x), 1)$, $\sigma'_{i,j}(x, t) = \sigma_{i,j}(x)$, $i \leq m$, and $\sigma'_{m',j}(x, t) = 0$, $j \in \{1, \dots, d\}$. Let $\hat{D} = D \times [0, T)$, and let $\tilde{\tau}'$ and τ' be the exit times of Y' and X' of \hat{D} respectively. Then, for $g(x, t) = \mathbb{1}(x \in D' \wedge t < T) + a$, $x \in \mathbb{R}^m$, $t \in \mathbb{R}$, $\tilde{Z}' = g(Y'_{\tilde{\tau}'})$, and $Z' = g(X'_{\tau'})$, we have $\tilde{Z} = \tilde{Z}'$ and $Z = Z'$.

Note that for τ' as in the above Remark we have $\tau' = \tau \wedge \lceil \frac{T}{h} \rceil$. Such τ' is the stopping time which we consider by default for IS for computing $p_{T,a}(x_0)$. A possible alternative would be to use τ , which, as discussed in Remark 41, would lead to not lower variance and mean cost for the cost variables being equal to the respective stopping times.

Let $\hat{\psi}_a$ be an unbiased estimator of ψ_a equal to $q_{AB,a}(x_0)$ or $p_{T,a}(x_0)$, i.e. $\mathbb{E}(\hat{\psi}_a) = \psi_a$. Then, the translated estimator $\hat{\psi}_{a,0} = \hat{\psi}_a - a$ is an unbiased estimator of ψ_0 equal to $q_{AB}(x_0)$ or $p_T(x_0)$ respectively, and $\text{Var}(\hat{\psi}_{a,0}) = \text{Var}(\hat{\psi}_a)$. The reason for considering such translated estimators of ψ_0 for nonzero added constants a is that using these estimators in adaptive IS procedures in our numerical experiments discussed in Section 9 led to lower variances and inefficiency constants than for $a = 0$.

Note that we have $q_{AB}(x_0) + q_{BA}(x_0) = 1$ and similarly for the diffusion case, so that if \hat{q} is an unbiased estimator of one of the quantities $q_{AB}(x_0)$ or $q_{BA}(x_0)$, then $1 - \hat{q}$ is such an estimator of the other quantity with the same variance and inefficiency constant. Therefore, given estimators \hat{q}_{AB} of $q_{BA}(x_0)$ and \hat{q}_{BA} of $q_{AB}(x_0)$, it seems reasonable to compute both quantities as above using the estimator leading to a lower inefficiency constant.

4.8 Diffusion in a potential

An important special case of a diffusion Y is a one in an appropriate differentiable potential $V : \mathbb{R}^m \mapsto \mathbb{R}$ for some temperature $\epsilon \in \mathbb{R}_+$, fulfilling

$$dY_t = -\nabla V(Y_t)dt + \sqrt{2\epsilon}dB_t, \quad Y_0 = x_0. \quad (108)$$

In such case, the IS drift (103) leading to zero-variance IS estimator and distribution $\tilde{\mathbb{V}}$ is

$$r^* = \sqrt{2\epsilon}(\nabla \ln(u)). \quad (109)$$

Let $F = -\epsilon \ln(u)$, $C_0 \in \mathbb{R}$, and let us define an optimally-tilted potential

$$V^* = V + 2F + C_0. \quad (110)$$

Then, (104) can be rewritten as

$$dY_t = -\nabla V^*(Y_t)dt + \sqrt{2\epsilon}d\tilde{B}_t, \quad Y_0 = x_0. \quad (111)$$

Thus, under $\tilde{\mathbb{V}}$, Y is a diffusion in potential V^* .

4.9 The special cases considered in our numerical experiments

Consider a three-well potential

$$V(x) = \frac{1}{200}(0.5x^6 - 15x^4 + 119x^2 + 28x + 50). \quad (112)$$

Such V restricted to $D := (a_1, a_2) = (-3.5, 3.5)$ is shown in Figure 1. For a




Figure 1: Three-well potential given by (112) on D .

temperature $\epsilon = 0.5$, consider a diffusion Y in such potential starting at some $x_0 \in D$. Let $\tilde{\tau}$ be the hitting time of Y of the boundary of D . Let $A = (-\infty, a_1)$, $B = (a_2, \infty)$, and let $\tilde{q}_{1,a}(x_0) = \tilde{q}_{AB,a}(x_0)$ and $\tilde{q}_{2,a}(x_0) = \tilde{q}_{BA,a}(x_0)$, which for $a = 0$ are denoted simply as $\tilde{q}_1(x_0)$ and $\tilde{q}_2(x_0)$. Let us also consider MGF $\widetilde{\text{mgf}}(x_0)$ for $p = \tilde{p} := 0.1$. We computed approximations of such $\tilde{q}_i(x)$ and $\widetilde{\text{mgf}}(x)$ in function of x using finite difference discretizations of PDEs given by (101) and (102). The results are shown in Figures 2a and 2b. The approximations of optimally tilted potentials (110) corresponding to zero-variance importance sampling distributions for MGF and committors $\tilde{q}_{i,a}$ for $a = 0$ and $a = \tilde{a} := 0.05$, $i = 1, 2$, are shown in figures 3a and 3b.

In our experiments we considered an Euler scheme X with a time step $h = 0.01$ corresponding to the above diffusion Y starting at $x_0 = 0$. We focused on estimating $\text{mgf}(x_0)$ for $p = \tilde{p}$, $q_i(x_0)$ for $i = 1, 2$, and $p_T(x_0)$ for $T = 10$. For some $M \in \mathbb{N}_+$, $\tilde{a}_1 = -3.6$, $\tilde{a}_2 = 3.6$, $\sigma = \frac{\tilde{a}_2 - \tilde{a}_1}{M-1}$, and $p_i = \tilde{a}_1 + (i-1)\sigma$, $i \in \{1, \dots, M\}$, consider Gaussian functions

$$\tilde{r}_i(x) = \frac{1}{\sqrt{\epsilon}} \exp\left(-\frac{(x - p_i)^2}{\sigma^2}\right), \quad i = 1, \dots, M. \quad (113)$$

In our IS experiments we used linear parametrization of IS drifts as in Section 4.5, using as IS basis functions the above functions for $M = 10$. For estimating $p_T(x_0)$, considering time-extended Euler scheme as in Remark 51, we additionally performed experiments using $2M$ time-dependent IS basis functions

$$\hat{r}_i(x, t) = \tilde{r}_i(x), \quad \hat{r}_{M+i}(x, t) = t^p \tilde{r}_i(x), \quad i = 1, \dots, M, \quad (114)$$

for different $p \in \mathbb{N}_+$, and for $M = 5$ and $M = 10$. See sections 6.1 and 9 for further details on our numerical experiments. Note that since the above \tilde{r}_i are




Figure 2: Committors and MGF for a diffusion in a three-well potential as in the main text.

continuous and D is bounded, from remarks 49, 50, and Theorem 48, in which one can take $v = \delta_1 = 1$, it follows that Condition 44 holds when estimating the MGF and committors as above. Using further the fact that $\mathbb{E}_{\tilde{U}}(\tilde{\tau}) < \infty$ (which follows e.g. from Lemma 7.4 in [30]), (107) holds.




Figure 3: Optimally tilted potentials as in (110) for MGF and committors. Fqi and $Fqia$ are the functions F for the i th committor with $a = 0$ and $a = \tilde{a}$ respectively. The constant C_0 for the i th committor for $i \in \{1, 2\}$ was chosen so that the tilted potential is equal to the original potential in point a_i .

5 The minimized functions and their estimators

In this section we define and investigate properties of various functions and their estimators, minimization of which will be discussed in the further sections.

5.1 The minimized functions

For some nonempty set A , consider a family of distributions as in Section 3.4 for which Condition 17 holds. Assuming Condition 23 and that

$$\mathbb{E}_{\mathbb{Q}_1}((Z \ln(L(b)))_-) < \infty, \quad (115)$$

we define cross-entropy as follows

$$ce(b) = \mathbb{E}_{\mathbb{Q}_1}(Z \ln(L(b))) \quad (116)$$

(see discussion in Section 1.2 regarding its name). If (115) holds for each $b \in A$, this defines a function $ce : A \rightarrow \mathbb{R} \cup \{\infty\}$. Let us discuss how $ce(b)$ is related to a certain f -divergence of $\mathbb{Q}(b)$ from the zero-variance IS distribution, assuming that such distribution exists. For some convex function $f : [0, \infty) \rightarrow \mathbb{R}$, an f -divergence $d(\mathbb{P}_1, \mathbb{P}_2)$ of a probability \mathbb{P}_2 from another one \mathbb{P}_1 such that $\mathbb{P}_2 \ll \mathbb{P}_1$ is given by formula

$$d(\mathbb{P}_1, \mathbb{P}_2) = \mathbb{E}_{\mathbb{P}_1}\left(f\left(\frac{d\mathbb{P}_2}{d\mathbb{P}_1}\right)\right). \quad (117)$$

Such f -divergence is also known as Csiszár f -divergence or Ali-Silvey distance [42, 2, 38]. From Jensen's inequality we have $d(\mathbb{P}_1, \mathbb{P}_2) \geq f(1)$, and if f is strictly convex then equality in this inequality holds only if $\mathbb{P}_1 = \mathbb{P}_2$. For example, for the strictly convex function $f(x) = x \ln(x)$ (which we assume to be zero for $x = 0$), $d(\mathbb{P}_1, \mathbb{P}_2)$ is called Kullback-Leibler divergence or cross-entropy distance (of \mathbb{P}_2 from \mathbb{P}_1), while for $f(x) = (x^2 - 1)$, $d(\mathbb{P}_1, \mathbb{P}_2)$ is called Pearson divergence.

For d denoting the cross-entropy distance, let us assume Condition 12, so that the zero-variance IS distribution \mathbb{Q}^* exists, $\mathbb{Q}^* \ll \mathbb{Q}(b)$, $b \in A$, and

$$\begin{aligned} d(\mathbb{Q}(b), \mathbb{Q}^*) &= \mathbb{E}_{\mathbb{Q}(b)}\left(\frac{d\mathbb{Q}^*}{d\mathbb{Q}(b)} \ln\left(\frac{d\mathbb{Q}^*}{d\mathbb{Q}(b)}\right)\right) \\ &= \mathbb{E}_{\mathbb{Q}^*}\left(\ln\left(\frac{d\mathbb{Q}^*}{d\mathbb{Q}(b)}\right)\right) \\ &= \mathbb{E}_{\mathbb{Q}_1}\left(\frac{Z}{\alpha}\left(\ln\left(\frac{d\mathbb{Q}^*}{d\mathbb{Q}_1}\right) + \ln(L(b))\right)\right), \end{aligned} \quad (118)$$

where in the last equality we used (28) and

$$\left(\frac{d\mathbb{Q}^*}{d\mathbb{Q}(b)}\right)_{Z \neq 0} = \frac{d\mathbb{Q}^*}{d\mathbb{Q}_1} L(b). \quad (119)$$

Assuming that $Z \geq 0$, we have

$$\mathbb{E}_{\mathbb{Q}_1}\left(Z \ln\left(\frac{d\mathbb{Q}^*}{d\mathbb{Q}_1}\right)\right) = \mathbb{E}_{\mathbb{Q}_1}(Z \ln(Z)) - \alpha \ln(\alpha). \quad (120)$$

From $x \ln(x) \geq -e^{-1}$ we have $\mathbb{E}_{\mathbb{Q}_1}(Z \ln(Z)) \geq -e^{-1}$. Assuming further that

$$\mathbb{E}_{\mathbb{Q}_1}(Z \ln(Z)) < \infty, \quad (121)$$

we receive from (118) that

$$d(\mathbb{Q}(b), \mathbb{Q}^*) = \alpha^{-1}(\mathbb{E}_{\mathbb{Q}_1}(Z \ln(Z)) - \alpha \ln(\alpha) + ce(b)). \quad (122)$$

If (122) holds as above for each $b \in A$, then $b \rightarrow ce(b)$ and $b \rightarrow d(\mathbb{Q}(b), \mathbb{Q}^*)$ are positively linearly equivalent (see Section 1.1). Note that from discussion leading to formula (122) and from $d(\mathbb{Q}(b), \mathbb{Q}^*) \geq 0$, a sufficient assumption for (115) to hold for each $b \in A$ is that we have $Z \geq 0$ and (121).

Let us define mean square (function) of the IS estimator as

$$\text{msq}(b) = \mathbb{E}_{\mathbb{Q}(b)}((ZL(b))^2) = \mathbb{E}_{\mathbb{Q}_1}(Z^2 L(b)), \quad b \in A, \quad (123)$$

and such variance (function) as

$$\text{var}(b) = \text{msq}(b) - \alpha^2, \quad b \in A. \quad (124)$$

Assuming that Condition 11 holds, for \mathbb{Q}^* denoting the optimal-variance IS distribution as in Section 3.2 and d denoting Pearson divergence, from (119) and (33) we have for $b \in A$

$$\begin{aligned} d(\mathbb{Q}(b), \mathbb{Q}^*) &= \mathbb{E}_{\mathbb{Q}(b)}\left(\left(\frac{|Z|}{\mathbb{E}_{\mathbb{Q}_1}(|Z|)} L(b)\right)^2 - 1\right) \\ &= \frac{1}{(\mathbb{E}_{\mathbb{Q}_1}(|Z|))^2} (\text{msq}(b) - (\mathbb{E}_{\mathbb{Q}_1}(|Z|))^2) \\ &= \frac{1}{(\mathbb{E}_{\mathbb{Q}_1}(|Z|))^2} (\text{var}(b) + \alpha^2 - (\mathbb{E}_{\mathbb{Q}_1}(|Z|))^2). \end{aligned} \quad (125)$$

Thus, in such case $b \in A \rightarrow d(\mathbb{Q}(b), \mathbb{Q}^*)$ is positively linearly equivalent to msq and var .

Let C be some $[0, \infty]$ -valued random variable on \mathcal{S}_1 , called (theoretical) cost variable. Let $c(b) = \mathbb{E}_{\mathbb{Q}(b)}(C)$ be such mean cost (function) under $\mathbb{Q}(b)$, $b \in A$.

Condition 52. For each $b \in A$, it does not hold $c(b) = \infty$ and $\text{var}(b) = 0$ or $c(b) = 0$ and $\text{var}(b) = \infty$.

Assuming Condition 52, let us define inefficiency constant (function) as

$$\text{ic}(b) = c(b) \text{var}(b), \quad b \in A. \quad (126)$$

5.2 Estimators of the minimized functions

Consider a family of distributions as in Section 3.4 and let us assume that conditions 17 and 18 hold. Let $f : \mathcal{S}(A) \rightarrow \mathcal{S}(\overline{\mathbb{R}})$ and for some $p \in \mathbb{N}_+$, consider

$$\widehat{\text{est}}_n : \mathcal{S}(A)^2 \otimes \mathcal{S}_1^n \rightarrow \mathcal{S}(\overline{\mathbb{R}}), \quad n \in \mathbb{N}_p, \quad (127)$$

called (parameter-dependent) estimators of f . In all this work, for $b' \in A$, we denote $\mathbb{Q}' = \mathbb{Q}(b')$ and $L' = L(b')$. We say that $\widehat{\text{est}}_n(b', b)$ as above is an unbiased estimator of $f(b)$ under $(\mathbb{Q}')^n$ if

$$f(b) = \mathbb{E}_{(\mathbb{Q}')^n}(\widehat{\text{est}}_n(b', b)). \quad (128)$$

Let us assume a following condition.

Condition 53. $\kappa_1, \kappa_2, \dots$, are i.i.d. $\sim \mathbb{Q}'$ and $\tilde{\kappa}_n = (\kappa_i)_{i=1}^n$, $n \in \mathbb{N}_+$.

Estimators $\widehat{\text{est}}_n(b', b)$, $n \in \mathbb{N}_p$, are called strongly consistent for $f(b)$ under $(\mathbb{Q}')^n$ if a.s.

$$\lim_{n \rightarrow \infty} \widehat{\text{est}}_n(b', b)(\tilde{\kappa}_n) = f(b). \quad (129)$$

For a function Y on Ω_1 (like e.g. Z or L), we define such functions Y_1, \dots, Y_n on Ω_1^n by formula

$$Y_i(\omega) = Y(\omega_i), \quad \omega = (\omega_i)_{i=1}^n \in \Omega_1^n, \quad (130)$$

and whenever Y takes values in some linear space we denote

$$\overline{(Y)}_n = \frac{1}{n} \sum_{i=1}^n Y_i. \quad (131)$$

For cross-entropy, under assumptions as in the previous section, we have

$$\text{ce}(b) = \mathbb{E}_{\mathbb{Q}'}(L' Z \ln(L(b))), \quad (132)$$

so for $n \in \mathbb{N}_+$, from Theorem 1, its unbiased strongly consistent estimators under $(\mathbb{Q}')^n$ are

$$\widehat{\text{ce}}_n(b', b) = \overline{(L' Z \ln(L(b)))}_n = \frac{1}{n} \sum_{i=1}^n L'_i Z_i \ln(L_i(b)). \quad (133)$$

For mean square we have

$$\text{msq}(b) = \mathbb{E}_{\mathbb{Q}'}(Z^2 L' L(b)), \quad (134)$$

so for $n \in \mathbb{N}_+$, its unbiased strongly consistent estimators under $(\mathbb{Q}')^n$ are

$$\widehat{\text{msq}}_n(b', b) = \overline{(Z^2 L' L(b))}_n. \quad (135)$$

As discussed in the Introduction, the above mean square estimators and estimators negatively linearly equivalent to the above cross-entropy estimators have been considered before in the literature. We shall now proceed to define some new estimators. If $\mathbb{Q} \ll \mathbb{Q}'$, then for variance we have for $n \in \mathbb{N}_2$,

$$\begin{aligned} \text{var}(b) &= \text{Var}_{\mathbb{Q}(b)}(ZL(b)) \\ &= \mathbb{E}_{(\mathbb{Q}(b))^n} \left(\frac{1}{n(n-1)} \sum_{i < j \in \{1, \dots, n\}} (Z_i L_i(b) - Z_j L_j(b))^2 \right) \\ &= \frac{1}{n(n-1)} \mathbb{E}_{(\mathbb{Q}')^n} \left(\sum_{i < j \in \{1, \dots, n\}} \left(\frac{d\mathbb{Q}(b)}{d\mathbb{Q}'} \right)_i \left(\frac{d\mathbb{Q}(b)}{d\mathbb{Q}'} \right)_j (Z_i L_i(b) - Z_j L_j(b))^2 \right). \end{aligned} \quad (136)$$

Let us assume conditions 22 and 23. Then, $\frac{d\mathbb{Q}(b)}{d\mathbb{Q}'} = \frac{L'}{L(b)}$ and from (136) we have the following unbiased estimators of $\text{var}(b)$ under $(\mathbb{Q}')^n$ for $n \in \mathbb{N}_2$

$$\begin{aligned} \widehat{\text{var}}_n(b', b) &= \frac{1}{n(n-1)} \sum_{i < j \in \{1, \dots, n\}} \frac{L'_i L'_j}{L_i(b) L_j(b)} (Z_i L_i(b) - Z_j L_j(b))^2 \\ &= \frac{1}{n(n-1)} \left(\sum_{i=1}^n \left(Z_i^2 L'_i L_i(b) \sum_{j \in \{1, \dots, n\}, j \neq i} \frac{L'_j}{L_j(b)} \right) - \sum_{i < j \in \{1, \dots, n\}} 2Z_i Z_j L'_i L'_j \right) \\ &= \frac{n}{n-1} \left(\widehat{\text{msq}}_n(b', b) \overline{\left(\frac{L'}{L(b)} \right)}_n - \overline{(ZL')}_n \right). \end{aligned} \quad (137)$$

Thus, $b \rightarrow \widehat{\text{var}}_n(b', b)$ is positively linearly equivalent to the following estimator of mean square

$$\widehat{\text{msq}}_n(b', b) = \widehat{\text{msq}}_n(b', b) \overline{\left(\frac{L'}{L(b)} \right)}_n, \quad (138)$$

which is in general not unbiased and which can be considered also for $n = 1$. From the facts that from SLLN, a.s.

$$\lim_{n \rightarrow \infty} \overline{(ZL')}_n(\tilde{\kappa}_n) = \mathbb{E}_{\mathbb{Q}'}(ZL') = \alpha \quad (139)$$

and

$$\lim_{n \rightarrow \infty} \overline{\left(\frac{L'}{L(b)} \right)}_n(\tilde{\kappa}_n) = \mathbb{E}_{\mathbb{Q}'} \left(\frac{L'}{L(b)} \right) = 1, \quad (140)$$

estimators $\widehat{\text{msq}}_n$ and $\widehat{\text{var}}_n(b', b)$ are strongly consistent. We shall sometimes call $\widehat{\text{msq}}_n$ the first or well-known mean square estimator and $\widehat{\text{msq}}_n$ the second or new one.

Strongly consistent and unbiased estimators of mean cost $c(b)$ under under $(\mathbb{Q}')^n$ are

$$\hat{c}_n(b', b) = \overline{\left(\frac{L'}{L(b)} C \right)}_n. \quad (141)$$

Let us assume Condition 52. Then, strongly consistent estimators of $\text{ic}(b)$ under $(\mathbb{Q}')^n$ are for $n \in \mathbb{N}_2$,

$$\widehat{\text{ic}}_n(b', b) = \widehat{c}_n(b', b) \cdot \widehat{\text{var}}_n(b', b) \quad (142)$$

(by which we mean that $\widehat{\text{ic}}_n(b', b) = 0$ whenever we multiply infinity by zero on the right and similarly in (144)), which are in general not unbiased. For each $n \in \mathbb{N}_3$, defining helper unbiased estimators of variance for $k = 1, \dots, n$

$$\widehat{\text{var}}_{n,k}(b', b) = \frac{1}{(n-1)(n-2)} \left(\sum_{i < j \in \{1, \dots, n\} \setminus \{k\}} \frac{L'_i L'_j}{L_i(b) L_j(b)} (Z_i L_i(b) - Z_j L_j(b))^2 \right), \quad (143)$$

we have a following unbiased estimator of $\text{ic}(b)$ under $(\mathbb{Q}')^n$

$$\widehat{\text{ic}}_2n(b', b) = \frac{1}{n} \sum_{k=1}^n \left(\frac{L'}{L(b)} C \right)_k \widehat{\text{var}}_{n,k}(b', b). \quad (144)$$

The fact that $\widehat{\text{ic}}_2n(b', b)$ are strongly consistent follows from an easy generalization of SLLN for U-statistics [26], analogous as in Theorem 1 for the ordinary SLLN.

5.3 Cross-entropy and its estimator for ECM

Consider the ECM setting as in Section 4.1. We have

$$\widehat{c}_n(b', b) = \overline{(ZL'(\Psi(b) - bX))}_n = \Psi(b) \overline{(ZL')}_n - b^T \overline{(ZL'X)}_n. \quad (145)$$

Let us assume Condition 32. Then,

$$\nabla_b \widehat{c}_n(b', b) = \nabla \Psi(b) \overline{(ZL')}_n - \overline{(ZL'X)}_n \quad (146)$$

and

$$\nabla_b^2 \widehat{c}_n(b', b) = \nabla^2 \Psi(b) \overline{(ZL')}_n. \quad (147)$$

Let us further assume Condition 31, so that $\nabla^2 \Psi$ is positive definite. Then, from (147), $b \rightarrow \widehat{c}_n(b', b)(\omega)$ has positive definite Hessian and thus it is strictly convex only for $\omega \in \Omega_1^n$ and $b' \in A$ such that

$$\overline{(ZL')}_n(\omega) > 0. \quad (148)$$

Furthermore, $b_n^* \in A$ is the unique minimum point of $b \rightarrow \widehat{c}_n(b', b)(\omega)$ only if (148) holds and

$$\nabla_b \widehat{c}_n(b', b_n^*)(\omega) = 0 \quad (149)$$

(where by $\nabla_b \widehat{c}_n(b', b_n^*)(\omega)$ we mean $\nabla_b(\widehat{c}_n(b', b)(\omega))_{b=b_n^*}$). Assuming (148), from (146), (149) holds only if

$$\mu(b_n^*) = \nabla \Psi(b_n^*) = \frac{\overline{(ZL'X)}_n(\omega)}{\overline{(ZL')}_n(\omega)}, \quad (150)$$

or from Theorem 36 only if $\frac{\overline{(ZL'X)}_n(\omega)}{\overline{(ZL')}_n(\omega)} \in \mu[A]$ and

$$b_n^* = \mu^{-1} \left(\frac{\overline{(ZL'X)}_n(\omega)}{\overline{(ZL')}_n(\omega)} \right). \quad (151)$$

Let us assume that

$$\mathbb{E}_{\mathbb{Q}_1}(|ZX_i|) < \infty, \quad i = 1, \dots, l. \quad (152)$$

Due to X having all mixed moments, from Hölder inequality, (152) holds e.g. when $\mathbb{E}_{\mathbb{Q}_1}(|Z|^p) < \infty$ for some $p > 1$. For cross-entropy we then have

$$\text{ce}(b) = \alpha \Psi(b) - b^T \mathbb{E}_{\mathbb{Q}_1}(ZX), \quad b \in A. \quad (153)$$

Thus, analogously as for the cross-entropy estimator above, ce has positive definite Hessian only if $\alpha > 0$, and ce has a unique minimum point only if $\alpha > 0$ and

$$\frac{\mathbb{E}_{\mathbb{Q}_1}(ZX)}{\alpha} \in \mu[A], \quad (154)$$

in which case such point is

$$b^* = \mu^{-1} \left(\frac{\mathbb{E}_{\mathbb{Q}_1}(ZX)}{\alpha} \right). \quad (155)$$

Remark 54. *Note that we can receive analogous conditions as above for cross-entropy and its estimator to have negative definite Hessians or have unique maximum points by replacing Z by $-Z$ in the above conditions. The formulas for the maximum points remain the same as for minimum points above. With some exceptions, in the further sections we shall focus on minimization of cross-entropy and its estimators and will be interested in checking the conditions as in the main text above. However, one can analogously perform their maximization, or jointly optimization if we consider alternatives of the above conditions.*

5.4 Some conditions in the LETS setting

Consider the LETS setting as in Section 4.3.4. For each real matrix-valued process $Y = (Y_k)_{k \in \mathbb{N}}$ on \mathcal{C} and $B \in \mathcal{E}$, let us define

$$\|Y\|_{\tau, B, \infty} = \text{ess sup}_{\cup} (\mathbb{1}_B \mathbb{1}(0 < \tau < \infty) \max(\|Y_0\|_{\infty}, \dots, \|Y_{\tau-1}\|_{\infty})), \quad (156)$$

which for $B = \Omega_1$ is denoted simply as $\|Y\|_{\tau, \infty}$. Let S be an $\overline{\mathbb{R}}$ -valued random variable on \mathcal{S}_1 . Further on in this work we will often assume the following conditions.

Condition 55. *It holds*

$$R := \|\Lambda\|_{\tau, S \neq 0, \infty} < \infty. \quad (157)$$

Condition 56. *For some $s \in \mathbb{N}_+$,*

$$\mathbb{1}(S \neq 0)\tau \leq s. \quad (158)$$

Note that conditions 55 or 56 hold for each possible random variable S as above only if they hold for some S such that $S(\omega) \neq 0$, $\omega \in \Omega_1$, that is only if

$$\|\Lambda\|_{\tau, \infty} < \infty \quad (159)$$

for Condition 55, or

$$\tau \leq s \in \mathbb{N}_+ \quad (160)$$

for Condition 56.

Remark 57. Note that Condition 56 implies Condition 43 for $Z = S$, while (160) implies Condition 44.

For each real matrix-valued function f on \mathbb{R}^m and $B \subset \mathbb{R}^m$, let us denote $\|f\|_{B,\infty} = \sup_{x \in B} \|f\|_\infty$. If τ is the exit time of an Euler scheme X of D , with $X_0 \in D$, and Λ is as in (95), then we have $\|\Lambda\|_{\tau,\infty} \leq \|\Theta\|_{D,\infty}$. In particular, if

$$\|\Theta\|_{D,\infty} < \infty, \quad (161)$$

then we have (159). Note that from (94), (161) is equivalent to $\|\tilde{\tau}_i\|_{D,\infty} < \infty$, $i = 1, \dots, l$. In particular, it holds in our numerical experiments as discussed in Section 4.9 both when using time-independent and time-dependent IS basis functions (where in the time-dependent case by $\tilde{\tau}_i$ we mean $\hat{\tau}_i$ as in Section 4.9 and we consider D equal to \hat{D} as in Remark 51). Let us discuss how one can enforce (160) if it is initially not fulfilled, as is the case of committors or MGF in our numerical experiments. Analogous reasonings as below can be applied also to more general stopped sequences or processes than in the LETS setting. Instead of τ and Z we can consider their terminated versions $\tau_s = \tau \wedge s$ and $Z_s = \mathbb{1}(\tau \leq s)Z + z_s \mathbb{1}(\tau > s)$ for some $z_s \in \mathbb{R}$, so that we focus on computing $\alpha_s = \mathbb{E}_\mathbb{U}(Z_s) = \mathbb{E}_\mathbb{U}(\mathbb{1}(\tau \leq s)Z) + z_s \mathbb{U}(\tau > s)$ rather than $\alpha = \mathbb{E}_\mathbb{U}(Z)$. If $\mathbb{U}(\tau = \infty) = 0$, or $\mathbb{U}(Z \neq 0, \tau = \infty) = 0$ and $\lim_{s \rightarrow \infty} z_s = 0$, then \mathbb{U} a.s. $Z_s \rightarrow Z$, so that assuming further that $\limsup_{s \rightarrow \infty} |z_s| < \infty$, from $|Z_s| \leq |Z| + |z_s|$ and Lebesgue dominated convergence, $\lim_{s \rightarrow \infty} \alpha_s = \alpha$. Thus, in such case, for sufficiently large s we will make arbitrarily small absolute error when approximating α by α_s . Let us provide some upper bounds on this error. If $\text{ess sup}_\mathbb{U}(|Z - z_s| \mathbb{1}(\tau > s)) \leq M_s \in [0, \infty)$, then

$$|\alpha - \alpha_s| = |\mathbb{E}_\mathbb{U}((Z - z_s) \mathbb{1}(\tau > s))| \leq M_s \mathbb{U}(\tau > s). \quad (162)$$

For the MGF example from Section 4.7 we can take $z_s = M_s = \frac{1}{2} \exp(-ph(s + 1))$, while for translated committors we can choose $z_s = a + \frac{1}{2}$ and $M_s = \frac{1}{2}$. The quantity $\mathbb{U}(\tau > s)$ can be estimated using IS from the same simulations as used to estimate α_s or in a separate IS MC procedure. Alternatively, if we have $\tau \leq \hat{\tau}$ for some random variable $\hat{\tau}$ with known distribution, we can use inequality $\mathbb{U}(\tau > s) \leq \mathbb{U}(\hat{\tau} > s)$ to bound the right side of (162) from above. For instance, if $\hat{\tau}$ has geometric distribution with parameter q as in Theorem 48, then we have $\mathbb{U}(\hat{\tau} > s) = (1 - q)^s$ and thus $|\alpha - \alpha_s| \leq M_s (1 - q)^s$.

5.5 Some conditions in the LETGS setting

Let us discuss some conditions and random conditions in the LETGS setting, which, as we shall discuss in the further sections, turn out to be necessary for existence of unique minimum points of cross-entropy and mean square or their estimators in this setting. Let Z be an \mathbb{R} -valued \mathcal{S}_1 -measurable random variable (where $\mathcal{S}_1 = (E, \mathcal{F}_\tau)$).

Definition 58. For some $b \in A = \mathbb{R}^l$, we define a random condition A_b on \mathcal{S}_1 as follows

$$A_b \equiv (Z \neq 0, 0 < \tau < \infty, \text{ and there exists } k \in \mathbb{N}, k < \tau, \text{ such that } \lambda_k(b) \neq 0). \quad (163)$$

Lemma 59. *If A_b does not hold and $Z \neq 0$, then for each $a \in \mathbb{R}^l$ and $t \in \mathbb{R}$*

$$L(a + tb) = L(a). \quad (164)$$

Proof. From (74), when $\tau = \infty$, then both sides of (164) are equal to ϵ . If A_b does not hold, $Z \neq 0$, and $0 < \tau < \infty$, then for each $0 \leq k < \tau$ we have $\lambda_k(b) = 0$, and thus for each $a \in \mathbb{R}^l$ and $t \in \mathbb{R}$, $\lambda_k(a + tb) = \lambda_k(a) + t\lambda_k(b) = \lambda_k(a)$, so that (164) also follows from (74). \square

Lemma 60. *For some $n \in \mathbb{N}_+$, the following random conditions on \mathcal{S}_1^n are equivalent.*

1. *For each $b \in \mathbb{R}^l$, $b \neq 0$, there exists $i \in \{1, \dots, n\}$ such that $(A_b)_i$ holds (where we use notation as in (130)).*
2. *For some (equivalently, for each) random variable K on Ω which is positive on $Z \neq 0$, $(\mathbb{1}(Z \neq 0)GK)_n$ is positive definite.*
3. *It holds $N := \sum_{i=1}^n \mathbb{1}(Z_i \neq 0, \tau_i < \infty)\tau_i > 0$. Let a matrix $B \in \mathbb{R}^{(dN) \times l}$ be such that for each $i \in \{1, \dots, n\}$ such that $0 < \tau_i < \infty$ and $Z_i \neq 0$, for each $k \in \{0, \dots, \tau_i - 1\}$ and $j \in \{1, \dots, d\}$ the $\sum_{v=1}^{i-1} \mathbb{1}(Z_v \neq 0, \tau_v < \infty)\tau_v + kd + j$ th row of B is equal to the j th row of $(\Lambda_k)_i$. Then, the columns of B are linearly independent.*

Proof. The fact that the second point above is a random condition follows from Sylvester's criterion. Equivalence of the first two conditions follows from the fact that for each $b \in \mathbb{R}^l$

$$b^T (\overline{K\mathbb{1}(Z \neq 0)G})_n b = \frac{1}{2n} \sum_{i=1}^n (\mathbb{1}(0 < \tau < \infty, Z \neq 0)K \sum_{k=0}^{\tau-1} |\lambda_k(b)|^2)_i, \quad (165)$$

and equivalence of the first and last condition is obvious. \square

Definition 61. *We define r_n to be one of the equivalent random conditions in Lemma 60.*

Condition 62. *The three below conditions are equivalent and we assume the condition under consideration to be one of them.*

1. *For each $b \in \mathbb{R}^l$, $b \neq 0$, we have $\mathbb{Q}_1(A_b) > 0$.*
2. *For each $b \in \mathbb{R}^l$, from $\mathbb{Q}_1(A_b) = 0$ it follows that $b = 0$.*
3. *Let $\tilde{\Lambda}_j = ((\Lambda_{k,i,j})_{i=1}^d)_{k \in \mathbb{N}} \in \mathcal{J}$, $j = 1, \dots, l$ (see Definition 38). Let \sim be a relation of equivalence on \mathcal{J} such that for $\psi_1, \psi_2 \in \mathcal{J}$, $\psi_1 \sim \psi_2$, only if \mathbb{Q}_1 a.s. when $0 < \tau < \infty$ and $Z \neq 0$ then $\psi_{1,i} = \psi_{2,i}$, $i = 0, \dots, \tau - 1$. Then, the equivalence classes $[\tilde{\Lambda}_1]_{\sim}, \dots, [\tilde{\Lambda}_l]_{\sim}$ are linearly independent in the linear space \mathcal{J}/\sim of equivalence classes of \sim , defined in a standard way.*

Remark 63. *Note that for a probability $\mathbb{S} \sim_{\tau < \infty} \mathbb{Q}_1$ we have $\mathbb{S}(A_b) > 0$ only if $\mathbb{Q}_1(A_b) > 0$, so that Condition 62 holds only if it holds for such \mathbb{S} in place of \mathbb{Q}_1 . Note also that $\mathbb{Q}_1(A_b) > 0$ only if for some $l \in \mathbb{N}_+$ and $k \in \mathbb{N}$, $k < l$, we have $\mathbb{Q}_1(Z \neq 0, \tau = l, \lambda_k(b) \neq 0) > 0$.*

Lemma 64. *Let for some probability $\mathbb{S} \sim_{\tau < \infty} \mathbb{Q}_1$, random variable K on \mathcal{S}_1 be \mathbb{S} a.s. positive on $Z \neq 0$, and let $\mathbb{1}(Z \neq 0)KG$ have \mathbb{S} -integrable entries. Then, $\mathbb{E}_{\mathbb{S}}(\mathbb{1}(Z \neq 0)KG)$ is positive definite only if Condition 62 holds.*

Proof. For each $b \in \mathbb{R}^l$, $b \neq 0$,

$$b^T \mathbb{E}_{\mathbb{S}}(\mathbb{1}(Z \neq 0)KG)b = \frac{1}{2} \mathbb{E}_{\mathbb{S}}(\mathbb{1}(Z \neq 0, 0 < \tau < \infty)K \sum_{k=0}^{\tau-1} |\lambda_k(b)|^2) \quad (166)$$

is greater than zero only if $\mathbb{S}(A_b) > 0$, so that from Remark 63 we receive the thesis. \square

Let $\text{Sym}_n(\mathbb{R})$ denote the subset of $\mathbb{R}^{n \times n}$ consisting of symmetric matrices, and let $m_n : \text{Sym}_n(\mathbb{R}) \rightarrow \mathbb{R}$ be such that for $A \in \text{Sym}_n(\mathbb{R})$, $m_n(A)$ is equal to the lowest eigenvalue of A , or equivalently

$$m_n(A) = \inf_{x \in \mathbb{R}^n, |x|=1} x^T A x. \quad (167)$$

Lemma 65. *m_n is Lipschitz from $(\text{Sym}_n(\mathbb{R}), \|\cdot\|_{\infty})$ to $(\mathbb{R}, |\cdot|)$ with Lipschitz constant 1.*

Proof. For $A, B \in \text{Sym}_n(\mathbb{R})$ and $x \in \mathbb{R}^n$, $|x| = 1$, we have $x^T A x = x^T B x + x^T(A - B)x$, so that

$$x^T B x - \|A - B\|_{\infty} \leq x^T A x \leq x^T B x + \|A - B\|_{\infty} \quad (168)$$

and thus

$$m_n(B) - \|A - B\|_{\infty} \leq m_n(A) \leq m_n(B) + \|A - B\|_{\infty} \quad (169)$$

and

$$|m_n(B) - m_n(A)| \leq \|A - B\|_{\infty}. \quad (170)$$

\square

Lemma 66. *If the entries of $M_n \in \text{Sym}_l(\mathbb{R})$, $n \in \mathbb{N}_+$, converge to the entries of a positive definite symmetric matrix $M \in \mathbb{R}^{l \times l}$, then for sufficiently large n , M_n is positive definite.*

Proof. This follows from the fact that $A \in \text{Sym}_l(\mathbb{R})$ is positive definite only if $m_l(A) > 0$, and from Lemma 65, $\lim_{n \rightarrow \infty} m_l(M_n) = m_l(M)$. \square

Theorem 67. *If Condition 62 holds, then under Condition 53, a.s. for sufficiently large n , $r_n(\tilde{\kappa}_n)$ holds for r_n as in Definition 61. In particular, $\lim_{n \rightarrow \infty} \mathbb{P}(r_n(\tilde{\kappa}_n)) = 1$.*

Proof. Let $K = \exp(-\max_{i,j=1,\dots,d} |G_{i,j}|)$. Then, $K > 0$ and entries of the matrix $\mathbb{1}(Z \neq 0)KG$ are bounded and thus \mathbb{Q}' -integrable. Thus, from Lemma 64 for $\mathbb{S} = \mathbb{Q}'$, $\mathbb{E}_{\mathbb{Q}'}(\mathbb{1}(Z \neq 0)KG)$ is positive definite. Let $A_n = (\mathbb{1}(Z \neq 0)KG)_n(\tilde{\kappa}_n)$. From SLLN, a.s.

$$\lim_{n \rightarrow \infty} A_n = \mathbb{E}_{\mathbb{Q}'}(\mathbb{1}(Z \neq 0)KG). \quad (171)$$

Thus, from Lemma 66, a.s. A_n is positive definite for sufficiently large n and the thesis follows from the second point of Lemma 60. \square

5.6 Discussion of Condition 62 in the Euler scheme case

In this section we reformulate Condition 62 in the special case of IS for Euler scheme with linear parametrization of IS drifts as in Section 4.5, and we provide some sufficient assumptions for this condition to hold.

Let us define a measure ν on $\mathcal{S}(\mathbb{R}^m)$ to be such that for each $B \in \mathcal{B}(\mathbb{R}^m)$

$$\begin{aligned} \nu(B) &= \mathbb{E}_{\mathbb{U}}(\mathbb{1}(Z \neq 0, 0 < \tau < \infty) \sum_{k=0}^{\tau-1} \mathbb{1}(X_k \in B)) \\ &= \sum_{l \in \mathbb{N}_+} \sum_{k=0}^{l-1} \mathbb{U}(Z \neq 0, \tau = l, X_k \in B) \\ &= \sum_{k=0}^{\infty} \mathbb{U}(Z \neq 0, k < \tau < \infty, X_k \in B). \end{aligned} \quad (172)$$

Remark 68. From the second line of (172), Remark 63, and (95), $\mathbb{Q}_1(A_b) = \mathbb{U}(A_b) = 0$ is equivalent to

$$\nu(\{\Theta b \neq 0\}) = 0 \quad (173)$$

(where $\{\Theta b \neq 0\} = \{x \in \mathbb{R}^m : \Theta(x)b \neq 0\}$).

Remark 69. Let $\tilde{\Theta}_i : \mathbb{R}^m \rightarrow \mathbb{R}^m$ be the i th column of Θ and $[\tilde{\Theta}_i]_{\approx}$ be the class of equivalence of $\tilde{\Theta}_i$ with respect to the relation \approx of equality ν a.e. on the set \mathcal{K} of measurable functions from $\mathcal{S}(\mathbb{R}^m)$ to $\mathcal{S}(\mathbb{R}^m)$. Then, Condition 62 is equivalent to $[\tilde{\Theta}_i]_{\approx}$, $i = 1, \dots, l$, being linearly independent in the linear space \mathcal{K}/\approx defined in a standard way.

Let us assume that $m = n + 1$ for some $n \in \mathbb{N}_+$. Consider a following condition on some functions \tilde{r}_i , which can be used as IS basis functions as in Section 4.5.

Condition 70. For some $m_1, m_2 \in \mathbb{N}_+$, functions $g_{1,i} : \mathbb{R} \rightarrow \mathbb{R}$, $i = 1, \dots, m_1$, and $g_{2,i} : \mathbb{R}^n \rightarrow \mathbb{R}^d$, $i = 1, \dots, m_2$, are such that for $K_1 = \{kh : k \in \{1, \dots, m_1\}\}$, $g_{1,i}|_{K_1}$, $i = 1, \dots, m_1$, are linearly independent and for each $i \in \{1, \dots, m_1\}$, for some open set $K_{2,i} \subset \mathbb{R}^n$, $g_{2,j}|_{K_{2,i}}$, $j = 1, \dots, m_2$, are continuous and linearly independent. Furthermore, for $M = m_1 m_2$, functions $\tilde{r}_i : \mathbb{R}^m \rightarrow \mathbb{R}^d$, $i = 1, \dots, M$, are such that, denoting $\pi(i, j) = m_2(i - 1) + j$, for each $x \in \mathbb{R}^n$ and $t \in \mathbb{R}$ we have $\tilde{r}_{\pi(i,j)}(x, t) = g_{1,i}(t)g_{2,j}(x)$, $i = 1, \dots, m_1$, $j = 1, \dots, m_2$.

Remark 71. As functions $g_{1,i}$ as in the above condition one can take for example polynomials $g_{1,i} = t^{i-1}$, $i = 1, \dots, m_1$. For $m_1 = 2$ one can also use $g_{1,1}(t) = 1$ and $g_{1,2} = t^p$ for some $p \in \mathbb{N}_+$. For $n = 1$ and arbitrary nonempty open sets $K_{2,i} \subset \mathbb{R}$, $i = 1, \dots, m_2$, as functions $g_{2,i}$ in the above condition one can take e.g. polynomials as above or Gaussian functions $g_{2,i}(x) = a_i \exp(\frac{(x-p_i)^2}{s})$ with $a_i \neq 0$, $s > 0$, and p_i different (for the proof of linear independence of the latter see [1]). In particular, for such $K_{2,i}$, Condition 70 holds for \tilde{r}_i equal to \hat{r}_i as in (114) or equal to \hat{r}_i such that $\hat{r}_i(x, t) = \tilde{r}_i(x)$ for \tilde{r}_i as in (113), where in the first case $m_1 = 2$, in the second $m_1 = 1$, and in both cases $n = 1$ and m_2 is equal to M as in Section 4.9.

Let λ denote the Lebesgue measure on \mathbb{R}^n .

Theorem 72. *If Condition 70 holds and*

$$\lambda \otimes \delta_{ih} \ll_{K_{2,i} \times \{ih\}} \nu, \quad i = 1, \dots, m_1, \quad (174)$$

then Condition 62 holds.

Proof. Let $b \in \mathbb{R}^M$ be such that $\mathbb{U}(A_b) = 0$. Then, from Remark 68, $\nu(\{\Theta b \neq 0\}) = 0$. Therefore, denoting for $i = 1, \dots, m_1$ and $k = 1, \dots, m_2$

$$a_{i,k} = \sum_{j=1}^{m_1} b_{\pi(j,k)} g_{1,j}(ih), \quad (175)$$

from (174) we have for $i = 1, \dots, m_1$, λ a.s. on $K_{2,i}$ that $\sum_{k=1}^{m_2} a_{i,k} g_{2,k} = 0$. Thus, for $i = 1, \dots, m_1$, from continuity and linear independence of $g_{2,k}|_{K_{2,i}}$, $k = 1, \dots, m_2$, we have $a_{i,k} = 0$, $k = 1, \dots, m_2$. Therefore, from (175), for $k = 1, \dots, m_2$, $\sum_{j=1}^{m_1} b_{\pi(j,k)} g_{1,j}|_{K_1} = 0$, so that from linear independence of $g_{1,j}|_{K_1}$, $j = 1, \dots, m_1$, we have $b = 0$. \square

Let us assume a following condition.

Condition 73. $\sigma_{m,i} = 0$, $i = 1, \dots, d$, $\mu_m = 1$, $x_{0,m} = 0$, and $\tilde{\sigma} : \mathbb{R}^m \rightarrow \mathbb{R}^{n \times d}$ is such that $\tilde{\sigma}_{i,j} = \sigma_{i,j}$, $i = 1, \dots, n$, $j = 1, \dots, d$.

Note that it now holds for $\tilde{X}_k = (X_{k,i})_{i=0}^n$, $k \in \mathbb{N}$, that $X_k = (\tilde{X}_k, kh)$, $k \in \mathbb{N}$. For $x \in \mathbb{R}^n$ and $k \in \mathbb{N}$ for which $\tilde{\sigma}(x, kh)$ has linearly independent rows, let $Q_k(x) = (h\tilde{\sigma}(x, kh)\tilde{\sigma}(x, kh)^T)^{-1}$, and for $y \in \mathbb{R}^n$, let

$$\rho_k(x, y) = \frac{\sqrt{\det(Q_k(x))}}{(2\pi)^{\frac{m}{2}}} \exp((y - x - h\mu(x))^T Q_k(y - x - h\mu(x))), \quad (176)$$

so that $\rho_k(x, \cdot)$ is the transition density of $\tilde{X} = (\tilde{X}_k)_{k \in \mathbb{N}}$ in x from its k th to its $k + 1$ st state.

Theorem 74. *Let $k \in \mathbb{N}_+$ and sets $B_1, B_2, \dots, B_k, C \in \mathcal{B}(\mathbb{R}^n)$ have positive Lebesgue measure. Let \mathbb{U} a.s. the fact that $\tilde{X}_i \in B_i$, $i = 1, \dots, k$ and $\tilde{X}_{k+1} \in C$ imply that $Z \neq 0$ and $k < \tau < \infty$. Let further $\tilde{\sigma}(x)$ have independent rows for each $x \in \{x_0\} \cup \bigcup_{i=1}^k B_i$. Then,*

$$\lambda \otimes \delta_{ih} \ll_{B_i \times \{ih\}} \nu, \quad i = 1, \dots, k. \quad (177)$$

Proof. It follows from the fact that for each $j \in \{1, \dots, k\}$, for each $D \subset \mathcal{B}(B_j)$ such that $\lambda(D) > 0$, for $\tilde{D} = \prod_{i=1}^{j-1} B_i \times D \times \prod_{i=j+1}^k B_i \times C$ we have

$$\begin{aligned} 0 &< \int_{\tilde{D}} \prod_{i=0}^k \rho_i(x_i, x_{i+1}) dx_1 dx_2 \dots dx_{k+1} \\ &= \mathbb{U}((\tilde{X}_i)_{i=1}^{k+1} \in \tilde{D}) \\ &= \mathbb{U}(Z \neq 0, k < \tau < \infty, (\tilde{X}_i)_{i=1}^{k+1} \in \tilde{D}) \\ &\leq \mathbb{U}(Z \neq 0, k < \tau < \infty, X_j \in D) \leq \nu(D). \end{aligned} \quad (178)$$

\square

Remark 75. Let us consider the problems of estimating MGF $\text{mgf}(x_0)$, translated committor $q_{AB,a}(x_0)$, and translated exit probability before a given time $p_{T,a}(x_0)$ as in Section 4.7. As x_0 , μ , and σ fulfilling the above Condition 73 let us consider x'_0 , μ' , and σ' as in Remark 51, and as an Euler scheme X above let us consider the time-extended process X' as in that remark. Note that X from Section 4.7 is now equal to \tilde{X} as above. Let $k \in \mathbb{N}_+$. Then for D and Z corresponding to the above expectations as in Section 4.7, assuming that $C \in \mathcal{B}(B)$ in case of estimation of a committor, or $C \in \mathcal{B}(D')$ for MGF or exit probability, and additionally $T > h(k+1)$ in case of the exit probability, for each $B \in \mathcal{B}(D)$ we have \cup a.s. that $\tilde{X}_i \in B$, $i = 1, \dots, k$ and $\tilde{X}_{k+1} \in C$ implies $Z \neq 0$ and $k < \tau < \infty$ (where for the exit probability rather than τ we mean τ' as in Remark 51). This holds also for Z and τ replaced by their terminated versions Z_s and τ_s for $s \in \mathbb{N}_+$, $s > k$, as in Section 5.4.

From remarks 71 and 75 and theorems 72 and 74 it easily follows that Condition 62 holds in all cases considered in our numerical experiments as in Section 4.9 if for the case of exit probability before a given time we assume that $T > h(m_2 + 1)$ for m_2 depending on basis functions used as in Remark 71. Furthermore, this condition also holds in such terminated cases as in Section 5.4 for each $s \in \mathbb{N}_+$, $s > m_2$.

5.7 Cross-entropy and its estimator in the LETGS setting

Consider the LETGS setting, assuming Condition 43. From (78),

$$\begin{aligned} \widehat{\text{ce}}_n(b', b) &= \overline{ZL'(b^T G b + H b + \mathbb{1}(\tau = \infty) \ln(\epsilon))}_n \\ &= b^T \overline{ZL'G}_n b + \overline{ZL'H}_n b + \ln(\epsilon) \overline{ZL'\mathbb{1}(\tau = \infty)}_n, \end{aligned} \quad (179)$$

so that

$$\nabla_b \widehat{\text{ce}}_n(b', b) = 2 \overline{ZL'G}_n b + \overline{ZL'H}_n. \quad (180)$$

Thus, $b \rightarrow \widehat{\text{ce}}_n(b', b)(\omega)$ has a unique minimum point $b_n^* \in A$ only for $\omega \in \Omega_1^n$ for which $\overline{ZL'G}_n(\omega)$ is positive definite (which for $Z \geq 0$ is the case only when $r_n(\omega)$ holds, see the second point of Lemma 60 for $K = ZL'$), in which case for $A_n(b') := 2 \overline{ZL'G}_n$ and $B_n(b') := -\overline{ZL'H}_n$ we have

$$b_n^* = (A_n(b'))^{-1}(\omega) B_n(b')(\omega). \quad (181)$$

Condition 76. ZG and ZH have \mathbb{Q}_1 -integrable (equivalently, \cup -integrable) entries.

Lemma 77. Let Condition 56 hold for $S = Z$, let for some $p > 1$, $\mathbb{E}_\cup(|Z|^p) < \infty$, and let for some $s \in \mathbb{N}_+$ and a random variable $\hat{\tau}$ with geometric distribution under \cup with parameter $q > 0$ it hold

$$\mathbb{1}(Z \neq 0) \tau \leq \tilde{\tau} := s + \hat{\tau}. \quad (182)$$

Then, for each $1 \leq u < p$, $\mathbb{E}_\cup(|ZH_i|^u) < \infty$ and $\mathbb{E}_\cup(|ZG_{i,j}|^u) < \infty$, $i, j \in \{1, \dots, l\}$. In particular, Condition 76 holds.

Proof. For $r \in (u, \infty)$ such that $\frac{u}{r} + \frac{u}{p} = 1$, using Hölder inequality and (182) we have

$$\mathbb{E}_{\mathbb{U}}(|Z||G|_{\infty}|^u) \leq \mathbb{E}_{\mathbb{U}}(|Z|\tau \frac{1}{2}R^2)^u \leq (\frac{1}{2}R^2)^u (\mathbb{E}_{\mathbb{U}}(|Z|^p))^{\frac{u}{p}} (\mathbb{E}_{\mathbb{U}}(\tau^r))^{\frac{u}{r}} < \infty \quad (183)$$

and

$$\mathbb{E}_{\mathbb{U}}((|Z||H|_{\infty})^u) \leq \mathbb{E}_{\mathbb{U}}(|Z|R \sum_{k=1}^{\tau} |\eta_k|)^u \leq R^u (\mathbb{E}_{\mathbb{U}}(Z^p))^{\frac{u}{p}} (\mathbb{E}_{\mathbb{U}}((\sum_{k=1}^{\tau} |\eta_k|)^r))^{\frac{u}{r}}. \quad (184)$$

Furthermore,

$$\mathbb{E}_{\mathbb{U}}((\sum_{k=1}^{\tilde{\tau}} |\eta_k|)^r) = \sum_{l=1}^{\infty} \mathbb{E}_{\mathbb{U}}(\mathbb{1}(\tilde{\tau} = l) (\sum_{k=1}^l |\eta_k|)^r) \leq \sum_{l=1}^{\infty} l^{r-1} \mathbb{E}_{\mathbb{U}}(\mathbb{1}(\tilde{\tau} = l) \sum_{k=1}^l |\eta_k|^r) \quad (185)$$

and from Schwartz inequality,

$$\mathbb{E}_{\mathbb{U}}(\mathbb{1}(\tilde{\tau} = l) \sum_{k=1}^l |\eta_k|^r) \leq \mathbb{U}(\tilde{\tau} = l)^{\frac{1}{2}} (\mathbb{E}_{\mathbb{U}}((\sum_{k=1}^l |\eta_k|^r)^2))^{\frac{1}{2}}. \quad (186)$$

It holds $\mathbb{U}(\tilde{\tau} = s + k) = q(1 - q)^{k-1}$, $k \in \mathbb{N}_+$, and $\mathbb{E}_{\mathbb{U}}((\sum_{k=1}^l |\eta_k|^r)^2) = l\mathbb{E}_{\mathbb{U}}(|\eta_1|^{2r} + l(l-1)(\mathbb{E}_{\mathbb{U}}(|\eta_1|^r))^2)$. The thesis easily follows from the above formulas. \square

Note that (182) in the above lemma holds e.g. for $s = 0$ and $\hat{\tau}$ as in theorem Theorem 48 if the assumptions of this theorem hold for $B = \{0\}$, or for $\hat{\tau} = 0$ for τ being an arbitrary stopping time terminated at s as in Section 5.4.

Let us assume Condition 76. Then, from Condition 43 and (78) we receive a following formula for cross-entropy

$$\text{ce}(b) = \mathbb{E}_{\mathbb{Q}_1}(Z \ln(L(b))) = b^T \mathbb{E}_{\mathbb{Q}_1}(ZG)b + \mathbb{E}_{\mathbb{Q}_1}(ZH)b. \quad (187)$$

Let $\tilde{A} = 2\mathbb{E}_{\mathbb{Q}_1}(ZG) = \nabla^2 \text{ce}(b)$, $b \in \mathbb{R}^l$, and $\tilde{B} = -\mathbb{E}_{\mathbb{Q}_1}(ZH)$. Then, we have $\nabla \text{ce}(b) = \tilde{A}b - \tilde{B}$. Thus, if $\mathbb{E}_{\mathbb{Q}_1}(ZG)$ is positive definite, then ce has a unique point $b^* \in A$, satisfying

$$b^* = \tilde{A}^{-1}\tilde{B}. \quad (188)$$

If $Z \geq 0$, then from Lemma 64 for $K = Z$, $\mathbb{E}_{\mathbb{Q}_1}(ZG)$ is positive definite only if Condition 62 holds. Remark 54 applies also to the above discussion in the LETGS setting.

5.8 Background theory for mean square, inefficiency constant, and their estimators

Let us present some background theory which shall be useful for proving different properties of mean square and inefficiency constant, as well as of their estimators. Some of the below theorems are modifications or slight extensions of well-known results (see e.g. the appendix of Chapter 1 in [51]). Let $A \in \mathcal{B}(\mathbb{R}^l)$ be nonempty. A function $f : A \rightarrow \bar{\mathbb{R}}$ is said to be lower semicontinuous in a point $b \in A$ if $\liminf_{x \rightarrow b} f(x) \geq f(b)$. It is said that f is lower semicontinuous if it is lower semicontinuous in each $b \in A$.

Lemma 78. *A lower semicontinuous function $f : A \rightarrow \overline{\mathbb{R}}$ such that $f > -\infty$ (i.e. $f(b) > -\infty, b \in A$) attains a minimum on each nonempty compact set $K \subset A$ (where such minimum may be equal to infinity).*

Proof. Let $m = \inf_{b \in K} f(b)$ and let $a_n \in K, n \in \mathbb{N}_+$, be such that $\lim_{n \rightarrow \infty} f(a_n) = m$. Consider a subsequence $(a_{n_k})_{k \in \mathbb{N}_+}$ of $(a_n)_{n \in \mathbb{N}_+}$, converging to some $b^* \in K$. Then, from lower semicontinuity, $m = \liminf_{k \rightarrow \infty} f(a_{n_k}) \geq f(b^*)$, so that $f(b^*) = m$. \square

Condition 79. *For a random variable Y with values in a measurable space \mathcal{S} , $h : \mathcal{S}(A) \otimes \mathcal{S} \rightarrow \mathcal{S}(\overline{\mathbb{R}})$ is such that a.s. $b \rightarrow h(b, Y)$ is lower semicontinuous on A and*

$$\mathbb{E}(\sup_{b \in A} (h(b, Y)_-)) < \infty. \quad (189)$$

For such h we define $b \in A \rightarrow f(b) = \mathbb{E}(h(b, Y))$.

Lemma 80. *Assuming Condition 79, we have $f > -\infty$ and f is lower semicontinuous on A .*

Proof. From (189), $f > -\infty$. For each $b \in A$ and $a_n \in A, n \in \mathbb{N}$, such that $\lim_{n \rightarrow \infty} a_n = b$, from Fatou lemma (which can be used thanks to (189)) and a.s. lower semicontinuity of $b \rightarrow h(b, Y)$,

$$\liminf_{n \rightarrow \infty} f(a_n) \geq \mathbb{E}(\liminf_{n \rightarrow \infty} h(a_n, Y)) \geq f(b). \quad (190)$$

\square

Let further in this section $A \subset \mathbb{R}^l$ be open. For $x \in A$, let $d_x = \inf_{y \in A^c} |y - x|$. For a sequence $x_n \in A, n \in \mathbb{N}_+$, let us write $x_n \uparrow A$ if $\max(\frac{1}{d_{x_n}}, |x_n|) \rightarrow \infty$ as $n \rightarrow \infty$, i.e. x_n in a sense tries to leave A . For $a \in \overline{\mathbb{R}}$ and $f : A \rightarrow \overline{\mathbb{R}}$, let us denote by $\lim_{x \uparrow A} f(x) = a$ the fact that $\lim_{n \rightarrow \infty} f(x_n) = a$ whenever $x_n \uparrow A$.

Condition 81. *A function $f : A \rightarrow \overline{\mathbb{R}}, f > -\infty$, is lower semicontinuous and $\lim_{x \uparrow A} f(x) = \infty$.*

Condition 82. *Condition 81 holds, the set B on which f is finite is nonempty and convex, and f is strictly convex on B .*

Lemma 83. *Under Condition 81, f attains a minimum on A and if Condition 82 holds, then the corresponding minimum point b^* is unique and $f(b^*) < \infty$.*

Proof. Under Condition 81, for sufficiently large $M > 0$, for a compact set

$$K = \{b \in A : |b| \leq M, d_b \geq \frac{1}{M}\}, \quad (191)$$

we have $\inf_{b \in A} f(b) = \inf_{b \in K} f(b)$. From Lemma 78 there exists a minimum point $b^* \in K$ of f on K and thus also on A . Under Condition 82 we have $f(b^*) < \infty$ and the uniqueness of b^* follows from strict convexity of f . \square

Lemma 84. *Assuming Condition 79, if with positive probability $\lim_{b \uparrow A} h(b, Y) = \infty$, then $\lim_{b \uparrow A} f(b) = \infty$.*

Proof. For $a_k \uparrow A$, from Fatou lemma

$$\liminf_{k \rightarrow \infty} f(a_k) \geq \mathbb{E}(\liminf_{k \rightarrow \infty} h(a_k, Y)) = \infty. \quad (192)$$

□

Lemma 85. *Under Condition 79, let us assume that A is convex, for some $b_0 \in A$, $f(b_0) < \infty$, and a.s. $b \rightarrow h(b, Y)$ is convex. Then, f is convex on the convex nonempty set $B \subset A$ on which it is finite. If further with positive probability $b \rightarrow h(b, Y) \in \mathbb{R}$ is strictly convex and $\lim_{b \uparrow A} h(b, Y) = \infty$, then f satisfies Condition 82.*

Proof. The (strict) convexity of f and convexity of B easily follow from $f(b) = \mathbb{E}(h(b, Y))$. The remaining points of Condition 82 follow from lemmas 80 and 84. □

Let us consider different functions and their estimators as in sections 5.1 and 5.2, under appropriate assumptions as in these sections.

Condition 86. *A is convex and $b \in A \rightarrow L(b)(\omega) \in \mathbb{R}$ is convex and continuous, $\omega \in \Omega_1$.*

From (135), if Condition 86 holds, then $b \rightarrow \widehat{\text{msq}}_n(b', b)$ is convex (for each $b' \in A$ and when evaluated on each $\omega \in \Omega_1^n$). For A open and convex, let the random condition p_{msq} on \mathcal{S}_1 hold only if $Z \neq 0$, $b \rightarrow L(b) \in \mathbb{R}_+$ is strictly convex, and $\lim_{b \uparrow A} L(b) = \infty$.

Remark 87. *If Condition 86 holds and for some $n \in \mathbb{N}_+$, $\omega = (\omega_i)_{i=1} \in \Omega_1^n$, and $i \in \{1, \dots, n\}$, $p_{\text{msq}}(\omega_i)$ holds, then for each $b' \in A$, $b \rightarrow \widehat{\text{msq}}_n(b', b)(\omega) \in \mathbb{R}$, is strictly convex, continuous, and $\lim_{b \uparrow A} \widehat{\text{msq}}_n(b', b)(\omega) = \infty$.*

It holds

$$\widehat{\text{msq}}_n(b', b) = \frac{1}{n^2} \sum_{i=1}^n \left(Z_i^2 L'_i L_i(b) \sum_{j \in \{1, \dots, n\}, j \neq i} \frac{L'_j}{L_j(b)} \right) + \frac{1}{n} \overline{((ZL')^2)}_n. \quad (193)$$

Thus, $b \rightarrow \widehat{\text{var}}_n(b', b)$ and $b \rightarrow \widehat{\text{msq}}_n(b', b)$ are positively linearly equivalent to $b \rightarrow f_{\text{var},n}(b', b)$ for

$$f_{\text{var},n}(b', b) = \sum_{i=1}^n \left(Z_i^2 L'_i \sum_{j \in \{1, \dots, n\}, j \neq i} \frac{L'_j L_i(b)}{L_j(b)} \right), \quad b', b \in A. \quad (194)$$

Condition 88. *For each $\omega_1, \omega_2 \in \Omega_1$, $b \rightarrow \frac{L(b)(\omega_1)}{L(b)(\omega_2)}$ is convex.*

Note that if Condition 88 holds, then $b \rightarrow f_{\text{var},n}(b', b)$ is convex and thus so are $b \rightarrow \widehat{\text{var}}_n(b', b)$ and $b \rightarrow \widehat{\text{msq}}_n(b', b)$.

Remark 89. *Let us assume Condition 53. Then, for each $n \in \mathbb{N}_+$, from $\widehat{\text{msq}}_n$ being nonnegative, Condition 79 holds for $Y = \tilde{\kappa}_n$ and $h(b, \cdot) = \widehat{\text{msq}}_n(b', b)(\cdot)$, for which $f = \text{msq}$ in that condition.*

Lemma 90. *Under Condition 86, if $\mathbb{Q}_1(p_{\text{msq}}) > 0$ and for some $b \in A$, $\text{msq}(b) < \infty$, then $f = \text{msq}$ satisfies Condition 82.*

Proof. It follows from Remark 87, Remark 89 for $n = 1$, and Lemma 85. \square

Condition 91. It holds $\inf_{b \in A} c(b) = c_{min} \in \mathbb{R}_+$.

Condition 92. For some $C_{min} \in \mathbb{R}_+$ we have $C(\omega) \geq C_{min}$, $\omega \in \Omega_1$.

Condition 92 implies Condition 91 for $c_{min} = C_{min}$.

Remark 93. Note that Condition 92 holds for $C_{min} = 1$ in the Euler scheme case as in Section 4.5 for $C = \tau$ being the exit time of the scheme of a set D such that $x_0 \in D$.

Under Condition 91, if $\text{var}(b) > 0$, then $\text{ic}(b) \geq c_{min} \text{var}(b)$, and thus if further $\lim_{b \uparrow A} \text{var}(b) = \infty$, then $\lim_{b \uparrow A} \text{ic}(b) = \infty$. Note also that if c and var are lower semicontinuous (which from Lemma 80 holds e.g. if $b \rightarrow L(b)(\omega)$ is continuous, $\omega \in \Omega_1$) then ic is lower semicontinuous as well. Thus, if further $\lim_{b \uparrow A} \text{ic}(b) = \infty$, then from Lemma 83, ic attains a minimum on A .

Remark 94. Let us assume that var has a unique minimum point $b^* \in A$. If for some $b \in A$ it holds $\text{ic}(b) < \text{ic}(b^*)$, then $b \neq b^*$ and thus $\text{var}(b^*) < \text{var}(b)$, so that we must have $c(b) < c(b^*)$. Note that if var , c , and ic are differentiable (some sufficient assumptions for which shall be discussed in Section 5.12) a sufficient condition for existence of $b \in A$ such that $\text{ic}(b) < \text{ic}(b^*)$ is that $\nabla \text{ic}(b^*) \neq 0$. Since $\nabla \text{var}(b^*) = 0$, we have $\nabla \text{ic}(b^*) = \text{var}(b^*) \nabla c(b^*)$, so $\nabla \text{ic}(b^*) \neq 0$ only if $\text{var}(b^*) \neq 0$ and $\nabla c(b^*) \neq 0$.

Remark 95. Let $c(b) > 0$, $b \in A$, let var have a unique minimum point b^* , and let $\text{var}(b^*) = 0$ and $c(b^*) < \infty$. Then, b^* is also the unique minimum point of ic and we have $\text{ic}(b^*) = 0$. If further A is open, msq and c are twice continuously differentiable, and $\nabla^2 \text{msq}(b^*)$ is positive definite, then from

$$\nabla^2 \text{ic}(b) = \nabla^2(c(b)) \text{var}(b) + c(b) \nabla^2 \text{msq}(b) + (\nabla c(b)) (\nabla \text{msq}(b))^T + (\nabla \text{msq}(b)) (\nabla c(b))^T, \quad (195)$$

we have

$$\nabla^2 \text{ic}(b^*) = c(b^*) \nabla^2 \text{msq}(b^*), \quad (196)$$

and thus $\nabla^2 \text{ic}(b^*)$ is positive definite.

5.9 Mean square and its estimators in the ECM setting

Let us consider the ECM setting as in Section 4.1. As discussed there, for each $\omega \in \Omega_1$, $b \rightarrow L(b)(\omega)$ is convex, and under Condition 29, $b \rightarrow L(b)(\omega)$ is strictly convex. Thus, under Condition 29, for each $\omega \in \Omega_1$, $p_{\text{msq}}(\omega)$ as in the previous section holds only if $Z(\omega) \neq 0$ and $\lim_{|b| \uparrow A} L(b)(\omega) = \infty$. Note that for X with non-degenerate normal distribution under \mathbb{Q}_1 , for each $\omega \in \Omega_1$, $\lim_{|b| \rightarrow \infty} L(b)(\omega) = \infty$, so that p_{msq} holds only if $Z \neq 0$. For X having distribution of a product of n exponentially tilted distributions from gamma family under \mathbb{Q}_1 , we have $\mathbb{Q}_1(X \in \mathbb{R}_+^n) = 1$, and for $\omega \in \Omega_1$ such that $X(\omega) \in \mathbb{R}_+^n$, we have $L(b)(\omega) \rightarrow \infty$ as $b \uparrow A$. Thus, for such ω , $p_{\text{msq}}(\omega)$ holds only if $Z(\omega) \neq 0$, and condition $\mathbb{Q}_1(p_{\text{msq}}) > 0$, appearing in Lemma 90, reduces to $\mathbb{Q}_1(Z \neq 0) > 0$. For X having Poisson distribution under \mathbb{Q}_1 , we have $\lim_{|b| \rightarrow \infty} L(b)(\omega) = \infty$ when $X(\omega) \in \mathbb{N}_+$, but not when $X(\omega) = 0$. Thus, p_{msq} holds when $Z \neq 0$ and $X \in \mathbb{N}_+$, but not when $X = 0$, and we have $\mathbb{Q}_1(p_{\text{msq}}) > 0$ only if $\mathbb{Q}_1(X \in \mathbb{N}_+, Z \neq 0) > 0$.

Remark 96. Let us assume Condition 32. Then, for each $n \in \mathbb{N}_+$, $b' \in A$, and $\omega \in \Omega_1^n$, $b \rightarrow \widehat{\text{msq}}_n(b', b)(\omega)$ is smooth and we have

$$\nabla_b \widehat{\text{msq}}_n(b', b) = \overline{(Z^2 L'(\mu(b) - X)L(b))}_n \quad (197)$$

and

$$\nabla_b^2 \widehat{\text{msq}}_n(b', b) = \overline{(Z^2 L'(\Sigma(b) + (\mu(b) - X)(\mu(b) - X)^T)L(b))}_n. \quad (198)$$

Let us further assume Condition 31, so that $\Sigma(b)$ is positive definite. Then, for $\omega \in \Omega_1^n$ such that $Z(\omega_i) \neq 0$ for some $i \in \{1, \dots, n\}$, $\nabla_b^2 \widehat{\text{msq}}_n(b', b)(\omega)$ is positive definite for each $b', b \in A$. Indeed, for each $v \in \mathbb{R}^l \setminus \{0\}$ we have

$$\begin{aligned} v^T \nabla_b^2 \widehat{\text{msq}}_n(b', b)v &= v^T \Sigma(b)v \overline{(Z^2 L'L(b))}_n + \overline{(Z^2 L'L(b)((\mu(b) - X)^T v)^2)}_n \\ &\geq v^T \Sigma(b)v \overline{(Z^2 L'L(b))}_n > 0. \end{aligned} \quad (199)$$

Note that Condition 88 holds for ECM since for each $\omega_1, \omega_2 \in \Omega_1$ we have

$$\frac{L(b)(\omega_1)}{L(b)(\omega_2)} = \exp(b^T(X(\omega_2) - X(\omega_1))). \quad (200)$$

In particular, as discussed in the previous section, the estimators variance and new estimators of mean square are convex. For each $\omega \in \Omega_1^n$ and $i, j \in \{1, \dots, n\}$, let us denote

$$v_{j,i}(\omega) = X(\omega_j) - X(\omega_i). \quad (201)$$

For each $n \in \mathbb{N}_2$, let function $g_{\text{var},n} : A \times \mathbb{R}^l \times \Omega_1^n \rightarrow \mathbb{R}$ be such that for each $b' \in A$, $b \in \mathbb{R}^l$, and $\omega \in \Omega_1^n$

$$g_{\text{var},n}(b', b)(\omega) = \sum_{i=1}^n \left((Z^2 L')(\omega_i) \sum_{j \in \{1, \dots, n\}, j \neq i} L'(\omega_j) \exp(b^T v_{j,i}(\omega)) \right). \quad (202)$$

Note that for each b' and ω as above, $b \in \mathbb{R}^l \rightarrow g_{\text{var},n}(b', b)(\omega)$ is convex and

$$g_{\text{var},n}(b', b) = f_{\text{var},n}(b', b), \quad b \in A. \quad (203)$$

For $A = \mathbb{R}^l$, we have $g_{\text{var},n} = f_{\text{var},n}$, but in some cases, like for gamma family of distributions in Section 6.9, we have $A \neq \mathbb{R}^l$ and $f_{\text{var},n}$ is only a restriction of $g_{\text{var},n}$. For each b', b , and ω as above it holds

$$\nabla_b g_{\text{var},n}(b', b)(\omega) = \sum_{i=1}^n \left((Z^2 L')(\omega_i) \sum_{j \in \{1, \dots, n\}, j \neq i} L'(\omega_j) v_{j,i}(\omega) \exp(b^T v_{j,i}(\omega)) \right) \quad (204)$$

and

$$\nabla_b^2 g_{\text{var},n}(b', b)(\omega) = \sum_{i=1}^n \left((Z^2 L')(\omega_i) \sum_{j \in \{1, \dots, n\}, j \neq i} L'(\omega_j) v_{j,i}(\omega) v_{j,i}(\omega)^T \exp(b^T v_{j,i}(\omega)) \right). \quad (205)$$

Let $n \in \mathbb{N}_2$ and $\omega \in \Omega_1^n$. Let $D(\omega) \in \mathbb{R}^{l \times n^2}$ be a matrix whose $(i-1)n + j$ th column, $i, j \in \{1, \dots, n\}$, is equal to $\mathbf{1}(Z \neq 0)(\omega_i) v_{j,i}(\omega)$ (which is equal to 0 for $i = j$).

Theorem 97. *If $D(\omega)$ has linearly independent rows, then for each $b \in \mathbb{R}^l$ and $b' \in A$, $\nabla_b^2 g_{\text{var},n}(b', b)(\omega)$ is positive definite.*

Proof. For each $t \in \mathbb{R}^l$, $t \neq 0$, there exist $i, j \in \{1, \dots, n\}$, $i \neq j$, such that $t^T \mathbb{1}(Z \neq 0)(\omega_i) v_{j,i}(\omega) \neq 0$, so that from (205),

$$t^T \nabla_b^2 g_{\text{var},n}(b', b)(\omega) t \geq (Z^2 L')(\omega_i) L'(\omega_j) (t^T v_{j,i}(\omega))^2 \exp(b^T v_{j,i}(\omega)) > 0. \quad (206)$$

□

Let for each $k \in \{1, \dots, n\}$ a matrix $\tilde{D}(k, \omega) \in \mathbb{R}^{l \times (n-1)}$ have consecutive columns equal to $v_{k,j}(\omega)$, $j \in \{1, \dots, n\} \setminus \{k\}$.

Lemma 98. *$D(\omega)$ has linearly independent rows only if for some $i \in \{1, \dots, n\}$, $Z(\omega_i) \neq 0$, and for some (equivalently, for each) $k \in \{1, \dots, n\}$, $\tilde{D}(k, \omega)$ has linearly independent rows.*

Proof. If $Z(\omega_i) = 0$, $i = 1, \dots, n$, then $D(\omega)$ has zero rows, so that they are linearly dependent. The dimensions of the linear spans of columns and vectors of a matrix are the same, so that matrices $D(\omega)$ or $\tilde{D}(k, \omega)$, $k \in \{1, \dots, n\}$, have linearly independent rows only if the dimension of the linear span of their columns is equal to l . Thus, the thesis follows from the easy to check fact that the linear span V of columns of $\tilde{D}(k, \omega)$ for different $k \in \{1, \dots, n\}$ is the same and if $Z(\omega_i) \neq 0$ for some $i \in \{1, \dots, n\}$, then the linear span of columns of $D(\omega)$ is equal to V . □

For a vector $v \in \mathbb{R}^m$, by $v \leq 0$ we mean that its coordinates are nonpositive.

Theorem 99. *If the system of linear inequalities*

$$D^T(\omega) b \leq 0, \quad b \in \mathbb{R}^l, \quad (207)$$

has only the zero solution, then for each $b' \in A$, $g_{\text{var},n}(b', b)(\omega) \rightarrow \infty$ as $|b| \rightarrow \infty$ and $\nabla_b^2 g_{\text{var},n}(b', b)(\omega)$ is positive definite, $b \in \mathbb{R}^l$. If b is a solution of (207), then for each $b' \in A$, $a \in \mathbb{R}^l$, and $t \in \mathbb{R}$, we have

$$g_{\text{var},n}(b', a + tb)(\omega) \leq g_{\text{var},n}(b', a)(\omega). \quad (208)$$

Proof. For $b \in \mathbb{R}^l$ for which (207) holds, for $i \in \{1, \dots, n\}$ such that $Z(\omega_i) \neq 0$, for $j \in \{1, \dots, n\}$, $i \neq j$, we have $b^T v_{j,i}(\omega) \leq 0$, so that (208) follows from (202). Let further (207) have only the zero solution. Then, $D(\omega)$ has linearly independent rows, and thus positive definiteness of $\nabla_b^2 g_{\text{var},n}(b', b)(\omega)$ follows from Theorem 97. Consider a function $b \in \mathbb{R}^l \rightarrow f(b) := \max\{b^T v_{j,i}(\omega) : Z(\omega_i) \neq 0, i, j \in \{1, \dots, n\}, i \neq j\}$. Then, for each $b \in \mathbb{R}^l$, $b \neq 0$, it holds $f(b) > 0$. Thus, from continuity of f we have $0 < \delta := \min\{f(b) : |b| = 1\}$ and for $0 < a := \min\{(Z^2 L')(\omega_i) L'(\omega_j) : Z(\omega_i) \neq 0, i \neq j \in \{1, \dots, n\}\}$, from (202)

$$g_{\text{var},n}(b', b)(\omega) \geq a \exp(\delta |b|) \rightarrow \infty \quad (209)$$

as $|b| \rightarrow \infty$. □

There exist numerical methods for finding the set of solutions of the system (207) and in particular for checking if it has only the zero solution, see [33]. Thanks to the below Theorem 100, if

$$Z(\omega_i) \neq 0, \quad i = 1 \dots, n, \quad (210)$$

then checking if (207) has only the zero solution is particularly simple.

Theorem 100. *Let us assume (210). Then, the system (207) has only the zero solution only if $D(\omega)$ has linearly independent rows, which from Lemma 98 holds only if for some (equivalently, for each) $k \in \{1, \dots, n\}$, $\tilde{D}(k, \omega)$ has linearly independent rows.*

Proof. Let $i, j \in \{1, \dots, n\}$. Since $v_{i,j} = -v_{j,i}$, we have

$$v_{i,j}^T b = -v_{j,i}^T b, \quad b \in \mathbb{R}^l. \quad (211)$$

From (210), the $i(n-1)+j$ th column of $D(\omega)$ is equal to $v_{i,j}$ and its $j(n-1)+i$ th column is $v_{j,i}$. Thus, from (211), if (207) holds for some $b \in \mathbb{R}^l$, then we must have $D^T(\omega)b = 0$. \square

5.10 Strongly convex functions and ϵ -minimizers

For some $A \subset \mathbb{R}^l$, consider a function $f : A \rightarrow \mathbb{R}$. For some $\epsilon \geq 0$, we say that $x^* \in A$ is an ϵ -minimizer of f , if

$$f(x^*) \leq \inf_{x \in A} f(x) + \epsilon. \quad (212)$$

Consider a convex set $S \subset A$, such that A is a neighbourhood of S (i.e. S is contained in some open set $D \subset A$). Then, f is said to be strongly convex on S (where we do not mention S if it is equal to A) with a constant $m > 0$, if f is twice differentiable on S and for each $b \in \mathbb{R}^l$ and $x \in S$

$$b^T \nabla^2 f(x) b \geq m|b|^2. \quad (213)$$

Let us discuss some properties of strongly convex functions f on S as above (see Section 9.1.2. in [8] for more details). It is well-known that f as above is strictly convex on S , and from Taylor's theorem it easily follows that for $x, y \in S$

$$f(y) \geq f(x) + (\nabla f(x))^T (y - x) + \frac{m}{2} |y - x|^2. \quad (214)$$

In particular, $f(y) \rightarrow \infty$ as $|y| \rightarrow \infty$, $y \in S$. The right-hand side of (214) in function of y is minimized by $\tilde{y} = x - \frac{1}{m} \nabla f(x)$, and thus we have

$$f(y) \geq f(x) + (\nabla f(x))^T (\tilde{y} - x) + \frac{m}{2} |\tilde{y} - x|^2 = f(x) - \frac{1}{2m} |\nabla f(x)|^2. \quad (215)$$

Let f have unique minimum point $b^* \in S$. Then, from (215) for $y = b^*$, for $x \in S$ we have

$$f(x) \leq f(b^*) + \frac{1}{2m} |\nabla f(x)|^2. \quad (216)$$

In particular, each $x \in S$ is a $\frac{1}{2m} |\nabla f(x)|^2$ -minimizer of f . Furthermore, from $\nabla f(b^*) = 0$ and (214) for $x = b^*$, we have for each $y \in S$

$$f(y) \geq f(b^*) + \frac{m}{2} |y - b^*|^2. \quad (217)$$

5.11 Mean square and its first estimator in the LETGS setting

Let us consider the LETGS setting and assume Condition 43.

Theorem 101. *For each $n \in \mathbb{N}_+$, $b \in \mathbb{R}^l \rightarrow f(b) := \widehat{\text{msq}}_n(b', b)(\omega)$ is convex and if $r_n(\omega)$ holds (see Definition 61), then f is strongly convex. If $r_n(\omega)$ does not hold, then there exists $b \in \mathbb{R}^l \setminus \{0\}$ such that*

$$f(a + tb) = f(a), \quad a \in \mathbb{R}^l, t \in \mathbb{R}. \quad (218)$$

Proof. It holds

$$\nabla^2 f(b) = \overline{(Z^2 L'(2G + (2Gb + H)(2Gb + H)^T)L(b))}_n(\omega), \quad (219)$$

which is positive semidefinite, so that f is convex. If $r_n(\omega)$ does not hold, then from the first point of Lemma 60 there exists $b \in \mathbb{R}^l$ such that each $i \in 1, \dots, n$, $A_b(\omega_i)$ does not hold, so that from Lemma 59 and (135) we receive (218). Let us assume that $r_n(\omega)$ holds. Then, from the second point of Lemma 60 for $K = Z^2 L'$, matrix $M := \overline{(Z^2 L'G)}_n(\omega)$ is positive definite. Let $m_1 > 0$ be such that $b^T M b \geq m_1 |b|^2$, $b \in \mathbb{R}^l$. For each $i \in \{1, \dots, n\}$ such that $0 < \tau(\omega_i) < \infty$, from Remark 46 we have

$$m_{2,i} := \inf_{b \in \mathbb{R}^l} L(b)(\omega_i) = \exp(\inf_{b \in \mathbb{R}^l} \ln(L(b)(\omega_i))) \in \mathbb{R}_+. \quad (220)$$

Let $m_2 = \min\{m_{2,i} : i \in \{1, \dots, n\}, 0 < \tau(\omega_i) < \infty\}$. Then, $m_2 \in \mathbb{R}_+$ and for each $a, b \in \mathbb{R}^l$ we have

$$a^T \nabla^2 f(b) a \geq 2a^T \overline{(Z^2 L'GL(b))}_n(\omega) a \geq 2m_2 a^T \overline{(Z^2 L'G)}_n(\omega) a \geq 2m_1 m_2 |a|^2. \quad (221)$$

□

Theorem 102. *If conditions 53 and 62 hold, then a.s. for sufficiently large n , $b \rightarrow \widehat{\text{msq}}_n(b', b)(\tilde{\kappa}_n)$ is strongly convex. In particular, the probability of this event converges to one as $n \rightarrow \infty$.*

Proof. It follows directly from theorems 67 and 101. □

Theorem 103. *If $\text{msq}(b_0) < \infty$ for some $b_0 \in \mathbb{R}^l$, then msq is convex on the convex nonempty set B on which it is finite and if further Condition 62 holds, then $f = \text{msq}$ satisfies Condition 82 (in particular, from Lemma 83, it has a unique minimum point). If Condition 62 does not hold, then there exists $b \in \mathbb{R}^l$, $b \neq 0$, such that*

$$\text{msq}(a + tb) = \text{msq}(a), \quad a \in \mathbb{R}^l, t \in \mathbb{R}. \quad (222)$$

Proof. The first part of the thesis follows from Theorem 102, properties of strongly convex functions discussed in Section 5.10, Remark 89, and Lemma 85. If Condition 62 does not hold, then there exists $b \in \mathbb{R}^l$, $b \neq 0$, such that $Q_1(A_b) = 0$, so that (222) follows from Lemma 59 and formula (123). □

5.12 Smoothness of functions

In this section we provide sufficient conditions for smoothness and certain properties of derivatives of functions defined in Section 5.1. Unless stated otherwise, we consider the LETS setting from Section 4.3.4, which contains the LETGS setting as a special case. As discussed at the end of Section 4.3.4, the ECM setting for $A = \mathbb{R}^l$ can be identified with the LETS setting for $\tau = 1$ and $\Lambda_1 = I_l$, so that it is easy to modify the below theory to deal also with such ECM setting.

Condition 104. For some $S : \mathcal{S}_1 \rightarrow \mathcal{S}(\overline{\mathbb{R}})$, conditions 55 and 56 hold and for each $\theta \in (\mathbb{R}^d)^s$

$$\mathbb{E}_{\mathbb{U}}(|S| \exp(\sum_{i=1}^s \theta_i^T \tilde{X}(\eta_i))) < \infty. \quad (223)$$

Note that Condition 104 implies that S is \mathbb{U} -integrable.

Remark 105. In the special case which can be identified with the ECM setting for $A = \mathbb{R}^l$ as discussed above, Condition 55 holds for $R = 1$ and Condition 56 holds for $s = 1$. Thus, for some $S : \mathcal{S}_1 \rightarrow \overline{\mathbb{R}}$, a counterpart of Condition 104 for the ECM setting for $A = \mathbb{R}^l$ reduces to demanding that

$$\mathbb{E}_{\mathbb{U}}(|S| \exp(\theta^T X)) < \infty, \quad \theta \in \mathbb{R}^l. \quad (224)$$

Remark 106. Since for $s \in \mathbb{N}_+$ and $\theta \in (\mathbb{R}^d)^s$, $\mathbb{E}_{\mathbb{U}}(\exp(\sum_{i=1}^s \theta_i^T \tilde{X}(\eta_i))) = \exp(\sum_{i=1}^s \tilde{\Psi}(\theta_i)) < \infty$, from Hölder inequality, (223) follows from $\mathbb{E}_{\mathbb{U}}(|S|^q) < \infty$ for some $q > 1$.

Condition 107. For some $t \in \mathbb{N}_+$, $f : (\mathbb{R}^l)^t \times E \rightarrow \mathbb{R}$ is such that for each $M \in \mathbb{R}_+$, for some $s, N \in \mathbb{N}_+$, $\phi \in ((\mathbb{R}^d)^s)^N$, and $u \in \mathbb{R}_+^N$, we have \mathbb{U} a.s.

$$\sup_{b \in (\mathbb{R}^l)^t : |b_i| \leq M, i=1, \dots, t} |f(b)| \leq \sum_{i=1}^N u_i \exp(\sum_{j=1}^s \phi_{i,j}^T \tilde{X}(\eta_j)) \quad (225)$$

(where $f(b) = f(b, \cdot)$).

Remark 108. Let Condition 107 hold and S satisfy Condition 104. Let $M \in \mathbb{R}_+$ and consider corresponding s, N, ϕ , and u as in Condition 107. Then, for each $\theta \in (\mathbb{R}^d)^s$

$$\mathbb{E}_{\mathbb{U}}\left(\sup_{b \in (\mathbb{R}^l)^t : |b_i| \leq M, i=1, \dots, t} |Sf(b) \exp(\sum_{j=1}^s \theta_j^T \tilde{X}(\eta_j))|\right) \leq \sum_{i=1}^N u_i \mathbb{E}_{\mathbb{U}}(S \exp(\sum_{j=1}^s (\phi_{i,j} + \theta_j)^T \tilde{X}(\eta_j))) < \infty. \quad (226)$$

In particular, $\mathbb{E}_{\mathbb{U}}(\sup_{b \in (\mathbb{R}^l)^t : |b_i| \leq M, i=1, \dots, t} |Sf(b)|) < \infty$. Furthermore, from the above $M \in \mathbb{R}_+$ being arbitrary, for each $b \in A$, $Sf(b)$ satisfies Condition 104.

Theorem 109. Let conditions 55 and 56 hold, let $t \in \mathbb{N}_+$, $r \in \mathbb{R}^t$, $w \in \mathbb{N}^{t \times l}$, $u \in \mathbb{N}^{t \times s \times d}$, $z \in \mathbb{N}^{t \times s \times d \times l}$, $y \in \mathbb{N}_+^{t \times s}$, $v \in \prod_{m=1}^t \prod_{i=1}^s (\mathbb{N}^l)^{y_{m,i}}$, and $q \in \prod_{m=1}^t \prod_{i=1}^s \mathbb{N}^{y_{m,i}}$. Let for each $b \in (\mathbb{R}^l)^t$

$$\begin{aligned} f(b) = & \mathbb{1}(S \neq 0) \prod_{m=1}^t (L(b_m))^{r_m} \prod_{i=1}^l b_{m,i}^{w_{m,i}} \prod_{i=1}^{\tau \wedge s} \prod_{j=1}^d (\prod_{i=1}^s \tilde{X}_j(\eta_i))^{u_{m,i,j}} \\ & \cdot \prod_{k=1}^l (\Lambda_{i-1})_{j,k}^{z_{m,i,j,k}} \prod_{j=1}^{y_{m,i}} (\partial_{v_{m,i,j}} \tilde{\Psi}(\lambda_{i-1}(b_m)))^{q_{m,i,j}}. \end{aligned} \quad (227)$$

Then, Condition 107 holds for such f .

Proof. Let $M \in \mathbb{R}_+$ and let $g(x) = e^x + e^{-x}$. For $p \in \mathbb{R}_+$ and $b \in \mathbb{R}^l$, $|b| \leq M$, we have \cup a.s.

$$\mathbb{1}(S \neq 0)L^p(b) \leq K(p) := \exp(ps\tilde{F}(RM)) \prod_{i=1}^s \prod_{j=1}^d g(pRM\tilde{X}_j(\eta_i)). \quad (228)$$

Let for $x \in [0, \infty)$ and $v \in \mathbb{N}^l$

$$U_v(x) = 1 + \sup_{b \in \mathbb{R}^l, |b| \leq x} |\partial_v \tilde{\Psi}(b)|, \quad (229)$$

which is finite thanks to Remark 33. Then, for each $b \in (\mathbb{R}^l)^t$, $|b_i| \leq M$, $i = 1, \dots, m$, we have \cup a.s.

$$\begin{aligned} |f(b)| &\leq \mathbb{1}(S \neq 0) \prod_{m=1}^t (K(r_m) M^{\sum_{i=1}^l w_{m,i}} \prod_{i=1}^s (\prod_{j=1}^d (g(\tilde{X}_j(\eta_i))^{u_{m,i,j}} (1+R)^{\sum_{k=1}^l z_{m,i,j,k}})) \\ &\quad \cdot \prod_{j=1}^{y_{m,i}} U_{v_{m,i,j}}(RM)^{q_{m,i,j}}). \end{aligned} \quad (230)$$

The right-hand side of (230) can be rewritten to have form as the right-hand side of (225). \square

Theorem 110. *If conditions 55 and 56 hold, then for each $t \in \mathbb{N}_+$, $p_1, p_2 \in \mathbb{R}^t$ such that $p_{2,i} \geq 1$, $i = 1, \dots, t$, $M \in \mathbb{R}_+$, $v \in (\mathbb{N}^l)^t$, and for $h_{p_1, p_2}(b, \omega) := (\mathbb{1}(S \neq 0) \prod_{i=1}^t |\partial_{v_i}(L^{p_{1,i}}(b_i))|^{p_{2,i}})(\omega)$, $b \in (\mathbb{R}^l)^t$, $\omega \in E$, Condition 107 holds for $f = h_{p_1, p_2}$.*

Proof. Since for $p_3 \in (\mathbb{N}_+)^t$ such that $p_{3,i} \geq p_{2,i}$, $i = 1, \dots, t$, we have

$$|h_{p_1, p_2}(b)| \leq \mathbb{1}(S \neq 0) \prod_{i=1}^t (1 + |\partial_{v_i} L^{p_{1,i}}(b_i)|)^{p_{3,i}}, \quad (231)$$

it is sufficient to prove the above theorem for $p_2 \in \mathbb{N}_+^t$. In such case $h_{p_1, p_2}(b)$ is a linear combination of a finite number of variables as in (227). Thus, the thesis follows from Theorem 109. \square

Theorem 111. *If Condition 104 holds, then for each $p \in \mathbb{R}_+$, for $g(b) = SL^p(b)$, $b \in \mathbb{R}^l \rightarrow f(b) = \mathbb{E}_\cup(g(b)) \in \mathbb{R}$ is smooth and we have $\partial_v f(b) = \mathbb{E}_\cup(\partial_v g(b))$, $v \in \mathbb{N}^l$, $b \in \mathbb{R}^l$.*

Proof. It follows from Theorem 110 for $p_1 = p$ and $p_2 = 1$ and from Remark 108 by induction using mean value and Lebesgue dominated convergence theorems. \square

Theorem 112. *If Condition 104 holds*

1. for $S = 1$, then $1 = \mathbb{E}_\cup(L^{-1}(b))$ and for each $v \in \mathbb{N}^l$, $v \neq 0$, $\mathbb{E}_\cup(\partial_v(L^{-1}(b))) = 0$,

2. for $S = Z^2$, then msq is smooth with $\partial_v \text{msq}(b) = \mathbb{E}_{\mathbb{U}}(Z^2 \partial_v L(b))$, $v \in \mathbb{N}^l$,
3. for $S = C$, then $b \rightarrow c(b)$ is smooth with $\partial_v c(b) = \mathbb{E}_{\mathbb{U}}(C \partial_v (L^{-1}(b)))$, $v \in \mathbb{N}^l$,
4. for S equal to Z^2 and C , then ic is smooth.

Proof. The first three points follow from Theorem 111 and from the fact that due to Remark 57 we have $1 = \mathbb{E}_{\mathbb{U}}(L(b)^{-1})$, $\text{msq}(b) = \mathbb{E}_{\mathbb{U}}(Z^2 L(b))$, and $c(b) = \mathbb{E}_{\mathbb{U}}(CL^{-1}(b))$. The last point is a consequence of points two and three. \square

Theorem 113. *In the ECM setting for $A = \mathbb{R}^l$, let us assume that $\mathbb{Q}_1(Z \neq 0) > 0$, conditions 31 and 32 hold, and we have (224) for $S = Z^2$. Then, $\nabla^2 \text{msq}(b)$ exists and is positive definite, $b \in \mathbb{R}^l$.*

Proof. From a counterpart of Theorem 109 and Remark 108 for ECM, $W = Z^2 L(b)(\mu(b) - X)(\mu(b) - X)^T$ has integrable entries. Thus, from the second point of a counterpart of Theorem 112 for ECM

$$\begin{aligned} \nabla^2 \text{msq}(b) &= \mathbb{E}_{\mathbb{Q}_1}(Z^2 L(b)(\nabla^2 \Psi(b) + (\mu(b) - X)(\mu(b) - X)^T)) \\ &= \nabla^2 \Psi(b) \text{msq}(b) + \mathbb{E}_{\mathbb{Q}_1}(W). \end{aligned} \quad (232)$$

For $v \in \mathbb{R}^l$, $v^T \mathbb{E}_{\mathbb{Q}_1}(W)v = \mathbb{E}_{\mathbb{Q}_1}(Z^2 L(b)((\nabla \Psi(b) - X)^T v)^2)$, so that $\mathbb{E}_{\mathbb{Q}_1}(W)$ is positive semidefinite. Thus, the thesis follows from the fact that as discussed in Section 4.1, $\nabla^2 \Psi(b)$ is positive definite and $\text{msq}(b) \in \mathbb{R}_+$, $b \in \mathbb{R}^l$. \square

Theorem 114. *In the LETGS setting, if Condition 104 holds for $S = Z^2$ and Condition 62 holds, then for a positive definite matrix*

$$M = \mathbb{E}_{\mathbb{Q}_1}(2GZ^2 \exp(-\frac{1}{2} \sum_{i=1}^{\tau} |\eta_i|^2)) \in \mathbb{R}^{l \times l}, \quad (233)$$

$\nabla^2 \text{msq}(b) - M$ is positive semidefinite, $b \in \mathbb{R}^l$. In particular, msq is strongly convex with constant m equal to the lowest eigenvalue of M .

Proof. From the second point of Theorem 112, we have

$$\nabla^2 \text{msq}(b) = \mathbb{E}_{\mathbb{Q}_1}(Z^2(2G + (2Gb + H)(2Gb + H)^T)L(b)). \quad (234)$$

From Theorem 109 and Remark 108, $Z^2 G$ and $W := Z^2(2Gb + H)(2Gb + H)^T L(b)$ have \mathbb{Q}_1 -integrable entries, and from $\mathbb{1}(Z \neq 0) |\exp(-\frac{1}{2} \sum_{i=1}^{\tau} |\eta_i|^2)| \leq 1$, so does $Z^2 G \exp(-\frac{1}{2} \sum_{i=1}^{\tau} |\eta_i|^2)$. Furthermore, $v^T \mathbb{E}_{\mathbb{Q}_1}(W)v = \mathbb{E}_{\mathbb{Q}_1}(Z^2 L(b)(2Gb + H)^T v)^2$, $v \in \mathbb{R}^l$, i.e. $\mathbb{E}_{\mathbb{Q}_1}(W)$ is positive semidefinite. From Lemma 64 for $K = 2Z^2 \exp(-\frac{1}{2} \sum_{i=1}^{\tau} |\eta_i|^2)$, M is positive definite. Furthermore, from Remark 46, $v^T \mathbb{E}_{\mathbb{Q}_1}(2GZ^2 L(b))v \geq v^T Mv$, $v \in \mathbb{R}^l$. \square

6 Minimization methods of estimators and their convergence properties

In this section we define various minimization methods of estimators and prove their appropriate strong convergence properties. Their asymptotic properties are discussed separately in Section 7.

6.1 A simple adaptive IS procedure used in our numerical experiments

Let us describe a simple framework of adaptive IS via minimization of estimators of various functions, shown in Scheme 1. A special case of this framework is used in numerical experiments in this work. In the further sections we discuss some modifications of this framework which ensure suitable convergence and asymptotic properties of the minimization results.

Let $b_0 \in A$, $k \in \mathbb{N}_+$, $n_i \in \mathbb{N}_+$, $i = 1, \dots, k$, and $N \in \mathbb{N}_+$. Let b_i , $i = 1, \dots, k$, be some A -valued random variables, defined in Scheme 1. Let us assume Condition 19 and let for $i = 1, \dots, k + 1$ and $j = 1, \dots, n_i$, $\beta_{i,j}$ be i.i.d. $\sim \mathbb{P}_1$ and $\chi_{i,j} = \xi(\beta_{i,j}, b_{i-1})$. Let us denote $\tilde{\chi}_i = (\chi_{i,j})_{j=1}^{n_i}$, $i = 1, \dots, k + 1$. For $k = 1$

Scheme 1 A scheme of adaptive IS

for $i := 1$ **to** k **do**

Minimize $b \rightarrow \widehat{\text{est}}_{n_i}(b_{i-1}, b)(\tilde{\chi}_i)$ over $b \in A$, e.g. using exact formulas or some numerical algorithm started at b_{i-1} . Let b_i be the minimization result.

end for

Approximate α with

$$\overline{(ZL(b_k))}_N(\tilde{\chi}_{k+1}). \quad (235)$$

we call the inside of the loop of Scheme 1 single-stage minimization (SSM) and denote $b' = b_0$, while for $k > 1$ we call this whole loop multi-stage minimization (MSM).

Let us now consider the LETGS setting for ξ as in (70). Then, using notation (72), (235) is equal to

$$\frac{1}{N} \sum_{i=1}^N (ZL(b_k))^{(b_k)}(\beta_{k+1,i}). \quad (236)$$

From discussion in Section 5.7, if $A_{n_i}(b_{i-1})(\tilde{\chi}_i) = \frac{1}{n_i} \sum_{j=1}^{n_i} 2(ZL(b_{i-1})G)^{(b_{i-1})}(\beta_{i,j})$ is positive definite, then for $B_n(b_{i-1})(\tilde{\chi}_i) = -\frac{1}{n_i} \sum_{j=1}^{n_i} (ZL((b_{i-1}))H)^{(b_{i-1})}(\beta_{i,j})$, the unique minimum point b_i of $b \rightarrow \widehat{\text{est}}_{n_i}(b_{i-1}, b)(\tilde{\chi}_i)$ is given by formula $b_i = (A_{n_i}(b_{i-1})(\tilde{\chi}_i))^{-1} B_n(b_{i-1})(\tilde{\chi}_i)$. Thus, in such case, finding b_i reduces to solving a linear system of equations. For $\widehat{\text{est}}$ replaced by msq , msq_2 , or ic , functions $b \rightarrow \widehat{\text{est}}_{n_i}(b_{i-1}, b)(\tilde{\chi}_i)$ can be minimized using some minimization algorithms which can utilise some formulas for their exact derivatives. Let us only provide formulas for such derivatives used in our numerical experiments. It holds

$$\nabla_b \widehat{\text{msq}}_{n_i}(b_{i-1}, b)(\tilde{\chi}_i) = \frac{1}{n_i} \sum_{j=1}^{n_i} (Z^2 L(b_{i-1})(2Gb + H)L(b))^{(b_{i-1})}(\beta_{i,j}), \quad (237)$$

$$\nabla_b^2 \widehat{\text{msq}}_{n_i}(b_{i-1}, b)(\tilde{\chi}_i) = \frac{1}{n_i} \sum_{j=1}^{n_i} (Z^2 L(b_{i-1})(2G + (2Gb + H)(2Gb + H)^T)L(b))^{(b_{i-1})}(\beta_{i,j}), \quad (238)$$

$$\begin{aligned}
\nabla_b \widehat{\text{msq}}_{2_{n_i}}(b_{i-1}, b)(\tilde{\chi}_i) &= \nabla_b \widehat{\text{msq}}_{n_i}(b_{i-1}, b)(\tilde{\chi}_i) \frac{1}{n_i} \sum_{j=1}^{n_i} \left(\frac{L(b_{i-1})}{L(b)} \right)^{(b_{i-1})} (\beta_{i,j}) \\
&\quad - \widehat{\text{msq}}_{n_i}(b_{i-1}, b)(\tilde{\chi}_i) \frac{1}{n_i} \sum_{j=1}^{n_i} \left((2Gb + H) \frac{L(b_{i-1})}{L(b)} \right)^{(b_{i-1})} (\beta_{i,j}),
\end{aligned} \tag{239}$$

and for

$$\widehat{\text{var}}_{n_i}(b_{i-1}, b)(\tilde{\chi}_i) = \frac{n_i}{n_i - 1} \left(\widehat{\text{msq}}_{2_{n_i}}(b_{i-1}, b)(\tilde{\chi}_i) - \left(\frac{1}{n_i} \sum_{j=1}^{n_i} (ZL(b_{i-1}))^{(b_{i-1})} (\beta_{i,j}) \right)^2 \right), \tag{240}$$

$$\begin{aligned}
\nabla_b \widehat{\text{ic}}_{n_i}(b_{i-1}, b) &= \frac{1}{n_i - 1} \sum_{j=1}^{n_i} \left(\frac{L(b_{i-1})}{L(b)} C \right)^{(b_{i-1})} (\beta_{i,j}) \nabla_b \widehat{\text{msq}}_{2_{n_i}}(b_{i-1}, b)(\tilde{\chi}_i) \\
&\quad - \frac{1}{n_i} \sum_{j=1}^{n_i} \left((2Gb + H) \frac{L(b_{i-1})}{L(b)} C \right)^{(b_{i-1})} (\beta_{i,j}) \widehat{\text{var}}_{n_i}(b_{i-1}, b)(\tilde{\chi}_i).
\end{aligned} \tag{241}$$

Formulas for the second derivatives of $\widehat{\text{msq}}_{2_{n_i}}(b_{i-1}, \cdot)$ and $\widehat{\text{ic}}_{n_i}(b_{i-1}, \cdot)$ can also be easily computed and used in minimization algorithms, but we did not apply them in our experiments. When evaluating the above expressions one can take advantage of formulas (76), (79), and (80).

6.2 Helper strong laws of large numbers

In this section we provide various uniform SLLNs needed further on. The following uniform SLLN is well-known, see Theorem A1, Section 2.6 in [47].

Theorem 115. *Let Y be a random variable with values in a measurable space \mathcal{S} , let $V \subset \mathbb{R}^l$ be nonempty and compact, and let $h : \mathcal{S}(V) \otimes \mathcal{S} \rightarrow \mathcal{S}(\mathbb{R})$ be such that a.s. $x \rightarrow h(x, Y)$ is continuous and $\mathbb{E}(\sup_{x \in V} |h(x, Y)|) < \infty$. Then, for Y_1, Y_2, \dots i.i.d. $\sim Y$, a.s. as $n \rightarrow \infty$, $x \in V \rightarrow \frac{1}{n} \sum_{i=1}^n h(x, Y_i)$ converges uniformly to a continuous function $x \in V \rightarrow \mathbb{E}(h(x, Y))$.*

For each $p \in [1, \infty]$ and $\overline{\mathbb{R}}$ -valued random variable U , let $|U|_p$ denote the norm of U in $L^p(\mathbb{P})$. We have a following well-known generalization of Hölder inequality which follows from it by induction.

Lemma 116. *For $\overline{\mathbb{R}}$ -valued random variables U_i and $q_i \in [0, \infty]$, $i = 1, \dots, n$, such that $\sum_{i=1}^n \frac{1}{q_i} = 1$, it holds*

$$\left| \prod_{i=1}^n U_i \right|_1 \leq \prod_{i=1}^n |U_i|_{q_i}. \tag{242}$$

Let further $n_i \in \mathbb{N}_+$, $i \in \mathbb{N}_+$, and $r \in \mathbb{N}_+$. To our knowledge, the SLLNs that follow are new.

Theorem 117. *Let $M_i \geq 0$, $i \in \mathbb{N}_+$, be such that*

$$\sum_{i=0}^{\infty} \frac{M_i}{n_i^r} < \infty. \tag{243}$$

Consider σ -fields $\mathcal{G}_i \subset \mathcal{F}$, $i \in \mathbb{N}_+$, and $\overline{\mathbb{R}}$ -valued random variables $\psi_{i,j}$, $j = 1 \dots, n_i$, $i \in \mathbb{N}_+$, which are conditionally independent given \mathcal{G}_i for the same i and different j , and we have $\mathbb{E}(\psi_{i,j}^{2r}) \leq M_i < \infty$ and $\mathbb{E}(\psi_{i,j} | \mathcal{G}_i) = 0$. Then, for $\hat{a}_i = \frac{1}{n_i} \sum_{j=1}^{n_i} \psi_{i,j}$ we have a.s. $\lim_{n \rightarrow \infty} \hat{a}_n = 0$.

Proof. From Borel-Cantelli lemma it is sufficient to prove that for each $\epsilon > 0$

$$\sum_{i=1}^{\infty} \mathbb{P}(|\hat{a}_i| > \epsilon) < \infty. \quad (244)$$

From Markov inequality we have

$$\mathbb{P}(|\hat{a}_i| > \epsilon) \leq \frac{\mathbb{E}(\hat{a}_i^{2r})}{\epsilon^{2r}}, \quad (245)$$

so that it is sufficient to prove that

$$\sum_{i=1}^{\infty} \mathbb{E}(\hat{a}_i^{2r}) < \infty. \quad (246)$$

Let us consider separately the easiest to prove case of $r = 1$. We have for $i \in \mathbb{N}_+$, and $j, l \in \{1, \dots, n_i\}$, $j \neq l$, from conditional independence

$$\mathbb{E}(\psi_{i,j} \psi_{i,l} | \mathcal{G}_i) = \mathbb{E}(\psi_{i,j} | \mathcal{G}_i) \mathbb{E}(\psi_{i,l} | \mathcal{G}_i) = 0, \quad (247)$$

and thus

$$\mathbb{E}(\psi_{i,j} \psi_{i,l}) = \mathbb{E}(\mathbb{E}(\psi_{i,j} \psi_{i,l} | \mathcal{G}_i)) = 0. \quad (248)$$

Thus, for $i \in \mathbb{N}_+$

$$\mathbb{E}(\hat{a}_i^2) = \frac{1}{n_i^2} \left(\sum_{j=1}^{n_i} \mathbb{E}(\psi_{i,j})^2 + \sum_{j < l \in \{1, \dots, n_i\}} 2\mathbb{E}(\psi_{i,j} \psi_{i,l}) \right) \leq \frac{M_i}{n_i}. \quad (249)$$

Now, (246) follows from (243).

For general $r \in \mathbb{N}_+$, denoting $J_i = \{v \in \mathbb{N}^{n_i} : \sum_{j=1}^{n_i} v_j = 2r\}$ and for $v \in J_i$, $\binom{2r}{v} = \frac{(2r)!}{\prod_{j=1}^{n_i} v_j!}$, we have for $v \in J_i$, from Lemma 116 for $n = n_i$, $U_j = \psi_{i,j}^{v_j}$, and $\frac{1}{q_i} = \frac{v_i}{2r}$,

$$\mathbb{E}(|\prod_{j=1}^{n_i} \psi_{i,j}^{v_j}|) \leq \prod_{j=1}^{n_i} (\mathbb{E}(\psi_{i,j}^{2r}))^{\frac{v_j}{2r}} \leq M_i. \quad (250)$$

Thus,

$$\mathbb{E}(\hat{a}_i^{2r}) = \frac{1}{n_i^{2r}} \sum_{v \in J_i} \binom{2r}{v} \mathbb{E}(\prod_{j=1}^{n_i} \psi_{i,j}^{v_j}) < \infty. \quad (251)$$

For $v \in J_i$ such that $v_k = 1$ for some $k \in \{1, \dots, n_i\}$, denoting $\psi_{i, \sim k} = \prod_{j \in \{1, \dots, n_i\}, j \neq k} \psi_{i,j}^{v_j}$, we have that $\psi_{i,k}$ and $\psi_{i, \sim k}$ are conditionally independent. Furthermore, from Lemma 116 for $n = n_i$, $U_j = \psi_{i,j}^{v_j}$ for $j \neq k$, $U_k = 1$, and $\frac{1}{q_i} = \frac{v_i}{2r}$, we have

$$\mathbb{E}(|\psi_{i, \sim k}|) \leq \prod_{j \in \{1, \dots, n_i\}, j \neq k} (\mathbb{E}(\psi_{i,j}^{2r}))^{\frac{v_j}{2r}} < \infty. \quad (252)$$

Thus,

$$\mathbb{E}\left(\prod_{j=1}^{n_i} \psi_{i,j}^{v_j} | \mathcal{G}_i\right) = \mathbb{E}\left(\mathbb{E}(\psi_{i,k} | \mathcal{G}_i) \mathbb{E}(\psi_{i,\sim k} | \mathcal{G}_i)\right) = 0, \quad (253)$$

and $\mathbb{E}\left(\prod_{j=1}^{n_i} \psi_{i,j}^{v_j}\right) = 0$. Therefore, for $\tilde{J}_i = \{v \in (\mathbb{N} \setminus \{1\})^{n_i} : \sum_{j=1}^{n_i} v_j = 2r\}$ we have

$$\mathbb{E}(\hat{a}_i^{2r}) = \frac{1}{n_i^{2r}} \sum_{v \in \tilde{J}_i} \binom{2r}{v} \mathbb{E}\left(\prod_{j=1}^{n_i} \psi_{i,j}^{v_j}\right). \quad (254)$$

Note that for $v \in \tilde{J}_i$ and $p(v) := |\{j \in \{1, \dots, n_i\} : v_j \neq 0\}|$, it holds $p(v) \leq r$, and thus for $J'_i = \{v \in \{0, 2, 3, \dots, 2r\}^{n_i} : p(v) \leq r\}$, we have $\tilde{J}_i \subset J'_i$. Therefore,

$$|\tilde{J}_i| \leq |J'_i| \leq \binom{n_i}{r} (2r)^r \leq n_i^r (2r)^r. \quad (255)$$

Furthermore,

$$\binom{2r}{v} \leq (2r)!. \quad (256)$$

From (254), (250), (255), and (256)

$$\mathbb{E}(\hat{a}_i^{2r}) \leq \frac{M_i}{n_i^r} (2r)! (2r)^r. \quad (257)$$

Inequality (246) follows from (257) and (243). \square

Let $A \in \mathcal{B}(\mathbb{R}^l)$ be nonempty, and let a family of probability distributions $\mathbb{Q}(b)$, $b \in A$, be as in Section 3.4. Let b_i , $i \in \mathbb{N}$, be A -valued random variables.

Condition 118. *Nonempty sets $K_i \in \mathcal{B}(A)$, $i \in \mathbb{N}_+$, are such that a.s. for sufficiently large i , $b_i \in K_i$.*

Condition 119. *For each $i \in \mathbb{N}_+$, $\chi_{i,j}$, $j = 1, \dots, n_i$, are conditionally independent given b_{i-1} and have conditional distribution $\mathbb{Q}(v)$ given $b_{i-1} = v$ (see page 420 in [17] or page 15 in [28] for definition of such conditional distribution).*

Condition 119 is implied by a following one.

Condition 120. *Condition 19 holds and for each $i \in \mathbb{N}_+$, $\beta_{i,j} \sim \mathbb{P}_1$, $j = 1, \dots, n_i$, are independent and independent of b_{i-1} . Furthermore, $\chi_{i,j} = \xi(\beta_{i,j}, b_{i-1})$ and $\tilde{\chi}_i = (\chi_{i,j})_{j=1}^{n_i}$, $i \in \mathbb{N}_+$.*

Let us further in this section assume conditions 118 and 119.

Condition 121. *Function $h : \mathcal{S}(A) \otimes \mathcal{S}_1 \rightarrow \mathcal{S}(\mathbb{R})$ is such that for each $v \in A$, $\mathbb{E}_{\mathbb{Q}(v)}(h(v, \cdot)) = 0$, and for*

$$M_i = \sup_{w \in K_{i-1}} \mathbb{E}_{\mathbb{Q}(w)}(h(w, \cdot)^{2r}), \quad i \in \mathbb{N}_+, \quad (258)$$

(243) holds.

Theorem 122. *Under Condition 121, for*

$$\widehat{b}_i = \frac{1}{n_i} \sum_{k=1}^{n_i} h(b_{i-1}, \chi_{i,k}), \quad i \in \mathbb{N}_+, \quad (259)$$

we have a.s.

$$\lim_{i \rightarrow \infty} \widehat{b}_i = 0. \quad (260)$$

Proof. Let for $i \in \mathbb{N}_+$, $h_i : A \times \Omega_1 \rightarrow \mathbb{R}$ be such that for each $x \in \Omega_1$, $h_i(v, x) = h(v, x)$ when $v \in K_{i-1}$ and $h_i(v, x) = 0$ when $v \in A \setminus K_{i-1}$. For

$$\widehat{a}_i = \frac{1}{n_i} \sum_{k=1}^{n_i} h_i(b_{i-1}, \chi_{i,k}), \quad (261)$$

from Condition 118 we have a.s. $\widehat{b}_i - \widehat{a}_i = \mathbb{1}(b_{i-1} \notin K_{i-1}) \widehat{b}_i \rightarrow 0$ as $i \rightarrow \infty$. Thus, to prove (260) it is sufficient to prove that a.s.

$$\lim_{i \rightarrow \infty} \widehat{a}_i = 0. \quad (262)$$

Let $\psi_{i,j} = h_i(b_{i-1}, \chi_{i,j})$, $i \in \mathbb{N}_+$, $j = 1, \dots, n_i$. From conditional Fubini's theorem (see Theorem 2, Section 22.1 in [17])

$$\begin{aligned} \mathbb{E}(\psi_{i,j}^{2r}) &= \mathbb{E}((\mathbb{E}_{\mathbb{Q}(v)}(h_i^{2r}(v, \cdot)))_{v=b_{i-1}}) \\ &= \mathbb{E}((\mathbb{1}(v \in K_{i-1}) \mathbb{E}_{\mathbb{Q}(v)}(h^{2r}(v, \cdot)))_{v=b_{i-1}}) \leq M_i. \end{aligned} \quad (263)$$

Furthermore, $\psi_{i,j}$, $j = 1, \dots, n_i$ are conditionally independent given $\mathcal{G}_i := \sigma(b_{i-1})$, and from well-known properties of conditional distribution (see Definition 1, Section 23.1 in [17]), we have

$$\begin{aligned} \mathbb{E}(\psi_{i,j} | \mathcal{G}_i) &= (\mathbb{E}_{\mathbb{Q}(v)}(h_i(v, \cdot)))_{v=b_{i-1}} \\ &= (\mathbb{1}(v \in K_{i-1}) \mathbb{E}_{\mathbb{Q}(v)}(h(v, \cdot)))_{v=b_{i-1}} = 0. \end{aligned} \quad (264)$$

Thus, (262) follows from Theorem 117. \square

Theorem 123. *If $g : \mathcal{S}(A) \otimes \mathcal{S}_1 \rightarrow \mathcal{S}(\mathbb{R})$ is such that $f(v) := \mathbb{E}_{\mathbb{Q}(v)}(g(v, \cdot)) \in \mathbb{R}$, $v \in A$, and for*

$$P_i = \sup_{v \in K_{i-1}} \mathbb{E}_{\mathbb{Q}(v)}(g(v, \cdot)^{2r}), \quad i \in \mathbb{N}_+, \quad (265)$$

we have

$$\sum_{i=1}^{\infty} \frac{P_i}{n_i^r} < \infty, \quad (266)$$

then Condition 121 holds for $h(v, y) = g(v, y) - f(v)$, $v \in A$, $y \in \Omega_1$.

Proof. Clearly, $\mathbb{E}_{\mathbb{Q}(v)}(h(v, \cdot)) = 0$, $v \in A$. Furthermore, for $v \in A$

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}(v)}(h(v, \cdot)^{2r}) &\leq \mathbb{E}_{\mathbb{Q}(v)}(|g(v, \cdot)| + |f(v)|)^{2r} \\ &\leq 2^{2r-1} \mathbb{E}_{\mathbb{Q}(v)}(g(v, \cdot)^{2r} + f(v)^{2r}) \\ &\leq 4^r \mathbb{E}_{\mathbb{Q}(v)}(g(v, \cdot)^{2r}), \end{aligned} \quad (267)$$

where in the second inequality we used the fact that $\frac{a+b}{2} \leq (\frac{a^p+b^p}{2})^{\frac{1}{p}}$, $a, b \in [0, \infty)$, $p \in [1, \infty)$, and in the last inequality we used conditional Jensen's inequality. Thus, $M_i \leq 4^r P_i$ and (243) follows from (266). \square

Condition 124. Condition 17 holds for Z replaced by some $S \in L^1(\mathbb{Q}_1)$ and for

$$P_i = \sup_{v \in K_{i-1}} \mathbb{E}_{\mathbb{Q}(v)}((SL(v))^{2r}) = \sup_{v \in K_{i-1}} \mathbb{E}_{\mathbb{Q}_1}(S^{2r}L(v)^{2r-1}), \quad i \in \mathbb{N}_+, \quad (268)$$

we have (266).

Theorem 125. Under Condition 124, a.s.

$$\lim_{k \rightarrow \infty} \overline{(SL(b_{k-1}))}_{n_k}(\tilde{\chi}_k) = \mathbb{E}_{\mathbb{Q}_1}(S). \quad (269)$$

Proof. This follows from Theorem 123 for $g(v, y) = (SL(v))(y)$, $v \in A$, $y \in \Omega_1$, in which $f(v) = \mathbb{E}_{\mathbb{Q}_1}(S)$, $v \in A$, as well as from Theorem 122. \square

For each \mathbb{R} -valued random variable Y on \mathcal{S}_1 and $q \geq 1$, let $\|Y\|_q = \mathbb{E}_{\mathbb{Q}_1}(|Y|^q)^{\frac{1}{q}}$.

Lemma 126. Let $p, q \in [1, \infty]$ be such that $\frac{1}{p} + \frac{1}{q} = 1$, let $S \in L^{2rp}(\mathbb{Q}_1)$, let Condition 17 hold for $Z = S$, and let for

$$R_i = \sup_{v \in K_{i-1}} \|\mathbb{1}(S \neq 0)L(v)^{2r-1}\|_q, \quad i \in \mathbb{N}_+, \quad (270)$$

it hold

$$\sum_{i=1}^{\infty} \frac{R_i}{n_i^r} < \infty. \quad (271)$$

Then, Condition 124 holds.

Proof. From Hölder inequality $\mathbb{E}_{\mathbb{Q}_1}(S^{2r}L(v)^{2r-1}) \leq \|S^{2r}\|_p \|\mathbb{1}(S \neq 0)L(v)^{2r-1}\|_q$, so that for P_i as in (268) we have $P_i \leq \|S^{2r}\|_p R_i$. Thus, from (271), (266) holds for such P_i . \square

The following uniform SLLN is a multi-stage version of Theorem 115, and its proof is similar to the proof of the latter (see the proof of Theorem A1, Section 2.6 in [47]).

Theorem 127. Let $V \subset \mathbb{R}^l$ be a nonempty compact set and let $h : \mathcal{S}_1 \otimes \mathcal{S}(V) \rightarrow \mathcal{S}(\mathbb{R})$ be such that for \mathbb{Q}_1 a.e. $\omega \in \Omega_1$, $b \rightarrow h(\omega, b)$ is continuous. Let $Y(\omega) = \sup_{b \in V} |h(\omega, b)|$, $\omega \in \Omega_1$, and let Condition 124 hold for $S = Y$. Then, a.s. as $k \rightarrow \infty$, $b \in V \rightarrow \hat{a}_k(b) := \frac{1}{n_k} \sum_{i=1}^{n_k} h(\chi_{k,i}, b)L(b_{k-1})(\chi_{k,i})$ converges uniformly to a continuous function $b \in V \rightarrow a(b) := \mathbb{E}_{\mathbb{Q}_1}(h(\cdot, b)) \in \mathbb{R}$.

Proof. Obviously,

$$|h(\omega, b)| \leq Y(\omega), \quad \omega \in \Omega_1, \quad b \in V, \quad (272)$$

and for each $b \in K_0$, for P_1 as in (268) for $S = Y$,

$$\mathbb{E}_{\mathbb{Q}_1}(Y) = \mathbb{E}_{\mathbb{Q}(b)}(YL(b)) \leq (\mathbb{E}_{\mathbb{Q}(b)}((YL(b))^{2r}))^{\frac{1}{2r}} \leq P_1^{\frac{1}{2r}} < \infty. \quad (273)$$

Thus, for each $v \in V$ and $v_k \in V$, $k \in \mathbb{N}_+$, such that $\lim_{k \rightarrow \infty} v_k = v$, from Lebesgue dominated convergence and \mathbb{Q}_1 a.s. continuity of $b \rightarrow h(\cdot, b)$,

$$\lim_{k \rightarrow \infty} a(v_k) = \mathbb{E}_{\mathbb{Q}_1}(\lim_{k \rightarrow \infty} h(\cdot, v_k)) = a(v) \in \mathbb{R}. \quad (274)$$

Thus, a is finite and continuous on V . Let $\epsilon > 0$. From uniform continuity of a on V , let $\delta > 0$ be such that

$$|a(x) - a(y)| < \epsilon, \quad x, y \in V, \quad |x - y| < \delta. \quad (275)$$

For each $y \in V$ and $n \in \mathbb{N}_+$, let $B_{n,y} = \{x \in V : |x - y| \leq \frac{1}{n}\}$, and let for each $\omega \in \Omega_1$

$$r_{n,y}(\omega) = \sup\{|h(\omega, x) - h(\omega, y)| : x \in B_{n,y}\}. \quad (276)$$

For \mathbb{Q}_1 a.e. ω for which $h(\omega, \cdot)$ is continuous, $\lim_{n \rightarrow \infty} r_{n,y}(\omega) = 0$, $y \in V$. Furthermore,

$$r_{n,y}(\omega) \leq 2Y(\omega), \quad \omega \in \Omega_1, \quad n \in \mathbb{N}_+, \quad y \in V, \quad (277)$$

so that from Lebesgue dominated convergence

$$\lim_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}_1}(r_{n,y}) = \mathbb{E}_{\mathbb{Q}_1}(\lim_{n \rightarrow \infty} r_{n,y}) = 0, \quad y \in V. \quad (278)$$

Thus, for each $y \in V$ there exists $n_y \in \mathbb{N}_+$, $n_y > \frac{1}{\delta}$, such that

$$\mathbb{E}_{\mathbb{Q}_1}(r_{n_y,y}) < \epsilon, \quad (279)$$

for which let us denote $W_y = B_{n_y,y}$. For each $x, y \in V$

$$|\hat{a}_k(x) - \hat{a}_k(y)| \leq \frac{1}{n_k} \sum_{i=1}^{n_k} L(b_{k-1})(\chi_{k,i}) |h(\chi_{k,i}, x) - h(\chi_{k,i}, y)|, \quad (280)$$

so that for each $y \in V$

$$\sup_{x \in W_y} |\hat{a}_k(x) - \hat{a}_k(y)| \leq \frac{1}{n_k} \sum_{i=1}^{n_k} L(b_{k-1})(\chi_{k,i}) r_{n_y,y}(\chi_{k,i}). \quad (281)$$

From (277), Condition 124 holds for $S = r_{n_y,y}$, so that from Theorem 125, the right-hand side of (281) converges a.s. to $\mathbb{E}_{\mathbb{Q}_1}(r_{n_y,y})$ as $k \rightarrow \infty$. Thus, from (279) and (281), for each $y \in V$, a.s. for sufficiently large k ,

$$\sup_{x \in W_y} |\hat{a}_k(x) - \hat{a}_k(y)| < \epsilon. \quad (282)$$

The family $\{W_y, y \in V\}$ is a cover of V . From compactness of V there exist finite number of points $y_1, \dots, y_m \in V$ such that $\{W_{y_i} : i = 1, \dots, m\}$ is a cover V and a.s. for sufficiently large k we have

$$\sup_{x \in W_{y_i}} |\hat{a}_k(x) - \hat{a}_k(y_i)| < \epsilon, \quad i = 1, \dots, m. \quad (283)$$

From (272), for each $x \in V$, Condition 124 holds for $S = h(\cdot, x)$, so that from Theorem 125, for each $x \in V$, a.s. $\lim_{k \rightarrow \infty} \hat{a}_k(x) = a(x)$. Thus, a.s. for sufficiently large k

$$|\hat{a}_k(y_i) - a(y_i)| < \epsilon, \quad i = 1, \dots, m. \quad (284)$$

Therefore, a.s. for sufficiently large k for which (283) and (284) hold, for each $x \in V$, for some $i \in \{1, \dots, m\}$ such that $|y_i - x| < \delta$,

$$|\hat{a}_k(x) - a(x)| \leq |\hat{a}_k(x) - \hat{a}_k(y_i)| + |\hat{a}_k(y_i) - a(y_i)| + |a(y_i) - a(x)| < 3\epsilon. \quad (285)$$

□

6.3 Locally uniform convergence of estimators

In this section we apply SLLNs from the previous section to provide sufficient conditions for single- and multi-stage a.s. locally uniform convergence of various estimators from Section 5.2 as well as their derivatives to the corresponding functions and their derivatives. Such convergence will be needed when proving convergence and asymptotic properties of minimization results of these estimators in the further sections. For some $A \subset \mathbb{R}^l$, we say that functions $f_n : A \rightarrow \mathbb{R}$, $n \in \mathbb{N}_+$, converge locally uniformly to some function $f : A \rightarrow \mathbb{R}$, which we denote $f_n \xrightarrow{loc} f$, if for each compact set $K \subset A$, $f_n|_K \xrightarrow{loc} f|_K$, i.e. f_n converges to f uniformly on K .

Lemma 128. *Let $D \subset \mathbb{R}^l$ be nonempty and compact, functions $f : D \rightarrow \mathbb{R}^m$ and $s : \mathbb{R}^m \rightarrow \mathbb{R}$ be continuous, and for some $f_n : D \rightarrow \mathbb{R}^m$, $n \in \mathbb{N}_+$, let $f_n \xrightarrow{loc} f$. Then, $s(f_n) \xrightarrow{loc} s(f)$. If further $s_n : \mathbb{R}^m \rightarrow \mathbb{R}$, $n \in \mathbb{N}_+$, are such that $s_n \xrightarrow{loc} s$, then $s_n(f_n) \xrightarrow{loc} s(f)$.*

Proof. For $M = \sup_{x \in D} |f(x)| < \infty$, let $K = \overline{B}_l(0, M + 1)$ and let $\epsilon > 0$. Since s is uniformly continuous on K , let us choose $0 < \delta < 1$ such that $|s(x) - s(y)| < \epsilon$ when $|x - y| < \delta$, $x, y \in K$. Let $N \in \mathbb{N}_+$ be such that for $n \geq N$, $|f_n(x) - f(x)| < \delta$, $x \in D$. Then, for $n \geq N$ we have $|s(f_n(x)) - s(f(x))| < \epsilon$, $x \in D$. Let further $M \in \mathbb{N}_+$, $M \geq N$, be such for $n \geq M$, $|s_n(y) - s(y)| < \epsilon$, $y \in K$. Then, for $n \geq M$ and $x \in D$

$$|s_n(f_n(x)) - s(f(x))| \leq |s_n(f_n(x)) - s(f_n(x))| + |s(f_n(x)) - s(f(x))| < 2\epsilon. \quad (286)$$

□

Until dealing with cross-entropy estimators at the end of this section, we shall consider the LETS setting. Similarly as in Section 5.12, this will allow us to cover the special case of the LETGS setting and it is straightforward to modify the below theory to deal with the ECM setting for $A = \mathbb{R}^l$.

Theorem 129. *Assuming Condition 53, if Condition 104 holds*

1. for $S = 1$, then a.s. (as $n \rightarrow \infty$) $b \rightarrow \overline{(L' \partial_v (L^{-1})(b))}_n(\tilde{\kappa}_n)$ converges locally uniformly to 0 for $v \in \mathbb{N}^l \setminus \{0\}$ and to 1 for $v = 0$,
2. for $S = Z^2$, then a.s. $b \rightarrow \partial_v \widehat{\text{msq}}_n(b', b)(\tilde{\kappa}_n) = \overline{(Z^2 L' \partial_v L(b))}_n(\tilde{\kappa}_n) \xrightarrow{loc} \partial_v \text{msq}$, $v \in \mathbb{N}^l$,
3. for $S = C$, then a.s. $b \rightarrow \partial_v \widehat{c}_n(b', b)(\tilde{\kappa}_n) \xrightarrow{loc} \partial_v c$, $v \in \mathbb{N}^l$,
4. both for S equal to Z^2 and 1, then a.s. $b \rightarrow \partial_v \widehat{\text{msq}}_n(b', b)(\tilde{\kappa}_n) \xrightarrow{loc} \partial_v \text{msq}$ and $b \rightarrow \partial_v \widehat{\text{var}}_n(b', b)(\tilde{\kappa}_n) \xrightarrow{loc} \partial_v \text{var}$, $v \in \mathbb{N}^l$,
5. for $S = C$, $S = Z^2$, and $S = 1$, then a.s. $b \rightarrow \partial_v \widehat{ic}_n(b', b)(\tilde{\kappa}_n) \xrightarrow{loc} \partial_v ic$, $v \in \mathbb{N}^l$.

Proof. The first three points follow from such points of Theorem 112, Theorem 110 for $p_2 = 1$ and appropriate p_1 , Remark 108, and from Theorem 115. The fourth point follows from the first two points, the fact that a.s. $\overline{(ZL')}_n(\tilde{\kappa}_n) \rightarrow \alpha$, the last line in (137), (138), and Lemma 128. The fifth point follows from points three, four, and Lemma 128. □

Let us further in this section assume a following condition.

Condition 130. $A = \mathbb{R}^l$, $r \in \mathbb{N}_+$, for each $i \in \mathbb{N}_+$, $n_i \in \mathbb{N}_+$, and for each $i \in \mathbb{N}$, $L_i \in [0, \infty)$ and $K_i = \{b \in \mathbb{R}^l : |b| \leq L_i\}$.

Consider the following conditions.

Condition 131. For each $a_1, a_2 \in \mathbb{R}_+$

$$\sum_{i=1}^{\infty} \frac{\exp(a_1 \tilde{F}(a_2 L_{i-1}))}{n_i^r} < \infty. \quad (287)$$

Condition 132. $\lim_{i \rightarrow \infty} L_i = \infty$.

Remark 133. Let us discuss possible choices of n_i and L_i such that conditions 131 and 132 hold for each $r \in \mathbb{N}_+$. Let $A_1 \in \mathbb{N}$, $A_2 \in \mathbb{N}_+$, $m \in \mathbb{N}_2$, $0 < \delta < 1$, and $B_1, B_2 \in \mathbb{R}_+$. Consider $\tilde{F}(x) = \frac{x^2}{2}$, which corresponds to the standard normal multivariate case (see Section 4.1). Then, one can take $n_i = A_1 + A_2 m^i$ and $L_i = (B_1 + B_2(i+1)^{1-\delta})^{\frac{1}{2}}$, or alternatively $n_i = A_1 + A_2 i!$ and $L_i = (B_1 + B_2(i+1))^{\frac{1}{2}}$. For some $a_1, a_2 \in \mathbb{R}_+$, denoting $b_i = \frac{\exp(a_1 \tilde{F}(a_2 L_{i-1}))}{n_i^r}$, $i \in \mathbb{N}_+$, in the first case we have $\lim_{i \rightarrow \infty} b_i^{\frac{1}{r}} = \frac{1}{m^r} < 1$ and in the second case, using Stirling's formula, we have $\lim_{i \rightarrow \infty} b_i^{\frac{1}{r}} = 0$. Thus, in both cases (287) follows from Cauchy's criterion. For $\tilde{F}(x) = \mu_0(\exp(x) - 1)$, which corresponds to the Poisson case with initial mean μ_0 , one can take e.g. $L_i = B_1 \ln(B_2 + \ln(i+1))$ and some n_i as for the normal case above.

Lemma 134. If conditions 55 and 56 hold, then for each $p \in [0, \infty)$

$$\mathbb{E}_{\mathbb{Q}_1}(\mathbb{1}(S \neq 0)L(b)^p) \leq \exp(s(p\tilde{F}(R|b) + \tilde{F}(Rp|b))). \quad (288)$$

In particular, if $p \geq 1$ then

$$\mathbb{E}_{\mathbb{Q}_1}(\mathbb{1}(S \neq 0)L(b)^p) \leq \exp(2ps\tilde{F}(Rp|b)). \quad (289)$$

Proof. From (77) we have \mathbb{Q}_1 a.s. that if $S \neq 0$ (and thus from Condition 56, $\tau \leq s$) then

$$\begin{aligned} L(b)^p &= \exp(p(U(b) + Hb)) \\ &= \exp(pU(b) + U(-bp)) \frac{1}{L(-bp)} \\ &\leq \exp(s(p\tilde{F}(R|b) + \tilde{F}(Rp|b))) \frac{1}{L(-bp)}. \end{aligned} \quad (290)$$

Now (288) follows from $\mathbb{E}_{\mathbb{Q}_1}(\mathbb{1}(S \neq 0) \frac{1}{L(-bp)}) = \mathbb{Q}(-bp)(S \neq 0) \leq 1$. \square

Condition 135. Conditions 55 and 56 hold and for some $p \in (1, \infty)$, $\mathbb{E}_{\mathbb{Q}_1}(S^{2pr}) < \infty$.

Theorem 136. If conditions 131 and 135 hold, then Condition 124 holds (for the same S , p , r , K_i , and n_i).

Proof. From Lemma 126 it is sufficient to check that for q as in that lemma corresponding to p from Condition 135, and for R_i as in (270), we have (271). From (289) in Lemma 134, it holds

$$\begin{aligned} R_i &= \sup_{b \in K_{i-1}} \|\mathbb{1}(S \neq 0)L(b)^{2r-1}\|_q \\ &\leq \sup_{b \in K_{i-1}} \exp(2s(2r-1)\tilde{F}(Rq(2r-1)|b|)) \\ &\leq \exp(2s(2r-1)\tilde{F}(Rq(2r-1)L_{i-1})), \end{aligned} \quad (291)$$

so that (271) follows from Condition 131 for $a_1 = 2s(2r-1)$ and $a_2 = Rq(2r-1)$. \square

Theorem 137. *If conditions 118, 119, and 131 hold and Condition 135 holds for $S = U$ (that is for S denoted as U), then for each $w \in \mathbb{R}$ and $v \in \mathbb{N}^l$, a.s. as $k \rightarrow \infty$, $b \in \mathbb{R}^l \rightarrow \hat{f}_k(b) := \overline{(UL(b_{k-1})\partial_v(L(b)^w))}_{n_k}(\tilde{\chi}_k)$ converges locally uniformly to $b \in \mathbb{R}^l \rightarrow f(b) := \mathbb{E}_{\mathbb{Q}_1}(U\partial_v(L(b)^w)) \in \mathbb{R}$.*

Proof. Let $M \in \mathbb{R}_+$, $V = \{x \in \mathbb{R}^l : |x| \leq M\}$, $h(\omega, b) = U(\omega)\partial_v(L(b)^w)(\omega)$, $\omega \in \Omega_1$, $b \in V$, and $W(\omega) = \sup_{b \in V} |h(\omega, b)|$, $\omega \in \Omega_1$. For some $1 < p' < p$, from Remark 106 for $S = U^{2rp'}$ and $q = \frac{p}{p'}$, Condition 104 holds for such S . Thus, from Theorem 110 for $p_1 = w$ and $p_2 = 2rp'$ and from Remark 108, we have

$$\mathbb{E}_{\mathbb{Q}_1}(W^{2rp'}) = \mathbb{E}_{\mathbb{Q}_1}(\sup_{|b| \leq M} (U\partial_v(L(b)^w))^{2rp'}) < \infty. \quad (292)$$

Furthermore, if $W \neq 0$ then also $U \neq 0$, so that Condition 135 holds for $S = W$ and $p = p'$. Thus, from theorems 127 and 136 we receive that a.s. $b \rightarrow \hat{f}_k(b)$ converges to f uniformly on V . \square

Theorem 138. *Let conditions 118, 119, and 131 hold. If Condition 135 holds,*

1. *then a.s. $\overline{(L(b_{k-1})S)}_{n_k}(\tilde{\chi}_k)$ converges to $\mathbb{E}_{\mathbb{Q}_1}(S)$ (as $k \rightarrow \infty$),*
2. *for $S = 1$, then a.s. $b \rightarrow \overline{(L(b_{k-1})\partial_v(L(b)^{-1}))}_{n_k}(\tilde{\chi}_k)$ converges locally uniformly to 0 for $v \in \mathbb{N}^l \setminus \{0\}$ and to 1 for $v = 0$,*
3. *for $S = Z^2$, then a.s. $b \rightarrow \partial_v \widehat{\text{msq}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k) \xrightarrow{\text{loc}} \partial_v \text{msq}$, $v \in \mathbb{N}^l$,*
4. *for $S = C$, then $b \rightarrow \partial_v \widehat{c}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k) \xrightarrow{\text{loc}} \partial_v c$, $v \in \mathbb{N}^l$,*
5. *both for $S = 1$ and $S = Z^2$, then a.s. $b \rightarrow \partial_v \widehat{\text{msq}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k) \xrightarrow{\text{loc}} \partial_v \text{msq}$ and $b \rightarrow \partial_v \widehat{\text{var}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k) \xrightarrow{\text{loc}} \partial_v \text{var}$, for $v \in \mathbb{N}^l$,*
6. *for $S = C$, $S = Z^2$, and $S = 1$, then a.s. $b \rightarrow \partial_v \widehat{\text{ic}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k) \xrightarrow{\text{loc}} \partial_v \text{ic}$, $v \in \mathbb{N}^l$.*

Proof. The first point follows directly from Theorem 137 for $v = 0$ and $w = 0$. Points two to four follow from Theorem 137 and points one to three of Theorem 112. The fifth point follows from point one for $S = Z$ as well as points two, three, and Lemma 128, similarly as in the proof of the fourth point of Theorem 129. The sixth point follows from points four, five, and Lemma 128. \square

Let us now discuss single- and multi-stage locally uniform convergence of cross-entropy estimators, for which we shall consider the ECM and LETGS settings separately.

Theorem 139. *In the ECM setting, let us assume Condition 53 and that we have (152). Then, a.s.*

$$\overline{(ZL')}_n(\tilde{\kappa}_n) \rightarrow \alpha \quad (293)$$

and

$$\overline{(ZL'X)}_n(\tilde{\kappa}_n) \rightarrow \mathbb{E}_{\mathbb{Q}_1}(ZX). \quad (294)$$

Assuming further Condition 32, we have a.s.

$$b \rightarrow \partial_v \widehat{\text{ce}}_n(b', b)(\tilde{\kappa}_n) \xrightarrow{\text{loc}} \partial_v \text{ce}, \quad v \in \mathbb{N}^l. \quad (295)$$

Proof. (293) and (294) follow from SLLN. From (145) and (153) we have for $v \in \mathbb{N}^l$

$$\partial_v \text{ce}(b) - \partial_v \widehat{\text{ce}}_n(b', b)(\tilde{\kappa}_n) = \partial_v \Psi(b)(\alpha - \overline{(ZL')}_n(\tilde{\kappa}_n)) - (\partial_v b^T)(\mathbb{E}_{\mathbb{Q}_1}(ZX) - \overline{(ZL'X)}_n(\tilde{\kappa}_n)). \quad (296)$$

Thus, for compact $K \subset A$, from (293) and (294),

$$\begin{aligned} \sup_{b \in K} |\partial_v \text{ce}(b) - \partial_v \widehat{\text{ce}}_n(b', b)(\tilde{\kappa}_n)| &\leq \sup_{b \in K} |\partial_v \Psi(b)|(\alpha - \overline{(ZL')}_n(\tilde{\kappa}_n)) \\ &\quad + \sup_{b \in K} |\partial_v b^T|(\mathbb{E}_{\mathbb{Q}_1}(ZX) - \overline{(ZL'X)}_n(\tilde{\kappa}_n)) \rightarrow 0. \end{aligned} \quad (297)$$

□

Theorem 140. *In the ECM setting, let us assume that $A = \mathbb{R}^l$, conditions 118, 119, and 131 hold, and for some $s > 2$ we have $\mathbb{E}(|Z|^{rs}) < \infty$. Then, a.s.*

$$\lim_{k \rightarrow \infty} \overline{(ZL(b_{k-1}))}_{n_k}(\tilde{\chi}_k) = \alpha, \quad (298)$$

$$\lim_{k \rightarrow \infty} \overline{(ZXL(b_{k-1}))}_{n_k}(\tilde{\chi}_k) = \mathbb{E}_{\mathbb{Q}_1}(ZX), \quad (299)$$

and assuming further Condition 32,

$$b \rightarrow \partial_v \widehat{\text{ce}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k) \xrightarrow{\text{loc}} \partial_v \text{ce}, \quad v \in \mathbb{N}^l. \quad (300)$$

Proof. From Hölder inequality, for each $2 < q < s$ we have $\mathbb{E}_{\mathbb{Q}_1}(|ZX_i|^{rq}) < \infty$, $i = 1, \dots, l$. Thus, (298) and (299) follow from the counterpart of the first point of Theorem 138 for ECM for $S = Z$ and $S = ZX$ respectively and (300) can be proved similarly as (295) in Theorem 139. □

Theorem 141. *In the LETGS setting, let us assume conditions 53 and 76. Then, a.s.*

$$\overline{(ZGL')}_n(\tilde{\kappa}_n) \rightarrow \mathbb{E}_{\mathbb{Q}_1}(ZG), \quad (301)$$

$$\overline{(ZHL')}_n(\tilde{\kappa}_n) \rightarrow \mathbb{E}_{\mathbb{Q}_1}(ZH), \quad (302)$$

and

$$b \rightarrow \partial_v \widehat{\text{ce}}_n(b', b)(\tilde{\kappa}_n) \xrightarrow{\text{loc}} \partial_v \text{ce}, \quad v \in \mathbb{N}^l. \quad (303)$$

Proof. (301) and (302) follow from SLLN. From (179) and (187), $\partial_v \text{ce}(b)$ and $\partial_v \widehat{\text{ce}}_n(b', b)(\tilde{\kappa}_n)$ can be nonzero only for $v \in \mathbb{N}^l$ such that $\sum_{i=1}^l v_i \leq 2$. It is easy to check that for such v , from (301) and (302), a.s.

$$\partial_v \text{ce}(b) - \partial_v \widehat{\text{ce}}_n(b', b)(\tilde{\kappa}_n) = \partial_v (b^T (\mathbb{E}_{\mathbb{Q}_1}(ZG) - \overline{(ZGL)'}_n(\tilde{\kappa}_n))b + (\mathbb{E}_{\mathbb{Q}_1}(ZH) - \overline{(ZHL)'}_n(\tilde{\kappa}_n))b) \xrightarrow{\text{loc}} 0. \quad (304)$$

□

Theorem 142. *In the LETGS setting, let us assume conditions 55, 56, 118, 119, and 131, and that for some $p > 2$ we have $\mathbb{E}(|Z|^{rp}) < \infty$. Then, a.s.*

$$\overline{(ZGL(b_{k-1}))}_{n_k}(\tilde{\chi}_k) \rightarrow \mathbb{E}_{\mathbb{Q}_1}(ZG), \quad (305)$$

$$\overline{(ZHL(b_{k-1}))}_{n_k}(\tilde{\chi}_k) \rightarrow \mathbb{E}_{\mathbb{Q}_1}(ZH), \quad (306)$$

and

$$b \rightarrow \partial_v \widehat{\text{ce}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k) \xrightarrow{\text{loc}} \partial_v \text{ce}, \quad v \in \mathbb{N}^l. \quad (307)$$

Proof. From Lemma 77, for each $2 < u < p$ and $i \in \{1, \dots, l\}$, we have $\mathbb{E}_{\mathbb{Q}_1}(|ZH_i|^{ru}) < \infty$ and $\mathbb{E}_{\mathbb{Q}_1}(|ZG_{i,j}|^{ru}) < \infty$. Thus, (305) and (306) follow from the first point of Theorem 138 for $S = ZH_i$ and $S = ZG_{i,j}$ respectively, and (307) can be proved similarly as (303) in Theorem 141. □

6.4 Exact minimization of estimators

In this section we define exact single- and multi-stage minimization methods of estimators, abbreviated as ESSM and EMSM. We also discuss their application to cross-entropy estimators in the ECM and LETGS settings.

Let $T \subset \mathbb{R}_+$ be unbounded and $B \in \mathcal{B}(\mathbb{R}^l)$ be nonempty. It will be convenient to view ESSM and EMSM as special cases of a following abstract method for exact minimization of random functions, called EM. In EM we assume a following condition.

Condition 143. *For each $t \in T$ we are given a function $\widehat{f}_t : \mathcal{S}(B) \otimes (\Omega, \mathcal{F}) \rightarrow \mathcal{S}(\mathbb{R})$, a set $G_t \in \mathcal{F}$, and a B -valued random variable d_t . Random variable $\widehat{f}_t(b, \cdot)$ is denoted shortly as $\widehat{f}_t(b)$.*

Furthermore, it assumed that for each $t \in T$ and $\omega \in G_t$, $d_t(\omega)$ is the unique minimum point of $b \rightarrow \widehat{f}_t(b, \omega)$.

Let us now define ESSM and EMSM. For some nonempty set $A \in \mathcal{B}(\mathbb{R}^l)$, $A \subset B$, and $p \in \mathbb{N}_+$, let us consider functions

$$\widehat{\text{est}}_n : \mathcal{S}(A) \otimes \mathcal{S}(B) \otimes \mathcal{S}_1^n \rightarrow \mathcal{S}(\overline{\mathbb{R}}), \quad n \in \mathbb{N}_p. \quad (308)$$

For $B = A$ these can be some estimators as in (127). We shall further often need a following condition.

Condition 144. *For each $n \in \mathbb{N}_p$, the set $D_n \in \mathcal{B}(A) \otimes \mathcal{F}_1^n$ is such that for each $(b', \omega) \in D_n$, function $b \in B \rightarrow \widehat{\text{est}}_n(b', b)(\omega)$ has a unique minimum point, denoted as $b_n^*(b', \omega)$.*

In some cases a following stronger condition will hold.

Condition 145. Condition 144 holds for $D_n = A \times \tilde{D}_n$ for some $\tilde{D}_n \in \mathcal{F}_1^n$, $n \in \mathbb{N}_p$.

In ESSM and EMSM we assume a following condition.

Condition 146. Condition 144 holds and for each $n \in \mathbb{N}_p$, for $\mathcal{F}'_n := \{D_n \cap D : D \in \mathcal{B}(A) \otimes \mathcal{F}_1^n\}$, the function $(b', \omega) \rightarrow b_n^*(b', \omega)$ is measurable from $\mathcal{S}'_n = (D_n, \mathcal{F}'_n)$ to $\mathcal{S}(B)$.

In ESSM we also assume Condition 53 and a following condition.

Condition 147. $\mathbb{N} \cup \{\infty\}$ -valued random variables N_t , $t \in T$, are such that a.s. (7) and (10) hold.

Remark 148. In Condition 147 one can take e.g. $T = \mathbb{N}_+$ and $\mathbb{N}_k = k$, $k \in \mathbb{N}_+$. Alternatively, one can take $T = \mathbb{R}_+$ and for some nonnegative random variable U on \mathcal{S}_1 , N_t can be given by formulas (5) or (6) but for $C_i = U(\kappa_i)$, $i \in \mathbb{N}_+$ (i.e. for $S_n = \sum_{i=1}^n U(\kappa_i)$, $n \in \mathbb{N}_+$). In such case sufficient conditions for (7) and (10) to hold a.s. were discussed in Section 2. For instance, such U can be some theoretical cost variable for generating a replicate under \mathbb{Q}' and doing helper computations needed for later estimator minimization (shortly, the cost variable of a step of SSM), defined analogously as such theoretical cost variable C for generating replicates in MC in Section 1.1. In such case, N_t can be interpreted as the number of steps of SSM corresponding to the approximate theoretical budget t . If one can take $U = C$, as is the case in our numerical experiments, then the (implementation dependent) proportionality constant p_U of U to the approximate practical cost of a step of SSM will be typically higher than an analogous constant p_C for C as in Section 1.1.

For each $t \in T$, in ESSM we define d_t to be a B -valued random variable such that on the event

$$G_t := \{(N_t = k \in \mathbb{N}_p) \wedge ((b', \tilde{\kappa}_k) \in D_k)\}, \quad (309)$$

we have

$$d_t = b_k^*(b', \tilde{\kappa}_k). \quad (310)$$

On $G'_t = \Omega \setminus G_t$ one can set e.g. $d_t = b'$, $t \in T$.

In EMSM we assume that conditions 118 and 119 hold for $n_k \in \mathbb{N}_p$, $k \in \mathbb{N}_+$. Furthermore, for each $k \in \mathbb{N}_+$, d_k is a B -valued random variable such that on the event

$$G_k := \{(b_{k-1}, \tilde{\chi}_k) \in D_{n_k}\} \quad (311)$$

we have

$$d_k = b_{n_k}^*(b_{k-1}, \tilde{\chi}_k). \quad (312)$$

On G'_k one can set e.g. $d_k = b_0$ or $d_k = b_{k-1}$.

Remark 149. ESSM and EMSM are special cases of EM for respective G_t and d_t as above, in ESSM for $\hat{f}_t(b, \omega) = \mathbb{1}(N_t = k \in \mathbb{N}_p) \widehat{\text{est}}_k(b', b)(\tilde{\kappa}_k(\omega))$, while in EMSM for $T = \mathbb{N}_+$ and $\hat{f}_k(b, \omega) = \widehat{\text{est}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k(\omega))$.

In EMSM the variables b_k , $k \in \mathbb{N}$, satisfying Condition 118 can be defined in various ways. An important possibility is when we are given some K_0 -valued random variable b_0 , and b_k , $k \in \mathbb{N}_+$, are as in the below condition.

Condition 150. For each $k \in \mathbb{N}_+$, if $d_k \in K_k$, then $b_k = d_k$, and otherwise $b_k = r_k$ for some K_k -valued random variable r_k .

Note that if $K_k \subset K_{k+1}$, $k \in \mathbb{N}$, then for $k \in \mathbb{N}_+$, in the above condition we can take e.g. $r_k = b_0$, $r_k = b_{k-1}$, or $r_k = \phi_k(d_k)$ for some $\phi_k : \mathcal{S}(B) \rightarrow \mathcal{S}(K_k)$.

Consider some function $f : A \rightarrow \mathbb{R}$ and let $b^* \in A$ be its unique minimum point. We will be interested in verifying when some of the below conditions hold for EM methods, like ESSM and EMSM under identifications as in Remark 149, or some other methods defined further on.

Condition 151. A. s. for sufficiently large $t \in T$, G_t holds.

Condition 152. It holds a.s. $\lim_{t \rightarrow \infty} d_t = b^*$.

Condition 153. It holds a.s. $\lim_{t \rightarrow \infty} \widehat{f}_t(d_t) = f(b^*)$.

Consider a following condition.

Condition 154. A is open and $K_i \in \mathcal{B}(A)$, $i \in \mathbb{N}$, are such that for each compact set $D \subset A$, for sufficiently large i , $D \subset K_i$.

Note that if conditions 130 and 132 hold, then Condition 154 holds.

Remark 155. For EMSM let us assume conditions 150 and 154. Then, if for some compact set $D \subset A$ a.s. $d_k \in D$ for sufficiently large k (which happens e.g. if a.s. $d_k \rightarrow b^*$ and D is some compact neighbourhood of b^*), then a.s. for sufficiently large k , $d_k = b_k$. In particular, if Condition 152 holds for EMSM, then it also holds for d_k replaced by b_k , and if further Condition 153 holds for EMSM, then it also holds for d_k replaced by b_k .

Let us now describe how ESSM and EMSM can be applied to $\widehat{\text{est}}_n = \widehat{\text{ce}}_n$ in the ECM and LETGS settings. Let us first consider ECM as in sections 4.1 and 5.3, assuming conditions 31 and 32, as well as that we have (152), $\alpha > 0$, and (154). Then, from discussion in Section 5.3, Condition 144 holds for

$$D_n = \{(b', \omega) \in A \times \Omega_1^n : \overline{(ZL')}_n(\omega) > 0 \wedge \frac{\overline{(ZL'X)}_n(\omega)}{\overline{(ZL')}_n(\omega)} \in \mu[A]\}, \quad (313)$$

and from formula (151), Condition 146 holds. In ESSM, from (293) and (294) in Theorem 139 as well as from $\alpha > 0$, a.s. for sufficiently large n we have $\overline{(ZL')}_n(\tilde{\kappa}_n) > 0$ and a.s.

$$\frac{\overline{(ZL'X)}_n(\tilde{\kappa}_n)}{\overline{(ZL')}_n(\tilde{\kappa}_n)} \rightarrow \frac{\mathbb{E}_{Q_1}(ZX)}{\alpha}. \quad (314)$$

Thus, using further (154), the fact that $\mu[A]$ is open, and Condition 147, a.s. for sufficiently large t , G_t as in (309) holds (i.e. Condition 151 holds for ESSM), in which case

$$d_t = \mu^{-1} \left(\frac{\overline{(ZL'X)}_k(\tilde{\kappa}_k)}{\overline{(ZL')}_k(\tilde{\kappa}_k)} \right). \quad (315)$$

From Condition 147, (155), (314), (315), and continuity of μ^{-1} , Condition 152 holds. For EMSM let us additionally make assumptions as in Theorem 140.

Then, from (298) and (299) in that theorem, by similar arguments as above for ESSM, conditions 151 and 152 hold for EMSM.

Consider the LETGS setting and using notations as in Section 5.7 let us assume that Condition 76 holds and \tilde{A} is positive definite. From discussion in that section, Condition 144 holds for $D_n = \{(b', \omega) \in A \times \Omega_1^n : A_n(b')(\omega) \text{ is positive definite}\}$, and if $Z \geq 0$, then Condition 145 holds for $\tilde{D}_n = \{\omega \in \Omega_1^n : r_n(\omega)\}$. From formula (181), Condition 146 holds. In ESSM, from SLLN a.s. $A_n(b')(\tilde{\kappa}_n) \rightarrow \tilde{A}$ and $B_n(b')(\tilde{\kappa}_n) \rightarrow \tilde{B}$. Thus, from Lemma 66 and Condition 147, a.s. for sufficiently large t , $N_t = n \in \mathbb{N}_+$ and $A_n(b')(\tilde{\kappa}_n)$ is positive definite (i.e. Condition 151 holds), in which case $d_t = (A_n(b')(\tilde{\kappa}_n))^{-1} B_n(b')(\tilde{\kappa}_n)$. Thus, from (188), Condition 152 holds. For EMSM, let us make additional assumptions as in Theorem 142, so that from (305), a.s. $A_{n_k}(b_{k-1})(\tilde{\chi}_k) \rightarrow \tilde{A}$, and from (306), a.s. $B_{n_k}(b_{k-1})(\tilde{\chi}_k) \rightarrow \tilde{B}$. Then, analogously as for ESSM above, conditions 151 and 152 hold for EMSM.

6.5 Helper theorems for proving convergence properties of minimization methods with gradient-based stopping criteria

Proof of the below lemma uses a similar reasoning as in the proof of consistency of M-estimators in Theorem 5.14 in [52].

Lemma 156. *Let Condition 79 hold, Y_1, Y_2, \dots be i.i.d. $\sim Y$, $b \in A \rightarrow \hat{f}_n(b) := \frac{1}{n} \sum_{i=1}^n h(b, Y_i)$, $K \subset A$ be a nonempty compact set, and m be the minimum of f on K (which exists due to lemmas 78 and 80). Then, for each $a \in (-\infty, m)$, a.s. for sufficiently large n , $\hat{f}_n(b) > a$, $b \in K$.*

Proof. Let $U_i = B_l(0, i^{-1})$, $i \in \mathbb{N}_+$. From a.s. lower semicontinuity of $b \rightarrow h(b, Y)$, for each $v \in K$, for $g_{l,v}(x) = \inf_{b \in v + U_l} h(b, x)$, we have a.s. $g_{l,v}(Y) \uparrow h(v, Y)$ as $l \rightarrow \infty$. Thus, from monotone convergence theorem, $\mathbb{E}(g_{l,v}(Y)) \uparrow f(v)$ as $l \rightarrow \infty$, $v \in K$. In particular, $\mathbb{E}(h_{l,v}(Y)) > a$ for $l \geq l_v$ for some $l_v \in \mathbb{N}_+$, $v \in K$. The family $\{K_v = v + U_{l_v} : v \in K\}$ is a cover of K . From compactness of K , let $\{K_{v_1}, \dots, K_{v_m}\}$ be its finite subcover. Then, from the generalized SLLN in Theorem 1 (which can be used thanks to (189)),

$$\inf_{b \in K} \hat{f}_n(b) \geq \min_{k \in \{1, \dots, m\}} \frac{1}{n} \sum_{i=1}^n g_{l_{v_k}, v_k}(Y_i) \xrightarrow{\text{a.s.}} \min_{k \in \{1, \dots, m\}} \mathbb{E}(g_{l_{v_k}, v_k}(Y)) > a. \quad (316)$$

□

Lemma 157. *Let Condition 79 hold for h equal to some nonnegative h_i , $i = 1, 2$. Let $g(b) = \mathbb{E}(h_1(b, Y))$ and $\mathbb{E}(h_2(b, Y)) = 1$, $b \in A$, let Y_1, Y_2, \dots be i.i.d. $\sim Y$, and let $b \in A \rightarrow \hat{f}_{i,n}(b) := \frac{1}{n} \sum_{j=1}^n h_i(b, Y_j)$, $i = 1, 2$, and $b \in A \rightarrow \hat{g}_n(b) := \hat{f}_{1,n}(b) \hat{f}_{2,n}(b)$, $n \in \mathbb{N}_+$. Let $K \subset A$ be a nonempty compact set and m be the minimum of g on K . Then, for each $a \in (-\infty, m)$, a.s. for sufficiently large n ,*

$$\hat{g}_n(b) > a, \quad b \in K. \quad (317)$$

Proof. It holds $\hat{g}_n(b) \geq 0$, $n \in \mathbb{N}_+$, $b \in A$, so that it is sufficient to consider the case when $m > 0$ and $0 < a < m$. Let $a < d < m$. Then, from Lemma 156, a.s.

for sufficiently large n , $\widehat{f}_{1,n}(b) > d$ and $\widehat{f}_{2,n}(b) > \frac{a}{d}$, $b \in K$, in which case (317) holds. \square

Condition 158. $b^* \in \mathbb{R}^l$ and $A \in \mathcal{B}(\mathbb{R}^l)$ is a neighbourhood of b^* . A function $f : A \rightarrow \overline{\mathbb{R}}$, $f > -\infty$, is lower semicontinuous and b^* is its unique minimum point (in particular, $f(b^*) < \infty$).

Condition 159. Condition 158 holds and $B \subset \mathbb{R}^l$ is such that $A \subset B$. Functions $f_n : B \rightarrow \mathbb{R}$, $n \in \mathbb{N}_+$, fulfill

$$\lim_{n \rightarrow \infty} f_n(b^*) = f(b^*). \quad (318)$$

Furthermore, for each compact set $K \subset A$, for m equal to the minimum of f on K , for each $a < m$, for sufficiently large n , $f_n(x) > a$, $x \in K$.

Remark 160. Let Condition 158 hold, $B \subset \mathbb{R}^l$, $A \subset B$, and $f_n : B \rightarrow \mathbb{R}$, $n \in \mathbb{N}_+$ be such that $f_n|_A \xrightarrow{loc} f$. Then, Condition 159 holds.

Remark 161. Assuming Condition 159, let $\epsilon \in \mathbb{R}_+$ be such that $\overline{B}_l(b^*, \epsilon) \in A$ and let c be the minimum of f on $S_l(b^*, \epsilon)$. From uniqueness of the minimum point b^* of f , we have $c > f(b^*)$. Let $\delta \in \mathbb{R}_+$ be such that $c > f(b^*) + \delta$. Then, for sufficiently large n

$$f_n(b) \geq f(b^*) + \delta, \quad b \in S_l(b^*, \epsilon) \quad (319)$$

and

$$f_n(b^*) \leq f(b^*) + \frac{\delta}{2}. \quad (320)$$

Theorem 162. Let us assume that Condition 159 holds for convex B and for f_n which are convex and continuous, $n \in \mathbb{N}_+$. Then, for sufficiently large n , f_n possesses a minimum point $a_n \in B$. Furthermore,

$$\lim_{n \rightarrow \infty} a_n = b^* \quad (321)$$

and

$$\lim_{n \rightarrow \infty} f_n(a_n) = f(b^*). \quad (322)$$

If further B is open, f_n are differentiable on B , and a sequence $b_n \in B$, $n \in \mathbb{N}_+$, is such that $\lim_{n \rightarrow \infty} |\nabla f_n(b_n)| = 0$, then

$$\lim_{n \rightarrow \infty} b_n = b^* \quad (323)$$

and

$$\lim_{n \rightarrow \infty} f_n(b_n) = f(b^*). \quad (324)$$

Proof. Let us consider $\epsilon, \delta \in \mathbb{R}_+$ as in Remark 161. From this remark, let $N \in \mathbb{N}_+$ be such that for $n > N$ we have (319) and (320). Then, for $n > N$, for each $b \in B$ such that $|b - b^*| \geq \epsilon$, from convexity of f_n

$$\begin{aligned} f_n(b) - f_n(b^*) &\geq \frac{|b - b^*|}{\epsilon} \left(f_n(b^* + \epsilon \frac{b - b^*}{|b - b^*|}) - f_n(b^*) \right) \\ &\geq \frac{|b - b^*| \delta}{2\epsilon} > 0. \end{aligned} \quad (325)$$

For $n > N$, from (325) and continuity of f_n we have $\inf_{x \in B} f_n(x) = \min_{x \in B_l(b^*, \epsilon)} f_n(x)$, and thus f_n has a minimum point a_n and

$$|a_n - b^*| < \epsilon. \quad (326)$$

This proves (321). For $n > N$, from (320) and $f_n(a_n) \leq f_n(b^*)$ we have

$$f_n(a_n) \leq f(b^*) + \frac{\delta}{2}. \quad (327)$$

From Condition 159, for some $N_1 > N$, for $n > N_1$,

$$f_n(b) \geq f(b^*) - \frac{\delta}{2}, \quad b \in \overline{B}_l(b^*, \epsilon). \quad (328)$$

Thus, for $n > N_1$, from (326), (328), and (327), we receive that $|f_n(a_n) - f(b^*)| \leq \frac{\delta}{2}$. Since we could have selected δ arbitrarily small, we receive (322).

Let B be open and f_n be differentiable. Then, for $b \in B$ such that $b \neq b^*$, for $v = \frac{b-b^*}{|b-b^*|}$, from convexity of f_n

$$|\nabla f_n(b)| \geq \nabla_v f_n(b) \geq \frac{f_n(b) - f_n(b^*)}{|b - b^*|}. \quad (329)$$

Thus, for each $b \in B$ for which $|b - b^*| \geq \epsilon$, for $n > N$, from (329) and (325)

$$|\nabla f_n(b)| \geq \frac{\delta}{2\epsilon}. \quad (330)$$

Let $N_2 > N$ be such that for $n > N_2$

$$|\nabla f_n(b_n)| < \frac{\delta}{2\epsilon}. \quad (331)$$

Then, from (330), for $n > N_2$

$$|b_n - b^*| < \epsilon, \quad (332)$$

which proves (323). For $n > N_1 \vee N_2$ we have

$$\begin{aligned} \frac{\delta}{2} &= \frac{\delta}{2\epsilon} \epsilon > |\nabla f_n(b_n)| |b_n - b^*| \geq f_n(b_n) - f_n(b^*) \\ &\geq f_n(b_n) - f(b^*) - \frac{\delta}{2} \geq -\delta, \end{aligned} \quad (333)$$

where in the first inequality we used (331) and (332), in the second (329), in the third (320), and in the last one (328). Thus, in such case

$$\delta \geq f_n(b_n) - f(b^*) \geq -\frac{\delta}{2}, \quad (334)$$

which proves (324). \square

Lemma 163. *Let $A \subset \mathbb{R}^l$ be open. If a twice continuously differentiable function $f : A \rightarrow \mathbb{R}$ has positive definite Hessian on A , then for each convex $U \subset A$ such that for some compact $K \subset A$, $U \subset K$, f is strongly convex on U . If further $\lim_{x \uparrow A} g(x) = \infty$, then for each $x_0 \in A$ as such U one can take the sublevel set $S = \{x \in A : g(x) \leq g(x_0)\}$.*

Proof. From Lemma 65, $b \in A \rightarrow m_l(f(b))$ is continuous and thus f is strongly convex on U with constant $\inf_{x \in K} m_l(f(x)) > 0$. From convexity of f , S as above is convex, and if $\lim_{x \uparrow A} g(x) = \infty$, then for sufficiently large M , for a compact set K as in (191) we have $S \subset K$. \square

6.6 Minimization of estimators with gradient-based stopping criteria

In this section we define single- and multi-stage minimization methods of estimators with gradient-based stopping criteria, abbreviated as GSSM and GMSM respectively. We also discuss their applications to the first mean square estimators in the LETGS setting and both mean square estimators in the ECM setting.

Consider some T and B as in Section 6.4 and let additionally B be open. GSSM and GMSM are special cases of a following abstract minimization method of random functions with gradient-based stopping criteria, abbreviated as GM. Let us assume Condition 143 and let $b \rightarrow \widehat{f}_t(b, \omega)$ be differentiable, $t \in T$, $\omega \in \Omega_1$. Consider $[0, \infty]$ -valued random variables ϵ_t , $t \in T$, such that a.s.

$$\lim_{t \rightarrow \infty} \epsilon_t = 0. \quad (335)$$

Then, in GM it is assumed that we are given B -valued random variables d_t , $t \in T$, such that

$$|\nabla_b \widehat{f}_t(d_t(\omega), \omega)| \leq \epsilon_t(\omega), \quad \omega \in G_t, \quad t \in T. \quad (336)$$

We shall further need the following conditions and lemmas.

Condition 164. *Condition 144 holds, for each $n \in \mathbb{N}_p$ and $(b', \omega) \in D_n$, $b \in B \rightarrow \widehat{\text{est}}_n(b', b)(\omega)$ is differentiable, and $b_n^*(b', \omega)$ is equal to the unique point $c \in B$ such that $\nabla_b \widehat{\text{est}}_n(b', c)(\omega) = 0$.*

Lemma 165. *Condition 164 implies Condition 146.*

Proof. Function $(b, (b', \omega)) \in B \times D_n \rightarrow g(b, (b', \omega)) := \nabla_b \widehat{\text{est}}_n(b', b)(\omega)$ is measurable from $\mathcal{S}(B) \otimes (D_n, \mathcal{F}'_n)$ to $\mathcal{S}(\mathbb{R}^l)$ and for each $D \in \mathcal{B}(B)$, $(b_n^*)^{-1}(D) = \{(b', \omega) \in D_n : g(c, b', \omega) = 0, \text{ for some } c \in D\}$ is a projection of $g^{-1}(0) \cap (D \times D_n) \in \mathcal{B}(B) \otimes \mathcal{F}'_n$ onto the second coordinate. Thus, $(b_n^*)^{-1}(D) \in \mathcal{F}'_n$, $D \in \mathcal{B}(B)$. \square

Condition 166. *B is convex. Furthermore, for each $n \in \mathbb{N}_p$, the set $D_n \in \mathcal{B}(A) \otimes \mathcal{F}'_1$ is such that for each $(b', \omega) \in D_n$, $b \in B \rightarrow g(b) := \widehat{\text{est}}_n(b', b)(\omega)$ is smooth with positive definite Hessian on B , and $\lim_{b \uparrow B} g(b) = \infty$.*

Lemma 167. *Condition 166 implies Condition 164 (for $b_n^*(b', \omega)$ as in Condition 144).*

Proof. It follows from lemmas 35 and 83. \square

Except for some differences mentioned below, GSSM and GMSM are defined in the same way as ESSM and EMSM in the previous section. The first difference is that in GSSM and GMSM we additionally assume that Condition 164 holds for open B as above and we consider variables ϵ_t , $t \in T$, as for GM above. Furthermore, in GSSM, for $t \in T$, on G_t as in (309), instead of (310) we require that $|\nabla_b \widehat{\text{est}}_k(b', d_t)(\tilde{\kappa}_k(\omega))| \leq \epsilon_t$, while in GMSM, for $k \in \mathbb{N}_+$, on G_k as in (311), instead of (312) we require that $|\nabla_b \widehat{\text{est}}_{n_k}(b_{k-1}, d_k)(\tilde{\chi}_k)| \leq \epsilon_k$.

Note that GSSM and GMSM are special cases of GM under identifications as in Remark 149. Such identifications shall be frequently considered below. From Lemma 165, for $\epsilon_t = 0$, $t \in T$, GSSM and GMSM become special cases of ESSM and EMSM methods respectively.

Remark 168. For $t \in T$, if $\omega \in G_t$ and $\epsilon_t(\omega) > 0$, then under some additional assumptions on For $t \in T$, if $\omega \in G_t$ and $\epsilon_t(\omega) > 0$, then under some additional assumptions on $b \rightarrow g(b) := \hat{f}_t(b, \omega)$, $d_t(\omega)$ in GSSM or GSM method can be a result of some globally convergent iterative minimization method (i.e. one for which gradients in subsequent points converge to zero), minimizing g , started at x_0 equal to b' in GSSM or $b_{t-1}(\omega)$ in GSM, and stopped in a first point $d_t(\omega)$ in which (336) holds. As such iterative method one could use damped Newton method, for the global and quadratic convergence of which it is sufficient if g is strongly convex on the sublevel set $S = \{x \in B : g(x) \leq g(x_0)\}$, g is twice continuously differentiable on some open neighbourhood of S , and the second derivative of g is Lipschitz continuous on S (see Section 9.5.3 in [8]). From Lemma 163, such assumptions hold in the above discussed GSSM or GSM methods for each $\omega \in G_t$ e.g. if Condition 166 holds. See [8] and [41] for other examples of globally convergent minimization methods requiring typically weaker assumptions. For most iterative minimization methods, including damped Newton method, if the same method is used for each ω as above, then the fact that the resulting d_t is measurable on $G_t \cap \{\epsilon_t > 0\}$ follows from the definition of the method, performing only measurable operations. On G'_t one can define d_t in similar ways as for ESSM or EMSM in the previous section. If $\epsilon_t(\omega) > 0$, $\omega \in \Omega$, then to avoid checking if G_t holds (i.e. if $\omega \in G_t$), in some cases one can perform minimization of g as above for each $\omega \in \Omega$. See Remark 172 for discussion of a situation in which this is possible. In such case measurability of d_t also typically follows from definition of the minimization method used.

Condition 169. B as above is an open convex neighbourhood of $b^* \in \mathbb{R}^l$ and $\epsilon \in \mathbb{R}_+$ is such that $\bar{B}_l(b^*, \epsilon) \in B$. Furthermore, for some $\widehat{\text{est}}_n$ as in (308), $b' \in A$ and $\omega \in \Omega_1^n$ are such that

$$\inf_{b \in S_1(b^*, \epsilon)} \widehat{\text{est}}_n(b', b)(\omega) > \widehat{\text{est}}_n(b', b^*)(\omega). \quad (337)$$

Remark 170. Consider $\widehat{\text{est}}_n = \widehat{\text{msq}}_n$, $n \in \mathbb{N}_+$, in the LETGS setting. Then, from Theorem 101, if Condition 169 holds, then we cannot have (218) for $a = b^*$ and

$$t = \frac{\epsilon(b - b^*)}{|b - b^*|}, \quad (338)$$

and thus $r_n(\omega)$ holds. Let us now consider the ECM setting and $\widehat{\text{est}}_n = g_{\text{var}, n}$, $n \in \mathbb{N}_2$, (see (203)). If Condition 169 holds, then we cannot have (208) for $a = b^*$, $b \in \mathbb{R}^l \setminus \{0\}$, and t as in (338). Thus, from Theorem 99, in such case system (207) has only the zero solution.

For GSM in all the below examples we assume conditions 130 and 131. Let us first discuss GSSM and GSM for $\widehat{\text{est}}_n = \widehat{\text{msq}}_n$, $n \in \mathbb{N}_+$, in the LETGS setting. From Theorem 101, we can and shall take in Condition 166 (and thus from Lemma 167 also in Condition 164), $\tilde{D}_n = \{\omega \in \Omega_1^n : r_n(\omega)\}$, $n \in \mathbb{N}_+$. Let us assume Condition 62 and that for some $b \in A$, $\text{msq}(b) < \infty$, so that from Theorem 103, we can and shall take in Condition 158, $f = \text{msq}$. In GSSM, from Condition 147 and Theorem 67, Condition 151 holds. Using further the fact that from SLLN and Lemma 156 for

$$h(b, x) = (Z^2 L' L(b))(x), \quad b \in A, \quad x \in \Omega_1, \quad (339)$$

and $Y_i = \kappa_i$, $i \in \mathbb{N}_+$, Condition 159 holds for $B = A$ and $f_n(b) = \widehat{\text{msq}}_n(b', b)(\tilde{\kappa}_n)$, from Theorem 162, (336), and (335), conditions 152 and 153 hold as well. For GMSM let us assume that Condition 135 holds for $S = Z^2$. Then, from the third point of Theorem 138 and remarks 160, 161, and 170, a.s. for sufficiently large k , $r_{n_k}(\tilde{\chi}_k)$ holds, i.e. Condition 151 holds. Using further (335), (336), and Theorem 162, conditions 152 and 153 hold too.

Let us now consider the ECM setting, assuming a following condition.

Condition 171. *Conditions 31 and 32 hold and $f = \text{msq}$ satisfies Condition 158.*

From Lemma 83, $f = \text{msq}$ satisfies Condition 158 for instance when $f = \text{msq}$ satisfies Condition 82, which due to Lemma 90 and $b \rightarrow L(b)(\omega)$ being strictly convex, $\omega \in \Omega_1$, holds e.g. if for some $b \in A$, $\text{msq}(b) < \infty$, and

$$\mathbb{Q}_1(p_{\text{msq}}) > 0. \quad (340)$$

Let us first consider the case of $\widehat{\text{est}}_n = \widehat{\text{msq}}_n$, $n \in \mathbb{N}_+$, for which let us assume (340). From remarks 87 and 96, we can and shall take in Condition 166, $\tilde{D}_n = \{\omega \in \Omega_1^n : p_{\text{msq}}(\omega_i) \text{ holds for some } i \in \{1, \dots, n\}\}$, $n \in \mathbb{N}_+$. In GSSM, from SLLN, a.s. $\lim_{n \rightarrow \infty} \overline{(p_{\text{msq}})_n}(\tilde{\kappa}_n) = \mathbb{Q}_1(p_{\text{msq}})$, so that from (340) and SLLN we have a.s. $\tilde{\kappa}_k \in \tilde{D}_k$ for sufficiently large k . Thus, from Condition 147, Condition 151 holds. Using further Lemma 156 for h as in (339) and Theorem 162, conditions 152 and 153 hold too. In GMSM, let $A = \mathbb{R}^l$. Since the counterpart of Condition 135 for ECM is fulfilled for $S = \mathbb{1}(p_{\text{msq}})$, from the first point of Theorem 138, a.s. $(L(b_{k-1})S)_{n_k}(\tilde{\chi}_k) \rightarrow \mathbb{Q}_1(p_{\text{msq}})$. Thus, from (340), Condition 151 holds. Let us assume Condition 135 for $S = Z^2$. Then, from the third point of Theorem 138 and Theorem 162, conditions 152 and 153 hold.

Let us now consider for each $n \in \mathbb{N}_2$

$$\widehat{\text{est}}_n(b', b) = \widehat{\text{msq}}_n(b', b) := \frac{1}{n^2} g_{\text{var}, n}(b', b) + \frac{1}{n} \overline{((ZL')^2)_n}, \quad b' \in A, b \in B = \mathbb{R}^l. \quad (341)$$

Then, from (138) and (194) we have

$$\widehat{\text{msq}}_n(b', b) = \widehat{\text{msq}}_n(b', b), \quad b', b \in A. \quad (342)$$

Note that $\frac{1}{n} \overline{((ZL')^2)_n}$ does not depend on b . Thus, from Theorem 99, we can and shall take in Condition 166, $\tilde{D}_n = \{\omega \in \Omega_1^n : \text{system (207) has only the zero solution}\}$. In GSSM, from SLLN and Lemma 157 for $h_1(b, y) = (Z^2 L' L(b))(y)$, $h_2(b, y) = \frac{L'}{L(b)}(y)$, $b \in \mathbb{R}^l$, $y \in \Omega_1$, and $Y_i = \kappa_i$, $i \in \mathbb{N}_+$, a.s. Condition 159 holds for $B = A$ and $f_n(b) = \widehat{\text{msq}}_n(b', b)(\tilde{\kappa}_n)$, so that it also holds a.s. for $B = \mathbb{R}^l$ and

$$f_n(b) = \widehat{\text{msq}}_n(b', b)(\tilde{\kappa}_n). \quad (343)$$

Thus, from remarks 161, 170, and Condition 147, Condition 151 holds. Using further Theorem 162 for f_n as in (343), conditions 152 and 153 hold as well.

In GMSM, let $A = \mathbb{R}^l$ and let us assume that the counterpart of Condition 135 for ECM holds for $S = Z^2$ (note that for $S = 1$ it holds automatically). Then, from the fifth point of Theorem 138 as well as from remarks 161 and 170, Condition 151 holds. Using further Theorem 162, conditions 152 and 153 hold as well.

Remark 172. Checking if G_t holds in possible practical realizations of GSSM or GSM methods, as discussed at the beginning of Remark 168, may be inconvenient. For instance, for $\widehat{\text{est}}_n = \widehat{\text{msq}}_n$ in the LETGS setting as above or $\widehat{\text{est}}_n$ as in (341) in the ECM setting, this typically cannot be done precisely due to numerical errors, and one has to make a rather arbitrary decision when such condition holds approximately. If $\epsilon_t(\omega) > 0$, $t \in T$, $\omega \in \Omega$, then, as already mentioned in Remark 168, in some cases one can avoid checking if G_t holds by performing minimization of g starting at x_0 as in that remark for each $\omega \in \Omega$. From Zoutendjik theorem (see Theorem 3.2 in [41]), for a number of line search minimization methods of a function g started at x_0 to be globally convergent, it is sufficient if g is bounded from below and continuously differentiable on some open neighbourhood $\mathcal{N} \subset B$ of the sublevel set $\mathcal{S} = \{x \in B : g(x) \leq g(x_0)\}$, and if ∇g is Lipschitz continuous on \mathcal{N} . In particular, it is sufficient if in addition to boundedness from below, g is twice differentiable and $\|\nabla^2 g\|$ is bounded on \mathcal{N} . One of methods for which this holds is gradient descent with step lengths satisfying Wolfe condition (see [41]). Note that from (202), (205), and $\|vv^T\| = \|v\|^2$, $v \in \mathbb{R}^l$, for $\omega \in \Omega_1^n$ and $K = \max_{i,j \in \{1, \dots, n\}} \|v_{j,i}(\omega)\|^2$, we have

$$\begin{aligned} \|\nabla_b^2 g_{\text{var},n}(b', b)(\omega)\| &\leq \sum_{i=1}^n (Z^2 L')(\omega_i) \sum_{j \in \{1, \dots, n\}, j \neq i} L'(\omega_j) \|v_{j,i}(\omega) v_{j,i}(\omega)^T\| \exp(b^T v_{j,i}(\omega)) \\ &\leq K g_{\text{var},n}(b', b)(\omega). \end{aligned} \tag{344}$$

Thus, for $\widehat{\text{est}}_n$ as in (341) it also holds $\|\nabla_b^2 \widehat{\text{est}}_n(b', b)(\omega)\| \leq K \widehat{\text{est}}_n(b', b)(\omega)$. From this it follows that for g as in Remark 168 corresponding to GSSM or GSM methods for such $\widehat{\text{est}}_n$, for some $\delta \in \mathbb{R}_+$, the assumptions of Zoutendjik theorem as above hold for each $x_0 \in \mathbb{R}^l$ and $\mathcal{N} = \{x \in \mathbb{R}^l : g(x) < g(x_0) + \delta\}$.

6.7 Helper theorems for proving convergence properties of multi-phase minimization methods

Theorem 173. Let $U \subset \mathbb{R}^l$ be an open ball with center b^* and $f : U \rightarrow \mathbb{R}$ be strongly convex with a constant $s \in \mathbb{R}_+$. Let $f_n : U \rightarrow \mathbb{R}$, $n \in \mathbb{N}_+$, be twice differentiable and such that $\nabla^2 f_n \rightrightarrows \nabla^2 f$. Then, for each $0 < m < s$, for sufficiently large n , f_n is strongly convex with constant m . Let further b^* be the minimum point of f and $\nabla f_n \rightrightarrows \nabla f$. Then, for sufficiently large n , f_n possess a unique minimum point a_n , which is equal to the unique point $x \in U$ for which $\nabla f_n(x) = 0$, and each $b \in U$ is a $\frac{1}{2m} |\nabla f_n(b)|^2$ -minimizer of f_n . Furthermore, $\lim_{n \rightarrow \infty} a_n = b^*$.

Proof. From Lemma 65, for sufficiently large n for which $\|f_n(x) - f(x)\|_\infty < s - m$, $x \in U$, we have $m_l(f_n(x)) > m$, $x \in U$, so that f_n is strongly convex with constant m . Let $h_n = f_n + f(b^*) - f_n(b^*)$, $n \in \mathbb{N}_+$. Then, $h_n(b^*) = f(b^*)$ and $\nabla h_n = \nabla f_n \rightrightarrows \nabla f$, so that $h_n \rightrightarrows f$. Furthermore, since $\nabla^2 h_n = \nabla^2 f_n$, $n \in \mathbb{N}_+$, h_n are strongly convex for sufficiently large n . Thus, from Remark 160 and Theorem 162, for sufficiently large n , h_n and thus also f_n possesses unique minimum point a_n and $\lim_{n \rightarrow \infty} a_n = b^*$. The rest of the thesis follows from the discussion at the end of Section 5.10. \square

Condition 174. $f : \mathbb{R}^l \rightarrow \mathbb{R}$ is continuous, functions $f_n : \mathbb{R}^l \rightarrow \mathbb{R}$, $n \in \mathbb{N}_+$, are such that $f_n \xrightarrow{loc} f$, and for a sequence $d_n \in \mathbb{R}^l$, $n \in \mathbb{N}_+$, we have

$$\lim_{n \rightarrow \infty} d_n = d^* \in \mathbb{R}^l. \quad (345)$$

We have a following easy-to-prove lemma.

Lemma 175. If Condition 174 holds, then $\lim_{n \rightarrow \infty} f_n(d_n) = f(d^*)$.

Theorem 176. Assuming condition 174, let for a bounded sequence $s_n \in \mathbb{R}^l$, $n \in \mathbb{N}_+$, it hold $f_n(s_n) \leq f_n(d_n)$, $n \in \mathbb{N}_+$. Then,

$$\limsup_{n \rightarrow \infty} f(s_n) \leq f(d^*). \quad (346)$$

Let further $d^* \in \mathbb{R}^l$ be the unique minimum point of f . Then,

$$\lim_{n \rightarrow \infty} f(s_n) = f(d^*). \quad (347)$$

If further f is convex, then

$$\lim_{n \rightarrow \infty} s_n = d^*. \quad (348)$$

Proof. Let $\epsilon > 0$. From boundedness of the set $D = \{s_n : n \in \mathbb{N}\}$ and $f_n \xrightarrow{loc} f$, let $N_1 \in \mathbb{N}_+$ be such that for $n \geq N_1$, $|f_n(x) - f(x)| < \frac{\epsilon}{2}$, $x \in D$. From Lemma 175, let $N_2 \geq N_1$ be such that for $n \geq N_2$, $|f_n(d_n) - f(d^*)| < \frac{\epsilon}{2}$. Then, for each $n \geq N_2$,

$$f(s_n) < f_n(s_n) + \frac{\epsilon}{2} \leq f_n(d_n) + \frac{\epsilon}{2} < f(d^*) + \epsilon, \quad (349)$$

which proves (346). Let d^* be the unique minimum point of f . Then, (347) follows from $f(s_n) \geq f(d^*)$, $n \in \mathbb{N}_+$, and (346). Let now f be convex and $\delta > 0$. Then, from continuity of f , there exists $x_0 \in S_l(d^*, \delta)$ such that $f(x_0) = \inf_{x \in S_l(0, \delta)} f(x)$. From uniqueness of d^* , $m = f(x_0) - f(d^*) > 0$. From convexity of f , for $x \in \mathbb{R}^l$ such that $|x - d^*| \geq \delta$ we have

$$f(x) - f(d^*) \geq \frac{|x - d^*|}{\delta} (f(d^* + \delta \frac{x - d^*}{|x - d^*|}) - f(d^*)) \geq m. \quad (350)$$

Thus, when (349) holds for $\epsilon \leq m$, then we must have $|s_n - d^*| < \delta$, which proves (348). \square

6.8 Two-phase minimization of estimators with gradient-based stopping criteria and constraints or function modifications

In this section we describe minimization methods in which two-phase minimization is used, in the first phase of which one can use some GM method as in Section 6.6 and in the second phase e.g. constrained minimization of an estimator or unconstrained minimization of a modified estimator, in both phases using gradient-based stopping criteria. The single- and multi-stage versions of these methods shall be abbreviated as CGSSM and CGMSM. We also discuss application of these minimization methods to the second mean square estimators in the LETGS setting and the inefficiency constant estimators in the ECM setting for $C = 1$.

Let us assume that $A = \mathbb{R}^l$ and that a following condition holds.

Condition 177. For some $\epsilon \in \mathbb{R}_+$, functions $g_1, g_2 : \mathbb{R}^l \rightarrow \mathcal{B}(\mathbb{R}^l)$ are such that for each $x \in \mathbb{R}^l$, $g_1(x)$ is open,

$$B_1(x, \epsilon) \subset g_1(x), \quad (351)$$

$$\overline{g_1(x)} \subset g_2(x), \quad (352)$$

and for each bounded set $B \subset \mathbb{R}^l$, the set $\bigcup_{x \in B} g_2(x)$ is bounded.

CGSSM and CGMSM will be defined as special cases of a following CGM method. Let us assume that for T as in Section 6.4, for each $t \in T$ we are given a $[0, \infty]$ -valued random variable $\tilde{\epsilon}_t$, an A -valued random variable d_t , a function $\tilde{d}_t : \Omega \rightarrow A$, and a random function $\tilde{f}_t : \mathcal{S}(A) \otimes (\Omega, \mathcal{F}) \rightarrow \mathcal{S}(\mathbb{R})$, such that $b \rightarrow \tilde{f}_t(b, \omega)$ is differentiable, $\omega \in \Omega$, we have

$$\tilde{d}_t \in g_2(d_t), \quad (353)$$

$$\tilde{f}_t(\tilde{d}_t) \leq \tilde{f}_t(d_t), \quad (354)$$

and if $\tilde{d}_t \in g_1(d_t)$, then

$$|\nabla_b \tilde{f}_t(\tilde{d}_t)| \leq \tilde{\epsilon}_t. \quad (355)$$

Remark 178. Functions \tilde{d}_t as above always exist, assuming that other variables as above are given. Indeed, without loss of generality let $\tilde{\epsilon}_t = 0$. Then, if \tilde{d}_t fulfilling (353), (354), and $\tilde{d}_t \notin g_1(d_t)$ does not exist, then \tilde{d}_t can be chosen to be a minimum point of $b \rightarrow \tilde{f}_t(b)$. Note that such \tilde{d}_t needs not to be a random variable.

For $k \in \mathbb{N}_p$, consider some functions $\tilde{\text{est}}_k$ as in (308) such that $b \rightarrow \tilde{\text{est}}_k(b', b)(\omega)$ is differentiable, $b' \in \mathbb{R}^l$, $\omega \in \Omega_1^k$.

Definition 179. CGSSM is defined as CGM in which for some GSSM as in Section 4.3.3 we take d_t as in it, and $\tilde{f}_t(b) = \mathbb{1}(N_t = k \in \mathbb{N}_p) \tilde{\text{est}}_k(b', b)(\tilde{\kappa}_k)$, $t \in T$. CGMSM is defined as CGM in which for some GMSM as in Section 4.3.3 (in particular, we take $T = \mathbb{N}_+$) we take d_k as in it and $\tilde{f}_k(b) = \tilde{\text{est}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k)$, $k \in \mathbb{N}_+$.

The following condition is needed e.g. if we want to investigate the asymptotic properties of \tilde{d}_t .

Condition 180. \tilde{d}_t , $t \in T$, are random variables (i.e. they are measurable functions from (Ω, \mathcal{F}) to $\mathcal{S}(\mathbb{R}^l)$).

Remark 181. Whenever dealing with some sets $D \in \Omega$ for which it is not immediately clear if $D \in \mathcal{F}$, when trying to prove that $\mathbb{P}(D) = 1$ and in particular $D \in \mathcal{F}$, we shall implicitly assume that we are working on a complete probability space, so that to achieve the goal it is sufficient to prove that $\mathbb{P}(E) = 1$ for some $E \in \mathcal{F}$ such that $E \subset D$. Such D will further typically appear when considering functions $g_t : \Omega \rightarrow \mathbb{R}^l$, like \tilde{d}_t above, without assuming that they are random variables. For instance, for some $b^* \in \mathbb{R}^l$, we will consider $D = \{\omega \in \Omega : \lim_{t \rightarrow \infty} g_t(\omega) \rightarrow b^*\}$ or $D = \{\omega \in \Omega : g_t(\omega) = b^* \text{ for sufficiently large } t\}$.

Condition 182. It holds $\tilde{\epsilon}_t(\omega) > 0$, $t \in T$, $\omega \in \Omega$, and we are given a function $R : \mathcal{S}(A) \rightarrow \mathcal{S}((\epsilon, \infty))$.

As the function R in the above condition one can take e.g. $R(x) = a|x| + b$ for some $a \in (0, \infty)$ and $b \in (\epsilon, \infty)$.

Remark 183. Let us assume Condition 182. Then, using e.g. boxes $g_1(x) = \{x \in \mathbb{R}^l : |x_i| \leq R(x), i = 1, \dots, l\}$, $g_2(x) = \bar{g}_1(x)$, $x \in A$, or appropriate balls in place of boxes, under some additional smoothness assumptions on $b \rightarrow \tilde{f}_t(\omega, b)$, $\omega \in \Omega$ (which in case of CGSSM and CGMSM reduce to appropriate such assumptions on $\tilde{\text{est}}_k$, $k \in \mathbb{N}_p$), \tilde{d}_t as above can be results of some constrained minimization methods of respective $\tilde{f}_t(b)$, started at d_t , constrained to $g_2(d_t)$, and stopped in the first point \tilde{d}_t in which the respective requirements for CGM as above are fulfilled. See e.g. [41, 12, 13] for examples of such constrained minimization algorithms (also called minimization methods with bounds when box constraints are used). In such case Condition 180 typically holds and can be proved using definition of the algorithm used.

Consider a following condition.

Condition 184. It holds $\delta \in \mathbb{R}_+$ and $h : A \rightarrow [0, \infty)$ is a twice continuously differentiable function such that $h(x) = 0$ for $x \in \bar{B}_1(0, 1)$ and $h(x) > 1$ for $|x| > 1 + \delta$.

An example of an easy to compute function fulfilling Condition 184 is $h(x) = 0$ for $|x| \leq 1$ and $h(x) = \frac{(|x|^2 - 1)^3}{\delta^3}$ for $|x| > 1$.

Remark 185. Let us assume conditions 182 and 184, and let \tilde{f}_t be nonnegative, $t \in T$. Let $g_1(x) = B_l(x, R(x))$ and $g_2(x) = \bar{B}_l(x, R(x)(1 + \delta))$, $x \in A$. Then, under some additional smoothness assumptions on $b \rightarrow \tilde{f}_t(b, \omega)$, $\omega \in \Omega$, rather than using constrained minimization as in Remark 183, to obtain \tilde{d}_t in CGM one can use some globally convergent unconstrained minimization method of a modified function

$$b \rightarrow h_t(b) = \tilde{f}_t(b) + \tilde{f}_t(d_t)h\left(\frac{|b - d_t|}{R(d_t)}\right). \quad (356)$$

Such method could start at d_t and stop in a first point \tilde{d}_t in which

$$h_t(\tilde{d}_t) \leq h_t(d_t) \quad (357)$$

and if $\tilde{d}_t \in g_1(d_t)$, then

$$|\nabla_b h_t(\tilde{d}_t)| \leq \tilde{\epsilon}_t. \quad (358)$$

Sufficient conditions for global convergence of a class of such minimization methods are given by Zoutendjik theorem, discussed in Remark 172. Such conditions are fulfilled in the above case if we have twice continuous differentiability of $b \rightarrow \tilde{f}_t(b, \omega)$ and h , which is why we assumed the latter in Condition 184.

Note that for \tilde{d}_t as above, if $\tilde{f}_t(d_t) = 0$, then from \tilde{f}_t being nonnegative, it holds $\nabla_b \tilde{f}_t(d_t) = 0$, and thus $\tilde{d}_t = d_t$ and we have (353). If $\tilde{f}_t(d_t) > 0$, then from (356) and Condition 184, $h_t(b) > h_t(d_t)$ for $|b - d_t| > (1 + \delta)R(d_t)$, and thus from (357) we also have (353). Furthermore, from (356), $\tilde{f}_t \leq h_t$, so that from (357) and $h_t(d_t) = \tilde{f}_t(d_t)$, we have (354). Finally, if $\tilde{d}_t \in g_1(d_t)$, then from (358) and from $\nabla_b h_t(x) = \nabla_b \tilde{f}_t(x)$, $x \in g_1(d_t)$, we have (355). Similarly as in Remark 183, Condition 180 typically holds for such constructed \tilde{d}_t .

Consider a following condition, which will be useful for proving asymptotic properties of minimization results of CGSSM, CGMSM, and some further methods.

Condition 186. *A. s. for sufficiently large t , (355) holds.*

The following theorem will be useful for proving convergence properties of CGM methods.

Theorem 187. *Let us assume that Condition 174 holds for $A = \mathbb{R}^l$ and f which is convex and has a unique minimum point d^* . Let $\tilde{d}_n \in g_2(d_n)$ be such that $f_n(\tilde{d}_n) \leq f_n(d_n)$, $n \in \mathbb{N}_+$. Then,*

$$\lim_{n \rightarrow \infty} \tilde{d}_n = d^* \quad (359)$$

and

$$\lim_{n \rightarrow \infty} f_n(\tilde{d}_n) = f(d^*). \quad (360)$$

Let further f be twice continuously differentiable with positive definite Hessian on A . Let f_n , $n \in \mathbb{N}_+$, be twice differentiable and whose i th derivatives for $i = 1, 2$, converge locally uniformly to such derivatives of f . Let $\epsilon_n \geq 0$, $n \in \mathbb{N}_+$, be such that $\lim_{n \rightarrow \infty} \epsilon_n = 0$. Let for $n \in \mathbb{N}_+$ it hold that if $\tilde{d}_n \in g_1(d_n)$ then

$$|\nabla f_n(\tilde{d}_n)| \leq \epsilon_n. \quad (361)$$

Let D be a bounded neighbourhood of d^* . Then, for sufficiently large n , (361) holds and $f_n|_D$ has a unique minimum point, which is equal to \tilde{d}_n if $\nabla f_n(\tilde{d}_n) = 0$.

Proof. From (345) and Condition 177, the set $\bigcup_{n \in \mathbb{N}_+} g(d_n)$ is bounded and so is the sequence $(\tilde{d}_n)_{n \in \mathbb{N}_+}$. Thus, (359) and (360) follow from Theorem 176. From (345) and (351), for sufficiently large n , $d_n \in \overline{B}_l(d^*, \frac{\epsilon}{2}) \subset g_1(d_n)$, and thus (361) holds. The rest of the thesis follows from Theorem 173, in which from Lemma 163 as U one can take any open ball with center d^* containing D , and as f and f_n in that theorem use restrictions to U of the above f and f_n . \square

For CGMSM in the below examples we assume conditions 130 and 131. By saying that counterparts of conditions 152 or 153 hold for CGSSM or CGMSM or that Condition 150 holds for CGMSM, we mean that these conditions hold for d_t and \hat{f}_t replaced by \tilde{d}_t and \tilde{f}_t , in counterpart of Condition 150 additionally assuming that Condition 180 holds.

Remark 188. *Note that we have a counterpart of Remark 155 with d_k replaced by \tilde{d}_k and conditions 152, 153, and 150 replaced by counterparts of such conditions for CGMSM.*

In the below CGSSM methods let us assume that Condition 104 holds for $S = Z^2$, while for CGMSM that Condition 135 holds for $S = Z^2$, in the LETGS setting additionally assuming these conditions for $S = 1$.

In the LETGS setting under Condition 62, let us consider CGSSM and CGMSM for $\widehat{\text{est}}_n = \widehat{\text{msq}}_n$ and $\widetilde{\text{est}}_n = \widetilde{\text{msq}}_n$. From Condition 152 holding for GSSM and GSM (see Section 6.6), Condition 147 and the fourth point of Theorem 129 for GSSM, the fifth point of Theorem 138 for GSM, as well as

from theorems 114, 187, and Remark 181, counterparts of conditions 152 and 153 and Condition 186 hold for CGSSM and CGMSM.

Let us now consider the ECM setting under Condition 171 and assuming that $C = 1$ as discussed in Remark 37, so that $\widehat{\text{ic}} = \widehat{\text{var}}$. Let $\widehat{\text{est}}_n = \widehat{\text{ic}}_n$ and $\widehat{\text{est}}_n = \widehat{\text{msq}}_n$ or $\widehat{\text{est}}_n = \widehat{\text{msq}}_n$, for the latter assuming (340). Then, from Condition 152 holding for GSSM and GSM, the fifth point of Theorem 129 for CGSSM, the sixth point of Theorem 138 for CGMSM, as well as theorems 113 and 187, counterparts of conditions 152 and 153 and Condition 186 hold for CGSSM and CGMSM.

6.9 Three-phase minimization of estimators with gradient-based stopping criteria and function modifications

In this section we define methods in which three-phase minimization with gradient-based stopping criteria and function modifications is used. In the first phase one performs some GM method as in Section 6.6, in the second a search of step lengths satisfying Wolfe conditions is carried out on a modified estimator, and in the third phase one can perform unconstrained minimization of the modified estimator using a gradient-based stopping criterion. The single- and multi-stage versions of these methods shall be abbreviated as MGSSM and MGMSM respectively. We also discuss application of such methods to the inefficiency constant estimators in the LETGS setting.

MGSSM and MGMSM will be defined as special cases of a following MGM method. Let Condition 184 hold, $0 < \alpha_1 < \alpha_2 < 1$, and $A = \mathbb{R}^l$. For T as in Section 6.4, for each $t \in T$, let d_t be an A -valued random variable and a random function $\tilde{f}_t : \mathcal{S}(A) \otimes (\Omega, \mathcal{F}) \rightarrow \mathcal{S}([0, \infty))$ be such that $b \rightarrow \tilde{f}_t(b, \omega)$ is continuously differentiable, $\omega \in \Omega$. Let $\tilde{\epsilon}_t$, $t \in T$, be as in the previous section and such that additionally a.s.

$$\lim_{t \rightarrow \infty} \tilde{\epsilon}_t = 0. \quad (362)$$

Let for each $t \in T$, r_t be an \mathbb{R}_+ -valued random variable,

$$h_t(b) = \tilde{f}_t(b) + \tilde{f}_t(b)h\left(\frac{|b - d_t|}{r_t}\right), \quad b \in A, \quad (363)$$

and a function $\tilde{d}_t : \Omega \rightarrow \mathbb{R}^l$ and a random variable d'_t be such that

$$\tilde{d}_t, d'_t \in \overline{B}_l(d_t, r_t(1 + \delta)). \quad (364)$$

For each $t \in T$, let for some $[0, \infty)$ -valued random variable p_t it hold

$$d'_t - d_t = -p_t \nabla h_t(d_t), \quad (365)$$

and let the following inequalities hold (which are Wolfe conditions on step length p_t when considering the steepest descent search direction, see e.g. (3.6) in [41])

$$h_t(d'_t) \leq h_t(d_t) - p_t \alpha_1 |\nabla h_t(d_t)|^2, \quad (366)$$

$$\nabla h_t(d'_t) \nabla h_t(d_t) \leq \alpha_2 |\nabla h_t(d_t)|^2. \quad (367)$$

Finally, let for each $t \in T$

$$h_t(\tilde{d}_t) \leq h_t(d'_t) \quad (368)$$

and

$$|\nabla h_t(\tilde{d}_t)| \leq \tilde{\epsilon}_t. \quad (369)$$

Let $\tilde{\text{est}}_k$, $k \in \mathbb{N}_p$, as in (308) be such that $b \rightarrow \tilde{\text{est}}_k(b', b)(\omega) \in [0, \infty)$ are continuously differentiable, $b' \in \mathbb{R}^l$, $\omega \in \Omega_1^k$, $k \in \mathbb{N}_p$. Then, MGSSM and MGMSM are defined as special cases of MGM method in the same way as CGSSM and CGMSM are defined as special cases of CGM in Definition 179.

Remark 189. For a given $t \in T$, d_t , and \tilde{f}_t , a possible construction of p_t , d'_t and \tilde{d}_t in MGM is as follows. If $\nabla h_t(d_t) = 0$, then we set $d'_t = \tilde{d}_t = d_t$. Consider further the case when $\nabla h_t(d_t) \neq 0$. Then, to obtain p_t and thus also d'_t , one can perform a line search of h_t in the steepest descent direction $-\nabla h_t(d_t)$ started in d_t and stopped when p_t and the corresponding d'_t start to satisfy Wolfe conditions (366) and (367). The line search can be performed e.g. using Algorithm 3.5 from [41]. If this algorithm is used for each $\omega \in \Omega$ such that $\nabla h_t(d_t)(\omega) \neq 0$, then such constructed d'_t is a random variable. When $\tilde{\epsilon}_t > 0$, then under some additional smoothness assumptions on $b \rightarrow \tilde{f}_t(b, \omega)$, $\omega \in \Omega$, to construct \tilde{d}_t one can use some convergent unconstrained minimization algorithm of h_t started in d'_t and stopped in a first point \tilde{d}_t in which (368) and (369) hold. See e.g. the assumptions of Zoutendjik theorem discussed in Remark 172. Note that if $\nabla h_t(d_t) \neq 0$, then $h_t(d_t) > 0$, and thus $h_t(x) > h_t(d_t)$ for $|x - d_t| > r_t(1 + \delta)$, so that from $h_t(\tilde{d}_t) \leq h_t(d'_t) \leq h_t(d_t)$, (364) holds. If $\tilde{\epsilon}_t(\omega) > 0$, $\omega \in \Omega$, and the same minimization algorithm is used for each $\omega \in \Omega$, then from the definition of the minimization method it typically follows that such constructed \tilde{d}_t is a random variable. On $\tilde{\epsilon}_t = 0$, \tilde{d}_t can be e.g. some (global) minimum point of h_t .

For $x \in \mathbb{R}^l$ and $B \subset \mathbb{R}^l$, let us denote

$$d(x, B) = \inf_{y \in B} |x - y|. \quad (370)$$

The following theorems will be useful for proving convergence properties of MGM methods.

Theorem 190. Let $K \subset \mathbb{R}^l$ be nonempty and compact and let $g_n : K \rightarrow \mathbb{R}$, $n \in \mathbb{N}_+$, converge uniformly to a continuous function $g : K \rightarrow \mathbb{R}$. Let m be the minimum of g and B be its set of minimum points. Then, for each sequence of points $d_n \in K$, $n \in \mathbb{N}_+$, such that $\lim_{n \rightarrow \infty} g_n(d_n) = m$, we have $\lim_{n \rightarrow \infty} d(d_n, B) = 0$.

Proof. Let $\epsilon \in \mathbb{R}_+$. From continuity of $x \in K \rightarrow d(x, B)$, $K_2 := \{x \in K : d(x, B) \geq \epsilon\}$ is a closed subset of K and thus it is compact. From continuity of g , for $w := \inf_{x \in K_2} g(x)$ we must have $\delta := w - m > 0$. For sufficiently large n for which $|g_n(x) - g(x)| < \frac{\delta}{2}$, $x \in K$, and $|g_n(d_n) - m| < \frac{\delta}{2}$, we have $|g(d_n) - m| \leq |g(d_n) - g_n(d_n)| + |g_n(d_n) - m| < \delta$ and thus $d_n \notin K_2$, i.e. $d(d_n, B) < \epsilon$. \square

Theorem 191. Let $f : \mathbb{R}^l \rightarrow \mathbb{R}$ and $f_n : \mathbb{R}^l \rightarrow \mathbb{R}$, $n \in \mathbb{N}_+$, be continuously differentiable and such that $f_n \xrightarrow{\text{loc}} f$ and $\nabla f_n \xrightarrow{\text{loc}} \nabla f$. Let further for some $d^* \in \mathbb{R}^l$, $s \in \mathbb{R}_+$, and $0 < w < r < \infty$ it hold

$$f(b) \geq f(d^*) + s, \quad b \in \mathbb{R}^l, \quad |b - d^*| \geq w, \quad (371)$$

and let $r_n \in \mathbb{R}_+$, $n \in \mathbb{N}_+$, be such that $\lim_{n \rightarrow \infty} r_n = r$. Let for a sequence $d_n \in \mathbb{R}^l$, $n \in \mathbb{N}_+$, it hold

$$\lim_{n \rightarrow \infty} d_n = d^*. \quad (372)$$

Let for each $n \in \mathbb{N}_+$, $h_n : \mathbb{R}^l \rightarrow \mathbb{R}$ be such that

$$h_n(b) = f_n(b) + f_n(d_n)h\left(\frac{|b - d_n|}{r_n}\right), \quad b \in \mathbb{R}^l. \quad (373)$$

Let $\epsilon_n \geq 0$, $n \in \mathbb{N}_+$, be such that $\lim_{n \rightarrow \infty} \epsilon_n = 0$, and let for each $n \in \mathbb{N}_+$, for some $p_n \in \mathbb{R}_+$, points $d'_n, \tilde{d}_n \in B_l(d_n, (1 + \delta)r_n)$ be such that

$$d'_n - d_n = -p_n \nabla h_n(d_n), \quad (374)$$

$$h_n(d'_n) \leq f_n(d_n) - p_n \alpha_1 |\nabla f_n(d_n)|^2, \quad (375)$$

$$\nabla h_n(d'_n) \nabla f_n(d_n) \leq \alpha_2 |\nabla f_n(d_n)|^2, \quad (376)$$

$$h_n(\tilde{d}_n) \leq h_n(d'_n), \quad (377)$$

and

$$|\nabla h_n(\tilde{d}_n)| \leq \epsilon_n. \quad (378)$$

Then, for sufficiently large n we have

$$d'_n, \tilde{d}_n \in B_l(d^*, w) \quad (379)$$

and

$$|\nabla f_n(\tilde{d}_n)| \leq \epsilon_n. \quad (380)$$

Let further

$$\phi(u) = \nabla f(d^* - u \nabla f(d^*)) \nabla f(d^*) - \alpha_2 |\nabla f(d^*)|^2, \quad u \in \mathbb{R}, \quad (381)$$

and $v = \inf\{u \geq 0 : \phi(u) = 0\}$. Then, if $|\nabla f(d^*)| = 0$ then $v = 0$ and if $|\nabla f(d^*)| \neq 0$ then $v \in (0, w)$. Furthermore, for

$$\mu = f(d^*) - v \alpha_1 |\nabla f(d^*)|^2 \quad (382)$$

we have

$$\limsup_{n \rightarrow \infty} f(d'_n) \leq \mu, \quad (383)$$

$$\limsup_{n \rightarrow \infty} f(\tilde{d}_n) \leq \mu, \quad (384)$$

for $E = \{x \in \mathbb{R}^l : f(x) \leq \mu\}$, we have

$$\lim_{n \rightarrow \infty} d(d'_n, E) = 0, \quad (385)$$

and for the set $D = \{x \in \mathbb{R}^l : \nabla f(x) = 0, f(x) \leq \mu\} \subset E$, containing the nonempty set of minimum points of f , we have

$$\lim_{n \rightarrow \infty} d(\tilde{d}_n, D) = 0. \quad (386)$$

Proof. Let $K = \overline{B}_l(d^*, w)$. Let $N_1 \in \mathbb{N}_+$ be such that for $n \geq N_1$, $|d_n - d^*| \leq \frac{r-w}{2}$ and $r - r_n \leq \frac{r-w}{2}$, in which case for $x \in K$ we have

$$|x - d_n| \leq |d^* - d_n| + |x - d^*| \leq \frac{r-w}{2} + w \leq \frac{r-w}{2} + w + \left(\frac{r-w}{2} - (r-r_n)\right) = r_n, \quad (387)$$

so that

$$K \subset \overline{B}_l(d_n, r_n). \quad (388)$$

From the set $F = \bigcup_{n=1}^{\infty} B_l(d_n, (1+\delta)r_n)$ being bounded, let $N_2 \geq N_1$ be such that for $n \geq N_2$

$$|f_n(x) - f(x)| \leq \frac{s}{2}, \quad x \in F. \quad (389)$$

From Lemma 175,

$$\lim_{n \rightarrow \infty} f_n(d_n) = f(d^*), \quad (390)$$

and thus let $N_3 \geq N_2$ be such that for $n \geq N_3$

$$|f_n(d_n) - f(d^*)| < \frac{s}{2}. \quad (391)$$

Then, for $n \geq N_3$ and $x \in F$ such that $|x - d^*| \geq w$, we have

$$h_n(x) \geq f_n(x) \geq f(x) - \frac{s}{2} \geq f(d^*) + \frac{s}{2} > f_n(d_n), \quad (392)$$

where in the first inequality we used (363) and Condition 184, in the second (389), in the third (371), and in the last (391). Thus, since from (375) and (377),

$$h_n(\tilde{d}_n) \leq h_n(d'_n) \leq f_n(d_n), \quad (393)$$

we must have (379). For $n \geq N_3$, from (388) and (379), we have $h_n(d'_n) = f_n(d'_n)$, $\nabla h_n(d'_n) = \nabla f_n(d'_n)$, and similarly for d'_n replaced by \tilde{d}_n , so that from (378), (380) holds, and from (393),

$$f_n(\tilde{d}_n) \leq f_n(d'_n) \leq f_n(d_n). \quad (394)$$

If $\nabla f(d^*) = 0$, then $v = 0$, in which case (383) and (384) follow from (394), the sequences $(d'_n)_{n \in \mathbb{N}_+}$ and $(\tilde{d}_n)_{n \in \mathbb{N}_+}$ being bounded, and Theorem 176. Let now $\nabla f(d^*) \neq 0$. Then, $\phi(0) = (1 - \alpha_2)|\nabla f(d^*)|^2 > 0$ and thus from continuity of ϕ , $v > 0$. The fact that $v < w$ follows from (371) and Lemma 3.1 in [41] about existence of steps $u > 0$ satisfying Wolfe conditions: $\phi(u) \leq 0$ and $f(d^* - \nabla f(d^*)v) \leq f(d^*) - u\alpha_1|\nabla f(d^*)|^2$. Let $1 < v' < v$. Then, from continuity of ϕ ,

$$\inf_{0 \leq u \leq v'} \phi(u) > 0. \quad (395)$$

For $n \in \mathbb{N}_+$, and $u \in \mathbb{R}$, let $\phi_n(u) = \nabla f_n(d_n - \nabla f_n(d_n)u) \nabla f(d^*) - \alpha_2 |\nabla f(d^*)|^2$. Since from Lemma 175

$$\lim_{n \rightarrow \infty} \nabla f_n(d_n) = \nabla f(d^*), \quad (396)$$

function $u \rightarrow d_n - u \nabla f_n(d_n)$ converges uniformly to $u \rightarrow d^* - u \nabla f(d^*)$ on $[0, v']$, and thus from Theorem 128, ϕ_n converges to ϕ uniformly on $[0, v']$. Thus, from

(395), let $N_4 \geq N_3$, be such that for $n \geq N_4$, $\inf_{u \in [0, v']} \phi_n(u) > 0$. For such n , from (374) and (376) it must hold $p_n > v'$ and from (375) we have

$$f_n(d'_n) \leq f_n(d_n) - v' \alpha_1 |\nabla f_n(d_n)|^2, \quad (397)$$

and thus

$$f(d'_n) \leq f(d'_n) - f_n(d'_n) + f_n(d_n) - v' \alpha_1 |\nabla f_n(d_n)|^2. \quad (398)$$

From (379) and the fact that f_n converges to f uniformly on K , we have $\lim_{n \rightarrow \infty} f(d'_n) - f_n(d'_n) = 0$. Thus, from (398), (390), and (396),

$$\limsup_{n \rightarrow \infty} f(d'_n) \leq f(d^*) - v' \alpha_1 |\nabla f(d^*)|^2. \quad (399)$$

Since this holds for each $v' < v$, we have (383), and from (394), we have (384). Due to (371), f attains minimum and for each its minimum point x_0 we have $\nabla f(x_0) = 0$ and from (383) and $f(x_0) \leq f(d'_n)$, $n \in \mathbb{N}_+$, we have $f(x_0) \leq \mu$. Thus, $x_0 \in D$. The minimum of $g := (f \vee \mu)|_K$ is μ and its set of minimum points is $E \subset K$. From (383), we have $\lim_{n \rightarrow \infty} f(d'_n) \vee \mu = \mu$, so that from (379) and Theorem 190 for such g and $g_n = g$, $n \in \mathbb{N}_+$, we receive (385). The minimum of $g := (|\nabla f| + f \vee \mu)|_K$ is μ and its set of minimum points is $D \subset K$. From $\epsilon_n \rightarrow 0$, (380), and (384), $\lim_{n \rightarrow \infty} |\nabla f_n(\tilde{d}_n)| + f(\tilde{d}_n) \vee \mu = \mu$. Thus, from (379) and Theorem 190 for such g and $g_n = (|\nabla f_n| + f \vee \mu)|_K$, $n \in \mathbb{N}_+$, we receive (386). \square

Let us discuss how MGSSM and MGMSM can be applied in the LETGS setting for $\widehat{\text{est}}_n = \widehat{\text{msq}}_n$ and $\widetilde{\text{est}}_n = \widehat{\text{ic}}_n$. We assume conditions 62, 91, and Condition 104 for $S = Z^2$. Then, from Theorem 114, msq has a unique minimum point d^* . Furthermore, for a positive definite matrix M and its lowest eigenvalue $m > 0$ as in that theorem, we have from (217) that

$$\text{var}(d^* + b) \geq \text{var}(d^*) + \frac{m}{2}|b|^2, \quad b \in \mathbb{R}^l, \quad (400)$$

and thus

$$\text{ic}(d^* + b) \geq c_{\min}(\text{var}(d^*) + \frac{m}{2}|b|^2), \quad b \in \mathbb{R}^l. \quad (401)$$

For some $\sigma_1, \sigma_2 \in \mathbb{R}_+$, $\sigma_1 < \sigma_2$, let us define

$$r = \sqrt{\frac{2}{m} \left(\frac{\text{ic}(d^*)}{c_{\min}} - \text{var}(d^*) \right)} + \sigma_2 \quad (402)$$

and

$$w = \sqrt{\frac{2}{m} \left(\frac{\text{ic}(d^*)}{c_{\min}} - \text{var}(d^*) \right)} + \sigma_1. \quad (403)$$

It holds $r > w > 0$ and from (401), for $b \in \mathbb{R}^l$, $|b| \geq w$,

$$\text{ic}(d^* + b) \geq c_{\min}(\text{var}(d^*) + \frac{mw^2}{2}) = \text{ic}(d^*) + \frac{m}{2}c_{\min}\sigma_1, \quad (404)$$

so that we have (371) for $f = \text{ic}$ and $s = \frac{m}{2}c_{\min}\sigma_1$.

Let us assume that Condition 104 holds for $S = C$ (in addition to this condition for $S = Z^2$ assumed above), so that from the fourth point of Theorem

112, ic is smooth. Let μ , E , and D be as in Theorem 191 for $f = \text{ic}$ and d^* as above. Note that we have $\mu < \text{ic}(d^*)$ only if $\nabla \text{ic}(d^*) \neq 0$, which from Remark 94 holds only if $\text{var}(d^*) \neq 0$ and $\nabla c(d^*) \neq 0$. Let for $n \in \mathbb{N}_+$ and $b \in \mathbb{R}^l$

$$\widehat{M}_n(b) = \left(2L(b)GZ^2 \exp\left(-\frac{1}{2} \sum_{i=1}^{\tau} |\eta_i|^2\right) \right)_n, \quad (405)$$

and let $\widehat{m}_n(b) = m_l(\widehat{M}_n(b))$ for m_l as in Section 5.5, i.e. $\widehat{m}_n(b)$ is the lowest eigenvalue of $\widehat{M}_n(b)$. For $n \in \mathbb{N}_2$, and $b, d \in \mathbb{R}^l$, we define $\widehat{r}_n(b, d) : \Omega_1^n \rightarrow \mathbb{R}$ to be such that for $\omega \in \Omega_1^n$ for which $\widehat{m}_n(b)(\omega) > 0$ and $\widehat{\text{ic}}(b, d)(\omega) - c_{\min} \widehat{\text{var}}_n(b, d)(\omega) > 0$

$$\widehat{r}_n(b, d)(\omega) = \sqrt{\frac{2}{\widehat{m}_n(b)(\omega)} \left(\frac{\widehat{\text{ic}}_n(b, d)(\omega)}{c_{\min}} - \widehat{\text{var}}_n(b, d)(\omega) \right)} + \sigma_2, \quad (406)$$

and otherwise $\widehat{r}_n(b, d)(\omega) = a$ for some $a \in \mathbb{R}_+$.

Let us further assume Condition 104 for $S = 1$. Then, from Theorem 129, a.s. $b \rightarrow \widehat{\text{var}}_n(b', b)(\tilde{\kappa}_n) \xrightarrow{\text{loc}} \text{var}$ and $b \rightarrow \widehat{\text{ic}}_n(b', b)(\tilde{\kappa}_n) \xrightarrow{\text{loc}} \text{ic}$. Thus, from the fact that for MGSSM, from discussion in Section 6.6 for GSSM, a.s. $d_t \rightarrow d^*$, from Lemma 175 and Condition 147, we have a.s. $\mathbb{1}(N_t = k \in \mathbb{N}_p) \widehat{\text{var}}_k(b', d_t)(\tilde{\kappa}_k) \rightarrow \text{var}(d^*)$ and $\mathbb{1}(N_t = k \in \mathbb{N}_p) \widehat{\text{ic}}_k(b', d_t)(\tilde{\kappa}_k) \rightarrow \text{ic}(d^*)$. Furthermore, from SLLN, a.s. $\widehat{M}_n(b')(\tilde{\kappa}_n) \rightarrow M$ and thus from Lemma 65, $\widehat{m}_n(b')(\tilde{\kappa}_n) \rightarrow m$. Therefore, for $r_t = \mathbb{1}(N_t = k \in \mathbb{N}_p) \widehat{r}_k(b', d_t)(\tilde{\kappa}_k)$, $t \in T$, we have a.s. $\lim_{t \rightarrow \infty} r_t = r$. Thus, from Theorem 191 and Remark 181 we receive that a following condition holds.

Condition 192. *Condition 186 holds, a.s. $\limsup_{t \rightarrow \infty} \text{ic}(d'_t) \leq \mu$, $\limsup_{t \rightarrow \infty} \text{ic}(\tilde{d}_t) \leq \mu$, $\lim_{t \rightarrow \infty} d(d'_t, E) = 0$, and*

$$\lim_{t \rightarrow \infty} d(\tilde{d}_t, D) = 0. \quad (407)$$

Furthermore, a.s. for sufficiently large t , $d'_t, \tilde{d}_t \in B_l(d^*, w)$.

For MGMSM let us assume that conditions 130 and 131 hold and that Condition 135 holds for $S = Z^2$, $S = C$, and $S = 1$. As discussed in Section 6.6 for GMSM, a.s. $d_k \rightarrow d^*$. From Theorem 138, a.s. $b \rightarrow \widehat{\text{var}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k) \xrightarrow{\text{loc}} \text{var}$ and $b \rightarrow \widehat{\text{ic}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k) \xrightarrow{\text{loc}} \text{ic}$. From Hölder inequality and Theorem 109 it easily follows that Condition 135 holds for S equal to different entries of $\widehat{M}_1(0)$. Thus, from the first point of Theorem 138 a.s. $\widehat{M}_{n_k}(b_{k-1})(\tilde{\chi}_k) \rightarrow M$, and thus $\widehat{m}_{n_k}(b_{k-1})(\tilde{\chi}_k) \rightarrow m$. Therefore, a.s. $\lim_{k \rightarrow \infty} \widehat{r}_k(b_{k-1}, d_k)(\tilde{\chi}_k) = r$. Thus, from Theorem 191 and Remark 181 it follows that Condition 192 holds for MGMSM.

Theorem 193. *Let functions var , c , and ic be as in Section 5.1 for A open. Let further var be lower semicontinuous and convex and have a unique minimum point $b^* \in A$, and let ic be continuous in b^* . Let for some $d_n \in A$, $n \in \mathbb{N}_+$,*

$$\limsup_{n \rightarrow \infty} \text{ic}(d_n) < \text{ic}(b^*). \quad (408)$$

Then,

$$\liminf_{n \rightarrow \infty} \text{var}(d_n) > \text{var}(b^*) \quad (409)$$

and

$$\limsup_{n \rightarrow \infty} c(d_n) < c(b^*). \quad (410)$$

Proof. For some $\text{ic}(b^*) > s > \limsup_{n \rightarrow \infty} \text{ic}(d_n)$, let $\epsilon \in \mathbb{R}_+$ be such that $\text{ic}(b) > s$ for $b \in B_l(b^*, \epsilon) \subset A$. Then, $d_n \in A \setminus B_l(b^*, \epsilon)$ for sufficiently large n . From semicontinuity of var , for some $b_0 \in S_l(b^*, \epsilon)$, $\text{var}(b_0) = \inf_{b \in S_l(b^*, \epsilon)} \text{var}(b) > \text{var}(b^*)$, and thus from convexity of var it holds $\text{var}(b) > \text{var}(b_0)$ for $b \in A \setminus B_l(b^*, \epsilon)$, and we have (409). Note that from (408), $\text{ic}(b^*) > 0$ and thus $\text{var}(b^*) > 0$. Therefore,

$$\limsup_{n \rightarrow \infty} c(d_n) \leq \frac{\limsup_{n \rightarrow \infty} \text{ic}(d_n)}{\liminf_{n \rightarrow \infty} \text{var}(d_n)} < \frac{\text{ic}(b^*)}{\text{var}(b^*)} = c(b^*). \quad (411)$$

□

If $\nabla \text{ic}(d^*) \neq 0$, so that $\mu < \text{ic}(d^*)$, then for $c_t = \tilde{d}_t$ or $c_t = d'_t$ as above for which we have a.s. $\limsup_{t \rightarrow \infty} \text{ic}(c_t) \leq \mu$, from Theorem 193 it also holds $\liminf_{t \rightarrow \infty} \text{var}(c_t) > \text{var}(d^*)$ and $\limsup_{t \rightarrow \infty} c(c_t) < c(d^*)$.

Condition 194. $D = \{b^*\}$ for some $b^* \in \mathbb{R}^l$.

Remark 195. Note that Condition 194 holds under assumptions as above e.g. if C is a positive constant or if $\text{var}(d^*) = 0$, and in both these cases $b^* = d^*$.

Remark 196. Let us assume Condition 194. Then, b^* is the unique minimum point of ic as above. Furthermore, under the above assumptions for MGSSM and MGMSM, from (407) and Lemma 175, counterparts of conditions 152 and 153 hold (by which we mean the same as above Remark 188).

Note that Remark 188 applies also to MGMSM.

6.10 Comparing the first-order asymptotic efficiency of minimization methods

Let $A \in \mathcal{B}(\mathbb{R}^l)$ be nonempty and $T \subset \mathbb{R}_+$ be unbounded. Consider a function $\phi : \mathcal{S}(A) \rightarrow \mathcal{S}(\overline{\mathbb{R}})$ and an A -valued stochastic process $d = (d_t)_{t \in T}$. For $t \in T$, d_t can be thought of as an adaptive random parameter trying to minimize ϕ for t being e.g. the simulation budget, the total number of steps in SSM methods, or the number of stages or simulations in MSM methods, used to compute d_t . We describe such possibilities in more detail in the below remark.

Remark 197. In various SSM methods, for some T as in Condition 147, we can consider d_t equal to d_t , \tilde{d}_t , or d'_t as in these methods, $t \in T$ (see also Remark 148). Let us further consider the case of MSM methods. Then, for variables p_k equal to d_k , \tilde{d}_k , or d'_k as in these methods, $k \in \mathbb{N}_+$, for some $\mathbb{N} \cup \{\infty\}$ -valued random variables N_t , $t \in T$, and some A -valued random variables p_0, p_∞ , one can set

$$d_t = p_{N_t} \mathbb{1}(N_t \neq \infty) + p_\infty \mathbb{1}(N_t = \infty), \quad t \in T. \quad (412)$$

The simplest choice would be to take $T = \mathbb{N}_+$ and $N_k = k$, so that $d_k = p_k$, $k \in \mathbb{N}_+$, i.e. k is the number of stages of MSM in which d_k is computed. If we want $t \in T$ to correspond to the total number of simulations performed to compute d_t , then for $s_k := \sum_{i=1}^k n_i$, $k \in \mathbb{N}_+$, and $T = \{s_k, k \in \mathbb{N}_+\}$, we can

take $N_{s_k} = k$, $k \in \mathbb{N}_+$. Alternatively, in MSM methods we can take $T = \mathbb{R}_+$ and for each $t \in T$, N_t can be the smallest number of stages using simulation budget t , or the highest such number before we exceed that budget. Let us discuss how one can model this. For some $[0, \infty)$ -valued random variables M_i modelling costs of minimization algorithms in the i th stage of MSM (we can set $M_i = 0$ if we do not want to consider them), $i \in \mathbb{N}_+$, and U being a nonnegative random variable on \mathcal{S}_1 modelling the cost of running a single simulation and doing helper computations for later estimator minimization similarly as for SSM in Remark 148, under Condition 119 we can take e.g.

$$N_t = \inf\{k \in \mathbb{N} : \sum_{i=1}^k (M_i + \sum_{j=1}^{n_i} U(\chi_{i,j})) \geq t\} \quad (413)$$

or

$$N_t = \sup\{k \in \mathbb{N} : \sum_{i=1}^k (M_i + \sum_{j=1}^{n_i} U(\chi_{i,j})) \leq t\}. \quad (414)$$

Note that if we have a.s. (7) and (10) and one of the following holds: a.s. $p_k \rightarrow b^*$, for some $f : A \rightarrow \mathbb{R}$ a.s. $f(p_k) \rightarrow f(b^*)$, or for some $m \in \mathbb{R}$, a.s. $\limsup_{k \rightarrow \infty} f(p_k) \leq m$, then we have respectively that a.s. $d_t \rightarrow b^*$ (compare with Remark 2), $f_{N_t}(d_t) \rightarrow f(b^*)$, or $\limsup_{t \rightarrow \infty} f_{N_t}(d_t) \leq m$. For N_t as in (413) or (414), (7) holds if $U < \infty$, $\mathbb{Q}(b)$ a.s., $b \in A$. Furthermore, (10) holds e.g. if Condition 92 holds for $C = U$, or from Theorem 122, if for some $K \in \mathcal{B}(A)$ such that $\inf_{b \in K} \mathbb{E}_{\mathbb{Q}(b)}(U) > 0$, Condition 118 holds for $K_i = K$, $i \in \mathbb{N}_+$, and the assumptions of Theorem 123 hold for $g(v, x) = U(x)$, $v \in A$, $x \in \Omega_1$.

For each $\overline{\mathbb{R}}$ -valued stochastic process $b = (b_t)_{t \in T}$, let us denote $\sigma_-(b) = \sup\{x \in \mathbb{R} : \lim_{t \rightarrow \infty} \mathbb{P}(b_t > x) = 1\}$ and $\sigma_+(b) = \inf\{x \in \mathbb{R} : \lim_{t \rightarrow \infty} \mathbb{P}(b_t < x) = 1\} = -\sigma_-(-b)$. Note that $\sigma_-(b) \leq \sigma_+(b)$ and $\sigma_-(b) = \sigma_+(b) = x \in \overline{\mathbb{R}}$ only if $b_t \xrightarrow{P} x$. Furthermore, if for some $x \in \overline{\mathbb{R}}$ a.s. $\limsup_{t \rightarrow \infty} b_t \leq x$, then $\sigma_+(b) \leq x$.

For $d = (d_t)_{t \in T}$ as above, let us denote $\phi(d) = (\phi(d_t))_{t \in T}$. For d' analogous as d we have $\sigma_-(\phi(d')) - \phi(d) \geq \sigma_-(\phi(d')) - \sigma_+(\phi(d))$. In particular, for each $\delta \in \mathbb{R}$ such that $\sigma_-(\phi(d')) - \sigma_+(\phi(d)) > \delta$, we have $\lim_{t \rightarrow \infty} \mathbb{P}(\phi(d'_t) - \phi(d_t) > \delta) = 1$, and such δ can be chosen positive if $\sigma_-(\phi(d')) > \sigma_+(\phi(d))$. Thus, we shall call d asymptotically not less efficient than d' for minimization of ϕ if $\sigma_+(\phi(d)) \leq \sigma_-(\phi(d'))$, and (asymptotically) more efficient for this purpose if this inequality is strict. If $\phi(d_t)$ and $\phi(d'_t)$ both converge in probability to the same real number, then d and d' shall be called equally efficient. We call such defined relations first-order asymptotic efficiency relations, to distinguish them from such second-order relations which will be defined in Section 7.2.

For instance, let for $d = (d_t)_{t \in T}$ denoting the parameters from single- or multi-stage minimization of some mean square estimators and d^* being the unique minimum point of mean square, it hold a.s. $d_t \rightarrow d^*$, and thus assuming further that ic is continuous in d^* , a.s. $\text{ic}(d_t) \rightarrow \text{ic}(d^*)$. Let further for $(d'_t)_{t \in T}$ denoting some parameters from minimization of inefficiency constant estimators it hold for μ as in Section 6.9 (for which $\mu \leq \text{ic}(d^*)$ and if $\nabla \text{ic}(d^*) \neq 0$, then $\mu < \text{ic}(d^*)$), that a.s. $\limsup_{t \rightarrow \infty} \text{ic}(d'_t) \leq \mu$. Then, d' is asymptotically not less efficient for minimization of ic than d and more efficient if $\nabla \text{ic}(d^*) \neq 0$. For MSM, under appropriate conditions as in Remark 197 we also have such

strong limits and asymptotic efficiency relations for the corresponding budget-dependent parameters.

Let now parameters $d = (d_t)_{t \in T}$ from single- or multi-stage minimization of cross-entropy estimators fulfill a.s. $d_t \rightarrow p^*$ for p^* being the unique minimum point of the cross-entropy. Let further $d' = (d'_t)_{t \in T}$ from minimization of mean square or inefficiency constant estimators for $C = 1$, fulfill a.s. $d_t \rightarrow b^*$ for b^* being the unique minimum point of msq, continuous in b^* . Then, d' is asymptotically not less efficient than d for minimization of msq, and more efficient if $b^* \neq p^*$.

6.11 Finding exactly the zero- or optimal-variance IS parameter

In this section we describe situations in which a.s. for sufficiently large t , minimization results d_t of our new estimators are equal to the zero-variance or optimal-variance IS parameter b^* as in Definition 26. This is the strongest form of convergence of such d_t to b^* considered in this work. When proving it below in the IS context we shall impose an assumption that we can find minimum or critical points of these estimators exactly. Even though such an assumption is unrealistic when minimization is performed using some iterative numerical methods, it is a frequent idealisation used to simplify the convergence analysis of stochastic counterpart methods, see e.g. [29, 51, 31]. Note also that such assumption is fulfilled for linearly parametrized control variates (at least when numerical errors occurring when solving a system of linear equations are ignored) when the zero-variance control variates parameter exists (see e.g. Chapter 5, Section 3 in [4]), in which case the below theory can be easily applied to prove that a.s. such parameter is found by the method after sufficiently many steps.

For a nonempty set $A \in \mathcal{B}(\mathbb{R}^l)$, consider a function

$$h : \mathcal{S}(A) \otimes \mathcal{S}_1 \rightarrow \mathcal{S}(\bar{\mathbb{R}}). \quad (415)$$

We will be most interested in cases corresponding to the below two conditions.

Condition 198. *Condition 18 holds and $h(b, \omega) = (ZL(b))(\omega)$, $\omega \in \Omega_1$, $b \in A$.*

Condition 199. *Condition 18 holds and $h(b, \omega) = (|Z|L(b))(\omega)$, $\omega \in \Omega_1$, $b \in A$.*

Let $b^* \in A$ and consider a family of distributions as in Section 3.4, satisfying Condition 22.

Condition 200. *For some $\beta \in \mathbb{R}_+$, \mathbb{Q}_1 a.s. (and thus also $\mathbb{Q}(b)$ a.s. for each $b \in A$)*

$$h(b^*, \cdot) = \beta. \quad (416)$$

Condition 201. *Condition 198 holds and b^* is the zero-variance IS parameter.*

Condition 202. *Condition 199 holds and b^* is the optimal-variance IS parameter.*

Remark 203. *From discussion in Section 3.2, under Condition 201, Condition 200 holds for $\beta = \alpha = \mathbb{E}_{\mathbb{Q}_1}(Z)$, and under Condition 202 — for $\beta = \mathbb{E}_{\mathbb{Q}_1}(|Z|)$.*

Remark 204. *If conditions 53 and 200 hold, then a.s.*

$$h(b^*, \kappa_i) = \beta, \quad i \in \mathbb{N}_+. \quad (417)$$

Lemma 205. *If conditions 119 and 200 hold, then a.s.*

$$h(b^*, \chi_{k,i}) = \beta, \quad i \in \{1, \dots, n_k\}, \quad k \in \mathbb{N}_+. \quad (418)$$

Proof. For each $k \in \mathbb{N}_+$ and $i \in \{1, \dots, n_k\}$

$$\mathbb{P}(h(b^*, \chi_{k,i}) = \beta) = \mathbb{E}(\mathbb{Q}(b_{k-1})(h(b^*, \cdot) = \beta)) = 1, \quad (419)$$

so that (418) holds a.s. as a conjunction of a countable number of conditions holding with probability one. \square

Let $D \in \mathcal{B}(A)$ be such that $b^* \in D$. For each $n \in \mathbb{N}_+$ and $\omega \in \Omega_1^n$, let us define

$$B_n(\omega) = \{b \in D : h(b, \omega_i) = h(b, \omega_j) \in \mathbb{R}_+, \quad i, j \in \{1, \dots, n\}\}. \quad (420)$$

For some $p \in \mathbb{N}_+$ and each $n \in \mathbb{N}_p$, let $\widehat{\text{est}}_n$ be as in (308). Consider a following condition.

Condition 206. *For each $n \in \mathbb{N}_p$ and $\omega \in \Omega_1^n$, if the set $B_n(\omega)$ is nonempty, then for each $d \in A$, $B_n(\omega)$ is the set of minimum points of $b \in D \rightarrow \widehat{\text{est}}_n(d, b)(\omega)$.*

Note that if Condition 206 holds, then it holds also for D replaced by its arbitrary subset (where D is replaced also in (420)).

Remark 207. *Let us assume Condition 23 and let $D = A$. It holds for $b', b \in A$*

$$\widehat{\text{msq}}_n(b', b) = \frac{1}{n^2} \sum_{i < j \in \{1, \dots, n\}} \frac{L'_i L'_j}{L_i(b) L_j(b)} (|Z_i| L_i(b) - |Z_j| L_j(b))^2 + ((|Z| |L'|)_n)^2. \quad (421)$$

Thus, under Condition 199, Condition 206 is satisfied for $p = 1$ and $\widehat{\text{est}}_n$ equal to $\widehat{\text{msq}}_n$ or msq_n (see (341) and (342)), or for $p = 2$ and $\widehat{\text{est}}_n$ equal to $\widehat{\text{var}}_n$ (which is positively linearly equivalent to $\widehat{\text{msq}}_n$ as discussed in Section 5.2). Assuming that $C > 0$, under Condition 198, Condition 206 is satisfied for $p = 2$ and $\widehat{\text{est}}_n = \widehat{\text{ic}}_n$ (see formulas (137) and (142)) or for $p = 3$ and $\widehat{\text{est}}_n = \widehat{\text{ic}}_n$ (see (144)).

Lemma 208. *If conditions 53 and 200 hold, then a.s. for each $k \in \mathbb{N}_p$, $b^* \in B_k(\tilde{\kappa}_k)$. If further Condition 206 holds, then for each $d \in A$, b^* is a minimum point of $b \in D \rightarrow \widehat{\text{est}}_k(d, b)(\tilde{\kappa}_k)$.*

Proof. It follows from Remark 204 and (420). \square

Condition 209. *Conditions 147 and 53 hold and functions $d_t : \Omega \rightarrow B$, $t \in T$, are such that a.s. for sufficiently large t , $b \in D \rightarrow \widehat{\text{est}}_{N_t}(b', b)(\tilde{\kappa}_{N_t})$ has a unique minimum point equal to d_t .*

Theorem 210. *If conditions 200, 206, and 209 hold, then a.s. for sufficiently large t , $d_t = b^*$ and $B_{N_t}(\tilde{\kappa}_{N_t}) = \{b^*\}$.*

Proof. It follows from Lemma 208 and Condition 209. \square

Lemma 211. *If Condition 119 holds for $n_k \in \mathbb{N}_p$, $k \in \mathbb{N}_+$, and Condition 200 holds, then a.s. for each $k \in \mathbb{N}_+$, $b^* \in B_{n_k}(\tilde{\chi}_k)$. If further Condition 206 holds, then for each $d \in A$, b^* is a minimum point of $b \in D \rightarrow \widehat{\text{est}}_{n_k}(d, b)(\tilde{\chi}_k)$.*

Proof. It follows from Lemma 205 and (420). \square

Condition 212. *Condition 119 holds for $n_k \in \mathbb{N}_p$, and functions $d_k : \Omega \rightarrow B$, $k \in \mathbb{N}_+$, are such that a.s. for sufficiently large k , $b \in D \rightarrow \widehat{\text{est}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k)$ has a unique minimum point equal to d_k .*

Theorem 213. *If conditions 200, 206, and 212 hold, then a.s. for sufficiently large k , $d_k = b^*$ and $B_{n_k}(\tilde{\chi}_k) = \{b^*\}$.*

Proof. It follows from Lemma 211 and Condition 212. \square

Let us consider the ECM setting and assume Condition 171 and that b^* is the optimal-variance IS parameter. Let us take $\widehat{\text{est}}_n = \widetilde{\text{msq}}_n$ as in (341), and $D = A$. As discussed in Section 6.6 for ESSM (that is for GSSM for $\epsilon_t = 0$, $t \in T$), conditions 151 and 152 hold. Thus, a.s. for sufficiently large t for which $\tilde{\kappa}_{N_t} \in \widetilde{D}_{N_t}$ and $d_t \in A$, $b \in D \rightarrow \widetilde{\text{msq}}_{N_t}(b', b)(\tilde{\kappa}_{N_t})$ has a unique minimum point equal to d_t , i.e. Condition 209 holds. Thus, from remarks 203, 207, and Theorem 210, a.s. for sufficiently large $t \in T$, $d_t = b^*$. For EMSM, under assumptions as in Section 6.6 which ensure that Condition 151 holds, Condition 212 holds and thus from remarks 203, 207, and Theorem 213, a.s. for sufficiently large $k \in \mathbb{N}_+$, $d_k = b^*$.

Let us now consider CGSSM and CGMSM in the LETGS setting for $\widetilde{\text{est}}_n = \widetilde{\text{msq}}_n$ and b^* being the optimal-variance IS parameter, or in the ECM setting for $\widetilde{\text{est}}_n = \widehat{\text{ic}}_n$, $C = 1$, and b^* being such zero-variance parameter. Let D be a bounded neighbourhood of b^* . Let us consider assumptions as in Section 6.8 for $\tilde{\epsilon}_t = 0$, $t \in T$ for appropriate T . Then, from Theorem 187 we receive that for $d_t = \tilde{d}_t$ and $\widehat{\text{est}}_n = \widetilde{\text{est}}_n$ Condition 209 holds for CGSSM and Condition 212 holds for CGMSM. Thus, from remarks 203, 207, and theorems 210 and 213, a.s. for sufficiently large $t \in T$, $\tilde{d}_t = b^*$ in CGSSM and CGMSM.

Let now b^* be the zero-variance IS parameter and consider MGSSM and MGMSM under assumptions as in Section 6.9 in the LETGS setting for $\widetilde{\text{est}}_n = \widehat{\text{ic}}_n$, $n \in \mathbb{N}_2$, and $\tilde{\epsilon}_t = 0$, $t \in T$. Then, from Remark 95, $\nabla^2 \text{ic}(b^*)$ is positive definite. Thus, from continuity of $\nabla^2 \text{ic}$, which follows from Theorem 112, and from Lemma 65, $\nabla^2 \text{ic}$ strongly convex on some open ball with center b^* . Therefore, from theorems 191 and 173, as well as remarks 195 and 196, conditions 209 and 212 hold for MGSSM and MGMSM respectively for $d_t = \tilde{d}_t$, $\widehat{\text{est}}_n = \widetilde{\text{est}}_n$, and $D \subset U$ being some neighbourhood of b^* . Therefore, by similar arguments as above, a.s. for sufficiently large t , $\tilde{d}_t = b^*$ in MGSSM and MGMSM.

7 Asymptotic properties of minimization methods

In this section we first discuss techniques for determining the asymptotic properties of minimization results of estimators as well as of functions of such results.

Then, we apply these techniques to minimization methods from the previous sections.

7.1 Helper theory for proving asymptotic properties of minimization results

Let $A \subset \mathbb{R}^l$ be an open set, $f : A \rightarrow \mathbb{R}$ be twice continuously differentiable, $b^* \in A$, and $H = \nabla^2 f(b^*)$.

Condition 214. $\nabla f(b^*) = 0$ and H is positive definite.

Condition 214 is implied e.g. by a following one.

Condition 215. H is positive definite and b^* is the unique minimum point of f .

Remark 216. Let us assume Condition 214. Then, from Lemma 65 and continuity of $\nabla^2 f$, for an open or closed ball $B \subset A$ with center b^* and sufficiently small positive radius, f is strongly convex on B and b^* is the unique minimum point of $f|_B$.

Let $T \subset \mathbb{R}_+$ be unbounded. Consider functions $f_t : \mathcal{S}(A) \otimes (\Omega, \mathcal{F}) \rightarrow \mathcal{S}(\mathbb{R})$, $t \in T$. Let $b \rightarrow f_t(b, \cdot)$ be a.s. twice continuously differentiable, $t \in T$. We shall denote $f_t(b) = f_t(b, \cdot)$ and $\nabla^i f_t(b) = \nabla_b^i f_t(b, \cdot)$, $i = 1, 2$.

Condition 217. For some neighbourhood $D \in \mathcal{B}(A)$ of b^* , $\nabla^2 f_t$ converges to $\nabla^2 f$ on D uniformly in probability (as $t \rightarrow \infty$), i.e. $\sup_{b \in D} \|\nabla^2 f_t(b) - \nabla^2 f(b)\|_\infty \xrightarrow{p} 0$.

Let $d_t, t \in T$, be A -valued random variables.

Condition 218. It holds $d_t \xrightarrow{p} b^*$.

For $g : T \rightarrow \mathbb{R}_+$, we write $X_t = o_p(g(t))$ if $\frac{X_t}{g(t)} \xrightarrow{p} 0$ (as $t \rightarrow \infty$). Let $r : T \rightarrow \mathbb{R}_+$ be such that

$$\lim_{t \rightarrow \infty} r(t) = \infty, \quad (422)$$

and let us denote $r_t = r(t)$.

Condition 219. For some nonnegative random variables $\delta_t, t \in T$, such that $\delta_t = o_p(r_t^{-1})$ and for some neighbourhood $U \in \mathcal{B}(A)$ of b^* , for the event A_t that d_t is a δ_t -minimizer of $f_t|_U$ (in particular, $d_t \in U$), we have

$$\lim_{t \rightarrow \infty} \mathbb{P}(A_t) = 1. \quad (423)$$

Condition 220. For some \mathbb{R}^l -valued random variable Y , $\sqrt{r_t} \nabla f_t(b^*) \Rightarrow Y$.

The below theorem is a consequence of Theorem 2.1 and discussion of its assumptions in sections 2 and 4 in [49], of implicit function theorem, and of the below Remark 223 (see also formula (4.14) and its discussion in [50]).

Theorem 221. Under conditions 214, 217, 218, 219, and 220, we have

$$\sqrt{r_t}(d_t - b^*) \Rightarrow -H^{-1}Y. \quad (424)$$

In particular, if $Y \sim \mathcal{N}(0, \Sigma)$ for some covariance matrix $\Sigma \in \mathbb{R}^{l \times l}$, then

$$\sqrt{r_t}(d_t - b^*) \Rightarrow \mathcal{N}(0, H^{-1}\Sigma H^{-1}). \quad (425)$$

We will need a following trivial remark.

Remark 222. Note that if for random variables \tilde{a}_t and a_t , $t \in T$, with probability tending to one (as $t \rightarrow \infty$) we have $a_t = \tilde{a}_t$, then for each $g : T \rightarrow \mathbb{R}$, $g(t)(\tilde{a}_t - a_t) \xrightarrow{p} 0$.

Remark 223. Theorem 2.1 in [49] uses $T = \mathbb{N}_+$ and $r_n = n$, $n \in T$, but its proof for general T and r as above is analogous. Let us assume conditions mentioned in Theorem 221. Let from Remark 216, B be a closed ball contained in sets D as in Condition 217 and in U as in Condition 219, and such that f is strongly convex on B and b^* is the unique minimum point of $f|_B$. From generalization of Theorem 2.1 in [49] to general T and r as above and discussion of assumptions of this theorem in [49], one easily receives the thesis (424) of Theorem 221 under additional assumptions that we have $d_t \in B$, $t \in T$, and Condition 219 holds with $A_t = \Omega$, $t \in T$. From Remark 222 and Condition 218, to prove Theorem 221 it is sufficient to prove (424) with d_t replaced by $\tilde{d}_t = \mathbb{1}(d_t \in B)d_t + \mathbb{1}(d_t \notin B)b^*$. For $C_t = A_t \cap \{d_t \in B\}$, let $\tilde{\delta}_t(\omega) = \delta_t(\omega)$, $\omega \in C_t$, and $\tilde{\delta}_t(\omega) = \infty$, $\omega \in C_t'$. Then, from Remark 222, Condition 218, and (423), the conditions of Theorem 221 are satisfied for d_t replaced by \tilde{d}_t and δ_t by $\tilde{\delta}_t$, and the above additional assumptions hold. Thus, (424) with d_t replaced by \tilde{d}_t follows from Theorem 2.1 in [49] as discussed above.

Condition 224. On some neighbourhood $K \in \mathcal{B}(A)$ of b^* , for $i = 1, 2$, the i th derivatives of $b \rightarrow f_t(b)$ (i.e. ∇f_t and $\nabla^2 f_t$) converge to such derivatives of f uniformly in probability.

Condition 224 is implied e.g. by the following one.

Condition 225. On some neighbourhood $K \in \mathcal{B}(A)$ of b^* , for $i = 1, 2$, a.s. the i th derivatives of $b \rightarrow f_t(b)$, converge uniformly to such derivatives of f .

Condition 226. It holds $|\nabla f_t(d_t)| = o_p(r_t^{-\frac{1}{2}})$.

Lemma 227. If conditions 214, 218, 224, and 226 hold, then Condition 219 holds.

Proof. From Remark 216, let U be an open ball with center b^* , contained in K as in Condition 224, and such that f is strongly convex on U with a constant $s \in \mathbb{R}_+$. Then, from Theorem 173, for each $0 < m < s$ there exist $\epsilon \in \mathbb{R}_+$ such that for each twice differentiable function $g : U \rightarrow \mathbb{R}$ for which

$$\sup_{x \in U} (|\nabla^2 f(x) - \nabla^2 g(x)|_\infty + |\nabla f(x) - \nabla g(x)|) < \epsilon, \quad (426)$$

each $b \in U$ is a $\frac{1}{2m}|\nabla g(b)|^2$ -minimizer of g . Thus, Condition 219 for the above U and $\delta_t = \frac{1}{2m}|\nabla f_t(d_t)|^2$ follows from conditions 218, 224, and 226. \square

From the above lemma we receive a following remark.

Remark 228. The assumption that conditions 217 and 219 hold in Theorem 221 can be replaced by the assumption that conditions 224 and 226 hold.

Theorem 229. *Let us assume conditions 214, 218, 224, and 226, and that a.s.*

$$\nabla f_t(b^*) = 0, \quad t \in T. \quad (427)$$

Then,

$$\sqrt{r_t}(d_t - b^*) \Rightarrow 0. \quad (428)$$

Proof. From (427), Condition 220 holds for $Y = 0$, so that the thesis follows from Remark 228 and Theorem 221. \square

Condition 230. *For some nonnegative random variables δ_t , $t \in T$, such that $\delta_t = o_p(r_t^{-\frac{1}{2}})$, with probability tending to one (as $t \rightarrow \infty$) we have*

$$|\nabla f_t(d_t)| \leq \delta_t. \quad (429)$$

Lemma 231. *Conditions 226 and 230 are equivalent.*

Proof. If Condition 226 holds, then Condition 230 holds for $\delta_t = |\nabla f_t(d_t)|$. Let us assume Condition 230. Then, for $\tilde{\delta}_t$ equal to δ_t if (429) holds and ∞ otherwise, we have $|\nabla f_t(d_t)| \leq \tilde{\delta}_t$ and from Remark 222, $\tilde{\delta}_t = o_p(r_t^{-\frac{1}{2}})$, from which Condition 226 follows. \square

7.2 Asymptotic properties of functions of minimization results and comparing the second-order asymptotic efficiency of minimization methods

Let us consider T and r as in the previous section. We shall further need the following theorem on delta method (see e.g. Theorem 3.1 and Section 3.3 in [52]).

Theorem 232. *Let $m, n \in \mathbb{N}_+$, let $D \in \mathcal{B}(\mathbb{R}^m)$ be a neighbourhood of $\theta \in \mathbb{R}^m$, and consider a function $\phi : D \rightarrow \mathbb{R}^n$. Let Y_t , $t \in T$, and Y be D -valued random variables such that we have $\sqrt{r_t}(Y_t - \theta) \Rightarrow Y$ (as $t \rightarrow \infty$). If ϕ is differentiable in θ with a differential $\phi'(\theta)$ (which we identify with its matrix), then*

$$\sqrt{r_t}(\phi(Y_t) - \phi(\theta)) \Rightarrow \phi'(\theta)Y. \quad (430)$$

If ϕ is twice differentiable in θ with the second differential $\phi''(\theta)$ and we have $\phi'(\theta) = 0$, then

$$r_t(\phi(Y_t) - \phi(\theta)) \Rightarrow \frac{1}{2}\phi''(\theta)(Y, Y). \quad (431)$$

Remark 233. *For $m \in \mathbb{N}_+$, let $\chi^2(m)$ denote the χ -squared distribution with m degrees of freedom. Let $m \in \mathbb{N}_+$, $S \sim \mathcal{N}(0, I_m)$, $B \in \text{Sym}_m(\mathbb{R})$, and $X = S^T B S$. Then, X has a special case of a generalized χ^2 distribution which we shall denote as $\tilde{\chi}^2(B)$. For B being a diagonal matrix $B = \text{diag}(v)$ for some $v \in \mathbb{R}^m$, $\tilde{\chi}^2(B)$ will be also denoted as $\tilde{\chi}^2(v)$. It holds $\mathbb{E}(X) = \text{Tr}(B)$. If $B = wI_m$ for some $w \in \mathbb{R}$, then we have $X \sim w\chi^2(m)$ (by which we mean that $X \sim wY$ for $Y \sim \chi^2(m)$). Let $\text{eig}(B) \in \mathbb{R}^m$ denote a vector of eigenvalues of B and let $\lambda = \text{eig}(B)$. Consider an orthogonal matrix $U \in \mathbb{R}^{m \times m}$ such that $B = U \text{diag}(\lambda) U^T$. Then, for $W = U^T S \sim \mathcal{N}(0, I_m)$ we have*

$$X = W^T \text{diag}(\lambda) W = \sum_{i=1}^m \lambda_i W_i^2, \quad (432)$$

and thus $\widetilde{\chi}^2(B) = \widetilde{\chi}^2(\lambda)$. Let $\Lambda = \{0\} \cup \{v \in (\mathbb{R} \setminus \{0\})^k : k \in \mathbb{N}_+, v_1 \leq v_2 \leq \dots \leq v_k\}$, i.e. Λ is the set of all real-valued vectors in different dimensions with ordered nonzero coordinates or having only one zero coordinate. Let for $v \in \mathbb{R}^m$, $\text{ord}(v) \in \Lambda$ be equal to $0 \in \mathbb{R}$ if $v = 0$ and otherwise result from ordering the coordinates of v in nondecreasing order and removing zero coordinates. Then, we have $\widetilde{\chi}^2(v) = \widetilde{\chi}^2(\text{ord}(v))$. For $Y \sim \chi^2(1)$ we have a moment-generating function $M_Y(t) := \mathbb{E}(\exp(tY)) = (1 - 2t)^{-\frac{1}{2}}$, $t < \frac{1}{2}$. Thus, for each $k \in \mathbb{N}_+$, $v \in \Lambda \cap \mathbb{R}^k$, $Y \sim \widetilde{\chi}^2(v)$, and $t \in \mathbb{R}$ such that $1 - 2v_i t > 0$, $i = 1, \dots, k$, we have $M_Y(t) = \prod_{i=1}^k (1 - 2v_i t)^{-\frac{1}{2}}$. Such M_Y is defined on some neighbourhood of 0 and it is different function for different $v \in \Lambda$. Thus, for $v_1, v_2 \in \Lambda$ such that $v_1 \neq v_2$, we have $\widetilde{\chi}^2(v_1) \neq \widetilde{\chi}^2(v_2)$. It follows that for two real symmetric matrices B_1, B_2 , we have $\widetilde{\chi}^2(B_1) = \widetilde{\chi}^2(B_2)$ only if $\text{ord}(\text{eig}(B_1)) = \text{ord}(\text{eig}(B_2))$.

Remark 234. Using notations as in Theorem 232, let $Y \sim \mathcal{N}(0, M)$ for some covariance matrix $M \in \mathbb{R}^{m \times m}$. Then, (430) implies that

$$\sqrt{r_t}(\phi(Y_t) - \phi(\theta)) \Rightarrow \mathcal{N}(0, \phi'(\theta)M\phi'(\theta)^T). \quad (433)$$

Let further $n = 1$. Then, (431) is equivalent to

$$r_t(\phi(Y_t) - \phi(\theta)) \Rightarrow R := \frac{1}{2}Y^T \nabla^2 \phi(\theta)Y. \quad (434)$$

For $S \sim \mathcal{N}(0, I_l)$ we have $Y \sim M^{\frac{1}{2}}S$. Thus, from Remark 233, for

$$B = \frac{1}{2}M^{\frac{1}{2}}\nabla^2 \phi(\theta)M^{\frac{1}{2}}, \quad (435)$$

we have $R \sim \widetilde{\chi}^2(B)$ and

$$\mathbb{E}(R) = \text{Tr}(B) = \frac{1}{2}\text{Tr}(\nabla^2 \phi(\theta)M). \quad (436)$$

Note that if θ is a local minimum point of ϕ , then $\nabla^2 \phi(\theta)$ is positive semidefinite and so is B .

Remark 235. If we have assumptions as in Theorem 221 leading to (425), then from Remark 234, for $M = H^{-1}\Sigma H^{-1}$ and

$$B = \frac{1}{2}M^{\frac{1}{2}}HM^{\frac{1}{2}} = \frac{1}{2}H^{-\frac{1}{2}}\Sigma H^{-\frac{1}{2}}, \quad (437)$$

we have

$$r_t(f(d_t) - f(b^*)) \Rightarrow \widetilde{\chi}^2(B). \quad (438)$$

Note that for $R \sim \widetilde{\chi}^2(B)$ it holds

$$\mathbb{E}(R) = \text{Tr}(B) = \frac{1}{2}\text{Tr}(\Sigma H^{-1}). \quad (439)$$

Let ϕ and $d = (d_t)_{t \in T}$ be as in Section 6.10 and such that for some probability μ on \mathbb{R} and $y \in \mathbb{R}$ we have

$$r_t(\phi(d_t) - y) \Rightarrow \mu. \quad (440)$$

Let further for some $d' = (d'_t)_{t \in T}$ and μ' analogous as above it hold $r_t(\phi(d'_t) - y) \Rightarrow \mu'$. Then, $\phi(d_t) \xrightarrow{P} y$ and similarly in the primed case, so that processes d and d' are equivalent from the point of view of first-order asymptotic efficiency for minimization of ϕ as discussed in Section 6.10. Their second-order asymptotic efficiency for this purpose can be compared by comparing the asymptotic distributions μ and μ' . For instance, if $\mu = \mu'$, then they can be considered equally efficient, while if for each $x \in \mathbb{R}$, $\mu((-\infty, x]) \geq \mu'((-\infty, x])$, then it is natural to consider the unprimed process to be not less efficient and more efficient if for some x this inequality is strict. The second-order asymptotic efficiency as above can be also compared using some moments like means or some quantiles like medians of the asymptotic distributions, where the process corresponding to lower such parameter can be considered more efficient. For $\mu = \widetilde{\chi}^2(B)$ and $\mu' = \widetilde{\chi}^2(B')$ for some symmetric matrices B and B' , which can arise e.g. from situations like in remarks 234 or 235, it may be convenient to compare the second-order efficiency of the corresponding processes using the means $\text{Tr}(B)$ and $\text{Tr}(B')$ of these distributions. In situations like in remarks 234 or 235, these means can be alternatively expressed by formulas (436) or (439), using which in some cases they can be estimated or even computed analytically (see e.g. Section 7.8).

Remark 236. *Under assumptions as above, let $\mu([0, \infty)) = 1$ and let for $X \sim \mu$ and $s \geq 1$ it hold $\mu' \sim sX$. Note that from Remark 233 this holds e.g. if for some symmetric positive semidefinite matrices B and B' we have $\mu = \widetilde{\chi}^2(B)$, $\mu' = \widetilde{\chi}^2(B')$, and $\text{ord}(\text{eig}(B')) = s \text{ord}(\text{eig}(B))$. If further $r_t = t$, $t \in T = \mathbb{R}_+$, then $t(\phi(d_t) - y)$ and $t(\phi(d'_{st}) - y)$ both converge in distribution to μ , but computing d'_{st} requires s times higher budget than d_t , $t \in T$ (assuming that such interpretation holds). Thus, the unprimed process can be called s times (asymptotically) more efficient for minimization of ϕ . Similarly, if $M' = sM$ for some nonzero covariance matrix $M \in \mathbb{R}^{l \times l}$, and for some $\theta \in A$, $\sqrt{t}(d_t - \theta) \Rightarrow \mathcal{N}(0, M)$ and $\sqrt{t}(d'_t - \theta) \Rightarrow \mathcal{N}(0, M')$, then d can be said to converge s times faster to θ than d' . In such case, if additionally ϕ is twice differentiable in θ with zero gradient and positive definite Hessian in this point, then from Remark 234, for B as in (435) we have $t(\phi(d_t) - y) \Rightarrow \widetilde{\chi}^2(B)$ and similarly for the primed process for $B' = sB$. Thus, the unprimed process is s times more efficient for minimization of ϕ .*

7.3 Some sufficient conditions for asymptotic properties in our methods

Consider a following composite condition consisting of conditions appearing in theorems 221, 229, and Remark 228.

Condition 237. *Conditions 214, 218, 224, and 226 hold.*

Let us discuss when, under appropriate identifications given below, Condition 237 holds for different variables in the LETGS and ECM settings for SSM methods from the previous sections, i.e. for ESSM, GSSM, CGSSM, and MGSSM, and for such MSM methods, i.e. for EMSM, GMSM, CGMSM, and MGMSM. We consider f equal to ce, msq, or ic, each defined on $A = \mathbb{R}^l$. We take T as for the minimization methods in the previous sections, in particular

for MSM methods $T = \mathbb{N}_+$. For EM and GM methods we take $f_t = \widehat{f}_t$ and d_t as in these methods, while for CGM and MGM methods $f_t = \widetilde{f}_t$ and $d_t = \widetilde{d}_t$, assuming Condition 180. Sufficient assumptions for Condition 215, implying Condition 214, to hold for f equal to ce in the ECM setting were provided in Section 5.3, and in the LETGS setting — in Section 5.7. From discussion in Section 5.10, for $A = \mathbb{R}^l$, Condition 215 follows from strong convexity of f , sufficient assumptions for which for f equal to msq or var in the LETGS setting were provided in Theorem 114. For f equal to msq in the ECM setting, sufficient conditions for it to have a unique minimum point were discussed in sections 5.8 (see e.g. Lemma 90) and 4.1, and for $\nabla^2 f$ to be positive definite — in Theorem 113. For $f = \text{ic}$, if C is a positive constant, then Condition 215 follows from such condition for $f = \text{var}$, and some other sufficient assumptions for Condition 214 to hold were discussed in Remark 95. From discussion in Section 6.3 we receive assumptions for which Condition 225, implying Condition 224, holds in the MSM methods as well as in the SSM methods for $T = \mathbb{N}_+$ and $N_k = t$, and thus from Condition 147 also in general case. For d_t as above, Condition 218 follows from Condition 152 and its counterparts.

Recall that from Lemma 231, conditions 226 and 230) are equivalent. For EM methods, if $b \rightarrow \widehat{\text{est}}_n(b', b)(\omega)$ is differentiable, $b' \in A$, $\omega \in \Omega_1^n$, $n \in \mathbb{N}_p$, and if Condition 151 holds, then Condition 230 holds in these methods even for $\delta_t = 0$, $t \in T$. For GM methods, if Condition 151 holds and $\epsilon_t = o_p(r_t^{-\frac{1}{2}})$ (which holds e.g. if $\epsilon_t = r_t^{-q}$ for some $\frac{1}{2} < q < \infty$), then Condition 230 holds for $\delta_t = \epsilon_t$. In CGM and MGM methods, if Condition 186 holds and $\tilde{\epsilon}_t = o_p(r_t^{-\frac{1}{2}})$, then Condition 230 holds for $\delta_t = \tilde{\epsilon}_t$.

7.4 Asymptotic properties of minimization results of the new estimators when the zero- or optimal-variance IS parameter exists

Condition 238. *We have assumptions as in Section 6.11, D is a neighbourhood of b^* , conditions 200 and 206 hold, and for each $b' \in A$, $k \in \mathbb{N}_p$, and $\omega \in \Omega_1^k$, $b \rightarrow \widehat{\text{est}}_k(b', b)(\omega)$ is differentiable in b^* .*

Theorem 239. *Let conditions 53, 147, and 238 hold, and let $b \in B \rightarrow f_t(b) := \mathbb{1}(N_t = k \in \mathbb{N}_p) \widehat{\text{est}}_k(b', b)(\tilde{\kappa}_k)$, $t \in T$. Then, a.s.*

$$\nabla f_t(b^*) = 0, \quad t \in T. \quad (441)$$

If further Condition 237 holds, then

$$\sqrt{r_t}(d_t - b^*) \Rightarrow 0, \quad t \rightarrow \infty. \quad (442)$$

Proof. From Lemma 208 we receive (441). Thus, (442) follows from Theorem 229. \square

Theorem 240. *Let conditions 119 and 238 hold, and let $b \in B \rightarrow f_k(b) := \widehat{\text{est}}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k)$, $k \in \mathbb{N}_+$. Then, for $T = \mathbb{N}_+$ we have a.s. (441). If further Condition 237 holds, then (442) holds.*

Proof. From Lemma 211 we have (441), so that (442) follows from Theorem 229. \square

Remark 241. Let conditions 18, 22, and 23 hold, A be a neighbourhood of b^* , and $b \rightarrow L(b)(\omega)$ be differentiable, $\omega \in \Omega_1$. Let $\widehat{\text{est}}_n$ be equal to $\widehat{\text{msq}}_n$ and b^* be the optimal-variance IS parameter, or $\widehat{\text{est}}_n$ be equal to $\widehat{\text{ic}}_n$ and b^* be the zero-variance IS parameter. Let further Conditions 53 and 147 hold for SSM or Condition 119 hold for MSM. Then, from remarks 203, 207, and the above theorems, we have (441) for f_t as in Theorem 239 for SSM, or as in Theorem 240 for MSM. If further Condition 237 holds (see Section 7.3 for some sufficient assumptions), then we also have (442) in these methods.

7.5 Asymptotic properties of SSM results

Let us assume that conditions 17, 18, 23, 53, and 147 hold, A is open, $b^* \in A$, and $b \in A \rightarrow L(b)(\omega)$ is differentiable, $\omega \in \Omega_1$. For some function $u \in A \rightarrow [0, \infty]$ such that $u(b') \in \mathbb{R}_+$, let us assume that

$$\frac{N_t}{t} \xrightarrow{p} \frac{1}{u(b')}. \quad (443)$$

Remark 242. Let N_t be given by some U as in Remark 148, and let $u(b) = \mathbb{E}_{\mathbb{Q}(b)}(U)$, $b \in A$. For such U being interpreted as the theoretical cost variable of a step of SSM as in Remark 148, $u(b')$ becomes such mean cost. If $u(b') \in \mathbb{R}_+$, then, as discussed in Section 2 (see (11)), we have a stronger fact than (443), namely that a.s. $\frac{N_t}{t} \rightarrow \frac{1}{u(b')}$. For the special case of $U = 1$, we have $u(b) = 1$, $b \in A$.

Below we shall prove that for g substituted by ce, msq, msq2, or ic, under appropriate assumptions, for some covariance matrix $\Sigma_g(b') \in \mathbb{R}^{l \times l}$ and

$$f_t(b) = \mathbb{1}(N_t = k \in \mathbb{N}_+) \widehat{g}_k(b', b)(\tilde{\kappa}_k), \quad t \in T, \quad (444)$$

we have

$$\sqrt{t} \nabla f_t(b^*) \Rightarrow \mathcal{N}(0, u(b') \Sigma_g(b')). \quad (445)$$

Remark 243. Let (445) hold for some g as above, and let Condition 237 hold for $r_t = t$ and for the minimized function f equal to msq if $g = \text{msq}2$, and to g otherwise. Then, from Theorem 221, denoting $H_f = \nabla^2 f(b^*)$ and

$$V_g(b') = H_f^{-1} \Sigma_g(b') H_f^{-1}, \quad (446)$$

we have

$$\sqrt{t}(d_t - b^*) \Rightarrow \mathcal{N}(0, u(b') V_g(b')). \quad (447)$$

Furthermore, from Remark 235, for

$$B_g(b') = \frac{1}{2} H_f^{-\frac{1}{2}} \Sigma_g(b') H_f^{-\frac{1}{2}}, \quad (448)$$

it holds

$$t(f(d_t) - f(b^*)) \Rightarrow \widetilde{\chi}^2(u(b') B_g(b')). \quad (449)$$

Let $u(b')$ be interpreted as the mean theoretical cost of a SSM step as in Remark 242. Let us consider different processes $d = (d_t)_{t \in T}$ from SSM methods for minimization of f as above, whose SSM methods have the same proportionality constant p_U of theoretical to approximate practical costs of SSM steps (see

Remark 148). Then, from discussion in Section 7.2 for $T = \mathbb{R}_+$ and $r_t = t$, $t \in T$, for $R \sim \widetilde{\chi}^2(u(b')B_g(b'))$, the second-order asymptotic efficiency of such processes can be compared using quantities

$$\mathbb{E}(R) = \frac{u(b')}{2} \text{Tr}(\Sigma_g(b')H_f^{-1}). \quad (450)$$

For instance, typically such quantities can be used for comparing the asymptotic efficiency of processes coming from SSM of different estimators or the same estimator for different b' . For different constants p_U , one can compare such quantities multiplied by such p_U .

Let $p(b) = \frac{1}{L(b)}$, consider likelihood function $l(b) = \ln(p(b)) = -\ln L(b)$, and score function

$$S(b) = \nabla l(b) = \frac{\nabla p(b)}{p(b)} = -\frac{\nabla L(b)}{L(b)}, \quad (451)$$

$b \in A$, where such terminology is used in maximum likelihood estimation, see e.g. [52]. Then,

$$\nabla_b \widehat{\text{ce}}_n(b', b) = -\overline{(ZL'S(b))}_n. \quad (452)$$

Thus, if for some $b^* \in A$

$$\mathbb{E}_{Q'}((ZL'S_i(b^*))^2) = \mathbb{E}_{Q_1}(L'(ZS_i(b^*))^2) < \infty, \quad i = 1, \dots, l, \quad (453)$$

(for which to hold in the LETS setting, from Theorem 110 and Remark 108, it is sufficient if Condition 104 holds for $S = Z^2$), and $\nabla \text{ce}(b^*) = -\mathbb{E}_{Q_1}(ZS(b)) = 0$, then, from Theorem 5, for

$$\Sigma_{\text{ce}}(b') = \mathbb{E}_{Q'}((ZL')^2 S(b^*) S(b^*)^T) = \mathbb{E}_{Q_1}(Z^2 L' S(b^*) S(b^*)^T), \quad (454)$$

we have (445) for $g = \text{ce}$ (and f_t as in (444) for such g).

It holds

$$\nabla_b \widehat{\text{msq}}_n(b', b) = \overline{(Z^2 L' \nabla L(b))}_n. \quad (455)$$

Thus, if

$$\mathbb{E}_{Q'}(Z^4 (L' \partial_i L(b^*))^2) = \mathbb{E}_{Q_1}(Z^4 L' (\partial_i L(b^*))^2) < \infty, \quad i = 1, \dots, l, \quad (456)$$

and

$$\nabla \text{msq}(b^*) = \mathbb{E}_{Q_1}(Z^2 \nabla L(b^*)) = 0, \quad (457)$$

then, from Theorem 5, for

$$\Sigma_{\text{msq}}(b') = \mathbb{E}_{Q_1}(L' Z^4 \nabla L(b^*) (\nabla L(b^*))^T), \quad (458)$$

we have (445) for $g = \text{msq}$.

Let us further in this section assume Condition 22. Consider the case of $g = \text{msq}$. Let $R_n(b', b) = \left(\frac{L'}{L(b)}\right)_n$, $n \in \mathbb{N}_+$, so that

$$\nabla_b \widehat{\text{msq}}_n 2_n = (\nabla_b \widehat{\text{msq}}_n) R_n + \widehat{\text{msq}}_n \nabla_b R_n. \quad (459)$$

Let

$$\begin{aligned} T_t(b') &= \sqrt{t} \mathbb{1}(N_t = k \in \mathbb{N}_+) (\nabla_b \widehat{\text{msq}}_k(b', b^*) + \text{msq}(b^*) \nabla_b R_k(b', b^*)) (\tilde{\kappa}_k) \\ &= \sqrt{t} \mathbb{1}(N_t = k \in \mathbb{N}_+) \overline{(L' \nabla L(b^*) (Z^2 - \text{msq}(b^*) L(b^*)^{-2}))}_k (\tilde{\kappa}_k) \end{aligned} \quad (460)$$

and

$$\begin{aligned}
Z_t(b') &= \sqrt{t} \mathbb{1}(N_t = k \in \mathbb{N}_+) \nabla_b \widehat{\text{msq}}_k(b', b^*)(\tilde{\kappa}_k) - T_t(b') \\
&= \mathbb{1}(N_t = k \in \mathbb{N}_+) (((R_k - 1) \sqrt{t} \nabla_b \widehat{\text{msq}}_k(b', b^*) \\
&\quad + (\widehat{\text{msq}}_k(b', b^*) - \text{msq}(b^*)) \sqrt{t} \nabla_b R_k(b', b^*)) (\tilde{\kappa}_k).
\end{aligned} \tag{461}$$

Let

$$0 = \nabla_b \mathbb{E}_{\mathbb{Q}'}(L'L^{-1}(b^*)) = -\mathbb{E}_{\mathbb{Q}'}(L'L^{-2}(b^*) \nabla L(b^*)) \tag{462}$$

(see e.g. the first point of Theorem 112 for sufficient conditions for this in the LETS setting) and $\mathbb{E}_{\mathbb{Q}_1}(L'L(b)^{-4}(\partial_i L(b))^2) < \infty$, $i = 1, \dots, l$. Then, from Theorem 5,

$$\sqrt{t} \mathbb{1}(N_t = k \in \mathbb{N}_+) \nabla_b R_k(b', b)(\tilde{\kappa}_k) \Rightarrow \mathcal{N}(0, u(b') \mathbb{E}_{\mathbb{Q}_1}(L'L(b)^{-4} \nabla L(b)(\nabla L(b))^T)). \tag{463}$$

Assuming further (456) and (457), from (445) for $g = \text{msq}$, (140), (463), the fact that from SLLN and Condition 147, a.s. $\mathbb{1}(N_t = k \in \mathbb{N}_+) \widehat{\text{msq}}_k(b', b^*)(\tilde{\kappa}_k) \rightarrow \text{msq}(b^*)$, as well as from the last line of (461) and Slutsky's lemma,

$$Z_t(b') \xrightarrow{P} 0. \tag{464}$$

Let

$$\Sigma_{\text{msq}2}(b') = \mathbb{E}_{\mathbb{Q}_1}((Z^2 - \text{msq}(b^*)L(b^*)^{-2})^2 L' \nabla L(b^*)(\nabla L(b^*))^T). \tag{465}$$

From Theorem 5, $T_t(b') \Rightarrow \mathcal{N}(0, u(b') \Sigma_{\text{msq}2}(b'))$, so that from (464), the first line of (461), and Slutsky's lemma, we receive (445).

Let us finally consider the case of $g = \text{ic}$. We have for $n \in \mathbb{N}_2$

$$\nabla_b \widehat{\text{ic}}_n = (\nabla_b \widehat{c}_n) \widehat{\text{var}}_n + \widehat{c}_n \nabla_b \widehat{\text{var}}_n, \tag{466}$$

where

$$\nabla_b \widehat{\text{var}}_n(b', b) = \frac{n}{n-1} ((\nabla_b \widehat{\text{msq}}_n(b', b)) \overline{(L'L^{-1}(b))}_n + \widehat{\text{msq}}_n(b', b) \overline{(L'L(b)^{-2} \nabla L(b))}_n). \tag{467}$$

Let for $D = (\mathbb{R} \times \mathbb{R}^l)^3 \times \mathbb{R}$ and $n \in \mathbb{N}_+$, $U_n(b') : \Omega_1^n \rightarrow D$ be equal to

$$(\widehat{c}_n(b', b^*), \nabla_b \widehat{c}_n(b', b^*), \widehat{\text{msq}}_n(b', b^*), \nabla_b \widehat{\text{msq}}_n(b', b^*), \overline{(L'L^{-1}(b^*))}_n, \overline{(L'L(b^*)^{-2} \nabla L(b^*))}_n, \overline{(ZL')}_n) \tag{468}$$

and let

$$\theta := (c(b^*), \nabla c(b^*), \text{msq}(b^*), \nabla \text{msq}(b^*), 1, 0, \alpha) = \mathbb{E}_{\mathbb{Q}'}(U_1(b')) \in D. \tag{469}$$

Let the coordinates of $U_1(b')$ be square-integrable under \mathbb{Q}' and

$$\Psi := \mathbb{E}_{\mathbb{Q}'}((U_1(b') - \theta)(U_1(b') - \theta)^T). \tag{470}$$

Then, from Theorem 5,

$$\sqrt{t} \mathbb{1}(N_t = k \in \mathbb{N}_+) (U_k(\tilde{\kappa}_k) - \theta) \Rightarrow \mathcal{N}(0, u(b') \Psi). \tag{471}$$

For $\phi : D \rightarrow \mathbb{R}^l$ such that

$$\phi((x_i)_{i=1}^7) = x_2(x_3x_5 - x_7^2) + x_1(x_4x_5 + x_3x_6), \tag{472}$$

we have for $n \in \mathbb{N}_2$

$$\nabla_{b\widehat{\text{ic}}_n}(b', b^*) = \frac{n}{n-1}\phi(U_n(b')) \quad (473)$$

and $\nabla \text{ic}(b^*) = \frac{n}{n-1}\phi(\theta)$. Let us assume that

$$\nabla \text{ic}(b^*) = \phi(\theta) = 0. \quad (474)$$

Using delta method from Theorem 232, as well as Remark 234 and (471), we receive that for

$$\Sigma_{\text{ic}}(b') = \phi'(\theta)\Psi(\phi'(\theta))^T \quad (475)$$

we have

$$\sqrt{t}\mathbb{1}(N_t = k \in \mathbb{N}_+) \phi(U_k(b')(\tilde{\kappa}_k)) \Rightarrow \mathcal{N}(0, u(b')\Sigma_{\text{ic}}(b')). \quad (476)$$

From $\lim_{n \rightarrow \infty} \frac{n}{n-1} = 1$, (473), (476), and Slutsky's lemma, we thus have (445). From (470) and (475), for $W(b') := \phi'(\theta)(U_1(b') - \theta)$

$$\Sigma_{\text{ic}}(b') = \mathbb{E}_{Q'}(W(b')W(b')^T). \quad (477)$$

We have

$$\begin{aligned} \phi'(\theta)((x_i)_{i=1}^7) &= \nabla \text{msq}(b^*)x_1 + \text{var}(b^*)x_2 + \nabla c(b^*)x_3 \\ &\quad + c(b^*)x_4 + (\nabla c(b^*) \text{msq}(b^*) + c(b^*)\nabla \text{msq}(b^*))x_5 \\ &\quad + c(b^*) \text{msq}(b^*)x_6 - 2\nabla c(b^*)\alpha x_7, \end{aligned} \quad (478)$$

so that

$$\begin{aligned} W(b') &= \nabla \text{msq}(b^*)(CL'L(b^*)^{-1} - c(b^*)) + \text{var}(b^*)(-CL'L^{-2}(b^*)\nabla L(b^*) - \nabla c(b^*)) \\ &\quad + \nabla c(b^*)(Z^2L'L(b^*) - \text{msq}(b^*)) + c(b^*)(Z^2L'\nabla L(b^*) - \nabla \text{msq}(b^*)) \\ &\quad + (\nabla c(b^*) \text{msq}(b^*) + c(b^*)\nabla \text{msq}(b^*))(L'L^{-1}(b^*) - 1) \\ &\quad - c(b^*) \text{msq}(b^*)L'L(b^*)^{-2}\nabla L(b^*) - 2\nabla c(b^*)\alpha(ZL' - \alpha). \end{aligned} \quad (479)$$

Remark 244. Note that for the special case of $C = 1$, we have $c(b) = 1$, $\nabla c(b) = 0$, $\text{ic}(b) = \text{var}(b)$, $b \in A$, and from (474), $\nabla \text{msq}(b^*) = \nabla \text{var}(b^*) = 0$, so that from (479),

$$W(b') = L'\nabla L(b^*)(Z^2 - (\text{var}(b^*) + \text{msq}(b^*))L(b^*)^{-2}), \quad (480)$$

and thus

$$\Sigma_{\text{ic}}(b') = \mathbb{E}_{Q_1}(L'(Z^2 - (\text{var}(b^*) + \text{msq}(b^*))L(b^*)^{-2})^2\nabla L(b^*)(\nabla L(b^*))^T). \quad (481)$$

7.6 A helper CLT

Consider a family of probability distributions as in Section 3.4, let $m \in \mathbb{N}_+$, $u : \mathcal{S}(A) \otimes \mathcal{S}_1 \rightarrow \mathcal{S}(\mathbb{R}^m)$, and $B \in \mathcal{B}(A)$.

Condition 245. For

$$f(b, M) := \mathbb{E}_{Q(b)}(|u(b, \cdot)|\mathbb{1}(|u(b, \cdot)| > M)), \quad b \in B, M \in \mathbb{R}, \quad (482)$$

and $R(M) := \sup_{b \in B} f(b, M)$, $M \in \mathbb{R}$, we have

$$\lim_{M \rightarrow \infty} R(M) = 0. \quad (483)$$

Note that the above condition is equivalent to saying that for random variables $\psi_b \sim \mathbb{Q}(b)$, $b \in B$, the family $\{|u(b, \psi_b)| : b \in B\}$ is uniformly integrable. In particular, similarly as for uniform integrability, using Hölder inequality one can prove a following criterion for the above condition to hold.

Lemma 246. *If for some $p > 1$,*

$$\sup_{b \in B} \mathbb{E}_{\mathbb{Q}(b)}(|u(b, \cdot)|^p) < \infty, \quad (484)$$

then Condition 245 holds.

For some $T \in \mathbb{N}_+ \cup \infty$, $\bar{\mathbb{R}}$ -valued random variables $(\psi_i)_{i=1}^T$ are said to be martingale differences for a filtration $(\mathcal{F}_i)_{i=0}^T$, if $M_n = \sum_{i=1}^n \psi_i$, $n \in \mathbb{N}$, (by which we mean that $M_0 = 0$) is a martingale for $(\mathcal{F}_i)_{i=0}^T$, that is if $\mathbb{E}(|\psi_i|) < \infty$, ψ_i is \mathcal{F}_i -measurable, and $\mathbb{E}(\psi_i | \mathcal{F}_{i-1}) = 0$, $i = 1, \dots, T$. The following martingale CLT is a special case of Theorem 8.2 with conditions II, page 442 in [39].

Theorem 247. *For each $n \in \mathbb{N}_+$, let $m_n \in \mathbb{N}_+$, $(\mathcal{F}_{n,k})_{k=0}^{m_n}$ be a filtration, and $(\psi_{n,k})_{k=1}^{m_n}$ be martingale differences for it such that $\mathbb{E}(\psi_{n,k}^2) < \infty$, $k = 1, \dots, m_n$. Let further*

1. *for each $\delta > 0$, $\sum_{k=1}^{m_n} \mathbb{E}(\psi_{n,k}^2 \mathbb{1}(|\psi_{n,k}| > \delta) | \mathcal{F}_{n,k-1}) \xrightarrow{p} 0$ (as $n \rightarrow \infty$),*
2. *for some $\sigma \in [0, \infty)$, $\sum_{k=1}^{m_n} \mathbb{E}(\psi_{n,k}^2 | \mathcal{F}_{n,k-1}) \xrightarrow{p} \sigma^2$.*

Then,

$$S_n = \sum_{k=1}^{m_n} \psi_{n,k} \Rightarrow \mathcal{N}(0, \sigma^2). \quad (485)$$

Let $r : \mathcal{S}(A) \otimes \mathcal{S}_1 \rightarrow \mathcal{S}(\mathbb{R}^m)$ be such that for each $b \in A$,

$$\mathbb{E}_{\mathbb{Q}(b)}(r(b, \cdot)) = 0. \quad (486)$$

Consider a matrix

$$\Sigma(b) = \mathbb{E}_{\mathbb{Q}(b)}(r(b, \cdot)r(b, \cdot)^T) \quad (487)$$

for $b \in A$ for which it is well-defined. Note that under Condition 245 for $u = |r|^2$, from (483) we have $R(M) < \infty$ for some $M > 0$, and thus $R(0) \leq M + R(M) < \infty$ and $\Sigma(b) \in \mathbb{R}^{m \times m}$, $b \in B$.

We will be most interested in the IS case in which we shall assume a following condition.

Condition 248. *For some \mathbb{R}^m -valued random variable Y on \mathcal{S}_1 , Condition 17 holds for Z replaced by Y , and we have*

$$r(b, \omega) = (YL(b))(\omega), \quad b \in A, \quad \omega \in \Omega_1. \quad (488)$$

Theorem 249. *Let us assume that Condition 119 holds and we have*

$$\lim_{k \rightarrow \infty} n_k = \infty. \quad (489)$$

Let further B as above be a neighbourhood of $b^ \in A$,*

$$b_n \xrightarrow{p} b^*, \quad (490)$$

Condition 245 hold for $u = |r|^2$, and $b \in B \rightarrow \Sigma(b)$ be continuous in b^ . Then, for $W_k := \frac{1}{\sqrt{n_k}} \sum_{i=1}^{n_k} r(b_{k-1}, \chi_{k,i})$, $k \in \mathbb{N}_+$, we have*

$$W_k \Rightarrow \mathcal{N}(0, \Sigma(b^*)). \quad (491)$$

Proof. For $D_k := \mathbb{1}(b_{k-1} \in B)W_k$, from (490) we have $W_k - D_k = \mathbb{1}(b_{k-1} \notin B)W_k \xrightarrow{p} 0$, so that from Slutsky's lemma it is sufficient to prove that $D_k \Rightarrow \mathcal{N}(0, \Sigma(b^*))$. Furthermore, using Cramér-Wold device it is sufficient to prove that for each $t \in \mathbb{R}^l$, for

$$v(b) = t^T \Sigma(b) t, \quad b \in \mathbb{R}^l, \quad (492)$$

and $S_k := t^T D_k$, we have $S_k \Rightarrow \mathcal{N}(0, v(b^*))$. For $t = 0$ this is obvious, so let us consider $t \neq 0$. It is sufficient to check that the assumptions of Theorem 247 hold for $m_k = n_k$, $\mathcal{F}_{k,i} = \sigma(b_{k-1}; \chi_{k,j} : j \leq i)$, $\psi_{k,i} = \frac{1}{\sqrt{n_k}} \mathbb{1}(b_{k-1} \in B) t^T r(b_{k-1}, \chi_{k,i})$, and $\sigma^2 = v(b^*)$. From Condition 245 (for $u = |r|^2$), we have

$$\begin{aligned} \mathbb{E}(\psi_{k,i}^2) &= \frac{1}{n_k} \mathbb{E}((\mathbb{E}_{\mathbb{Q}(b)}(\mathbb{1}(b \in B)(t^T r(b, \cdot))^2))_{b=b_{k-1}}) \\ &\leq \frac{|t|^2}{n_k} \mathbb{E}((\mathbb{E}_{\mathbb{Q}(b)}(\mathbb{1}(b \in B)|r(b, \cdot)|^2))_{b=b_{k-1}}) \\ &\leq \frac{|t|^2}{n_k} \mathbb{E}(\mathbb{1}(b_{k-1} \in B) f(b_{k-1}, 0)) \\ &\leq \frac{|t|^2}{n_k} R(0) < \infty. \end{aligned} \quad (493)$$

For $\delta > 0$, from Condition 245 and (489), we have

$$\begin{aligned} \sum_{i=1}^{m_k} \mathbb{E}(\psi_{k,i}^2 \mathbb{1}(|\psi_{k,i}| > \delta) | \mathcal{F}_{k,i-1}) &= (\mathbb{1}(b \in B) \mathbb{E}_{\mathbb{Q}(b)}(|t^T r(b, \cdot)|^2 \mathbb{1}(|t^T r(b, \cdot)| > \sqrt{n_k \delta})))_{b=b_{k-1}} \\ &\leq |t|^2 (\mathbb{1}(b \in B) \mathbb{E}_{\mathbb{Q}(b)}(|r(b, \cdot)|^2 \mathbb{1}(|r(b, \cdot)| > \sqrt{n_k \delta})))_{b=b_{k-1}} \\ &= |t|^2 \mathbb{1}(b_{k-1} \in B) f(b_{k-1}, n_k (\frac{\delta}{|t|})^2) \\ &\leq |t|^2 R(n_k (\frac{\delta}{|t|})^2) \rightarrow 0, \quad k \rightarrow \infty. \end{aligned} \quad (494)$$

To prove the second point of Theorem 247 we notice that

$$\begin{aligned} \sum_{i=1}^{m_k} \mathbb{E}(\psi_{k,i}^2 | \mathcal{F}_{k,i-1}) &= (\mathbb{1}(b \in B) \mathbb{E}_{\mathbb{Q}(b)}(|t^T r(b, \cdot)|^2))_{b=b_{k-1}} \\ &= (\mathbb{1}(b \in B) v(b))_{b=b_{k-1}} \xrightarrow{p} v(b^*) \end{aligned} \quad (495)$$

from (490) and continuity of $b \rightarrow \mathbb{1}(b \in B)v(b)$ in b^* . \square

Lemma 250. *Let us assume Condition 248, and that for $F = \sup_{b \in B} \mathbb{1}(Y \neq 0)L(b)$ we have*

$$\mathbb{E}_{\mathbb{Q}_1}(|Y|^2 F) < \infty. \quad (496)$$

Then, Condition 245 holds for $u = |r|^2$. If further \mathbb{Q}_1 a.s. $b \rightarrow L(b)$ is continuous, then $b \rightarrow \Sigma(b)$ is continuous on B .

Proof. For each $M \in \mathbb{R}$ and $b \in B$

$$\begin{aligned} f(b, M) &= \mathbb{E}_{\mathbb{Q}(b)}(|YL(b)|^2 \mathbb{1}(|YL(b)| > \sqrt{M})) \\ &= \mathbb{E}_{\mathbb{Q}_1}(|Y|^2 L(b) \mathbb{1}(|YL(b)| > \sqrt{M})) \\ &\leq \mathbb{E}_{\mathbb{Q}_1}(|Y|^2 F \mathbb{1}(|YF| > \sqrt{M})), \end{aligned} \quad (497)$$

so that

$$R(M) \leq \mathbb{E}_{\mathbb{Q}_1}(|Y|^2 F \mathbb{1}(|YF| > \sqrt{M})) \leq \mathbb{E}_{\mathbb{Q}_1}(|Y|^2 F) < \infty. \quad (498)$$

From (496) we have $0 = \mathbb{P}(|Y|^2 F = \infty) = \mathbb{P}(|Y|F = \infty)$. Thus, as $M \rightarrow \infty$, we have $\mathbb{1}(|YF| > \sqrt{M}) \downarrow 0$ and thus from Lebesgue dominated convergence also $\mathbb{E}_{\mathbb{Q}_1}(|Y|^2 F \mathbb{1}(|YF| > \sqrt{M})) \downarrow 0$ and $R(M) \downarrow 0$, i.e. Condition 245 holds. We have

$$\Sigma_{i,j}(b) = \mathbb{E}_{\mathbb{Q}(b)}(Y_i Y_j L(b)^2) = \mathbb{E}_{\mathbb{Q}_1}(Y_i Y_j L(b)) \quad (499)$$

and $|Y_i Y_j L(b)| \leq |Y|^2 F$, $b \in B$. Thus, continuity of $b \rightarrow \Sigma_{i,j}(b)$ (and thus also of $b \rightarrow \Sigma(b)$) on B follows from Lebesgue dominated convergence. \square

Remark 251. *From Theorem 110 and Remark 108, in the LETS setting, for B bounded, under Condition 248, and for F as in Lemma 250, (496) holds if Condition 104 holds for $S = |Y|^2$.*

7.7 Asymptotic properties of MSM results

Consider following conditions.

Condition 252. $d^* \in A \in \mathcal{B}(\mathbb{R}^l)$ and A -valued random variables b_k , $k \in \mathbb{N}_+$, are such that

$$b_k \xrightarrow{p} d^*. \quad (500)$$

Condition 253. *Conditions 150, 152, and 154 hold.*

Remark 254. *From Remark 155 and its counterpart for CGMSM and MGMSM as discussed in Remark 188, under Condition 253 (in which for CGMSM and MGMSM instead of conditions 150 and 152 we take their counterparts as discussed above Remark 188), we have a.s. for sufficiently large k , $b_k = d_k$ for EMSM and GMSM, and $b_k = \tilde{d}_k$ for CGMSM and MGMSM. In particular, it follows that a.s. $\lim_{k \rightarrow \infty} b_k = b^*$ and thus Condition 252 holds for $b^* = d^*$.*

Let us further in this section assume conditions 119 and 252. Using analogous reasonings and assumptions as in Section 7.5, but using CLT from Theorem 249 for b^* replaced by d^* instead of Theorem 5, we receive that under appropriate assumptions as in this theorem we have for g in the below formulas substituted by ce, msq, or ic, that for $f_k(b) = \hat{g}_{n_k}(b_{k-1}, b)(\tilde{\chi}_k)$

$$\sqrt{n_k} \nabla f_k(b^*) \Rightarrow \mathcal{N}(0, \Sigma_g(d^*)). \quad (501)$$

To prove (501) for $g = \text{msq}$ 2 using a reasoning analogous as in Section 7.5 we additionally need the facts that $\widehat{\text{msq}}_{n_k}(b_{k-1}, b^*)(\tilde{\chi}_k) \xrightarrow{p} \text{msq}(b^*)$ and $\widehat{R}_k(b_{k-1}, b^*)(\tilde{\chi}_k) \xrightarrow{p} 1$ which, under appropriate assumptions, follow from convergence in distribution of $\sqrt{n_k}(\widehat{\text{msq}}_{n_k}(b_{k-1}, b^*)(\tilde{\chi}_k) - \text{msq}(b^*))$ and $\sqrt{n_k}(\widehat{R}_k(b_{k-1}, b^*)(\tilde{\chi}_k) - 1)$ which can be proved using Theorem 249 as above. For different g as above, assuming (501) and that Condition 237 holds for $T = \mathbb{N}_+$, $r_k = n_k$, f_k as above, and f corresponding to g as in Remark 243 (see Section 7.3 for some sufficient assumptions), for V_g and B_g as in hat remark we have from Theorem 221

$$\sqrt{n_k}(d_k - b^*) \Rightarrow \mathcal{N}(0, V_g(d^*)), \quad (502)$$

and from Remark 235

$$n_k(f(d_k) - f(b^*)) \Rightarrow \tilde{\chi}^2(B_g(d^*)). \quad (503)$$

If for s_k denoting the number of samples generated till the k th stage of MSM, i.e. $s_k := \sum_{i=1}^k n_i$, $k \in \mathbb{N}_+$, we have

$$\lim_{k \rightarrow \infty} \frac{s_k}{n_k} = \gamma \in [1, \infty), \quad (504)$$

then from (502) it follows that

$$\sqrt{s_k}(d_k - b^*) \Rightarrow \mathcal{N}(0, \gamma V_g(d^*)), \quad (505)$$

while from (503) — that

$$s_k(f(d_k) - f(b^*)) \Rightarrow \widetilde{\chi}^2(\gamma B_g(d^*)). \quad (506)$$

For instance, for $n_k = A_1 + A_2 m^k$ as in Remark 133, we have $s_k = A_1 k + A_2 \frac{m^{k+1} - 1}{m - 1}$, so that $\gamma = \frac{m}{m-1}$. For $n_k = A_1 + A_2 k!$ as in that remark, we have

$$\frac{s_k}{n_k} \leq \frac{s_k}{A_2 k!} = \frac{A_1}{A_2(k-1)!} + 1 + \frac{1}{k} + \frac{1}{k(k-1)} + \dots + \frac{1}{k!} \leq \frac{A_1}{A_2(k-1)!} + 1 + \frac{2}{k}, \quad (507)$$

so that $\gamma = 1$.

Remark 255. *Let us assume that (505) and (506) hold, and let $p_{s_k} = d_k$, $k \in \mathbb{N}_+$. Consider now d_k as in SSM in Section 7.5 for $T = \mathbb{N}_+$, $N_k = k$, $k \in T$, and $b' = d^*$, so that we have (443) for $u(b') = u(d^*) = 1$. Let us assume that (447) and (449) hold for such d_k , and let $p'_{s_k} = d_{s_k}$, $k \in \mathbb{N}_+$. Let further $T = \{s_k : k \in \mathbb{N}_+\}$. Then, $t(f(p_t) - f(b^*)) \Rightarrow \widetilde{\chi}^2(\gamma B_g(d^*))$ and $t(f(p'_t) - f(b^*)) \Rightarrow \widetilde{\chi}^2(B_g(d^*))$, $t \rightarrow \infty$, $t \in T$. Thus, for $\gamma = 1$ or $B_g(d^*) = 0$ the processes p and p' can be considered equally efficient for minimization of f in the second-order sense as discussed in section 7.2, while for $\gamma > 1$ and $B_g(d^*) \neq 0$ the process from SSM is more efficient in such sense.*

Remark 256. *From discussion in Section 7.2 for $T = \mathbb{N}_+$ and $r_k = n_k$, $k \in T$, for $R \sim \widetilde{\chi}^2(B_g(d^*))$, the second-order asymptotic inefficiency for minimization of f of processes $d = (d_k)_{k \in \mathbb{N}_+}$ from MSM and satisfying (503) like above, e.g. minimizing different estimators or using different d^* but the same b^* and n_k , can be quantified using*

$$\mathbb{E}(R) = \frac{1}{2} \text{Tr}(\Sigma_g(d^*) H_f^{-1}). \quad (508)$$

Let further (506) hold and consider a process p indexed by the number of simulations performed and corresponding to some d as above as in Remark 256. Then, the asymptotic inefficiency of p for minimizing f can be quantified using

$$\gamma \frac{1}{2} \text{Tr}(\Sigma_g(d^*) H_f^{-1}). \quad (509)$$

Using (509) one can compare the efficiency of p as above e.g. with that of processes p' from SSM as in Remark 256 but this time without assuming that $b' = d^$.*

If the mean cost $u(b)$ of MSM steps, analogous as in Remark 242 for SSM, depends on b , and $b \rightarrow u(b)$ is continuous in d^ , then it might be reasonable to*

favour MSM processes d as above with smaller values of $u(d^*)$ if the constants p_U as in Remark 148 for the compared processes are the same, or with smaller $p_U u(d^*)$ if these constants are different. By analogy to formula (450) for SSM, one can do this heuristically e.g. by using instead of (508) as a measure of asymptotic inefficiency of such processes the quantity

$$u(d^*) \frac{1}{2} \text{Tr}(\Sigma_g(d^*) H_f^{-1}), \quad (510)$$

or such quantity multiplied by p_U respectively.

A more desirable possibility than having Condition 252 satisfied for $d^* = b^*$ as in Remark 254, may be to have it fulfilled for d^* minimizing some measure of asymptotic inefficiency of d_k for minimization of f , like (508) or (510) in Remark 256. See Section 10 for further discussion of this idea. From Remark 261, which we shall prove in Section 7.8, it follows that for d^* minimizing (508) it may hold $d^* \neq b^*$.

7.8 Properties of matrices characterizing the asymptotic distributions when the optimal- or zero-variance IS parameter exists

In all this section we assume conditions 22 and 23. Consider matrix-valued functions Σ_g , V_g , and B_g , given by formulas from Section 7.5 and considered on subsets of A on which these formulas make sense.

From reasonings as in Section 7.5, for each $b' \in A$, for g replaced by msq2 or ic, for $f_n(b) = \hat{g}_n(b', b^*)(\tilde{\kappa}_n)$, $n \in \mathbb{N}_+$, under appropriate assumptions $\Sigma_g(b')$ is the asymptotic covariance matrix of $\sqrt{n} \nabla f_n(b)$. From Remark 241, for b^* being the optimal-variance IS parameter in case of $g = \text{msq2}$ or the zero-variance one for $g = \text{ic}$, we have a.s. $\nabla f_n(b^*) = 0$, $n \in \mathbb{N}_+$. Thus, in such case $\Sigma_g(b') = 0$, $b' \in A$. This can be also verified by direct calculation. For b^* being the optimal-variance IS parameter, from (35), \mathbb{Q}_1 a.s. (and thus from Condition 22 also $\mathbb{Q}(b)$ a.s., $b \in A$)

$$(ZL(b^*))^2 = \text{msq}(b^*), \quad (511)$$

and thus from (465), $\Sigma_{\text{msq2}} = 0$. Let now b^* be the zero-variance IS parameter. Then, we have $\text{var}(b^*) = 0$ and under appropriate smoothness assumptions $\nabla \text{msq}(b^*) = 0$. Using further (511) and the fact that \mathbb{Q}_1 a.s. $ZL(b^*) = \alpha$, from (479) we receive that for each $b' \in A$, \mathbb{Q}' a.s.

$$\begin{aligned} W(b') &= \nabla c(b^*) (Z^2 L' L(b^*) - \text{msq}(b^*) + \text{msq}(b^*) (L' L^{-1}(b^*) - 1) - 2\alpha (ZL' - \alpha)) \\ &\quad + c(b^*) \nabla L(b^*) L' (Z^2 - \text{msq}(b^*) L(b^*)^{-2}) = 0. \end{aligned} \quad (512)$$

Thus, from (477), $\Sigma_{\text{ic}} = 0$. Using notations as in Remark 243, if H_{msq} is positive definite for $g = \text{msq2}$ or H_{ic} is positive definite for $g = \text{ic}$, and we have $\Sigma_g = 0$ as above, then it also holds $V_g = B_g = 0$.

Using notations as in Section 7.5, let us define Fisher information matrix

$$I(b) = \mathbb{E}_{\mathbb{Q}(b)}(S(b)S(b)^T), \quad b \in A \quad (513)$$

(assuming that it is well-defined). It is well known that under appropriate assumptions, allowing one to move derivatives under expectations in the below derivation, we have

$$I(b) = -\mathbb{E}_{\mathbb{Q}(b)}(\nabla^2 l(b)). \quad (514)$$

The following derivation is as on page 63 in [52]. From $\mathbb{E}_{\mathbb{Q}_1}(p(b)) = 1$ we have $\mathbb{E}_{\mathbb{Q}_1}(\nabla p(b)) = 0$ and $0 = \mathbb{E}_{\mathbb{Q}_1}(\nabla^2 p(b)) = \mathbb{E}_{\mathbb{Q}(b)}(\frac{\nabla^2 p(b)}{p(b)})$, so that taking expectations with respect to $\mathbb{Q}(b)$ of

$$\nabla^2 l(b) = \frac{\nabla^2 p(b)}{p(b)} - \frac{\nabla p(b)(\nabla p(b))^T}{p^2(b)}, \quad (515)$$

we receive (514).

Let us define

$$R(a, b) = \mathbb{E}_{\mathbb{Q}(a)}\left(\frac{L(b)}{L(a)} S(a) S(a)^T\right), \quad a, b \in A. \quad (516)$$

Note that

$$R(b, b) = I(b). \quad (517)$$

Remark 257. For some $a, b \in A$, let $R(a, b)$ and $I(a)$ have real-valued entries. Then, $R(a, b)$ is positive definite only if for each $v \in \mathbb{R}^l$, $v \neq 0$, $\mathbb{E}_{\mathbb{Q}(a)}\left(\frac{L(b)}{L(a)} |S^T(a)v|^2\right) > 0$, which holds only if $\mathbb{Q}(a)(|S^T(a)v| \neq 0) > 0$, $v \in \mathbb{R}^l$, $v \neq 0$, and thus only if $I(a)$ is positive definite.

Let b^* be the optimal-variance IS parameter. Then, from (511)

$$\begin{aligned} \Sigma_{\text{ce}}(b) &= \mathbb{E}_{\mathbb{Q}_1}(Z^2 L(b) S(b^*) S(b^*)^T) \\ &= \text{msq}(b^*) \mathbb{E}_{\mathbb{Q}_1}\left(\frac{L(b)}{L(b^*)^2} S(b^*) S(b^*)^T\right) \\ &= \text{msq}(b^*) R(b^*, b) \end{aligned} \quad (518)$$

and

$$\begin{aligned} \Sigma_{\text{msq}}(b) &= \mathbb{E}_{\mathbb{Q}_1}(Z^4 L(b) L(b^*)^2 S(b^*) S(b^*)^T) \\ &= \text{msq}(b^*)^2 \mathbb{E}_{\mathbb{Q}_1}\left(\frac{L(b)}{L(b^*)^2} S(b^*) S(b^*)^T\right) \\ &= \text{msq}(b^*)^2 R(b^*, b). \end{aligned} \quad (519)$$

It holds (assuming that one can move derivatives under the expectation)

$$\nabla^2 \text{ce}(b) = -\nabla^2 \mathbb{E}_{\mathbb{Q}_1}(Zl(b)) = -\mathbb{E}_{\mathbb{Q}_1}(Z \nabla^2 l(b)). \quad (520)$$

For cross-entropy, let us assume that b^* is the zero-variance IS parameter, so that \mathbb{Q}_1 a.s. we have $ZL(b^*) = \alpha$. Then, from (520) and (514)

$$\begin{aligned} H_{\text{ce}} &= \nabla^2 \text{ce}(b^*) = -\alpha \mathbb{E}_{\mathbb{Q}_1}\left(\frac{1}{L(b^*)} \nabla^2 l(b)\right) \\ &= \alpha I(b^*). \end{aligned} \quad (521)$$

Assuming that $I(b^*)$ is positive definite and $\alpha \neq 0$, from $\text{msq}(b^*) = \alpha^2$, (446), (518), and (521) we have

$$V_{\text{ce}}(b) = H_{\text{ce}}^{-1} \Sigma_{\text{ce}}(b) H_{\text{ce}}^{-1} = I(b^*)^{-1} R(b^*, b) I(b^*)^{-1}, \quad (522)$$

which is positive definite from Remark 257.

Remark 258. From (517) and (522) we have $V_{\text{ce}}(b^*) = I(b^*)^{-1}$. Note that this is the asymptotic covariance matrix of maximum likelihood estimators for b^* being the true parameter, see e.g. page 63 in [52]. This should be the case, since under Condition 53 for $b' = b^*$, a.s.

$$\widehat{\text{ce}}_n(b^*, b)(\tilde{\kappa}_n) = \overline{(ZL(b^*) \ln(L(b)))}_n(\tilde{\kappa}_n) = -\alpha \overline{\ln(p(b))}_n(\tilde{\kappa}_n), \quad (523)$$

and $\overline{\ln(p(b))}_n(\tilde{\kappa}_n) = \frac{1}{n} \sum_{i=1}^n \ln(p(b)(\kappa_i))$ is the estimator maximized in maximum likelihood estimation for b^* being the true parameter.

Under assumptions as above and for $\alpha > 0$, from (448), (518), and (521)

$$B_{\text{ce}}(b) = \frac{1}{2} H_{\text{ce}}^{-\frac{1}{2}} \Sigma_{\text{ce}}(b) H_{\text{ce}}^{-\frac{1}{2}} = \frac{1}{2} \alpha I(b^*)^{-\frac{1}{2}} R(b^*, b) I(b^*)^{-\frac{1}{2}}, \quad (524)$$

which is positive definite. Note that $B_{\text{ce}}(b^*) = \frac{1}{2} \alpha I_l$ and $\text{Tr}(B_{\text{ce}}(b^*)) = \frac{l\alpha}{2}$.

It holds

$$\begin{aligned} \nabla^2 \text{msq}(b) &= -\nabla \mathbb{E}_{\mathbb{Q}_1}(Z^2 L(b) S(b)^T) \\ &= \mathbb{E}_{\mathbb{Q}_1}(Z^2 L(b)(S(b)S(b)^T - \nabla^2 l(b))). \end{aligned} \quad (525)$$

Let us assume that b^* is the optimal-variance IS parameter. Then, from (511), (514), and (525)

$$\begin{aligned} H_{\text{msq}} &= \text{msq}(b^*) \mathbb{E}_{\mathbb{Q}_1} \left(\frac{1}{L(b^*)} (S(b^*)S(b^*)^T - \nabla^2 l(b^*)) \right), \\ &= \text{msq}(b^*) \mathbb{E}_{\mathbb{Q}(b^*)} (S(b^*)S(b^*)^T - \nabla^2 l(b^*)), \\ &= \text{msq}(b^*) 2I(b^*). \end{aligned} \quad (526)$$

Remark 259. Recall that under appropriate assumptions as in Section 5.1, when b^* is the zero-variance IS parameter, then cross-entropy is linearly equivalent to cross-entropy distance of $\mathbb{Q}(b^*)$ from $\mathbb{Q}(b)$ with linear proportionality constant α , and when b^* is the optimal-variance IS parameter, then mean square is linearly equivalent to Pearson's divergence of $\mathbb{Q}(b^*)$ from $\mathbb{Q}(b)$ with such constant $\text{msq}(b^*)$. Thus, the above formulas for second derivatives of cross-entropy and mean square follow from the well-known fact that the second derivative of such cross-entropy distance is equal to Fisher information matrix, and the fact that the second derivative of such Pearson's divergence in b^* is twice such derivative of the cross-entropy distance.

Assuming that $I(b^*)$ is positive definite and $\text{msq}(b^*) \neq 0$, from (446), (519), and (526),

$$\begin{aligned} V_{\text{msq}}(b) &= H_{\text{msq}}^{-1} \Sigma_{\text{msq}}(b) H_{\text{msq}}^{-1} \\ &= \frac{1}{4} I(b^*)^{-1} R(b^*, b) I(b^*)^{-1}. \end{aligned} \quad (527)$$

which is positive definite from Remark 257. Thus, assuming further that b^* is the zero-variance IS parameter and we have (522), it holds

$$V_{\text{msq}}(b) = \frac{1}{4} V_{\text{ce}}(b). \quad (528)$$

From (527), we have $V_{\text{msq}}(b^*) = \frac{1}{4}I(b^*)^{-1}$. Furthermore, from (448)

$$\begin{aligned} B_{\text{msq}}(b) &= \frac{1}{2}H_{\text{msq}}^{-\frac{1}{2}}\Sigma_{\text{msq}}(b)H_{\text{msq}}^{-\frac{1}{2}} \\ &= \frac{\text{msq}(b^*)}{4}I(b^*)^{-\frac{1}{2}}R(b^*, b)I(b^*)^{-\frac{1}{2}}, \end{aligned} \quad (529)$$

which is positive definite. Note that $B_{\text{msq}}(b^*) = \frac{\text{msq}(b^*)}{4}I_I$ and $\text{Tr}(B_{\text{msq}}(b^*)) = \frac{l \text{msq}(b^*)}{4}$.

Remark 260. Let A, T, d_t , and r_t be as in Section 7.1, let $b \in A$, and $u : A \rightarrow \bar{\mathbb{R}}$ be such that $u(b) \in \mathbb{R}_+$ and $r_t(d_t - b^*) \Rightarrow \mathcal{N}(0, u(b)V_{\text{ce}}(b))$ (see Section 7.5 for sufficient conditions for this to hold for $b = b'$ and $r_t = t$ for SSM and Section 7.7 for $b = d^*$, $u(d^*) = 1$, and $r_k = n_k$ for MSM). Let further msq be twice differentiable in b^* with $\nabla \text{msq}(b^*) = 0$ and $H_{\text{msq}} = \nabla^2 \text{msq}(b^*)$. Then, from Remark 234, for

$$B_{\text{ce,msq}}(b) := \frac{1}{2}V_{\text{ce}}(b)^{\frac{1}{2}}H_{\text{msq}}V_{\text{ce}}(b)^{\frac{1}{2}}, \quad (530)$$

we have

$$r_t(\text{msq}(d_t) - \text{msq}(b^*)) \Rightarrow \widetilde{\chi}^2(u(b)B_{\text{ce,msq}}(b)). \quad (531)$$

For b^* being the zero-variance IS parameter, $I(b^*)$ being positive definite, and $\alpha \neq 0$, from (530), (526), (522), and (529)

$$\begin{aligned} B_{\text{ce,msq}}(b) &= \text{msq}(b^*)I(b^*)^{-\frac{1}{2}}R(b^*, b)I(b^*)^{-\frac{1}{2}} \\ &= 4B_{\text{msq}}(b). \end{aligned} \quad (532)$$

Consider now the ECM setting as in Section 4.1. Then,

$$I(b) = \mathbb{E}_{\mathbb{Q}(b)}((X - \mu(b))(X - \mu(b))^T) = \Sigma(b). \quad (533)$$

Furthermore,

$$\begin{aligned} R(a, b) &= \mathbb{E}_{\mathbb{Q}_1}\left(\frac{L(b)}{L(a)^2}S(a)S(a)^T\right) \\ &= \exp(\Psi(b) - 2\Psi(a))\mathbb{E}_{\mathbb{Q}_1}(\exp((2a - b)^T X)(X - \mu(a))(X - \mu(a))^T) \\ &= \exp(\Psi(b) + \Psi(2a - b) - 2\Psi(a))\mathbb{E}_{\mathbb{Q}(2a - b)}((X - \mu(a))(X - \mu(a))^T) \\ &= \exp(\Psi(b) + \Psi(2a - b) - 2\Psi(a))(\Sigma(2a - b) \\ &\quad + (\mu(2a - b) - \mu(a))(\mu(2a - b) - \mu(a))^T). \end{aligned} \quad (534)$$

For a positive definite matrix $B \in \text{Sym}(\mathbb{R})$ and $b \in \mathbb{R}^l$, let $|\cdot|_B = \sqrt{b^T B b}$. $|\cdot|_B$ is a norm on \mathbb{R}^l . For $X \sim \mathcal{N}(\mu_0, M)$ under \mathbb{Q}_1 for some positive definite covariance matrix M , we have from (534) and discussion in Section 4.1

$$R(a, b) = \exp(|a - b|_M^2)(M + M(a - b)(a - b)^T M). \quad (535)$$

In particular, for b^* being the zero-variance IS parameter we have

$$V_{\text{ce}}(b) = \exp(|b^* - b|_M^2)(M^{-1} + (b^* - b)(b^* - b)^T) = 4V_{\text{msq}}(b), \quad (536)$$

and

$$B_{\text{msq}}(b) = \frac{\text{msq}(b^*)}{4} \exp(|b^* - b|_M^2) (I_l + M^{\frac{1}{2}}(b^* - b)(b^* - b)^T M^{\frac{1}{2}}). \quad (537)$$

Thus, the mean of $\widetilde{\chi}^2(B_{\text{msq}}(b))$ is

$$\text{Tr}(B_{\text{msq}}(b)) = \frac{\text{msq}(b^*)}{4} \exp(|b^* - b|_M^2) (l + |b^* - b|_M^2). \quad (538)$$

Note that b^* is the unique minimum point of $b \rightarrow \text{Tr}(B_{\text{msq}}(b))$. From the below remark it follows that for X having a different distribution under \mathbb{Q}_1 this may be not the case.

Remark 261. Consider the ECM setting for $l = 1$. Then, $R(a, b) = \mathbb{E}_{\mathbb{Q}(a)}(\frac{L(b)}{L(a)}(S(a))^2)$. Assuming that one can move derivative under expectation in the below formula, which from theorems 110 and 111 is the case when $A = \mathbb{R}$, we have

$$\begin{aligned} (\nabla_b R(a, b))_{b=a} &= -\mathbb{E}_{\mathbb{Q}(a)}(S(a)^3) \\ &= -\mathbb{E}_{\mathbb{Q}(a)}((X - \mu(a))^3). \end{aligned} \quad (539)$$

In such case, from convexity of $b \rightarrow R(a, b)$ (which follows from such convexity of $b \rightarrow L(b)$), a necessary and sufficient condition for $b = a$ to be its minimum point (and thus for the zero-variance IS parameter $b^* = a$ to be the minimum point of $b \rightarrow B_{\text{msq}}(b)$ as in (529)), is that X has zero third central moment under $\mathbb{Q}(a)$. This does not hold e.g. for $X \sim \text{Pois}(\mu(a))$ under $\mathbb{Q}(a)$ as in Section 4.1, for which $\mathbb{E}_{\mathbb{Q}(a)}((X - \mu(a))^3) = \mu(a) > 0$.

In the LETGS setting, under appropriate integrability assumptions, from (78) and (514), $I(b) = 2\mathbb{E}_{\mathbb{Q}_1}(G)$, which, from Lemma 64, is positive definite only if Condition 62 holds for $Z = 1$.

7.9 Analytical example of a symmetric three-point distribution

Let us consider ECM as in Section 4.1 for $l = 1$, assuming that $\mathbb{Q}_1(X = -1) = \mathbb{Q}_1(X = 0) = \mathbb{Q}_1(X = 1) = \frac{1}{3}$. Then, we have $A = \mathbb{R}$, $\Phi(b) = \frac{e^b + e^{-b} + 1}{3}$, $L(b) = \Phi(b) \exp(-bX)$, $\Psi(b) = \ln(\Phi(b))$, $\mu(b) = \nabla \Psi(b) = \frac{e^b - e^{-b}}{e^b + e^{-b} + 1}$, and $\nabla^2 \Psi(b) = \frac{e^b + e^{-b} + 4}{(e^b + e^{-b} + 1)^2}$. For some $d \in \mathbb{R}$, let $Z = \mathbb{1}(X = 0) + d\mathbb{1}(X \neq 0)$. We have $\alpha = \frac{1+2d}{3}$ and

$$\text{msq}(b) = \mathbb{E}_{\mathbb{Q}_1}(Z^2 L(b)) = \frac{1}{9}(1 + e^b + e^{-b})(1 + d^2(e^b + e^{-b})). \quad (540)$$

The unique minimizer of msq is 0, which corresponds to CMC. It holds $\text{msq}(0) = \frac{1+2d^2}{3}$ and $\text{var}(0) = \frac{1+2d^2}{3} - (\frac{1+2d}{3})^2 = \frac{2}{9}(1-d)^2$. Thus, there exists zero-variance IS parameter only if $d = 1$, in which case this parameter is $b^* := 0$. We have $\mathbb{E}_{\mathbb{Q}_1}(ZX) = 0$, so that from (153), $\text{ce}(b) = \alpha \Psi(b)$. Thus, if $d \neq -\frac{1}{2}$, then ce has unique optimum point $0 = b^*$, which is minimum point if $\alpha > 0$ and maximum point if $\alpha < 0$. We have $\nabla^2 \Psi(0) = \frac{2}{3}$ and

$$H_{\text{ce}} = \nabla^2 \text{ce}(0) = \alpha \nabla^2 \Psi(0) = \frac{2(1+2d)}{9}. \quad (541)$$

It holds $\nabla L(b) = -X \exp(-bX)\Phi(b) + \exp(-bX)\frac{e^b - e^{-b}}{3}$, so that $\nabla L(0) = -X$. Thus, from (454), for

$$g(b) := (1 + e^b + e^{-b})(e^b + e^{-b}), \quad (542)$$

we have

$$\Sigma_{ce}(b) = \mathbb{E}_{\mathbb{Q}_1}(Z^2 L(b)X^2) = \frac{d^2}{9}g(b) \quad (543)$$

and

$$V_{ce}(b) = H_{ce}^{-2}\Sigma_{ce}(b) = \left(\frac{3d}{2(1+2d)}\right)^2 g(b). \quad (544)$$

We have

$$H_{msq} = \nabla^2 \text{msq}(0) = \frac{2}{9}(1 + 5d^2), \quad (545)$$

from (458)

$$\Sigma_{msq}(b) = \mathbb{E}_{\mathbb{Q}_1}(L(b)Z^4 X^2) = \frac{d^4}{9}g(b), \quad (546)$$

and thus

$$V_{msq}(b) = H_{msq}^{-2}\Sigma_{msq}(b) = \left(\frac{3d^2}{2(1+5d^2)}\right)^2 g(b). \quad (547)$$

From (465)

$$\Sigma_{msq2}(b) = \mathbb{E}_{\mathbb{Q}_1}((Z^2 - \text{msq}(0))^2 L(b)X^2) = \frac{1}{3^4}(d^2 - 1)^2 g(b), \quad (548)$$

so that

$$V_{msq2}(b) = H_{msq}^{-2}\Sigma_{msq2}(b) = \left(\frac{d^2 - 1}{2(1+5d^2)}\right)^2 g(b). \quad (549)$$

From (481)

$$\Sigma_{ic}(b) = \mathbb{E}_{\mathbb{Q}_1}((Z^2 - \text{msq}(0) - \text{var}(0))^2 L(b)X^2) = \frac{1}{3^6}((d-1)(d+5))^2 g(b), \quad (550)$$

and thus

$$V_{ic}(b) = H_{msq}^{-2}\Sigma_{ic}(b) = \left(\frac{(d-1)(d+5)}{6(1+5d^2)}\right)^2 g(b). \quad (551)$$

For s substituted by ce , msq , $msq2$, and ic , let us further write $V_s(b, d)$ rather than $V_s(b)$ to mark its dependence on d . Let for such different s , $f_s(d) = 2(1 + 5d^2)\sqrt{\frac{V_s(b, d)}{g(b)}}$, so that $f_{ce}(d) = \left|\frac{3d(1+5d^2)}{1+2d}\right|$, $f_{msq}(d) = 3d^2$, $f_{msq2}(d) = |d^2 - 1|$, and $f_{ic}(d) = \frac{1}{3}|(d-1)(d+5)|$. It holds $\frac{V_{s_1}}{V_{s_2}}(b, d) = \left(\frac{f_{s_1}}{f_{s_2}}(d)\right)^2$ whenever $f_{s_2}(d) \neq 0$. The graphs of functions f_s are shown in Figure 4. Note that $b \rightarrow g(b)$ is positive and has a unique minimum point in $0 = b^*$. Thus, if $f_s(d) \neq 0$, then $b \rightarrow V_s(b, d) = g(b)(f_s(d)2(1+5d^2))^2$ also has a unique minimum point in b^* . It holds $f_{ce}(0) = f_{msq}(0) = 0$. For $d \neq 0$ and $d \neq -\frac{1}{2}$, for $r(d) := \frac{f_{ce}}{f_{msq}}(d) = \left|\frac{1+5d^2}{d(1+2d)}\right|$, we have $|r'(d)| = \left|\frac{(d-1)(d+5)}{(d(1+2d))^2}\right| = 0$ only for $d = -5$ or $d = 1$, which correspond to minimum points of r and in which $r(-5) = \frac{42}{15}$ and $r(1) = 2$. In particular, $f_{ce}(d) \geq 2f_{msq}(d)$ and thus $V_{ce}(b, d) \geq 4V_{msq}(b, d)$, with equality holding only for d equal to 0 and 1, the latter being in agreement

with theory of Section 7.8. From an easy calculation we have $f_{s_1} < f_{s_2}$ on D , $f_{s_1} = f_{s_2}$ on ∂D , and $f_{s_1} > f_{s_2}$ otherwise for $s_1 = \text{msq}$, $s_2 = \text{msq} 2$, and $D = (-\frac{1}{2}, \frac{1}{2})$, $s_1 = \text{msq}$, $s_2 = \text{ic}$, and $D = (\frac{-2-3\sqrt{2}}{10}, \frac{-2+3\sqrt{2}}{10})$, or $s_1 = \text{msq} 2$, $s_2 = \text{ic}$, and $D = (-2, 1)$. Since for s equal to $\text{msq} 2$ or ic , f_s is continuous and $f_s(0) > 0$, such f_s are smaller than f_{ce} on some neighbourhood of 0. Note that in agreement with theory of Section 7.8, for $d = 1$, for which b^* is zero-variance IS parameter, the new estimators lead to zero values of $V_s(b, d)$ and the new mean square estimator does this also for $d = -1$, for which b^* is the optimal-variance IS parameter. For $B_s(b, d) = \frac{1}{2}V_s(b, d)H_{\text{msq}}$ for s equal to msq , $\text{msq} 2$, or ic , and for $B_{\text{ce,msq}}(b, d) = \frac{1}{2}V_{\text{ce}}(b, d)H_{\text{msq}}$ for $s = \text{ce}$ as in the previous sections, we have analogous relations as for different $V_s(b, d)$ above.




Figure 4: Functions f_s as in the main text for s equal to ce , msq , $\text{msq} 2$, and ic .

8 Two-stage estimation of the quantity of interest

In this section we shortly discuss a two-stage adaptive method for estimation, in the first stage of which some parameter vector d is computed, with the help of which in the second stage an estimator of the quantity of interest α is calculated. Detailed analysis of such approach, including its various limiting properties, is relegated to a future paper. Some alternative estimation approaches are discussed in the below Remark 265 and in Section 10.

For a nonempty parameter set $A \in \mathcal{B}(\mathbb{R}^l)$ and a family of distributions as in Section 3.4, let a measurable function h as in (415) be such that for each $b \in A$,

$$\mathbb{E}_{\mathbb{Q}(b)}(h(b, \cdot)) = \alpha. \quad (552)$$

Let $\text{var}(b) = \text{Var}_{\mathbb{Q}(b)}(h(b, \cdot))$, $b \in A$. Consider some cost variable C and functions c and ic as in Section 5.1 but using the above var (in particular, for ic we assume Condition 52). In the IS case conditions 17 and 198 hold, but the above setting is more general and can also describe e.g. control variates or control variates

in conjunction with IS. The parameter vector d computed in the first stage is an A -valued random variable, which can be obtained e.g. using some adaptive algorithm like single- or multi-stage minimization methods of some estimators described in the previous sections. This can be done in many different ways as discussed in the below remark.

Remark 262. *One possibility is to use as d some parameter d_t corresponding to some first-stage budget t as in Remark 197. Alternatively, in methods in which to compute some parameter p_k one first needs to compute p_l , $l = 1, \dots, k-1$, like in MSM methods from the previous sections for p_k as in Remark 197, one can set $d = p_\tau \mathbb{1}(\tau \neq \infty) + p_\infty \mathbb{1}(\tau = \infty)$ for some $p_\infty \in A$ and an a.s. finite $\mathbb{N}_+ \cup \{\infty\}$ -valued stopping time τ for the filtration $\mathcal{F}_n = \sigma\{p_l : l \leq n\}$, $n \in \mathbb{N}_+$. For instance, if a.s. $p_k \rightarrow b^*$, then τ can be the moment when the change of p_k from p_{k-1} , or such relative change if $b^* \neq 0$, becomes smaller than some $\epsilon \in \mathbb{R}_+$ (see e.g. Remark 10 in [42]). For different p_k in MSM methods, the fact that a.s. $p_k \rightarrow b^*$ follows from Condition 152. When for some functions $f_k : \mathcal{S}(A) \otimes \mathcal{S}_1 \rightarrow \mathcal{S}(\mathbb{R})$, $k \in \mathbb{N}_+$, (which can be e.g. the minimized estimators) and $f : A \rightarrow \mathbb{R}$ we have a.s. $f_k(p_k) \rightarrow f(b^*)$, then the stopping time can be based on the behaviour of $f_k(p_k)$, similarly as for p_k above. Assuming that a.s. $f_k \xrightarrow{loc} f$ (see Section 6.3 for sufficient assumptions), if a.s. $p_k \rightarrow b^*$, then from Lemma 175 a.s. $f_k(p_k) \rightarrow f(b^*)$.*

One way to model the second stage of a two-stage method is to consider it on a different probability space than the first one, for a fixed computed value v of the variable d from the first stage. In such case, for some $\kappa_1, \kappa_2, \dots$, i.i.d. $\sim \mathbb{Q}(v)$, in the second stage we perform a MC procedure as in Section 2 but using $Z_i = h(v, \kappa_i)$, $i \in \mathbb{N}_+$, e.g. using a fixed number of samples or a fixed approximate budget. We can also construct asymptotic confidence intervals for α as in that section. From discussion in Section 2, the inefficiency of such procedure can be quantified using inefficiency constant $\text{ic}(v)$. This justifies comparing the asymptotic efficiency of methods for finding adaptive parameters in the first stage by comparing their first- and, if applicable, second-order asymptotic efficiency for minimization of ic as discussed in sections 6.10 and 7.2.

An alternative way to model a two-stage method is to consider both stages on the same probability space. Let us assume a following condition.

Condition 263. *Random variables ϕ_i , $i \in \mathbb{N}_+$, are conditionally independent given d and have the same conditional distribution $\mathbb{Q}(b)$ given $d = b$.*

Condition 263 is implied by the following one.

Condition 264. *Condition 19 holds, and for β_1, β_2, \dots , i.i.d. $\sim \mathbb{P}_1$ and independent of d we have $\tilde{\phi} = (\phi_i)_{i \in \mathbb{N}_+} = (\xi(d, \beta_i))_{i \in \mathbb{N}_+}$.*

In the second stage of the considered method one computes an estimator

$$\hat{\alpha}_n = \frac{1}{n} \sum_{i=1}^n h(d, \chi_i). \quad (553)$$

Similarly as above, the number n of samples can be deterministic or random. In the first case the resulting estimator is unbiased, while in the second this needs not to be true. Random n can correspond e.g. to a fixed approximate

computational budget and be given by definition (5) or (6) but for C_i replaced by $C(\phi_i)$, $i \in \mathbb{N}_+$.

Remark 265. *A possible alternative to the above discussed two-stage estimation method is the same as its second model above except that for computation of $\hat{\alpha}_n$ in the second stage one uses variables $\phi_i = \xi(d, \beta_i)$ as in Condition 264 but without assuming that β_i are independent of d . In such case, Condition 263 may not hold. For example, one could reuse the i.i.d. random variables with distribution \mathbb{P}_1 generated for estimation of d in the first stage as some (potentially all) variables β_i used for computation of $\hat{\alpha}_n$, which could save computation time. Under appropriate identifications, such approach using exactly the same β_i , $i = 1, \dots, n$ in ESSM to compute d and then (553) is a common one in multiple control variates (see [4, 21]), while for IS it was considered in [29]. In such reusing approach, $\hat{\alpha}_n$ as in (553) needs not to be unbiased even for n deterministic. Furthermore, one needs to store a potentially large random number of generated values of random variables, which may be more difficult to implement and requires additional computer memory. Finally, in a number of situations, like in case of our numerical experiments, generating the required parts of variables β_i forms only a small fraction of the computation time needed for computing $h(d, \phi_i) = (ZL(d))(\xi_i(d, \beta_i))$ in the second stage, so that reusing some β_i from the first stage would not lead to considerable time savings.*

9 Numerical experiments

In this section we describe our numerical experiments, already briefly discussed in Section 1.2. Our experiments were carried out using programs written in matlab2012a. Unless stated otherwise, we used the simulation parameters, variables, and IS basis functions as for the problems of estimation of expectations $\text{mgf}(x_0)$, $q_{1,a}(x_0)$, $q_{2,a}(x_0)$, and $p_{T,a}(x_0)$, described in Section 4.9. In some of our experiments we performed SSM or MSM of estimators $\widehat{\text{est}}$ (where for short we write $\widehat{\text{est}}$ rather than $\widehat{\text{est}}_n$, $n \in \mathbb{N}_+$) equal to \widehat{c}_e , $\widehat{\text{msq}}$, $\widehat{\text{msq}2}$, and \widehat{ic} , as discussed in Section 6.1. In MSM we used in each case $b_0 = 0$. For minimization of $\widehat{\text{msq}2}$ and \widehat{ic} in these methods we used the matlab `fminunc` unconstrained minimization function with default settings and exact gradient, while for $\widehat{\text{msq}}$ additionally using its exact Hessian matrix. The minimum points of \widehat{c}_e were found by solving linear systems of equations as discussed in Section 6.1. Both for CMC and when using IS, the computation times of simulations on a laptop were typically approximately proportional to the exit times τ in a given estimation problem (where when considering $p_{T,a}(x_0)$ by exit time we mean τ' as in Remark 51) with some proportionality constant p_C (see Section 1.1). Such p_C was different for CMC and for different basis functions used in IS.

The remainder of this section is organized as follows. In Section 9.1 we discuss some methods for testing hypotheses and conventions used for interpreting and describing the results of our numerical experiments. In Section 9.2 we describe two-stage estimation experiments as in Section 6.1, performing MSM in the first stages and in the second stages estimating expectations of functionals of Euler scheme as above. In the second stages we also estimated some other quantities, like inefficiency constants, variances, mean exit times, and proportionality constants p_C as above. We use these quantities to compare

the efficiency of applying in IS MC method the IS parameters obtained from MSM of different estimators, as well as of using different parameters a and IS basis functions in such adaptive IS procedures. In Section 9.3 we compare the spreads of IS drifts coming from SSM of different estimators and using different parameters b' . In Section 9.4 we provide some intuitions behind the results of our numerical experiments.

9.1 Testing statistical hypotheses

Let $\mu_X, \mu_Y \in \mathbb{R}$, $\sigma_X, \sigma_Y \in \mathbb{R}_+$, and let \mathbb{R} -valued random variables X_n and Y_n , $n \in \mathbb{N}_+$, be such that $\tilde{X}_n := \sqrt{n}(X_n - \mu_X) \Rightarrow \mathcal{N}(0, \sigma_X^2)$, $\tilde{Y}_n := \sqrt{n}(Y_n - \mu_Y) \Rightarrow \mathcal{N}(0, \sigma_Y^2)$, and for each $j, k \in \mathbb{N}_+$, X_j is independent of Y_k . Let $\hat{\sigma}_{X,n}$ and $\hat{\sigma}_{Y,n}$, $n \in \mathbb{N}_+$, be $[0, \infty)$ -valued random variables such that $\hat{\sigma}_{X,n} \xrightarrow{p} \sigma_X$ and $\hat{\sigma}_{Y,n} \xrightarrow{p} \sigma_Y$. Let $a_n, b_n \in \mathbb{N}_+$, $n \in \mathbb{N}_+$, be such that $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = \infty$ and

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \rho \in \mathbb{R}_+. \quad (554)$$

Let

$$t_n = \frac{X_{a_n} - Y_{b_n}}{\sqrt{\frac{\hat{\sigma}_{X,a_n}^2}{a_n} + \frac{\hat{\sigma}_{Y,b_n}^2}{b_n}}} = \frac{\sqrt{a_n}(X_{a_n} - Y_{b_n})}{\sqrt{\hat{\sigma}_{X,a_n}^2 + \frac{a_n}{b_n}\hat{\sigma}_{Y,b_n}^2}} \quad (555)$$

and $H_n = \frac{\sqrt{a_n}(\mu_Y - \mu_X)}{\sqrt{\hat{\sigma}_{X,a_n}^2 + \frac{a_n}{b_n}\hat{\sigma}_{Y,b_n}^2}}$.

Lemma 266. *Under assumptions as above, we have*

$$t_n + H_n = \frac{\tilde{X}_{a_n} - \sqrt{\frac{a_n}{b_n}}\tilde{Y}_{b_n}}{\sqrt{\hat{\sigma}_{X,a_n}^2 + \frac{a_n}{b_n}\hat{\sigma}_{Y,b_n}^2}} \Rightarrow \mathcal{N}(0, 1). \quad (556)$$

Proof. From the asymptotic properties of \tilde{X}_i and \tilde{Y}_j as above and their independence, we receive, e.g. using Fubini's theorem and the fact that convergence in distribution is equivalent to pointwise convergence of characteristic functions, that

$$\tilde{X}_{a_n} + \sqrt{\rho}\tilde{Y}_{b_n} \Rightarrow \mathcal{N}(0, \sigma_X^2 + \rho\sigma_Y^2). \quad (557)$$

Thus, from $G_n := \tilde{Y}_{b_n}(\sqrt{\rho} - \sqrt{\frac{a_n}{b_n}}) \xrightarrow{p} 0$,

$$\tilde{X}_{a_n} - \sqrt{\frac{a_n}{b_n}}\tilde{Y}_{b_n} = \tilde{X}_{a_n} - \sqrt{\rho}\tilde{Y}_{b_n} + G_n \Rightarrow \mathcal{N}(0, \sigma_X^2 + \rho\sigma_Y^2). \quad (558)$$

Furthermore, from the continuous mapping theorem,

$$\sqrt{\hat{\sigma}_{X,a_n}^2 + \hat{\sigma}_{Y,b_n}^2 \frac{a_n}{b_n}} \xrightarrow{p} \sqrt{\sigma_X^2 + \rho\sigma_Y^2}. \quad (559)$$

Now, (556) follows from (558), (559), and Slutsky's lemma. \square

If $\mu_X \leq \mu_Y$, then from (556) and $H_n \geq 0$, for each $\alpha \in (0, 1)$ and $z_{1-\alpha}$ as in Remark 7,

$$\limsup_{n \rightarrow \infty} \mathbb{P}(t_n > z_{1-\alpha}) \leq \lim_{n \rightarrow \infty} \mathbb{P}(t_n + H_n > z_{1-\alpha}) = \alpha, \quad (560)$$

i.e. the tests of the null hypothesis $\mu_X \leq \mu_Y$ with regions of rejection $t_n > z_{1-\alpha}$ are pointwise asymptotically level α (see Definition 11.1.1 in [36]). We shall further use such tests for $z_{1-\alpha} = 3$, so that $\alpha \approx 0.00270$. If for some selected n we have $t_n \geq 3$, i.e. the null hypothesis as above can be rejected, then we shall informally say that the estimate $X_{a_n} \pm \frac{\hat{\sigma}_{X,a_n}}{\sqrt{a_n}}$ of μ_X is (statistically significantly) higher than such estimate $Y_{b_n} \pm \frac{\hat{\sigma}_{Y,b_n}}{\sqrt{b_n}}$ of μ_Y .

Most frequently, for some i.i.d. square integrable random variables $X'_i, i \in \mathbb{N}_+$, and such variables $Y'_i, i \in \mathbb{N}_+$, independent of X'_j , in such tests we shall use $X_n = \frac{1}{n} \sum_{i=1}^n X'_i$ and $\hat{\sigma}_{X,n} = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (X'_i - X_n)^2}$, and analogously for Y_n and $\hat{\sigma}_{Y,n}$. In such case $\frac{\hat{\sigma}_{X,n}}{\sqrt{n}}$ shall be called an estimate of the standard deviation of the mean X_n .

9.2 Estimation experiments

We first performed k -stage minimization methods of different estimators and for different estimation problems, using $n_i = 50 \cdot 2^{i-1}$ samples in the i th stage for $i = 1, \dots, k$ (see Section 6.1). We chose $k = 3$ for the problem of estimating $q_{1,a}(x_0)$, $k = 5$ for $\text{mgf}(x_0)$, and $k = 6$ for $p_{T,a}(x_0)$ and $q_{2,a}(x_0)$. We first used $a = 0.05$ and $M = 10$ time-independent IS basis functions as in (113). For $i = 1, 2, \dots, 6$, the IS drifts $r(b_i)$ corresponding to the minimization results b_i from the i th stage of MSM of different estimators for estimating translated committor $q_{2,a}(x_0)$ are shown in Figure 5. The final drifts from MSM for estimation of all the expectations are shown in Figure 6. In figures 5 and 6 we also show for comparison approximations of the zero-variance IS drifts r^* for the diffusion problems for translated committors and MGF, computed from formula (109) using finite differences instead of derivatives and finite difference approximation of u in that formula computed as in Section 4.9. In Figure 5, the IS drifts from the consecutive stages of MSM of $\widehat{\text{msq}}_2$ and $\widehat{\text{ic}}$ seem to converge fastest to some limiting drift close to (the approximation of) r^* , from MSM of $\widehat{\text{msq}}$ — slower, and of $\widehat{\text{c}}_e$ — slowest. See Section 9.4 for some intuitions behind these results.

Consider a numerical experiment in which, for a given IS parameter $v \in A$, we compute unbiased estimates of mean exit times $\mathbb{E}_{\mathbb{Q}(v)}(\tau)$ as well as of (theoretical) inefficiency constants and variances of IS estimators of the expectation of interest, using estimators (133), (144), and (137) respectively for $b = b' = v$ and $n = 10$. For some β_1, \dots, β_n i.i.d. $\sim \mathcal{U}$, these estimators are evaluated on $(\xi(\beta_i, v))_{i=1}^n$, so that the computations involve simulating n independent Euler schemes with additional drift $r(v)$ as in (92). Such experiments for different estimation problems and for v equal to the final results of MSM of $\widehat{\text{c}}_e$, $\widehat{\text{msq}}$, $\widehat{\text{msq}}_2$, and $\widehat{\text{ic}}$ as above, and for $v = 0$ for CMC, were repeated independently K times in an outer MC loop for different K . For the problem of estimation of $q_{2,a}(x_0)$ we additionally used as v the minimization result of the third MSM step. For MGF we made in all cases $K = 75000$ repetitions. For translated committors, when using CMC or IS with IS parameter v from 3-stage MSM of estimators other than $\widehat{\text{c}}_e$, we took $K = 2000$, while in other cases we chose $K = 5000$. For $p_{T,a}$ we made $K = 20000$ repetitions both for CMC and when using v from MSM of $\widehat{\text{c}}_e$, and $K = 2 \cdot 10^5$ repetitions for v from MSM of $\widehat{\text{msq}}$, $\widehat{\text{msq}}_2$, and $\widehat{\text{ic}}$. The MC means of inefficiency constant and variance estimators from the outer loops for translated committors and MGF are given in Table 1,




Figure 5: IS drifts from different stages of MSM for estimating $q_{2,a}$, minimizing \widehat{ce} in (a), \widehat{msq} in (b), \widehat{msq}^2 in (c), and \widehat{ic} in (d). 'Optimal' denotes an approximation of the zero-variance IS drift r^* .

along with estimates of standard deviations of such means. For $p_{T,a}(x_0)$, the estimates of inefficiency constants, variances, and mean exit times are given in Table 2.

Remark 267. *Note that due to Remark 49, τ as above have all moments (and thus also variance) finite under $\mathbb{Q}(v)$, $v \in A$. For $v = 0$ (i.e. for CMC), from boundedness of the considered Z , we have finiteness of variances and inefficiency constants of the estimators of quantities of interest as well as of variances of estimators of such variances and inefficiency constants. For general v and τ bounded (as is the case for τ equal to τ' when estimating $p_{T,a}$), finiteness of quantities as in the previous sentence follows from the corresponding Z being bounded, as well as from Theorem 110 and Remark 108. In cases when τ is not bounded (like for MGF and translated committors above), one can ensure such boundedness by terminating the simulations at some fixed time as discussed in Section 5.4. We did not terminate our simulations, but still our results can be interpreted as coming from simulations terminated at some time larger than any of simulation times encountered in our experiments.*

From tables 1 and 2 we can see that using the IS parameters from MSM of \widehat{ic}




Figure 6: Final IS drifts from different MSM experiments for $q_{1,a}(x_0)$ in (a), $q_{2,a}(x_0)$ in (b), $\text{mgf}(x_0)$ in (c), and $p_{T,a}(x_0)$ in (d). “Optimal“ denotes an approximation of the zero-variance IS drift.

and $\widehat{\text{msq}}_2$ led in each case to the lowest estimates of variances and (theoretical) inefficiency constants, followed by the ones from using parameters from MSM of $\widehat{\text{msq}}$, and finally $\widehat{c\epsilon}$. Using CMC led in each case to the highest such estimates. For $q_{2,a}(x_0)$ and each of $\widehat{c\epsilon}$, $\widehat{\text{msq}}$, and $\widehat{\text{msq}}_2$, using the IS parameter from the sixth stage of MSM led to lower estimate of variance and inefficiency constant than using such parameter from the third stage. Note also that the estimates of inefficiency constants and variances for $q_{2,a}(x_0)$ when using the IS parameters from the third stage of MSM of $\widehat{\text{msq}}_2$ and \widehat{ic} are lower than when using the parameters from the sixth stage of MSM of $\widehat{c\epsilon}$ and $\widehat{\text{msq}}$ (though for estimates of inefficiency constants for $\widehat{\text{msq}}_2$ and $\widehat{\text{msq}}$ we cannot confirm this at a desired significance level as in Section 9.1). For $p_{T,a}(x_0)$, the estimates of inefficiency constants, variances, and mean exit times are respectively lower, higher, and lower when minimizing \widehat{ic} than $\widehat{\text{msq}}_2$. Some intuitions behind these results are given by Theorem 193 and Remark 94, see also discussion in Section 9.4.

Using the IS parameters from MSM of \widehat{ic} and averaging estimates from nK simulations available in each case we computed estimates of mean exit times $\mathbb{E}_{Q(v)}(\tau)$, as well as of quantities of interest: $\text{mgf}(x_0)$, and using translated

	CMC	\hat{c}_e	\widehat{msq}	\widehat{msq}^2	\hat{ic}
Estimates of inefficiency constants					
$q_{1,a} (\cdot 10^{-3})$	7204 ± 92	4855 ± 649	507 ± 10	129.1 ± 3.8	132.9 ± 3.5
$q_{2,a} (\cdot 10^{-3}), k = 3$	7300 ± 88	553 ± 26	60.8 ± 1.2	53.4 ± 1.0	50.5 ± 1.1
$q_{2,a} (\cdot 10^{-3}), k = 6$		73.1 ± 1.0	56.10 ± 0.69	48.78 ± 0.59	47.99 ± 0.61
mgf ($\cdot 10^{-3}$)	2041 ± 15	6.276 ± 0.055	3.691 ± 0.035	3.177 ± 0.029	3.163 ± 0.028
Estimates of variances					
$q_{1,a} (\cdot 10^{-3})$	174.0 ± 1.7	126 ± 14	12.36 ± 0.20	3.155 ± 0.083	3.208 ± 0.073
$q_{2,a} (\cdot 10^{-3}), k = 3$	177.43 ± 1.7	16.35 ± 0.79	1.437 ± 0.021	1.268 ± 0.019	1.223 ± 0.020
$q_{2,a} (\cdot 10^{-3}), k = 6$		1.794 ± 0.021	1.347 ± 0.013	1.164 ± 0.011	1.169 ± 0.012
mgf ($\cdot 10^{-3}$)	49.41 ± 0.32	0.9040 ± 0.0072	0.3732 ± 0.0029	0.3195 ± 0.0024	0.3209 ± 0.0024

Table 1: Estimates of inefficiency constants and variances of estimators of translated committors for $a = 0.05$ and MGF when using CMC or IS with IS parameters from MSM of different estimators. For $q_{2,a}$ we consider using IS parameters from the k th stages of MSM for $k \in \{3, 6\}$.

	CMC	\hat{c}_e	\widehat{msq}	\widehat{msq}^2	\hat{ic}
ic ($\cdot 10^{-3}$)	1391 ± 5	65.22 ± 0.30	63.332 ± 0.09	62.677 ± 0.090	61.863 ± 0.091
var ($\cdot 10^{-3}$)	150.78 ± 0.58	10.258 ± 0.042	10.036 ± 0.013	9.8937 ± 0.0126	9.9491 ± 0.0130
$\mathbb{E}_{Q(v)}(\tau)$	9.227 ± 0.004	6.3608 ± 0.0066	6.3117 ± 0.0021	6.3355 ± 0.0021	6.2173 ± 0.0021

Table 2: Estimates of inefficiency constants and variances of estimators of $p_{T,a}$ for $a = 0.05$ as well as of mean exit times ($\mathbb{E}_{Q(v)}(\tau)$), when using CMC or IS with IS parameters v from MSM of different estimators.

estimators as in Section 4.7 also of $p_T(x_0)$ and $q_i(x_0)$, $i = 1, 2$. The results are presented in Table 3. For comparison, in Table 3 we also provide an estimate of mean exit time for CMC, computed from $7.5 \cdot 10^5$ simulations.

i	$q_1(x_0)$	$q_2(x_0)$	mgf(x_0)	$p_T(x_0)$	CMC
Results	0.7751 ± 0.0004	0.22597 ± 0.00025	$0.16682 \pm (6 \cdot 10^{-5})$	$0.18396 \pm (7 \cdot 10^{-5})$	
$\mathbb{E}_{Q(v)}(\tau)$	41.26 ± 0.28	41.18 ± 0.28	9.89 ± 0.03	6.2173 ± 0.0021	41.44 ± 0.15

Table 3: Estimates of different expectations using IS with IS parameters v from MSM of \hat{ic} , and the corresponding mean exit times. In the last column we provide for comparison the mean exit time in CMC.

We have $q_1(x_0) = 1 - q_2(x_0) \approx 0.78$, and the estimates of inefficiency constants in Table 1 for estimating the lower value committor $q_2(x_0)$ are lower. Thus, it seems reasonable to use the translated IS estimator for $q_2(x_0)$ also for computing $q_1(x_0)$ as discussed in Section 4.7. Note that the estimates of mean exit times in Table 3 are lower under the IS parameter from MSM for computing mgf(x_0) and $p_{T,a}(x_0)$ than for CMC. As discussed in Section 1.2, an intuition behind these results is provided by Theorem 16.

We also performed two-stage experiments similar as above for $q_{2,a}(x_0)$ and $p_{T,a}(x_0)$ for several different added constants $a \in \mathbb{R}_+$ other than $a = 0.05$ considered above, for $q_{2,a}(x_0)$ using IS basis functions as above, while for $p_{T,a}(x_0)$ also using time-dependent basis functions as in (114) for $M = 5$ or $M = 10$ and various $p \in \mathbb{N}_+$. This time in the first stages we performed MSM only of \hat{ic} for $k = 3$ and $n_i = 400 \cdot 2^{i-1}$, $i = 1, \dots, k$, so that the number of samples $n_k = 1600$ used in the final stages of MSM was the same as for $a = 0.05$ above. In the

second stages we estimated inefficiency constants, mean hitting times, and variances in an external loop like above. For $q_{2,a}(x_0)$ we made $K = 3000$ repetitions in such a loop, while for $p_{T,a}(x_0)$ — $K = 10000$ for basis functions as in (113), as well as $K = 50000$ for basis functions as in (114) for $M = 5$ and $K = 30000$ for $M = 10$. The results are presented in tables 4, 5, and 6, along with the results for the case of $a = 0.05$ considered before. The smallest estimates of inefficiency constants and variances for $q_{2,a}(x_0)$ were obtained for $a = 0.05$. For $p_{T,a}(x_0)$ and basis functions as in (113), we obtained the smallest variance for $a = 0.2$ and the lowest inefficiency constants for $a = 0.1$ and $a = 0.2$. Among all cases for $p_{T,a}$, the smallest variances and theoretical inefficiency constants were received for $a = 0$ and when using time-dependent IS basis functions (114) for $M = 10$ and $p = 3$.

$a =$	0	0.05	0.1	0.2
ic ($\cdot 10^{-3}$)	82.6 ± 2.3	47.99 ± 0.61	51.89 ± 0.80	55.52 ± 0.86
var ($\cdot 10^{-3}$)	2.008 ± 0.058	1.169 ± 0.012	1.252 ± 0.016	1.359 ± 0.016

Table 4: Estimates of inefficiency constants and variances of IS estimators of $q_{2,a}$ for different a , using the IS parameters from MSM of \hat{ic} .

$a =$	0	0.05	0.1	0.2	0.3
ic ($\cdot 10^{-3}$)	100.8 ± 0.7	61.86 ± 0.09	55.95 ± 0.35	56.44 ± 0.39	61.43 ± 0.48
var ($\cdot 10^{-3}$)	17.88 ± 0.10	9.949 ± 0.013	8.224 ± 0.047	7.544 ± 0.050	7.794 ± 0.058
$\mathbb{E}_{Q(v)}(\tau)$	5.641 ± 0.009	6.2173 ± 0.0021	6.803 ± 0.009	7.489 ± 0.009	7.874 ± 0.009

Table 5: Estimates of inefficiency constants and variances of estimators of $p_{T,a}$ for different a , and estimates of mean exit times ($\mathbb{E}_{Q(v)}(\tau)$), for IS with parameters v from MSM of \hat{ic} .

	$M = 5$				$M = 10, a = 0$		
	$t, a = 0$	$t, a = 0.05$	$t^2, a = 0$	$t^3, a = 0$	t	t^2	t^3
ic ($\cdot 10^{-3}$)	63.01 ± 0.39	93.44 ± 0.45	48.18 ± 0.28	46.52 ± 0.23	34.11 ± 0.16	22.95 ± 0.17	17.34 ± 0.12
var ($\cdot 10^{-3}$)	10.544 ± 0.064	13.60 ± 0.06	7.975 ± 0.045	7.629 ± 0.036	5.714 ± 0.025	3.899 ± 0.027	2.951 ± 0.019
$\mathbb{E}_{Q(v)}(\tau)$	5.978 ± 0.003	6.871 ± 0.004	6.037 ± 0.003	6.086 ± 0.003	5.966 ± 0.004	5.878 ± 0.004	5.887 ± 0.004

Table 6: Estimates of inefficiency constants and variances of IS estimators of $p_{T,a}$ for different a , as well as of mean exit times ($\mathbb{E}_{Q(v)}(\tau)$), for IS parameters v from MSM of \hat{ic} , using time-dependent IS basis functions as in (114) for different M and p in t^p .

In our experiments, when using CMC and IS MC with different sets of IS basis functions, the approximate proportionality constants p_C as in Section 1.1 were considerably different. Thus, to compare the efficiency of MC methods using estimators corresponding to these different bases, one should compare their practical rather than theoretical inefficiency constants. We performed separate experiments approximating some p_C as above and computing the corresponding practical inefficiency constants (equal to products of such p_C and the respective theoretical inefficiency constants). For $n = 10^5$, we run n -step CMC and IS MC procedures for estimating $q_{2,0.05}(x_0)$ using IS basis functions as in (113), and for estimating $p_{T,a}$: for $a = 0.05$ for IS basis functions as in (113) for $M = 10$,

and for $a = 0$ for IS basis functions as in (114): for $M = 5$ and $p = 1$, and for $M = 10$ and $p \in \{1, 3\}$. For τ_i being the exit time in the i th step of a given MC procedure and $\tilde{\tau}_i$ being its practical cost equal to its computation time calculated using matlab tic and toc functions, as an approximation of p_C we used the ratio $\hat{p}_{C,n} = \frac{\sum_{i=1}^n \tilde{\tau}_i}{\sum_{i=1}^n \tau_i}$. Treating $(\tilde{\tau}_i, \tau_i)$, $i = 1, 2, \dots$, as i.i.d. random vectors with square-integrable coordinates, for $p_C := \frac{\mathbb{E}(\tilde{\tau}_1)}{\mathbb{E}(\tau_1)}$, from delta method it easily follows that for

$$\sigma := p_C \sqrt{\frac{\text{Var}(\tau_1)}{(\mathbb{E}(\tau_1))^2} + \frac{\text{Var}(\tilde{\tau}_1)}{(\mathbb{E}(\tilde{\tau}_1))^2} - 2 \frac{\text{Cov}(\tau_1, \tilde{\tau}_1)}{\mathbb{E}(\tilde{\tau}_1)\mathbb{E}(\tau_1)}} \quad (561)$$

we have $\sqrt{n}(\hat{p}_{C,n} - p_C) \Rightarrow \mathcal{N}(0, \sigma^2)$. For $\hat{\sigma}_n$ being an estimate of σ in which instead of means, variances, and covariances one uses their standard unbiased estimators computed using $(\tau_i, \tilde{\tau}_i)_{i=1}^n$, we have a.s. $\frac{\hat{\sigma}_n}{\sigma} \rightarrow 1$. Thus, from Slutsky's lemma, $\frac{\sqrt{n}}{\hat{\sigma}_n}(\hat{r}_n - r) \Rightarrow \mathcal{N}(0, 1)$, which can be used for constructing asymptotic confidence intervals for p_C . In Table (7) we provide the computed estimates in form $\hat{p}_{C,n} \pm \frac{\hat{\sigma}_n}{\sqrt{n}}$. It can be seen that these approximations of p_C are close for $q_{2,a}(x_0)$ and $p_{T,a}(x_0)$ when using in both cases CMC or IS with basis functions (113), and for $p_{T,a}(x_0)$ when using basis functions (114) for $M = 10$ and different p . However, such p_C differ significantly for other pairs of MC methods. In Table (7) we also provide estimates of practical inefficiency constants ic obtained by multiplying appropriate theoretical inefficiency constants computed earlier by the received approximations of p_C . From this table we can see that using IS led to considerable practical inefficiency constant reductions over using CMC.

	$q_{2,a}$		$p_{T,a}$				
	CMC	$M = 10$	CMC	$M = 10$	$M = 5, t$	$M = 10, t$	$M = 10, t^3$
p_C ($\cdot 10^{-6}s$)	33.612 ± 0.003	137.06 ± 0.01	33.998 ± 0.004	138.75 ± 0.01	96.36 ± 0.01	148.92 ± 0.01	149.74 ± 0.01
ic ($\cdot 10^{-3}s$)	245.4 ± 3.0	6.578 ± 0.084	47.3 ± 0.2	8.584 ± 0.013	6.072 ± 0.038	5.080 ± 0.024	2.597 ± 0.018

Table 7: Estimates of p_C and ic for computing $q_{2,1}$ and $p_{T,a}$ for various a , and basis functions as discussed in the main text.

9.3 Experiments comparing the spread of IS drifts.

In experiments described in this section we consider the assumptions as for estimation of $q_{2,a}(x_0)$ in Section 4.9. For estimators $\widehat{\text{est}}$ equal to each of $\widehat{\text{c\hat{e}}}$, $\widehat{\text{msq}}$, $\widehat{\text{msq}^2}$, and $\widehat{\text{ic}}$, we performed 20 independent SSM experiments for $n_1 = 100$ and $b' = 0$ as in Section 6.1, i.e. minimizing $b \rightarrow \widehat{\text{est}}_{n_1}(b', b)(\tilde{\chi}_1)$ for some $\tilde{\chi}_1$ as in that section and starting at $b' = 0$. For each such experiment for $\widehat{\text{est}} = \widehat{\text{ic}}$, for the same $\tilde{\chi}_1$ as in that experiment, we additionally carried out a two-phase minimization, in its first phase we minimizing $b \rightarrow \widehat{\text{msq}}_n(b', b)(\tilde{\chi}_1)$, and in the second $b \rightarrow \widehat{\text{ic}}_n(b', b)(\tilde{\chi}_1)$ using the first-phase minimization result as a starting point.

The IS drifts corresponding to IS parameters computed in the above experiments are shown in Figure 7, in which we also show (approximation of) the zero-variance IS drift r^* for the corresponding diffusion problem as in the previous section.




Figure 7: Drifts from SSM for estimating $q_{2,a}$, minimizing \hat{c}_e in (a), \widehat{msq} in (b), \widehat{msq}^2 in (c), and \hat{c} in (d). "Optimal" IS drift is a finite-difference approximation of the zero-variance drift r^* . 'msqic' in (d) corresponds to two-phase minimization and 'ic' to single-phase minimization as discussed in the main text.

In Figure 7 (d) it can be seen that ordinary (i.e. single-phase) SSM using \hat{c} yielded in three experiments IS drifts far from r^* , while in other 17 experiments and in all 20 experiments when using two-phase minimization we received drifts close to r^* . In experiments in which ordinary minimization led to drifts far from r^* , the value of $\hat{c}_n(b', b)(\tilde{\chi}_1)$ in the minimization result b was several times smaller than when using two-phase minimization, while in other cases these values were very close (e.g. the absolute value of difference of such values divided by smaller of them was in each case below 1%). In Figure 7, the IS drifts from SSM of \widehat{msq}^2 and two-phase minimization minimizing \hat{c} in the second phase as above seem to be the least spread, followed by the ones from minimization of \widehat{msq} , and finally \hat{c}_e .

From the below remark one can expect that for sufficiently large n_1 , IS drifts from SSM experiments like above should have approximately normal distribution in each point.

Remark 268. Let A , T , d_t , and r_t be as in Section 7.1 and let for some covariance matrix $D \in \mathbb{R}^{l \times l}$, $r_t(d_t - b^*) \Rightarrow \mathcal{N}(0, D)$. Note that this holds e.g.

for d_t being the SSM or MSM results of estimators \hat{g} for g replaced by \widehat{ce} , \widehat{msq} , $\widehat{msq2}$, or \widehat{ic} , for SSM under assumptions as in Section 7.5 for $D = u(b')V_g(b')$ and $r_t = t$, while for MSM under assumptions as in Section 7.7 for $D = V(d^*)$, $T = \mathbb{N}_+$, and $r_k = n_k$. Let us assume a linear parametrization of IS drifts as in (93), let $x \in \mathbb{R}^m$, and let $B \in \mathbb{R}^{d \times l}$ be such that $B_{i,j} = \tilde{r}_{i,j}(x)$, $i = 1, \dots, d$, $j = 1 \dots, l$. Then, we have

$$\sqrt{r_t}(r(d_t) - r(b^*))(x) \Rightarrow \mathcal{N}(0, BDB^T). \quad (562)$$

In the further experiments, to be able to carry out more simulations in a reasonable time, we changed the model considered by increasing the temperature 10 times. For such a new temperature we received an estimate 1.468 ± 0.002 of mean hitting time in CMC, as compared to 41.44 ± 0.15 under the original temperature as in Table 3. We carried out an MSM procedure of $\widehat{msq2}$ for $k = 6$ and $\underline{n}_i = 50 \cdot 2^{i-1}$, $i = 1, \dots, k$, receiving the final minimization result b_{MSM} . For est equal to each of different estimators as above and $b' = 0$ or $b' = b_{MSM}$, we carried out independently $N = 5000$ times SSM of \widehat{est} for $n_1 = 200$, in the i th SSM receiving result a_i and then computing $r(a_i)(0)$, i.e. the corresponding IS drift at zero, $i = 1, \dots, N$. The histograms of such IS drifts at zero for $b' = 0$, with fitted Gaussian functions, are shown in Figure 8.




Figure 8: Histograms of drifts at zero from SSM for computing $q_{2,a}$, minimizing \widehat{ce} in (a), \widehat{msq} in (b), $\widehat{msq2}$ in (c), and \widehat{ic} in (d).

The figure suggests that the distributions of drifts at zero are approximately

normal, as could be expected from Remark 268. Furthermore, the (empirical) distribution of drifts at zero for $b' = 0$ is in a sense the least spread when minimizing $\widehat{\text{msq}}^2$ and $\widehat{\text{ic}}$, followed by $\widehat{\text{msq}}$, and finally $\widehat{\text{ce}}$. The same observations can be made from inspection of histograms for the case of $b' = b_{MSM}$, which are not shown. We shall now compare quantitatively the spreads of empirical distributions of IS drifts at zero in the above experiments for different estimators and b' using interquartile ranges (IQRs), the definition and some required properties of which are provided in the below remark.

Remark 269. For i.i.d. random variables X_1, X_2, \dots , and $k, n \in \mathbb{N}_+$, let $X_{k:n}$ be the k -th coordinate of $\tilde{X}_n = (X_i)_{i=1}^n$ in the nondecreasing order. Let us define for $n \geq 4$, the interquartile range (IQR) of the coordinates of \tilde{X}_n as $\widehat{\text{IQR}}_n = X_{\lfloor \frac{3n}{4} \rfloor:n} - X_{\lfloor \frac{n}{4} \rfloor:n}$. Let further $X_1 \sim \mathcal{N}(\mu, \sigma^2)$ for some $\mu \in \mathbb{R}$ and $\sigma \in \mathbb{R}_+$, and let q denote the IQR of $\mathcal{N}(\mu, \sigma^2)$ (i.e. the difference of its third and first quartile). Then, for $d \approx 1.36$ we have $\sqrt{n}(\widehat{\text{IQR}}_n - q) \Rightarrow \mathcal{N}(0, dq^2)$ (see page 327 in [14]). Thus, for $\hat{\sigma}_n = \sqrt{d}\widehat{\text{IQR}}_n$ we have $\frac{\sqrt{n}}{\hat{\sigma}_n}(\widehat{\text{IQR}}_n - q) \Rightarrow \mathcal{N}(0, 1)$, which can be used for constructing asymptotic confidence intervals for q . Consider further X'_1, X'_2, \dots , i.i.d., $X'_1 \sim \mathcal{N}(\mu', (\sigma')^2)$, for some $\sigma' \in \mathbb{R}_+$, let $(X'_i)_{i \in \mathbb{N}_+}$ be independent of $(X_i)_{i \in \mathbb{N}_+}$, and let $q' \in \mathbb{R}_+$ be the IQR of $\mathcal{N}(\mu', (\sigma')^2)$. Then, for $\widehat{\text{IQR}}'_n$ analogous as above we have $(\widehat{\text{IQR}}_n - q, \widehat{\text{IQR}}'_n - q') \Rightarrow \mathcal{N}(0, d \text{diag}(q^2, (q')^2))$. Thus, for $R = \frac{q}{q'}$, $\hat{R}_n = \frac{\widehat{\text{IQR}}_n}{\widehat{\text{IQR}}'_n}$, and $\sigma_R = R\sqrt{2d}$, from delta method we have $\sqrt{n}(\hat{R}_n - R) \Rightarrow \mathcal{N}(0, \sigma_R^2)$. Therefore, for $\hat{\sigma}_{R,n} = \hat{R}_n\sqrt{2d}$ we have $\frac{\sqrt{n}}{\hat{\sigma}_{R,n}}(\hat{R}_n - R) \Rightarrow \mathcal{N}(0, 1)$, which can be used for constructing asymptotic confidence intervals for R .

For $X_i = r(a_i)(0)$, $i = 1, \dots, N$, received from SSM of different estimators as above, we computed the estimates $\widehat{\text{IQR}}_N$ of IQRs of drifts at zero and values $\frac{\hat{\sigma}_N}{\sqrt{N}}$ as in Remark 269. The results are provided in Table 8.

	ce	msq	msq 2	ic
$b' = 0$	0.736 ± 0.012	0.3770 ± 0.0062	0.1039 ± 0.0017	0.1006 ± 0.0017
$b' = b_{MSM}$	0.546 ± 0.009	0.2910 ± 0.0048	0.0932 ± 0.0015	0.0929 ± 0.0015

Table 8: Estimates of IQRs of drifts at zero from SSM for estimating $q_{2,a}(x_0)$, using different parameters b' .

From this table we can see that for both values of b' , the computed estimates of IQRs from minimization of $\widehat{\text{ic}}$ and $\widehat{\text{msq}}^2$ are the lowest, followed by such estimates from minimization of $\widehat{\text{msq}}$, and finally $\widehat{\text{ce}}$. The ratio of estimates of IQRs from minimization of $\widehat{\text{ce}}$ to $\widehat{\text{msq}}$ is 1.951 ± 0.045 for $b' = 0$ and 1.8771 ± 0.044 for $b' = b_{MSM}$ (where the results are provided in form $\hat{R}_n \pm \frac{\hat{\sigma}_{R,N}}{\sqrt{N}}$ under appropriate identifications with variables from Remark 269). Note that these ratios are close to 2. Intuitions supporting the above results are given in Section 9.4. Note also that the estimates of IQRs are lower when using $b' = b_{MSM}$ than $b' = 0$.

9.4 Intuitions behind some results of our numerical experiments

Recall that in numerical experiments in Section 9.2 we observed the fastest convergence of IS drifts in the MSM results and in Section 9.3 the lowest spreads of such drifts in the SSM results when minimizing $\widehat{\text{msq2}}$ and $\widehat{\text{ic}}$, followed by $\widehat{\text{msq}}$, and finally $\widehat{\text{ce}}$. Furthermore, in Section 9.3 the IQRs of values at zero of IS drifts corresponding to SSM results were approximately two times higher when minimizing $\widehat{\text{ce}}$ than $\widehat{\text{msq}}$. In this section we provide some intuitions behind these and some other experimental results. We will need a following remark.

Remark 270. *Let us assume that for b^* being the zero-variance IS parameter, similarly as in Section 7.8, for each $b \in \mathbb{R}^l$ we have $V_{\text{msq2}}(b) = V_{\text{ic}}(b) = 0$, $V_{\text{ce}}(b) = 4V_{\text{msq}}(b)$, and $V_{\text{ce}}(b)$ is positive definite. Let further, similarly as in Remark 268, for $d = 1$, for g replaced by each of ce , msq , msq2 , and ic , for $v_g = u(b')r(x)^T V_g(b')r(x)$ and $r_t = t$ for SSM or $v_g = r(x)^T V_g(b^*)r(x)$ and $r_k = n_k$ for MSM, the IS drifts corresponding to SSM and MSM results d_t of estimators \hat{g} fulfill*

$$\sqrt{r_t}(r(d_t) - r(b^*))(x) \Rightarrow \mathcal{N}(0, v_g). \quad (563)$$

Then, for g replaced by msq2 or ic we have $v_g = 0$ and the distribution $\mathcal{N}(0, v_g)$ has zero IQR. Furthermore, for $r(x) \neq 0$ we have $0 < v_{\text{ce}} = 4v_{\text{msq}}$, so that $\mathcal{N}(0, v_{\text{ce}})$ has positive IQR which is exactly two times higher than the IQR of $\mathcal{N}(0, v_{\text{msq}})$.

A possible reason why we received the above mentioned experimental results is that we can have approximately the same relations as in the above remark between the matrices $V_g(b)$ for $b = b'$ for SSM or $b = b^*$ for different b^* for MSM in our experiments, that is the entries of $V_g(b)$ can be much smaller for g equal to msq2 or ic than msq or ce and we can have $V_{\text{ce}}(b) \approx 4V_{\text{msq}}(b)$. This would lead to approximately the same relations between the asymptotic variances v_g and IQRs as in the above remark.

Such approximate relations between the matrices $V_g(b)$ can be a consequence of the IS distributions and densities corresponding to minimum points of the minimized functions being close to the zero-variance ones, thanks to which derivations as in Section 7.8 can be carried out approximately. For the estimation problems for whose diffusion counterparts there exist zero-variance IS drifts, like for the case of translated committors and MGF, we also have a following possible intuition behind the hypothesized approximate relations between matrices $V_g(b)$ as above. For the diffusion counterparts of these estimation problems, the zero-variance IS drifts minimize mean square, inefficiency constant, and cross-entropy among all appropriate drifts. Furthermore, as evidenced in Figure 6, the diffusion zero-variance IS drifts can be approximated very well using linear combinations of the IS basis functions considered. Thus, the diffusion IS drifts corresponding to minimizers of the functions considered are likely to be close to the zero-variance ones. Therefore, using such drifts in place of the zero-variance ones, the derivations as in Section 7.8 can be carried out approximately and we should have approximately the same relations between the matrices $V_g(b)$ for the diffusion case as in Remark 270. For small stepsizes h , like the ones used in our numerical experiments, the matrices $V_g(b)$ for the Euler scheme case can

be expected to be close to their diffusion counterparts and thus we should also have approximately the same relations between them as above.

For small stepsizes we can also expect the IS drifts corresponding to the minimizers of the functions considered for the Euler scheme case to be close to their diffusion counterparts, and thus, from the above discussion, also close to the diffusion zero-variance IS drifts. This would provide an intuition why in Figure 6 the IS drifts from minimization of various estimators of the functions considered are close to the approximations of the zero-variance IS drifts for the diffusion case.

In experiments from Section 9.2 for computing $p_{T,a}(x_0)$, the MSM results of \hat{ic} led to lower inefficiency constants than these of $\widehat{msq}2$, at the same time yielding higher variance and lower mean escape time. A possible intuition behind this result is that from Theorem 193, under appropriate assumptions a.s. we eventually should have such relations for some parameters converging a.s. to the minimum point of mean square and ones minimizing inefficiency constant (see Section 6.9 for some sufficient assumptions). Note, however, that this intuition fails when comparing variances in minimization results of \widehat{msq} and \hat{ic} , as the latter were smaller in all of our estimation experiments. A possible factor that could have contributed to the fact that in Section 9.2 we obtained the lowest inefficiency constants and variances when minimizing the new estimators \hat{ic} and \widehat{msq} , followed by \widehat{msq} , and then \hat{ce} , is that from the above hypothesis on the approximate relations of matrices $V_g(b)$ we may have the lowest spread of distributions of minimization results of the new estimators around the minimum points of variance and inefficiency constants, followed by such results for \widehat{msq} , and then \hat{ce} . We suspect that if sufficiently long minimization methods are performed (i.e. for sufficiently large n_1 for SSM or k for MSM), so that distributions of minimization results of the estimators considered become much less spread around the minimum points of their corresponding functions, then, as suggested by Theorem 193, the minimization results of \widehat{msq} should typically lead to lower variance than these of \hat{ic} . However, if the above hypothesis on the entries of $V_{msq}(b)$ being much smaller than these of $V_{msq2}(b)$ is correct, then, for longer minimization, the minimization results of $\widehat{msq}2$ should still typically lead to lower variance than these of \widehat{msq} . This is because such results d_t for $\widehat{msq}2$ would be asymptotically much more efficient for minimization of variance in different second-order senses as discussed in Section 7.2, e.g. in the sense of the mean of asymptotic distribution of $r_t(msq(d_t) - msq(b^*))$, equal to $\frac{u(b')}{2} \text{Tr}(V_g(b')H_{msq})$ for SSM or $\frac{1}{2} \text{Tr}(V_g(d^*)H_{msq})$ for MSM, being much smaller for g equal to $\widehat{msq}2$ than \widehat{msq} . Apart from the highest spread of distributions of minimization results when minimizing \hat{ce} , another factor that could have contributed to the higher variances in minimization results of \hat{ce} than for mean square estimators is that the minimum point of cross-entropy is likely to be different than that of mean square, so that, as discussed in Section 6.10, minimizing the mean square estimators can be more efficient in the first-order sense.

10 Conclusions and further ideas

In this work we developed methods for obtaining IS parameters via single- or multi-stage minimization of estimators of cross-entropy, mean square, and inefficiency constant, ensuring their various convergence and asymptotic properties

under appropriate assumptions in the ECM and LETGS settings. It would be interesting to prove such properties of our methods, or some their modifications, using other parametric families of IS distributions (see e.g. [37, 46] for some examples). We proposed ways of comparing the first- and second-order asymptotic efficiency of minimization methods of functions, which for such functions being equal to inefficiency constants can be used for comparing efficiency of methods for finding adaptive parameters in the first stage of a two-stage estimation method. The analysis of various asymptotic properties of such two-stage methods and discussion how to compare their asymptotic efficiency for estimation is relegated to a future paper.

Let us now discuss some problems which one can face when trying to implement and use in practice the methods for which we proved a.s. convergence or asymptotic properties of minimization results, as well as possible solutions to these problems. When using gradient-based stopping criteria in some of these methods, one should choose some nonnegative random bounds ϵ_i or $\tilde{\epsilon}_i$ on norms of gradients in minimization results, converging to zero a.s. (or equivalently, ensure that these gradients converge to zero a.s.). If chosen too large, such bounds can make the minimization algorithm perform no steps at all, and if taken too small, they can make the algorithm run longer than it can be afforded. To ensure a.s. convergence of gradients to zero in MSM methods and that a reasonable computational effort is made by the minimization algorithm in each stage, one can e.g. use at least a fixed number of steps of the minimization algorithm plus additional number needed to make the norm of gradient at least $q > 1$ times smaller than in the most recent step in which the final gradient was nonzero (if such step exists).

In MSM methods in which Condition 253 holds, which is useful for ensuring Condition 252 for $d^* = b^*$ as discussed in Remark 254, one should choose appropriate sets K_i such that Condition 154 holds. From $b_i \in K_i$, if for some $m \in \mathbb{N}_+$ the sets K_i contain b^* only for $i \geq m$, then convergence of b_i to b^* in Condition 252 may be very slow until i exceeds such m . One can try to deal with this problem for example by choosing all the sets K_i very large in the hope that they will all contain b^* , or by performing some preliminary MSM until the sequence of minimization results has approximately converged to b^* and then taking in a new MSM all the sets K_i containing some neighbourhood of the computed approximation of b^* .

As discussed in Section 7.7, as an alternative to using b_k converging to b^* in MSM, it may be reasonable to choose such b_k converging to d^* minimizing some measures of asymptotic inefficiency of d_k for minimization of f . Such b_k could be obtained e.g. using stochastic counterpart method minimizing estimators of such measures. An analogous idea is to use as the parameter b' in SSM an estimate of d^* minimizing the measure of inefficiency (450), which can be obtained e.g. using the stochastic counterpart method.

For IS in which mean theoretical cost is not constant in function of the IS parameter, minimizing the inefficiency constant estimator can be asymptotically the best option as under appropriate assumptions it can outperform minimization of other estimators in terms of the first-order asymptotic efficiency for minimization of inefficiency constant (see Section 6.10). However, if the mean cost is constant so that inefficiency constant is proportional to variance, then minimization results of mean square and inefficiency constant estimators considered can all converge to the minimum point of variance, so that their minimization

is asymptotically equally efficient in the first-order sense. In such case it may be reasonable to choose the estimator whose minimization results are the most efficient for minimization of variance in the second-order sense, as discussed in Section 7.2, e.g. the one minimizing mean of asymptotic distribution of variance as in that section. A possible idea is to estimate such asymptotic mean of different estimators, which can be combined with estimation of the parameters d^* minimizing such means as discussed above. An estimator, and potentially also an estimate of d^* as above, leading to the lowest estimates of asymptotic means, can be later used in SSM or can be chosen adaptively to construct the parameters b_k and minimization results in MSM.

In our numerical experiments, using different IS basis functions and added constants a led to considerably different inefficiency constants of the adaptive IS estimators. It would be interesting to develop adaptive methods for choosing such basis functions and constants. For instance, the added constant a can be chosen adaptively via minimization of estimators of variance or inefficiency constant in which a is treated as an additional minimization parameter.

In MSM, an alternative approach to minimization of estimators constructed using only the samples from the last stage, as in this work, would be to minimize some weighted average of such estimators from all previous stages. In our initial numerical experiments, minimizing such averages typically yielded drifts farther from the approximations of the zero-variance drift r^* for the corresponding diffusion than the approach from this work (data not shown), which is why we focused on the current approach. Similarly, α could be estimated using a weighted average of estimators from different stages, which closely resembles a purely adaptive approach used in stochastic approximation methods [31, 3, 37, 35]. For instance, under assumptions as in Section 6.1 and denoting $s_k = \sum_{i=1}^k n_i$, such estimator from the k th stage would be $\frac{1}{s_k} \sum_{l=1}^k \sum_{i=1}^{n_l} (ZL(b_{l-1}))(\chi_{l,i})$. A SLLN and CLT for such estimator can be proved similarly as in [35].

The model which we used for numerical experiments in this work is only a toy one. It would be interesting to test and compare performance of minimization of different estimators from this work on some realistic molecular models, as well as on models arising in other application areas of IS sampling, like computational finance and queueing theory. When using our methods for rare event simulation in practice one should take care to choose the IS parameter b equal to b' in SSM or b_0 in MSM so that the considered event is not too rare under the IS distribution $Q(b)$. This is because if such event was too rare, it could typically not happen at all in a reasonable simulation time. To find such b adaptively one can use e.g. some MSM method in which the problem is modified in the initial stages to make the rare event less rare in these stages as discussed at the end of Section 1.1.

Note that one can combine our methods with other EITs like stratified sampling or control variates. Note also that instead of using stochastic counterpart approach one can minimize inefficiency constants using stochastic approximation methods.

References

- [1] <http://math.stackexchange.com/questions/655825/proving-linear-independence-of-gaussian-functions>.

- [2] S. M. Ali and S. D. Silvey. A general class of coefficients of divergence of one distribution from another. *J. Roy. Statist. Soc. Ser. A*, 28(1):131–142, 1966.
- [3] B. Arouna. Adaptative Monte Carlo method. *Monte Carlo Methods Appl.*, 10(1):1–24, 2004.
- [4] S. Asmussen and P. W. Glynn. *Stochastic simulation: algorithms and analysis*. Stochastic Modelling and Applications. Springer, 2007.
- [5] T. Badowski. Variance-based sensitivity analysis for stochastic chemical kinetics. Master’s thesis, Faculty of Mathematics, Informatics and Mechanics, University of Warsaw, 2011.
- [6] T. Badowski. Variance-based sensitivity analysis and orthogonal approximations for stochastic models. Master’s thesis, Faculty of Physics, University of Warsaw, 2013.
- [7] P. Billingsley. *Probability and measure*. Wiley series in probability and mathematical statistics. Probability and mathematical statistics. Wiley, 1979.
- [8] S. Boyd and L. Vandenberghe. *Convex Optimization*. Cambridge University Press, New York, NY, USA, 2004.
- [9] J. A. Bucklew. *Introduction to rare event simulation*. Springer series in statistics. Springer, New York, 2004.
- [10] J. C. C. Chan, P. W. Glynn, and D. P. Kroese. A comparison of cross-entropy and variance minimization strategies. *Ann. Appl. Probab.*, 48A(3):183–194, 2011.
- [11] K. L. Chung. *A Course in Probability Theory*. Academic Press, 2001.
- [12] T. F. Coleman and Y. Li. On the convergence of interior-reflective Newton methods for nonlinear minimization subject to bounds. *Math. Program.*, 67(1-3):189–224, 1994.
- [13] T. F. Coleman and Y. Li. An interior trust region approach for nonlinear minimization subject to bounds. *SIAM J. Optim.*, 6(2):418–445, 1996.
- [14] A. DasGupta. *Probability for Statistics and Machine Learning: Fundamentals and Advanced Topics*. Springer Texts in Statistics. Springer, 2011.
- [15] P. Dupuis, K. Spiliopoulos, and H. Wang. Importance sampling for multiscale diffusions. *Multiscale Model. Simul.*, 10(1):1–27, July 2011.
- [16] P. Dupuis, K. Spiliopoulos, and X. Zhou. Escaping from an attractor: Importance sampling and rest points., 2012. <http://arxiv.org/abs/1303.0450>.
- [17] B. E. Fristedt and L. F. Gray. *A Modern Approach to Probability Theory*. Probability and Its Applications. Birkhäuser Boston, 1997.
- [18] P. Glasserman. *Monte Carlo Methods in Financial Engineering*. Applications of mathematics : stochastic modelling and applied probability. Springer, 2004.
- [19] P. W. Glynn. Efficiency improvement techniques. *Ann. Oper. Res.*, 53(1):175–197, 1994.
- [20] P. W. Glynn, H. P. Awad, and R. Y. Rubinstein. Zero-variance importance sampling estimators for Markov process expectations. *Math. Oper. Res., Articles in advance*, pages 1–30, 2012.
- [21] P. W. Glynn and R. Szechtman. Some new perspectives on the method of control variates. In K. Fang, H. Niederreiter, and F. J. Hickernell, editors, *Monte Carlo and Quasi-Monte Carlo Methods 2000*, pages 27–49. Springer Berlin Heidelberg, 2002.
- [22] P. W. Glynn and W. Whitt. The asymptotic efficiency of simulation estimators. *Oper. Res.*, 40(3):505–520, 1992.
- [23] E. Gobet and S. Menozzi. Stopped diffusion processes: Boundary corrections and overshoot. *Stochastic Process. Appl.*, 120(2):130 – 162, 2010.
- [24] C. Hartmann, R. Banisch, M. Sarich, T. Badowski, and C. Schütte. Characterization of rare events in molecular dynamics. *Entropy*, 16(1):350–376, 2014.
- [25] C. Hartmann and C. Schütte. Efficient rare event simulation by optimal nonequilibrium forcing. *J. Stat. Mech.*, 2012(11):P11004, 2012.
- [26] W. Hoeffding. The strong law of large numbers for U-statistics. Technical Report 302, Department of Statistics, Universty of North Carolina, 1961. Mimeograph series.
- [27] T. Homem-de Mello and R. Y. Rubinstein. Rare event estimation for static models via cross-entropy and importance sampling, 2002.

- [28] N. Ikeda and S. Watanabe. *Stochastic Differential Equations and Diffusion Processes*. North-Holland Pub. Co., 1981.
- [29] B. Jourdain and J. Lelong. Robust adaptive importance sampling for normal random vectors. *Ann. Appl. Probab.*, 19(5):1687–1718, 2009.
- [30] I. Karatzas and S. E. Shreve. *Brownian Motion and Stochastic Calculus*. Graduate Texts in Mathematics. Springer New York, 1991.
- [31] S. Kim and S. G. Henderson. Adaptive control variates for finite-horizon simulation. *Math. Oper. Res.*, 32(3):508–527, 2007.
- [32] P. E. Kloeden and E. Platen. *Numerical Solution of Stochastic Differential Equations*. Applications of mathematics : stochastic modelling and applied probability. Springer, 1992.
- [33] K. S. Kölbig and F. Schwartz. A program for solving systems of homogeneous linear inequalities. *Comput. Phys. Commun.*, pages 375–382, 1979.
- [34] D. P. Kroese, T. Taimre, and Z. I. Botev. *Handbook of Monte Carlo methods*. Wiley series in probability and statistics. Hoboken, N.J. Wiley, 2011.
- [35] B. Lapeyre and J. Lelong. A framework for adaptive Monte Carlo procedures. *Monte Carlo Methods Appl.*, 17(1):1–98, 2011.
- [36] E. L. Lehmann and J. P. Romano. *Testing statistical hypotheses*. Springer Texts in Statistics. Springer, New York, third edition, 2005.
- [37] V. Lemaire and G. Pages. Unconstrained recursive importance sampling. *Ann. Appl. Probab.*, 20(3):1–24, 2010.
- [38] F. Liese and I. Vajda. On divergences and informations in statistics and information theory. *IEEE Trans. Inform. Theory*, 52(10):4394–4412, 2006.
- [39] R. S. Lipster and A. N. Shiryaev. *Theory of Martingales*. Kluwer Academic, 1989.
- [40] B. L. Nelson. Control variate remedies. *Oper. Res.*, 38(6):974–992, 1990.
- [41] J. Nocedal and S. Wright. *Numerical Optimization*. Springer Series in Operations Research and Financial Engineering. Springer New York, 2006.
- [42] M. Pupashenko. Sampling methods for rare event simulation. Master’s thesis, Department of Mathematics, University of Kaiserslautern, 2011.
- [43] D. Revuz and M. Yor. *Continuous Martingales and Brownian Motion*. Grundlehren der mathematischen Wissenschaften A series of comprehensive studies in mathematics. Springer, 1999.
- [44] R. Rubinstein. The cross-entropy method for combinatorial and continuous optimization. *Methodol. Comput. Appl. Probab.*, 1:127–190, 1999.
- [45] R. Y. Rubinstein. Optimization of computer simulation models with rare events. *Eur. J. Oper. Res.*, 99:89–112, 1997.
- [46] R. Y. Rubinstein and D. P. Kroese. *The Cross Entropy Method: A Unified Approach To Combinatorial Optimization, Monte-Carlo Simulation and Machine Learning*. Information Science and Statistics. Springer-Verlag New York, Inc., Secaucus, NJ, USA, 2004.
- [47] R. Y. Rubinstein and A. Shapiro. *Discrete event systems: sensitivity analysis and stochastic optimization by the score function method*. Wiley series in probability and mathematical statistics. Wiley, 1993.
- [48] W. Rudin. *Real and Complex Analysis*. McGraw-Hill Book Co., New York, 1970.
- [49] A. Shapiro. Asymptotic behaviour of optimal solutions in stochastic programming. *Math. Oper. Res.*, 18(4):829–845, 1993.
- [50] A. Shapiro. Simulation-based optimization—convergence analysis and statistical inference. *Comm. Statist. Stochastic Models*, 12(3):425–454, 1996.
- [51] A. Shapiro. *Stochastic Programming*, chapter Monte Carlo sampling methods. Handbooks in Operations Research and Management Science. Elsevier, 2003.
- [52] A. W. van der Vaart. *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2000.
- [53] W. Zhang, H. Wang, C. Hartmann, M. Weber, and C. Schütte. Applications of the cross-entropy method to importance sampling and optimal control of diffusions. *SIAM J. Sci. Comput.*, 36(6):A2654–A2672, 2014.