

**ON NUMERICAL SEMIGROUPS CLOSED WITH RESPECT TO
THE ACTION OF CERTAIN AFFINE MAPS**

S. UGOLINI

ABSTRACT. In this paper we study numerical semigroups containing a positive integer and closed with respect to the action of an affine map of the form $x \mapsto ax + 1$, where a is a positive integer.

1. INTRODUCTION

A numerical semigroup G is a subsemigroup of the semigroup of non-negative integers $(\mathbf{N}, +)$ containing 0 and such that $\mathbf{N} \setminus G$ is finite.

For any $a \in \mathbf{N}^* := \mathbf{N} \setminus \{0\}$ and $b \in \mathbf{N}$ we define the affine map

$$\begin{aligned} \vartheta_{a,b} : \mathbf{N} &\rightarrow \mathbf{N} \\ x &\mapsto ax + b. \end{aligned}$$

We say that a numerical semigroup G is a $\vartheta_{a,b}$ -semigroup if $G^* := G \setminus \{0\}$ is closed with respect to the action of the map $\vartheta_{a,b}$, namely if $\vartheta_{a,b}(y) \in G^*$ for any $y \in G^*$.

The problem we will deal with along the paper consists in finding the smallest $\vartheta_{a,1}$ -semigroup containing a positive integer c , where a is a positive integer.

We notice in passing that Thabit numerical semigroups [2], namely numerical semigroups defined for any $n \in \mathbf{N}$ as

$$T(n) = \langle \{3 \cdot 2^{n+i} - 1 : i \in \mathbf{N}\} \rangle,$$

are $\vartheta_{2,1}$ -semigroups. More precisely, if we set $c := 3 \cdot 2^n$, then $T(n)$ is the smallest $\vartheta_{2,1}$ -semigroup containing c .

Also Mersenne numerical semigroups [1], namely numerical semigroups defined for any $n \in \mathbf{N}^*$ as

$$\langle \{2^{n+i} - 1 : i \in \mathbf{N}\} \rangle,$$

are $\vartheta_{2,1}$ -semigroups. In this case we set $c := 2^n$.

The paper is organized as follows: after a preliminary section where some notations are introduced, in Section 3 we deal with the aforementioned problem and show how to construct $\vartheta_{a,1}$ -semigroups.

2. NOTATIONS

We introduce some notations, we will use repeatedly along the paper.

2010 *Mathematics Subject Classification.* 11A05, 11A63, 11D04, 11D07, 20M14.

Key words and phrases. Numerical semigroups, Diophantine equations, Frobenius problem.

Definition 2.1. If y and z are two non-negative integers such that $y < z$, then

$$\begin{aligned} [y, z]_{\mathbf{N}} &:= \{x \in \mathbf{R} : y \leq x < z\} \cap \mathbf{N}; \\ [y, z]_{\mathbf{N}} &:= \{x \in \mathbf{R} : y \leq x \leq z\} \cap \mathbf{N}; \\ [y, +\infty]_{\mathbf{N}} &:= \{x \in \mathbf{R} : y \leq x < +\infty\} \cap \mathbf{N}. \end{aligned}$$

If $\{a, c\} \subseteq \mathbf{N}^*$ and $k \in \mathbf{N}$, then we define the sum

$$s_k(a) := \begin{cases} 0 & \text{if } k = 0, \\ \sum_{i=0}^{k-1} a^i & \text{otherwise,} \end{cases}$$

and accordingly

$$t_k(a, c) := a^k c + s_k(a).$$

Moreover, we define the set

$$S(a, c) := \{t_k(a, c) : k \in \mathbf{N}\}$$

and, for any non-negative integer \tilde{k} ,

$$S_{\tilde{k}}(a, c) := \{t_k(a, c) : k \in \mathbf{N} \text{ and } 0 \leq k \leq \tilde{k}\}.$$

Finally, we define

$$H(a, c) := \langle S(a, c) \rangle$$

and denote by $G_{a,1}(c)$ the smallest $\vartheta_{a,1}$ -semigroup containing c .

3. STRUCTURE AND CONSTRUCTION OF $\vartheta_{a,1}$ -SEMIGROUPS

We fix once for all this section a subset $\{a, c\} \subseteq \mathbf{N}^*$.

In the first part of this section we prove some technical lemmas which let us prove in Theorem 3.5 that

$$G_{a,1}(c) = H(a, c).$$

Lemma 3.1. *We have that $S(a, c) \subseteq G_{a,1}(c)$.*

Proof. We prove the claim by induction on $k \in \mathbf{N}$.

If $k = 0$, then $t_0(a, c) = c \in G_{a,1}(c)$.

Suppose now that $t_k(a, c) \in G_{a,1}(c)$ for some non-negative integer k . Then,

$$t_{k+1}(a, c) = a \cdot t_k(a, c) + 1 = \vartheta_{a,1}(t_k(a, c)) \in G_{a,1}(c).$$

□

Lemma 3.2. *$H(a, c)$ is a subsemigroup of $(\mathbf{N}, +)$ closed with respect to the action of the map $\vartheta_{a,1}$.*

Proof. By definition, $H(a, c)$ is a subsemigroup of $(\mathbf{N}, +)$.

We prove that $H(a, c)$ is closed with respect to the action of the map $\vartheta_{a,1}$.

Consider an element

$$y = \sum_{k \in K} j_k \cdot t_k(a, c) \in H(a, c),$$

where K is a non-empty finite subset of \mathbf{N} and $\{j_k\}_{k \in K}$ is a set of positive integers. Let \tilde{k} be a chosen element of K . Then,

$$\begin{aligned}
ay + 1 &= \sum_{k \in K} aj_k \cdot t_k(a, c) + 1 \\
&= \sum_{\substack{k \in K \\ k \neq \bar{k}}} aj_k \cdot t_k(a, c) + a(j_{\bar{k}} - 1) \cdot t_{\bar{k}}(a, c) + a \cdot t_{\bar{k}}(a, c) + 1 \\
&= \sum_{\substack{k \in K \\ k \neq \bar{k}}} aj_k \cdot t_k(a, c) + a(j_{\bar{k}} - 1) \cdot t_{\bar{k}}(a, c) + (t_{\bar{k}+1}(a, c) - 1) + 1 \\
&= \sum_{\substack{k \in K \\ k \neq \bar{k}}} aj_k \cdot t_k(a, c) + a(j_{\bar{k}} - 1) \cdot t_{\bar{k}}(a, c) + t_{\bar{k}+1}(a, c).
\end{aligned}$$

Since this latter is a linear combination of elements in $S(a, c)$ with coefficients in \mathbf{N} , we conclude that $ay + 1 \in H(a, c)$. \square

Lemma 3.3. *If $\{m, n\} \subseteq \mathbf{N}$, where $n \geq c - 1$, and*

$$[mc, mc + n]_{\mathbf{N}} \subseteq H(a, c),$$

then

$$[mc, +\infty[_{\mathbf{N}} \subseteq H(a, c).$$

Proof. According to Lemma 3.2, $H(a, c)$ is a subsemigroup of $(\mathbf{N}, +)$ closed with respect to the action of the map $\vartheta_{a,1}$.

If y is an integer such that $y \geq mc$, then

$$\begin{cases} y - mc = cq + r \\ 0 \leq r < c \end{cases}$$

for some non-negative integers q and r , where $r \leq n$.

Since

$$mc + r \in [mc, mc + n]_{\mathbf{N}} \subseteq H(a, c)$$

and $qc \in H(a, c)$, we conclude that $y \in H(a, c)$. Therefore,

$$[mc, +\infty[_{\mathbf{N}} \subseteq H(a, c).$$

\square

Lemma 3.4. *For any $k \in \mathbf{N}$ we have that $[a^k c, t_k(a, c)]_{\mathbf{N}} \subseteq H(a, c)$.*

Proof. We prove the claim by induction on k .

Proving the base step is trivial, since $c \in H(a, c)$ and

$$[a^0 c, t_0(a, c)]_{\mathbf{N}} = [c, c]_{\mathbf{N}} \subseteq H(a, c).$$

Suppose now that $[a^k c, t_k(a, c)]_{\mathbf{N}} \subseteq H(a, c)$ for some $k \in \mathbf{N}$. Then,

$$a \cdot (a^k c + r) = a^{k+1} c + ar \in H(a, c)$$

for any $r \in [0, s_k(a)]_{\mathbf{N}}$. Moreover,

$$a^{k+1} c + ar + j \in H(a, c)$$

for any $r \in [0, s_k(a)]_{\mathbf{N}}$ and any $j \in [1, a]_{\mathbf{N}}$, since

$$a^{k+1} c + ar + j = (a - j) \cdot (a^k c + r) + j \cdot (a^k c + r + 1).$$

Therefore,

$$[a^{k+1} c, a^{k+1} c + s_k(a) \cdot a]_{\mathbf{N}} \subseteq H(a, c).$$

Finally,

$$\begin{aligned} a^{k+1}c + s_k(a) \cdot a &= a^{k+1}c + \sum_{i=1}^k a^i = a \cdot t_k(a, c) \in H(a, c), \\ t_{k+1}(a, c) &= \vartheta_{a,1}(t_k(a, c)) \in H(a, c). \end{aligned}$$

All considered, we get that $[a^{k+1}c, t_{k+1}(a, c)]_{\mathbf{N}} \subseteq H(a, c)$ and the inductive step is proved. \square

Theorem 3.5. *We have that $G_{a,1}(c) = H(a, c)$.*

Proof. First, we notice that $H(a, c) \subseteq G_{a,1}(c)$ according to Lemma 3.1.

According to Lemma 3.2, $H(a, c)$ is a subsemigroup of $(\mathbf{N}, +)$ closed with respect to the action of the map $\vartheta_{a,1}$.

Let k be a positive integer such that $s_k(a) \geq c - 1$.

According to Lemma 3.4 we have that

$$[a^k c, t_k(a, c)]_{\mathbf{N}} \subseteq H(a, c).$$

Moreover, according to Lemma 3.3, $\mathbf{N} \setminus H(a, c)$ is finite too.

All considered, $H(a, c)$ is a numerical semigroup and $G_{a,1}(c) = H(a, c)$. \square

The following technical lemma let us prove Theorem 3.7. As a consequence of such a theorem we deduce Corollary 3.8, where the embedding dimension $e(G_{a,1}(c))$ of $G_{a,1}(c)$, namely the cardinality of the minimal set of generators of $G_{a,1}(c)$, is given.

Lemma 3.6. *If $k \in \mathbf{N}^*$ and $\{j_i\}_{i \in K}$ is a set of positive integers indexed on a non-empty set $K \subseteq [0, k-1]_{\mathbf{N}}$ such that $\sum_{i \in K} j_i \cdot a^i \leq a^k$, then*

$$\sum_{i \in K} j_i \cdot t_i(a, c) < t_k(a, c).$$

Proof. We distinguish three different cases.

- If $K = \{0\}$, then $j_0 \leq a^k$ and

$$\sum_{i \in K} j_i \cdot t_i(a, c) = j_0 \cdot c < a^k \cdot c + s_k(a) = t_k(a, c).$$

- If $K \neq \{0\}$ but the cardinality of K is still one, then

$$\sum_{i \in K} j_i \cdot t_i(a, c) = j_r \cdot t_r(a, c)$$

for some $r \in [0, k-1]_{\mathbf{N}}$ and some $j_r \in [0, a^{k-r}]_{\mathbf{N}}$. Therefore,

$$\begin{aligned} \sum_{i \in K} j_i \cdot t_i(a, c) &= j_r \cdot (a^r c + s_r(a)) \\ &\leq a^k c + a^{k-r} \cdot s_r(a) = a^k c + \sum_{i=k-r}^{k-1} a^i \\ &< t_k(a, c). \end{aligned}$$

- If the cardinality of K is greater than one, then $a \neq 1$. Indeed,

$$\sum_{i \in K} j_i 1^i \leq 1^k$$

if and only if $|K| = 1$ and the only integer j_i is equal to 1.
Therefore, assuming $a \neq 1$, we have that

$$\begin{aligned}
\sum_{i \in K} j_i \cdot t_i(a, c) &= \sum_{i \in K} j_i \cdot a^i c + \sum_{i \in K} j_i \cdot s_i(a) \\
&\leq a^k \cdot c + \left(\sum_{i \in K} j_i \cdot \frac{a^i - 1}{a - 1} \right) \\
&= a^k \cdot c + \frac{1}{a - 1} \cdot \left(\sum_{i \in K} j_i \cdot a^i - \sum_{i \in K} j_i \right) \\
&\leq a^k \cdot c + \frac{1}{a - 1} \cdot \left(a^k - \sum_{i \in K} j_i \right) \\
&\leq a^k \cdot c + \frac{1}{a - 1} \cdot (a^k - 2) \\
&< a^k \cdot c + \frac{a^k - 1}{a - 1} = t_k(a, c).
\end{aligned}$$

□

Theorem 3.7. Let $\tilde{k} := \min\{k \in \mathbf{N} : s_k(a) \geq c - 1\}$.

Then, the following hold.

- (1) If $s_{\tilde{k}}(a) = c - 1$, then $S_{\tilde{k}}(a, c)$ is a minimal set of generators for $G_{a,1}(c)$.
- (2) If $s_{\tilde{k}}(a) > c - 1$, then $S_{\tilde{k}-1}(a, c)$ is a minimal set of generators for $G_{a,1}(c)$.

Proof. Let k be a positive integer such that $s_k(a) < c$. Then,

$$t_k(a, c) \in [a^k c, (a^k + 1)c]_{\mathbf{N}}.$$

Consider now an element

$$\sum_{i \in K} j_i \cdot t_i(a, c) \in [0, (a^k + 1)c]_{\mathbf{N}},$$

where $\{j_i\}_{i \in K}$ is a set of positive integers indexed on a set $K \subseteq [0, k]_{\mathbf{N}}$.

We have that

$$\sum_{i \in K} j_i \cdot a^i c \leq \sum_{i \in K} j_i \cdot t_i(a, c) < (a^k + 1) \cdot c.$$

As a consequence,

$$\sum_{i \in K} j_i \cdot a^i \leq a^k.$$

Hence, in accordance with Lemma 3.6, we have that

$$\sum_{i \in K} j_i \cdot t_i(a, c) < t_k(a, c).$$

All considered, we conclude that

$$t_k(a, c) \notin \langle \{t_i(a, c) : 0 \leq i < k\} \rangle$$

for any positive integer k such that $s_k(a) < c$.

We prove now separately the 2 statements in the claim of the theorem.

(1) If $s_{\tilde{k}}(a) = c - 1 < c$, then

$$t_k(a, c) \notin \langle \{t_i(a, c) : 0 \leq i < k\} \rangle$$

for any positive integer $k \leq \tilde{k}$.

Nevertheless, $S_{\tilde{k}}(a, c)$ is a minimal set of generators for $G_{a,1}(c)$, since

$$[a^{\tilde{k}}c, +\infty[_{\mathbf{N}} \subseteq G_{a,1}(c)$$

according to Lemma 3.3.

(2) If $s_{\tilde{k}}(a) > c - 1$, then

$$t_k(a, c) \notin \langle \{t_i(a, c) : 0 \leq i < k\} \rangle$$

for any positive integer $k < \tilde{k}$, while

$$t_{\tilde{k}}(a, c) \in \langle \{t_i(a, c) : 0 \leq i < \tilde{k}\} \rangle.$$

The latter assertion holds since

$$\begin{cases} t_{\tilde{k}}(a, c) - a^{\tilde{k}}c = qc + r \\ 0 \leq r < c \end{cases}$$

for some non-negative integers q and r . Since

$$a^{\tilde{k}}c \leq a^{\tilde{k}}c + r < t_{\tilde{k}}(a, c)$$

we have that

$$a^{\tilde{k}}c + r \in G_{a,1}(c)$$

according to Lemma 3.4. Indeed,

$$\{a^{\tilde{k}}c + r, qc\} \subseteq \langle S_{\tilde{k}-1}(a, c) \rangle.$$

Therefore,

$$t_{\tilde{k}}(a, c) \in \langle S_{\tilde{k}-1}(a, c) \rangle,$$

namely $S_{\tilde{k}-1}(a, c)$ is a minimal set of generators for $G_{a,1}(c)$. □

Corollary 3.8. *Let $\tilde{k} := \min\{k \in \mathbf{N} : s_k(a) \geq c - 1\}$.*

Then, the following hold.

(1) *If $s_{\tilde{k}}(a) = c - 1$, then $e(G_{a,1}(c)) = \tilde{k} + 1$.*

(2) *If $s_{\tilde{k}}(a) > c - 1$, then $e(G_{a,1}(c)) = \tilde{k}$.*

Proof. The result follows immediately from Theorem 3.7 since $|S_k(a, c)| = k + 1$ for any $k \in \mathbf{N}$. □

Remark 3.9. In [2, Theorem 6] it is claimed that $e(T(n)) = n + 2$ for any $n \in \mathbf{N}$. Indeed, if we set

$$c := 3 \cdot 2^n - 1,$$

then

$$c - 1 = 3 \cdot 2^n - 2 = 2^{n+1} + 2^n - 2.$$

If $k \in \mathbf{N}^*$, then

$$s_k(2) = \sum_{i=0}^{k-1} 2^i > c - 1$$

if and only if

$$k \geq n + 2.$$

Adopting the notations introduced in the current paper, $a = 2$ and $T(n) = G_{2,1}(c)$. Therefore, according to Corollary 3.8, $\tilde{k} = n + 2$ and $s_{\tilde{k}}(2) > c - 1$. All considered,

$$e(G_{2,1}(c)) = e(T(n)) = n + 2$$

in accordance with [2, Theorem 6].

In the following theorem we estimate the value for the Frobenius number $F(G_{a,1}(c))$, which is the greatest integer which does not belong to $G_{a,1}(c)$.

Theorem 3.10. *Let $\tilde{k} := \min\{k \in \mathbf{N} : s_k(a) \geq c - 1\}$.*

Then,

$$a^{\tilde{k}-1}c + ac - 1 \leq F(G_{a,1}(c)) \leq a^{\tilde{k}}c - 1.$$

Proof. According to Lemma 3.3, we have that

$$[a^{\tilde{k}}c, +\infty[_{\mathbf{N}} \subseteq G_{a,1}(c).$$

Hence,

$$F(G_{a,1}(c)) \leq a^{\tilde{k}}c - 1.$$

Consider now the integer

$$a^{\tilde{k}-1}c + ac - 1 = (a^{\tilde{k}-1} + 1)c - 1 + (a - 1)c.$$

We notice that

$$(a^{\tilde{k}-1} + 1)c - 1 \in [t_{\tilde{k}-1}(a, c) + 1, (a^{\tilde{k}} + 1)c[_{\mathbf{N}}.$$

Let

$$T := \sum_{i \in K} j_i \cdot t_i(a, c),$$

where $K \subseteq [0, \tilde{k} - 1]_{\mathbf{N}}$ and $\{j_i\}_{i \in K}$ is a set of positive integers.

If $\tilde{k} - 1 \in K$, then either $T = t_{\tilde{k}-1}(a, c)$ or $T \geq t_{\tilde{k}-1}(a, c) + c$. In both cases, $T \neq (a^{\tilde{k}-1} + 1)c - 1$.

Suppose now that $K \subseteq [0, \tilde{k} - 2]_{\mathbf{N}}$.

If $\sum_{i \in K} j_i \cdot a^i \leq a^{\tilde{k}-1}$, then

$$\sum_{i \in K} j_i \cdot t_i(a, c) < t_{\tilde{k}-1}(a, c)$$

according to Lemma 3.6.

If $\sum_{i \in K} j_i \cdot a^i > a^{\tilde{k}-1}$, then

$$\sum_{i \in K} j_i \cdot t_i(a, c) \geq (a^{\tilde{k}} + 1)c.$$

All considered,

$$a^{\tilde{k}-1}c + ac - 1 = (a^{\tilde{k}-1} + 1)c - 1 + (a - 1)c \notin \langle S_{\tilde{k}-1}(a, c) \rangle.$$

Therefore,

$$a^{\tilde{k}-1}c + ac - 1 \leq F(G_{a,1}(c)).$$

□

Example 3.11. In this example we list all the elements of $G_{3,1}(3)$.

Adopting the notations introduced above we have that

$$a = 3, \quad c = 3, \quad \tilde{k} = 2$$

since

$$\min\{k \in \mathbf{N} : s_k(3) \geq 2\} = 2.$$

According to Theorem 3.7 we have that

$$G_{3,1}(3) = \langle \{t_0(3, 3), t_1(3, 3)\} \rangle.$$

Moreover,

$$[3^2 \cdot 3, +\infty[_{\mathbf{N}} \subseteq G_{3,1}(3)$$

according to Lemma 3.3 and Lemma 3.4.

The non-negative integers belonging to $G_{3,1}(3)$ are listed in the following table (the numbers in bold are the generators of $G_{3,1}(3)$):

0	3	6	9	12	15	18	21	24
			10	13	16	19	22	25
						20	23	26

Finally we note that

$$F(G_{3,1}(3)) = 17 = a^{\tilde{k}-1}c + ac - 1.$$

Example 3.12. In this example we list all the elements of $G_{3,1}(5)$.

We have that

$$a = 3, \quad c = 5, \quad \tilde{k} = 2.$$

Since $s_2(3) = c - 1$, we have that

$$G_{3,1}(5) = \langle \{t_0(3, 5), t_1(3, 5), t_2(3, 5)\} \rangle$$

according to Theorem 3.7.

Moreover,

$$[3^2 \cdot 5, +\infty[_{\mathbf{N}} \subseteq G_{3,1}(5)$$

according to Lemma 3.3 and Lemma 3.4.

The non-negative integers belonging to $G_{3,1}(5)$ are listed in the following table (the numbers in bold are the generators of $G_{3,1}(5)$):

0	5	10	15	20	25	30	35	40	45
			16	21	26	31	36	41	46
						32	37	42	47
									48
									49

Finally we note that

$$F(G_{3,1}(5)) = 44 = a^{\tilde{k}}c - 1.$$

REFERENCES

1. J. C. Rosales, M.B. Branco, and D. Torrão, *The Frobenius problem for Mersenne numerical semigroups*, Preprint.
2. ———, *The Frobenius problem for Thabit numerical semigroups*, J. Number Theory **155** (2015), 85–99.

E-mail address: sugolini@gmail.com