

Equivariant algebraic K-theory

Mona Merling

Abstract

A group action on the input ring or category induces an action on the algebraic K -theory spectrum. However, a shortcoming of this naive approach to equivariant algebraic K -theory is, for example, that the map of spectra with G -action induced by a G -map of G -rings is not equivariant. We define a version of equivariant algebraic K -theory which encodes a group action on the input in a functorial way to produce a *genuine* algebraic K -theory G -spectrum for a finite group G . The main technical work lies in studying coherent actions on the input category. A payoff of our approach is that it builds a unifying framework for equivariant topological K -theory, Atiyah's Real K -theory, and existing statements about algebraic K -theory spectra with G -action. We recover the map from the Quillen-Lichtenbaum conjecture and the representational assembly map studied by Carlsson and interpret them from the perspective of equivariant stable homotopy theory. We also give a definition of an equivariant version of Waldhausen's A -theory of a G -space.

Contents

1	Introduction	2
2	Categorical homotopy fixed points	7
3	Pseudo equivariance	15
4	Homotopy fixed points of module categories	18
5	Equivariant algebraic K-theory of G-rings	26
6	Equivariant K-theory of Galois extensions	39
7	Equivariant algebraic K-theory of topological rings	45
8	Definition of equivariant A-theory	47
9	Transfer maps and relationship to spectral Mackey functor approach	52

1 Introduction

Algebraic K -theory is the meeting ground for various subjects such as algebraic geometry, number theory, and algebraic topology. Despite computations being very hard, algebraic K -theory is intensely studied precisely because of the deep connections it has to other fields. For example, Quillen's algebraic K -groups of number rings can be used to formulate a generalization of the class number formula, and knowledge of the K -theory of \mathbb{Z} would settle the Kummer-Vandiver conjecture, whereas Waldhausen's A -theory, an extension of algebraic K -theory to spaces, encodes information about pseudoisotopies of manifolds.

Often enough the objects whose K -theory one is interested in understanding (such as rings, ring spectra, spaces) have inherent group actions on them, and by functoriality of the K -theory construction, their associated K -theory spectra are spectra with group action. Nevertheless, equivariant stable homotopy theory is more intricate than just the study of group actions on spectra. Spectra with group action, which are suggestively called *naive* G -spectra, do not allow for the usual duality phenomena of stable homotopy theory that one has nonequivariantly (such as Poincaré duality for G -manifolds.) In order to rectify this, one needs to allow suspension by representation spheres S^V , which are one point compactifications of representations V of the group of equivariance G . This leads to the notion of a *genuine* G -spectrum, which is indexed on representations of G and in turn represents a cohomology theory graded on representations. The aim of this paper is to study algebraic K -theory from the genuine equivariant perspective: we construct and study a genuine algebraic K -theory G -spectrum in the case when a finite group G acts on the input.

Equivariant stable homotopy theory is a very fascinating subject in its own right (and has been studied as such, for example, in the early work of [LMS86]), but what has really driven its development is the fact that the powerful tools of genuine equivariant homotopy theory can shed light on nonequivariant problems. An early example of such an application is Carlsson's proof of the Segal conjecture, which states that the zeroth stable cohomotopy group of the classifying space of a group is isomorphic to a completion of the Burnside ring of the group. The proof of this nonequivariant statement requires heavy use of equivariant theory (see [Car84]). A more recent example is the Hill-Hopkins-Ravenel solution to the Kervaire invariant one problem, which is ultimately a statement about manifolds – the proof, however, unexpectedly relies on sophisticated equivariant machinery (see [HHR]). The solution to the Kervaire invariant one problem has reinvigorated interest in equivariant homotopy theory, since it demonstrates the power of equivariant techniques for solving problems in nonequivariant homotopy theory.

Moreover, the study of algebraic K -theory is linked to equivariant stable homotopy theory: one of the main computational approaches is approximating K -theory by topological cyclic homology, which in many cases can be computed using tools of genuine S^1 equivariant stable homotopy theory. However, these methods do not take into account an inherent action on the input ring (or category), which is what we want to capture. On the other hand, Galois group actions have provided organizing principles for studying algebraic K -theory. It has long been suspected that the K -theory of a field should be computable in terms of the K -theory of the algebraic closure and the action of the absolute Galois group – one of the early Quillen-Lichtenbaum conjectures was that the map from fixed points to homotopy fixed points of naive G -spectra is an equivalence after p -completion. Thomason later showed that in order to obtain

an equivalence one needs to invert a “Bott” element in K -theory and reduce mod a prime power. The concept of descent and the Quillen-Lichtenbaum conjecture has motivated Carlsson’s program to study the K -theory of fields in terms of the representational assembly map for a Galois extension E/F induced by tensoring a G -representations over F with E . The aim of this paper is provide the framework that allows us to interpret these maps as maps of genuine G -spectra or their fixed points, thus making the tools of stable equivariant homotopy theory directly available for the study of K -theory.

We describe our philosophy for defining equivariant algebraic K -theory: If the input has a G -action, this induces a G -action on the category that one builds algebraic K -theory out of. For example, if R is a G -ring, then the category of finitely generated projective R -modules and isomorphisms $\text{iso } \mathcal{P}(R)$ has a G -action: for a module M , gM is defined by twisting the scalar multiplication on R by g . One similarly obtains a G -action on the category of modules over a G -ring spectrum R . If X is a G -space, then the category $R(X)$ of retractive spaces over X has a G -action. For a retractive space Y , gY is defined by precomposing the inclusion map by g^{-1} and postcomposing the retraction map by g . However, by applying the nonequivariant constructions to these categories with G -action, we obtain just a spectra with G -action, and not genuine G -spectra – the K -theory G -spaces we obtain have deloopings with respect to all spheres S^n with trivial G -action, but it does not deloop with respect to representation spheres S^V . We need to modify these categories with G -action to turn them into “genuine” G -categories, very loosely speaking. We try to make this more precise in the next paragraphs.

Nonequivariantly, the algebraic K -theory space of R is defined as the group completion of the classifying space of the symmetric monoidal category $\text{iso } \mathcal{P}(R)$, and this space is delooped using an infinite loop space machine such as the operadic one developed by May in [May72] or the one based on Γ -spaces developed by Segal in [Seg74]. These nonequivariant machines are equivalent by a celebrated theorem of May and Thomason [MT78]. The Segalic infinite loop space machine has been generalized equivariantly by Shimakawa in [Shi89], and the operadic infinite loop space machine has been generalized equivariantly by Guillou and May in [GMa], to give *genuine* Ω - G -spectra with zeroth space the group completion of the input category. We describe these machines in §5 where we use them, and we note that we have shown in [MMO] that when fed equivalent input, they produce equivalent G -spectra. But, the input these equivariant infinite loop space machines take is not simply symmetric monoidal categories with G -action – their input is *genuine symmetric monoidal G -categories*. Genuine permutative G -categories have been defined in [GMa] as algebras over an equivariant version of the Barrat-Eccles operad, and we have defined genuine symmetric monoidal G -categories as pseudo algebras over the same operad in [GMMO]. We will not dwell on this since all the genuine symmetric monoidal G -categories we consider in this paper arise in the concrete way described in the following paragraph.

Definition 1.1. Let \tilde{G} be the category with objects the elements of G and a unique morphism between any two objects, with G acting by translation on the objects and diagonally on the morphisms. For a G -category \mathcal{C} , let $\mathcal{C}at(\tilde{G}, \mathcal{C})$ be the category of all functors and all natural transformations, with G acting by conjugation.

Note that \tilde{G} is G -isomorphic to the translation category of G , and its classifying space $B\tilde{G}$ is equivalent to EG . If \mathcal{C} is a symmetric monoidal category with G -action, then it turns out that $\mathcal{C}at(\tilde{G}, \mathcal{C})$ is an example of a genuine symmetric monoidal G -category, and therefore, it is input for the equivariant infinite loop space machines. We will show that replacing a

symmetric monoidal category with G -action \mathcal{C} with $\mathcal{C}at(\tilde{G}, \mathcal{C})$ not only makes it a genuine symmetric monoidal category, but it also fixes coherence issues that arise equivariantly. Even if the action does not preserve the symmetric monoidal structure strictly, but only up to coherent isomorphism, this can be rectified after applying $\mathcal{C}at(\tilde{G}, -)$.

We define the equivariant algebraic K -theory G -spectrum $\mathbf{K}_G(R)$ of a G -ring R as the Ω - G -spectrum obtained by applying one of the equivariant infinite loop space machines to the category $\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))$. We summarize some of the properties of $\mathbf{K}_G(R)$ that we prove.

Theorem 1.2. *For finite groups G , the assignment*

$$R \mapsto \mathbf{K}_G(R)$$

can be extended to a functor from G -rings and G -maps to genuine (connective) Ω - G spectra, with the following properties

1. *For the topological rings \mathbb{C} and \mathbb{R} with trivial G -action for any finite group G ,*

$$\mathbf{K}_G(\mathbb{C}) \simeq ku_G \text{ and } \mathbf{K}_G(\mathbb{R}) \simeq ko_G,$$

where ku_G and ko_G are connective versions of equivariant topological K -theory;

2. *For the topological ring \mathbb{C} with $\mathbb{Z}/2$ conjugation action*

$$\mathbf{K}_{\mathbb{Z}/2}(\mathbb{C}) \simeq kr,$$

where kr is a connective version of Atiyah's Real K -theory;

3. *If $|H|^{-1} \in R$, then*

$$\mathbf{K}_G(R)^H \simeq \mathbf{K}(R_H[H]),$$

where $R_H[H]$ is the twisted group ring;

4. *For a Galois extension of rings $R \rightarrow S$ with Galois group G ,*

$$\mathbf{K}_G(S)^G \simeq \mathbf{K}(R);$$

5. *\mathbf{K}_G is invariant under a suitable notion of equivariant Morita equivalence;*

6. *For a finite Galois extension with group G , the map from fixed points to homotopy fixed points of **genuine** G -spectra*

$$\mathbf{K}_G(E)^G \longrightarrow \mathbf{K}_G(E)^{hG}$$

*is equivalent to the map from fixed points to homotopy fixed points of **naive** G -spectra $\mathbf{K}(F) \rightarrow \mathbf{K}(E)^G$ from the Quillen-Lichtenbaum conjecture.*

7. *For a finite Galois extension with group G , the representational assembly map defined by Carlsson*

$$\mathbf{K}(\text{Rep}_F[G]) \rightarrow \mathbf{K}F$$

is the fixed point map of G -map of genuine G -spectra

$$\mathbf{K}_G(F) \longrightarrow \mathbf{K}_G(E).$$

In order to deduce the first two results about topological rings, we connect the definition that we give of equivariant algebraic K -theory to equivariant bundle theory. Nonequivariantly, Quillen's plus construction $BGL(R)^+$ is the zeroth component of the group completion of the monoid of classifying spaces of principal $GL_n(R)$ -bundles. Equivariantly, we show that the K -theory space (which we define in terms of the G -category of projective modules) is also equivalent to the equivariant group completion of the monoid of, in this case, equivariant $GL_n(R)$ -bundles. For this we use the models for equivariant bundles that we have found in [GMM]. This connection allows us to recover equivariant topological real and complex K -theory, and Atiyah's Real K -theory as examples of our construction.

For the rest of the results stated above, we need to analyze the fixed point spectrum $\mathbf{K}_G(R)^H$ for subgroups $H \subseteq G$. One of the formal properties of the equivariant infinite loop space machines is that they commute with fixed points, so our task amounts to studying the fixed point categories $\mathcal{C}at(\tilde{G}, \mathcal{C})^H$ for suitable G -categories \mathcal{C} . By analogy with the homotopy fixed point set of a G -space, we define the *homotopy fixed points* of a G -category \mathcal{C} as the fixed point category $\mathcal{C}at(\tilde{G}, \mathcal{C})^G$, which we introduce and study in §2. This is the category of G -equivariant functors and natural transformations, which Thomason called the lax limit of the category \mathcal{C} in [Tho83]. However, we shift perspective: our philosophy is to work with the equivariant object $\mathcal{C}at(\tilde{G}, \mathcal{C})$, as opposed to just restricting attention to its fixed points. We don't merely study the H -fixed points of $\mathcal{C}at(\tilde{G}, \mathcal{C})$, which are the H -homotopy fixed points of \mathcal{C} , but we also study how homotopy fixed point categories relate, and for this it is convenient to study G -maps between the G -categories $\mathcal{C}at(\tilde{G}, \mathcal{C}) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{D})$.

In §4 we study the homotopy fixed point categories of module categories of G -rings and then we exploit these results in §5, §6, and §7 to draw the conclusions about the equivariant algebraic K -theory of G -rings described above. As an accidental corollary of our results about homotopy fixed points of module categories, we obtain an alternative proof of Serre's generalization of Hilbert's theorem 90, which we give in §6.2. The proof in the same spirit of Deligne's alternative proof of the original statement of Hilbert 90 from [Del77] using faithfully flat descent.

One property that one would expect of the homotopy fixed points of a category (which also justifies the name) is that they are homotopy invariant. We show that this is so.

Proposition 1.3. *A G -map, which is a nonequivariant equivalence of categories, induces equivalences of categories on homotopy fixed points.*

Another property of the $\mathcal{C}at(\tilde{G}, -)$ construction, which is more surprising maybe, and which is at the heart of our results is that it turns maps for which equivariance holds up to isomorphism into on the nose equivariant maps. In §3, we define the notion of a pseudo equivariant functor between G -categories as a functor which commutes with the G -action only up to coherent isomorphism. Very precisely, if one regards G -categories as functors $G \rightarrow \mathcal{C}at$, then an equivariant map of G -categories is a natural transformation between these and a pseudo equivariant map is a pseudo natural transformation. The main result of that section is as follows.

Proposition 1.4. *Given a pseudo equivariant functor of G -categories $\mathcal{C} \rightarrow \mathcal{D}$, there is an*

induced on the nose equivariant functor $\mathcal{C}at(\tilde{G}, \mathcal{C}) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{D})$, so there are induced maps on homotopy fixed point categories $\mathcal{C}^{hH} \rightarrow \mathcal{D}^{hH}$.

We showcase some of the applications of the result about pseudo equivariant functors. For example, the extension of scalars map between the module categories of G -rings (with actions defined a few paragraphs above) along a G -map of rings, is not equivariant, but only pseudo equivariant. Because this allows us to construct an on the nose equivariant functor after applying $\mathcal{C}at(\tilde{G}, -)$ to our module categories, we can ensure that we actually get a functor from the category of G -rings to the category of G -spectra. The definition of equivariant Morita equivalence given in §4.6, which equivariant algebraic K -theory is invariant under, is also in terms of a pseudo equivariant functor. We have claimed above that applying $\mathcal{C}at(\tilde{G}, -)$ rectifies an action that preserves the symmetric monoidal structure of a category \mathcal{C} to an action that preserves it strictly. This is also an application of the same result: the functor $\mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ that gives the symmetric monoidal structure is pseudo equivariant.

In §8 we give a definition of equivariant A -theory of a G -space X as the S_\bullet -construction of $B\mathcal{C}at(\tilde{G}, R(X))$, where the category of retractive spaces over X , $R(X)$, has the G -action we have defined above. We show that the homotopy fixed point category $R(X)^{hH}$ is precisely the category of retractive H -spaces with equivariant inclusion and retraction maps, and the H -fixed points of $A_G(X)$ are therefore the K -theory of this category. We point out that the weak equivalences in the fixed point subcategories are coarse: they are nonequivariant equivalences that are G -equivariant maps. An observation that we owe to discussions with C. Malkiewich and D. Schäppi is that the K -theory space we define for Waldhausen G -categories can be delooped by an equivariant infinite loop space machine. Nonequivariantly, it is an observation of Waldhausen that it is enough to apply the S_\bullet -construction once and then use an infinite loop space machine with the group completion property. We refer the reader to [Mal] for an explicit proof of Waldhausen’s observation. We conjecture that the deloopings of the equivariant A -theory space with respect to representations are given by iterations of the S_\bullet -construction on representations - this would ultimately follow from an equivariant version of the additivity theorem and work of Dotto and Moi on homotopy limits of G -diagrams [DM]. In forthcoming work with Emanuele Dotto we aim to prove an equivariant additivity theorem and produce the deloopings with respect to representations using an equivariant version of the S_\bullet -construction.

In order to study Waldhausen G -categories in §8, the result about pseudo equivariant functors becomes absolutely essential. The fixed point categories of a Waldhausen category \mathcal{C}^H are not Waldhausen, because the action does not preserve the zero object or the pushouts strictly. Using the result that a pseudo equivariant functor can be strictified to an equivariant functor after applying $\mathcal{C}at(\tilde{G}, -)$, we show that the homotopy fixed point categories \mathcal{C}^{hH} are Waldhausen categories. Moreover, using the same result we can see how one gets transfer “wrong way” maps between the homotopy fixed point categories. We explain this in §9 – this is the main ingredient in relating our definition of equivariant algebraic K -theory to the alternative definition of C. Barwick from [Bar] using “spectral Mackey functors.” His perspective seems very fruitful, especially in the setting of ∞ -categories, and we hope we will be able to give a more precise comparison in the future.

We conclude the introduction with two technical remarks. Note that everywhere we need to take classifying spaces of categories that are clearly not small. Nonequivariantly, it is always

assumed in K -theory that when we take the classifying space of a category which is not small, such as $\mathcal{P}(R)$, $\mathcal{F}(R)$, or $\text{Mod}(R)$, we are tacitly replacing the category by a small category, which is equivalent to it, such as its skeleton. The situation is a little trickier equivariantly, because we do not have an equivariant equivalence between a G -category and its skeleton. This is too much to hope for; however, we show in §4.4 that there is a weak G -equivalence $\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{F}(R)) \simeq \mathcal{C}at(\tilde{G}, \mathcal{GL}(R))$. In §4.5 we generalize this to any G -category – this is even more subtle because unlike in the case of free modules where we show that $R^n \cong gR^n$, in general, an object C is not necessarily isomorphic to gC . We show that for a G -category \mathcal{C} , we can put a G -action on the skeletal category $\text{sk } \mathcal{C}$, such that we get a weak G -equivalence $\mathcal{C}at(\tilde{G}, \mathcal{C}) \rightarrow \mathcal{C}at(\tilde{G}, \text{sk } \mathcal{C})$. This suffices for our applications, because in equivariant algebraic K -theory we are only taking classifying spaces of G -categories of the form $\mathcal{C}at(\tilde{G}, \mathcal{C})$.

We end with a remark about the group G . All of our categorical work on homotopy fixed points of G -categories works for any topological group G . However, in order to pass the statements to the spectrum level, we have to restrict to finite groups because of the limitations of the equivariant infinite loop space machines. We do hope that these limitations can be overcome in the near future, at least for profinite groups.

Acknowledgements

We would like to thank Peter May for suggesting equivariant algebraic K -theory as a thesis topic; Bertrand Guillou, Peter May, and Angélica Osorno for all the joint work that has fed into this project; Angélica Osorno for adding categorical sophistication to some of this work, and in particular for suggesting using the name “pseudo equivariant functor” as opposed to the name “weakly equivariant functor” which the author was initially using; Andrew Blumberg, Lars Hesselholt and Mike Hill, for sharing their invaluable insights on innumerable occasions; Dan Grayson, for sharing his intuition and explaining alternative approaches to some of our results, and Clark Barwick for sharing his perspective on equivariant algebraic K -theory and for the many discussions about how our approaches compare. We would also like to thank Emanuele Dotto and Daniel Schäppi for the illuminating discussions about Waldhausen G -categories which led to the acknowledgement of a mistake in the author’s thesis and the way of fix it. Lastly, we thank Cary Malkiewich and Lennart Meier for many great conversations on the topics of this paper and related work in progress.

2 Categorical homotopy fixed points

By analogy with the homotopy fixed points for a G -space we define the homotopy fixed points of a G -category. These were also studied by Thomason under the name “*lax limit*” in [Tho83]. However, we take an equivariant point of view: for us, the homotopy fixed points are the actual fixed points of a G -category, and we study this equivariant object as opposed to just restricting attention to the fixed points.

2.1 Preliminaries on G -categories

Concisely, a G -category can be defined as a functor $G \rightarrow \mathcal{Cat}$. Explicitly, the data of such a functor is a category \mathcal{C} , and for each $g \in G$, an endofunctor $(g\cdot): \mathcal{C} \rightarrow \mathcal{C}$ such that $(e\cdot) = \text{id}_{\mathcal{C}}$ and $(g\cdot) \circ (h\cdot) = (gh\cdot)$. By slight abuse, we will often call the category \mathcal{C} a G -category, which means we are implicitly thinking of the action endofunctors $(g\cdot)$. Sometimes we might omit the “.” from the notation and write gC or gf to denote the action of g on an object C or a morphism f . A natural transformation of functors $G \rightarrow \mathcal{Cat}$ translates to a functor between the two G -categories which commutes with the G -action. We denote the category of G -categories and G -equivariant functors by $G\mathcal{Cat}$.

For subgroups $H \subseteq G$, we define the H -fixed point category \mathcal{C}^H of a G -category \mathcal{C} as the subcategory with objects those $C \in \mathcal{C}$ such that $hC = C$ and morphisms those $f \in \mathcal{C}$ such that $hf = f$ for all $h \in H$. This definition coincides with the categorical definition as $\lim_H \mathcal{C}$ when we think of \mathcal{C} as a functor $G \rightarrow \mathcal{Cat}$. A crucial fact is that the classifying space functor $B: \mathcal{Cat} \rightarrow \text{Top}$ commutes with fixed points, namely

$$B(\mathcal{C}^H) = (B\mathcal{C})^H. \quad (1)$$

Definition 2.1. A functor between G -categories $F: \mathcal{C} \rightarrow \mathcal{D}$ is a *weak G -equivalence* if it induces a weak G -equivalence on classifying spaces $BF: B\mathcal{C} \rightarrow B\mathcal{D}$.

2.2 $G\mathcal{Cat}$ as a 2-category

We may view \mathcal{Cat} as the 2-category of categories, with 0-cells, 1-cells, and 2-cells the categories, functors, and natural transformations. From that point of view, \mathcal{Cat} is enriched over itself: the internal hom, $\mathcal{Cat}(\mathcal{A}, \mathcal{B})$, is the category whose objects are the functors $\mathcal{A} \rightarrow \mathcal{B}$ and whose morphisms are the natural transformations between them.

Similarly, we may view $G\mathcal{Cat}$ as the underlying 2-category of a category enriched over $G\mathcal{Cat}$. The 0-cells are G -categories, and the internal hom between them is the G -category $\mathcal{Cat}(\mathcal{A}, \mathcal{B})$. Its underlying category is $\mathcal{Cat}(\mathcal{A}, \mathcal{B})$, and G acts by conjugation on functors and natural transformations. Thus, for $F: \mathcal{A} \rightarrow \mathcal{B}$, $g \in G$, and A either an object or a morphism of \mathcal{A} , $(gF)(A) = gF(g^{-1}A)$. Similarly, for a natural transformation $\eta: E \rightarrow F$ and an object A of \mathcal{A} ,

$$(g\eta)_A = g\eta_{g^{-1}A}: gE(g^{-1}A) \rightarrow gF(g^{-1}A).$$

The category $G\mathcal{Cat}(\mathcal{A}, \mathcal{B})$ of G -equivariant functors and G -equivariant natural transformations is the same as the G -fixed category $\mathcal{Cat}(\mathcal{A}, \mathcal{B})^G$.

Remark 2.2. We can topologize the definitions so far, starting with the 2-category of categories internal to the category Top , together with continuous functors and continuous natural transformations. A *topological G -category* \mathcal{A} is a category internal to the cartesian monoidal category $G\text{Top}$. It has object and morphism G -spaces and continuous G -equivariant source, target, identity and composition structure maps. These maps are denoted S , T , I , and C , and the usual category axioms must hold. These are more general than (small) topologically enriched categories, which have discrete sets of objects.

2.3 The functor $\mathcal{C}at(\tilde{G}, -)$ and categorical homotopy fixed points

Definition 2.3. For a topological group G , define \tilde{G} to be the topological G -groupoid with object space G and morphism space $G \times G$. The source and target maps are the projections onto the two factors.

Thus the objects of \tilde{G} are the elements of G and there is a unique morphism between any two objects. We choose to label the unique morphism $g \rightarrow h$ by the pair (h, g) in order to be consistent with [GMM]. The idea is that reversing the order of source and target makes the notation for composition more transparent: $(g, h) \circ (h, k) = (g, k)$. The G -action on \tilde{G} is given by translation on the objects, which forces it to be diagonal on morphisms, since $g(h \rightarrow k)$, namely $g(k, h)$ must be the unique map $gh \rightarrow gk$, namely (gk, gh) .

Definition 2.4. Define the translation category of \underline{G} of G in the standard way as having object space G and morphism space $G \times G$, with the morphism $h \rightarrow gh$ labeled by (g, h) .

Again, since there is a unique morphism between any two objects, the G -action on objects by translation completely determines the action on the morphism space: G acts on the second coordinate of $G \times G$. The following lemma follows immediately from the fact that $G \times G$ with G acting diagonally and $G \times G$ with G acting on the second coordinate are G -homeomorphic.

Lemma 2.5. *The translation category \underline{G} is G -isomorphic to the category \tilde{G} .*

Remark 2.6. The category \tilde{G} is an instance of the more general concept of *chaotic category* corresponding to a space. There is a *chaotic category functor* from spaces to categories (actually, to groupoids), sending a space X to the category \tilde{X} with object space X and morphism space $X \times X$; there is a unique morphism between any two objects in \tilde{X} . The relevant point is that the object functor is right adjoint to the chaotic category functor, and in particular, we have a homeomorphism between the mapping spaces

$$\mathcal{C}at(\mathcal{C}, \tilde{X}) \cong \text{Map}(\mathcal{Ob}\mathcal{C}, X). \quad (2)$$

Similarly, the translation category \underline{G} of G is an instance of the more general notion of translation category of a G -space. For a G -set, or a G -space X , we denote by \underline{X} the translation category of X with objects the points of X and morphisms $(g, x) : x \rightarrow gx$. However, as we have seen, the concepts of chaotic and translation category agree for G up to G -isomorphism. Thus, it is harmless to think of \tilde{G} as the translation category of G . For a more comprehensive treatment of both chaotic and translation categories, we refer the reader to [GMM].

We make the following crucial observation.

Observation 2.7. The classifying space $B\tilde{G}$ is G -equivalent to the universal principal G -bundle EG since \tilde{G} is a contractible category (every object is initial and terminal) and it has a free G -action.

We have a functor $\mathcal{C}at(\tilde{G}, -)$ from G -categories to G -categories, which sends a G -category \mathcal{C} to the category of functors and natural transformations $\mathcal{C}at(\tilde{G}, \mathcal{C})$, with G -action by conjugation, as described in section 2.2. This is a topological category when \mathcal{C} is such. In view of Section 2.2,

$\mathcal{C}at(\tilde{G}, -)$ can be viewed as a 2-functor. Observe that the functor $\mathcal{C}at(\tilde{G}, -)$ is corepresented and is thus a right adjoint. Therefore it preserves all limits; in particular it preserves products, which will be crucial to our applications.

The equivariant projection $\tilde{G} \rightarrow *$ to the trivial G -category induces a natural G -map

$$\iota: \mathcal{A} \simeq \mathcal{C}at(*, \mathcal{A}) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{A}), \quad (3)$$

which is always a nonequivariant equivalence of G -categories, but not usually a G -equivalence. However, as observed in [GMa], the functor $\mathcal{C}at(\tilde{G}, -)$ is idempotent:

Lemma 2.8. *For any G -category \mathcal{A} ,*

$$\iota: \mathcal{C}at(\tilde{G}, \mathcal{A}) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{C}at(\tilde{G}, \mathcal{A}))$$

is an equivalence of G -categories.

By analogy with the definition of homotopy fixed points of G -spaces, we make the following definition.

Definition 2.9. The *homotopy fixed points* of a G -category \mathcal{C} , denoted by \mathcal{C}^{hG} , are defined as $\mathcal{C}at(\tilde{G}, \mathcal{C})^G$, namely the G -equivariant functors $\tilde{G} \rightarrow \mathcal{C}$ and the G -natural transformations between these.

Observation 2.10. Note that \tilde{H} and \tilde{G} are equivalent as H -categories since they are both H -free contractible categories. Therefore we can identify

$$\mathcal{C}at(\tilde{G}, \mathcal{C})^H = H\mathcal{C}at(\tilde{G}, \mathcal{C}) \simeq H\mathcal{C}at(\tilde{H}, \mathcal{C}) = \mathcal{C}at(\tilde{H}, \mathcal{C})^H.$$

Consequently, for any $H \subseteq G$, we can unambiguously define the H -homotopy fixed points of a G -category \mathcal{C} as either $\mathcal{C}at(\tilde{G}, \mathcal{C})^H$ or by applying the definition above of homotopy fixed points to \mathcal{C} , regarded as an H -category. And conveniently, for any statement that we wish to prove holds for H -fixed points $\mathcal{C}at(\tilde{G}, \mathcal{C})^H$ for any $H \subseteq G$, it is enough to prove it for G -fixed points $\mathcal{C}at(\tilde{G}, \mathcal{C})^G$ as long as G is arbitrary.

2.4 Explicit description of homopy fixed point categories

We describe explicitly the category of equivariant functors and equivariant natural transformations $G\mathcal{C}at(\tilde{G}, \mathcal{C})$. Any G -fixed functor $F: \tilde{G} \rightarrow \mathcal{C}$ is determined on objects by where the identity e of G gets mapped to since $F(g) = g \cdot F(e)$. On morphisms, F is determined by where it sends morphisms of the type (g, e) since $F(g, h) = h \cdot F(h^{-1}g, e)$. We have that $F(e, e) = id_C$, where id_C is the identity morphism of the object $C \in \mathcal{C}$ and $F(e) = C$. The following cocycle condition is also satisfied:

$$F(gh, e) = F(gh, g)F(g, e) = g \cdot F(h, e)F(g, e).$$

We summarize this discussion in the following result, which gives an explicit description of

the homotopy fixed point category of a G -category \mathcal{C} .¹

Proposition 2.11. *The objects of the homotopy fixed point category $\mathcal{C}^{hG} = G\mathcal{C}at(\tilde{G}, \mathcal{C})$ are pairs (C, f) where C is an object of \mathcal{C} and $f : G \rightarrow \text{Mor}(\mathcal{C})$ is a map from G to morphisms of \mathcal{C} such that $f(g) : C \rightarrow g \cdot C$ and f satisfies the condition $f(e) = id_C$ and the cocycle condition*

$$f(gh) = (g \cdot f(h))f(g). \quad (4)$$

A morphism $(C, f) \rightarrow (C', f')$ is given by a morphism $\alpha : C \rightarrow C'$ in \mathcal{C} such that the following diagram commutes for any $g \in G$:

$$\begin{array}{ccc} C & \xrightarrow{f(g)} & g \cdot C \\ \alpha \downarrow & & \downarrow g \cdot \alpha \\ C' & \xrightarrow{f'(g)} & g \cdot C' \end{array}$$

However, the alternative cocycle condition

$$f(gh) = f(g)(g \cdot f(h)) \quad (5)$$

is the standard one, which will appear in all of our applications. For instance, this is the condition that yields a crossed homomorphism when \mathcal{C} is a group, whereas condition (4) yields a crossed antihomomorphism, which is less customary.

We show that changing condition (4) to the usual cocycle condition (5) is inoffensive since it yields an equivalent category. The proof is a straightforward generalization of the proof of Lemma 4.13 in [GMM], which is the special case where the category \mathcal{C} is a group.

Proposition 2.12. *There is an isomorphism of categories between the homotopy fixed point category $\mathcal{C}^{hG} = G\mathcal{C}at(\tilde{G}, \mathcal{C})$ and the category described as follows. The objects are pairs (C, f) where C is an object of \mathcal{C} and $f : G \rightarrow \text{Mor}(\mathcal{C})$ is a map from G to morphisms of \mathcal{C} such that $f(g) : g \cdot C \rightarrow C$ and f satisfies the condition $f(e) = id_C$ and the cocycle condition*

$$f(gh) = f(g)(g \cdot f(h)).$$

A morphism $(C, f) \rightarrow (C', f')$ is given by a morphism $\alpha : C \rightarrow C'$ in \mathcal{C} such that the following diagram commutes for any $g \in G$:

$$\begin{array}{ccc} g \cdot C & \xrightarrow{f(g)} & C \\ g \cdot \alpha \downarrow & & \downarrow \alpha \\ g \cdot C' & \xrightarrow{f'(g)} & C' \end{array}$$

Proof. We explicitly construct the isomorphism between the category described in Proposition 2.12 to the category described in Proposition 2.11. The construction of the inverse isomor-

¹This explicit description is also given in more concise terms in [Tho85].

phism is similar.

Let $f: G \rightarrow \text{Mor}(\mathcal{C})$ be such that $f(g): g \cdot C \rightarrow C$ and suppose that f satisfies $f(e) = id_C$ and condition (5). Note that $f(g)$ is an isomorphism with inverse $g \cdot f(g^{-1})$ for all $g \in G$.

Define $\bar{f}: G \rightarrow \text{Mor}(\mathcal{C})$ by

$$\bar{f}(g) = g \cdot f(g^{-1}),$$

so that $\bar{f}: C \rightarrow g \cdot C$. Then

$$\bar{f}(gh) = (gh) \cdot f(h^{-1}g^{-1}) = g \cdot h \cdot (f(h^{-1})(h^{-1} \cdot f(g^{-1}))) = (g \cdot \bar{f}(h))(\bar{f}(g)),$$

so that \bar{f} is satisfies condition (4).

Let $f, f': G \rightarrow \text{Mor}(\mathcal{C})$, such that $f(g): g \cdot C \rightarrow C$ and $f'(g): g \cdot C' \rightarrow C'$. Suppose that f and f' satisfy (5), and $\alpha: C \rightarrow C'$ is a morphism in \mathcal{C} for which the diagram

$$\begin{array}{ccc} g \cdot C & \xrightarrow{f(g)} & C \\ g \cdot \alpha \downarrow & & \downarrow \alpha \\ g \cdot C' & \xrightarrow{f'(g)} & C' \end{array}$$

commutes for all $g \in G$. Then

$$\bar{f}'(g) \alpha = (g \cdot f'(g^{-1})) \alpha = g \cdot (f'(g^{-1})(g^{-1} \cdot \alpha)) = g \cdot (\alpha f(g^{-1})) = (g \cdot \alpha) \bar{f}(g),$$

i.e. the diagram

$$\begin{array}{ccc} C & \xrightarrow{\bar{f}(g)} & g \cdot C \\ \alpha \downarrow & & \downarrow g \cdot \alpha \\ C' & \xrightarrow{\bar{f}'(g)} & g \cdot C' \end{array}$$

commutes for all $g \in G$. □

Remark 2.13. Note that if G acts trivially on \mathcal{C} , then $\mathcal{C}at(\tilde{G}, \mathcal{C})^G \cong \mathcal{C}at(G, \mathcal{C})$, the category of functors $G \rightarrow \mathcal{C}$, i.e., \mathcal{C}^{hG} is the category of representations of G in \mathcal{C} .

We emphasize that the homotopy fixed points do not in general commute with the classifying space functor. However, if \mathcal{C} is a *discrete* G -groupoid, then the comparison map $B\mathcal{C}at(\tilde{G}, \mathcal{C}) \rightarrow \text{Map}(EG, B\mathcal{C})$ is a weak G -equivalence, and we have

$$B(\mathcal{C}^{hH}) \simeq (B\mathcal{C})^{hH}$$

for any $H \subseteq G$ (see [GMM, §5] for a proof). However, in some of our examples of interest, even when the category \mathcal{C} is a groupoid, we need it to be topological, and therefore cannot assume this commutation.

2.5 Homotopy fixed points of a group

The homotopy fixed point category \mathcal{C}^{hG} simplifies when $\mathcal{C} = \Pi$, a topological group regarded as a topological category with one object, with G -action. In that case, the homotopy fixed points can be interpreted in terms of the well-known notion of *crossed group homomorphisms*. The category $G\mathcal{C}at(\tilde{G}, \Pi)$ has been studied extensively in [GMM], where it was shown that it has the following interpretation.

Theorem 2.14 ([GMM], 4.15). *Suppose Π is a group with G -action. The homotopy fixed point category Π^{hG} is equivalent to the crossed functor category $\mathcal{C}at_{\times}(G, \Pi)$ whose objects are the continuous crossed homomorphisms $G \rightarrow \Pi$ and whose morphisms $\sigma: \alpha \rightarrow \beta$ are the elements $\sigma \in \Pi$ such that*

$$\beta(g)(g \cdot \sigma) = \sigma \alpha(g).$$

This interpretation leads to the following condition for when the nonequivariant equivalence

$$\iota: \Pi \simeq \mathcal{C}at(*, \Pi) \rightarrow \mathcal{C}at(\tilde{G}, \Pi)$$

is a weak G -equivalence.

Proposition 2.15 ([GMM], 4.19). *The functor $\iota^H: \Pi^H \rightarrow \mathcal{C}at(\tilde{G}, \Pi)^H$ is an equivalence of categories if and only if the first nonabelian cohomology set $H^1(H; \Pi)$ is trivial.*

2.6 Homotopy invariance of homotopy fixed points

The following lemma is inspired by the analogous result for homotopy fixed points of G -spaces or naive G -spectra.

Proposition 2.16. *If $\Theta: \mathcal{C} \rightarrow \mathcal{D}$ is a G -functor that is a nonequivariant equivalence of categories then the functor induced by post composition*

$$\mathcal{C}at(\tilde{G}, \mathcal{C}) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{D})$$

is a weak G -equivalence of categories.

Proof. From Observation 2.7, we see that it is enough to prove that we get an equivalence on G -fixed points. The map

$$\mathcal{C}at(\tilde{G}, \mathcal{C})^G \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{D})^G$$

is faithful since it is the restriction of a faithful map to a subcategory. We show that is essentially surjective and full. We use the explicit description of fixed points given above.

Pick an object (D, f) in $\mathcal{C}at(\tilde{G}, \mathcal{D})^G$. Since Θ is essentially surjective, there exists a nonequivariant isomorphism $\psi: D \xrightarrow{\cong} \Theta(C)$ for some $C \in \mathcal{C}$. By applying $g \cdot$ we get $g\psi: gD \xrightarrow{\cong} g\Theta(C) = \Theta(gC)$. Since Θ is fully faithful, for every $f(g): D \xrightarrow{\cong} gD$ there exists a unique

map $f'(g): C \xrightarrow{\cong} gC$ such that $\Theta(f'(g))$ is the composite

$$\Theta(C) \xrightarrow{\psi^{-1}} D \xrightarrow{f(g)} gD \xrightarrow{g\psi} g\Theta(C),$$

and $f'(g)$ is an isomorphism since $f(g)$ and ψ are. We need to check the cocycle condition on f' .

We will read it off from the following commutative diagram

$$\begin{array}{ccccc} D & \xrightarrow{f(g)} & gD & \xrightarrow{gf(h)} & ghD \\ \psi \downarrow & & \downarrow g\psi & & \downarrow gh\psi \\ \Theta(C) & \xrightarrow{\Theta(f'(g))} & g\Theta(C) & \xrightarrow{g\Theta(f'(h))} & gh\Theta(C) \end{array}$$

The top composite is $f(gh)$ since f satisfies the cocycle condition. Thus the bottom map must be $\Theta(f(gh))$. By the commutation of g with Θ , the bottom map is just Θ applied to the composite

$$C \xrightarrow{f'(g)} gC \xrightarrow{gf'(h)} ghC.$$

Thus f' satisfies the cocycle condition.

We are left to show fullness. Suppose we have a morphism in $\mathcal{C}at(\tilde{G}, \mathcal{D})$ from $(\Theta(C), \Theta(f))$ to $(\Theta(C'), \Theta(f'))$ given by the diagrams

$$\begin{array}{ccc} \Theta(C) & \xrightarrow{\Theta(f(g))} & g\Theta(C) \\ \alpha \downarrow & & \downarrow g\alpha \\ \Theta(C') & \xrightarrow{\Theta(f'(g))} & g\Theta(C') \end{array}$$

Since Θ is full, there exists a map $C \xrightarrow{\alpha'} C'$ such that $\Theta(\alpha') = \alpha$. Thus there is a map in $\mathcal{C}at(\tilde{G}, \mathcal{C})$

$$\begin{array}{ccc} C & \xrightarrow{f(g)} & gC \\ \alpha' \downarrow & & \downarrow g\alpha' \\ C' & \xrightarrow{f'(g)} & gC' \end{array}$$

whose image is the map above. This gives fullness. \square

The proposition shows that the homotopy fixed point construction for a G -category \mathcal{C} is homotopy invariant.

3 Pseudo equivariance

3.1 Pseudo equivariant functors

Let \mathcal{C} and \mathcal{D} be G -categories. We define the notion of a *pseudo equivariant* functor $\Theta: \mathcal{C} \rightarrow \mathcal{D}$, and we then show that such a functor induces an on the nose equivariant functor

$$\mathcal{C}at(\tilde{G}, \mathcal{C}) \longrightarrow \mathcal{C}at(\tilde{G}, \mathcal{D}).$$

Thus it induces maps on fixed points

$$\mathcal{C}at(\tilde{G}, \mathcal{C})^H \longrightarrow \mathcal{C}at(\tilde{G}, \mathcal{D})^H$$

for all subgroups $H \subseteq G$. Pseudo equivariance will be absolutely central to a lot of our K -theory results because often enough the maps between the G -categories we will consider in equivariant algebraic K -theory are not on the nose equivariant, but pseudo equivariant. The construction of equivariant algebraic K -theory will ensure that this is enough to get actual equivariant maps on the spectrum level. Moreover, this result will be what allows us to rectify a G -action on a symmetric monoidal or Waldhausen category which does not preserve the structure.

Definition 3.1. A *pseudo equivariant* functor between G -categories \mathcal{C} and \mathcal{D} is a functor $\Theta: \mathcal{C} \rightarrow \mathcal{D}$, together with natural isomorphisms of functors θ_g for all $g \in G$

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{g \cdot} & \mathcal{C} \\ \Theta \downarrow & \Downarrow \theta_g & \downarrow \Theta \\ \mathcal{D} & \xrightarrow{g \cdot} & \mathcal{D} \end{array} .$$

such that $\theta_e = \text{id}$ and for $g, h \in G$ we have an equality of natural transformations, where on the left hand side we are considering the composite of natural transformations.

$$\begin{array}{ccccc} \mathcal{C} & \xrightarrow{h \cdot} & \mathcal{C} & \xrightarrow{g \cdot} & \mathcal{C} \\ \Theta \downarrow & \Downarrow \theta_h & \downarrow \Theta & \Downarrow \theta_g & \downarrow \Theta \\ \mathcal{D} & \xrightarrow{h \cdot} & \mathcal{D} & \xrightarrow{g \cdot} & \mathcal{D} \end{array} = \begin{array}{ccc} \mathcal{C} & \xrightarrow{gh \cdot} & \mathcal{C} \\ \Theta \downarrow & \Downarrow \theta_{gh} & \downarrow \Theta \\ \mathcal{D} & \xrightarrow{gh \cdot} & \mathcal{D} \end{array}$$

Note that requiring this equality makes sense because the outer right down and down right composites in the two diagrams are equal. Explicitly, for C an object of \mathcal{C} , this means that the following diagram commutes:

$$\begin{array}{ccccc} & & \theta_{gh}(C) & & \\ & & \curvearrowright & & \\ \Theta(ghC) & \xrightarrow{\theta_g(hC)} & g\Theta(hC) & \xrightarrow{g\theta_h(C)} & gh\Theta(C) \end{array}$$

Remark 3.2. If θ_g are equalities for all $g \in G$, then Θ is actually an equivariant functor.

We explain the choice of nomenclature. Recall that a G -category is a functor $G \rightarrow \mathcal{Cat}$, and an equivariant map between G -categories is then just a natural transformations of such functors. A *pseudo equivariant* map between G -categories is a *pseudo natural transformation*. We prove next that a pseudo equivariant functor $\Theta: \mathcal{C} \rightarrow \mathcal{D}$ naturally induces an on the nose equivariant map after applying the $\mathcal{Cat}(\tilde{G}, -)$ functor.

Proposition 3.3. *A pseudo equivariant functor $\Theta: \mathcal{C} \rightarrow \mathcal{D}$ naturally induces an equivariant functor*

$$\tilde{\Theta}: \mathcal{Cat}(\tilde{G}, \mathcal{C}) \longrightarrow \mathcal{Cat}(\tilde{G}, \mathcal{D}).$$

Proof. Clearly post composing a functor $F: \tilde{G} \rightarrow \mathcal{C}$ with Θ does not yield an equivariant functor, but we can use the natural isomorphisms θ_g to create one. We define

$$\tilde{\Theta}(F)(g) = g\Theta((g^{-1}F)(e)) = g\Theta(g^{-1}F(g))$$

Recall that there is a unique map in \tilde{G} from g to g' , which we denote by (g', g) . Applying $\Theta \circ F$ we get a map $\Theta(F(g)) \xrightarrow{\Theta(F(g',g))} \Theta(F(g'))$ in \mathcal{D} . We define $\tilde{\Theta}(g', g)$ to be the composite

$$g\Theta(g^{-1}F(g)) \xrightarrow{\theta_g^{-1}} \Theta(gg^{-1}F(g)) \xrightarrow{\Theta(F(g',g))} \Theta(g'g'^{-1}F(g')) \xrightarrow{\theta_{g'}} g'\Theta(g'^{-1}F(g')).$$

It is very tedious, but not very hard to check that with these definitions $\tilde{\Theta}$ is an equivariant functor. We omit these details here, but we show below explicitly what the induced map on fixed points is. \square

Corollary 3.4. *A pseudo equivariant functor $\Theta: \mathcal{C} \rightarrow \mathcal{D}$, induces functors $\tilde{\Theta}^H: \mathcal{C}^{hH} \rightarrow \mathcal{D}^{hH}$ on homotopy fixed points for all $H \subseteq G$.*

Note that the definition of $\tilde{\Theta}$ on objects makes sense for any functor Θ and $\tilde{\Theta}$ is equivariant on objects. However, without the isomorphisms encoded in the pseudo equivariance condition for Θ , it is not possible to define the map $\tilde{\Theta}$ on morphisms, and thus the 2-categorical point of view is forced upon us. We write down explicitly the fixed point map $\tilde{\Theta}^H: \mathcal{Cat}(\tilde{G}, \mathcal{C})^H \rightarrow \mathcal{Cat}(\tilde{G}, \mathcal{D})^H$ induced from a pseudo equivariant functor $\Theta: \mathcal{C} \rightarrow \mathcal{D}$, because it sheds light on how the 2-cells come in, and because instances of this induced fixed point map are relevant in equivariant algebraic K -theory. We will encounter interesting maps in K -theory which turn out to be fixed point maps of equivariant K -theory maps that arise from pseudo equivariant functors on the categorical level.

Recall the explicit description of homotopy fixed points given in Proposition 2.12. Let (C, f) be an object in $\mathcal{Cat}(\tilde{G}, \mathcal{C})^H$. Under the induced map on H -fixed points

$$\tilde{\Theta}^H: \mathcal{Cat}(\tilde{G}, \mathcal{C})^H \longrightarrow \mathcal{Cat}(\tilde{G}, \mathcal{D})^H,$$

this gets sent to $(\Theta(C), f_\theta)$ where $f_\theta(g)$ is defined as the composite

$$\Theta(C) \xrightarrow[\cong]{\Theta(f(g))} \Theta(gC) \xrightarrow[\cong]{\theta_g} g\Theta(C).$$

Since $f(e) = \text{id}$ and $\theta_e = \text{id}$, it follows immediately that $f_\theta(e) = \text{id}$. To show that f_θ satisfies the cocycle condition, we use the fact that f satisfies it, together with the diagram in Definition 3.1. By that diagram, the maps in the following composite themselves factor as composites:

$$\begin{array}{ccccc}
 \Theta(C) & \xrightarrow{\Theta(f(gh))} & \Theta(ghC) & \xrightarrow{\theta_{gh}(C)} & gh\Theta(C) \\
 \searrow \Theta(f(g)) & & \nearrow \Theta(gf(h)) & & \nearrow g\theta_h(C) \\
 & & \Theta(gC) & \xrightarrow{\theta_g(hC)} & g\Theta(hC)
 \end{array}$$

We can use the naturality diagram for θ_g

$$\begin{array}{ccc}
 \Theta(gC) & \xrightarrow{\Theta(gf(h))} & \Theta(ghC) \\
 \theta_g(C) \downarrow & & \downarrow \theta_g(hC) \\
 g\Theta(C) & \xrightarrow{g\Theta(f(h))} & g\Theta(hC)
 \end{array}$$

to replace the middle maps in the diagram above and we get that

$$\Theta(C) \xrightarrow{\Theta(f(gh))} \Theta(ghC) \xrightarrow{\theta_{gh}(C)} gh\Theta(C)$$

is the same as

$$\Theta(C) \xrightarrow{\Theta(f(g))} \Theta(gC) \xrightarrow{\theta_g(C)} g\Theta(C) \xrightarrow{g\Theta(f(h))} g\Theta(hC) \xrightarrow{g\theta_h(C)} gh\Theta(C).$$

Thus $f_\theta(gh) = (g \cdot f_\theta(h))f_\theta(g)$.

Question 3.5. Does every equivariant functor $\mathcal{C}at(\tilde{G}, \mathcal{C}) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{D})$ come from a pseudo equivariant functor $\mathcal{C} \rightarrow \mathcal{D}$?

3.2 Homotopy invariance of homotopy fixed points revisited

We can use pseudo equivariance to weaken the hypothesis of Proposition 2.16 from requiring the functor to be on the nose equivariant to requiring it to be pseudo equivariant. Surprisingly, we get the same conclusion.

Proposition 3.6. *Let $\Theta: \mathcal{C} \rightarrow \mathcal{D}$ be a pseudo equivariant functor which is a nonequivariant equivalence. Then the induced functor*

$$\mathcal{C}at(\tilde{G}, \mathcal{C}) \longrightarrow \mathcal{C}at(\tilde{G}, \mathcal{D})$$

is a weak G -weak equivalence.

Proof. The equivariant map

$$\tilde{\Theta}: \mathcal{C}at(\tilde{G}, \mathcal{C}) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{D}),$$

given in Proposition 3.3 is a nonequivariant equivalence with inverse $\widetilde{\Theta}^{-1}$.

We have a commutative diagram:

$$\begin{array}{ccc} \mathcal{C}at(\tilde{G}, \mathcal{C}) & \xrightarrow{\tilde{\Theta}} & \mathcal{C}at(\tilde{G}, \mathcal{D}) \\ \downarrow \iota & & \downarrow \iota \\ \mathcal{C}at(\tilde{G}, \mathcal{C}at(\tilde{G}, \mathcal{C})) & \xrightarrow{\mathcal{C}at(\tilde{G}, -)(\tilde{\Theta})} & \mathcal{C}at(\tilde{G}, \mathcal{C}at(\tilde{G}, \mathcal{D})) \end{array}$$

By Proposition 2.16, the bottom map is a weak G -equivalence, and by Lemma 2.8 the vertical maps are G -equivalences. Therefore the top map is a weak G -equivalence. □

Corollary 3.7. *A pseudo equivariant functor $\Theta: \mathcal{C} \rightarrow \mathcal{D}$ which is a nonequivariant equivalence induces equivalences of homotopy fixed points*

$$\mathcal{C}^{hH} \rightarrow \mathcal{D}^{hH}$$

for all $H \subseteq G$.

4 Homotopy fixed points of module categories

4.1 G -rings and twisted group rings

A G -ring is a ring R with a left action of G by ring automorphisms. If R is a topological ring, we ask for the action to be through continuous ring automorphisms. We have a homomorphism $G \rightarrow \text{Aut}(R)$, and we write $g(r) = r^g$ for the automorphism $g: R \rightarrow R$ determined by $g \in G$. Then $r^{gh} = g(h(r)) = (r^h)^g$.

Note that when R is a subquotient of \mathbb{Q} , the only automorphism of R is the identity and the action of G must be trivial. However, we will see that even trivial G -actions on rings yield nontrivial equivariant algebraic K -theory. For example, we will show that the equivariant algebraic K -theory of the topological rings \mathbb{R} and \mathbb{C} is equivariant topological real and complex K -theory. Nevertheless, we are interested in many examples where the group action on the ring is nontrivial such as the Galois extensions of rings, or the topological ring \mathbb{C} with $\mathbb{Z}/2\mathbb{Z}$ conjugation action.

Suppose that R is a commutative G -ring with action given by $\theta: G \rightarrow \text{Aut}(R)$. Observe that R is an R^G -algebra, where R^G is the subring of G -invariants. We can reinterpret θ as a group homomorphism $\theta: G \rightarrow \text{End}_{R^G} R$, and ask the question of when we can extend this to a ring

map. More precisely, we seek to put a ring structure on the underlying abelian group of the group ring $R[G]$, for which the map θ extends to a ring map.

This naturally leads to the definition of *twisted group ring* (or *skew group ring*), which we will denote by $R_G[G]$ (it is variously denoted in the literature also as $R \rtimes G$ or $R * G$). A more precise notation that takes into the action of G on R given by the homomorphism $\theta: G \rightarrow \text{Aut}(R)$ would be $R_\theta[G]$. However, the action of G on R will many times be implicit, so we will not adopt this more pendantic notation.

Definition 4.1. As an R -module, the *twisted, or skew, group ring* $R_G[G]$ is the same as the group ring $R[G]$, which is the case when G acts trivially on R . We define the product on $R_G[G]$ by R^G -linear (not R -linear) extension of the relation

$$(rg)(sh) = rs^g gh$$

for $r, s \in R$ and $g, h \in G$.

Thus moving g past s , “twists” the ring element by the group action. Note that R and $R^G[G]$ are subrings of $R_G[G]$ and

$$gr = r^g g.$$

Observe that the definition of the twisted multiplication in $R_G[G]$ is precisely what enables us to extend the group homomorphism $\theta: G \rightarrow \text{End}_{R^G} R$ to a ring homomorphism

$$\theta: R_G[G] \rightarrow \text{End}_{R^G} R, \quad (rg) \mapsto (s \mapsto r s^g).$$

4.2 Modules over twisted group rings

Definition 4.2. We call (left) $R_G[G]$ -modules *G-ring modules* or *skew G-modules*.

Note that $(rs)^g = (r^g)(s^g)$ for all $r, s \in R$, thus R is an example of an $R_G[G]$ -module.

Observation 4.3. An $R_G[G]$ -module M is a left R -module with a *semilinear G-action*, i.e., $g(rm) = r^g(gm)$ for $m \in M$. If the action of G on R is trivial, then an $R[G]$ -module is a left R -module M with *linear G-action*, namely, $g(rm) = r(gm)$. From this point of view an $R_G[G]$ -linear map of $R_G[G]$ -modules $f: M \rightarrow N$ is a map of R -modules, which commutes with the G -action.

If G is finite and $|G|$ is invertible in R , we obtain the following characterization of projective modules over $R_G[G]$, which will be crucial in our applications to K -theory of G -rings.

Proposition 4.4. *If G is finite and $|G|^{-1} \in R$, then an $R_G[G]$ -module is projective if and only if it is projective as an R -module.*

Proof. An $R_G[G]$ -module M is projective if and only if the functor

$$\text{Hom}_{R_G[G]}(M, -) : \text{Mod}(R_G[G]) \rightarrow \text{Mod}(R_G[G])$$

is exact. This functor is always left exact, and it is also right exact precisely when M is projective. Let M and N be $R_G[G]$ -modules. As noted in Observation 4.3, M and N are R -modules with semilinear G -action. Then the $R_G[G]$ -module $\text{Hom}_{R_G[G]}(M, N)$ is the R -module $\text{Hom}_R(M, N)$ with semilinear G -action given by conjugation, i.e., for an R -linear map $f: M \rightarrow N$, $gf(m) = g(f(g^{-1}m))$. Again from Observation 4.3, we have that

$$\text{Hom}_{R_G[G]}(M, N) \cong \text{Hom}_R(M, N)^G.$$

The fixed point functor $(-)^G$ on $R_G[G]$ -modules is right exact when the order of G is invertible in R . Thus when $|G|^{-1} \in R$, the functor $\text{Hom}_{R_G[G]}(M, -)$ is exact precisely when the functor $\text{Hom}_R(M, -)$ is exact.

□

Of course, we do not have a similar statement for free modules. Clearly, a free $R_G[G]$ -module is free over R , but the converse is not true: Freeness over R definitely does not imply freeness over $R_G[G]$. For a set A , let $R[A]$ denote the free R -module on the basis A . The following proposition shows how we can put an $R_G[G]$ -module structure on $R[A]$ if A is a G -set; this is equivalent to specifying a semilinear G -action on $R[A]$.

Proposition 4.5. *Let A be a G -set and define*

$$g\left(\sum_a r_a a\right) = \sum_a r_a^g g a$$

for $g \in G$, $r_a \in R$, and $a \in A$. Then $R[A]$ is an $R_G[G]$ -module.

In [GMM, 6.8], following [Kaw86, 5.1] we gave a classification of $R_G[G]$ -module structures on free rank n R -modules in terms of the homotopy fixed point category of the group $GL_n(R)$, regarded as a single object groupoid. It inherits a G -action from the G -action on R .

Theorem 4.6. *Let R be a G -ring. Then the set of isomorphism classes of $R_G[G]$ -module structures on the R -module R^n is in canonical bijective correspondence with the isomorphism classes of objects in the homotopy fixed point category $GL_n(R)^{hG} = \mathcal{C}at(\tilde{G}, GL_n(R))^G$.*

4.3 The category $\mathcal{C}at(\tilde{G}, \text{Mod}(R))$

For a G -ring R , the category of finitely generated R -modules $\text{Mod}(R)$ becomes a G -category with action defined in the following way. Let M be an R -module with action map $\gamma: R \times M \rightarrow M$. Then we let $gM = M$ as abelian groups, and we define the action map by pulling back the action on M along $g: R \rightarrow R$:

$$\gamma_g: R \times M \xrightarrow{g \times \text{id}} R \times M \xrightarrow{\gamma} M.$$

This twists the R -action on M by the action of G on R . Explicitly, the R action on gM , which we will denote by \cdot_g to differentiate from the R -action on M , is given by

$$r \cdot_g m := r^g m,$$

where on the right hand side of the equation we are using the action of R on M .

We note that

$$R_G[G] \otimes_R M \cong \bigoplus_{g \in G} gM.$$

For a morphism $f: M \rightarrow N$, we define $gf: gM \rightarrow gN$ by $(gf)(m) = f(m)$. Thus gf is the same as f as a homomorphism of abelian groups, but it interacts differently from f with the scalar multiplication.

Note that in general M is not necessarily isomorphic to gM as R -modules. The identity of abelian groups $M = gM$ is not an R -linear map, since the R -action is different on the two sides of the equality. However, we do have an isomorphism of free R -modules $R^n \cong gR^n$, which plays an important role.

Lemma 4.7. *The R -modules R^n and gR^n are isomorphic.*

Proof. Let $\{e_i\}$ be the standard basis for R^n . Note that this is also a basis for gR^n : if $r_1^g e_1 + \dots + r_n^g e_n = 0$, then $r_i^g = 0$ for all i , so $r_i = 0$ since G acts by ring automorphisms. Also, every element in gR^n can be written as

$$(r_1, \dots, r_n) = (r_1^{(g^{-1})})^g e_1 + \dots + (r_n^{(g^{-1})})^g e_n.$$

Now just define a map on basis elements as the identity $e_i \mapsto e_i$ and extend linearly, i.e. $re_i \mapsto r^g e_i$. \square

We emphasize that the objects of the category $\text{Mod}(R)$ are R -modules M , which know nothing about the G -action on R . We used this action to define a G -action on the category $\text{Mod}(R)$, and now we will show how the G -category $\text{Mod}(R)$ relates to the category of modules over the twisted group ring $R_G[G]$, which by Observation 4.3 is the same as the category of R -modules with semilinear G -action.

Proposition 4.8. *The homotopy fixed point category $\text{Mod}(R)^{hG}$ is equivalent to the category $\text{Mod}(R_G[G])$.*

Proof. From the description of homotopy fixed point categories given in Proposition 2.11, the objects of the homotopy fixed point category $\mathcal{C}at(\tilde{G}, \text{Mod}(R))^G$ are R -modules M together with compatible isomorphisms $f(g): gM \xrightarrow{\cong} M$, one for each element $g \in G$, for which $f(e) = \text{id}_M$ and which make the diagrams

$$\begin{array}{ccc} (gh)M & \xrightarrow{gf(h)} & gM \\ & \searrow f(gh) & \swarrow f(g) \\ & M & \end{array}$$

commute.

Define an action of G on M by $g \cdot m = f(g)(m)$. This is indeed an action since $f(e) = \text{id}_M$ and

$$\begin{aligned} (gh) \cdot m &= f(gh)(m) \\ &= f(g)gf(h)(m) \\ &= f(g)f(h)(m) \\ &= g \cdot (h \cdot m). \end{aligned}$$

The second to last identification is just the definition of the G -action on morphisms of modules in $\text{Mod}(R)$; the morphism $gf(h)$ is the same as $f(h)$ as a morphism of abelian groups.

Now note that this action is indeed semilinear:

$$g \cdot (rm) = f(g)(r \cdot_g m) = r^g f(g)(m) = r^g (g \cdot m).$$

Via this identification, the morphisms in the homotopy fixed point category are precisely the G -equivariant maps of G -modules.

Thus we have shown that the homotopy fixed point category $\text{Mod}(R)^{hG}$ can be identified with the category of modules with semilinear G -action. Combining this with Observation 4.3, we obtain the desired result. \square

By Observation 2.10, we immediately get the following corollary.

Corollary 4.9. *The homotopy fixed point category $\text{Mod}(R)^{hH}$ is equivalent to the category $\text{Mod}(R_H[H])$ for all subgroups $H \subseteq G$.*

Therefore, the G -category $\mathcal{C}at(\tilde{G}, \text{Mod}(R))$ encodes the module categories over the twisted group rings for all subgroups H as fixed point subcategories. Thus by studying the equivariant object $\mathcal{C}at(\tilde{G}, \text{Mod}(R))$ we are implicitly studying the representation theory of all the subgroups at once.

Let $\mathcal{P}(R)$ be the category of finitely generated projective R -modules. This becomes a G -category in the same way that $\text{Mod}(R)$ does since gP is projective if P is so: if $P \oplus Q \cong R^n$, then $gP \oplus gQ \cong gR^n \cong R^n$. The proof of Proposition 4.8 goes through to show that the category $\mathcal{C}at(\tilde{G}, \mathcal{P}(R))$ is equivalent to the category of finitely generated projective R -modules with semilinear G -action. Therefore, by Proposition 4.4, if G is finite and the order of G is invertible in R , we obtain Proposition 4.8 and its corollary if we restrict to the category of finitely generated R -modules.

Proposition 4.10. *Suppose G is finite and $|G|^{-1} \in R$. The homotopy fixed point category $\mathcal{P}(R)^{hG}$ is equivalent to the category $\mathcal{P}(R_G[G])$.*

Corollary 4.11. *Suppose G is finite and $|G|^{-1} \in R$. The homotopy fixed point category $\mathcal{P}(R)^{hH}$ is equivalent to the category $\mathcal{P}(R_H[H])$ for all subgroups $H \subseteq G$.*

4.4 The equivariant skeleton of free modules

If $M \cong R^n$, then $gM \cong gR^n \cong R^n$, so the G -action on $\text{Mod}(R)$ restricts to an action on the category $\mathcal{F}(R)$ of finitely generated free R -modules.

Definition 4.12. Let $\mathcal{GL}(R)$ be the category with objects the based free R -modules R^n and morphism spaces

$$\text{Mor}_{\mathcal{GL}(R)}(R^n, R^m) = \begin{cases} \emptyset & \text{if } n \neq m \\ GL_n(R) & \text{if } n = m. \end{cases}$$

This is the same as the disjoint union of the one object categories $GL_n(R)$, i.e.,

$$\mathcal{GL}(R) = \coprod_{n \geq 0} GL_n(R),$$

and it is a skeleton of the category of $\text{iso } \mathcal{F}(R)$ of finitely generated free R -modules and isomorphisms.

We note that in general, even if \mathcal{C} is a G -category, the skeleton $\text{sk } \mathcal{C}$ is not closed under the G -action². However, if R is a G -ring, we have an obvious action on $\mathcal{GL}(R)$: it is trivial on objects and on morphisms g acts entrywise. Clearly, the inclusion of the skeleton

$$i: \mathcal{GL}(R) \longrightarrow \text{iso } \mathcal{F}(R)$$

is not an equivariant map since the object R^n is fixed in $\mathcal{GL}(R)$ but not in $\text{iso } \mathcal{F}(R)$. However, we can define an inverse to it which is equivariant. Fix isomorphisms $\gamma_M: M \xrightarrow{\cong} R^k$ for all finitely generated free modules M , i.e., fix a basis $\{\gamma_M^{-1}(e_i) = m_i\}$ for all M such that $\gamma_M = \gamma_{gM}$ as isomorphisms of abelian groups. In other words, we pick the same basis for M and gM ; recall that M and gM are equal as abelian groups. We define the inverse equivalence i^{-1} by $M \mapsto R^k$ on objects. Given an isomorphism $M \rightarrow N$ in $\text{iso } \mathcal{F}(R)$, it maps to the composite

$$R^k \xrightarrow{\gamma_M^{-1}} M \xrightarrow{f} N \xrightarrow{\gamma_N} R^k.$$

We show that the map i^{-1} is equivariant. Clearly, it commutes with the G -action on objects, since the action is trivial in $\mathcal{GL}(R)$ and if M has dimension k so does gM . Now let $f: M \rightarrow N$ be an isomorphism in $\text{iso } \mathcal{F}(R)$, and suppose that

$$f(m_i) = r_{i1}n_1 + \cdots + r_{ik}n_k.$$

The morphism $gM \xrightarrow{gf} gN$ maps to

$$R^k \xrightarrow{\gamma_M^{-1}} gM \xrightarrow{gf} gN \xrightarrow{\gamma_N} R^k.$$

On basis elements, this is

$$(gf)(e_i) = r_{i1} \cdot_g e_1 + \cdots + r_{ik} \cdot_g e_k = r_{i1}^g e_1 + \cdots + r_{ik}^g e_k.$$

²We treat this general case in the appendix.

Therefore, we get entrywise action by g on the matrix representing f , and the map

$$i^{-1}: \text{iso } \mathcal{F}(R) \longrightarrow \mathcal{GL}(R)$$

is G -equivariant. It is a nonequivariant equivalence, thus by Proposition 2.16 we obtain the following result.

Proposition 4.13. *Suppose R is a G -ring. Then there is a weak G -equivalence*

$$\mathcal{Cat}(\tilde{G}, \text{iso } \mathcal{F}(R)) \longrightarrow \mathcal{Cat}(\tilde{G}, \mathcal{GL}(R)).$$

This is very useful because it will allow us to use the skeleton $\mathcal{GL}(R)$ in equivariant algebraic K -theory without losing information about the entire category of free modules with its induced action of G .

4.5 Equivariant skeleta

Nonequivariantly, it is always assumed in K -theory that when we take the classifying space of a category which is not small, such as $\mathcal{P}(R)$, $\mathcal{F}(R)$, or $\text{Mod}(R)$, we are tacitly replacing the category by a small category which is equivalent to it, such as its skeleton.

As we have seen in Section 4.4, the situation is a little trickier equivariantly, because we do not have an equivariant equivalence between a G -category and its skeleton. This is too much to hope for; however, we show that the discussion in Section 4.4 generalizes. What makes the general case trickier is the fact that unlike in the case of free modules where we showed that $R^n \cong gR^n$, in general, an object C is not necessarily isomorphic to gC .

We show that for a G -category \mathcal{C} , we can put a G -action on the skeletal category $\text{sk } \mathcal{C}$, such that the inverse of the inclusion of the skeleton $i: \text{sk } \mathcal{C} \rightarrow \mathcal{C}$ is a G -map which is a nonequivariant equivalence. This implies by Proposition 2.16 that the map

$$\mathcal{Cat}(\tilde{G}, \mathcal{C}) \longrightarrow \mathcal{Cat}(\tilde{G}, \text{sk } \mathcal{C})$$

is a weak G -equivalence. This suffices for our applications, because in equivariant algebraic K -theory we are only taking classifying spaces of categories of the form $\mathcal{Cat}(\tilde{G}, \mathcal{C})$.

For an object $C \in \mathcal{C}$, denote by C^{rep} the representative of the isomorphism class of C in $\text{sk } \mathcal{C}$, so that if $C \cong D$, then $C^{rep} = D^{rep}$. We fix isomorphisms $\gamma_C: C \xrightarrow{\cong} C^{rep}$. The map

$$i^{-1}: \mathcal{C} \longrightarrow \text{sk } \mathcal{C}$$

is defined on objects as $C \mapsto C^{rep}$ and on morphisms as

$$(C \xrightarrow{f} D) \mapsto (C^{rep} \xrightarrow{\gamma_C^{-1}} C \xrightarrow{f} D \xrightarrow{\gamma_D} D^{rep}).$$

We define a G -action on $\text{sk } \mathcal{C}$ in the following way. On objects,

$$gC^{rep} := (gC)^{rep}.$$

We remark that there is no way to consistently pick the representatives such that $gC^{rep} = (gC)^{rep}$ is an equality in \mathcal{C} . However, we do have isomorphisms in \mathcal{C}

$$(gC)^{rep} \xrightarrow[\gamma_g^{-1}]{\cong} gC \xrightarrow[g\gamma_C]{\cong} gC^{rep}.$$

We define the action on morphisms of $\text{sk } \mathcal{C}$. We defined $g(C^{rep} \xrightarrow{f} D^{rep})$ as

$$(gC)^{rep} \xrightarrow[\gamma_g^{-1}]{\cong} gC \xrightarrow[g\gamma_C]{\cong} gC^{rep} \xrightarrow{gf} gD^{rep} \xrightarrow[g\gamma_D]{\cong} gD \xrightarrow[\gamma_{gD}]{\cong} (gD)^{rep}.$$

Now the map i^{-1} is clearly equivariant on objects. We show it is also equivariant on morphisms. Let $f: C \rightarrow D$ be a morphism in \mathcal{C} , which gets mapped by i^{-1} to $C^{rep} \xrightarrow{\gamma_C^{-1}} C \xrightarrow{f} D \xrightarrow{\gamma_D} D^{rep}$ in $\text{sk } \mathcal{C}$. Acting by g , we get

$$(gC)^{rep} \xrightarrow[\gamma_g^{-1}]{\cong} gC \xrightarrow[g\gamma_C]{\cong} gC^{rep} \xrightarrow[g\gamma_C^{-1}]{\cong} gC \xrightarrow{gf} gD \xrightarrow[g\gamma_D]{\cong} gD^{rep} \xrightarrow[g\gamma_D]{\cong} gD \xrightarrow[\gamma_{gD}]{\cong} (gD)^{rep}.$$

By composing the inverse isomorphism, this is the same as

$$(gC)^{rep} \xrightarrow[\gamma_g^{-1}]{\cong} gC \xrightarrow{gf} gD \xrightarrow[\gamma_{gD}]{\cong} (gD)^{rep},$$

which is just i^{-1} applied to $gC \xrightarrow{gf} gD$, and therefore, the map $i^{-1}: \mathcal{C} \rightarrow \text{sk } \mathcal{C}$ is a G -map for the action we defined on $\text{sk } \mathcal{C}$.

Since the equivariant algebraic K -theory construction involves replacing the usually non-small G -category of interest \mathcal{C} with $\mathcal{C}at(\tilde{G}, \mathcal{C})$, we can with clear conscience assume use of equivariantly skeletally small models when we apply the classifying space functor B .

4.6 Equivariant Morita theory

We give a definition of *equivariant Morita equivalence* of G -rings; the philosophy is that this notion should capture Morita equivalences of twisted group rings.

Definition 4.14. Two G -rings R and S are *equivariantly Morita equivalent* if they are nonequivariantly Morita equivalent and the equivalence

$$\text{Mod}(R) \rightarrow \text{Mod}(S)$$

is pseudo equivariant.

In [Bil12], Biland gives a definition of equivariant Morita equivalence, and it is easy to see

that his definition agrees with ours³. Biland shows that Definition 4.14 is equivalent to having a G -equivariant bimodule, which provides the equivariant Morita equivalence. For the definition of G -bimodule and the details of the equivalence of the two statements we refer the reader to Biland's preprint [Bil12, Thm. A].

Note that a consequence of our definition of equivariant Morita equivalence and Proposition 3.6 is the following proposition.

Proposition 4.15. *If two G -rings R and S are equivariantly Morita equivalent, then there is an equivariant weak equivalence*

$$\mathcal{C}at(\tilde{G}, \text{Mod}(R)) \longrightarrow \mathcal{C}at(\tilde{G}, \text{Mod}(S)).$$

Thus we have a G -map which induces an equivalence on all fixed points

$$\mathcal{C}at(\tilde{G}, \text{Mod}(R))^H \longrightarrow \mathcal{C}at(\tilde{G}, \text{Mod}(S))^H.$$

As we have shown in Proposition 4.8 this ensures that the twisted group rings $R_H[H]$ and $S_H[H]$ are Morita equivalent in the classical sense for all $H \subseteq G$.

We end with a consequence of equivariant Morita equivalence, which will be relevant in algebraic K -theory. Recall that a nonequivariant Morita equivalence $\text{Mod}(R) \rightarrow \text{Mod}(S)$ restricts to an equivalence $\mathcal{P}(R) \rightarrow \mathcal{P}(S)$ on the categories of finitely generated projective modules (for example, see [Wei13, II, 2.7.]).

Lemma 4.16. *If R and S are equivariantly Morita equivalent, then there is a weak G -equivalence*

$$\mathcal{C}at(\tilde{G}, \mathcal{P}(R)) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{P}(S))$$

which induces equivalences of the homotopy fixed point categories of finitely generated projective modules $\mathcal{P}(R)^{hH} \rightarrow \mathcal{P}(S)^{hH}$ for all $H \subseteq G$.

Proof. Since R and S are equivariantly Morita equivalent, by definition we have a nonequivariant Morita equivalence $\text{Mod}(R) \rightarrow \text{Mod}(S)$, which is pseudo equivariant. This restricts to an equivalence $\mathcal{P}(R) \rightarrow \mathcal{P}(S)$, which is pseudo equivariant, and we get the result by applying Proposition 3.6. The second statement follows by passing to fixed points. \square

5 Equivariant algebraic K -theory of G -rings

Nonequivariantly, the algebraic K -theory space of R is defined as the group completion of the classifying space of the symmetric monoidal category $\text{iso } \mathcal{P}(R)$ of finitely generated projective modules and isomorphisms, and this space is delooped using an infinite loop space machine such as the operadic one developed by May in [May72] or the one based on Γ -spaces developed by Segal in [Seg74].

³The only difference in the definitions is that Biland does not require $\theta_e = \text{id}$, but we suspect that is a typo in his preprint. Also, we note that his notion of equivariance is not the standard one; it corresponds to our notion of pseudo equivariance.

The category $\text{iso } \mathcal{P}(R)$ is a G -category with action defined as in the previous section, and it is not hard to see that it yields a *naive* Ω - G -spectrum, i.e., an Ω -spectrum with G -action. The Segalic infinite loop space machine has been generalized equivariantly by Shimakawa in [Shi89], and the operadic infinite loop space machine has been generalized equivariantly by Guillou and May in [GMa], to give *genuine* Ω - G -spectra. We show in [MMO] that the equivariant generalizations yield equivalent genuine G -spectra when fed equivalent input, so we can use either machine to deloop equivariant algebraic K -theory. We describe these machines and the input they take in section 5.4. It turns out that a symmetric monoidal category with G -action such as $\text{iso } \mathcal{P}(R)$ is inadequate input for these machines, but $\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))$ is the “genuine” kind of input these machines take to produce a genuine Ω - G -spectrum with zeroth space the equivariant group completion of $B\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))$.

The equivariant infinite loop space machines will provide a functorial model for the equivariant group completion of $B\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))$; however, we will first define the equivariant algebraic K -theory space of a G -ring R via an explicit model of the equivariant group completion of the classifying space of a symmetric monoidal G -category, namely the equivariant version of Quillen’s $S^{-1}S$ -construction. This model allows us to run an equivariant version of the first part of Quillen’s “plus=Q” proof relating the group completion of the symmetric monoidal category of finitely generated projective R -modules and isomorphisms to the group completion of the topological monoid $\coprod BGL_n(R)$ of classifying spaces of principal $GL_n(R)$ -bundles. Using this model, we show that the equivariant algebraic K -theory space of a G -ring R is equivalent on higher homotopy groups with an equivariant interpretation of the “plus” construction, namely the group completion of a topological G -monoid of G -equivariant $GL_n(R)$ -bundles.

5.1 Symmetric monoidal G -categories

We define a *symmetric monoidal G -category* as a strict symmetric monoidal category \mathcal{C} with G -action which commutes with the symmetric monoidal structure. Concisely, this is a functor $G \rightarrow \text{Sym}\mathcal{C}at^{\text{strict}}$, from G to the category of symmetric monoidal categories and *strict* monoidal functors. However, in practice, some symmetric monoidal categories that we care about have a G -action which preserves the monoidal structure only up to isomorphism, i.e., they are functors $G \rightarrow \text{Sym}\mathcal{C}at^{\text{strong}}$, from G to the category of strict symmetric monoidal categories and *strong* monoidal functors.

The problem with the latter is that if the G -action preserves the monoidal structure only up to isomorphism, the fixed point subcategories \mathcal{C}^H are not necessarily closed under the monoidal structure. We show that applying the functor $\mathcal{C}at(\tilde{G}, -)$ rectifies symmetric monoidal categories for which the action functors $g \cdot$ are only strong symmetric monoidal to symmetric monoidal G -categories with action that preserves the monoidal structure on the nose. Therefore the homotopy fixed point categories \mathcal{C}^{hH} are closed under the symmetric monoidal structure.⁴

Suppose that \mathcal{C} is defined by a functor $G \rightarrow \text{Sym}\mathcal{C}at^{\text{strong}}$. Then \mathcal{C} is a symmetric monoidal category for which the symmetric monoidal structure map

$$\mathcal{C} \times \mathcal{C} \xrightarrow{\oplus} \mathcal{C}$$

⁴This issue becomes even more subtle for Waldhausen categories as explained in Section 8.1.

is pseudoequivariant, where the G -action on $\mathcal{C} \times \mathcal{C}$ is diagonal, and for which $gI \cong I$ for every $g \in G$, where I is the unit object of \mathcal{C} . By Proposition 3.3, we get an on the nose equivariant functor

$$\oplus: \mathcal{C}at(\tilde{G}, \mathcal{C} \times \mathcal{C}) \cong \mathcal{C}at(\tilde{G}, \mathcal{C}) \times \mathcal{C}at(\tilde{G}, \mathcal{C}) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{C}).$$

For F_1, F_2 , the functor $F_1 \oplus F_2$ is defined on objects as

$$(F_1 \oplus F_2)(g) = g(g^{-1}F_1(g) \oplus g^{-1}F_2(g)),$$

which, of course, is the same as $F_1(g) \oplus F_2(g)$ when the G -action on \mathcal{C} preserves \oplus strictly. On a morphism (g', g) , $F_1 \oplus F_2$ is defined as

$$(F_1 \oplus F_2)(g) \xrightarrow{\cong} F_1(g) \oplus F_2(g) \xrightarrow{F_1(g', g) \oplus F_2(g', g)} F_1(g') \oplus F_2(g') \xrightarrow{\cong} (F_1 \oplus F_2)(g').$$

It is not hard to see that the functor $F_I: \tilde{G} \rightarrow \mathcal{C}$ defined by $F_I(g) = gI$, where I is the unit of \mathcal{C} is a unit for the symmetric monoidal structure defined above. Therefore, even when the G -action on \mathcal{C} does not preserve the symmetric monoidal structure strictly, $\mathcal{C}at(\tilde{G}, \mathcal{C})$ does become a symmetric monoidal G -category for which the action commutes with the symmetric monoidal structure.

5.2 The equivariant group completion of the classifying space of a symmetric monoidal G -category

A Hopf G -space is a Hopf space with equivariant multiplication map and for which multiplying by the identity element is G -homotopic to the identity map such as, for example, ΩX for a G -space X . The equivariant notion of group completion is captured by the fixed point maps being group completions.

Definition 5.1. A G -map $X \rightarrow Y$ of homotopy associative and commutative Hopf G -spaces is an *equivariant group completion* if the fixed point maps $X^H \rightarrow Y^H$ are group completions for all $H \subseteq G$.

Remark 5.2. For a homotopy commutative topological G -monoid, since the classifying space functor B and the loop functor Ω have the wonderful virtue of commuting with fixed points, the nonequivariant group completion map $M \rightarrow \Omega BM$ ([MS76], [May75]) is an equivariant group completion.

Note that the classifying space $B\mathcal{C}$ of a symmetric monoidal G -category is a Hopf G -space. We give a functorial construction of the group completion of $B\mathcal{C}$, following [Qui73]. The idea is to define the group completion on the category level. We recall the model for the categorical group completion in the nonequivariant case, and then we observe that the theory carries through equivariantly *as long as the G -action preserves the symmetric monoidal structure strictly*.

Definition 5.3 ([Qui73]). Let S be a symmetric monoidal category. The category $S^{-1}S$ has objects pairs (m, n) of objects in S . A morphism $(m, n) \rightarrow (p, q)$ in $S^{-1}S$ is an equivalence class

of triples

$$(r, r \oplus m \xrightarrow{f} p, r \oplus n \xrightarrow{g} q)$$

where two triple are equivalent if there is an isomorphism of the first entries that makes the relevant diagrams commute. Composition for a pair of morphisms is defined as

$$(r, r \oplus m \xrightarrow{f} p, r \oplus n \xrightarrow{g} q) \circ (s, s \oplus p \xrightarrow{\phi} u, s \oplus q \xrightarrow{\psi} v) = (s \oplus r, s \oplus r \oplus m \xrightarrow{\phi \circ (s \oplus f)} u, s \oplus r \oplus n \xrightarrow{\psi \circ (s \oplus g)} v).$$

Note that $S^{-1}S$ is symmetric monoidal with $(m, n) \oplus (p, q) = (m \oplus p, n \oplus q)$, and there is a strict monoidal functor $S \rightarrow S^{-1}S$ given by $m \mapsto (m, 0)$, where 0 is the unit of S . This induces a map of Hopf spaces

$$BS \longrightarrow BS^{-1}S, \quad (6)$$

which, subject to a mild condition, was shown by Quillen to be a group completion when S is a groupoid.

Theorem 5.4 ([Qui73]). *Let S be a symmetric monoidal groupoid such that translations are faithful. i.e.,*

$$\text{Aut}(s) \rightarrow \text{Aut}(s \oplus t)$$

is injective for all $s, t \in S$. Then the map $BS \rightarrow BS^{-1}S$ is a group completion.

Now if S is a symmetric monoidal G -category with G -action that preserves \oplus , then $S^{-1}S$ is also a symmetric monoidal G -category with diagonal action on objects. On morphisms,

$$g((m, n) \xrightarrow{(r, f, f')} (p, q)) = (gm, gn) \xrightarrow{(gr, gf, gf')} (gp, gq).$$

Note that this only works, because the action of G commutes with \oplus . The fixed point subcategory S^H is also a symmetric monoidal category, thus we can form $(S^H)^{-1}(S^H)$, and it is not hard to see that the construction commutes with fixed points.

Lemma 5.5. *Let S be a symmetric monoidal G -category. Then*

$$(S^H)^{-1}(S^H) \cong (S^{-1}S)^H$$

for all $H \subseteq G$.

Also, note that if translations are faithful in S , i.e., if $\text{Aut}(s) \rightarrow \text{Aut}(s \oplus t)$ is injective for all $s, t \in S$, then the same holds for the fixed point subcategories S^H . This has the following immediate consequence.

Proposition 5.6. *Let S be a symmetric monoidal G -groupoid such that translations are faithful. Then the map $BS \rightarrow BS^{-1}S$ is an equivariant group completion.*

We note the following useful result stating that restricting to the subcategory of isomorphisms commutes with applying the $\mathcal{C}at(\tilde{G}, -)$ functor.

Lemma 5.7. *For any G -category \mathcal{C} , we have an identification*

$$\text{iso } \mathcal{C}at(\tilde{G}, \mathcal{C}) \cong \mathcal{C}at(\tilde{G}, \text{iso } \mathcal{C}).$$

Proof. Note that a functor $F: \tilde{G} \rightarrow \mathcal{C}$ actually lands in $F: \tilde{G} \rightarrow \text{iso } \mathcal{C}$ since every morphism in \tilde{G} is an isomorphism. Therefore the objects of $\text{iso } \mathcal{C}at(\tilde{G}, \mathcal{C})$ and $\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{C})$ are the same. Now a morphism in $\text{iso } \mathcal{C}at(\tilde{G}, \mathcal{C})$ is a natural transformation whose component maps are all isomorphisms, which is the same with a morphism in $\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{C})$. \square

Now, using the explicit model for group completion we make the following definition of the equivariant algebraic K -theory space of a G -ring R .

Definition 5.8. The equivariant algebraic K -theory space of a G -ring R is the G -space $K_G(R) = B(S^{-1}S)$, where S is the symmetric monoidal G -category $\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))$.

We define the equivariant K -groups as the equivariant homotopy groups of this space. Once we deloop this space equivariantly to a genuine Ω - G -spectrum, it will turn out that these are the homotopy groups of that spectrum, and therefore, they have Mackey functor structure. Recall that for a subgroup $H \subseteq G$ and a G -space X , we have

$$\pi_i^H(X) = [(G/H)_+ \wedge S^i, X]_G \cong [S^i, X^H] = \pi_i(X^H), \quad (7)$$

where $[X, Y]_G$ denotes the set of homotopy classes of based G -maps $X \rightarrow Y$ between based G -spaces, and X_+ denotes the union of X with a disjoint basepoint. We define the K -groups for $i \geq 0$.

Definition 5.9. The algebraic K -theory groups are given by

$$K_i^H(R) = \pi_i^H(K_G(R)).$$

Remark 5.10. We spell out what the equivariant K_0 is so that it is clear how it relates back to the nonequivariant algebraic definition. Nonequivariantly, $K_0(R)$ is the group completion of the abelian monoid $\mathcal{P}(R)$ of finitely generated projective R -modules, i.e., it is π_0 of the topological group completion of $B(\text{iso } \mathcal{P}(R))$, the classifying space of the category of finitely generated projective modules and isomorphisms. Equivariantly, $K_0^H(R) \cong \pi_0(K_G(R)^H)$, and $K_G(R)^H$ is the group completion of $B\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))^H$. Therefore $K_0^H(R)$ is the group completion of the abelian monoid of isomorphism classes of objects in the homotopy fixed point category $\text{iso } \mathcal{P}(R)^{hH} = \text{iso } \mathcal{P}(R_H(H))$, and thus it agrees with K_0 of the twisted group ring $R_H(H)$.

5.3 “Plus” construction interpretation and connection to equivariant bundle theory

Quillen’s first definition of higher algebraic K -groups was as the homotopy groups of the space $BGL(R)^+$, which he showed to be homotopy equivalent to the basepoint component of the group completion of the topological monoid $B(\coprod_n GL_n(R)) \cong \coprod_n BGL_n(R)$. Note that this is the monoid of classifying spaces of principal $GL_n(R)$ -bundles under Whitney sum. Fiedorowicz, Hauschild and May gave a first definition of equivariant algebraic K -groups of a ring R with *trivial* G -action in [FHM82] by replacing this space with the monoid of classifying spaces of equivariant bundles. However, in their definition, since G does not act on R , the equivariance group G does not act on the structure group $GL_n(R)$ of the bundles; they are considering equivariant bundles that have commuting actions of G and $GL_n(R)$ on the total space.

We generalize the definition of Fiedorowicz, Hauschild and May so as to allow nontrivial action of G on the ring R . Instead of using the classifying spaces of equivariant $(G, GL_n(R))$ -bundles, which correspond to a trivial group extension

$$1 \rightarrow GL_n(R) \rightarrow GL_n(R) \times G \xrightarrow{q} G \rightarrow 1,$$

we will use the classifying spaces of $(G, GL_n(R) \rtimes G)$ -bundles, which correspond to split extensions

$$1 \rightarrow GL_n(R) \rightarrow GL_n(R) \rtimes G \xrightarrow{q} G \rightarrow 1.$$

For a precise definition of such equivariant bundles, see [GMM], or any of the earlier sources cited there. Suitable categorical models for classifying space of $(G, GL_n(R) \rtimes G)$ -bundles have been constructed in [GMM], and these are central to our definition. The relevant theorem is the following.

Theorem 5.11 ([GMM]). *If G is discrete and Π is discrete or compact Lie, then $B\mathcal{C}at(\tilde{G}, \Pi)$ is a classifying space for $(G, \Pi \rtimes G)$ -bundles.*

Recall from Definition 4.12 that $\mathcal{GL}(R) = \coprod_n GL_n(R)$. By Theorem 5.11, the monoid of classifying spaces of $(G, GL_n(R)_G)$ -bundles is $B\mathcal{C}at(\tilde{G}, \mathcal{GL}(R))$, and a model for the group completion is, by Remark 5.2,

$$\Omega BB\mathcal{C}at(\tilde{G}, \mathcal{GL}(R)).$$

We proceed to show that we have an equivalence

$$K_G(R)^H \simeq \Omega BB\mathcal{C}at(\tilde{G}, \mathcal{GL}(R))^H$$

on basepoint components, so these spaces have the same higher homotopy groups. This shows that the definition of [FHM82] of higher equivariant K -groups for a ring with trivial G -action agrees with the one given in Definition 5.9.

Again, we will follow Quillen's nonequivariant proof. We recall the definition of cofinality and then state the result that leads to showing that cofinality gives an equivalence on higher K -theory. The proof is in [Qui73] and a great exposition is also given in [Gui].

Definition 5.12. A monoidal functor $F: S \rightarrow T$ is cofinal if for every $t \in T$ there is $t' \in T$ and $s \in S$ such that $t \oplus t' = F(s)$.

Proposition 5.13 ([Qui73]). *If $F: S \rightarrow T$ is cofinal and $\text{Aut}_S(s) \cong \text{Aut}_T(F(s))$ for all $s \in S$, then the map $B(S^{-1}S) \rightarrow B(T^{-1}T)$ induces an equivalence of basepoint components.*

Nonequivariantly this is applied to the cofinal inclusion $\text{iso } \mathcal{F}(R) \hookrightarrow \text{iso } \mathcal{P}(R)$, which is in fact the idempotent completion. We wish to apply it to the fixed point maps of the inclusions

$$\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{F}(R))^H \longrightarrow \mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))^H.$$

For this we recall from section 4.3 that the homotopy fixed point category $\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{F}(R))^H$ is the category of finitely generated free R -modules with semilinear H -action and H -isomorphism. Similarly, $\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))^H$ is the category of finitely generated projective R -modules with

semilinear H -action. This inclusion is cofinal; however it is not an idempotent completion anymore. One way to see this is to consider the case when G is finite and $|G|^{-1} \in R$, so that, by Corollary 4.11, $\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{F}(R))^H \simeq \mathcal{P}(R_H[H])$. We have a commutative diagram of inclusions:

$$\begin{array}{ccc} \mathcal{C}at(\tilde{G}, \text{iso } \mathcal{F}(R))^H & \longrightarrow & \mathcal{P}(R_H[H]) \\ \uparrow & \nearrow & \\ \mathcal{F}(R_H[H]) & & \end{array}$$

The diagonal map is an idempotent completion, but the map going straight up is not an equivalence, since free $R_H[H]$ -modules do not coincide with modules with semilinear G -action which are free as R -modules. Therefore, the top map is not the idempotent completion; it just factors it.

Theorem 5.14. *There is an equivalence on connected basepoint components*

$$\Omega BB\mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(R))^H \simeq_0 K_G(R)^H.$$

(We used the notation \simeq_0 instead of \simeq in order to emphasize that this equivalence only holds on basepoint components and to avoid possible misinterpretation.)

Proof. By Proposition 4.13, the inverse of the nonequivariant equivalence given by the inclusion of the skeleton $i: \mathcal{G}\mathcal{L}(R) \rightarrow \text{iso } \mathcal{F}(R)$ induces a weak G -equivalence

$$\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{F}(R)) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(R)).$$

Now

$$\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{F}(R))^H \longrightarrow \mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))^H$$

is cofinal. Therefore, by applying Proposition 5.13, we get a weak G -equivalence of basepoint components

$$B(S^{-1}S) \rightarrow B(T^{-1}T)$$

for $S = \mathcal{C}at(\tilde{G}, \text{iso } \mathcal{F}(R))^H$ and $T = \mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))^H$. □

Remark 5.15. We note that equivariantly there is no meaningful way to write down a decomposition of the K -theory space as a product of K_0 and a connected component, analogous to the widely used nonequivariant one, which is

$$K(R) \simeq K_0(R) \times \Omega_0 BB\mathcal{G}\mathcal{L}(R).$$

The reason is that taking basepoint components does not commute with taking fixed points, so if we split off the basepoint component we change the equivariant homotopy type of the space. However, even nonequivariantly, this decomposition is not functorial, so it is technically more correct to define the K -theory space via a functorial model for the group completion of the

classifying space of the symmetric monoidal category of finitely generated projective modules and isomorphisms.

5.4 Equivariant delooping of the K -theory space

We describe the May and Segal equivariant infinite loop space machines and the input they take. By a celebrated theorem of May and Thomason, the nonequivariant infinite loop space machines are equivalent. Their proof does not generalize equivariantly, but we have shown in [MMO] through a surprising chain of equivalences that the equivariant generalizations of the machines also produce equivalent Ω - G -spectra. So, up to equivalence, we could use either machine to define the equivariant algebraic K -theory spectrum of a G -ring. The construction of each machine has its own advantages, and in some applications we have in mind we will need the specifics of one machine over the other. However, for the rest of this paper, we will study the equivariant homotopy type of the spectrum we get, and we will not need the specifics of either of these machines. We choose to define equivariant algebraic K -theory of a G -ring using the equivariant May machine, but we describe the alternative construction using the equivariant Segal machine and invoke the theorem by which they are equivalent.

5.4.1 Equivariant May infinite loop space machine

Algebras over the Barratt-Eccles operad \mathcal{O} in \mathcal{Cat} with $\mathcal{O}(j) = \tilde{\Sigma}_j$ are symmetric monoidal categories with strict unit and strict associativity, which are also known as *permutative categories* (see [May74]). By analogy, having an E_∞ -operad in $G\mathcal{Cat}$ allows one to define *genuine permutative G -categories* as algebras over it, and the classifying spaces of these turn out to be, after group completion, equivariant infinite loop spaces. This is carried out in the program started by Guillou and May in [GMa].

Of course, there are permutative categories, i.e., algebras over the Barratt-Eccles operad \mathcal{O} , which are also G -categories, and [GMa] reserves the name *naive permutative G -categories* for those. The reason is that their classifying spaces are G -spaces, which are naive equivariant infinite loop spaces, i.e., they have deloopings with respect to all spheres with trivial G -action, but not necessarily with respect to representation spheres. We note that in this light, what we defined as symmetric monoidal G -categories are *naive symmetric monoidal G -categories*.

Definition 5.16. The operad \mathcal{O}_G in $G\mathcal{Cat}$ defined by $\mathcal{O}_G(j) = \mathcal{Cat}(\tilde{G}, \tilde{\Sigma}_j)$ is an E_∞ - G -operad. A *genuine permutative G -category* is defined to be an \mathcal{O}_G -algebra.

If we take any *naive permutative category* \mathcal{C} , i.e., a permutative category with a G -action, since it is an algebra over the Barratt-Eccles operad \mathcal{O} with $\mathcal{O}(j) = \tilde{\Sigma}_j$, there are maps

$$\mathcal{O}(j) \times \mathcal{C}^j \longrightarrow \mathcal{C}$$

compatible with the operad structure maps. Since $\mathcal{Cat}(\tilde{G}, -)$ is a product preserving functor, these maps yield maps

$$\mathcal{Cat}(\tilde{G}, \tilde{\Sigma}_j) \times \mathcal{Cat}(\tilde{G}, \mathcal{C})^j \longrightarrow \mathcal{Cat}(\tilde{G}, \mathcal{C}),$$

and all the necessary diagrams still commute, so $\mathcal{C}at(\tilde{G}, \mathcal{C})$ is a genuine permutative category, and surprisingly, the only examples of genuine permutative categories we know arise in this way.

Example 5.17. Recall Definition 4.12 of the category $\mathcal{G}\mathcal{L}(R)$. It is a skeleton of the category of finitely generated free R -modules $\mathcal{F}(R)$. The category $\mathcal{G}\mathcal{L}(R)$ is permutative under direct sum of modules and block sum of matrices $\oplus : GL_n(R) \times GL_m(R) \rightarrow GL_{n+m}(R)$, since associativity and the unit are strict and commutativity holds only up to isomorphism (reordering of the basis elements by conjugation). It is a naive permutative G -category with trivial G -action on objects and entrywise G -action on matrices. Therefore the category $\mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(R))$ is a genuine permutative G -category.

The original May infinite loop space machine, which we will denote by \mathbb{K} , was developed in [May72]; it takes as input a permutative category and produces Ω -spectra with zeroth space the group completion of the classifying space of the input permutative category. An equivariant version of May's operadic infinite loop space machine is developed in [GMa]⁵. It takes as input an \mathcal{O}_G -category \mathcal{C} , i.e., a genuine permutative G -category, and produces a genuine orthogonal Ω - G -spectrum with zeroth space the equivariant group completion of $B\mathcal{C}$. We give a brief overview of the machine. As explained in [GMa], we need to use not only an E_∞ operad \mathcal{C}_G in $G\text{Top}$ (such as $B\mathcal{C}_G$), but also the Steiner operads \mathcal{K}_V indexed over finite dimensional subspaces of a complete G -universe U , because these act on V -fold loop spaces. These operads are described in detail in [GMa, Appendix]. Intuitively, they are a generalization of the little disks operad, which is compatible with suspension: instead of considering a tuple of embeddings of V into V , one considers a tuple of paths of embeddings of V into V , which at time 0 are the identity and at time 1 are disjoint. So the picture of an element in the Steiner operad would look like a cylinder with V at one end and an element of the little disks operad at the other end. We define the product operad

$$\mathcal{C}_V = \mathcal{C}_G \times \mathcal{K}_V.$$

A \mathcal{C}_G -space can be viewed as an \mathcal{C}_V -space for any V , and this has the advantage that \mathcal{C}_V acts on V -fold loop spaces via its projection onto \mathcal{K}_V . Let \mathbf{C}_V be the monad of based G -spaces associated to the operad \mathcal{C}_V .

For a genuine permutative G -category \mathcal{A} , the orthogonal G -spectrum $\mathbb{K}_G(\mathcal{A})$ has spaces given by the monadic bar constructions

$$\mathbb{K}_G \mathcal{A}(V) = B(\Sigma^V, \mathbf{C}_V, B\mathcal{C}).$$

The structure maps for $V \subset W$ are given by

$$\Sigma^{W-V} B(\Sigma^V, \mathbf{C}_V, X) \cong B(\Sigma^W, \mathbf{C}_V, X) \rightarrow B(\Sigma^W, \mathbf{C}_W, X).$$

Theorem 5.18 ([GMa]). *For a genuine permutative G -category \mathcal{C} , the spectrum $\mathbb{K}_G \mathcal{A}$ is a genuine Ω - G -spectrum and there is a group completion map $B\mathcal{A} \rightarrow (\mathbb{K}_G \mathcal{A})(0)$.*

The essential formal properties of the machine, which we will need, are the following theorems from [GMa].

⁵On the G -space level, operadic infinite loop space theory was first developed in unpublished work of Costenoble, Haushild, May, and Waner in the early 1980's.

Theorem 5.19 ([GMa]). *Let \mathcal{A} and \mathcal{B} be \mathcal{O}_G -categories. Then the map*

$$\mathbb{K}_G(\mathcal{A} \times \mathcal{B}) \rightarrow \mathbb{K}_G \mathcal{A} \times \mathbb{K}_G \mathcal{B}$$

induced by the projections is a weak equivalence of G -spectra.

Theorem 5.20 ([GMa]). *For \mathcal{O}_G -categories \mathcal{A} , there is a natural weak equivalence of spectra*

$$\mathbb{K}(\mathcal{A}^G) \rightarrow (\mathbb{K}_G \mathcal{A})^G.$$

The inclusion $\iota: \mathcal{O} \rightarrow \mathcal{O}_G$ induces a forgetful functor ι^* from genuine to naive permutative G -categories. Also, we have a forgetful functor i^* from genuine to naive G -spectra.

Theorem 5.21 ([GMa]). *For \mathcal{O}_G -categories \mathcal{A} , there is a natural weak equivalence of naive G -spectra $\mathbb{K} \iota^* \mathcal{A} \rightarrow i^* \mathbb{K}_G \mathcal{A}$.*

By Proposition 4.13, since $\mathcal{F}(F) = \mathcal{P}(F)$, the K -theory space of a field F with G -action is the equivariant group completion of $B\mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(F))$, and as we have seen in the example above, $\mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(F))$ is a genuine permutative G -category. Therefore, we can define

$$\mathbf{K}_G(F) = \mathbb{K}_G(\mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(F))).$$

Nonequivariantly, it is well known that using a construction of MacLane from [ML63], any symmetric monoidal category \mathcal{C} can be strictified to an equivalent permutative category \mathcal{C}^{str} , and therefore we can apply the nonequivariant infinite loop space machine \mathbb{K} to a symmetric monoidal category by implicitly doing this replacement first. The category \mathcal{C}^{str} has objects given by strings (c_1, \dots, c_n) of objects in \mathcal{C} , and morphisms

$$(c_1, \dots, c_n) \rightarrow (d_1, \dots, d_m)$$

given by morphisms

$$c_1 \oplus \dots \oplus c_n \rightarrow d_1 \oplus \dots \oplus d_m$$

in \mathcal{C} . The symmetric monoidal structure is given by concatenation and the identity is given by the empty string $()$.

This carries through equivariantly: if \mathcal{C} is a symmetric monoidal G -category, then \mathcal{C}^{str} is naturally also a symmetric monoidal G -category, with G -action given on objects by $g(c_1, \dots, c_n) = (gc_1, \dots, gc_n)$. Since G commutes with \oplus , we can define the action on morphisms by

$$g((c_1, \dots, c_n) \xrightarrow{f} (d_1, \dots, d_m)) = (gc_1, \dots, gc_n) \xrightarrow{gf} (gd_1, \dots, gd_m).$$

It is not hard to see the inverse functors in the equivalence $C \simeq C^{str}$ are G -equivariant. Therefore, given a symmetric monoidal G -category \mathcal{C} , the naive permutative G -category \mathcal{C}^{str} is G -equivalent to it.

Using this strictification implicitly, we can use the operadic machine on symmetric monoidal G -categories. We give the following definition for all G -rings.

Definition 5.22. We define the equivariant algebraic K -theory spectrum of R as

$$\mathbf{K}_G(R) = \mathbb{K}_G(\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))),$$

with the understanding that we have replaced the symmetric monoidal G -category $\mathcal{P}(R)$ with an equivalent naive permutative G -category.

We note that since the zeroth space $\mathbf{K}_G(R)(0)$ is the group completion of $B\mathcal{C}at(\tilde{G}, \text{iso } \mathcal{P}(R))$, this is equivalent to our previous definition of the K -theory space $K_G(R)$. Alternatively, we can use the equivariant Segal machine for delooping this space, which we address in the next section.

5.4.2 Equivariant Segal infinite loop space machine

Segal developed an alternative delooping machine to the operadic May machine in the celebrated paper [Seg74], which we will denote as \mathbb{S} . The input is a Γ -space, which is just a functor

$$X: \mathcal{F} \rightarrow \text{Top}, \quad \mathbf{n} \mapsto X_{\mathbf{n}},$$

where \mathcal{F} is a skeleton of the category of based finite sets⁶. A Γ -space is *special* if the map $\delta: X_{\mathbf{n}} \rightarrow X_{\mathbf{1}^n}$, induced by the projections $\delta_i: \mathbf{n} \rightarrow \mathbf{1}$, is an equivalence. From a Γ -space, Segal produces a spectrum, and he shows that for a special Γ -space, the spectrum is Ω , with zeroth space the group completion of $X_{\mathbf{1}}$.

One can start with a Γ -category instead, i.e., a functor $\mathcal{F} \rightarrow \mathcal{C}at$ and define it to be special if the Γ -space obtained by applying the classifying space functor levelwise is a special Γ -space. Segal gives a construction of a special Γ -category X from a symmetric monoidal category \mathcal{C} , with $X_{\mathbf{1}} \simeq \mathcal{C}$. Therefore, $\mathbb{S}(\mathcal{C})$, the spectrum obtained from the special Γ -space associated to the symmetric monoidal category \mathcal{C} , is Ω , with zeroth space the group completion of $B\mathcal{C}$.

Shimakawa has generalized Segal's machine in [Shi89] to produce an orthogonal genuine Ω - G -spectrum starting from a special Γ_G -space. A Γ_G -space is a functor

$$X: \mathcal{F}_G \rightarrow \text{Top}_G, \quad A \mapsto X(A),$$

where \mathcal{F}_G is the category of finite G -sets and Top_G is the category of G -spaces and nonequivariant based maps; G acts by conjugation on morphism sets. For any $A \in \mathcal{F}_G$, we have a projection $\delta_a: A \rightarrow \mathbf{1}$, which sends all the nonbasepoint elements of A to 1 and the basepoint to 0. A Γ_G -space is special if the map $\delta_A: X(A) \rightarrow \text{Map}(A, X_{\mathbf{1}})$ induced by these projections is a G -equivalence for all $A \in \mathcal{F}_G$. We note that this map turns out to be a G -map even though the individual maps δ_a are generally not G -maps.

Given a Γ_G -space X , Shimakawa constructs a spectrum $\mathbb{S}_G X$ with V th space given by the two-sided bar construction $B((S^V)^\bullet, \mathcal{F}_G, X)$, where $(S^V)^\bullet$ is the contravariant functor $\mathcal{F}_G \rightarrow \text{Top}_G$ defined on objects as $A \mapsto \text{Map}(A, S^V)$. It is not hard to see that there are structure maps

$$S^W \wedge B((S^V)^\bullet, \mathcal{F}_G, X) \rightarrow B((S^{V \oplus W})^\bullet, \mathcal{F}_G, X).$$

⁶The opposite of Segal's original category Γ turns out to be just \mathcal{F} .

The following is the main theorem in [Shi89].

Theorem 5.23 ([Shi89]). *For a special Γ_G -space X , the spectrum $\mathbb{S}_G X$ is a genuine Ω - G -spectrum, for which $X_1 \simeq (\mathbb{S}_G X)(0)$ if and only if X_1 is grouplike.*

Essential to our applications is that in general there is a group completion map $X_1 \rightarrow (\mathbb{S}_G X)(0)$, which Shimakawa does not prove, but we fill this gap in [MMO].

A Γ_G -category is a functor $\mathcal{F}_G \rightarrow \mathcal{Cat}_G$, where \mathcal{Cat}_G is the category of G -categories and nonequivariant functors. It is special if the Γ_G -space obtained by applying the classifying space functor levelwise is special. Shimakawa generalizes Segal's combinatorial way of constructing a Γ -category from a symmetric monoidal category to construct a Γ_G -category from a symmetric monoidal G -category \mathcal{C} . This Γ_G -category is not necessarily special, but Shimakawa shows that replacing \mathcal{C} by the symmetric monoidal G -category $\mathcal{Cat}(\tilde{G}, \mathcal{C})$ does yield a special Γ_G -category, and therefore, $\mathbb{S}_G(\mathcal{Cat}(\tilde{G}, \mathcal{C}))$, the machine applied to the special Γ_G -category obtained from the symmetric monoidal G -category $\mathcal{Cat}(\tilde{G}, \mathcal{C})$, is a genuine orthogonal Ω - G -spectrum with zeroth space the group completion of $B\mathcal{Cat}(\tilde{G}, \mathcal{C})$.

In [MMO], we prove that the two equivariant delooping machines agree; in particular, we have the following result.

Theorem 5.24. [MMO] *For a symmetric monoidal G -category \mathcal{C} we have an equivalence of orthogonal G -spectra $\mathbb{K}_G(\mathcal{Cat}(\tilde{G}, \mathcal{C})) \simeq \mathbb{S}_G(\mathcal{Cat}(\tilde{G}, \mathcal{C}))$.*

Corollary 5.25. *There is an equivalence of orthogonal Ω - G -spectra*

$$\mathbf{K}_G(R) \simeq \mathbb{S}_G(\mathcal{Cat}(\tilde{G}, \text{iso } \mathcal{P}(R))).$$

5.5 Functoriality of \mathbf{K}_G

Now we address functoriality of the construction. Even nonequivariantly, the assignment $R \mapsto \mathcal{P}(R)$ is not a functor, but just a pseudo functor, since composition is not preserved strictly. One way to rectify any pseudo functor landing in \mathcal{Cat} to an actual functor is using Street's first construction from [Str72]. This generalizes equivariantly to strictify a pseudo functor landing in $G\mathcal{Cat}$ to an actual functor. We will tacitly assume this strictification from now on, and address the new subtleties that arise and are specific to the equivariant case. The issue that arises is that for a G -map of G -rings $R \rightarrow S$, the functor $\mathcal{P}(R) \rightarrow \mathcal{P}(S)$ is *not* equivariant, so it is not a morphism in $G\mathcal{Cat}$. So for sure the assignment $R \mapsto \mathcal{P}(R)$ is not a functor or even a pseudo functor. We show that this gets rectified after applying $\mathcal{Cat}(\tilde{G}, -)$.

Theorem 5.26. *The assignment $R \mapsto \mathbf{K}_G(R)$ is a functor from the category of G -rings and G -maps to genuine orthogonal G -spectra.*

Proof. The equivariant infinite loop space machine \mathbb{K}_G is a functor from the category of genuine permutative G -categories and G -maps between them to the category of genuine orthogonal G -spectra. Thus it suffices to show that having a map of G -rings $R \rightarrow S$ yields an *equivariant* map $\mathcal{Cat}(\tilde{G}, \mathcal{P}(R)) \rightarrow \mathcal{Cat}(\tilde{G}, \mathcal{P}(S))$.

Suppose $f: R \rightarrow S$ is a G -map of G -rings, and consider the functor $\mathcal{P}(R) \rightarrow \mathcal{P}(S)$ defined as $M \mapsto M \otimes_R S$. Note that certainly $gM \otimes_R S \neq g(M \otimes_R S)$ since the scalar multiplication is different in the two modules; however we go on to define an isomorphism

$$gM \otimes_R S \cong g(M \otimes_R S).$$

Recall that in gM , the scalar multiplication is defined as $r \cdot_g m = r^g m$, where $(-)^g$ denotes the action of g on R . Define

$$gM \otimes_R S \rightarrow g(M \otimes_R S), \quad m \otimes s \mapsto m \otimes s^g.$$

First of all, we use the assumption that f is a G -map to show that this assignment is well-defined. Note that for $t \in R$, we have the following identification in $gM \otimes_R S$:

$$m \otimes ts \sim f(t) \cdot_g m \otimes s = f(t)^g m \otimes s,$$

Now

$$m \otimes ts \mapsto m \otimes (ts)^g = m \otimes t^g s^g \sim f(t^g) m \otimes s^g,$$

and

$$f(t)^g m \otimes s \mapsto f(t)^g m \otimes s^g,$$

but these are equal since $f(t^g) = f(t)^g$.

We check next that the assignment is S -linear: for $t \in S$,

$$t(m \otimes s) = m \otimes ts \mapsto m \otimes (ts)^g = m \otimes t^g s^g = t \cdot_g (m \otimes s^g).$$

Similarly, we can check that the inverse map

$$g(M \otimes_R S) \rightarrow gM \otimes_R S, \quad m \otimes s \mapsto m \otimes s^{(g^{-1})}$$

is well-defined and S -linear, so that we have the claimed isomorphism.

It is not hard to see that these isomorphisms make the functor $- \otimes_R S$ pseudo equivariant, and Proposition 3.3 provides the desired G -map

$$\mathcal{C}at(\tilde{G}, \mathcal{P}(R)) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{P}(S)),$$

which in turn gives a map of genuine G -spectra

$$\mathbf{K}_G(R) \rightarrow \mathbf{K}_G(S)$$

by the functoriality of the equivariant infinite loop space machine \mathbb{K}_G . □

5.6 Properties of the equivariant algebraic K -theory spectrum $\mathbf{K}_G(R)$

Now we can exploit the results that we have proved in section 4 about the homotopy fixed points of module categories. An immediate consequence Theorem 5.20 and Proposition 4.10 is the following theorem, which says that we recover the classical nonequivariant K -theory of twisted group rings as the fixed points of our construction.

Theorem 5.27. *If $H \subseteq G$ and $|H|^{-1} \in R$, there is an equivalence of spectra*

$$\mathbf{K}_G(R)^H \simeq \mathbf{K}(R_H[H]).$$

By Lemma 4.16, we immediately get that the equivariant algebraic K -theory of G -rings is equivariantly Morita invariant.

Proposition 5.28. *If R and S are equivariantly Morita equivalent, then there is a G -equivalence*

$$K_G(R) \simeq K_G(S).$$

6 Equivariant K -theory of Galois extensions

The algebraic K -theory of Galois extensions behaves particularly nicely as a result of faithfully flat descent and the fact that for G -Galois extensions the category of descent data has an interpretation in terms of modules with semilinear G -action.

6.1 Galois extensions of rings

Galois extensions of rings have been introduced and first studied by Auslander and Golman in [AG60]. For a ring extension $R \rightarrow S$, let $\text{Aut}_R(S)$ be the group of automorphisms of S fixing R . We recall the definition.

Definition 6.1. Let $R \rightarrow S$ be a faithfully flat ring extension and suppose that G is a finite subgroup of $\text{Aut}_R(S)$. The extension $R \rightarrow S$ is Galois with Galois group G if the map

$$\gamma: S \otimes_R S \rightarrow \prod_G S, \quad a \otimes b \mapsto ((g \cdot a)b)_{g \in G}. \quad (8)$$

is an S -algebra isomorphism.

It is an easy exercise to see that $R = S^G$. The wonderful fact about Galois extensions is that if $R \rightarrow S$ is a Galois ring extension with Galois group G , then the category of S -modules with semilinear G -action is equivalent to the category of S -modules with descent data, and in turn, by faithfully flat descent this is equivalent to the category of modules over R .

In the proof of Proposition 4.8 we showed that for a G -ring S , the category of S -modules with semilinear G -action is equivalent to the homotopy fixed point category $\mathcal{C}at(\widetilde{G}, \text{Mod}(S))^G$. Thus we have the following proposition.

Proposition 6.2. *Suppose $R \rightarrow S$ is a Galois ring extension with Galois group G . Then there is an equivalence of categories*

$$\mathrm{Mod}(R) \simeq \mathcal{C}at(\tilde{G}, \mathrm{Mod}(S))^G.$$

Suppose $R \rightarrow S$ is faithfully flat. Then an R -module M is finitely generated projective if and only if the S -module $M \otimes_R S$ is finitely generated projective (see [Cha13, Prop. 2.12.]). Therefore, the equivalence of categories mentioned above restrict to the corresponding categories of finitely generated projective modules, and we obtain the following analogue of Proposition 6.2.

Proposition 6.3. *Suppose $R \rightarrow S$ is a Galois ring extension with Galois group G . Then there is an equivalence of categories*

$$\mathcal{P}(R) \simeq \mathcal{C}at(\tilde{G}, \mathcal{P}(S))^G.$$

This leads to the following theorem, which says that for a G -Galois extension $R \rightarrow S$, the G -fixed point spectrum of the G -equivariant K -theory of S is the same as the nonequivariant K -theory spectrum of the fixed ring $S^G = R$.

Theorem 6.4. *Let $R \rightarrow S$ be a Galois extension of rings with Galois group G . Then there is an equivalence of orthogonal spectra*

$$\mathbf{K}_G(S)^G \simeq \mathbf{K}(R).$$

Proof. By Proposition 6.3, we have an equivalence of categories

$$\mathcal{P}(R) \simeq \mathcal{C}at(\tilde{G}, \mathcal{P}(S))^G.$$

By Theorem 5.20, we have

$$\mathbb{K}_G(\mathcal{C}at(\tilde{G}, \mathcal{P}(S)))^G \simeq \mathbb{K}(\mathcal{C}at(\tilde{G}, \mathcal{P}(S))^G).$$

Therefore,

$$\mathbf{K}_G(S)^G \simeq \mathbf{K}(R).$$

□

Example 6.5. For any finite G -Galois extension of fields E/F , $\mathbf{K}_G(E)^G \simeq \mathbf{K}(F)$. In particular, this recovers $\mathbf{K}(\mathbb{Q})$ as the fixed point spectrum of the genuine equivariant K -theory spectrum of any finite Galois extension of \mathbb{Q} .

Example 6.6. For any ring R , the diagonal map $R \rightarrow R \times R$ is a Galois extension with group $\mathbb{Z}/2\mathbb{Z}$, where the nontrivial element acts on $R \times R$ by interchanging the factors. Thus,

$$\mathbf{K}_{\mathbb{Z}/2\mathbb{Z}}(R \times R)^{\mathbb{Z}/2\mathbb{Z}} \simeq \mathbf{K}(R).$$

6.2 Strong form of Hilbert's theorem 90

As an accidental corollary of our interpretations of homotopy fixed point categories of modules, we obtain a new proof of Serre's generalization of Hilbert's theorem 90. The original Hilbert

90 theorem states that for a Galois extension E/F with Galois group G , the cohomology group $H^1(G, E^\times)$ is trivial. This theorem is reproved in [Del77] using faithfully flat descent. In the same spirit, we use Proposition 6.2 to give an alternative proof of the generalization of Hilbert's theorem 90, which is due to Serre:

Theorem 6.7 (Serre). *The nonabelian cohomology $H^1(G, GL_n(E))$ is trivial.*

Proof. Faithfully flat descent gives an equivalence of categories between the category of F -modules and the category of descent data, which in turn is equivalent to the category of E -modules with semilinear G -action. From the proof of Proposition 4.8, the latter is just $\mathcal{C}at(\tilde{G}, \text{iso Mod}(E))^G$. Note that $\coprod_n GL_n(E)$ is a skeleton of $\text{iso Mod}(E)$, and by Proposition 2.16 we have a weak equivalence

$$\mathcal{C}at(\tilde{G}, \text{iso Mod}(E))^G \simeq \mathcal{C}at(\tilde{G}, \coprod_n GL_n(E))^G.$$

Using that $\coprod_n GL_n(F)$ is a skeleton for $\text{iso Mod}(F)$ and the description of the fixed points of $\mathcal{C}at(\tilde{G}, GL_n(E))$ given in Theorem 2.14, we have

$$\coprod_n GL_n(F) \simeq \coprod_n \mathcal{C}at_\times(G, GL_n(E)),$$

where $\mathcal{C}at_\times(G, GL_n(E))$ is the crossed functor category defined in Theorem 2.14. Therefore, for any summand in the coproduct, there is only one isomorphism class of objects. Now we note that the isomorphism set of objects in $\mathcal{C}at_\times(G, GL_n(E))$ is precisely the first nonabelian cohomology set $H^1(G, GL_n(E))$. \square

This gives the following result, which we could have used directly to conclude that for a finite Galois extension of fields E/F with Galois group G we have an equivalence $\mathbf{K}_G(E)^G \simeq \mathbf{K}(F)$.

Proposition 6.8. *There is a symmetric monoidal weak G -equivalence*

$$\iota: \mathcal{G}\mathcal{L}(E) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(E)).$$

Proof. By Proposition 2.15, we have that the map

$$\iota: \mathcal{G}\mathcal{L}(E)^H \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(E))^H$$

is an equivalence precisely when $H^1(G, GL_n(E))$ is trivial, and this is true in this case by Theorem 6.7. Now also note that

$$\mathcal{G}\mathcal{L}(E)^H = \mathcal{G}\mathcal{L}(E^H).$$

This completes the proof. \square

6.3 Remarks on profinite Galois extensions

Let k be a field with separable closure \bar{k} and absolute Galois group $\text{Gal}(\bar{k}/k)$. Then

$$\text{Gal}(\bar{k}/k) = \lim_{\leftarrow} \text{Gal}(L/k),$$

where the limit runs over all finite Galois extensions L/k . The absolute Galois group is a profinite group with the profinite group topology, and it acts continuously through group homomorphisms on the discrete group \bar{k}^\times , or more generally, on $GL_n(\bar{k})$.

There are inclusions

$$H^1(\text{Gal}(L/k), GL_n(L)) \hookrightarrow H^1(\text{Gal}(\bar{k}/k), GL_n(\bar{k}))$$

coming from the inclusions of cocycles, which by definition need to be *continuous* crossed homomorphisms. We get

$$H^1(\text{Gal}(\bar{k}/k), GL_n(\bar{k})) = \text{colim} H^1(\text{Gal}(L/k), GL_n(\bar{k})),$$

where the colimit runs over all finite Galois extensions.

Thus we get an equivalence analogous to the one in proposition Proposition 6.8:

$$\mathcal{GL}(k) \xrightarrow{\simeq} \mathcal{C}at(\text{Gal}(\bar{k}/k), \mathcal{GL}(\bar{k}))^{\text{Gal}(\bar{k}, k)}.$$

However, the equivariant infinite loop space machines, as currently developed, do not apply to profinite groups, so we cannot pass from this statement to a spectrum level statement. We do hope, though, that in future work we will be able to generalize the delooping machines to profinite groups.

6.4 Quillen-Lichtenbaum formulation

Let E/F be a finite Galois extension of fields with Galois group $G = \text{Gal}(E/F)$ on E . Since G acts on E , it acts by functoriality on the spectrum $\mathbf{K}E$, so $\mathbf{K}E$ is a naive G -spectrum. The fixed points of this naive G -spectrum are easily seen to be $(\mathbf{K}E)^G \simeq \mathbf{K}F$. The initial Quillen-Lichtenbaum conjecture was that the map of spectra

$$\mathbf{K}F \simeq \mathbf{K}E^G \longrightarrow \mathbf{K}E^{hG}$$

is an equivalence, where $\mathbf{K}E^{hG}$ denotes the homotopy fixed points of the naive G -spectrum $\mathbf{K}E$. However, low dimensional examples disprove this conjecture as stated even after p -adic completion for a prime p . Thomason showed that this map only becomes an equivalence after reducing mod a prime power and inverting the Bott element.

One might ask the same question of the map from fixed points to homotopy fixed points of the genuine K -theory G -spectrum of E and hope that it becomes an equivalence there, but we show that that map is equivalent to the one in terms of naive spectra from the original

Quillen-Lichtenbaum conjecture. We review the definition of homotopy fixed points of a fibrant genuine orthogonal G -spectrum X . Note that a fibrant spectrum is an Ω - G -spectrum.

Definition 6.9. Let X be a fibrant genuine orthogonal G -spectrum. Then X^{hG} is defined to be the fixed point spectrum

$$\mathrm{Map}_*(EG_+, X)^G = (i^* \mathrm{Map}_*(EG_+, X))^G \simeq \mathrm{Map}_*(EG_+, i^* X)^G,$$

where i^* is the forgetful functor from genuine to naive G -spectra.

Just as for G -spaces and naive G -spectra, we have a natural map $X^G \rightarrow X^{hG}$, induced by the projection $EG_+ \rightarrow S^0$.

Proposition 6.10. *Let E/F be a finite Galois extension with Galois group G . The map from fixed points to homotopy fixed points of genuine G -spectra*

$$\mathbf{K}_G(E)^G \longrightarrow \mathbf{K}_G(E)^{hG}$$

is equivalent to the map from fixed points to homotopy fixed points of naive G -spectra

$$\mathbf{K}F \longrightarrow \mathbf{K}^{hG}.$$

From Example 6.5, we have that $\mathbf{K}F \simeq \mathbf{K}_G(E)^G$. We show in the next proposition that $\mathbf{K}_G(E)^{hG} \simeq \mathbf{K}(E)^{hG}$, where on the left hand side we are taking homotopy fixed points of a genuine G -spectrum, and on the right hand side we are taking homotopy fixed points of a naive G -spectrum. This proves the above proposition, and thus we recover the initial form of the Quillen-Lichtenbaum conjecture as a statement about genuine G -spectra.

Lemma 6.11. *For a Galois extension E/F with Galois group G , the homotopy fixed points of the naive G -spectrum $\mathbf{K}E$ and the homotopy fixed points of the genuine G -spectrum $\mathbf{K}_G(E)$ are equivalent.*

Proof. We will show that we have an equivalence of naive G -spectra $i^* \mathbf{K}_G(E) \simeq \mathbf{K}E$, which will imply the result. Recall that $\mathbf{K}E$ is defined as the K theory of the naive permutative G -category $\mathcal{G}\mathcal{L}(E)$, while $\mathbf{K}_G(E)$ is the equivariant algebraic K -theory of the genuine permutative G -category $\mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(E))$. We have a map

$$\mathbf{K}(\mathcal{G}\mathcal{L}(E)) \rightarrow \mathbf{K}(i^*(\mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(E)))) \xrightarrow{\simeq} i^* \mathbf{K}_G(\mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(E))),$$

where the second map is shown to be an equivalence in [GMa].

By Proposition 6.8, there is a symmetric monoidal weak G -equivalence

$$\mathcal{G}\mathcal{L}(E) \simeq i^*(\mathcal{C}at(\tilde{G}, \mathcal{G}\mathcal{L}(E))),$$

so the first map is also an equivalence. □

6.5 Carlsson's assembly map from the equivariant perspective

There has been a long standing program initiated and lead by G. Carlsson of studying the K -theory of fields motivated by the concept of descent and the Quillen-Lichtenbaum conjecture. Suppose that E/F is a Galois extension with Galois group G . We can consider the *assembly map* induced in K -theory by extension of scalars

$$\mathrm{Rep}_F[G] \xrightarrow{E \otimes_F -} V^G(E), \quad (9)$$

from the category of continuous finite dimensional G -representations in F , which is denoted by $\mathrm{Rep}_F[G]$ in [Car11], to the category of finite dimensional E -vector spaces with semilinear G -action, or equivalently, the category of E -vector spaces with descent data for the faithfully flat extension E/F , denoted by $V^G(E)$ in [Car11], and which is equivalent to the category $\mathrm{Vect}(F)$ of finite dimensional F -vector spaces by Proposition 6.2. Since all these categories are nonequivariant symmetric monoidal categories, their K -theory spectra are defined by using standard nonequivariant infinite loop space machines such as the May [May72] or the Segal machine [Seg74].

Carlsson conjectured that for the Galois extension \bar{F}/F with absolute Galois group G and an algebraically closed subfield $k \hookrightarrow F$ the composite

$$\mathrm{Rep}_k[G] \rightarrow \mathrm{Rep}_F[G] \xrightarrow{\bar{F} \otimes_F -} V^G(E) \simeq \mathrm{Vect}(F). \quad (10)$$

induces an equivalence on K -theory after derived completion, i.e.,

$$\mathbf{K}(\mathrm{Rep}_k[G])_{\alpha_p}^{\wedge} \longrightarrow \mathbf{K}(F)_{\alpha_p}^{\wedge} \simeq \mathbf{K}(F)_p^{\wedge}.$$

Carlsson defines *the derived completion* of a ring spectrum in [Car08] and shows that for $\mathbf{K}F$ the derived completion agrees with the Bousfield-Kan p completion, which accounts for the identification made above on the right hand side. This conjecture has recently been proved by G. Carlsson for pro- l absolute Galois groups in [Car] and [CR]. Attacking the problem from the perspective of full-fledged equivariant spectra might eventually shed some light on the general case. C. Barwick has announced such a proof from an ∞ -categorical point of view using spectral Mackey functors.

The key to any equivariant point of view is, of course, to interpret the assembly map as the fixed point map of an equivariant map between genuine G -spectra. We show how to construct such an equivariant map. On the categorical level our definition makes sense for the separable Galois extension of a field and the profinite Galois group; however, we only know how to obtain a map of genuine G -spectra for a finite Galois extension at the moment, because of the limitations of the equivariant infinite loop space machines. It is very easy to see that the source and target of the assembly map are instances of fixed points of our construction of equivariant algebraic K -theory, but it is non-trivial to see that there is an equivariant map which restricts to the assembly map on fixed points. This relies on the fact that our construction of equivariant algebraic K -theory turns pseudo-equivariant maps to on the nose equivariant maps.

Proposition 6.12. *Suppose E/F is a finite Galois extension with Galois group G . There is a*

G-map of genuine *G*-spectra

$$\mathbf{K}_G(F) \longrightarrow \mathbf{K}_G(E),$$

which, on *G*-fixed points, restricts to the assembly map

$$\mathbf{K}(\mathrm{Rep}_F[G]) \longrightarrow \mathbf{K}(V^G(E)).$$

Proof. It is easy to identify the source and target. From Proposition 4.8, the category $V^G(E)$ of *E*-vector spaces with semilinear *G*-action is isomorphic to $\mathcal{C}at(\tilde{G}, \mathrm{Vect}(F))^G$. As we have remarked before, if *G* is acting trivially on *F*, $\mathcal{C}at(\tilde{G}, \mathrm{Vect}(F))^G \cong \mathcal{C}at(G, \mathrm{Vect}(F))$, the category of *F*-vector spaces with *G*-linear action, i.e., the category of continuous *G*-representations over *F*. Thus $\mathrm{Rep}_F[G] \cong \mathcal{C}at(\tilde{G}, \mathrm{Vect}(F))^G$. Therefore, the map from equation (9), translates to a map

$$\mathcal{C}at(\tilde{G}, \mathrm{Vect}(F))^G \longrightarrow \mathcal{C}at(\tilde{G}, \mathrm{Vect}(E))^G.$$

It remains to show that we indeed have an equivariant map

$$\mathcal{C}at(\tilde{G}, \mathrm{Vect}(F)) \longrightarrow \mathcal{C}at(\tilde{G}, \mathrm{Vect}(E)),$$

inducing this map on fixed points.

Note that the extension of scalars map $\mathrm{Vect}(F) \xrightarrow{E \otimes_F -} \mathrm{Vect}(E)$ is not a *G*-map. The action of *G* on $\mathrm{Vect}(F)$ is trivial; however, for $V \in \mathrm{Vect}(F)$, the object $E \otimes_F V$ is not *G*-fixed:

$$E \otimes_F V \neq g(E \otimes_F V)$$

since they have different scalar multiplication. However, we have shown that for free *R*-modules *M* over a *G*-ring *R*, we have that $gM \cong M$. Thus in this case,

$$E \otimes_F V \cong g(E \otimes_F V)$$

and the extension of scalars functor is pseudo equivariant. By Proposition 3.3 this induces an equivariant map $\mathcal{C}at(\tilde{G}, \mathrm{Vect}(F)) \rightarrow \mathcal{C}at(\tilde{G}, \mathrm{Vect}(E))$, which on application of the equivariant infinite loop space machine \mathbf{K}_G yields a map

$$\mathbf{K}_G(F) \longrightarrow \mathbf{K}_G(E),$$

which restricts to the assembly map on fixed points. □

7 Equivariant algebraic *K*-theory of topological rings

We describe how our construction of equivariant algebraic *K*-theory recovers the connective covers of the the well-known equivariant topological real and complex *K*-theories KU_G and KO_G , defined in [Seg68] and Atiyah's Real *K*-theory KR , defined in [Ati66]. When Atiyah introduced KR , he described it as a mixture of real *K*-theory KO and equivariant topological

K -theory KU_G and KO_G . We show that they all fit under the unifying framework of equivariant K -theory developed in this paper.

In this section, any time we refer to topological K -theory, we mean the connective version. We denote by ku_G , ko_G and kr the connective covers of KU_G , KO_G , and KR , respectively. Whereas the first two are well studied, the latter is not so well-known. A construction of the connective cover of KR is given, for example, in [Dug05].

Recall that as topological groups $GL_n(\mathbb{C}) \simeq U_n$ and $GL_n(\mathbb{R}) \simeq O_n$, and if one takes the topology into account when forming the bar construction, $BGL_n(\mathbb{C})$ and $BGL_n(\mathbb{R})$ are equivalent to the Grassmanians BU_n and BO_n . We recall that the representing spaces for complex and real topological K -theory, namely $BU \times \mathbb{Z}$ and $BO \times \mathbb{Z}$, are the group completions of the topological monoids $\coprod BU_n$ and $\coprod BO_n$, respectively. Therefore,

$$\mathbf{K}^{top}(\mathbb{C}) \simeq ku \quad \text{and} \quad \mathbf{K}^{top}(\mathbb{R}) \simeq ko,$$

where \mathbf{K}^{top} is algebraic K -theory for which the topology on the ring is taken into account when forming the bar construction.

We note that ku_G and ko_G are represented by the G -spaces which are the group completions of the monoids of equivariant bundles corresponding to split extensions

$$1 \rightarrow U_n \rightarrow U_n \times G \rightarrow G \rightarrow 1 \tag{11}$$

and

$$1 \rightarrow O_n \rightarrow O_n \times G \rightarrow G \rightarrow 1, \tag{12}$$

respectively.

Consider the topological rings \mathbb{C} and \mathbb{R} with trivial G -action for any finite group G . Then, by definition, $\mathbf{K}_G(\mathbb{C})$ and $\mathbf{K}_G(\mathbb{R})$ are genuine Ω - G -spectra with zeroth spaces given by the group completions of $\coprod B\mathcal{C}at(\tilde{G}, GL_n(\mathbb{C}))$, and $\coprod B\mathcal{C}at(\tilde{G}, GL_n(\mathbb{R}))$, respectively, where the topology of $GL_n(\mathbb{C})$, and $GL_n(\mathbb{R})$ respectively, is taken into account when forming the classifying space. By Theorem 5.11, these are the monoids of classifying spaces of (G, U_n) -bundles, and (G, O_n) -bundles, respectively, under Whitney sum. Note that here it was crucial that in the hypotheses of Theorem 5.11, even though the group of equivariance G has to be discrete or finite, the structure group of the bundle is allowed to be compact Lie. Therefore, we obtain the following theorem, where \mathbf{K}_G^{top} is G -equivariant algebraic K -theory for which the topology of the ring is taken into account.

Theorem 7.1. *Consider the topological rings \mathbb{C} and \mathbb{R} with trivial G -action for any finite group G . We have equivalences of connective Ω - G -spectra*

$$\mathbf{K}_G^{top}(\mathbb{C}) \simeq ku_G \quad \text{and} \quad \mathbf{K}_G^{top}(\mathbb{R}) \simeq ko_G.$$

In the definition of KR , the bundles corresponding to split exact sequences (11) are replaced by equivariant $(C_2, U_n \times C_2)$ -bundles corresponding to split exact sequences

$$1 \rightarrow U_n \rightarrow U_n \times C_2 \rightarrow C_2 \rightarrow 1, \tag{13}$$

where the cyclic group of order 2, C_2 , acts on U_n by complex conjugation.

Atiyah shows a “Real” version of Bott periodicity, which gives that the representing space for KR has deloopings with respect to C_2 -representations, and thus KR represents a genuine Ω - C_2 -spectrum. Of course, [Ati66] does not mention spectra and instead states the result in terms of a periodic $RO(C_2)$ -graded cohomology theory.

The zeroth space of the connective spectrum kr is the group completion of the topological C_2 -monoid of $(C_2, U_n \rtimes C_2)$ -bundles, which by Theorem 5.11 and because the equivalence of topological groups $GL(\mathbb{C}) \simeq U_n$ is C_2 -equivariant, is equivalent to $\coprod B\mathcal{C}at(\tilde{G}, GL_n(\mathbb{C}))$. Therefore, we get the following theorem.

Theorem 7.2. *Let \mathbb{C} be the topological ring of complex numbers with conjugation action by C_2 . Then there is an equivalence of connective Ω - C_2 -spectra*

$$\mathbf{K}_{C_2}^{top}(\mathbb{C}) \simeq kr.$$

8 Definition of equivariant A -theory

8.1 Waldhausen G -categories and their subtleties

A Waldhausen G -category is a Waldhausen category \mathcal{C} , with G -action that respects the Waldhausen structure. Precisely, it is a functor $G \rightarrow \text{Wald}\mathcal{C}at$ from G to the category of Waldhausen categories and exact functors.

The fact that for each $g \in G$ the functor $g \cdot : \mathcal{C} \rightarrow \mathcal{C}$ is exact means that it preserves the zero object, cofibrations, weak equivalences and pushouts along cofibrations. Actually, since the functor $g \cdot$ is both left and right adjoint to the functor $g^{-1} \cdot$, it automatically preserves pushouts. However, we emphasize that $g \cdot$ preserves the zero object and pushouts only *up to unique isomorphism*, and not on the nose. This makes the equivariant theory subtle, as we go on to explain.

Observation 8.1. We make the crucial observation that the fixed point category \mathcal{C}^H is *not* a Waldhausen category. The point is precisely that since pushouts are preserved by the G -action only up to isomorphism, the category \mathcal{C}^H is not closed under pushouts; there is no choice of pushouts that makes this possible. However, applying the functor $\mathcal{C}at(\tilde{G}, -)$ to \mathcal{C} rectifies this, and the homotopy fixed point category \mathcal{C}^{hH} is a Waldhausen category. The author thanks Emanuele Dotto and Daniel Schäppi for illuminating conversations on the topic of Waldhausen G -categories which led to the clarification of this subtlety.

Note that if \mathcal{C} is a Waldhausen category, then one can naively define a Waldhausen structure on $\mathcal{C}at(\tilde{G}, \mathcal{C})$ by defining the cofibrations and weak equivalences pointwise. More precisely, one can specify that for $F_1, F_2 \in \mathcal{C}at(\tilde{G}, \mathcal{C})$,

$$F_1 \xrightarrow{\eta} F_2$$

is a cofibration or a weak equivalence if for every $g \in \tilde{G}$, the map $F_1(g) \rightarrow F_2(g)$ is a cofibration or a weak equivalence, respectively, in \mathcal{C} . The category $\mathcal{C}at(\tilde{G}, \mathcal{C})$ is clearly a Waldhausen

category with zero object defined as the functor with constant value 0 for any chosen zero object 0 of \mathcal{C} and pushouts defined pointwise, but the fixed point subcategories \mathcal{C}^H are not closed under this zero object and these pushouts. However, we show that there is a choice of zero object and pushouts in $\mathcal{Cat}(\tilde{G}, \mathcal{C})$, which do lie in the fixed point subcategories.

Lemma 8.2. *There is a zero object in $\mathcal{Cat}(\tilde{G}, \mathcal{C})$, which is G -fixed.*

Proof. Consider the functor $Z : * \rightarrow \mathcal{C}$ from the one object category $*$ to \mathcal{C} , which picks out the zero object 0 of \mathcal{C} . Note that this functor is not equivariant since $0 \neq g \cdot 0$, but for every g we have a unique isomorphism $\theta_g : 0 \xrightarrow{\cong} g \cdot 0$. Since these isomorphisms are unique, it must be that the isomorphisms $0 \xrightarrow{\theta_g} g \cdot 0 \xrightarrow{g\theta_h} (gh) \cdot 0$ and $0 \xrightarrow{\theta_{gh}} (gh) \cdot 0$ coincide, and therefore Z is pseudo equivariant.

Note that the categories $\mathcal{Cat}(\tilde{G}, *)$ and $*$ are trivially G -isomorphic. As shown in Proposition 3.3, since Z is pseudo equivariant, there is an induced on the nose equivariant functor $* \cong \mathcal{Cat}(\tilde{G}, *) \rightarrow \mathcal{Cat}(\tilde{G}, \mathcal{C})$, which sends the one object of $*$ to the functor $F_0 \in \mathcal{Cat}(\tilde{G}, \mathcal{C})$ defined on objects by $F_0(g) = g \cdot 0$, and defined on the unique morphism from g to h by composing the unique isomorphisms $0 \cong g \cdot 0$ and $0 \cong h \cdot 0$ to get an isomorphism $g \cdot 0 \xrightarrow{\cong} h \cdot 0$ in \mathcal{C} . Since the functor $* \rightarrow \mathcal{Cat}(\tilde{G}, \mathcal{C})$ with value F_0 is equivariant by Proposition 3.3, the object F_0 of $\mathcal{Cat}(\tilde{G}, \mathcal{C})$ lies in the G -fixed point subcategory.

This is indeed a zero object in $\mathcal{Cat}(\tilde{G}, \mathcal{C})$: for any $g \in G$, there is a map from $F_0(g)$ to $F(g)$ defined as the composite of the unique isomorphism $g \cdot 0 \cong 0$ and the unique map $0 \rightarrow F(g)$, and it is not hard to see that the uniqueness of these maps forces the composite to also be unique. So there is a unique map $F_0 \rightarrow F$ and similarly, there is a unique map $F \rightarrow F_0$ for any $F \in \mathcal{Cat}(\tilde{G}, \mathcal{C})$.

□

Lemma 8.3. *There exist pushouts along cofibrations in $\mathcal{Cat}(\tilde{G}, \mathcal{C})$, so that pushouts of H -fixed diagrams are H -fixed.*

Proof. The same argument as in the previous proof applies: if one considers the category of pushout diagrams along cofibrations and a functor, which assigns to each pushout diagram a choice of pushout, this functor is not equivariant, but the unique isomorphisms $P \cong g \cdot P$ that exist for any pushout P in \mathcal{C} and any $g \in G$, ensure that the functor is pseudo equivariant and Proposition 3.3 applies to give an on the nose equivariant functor after applying $\mathcal{Cat}(\tilde{G}, -)$.

It is not hard to see that $\mathcal{Cat}(\tilde{G}, -)$ commutes with taking the category of pushout diagrams along cofibrations and we get an on the nose equivariant functor from the category of pushout diagrams along cofibrations in $\mathcal{Cat}(\tilde{G}, \mathcal{C})$ to $\mathcal{Cat}(\tilde{G}, \mathcal{C})$, which assigns to each diagram a pushout such that for an H -fixed diagram the pushout is now also H -fixed. □

From the construction of the corresponding equivariant functor from a pseudo equivariant functor in the proof of Proposition 3.3, we get an explicit description for the pushouts in

$\mathcal{C}at(\tilde{G}, \mathcal{C})$. For a diagram

$$\begin{array}{ccc} F_1 & \xrightarrow{\quad} & F_2 \\ \downarrow & & \\ F_3 & & \end{array}$$

in $\mathcal{C}at(\tilde{G}, \mathcal{C})$, the pushout $\mathbf{P}: \tilde{G} \rightarrow \mathcal{C}$ is defined on objects by

$$\mathbf{P}(g) = g \cdot (g^{-1}F_3(g) \coprod_{g^{-1}F_1(g)} g^{-1}F_2(g)).$$

If the pushout diagram is G -fixed, then the pushout \mathbf{P} is defined by $\mathbf{P}(e) = P$, where P is a pushout of the above diagram evaluated at e , and $\mathbf{P}(g) = g \cdot P$. On morphisms, $\mathbf{P}(g, g')$ is the composite of the unique isomorphisms $P \cong g \cdot P$ and $P \cong g' \cdot P$.

Theorem 8.4. *Let \mathcal{C} be a G -equivariant Waldhausen category, and let H be a subgroup of G . Then \mathcal{C}^{hH} is a Waldhausen category with cofibrations and weak equivalences the H -fixed cofibrations and weak equivalences in $\mathcal{C}at(\tilde{G}, \mathcal{C})$.*

Proof. Note that composition of H -fixed maps is H -fixed, thus the classes of cofibrations and weak equivalences in $\mathcal{C}at(\tilde{G}, \mathcal{C})^H$ are closed under composition, and an H -fixed isomorphism is in particular a H -fixed cofibration and weak equivalence.

By Lemma 8.2, there is a zero object F_0 in $\mathcal{C}at(\tilde{G}, \mathcal{C})^H$. Moreover, for any functor F in $\mathcal{C}at(\tilde{G}, \mathcal{C})$, each map $F_0(g) \rightarrow F(g)$ is a cofibration since it is the composite of $g \cdot 0 \cong 0$ and the unique map $0 \rightarrow F(g)$, which are both cofibrations. Thus the map $F_0 \rightarrow F$ is by definition a cofibration.

By Lemma 8.3, for a pushout diagram along a cofibration in $\mathcal{C}at(\tilde{G}, \mathcal{C})^H$, there exists a pushout in this fixed point subcategory. The glueing axiom for weak equivalences is inherited from \mathcal{C} . \square

The G -action on $\mathcal{C}at(\tilde{G}, \mathcal{C})$ induces a G -action on the simplicial Waldhausen category $S_\bullet \mathcal{C}at(\tilde{G}, \mathcal{C})$. It is not hard to see that the S_\bullet -construction commutes with fixed points.

Lemma 8.5. *Let \mathcal{C} be a G -equivariant Waldhausen category and let H be a subgroup of G . Then we have an isomorphism of simplicial Waldhausen categories $(S_\bullet \mathcal{C}at(\tilde{G}, \mathcal{C}))^H \cong S_\bullet(\mathcal{C}at(\tilde{G}, \mathcal{C})^H)$.*

8.2 Equivariant K -theory of a Waldhausen category

We define the equivariant algebraic K -theory space for a Waldhausen G -category \mathcal{C} in the same spirit of the definition for symmetric monoidal categories, namely by replacing the category \mathcal{C} with $\mathcal{C}at(\tilde{G}, \mathcal{C})$ and applying the nonequivariant construction.

Definition 8.6. The equivariant algebraic K -theory space of a Waldhausen G -category \mathcal{C} is

$$K_G(\mathcal{C}) = \Omega |wS_\bullet \mathcal{C}at(\tilde{G}, \mathcal{C})|.$$

Nonequivariantly, it is an observation of Waldhausen that it is enough to apply the S_\bullet -construction once and then use an infinite loop space machine with the group completion property to obtain the K -theory Ω -spectrum of a Waldhausen category. The argument Waldhausen alludes to is written out explicitly in [Mal]. The equivariant generalization of this observation that we owe to discussions with C. Malkiewich and D. Schäppi is that the K -theory space we define for Waldhausen G -categories can be delooped by an equivariant infinite loop space machine. Work in progress with E. Dotto is aimed at showing that the deloopings of the equivariant A -theory space with respect to representations can be obtained by an equivariant version of the S_\bullet -construction, which uses iterations of the S_\bullet -construction on representations. The proof that the resulting spectrum is a genuine Ω - G -spectrum would ultimately follow from an equivariant version of the additivity theorem and work of Dotto and Moi on homotopy limits of G -diagrams [DM]. In forthcoming work with E. Dotto we aim to prove an equivariant additivity theorem and produce the deloopings with respect to representations using an equivariant version of the S_\bullet -construction.

We emphasize that by Observation 8.1, unless we apply the functor $\mathcal{C}at(\tilde{G}, -)$ to our Waldhausen category first, it we could not apply S_\bullet to the fixed point category \mathcal{C}^H , and asking for a commutation as in Lemma 8.5 would be meaningless. However, with the given definition, by Lemma 8.5, $K_G(\mathcal{C})^H \simeq K(\mathcal{C}^{hH})$.

Generalizing the previous definitions, the equivariant algebraic K -groups of a Waldhausen G -category are

$$K_i^H(\mathcal{C}) = \pi_i^H(K_G(\mathcal{C})).$$

It is not hard to see that this agrees with Kuku's definition of equivariant algebraic K -groups of a Waldhausen category \mathcal{C} with *trivial* G -action given in [Kuk06]. He defines

$$K_i^G(X, \mathcal{C}) = K_i(\mathcal{C}at(\underline{X}, \mathcal{C})),$$

where X is a G -set and \underline{X} is its translation category.

Lemma 8.7. (*Agreement with Kuku's definition*) *Let \mathcal{C} be a Waldhausen category with trivial G -action, and let H be a subgroup of G . Then*

$$K_i^H(\mathcal{C}) \cong K_i^G(\underline{G/H}, \mathcal{C}).$$

Proof. We claim that we have an equivalence

$$K_G(\mathcal{C})^H \simeq \Omega |S_\bullet \text{ iso } \mathcal{C}at(\underline{G/H}, \mathcal{C})|.$$

Note that since Ω , S_\bullet , geometric realization and restricting to isomorphisms commute with fixed points, the left hand side is $\Omega |S_\bullet \text{ iso } \mathcal{C}at(\tilde{G}, \mathcal{C})^H|$, so it is enough to show we have an equivalence of Waldhausen categories

$$\mathcal{C}at(\tilde{G}, \mathcal{C})^H \simeq \mathcal{C}at(\underline{G/H}, \mathcal{C}).$$

Since $\underline{H/H}$ is a full subcategory of $\underline{G/H}$, and any object in $\underline{G/H}$ is isomorphic to H , the categories $\underline{H/H}$ and $\underline{G/H}$ are equivalent, and thus we have an equivalence of categories $\mathcal{C}at(\underline{G/H}, \mathcal{C}) \simeq \mathcal{C}at(\underline{H/H}, \mathcal{C})$. But $\underline{H/H}$ can be identified with H , so we have

$\mathcal{C}at(\underline{G/H}, \mathcal{C}) \simeq \mathcal{C}at(H, \mathcal{C})$. Also, we have shown that $\mathcal{C}at(\tilde{G}, \mathcal{C})^H \simeq \mathcal{C}at(H, \mathcal{C})$ since the action on \mathcal{C} is trivial, and the claim follows.

The conclusion of the lemma follows immediately since

$$K_i^H(\mathcal{C}) \cong \pi_i(K_G(\mathcal{C})^H) \cong \pi_i(\Omega | S_{\bullet} \text{ iso } \mathcal{C}at(\underline{G/H}, \mathcal{C})) = K_i(\mathcal{C}at(\underline{G/H}, \mathcal{C})) = K_i^G(\underline{G/H}, \mathcal{C}).$$

□

8.3 Equivariant A -theory definition

Nonequivariantly, A -theory was defined in Waldhausen's seminal paper [Wal85] as the algebraic K -theory of the Waldhausen category $R(X)$ of retractive spaces over X (i.e., spaces Y with maps $X \xrightarrow{i} Y \xrightarrow{r} X$ such that $r \circ i = \text{id}$) which are finite relative to X . Morphisms in $R(X)$ are continuous maps $Y \rightarrow Y'$ making the obvious diagram commute.

Let X be a G -space. We show that the Waldhausen category $R(X)$ of retractive spaces over X , i.e., spaces Y (*not G -spaces!*) naturally inherits a G -action

$$g(X \xrightarrow{i} Y \xrightarrow{r} X) = X \xrightarrow{g^{-1}} X \xrightarrow{i} Y \xrightarrow{r} X \xrightarrow{g} X.$$

For a morphism f , namely the data of a commuting diagram

$$\begin{array}{ccc} & Y & \\ & \nearrow i & \searrow r \\ X & & X \\ & \searrow i' & \nearrow r' \\ & Y' & \end{array}$$

we define $(gf)(y) = f(y)$ for all $y \in Y$. This is indeed a morphism

$$g(X \xrightarrow{i} Y \xrightarrow{r} X) \longrightarrow g(X \xrightarrow{i'} Y' \xrightarrow{r'} X),$$

since the diagram

$$\begin{array}{ccc} & Y & \\ & \nearrow g^{-1} \circ i & \searrow r \circ g \\ X & & X \\ & \searrow g^{-1} \circ i' & \nearrow r' \circ g \\ & Y' & \end{array}$$

then also commutes.

We define equivariant A_G -theory as the equivariant algebraic K -theory of the Waldhausen G -category $R(X)$.

Definition 8.8. Let X be a G -space. We define the equivariant A_G -theory space of X as

$$A_G(X) = \Omega|S_\bullet \mathcal{C}at(\tilde{G}, \text{iso } R(X))|,$$

where $R(X)$ is the category of (nonequivariant) retractive spaces over X with the G -action defined above.

The following proposition shows that the fixed points of the equivariant A_G -theory of a G -space X are the nonequivariant A -theory of the category of retractive G -spaces and equivariant maps between these.

Proposition 8.9. *The homotopy fixed point category $R(X)^{hG}$ is the category of retractive G -spaces over X with equivariant inclusion and retraction maps, and equivariant maps between these.*

Proof. The homotopy fixed points $R(X)^{hG} = \mathcal{C}at(\tilde{G}, R(X))^G$ are, by Proposition 2.11, spaces Y together with isomorphisms $f(g): Y \xrightarrow{\cong} Y$ for all g making the following diagram commute:

$$\begin{array}{ccccc} X & \xrightarrow{i} & Y & \xrightarrow{r} & X \\ \downarrow g^{-1} & & \downarrow f(g) & & \uparrow g \\ X & \xrightarrow{i} & Y & \xrightarrow{r} & X \end{array}$$

We also have that $f(e) = \text{id}$ and the cocycle condition $f(gh) = f(g)gf(h)$ is satisfied. We can define an action of Y by $g \cdot y = f(g)(y)$. This is indeed an action since $f(e) = \text{id}$ and

$$\begin{aligned} (gh) \cdot y &= f(gh)(y) \\ &= f(g)(gf(h)(y)) \\ &= f(g)(f(h)(y)) \\ &= g \cdot (h \cdot y). \end{aligned}$$

This action on Y then makes i and r equivariant maps by the commutativity of the above diagram. Thus the category of homotopy fixed points $R(X)^{hG}$ can be identified with the category with objects $X \xrightarrow{i} Y \xrightarrow{r} X$, where Y is a G -space and i and r are G -maps. \square

9 Transfer maps and relationship to spectral Mackey functor approach

A different approach to defining equivariant K -theory and equivariant A -theory is given by C. Barwick, who defines a G -spectrum from a G -Waldhausen category in the ∞ -categorical

setting in [Bar] using spectral Mackey functors. These are a model of G -spectra as diagrams of nonequivariant spectra inspired by [GMb], where Guillou and May describe the category of G -spectra as spectrally enriched functors on a spectrally enriched Burnside category.

A. M. Bohmann and A. Osorno have developed input to the Guillou-May theorem in the form of categorical Mackey functors of permutative categories in [BO], and more recently, of Waldhausen categories, and they are using the Guillou-May theorem directly to produce genuine G -spectra from these.

We describe the intuition of how one could recover our construction of equivariant K -theory or equivariant A -theory, as one of these “spectral Mackey functors.” We thank C. Barwick for all the discussions over the years about how the approaches relate.

For a G -ring, the “categorical Mackey functor” of symmetric monoidal categories would have values

$$G/H \mapsto \mathcal{P}(R)^{hH},$$

namely the homotopy fixed points of the category of finitely generated projective R -modules with the action defined in section 4.3, which by Corollary 4.11 is equivalent to $\mathcal{P}(R_H[H])$, the category of finitely generated projective R -modules over the twisted group ring $R_H[H]$ when $|H|^{-1} \in R$.

For a G -space X , the values would be given by

$$G/H \mapsto R(X)^{hH},$$

the homotopy fixed point category of the category of retractive spaces over X with the action defined in section 8.3, which by Proposition 8.9 is the category of retractive H -spaces over X with equivariant inclusion and retraction maps.

More generally, if \mathcal{C} is a Waldhausen G -category, the “categorical Mackey functor” of Waldhausen categories would have values

$$G/H \mapsto \mathcal{C}^{hH},$$

since H -fixed points of our equivariant K -theory construction are given by the nonequivariant K -theory of the homotopy fixed point category \mathcal{C}^{hH} .

What is essential in order for these assignments to potentially define these “categorical Mackey functors” is that for $H \subseteq K \subseteq G$, there are not only restriction maps $\mathcal{C}^{hK} \rightarrow \mathcal{C}^{hH}$, but also wrong-way transfer maps $\mathcal{C}^{hH} \rightarrow \mathcal{C}^{hK}$, suitably related. We show that these maps exist. For simplicity, we construct the transfer from $\mathcal{C}^{hH} \rightarrow \mathcal{C}^{hG}$; this is easy to generalize to other subgroups.

Proposition 9.1. *Let \mathcal{C} be a G -category with sum \oplus that is preserved by the G -action up to coherent isomorphism (for example, a symmetric monoidal G -category or a Waldhausen G -category with $\vee = \oplus$). Then there are transfer maps*

$$\mathcal{C}^{hH} \rightarrow \mathcal{C}^{hG}.$$

We note not only that there are no transfer maps in general between the fixed point categories \mathcal{C}^H and \mathcal{C}^G , but we have seen that if \mathcal{C} is symmetric monoidal and the G -action does not preserve \oplus strictly, or if \mathcal{C} is Waldhausen, then the fixed point categories \mathcal{C}^H are not symmetric monoidal, or Waldhausen, respectively.

Proof. This will be yet another application of pseudo equivariance. Define

$$\mathcal{C}at(G/H, \mathcal{C}) \cong \prod_{G/H} \mathcal{C} \longrightarrow \mathcal{C}$$

by

$$(c_1, \dots, c_j) \mapsto c_1 \oplus \dots \oplus c_j.$$

This is not necessarily equivariant even when the G -action preserves \oplus strictly since

$$gc_1 \oplus \dots \oplus gc_j \neq gc_{\sigma_g^{-1}(1)} \oplus \dots \oplus gc_{\sigma_g^{-1}(j)},$$

where σ_g here is the permutation in Σ_j , where $j = |G/H|$, which gives the action of g on G/H .

However,

$$gc_1 \oplus \dots \oplus gc_j \cong gc_{\sigma_g^{-1}(1)} \oplus \dots \oplus gc_{\sigma_g^{-1}(j)},$$

and this is another instance of a pseudo equivariant functor, and therefore, by Proposition 3.3 it induces an on the nose equivariant functor

$$\mathcal{C}at(\tilde{G}, \mathcal{C}at(G/H, \mathcal{C})) \cong \mathcal{C}at(G/H, \mathcal{C}at(\tilde{G}, \mathcal{C})) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{C}).$$

By taking G -fixed points of the G -map $\mathcal{C}at(G/H, \mathcal{C}at(\tilde{G}, \mathcal{C})) \rightarrow \mathcal{C}at(\tilde{G}, \mathcal{C})$, we get a transfer map

$$\mathcal{C}^{hH} \rightarrow \mathcal{C}^{hG}.$$

□

References

- [AG60] Maurice Auslander and Oscar Goldman, *The Brauer group of a commutative ring*, Trans. Amer. Math. Soc. **97** (1960), 367–409. MR 0121392 (22 #12130)
- [Ati66] M. F. Atiyah, *K-theory and reality*, Quart. J. Math. Oxford Ser. (2) **17** (1966), 367–386. MR 0206940 (34 #6756)
- [Bar] C. Barwick, *Spectral Mackey functors and equivariant algebraic K-theory (i)*, arXiv:1404.0108.
- [Bil12] E. Biland, *An application of equivariant morita theory*.
- [BO] Anna Marie Bohmann and Angélica M. Osorno, *Constructing equivariant spectra via categorical Mackey functors*, arXiv:1405.6126v2.

- [Car] G. Carlsson, *A geometric reformulation of the representational assembly conjecture for the k -theory of fields with pro- l absolute galois group*, arXiv:1309.5649.
- [Car84] Gunnar Carlsson, *Equivariant stable homotopy and Segal's Burnside ring conjecture*, *Ann. of Math. (2)* **120** (1984), no. 2, 189–224. MR 763905 (86f:57036)
- [Car08] ———, *Derived completions in stable homotopy theory*, *J. Pure Appl. Algebra* **212** (2008), no. 3, 550–577. MR 2365333 (2008m:55018)
- [Car11] Gunnar E. Carlsson, *Derived representation theory and the algebraic K -theory of fields*, *J. Topol.* **4** (2011), no. 3, 543–572. MR 2832568
- [Cha13] Zihua Chang, *Descent theory with an application to Infinite Dimensional Lie Algebras*, Notes on lectures by Arturo Pianzola (2013).
- [CR] G. Carlsson and J. Ray, *Galois descent for completed algebraic k -theory*, arXiv:1309.6045.
- [Del77] P. Deligne, *Cohomologie étale*, Lecture Notes in Mathematics, Vol. 569, Springer-Verlag, Berlin, 1977, Séminaire de Géométrie Algébrique du Bois-Marie SGA 41øer2, Avec la collaboration de J. F. Boutot, A. Grothendieck, L. Illusie et J. L. Verdier. MR 0463174 (57 #3132)
- [DM] Emanuele Dotto and Kristian Moi, *Homotopy theory of g -diagrams and equivariant excision*, arXiv:1403.6101v2.
- [Dug05] Daniel Dugger, *An Atiyah-Hirzebruch spectral sequence for KR -theory*, *K-Theory* **35** (2005), no. 3-4, 213–256 (2006). MR 2240234 (2007g:19004)
- [FHM82] Z. Fiedorowicz, H. Hauschild, and J. P. May, *Equivariant algebraic K -theory*, Algebraic K -theory, Part II (Oberwolfach, 1980), Lecture Notes in Math., vol. 967, Springer, Berlin, 1982, pp. 23–80. MR 689388 (84h:57023)
- [GMa] B. J. Guillou and J. P. May, *Permutative G -categories in equivariant infinite loop space theory*, arXiv:1207.3459v2.
- [GMb] Bertrand Guillou and J. P. May, *Models of G -spectra as presheaves of spectra*, arXiv:1110.3571.
- [GMM] B. J. Guillou, J. P. May, and M. Merling, *Categorical models for equivariant classifying spaces*, arXiv:1201.5178v2.
- [GMMO] Bertrand Guillou, J. Peter May, Mona Merling, and Angelica Osorno, *Categorical multiplicative infinite loop space theory*, in preparation.
- [Gui] B. J. Guillou, *Algebraic K -theory*, expository note.
- [HHR] Michael A. Hill, Michael J. Hopkins, and Douglas C. Ravenel, *On the non-existence of elements of Kervaire invariant one*, arXiv:0908.3724.
- [Kaw86] K. Kawakubo, *Induction theorems for equivariant K -theory and J -theory*, *J. Math. Soc. Japan* **38** (1986), 173–198.

- [Kuk06] Aderemi Kuku, *Equivariant higher algebraic K-theory for Waldhausen categories*, *Beiträge Algebra Geom.* **47** (2006), no. 2, 583–601. MR 2310576 (2008c:19002)
- [LMS86] L. G. Lewis, Jr., J. P. May, and M. Steinberger, *Equivariant stable homotopy theory*, *Lecture Notes in Mathematics*, vol. 1213, Springer-Verlag, Berlin, 1986, With contributions by J. E. McClure. MR 866482 (88e:55002)
- [Mal] C. Malkiewich, *Coassembly and the K-theory of finite groups*, arXiv:1503.06504.
- [May72] J. P. May, *The geometry of iterated loop spaces*, Springer-Verlag, Berlin, 1972, *Lectures Notes in Mathematics*, Vol. 271. MR 0420610 (54 #8623b)
- [May74] ———, *E_∞ spaces, group completions, and permutative categories*, *New developments in topology (Proc. Sympos. Algebraic Topology, Oxford, 1972)*, Cambridge Univ. Press, London, 1974, pp. 61–93. *London Math. Soc. Lecture Note Ser.*, No. 11. MR 0339152 (49 #3915)
- [May75] J. Peter May, *Classifying spaces and fibrations*, *Mem. Amer. Math. Soc.* **1** (1975), no. 1, 155, xiii+98. MR 0370579 (51 #6806)
- [ML63] Saunders Mac Lane, *Natural associativity and commutativity*, *Rice Univ. Studies* **49** (1963), no. 4, 28–46. MR 0170925 (30 #1160)
- [MMO] J. P. May, M. Merling, and A. Osorno, *Comparison of equivariant infinite loop space machines*, In preparation.
- [MS76] D. McDuff and G. Segal, *Homology fibrations and the “group-completion” theorem*, *Invent. Math.* **31** (1975/76), no. 3, 279–284. MR 0402733 (53 #6547)
- [MT78] J. P. May and R. Thomason, *The uniqueness of infinite loop space machines*, *Topology* **17** (1978), no. 3, 205–224. MR 508885 (80g:55015)
- [Qui73] Daniel Quillen, *Higher algebraic K-theory. I*, *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, Springer, Berlin, 1973, pp. 85–147. *Lecture Notes in Math.*, Vol. 341. MR 0338129 (49 #2895)
- [Seg68] Graeme Segal, *Equivariant K-theory*, *Inst. Hautes Études Sci. Publ. Math.* (1968), no. 34, 129–151. MR 0234452 (38 #2769)
- [Seg74] ———, *Categories and cohomology theories*, *Topology* **13** (1974), 293–312. MR MR0353298 (50 #5782)
- [Shi89] Kazuhisa Shimakawa, *Infinite loop G-spaces associated to monoidal G-graded categories*, *Publ. Res. Inst. Math. Sci.* **25** (1989), no. 2, 239–262. MR 1003787 (90k:55017)
- [Str72] Ross Street, *Two constructions on lax functors*, *Cahiers Topologie Géom. Différentielle* **13** (1972), 217–264. MR 0347936 (50 #436)
- [Tho83] R. W. Thomason, *The homotopy limit problem*, *Proceedings of the Northwestern Homotopy Theory Conference (Evanston, Ill., 1982) (Providence, R.I.)*, *Contemp. Math.*, vol. 19, Amer. Math. Soc., 1983, pp. 407–419. MR 711065 (84j:18012)
- [Tho85] ———, *Algebraic K-theory and étale cohomology*, *Ann. Sci. École Norm. Sup. (4)* **18** (1985), no. 3, 437–552. MR 826102 (87k:14016)

- [Wal85] Friedhelm Waldhausen, *Algebraic K-theory of spaces*, Algebraic and geometric topology (New Brunswick, N.J., 1983), Lecture Notes in Math., vol. 1126, Springer, Berlin, 1985, pp. 318–419. MR 802796 (86m:18011)
- [Wei13] Charles A. Weibel, *The K-book*, Graduate Studies in Mathematics, vol. 145, American Mathematical Society, Providence, RI, 2013, An introduction to algebraic K -theory. MR 3076731